

Derivation of Euler equations from quantum and classical microscopic dynamics

Amirali Hannani¹ and François Huveneers²

¹ ICERM, Brown University, 121 S Main St, Providence, RI 02903, United States

² Ceremade, UMR-CNRS 7534, Université Paris Dauphine, PSL Research University, 75775 Paris cedex 16, France

September 15, 2022

Abstract

We derive Euler equations from a Hamiltonian microscopic dynamics. The microscopic system is a one-dimensional disordered harmonic chain, and the dynamics is either quantum or classical. This chain is an Anderson insulator with a symmetry protected mode: Thermal fluctuations are frozen while the low modes ensure the transport of elongation, momentum and mechanical energy, that evolve according to Euler equations in an hyperbolic scaling limit. In this paper, we strengthen considerably the results in [3] and [12], where we established a limit in mean starting from a local Gibbs state: We now control the second moment of the fluctuations around the average, yielding a limit in probability, and we enlarge the class of admissible initial states.

1 Introduction and results

The deep understanding of transport properties of materials is of fundamental physical interest. It is also an utmost challenging task from a mathematical perspective, and only some specific cases have been successfully investigated so far. Here we focus on the derivation of *Euler* system of conservation laws in an appropriate scaling limit, the so-called hyperbolic space-time rescaling, where both space and time are rescaled by a common factor $n \rightarrow \infty$.

The usual route to Euler equations rests on some form of *ergodicity*: The replacement of time averages by a spatial averages allows to close the equations of motion for the conserved fields. These fields usually involve at least the number of particles, conveniently replaced by elongation in 1-d systems, momentum and energy. However, while the common belief is that “typical” anharmonic dynamics are ergodic in the large volume limit, it has so far not been possible to establish it at a mathematical level of rigor. Instead, most rigorous results

rely on the introduction of some noise that mimics the effect of anharmonic interactions and ensure ergodicity [18, 7, 14]. In addition, recently, the macroscopic evolution of conserved charges was derived rigorously for a class of 1-d quantum systems in the hyperbolic scaling [11], but the number of charges remains unknown as a consequence of the lack of knowledge on the ergodicity.

In [3] and [12], we have introduced a completely new approach to this question, that we further investigate in this paper. Indeed, we consider an *integrable*, hence strongly non-ergodic, dynamics and show the validity of Euler equations in an hyperbolic scaling limit. While the integrability and some particular features of our model render possible a full mathematical treatment, our primary objective is not to exploit these specific aspects. Instead, we want to convey the message that ergodicity needs not to be the crucial ingredient for the emergence of the hydrodynamic behavior described by Euler equations. In our work, the macroscopic description is derived from the fact that mechanical and thermal modes, corresponding respectively to low and high modes, evolve on strikingly different time scales. We live to further investigations the possibility of implementing our strategy in the presence of anharmonic interactions.

Model. We study the disordered harmonic chain described by the Hamiltonian

$$H = \sum_{x=1}^n \frac{p_x^2}{2m_x} + g \frac{(q_{x+1} - q_x)^2}{2}$$

where n is the number of particles, with canonical coordinates (p_x, q_x) (boundary conditions will be specified later on and play no crucial role). We will further fix energy units so that $g = 1$. The equations of motion are given by

$$m_x \frac{dq_x}{dt} = p_x, \quad \frac{dp_x}{dt} = (\Delta q)_x$$

where Δ denotes the lattice Laplacian. We consider both classical and quantum dynamics, and these equations of motion should thus be interpreted either as Hamilton equations, or as Heisenberg equations.

Crucially, the masses m_x are assumed to be random and independent. This guarantees that the spectrum of $M^{-1}\Delta$ is *Anderson localized*, with a localization length diverging as one approaches the ground state, where M is the diagonal matrix with $M_{x,x} = m_x$, cf. [8, 22, 10, 2]. The ground state, explicitly given by $\psi(x) = 1/\sqrt{N}$ for $x = 1, \dots, n$, is an example of a symmetry protected mode, here issuing from the conservations of momentum. Anderson localization guarantees that thermal fluctuations are frozen on all time scales, so that the evolution of the mechanical energy is entirely slaved to the evolution of momentum and elongation (cf. below).

The system has three obvious locally conserved quantities (that may be destroyed at the boundary depending on the choice of boundary conditions): energy, momentum and elongation:

$$H_n = \sum_{x=1}^n \frac{1}{2} \left(\frac{p_x^2}{m_x} + r_x^2 \right) =: \sum_{x=1}^n e_x, \quad P_n = \sum_{x=1}^n p_x, \quad R_n = \sum_{x=1}^n r_x, \quad (1.1)$$

with

$$r_x = q_{x+1} - q_x.$$

Since the dynamics is integrable, there exists actually a full set of conserved quantities. As said though, we are not primarily interested in studying peculiarities of integrable systems; these other quantities will actually remain frozen in the macroscopic limit for the type of initial states that we consider, cf. [3, 12].

Our choice of working with a one-dimensional lattice is mainly motivated by technical considerations: No detailed description of the spectrum of $M^{-1}\Delta$ is available in higher dimensions, and it is no longer possible to use the convenient elongation variables r_x instead of q_x if $d > 1$. We expect however that our approach extends to higher dimensions, though one may not be able to carry over a full mathematical proof. In particular, in dimension 3 at moderate values of the disorder strength, high modes presumably become diffusive instead of staying localized. Since diffusive motion is too slow to be observed in the hyperbolic scaling limit, thermal fluctuations would appear frozen as well and we reckon that the hydrodynamic behavior will still be determined by the evolution of mechanical modes. The same phenomenology is at play for clean harmonic chains perturbed by a noise [7, 14, 15].

Initial states. We assume that the system is initially in a local equilibrium state, that is presumably the most common state of matter in usual circumstances. In our case, it is characterized by three macroscopic (smooth) profiles: the inverse temperature profile $\beta(y)$, the momentum profile $\bar{p}(y)$ and the elongation profile $\bar{r}(y)$, with $y \in [0, 1]$. This means that, at the microscopic site x , the local inverse temperature is $\beta_x := \beta(x/n)$, the average momentum is $\bar{p}_x := \bar{p}(x/n)$ and the average elongation is $\bar{r}_x := \bar{r}(x/n)$.

For a classical dynamics, this state is given by a local Gibbs state, see (2.6) in Section 2.1. If the system is quantum, the precise definition is provided by Assumptions 1-3 in Section 3.1, that involve some clustering conditions. We notice that a local Gibbs state analogous to the classical local Gibbs state satisfies these conditions, see the definition (3.14) and Theorem 3.2 in Section 3.1. However, our definition is more general and may in particular be satisfied by pure states. Actually, the eigenstate thermalization hypothesis and the notion of quantum typicality [9][21][20] suggest that, if the system is prepared through some thermal process, the state of the system will share many similarities with a thermal state, and may satisfy the required conditions.

Outline of the results. On the macroscopic level, the evolution of the three conserved fields H_n , P_n and R_n is described by the following system of conservation laws, known as Euler equations:

$$\partial_t \mathbf{r}(y, t) = \frac{1}{\bar{m}} \partial_y \mathbf{p}(y, t), \quad \partial_t \mathbf{p}(y, t) = \partial_y \mathbf{r}(y, t), \quad \partial_t \mathbf{e}(y, t) = \frac{1}{\bar{m}} \partial_y (\mathbf{r}(y, t) \mathbf{p}(y, t)) \quad (1.2)$$

where \bar{m} denotes the average mass (one may not immediately recognize Euler equations, because they are here written in Lagrangian coordinates, through the use of the elongation variable r_x). More precisely, we have established in [3, 12] that, given a smooth test function f on $[0, 1]$, the following limit on average holds:

$$\frac{1}{n} \sum_{x=1}^n f\left(\frac{x}{n}\right) \langle z_x(nt) \rangle_\rho \rightarrow \int_0^1 f(y) \mathbf{z}(y, t) dy \quad (1.3)$$

as $n \rightarrow \infty$ almost surely w.r.t. the distribution of the masses, uniformly for any time t smaller than some arbitrary time horizon T , where z stands for r , p or e , where \mathfrak{z} stands for \mathfrak{r} , \mathfrak{p} or \mathfrak{e} respectively, and where $\langle \cdot \rangle_\rho$ denotes the averaging w.r.t. a local Gibbs state.

It is interesting to observe that the solution $\mathfrak{e}(y, t)$ to the above Euler equations reads

$$\mathfrak{e}(y, t) = \frac{\mathfrak{p}^2(y, t)}{2\bar{m}} + \frac{\mathfrak{r}^2(y, t)}{2} + C(y)$$

where $C(y)$ is a constant that depends on the initial conditions and is simply equal to the local temperature in the classical set-up. This means that the macroscopic evolution of the energy is *slaved* to the macroscopic evolution of the elongation and momentum, with a functional dependence that is the same as on the microscopic level. We expect this to be more general: If $O_x = f(r_x, p_x)$ is a microscopic local observable, its macroscopic description would be simply given by $f(\mathfrak{r}(y, t), \mathfrak{p}(y, t))$ (we have here taken a local observable depending on a single site but this restriction can be dropped). We leave this to further investigations.

In this paper, we extend our previous results by proving a convergence in quadratic variation and allowing for a broader class of initial states in the quantum set-up. In Theorem 2.1 and Theorem 3.1 below, devoted respectively to the classical and quantum dynamics, we establish that

$$\begin{aligned} \left\langle \left(\frac{1}{n} \sum_{x=1}^n f\left(\frac{x}{n}\right) r_x(nt) - \int_0^1 f(y) \mathfrak{r}(y, t) dy \right)^2 \right\rangle_\rho &\rightarrow 0, \\ \left\langle \left(\frac{1}{n} \sum_{x=1}^n f\left(\frac{x}{n}\right) p_x(nt) - \int_0^1 f(y) \mathfrak{p}(y, t) dy \right)^2 \right\rangle_\rho &\rightarrow 0, \\ \left\langle \left(\frac{1}{n} \sum_{x=1}^n f\left(\frac{x}{n}\right) e_x(nt) - \int_0^1 f(y) \mathfrak{e}(y, t) dy \right)^2 \right\rangle_\rho &\rightarrow 0. \end{aligned}$$

as $n \rightarrow \infty$, almost surely w.r.t. the distribution of the masses, and uniformly for any time t smaller than some arbitrary time horizon T . In addition, $\langle \cdot \rangle_\rho$ refers now to any local equilibrium state as described above.

The convergence in quadratic variation implies a strong concentration around the average behavior and provides concrete support to the claim that ergodic averages suppress fluctuations at the macroscopic scale. It is also a first step towards a more global control over out-of-equilibrium fluctuations. Indeed, in macroscopic fluctuation theories [4], the Euler equation would appear as the minimizer of an action functional and macroscopic fluctuations are expected to be exponentially suppressed as a function of the system size. We hope that the techniques developed in this paper will prove useful in establishing such a global picture.

As they stand, our bounds already yield a convergence in probability of the empirical distributions to their classical limit: For any $\delta > 0$,

$$\mathbb{P}_\rho \left(\left| \frac{1}{n} \sum_{x=1}^n f\left(\frac{x}{n}\right) z_x(nt) - \int_0^1 f(y) \mathfrak{z}(y, t) dy \right| > \delta \right) \rightarrow 0$$

as $n \rightarrow \infty$, with $z = r$, $z = p$ or $z = e$. In the classical set-up, \mathbf{P}_ρ denotes the probability induced by the average $\langle \cdot \rangle_\rho$, and the claim follows from Markov inequality. Indeed, writing

$$O_n = \frac{1}{n} \sum_{x=1}^n f\left(\frac{x}{n}\right) z_x(nt) - \left(\int_0^1 f(y) \mathfrak{z}(y, t) dy \right),$$

we derive

$$\mathbf{P}_\rho(|O_n| > \delta) = \mathbf{P}_\rho(O_n^2 > \delta^2) \leq \frac{\langle O_n^2 \rangle_\rho}{\delta^2}$$

that converges to 0 as $n \rightarrow \infty$.

In the quantum set-up, one first needs to properly define the probability \mathbf{P}_ρ on the left-hand side. Let us define the operators

$$O_n = \frac{1}{n} \sum_{x=1}^n f\left(\frac{x}{n}\right) z_x(nt) - \left(\int_0^1 f(y) \mathfrak{z}(y, t) dy \right) \text{Id}$$

for any $n \geq 1$. Let also P_δ^n be the projector on the spectral subspace spanned by all normalized states f such that $\|O_n^2 f\|_2 \leq \delta^2$. Following the basic postulates of quantum mechanics, we define

$$\mathbf{P}_\rho(|O_n| > \delta) = \langle \text{Id} - P_\delta^n \rangle_\rho$$

and we obtain

$$\langle \text{Id} - P_\delta^n \rangle_\rho \leq \frac{\langle (\text{Id} - P_\delta^n) O_n^2 (\text{Id} - P_\delta^n) \rangle_\rho}{\delta^2} \leq \frac{\langle O_n^2 \rangle_\rho}{\delta^2}$$

that converges here as well to 0 as $n \rightarrow \infty$. To our knowledge, it is the first time that such a limit in probability is shown starting from a fully quantum system. Reminding that ρ may be a pure state, the result shows how the probabilistic quantum description may lead to a purely classical and deterministic behavior in the macroscopic limit.

Remarks on the proofs. The proofs of Theorem 2.1 and Theorem 3.1 rely first on our previous result, the convergence on average (1.3), and on the methods developed in [3, 12]. We refer to these papers for a heuristic description of these methods. Let us here comment on the principal difficulty that needed to be overcome in order to establish the Theorems 2.1 and 3.1. The main technical step was to derive a sufficient decay of correlations for local observables at the hyperbolic time scale, cf. (2.36) for the classical case, and (3.42) for the quantum case. Indeed, let us take the observable p_x as an example. For a finite time t , $\langle p_x(t) p_y(t) \rangle_\rho$ decays sufficiently fast as a function of $|x - y|$, assuming that the initial state has suitable decaying properties, as we do. This decay follows from analyticity and holds independently of localization. However, this argument breaks as soon as we rescale the time t with the length of the system n , since faraway observables may in principle become entangled and correlated. But correlation is suppressed by localization in our case. More specifically, when $|x - y| \geq n^\theta$ for some suitable $0 < \theta < 1$, localization estimates (2.18) in Lemma 2.2 provide the desired decay for the high modes of $p_x(nt)$ at any time-scale, cf. (2.31) and (3.37). We prove that this decay holds and that the contributions from the low modes and from the sites with $|x - y| < n^\theta$ is vanishing.

Plan of the paper. In the rest of this paper, the above claims will be made mathematically rigorous. First, we deal with the classical dynamics in Section 2, where we state and prove Theorem 2.1. Second, Section 3 is devoted to the study of the quantum dynamics; Theorems 3.1 and 3.2 are shown there.

2 Classical case

2.1 Model and Result

We consider a classical disordered chain of n harmonic oscillators in one dimension. The phase space is given by $\{(p, q) \in \mathbb{R}^{2n}\}$, where $p = (p_1, \dots, p_n)$, $q = (q_1, \dots, q_n) \in \mathbb{R}^n$ denote the momentum and position vectors, respectively. We label particles by $x \in \mathbb{I}_n = \{1, \dots, n\}$. Let us define the discrete gradients $\nabla_- : \mathbb{R}^{n-1} \rightarrow \mathbb{R}^n$ and $\nabla_+ : \mathbb{R}^n \rightarrow \mathbb{R}^{n-1}$ such that, for any $v \in \mathbb{R}^{n-1}$, $w \in \mathbb{R}^n$, $x \in \mathbb{I}_n$, and $y \in \mathbb{I}_{n-1}$, we have

$$(\nabla_- v)_x = v_x - v_{x-1}, \quad (\nabla_+ w)_y = w_{y+1} - w_y, \quad (2.1)$$

with the convention $v_n = 0$. Let Δ denote the discrete Laplacian with free boundary conditions: For any $v \in \mathbb{R}^n$ and $x \in \mathbb{I}_n$,

$$\Delta = \nabla_- \nabla_+, \quad (\Delta v)_x = v_{x+1} - 2v_x + v_{x-1}, \quad (2.2)$$

with the conventions $v_{n+1} = v_n$ and $v_0 = v_1$.

The Hamiltonian H_n is defined on \mathbb{R}^{2n} as:

$$H_n(p, q) = \frac{1}{2} \sum_{x=1}^n \left(\frac{p_x^2}{m_x} + (q_{x+1} - q_x)^2 \right), \quad (2.3)$$

where $\{m_x\}_{x=1}^\infty$ are i.i.d random variables defined on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. We denote the expectation w.r.t \mathbb{P} with \mathbb{E} . We let also

$$\bar{m} = \mathbb{E}(m_x).$$

Furthermore, we assume the law of the random variables to be smooth and compactly supported in $[m_-, m_+]$ with $0 < m_- < m_+ < \infty$. These assumptions are the same as in [2][3] and allow to exploit results on Anderson localization. Notice that we assume free boundary conditions $q_0 = q_1$ and $q_n = q_{n+1}$ in (2.3), for the sake of concreteness.

Let us define the elongation variable r as follows: for any $x \in \mathbb{I}_{n-1}$,

$$r_x = (\nabla_+ q)_x. \quad (2.4)$$

Free boundary conditions in terms of elongation variable read $r_0 = r_n = 0$. From now on, we describe our model in terms of these variables. The equations of motion (Hamiltonian dynamics) in this coordinate is as follows:

$$\dot{r}_x = (\nabla_+ M^{-1} p)_x, \quad x \in \mathbb{I}_{n-1}, \quad \dot{p}_x = (\nabla_- r)_x, \quad x \in \mathbb{I}_n, \quad (2.5)$$

where $M = \text{diag}(m_1, \dots, m_n)$ is the diagonal matrix of masses, in the first equation $r_0 = r_n = 0$ is considered, and \dot{a} denotes the time derivative, i.e. $\dot{a} = da/dt$.

Despite the existence of a full set of conserved quantities, we focus on the evolution of the locally conserved quantities H_n , P_n and R_n defined in (1.1) (let us notice that while H_n and P_n are also globally conserved, the conservation of R_n is broken at the boundaries). As initial state of the chain, we take a locally Gibbs state parametrized by the local values of these three quantities. This state is thus parameterized by a temperature profile $\beta \in C^0([0, 1])$ with $0 < \beta_- \leq \beta(y) \leq \beta_+ < \infty$, a momentum profile $\bar{p} \in C^1([0, 1])$ and an elongation profile $\bar{r} \in C^1([0, 1])$ with $\bar{r}(0) = \bar{r}(1) = 0$. Given β, \bar{p}, \bar{r} , as above, the locally Gibbs state is a probability distribution defined on \mathbb{R}^{2n-1} , with the following probability density $\rho_{\beta, \bar{p}, \bar{r}}^n$:

$$\rho_{\beta, \bar{p}, \bar{r}}^n(r, p) = \frac{1}{Z_n} \exp \left(-\frac{1}{2} \sum_{x=1}^n \left[\frac{\beta_x}{m_x} \left(p_x - \bar{p}_x \frac{m_x}{\bar{m}} \right)^2 + \beta_x (r_x - \bar{r}_x)^2 \right] \right), \quad (2.6)$$

where we used the shorthand notation $\beta_x := \beta(\frac{x}{n})$, $\bar{p}_x := \bar{p}(\frac{x}{n})$, $\bar{r}_x := \bar{r}(\frac{x}{n})$ and where Z_n is a normalizing constant. We will drop the subscripts and superscripts from ρ whenever there is no risk of confusion. We denote the expectation w.r.t ρ by $\langle \cdot \rangle_\rho$. For any suitable observable $O(r, p) : \mathbb{R}^{2n-1} \rightarrow \mathbb{R}$, we define

$$\langle O \rangle_\rho := \int_{\mathbb{R}^{2n-1}} O(r, p) \rho_{\beta, \bar{p}, \bar{r}}^n(r, p) dr dp. \quad (2.7)$$

At the macroscopic level, fix the macroscopic time $T > 0$ and let the initial profile of elongation and momentum evolve according to the following system of conservation laws:

$$\partial_t \mathbf{r}(y, t) = \frac{1}{\bar{m}} \partial_y \mathbf{p}(y, t), \quad \partial_t \mathbf{p}(y, t) = \partial_y \mathbf{r}(y, t), \quad \partial_t \mathbf{e}(y, t) = \frac{1}{\bar{m}} \partial_y (\mathbf{r}(y, t) \mathbf{p}(y, t)), \quad (2.8)$$

where $\mathbf{r}, \mathbf{p} \in C^1([0, 1] \times [0, T])$, with the following initial and boundary conditions:

$$\begin{aligned} \mathbf{r}(y, 0) &= \bar{r}(y), \quad \mathbf{p}(y, 0) = \bar{p}(y), \quad \mathbf{e}(y, 0) = \frac{\bar{p}^2(y)}{2\bar{m}} + \frac{\bar{r}^2(y)}{2} + \frac{1}{\beta(y)}, \\ \mathbf{r}(0, t) &= \mathbf{r}(1, t) = 0 \quad \forall t \in [0, T]. \end{aligned} \quad (2.9)$$

The hydrodynamic limit on average (1.3) was proven in [3] and we now improve this result into

Theorem 2.1. *Let $f \in C^0([0, 1])$ be an arbitrary test function. Fix $T > 0$. Initially let the chain to be in the locally Gibbs state $\rho_{\beta, \bar{p}, \bar{r}}^n$ (2.6), corresponding to the profiles β, \bar{p}, \bar{r} satisfying the assumptions stated in the definition of (2.6), and recall $\langle \cdot \rangle_{\rho^n}$ to be the average w.r.t this state (2.7). Denote the solution to the microscopic evolution equation (2.5) by $(r(t), p(t), e(t))$. Moreover, let $\mathbf{r}(y, t), \mathbf{p}(y, t), \mathbf{e}(y, t)$ denote the solution to the macroscopic evolution equation (2.8) with initial and boundary conditions (2.9). Then for any $t \in [0, T]$ we have:*

$$\left\langle \left(\frac{1}{n} \sum_{x=1}^n f\left(\frac{x}{n}\right) r_x(nt) - \int_0^1 f(y) \mathbf{r}(y, t) dy \right)^2 \right\rangle_{\rho^n} \rightarrow 0, \quad (2.10)$$

$$\left\langle \left(\frac{1}{n} \sum_{x=1}^n f\left(\frac{x}{n}\right) p_x(nt) - \int_0^1 f(y) \mathbf{p}(y, t) dy \right)^2 \right\rangle_{\rho^n} \rightarrow 0, \quad (2.11)$$

$$\left\langle \left(\frac{1}{n} \sum_{x=1}^n f\left(\frac{x}{n}\right) e_x(nt) - \int_0^1 f(y) \mathfrak{e}(y, t) dy \right)^2 \right\rangle_{\rho^n} \rightarrow 0, \quad (2.12)$$

as $n \rightarrow \infty$ almost surely w.r.t the distribution of the masses.

Before proceeding, let us briefly mention that the usual mathematical tool to control moments similar to the above ones is the relative entropy method (cf. [23][18]). However, it is not clear that this method could be applied here, and we are able to handle them with a novel approach, i.e. by using localization estimates. We devote the rest of this section to the proof of Theorem 2.1.

2.2 Solution to the equation of motion and localization

The proof of Theorem 2.1 rests on localization of the high eigenmodes of the chain. In this section, we solve the equation of motion (2.5) explicitly and state the localization estimates. Consequently, we can obtain proper decay estimates.

2.2.1 Solution to the equation of motion

We denote the inner product in \mathbb{R}^n by $\langle \cdot, \cdot \rangle_n$, and we drop the subscript whenever it is convenient. Recall the definition of matrices ∇_- , ∇_+ , Δ , and M (2.1), (2.2), (2.5). Taking the time derivative of (2.5) we get:

$$\ddot{r}_x = (\nabla_+ M^{-1} \nabla_- r)_x, \quad 1 \leq x \leq n-1; \quad \ddot{p}_x = (\Delta M^{-1} p)_x, \quad 1 \leq x \leq n, \quad (2.13)$$

where one should recall $r_0 = r_n$ in the first equation, and free boundary conditions ($p_0/m_0 = p_1/m_1$, $p_n/m_n = p_{n+1}/m_{n+1}$) in the second equation.

Notice that $\Delta^\dagger = \Delta$ and $\nabla_+^\dagger = -\nabla_-$. Define

$$A_p := M^{-\frac{1}{2}}(-\Delta)M^{-\frac{1}{2}} \in \mathbb{R}^{n \times n}, \quad A_r := -\nabla_+ M^{-1} \nabla_- \in \mathbb{R}^{(n-1) \times (n-1)}. \quad (2.14)$$

A_p is symmetric, positive semidefinite, and it has a non-degenerate spectrum almost surely (cf. proposition II.1 of [16]). We denote this spectrum by $0 = \omega_0^2 < \omega_1^2 < \dots < \omega_{n-1}^2$, and designate their corresponding eigenvectors by $\{\varphi^k\}_{k=0}^{n-1}$, such that they form an orthonormal basis for \mathbb{R}^n . In particular, $\langle \varphi^k, \varphi^j \rangle = \delta_{k,j}$.

On the other hand, one can check that A_r is a positive symmetric matrix. Furthermore, if for $k \in \mathbb{I}_{n-1}$, we denote $\phi^k := \frac{1}{\omega_k} \nabla_+ M^{-\frac{1}{2}} \varphi^k$, then by using the fact that $\Delta = \nabla_- \nabla_+$, it is straightforward to observe that $\{\phi^k\}_{k=1}^{n-1}$ is a complete set of eigenvectors for A_r with eigenvalues $\omega_1^2 < \omega_2^2 < \dots < \omega_{n-1}^2$ similar to the eigenvalues of A_p , and they form an orthonormal basis for \mathbb{R}^{n-1} i.e.,

$$A_r \phi^k = \omega_k^2 \phi^k, \quad 1 \leq k \leq n-1; \quad \langle \phi^k, \phi^j \rangle_{n-1} = \delta_{k,j}, \quad 1 \leq k, j \leq n.$$

We obtain the normal modes of the chain as follows: define \hat{p}_k for $k \in \mathbb{I}_{n-1}^o = \{0, \dots, n-1\}$, and \hat{r}_k for $k \in \mathbb{I}_{n-1}$ as

$$\hat{p}_k := \langle \varphi^k, M^{-\frac{1}{2}} p \rangle_n = \langle \tilde{\varphi}^k, p \rangle_n, \quad \hat{r}_k := \langle \phi^k, r \rangle_{n-1}. \quad (2.15)$$

Notice that we implicitly defined $\tilde{\varphi}^k := M^{-\frac{1}{2}}\varphi^k$. We take $\hat{r}_0 = 0$ by convention. Let us mention that from the definition we have $\varphi_x^0 = m_x^{1/2}(\sum_{y=1}^n m_y)^{-1/2}$ for all $x \in \mathbb{I}_n$.

From equation of motion (2.13), and definition of \hat{p}_k, \hat{r}_k (2.15), thanks to the fact that ϕ^k, φ^k , are eigenvectors of A_r, A_p with eigenvalue ω_k^2 respectively, we obtain:

$$\ddot{\hat{p}}_k = -\omega_k^2 \hat{p}_k, \quad \ddot{\hat{r}}_k = -\omega_k^2 \hat{r}_k. \quad (2.16)$$

Solving the latter yields: for $k \in \mathbb{I}_{n-1}^o$ (recall the convention $\hat{r}_0 = 0$),

$$\begin{aligned} \hat{p}_k(t) &= \hat{p}_k(0) \cos(\omega_k t) - \hat{r}_k(0) \sin(\omega_k t) = \langle M^{-\frac{1}{2}}\varphi^k, p(0) \rangle_n \cos(\omega_k t) - \langle \phi^k, r(0) \rangle_{n-1} \sin(\omega_k t), \\ \hat{r}_k(t) &= \hat{r}_k(0) \cos(\omega_k t) + \hat{p}_k(0) \sin(\omega_k t) = \langle \phi^k, r(0) \rangle_{n-1} \cos(\omega_k t) + \langle M^{-\frac{1}{2}}\varphi^k, p(0) \rangle_n \sin(\omega_k t). \end{aligned}$$

Therefore, thanks to the above expressions, by using the inverse of (2.15), time evolution of r, p has the following explicit expression:

$$\begin{aligned} p(t) &= \sum_{k=0}^{n-1} M^{\frac{1}{2}}\varphi^k \hat{p}_k(t) = \sum_{k=0}^{n-1} (\cos(\omega_k t) \hat{p}_k(0) - \sin(\omega_k t) \hat{r}_k(0)) M^{\frac{1}{2}}\varphi^k, \\ r(t) &= \sum_{k=1}^{n-1} \phi^k \hat{r}_k(t) = \sum_{k=1}^{n-1} (\cos(\omega_k t) \hat{r}_k(0) + \sin(\omega_k t) \hat{p}_k(0)) \phi^k. \end{aligned} \quad (2.17)$$

2.2.2 Localization

The proof of the limit on average (1.3) relies on localization properties of $\tilde{\varphi}^k = M^{-\frac{1}{2}}\varphi^k$ for $k \gg \sqrt{n}$. We use similar properties in order to obtain proper decay estimates and deduce (2.10)- (2.12). Therefore, we express the desired localization estimates directly from [3] and [12] (cf. [1][22] for general theory, and [2][22] for more precise estimates).

Lemma 2.2. *Recall the definition of the vectors φ^k , as ordered eigenvectors of A_p (2.14) and $\tilde{\varphi}^k = M^{-\frac{1}{2}}\varphi^k$. Let $0 < \alpha < \frac{1}{2}$, and define $I(\alpha) :=]n^{(1-\alpha)}, n] \cap \mathbb{Z}$. There exists $C, c > 0$ independent of n such that*

$$\mathbb{E} \left(\sum_{k \in I(\alpha)} |\tilde{\varphi}_x^k \tilde{\varphi}_y^k| \right) \leq C \exp\left(-\frac{c|x-y|}{\xi(\alpha)}\right), \quad \text{with } \xi(\alpha) = n^{2\alpha}, \quad (2.18)$$

for $x, y \in \mathbb{I}_n$, where we remind that \mathbb{E} is the expectation with respect to the distribution of the masses. Notice that thanks to the fact that masses are compactly supported, the same estimate holds for φ^k with possibly different constants $c, C > 0$ independent of n .

The above lemma is enough to deal with the terms involving p variables. The terms with r variables involve the vectors ϕ^k but, instead of directly establishing a localization estimate for these, we will use Lemma 6.3 in [12]:

Lemma 2.3. *Recall the definition of $A_p = M^{-\frac{1}{2}}(-\Delta)M^{\frac{1}{2}}$ (2.14), and its ordered eigenvalues $0 = \omega_0 < \omega_1 < \dots < \omega_{n-1}$. Fix $\alpha, \gamma > 0$, such that $0 < 2\alpha < \gamma < 1$. There exists almost surely $n_0 \in \mathbb{N}$ such that $\forall n > n_0$, and for $k \in I(\alpha) =]n^{(1-\alpha)}, n-1] \cap \mathbb{Z}$ we have:*

$$\omega_k^{-1} \leq cn^{\frac{3\gamma}{2}}, \quad (2.19)$$

where c is a constant independent of n .

Let us remark that the estimate (2.19) is not optimal; we expect indeed that $1/\omega_k \leq cn$ for all $k \in \mathbb{I}_{n-1}$. However, (2.19) is sufficient for our purposes.

2.3 Evolution of r and p

In this section, we prove the two limits (2.10) and (2.11) featuring in Theorem 2.1. In the proof, we take advantage of the limit on average (1.3) which have been proved in [3]. We also use a couple of bounds introduced in [3].

Proof of Theorem 2.1: (2.10), (2.11). We define the *fluctuation* variables $\tilde{r}_x(nt)$ and $\tilde{p}_x(nt)$ for any $x \in \mathbb{I}_n$ as follows:

$$\tilde{r}_x(nt) := r_x(nt) - \langle r_x(nt) \rangle_{\rho^n}, \quad \tilde{p}_x(nt) := p_x(nt) - \langle p_x(nt) \rangle_{\rho^n}. \quad (2.20)$$

Adding and subtracting $\frac{1}{n} \sum_{x=1}^n f(\frac{x}{n}) \langle p_x(nt) \rangle_{\rho^n}$ inside the square in (2.11), and using the fact that $\langle \tilde{p}_x(nt) \rangle_{\rho} = 0$ we get:

$$\begin{aligned} \left\langle \left(\frac{1}{n} \sum_{x=1}^n f(\frac{x}{n}) p_x(nt) - \int_0^1 f(y) \mathfrak{p}(y, t) dy \right)^2 \right\rangle_{\rho^n} &= \left\langle \left(\frac{1}{n} \sum_{x=1}^n f(\frac{x}{n}) \tilde{p}_x(nt) \right)^2 \right\rangle_{\rho^n} \\ &+ \left\langle \left(\frac{1}{n} \sum_{x=1}^n f(\frac{x}{n}) \langle p_x(nt) \rangle_{\rho^n} - \int_0^1 f(y) \mathfrak{p}(y, t) dy \right)^2 \right\rangle_{\rho^n}. \end{aligned} \quad (2.21)$$

Notice that the last term converges to zero almost surely w.r.t. the distribution of the masses thanks to the limit on average (1.3). One can do the same for r as well. Therefore, in order to prove (2.10), and (2.11) it is sufficient to prove:

$$\mathcal{R}_n := \left\langle \left(\frac{1}{n} \sum_{x=1}^n f(\frac{x}{n}) \tilde{r}_x(nt) \right)^2 \right\rangle_{\rho^n} \rightarrow 0, \quad \mathcal{P}_n := \left\langle \left(\frac{1}{n} \sum_{x=1}^n f(\frac{x}{n}) \tilde{p}_x(nt) \right)^2 \right\rangle_{\rho^n} \rightarrow 0, \quad (2.22)$$

as $n \rightarrow \infty$, almost surely w.r.t the distribution of the masses.

Before proceeding, notice that the evolution equation (2.5) is linear. Therefore, $\langle r_x \rangle_{\rho^n}$, $\langle p_x \rangle_{\rho^n}$ satisfy the same set of equations with modified initial data. This is still true for \tilde{r}_x and \tilde{p}_x . Consequently, one can solve the equation of motion featuring \tilde{r}_x and \tilde{p}_x as in (2.17). More precisely, denote the vectors $\tilde{p} = p - \langle p \rangle_{\rho^n}$, $\tilde{r} = r - \langle r \rangle_{\rho^n}$ and define

$$\hat{\tilde{p}}_k = \hat{\tilde{p}}_k(0) := \langle \varphi^k, M^{-\frac{1}{2}} \tilde{p} \rangle_n, \quad \hat{\tilde{r}}_k := \langle \phi^k, \tilde{r} \rangle_{n-1}, \quad (2.23)$$

then $\tilde{r}_x(t)$, $\tilde{p}_x(t)$ is given by:

$$\begin{aligned} \tilde{r}_x(t) &= \sum_{k=1}^{n-1} \phi_x^k \hat{\tilde{r}}_k(t) = \sum_{k=1}^{n-1} \left(\cos(\omega_k t) \hat{\tilde{r}}_k(0) + \sin(\omega_k t) \hat{\tilde{p}}_k(0) \right) \phi_x^k, \\ \tilde{p}_x(t) &= \sum_{k=0}^{n-1} \sqrt{m_x} \varphi_x^k \hat{\tilde{p}}_k(t) = \sum_{k=0}^{n-1} \left(\cos(\omega_k t) \hat{\tilde{p}}_k(0) - \sin(\omega_k t) \hat{\tilde{r}}_k(0) \right) \sqrt{m_x} \varphi_x^k. \end{aligned} \quad (2.24)$$

In order to prove (2.22), we divide each sum into three parts, and prove that each part converges to zero almost surely w.r.t the distribution of the masses. Based on this, we divide the rest of the proof into three steps.

Step 1: Low modes. Lemma 2.2 only provides localization of the high modes. Hence we treat the contribution of the low modes separately. For proper γ such that $0 < \gamma < \frac{1}{2}$, define $\tilde{p}^o(nt)$, and $\tilde{r}^o(nt)$ as the “low mode portion” of p , and r : for $x \in \mathbb{I}_n$ let

$$\tilde{p}_x^o(nt) := \sum_{k \in \mathbb{Z} \cap [0, n^{(1-\gamma)}]} \hat{p}_k(nt) \sqrt{m_x} \varphi_x^k, \quad \tilde{r}_x^o(nt) := \sum_{k \in \mathbb{Z} \cap [1, n^{(1-\gamma)}]} \hat{r}_k(nt) \phi_x^k. \quad (2.25)$$

In this step, we prove for any $t \in [0, T]$:

$$\mathcal{L}_n := \left\langle \left(\frac{1}{n} \sum_{x=1}^n f\left(\frac{x}{n}\right) \tilde{r}_x^o(nt) \right)^2 \right\rangle_{\rho^n} \rightarrow 0, \quad \mathcal{L}'_n := \left\langle \left(\frac{1}{n} \sum_{x=1}^n f\left(\frac{x}{n}\right) \tilde{p}_x^o(nt) \right)^2 \right\rangle_{\rho^n} \rightarrow 0. \quad (2.26)$$

for any realization of the masses.

First, notice that ρ^n (2.6) is a Gaussian probability density function, and $\{r_x\}_{x=1}^{n-1}$ and $\{p_x\}_{x=1}^n$ are independently distributed. Therefore, we have for proper $x, y \in \mathbb{I}_n$:

$$\langle \tilde{r}_x \tilde{r}_y \rangle_{\rho^n} = \frac{\delta_{x,y}}{\beta(\frac{x}{n})}, \quad \langle \tilde{p}_x \tilde{p}_y \rangle_{\rho^n} = \delta_{x,y} \frac{m_x}{\beta(\frac{x}{n})}, \quad \langle \tilde{r}_x \tilde{p}_y \rangle = 0. \quad (2.27)$$

Recall the definition of \hat{r}_k, \hat{p}_k (2.23). For any k , there exists $c > 0$ independent of n , and independent of the realization of the masses such that

$$\begin{aligned} \langle \hat{p}_k^2 \rangle_{\rho^n} &= \sum_{x,y=1}^n \varphi_x^k \varphi_y^k \frac{\langle \tilde{p}_x \tilde{p}_y \rangle_{\rho^n}}{\sqrt{m_x m_y}} = \sum_{x=1}^n \beta^{-1}\left(\frac{x}{n}\right) (\varphi_x^k)^2 \leq c, \\ \langle \hat{r}_k^2 \rangle_{\rho^n} &= \sum_{x=1}^{n-1} \beta^{-1}\left(\frac{x}{n}\right) (\phi_x^k)^2 \leq c, \quad \langle \hat{p}_k \hat{r}_{k'} \rangle_{\rho^n} = 0, \end{aligned} \quad (2.28)$$

where first we exploit (2.27), then we take advantage of the fact that $0 < \beta_- < \beta(y)$ (in fact, we can take $c = \beta_-^{-1}$), and finally we use the fact that ϕ^k , and φ^k are members of two orthonormal basis. Recall $\hat{p}_k(nt), \hat{r}_k(nt)$ from (2.24), then from (2.28) it is evident that for any $t \in [0, T]$, and any $k \in \mathbb{I}_{n-1}^o$

$$\langle \hat{p}_k^2(nt) \rangle_{\rho^n} \leq c, \quad \langle \hat{r}_k^2(nt) \rangle_{\rho^n} \leq c, \quad (2.29)$$

where, as before c is independent of n and the realization of the masses.

Let $f_x := f(\frac{x}{n})$ and $\tilde{I}(\gamma) := [0, n^{(1-\gamma)}] \cap \mathbb{Z}$. We bound \mathcal{L}'_n defined in (2.26) as follows:

$$\begin{aligned}
0 \leq \mathcal{L}'_n &= \frac{1}{n^2} \sum_{x,y=1}^{n-1} f_x f_y \langle \tilde{p}_x^o(nt) \tilde{p}_y^o(nt) \rangle_{\rho^n} \leq \frac{C}{n^2} \sum_{x,y=1}^n \frac{|\langle \tilde{p}_x^o(nt) \tilde{p}_y^o(nt) \rangle_{\rho^n}|}{\sqrt{m_x} \sqrt{m_y}} \\
&\leq \frac{C}{n^2} \sum_{x,y=1}^n \frac{1}{\sqrt{m_x m_y}} \langle (\tilde{p}_x^o(nt))^2 \rangle_{\rho^n}^{\frac{1}{2}} \langle (\tilde{p}_y^o(nt))^2 \rangle_{\rho^n}^{\frac{1}{2}} = C \left(\frac{1}{n} \sum_{x=1}^n \frac{1}{\sqrt{m_x}} \langle (\tilde{p}_x^o(nt))^2 \rangle_{\rho^n}^{\frac{1}{2}} \right)^2 \\
&\leq \frac{C}{n} \sum_{x=1}^n \frac{\langle (\tilde{p}_x^o(nt))^2 \rangle_{\rho^n}}{m_x} = \frac{C}{n} \sum_{x=1}^n \left\langle \sum_{k,k' \in \tilde{I}(\gamma)} \hat{p}_k(nt) \hat{p}_{k'}(nt) \varphi_x^k \varphi_x^{k'} \right\rangle_{\rho^n} \quad (2.30) \\
&\leq \frac{C}{n} \sum_{k,k' \in \tilde{I}(\gamma)} \langle \hat{p}_k(nt) \hat{p}_{k'}(nt) \rangle_{\rho^n} \sum_{x=1}^n \varphi_x^k \varphi_x^{k'} = \frac{C}{n} \sum_{k \in \tilde{I}(\gamma)} \langle (\hat{p}_k(nt))^2 \rangle_{\rho^n} \\
&\leq cC \frac{|\tilde{I}(\gamma)|}{n} = cC \frac{n^{(1-\gamma)}}{n} \rightarrow 0
\end{aligned}$$

as $n \rightarrow \infty$, for any realization of the masses; where, in the first line we bounded $|f_x f_y| \sqrt{m_x m_y}$ by $C = \|f\|_\infty^2 m_+ > 0$ which is independent of n (recall f is continuous on $[0, 1]$, and the distribution of the masses is supported in $[m_-, m_+]$), in the second line we take advantage of the Cauchy-Schwartz inequality i.e. $|\langle AB \rangle_{\rho^n}|^2 \leq \langle A^2 \rangle_{\rho^n} \langle B^2 \rangle_{\rho^n}$, in the third line we used another Cauchy-Schwartz inequality ($|\langle a, b \rangle_n|^2 \leq |a|^2 |b|^2$ for $a, b \in \mathbb{R}^n$), in the third line, we also used the definition of $\tilde{p}_x^o(nt)$ from (2.25) and obtained a double sum by squaring this definition, in the fourth line we used the linearity of $\langle \cdot \rangle_{\rho^n}$ as well as the fact that φ^k is an orthonormal basis ($\langle \varphi^k, \varphi^{k'} \rangle_n = \delta_{k,k'}$), finally in the last inequality we used the second bound in (2.29) ($\langle \hat{p}_k^2(nt) \rangle_{\rho^n} \leq c$).

We can prove $\mathcal{L}_n \rightarrow 0$ deterministically in a similar way with some small modifications: we should use the definition of \tilde{p}^o instead of \tilde{r}^o , then use the fact that $\langle \phi^k, \phi^{k'} \rangle_{n-1} = \delta_{k,k'}$, and finally take advantage of the second bound in (2.28) ($\langle \hat{r}_k^2(nt) \rangle_{\rho^n} \leq c$).

Step2: High modes. Recall $0 < \gamma < \frac{1}{2}$, $I(\gamma) =]n^{(1-\gamma)}, n] \cap \mathbb{Z}$, and (2.25), then define the “high mode portion” of p , and r as follows:

$$\begin{aligned}
\tilde{p}_x^\bullet(nt) &:= \tilde{p}_x(nt) - \tilde{p}_x^o(nt) = \sum_{k \in I(\gamma)} \hat{p}_k(nt) \sqrt{m_x} \varphi_x^k, \\
\tilde{r}_x^\bullet(nt) &:= \tilde{r}_x(nt) - \tilde{r}_x^o(nt) = \sum_{k \in I(\gamma)} \hat{r}_k(nt) \phi_x^k.
\end{aligned} \quad (2.31)$$

In this step we prove that

$$\mathcal{U}'_n := \left\langle \left(\frac{1}{n} \sum_{x=1}^n f\left(\frac{x}{n}\right) \tilde{r}_x^\bullet(nt) \right)^2 \right\rangle_{\rho^n} \rightarrow 0, \quad \mathcal{U}_n := \left\langle \left(\frac{1}{n} \sum_{x=1}^n f\left(\frac{x}{n}\right) \tilde{p}_x^\bullet(nt) \right)^2 \right\rangle_{\rho^n} \rightarrow 0, \quad (2.32)$$

almost surely w.r.t the distribution of the masses. Take $0 < 2\gamma < \theta < 1$. Expanding the sum and bounding $|f(y)f(y')m_x m_y| \leq C = \|f\|_\infty^2 m_+$, we have for n sufficiently large:

$$0 \leq \mathcal{U}_n \leq \frac{C}{n^2} \sum_{x,y=1}^n \left| \frac{\langle \tilde{p}_x^\bullet(nt) \tilde{p}_y^\bullet(nt) \rangle_{\rho^n}}{\sqrt{m_x m_y}} \right| =$$

$$C \left(\underbrace{\frac{1}{n^2} \sum_{|x-y| \leq 2n^\theta} \left| \frac{\langle \tilde{p}_x^\bullet(nt) \tilde{p}_y^\bullet(nt) \rangle_{\rho^n}}{\sqrt{m_x m_y}} \right|}_{:=\mathcal{U}_n^<} + \underbrace{\frac{1}{n^2} \sum_{|x-y| > 2n^\theta} \left| \frac{\langle \tilde{p}_x^\bullet(nt) \tilde{p}_y^\bullet(nt) \rangle_{\rho^n}}{\sqrt{m_x m_y}} \right|}_{:=\mathcal{U}_n^>} \right) \quad (2.33)$$

Step 2.1: High modes I ($|x-y| > 2n^\theta$). We defined $\mathcal{U}_n^<, \mathcal{U}_n^>$ in (2.33) implicitly. First, we prove $\mathcal{U}_n^> \rightarrow 0$, almost surely w.r.t the distribution of the masses. Before proceeding, recall the definition of $\hat{p}_k(0) \equiv \hat{p}_k$, and $\hat{r}_k(0) \equiv \hat{r}_k$ from (2.23). By using (2.27), observe that for proper k, k' (we use the shorthand notation $\beta^{-1}(\frac{x}{n}) =: b_x$):

$$\langle \hat{p}_k \hat{p}_{k'} \rangle_{\rho^n} = \sum_{z, z'} \frac{\varphi_z^k \varphi_{z'}^{k'}}{\sqrt{m_z m_{z'}}} \langle \tilde{p}_z \tilde{p}_{z'} \rangle_{\rho^n} = \sum_{z=1}^n b_z \varphi_z^k \varphi_z^{k'}, \quad \langle \hat{r}_k \hat{r}_{k'} \rangle_{\rho^n} = \sum_{z=1}^{n-1} b_z \phi_z^k \phi_z^{k'}. \quad (2.34)$$

Thanks to (2.31) we have:

$$\mathcal{U}_n^> = \frac{1}{n^2} \sum_{|x-y| > 2n^\theta} \left| \sum_{k, k' \in I(\gamma)} \langle \hat{p}_k(nt) \hat{p}_{k'}(nt) \rangle_{\rho^n} \varphi_x^k \varphi_y^{k'} \right| =$$

$$\frac{1}{n^2} \sum_{|x-y| > 2n^\theta} \left| \sum_{k, k' \in I(\gamma)} \varphi_x^k \varphi_y^{k'} \left(\langle \hat{p}_k \hat{p}_{k'} \rangle_{\rho^n} \cos(\omega_k nt) \cos(\omega_{k'} nt) + \langle \hat{r}_k \hat{r}_{k'} \rangle_{\rho^n} \sin(\omega_k nt) \sin(\omega_{k'} nt) \right) \right| \leq$$

$$\frac{1}{n^2} \sum_{|x-y| > 2n^\theta} \left| \sum_{z=1}^n b_z \left(\sum_{k \in I(\gamma)} \varphi_x^k \varphi_z^k \cos(\omega_k nt) \right) \left(\sum_{k' \in I(\gamma)} \varphi_y^{k'} \varphi_z^{k'} \cos(\omega_{k'} nt) \right) \right| +$$

$$\frac{1}{n^2} \sum_{|x-y| > 2n^\theta} \left| \sum_{z=1}^{n-1} b_z \left(\sum_{k \in I(\gamma)} \varphi_x^k \phi_z^k \sin(\omega_k nt) \right) \left(\sum_{k' \in I(\gamma)} \varphi_y^{k'} \phi_z^{k'} \sin(\omega_{k'} nt) \right) \right| \leq$$

$$\underbrace{\frac{C'}{n^2} \sum_{|x-y| > 2n^\theta} \sum_{z=1}^n \left(\sum_{k \in I(\gamma)} |\varphi_x^k| |\varphi_z^k| \right) \left(\sum_{k' \in I(\gamma)} |\varphi_y^{k'}| |\varphi_z^{k'}| \right)}_{:=\mathcal{U}_n^{>,p}} +$$

$$\underbrace{\frac{C'}{n^2} \sum_{|x-y| > 2n^\theta} \sum_{z=1}^n \left(\sum_{k \in I(\gamma)} |\varphi_x^k| |\phi_z^k| \right) \left(\sum_{k' \in I(\gamma)} |\varphi_y^{k'}| |\phi_z^{k'}| \right)}_{:=\mathcal{U}_n^{>,r}}, \quad (2.35)$$

where in the second line we used the definition of $\hat{p}_k(nt)$ (2.24), and the fact that $\langle \hat{r}_k \hat{p}_{k'} \rangle_{\rho^n} = 0$ (2.28), in the third line we take advantage of (2.34), and in the fourth line we bounded $|\sin(\cdot)|, |\cos(\cdot)| \leq 1$, and $|b_z| = \beta^{-1}(\frac{z}{n}) \leq C' = \beta_-^{-1}$ uniform in n .

For any $x \in \mathbb{I}_n$, denote $J(x) := \{z \in \mathbb{I}_n \mid |x - z| \leq n^\theta\}$. Notice the definition of $\mathcal{U}_n^{<,p}$, $\mathcal{U}_n^{>,p}$ from the last lines of (2.35). Take a single term from the main sum featuring $\mathcal{U}_n^{<,p}$ and observe for n sufficiently large (recall \mathbb{E} is the expectation w.r.t the distribution of the masses and recall that here $|x - y| > n^{2\theta}$):

$$\begin{aligned} \mathbb{E} \left(\sum_{z=1}^n \left(\sum_{k \in I(\gamma)} |\varphi_x^k| |\varphi_z^k| \right) \left(\sum_{k' \in I(\gamma)} |\varphi_y^{k'}| |\varphi_z^{k'}| \right) \right) &= \sum_{z \in J(x)} \mathbb{E} \left(\left(\sum_{k \in I(\gamma)} |\varphi_x^k| |\varphi_z^k| \right) \left(\sum_{k' \in I(\gamma)} |\varphi_y^{k'}| |\varphi_z^{k'}| \right) \right) + \\ &\sum_{z \notin J(x)} \mathbb{E} \left(\left(\sum_{k \in I(\gamma)} |\varphi_x^k| |\varphi_z^k| \right) \left(\sum_{k' \in I(\gamma)} |\varphi_y^{k'}| |\varphi_z^{k'}| \right) \right) \leq \sum_{z \in J(x)} \mathbb{E} \left(\sum_{k' \in I(\gamma)} |\varphi_y^{k'}| |\varphi_z^{k'}| \right) + \\ &\sum_{z \notin J(x)} \mathbb{E} \left(\sum_{k \in I(\gamma)} |\varphi_x^k| |\varphi_z^k| \right) \leq C \sum_{z \in J(x)} \exp(-c|y - z|/n^{2\gamma}) + C \sum_{z \notin J(x)} \exp(-c|x - z|/n^{2\gamma}) \leq \\ &4Cn^\theta \exp(-c(n^\theta - n^{2\gamma})) + Cn \exp(-c(n^\theta - n^{2\gamma})) \leq 4C \exp(-\frac{c}{4}(n^\theta)), \end{aligned} \quad (2.36)$$

where in the first inequality we bounded:

$$\sum_{k \in I(\gamma)} |\varphi_x^k| |\varphi_z^k| \leq \left(\sum_{k \in I(\gamma)} |\varphi_x^k|^2 \right)^{\frac{1}{2}} \left(\sum_{k \in I(\gamma)} |\varphi_z^k|^2 \right)^{\frac{1}{2}} \leq \left(\sum_{k=0}^{n-1} |\varphi_x^k|^2 \right)^{\frac{1}{2}} \left(\sum_{k=0}^{n-1} |\varphi_z^k|^2 \right)^{\frac{1}{2}} = 1, \quad (2.37)$$

thanks to a Cauchy-Schwartz inequality and the fact that φ^k is an orthonormal basis. We bounded $\sum_{k' \in I(\gamma)} |\varphi_y^{k'}| |\varphi_z^{k'}| \leq 1$ similarly. The second inequality in (2.36) is deduced from Lemma 2.2 (modification of (2.18) by a constant). Notice that this is the crucial place where we use localization. The fourth inequality in (2.36) is deduced from the definition of $J(x)$ and the fact that $|x - y| > 2n^\theta$: more precisely, for $z \notin J(x)$, $|x - z| > n^\theta$, and for $z \in J(x)$ (i.e. $|z - x| \leq n^\theta$), we have $|y - z| > n^\theta$ thanks to the choice of y : $|x - y| > 2n^\theta$. In the last inequality we use the choice of θ : $0 < 2\gamma < \theta < 1$, and that n is sufficiently large.

Recalling the definition of $\mathcal{U}_n^{<,p}$ (2.35), thanks to the latter estimate (2.36) there exists $c_1, C > 0$ independent of n such that for n sufficiently large we have:

$$\mathbb{E}(|\mathcal{U}_n^{>,p}|) \leq C \exp(-c_1 n^\theta) \implies \sum_n \mathbb{E}(|\mathcal{U}_n^{>,p}|) < \infty \implies \mathcal{U}_n^{>,p} \rightarrow 0, \quad (2.38)$$

almost surely w.r.t the distribution of the masses, where we used the fact that $\theta > 0$, which means that the above series is summable, and finally we conclude thanks to Borel-Cantelli's lemma.

Recall $\mathcal{U}_n^{>,r}$ from (2.35). This expression can be treated similar to $\mathcal{U}_n^{>,p}$ thanks to Lemma 2.3: First, recall for $z \in \mathbb{I}_{n-1}$, $\phi_z^k = \frac{1}{\omega_k}(\frac{\varphi_{z+1}^k}{m_{z+1}} - \frac{\varphi_z^k}{\sqrt{m_z}})$. Let us denote $\tilde{\omega} := \omega_{k_*}$ with

$k_* := \min\{I(\gamma)\}$. For any n define the sequence $a_n := \tilde{\omega}n^{\frac{3\gamma}{2}}$. Since $\tilde{\omega} \leq \omega_k$ for any $k \in I(\omega)$ we have $C, c > 0$ independent of n such that:

$$\mathbb{E} \left(a_n \sum_{k \in I(\gamma)} |\varphi_x^k| |\phi_z^k| \right) \leq n^{\frac{3\gamma}{2}} 2C \exp\left(-\frac{c}{2}|z-x|/n^{2\gamma}\right), \quad (2.39)$$

where we used the definition of ϕ^k , and estimate (2.18).

Having above estimate, by doing a computation similar to (2.36), and (2.37) we can deduce for n sufficiently large there exist $c', C > 0$ independent of n :

$$\mathbb{E}(|a_n \mathcal{U}_n^{>,r}|) \leq C n^{\frac{3\gamma}{2}} \exp(-c'n^\theta) \leq C \exp\left(-\frac{c'}{2}n^\theta\right). \quad (2.40)$$

Therefore, by Borel-Cantelli's lemma we have $a_n \mathcal{U}_n^{>,r} \rightarrow 0$, almost surely w.r.t the distribution of the masses. However, thanks to Lemma 2.3 (i.e. (2.19)), we know there exists almost surely n_0 , such that for $n > n_0$, we have $a_n \geq \frac{1}{c}$ with $c > 0$. Using these two facts we deduce that $\mathcal{U}_n^{>,r} \rightarrow 0$, almost surely w.r.t distribution of the masses. Combining $\mathcal{U}_n^{>,r} \rightarrow 0$, with (2.38) ($\mathcal{U}_n^{>,p} \rightarrow 0$) we get $\mathcal{U}_n^{>} \rightarrow 0$ almost surely w.r.t the distribution of the masses thanks to (2.35).

Recall the definition of \mathcal{U}'_n (2.32). We can define $\mathcal{U}'_n^{<}, \mathcal{U}'_n^{>}$ corresponding to r variable similar to $\mathcal{U}_n^{<}, \mathcal{U}_n^{>}$ as in (2.33):

$$\mathcal{U}'_n^{<} := \frac{1}{n^2} \sum_{|x-y| \leq 2n^\theta} |\langle \tilde{r}_x^\bullet(nt) \tilde{r}_y^\bullet(nt) \rangle_{\rho^n}|, \quad \mathcal{U}'_n^{>} := \frac{1}{n^2} \sum_{|x-y| > 2n^\theta} |\langle \tilde{r}_x^\bullet(nt) \tilde{r}_y^\bullet(nt) \rangle_{\rho^n}|. \quad (2.41)$$

Then we can prove $\mathcal{U}'_n^{>} \rightarrow 0$ almost surely, with respect to the distribution of the masses as we did for $\mathcal{U}_n^{>}$. We only sketch the former, since these proofs are almost identical. First, one can compute $\mathcal{U}'_n^{>}$ similar to (2.35) and decompose the r and p contribution and define $\mathcal{U}'_n^{>,r}, \mathcal{U}'_n^{>,p}$ similar to (2.35). Then one can observe that $\mathcal{U}'_n^{>,p} = \mathcal{U}_n^{>,p} \rightarrow 0$ almost surely. Proving $\mathcal{U}'_n^{>,r} \rightarrow 0$, almost surely, is similar to the previous terms: First we decompose the sum in $\mathcal{U}'_n^{>,r}$ into two parts similar to (2.36), then we use proper estimates. The only difference lies in the following estimate: Define $b_n = \tilde{\omega}^2 n^{3\gamma}$ (recall $\tilde{\omega} = \omega_{k_*}$ with $k_* = \min\{I(\gamma)\}$). Then similar to (2.39), one can observe that there exist c, C such that

$$\mathbb{E} \left(b_n \sum_{k \in I(\gamma)} |\phi_x^k \phi_z^k| \right) \leq n^{3\gamma} C \exp(-c|x-z|/n^{2\gamma}). \quad (2.42)$$

Consequently, one can deduce that $b_n \mathcal{U}'_n^{>,r} \rightarrow 0$ almost surely, as we did for $a_n \mathcal{U}_n^{>,r}$ in (2.40). Therefore, thanks to Lemma 2.19 we can deduce there exists almost surely n_0 such that for $n > n_0$, we have $b_n \geq \frac{1}{c^2}$. Hence, we have $\mathcal{U}'_n^{>,r} \rightarrow 0$ almost surely. This finishes the proof of the fact that $\mathcal{U}'_n^{>} \rightarrow 0$ almost surely w.r.t the distribution of the masses.

Step 2.2: High modes II ($|x-y| \leq 2n^\theta$). In this step, we prove $\mathcal{U}'_n^{<}$ and $\mathcal{U}'_n^{>}$ defined in (2.33) and (2.41) respectively, converge to zero almost surely. Similarly to the computation

in (2.36), thanks to the definition of $\tilde{p}_x^\bullet(nt)$ (2.31), and identities (2.28), (2.34), we have for any $x \in \mathbb{I}_n$ (recall $I(\gamma) =]n^{1-\gamma}, n] \cap \mathbb{Z}$, $b_z = \beta^{-1}(\frac{z}{n})$):

$$\begin{aligned}
\left\langle \frac{(\tilde{p}_x^\bullet(nt))^2}{m_x} \right\rangle_{\rho^n} &= \sum_{k,k' \in I(\gamma)} \varphi_x^k \varphi_x^{k'} \langle \hat{p}_k(nt) \hat{p}_{k'}(nt) \rangle_{\rho^n} = \\
&= \sum_{z=1}^n b_z \left(\sum_{k \in I(\gamma)} \cos(\omega_k nt) \varphi_x^k \varphi_z^k \right)^2 + \sum_{z=1}^{n-1} b_z \left(\sum_{k \in I(\gamma)} \sin(\omega_k nt) \varphi_x^k \varphi_z^k \right)^2 \leq \\
&= C \sum_{z=1}^n \left(\sum_{k \in I(\gamma)} \cos(\omega_k nt) \varphi_x^k \varphi_z^k \right)^2 + C \sum_{z=1}^{n-1} \left(\sum_{k \in I(\gamma)} \sin(\omega_k nt) \varphi_x^k \varphi_z^k \right)^2 = \\
&= C \sum_{k,k' \in I(\gamma)} \cos(\omega_k nt) \cos(\omega_{k'} nt) \varphi_x^k \varphi_x^{k'} \sum_{z=1}^n \varphi_z^k \varphi_z^{k'} + C \sum_{k,k' \in I(\gamma)} \sin(\omega_k nt) \sin(\omega_{k'} nt) \varphi_x^k \varphi_x^{k'} \sum_{z=1}^{n-1} \varphi_z^k \varphi_z^{k'} = \\
&= C \sum_{k \in I(\gamma)} (\cos^2(\omega_k nt) + \sin^2(\omega_k nt)) (\varphi_x^k)^2 \leq C \sum_{k=0}^{n-1} (\varphi_x^k)^2 = C,
\end{aligned} \tag{2.43}$$

where in the first inequality we bounded b_z by $C = \beta^{-1}$ (which is independent of n and the realization of the masses). Notice that in the first inequality, we used the fact that all the terms in the sum are positive (i.e. $b_z > 0$). In the last two lines we used the fact that φ^k, ϕ^k are orthonormal basis ($\langle \varphi^k, \varphi^{k'} \rangle_n = \delta_{k,k'}$, $\langle \phi^k, \phi^{k'} \rangle_{n-1} = \delta_{k,k'}$, $\sum_z \varphi_z^k \varphi_z^{k'} = \sum_z \phi_z^k \phi_z^{k'} = \delta_{k,k'}$).

Recall the definition of $\mathcal{U}_n^<$ (2.33) with $\theta < 1$, then thanks to (2.43) we have:

$$\begin{aligned}
\mathcal{U}_n^< &= \frac{1}{n^2} \sum_{|x-y| \leq 2n^\theta} \left| \left\langle \frac{\tilde{p}_x^\bullet(nt) \tilde{p}_y^\bullet(nt)}{\sqrt{m_x m_y}} \right\rangle_{\rho^n} \right| \leq \frac{1}{n^2} \sum_{|x-y| \leq 2n^\theta} \left\langle \frac{(\tilde{p}_x^\bullet(nt))^2}{m_x} \right\rangle_{\rho^n}^{\frac{1}{2}} \left\langle \frac{(\tilde{p}_y^\bullet(nt))^2}{m_y} \right\rangle_{\rho^n}^{\frac{1}{2}} \leq \\
&\leq \frac{1}{n^2} \sum_{|x-y| < 2n^\theta} C \leq 8C \frac{n^{1+\theta}}{n^2} \rightarrow 0,
\end{aligned} \tag{2.44}$$

where in the first line we bounded $\mathcal{U}_n^<$ by Cauchy-Schwartz inequality, and in the second line we used (2.43). Recall the decomposition $\mathcal{U}_n \leq C(\mathcal{U}_n^< + \mathcal{U}_n^>)$ (2.33). In Step 2.1, we observed that $\mathcal{U}_n^> \rightarrow 0$, almost surely. Combining this fact with (2.44), we deduce that $\mathcal{U}_n \rightarrow 0$, almost surely w.r.t the distribution of the masses.

Recall the definition of $\mathcal{U}_n'^<$ (2.41). Exactly similar to (2.43), one can bound $\langle (\tilde{r}_x^\bullet(nt))^2 \rangle_{\rho^n}$ thanks to its definition (2.31), and averages (2.28), (2.34): for any $x \in \mathbb{I}_n$ there exists $\beta_-^{-1} = C > 0$ independent of n and realization of the masses such that

$$\langle (\tilde{r}_x^\bullet(nt))^2 \rangle_{\rho^n} \leq C. \tag{2.45}$$

Similar to (2.44), by using a Cauchy-Schwartz inequality and the bound (2.45) we deduce, that $\mathcal{U}_n'^< \rightarrow 0$. Having the latter, we also proved $\mathcal{U}_n'^> \rightarrow 0$, almost surely in the previous step. Therefore, thanks to the bound $\mathcal{U}_n' \leq C(\mathcal{U}_n'^> + \mathcal{U}_n'^<)$ we conclude $\mathcal{U}_n' \rightarrow 0$ almost surely w.r.t distribution of the masses. This finishes the proof of (2.32).

Step 3: Summing up. Recall the definition of \mathcal{P}_n , \mathcal{L}'_n , and \mathcal{U}_n (2.22), (2.26), and (2.32). Since $\tilde{p}_x(nt) = \tilde{p}_x^o(nt) + \tilde{p}_x^\bullet(nt)$ (cf. (2.31), (2.25)), one can observe that $\mathcal{P}_n \leq \mathcal{L}'_n + \mathcal{U}_n + (\mathcal{L}'_n)^{\frac{1}{2}}(\mathcal{U}_n)^{\frac{1}{2}}$ thanks to a Cauchy-Schwartz inequality ($|\langle AB \rangle_{\rho^n}| \leq \langle A^2 \rangle_{\rho^n}^{\frac{1}{2}} \langle B^2 \rangle_{\rho^n}^{\frac{1}{2}}$). Hence we have $\mathcal{P}_n \rightarrow 0$, almost surely w.r.t the distribution of the masses, since $\mathcal{L}'_n \rightarrow 0$, and $\mathcal{U}_n \rightarrow 0$ almost surely thanks to (2.26), and (2.32). We can deduce $\mathcal{R}_n \rightarrow 0$ (2.22) almost surely similarly, by using the fact that $\tilde{r}_x(nt) = \tilde{r}_x^o(nt) + \tilde{r}_x^\bullet(nt)$, and $\mathcal{L}_n \rightarrow 0$, $\mathcal{U}'_n \rightarrow 0$, almost surely. This finishes the proof of (2.22), and we conclude first two limit of Theorem 2.1: (2.10), and (2.11). \square

Remark 2.1. The bound (2.43) is true for \tilde{p}_x as well. Moreover, similar to (2.43), we can bound $\langle \tilde{p}_x^2(nt)/m_x \rangle_{\rho^n}$ by β_+^{-1} from below. This means the temperature at each point is bounded by the maximum and minimum of the temperature profile at each time.

Remark 2.2. Notice that in the above proof, for proving $\mathcal{P}_n \rightarrow 0$, $\mathcal{R}_n \rightarrow 0$ almost surely w.r.t distribution of the masses, (2.22), we only used the fact that f is bounded. Therefore, it is straightforward to observe that (2.22) holds for bounded f as well.

2.4 Evolution of e

In this section, we prove (2.12). Again, our proof rests on the limit on average (1.3).

Proof of Theorem 2.1: (2.12). Similarly to (2.20), let us define:

$$\tilde{e}_x(nt) := e_x(nt) - \langle e_x(nt) \rangle_{\rho^n} = \frac{1}{2} \left(\frac{p_x^2(nt)}{m_x} + r_x^2(nt) - \frac{\langle p_x^2(nt) \rangle_{\rho^n}}{m_x} - \langle r_x^2(nt) \rangle_{\rho^n} \right). \quad (2.46)$$

Similarly to (2.22), one can add and subtract $\frac{1}{n} \sum_{x=1}^n f(\frac{x}{n}) \langle e_x \rangle_{\rho^n}$ inside the square in (2.12). Then using the limit on average (1.3), it is clear that in order to obtain (2.12) it is sufficient to prove

$$\mathcal{E}_n := \left\langle \left(\frac{1}{n} \sum_{x=1}^n f\left(\frac{x}{n}\right) \tilde{e}_x(nt) \right)^2 \right\rangle_{\rho^n} \rightarrow 0, \quad (2.47)$$

almost surely w.r.t the distribution of the masses. Plugging the definition of (2.46) into (2.47), and taking advantage of Cauchy-Schwartz inequality ($|\langle AB \rangle_{\rho^n}| \leq \langle A^2 \rangle_{\rho^n}^{\frac{1}{2}} \langle B^2 \rangle_{\rho^n}^{\frac{1}{2}}$), we observe that for deducing (2.47) it is enough to show:

$$\begin{aligned} \mathfrak{P}_n &:= \left\langle \left(\frac{1}{n} \sum_{x=1}^n f\left(\frac{x}{n}\right) \left(\frac{p_x^2(nt)}{m_x} - \langle \frac{p_x^2(nt)}{m_x} \rangle_{\rho^n} \right) \right)^2 \right\rangle_{\rho^n} \rightarrow 0, \\ \mathfrak{R}_n &:= \left\langle \left(\frac{1}{n} \sum_{x=1}^n f\left(\frac{x}{n}\right) (r_x^2(nt) - \langle r_x^2(nt) \rangle_{\rho^n}) \right)^2 \right\rangle_{\rho^n} \rightarrow 0, \end{aligned} \quad (2.48)$$

almost surely w.r.t the distribution of the masses. In the following, we prove $\mathfrak{P}_n \rightarrow 0$ almost surely. The proof of $\mathfrak{R}_n \rightarrow 0$ goes in parallel, and we omit it for the sake of brevity. Let us denote

$$\bar{p}_x(nt) := \langle p_x(nt) \rangle_{\rho^n},$$

and recall the definition of $\tilde{p}_x(nt) = p_x(nt) - \bar{p}_x(nt)$ (2.20). Since $\langle \tilde{p}_x(nt) \rangle_{\rho^n} = 0$, after a straightforward computation, we have for $x, y \in \mathbb{I}_n$:

$$\begin{aligned} & \langle (p_x^2(nt) - \langle p_x^2(nt) \rangle_{\rho^n}) (p_y^2(nt) - \langle p_y^2(nt) \rangle_{\rho^n}) \rangle_{\rho^n} = \\ & \langle \tilde{p}_x^2(nt) \tilde{p}_y^2(nt) \rangle_{\rho^n} - \langle \tilde{p}_x^2(nt) \rangle_{\rho^n} \langle \tilde{p}_y^2(nt) \rangle_{\rho^n} + 4\bar{p}_x(nt)\bar{p}_y(nt)\langle \tilde{p}_x(nt)\tilde{p}_y(nt) \rangle_{\rho^n} + \\ & 2\bar{p}_x(nt)\langle \tilde{p}_x(nt)\tilde{p}_y^2(nt) \rangle_{\rho^n} + 2\bar{p}_y(nt)\langle \tilde{p}_x^2(nt)\tilde{p}_y(nt) \rangle_{\rho^n}. \end{aligned} \quad (2.49)$$

Notice that owing to the definition of ρ^n (2.6), for any realization of the masses $(p(0), r(0))$ is a Gaussian random vector, where the randomness refers to the probability distribution ρ^n . In addition, the evolution equation (2.5) is linear, and the vector $(p(nt), r(nt))$ can be written as $(p(nt), r(nt))^\dagger = O(t)(p(0), r(0))^\dagger$ for a linear transformation $O(t)$, as we observed in (2.17). Therefore, the vector $(p(nt), r(nt))$ is another Gaussian vector. Consequently, $(\tilde{p}(nt), \tilde{r}(nt))$ is a zero mean Gaussian vector (equivalently if $\rho^n(t)$ denotes the solution to the backward equation, $\rho^n(t)$ will be Gaussian). Hence we can use Isserlis' theorem (Wick's theorem cf. [13]) and observe that any odd moments of \tilde{p}, \tilde{r} is zero. Moreover, for any even moment we have pairing. More precisely, for k odd $\langle \tilde{p}_{x_1}(nt) \dots \tilde{p}_{x_k}(nt) \rangle_{\rho^n} = 0$, for any $x_1, \dots, x_k \in \mathbb{I}_k$. For even k we have

$$\begin{aligned} \langle \tilde{p}_{x_1}(nt) \dots \tilde{p}_{x_k}(nt) \rangle_{\rho^n} &= \sum_{\pi \in \Pi} \prod_{\{i,j\} \in \pi} \langle \tilde{p}_{x_i}(nt) \tilde{p}_{x_j}(nt) \rangle_{\rho^n} \implies \\ \langle \tilde{p}_x^2(nt) \tilde{p}_y^2(nt) \rangle_{\rho^n} - \langle \tilde{p}_x^2(nt) \rangle_{\rho^n} \langle \tilde{p}_y^2(nt) \rangle_{\rho^n} &= 2\langle \tilde{p}_x(nt) \tilde{p}_y(nt) \rangle_{\rho^n}^2; \quad \langle \tilde{p}_x(nt) \tilde{p}_y^2(nt) \rangle_{\rho^n} = 0 \end{aligned} \quad (2.50)$$

where π denotes a pairing of \mathbb{I}_k i.e. a partition of \mathbb{I}_n into pairs $\{i, j\}$, and Π denotes the set of all these pairings; the product here is over all the pairs inside a pairing.

As a result of (2.50), we rewrite (2.49) as:

$$\begin{aligned} & \langle (p_x^2(nt) - \langle p_x^2(nt) \rangle_{\rho^n}) (p_y^2(nt) - \langle p_y^2(nt) \rangle_{\rho^n}) \rangle_{\rho} \\ &= 2\langle \tilde{p}_x(nt) \tilde{p}_y(nt) \rangle_{\rho^n}^2 + 4\bar{p}_x(nt)\bar{p}_y(nt)\langle \tilde{p}_x(nt) \tilde{p}_y(nt) \rangle_{\rho^n}. \end{aligned} \quad (2.51)$$

Before proceeding, let us recall the estimate (3.6) from [3]: There exists a constant $C > 0$ independent of n and the realization of the masses such that, for any n and $x \in \mathbb{I}_n$:

$$|\langle p_x(nt) \rangle_{\rho^n}| \leq C, \quad |\langle r_x(nt) \rangle_{\rho^n}| \leq C. \quad (2.52)$$

Furthermore, following the exact same line of (2.43) (with only replacing $\tilde{p}_x^\bullet(nt)$ with $\tilde{p}_x(nt)$ and summing over \mathbb{I}_n^o instead of $I(\gamma)$), it is evident that there exist $C > 0$ independent of n and realization of the masses such that for any n and $x, y \in \mathbb{I}_n$, we have (in fact one can take $C = \beta_-^{-1}$):

$$\langle \frac{\tilde{p}_x^2(nt)}{m_x} \rangle_{\rho^n} \leq C, \quad \langle \tilde{r}_x^2(nt) \rangle_{\rho^n} \leq C \implies \left| \frac{\langle \tilde{p}_x(nt) \tilde{p}_y(nt) \rangle_{\rho^n}}{\sqrt{m_x m_y}} \right| \leq C, \quad (2.53)$$

where we used a Cauchy-Schwartz inequality.

Recall \mathfrak{P}_n from (2.48). Expanding the square, and using the identity (2.51) we have, with $f_x := f(\frac{x}{n})$,

$$\begin{aligned} 0 \leq \mathfrak{P}_n &= \frac{2}{n^2} \sum_{x,y=1}^n f_x f_y \left(\frac{\langle \tilde{p}_x(nt) \tilde{p}_y(nt) \rangle_{\rho^n}^2}{m_x m_y} + 2 \frac{\bar{p}_x(nt) \bar{p}_y(nt) \langle \tilde{p}_x(nt) \tilde{p}_y(nt) \rangle_{\rho^n}}{m_x m_y} \right) \leq \\ &= \frac{2c_1}{n^2} \sum_{x,y=1}^n \left| \frac{\langle \tilde{p}_x(nt) \tilde{p}_y(nt) \rangle_{\rho^n}}{\sqrt{m_x m_y}} \right| \left(\left| \frac{\langle \tilde{p}_x(nt) \tilde{p}_y(nt) \rangle_{\rho^n}}{\sqrt{m_x m_y}} \right| + 2 \left| \frac{\bar{p}_x(nt) \bar{p}_y(nt)}{\sqrt{m_x m_y}} \right| \right) \leq \underbrace{\frac{c_2}{n^2} \sum_{x,y=1}^n \left| \frac{\langle \tilde{p}_x(nt) \tilde{p}_y(nt) \rangle_{\rho^n}}{\sqrt{m_x m_y}} \right|}_{=:\mathfrak{P}'_n}, \end{aligned} \quad (2.54)$$

where c_1, c_2 are independent of n and realization of the masses. In the first inequality we bounded $|f(y)f(y')| \leq c_1$, since f is bounded, and in the second inequality we take advantage of (2.53) to bound the first term, and (2.52) as well as the fact that distribution of the masses is compactly supported in $[m_-, m_+]$ to bound the second term.

Gathering a couple of results from Sec. 2.3, it is straightforward to observe that $\mathfrak{P}'_n \rightarrow 0$ almost surely, where \mathfrak{P}'_n is defined in the last line of (2.54): Recall $\tilde{p}_x^\bullet(nt)$, $\tilde{p}_x^o(nt)$ (2.31), (2.25), where $\tilde{p}_x(nt) = \tilde{p}_x^\bullet(nt) + \tilde{p}_x^o(nt)$ then we have:

$$\begin{aligned} 0 \leq \mathfrak{P}'_n &\leq \underbrace{\frac{c_2}{n^2} \sum_{x,y=1}^n \left| \frac{\langle \tilde{p}_x^o(nt) \tilde{p}_y^o(nt) \rangle_{\rho^n}}{\sqrt{m_x m_y}} \right|}_{:=\mathfrak{P}_n^1} + \underbrace{\frac{c_2}{n^2} \sum_{x,y=1}^n \left| \frac{\langle \tilde{p}_x^\bullet(nt) \tilde{p}_y^\bullet(nt) \rangle_{\rho^n}}{\sqrt{m_x m_y}} \right|}_{:=\mathfrak{P}_n^2} + \\ &\quad \underbrace{\frac{c_2}{n^2} \sum_{x,y=1}^n \left(\left| \frac{\langle \tilde{p}_x^\bullet(nt) \tilde{p}_y^o(nt) \rangle_{\rho^n}}{\sqrt{m_x m_y}} \right| + \left| \frac{\langle \tilde{p}_x^o(nt) \tilde{p}_y^\bullet(nt) \rangle_{\rho^n}}{\sqrt{m_x m_y}} \right| \right)}_{:=\mathfrak{P}_n^3} \end{aligned} \quad (2.55)$$

Observe the definition of $\mathfrak{P}_n^1, \mathfrak{P}_n^2, \mathfrak{P}_n^3$ in the above expression. $\mathfrak{P}_n^1 \rightarrow 0$ as we proved in (2.30). $\mathfrak{P}_n^2 \leq c'(\mathcal{U}_n^> + \mathcal{U}_n^<)$ thanks to (2.33). However, we observed that $\mathcal{U}_n^> \rightarrow 0$ almost surely w.r.t the distribution of the masses after (2.40). Moreover, we showed $\mathcal{U}_n^< \rightarrow 0$ in (2.44). Hence, $\mathfrak{P}_n^2 \rightarrow 0$ almost surely w.r.t the distribution of the masses. Therefore, it only remains to prove that $\mathfrak{P}_n^3 \rightarrow 0$. Thanks to Cauchy-Schwartz inequality ($|\langle AB \rangle| \leq \langle A^2 \rangle^{\frac{1}{2}} \langle B^2 \rangle^{\frac{1}{2}}$) we have:

$$0 \leq \mathfrak{P}_n^3 \leq 2c_2 \left(\frac{1}{n} \sum_{x=1}^n \frac{\langle (\tilde{p}_x^o(nt))^2 \rangle_{\rho^n}^{\frac{1}{2}}}{\sqrt{m_x}} \right) \left(\frac{1}{n} \sum_{y=1}^n \frac{\langle (\tilde{p}_y^\bullet(nt))^2 \rangle_{\rho^n}^{\frac{1}{2}}}{\sqrt{m_y}} \right) \leq C' \underbrace{\left(\frac{1}{n} \sum_{x=1}^n \frac{\langle (\tilde{p}_x^o(nt))^2 \rangle_{\rho^n}^{\frac{1}{2}}}{\sqrt{m_x}} \right)}_{:=\mathcal{L}_n'''} \rightarrow 0, \quad (2.56)$$

where we bounded the second sum thanks to the bound $\langle \frac{(\tilde{p}_x^\bullet(nt))^2}{m_x} \rangle_{\rho^n} \leq C$ (2.43); moreover, the last limit is deduced from (2.30), where we proved $(\mathcal{L}_n''')^2 \rightarrow 0$ (\mathcal{L}_n''' is positive). Therefore, we conclude that $\mathfrak{P}'_n \rightarrow 0$ almost surely thanks to (2.55), and this finishes the proof of (2.48) for \mathfrak{P}_n thanks to (2.54). As we mentioned the proof of the fact that $\mathfrak{R}_n \rightarrow 0$ almost surely w.r.t distribution of the masses is exactly similar, where, one should take into account the corresponding bound for r in (2.52), (2.53). This finish the proof of (2.48) which finishes the proof of (2.12) and Theorem 2.1. \square

3 Quantum Case

3.1 Model and Results

The quantum disordered harmonic chain of size n is defined as follows: Denote the space variable by $\xi \in \mathbb{R}^{n-1}$, let the state space be given by the Hilbert space $\mathcal{H}_n := L^2(\mathbb{R}^{n-1}, d\xi)$, denote the elements of \mathcal{H}_n by bold ket notation $|\psi\rangle$, and denote the inner product in \mathcal{H}_n by $\langle\phi|\psi\rangle$. Denote also the Schwartz space by $\mathcal{S}(\mathbb{R}^{n-1}; \mathbb{C})$, which is dense in \mathcal{H}_n . The elongation operator r_x for any $x \in \mathbb{I}_{n-1}$ is defined on $\mathcal{S}(\mathbb{R}^{n-1}, \mathbb{C})$ as follows: for any $|\psi\rangle \in \mathcal{S}(\mathbb{R}^{n-1}; \mathbb{C})$

$$r_x|\psi(\xi_1, \dots, \xi_x, \dots, \xi_{n-1})\rangle = \xi_x|\psi(\xi_1, \dots, \xi_x, \dots, \xi_{n-1})\rangle,$$

Correspondingly, for $x \in \mathbb{I}_n$, p_x denotes the momentum operator of particle x , and it is defined as follows: for any $|\psi\rangle \in \mathcal{S}(\mathbb{R}^{n-1}; \mathbb{C})$

$$p_x|\psi(\xi_1, \dots, \xi_x, \dots, \xi_{n-1})\rangle = -i\left(\frac{\partial}{\partial \xi_{x-1}} - \frac{\partial}{\partial \xi_x}\right)|\psi(\xi_1, \dots, \xi_x, \dots, \xi_{n-1})\rangle.$$

We assume free boundary conditions $r_0 = r_n = 0$, which means $p_1 = i\partial/\partial \xi_1$, and $p_n = -i\partial/\partial \xi_{n-1}$ ($\partial/\partial \xi_0 = \partial/\partial \xi_n = 0$, by convention). This means that $\sum_{x=1}^n p_x = 0$, corresponding to the fact that we define our model from a center of mass observer's viewpoint, without loss of generality. The canonical commutation relations (CCR) read $([a, b] := ab - ba)$:

$$[r_x, r_y] = [p_x, p_y] = 0, \quad [r_x, p_y] = i(\delta_{x, (y-1)} - \delta_{x, y}), \quad \forall x \in \mathbb{I}_{n-1}, y \in \mathbb{I}_n. \quad (3.1)$$

The Hamiltonian operator \mathfrak{H}_n is defined on $\mathcal{S}(\mathbb{R}^{n-1}; \mathbb{C})$ as

$$\mathfrak{H}_n = \frac{1}{2} \sum_{x=1}^n \left(\frac{p_x^2}{m_x} + r_x^2 \right) =: \sum_{x=1}^n e_x, \quad (3.2)$$

where $\{m_x\}_{x=1}^\infty$ are the i.i.d random variables defined after (2.3). The operators p_x , r_x , and consequently e_x and \mathfrak{H}_n are essentially self-adjoint on \mathcal{H}_n (cf. [19], [17]). Therefore, we consider their closure on \mathcal{H}_n , that we denote with the same symbols. The dense domain of \mathfrak{H}_n is denoted by $\mathcal{D}(\mathfrak{H}_n)$.

The chain evolves in time according to the Heisenberg dynamics generated by \mathfrak{H}_n : For any $t \in \mathbb{R}$, $e^{it\mathfrak{H}_n}$ is well define thanks to the spectral theorem, since \mathfrak{H}_n is self adjoint. Therefore, using Stone's theorem we can define the one-parameter group of authomorphism τ_t^n , on $\mathcal{B}(\mathcal{H}_n)$ (set of bounded operator on \mathcal{H}_n) as

$$a(t) := \tau_t^n(a) = e^{it\mathfrak{H}_n} a e^{-it\mathfrak{H}_n}, \quad a \in \mathcal{B}(\mathcal{H}_n).$$

Notice that $a(t)$ is the solution to the Heisenberg equation

$$\dot{a} = i[\mathfrak{H}_n, a], \quad a(0) = a_0, \quad (3.3)$$

where $\dot{a} = \partial_t a(t)$. This equation holds in the strong sense on the proper domain. Appealing to Stone's theorem, we can extend the domain of τ_t^n to certain unbounded operators such as r_x , p_x , e_x (cf. [5][6][17][12][19]). Denoting $\tau_t^n(r)$, $\tau_t^n(p)$, $\tau_t^n(e)$ by $r_x(t)$, $p_x(t)$, $e_x(t)$

respectively, one can observe that, thanks to the CCR (3.1), $\mathbf{r}_x(t)$ and $\mathbf{p}_x(t)$ are the strong solutions to the Heisenberg equations

$$\begin{aligned}\dot{\mathbf{p}}_x &= i[\mathfrak{H}_n, \mathbf{p}_x] = (\nabla_- \mathbf{r})_x = \mathbf{r}_x(t) - \mathbf{r}_{x-1}(t), \quad x \in \mathbb{I}_n, \\ \dot{\mathbf{r}}_x &= i[\mathfrak{H}_n, \mathbf{r}_x] = (\nabla_+ M^{-1} \mathbf{p})_x = \frac{\mathbf{p}_{x+1}}{m_{x+1}} - \frac{\mathbf{p}_x}{m_x}, \quad x \in \mathbb{I}_n,\end{aligned}\tag{3.4}$$

on a proper domain, where ∇_- , ∇_+ and M have been defined in (2.1) and below (2.5). Notice that \mathbf{r} , \mathbf{p} in (3.4) should be understood as vectors of elongation and momentum operators (and by abusing the notation matrix multiplication $\nabla_- \mathbf{r}$ should be understood as multiplication of a finite dimensional matrix and a vector of operators). Let us emphasize that since our system is quadratic, the evolution equation for \mathbf{r} and \mathbf{p} in the quantum case (3.4) is similar to the classical case (2.5).

Given initial smooth profiles $\bar{p}, \bar{r} \in C^1([0, 1])$, such that $\bar{r}(0) = \bar{r}(1) = 0$, $\int_0^1 \bar{p}(y) dy = 0$, and $\bar{b} \in C^0([0, 1])$ with $0 < b_- < \bar{b}(y) < b_+$, we define the state (density matrix) $\varrho_n^{\bar{r}, \bar{p}, \bar{b}} \in \mathcal{B}(\mathcal{H}_n)$ such that for each n , $\varrho_n^{\bar{r}, \bar{p}, \bar{b}}$ is trace-class and bounded with $\text{Tr}(\varrho_n^{\bar{r}, \bar{p}, \bar{b}}) = 1$ (as in the classical case, superscripts will be omitted whenever possible). In addition, for any operator a such that $a\varrho_n^{\bar{r}, \bar{p}, \bar{b}}$ be trace-class we define:

$$\langle a \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} := \text{Tr}(\varrho_n^{\bar{r}, \bar{p}, \bar{b}} a).\tag{3.5}$$

Finally we impose that $\varrho_n^{\bar{r}, \bar{p}, \bar{b}}$ represents a local equilibrium state, by imposing the following set of conditions:

Assumptions on $\varrho_n^{\bar{r}, \bar{p}, \bar{b}}$:

1. Local mechanical equilibrium: For any $y \in [0, 1]$,

$$\langle \mathbf{r}_{[ny]} \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} = \bar{r}\left(\frac{[ny]}{n}\right), \quad \frac{1}{m_{[ny]}} \langle \mathbf{p}_{[ny]} \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} = \frac{1}{\bar{m}} \bar{p}\left(\frac{[ny]}{n}\right) + \epsilon_n^{[ny]},\tag{3.6}$$

where for all $y \in [0, 1]$, $\epsilon_n^{[ny]} \rightarrow 0$ as $n \rightarrow \infty$ almost surely w.r.t the distribution of the masses.

2. Local thermal equilibrium: Let

$$\tilde{\mathbf{p}}_x := \mathbf{p}_x - \langle \mathbf{p}_x \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}, \quad \tilde{\mathbf{r}}_x := \mathbf{r}_x - \langle \mathbf{r}_x \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}, \quad \check{\mathbf{e}}_x = \frac{1}{2} \left(\frac{\tilde{\mathbf{p}}_x^2}{m_x} + \tilde{\mathbf{r}}_x^2 \right).\tag{3.7}$$

For any test function $f \in C^0([0, 1])$,

$$\frac{1}{n} \sum_{x=1}^n f\left(\frac{x}{n}\right) \langle \check{\mathbf{e}}_x \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \rightarrow \int_0^1 f(y) \bar{b}(y) dy,\tag{3.8}$$

almost surely w.r.t to the distribution of the masses. Contrary to the classical case, $\bar{b}(y)$ cannot longer be identified with a local temperature $\beta^{-1}(y)$, cf. (3.16) below.

3. Polynomial clustering: For $x \in \mathbb{I}_n$, let us use the notations $O_x^r := \tilde{r}_x$ and $O_x^p := \tilde{p}_x$, see (3.7). There exists $C > 0$ such that for any n , any $x_1, x_2, x_3, x_4 \in \mathbb{I}_n$, any $\sharp_1, \sharp_2, \sharp_3, \sharp_4 \in \{r, p\}$, and any realization of the masses, we have:

$$\begin{aligned} \left| \langle O_{x_1}^{\sharp_1} O_{x_2}^{\sharp_2} \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right| &\leq \frac{C}{|x_1 - x_2|^{a_1}}, \quad \text{with } a_1 \geq 2, \\ \left| \langle O_{x_1}^{\sharp_1} O_{x_2}^{\sharp_2} O_{x_3}^{\sharp_3} \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right| &\leq \frac{C}{|\max\{|x_2 - x_1|, |x_3 - x_2|\}|^{a_2}}, \quad \text{with } a_2 \geq 3, \\ \left| \langle O_{x_1}^{\sharp_1} O_{x_2}^{\sharp_2} O_{x_3}^{\sharp_3} O_{x_4}^{\sharp_4} \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} - \langle O_{x_1}^{\sharp_1} O_{x_2}^{\sharp_2} \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \langle O_{x_3}^{\sharp_3} O_{x_4}^{\sharp_4} \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right| &\leq \\ \frac{C}{|x_2 - x_3|^{a_3} |x_1 - x_4|^{a_3}} + \frac{C}{|x_1 - x_3|^{a_3} |x_2 - x_4|^{a_3}}, &\quad \text{with } a_3 \geq 2 \end{aligned} \quad (3.9)$$

with the convention $\frac{1}{0} = 1$.

Initially, we prepare the chain in the state $\varrho_n^{\bar{r}, \bar{p}, \bar{b}}$ having the above properties. Let us emphasize that this state could be either a mixed state or a pure state i.e., $\varrho_n^{\bar{r}, \bar{p}, \bar{b}} = |\psi\rangle\langle\psi|$ for a suitable wave function $\psi \in \mathcal{H}_n$.

The evolution at the macroscopic level is still given by (2.8), where one should modify the initial datum (2.9) by replacing $\beta^{-1}(y)$ with $\bar{b}(y)$ featuring in (3.8):

$$\begin{aligned} \mathbf{r}(y, 0) &= \bar{r}(y), \quad \mathbf{p}(y, 0) = \bar{p}(y), \quad \mathbf{e}(y, 0) = \frac{\bar{p}^2(y)}{2\bar{m}} + \frac{\bar{r}^2(y)}{2} + \bar{b}(y), \\ \mathbf{r}(0, t) &= \mathbf{r}(1, t) = 0 \quad \forall t \in [0, T]. \end{aligned} \quad (3.10)$$

We prove the following result:

Theorem 3.1. *Let $f \in C^0([0, 1])$, and fix $T > 0$. We assume that the initial state $\varrho_n^{\bar{r}, \bar{p}, \bar{b}}$, parametrized by the smooth profiles $\bar{p}, \bar{r}, \bar{b}$ satisfies the conditions (3.6) to (3.9). Let $(\mathbf{r}(nt), \mathbf{p}(nt), \mathbf{e}(nt))$ be the vector of evolved operators according to the Heisenberg evolution (3.4). In addition, let $(\mathbf{r}(y, t), \mathbf{p}(y, t), \mathbf{e}(y, t))$ be solutions to the macroscopic evolution equations (2.8) with initial datum (3.10). Then for any $t \in [0, T]$,*

$$\left\langle \left(\frac{1}{n} \sum_{x=1}^n f\left(\frac{x}{n}\right) \mathbf{r}_x(nt) - \int_0^1 f(y) \mathbf{r}(y, t) dy \right)^2 \right\rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \rightarrow 0, \quad (3.11)$$

$$\left\langle \left(\frac{1}{n} \sum_{x=1}^n f\left(\frac{x}{n}\right) \mathbf{p}_x(nt) - \int_0^1 f(y) \mathbf{p}(y, t) dy \right)^2 \right\rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}^n}} \rightarrow 0, \quad (3.12)$$

$$\left\langle \left(\frac{1}{n} \sum_{x=1}^n f\left(\frac{x}{n}\right) \mathbf{e}_x(nt) - \int_0^1 f(y) \mathbf{e}(y, t) dy \right)^2 \right\rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}^n}} \rightarrow 0, \quad (3.13)$$

as $n \rightarrow \infty$ almost surely w.r.t the distribution of the masses.

In order to make the above theorem more concrete, let us give an example of state that satisfies conditions (3.6) to (3.9) stated above. Given momentum and elongation profiles \bar{p}

and $\bar{r} \in C^1([0, 1])$ as before, and a profile $\beta \in C^0([0, 1])$ such that $0 < \beta_- \leq \beta(y) \leq \beta_+$, we define the *locally Gibbs state* as the density operator acting on \mathcal{H}_n given by

$$\varrho_{\bar{r}, \bar{p}, \beta}^n = \frac{1}{Z_n} \exp\left(-\frac{1}{2} \sum_{x=1}^n \left[\frac{\beta(\frac{x}{n})}{m_x} (\mathbf{p}_x - \bar{p}(\frac{x}{n}) \frac{m_x}{\bar{m}})^2 + \beta(\frac{x}{n}) (\mathbf{r}_x - \bar{r}(\frac{x}{n}))^2 \right]\right) =: \frac{1}{Z_n} \exp(-H_\beta^n), \quad (3.14)$$

with Z_n a normalizing constant such that $\text{Tr}(\varrho_{\bar{r}, \bar{p}, \beta}^n) = 1$. The operator H_β^n implicitly defined by (3.14) is essentially self-adjoint on $\mathcal{S}(\mathbb{R}^{n-1}, \mathbb{C})$. Its closure is self-adjoint and we denote it with the same symbol. Since H_β^n has a discrete spectrum with non-negative eigenvalues, one can observe that $\varrho_{\bar{r}, \bar{p}, \beta}^n$ is a well-defined density operator (bounded and trace-class, cf. [12]). Let us denote the average with respect to $\varrho_{\bar{r}, \bar{p}, \beta}^n$ by

$$\langle a \rangle_\beta := \text{Tr}(\varrho_{\bar{r}, \bar{p}, \beta}^n a) \quad (3.15)$$

for any suitable operator a . Thanks to the properties of $\varrho_{\bar{r}, \bar{p}, \beta}^n$, it is straightforward to observe that all the averages appearing subsequently are well defined and bounded (cf. [12]). Given the distribution of the masses μ , and temperature profile β , we define the macroscopic thermal energy profile as follows:

$$\mathfrak{f}_\beta^\mu(y) := \lim_{n \rightarrow \infty} \mathbb{E}(\langle \check{\mathfrak{e}}_{[ny]} \rangle_\beta). \quad (3.16)$$

with $\check{\mathfrak{e}}_x$ defined in (3.7). In [12], it is proven that \mathfrak{f}_β^μ is well-defined and continuous, and $\mathfrak{f}_\beta^\mu(y) = \mathfrak{f}^\mu(\beta(y))$, where $\mathfrak{f}^\mu(\beta_{eq})$ is the corresponding function for a constant profile of temperature $\beta(y) = \beta_{eq}$, $\forall y$ (cf. [12] Appendix A).

Theorem 3.2. *The state $\varrho_{\bar{r}, \bar{p}, \beta}^n$ defined in (3.14) satisfies conditions (3.6) to (3.9) with $\bar{p}, \bar{r}, \bar{b}(y) = \mathfrak{f}_\beta^\mu(y)$.*

Before proceeding, let us briefly mention that in the assumption (3.9), the estimates need to hold for all realizations of the masses. However, it is possible to show that they could be replaced by estimates on the expectations (w.r.t μ) of the same quantities, at the cost of stronger decay rates.

3.2 Evolution of r, p

3.2.1 Time evolution

Since our system is harmonic, equation of motion (3.4) can be solved similarly to the classical case (2.17). Let us briefly recall the desired transformations in this case from [12], mainly to fix proper notations. Recall the random matrices A_p , and A_r (2.14), and their corresponding set of eigenvalues and eigenvectors ($0 < \omega_1^2 < \dots < \omega_{n-1}^2$ and $\{\varphi^k\}_{k=0}^n$ for the former and $\omega_1^2 < \dots < \omega_{n-1}^2$ and $\{\phi^k\}_{k=1}^n$ for the latter). Define the following set of operators (recall that $\langle \cdot, \cdot \rangle_n$ is the inner product in \mathbb{R}^n):

$$\hat{\mathbf{p}}_k := \langle \varphi^k, M^{-\frac{1}{2}} \mathbf{p} \rangle_n = \sum_{x=1}^n \frac{\varphi_x^k}{\sqrt{m_x}} \mathbf{p}_x, \quad \forall k \in \mathbb{I}_{n-1}; \quad \hat{\mathbf{r}}_k := \langle \phi^k, \mathbf{r} \rangle_{n-1} = \sum_{x=1}^{n-1} \phi_x^k \mathbf{r}_x, \quad \forall k \in \mathbb{I}_{n-1}. \quad (3.17)$$

After a straightforward computation (cf. Sect.3 of [12]), we have the following commutation relations for $k, k' \in \mathbb{I}_{n-1}^o$ (by convention $\hat{r}_0 = 0$):

$$[\hat{r}_k, \hat{r}_{k'}] = [\hat{p}_k, \hat{p}_{k'}] = 0, \quad [\hat{r}_k, \hat{p}_{k'}] = i\omega_k \delta_{k,k'}. \quad (3.18)$$

Consequently, we define the corresponding bosonic operators $\hat{\mathbf{b}}_k, \hat{\mathbf{b}}_k^*$, having suitable commutation relations (thanks to (3.18)) as follows: for any $k, k' \in \mathbb{I}_{n-1}$:

$$\begin{aligned} \hat{\mathbf{b}}_k &:= \frac{1}{\sqrt{2\omega_k}}(\hat{r}_k + i\hat{p}_k), \quad \hat{\mathbf{b}}_k^* := \frac{1}{\sqrt{2\omega_k}}(\hat{r}_k - i\hat{p}_k), \\ [\hat{\mathbf{b}}_k, \hat{\mathbf{b}}_{k'}^*] &= \delta_{k,k'}, \quad [\hat{\mathbf{b}}_k, \hat{\mathbf{b}}_{k'}] = [\hat{\mathbf{b}}_k^*, \hat{\mathbf{b}}_{k'}^*] = 0. \end{aligned} \quad (3.19)$$

Thanks to (3.17) and (3.19), and by using the definition of φ^k, ϕ^k , we can rewrite the Hamiltonian \mathfrak{H}_n in terms of bosonic operators:

$$\mathfrak{H}_n = \frac{1}{2} \sum_{x=1}^n \left(\frac{\mathbf{p}_x^2}{m_x} + r_x^2 \right) = \frac{1}{2} \hat{\mathbf{p}}_0 + \frac{1}{2} \sum_{k=1}^{n-1} (\hat{\mathbf{p}}_k^2 + \hat{r}_k^2) = \frac{1}{2} \sum_{k=1}^{n-1} (\hat{\mathbf{p}}_k^2 + \hat{r}_k^2) = \sum_{k=1}^{n-1} \omega_k (\hat{\mathbf{b}}_k^* \hat{\mathbf{b}}_k + \frac{1}{2}), \quad (3.20)$$

where we used the fact that $\mathbf{p}_0 = \sum_{x=1}^n \mathbf{p}_x = 0$. Notice that thanks to the above expression, one could obtain the spectrum of \mathfrak{H}_n explicitly, which is a discrete spectrum (cf. Sect.3 of [12]). In addition, thanks to the above representation of \mathfrak{H}_n , and commutation relations (3.19), the time evolution of the bosonic operators (action of τ_t^n i.e., solution to the Heisenberg equation (3.3)) is given by:

$$\hat{\mathbf{b}}_k(t) = e^{-i\omega_k t} \hat{\mathbf{b}}_k(0), \quad \hat{\mathbf{b}}_k^*(t) = e^{i\omega_k t} \hat{\mathbf{b}}_k^*(0). \quad (3.21)$$

By using the inverse of (3.19), we get for $k \in \mathbb{I}_{n-1}$:

$$\hat{\mathbf{p}}_k(t) = \cos(\omega_k t) \hat{\mathbf{p}}_k(0) - \sin(\omega_k t) \hat{r}_k(0), \quad \hat{r}_k(t) = \cos(\omega_k t) \hat{r}_k(0) + \sin(\omega_k t) \hat{\mathbf{p}}_k(0) \quad (3.22)$$

Plugging the latter inside the inverse of (3.17), the time evolution of p_x, r_x is given by (recall the definition of $\hat{\mathbf{p}}, \hat{r}$ (3.17)):

$$\begin{aligned} \mathbf{p}(t) &= \sum_{k=0}^{n-1} M^{\frac{1}{2}} \varphi^k \hat{\mathbf{p}}_k(t) = \sum_{k=0}^{n-1} (\cos(\omega_k t) \hat{\mathbf{p}}_k(0) - \sin(\omega_k t) \hat{r}_k(0)) M^{\frac{1}{2}} \varphi^k, \\ \mathbf{r}(t) &= \sum_{k=1}^{n-1} \phi^k \hat{r}_k(t) = \sum_{k=1}^{n-1} (\cos(\omega_k t) \hat{r}_k(0) + \sin(\omega_k t) \hat{\mathbf{p}}_k(0)) \phi^k. \end{aligned} \quad (3.23)$$

These equations are formally similar to their classical counterpart (2.17), but \mathbf{r} and \mathbf{p} are (non-commuting) operators rather than numbers.

3.2.2 Proof for r, p

In this section, we prove the first two limits in Theorem 3.1, i.e. (3.11) and (3.12). Similarly to the classical case we use the explicit time evolution (3.23), as well as the localization estimates (2.18). However, our state is no longer Gaussian and simple correlation identities (2.27), (2.34) need to be replaced by (3.9).

Proof of (3.11) and (3.12). Recall that the state $\varrho_n^{\bar{r}, \bar{p}, \bar{b}}$ satisfies assumptions (3.6) to (3.9). Let us define the *average* variables $\bar{\mathbf{p}}_x(nt)$, $\bar{\mathbf{r}}_x(nt)$, and *fluctuation* operators $\tilde{\mathbf{p}}_x(nt)$, $\tilde{\mathbf{r}}_x(nt)$ as follows for any $x \in \mathbb{I}_n$:

$$\begin{aligned}\bar{\mathbf{p}}_x(nt) &:= \langle \mathbf{p}_x(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}, & \bar{\mathbf{r}}_x(nt) &:= \langle \mathbf{r}_x(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}; \\ \tilde{\mathbf{p}}_x(nt) &:= \mathbf{p}_x(nt) - \bar{\mathbf{p}}_x(nt), & \tilde{\mathbf{r}}_x(nt) &:= \mathbf{r}_x(nt) - \bar{\mathbf{r}}_x(nt).\end{aligned}\tag{3.24}$$

Thanks to the linearity of the trace, and the evolution equation (3.4), $(\bar{\mathbf{p}}(t), \bar{\mathbf{r}}(t)) \in \mathbb{R}^{2n-1}$ satisfies the following system of ODEs:

$$\frac{d}{dt}\bar{\mathbf{p}}_x(t) = \bar{\mathbf{r}}_x(t) - \bar{\mathbf{r}}_{x-1}(t), \quad \frac{d}{dt}\bar{\mathbf{r}}_x(t) = \frac{\bar{\mathbf{p}}_{x+1}(t)}{m_{x+1}} - \frac{\bar{\mathbf{p}}_x(t)}{m_x},\tag{3.25}$$

where the initial data are given by $\bar{\mathbf{r}}_x(0) = \bar{\mathbf{r}}(\frac{x}{n})$, and $\bar{\mathbf{p}}_x(0) = \frac{m_x}{\bar{m}}\bar{\mathbf{p}}(\frac{x}{n}) + \epsilon_n^x$ for all x , thanks to (3.6). In [12], it is shown that

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{x=1}^n \bar{\mathbf{p}}_x(nt) f\left(\frac{x}{n}\right) \rightarrow \int_0^1 \mathbf{p}(y, t) f(y) dy, \quad \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{x=1}^n \bar{\mathbf{r}}_x(nt) f\left(\frac{x}{n}\right) \rightarrow \int_0^1 \mathbf{r}(y, t) f(y) dy,\tag{3.26}$$

i.e. the limit on average (1.3). The initial conditions in [12] are not the same as here, but the difference vanishes in the limit. As in the classical case, we can add and subtract $\frac{1}{n} \sum_{x=1}^n f(\frac{x}{n}) \langle \mathbf{p}_x(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}$ inside the square in (3.12), and similarly for \mathbf{r} . Then by using the basic properties of the trace (linearity and scalar multiplication), and limits (3.26), the proof of (3.11) and (3.12) boils down to proving that

$$\mathbf{R}_n := \left\langle \left(\frac{1}{n} \sum_{x=1}^n f\left(\frac{x}{n}\right) \tilde{\mathbf{r}}_x(nt) \right)^2 \right\rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \rightarrow 0, \quad \mathbf{P}_n := \left\langle \left(\frac{1}{n} \sum_{x=1}^n f\left(\frac{x}{n}\right) \tilde{\mathbf{p}}_x(nt) \right)^2 \right\rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \rightarrow 0,\tag{3.27}$$

as $n \rightarrow \infty$, almost surely w.r.t the distribution of the masses, where one should recall the fluctuation operators $\tilde{\mathbf{p}}_x, \tilde{\mathbf{r}}_x$ (3.24).

The evolution of $\tilde{\mathbf{p}}(nt)$ and $\tilde{\mathbf{r}}(nt)$ can be obtained explicitly from (3.4) and (3.25). For $k \in \mathbb{I}_n^o$, let

$$\hat{\tilde{\mathbf{p}}}_k = \hat{\tilde{\mathbf{p}}}_k(0) := \langle \varphi^k, M^{-\frac{1}{2}} \tilde{\mathbf{p}} \rangle_n, \quad \hat{\tilde{\mathbf{r}}}_k := \langle \phi^k, \tilde{\mathbf{r}} \rangle_{n-1},\tag{3.28}$$

where the vectors $\tilde{\mathbf{p}}$ and $\tilde{\mathbf{r}}$ are defined in (3.7), (3.24). Then $\tilde{\mathbf{r}}_x(t)$, $\tilde{\mathbf{p}}_x(t)$ is given by:

$$\begin{aligned}\tilde{\mathbf{r}}_x(t) &= \sum_{k=1}^{n-1} \phi_x^k \hat{\tilde{\mathbf{r}}}_k(t) = \sum_{k=1}^{n-1} \left(\cos(\omega_k t) \hat{\tilde{\mathbf{r}}}_k(0) + \sin(\omega_k t) \hat{\tilde{\mathbf{p}}}_k(0) \right) \phi_x^k, \\ \tilde{\mathbf{p}}_x(t) &= \sum_{k=0}^{n-1} \sqrt{m_x} \varphi_x^k \hat{\tilde{\mathbf{p}}}_k(t) = \sum_{k=0}^{n-1} \left(\cos(\omega_k t) \hat{\tilde{\mathbf{p}}}_k(0) - \sin(\omega_k t) \hat{\tilde{\mathbf{r}}}_k(0) \right) \sqrt{m_x} \varphi_x^k,\end{aligned}\tag{3.29}$$

where $\hat{\tilde{\mathbf{r}}}_k(t)$, and $\hat{\tilde{\mathbf{p}}}_k(t)$ are implicitly defined.

Before proceeding, we need bounds similar to (2.29) in the quantum case. This can be done thanks to the first bound in (3.9):

$$\begin{aligned} \langle \hat{\mathbf{p}}_k^2 \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} &= \sum_{x,y=1}^n \frac{\varphi_x^k \varphi_y^k}{\sqrt{m_x m_y}} \langle \tilde{\mathbf{p}}_x \tilde{\mathbf{p}}_y \rangle \leq \sum_{x=1}^n \frac{(\varphi_x^k)^2}{m_x} \langle \tilde{\mathbf{p}}_x^2 \rangle + \sum_{l=1}^{n-1} \sum_{|x-y|=l} \left| \frac{\varphi_x^k \varphi_y^k}{\sqrt{m_x m_y}} \right| |\langle \tilde{\mathbf{p}}_x \tilde{\mathbf{p}}_y \rangle| \\ &\leq C_1 \sum_{x=1}^n (\varphi_x^k)^2 + \sum_{l=1}^{n-1} \frac{C_2}{l^2} \sum_{|x-y|=l} |\varphi_x^k \varphi_y^k| \leq C_1 + \sum_{l=1}^{n-1} \frac{C_3}{l^2} \leq C, \end{aligned} \quad (3.30)$$

where we have dropped the subscript $\varrho_n^{\bar{r}, \bar{p}, \bar{b}}$ on the r.h.s. To get this, we used the definition of $\hat{\mathbf{p}}_k$ (3.28), we used (3.9) in the second line, we took advantage of the following Cauchy-Schwartz inequality,

$$\sum_{|x-y|=l} |\varphi_x^k \varphi_y^k| = 2 \sum_{x=1}^{n-l} |\varphi_x^k \varphi_{x+l}^k| \leq 2 \left(\sum_{x=1}^{n-l} (\varphi_x^k)^2 \right)^{\frac{1}{2}} \left(\sum_{x=1}^{n-l} (\varphi_{x+l}^k)^2 \right)^{\frac{1}{2}} \leq 2 \sum_{x=1}^n (\varphi_x^k)^2 \leq 2,$$

and finally, in the last inequality of (3.30), we used the fact that the featuring series is summable. Notice that the constant $C > 0$ in (3.30) is independent of n and the realization of the masses. Similarly for r , by an exact identical computation, and using corresponding definition and bounds (3.28), (3.9) we have: there exist $C' > 0$ such that for all n and any realization of the masses we have:

$$\langle \hat{\mathbf{r}}_k^2 \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \leq C'. \quad (3.31)$$

In the proceeding, we follow somewhat similar steps as in the classical setup, where each step is modified due to the quantum nature:

Step1: Low modes. For a proper γ with $0 < \gamma < \frac{1}{2}$, define $\tilde{\mathbf{p}}^o(nt)$, and $\tilde{\mathbf{r}}^o(nt)$ as low modes portion of $\mathbf{p}(nt)$, and $\mathbf{r}(nt)$ (we will choose γ opportunely later): for $x \in \mathbb{I}_n$:

$$\tilde{\mathbf{p}}_x^o(nt) := \sum_{k \in \mathbb{Z} \cap [0, n^{(1-\gamma)}]} \hat{\mathbf{p}}_k(nt) \sqrt{m_x} \varphi_x^k, \quad \tilde{\mathbf{r}}_x^o(nt) := \sum_{k \in \mathbb{Z} \cap [1, n^{(1-\gamma)}]} \hat{\mathbf{r}}_k(nt) \phi_x^k, \quad (3.32)$$

where definition of $\hat{\mathbf{p}}_k(nt)$, $\hat{\mathbf{r}}_k(nt)$ should be recalled from (3.29), and by convention we set $\tilde{\mathbf{r}}_n(nt) = 0$. Here we prove that:

$$\mathbf{L}_n := \left\langle \frac{1}{n} \left(\sum_{x=1}^n f\left(\frac{x}{n}\right) \tilde{\mathbf{p}}_x^o(nt) \right)^2 \right\rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \rightarrow 0, \quad \mathbf{L}'_n := \left\langle \frac{1}{n} \left(\sum_{x=1}^n f\left(\frac{x}{n}\right) \tilde{\mathbf{r}}_x^o(nt) \right)^2 \right\rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \rightarrow 0, \quad (3.33)$$

almost surely w.r.t the distribution of the masses.

Before proceeding, thanks to the above mentioned assumptions, one can check that $(\varrho_n^{\bar{r}, \bar{p}, \bar{b}})^{\frac{1}{2}} a$ is a Hilbert-Schmidt operator for any operator a that is a linear or quadratic function of \mathbf{p}_x and \mathbf{r}_x . Therefore, it is straightforward to see that we can use the following form of Cauchy-Schwartz inequality for our desired operators:

$$\left| \langle ab^* \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right|^2 \leq \langle aa^* \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \langle bb^* \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}. \quad (3.34)$$

Later, we will simply use this inequality, and we will not mention its legitimacy anymore which is easy to check in each case.

Following exactly the same steps as in (2.30) yields:

$$0 \leq \mathbb{L}_n \leq \frac{C}{n} \sum_{k \in \tilde{I}(\gamma)} \langle (\hat{\mathbf{p}}_k(nt))^2 \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}, \quad (3.35)$$

where $C > 0$ is a constant independent of n and the realization of the masses. Notice that the only difference between (3.35) and (2.30) is the fact that in the former we should use the operator form of the Cauchy-Schwartz inequality (3.34). By using the definition of $\hat{\mathbf{p}}_k(nt)$ from (3.29), thanks to (3.30), (3.31) and a Cauchy-Schwartz inequality, we have for any $k \in \mathbb{I}_{n-1}^o$

$$\langle (\hat{\mathbf{p}}_k(nt))^2 \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \leq C, \quad \langle (\hat{\mathbf{r}}_k(nt))^2 \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \leq C, \quad (3.36)$$

where $C > 0$ is independent of n , and the distribution of the masses. Combining (3.36) with (3.32) and recalling the fact that $\tilde{I}(\gamma) = [0, n^{1-\gamma}] \cap \mathbb{Z}$ with $\gamma > 0$ we deduce that $\mathbb{L}_n \rightarrow 0$ as $n \rightarrow \infty$. Again, a similar computation, using the definition of $\hat{\mathbf{r}}^o$, the fact that ϕ^k is an orthonormal basis, and the second bound in (3.36) yield $\mathbb{L}'_n \rightarrow 0$ deterministically. This finishes the proof of this step, i.e. (3.33).

Step2: High modes. As in (2.31), define the “high mode operators” for $x \in \mathbb{I}_n$:

$$\tilde{\mathbf{p}}_x^\bullet(nt) := \tilde{\mathbf{p}}_x(nt) - \tilde{\mathbf{p}}_x^o(nt) = \sum_{k \in I(\gamma)} \hat{\mathbf{p}}_k(nt) \sqrt{m_x} \varphi_x^k, \quad \tilde{\mathbf{r}}_x^\bullet(nt) := \tilde{\mathbf{r}}_x(nt) - \tilde{\mathbf{r}}_x^o(nt) = \sum_{k \in I(\gamma)} \hat{\mathbf{r}}_k(nt) \sqrt{m_x} \phi_x^k, \quad (3.37)$$

where we choose $0 < \gamma < \frac{1}{2}$ opportunely later, and $I(\gamma) =]n^{(1-\gamma)}, n] \cap \mathbb{Z}$. Moreover, by convention we have $\tilde{\mathbf{r}}_n(nt) = 0$. Following the lines of the classical case, for a proper choice of $0 < \gamma < \frac{1}{2}$ we prove:

$$\mathbb{U}_n := \left\langle \left(\frac{1}{n} \sum_{x=1}^n f\left(\frac{x}{n}\right) \tilde{\mathbf{p}}_x^\bullet(nt) \right)^2 \right\rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \rightarrow 0, \quad \mathbb{U}'_n := \left\langle \left(\frac{1}{n} \sum_{x=1}^n f\left(\frac{x}{n}\right) \tilde{\mathbf{r}}_x^\bullet(nt) \right)^2 \right\rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \rightarrow 0, \quad (3.38)$$

almost surely, w.r.t distribution of the masses.

Step 2.1: High modes $|x - y| > 4n^{\theta'}$. Let $0 < 2\gamma < \theta < \theta' < 1$. Fix n , and consider $x, y \in \mathbb{I}_n$ such that $|x - y| > 4n^{\theta'}$. Let us bound $|\langle \tilde{\mathbf{p}}_x^\bullet(nt) \tilde{\mathbf{p}}_y^\bullet(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}| / m_x m_y$ (recall

$$\tilde{\varphi}_x^k = \frac{\varphi_x^k}{\sqrt{m_x}}):$$

$$\begin{aligned} \frac{|\langle \tilde{\mathbf{p}}_x^\bullet(nt) \tilde{\mathbf{p}}_y^\bullet(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}}{m_x m_y} &= \left| \sum_{k, k' \in I(\gamma)} \langle \hat{\mathbf{p}}_k(nt) \hat{\mathbf{p}}_{k'}(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \tilde{\varphi}_x^k \tilde{\varphi}_y^{k'} \right| \leq \\ &\underbrace{\left| \sum_{k, k' \in I(\gamma)} \sin(\omega_k nt) \sin(\omega_{k'} nt) \tilde{\varphi}_x^k \tilde{\varphi}_y^{k'} \langle \hat{\mathbf{r}}_k \hat{\mathbf{r}}_{k'} \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right|}_{=: \mathbf{C}_{xy}^{rr}} + \underbrace{\left| \sum_{k, k' \in I(\gamma)} \cos(\omega_k nt) \sin(\omega_{k'} nt) \tilde{\varphi}_x^k \tilde{\varphi}_y^{k'} \langle \hat{\mathbf{p}}_k \hat{\mathbf{r}}_{k'} \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right|}_{=: \mathbf{C}_{xy}^{pr}} \\ &\underbrace{\left| \sum_{k, k' \in I(\gamma)} \sin(\omega_k nt) \cos(\omega_{k'} nt) \tilde{\varphi}_x^k \tilde{\varphi}_y^{k'} \langle \hat{\mathbf{r}}_k \hat{\mathbf{p}}_{k'} \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right|}_{=: \mathbf{C}_{xy}^{rp}} + \underbrace{\left| \sum_{k, k' \in I(\gamma)} \cos(\omega_k nt) \cos(\omega_{k'} nt) \tilde{\varphi}_x^k \tilde{\varphi}_y^{k'} \langle \hat{\mathbf{p}}_k \hat{\mathbf{p}}_{k'} \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right|}_{=: \mathbf{C}_{xy}^{pp}}, \end{aligned} \quad (3.39)$$

where we used the definition of $\tilde{\mathbf{p}}_x^\bullet(nt)$ (3.37), as well as the definition of $\hat{\mathbf{p}}_k(nt)$ (3.29). Notice the definition of \mathbf{C}_{xy}^{pp} , \mathbf{C}_{xy}^{pr} , \mathbf{C}_{xy}^{rp} , and \mathbf{C}_{xy}^{rr} in the above expression. Let us bound \mathbf{C}_{xy}^{pp} . The other terms can be treated similarly. Define

$$\tilde{J}(x, y) := \{(z, z') \in \mathbb{I}_n \times \mathbb{I}_n \mid |x - z| \leq n^\theta, |y - z'| \leq n^\theta\}.$$

Plugging the definition of $\hat{\mathbf{p}}_k$ in \mathbf{C}_{xy}^{pp} , rearranging the terms, using the linearity of trace, and bounding $|\cos(\cdot)|$ by one we have:

$$\begin{aligned} 0 \leq \mathbf{C}_{xy}^{pp} &= \left| \sum_{z, z'=1}^n \left(\sum_{k \in I(\gamma)} \cos(\omega_k nt) \tilde{\varphi}_x^k \tilde{\varphi}_z^k \right) \left(\sum_{k' \in I(\gamma)} \cos(\omega_{k'} nt) \tilde{\varphi}_y^{k'} \tilde{\varphi}_{z'}^{k'} \right) \langle \tilde{\mathbf{p}}_z \tilde{\mathbf{p}}_{z'} \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right| \leq \\ &\underbrace{\sum_{(z, z') \in \tilde{J}(x, y)} \left(\sum_{k \in I(\gamma)} |\tilde{\varphi}_x^k \tilde{\varphi}_z^k| \right) \left(\sum_{k' \in I(\gamma)} |\tilde{\varphi}_y^{k'} \tilde{\varphi}_{z'}^{k'}| \right) |\langle \tilde{\mathbf{p}}_z \tilde{\mathbf{p}}_{z'} \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}|}_{\mathbf{C}_{xy}^{pp, 1}} + \\ &\underbrace{\sum_{(z, z') \notin \tilde{J}(x, y)} \left(\sum_{k \in I(\gamma)} |\tilde{\varphi}_x^k \tilde{\varphi}_z^k| \right) \left(\sum_{k' \in I(\gamma)} |\tilde{\varphi}_y^{k'} \tilde{\varphi}_{z'}^{k'}| \right) |\langle \tilde{\mathbf{p}}_z \tilde{\mathbf{p}}_{z'} \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}|}_{\mathbf{C}_{xy}^{pp, 2}} \end{aligned} \quad (3.40)$$

Let us first bound $\mathbf{C}_{xy}^{pp, 1}$. Each term involving sum over k and k' is bounded by 1 as we observed in (2.37). Hence we have:

$$0 \leq \mathbf{C}_{xy}^{pp, 1} \leq \sum_{(z, z') \in \tilde{J}(x, y)} \left| \langle \tilde{\mathbf{p}}_z \tilde{\mathbf{p}}_{z'} \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right| \leq \sum_{(z, z') \in \tilde{J}(x, y)} \frac{C}{|z - z'|^2} \leq \sum_{(z, z') \in \tilde{J}(x, y)} \frac{C}{n^{2\theta'}} = |\tilde{J}(x, y)| \frac{C}{n^{\theta'}} \leq \frac{8C}{n^{2(\theta' - \theta)}}, \quad (3.41)$$

where in the third inequality we take advantage of the first bound in (3.9); moreover, in the fourth inequality we use the fact that $|x - y| > 4n^{\theta'}$, and thanks to the definition of

$\tilde{J}(x, y)$, for each term in the above sum $|z - z'| > n^{\theta'}$. In the last inequality the definition of $\tilde{J}(x, y)$ is used.

Now we treat $\mathbb{C}_{xy}^{pp,2}$. Recall the definition of $J(x) = \{z \in \mathbb{I}_n \mid |x - z| \leq n^\theta\}$. Thus for n sufficiently large we get

$$\begin{aligned} 0 \leq \mathbb{E}(\mathbb{C}_{xy}^{pp,2}) &\leq C\mathbb{E}\left(\sum_{z \notin J(x)} \sum_{z'=1}^n \left(\sum_{k \in I(\gamma)} |\tilde{\varphi}_x^k \tilde{\varphi}_z^k|\right) \left(\sum_{k' \in I(\gamma)} |\tilde{\varphi}_y^{k'} \tilde{\varphi}_{z'}^{k'}|\right)\right) + \\ &C\mathbb{E}\left(\sum_{z \in J(x)} \sum_{z' \notin J(y)} \left(\sum_{k \in I(\gamma)} |\tilde{\varphi}_x^k \tilde{\varphi}_z^k|\right) \left(\sum_{k' \in I(\gamma)} |\tilde{\varphi}_y^{k'} \tilde{\varphi}_{z'}^{k'}|\right)\right) \leq C\mathbb{E}\left(\sum_{(z,z') \in J^c(x) \times \mathbb{I}_n} \left(\sum_{k \in I(\gamma)} |\tilde{\varphi}_x^k \tilde{\varphi}_z^k|\right)\right) + \\ &C\mathbb{E}\left(\sum_{(z,z') \in J(x) \times J^c(y)} \left(\sum_{k' \in I(\gamma)} |\tilde{\varphi}_y^{k'} \tilde{\varphi}_{z'}^{k'}|\right)\right) \leq 2n^2 C' \exp(-c(n^{\theta-n^{2\gamma}})) \leq C' \exp(-\frac{c}{2}n^\theta), \end{aligned} \quad (3.42)$$

where first we bounded $|\langle \tilde{\mathbf{p}}_z \tilde{\mathbf{p}}_{z'} \rangle_{\tilde{\varrho}_n^{\bar{r}, \bar{p}, \bar{b}}}| \leq C$, thanks to assumption (3.9), where $C > 0$ is independent of n and the realization of the masses; in the second line we take advantage of (2.37). In the third line we used the properties of $J(x)$, $J(y)$ and the bound (2.18) from Lemma 2.2. Finally, in the last inequality we take advantage of the choice of $0 < 2\gamma < \theta < 1$ and the fact that n is sufficiently large. Notice that $C' > 0$ and $c > 0$ are independent of n .

Thanks to (3.42), for n sufficiently large we have:

$$\mathbb{E}\left(\frac{1}{n^2} \sum_{|x-y| > 4n^{\theta'}} \mathbb{C}_{xy}^{pp,2}\right) \leq C' \exp(-\frac{cn^\theta}{2}) \implies \frac{1}{n^2} \sum_{|x-y| > 4n^{\theta'}} \mathbb{C}_{xy}^{pp,2} \rightarrow 0, \quad (3.43)$$

almost surely w.r.t the distribution of the masses, thanks to Borel-Cantelli lemma (recall that $C > 0$, $c' > 0$ are independent of n). Combining (3.41) and (3.43), using the fact that $\theta' > \theta$, and recalling the expression of \mathbb{C}_{xy}^{pp} (3.40) we deduce:

$$\frac{1}{n^2} \sum_{|x-y| > n^{4\theta'}} \mathbb{C}_{xy}^{pp} \rightarrow 0, \quad (3.44)$$

almost surely w.r.t the distribution of the masses. We can prove similar identities for $\sum \mathbb{C}_{xy}^{pr}$, $\sum \mathbb{C}_{xy}^{pr}$, and $\sum \mathbb{C}_{xy}^{rr}$. To this end first we should decompose them as in (3.40). Then $\sum \mathbb{C}_{xy}^{pr,1}$ (and other similar terms) can be treated exactly similar to (3.41). However, the other terms such as $\sum \mathbb{C}_{xy}^{pr,2}$ need more attention: we should use Lemma 2.3 to treat this terms as we did in (2.39). Since these lengthy computation follow the exact same lines of the previous cases, and they do not contain any novel point, we omit them for the sake of brevity. Therefore, thanks to the (3.39), by bounding $f(\frac{x}{n})$ and m_x by uniform constants, we have:

$$\left| \frac{1}{n^2} \sum_{|x-y| > 4n^{\theta'}} f\left(\frac{x}{n}\right) f\left(\frac{y}{n}\right) \langle \tilde{\mathbf{p}}_x^\bullet(nt) \tilde{\mathbf{p}}_y^\bullet(nt) \rangle_{\tilde{\varrho}_n^{\bar{r}, \bar{p}, \bar{b}}} \right| \leq \frac{C}{n^2} \sum_{|x-y| > 4n^{\theta'}} (\mathbb{C}_{xy}^{pp} + \mathbb{C}_{xy}^{pr} + \mathbb{C}_{xy}^{rp} + \mathbb{C}_{xy}^{rr}) \rightarrow 0, \quad (3.45)$$

almost surely w.r.t the distribution of the masses. Following same steps one can show that

$$\left| \frac{1}{n^2} \sum_{|x-y|>4n^{\theta'}} f\left(\frac{x}{n}\right) f\left(\frac{y}{n}\right) \langle \tilde{\mathbf{r}}_x^\bullet(nt) \tilde{\mathbf{r}}_y^\bullet(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right| \rightarrow 0 \quad (3.46)$$

almost surely w.r.t. the distribution of the masses. Since this proof is identical, we do not bring it here. This finishes the proof of this step.

Step 2.2: High modes $|x - y| \leq 4n^{\theta'}$. Let us bound $\frac{1}{m_x} \langle (\tilde{\mathbf{p}}_x^\bullet(nt))^2 \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}$: recall the expression (3.39), let $x = y$, and bound $(m_x)^{-1}$ by a constant; then we have:

$$\frac{\langle (\tilde{\mathbf{p}}_x^\bullet(nt))^2 \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}}{m_x} \leq m_x (C_{xx}^{pp} + C_{xx}^{pr} + C_{xx}^{rp} + C_{xx}^{rr}), \quad (3.47)$$

where definition of $C_{xx}^{\sharp_1 \sharp_2}$ for $\sharp_1, \sharp_2 \in \{r, p\}$ should be recalled from (3.39). Let us bound $m_x C_{xx}^{pr}$ by a constant uniform in n and realization of the masses: Similar to (3.40), by using the definition of $\hat{\mathbf{p}}_k$, and $\hat{\mathbf{r}}_k$ we get (recall the convention $\tilde{\mathbf{r}}_n = \phi_n^k = 0$)

$$m_x C_{xx}^{pr} = \left| \sum_{z, z'=1}^n \underbrace{\left(\sum_{k \in I(\gamma)} \cos(\omega_k nt) \varphi_x^k \varphi_z^k \right)}_{=: \mathbf{a}_z} \underbrace{\left(\sum_{k' \in I(\gamma)} \sin(\omega_{k'} nt) \varphi_x^{k'} \phi_{z'}^{k'} \right)}_{=: \mathbf{b}_{z'}} \left\langle \frac{\tilde{\mathbf{p}}_z}{\sqrt{m_z}} \tilde{\mathbf{r}}_{z'} \right\rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right| \quad (3.48)$$

Notice the definition of \mathbf{a}_z , \mathbf{b}_z in the above expression. Then we have:

$$\begin{aligned} m_x C_{xx}^{pr} &\leq \sum_{z=1}^n |\mathbf{a}_z| |\mathbf{b}_z| \left| \left\langle \frac{\tilde{\mathbf{p}}_z}{\sqrt{m_z}} \tilde{\mathbf{r}}_{z'} \right\rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right| + \sum_{l=1}^{n-1} \sum_{|z-z'|=l} \left| \left\langle \frac{\tilde{\mathbf{p}}_z}{\sqrt{m_z}} \tilde{\mathbf{r}}_{z'} \right\rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right| |\mathbf{a}_z| |\mathbf{b}_{z'}| \leq \\ &C \sum_{z=1}^n |\mathbf{a}_z| |\mathbf{b}_z| + \sum_{l=1}^{n-1} \frac{C}{l^2} \sum_{z=1}^{n-l} (|\mathbf{a}_z| |\mathbf{b}_{z+l}| + |\mathbf{a}_{z+l}| |\mathbf{b}_z|) \leq C \left(\sum_{z=1}^n |\mathbf{a}_z|^2 \right)^{\frac{1}{2}} \left(\sum_{z=1}^n |\mathbf{b}_z|^2 \right)^{\frac{1}{2}} + \\ &\sum_{l=1}^{n-1} \frac{C}{l} \left(\left(\sum_{z=1}^{n-l} |\mathbf{a}_z|^2 \right)^{\frac{1}{2}} \left(\sum_{z=1}^{n-l} |\mathbf{b}_{z+l}|^2 \right)^{\frac{1}{2}} + \left(\sum_{z=1}^{n-l} |\mathbf{a}_{z+l}|^2 \right)^{\frac{1}{2}} \left(\sum_{z=1}^{n-l} |\mathbf{b}_z|^2 \right)^{\frac{1}{2}} \right) \leq \\ &C \left(\sum_{z=1}^n |\mathbf{a}_z|^2 \right)^{\frac{1}{2}} \left(\sum_{z=1}^n |\mathbf{b}_z|^2 \right)^{\frac{1}{2}} + \sum_{l=1}^{n-1} \frac{2C}{l^2} \left(\sum_{z=1}^n |\mathbf{a}_z|^2 \right)^{\frac{1}{2}} \left(\sum_{z=1}^n |\mathbf{b}_z|^2 \right)^{\frac{1}{2}}, \end{aligned} \quad (3.49)$$

where in the second line we used the assumption (3.9), and then we take advantage of two Cauchy-Schwartz inequality. Recalling the definition of \mathbf{b}_z (3.48) we have (recall the convention $\phi_n^k = 0$):

$$\begin{aligned} \sum_{z=1}^n \mathbf{b}_z^2 &= \sum_{z=1}^n \sum_{k, k' \in I(\gamma)}^{n-1} \sin(\omega_k nt) \sin(\omega_{k'} nt) \varphi_x^k \varphi_x^{k'} \phi_z^k \phi_z^{k'} = \sum_{k, k' \in I(\gamma)} \sin(\omega_k nt) \sin(\omega_{k'} nt) \varphi_x^k \varphi_x^{k'} \sum_{z=1}^{n-1} \phi_z^k \phi_z^{k'} = \\ &\sum_{k \in I(\gamma)} \sin^2(\omega_k nt) (\varphi_x^k)^2 \leq \sum_{k=0}^{n-1} (\varphi_x^k)^2 = 1, \end{aligned} \quad (3.50)$$

where we used the fact that ϕ^k and φ^k are orthonormal basis ($\sum_{z=1}^{n-1} \phi_z^k \phi_z^{k'} = \delta_{k,k'}$, $\sum_{k=0}^{n-1} (\varphi_x^k)^2 = 1$). Similarly, we have $\sum_{z=1}^n \mathbf{a}_z^2 \leq 1$. Plugging the latter and (3.50) inside (3.49), we get:

$$m_x \mathbf{C}_{xx}^{pr} \leq C + \sum_{l=1}^{n-1} \frac{2C}{l^2} \leq c', \quad (3.51)$$

where as usual $c' > 0$ is a constant independent of n and the realization of the masses. Similarly, we can bound other terms in (3.47) by constants independent of n and the realization of the masses. Notice that this constants can be chosen to be independent of x as well. Moreover, the exact same argument works for $\langle (\tilde{\mathbf{r}}_x(nt))^{\bullet} \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}$. Hence, there exists a constant $C > 0$ such that for n , any $x \in \mathbb{I}_n$, and any realization of the masses we have:

$$\frac{\langle (\tilde{\mathbf{p}}_x^{\bullet}(nt))^2 \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}}{m_x} \leq C, \quad \langle (\tilde{\mathbf{r}}_x^{\bullet}(nt))^2 \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \leq C. \quad (3.52)$$

Before proceeding, notice that in the above argument, we can replace $\tilde{\mathbf{p}}_x^{\bullet}(nt)$ with $\tilde{\mathbf{p}}_x(nt)$ and all the steps work. Therefore, exists $C > 0$ such that for n , any $x \in \mathbb{I}_n$, and any realization of the masses we have:

$$\frac{\langle (\tilde{\mathbf{p}}_x(nt))^2 \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}}{m_x} \leq C, \quad \langle (\tilde{\mathbf{r}}_x(nt))^2 \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \leq C. \quad (3.53)$$

Having (3.52), bounding $f(x/n)/(m_x)^{\frac{1}{2}}$, and a Cauchy-Schwartz (3.34) yields:

$$\begin{aligned} & \left| \frac{1}{n^2} \sum_{|x-y| \leq 4n^{\theta'}} f\left(\frac{x}{n}\right) f\left(\frac{y}{n}\right) \langle \tilde{\mathbf{p}}_x^{\bullet}(nt) \tilde{\mathbf{p}}_y^{\bullet}(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right| \leq \frac{C}{n^2} \sum_{|x-y| \leq 4n^{\theta'}} \left| \frac{\langle \tilde{\mathbf{p}}_x^{\bullet}(nt) \tilde{\mathbf{p}}_y^{\bullet}(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}}{\sqrt{m_x m_y}} \right| \leq \\ & \frac{C'}{n^2} \sum_{|x-y| \leq 4n^{\theta'}} \left(\frac{\langle (\tilde{\mathbf{p}}_x^{\bullet}(nt))^2 \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}}{m_x} \right)^{\frac{1}{2}} \left(\frac{\langle (\tilde{\mathbf{p}}_y^{\bullet}(nt))^2 \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}}{m_y} \right)^{\frac{1}{2}} \leq \frac{C'}{n^2} \sum_{|x-y| \leq 4n^{\theta'}} C \leq \frac{C''}{n^2} \times cn^{1+\theta'} \rightarrow 0, \end{aligned} \quad (3.54)$$

as $n \rightarrow \infty$, thanks to the choice of $0 < \theta' < 1$. Similar identity holds for \mathbf{r} :

$$\left| \frac{1}{n^2} \sum_{|x-y| \leq 4n^{\theta'}} f\left(\frac{x}{n}\right) f\left(\frac{y}{n}\right) \langle \tilde{\mathbf{r}}_x^{\bullet}(nt) \tilde{\mathbf{r}}_y^{\bullet}(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right| \rightarrow 0. \quad (3.55)$$

Combining (3.45), and (3.54) we can deduce $\mathbf{U}_n \rightarrow 0$ almost surely w.r.t the distribution of the masses, similarly (3.46), and (3.55) gives us $\mathbf{U}'_n \rightarrow 0$ almost surely. This conclude the proof of (3.38) and this step.

Step 3: Summing up. Recall \mathbf{P}_n (3.27). By using the fact that $\tilde{\mathbf{p}}_x(nt) = \tilde{\mathbf{p}}_x^o(nt) + \tilde{\mathbf{p}}_x^{\bullet}(nt)$, and a Cauchy-Schwartz ($|\langle AB \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}| \leq \langle AA^* \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}^{\frac{1}{2}} \langle BB^* \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}^{\frac{1}{2}}$) inequality we observe that $\mathbf{P}_n \leq \mathbf{L}_n + \mathbf{U}_n + (\mathbf{L}_n)^{\frac{1}{2}} (\mathbf{U}_n)^{\frac{1}{2}}$. Combining this with (3.33), and (3.38) we deduce that $\mathbf{P}_n \rightarrow 0$ almost surely w.r.t the distribution of the masses. Similarly, we can deduce that $\mathbf{R}_n \rightarrow 0$ almost surely w.r.t the distribution of the masses. This finishes the proof of (3.27) and consequently (3.11), (3.12). \square

3.3 Evolution of e

Recall average variables $\bar{\mathbf{p}}_x(nt)$, $\bar{\mathbf{r}}_x(nt)$ in (3.24). As we mentioned, they satisfy the same system of ODEs as in [12] up to a vanishing constant. Therefore, by using the regularity of initial profiles \bar{r} , \bar{p} , we can adapt the argument of [12], and deduce the following (cf. [12], Sect.4, (4.29)) for any $y \in (0, 1)$:

$$\bar{\mathbf{r}}_{[ny]}(nt) \rightarrow \mathbf{r}(y, t), \quad \frac{\bar{\mathbf{p}}_{[ny]}}{m_{[ny]}} \rightarrow \frac{\mathbf{p}(y, t)}{\bar{m}}, \quad (3.56)$$

almost surely w.r.t the distribution of the masses. Therefore, by using the averaging lemma of [12] (cf. (4.13) in [12]) we can deduce the convergence of the average of mechanical energy in the following sense (cf. Lemma 6.1 in [12]): for any $f \in C^0([0, 1])$

$$\frac{1}{n} \sum_{x=1}^n f\left(\frac{x}{n}\right) \left(\frac{\langle \mathbf{p}_x(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}}{2m_x} + \frac{\langle \mathbf{r}_x(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}}{2} \right) \rightarrow \int_0^1 f(y) \left(\frac{\mathbf{p}(y, t)}{2\bar{m}} + \frac{\mathbf{r}^2(y, t)}{2} \right). \quad (3.57)$$

On the other hand, the evolution of fluctuation variables (3.29) is identical to the one in [12]. Although, the ensemble average here is different from [12], one can observe that having the assumption (3.8), and the bounds (3.30), (3.31), and (3.45) are sufficient for using the result of [12]. More precisely, recall the fluctuation operators (3.24) $\tilde{\mathbf{p}}_x(nt)$, $\tilde{\mathbf{r}}_x(nt)$; having the aforementioned bounds we can apply Lemma 6.4. of [12] and deduce for any $f \in C^0([0, 1])$:

$$T_n(t) - T_n(0) := \frac{1}{n} \sum_{x=1}^n f\left(\frac{x}{n}\right) \left(\frac{\langle \tilde{\mathbf{p}}_x^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}}{2m_x} + \frac{\langle \tilde{\mathbf{r}}_x^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}}{2} \right) - \frac{1}{n} \sum_{x=1}^n f\left(\frac{x}{n}\right) \left(\frac{\langle \tilde{\mathbf{p}}_x^2(0) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}}{2m_x} + \frac{\langle \tilde{\mathbf{r}}_x^2(0) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}}{2} \right) \rightarrow 0, \quad (3.58)$$

almost surely w.r.t. the distribution of the masses. Combining the latter with (3.8) yields $T_n(t) \rightarrow \int_0^1 f(y) \bar{b}(y) dy$. Combining this with (3.57), adding and subtracting the average and comparing the obtained expression with explicit solution to the macroscopic evolution equation (1.2), i.e. $2\mathbf{e}(y, t) = \mathbf{p}^2(y, t)/\bar{m} + \mathbf{r}^2(y, t) + 2\bar{b}(y)$, we have the quantum counterpart of (1.3) for the energy:

$$\frac{1}{n} \sum_{x=1}^n f\left(\frac{x}{n}\right) \langle \mathbf{e}_x(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} = \frac{1}{n} \sum_{x=1}^n f\left(\frac{x}{n}\right) \left(\frac{\langle \mathbf{p}_x^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}}{2m_x} + \frac{\langle \mathbf{r}_x^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}}{2} \right) \rightarrow \int_0^1 f(y) \mathbf{e}(y, t) dy, \quad (3.59)$$

almost surely w.r.t the distribution of the masses. Having (3.59), we can complete the proof of Theorem 3.1.

Before proceeding, notice that thanks to (3.1) and (3.23) we have for all $k, k' \in \mathbb{I}_n^o$, and all $x, y \in \mathbb{I}_n$:

$$\begin{aligned} [\hat{\mathbf{p}}_k(nt), \hat{\mathbf{p}}_{k'}(nt)] &= [\hat{\mathbf{r}}_k(nt), \hat{\mathbf{r}}_{k'}(nt)] = 0, \implies [\tilde{\mathbf{p}}_x(nt), \tilde{\mathbf{p}}_y(nt)] = [\tilde{\mathbf{r}}_x(nt), \tilde{\mathbf{r}}_y(nt)] = 0, \\ [\tilde{\mathbf{p}}_x^o(nt), \tilde{\mathbf{p}}_y^o(nt)] &= [\tilde{\mathbf{r}}_x^o(nt), \tilde{\mathbf{r}}_y^o(nt)] = [\tilde{\mathbf{p}}_x^\bullet(nt), \tilde{\mathbf{p}}_y^\bullet(nt)] = [\tilde{\mathbf{r}}_x^\bullet(nt), \tilde{\mathbf{r}}_y^\bullet(nt)] = 0, \\ [\tilde{\mathbf{p}}_x^\bullet(nt), \tilde{\mathbf{p}}_y^o(nt)] &= [\tilde{\mathbf{r}}_x^\bullet(nt), \tilde{\mathbf{r}}_y^o(nt)] = 0. \end{aligned} \quad (3.60)$$

In the following, we will use above identities without mentioning them.

Proof of Theorem 3.1: (3.13). Let us define the following operator:

$$\tilde{\mathbf{e}}_x(nt) := \mathbf{e}_x(nt) - \langle \mathbf{e}_x(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}. \quad (3.61)$$

At $t = 0$, notice the difference among $\tilde{\mathbf{e}}_x$ and $\check{\mathbf{e}}_x$ (3.7). As we did before, adding and subtracting the average term, and using (3.59), proof of (3.13) boils down to proving that

$$\mathbf{E}_n := \left\langle \left(\frac{1}{n} \sum_{x=1}^n f\left(\frac{x}{n}\right) \tilde{\mathbf{e}}_x(nt) \right)^2 \right\rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \rightarrow 0, \quad (3.62)$$

almost surely w.r.t the distribution of the masses. By using the definition of $\tilde{\mathbf{e}}_x(nt)$ and a Cauchy-Schwartz inequality (3.34), it is sufficient to prove:

$$\begin{aligned} \mathbf{P}_n &:= \left\langle \left(\frac{1}{n} \sum_{x=1}^n f\left(\frac{x}{n}\right) \left(\frac{\mathbf{p}_x^2(nt)}{m_x} - \left\langle \frac{\mathbf{p}_x^2(nt)}{m_x} \right\rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right) \right)^2 \right\rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \rightarrow 0, \\ \mathbf{R}_n &:= \left\langle \left(\frac{1}{n} \sum_{x=1}^n f\left(\frac{x}{n}\right) \left(\mathbf{r}_x^2(nt) - \langle \mathbf{r}_x^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right) \right)^2 \right\rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \rightarrow 0, \end{aligned} \quad (3.63)$$

almost surely w.r.t the distribution of the masses. First, recall average variables and fluctuation operators (3.24), observe that computation of (2.49) is valid in the quantum case as well:

$$\begin{aligned} &\left\langle \left(\mathbf{p}_x^2(nt) - \langle \mathbf{p}_x^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right) \left(\mathbf{p}_y^2(nt) - \langle \mathbf{p}_y^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right) \right\rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} = \\ &\langle \tilde{\mathbf{p}}_x^2(nt) \tilde{\mathbf{p}}_y^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} - \langle \tilde{\mathbf{p}}_x^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \langle \tilde{\mathbf{p}}_y^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} + 4\bar{\mathbf{p}}_x(nt)\bar{\mathbf{p}}_y(nt) \langle \tilde{\mathbf{p}}_x(nt) \tilde{\mathbf{p}}_y(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} + \\ &2\bar{\mathbf{p}}_x(nt) \langle \tilde{\mathbf{p}}_x(nt) \tilde{\mathbf{p}}_y^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} + 2\bar{\mathbf{p}}_y(nt) \langle \tilde{\mathbf{p}}_x^2(nt) \tilde{\mathbf{p}}_y(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}. \end{aligned} \quad (3.64)$$

In contrast to the classical case, we cannot simply do pairings in $\langle \cdot \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}$. Therefore, controlling each term in (3.64) is more involved. We control each term separately: decompose them into low and high modes, and take advantage of the localization for high modes with $|x - y|$ large enough, and control the remainder.

We only prove (3.63) for \mathbf{P}_n , thanks to symmetry in our assumption, the proof for \mathbf{R}_n is identical, up to small modifications which we will omit. By using a Cauchy-Schwartz inequality and recalling the definition of $\bar{\mathbf{p}}_x(nt)$, $\tilde{\mathbf{p}}_x(nt)$ (3.24), for obtaining (3.63) it is sufficient to prove:

$$\begin{aligned} \mathbf{P}'_n &:= \left\langle \left(\frac{1}{n} \sum_{x=1}^n f\left(\frac{x}{n}\right) \left(\frac{\tilde{\mathbf{p}}_x^2(nt)}{m_x} - \frac{\langle \tilde{\mathbf{p}}_x^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}}{m_x} \right) \right)^2 \right\rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \rightarrow 0, \\ \mathbf{P}''_n &:= \left\langle \left(\frac{1}{n} \sum_{x=1}^n f\left(\frac{x}{n}\right) \frac{\bar{\mathbf{p}}_x(nt) \tilde{\mathbf{p}}_x(nt)}{m_x} \right)^2 \right\rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \rightarrow 0, \end{aligned} \quad (3.65)$$

as $n \rightarrow \infty$ almost surely w.r.t distribution of the masses. The proof of second identity is straightforward: thanks to (3.56), and regularity of the macroscopic equation, $\bar{\mathbf{p}}_x/m_x$ is

uniformly (in n and distribution of the masses) bounded (cf. [12] (4.11)). On the other hand, proof of (3.27) only rests on the fact that f is bounded and does not use its regularity. Therefore, we absorb \bar{p}_x/m_x inside f and follow similar steps and thanks to (3.27) deduce that:

$$P''_n \rightarrow 0, \quad (3.66)$$

almost surely w.r.t the distribution of the masses.

Fix $0 < 2\gamma < \theta < \theta' < 1$ (we choose them opportunely later), and recall the definition of $\tilde{p}_x^o(nt)$, $\tilde{p}_x^\bullet(nt)$ (3.32), (3.37) (recall $I(\gamma) =]n^{1-\gamma}, n] \cap \mathbb{Z}$, $\tilde{I}(\gamma) = [1, n^{1-\gamma}] \cap \mathbb{Z}$). Thanks to a Cauchy-Schwartz inequality proof of the first limit in (3.65) reduces to the following :

$$\begin{aligned} \Pi_n^o &:= \left\langle \left(\frac{1}{n} \sum_{x=1}^n \frac{f(\frac{x}{n})}{m_x} \left((\tilde{p}_x^o)^2(nt) - \langle (\tilde{p}_x^o)^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right) \right)^2 \right\rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \rightarrow 0, \\ \Pi_n^\bullet &:= \left\langle \left(\frac{1}{n} \sum_{x=1}^n \frac{f(\frac{x}{n})}{m_x} \left((\tilde{p}_x^\bullet)^2(nt) - \langle (\tilde{p}_x^\bullet)^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right) \right)^2 \right\rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \rightarrow 0, \\ \Pi_n^{\bullet\bullet} &:= \left\langle \left(\frac{1}{n} \sum_{x=1}^n \frac{f(\frac{x}{n})}{m_x} \left(\tilde{p}_x^o(nt) \tilde{p}_x^\bullet(nt) - \langle \tilde{p}_x^o(nt) \tilde{p}_x^\bullet(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right) \right)^2 \right\rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \rightarrow 0, \end{aligned} \quad (3.67)$$

as $n \rightarrow \infty$ almost surely w.r.t the distribution of the masses. Let us begin with the first limit. To this end, we need following bounds: Thanks to assumption 3 (3.9), and by a computation similar to (3.30) we have:

$$\begin{aligned} \langle \hat{\tilde{p}}_k^4 \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} &\leq c \sum_{z_1, z_2, z_3, z_4} |\varphi_{z_1}^k \varphi_{z_2}^k \varphi_{z_3}^k \varphi_{z_4}^k| \left| \langle \tilde{p}_{z_1} \tilde{p}_{z_2} \tilde{p}_{z_3} \tilde{p}_{z_4} \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right| \leq \\ &c \sum_{\sigma} \sum_{z_{\sigma_1}, z_{\sigma_2}} \frac{C}{|z_{\sigma_1} - z_{\sigma_2}|^2} |\varphi_{z_{\sigma_1}}^k \varphi_{z_{\sigma_2}}^k| \sum_{z_{\sigma_3}, z_{\sigma_4}} \frac{C}{|z_{\sigma_3} - z_{\sigma_4}|^2} |\varphi_{z_{\sigma_3}}^k \varphi_{z_{\sigma_4}}^k| \leq C, \end{aligned} \quad (3.68)$$

where in the first line we used (3.9), in the second line we sum over three following permutations $\sigma : \{1, 2, 3, 4\} \rightarrow \{1, 2, 3, 4\}$: (12)(34), (13)(24), and (14)(23). The last inequality is obtained similar to (3.30). Notice that C is independent of n , and realization of the masses. Thanks to symmetry of our assumption, we can deduce similar bounds for $\langle (\hat{\tilde{p}}_k)^j (\hat{\tilde{r}}_k)^{4-j} \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}$, for $j \in \{0, 1, 2, 3, 4\}$. Therefore, we can deduce similar bounds at time nt : there exists $C > 0$ such that for any n , $k \in \mathbb{I}_n^o$, and any realization of the masses:

$$\langle \hat{\tilde{p}}_k^4(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \leq C, \quad \langle \hat{\tilde{r}}_k^4(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \leq C. \quad (3.69)$$

Recall $\tilde{I}(\gamma) = [0, n^{1-\gamma}] \cap \mathbb{Z}$ and observe for any constant $C > 0$:

$$\begin{aligned}
& \frac{C}{n^2} \sum_{x,y=1}^n \left| \langle (\tilde{\mathbf{p}}_x^o)^2(nt) (\tilde{\mathbf{p}}_y^o)^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} - \langle (\tilde{\mathbf{p}}_x^o)^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \langle (\tilde{\mathbf{p}}_y^o)^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right| \leq \\
& \frac{C'}{n^2} \sum_{x,y=1}^n \frac{1}{m_x m_y} \left(\langle (\tilde{\mathbf{p}}_x^o)^2(nt) (\tilde{\mathbf{p}}_y^o)^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} + \langle (\tilde{\mathbf{p}}_x^o)^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \langle (\tilde{\mathbf{p}}_y^o)^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right) \leq \\
& \frac{C''}{n^2} \sum_{x,y=1}^n \sum_{k_1, k_2, k_3, k_4 \in \tilde{I}(\gamma)} \varphi_x^{k_1} \varphi_x^{k_2} \varphi_y^{k_3} \varphi_y^{k_4} \left(\langle \hat{\mathbf{p}}_{k_1}(nt) \hat{\mathbf{p}}_{k_2}(nt) \hat{\mathbf{p}}_{k_3}(nt) \hat{\mathbf{p}}_{k_4}(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} + \right. \\
& \left. \langle \hat{\mathbf{p}}_{k_1}(nt) \hat{\mathbf{p}}_{k_2}(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \langle \hat{\mathbf{p}}_{k_3}(nt) \hat{\mathbf{p}}_{k_4}(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right) = \\
& \frac{C''}{n^2} \sum_{k_1, k_3 \in \tilde{I}(\gamma)} \left(\langle (\hat{\mathbf{p}}_{k_1}(nt))^2 (\hat{\mathbf{p}}_{k_3}(nt))^2 \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} + \langle (\hat{\mathbf{p}}_{k_1}(nt))^2 \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \langle (\hat{\mathbf{p}}_{k_3}(nt))^2 \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right) \leq \frac{\tilde{C} n^{2-2\gamma}}{n^2} \rightarrow 0,
\end{aligned} \tag{3.70}$$

as $n \rightarrow \infty$, where we used the fact that $\sum_x \varphi_x^k \varphi_x^{k'} = \delta_{k,k'}$, a Cauchy-Schwartz inequality and bounds (3.69), (3.36), as well as the fact that $\gamma > 0$. Expanding the square in Π_n^o , bounding $|f(\frac{x}{n})/m_x|$, and taking advantage of (3.70) we deduce that $\Pi_n^o \rightarrow 0$.

In order to prove two other limits, we need the following bound: there exists $C > 0$ such that for any n , $x \in \mathbb{I}_n$, and any realization of the masses:

$$\begin{aligned}
\langle \tilde{\mathbf{p}}_x^4(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} &\leq C, \quad \langle (\tilde{\mathbf{p}}_x^o)^4(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \leq C, \quad \langle (\tilde{\mathbf{p}}_x^\bullet)^4(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \leq C, \\
\langle \tilde{\mathbf{r}}_x^4(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} &\leq C, \quad \langle (\tilde{\mathbf{r}}_x^o)^4(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \leq C, \quad \langle (\tilde{\mathbf{r}}_x^\bullet)^4(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \leq C,
\end{aligned} \tag{3.71}$$

we only show the above bound for $\langle \tilde{\mathbf{p}}_x^4(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}$, other bound can be obtained similarly. Notice that we have:

$$\frac{\langle \tilde{\mathbf{p}}_x^4(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}}{m_x} = \sum_{k_1, k_2, k_3, k_4} \varphi_x^{k_1} \varphi_x^{k_2} \varphi_x^{k_3} \varphi_x^{k_4} \langle \hat{\mathbf{p}}_{k_1}(nt) \hat{\mathbf{p}}_{k_2}(nt) \hat{\mathbf{p}}_{k_3}(nt) \hat{\mathbf{p}}_{k_4}(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}. \tag{3.72}$$

Considering the definition of $\hat{\mathbf{p}}_{k_1}(nt)$ (3.29), the last average in the above expression contains 16 terms. However, thanks to symmetry of our assumption it is sufficient to bound one of them and the rest are treated similarly. Let us take the term involving only \mathbf{p} terms. Let us denote

$$\mathbf{a}_z^x := \sum_k \varphi_x^k \varphi_z^k \cos(\omega_k nt). \tag{3.73}$$

Then by using the definition of $\hat{\mathbf{p}}_k$ we have:

$$\begin{aligned}
& \sum_{k_1, k_2, k_3, k_4} \varphi_x^{k_1} \varphi_x^{k_2} \varphi_x^{k_3} \varphi_x^{k_4} \cos(\omega_{k_1} nt) \cos(\omega_{k_2} nt) \cos(\omega_{k_3} nt) \cos(\omega_{k_4} nt) \langle \hat{\mathbf{p}}_{k_1} \hat{\mathbf{p}}_{k_2} \hat{\mathbf{p}}_{k_3} \hat{\mathbf{p}}_{k_4} \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} = \\
& \sum_{z_1, z_2, z_3, z_4=1}^n \left\langle \frac{\tilde{\mathbf{p}}_{z_1}}{\sqrt{m_{z_1}}} \frac{\tilde{\mathbf{p}}_{z_2}}{\sqrt{m_{z_2}}} \frac{\tilde{\mathbf{p}}_{z_3}}{\sqrt{m_{z_3}}} \frac{\tilde{\mathbf{p}}_{z_4}}{\sqrt{m_{z_4}}} \right\rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \mathbf{a}_{z_1}^x \mathbf{a}_{z_2}^x \mathbf{a}_{z_3}^x \mathbf{a}_{z_4}^x \leq \\
& \sum_{\sigma} \sum_{z_{\sigma_1}, z_{\sigma_2}} \frac{C}{|z_{\sigma_1} - z_{\sigma_2}|^2} |\mathbf{a}_{z_{\sigma_1}}| |\mathbf{a}_{z_{\sigma_2}}| \sum_{z_{\sigma_3}, z_{\sigma_4}} \frac{C}{|z_{\sigma_3} - z_{\sigma_4}|^2} |\mathbf{a}_{z_{\sigma_3}}| |\mathbf{a}_{z_{\sigma_4}}| \leq C,
\end{aligned} \tag{3.74}$$

where \sum_{σ} is a summation over the three above-mentioned permutations; in the second line we used the assumption (3.9). Comparing each sum ($\sum_{z_{\sigma_i}, z_{\sigma_j}}$) with the expression appearing in (3.48), and using the bound (3.51) it is straightforward to observe that the above expression is bounded by a constant. Notice that as in (3.48) our argument can be adapted for $\tilde{\mathbf{p}}^{\bullet}$, and $\tilde{\mathbf{p}}^{\circ}$. Hence, we can deduce (3.71).

Recall $0 < 2\gamma < \theta < \theta' < 1$. Bounding $f(\frac{x}{n})/m_x$, taking advantage of the Cauchy-Schwartz inequality (3.34), and using the bounds (3.52), and (3.71) we have:

$$\left| \frac{1}{n^2} \sum_{|x-y| \leq 16n^{\theta'}} \frac{f(\frac{x}{n})f(\frac{y}{n})}{m_x m_y} \left((\tilde{\mathbf{p}}_x^{\bullet})^2(nt) - \langle (\tilde{\mathbf{p}}_x^{\bullet})^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right) \left((\tilde{\mathbf{p}}_y^{\bullet})^2(nt) - \langle (\tilde{\mathbf{p}}_y^{\bullet})^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right) \right| \leq C \frac{n^{\theta'}}{n} \rightarrow 0, \quad (3.75)$$

as $n \rightarrow \infty$. Now take $|x - y| > 4n^{\theta'}$, and observe:

$$\left| \langle (\tilde{\mathbf{p}}_x^{\bullet})^2(nt) (\tilde{\mathbf{p}}_y^{\bullet})^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} - \langle (\tilde{\mathbf{p}}_x^{\bullet})^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \langle (\tilde{\mathbf{p}}_y^{\bullet})^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right| = \left| \sum_{k_1, k_2, k_3, k_4 \in I(\gamma)} \varphi_x^{k_1} \varphi_x^{k_2} \varphi_y^{k_3} \varphi_y^{k_4} \times \right. \\ \left. \langle \hat{\mathbf{p}}_{k_1}(nt) \hat{\mathbf{p}}_{k_2}(nt) \hat{\mathbf{p}}_{k_3}(nt) \hat{\mathbf{p}}_{k_4}(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} - \langle \hat{\mathbf{p}}_{k_1}(nt) \hat{\mathbf{p}}_{k_2}(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \langle \hat{\mathbf{p}}_{k_3}(nt) \hat{\mathbf{p}}_{k_4}(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right|. \quad (3.76)$$

Using the definition of $\hat{\mathbf{p}}_k(nt)$, we obtain 16 different terms. However, thanks to the symmetry of (3.9), and Lemma 2.19, all these terms can be treated similarly. Therefore, we take the terms involving only $\tilde{\mathbf{p}}$. Let us denote

$$\tilde{\mathbf{a}}_z^x := \sum_{k \in I(\gamma)} \cos(\omega_k nt) \varphi_x^k \varphi_z^k. \quad (3.77)$$

Then by using the definition of $\hat{\mathbf{p}}_k$ we have:

$$\tilde{\mathbf{C}}_{xy} := \left| \sum_{k_1, k_2, k_3, k_4 \in I(\gamma)} \varphi_x^{k_1} \varphi_x^{k_2} \varphi_y^{k_3} \varphi_y^{k_4} \cos(\omega_{k_1} nt) \cos(\omega_{k_1} nt) \cos(\omega_{k_2} nt) \cos(\omega_{k_3} nt) \cos(\omega_{k_4} nt) \right. \\ \left. \langle \hat{\mathbf{p}}_{k_1} \hat{\mathbf{p}}_{k_2} \hat{\mathbf{p}}_{k_3} \hat{\mathbf{p}}_{k_4} \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} - \langle \hat{\mathbf{p}}_{k_1} \hat{\mathbf{p}}_{k_2} \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \langle \hat{\mathbf{p}}_{k_3} \hat{\mathbf{p}}_{k_4} \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right| = \\ \left| \sum_{z_1, z_2, z_3, z_4=1}^n \left(\langle \frac{\tilde{\mathbf{p}}_{z_1}}{\sqrt{m_{z_1}}} \frac{\tilde{\mathbf{p}}_{z_2}}{\sqrt{m_{z_2}}} \frac{\tilde{\mathbf{p}}_{z_3}}{\sqrt{m_{z_3}}} \frac{\tilde{\mathbf{p}}_{z_4}}{\sqrt{m_{z_4}}} \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} - \langle \frac{\tilde{\mathbf{p}}_{z_1}}{\sqrt{m_{z_1}}} \frac{\tilde{\mathbf{p}}_{z_2}}{\sqrt{m_{z_2}}} \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \langle \frac{\tilde{\mathbf{p}}_{z_3}}{\sqrt{m_{z_3}}} \frac{\tilde{\mathbf{p}}_{z_4}}{\sqrt{m_{z_4}}} \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right) \times \right. \\ \left. \tilde{\mathbf{a}}_{z_1}^x \tilde{\mathbf{a}}_{z_2}^x \tilde{\mathbf{a}}_{z_3}^x \tilde{\mathbf{a}}_{z_4}^x \right| \leq \sum_{z_1, z_3} \frac{c}{|z_1 - z_3|^2} |\tilde{\mathbf{a}}_{z_1}^x| |\tilde{\mathbf{a}}_{z_3}^y| \sum_{z_2, z_4} \frac{c}{|z_2 - z_4|^2} |\tilde{\mathbf{a}}_{z_2}^x| |\tilde{\mathbf{a}}_{z_4}^y| + \\ \sum_{z_1, z_4} \frac{c}{|z_1 - z_4|^2} |\tilde{\mathbf{a}}_{z_1}^x| |\tilde{\mathbf{a}}_{z_4}^y| \sum_{z_2, z_3} \frac{c}{|z_2 - z_3|^2} |\tilde{\mathbf{a}}_{z_2}^x| |\tilde{\mathbf{a}}_{z_3}^y| \quad (3.78)$$

where we used the assumption (3.9). We can decompose each sum in the above expression as we did in (3.40). In fact, comparing above sums, with the expression in (3.40), thanks

to (3.42), (3.43) it is straightforward to observe that

$$\sum_{|x-y|>4n^{\theta'}} \tilde{C}_{xy} \rightarrow 0, \quad (3.79)$$

almost surely w.r.t the distribution of the masses, where one should recall (3.42), and (3.41). Combining the above symmetry argument, (3.76) (3.79), and (3.78), we have:

$$\left| \frac{1}{n^2} \sum_{|x-y|>4n^{\theta'}} \frac{f(\frac{x}{n})f(\frac{y}{n})}{m_x m_y} \left((\tilde{\mathbf{p}}_x^\bullet)^2(nt) - \langle (\tilde{\mathbf{p}}_x^\bullet)^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right) \left((\tilde{\mathbf{p}}_y^\bullet)^2(nt) - \langle (\tilde{\mathbf{p}}_y^\bullet)^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right) \right| \rightarrow 0, \quad (3.80)$$

almost surely w.r.t the distribution of the masses. Finally, having (3.80), (3.75) we conclude $\Pi_n^\bullet \rightarrow 0$ almost surely w.r.t the distribution of the masses.

Now we prove the last limit of (3.67) (we abbreviate: $a(nt)b(nt) = (ab)(nt)$):

$$\begin{aligned} \Pi_n^\bullet &\leq \frac{C}{n^2} \sum_{x,y=1}^n \left| \langle (\tilde{\mathbf{p}}_x^o \tilde{\mathbf{p}}_y^o \tilde{\mathbf{p}}_x^\bullet \tilde{\mathbf{p}}_y^\bullet)(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right| + \frac{C}{n^2} \sum_{x,y=1}^n \left| \langle (\tilde{\mathbf{p}}_x^o \tilde{\mathbf{p}}_x^\bullet)(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right| \left| \langle (\tilde{\mathbf{p}}_y^o \tilde{\mathbf{p}}_y^\bullet)(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right| \\ &\leq \frac{C}{n^2} \sum_{x,y=1}^n \langle (\tilde{\mathbf{p}}_x^o \tilde{\mathbf{p}}_y^o)^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}^{\frac{1}{2}} \langle (\tilde{\mathbf{p}}_x^\bullet \tilde{\mathbf{p}}_y^\bullet)^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}^{\frac{1}{2}} + \\ &\quad \frac{C}{n^2} \sum_{x,y=1}^n \langle (\tilde{\mathbf{p}}_x^o)^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}^{\frac{1}{2}} \langle (\tilde{\mathbf{p}}_y^o)^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}^{\frac{1}{2}} \langle (\tilde{\mathbf{p}}_x^\bullet)^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}^{\frac{1}{2}} \langle (\tilde{\mathbf{p}}_y^\bullet)^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}^{\frac{1}{2}} \leq \\ &\quad \frac{C'}{n^2} \sum_{x,y=1}^n \langle (\tilde{\mathbf{p}}_x^o)^2(nt) (\tilde{\mathbf{p}}_y^o)^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}^{\frac{1}{2}} + \frac{C'}{n^2} \sum_{x,y=1}^n \langle (\tilde{\mathbf{p}}_x^o)^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}^{\frac{1}{2}} \langle (\tilde{\mathbf{p}}_y^o)^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}}^{\frac{1}{2}} \leq \\ &\quad c' \left(\frac{C}{n^2} \sum_{x,y=1}^n \frac{1}{m_x m_y} \langle (\tilde{\mathbf{p}}_x^o)^2(nt) (\tilde{\mathbf{p}}_y^o)^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right)^{\frac{1}{2}} + \\ &\quad c' \left(\frac{C}{n^2} \sum_{x,y=1}^n \frac{1}{m_x m_y} \langle (\tilde{\mathbf{p}}_x^o)^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \langle (\tilde{\mathbf{p}}_y^o)^2(nt) \rangle_{\varrho_n^{\bar{r}, \bar{p}, \bar{b}}} \right)^{\frac{1}{2}} \leq c'' \frac{n^{1-\gamma}}{n} \rightarrow 0, \end{aligned} \quad (3.81)$$

as $n \rightarrow \infty$, where in the above computation we take advantage of the fact that above operators commute (3.60); in the first line we bounded f/m_x , in the second inequality we performed Cauchy-Schwartz (3.34), in the third inequality we performed another Cauchy-Schwartz and used the bounds on "second and fourth moments" (3.71), (3.52). In the fourth inequality we take advantage of the Jensen inequality thanks to the fact that $f(z) = z^{\frac{1}{2}}$ is concave. Finally, the last inequality is deduced from (3.70).

This finishes proof of (3.67), which yields (3.65). This means $\mathbf{P}_n \rightarrow 0$ almost surely. As we mentioned the proof of the other limit in (3.63) is similar. Hence we conclude (3.62), and this finishes the proof of (3.13) which complete the proof of Theorem 3.1. \square

3.4 Locally Gibbs state

In this section, we me prove Theorem (3.2). Let us emphasize that this theorem can be deduced from tools developed in Sect. 5 of [12]. Here we only mention the main critical steps and refer reader to [12] for more details. Before proceeding, let us recall basic notations, definitions, and identities related to $\varrho_{\bar{r}, \bar{p}, \beta}^n$ mostly from [12]:

Recall the definition of $\varrho_{\bar{r}, \bar{p}, \beta}^n$ (3.14). Thanks to (3.41), (3.42) of [12] up to an error that vanishes in the limit,

$$\tilde{p}_x := p_x - \langle p_x \rangle_{\varrho_{\bar{r}, \bar{p}, \beta}^n} = p_x - \bar{p} \left(\frac{x}{n} \right) \left(\frac{m_x}{\bar{m}} \right), \quad \tilde{r}_x := r_x - \langle r_x \rangle_{\varrho_{\bar{r}, \bar{p}, \beta}^n} = r_x - \bar{r} \left(\frac{x}{n} \right). \quad (3.82)$$

Therefore, $\varrho_{\bar{r}, \bar{p}, \beta}^n$ can be written as $\varrho_{\bar{r}, \bar{p}, \beta}^n = \frac{1}{Z_n} \exp(-H_n^\beta)$, with

$$H_n^\beta = \frac{1}{2} \sum_{x=1}^n \left(\frac{\beta(\frac{x}{n})}{m_x} \tilde{p}_x^2 + \beta(\frac{x}{n}) \tilde{r}_x^2 \right). \quad (3.83)$$

Let us diagonalize H_n^β . We begin by a couple of definitions: Let $M_\beta = M \tilde{\beta}^{-1}$, with $\tilde{\beta} := \text{diag}(\beta(\frac{1}{n}), \dots, \beta(\frac{n}{n}))$ and $\beta^o := \text{diag}(\beta(\frac{1}{n}), \dots, \beta(\frac{n-1}{n}))$. Define A_p^β and A_r^β as:

$$A_p^\beta = M_\beta^{-\frac{1}{2}} (-\nabla_- \beta^o \nabla_+) M_\beta^{-\frac{1}{2}}, \quad A_r^\beta = (\beta^o)^{\frac{1}{2}} (-\nabla_+ M_\beta^{-1} \nabla_-) (\beta^o)^{\frac{1}{2}}. \quad (3.84)$$

A_p^β is symmetric positive semidefinite with almost sure non-degenerate spectrum. Denote $\{\psi^k\}_{k=0}^{n-1}$ to be its orthonormal set of eigenvectors, forming a basis for \mathbb{R}^n . Denote the corresponding set of increasing eigenvalues by $\{0 = \gamma_0^2 < \gamma_1^2 < \dots < \gamma_{n-1}^2\}$. Notice $\langle \psi^k, \psi^{k'} \rangle = \sum_{x=1}^n \psi_x^k \psi_x^{k'} = \delta_{k,k'}$, and $\sum_{k=0}^{n-1} \psi_x^k \psi_y^{k'} = \delta_{x,y}$. A_r^β is symmetric positive definite. Define for $k \in \mathbb{I}_{n-1}$,

$$\tilde{\psi}^k := \frac{1}{\gamma_k} (\beta^o)^{\frac{1}{2}} \nabla_+ M_\beta^{-\frac{1}{2}} \psi^k,$$

observe that $\{\tilde{\psi}^k\}_{k=1}^{n-1}$ is the set of eigenvectors of A_r^β with similar eigenvalues $\gamma_1^2 < \dots < \gamma_{n-1}^2$. We define another set of coordinates: for $k \in \mathbb{I}_n$ (convention: $r_n = 0$):

$$p_k := \langle \psi^k, M_\beta^{-\frac{1}{2}} \tilde{p} \rangle_n, \quad r_k := \langle \tilde{\psi}^k, \tilde{r} \rangle_{n-1}, \quad (3.85)$$

Correspondingly, we define the following set of bosonic operators for $k \in \mathbb{I}_{n-1}$:

$$b_k := \frac{1}{2\gamma_k} (r_k + ip_k), \quad b_k^* := \frac{1}{2\gamma_k} (r_k - ip_k). \quad (3.86)$$

From (3.85), and (3.86), by using properties of $\psi^k, \tilde{\psi}^k$, we have upto a vanishing error:

$$H_n^\beta = \frac{1}{2} \sum_{k=1}^{n-1} (p_k^2 + r_k^2) = \sum_{k=1}^{n-1} \gamma_k (b_k^* b_k + \frac{1}{2}), \quad (3.87)$$

where we have following commutation relations for $k, k' \in \mathbb{I}_n$ thanks to (3.1) and above definitions:

$$\begin{aligned} [p_k, p_{k'}] &= [r_k, r_{k'}] = 0, & [r_k, p_{k'}] &= i\gamma_k \delta_{k,k'}, \\ [b_k, b_{k'}] &= [b_k^*, b_{k'}^*] = 0, & [b_k^*, b_{k'}] &= \delta_{k,k'}. \end{aligned} \quad (3.88)$$

The detailed computation of above identities can be found in sec. 3 of [12], in particular (3.30)-(3.36).

Having (3.87) and (3.88) the discrete spectrum of H_β^n is understood. Moreover, thanks to spectral theorem we can compute average of certain observables w.r.t $\varrho_{\bar{r},\bar{p},\beta}^n$. In particular we have (cf. [12] Sect.3 (3.38), (3.48) and Appendix C): for $k, k' \in \mathbb{I}_{n-1}$, and $x, y \in \mathbb{I}_n$:

$$\begin{aligned} \langle b_k \rangle_{\varrho_{\bar{r},\bar{p},\beta}^n} &= \langle b_k^* \rangle_{\varrho_{\bar{r},\bar{p},\beta}^n} = 0, \quad \langle b_k^* b_{k'} \rangle_{\varrho_{\bar{r},\bar{p},\beta}^n} = \frac{\delta_{k,k'}}{e^{\gamma_k} - 1}, \implies \\ \langle p_k \rangle_{\varrho_{\bar{r},\bar{p},\beta}^n} &= \langle r_k \rangle_{\varrho_{\bar{r},\bar{p},\beta}^n} = 0, \quad \langle p_k p_{k'} \rangle_{\varrho_{\bar{r},\bar{p},\beta}^n} = \langle r_k r_{k'} \rangle_{\varrho_{\bar{r},\bar{p},\beta}^n} = \delta_{k,k'} \frac{\gamma_k}{2} \coth\left(\frac{\gamma_k}{2}\right). \end{aligned} \quad (3.89)$$

Finally, thanks to ladder structure of bosonic operators, by using spectral theorem we can observe that b_k has pairing structure in $\varrho_{\bar{r},\bar{p},\beta}^n$. Denote $b_k^1 := b_k$, $b_k^2 := b_k^*$. Then we have for any k_1, k_2, k_3, k_4 , and any $\sharp_1, \sharp_2, \sharp_3, \sharp_4 \in \{1, 2\}$:

$$\begin{aligned} \langle b_{k_1}^{\sharp_1} b_{k_2}^{\sharp_2} b_{k_3}^{\sharp_3} \rangle_{\varrho_{\bar{r},\bar{p},\beta}^n} &= 0, \\ \langle b_{k_1}^{\sharp_1} b_{k_2}^{\sharp_2} b_{k_3}^{\sharp_3} b_{k_4}^{\sharp_4} \rangle_{\varrho_{\bar{r},\bar{p},\beta}^n} &= \langle b_{k_1}^{\sharp_1} b_{k_3}^{\sharp_3} \rangle_{\varrho_{\bar{r},\bar{p},\beta}^n} \langle b_{k_2}^{\sharp_2} b_{k_4}^{\sharp_4} \rangle_{\varrho_{\bar{r},\bar{p},\beta}^n} + \langle b_{k_1}^{\sharp_1} b_{k_2}^{\sharp_2} \rangle_{\varrho_{\bar{r},\bar{p},\beta}^n} \langle b_{k_3}^{\sharp_3} b_{k_4}^{\sharp_4} \rangle_{\varrho_{\bar{r},\bar{p},\beta}^n} + \\ &\langle b_{k_1}^{\sharp_1} b_{k_4}^{\sharp_4} \rangle_{\varrho_{\bar{r},\bar{p},\beta}^n} \langle b_{k_2}^{\sharp_2} b_{k_3}^{\sharp_3} \rangle_{\varrho_{\bar{r},\bar{p},\beta}^n}. \end{aligned} \quad (3.90)$$

Now we state the proof of Theorem 3.2.

Proof of theorem 3.2. First notice that thanks to the pairing structure (3.90), the pairing structure remains true for $\tilde{p}_x(nt)$, $\tilde{r}_x(nt)$: similar to (2.50) with proper modification (since they are linear combination of bosonic operators, it is a matter of lengthy but straightforward computation to see this pairing structure persists). Therefore, $\varrho_{\bar{r},\bar{p},\beta}^n$ satisfies the third bound in (3.9), since odd moments are zero. Moreover, thanks to the pairing structure, it is sufficient to prove the first bound in (3.9). Finally, we prove the first bound for \tilde{p}_x , \tilde{p}_y . Adapting the proof to the case of \tilde{r}_x , \tilde{r}_y and cross terms is straightforward.

Inverse of (3.85) reads ($\beta_x := \beta(\frac{x}{n})$):

$$\tilde{p}_x = \sqrt{\frac{m_x}{\beta_x}} \sum_{k=1}^{n-1} \psi_x^k p_k. \quad (3.91)$$

Hence, by using (3.89) we have:

$$\langle \tilde{p}_x \tilde{p}_y \rangle_{\varrho_{\bar{r},\bar{p},\beta}^n} = \sqrt{\frac{m_x m_y}{\beta_x \beta_y}} \sum_{k,k'=1}^{n-1} \psi_x^k \psi_y^{k'} \langle p_k p_{k'} \rangle_{\varrho_{\bar{r},\bar{p},\beta}^n} = \sqrt{\frac{m_x m_y}{\beta_x \beta_y}} \sum_{k=1}^{n-1} \frac{\gamma_k}{2} \coth\left(\frac{\gamma_k}{2}\right) \psi_x^k \psi_y^k. \quad (3.92)$$

Recall the matrix A_p^β , and the fact that ψ^k is their eigenvectors. Denoting

$$f(z) = \begin{cases} z^{\frac{1}{2}} \coth(z^{\frac{1}{2}}), & z \neq 0, \\ 1, & z = 0. \end{cases} \quad (3.93)$$

by above expression (3.92) it is clear that we have (up to a vanishing error as $n \rightarrow \infty$):

$$\langle \tilde{p}_x \tilde{p}_y \rangle_{\varrho_{\bar{r},\bar{p},\beta}^n} = \langle x, f(A_p^\beta) y \rangle_n, \quad (3.94)$$

where $|x\rangle = (0, 0, \dots, 1, 0, \dots, 0) \in \mathbb{R}^n$ with 1 located at x th position.

The poles of the function $z^{\frac{1}{2}} \coth(z^{\frac{1}{2}})$ is the following set: $\{z \in \mathbb{C} | z = -k^2\pi^2, k \in \mathbb{Z}\}$, and this function is analytic on the rest of the complex plane. The point zero is a removable pole, and by redefining the function at zero, we can remove this pole: $\coth(z)$ has the following Taylor series expression for $0 < |z| < \pi$: $\coth(z) = z^{-1} + \sum_{n=1}^{\infty} a_n z^{2n-1}$, where $a_n = \frac{2^{2n} B_{2n}}{(2n)!}$, and B_{2n} are Bernoulli numbers. Hence, $z \coth(z) = 1 + \sum_{n=1}^{\infty} a_n z^{2n}$, and $z^{\frac{1}{2}} \coth(z^{\frac{1}{2}})$ is given by the following Taylor series: $1 + \sum_{n=1}^{\infty} a_n z^n$ for $0 < |z| < \pi^2$. Consequently, the pole of \mathfrak{f} is given by the set $\{z \in \mathbb{C} | z = -k^2\pi^2, k \in \mathbb{N}, k > 0\}$. Notice that by definition of A_p^β , thanks to properties of masses and β there is a constant $c_0 > 0$, uniform in n^1 , such that for any configuration of the masses, we have $\|A_p^\beta\|_2 < c_0$ ($\|\cdot\|_2$ denotes the usual operator norm of the matrix). Define $\alpha := \frac{1}{2}(c_0 + 1)$, let $\mathcal{R} := \alpha + \pi^2$, by the above argument $\mathfrak{f}(z)$ is analytic in the open disk $|z - \alpha| < \mathcal{R}$, and \mathcal{R} is the radius of convergence for the Taylor expansion of \mathfrak{f} , $\mathfrak{f}(z) = \sum_{k=0}^{\infty} a_k (z - \alpha)^k$. Moreover, by the choice of α and c_0 , the eigenvalues of A_p^β and A_r^β lies in the disk $|z - \alpha| < \mathcal{R}$. So we have the following Taylor expansions for $\mathfrak{f}(A_p^\beta)$ (cf. [12] sect. 5 for more details):

$$\mathfrak{f}(A_p^\beta) = \sum_{k=0}^{\infty} a_k (A_p^\beta - \alpha I_n)^k. \quad (3.95)$$

By a random walk representation argument(cf. (5.11) of [12]), since A_p^β is tridiagonal one can observe that for any $k < |x - y| - 1$, we have:

$$\langle x, (A_p^\beta)^k y \rangle_n = 0. \quad (3.96)$$

Plugging the Taylor series (3.95) inside the expression (3.94) and using (3.96) we have:

$$|\langle \tilde{\mathbf{p}}_x \tilde{\mathbf{p}}_y \rangle_{\mathcal{Q}_{\tilde{\mathbf{r}}, \tilde{\mathbf{p}}, \beta}^n}| \leq \left\| \sum_{k > |x-y|-2} a_k (A_p^\beta - \alpha I_n)^k \right\|_2 \leq \sum_{k > |x-y|-2} |a_k| \| (A_p^\beta - \alpha I_n) \|_2^k \leq C \mathfrak{q}^{|x-y|}, \quad (3.97)$$

for some $0 < \mathfrak{q} < 1$, where we used the fact that $\|(A_p^\beta - \alpha I_n)\|_2$ is inside the radius of convergence of the Taylor series, and we take advantage of properties of Taylor series (cf. (5.25) of [12]). The latter estimate gives us the first bound in (3.9). In fact, it provides much stronger exponential decay. This finishes proof of Theorem 3.2. \square

Acknowledgement

This work was partially supported by ANR-15-CE40-0020-01 LSD of the French National Research Agency. A.H. would like to thank prof. Gigliola Staffilani and M.I.T math department for their hospitality during this project.

¹This constant can be taken equal to $4 \frac{\beta_{max}^2}{m_{min}}$.

References

- [1] M Aizenman and G M Graf. Localization bounds for an electron gas. *Journal of Physics A: Mathematical and General*, 31(32):6783–6806, aug 1998.
- [2] Oskari Ajanki and François Huveneers. Rigorous scaling law for the heat current in disordered harmonic chain. *Communications in Mathematical Physics*, 301(3):841–883, Feb 2011.
- [3] Cédric Bernardin, François Huveneers, and Stefano Olla. Hydrodynamic limit for a disordered harmonic chain. *Communications in Mathematical Physics*, 365(1):215–237, Jan 2019.
- [4] Lorenzo Bertini, Alberto De Sole, Davide Gabrielli, Giovanni Jona-Lasinio, and Claudio Landim. Macroscopic fluctuation theory. *Rev. Mod. Phys.*, 87:593–636, Jun 2015.
- [5] O. Bratteli and D.W. Robinson. *Operator Algebras and Quantum Statistical Mechanics 1: C^* - and W^* -Algebras. Symmetry Groups. Decomposition of States.* Operator Algebras and Quantum Statistical Mechanics. Springer, 1987.
- [6] O. Bratteli and D.W. Robinson. *Operator Algebras and Quantum Statistical Mechanics: Equilibrium States. Models in Quantum Statistical Mechanics.* Theoretical and Mathematical Physics. Springer Berlin Heidelberg, 2003.
- [7] Nadine Braxmeier-Even and Stefano Olla. Hydrodynamic limit for a hamiltonian system with boundary conditions and conservative noise. *Archive for Rational Mechanics and Analysis*, 213(2):561–585, Aug 2014.
- [8] A. Casher and J. L. Lebowitz. Heat Flow in Regular and Disordered Harmonic Chains. *Journal of Mathematical Physics*, 12(8):1701–1711, August 1971.
- [9] J. M. Deutsch. Quantum statistical mechanics in a closed system. *Physical Review A*, 43:2046, 1991.
- [10] Abhishek Dhar. Heat conduction in the disordered harmonic chain revisited. *Phys. Rev. Lett.*, 86:5882–5885, Jun 2001.
- [11] Benjamin Doyon. Hydrodynamic projections and the emergence of linearised euler equations in one-dimensional isolated systems. *Communications in Mathematical Physics*, January 2022.
- [12] Amirali Hannani. Hydrodynamic limit for a disordered quantum harmonic chain. *Communications in Mathematical Physics*, 390:349–423, January 2022.
- [13] L. Isserlis. On a formula for the product-moment coefficient of any order of a normal frequency distribution in any number of variables. *Biometrika*, 12(1/2):134–139, 1918.
- [14] Tomasz Komorowski and Stefano Olla. Ballistic and superdiffusive scales in the macroscopic evolution of a chain of oscillators. *Nonlinearity*, 29(3):962–999, feb 2016.

- [15] Tomasz Komorowski, Stefano Olla, and Marielle Simon. Macroscopic evolution of mechanical and thermal energy in a harmonic chain with random flip of velocities. *Kinetic and Related Models*, 11(3):615–645, 2018.
- [16] Hervé Kunz and Bernard Souillard. Sur le spectre des opérateurs aux différences finies aléatoires. *Communications in Mathematical Physics*, 78(2):201–246, Dec 1980.
- [17] Bruno Nachtergaele, Robert Sims, and Günter Stolz. Quantum harmonic oscillator systems with disorder. *Journal of Statistical Physics*, 149(6):969–1012, Dec 2012.
- [18] S. Olla, S. R. S. Varadhan, and H.-T. Yau. Hydrodynamical limit for a hamiltonian system with weak noise. *Comm. Math. Phys.*, 155(3):523–560, 1993.
- [19] M. Reed and B. Simon. *Methods of Modern Mathematical Physics: Functional analysis*. Methods of Modern Mathematical Physics. Academic Press, 1980.
- [20] Marcos Rigol, Vanja Dunjko, and Maxim Olshanii. Thermalization and its mechanism for generic isolated quantum systems. *Nature*, 452:854–858, 2008.
- [21] Mark Srednicki. Chaos and quantum thermalization. *Physical Review E*, 50:888, 1994.
- [22] Theo Verheggen. Transmission coefficient and heat conduction of a harmonic chain with random masses: Asymptotic estimates on products of random matrices. *Communications in Mathematical Physics*, 68(1):69–82, Feb 1979.
- [23] Horng-Tzer Yau. Relative entropy and hydrodynamics of ginzburg-landau models. *Letters in Mathematical Physics*, 22(1):63–80, May 1991.