

Vector dark boson mediated feeble interaction between fermionic dark matter and strange quark matter in quark stars

Debashree Sen,^{1★} Atanu Guha,^{2†}

¹*Physics Group, Variable Energy Cyclotron Centre, 1/AF Bidhan Nagar, Kolkata 700064, India*

²*Department of Physics, Chungnam National University, 99, Daehak-ro, Yuseong-gu, Daejeon-34134, South Korea*

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

We study the structural properties like the gravitational mass, radius and tidal deformability of dark matter (DM) admixed strange quark stars (SQSs). For the purpose we consider the vector MIT Bag model to describe the strange quark matter (SQM) and investigate the possible presence of accreted DM in the SQSs consequently forming DM admixed SQSs. We introduce feeble interaction between SQM and the accreted fermionic DM via a vector dark boson mediator. Considering the present literature, in the context of possible presence of DM in SQSs, this work is the first to consider interaction between DM and SQM in the DM admixed SQSs. The mass of the DM fermion (m_χ) and the vector mediator (m_ξ) and the coupling (y_ξ) between them are determined in accordance with the constraint from Bullet cluster and the present day relic abundance, respectively. We find that the presence of DM reduces both the mass and radius of the star compared to the no-DM case. The massive the DM fermion, the lower the values of maximum mass and radius of the DM admixed SQSs. For the chosen values of m_χ and corresponding values of m_ξ and y_ξ , the computed structural properties of the DM admixed SQSs satisfy all the various present day astrophysical constraints. We obtain massive DM admixed SQSs configurations consistent with the GW190814 observational data. Hence the secondary compact object associated with this event may be a DM admixed SQS.

Key Words: (cosmology:) dark matter; gravitational waves; dense matter; equation of state; stars: massive.

1 INTRODUCTION

Several unknown and inconclusive facets of compact stars make them one of the most interesting objects in the Universe. One such inconclusive fact is their composition at such high density (5–10 times the nuclear density). So the dense matter composition and interactions at conditions relevant to compact stars are at present best understood by theoretical modeling of compact star matter. The corresponding interactions and the equation of state (EoS) are thus obtained based on speculative studies on theoretical modeling of compact star matter. Consequently, the presence of quark matter (QM) in compact stars is still experimentally unknown and a topic of current research. Theoretical speculations have predicted the possible existence of strange quark stars (SQSs) [Olinto \(1987\)](#) based on the Bodmer-Witten conjecture which states that strange QM (SQM), being composed of u , d and s quarks, have a lower energy per baryon number than the pure nucleonic system [Bodmer \(1971\)](#); [Chin & Kerman \(1979\)](#); [Witten \(1984\)](#). Ref. [Farhi & Jaffe \(1984\)](#) also ensured the stability of SQM at large baryon number and no external pressure with certain QCD parameters. Consequently, several theoretical works attempted the modeling of SQM in order to establish the possible existence of SQSs. The first and one of the most widely adopted model is the original MIT Bag model [Chodos et al. \(1974\)](#) which was later modified in [Fraga](#)

[et al. \(2001\)](#); [Alford et al. \(2005\)](#) as the non-ideal bag model. Further, repulsive interaction between the quarks was introduced via a parameter α_4 [Fraga et al. \(2001\)](#); [Alford et al. \(2005\)](#); [Glendenning \(2000\)](#); [Weissenborn et al. \(2011\)](#). The repulsive effect was also included by introducing vector meson as mediator (vBag model) [Klahn & Fischer \(2015\)](#); [Cierniak et al. \(2019\)](#); [Franzon et al. \(2016\)](#); [Wei et al. \(2019\)](#); [Lopes et al. \(2021\)](#); [Kumar et al. \(2022\)](#).

The bag constant B associated with the MIT Bag model represent the difference in energy density between the perturbative vacuum and the true vacuum. The bag constant B is still not well known. Model dependent analysis with respect to GW170817 data constrained $B^{1/4} = (134.1 - 141.4)$ MeV with low-spin prior and $B^{1/4} = (126.1 - 141.4)$ MeV with high-spin prior for SQSs [Zhou et al. \(2018\)](#) while for hybrid stars [Nandi & Char \(2018\)](#); [Nandi & Pal \(2021\)](#) estimated similar range of B . Ref. [Aziz et al. \(2019\)](#) also obtained the allowed range of $B^{1/4} = (133.68 - 222.53)$ MeV for SQSs while [Yang et al. \(2020\)](#) constrained $B^{1/4} = (141.3 - 150.9)$ MeV for SQSs with modified Tolman-Oppenheimer-Volkoff (TOV) formalism. It is also seen that the stability of SQS in terms of binding energy per baryon ε/ρ_B is controlled by B [Farhi & Jaffe \(1984\)](#); [Torres & Menezes \(2013\)](#); [Ferrer et al. \(2015\)](#). Ref. [Torres & Menezes \(2013\)](#) estimated the allowed range of B with respect to the stability condition of SQSs demanding that $\varepsilon/\rho_B \leq 930$ MeV, where ρ_B is the baryon density. In β equilibrated SQM the upper bound on B is set by considering charge neutral 3 flavor SQM in presence of electrons while the lower bound is obtained with 2 flavor QM [Torres & Menezes \(2013\)](#). Recently, [Lopes et al. \(2021\)](#) calculated the same for

the vBag model and established the allowed range of B for different vector couplings G_V . In the present work we adopt the same vBag model following [Lopes et al. \(2021\)](#) and calculate the stability window of B following the same criteria as $\varepsilon/\rho_B \leq m_n$. To calculate the EoS of SQM we consider an average value of B between B_{max} and B_{min} for a particular value of G_V .

Certain observational evidences like the rotation curves of the galaxies, observation of gravitational lensing, X-ray analysis of Bullet cluster [Bertone et al. \(2005\)](#); [Aghanim et al. \(2020\)](#) support the existence of dark matter (DM) in the Universe and compact stars like SQSs, being highly gravitating objects, are capable of accreting DM onto the system forming DM admixed SQSs. The exact nature and properties and interaction of DM particle candidates are unknown. The most suitable DM particle candidates are the Weakly Interacting Massive Particles (WIMPs) whose direct detection is being attempted in various experiments like super-CDMS [Agnese et al. \(2018\)](#), XENON100 [Aprile et al. \(2012\)](#), XENON1T [Aprile et al. \(2018\)](#), LUX [Akerib et al. \(2013\)](#), PANDAX-II [Wang et al. \(2020\)](#), DARKSIDE-50 [Agnes et al. \(2018\)](#), SENSEI [Crisler et al. \(2018\)](#) and very recently the LUX-ZEPLIN (LZ) [Aalbers et al. \(2022\)](#) etc. However, the exclusion bounds prescribed by such direct detection experiments are dependent on the local DM density around the solar neighborhood which do not affect the density of DM in the NS/SQS environment. The Cosmic Microwave Background (CMB) anisotropy maps, obtained from the Wilkinson Microwave Anisotropy Probe (WMAP) data [Bennett et al. \(2013\)](#), furnishes the present day thermal relic abundances of DM to be $\sim \Omega h^2 \approx 0.12$ [Tanabashi et al. \(2018\)](#); [Bauer & Plehn \(2019\)](#); [Cannoni \(2016\)](#). Thus any feasible DM model has to successfully reproduce the observed non-baryonic relic density. This sets strong constraints on the model parameters. On being accreted to the SQS system, the DM particles suffer collisions with the quarks and thereby lose kinetic energy and eventually become gravitationally bound to the star. At some point of time the accretion stops and the DM particles attain thermal equilibrium among themselves due to the self interactions. This justifies the DM particle density to be almost constant as considered by [Panotopoulos & Lopes \(2017b\)](#); [Guha & Sen \(2021\)](#); [Sen & Guha \(2021\)](#) in case of DM admixed neutron stars (NSs). The accreted DM particles therefore remain confined within a region of small radius inside the star. Several works have successfully explained the possible existence of DM admixed SQSs and DM admixed NSs in which the SQM or the NS matter do not interact with the DM [Lopes & Panotopoulos \(2018\)](#); [Ellis et al. \(2018\)](#); [Li et al. \(2012\)](#); [Tolos & Schaffner-Bielich \(2015\)](#); [Deliyergiyev et al. \(2019\)](#); [Rezaei \(2017\)](#); [Mukhopadhyay et al. \(2017\)](#); [Mukhopadhyay & Schaffner-Bielich \(2016\)](#); [Panotopoulos & Lopes \(2017a\)](#); [Jiménez & Fraga \(2022\)](#); [Panotopoulos & Lopes \(2018b\)](#); [Leung et al. \(2022\)](#); [Karkevandi et al. \(2022\)](#); [Lourenço et al. \(2022a\)](#); [Gleason et al. \(2022\)](#); [Dengler et al. \(2022\)](#); [Karkevandi et al. \(2022\)](#); [Panotopoulos & Lopes \(2018a, 2017c\)](#); [Miao et al. \(2022\)](#) and the two fluid approach is mostly adopted in such works. In this context it is worth mentioning that the interaction between DM and SQM in the DM admixed SQSs is not considered before in the existing literature to the best of our knowledge.

The accreted DM may eventually lead to the collapse of the star into a black hole. To prevent this, the interaction between

DM and the standard model constituents of the star matter must be extremely weak [Zheng & Chen \(2016\)](#). Therefore in the present work we invoke feeble interaction between SQM and fermionic DM χ . As stated earlier that for the description of SQM, in the present work we consider the vbag model that introduces a vector mediator (ω meson) to ensure quark interaction. Motivated by this fact, in the present work we include the vector new physics mediator ξ to study the effects of the feeble DM-SQM interaction on the DM admixed SQS properties. We choose the dark boson mediator ξ in the dark sector to be of vector type in order to maintain similarity with the pure SQM that involves the ω meson which is a vector mediator. Moreover, the vector ω meson and the vector dark boson ξ carry the same net spin. It is well known that the net spin of ω meson is 1 while that of the vector dark boson is also 1 [Hambye & Vanderheyden \(2020\)](#); [Arcadi et al. \(2020\)](#); [Gabrielli et al. \(2015\)](#). However, one can also consider a scalar mediator from the dark sector. Several works [Panotopoulos & Lopes \(2017b\)](#); [Bertoni et al. \(2013\)](#); [Nelson et al. \(2019\)](#); [Bhat & Paul \(2020\)](#); [Lourenço et al. \(2022b\)](#); [Quddus et al. \(2020\)](#); [Das et al. \(2019, 2020, 2021\)](#) have considered interaction between dark and hadronic matter via Higgs boson and have successfully computed the structural properties of DM admixed NSs. In our earlier works we introduced feeble interaction between DM fermion and β equilibrated hadronic NS matter (described by the hadronic relativistic mean field model [Sen \(2021\)](#)) via a dark scalar mediator ϕ [Sen & Guha \(2021\)](#) while in [Guha & Sen \(2021\)](#) we included same dark vector mediator ξ (as chosen in the present work) along with the scalar dark mediator ϕ in order to determine the structural properties of the DM admixed NSs. In the present work we consider only the vector dark mediator ξ in order to be consistent with the SQM sector which involves only vector mediator and also to match the net spin of the ω and ξ vector mediators. Similar to [Guha & Sen \(2021\)](#), in the present work the mass of DM fermion m_χ , mass of vector mediator m_ξ and the coupling y_ξ between them are determined in consistency with the self-interaction constraint from Bullet cluster [Tulin et al. \(2013\)](#); [Tulin & Yu \(2018\)](#); [Hambye & Vanderheyden \(2020\)](#) and from present day relic abundance [Belanger et al. \(2013\)](#); [Gondolo & Gelmini \(1991\)](#); [Guha et al. \(2019\)](#). In the present work we study the effects of variation of these DM parameters m_χ , m_ξ and y_ξ on the DM admixed SQSs in the light of various astrophysical constraints like the lower bound on maximum mass from PSR J0740+6620 [Fonseca et al. \(2021\)](#), the constraint on tidal deformability of a $1.4 M_\odot$ from GW170817 [Abbott et al. \(2018\)](#) and the NICER data for PSR J0030+0451 [Riley et al. \(2019\)](#); [Miller et al. \(2019\)](#). These astrophysical constraints on the structural properties of compact stars help us to obtain realistic EoS of compact stars to certain extent and to understand the possible composition of matter at such conditions. The nature of the massive secondary compact object associated with the detection of GW190814 [Abbott et al. \(2020\)](#) has remained inconclusive whether this object is a black hole (BH) or a NS. This is because any further information related to GW190814 like its electromagnetic counterpart or the tidal deformability of this compact object is still not obtained.

The paper is organized as follows. In the next section 2 we depict the formalism of the vbag model for SQSs in the presence of DM. In the following section 3 we present our

results along with relevant discussions. We finally conclude in the final section 4 of the paper.

2 FORMALISM

2.1 vBag Model with Dark Matter

We adopt the vector MIT Bag (vBag) model with the u , d and s quarks in presence of electrons. The mass of u and d quarks is very small compared to that of the s quark ($m_s=95$ MeV). The formalism of obtaining the EoS of such a system is well depicted in Lopes et al. (2021); Kumar et al. (2022). In addition to SQM in the model, we consider the presence of accreted DM. The interaction between the quarks and fermionic DM χ is mediated by the vector dark boson ξ of mass m_ξ . The complete modified Lagrangian of such a system is given as

$$\begin{aligned} \mathcal{L} = & \sum_f \left[\bar{\psi}_f \left(i\gamma_\mu \partial^\mu - g_\xi \gamma_\mu \xi^\mu - m_f \right) \psi_f - B \right] \Theta(\bar{\psi}_f \psi_f) \\ & - \sum_f g_{qqV} \left[\bar{\psi}_f \left(\gamma_\mu V^\mu \right) \psi_f \right] \Theta(\bar{\psi}_f \psi_f) + \frac{1}{2} m_V^2 V_\mu V^\mu - \frac{1}{4} V_{\mu\nu} V^{\mu\nu} \\ & + \bar{\psi}_l \left(i\gamma_\mu \partial^\mu - m_l \right) \psi_l - \frac{1}{4} F'_{\mu\nu} F'^{\mu\nu} + \frac{1}{2} m_\xi^2 \xi_\mu \xi^\mu \\ & + \bar{\chi} \left[i\gamma_\mu \partial^\mu - y_\xi \gamma_\mu \xi^\mu \right] \chi \end{aligned} \quad (1)$$

where, $f = u, d$ and s and the lepton $l=e$ is the electrons. B is the Bag constant and the Heaviside function $\Theta=1$ inside the bag. Here the quark interaction is mediated by the repulsive vector ω meson channel and thus $m_V=783$ MeV with g_{qqV} as the coupling strength. For simplicity, in the present work we consider universal coupling scheme i.e., $g_{uuV} = g_{ddV} = g_{ssV} = g_{qqV}$. The scaled coupling is defined as $G_V = (g_{qqV}/m_V)^2$. So taking $G_V=0$ reduces to the original form of the MIT Bag model without interactions. Refs. Lopes et al. (2021); Kumar et al. (2022) also introduced the self-interaction of the vector ω field via its quartic contribution in terms of a parameter b_4 that regulates the increment/decrement of the vacuum expectation value (V_0) of the ω field. This correction term also mimics the Dirac sea contribution of the quarks. However, in the present work we do not consider this self-interaction of the ω field since we intend to study the exclusive effects of DM and the DM parameters on the structural properties of the DM admixed SQSs. We consider the values of $G_V=0.3, 0.5$ and 0.7 . As discussed in the Introduction section 1, in order to calculate the stability window for B , we adopt the same procedure as Torres & Menezes (2013); Lopes et al. (2021); Ferrer et al. (2015) based on the criteria as $\varepsilon/\rho_B \leq m_n$ for the upper bound of B with 3 flavor SQM while the lower bound is obtained with the 2 flavor QM. Here we consider, $m_n=939$ MeV to obtain the stability window for B . We present the calculated allowed range of B for the considered values of G_V in table 1. Our calculated allowed range of B is slightly different from that of Torres & Menezes (2013); Lopes et al. (2021) since unlike them we have considered $m_n=939$ MeV. Consistent with Lopes et al. (2021), we find that the value of both B_{min} and B_{max} decrease with increase of G_V . To calculate the EoS given using Eqs. (7) and (8), we consider an average value of B between B_{max} and B_{min} for a particular value of G_V .

The terms $\bar{\psi}_f (g_\xi \gamma_\mu \xi^\mu) \psi_f$ of Eq. (1) indicate the interaction

Table 1. Stability window obtained for the vector MIT Bag model with $X_V = 1$ and $m_n=939$ MeV.

G_V	$B_{min}^{1/4}$ (MeV)	$B_{max}^{1/4}$ (MeV)
0.3	138	148
0.5	134	143
0.7	131	139

of the the quark fields ψ_f with the vector new physics mediator ξ from dark sector with a very feeble coupling strength $g_\xi \sim 10^{-4}$ as we assumed in our previous works Guha & Sen (2021); Sen & Guha (2021). We have checked that this coupling, being extremely small, the change in its order below 10^{-4} do not bring any significant change to the EoS and structure of DM admixed SQSs. The interaction between the fermionic DM χ and ξ is depicted in the last line of Eq. (1). The corresponding coupling between the two is denoted by y_ξ . The terms $\frac{1}{4} V_{\mu\nu} V^{\mu\nu}$ and $\frac{1}{4} F'_{\mu\nu} F'^{\mu\nu}$ in Eq. (1) denote the kinetic terms of the vectors fields ω and ξ , respectively. In the present work the higher order self-interaction terms of the dark mediator ξ has been neglected and has been chosen up to second order for simplicity. It is seen that the inclusion of the higher order terms do not contribute much and do not bring any substantial change to the results presented in the present work. The mass of the fermionic DM (m_χ) and the corresponding value of m_ξ is chosen in consistence with the self-interaction constraint from Bullet cluster Tulin et al. (2013); Tulin & Yu (2018); Hambye & Vanderheyden (2020) while the corresponding value of y_ξ is chosen in accordance to the present day relic abundance Belanger et al. (2013); Gondolo & Gelmini (1991); Guha et al. (2019). In our earlier work Guha & Sen (2021) we have already discussed in details the calculations of m_χ , m_ξ and y_ξ consistent with the mentioned constraints. However, for the sake of completeness, we discuss the DM parameters and their calculations. Bullet cluster observational data suggests an estimate of the self-interaction of the DM particles Randall et al. (2008); Bradac et al. (2006). The self-scattering transfer cross-section of DM fermions is typically in the range $\sigma_T/m_\chi \approx (0.1-10)$ cm²/gm Randall et al. (2008); Bradac et al. (2006); Dawson et al. (2012); Dave et al. (2001); Vogelsberger et al. (2012); Kahlhoefer et al. (2015). For the DM fermions of mass m_χ , self-scattered through the light vector mediator of mass m_ξ , the bullet cluster data estimates for the transfer cross-section as $\sigma_T/m_\chi \leq 1.25$ cm²/gm Randall et al. (2008); Robertson et al. (2017). The DM parameters involved in the present work viz. the mass (m_χ) of the DM fermions χ and the mass (m_ξ) of the light mediator ξ satisfy the self-interaction constraints from bullet cluster Tulin et al. (2013); Tulin & Yu (2018); Hambye & Vanderheyden (2020) as seen from Fig. 1. The coupling constant (y_ξ) for the interaction between the DM fermion χ and the light vector mediators from the hidden sector ξ is determined by satisfying the present day thermal relic abundances of DM. For detailed calculations one can refer to Belanger et al. (2013); Gondolo & Gelmini (1991); Guha et al. (2019). The chosen parameter sets of the dark sector are listed in table 2.

Applying mean field approximation, the equation of motion of the vector field in terms of V_0 reads

Table 2. Chosen values of self interacting DM m_χ and corresponding values of m_ξ from the constraints obtained from Bullet cluster. y_ξ have been fixed from observed relic abundance.

m_χ (GeV)	m_ξ (MeV)	y_ξ
25	50	0.27
50	60	0.32
75	20	0.40
100	10	0.46
150	6	0.50
200	4	0.52




Figure 1. Combination of m_χ and m_ξ satisfying the self-interaction constraint from bullet cluster [Randall et al. \(2008\)](#); [Tulin et al. \(2013\)](#). The color coding denotes the values of DM self-interaction transfer cross-section $\sigma_T/m_\chi = (0.1-1) \text{ cm}^2/\text{gm}$ (purple) and $(1-10) \text{ cm}^2/\text{gm}$ (sea-green).

$$V_0 = \frac{g_{qqV}}{m_V^2} \rho \quad (2)$$

where, the total quark density is given as

$$\rho = \langle \psi_f^\dagger \psi_f \rangle = (\rho_u + \rho_d + \rho_s) = \frac{\gamma_f}{6\pi^2} \sum_f k_{Ff}^3 \quad (3)$$

Here k_{Ff} is the Fermi momenta of the quarks and $\gamma_f = 6$ for quarks.

The vacuum expectation value (ξ_0) of the vector dark boson is

$$\xi_0 = \frac{g_\xi \rho + y_\xi \rho_\chi}{m_\xi^2} \quad (4)$$

where, the DM density is

$$\rho_\chi = \frac{\gamma_f}{6\pi^2} k_{F\chi}^3 \quad (5)$$

Based on the discussion given in [Panotopoulos & Lopes](#)

(2017b); [Guha & Sen \(2021\)](#), for the present work a constant number density of the fermionic DM has been assumed throughout the radius profile of the star. This estimate of constant number density (ρ_χ) of fermionic DM is ~ 1000 times smaller than the average baryon number density of the SQM. In absence of any proper experimental or observational evidence regarding the presence and distribution of DM in compact stars, many recent works [Bhat & Paul \(2020\)](#); [Sen & Guha \(2021\)](#); [Guha & Sen \(2021\)](#); [Quddus et al. \(2020\)](#); [Das et al. \(2019, 2020, 2021\)](#) have also considered the same assumption of constant number density of DM throughout the density profile of NSs. Going with this assumption, the Fermi momentum of the DM fermions $k_{F\chi}$ also turns out to be constant. Therefore in the present work we fix $k_{F\chi} = 0.03 \text{ GeV}$.

The quark chemical potential in presence of DM is modified as

$$\mu_f = \sqrt{k_f^2 + m_f^2} + g_{qqV} V_0 + g_\xi \xi_0 \quad (6)$$

The chemical equilibrium conditions and the charge neutrality condition for the quarks are additionally imposed.

The complete EoS is given as follows. The energy density is

$$\begin{aligned} \varepsilon = & \frac{1}{2} \frac{g_{qqV}^2}{m_V^2} \rho^2 + \frac{1}{2} \frac{(g_\xi \rho + y_\xi \rho_\chi)^2}{m_\xi^2} + \frac{\gamma_f}{2\pi^2} \sum_f \int_0^{k_{Ff}} \sqrt{k_f^2 + m_f^2} k_f^2 dk_f \\ & + \frac{\gamma_l}{2\pi^2} \int_0^{k_{Fl}} \sqrt{k_l^2 + m_l^2} k_l^2 dk_l + \frac{\gamma_\chi}{2\pi^2} \int_0^{k_{F\chi}} \sqrt{k_\chi^2 + m_\chi^2} k_\chi^2 dk_\chi + B \end{aligned} \quad (7)$$

while the pressure is

$$\begin{aligned} P = & \frac{1}{2} \frac{g_{qqV}^2}{m_V^2} \rho^2 + \frac{1}{2} \frac{(g_\xi \rho + y_\xi \rho_\chi)^2}{m_\xi^2} + \frac{\gamma_f}{6\pi^2} \sum_f \int_0^{k_{Ff}} \frac{k_f^4 dk_f}{\sqrt{k_f^2 + m_f^2}} \\ & + \frac{\gamma_l}{6\pi^2} \int_0^{k_{Fl}} \frac{k_l^4 dk_l}{\sqrt{k_l^2 + m_l^2}} + \frac{\gamma_\chi}{6\pi^2} \int_0^{k_{F\chi}} \frac{k_\chi^4 dk_\chi}{\sqrt{k_\chi^2 + m_\chi^2}} - B \end{aligned} \quad (8)$$

With the obtained EoS we proceed to calculate the structural properties of the SQSs.

2.2 Structural Properties of Quark Stars

With the obtained EoS, the structural properties like the gravitational mass (M) and the radius (R) of the DM admixed SQSs are computed by integrating the following Tolman-Oppenheimer-Volkoff (TOV) equations [Tolman \(1939\)](#); [Oppenheimer & Volkoff \(1939\)](#) based on the hydrostatic equilibrium between gravity and the internal pressure of the star.

$$\frac{dP}{dr} = -\frac{G}{r} \frac{(\varepsilon + P)(M + 4\pi r^3 P)}{(r - 2GM)}, \quad (9)$$

$$\frac{dM}{dr} = 4\pi r^2 \varepsilon, \quad (10)$$

The deformation of the metric $h_{\alpha\beta}$ in Regge-Wheeler gauge is given as [Hinderer \(2008\)](#); [Hinderer et al. \(2010\)](#)

$$h_{\alpha\beta} = \text{diag}[e^{-\nu(r)} H_0, e^{\lambda(r)} H_2, r^2 K(r), r^2 \sin^2 \theta K(r)] Y_{2m}(\theta, \phi) \quad (11)$$

The tidal Love number k_2 is obtained in terms of the compactness ($C = M/R$) and a quantity y which in case of QSs is defined as [Hinderer et al. \(2010\)](#); [Kumar et al. \(2022\)](#)

$$y = \frac{RH'(R)}{H(R)} - \frac{4\pi R^3 \varepsilon_s}{M(R)} \quad (12)$$

where, ε_s is the energy density at the surface of the QS. The tidal deformability parameter λ in terms of k_2 is given as

$$\lambda = \frac{2}{3} k_2 R^5 \quad (13)$$

The dimensionless tidal deformability Λ is then calculated as a function of Love number, gravitational mass and radius [Hinderer \(2008\)](#); [Hinderer et al. \(2010\)](#) as

$$\Lambda = \frac{2}{3} k_2 (R/M)^5 \quad (14)$$

3 RESULTS

We compute the EoS of SQS both in presence and absence of DM. For the purpose we consider the values of $G_V=0.3, 0.5$ and 0.7 . In the present work, for each value of G_V we calculate the EoS with a corresponding value of B which is the average of B_{max} and B_{min} presented in table 1. The DM admixed SQM EoS is computed for three values of $m_\chi=50, 75$ and 100 GeV. For each value of m_χ , m_ξ and y_ξ are calculated following [Guha & Sen \(2021\)](#) in order to be consistent with the constraints from Bullet cluster and present day relic density bound. With the obtained EoS for SQM and DM admixed SQM, we obtain the structural properties like the mass M , radius R and tidal deformability Λ of both SQSs and DM admixed SQSs following the formalism mentioned in section 2.2.

In Fig. 2 we show our results of mass and radius (Fig. 2a) and tidal deformability (Fig. 2b) with a lower value of $G_V=0.3$ and corresponding value of B for both SQSs and DM admixed SQSs. From Fig. 2a we find that the maximum mass M_{max} of the star decreases as more massive DM fermion is considered. Compared to the no-DM (SQS) case, the constraint on mass from the secondary component of GW190814 [Abbott et al. \(2020\)](#) is hardly satisfied in case of DM admixed SQSs. However, it is still debatable in literature whether this object is a compact star or a black hole. For both SQSs and DM admixed SQSs the $M-R$ constraints from GW170817 [Abbott et al. \(2018\)](#) and PSR J0740+6620 [Fonseca et al. \(2021\)](#); [Miller et al. \(2021\)](#); [Riley et al. \(2021\)](#) are well satisfied. However, the constraint on the $M-R$ plane from NICER data for PSR J0030+0451 [Riley et al. \(2019\)](#); [Miller et al. \(2019\)](#), though satisfied by SQS, is hardly satisfied by the DM admixed SQSs configurations. In Fig. 2b we find that for the low value of $G_V(=0.3)$ the constraints on $\Lambda_{1.4}$ from both GW170817 and GW190814 are satisfied with $m_\chi=75$ and 100 GeV. For $m_\chi=50$ GeV and the no-DM only the constraint from GW190814 is satisfied. For the no-DM case our result is consistent with [Kumar et al. \(2022\)](#).

We next investigate the effect on the structural properties of SQSs and DM admixed SQSs in Fig. 3 with an intermediate value of $G_V=0.5$ and the corresponding value of B .





Figure 2. (a) Mass-radius relationship of dark matter admixed quark stars for different $G_V=0.3$. Observational limits imposed from the most massive pulsar PSR J0740+6620 [Fonseca et al. \(2021\)](#); [Miller et al. \(2021\)](#); [Riley et al. \(2021\)](#) are also indicated. Mass of secondary component of GW190814 [Abbott et al. \(2020\)](#) is also shown. The constraints on $M-R$ plane prescribed from GW170817 [Abbott et al. \(2018\)](#) and NICER experiment for PSR J0030+0451 [Riley et al. \(2019\)](#); [Miller et al. \(2019\)](#) are also compared. (b) Corresponding variation of tidal deformability with respect to mass. Constraint on $\Lambda_{1.4}$ from GW170817 [Abbott et al. \(2018\)](#) and GW190814 [Abbott et al. \(2020\)](#) observations are also shown.




Figure 3. Same as Fig. 2 but for $G_V = 0.5$.

For DM admixed SQSs the same values of m_χ are considered same as that in case of Fig. 2. Comparing Figs. 2a and 3a we find that both $M_{\max}$ and the corresponding radius of SQSs increase with increasing values of G_V . This is also consistent with Lopes et al. (2021); Kumar et al. (2022). This fact is also true for the DM admixed SQSs for any particular value of m_χ . For DM admixed SQSs the various constraints on the $M-R$ plane are better satisfied with a higher value of G_V . From Fig. 3a we find that the constraint from NICER

data for PSR J0030+0451, which was hardly satisfied with $G_V = 0.3$, is now well satisfied for $G_V = 0.5$ with all the three chosen values of m_χ . We also note that unlike the DM admixed SQSs obtained with $G_V = 0.3$, constraint on mass from the secondary component of GW190814 is well satisfied by that with $G_V = 0.5$. Similar to the case of the lower value of G_V , our results with $G_V = 0.5$ are in good agreement with the other constraints from GW170817 and PSR J0740+6620 on the $M-R$ relationship of compact stars. In Fig. 3b we find that for the moderate value of $G_V (=0.5)$ the constraints on $\Lambda_{1.4}$ from both GW170817 and GW190814 are satisfied with $m_\chi = 100$ GeV. The rest including that for the no-DM case satisfy only the one from GW190814.

We finally present our results with a high value of $G_V = 0.7$ in Fig. 4. From Fig. 4a we find that $M_{\max}$ offshoots the theoretical limit on the upper bound of compact stars ($> 3 M_\odot$) in case of the pure SQS without DM. The presence of DM with all the chosen values of m_χ reduces the maximum mass to an acceptable range. All the constraints on the mass-radius relationship of compact stars are well fulfilled by all the DM admixed SQSs configurations. Fig. 4b suggests that for the high value of $G_V (=0.7)$ the constraints on $\Lambda_{1.4}$ from GW170817 is satisfied by none of the configurations. The constraint from GW190814 is satisfied with only $m_\chi = 75$ and 100 GeV.

Comparing Figs. 2a, 3a and 4a, we find that for any particular value of the m_χ , both mass and radius of the DM admixed SQSs increase with increasing values of G_V . Thus we find that the NICER data for PSR J0030+0451 is better satisfied with $G_V > 0.3$. With $G_V = 0.3$ this constraint is marginally satisfied only for $m_\chi = 50$ GeV. For $G_V < 0.3$ the aforesaid constraint is not satisfied with any of the chosen values of m_χ . We therefore chose values of $G_V > 0.3$ in the present work. Overall, in the present work we obtain very massive SQS configurations even in the presence of DM. In most cases the maximum mass is high enough to be comparable with that of the secondary component of GW190814 merger event. Although it is at present debatable whether it is a BH or NS, the results of this work suggest that this compact object may be a DM admixed SQS.

We next study the variation of $R_{1.4}$ and $\Lambda_{1.4}$ with respect to m_χ . In Fig. 5 we show the same. For all the values of G_V , both $R_{1.4}$ and $\Lambda_{1.4}$ increase with decreasing values of m_χ . We also notice that for very high values of m_χ the constraint on $\Lambda_{1.4}$ from GW170817 is not well satisfied. This effect becomes more pronounced for lower values of G_V . Each line in Figs. 5a and 5b are the fitted function for the variation of $R_{1.4}$ and $\Lambda_{1.4}$, respectively with respect to m_χ for the corresponding value of G_V . In absence of any direct relation between $R_{1.4}$ and $\Lambda_{1.4}$ with respect to m_χ , we have obtained the specific (fitted) relationship between m_χ and $R_{1.4}$ and $\Lambda_{1.4}$ individually within the scope of the model. From these fits, the particular relation between $R_{1.4}$ and m_χ for different G_V is obtained as

$$R_{1.4} = 7.677 + 4.346 \exp\left(-\frac{m_\chi}{542.118}\right), \text{ for } G_V = 0.3 \quad (15)$$

$$R_{1.4} = 8.321 + 4.138 \exp\left(-\frac{m_\chi}{459.488}\right), \text{ for } G_V = 0.5 \quad (16)$$

and





Figure 4. Same as Fig. 2 but for $G_V=0.7$.

$$R_{1.4} = 9.471 + 3.665 \exp\left(-\frac{m_\chi}{319.798}\right), \text{ for } G_V = 0.7 \quad (17)$$

while that between $\Lambda_{1.4}$ and m_χ are

$$\Lambda_{1.4} = 257.08 + 553.23 \exp\left(-\frac{m_\chi}{139.95}\right), \text{ for } G_V = 0.3 \quad (18)$$

$$\Lambda_{1.4} = 374.21 + 726.70 \exp\left(-\frac{m_\chi}{106.13}\right), \text{ for } G_V = 0.5 \quad (19)$$





Figure 5. Variations of (a) $R_{1.4}$ and (b) $\Lambda_{1.4}$ with respect to m_χ for different values of G_V .

and

$$\Lambda_{1.4} = 558.56 + 1067.29 \exp\left(-\frac{m_\chi}{64.80}\right), \text{ for } G_V = 0.7 \quad (20)$$

It can be seen from Figs. 5a and 5b that compared to the constraint on $\Lambda_{1.4}$, that on $R_{1.4}$ is better satisfied with the chosen values of m_χ for different G_V . Moreover, as mentioned earlier, that for all the chosen values of m_χ , the NICER data for PSR J0030+0451 is better satisfied with $G_V > 0.3$. However, there are still many uncertainties pertaining to compact star physics viz. the composition and EoS, presence of DM and its interaction with the standard model particles. Moreover, the SQM EoS is also obtained with the vbag model in the present work but there are various other well-known models to describe the SQM properties. Under such circumstances, prescribing any specific bound on the DM mass with respect to the various astrophysical constraints is beyond the scope of this work. However, within the scope of the present study, we can roughly conclude that with an average value of $m_\chi \sim 100$ GeV the present day astrophysical constraints on

compact star structural properties could be satisfied. Therefore instead of providing any specific bound on m_χ , we have focused on obtaining the specific (fitted) relationship and dependence of $R_{1.4}$ and $\Lambda_{1.4}$ on m_χ for a specific value of G_V . Within the scope of this study the relations presented in Eqs. (15), (16) and (17) show the specific (fitted) relationship between $R_{1.4}$ and m_χ while (18), (19) and (20) show the same between $\Lambda_{1.4}$ and m_χ .

4 SUMMARY AND CONCLUSION

We have computed the structural properties of DM admixed SQSs. We consider the vBag model to describe the SQM with the bag constant been determined by following the stability condition of the SQM in terms of the energy density per baryon. With the notion such compact stars, being highly gravitating objects, we study the possible presence of accreted DM in SQSs thereby forming DM admixed SQSs. We introduce a feeble interaction between pure SQM and the accreted fermionic DM via a vector dark boson mediator. The masses of the DM fermion, mass of vector mediator and the coupling between them are determined in consistence with the self-interaction constraint from Bullet cluster and from present day relic abundance, respectively.

Overall, we find that the presence of DM reduces both the mass and radius of the star compared to the no-DM case. We find that the presence of more massive DM particle leads to the less massive DM admixed SQSs with smaller radius. It is also seen that the various astrophysical constraints on the mass-radius relationship of the DM admixed SQSs is better satisfied with higher values of the vector coupling (G_V) between the quarks. With the chosen values of m_χ and corresponding values of m_ξ and y_ξ , the calculated structural properties (gravitational mass, radius and tidal deformability) of the DM admixed SQS configurations satisfy all the various present day astrophysical constraints obtained from the pulsars like PSR J0740+6620 and PSR J0030+0451 and the gravitational wave data of GW170817 and GW190814.

ACKNOWLEDGMENTS

Work of A.G. is supported by the National Research Foundation of Korea (NRF-2019R1C1C1005073).

DATA AVAILABILITY

The data underlying this article are available within the article.

REFERENCES

Aalbers J., et al., 2022, arxiv, 2207.03764
 Abbott B. P., et al., 2018, *Phys. Rev. Lett.*, 121, 161101
 Abbott R., et al., 2020, *Astrophys. J. Lett.*, 896, L44
 Aghanim N., et al., 2020, *Astron. Astrophys.*, 641, A6
 Agnes P., et al., 2018, *Phys. Rev. Lett.*, 121, 111303
 Agnese R., et al., 2018, *Phys. Rev. Lett.*, 121, 051301
 Akerib D. S., et al., 2013, *Astropart. Phys.*, 45, 34
 Alford M., Braby M., Paris M. W., Reddy S., 2005, *Astrophys. J.*, 629, 969

Aprile E., et al., 2012, *Phys. Rev. Lett.*, 109, 181301
 Aprile E., et al., 2018, *Phys. Rev. Lett.*, 121, 111302
 Arcadi G., Djouadi A., Kado M., 2020, *Phys. Lett. B*, 805, 135427
 Aziz A., Ray S., Rahaman F., Khlopov M., Guha B. K., 2019, *Int. J. Mod. Phys. D*, 28, 1941006
 Bauer M., Plehn T., 2019, Yet Another Introduction to Dark Matter: The Particle Physics Approach. Lecture Notes in Physics Vol. 959, Springer ([arXiv:1705.01987](https://arxiv.org/abs/1705.01987)), doi:10.1007/978-3-030-16234-4
 Belanger G., Boudjema F., Pukhov A., Semenov A., 2013, *Comput. Phys. Commun.*, 185
 Bennett C. L., et al., 2013, *Astrophys. J. Suppl.*, 208, 20
 Bertone G., Hooper D., Silk J., 2005, *Phys. Rept.*, 405, 279
 Bertoni B., Nelson A. E., Reddy S., 2013, *Phys. Rev. D*, 88, 123505
 Bhat S. A., Paul A., 2020, *Eur. Phys. J. C*, 80, 544
 Bodmer A. R., 1971, *Phys. Rev. D*, 4, 1601
 Bradac M., et al., 2006, *Astrophys. J.*, 652, 937
 Cannoni M., 2016, *Eur. Phys. J. C*, 76, 137
 Chin S. A., Kerman A. K., 1979, *Phys. Rev. Lett.*, 43, 1292
 Chodos A., Jaffe R. L., Johnson K., Thorn C. B., Weisskopf V. F., 1974, *Phys. Rev. D*, 9, 3471
 Cierniak M., Fischer T., Bastian N.-U., Klähn T., Salinas M., 2019, *Universe*, 5, 186
 Crisler M., Essig R., Estrada J., Fernandez G., Tiffenberg J., Sofor haro M., Volansky T., Yu T.-T., 2018, *Phys. Rev. Lett.*, 121, 061803
 Das A., Malik T., Nayak A. C., 2019, *Phys. Rev. D*, 99, 043016
 Das H. C., Kumar A., Kumar B., Kumar Biswal S., Nakatsukasa T., Li A., Patra S. K., 2020, *Mon. Not. Roy. Astron. Soc.*, 495, 4893
 Das H. C., Kumar A., Patra S. K., 2021, *Phys. Rev. D*, 104, 063028
 Dave R., Spergel D. N., Steinhardt P. J., Wandelt B. D., 2001, *Astrophys. J.*, 547, 574
 Dawson W. A., et al., 2012, *Astrophys. J. Lett.*, 747, L42
 Deliyergiyev M., Del Popolo A., Tolos L., Le Delliou M., Lee X., Burgio F., 2019, *Phys. Rev. D*, 99, 063015
 Dengler Y., Schaffner-Bielich J., Tolos L., 2022, *Phys. Rev. D*, 105, 043013
 Ellis J., Hütsi G., Kannike K., Marzola L., Raidal M., Vaskonen V., 2018, *Phys. Rev. D*, 97, 123007
 Farhi E., Jaffe R. L., 1984, *Phys. Rev. D*, 30, 2379
 Ferrer E. J., de la Incera V., Paulucci L., 2015, *Phys. Rev. D*, 92, 043010
 Fonseca E., et al., 2021, *Astrophys. J. Lett.*, 915, L12
 Fraga E. S., Pisarski R. D., Schaffner-Bielich J., 2001, *Phys. Rev. D*, 63, 121702
 Franzon B., Gomes R. O., Schramm S., 2016, *Mon. Not. Roy. Astron. Soc.*, 463, 571
 Gabrielli E., Marzola L., Raidal M., Veermäe H., 2015, *JHEP*, 08, 150
 Gleason T., Brown B., Kain B., 2022, *Phys. Rev. D*, 105, 023010
 Glendenning N. K., 2000, Compact stars: Nuclear physics, particle physics, and general relativity. Springer-Verlag, New York
 Gondolo P., Gelmini G., 1991, *Nucl. Phys. B*, 360, 145
 Guha A., Sen D., 2021, *JCAP*, 09, 027
 Guha A., Dev P. S. B., Das P. K., 2019, *JCAP*, 02, 032
 Hambye T., Vanderheyden L., 2020, *JCAP*, 05, 001
 Hinderer T., 2008, *Astrophys. J.*, 677, 1216
 Hinderer T., Lackey B. D., Lang R. N., Read J. S., 2010, *Phys. Rev. D*, 81, 123016
 Jiménez J. C., Fraga E. S., 2022, *Universe*, 8, 34
 Kahlhoefer F., Schmidt-Hoberg K., Kummer J., Sarkar S., 2015, *Mon. Not. Roy. Astron. Soc.*, 452, L54
 Karkevandi D. R., Shakeri S., Sagun V., Ivanytskyi O., 2022, *Phys. Rev. D*, 105, 023001
 Klähn T., Fischer T., 2015, *Astrophys. J.*, 810, 134
 Kumar A., Thapa V. B., Sinha M., 2022, *Mon. Not. Roy. Astron. Soc.*, 513, 3788

- Leung K.-L., Chu M.-c., Lin L.-M., 2022, *Phys. Rev. D*, 105, 123010
- Li A., Huang F., Xu R.-X., 2012, *Astropart. Phys.*, 37, 70
- Lopes I., Panotopoulos G., 2018, *Phys. Rev. D*, 97, 024030
- Lopes L. L., Biesdorf C., Menezes D. e. P., 2021, *Phys. Scripta*, 96, 065303
- Lourengo O., Frederico T., Dutra M., 2022a, *Phys. Rev. D*, 105, 023008
- Lourengo O., Lenzi C. H., Frederico T., Dutra M., 2022b, *Phys. Rev. D*, 106, 043010
- Miao Z., Zhu Y., Li A., Huang F., 2022, *Astrophys. J.*, 936, 69
- Miller M. C., et al., 2019, *Astrophys. J. Lett.*, 887, L24
- Miller M. C., et al., 2021, *Astrophys. J. Lett.*, 918, L28
- Mukhopadhyay P., Schaffner-Bielich J., 2016, *Phys. Rev. D*, 93, 083009
- Mukhopadhyay S., Atta D., Imam K., Basu D. N., Samanta C., 2017, *Eur. Phys. J. C*, 77, 440
- Nandi R., Char P., 2018, *Astrophys. J.*, 857, 12
- Nandi R., Pal S., 2021, *Eur. Phys. J. ST*, 230, 551
- Nelson A., Reddy S., Zhou D., 2019, *JCAP*, 07, 012
- Olinto A. V., 1987, *Phys. Lett. B*, 192, 71
- Oppenheimer J. R., Volkoff G. M., 1939, *Phys. Rev.*, 55, 374
- Panotopoulos G., Lopes I., 2017a, *Phys. Rev. D*, 96, 023002
- Panotopoulos G., Lopes I., 2017b, *Phys. Rev. D*, 96, 083004
- Panotopoulos G., Lopes I., 2017c, *Phys. Rev. D*, 96, 083013
- Panotopoulos G., Lopes I., 2018a, *Int. J. Mod. Phys. D*, 27, 1850093
- Panotopoulos G., Lopes I., 2018b, *Phys. Rev. D*, 98, 083001
- Quddus A., Panotopoulos G., Kumar B., Ahmad S., Patra S. K., 2020, *J. Phys. G*, 47, 095202
- Randall S. W., Markevitch M., Clowe D., Gonzalez A. H., Bradac M., 2008, *Astrophys. J.*, 679, 1173
- Rezaei Z., 2017, *Astrophys. J.*, 835, 33
- Riley T. E., et al., 2019, *Astrophys. J. Lett.*, 887, L21
- Riley T. E., et al., 2021, *Astrophys. J. Lett.*, 918, L27
- Robertson A., Massey R., Eke V., 2017, *Mon. Not. Roy. Astron. Soc.*, 465, 569
- Sen D., 2021, *J. Phys. G*, 48, 025201
- Sen D., Guha A., 2021, *Mon. Not. Roy. Astron. Soc.*, 504, 3354
- Tanabashi M., et al., 2018, *Phys. Rev. D*, 98, 030001
- Tolman R. C., 1939, *Phys. Rev.*, 55, 364
- Tolos L., Schaffner-Bielich J., 2015, *Phys. Rev. D*, 92, 123002
- Torres J. R., Menezes D. P., 2013, *EPL*, 101, 42003
- Tulin S., Yu H.-B., 2018, *Phys. Rept.*, 730, 1
- Tulin S., Yu H.-B., Zurek K. M., 2013, *Phys. Rev. D*, 87, 115007
- Vogelsberger M., Zavala J., Loeb A., 2012, *Mon. Not. Roy. Astron. Soc.*, 423, 3740
- Wang Q., et al., 2020, *Chin. Phys. C*, 44, 125001
- Wei W., Irving B., Klähn T., Jaikumar P., 2019, *Astrophys. J.*, 887, 151
- Weissenborn S., Sagert I., Pagliara G., Hempel M., Schaffner-Bielich J., 2011, *Astrophys. J. Lett.*, 740, L14
- Witten E., 1984, *Phys. Rev. D*, 30, 272
- Yang S.-H., PI C.-M., Zheng X.-P., Weber F., 2020, *Astrophys. J.*, 902, 32
- Zheng H., Chen L.-W., 2016, *Astrophys. J.*, 831, 127
- Zhou E.-P., Zhou X., Li A., 2018, *Phys. Rev. D*, 97, 083015