

Quantum Correction to Hawking Temperature and Thermodynamics of Schwarzschild AdS black holes

Prosenjit Paul^{*1}

¹Indian Institute Of Engineering Science and Technology (IIST), Shibpur-711103, WB, India

Abstract

We investigate quantum corrections to $4D$ Schwarzschild Anti de Sitter black hole using an effective field theory approach. We compute the quantum corrected Schwarzschild AdS spacetime upto order \mathcal{G}^2 . The Hawking temperature of Schwarzschild AdS black hole is computed upto order $\mathcal{O}(G^2)$ when the emitted object is massless and massive. We use the expression of the modified horizon radius to compute the quantum corrected entropy upto order $\mathcal{O}(G^2)$. Finally, we study quantum corrected thermodynamic parameters specific heat, free energy and pressure.

^{*}prosenjitpaul629@gmail.com

Contents

1	Introduction	2
2	Quantum correction to schwarzschild AdS metric	2
3	Quantum correction to Hawking temperature for Schwarzschild AdS black hole	4
3.1	Quantum correction to Hawking temperature	5
3.1.1	Massless particles	5
3.1.2	Massive particles	7
4	Quantum correction to thermodynamics of Schwarzschild AdS black hole	8
4.1	Quantum correction to entropy	8
4.2	Quantum correction to free energy and pressure	9
5	Conclusions	10

1 Introduction

Black hole are fascinating and elegant objects. They are described very accurately by a small number of macroscopic parameters (e.g., mass, angular momentum, and charge). The surprising discovery that black holes behave as thermodynamic objects has radically affected our understanding of general relativity and its relationship to quantum field theory. In the early 1970s, Bekenstein [1, 2] showed that black holes entropy proportional to area of the event horizon. In 1974, Hawking’s original discovery [3, 4] has showed that black holes can emit thermal radiation at a temperature inversely proportional to their mass. In particular there have been detailed studies [5, 6, 7, 8, 9] of Hawking temperature and radiation. Particle emission from a non-rotating and rotating black hole has calculated by Page in refs. [5, 6] In 1999, Wilczek and Parikh showed that Hawking radiation can be viewed as a tunnelling process [7], in which particles pass through the contracting horizon of the black hole, lending formal justification to the most intuitive picture we have of black hole evaporation.

In general, any correction to the black hole entropy and temperature leads to a correction of the black hole thermodynamics. As the black hole entropy is obtained using area of the event horizon and it is natural to expect that the quantum corrections modify the black hole metric as to reproduce the corrected entropy. Using Effective field theory one can obtain the quantum corrected metric. The quantum corrected metric of Schwarzschild black holes has been obtained recently [10] using Effective field theory. The effects of quantum correction on the Hawking temperature, thermodynamics and greybody factors of Schwarzschild black holes studied in [11]. Quantum correction on Reissner-Nordstrom black holes using Effective field theory approach studied in Refs. [12, 13].

The main aim of the original paper [10] is to describe the metric inside and outside of a static and spherical star, taking into account quantum effects. However, the authors also proposed that the exterior metric may provide a good model of quantum black holes. The aim of this paper is to consider quantum correction to Hawking temperature (for massless & massive particles) and thermodynamics of a static and spherical Schwarzschild AdS black hole. In section 2, we introduce the effective field theory approach to quantum gravity and obtain the solution of the equations of motion that represents the quantum corrected metric of a Schwarzschild AdS black hole up to order $\mathcal{O}(G^2)$. In section 3, we compute quantum corrected Hawking temperature of Schwarzschild AdS black hole for massless and massive particles up to order $\mathcal{O}(G^2)$. In section 4, we compute quantum correction to entropy, specific heat and Helmholtz free energy and pressure. Finally, in section 5, we summarise our results. We take $c = k_B = \hbar = 1$ throughout.

2 Quantum correction to schwarzschild AdS metric

In this section, we study quantum corrected Schwarzschild AdS metric up to order $\mathcal{O}(G^2)$ obtained from Effective Field Theory (EFT). The quantum correction to Schwarzschild metric is given by [14]. The action of the theory is

$$\Gamma[g] = \Gamma_L[g] + \Gamma_{NL}[g]. \quad (2.1)$$

where the local part of the action is given by

$$\Gamma_L[g] = \int d^4x \sqrt{-g} \left\{ \frac{R + 2\Lambda}{16\pi G} + c_1(\mu)R^2 + c_2(\mu)R_{\mu\nu}R^{\mu\nu} + c_3(\mu)R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} \right\}, \quad (2.2)$$

where Λ is cosmological constant and μ is an energy scale. The exact values of the coefficients c_1 , c_2 and c_3 are unknown, as they depend on the nature of the ultra-violet theory of quantum gravity. The non-local part of the action is

$$\Gamma_{NL}(g) = - \int d^4x \sqrt{-g} \left\{ \alpha R \ln\left(\frac{\square}{\mu^2}\right) R + \beta R_{\mu\nu} \ln\left(\frac{\square}{\mu^2}\right) R^{\mu\nu} + \gamma R_{\mu\nu\rho\sigma} \ln\left(\frac{\square}{\mu^2}\right) R^{\mu\nu\rho\sigma} \right\}, \quad (2.3)$$

the parameters α , β and γ are non-local Wilson coefficients for fields. The values of the parameters for different fields have been listed in ref. [10]. $\square = g^{\mu\nu} \nabla_\mu \nabla_\nu$ and the operator $\ln(\frac{\square}{\mu^2})$ has integral representation[12]

$$\ln\left(\frac{\square}{\mu^2}\right) = \int_0^\infty ds \left(\frac{1}{\mu^2 + s} - \frac{1}{\square + s} \right). \quad (2.4)$$

Using both the local and non-local Gauss-Bonnet identities [14], it is possible to eliminate the Riemann tensor by redefining the coefficients as

$$\begin{aligned} c_1 &\rightarrow \bar{c}_1 = c_1 - c_3, & c_2 &\rightarrow \bar{c}_2 = c_2 + 4c_3, & c_3 &\rightarrow \bar{c}_3 = 0, \\ \alpha &\rightarrow \bar{\alpha} = \alpha - \gamma, & \beta &\rightarrow \bar{\beta} = \beta + 4\gamma, & \gamma &\rightarrow \bar{\gamma} = 0. \end{aligned} \quad (2.5)$$

The quantum corrected Einstein equations can be obtained by varying action 2.1 with respect to the metric,

$$(G_{\mu\nu} - \Lambda g_{\mu\nu}) + 16\pi G (H_{\mu\nu}^L + H_{\mu\nu}^{NL}) = 0. \quad (2.6)$$

where $G_{\mu\nu}$ is Einstein tensor given by

$$G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2} R g_{\mu\nu}, \quad (2.7)$$

The local part of the equation of motion is given by

$$\begin{aligned} H_{\mu\nu}^L &= \bar{c}_1 \left(2R R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R^2 - 2\nabla_\mu \nabla_\nu R + 2g_{\mu\nu} \square R \right) \\ &+ \bar{c}_2 \left(2R^{\rho\sigma} R_{\mu\rho\nu\sigma} - \frac{1}{2} g_{\mu\nu} R_{\rho\sigma} R^{\rho\sigma} - \nabla_\mu \nabla_\nu R + \square R_{\mu\nu} + \frac{1}{2} g_{\mu\nu} \square R \right), \end{aligned} \quad (2.8)$$

The non-local contribution is

$$\begin{aligned} H_{\mu\nu}^{NL} &= -2\bar{\alpha} \left(R_{\mu\nu} - \frac{1}{4} g_{\mu\nu} R + g_{\mu\nu} \square - \nabla_\mu \nabla_\nu \right) \ln\left(\frac{\square}{\mu^2}\right) R - \bar{\beta} \left(\delta_\mu^\rho R_{\nu\sigma} + \delta_\nu^\rho R_{\mu\sigma} - \frac{1}{2} g_{\mu\nu} R_\sigma^\rho + \delta_\mu^\rho g_{\nu\sigma} \square \right. \\ &\quad \left. + g_{\mu\nu} \nabla^\rho \nabla_\sigma - \delta_\mu^\rho \nabla_\rho \nabla_\nu - \delta_\nu^\rho \nabla_\sigma \nabla_\mu \right) \ln\left(\frac{\square}{\mu^2}\right) R_\rho^\sigma. \end{aligned} \quad (2.9)$$

Classical four dimension Schwarzschild AdS black hole metric is[15]

$$ds^2 = \left(1 - \frac{2GM}{r} + \frac{r^2}{a^2} \right) dt^2 - \left(1 - \frac{2GM}{r} + \frac{r^2}{a^2} \right)^{-1} dr^2 + r^2 d\Omega^2, \quad (2.10)$$

where $d\Omega^2 = d\theta^2 + \sin^2 \theta d\phi^2$, a is the AdS radius & it is related to the cosmological constant

$$a = \sqrt{\frac{3}{|\Lambda|}}. \quad (2.11)$$

Now we perturb metric,

$$\bar{g}_{\mu\nu} = g_{\mu\nu} + g_{\mu\nu}^q. \quad (2.12)$$

We now consider a small perturbation of order $\mathcal{O}(G)$. The equation of motion becomes

$$\left(G_{\mu\nu}^L[g^q] - \Lambda g_{\mu\nu}[g^q]\right) + 16\pi G \left(H_{\mu\nu}^L[q] + H_{\mu\nu}^{NL}[q]\right) = 0. \quad (2.13)$$

In case of Schwarzschild AdS solution Ricci scalar and tensor, both are non zero. The action of $\ln(\frac{\square}{\mu^2})$ on several radial functions are collected in [10, 16, 13]. The final solution of the quantum corrected equations of motion (2.13) outside ($r > R$) the star is

$$ds^2 = -f(r)dt^2 + g(r)dr^2 + r^2 d\Omega^2, \quad (2.14)$$

where $d\Omega^2 = d\theta^2 + \sin^2\theta d\phi^2$ and the metric function is given by

$$f(r) = \left(1 - \frac{2GM}{r} + \frac{r^2}{a^2}\right) + \tilde{\alpha} \frac{2G^2 M}{R^3} \left(2\frac{R}{r} + \ln\left(\frac{r-R}{r+R}\right)\right) + \mathcal{O}(G^3), \quad (2.15)$$

$$g(r) = \left(1 - \frac{2GM}{r} + \frac{r^2}{a^2}\right)^{-1} + \tilde{\beta} \frac{2G^2 M}{r(r^2 - R^2)} + \mathcal{O}(G^3), \quad (2.16)$$

with

$$\tilde{\alpha} = 96\pi(\alpha + \beta + 3\gamma), \quad (2.17)$$

$$\tilde{\beta} = 192\pi(\gamma - \alpha). \quad (2.18)$$

and M is the mass of black hole. The correction of the metric include two parts : one is from local part i.e. eq., (2.2) & other is from non-local part i.e.(2.3). The correction from local part is trivially 0. Inside the star this is not the case. However, these corrections turn out to be $\mathcal{O}(G^3)$, and thus sub-leading. Therefore the local part in the equations of motion (2.8) does not contribute. The two terms $\tilde{\alpha}$ and $\tilde{\beta}$ are the contributions of non-local part of i.e., eq.(2.3).

3 Quantum correction to Hawking temperature for Schwarzschild AdS black hole

In this section, we study the semiclassical tunnelling of particles through the horizon of Schwarzschild black holes, in line with the analysis of [17]. The rate of emission Γ will have exponential part given by

$$\Gamma \sim \exp(-2\text{Im}S), \quad (3.1)$$

where S is the tunnelling action of particles and $\text{Im}S$ is the imaginary part of the tunnelling action. According to Plank radiation law, the emitted rate Γ of particles with frequency ω can be written as

$$\Gamma \sim \exp\left(-\frac{\omega}{T_{BH}}\right). \quad (3.2)$$

Hence the temperature at which black hole radiates given by

$$T_{BH} = \frac{\omega}{2\text{Im}S}. \quad (3.3)$$

In this section, we will study the process in Painlevé-Gullstrand coordinates. Firstly, we need to rewrite the corrected metric in Painlevé-Gullstrand coordinates to order G^2 . We define

$$t_r = t - a(r), \quad (3.4)$$

where $a(r)$ is function of r . Then

$$dt^2 = dt_r^2 + a'^2(r)dr^2 + 2a'(r)dt_r dr, \quad (3.5)$$

where $a'(r) = \frac{da(r)}{dr}$. Using equation (3.5) & (2.14) we have

$$ds^2 = -f(r)dt_r^2 + \{g(r) - f(r)a'^2(r)\}dr^2 - 2a'(r)f(r)dt_r dr + r^2 d\Omega^2. \quad (3.6)$$

We require

$$\{g(r) - f(r)a'^2(r)\} = 1, \quad (3.7)$$

we have $a'^2 = \frac{g(r)-1}{f(r)}$. We take $a' = -\sqrt{\frac{g(r)-1}{f(r)}}$. Then

$$-2a'f(r) = 2\sqrt{f(r)\{g(r) - 1\}}, \quad (3.8)$$

For classical Schwarzschild AdS metric, $f(r) = (1 - \frac{2GM}{r} + \frac{r^2}{a^2})$ and $g(r) = (1 - \frac{2GM}{r} + \frac{r^2}{a^2})^{-1}$, we have

$$ds^2 = -(1 - \frac{2GM}{r} + \frac{r^2}{a^2})dt_r^2 + dr^2 + 2\sqrt{(1 + \frac{r^2}{a^2})\frac{2GM}{r} - (1 - \frac{2GM}{r} + \frac{r^2}{a^2})\frac{r^2}{a^2}} dt_r dt + r^2 d\Omega^2. \quad (3.9)$$

For the quantum corrected Schwarzschild AdS metric, $f(r)$ and $g(r)$ have been given in equations (2.15) and (2.16), then we have

$$f(r)(g(r) - 1) = (1 + \frac{r^2}{a^2})\frac{2GM}{r} - (1 - \frac{2GM}{r} + \frac{r^2}{a^2})\frac{r^2}{a^2} + (1 + \frac{r^2}{a^2})\beta - \frac{r^2}{a^2}\alpha, \quad (3.10)$$

$$-2a'f(r) = 2\sqrt{f(r)\{g(r) - 1\}} = 2\sqrt{(1 + \frac{r^2}{a^2})\frac{2GM}{r} - (1 - \frac{2GM}{r} + \frac{r^2}{a^2})\frac{r^2}{a^2} + (1 + \frac{r^2}{a^2})\beta - \frac{r^2}{a^2}\alpha}, \quad (3.11)$$

Therefore, to order $\mathcal{O}(G^2)$ the quantum corrected Schwarzschild AdS metric can be written as,

$$ds^2 = -f(r)dt_r^2 + dr^2 + 2\sqrt{(1 + \frac{r^2}{a^2})\frac{2GM}{r} - (1 - \frac{2GM}{r} + \frac{r^2}{a^2})\frac{r^2}{a^2} + (1 + \frac{r^2}{a^2})\beta - \frac{r^2}{a^2}\alpha} dt_r dr + r^2 d\Omega^2, \quad (3.12)$$

Dropping the subscript r for convenience, i.e.

$$ds^2 = -f(r)dt^2 + dr^2 + 2\sqrt{(1 + \frac{r^2}{a^2})\frac{2GM}{r} - (1 - \frac{2GM}{r} + \frac{r^2}{a^2})\frac{r^2}{a^2} + (1 + \frac{r^2}{a^2})\beta - \frac{r^2}{a^2}\alpha} dt dr + r^2 d\Omega^2. \quad (3.13)$$

The action for a particle moving freely in a curved background can be written as

$$S = \int p_\mu dx^\mu, \quad (3.14)$$

where,

$$p_\mu = mg_{\mu\nu} \frac{dx^\nu}{d\sigma}. \quad (3.15)$$

where σ is an affine parameter along the worldline of the particle, chosen so that p_μ coincides with the physical 4-momentum of the particle. For a massive particle, this requires that $d\sigma = \frac{d\tau}{m}$, with τ is the proper time.

3.1 Quantum correction to Hawking temperature

3.1.1 Massless particles

The radial dynamics of massless particles in Schwarzschild spacetime are determined by the equations

$$\dot{r}^2 + 2\sqrt{(1 + \frac{r^2}{a^2})\frac{2GM}{r} + (1 + \frac{r^2}{a^2})\beta - \frac{r^2}{a^2}f(r)} \dot{t} \dot{x} - f(r)\dot{t}^2 = 0. \quad (3.16)$$

From equation (3.15) taking $\mu = t$

$$p_t = g_{tt}\dot{t} + g_{tr}\dot{r}, \quad (3.17)$$

$p_t = -\omega$ and ω has the interpretation of the energy of the particle as measured at infinity. Equation (3.16) can be factorised to the equation:

$$\dot{r} = \left[\pm \left(1 - \sqrt{\frac{2GM}{r}} \right) \left(1 + \frac{r^2}{2a^2} \right) - \sqrt{\frac{2GM}{r}} \frac{\tilde{\beta}G}{2(r^2 - R^2)} \left(1 + \frac{r^2}{a^2} \right) \right] \dot{t}, \quad (3.18)$$

with the positive sign for outgoing and negative sign for ingoing geodesics. We only consider the case of outgoing particles. From equations (3.17) and (3.18) we have

$$\dot{t} = \frac{\omega}{1 - \sqrt{\frac{2GM}{r}} \left(1 + \frac{r^2}{a^2} \right) - \sqrt{\frac{2GM}{r}} \frac{\tilde{\beta}G}{2(r^2 - R^2)} \left(1 + \frac{r^2}{a^2} \right)}, \quad (3.19)$$

From equation (3.14), (3.15) & (3.19) we have

$$ImS = Im \int p_r dr = Im \int \dot{t} dr. \quad (3.20)$$

Define

$$1 - \sqrt{\frac{2GM}{r}} = 1 - \sqrt{\frac{2GM}{r}} \left(1 + \frac{r^2}{a^2} \right) - \sqrt{\frac{2GM}{r}} \frac{\tilde{\beta}G}{2(r^2 - R^2)} \left(1 + \frac{r^2}{a^2} \right) \quad (3.21)$$

,

Then equation (3.20) becomes

$$ImS = Im \int \dot{t} dr = \int \frac{\omega dr}{1 - \sqrt{\frac{2GM}{r}}}. \quad (3.22)$$

The integrand has a pole at $r = 2\tilde{M}$, the horizon. Choosing the prescription to integrate clockwise around this pole (into the upper-half complex-r plane) [17], [11] we find

$$ImS = 4\pi G\omega\tilde{M}. \quad (3.23)$$

Then the temperature at which the black hole radiates, equation (3.3) becomes

$$T_{BH} = \frac{\omega}{2ImS} = \frac{1}{8\pi G\tilde{M}}. \quad (3.24)$$

From equation (3.21)

$$\tilde{M} = M \left[1 + \frac{\tilde{\beta}G}{2(r^2 - R^2)} \right]^2 \left(1 + \frac{r^2}{a^2} \right)^2. \quad (3.25)$$

Therefor

$$T_{BH} = \frac{1}{8\pi GM} \left(1 - \frac{\tilde{\beta}G}{(r^2 - R^2)} - \frac{\tilde{\beta}^2 G^2}{4(r^2 - R^2)^2} \right) \left(1 - 2\frac{r^2}{a^2} - \frac{r^4}{a^4} \right). \quad (3.26)$$

3.1.2 Massive particles

For massive point-particles, the dynamics are governed by

$$\dot{r}^2 + 2\sqrt{\left(1 + \frac{r^2}{a^2}\right)\frac{2GM}{r} + \left(1 + \frac{r^2}{a^2}\right)\beta - \frac{r^2}{a^2}f(r)} \dot{t} + (1 - f(r)\dot{t}^2) = 0. \quad (3.27)$$

From equation (3.15) taking $\mu = t$

$$p_t = g_{tt}\dot{t} + g_{tr}\dot{r}. \quad (3.28)$$

$p_t = -\omega$ and ω has the interpretation of the energy of the particle as measured at infinity. Equation (3.27) can be factorised to the equation

$$\dot{r} = \left[\pm \left(1 - \sqrt{\frac{2GM}{r}}\right)\left(1 + \frac{r^2}{2a^2}\right) - \sqrt{\frac{2GM}{r}} \frac{\tilde{\beta}G}{2(r^2 - R^2)} \left(1 + \frac{r^2}{a^2}\right) \right] \dot{t}, \quad (3.29)$$

with the positive sign for outgoing and negative sign for ingoing geodesics. We only consider the case of outgoing particles. From equations (3.28) & (3.29) we have

$$\dot{t} = \frac{\omega}{1 - \sqrt{\frac{2GM}{r}}\left(1 + \frac{r^2}{a^2}\right) - \sqrt{\frac{2GM}{r}} \frac{\tilde{\beta}G}{2(r^2 - R^2)} \left(1 + \frac{r^2}{a^2}\right)}, \quad (3.30)$$

From equation (3.14), (3.15) & (3.30), we have

$$ImS = Im \int p_r dr = Im \int \dot{t} dr. \quad (3.31)$$

Define

$$1 - \sqrt{\frac{2G\tilde{M}}{r}} = 1 - \sqrt{\frac{2GM}{r}}\left(1 + \frac{r^2}{a^2}\right) - \sqrt{\frac{2GM}{r}} \frac{\tilde{\beta}G}{2(r^2 - R^2)} \left(1 + \frac{r^2}{a^2}\right), \quad (3.32)$$

Then equation (3.31) becomes

$$ImS = Im \int \dot{t} dr = \int \frac{\omega dr}{1 - \sqrt{\frac{2G\tilde{M}}{r}}}. \quad (3.33)$$

The integrand has a pole at $r = 2\tilde{M}$, the horizon. Choosing the prescription to integrate clockwise around this pole (into the upper-half complex- r plane) [17], [11] we find

$$ImS = 4\pi G\omega\tilde{M}. \quad (3.34)$$

Then the temperature at which the black hole radiates, equation (3.3) becomes

$$T_{BH} = \frac{\omega}{2ImS} = \frac{1}{8\pi G\tilde{M}}. \quad (3.35)$$

From equation (3.32)

$$\tilde{M} = M \left[1 + \frac{\tilde{\beta}G}{2(r^2 - R^2)} \right]^2 \left(1 + \frac{r^2}{a^2} \right)^2, \quad (3.36)$$

Therefore

$$T_{BH} = \frac{1}{8\pi GM} \left(1 - \frac{\tilde{\beta}G}{(r^2 - R^2)} - \frac{\tilde{\beta}^2 G^2}{4(r^2 - R^2)^2} \right) \left(1 - 2\frac{r^2}{a^2} - \frac{r^4}{a^4} \right). \quad (3.37)$$

Same as equation (3.26).

4 Quantum correction to thermodynamics of Schwarzschild AdS black hole

In this section we study the quantum corrected black hole entropy and various thermodynamics parameters. We find entropy of Schwarzschild AdS black hole upto order $\mathcal{O}(G^2)$ and plot the classical & quantum entropy. Next we study specific heat and its comes out positive. Finally we compute Helmholtz free energy and pressure of quantum corrected Schwarzschild AdS black hole and we plot it's classical and quantum version.

4.1 Quantum correction to entropy

The entropy of the system is defined as

$$S = \frac{\pi r_h^2}{G} = \frac{\pi(2G\tilde{M})^2}{G}, \quad (4.1)$$

$$S = 4\pi GM^2 \left[1 + \frac{\tilde{\beta}G}{(r^2 - R^2)} + \frac{\tilde{\beta}^2 G^2}{4(r^2 - R^2)^2} \right]^2 \left(1 + \frac{2r^2}{a^2} + \frac{r^4}{a^4} \right)^2, \quad (4.2)$$

Now substituting M from equation (3.37) and taking term upto order $\mathcal{O}(G^2)$ we have

$$S = \frac{1}{16\pi T_{BH}^2} \left[1 + \frac{2\tilde{\beta}G}{(r^2 - R^2)} + \frac{3\tilde{\beta}^2 G^2}{2(r^2 - R^2)^2} \right] \left[1 - \frac{2\tilde{\beta}G}{(r^2 - R^2)} + \frac{\tilde{\beta}^2 G^2}{2(r^2 - R^2)^2} \right] \left(1 + \frac{2r^2}{a^2} + \frac{r^4}{a^4} \right)^2 \left(1 - \frac{2r^2}{a^2} - \frac{r^4}{a^4} \right)^2, \quad (4.3)$$

$$S = \frac{1}{16\pi T_{BH}^2} \left[1 - \frac{2\tilde{\beta}^2 G^2}{(r^2 - R^2)^2} \right] \left(1 + \frac{2r^2}{a^2} + \frac{r^4}{a^4} \right)^2 \left(1 - \frac{2r^2}{a^2} - \frac{r^4}{a^4} \right)^2. \quad (4.4)$$




Figure 1

Figure: (a) Entropy of Classical Schwarzschild AdS black hole with $G = 1$, $r = 3R$, $M = 2$ and $a = 1.5$. **Figure: (b)** Entropy of quantum corrected Schwarzschild AdS black hole with $G = 1$, $r = 3R$, $M = 2$ and $a = 1.5$. Numerical values of α , β and γ taken from [10]. All fields (Scalar, Fermionic, Vector and Gravitational) give rise to approximately the same behaviour.

The specific heat of black hole is

$$C = \frac{M}{T_{BH}} \left(\frac{dS}{dT_{BH}} \right), \quad (4.5)$$

$$C = \frac{G \left[1 + \frac{2\tilde{\beta}G}{(r^2-R^2)} + \frac{3\tilde{\beta}^2 G^2}{2(r^2-R^2)^2} \right] \left[1 - \frac{2\tilde{\beta}G}{(r^2-R^2)} + \frac{\tilde{\beta}^2 G^2}{2(r^2-R^2)^2} \right] \left(1 + \frac{2r^2}{a^2} + \frac{r^4}{a^4} \right)^2 \left(\frac{2r^2}{a^2} + \frac{r^4}{a^4} - 1 \right)}{T_{BH} \left[1 - \frac{\tilde{\beta}G}{(r^2-R^2)} - \frac{\tilde{\beta}^2 G^2}{4(r^2-R^2)^2} \right]}. \quad (4.6)$$

Equations (3.36), (3.37), (4.2) and (4.6) depends on R . Since the quantum modification are tiny as the term $\frac{\tilde{\beta}G}{(r^2-R^2)}$ is very small. Classical entropy ($S = \frac{A}{4G}$) of black hole get modified, as there is a shift in the position of horizon.

4.2 Quantum correction to free energy and pressure

The Helmholtz free energy is

$$F = \frac{GM \left[1 + \frac{2\tilde{\beta}G}{(r^2-R^2)} + \frac{3\tilde{\beta}^2 G^2}{2(r^2-R^2)^2} \right] \left[1 - \frac{2\tilde{\beta}G}{(r^2-R^2)} + \frac{\tilde{\beta}^2 G^2}{2(r^2-R^2)^2} \right] \left(1 + \frac{2r^2}{a^2} + \frac{r^4}{a^4} \right)^2 \left(\frac{2r^2}{a^2} + \frac{r^4}{a^4} - 1 \right)}{2 \left[1 - \frac{\tilde{\beta}G}{(r^2-R^2)} - \frac{\tilde{\beta}^2 G^2}{4(r^2-R^2)^2} \right]} + M. \quad (4.7)$$

Quantum effects also produce modification on the pressure. For a schwarzschild AdS black hole pressure is

$$P = -\frac{T_{BH} \frac{dS}{dM} - 1}{\frac{dV}{dM}} \quad (4.8)$$

$$P = \frac{\left[1 - \frac{\tilde{\beta}G}{(r^2-R^2)} - \frac{\tilde{\beta}^2 G^2}{4(r^2-R^2)^2} \right] \left[1 + \frac{\tilde{\beta}G}{(r^2-R^2)} + \frac{\tilde{\beta}^2 G^2}{4(r^2-R^2)^2} \right]^2 \left(\frac{2r^2}{a^2} + \frac{r^4}{a^4} - 1 \right) \left(1 + 2\frac{r^2}{a^2} + \frac{r^4}{a^4} \right)^2 - 1}{32\pi G^3 M^2 \left[1 + \frac{2\tilde{\beta}G}{2(r^2-R^2)^2} \right]^6 \left(1 + 2\frac{r^2}{a^2} + \frac{r^4}{a^4} \right)^4}. \quad (4.9)$$

Next, we will discuss the variation of Hawking temperature for classical and quantum corrected schwarzschild AdS black holes.




Figure 2

Figure: (a) Hawking temperature of Classical Schwarzschild AdS black hole with $G = 1$, $r = 3R$, $M = 2$ and $a = 1.5$. **Figure:** (b) Hawking temperature of quantum corrected Schwarzschild AdS black hole with $G = 1$, $r = 3R$, $M = 2$ and $a = 1.5$. Dash dot red: Quantum corrected Schwarzschild AdS with scalar field

($\alpha = 0.0010994$, $\beta = -0.00001759$, $\gamma = 0.00001759$). Dash dot black: Quantum corrected Schwarzschild AdS with fermionic field ($\alpha = -0.000043976$, $\beta = 0.000070361$, $\gamma = 0.000061566$). Solid green: Quantum corrected Schwarzschild AdS with vector field ($\alpha = -0.000433976$, $\beta = 0.0015480$, $\gamma = -0.00022867$). Solid gold: Quantum corrected Schwarzschild AdS with graviton field ($\alpha = 0.0037819$, $\beta = -0.012700$, $\gamma = 0.0037292$). Numerical values of α , β and γ taken from [10].

5 Conclusions

In this paper, we consider four-dimensional Anti de Sitter Schwarzschild black holes. In ref. [10] the authors used EFT methods to obtain the metric inside and outside a static and spherically symmetric star, taking into account quantum effects. Authors also proposed that the exterior metric may provide a good model of quantum black holes. In section 2, we used EFT method to obtain quantum correction to the four-dimensional Anti de Sitter Schwarzschild black holes. In section 3, we studied quantum correction of Hawking temperature for massless and massive particle. First we studied the radial dynamic equations of massless and massive particle in quantum corrected Schwarzschild AdS spacetime. Then, following the spirit of ref.[17, 11], we obtained the modified Hawking temperature. By computing the modified Hawking entropy, we found that the famous area law of black holes $S = \frac{A}{4G}$ need to be corrected. In section 4, we compute the quantum correction to entropy, specific heat, free energy and pressure.

References

- [1] Jacob D Bekenstein. “Black holes and the second law”. In: *JACOB BEKENSTEIN: The Conservative Revolutionary*. World Scientific, 2020, pp. 303–306.
- [2] Jacob D Bekenstein. “Black holes and entropy”. In: *JACOB BEKENSTEIN: The Conservative Revolutionary*. World Scientific, 2020, pp. 307–320.
- [3] Stephen W Hawking. “Black hole explosions?”. In: *Nature* 248.5443 (1974), pp. 30–31.
- [4] Stephen W Hawking. “Particle creation by black holes”. In: *Commun. Math. Phys.* 43.5443 (1975), pp. 199–220.
- [5] Don N. Page. “Particle emission rates from a black hole: Massless particles from an uncharged, nonrotating hole”. In: *Phys. Rev. D* 13 (2 Jan. 1976), pp. 198–206.
- [6] Don N. Page. “Particle emission rates from a black hole. II. Massless particles from a rotating hole”. In: *Phys. Rev. D* 14 (12 Dec. 1976), pp. 3260–3273.
- [7] Maulik K. Parikh and Frank Wilczek. “Hawking Radiation As Tunneling”. In: *Phys. Rev. Lett.* 85 (24 Dec. 2000), pp. 5042–5045.
- [8] Roberto Emparan, Gary T. Horowitz, and Robert C. Myers. “Black Holes Radiate Mainly on the Brane”. In: *Phys. Rev. Lett.* 85 (3 July 2000), pp. 499–502.
- [9] Samuli Hemming and Esko Keski-Vakkuri. “Hawking radiation from AdS black holes”. In: *Phys. Rev. D* 64 (4 July 2001), p. 044006.
- [10] Xavier Calmet, Roberto Casadio, and Folkert Kuipers. “Quantum gravitational corrections to a star metric and the black hole limit”. In: *Physical Review D* 100.8 (2019), p. 086010.
- [11] Yang Liu. “The effect of quantum correction on Hawking radiation for Schwarzschild black holes”. In: *arXiv preprint arXiv:2201.00599* (2022).
- [12] John F Donoghue and Basem Kamal El-Menoufi. “Covariant non-local action for massless QED and the curvature expansion”. In: *Journal of High Energy Physics* 2015.10 (2015), pp. 1–24.
- [13] Behnam Pourhassan and Ruben Campos Delgado. “Quantum Gravitational Corrections to the Geometry of Charged AdS Black Holes”. In: *arXiv preprint arXiv:2205.00238* (2022).
- [14] Xavier Calmet. “Vanishing of quantum gravitational corrections to vacuum solutions of general relativity at second order in curvature”. In: *Physics Letters B* 787 (2018), pp. 36–38.
- [15] Miguel Socolovsky. “Schwarzschild black hole in anti-de Sitter space”. In: *arXiv preprint arXiv:1711.02744* (2017).
- [16] Ruben Campos Delgado. “Quantum gravitational corrections to the entropy of a Reissner–Nordström black hole”. In: *The European Physical Journal C* 82.3 (2022), pp. 1–8.
- [17] George Johnson and John March-Russell. “Hawking radiation of extended objects”. In: *Journal of High Energy Physics* 2020.4 (2020), pp. 1–16.