

Correlation energy of the paramagnetic electron gas at the thermodynamic limit

Sam Azadi,^{1,*} N. D. Drummond,² and S. M. Vinko¹

¹*Department of Physics, Clarendon Laboratory, University of Oxford,
Parks Road, Oxford OX1 3PU, United Kingdom*

²*Department of Physics, Lancaster University, Lancaster LA1 4YB, United Kingdom*
(Dated: September 22, 2022)

The variational and diffusion quantum Monte Carlo methods are used to calculate the correlation energy of the paramagnetic three-dimensional homogeneous electron gas at intermediate to high density. Ground state energies in finite cells are determined using Slater-Jastrow-backflow trial wave functions, and finite-size errors are removed using twist-averaged boundary conditions and extrapolation of the energy per particle to the thermodynamic limit of infinite system size. Our correlation energies in the thermodynamic limit are lower (i.e., more negative, and therefore more accurate according to the variational principle) than previous results, and can be used for the parameterization of density functionals to be applied to high-density systems.

CONTENTS

I. Introduction	1
II. Trial wave function	2
III. Finite-size effects	3
IV. Correlation energies	3
V. Conclusion	4
Acknowledgments	5
References	5

I. INTRODUCTION

The pairwise Coulomb repulsion between electrons results in many-body correlations in electronic systems such as the homogeneous electron gas (HEG) [1, 2]. The so-called correlation energy is a negative correction to the mean-field Hartree-Fock energy. Although the correlation energy is usually only a small percentage of the total energy of an electronic system, it is crucial for an accurate description of chemical and electronic properties [3–5]. Unfortunately it is also the most complicated part of the energy to calculate accurately.

The three-dimensional (3D) HEG plays a crucial role in our understanding of the nature of electronic correlation in real materials [6–8]. Moreover, the HEG is one of the most important models for our understanding of bulk systems under extreme conditions, such as warm dense matter, which is an exotic, highly compressed state of matter that exists between solid and plasma phases at high temperatures [9–11]. The correlation energy of the

3D HEG as a function of density [12–14] is a fundamental element in the description of the electronic properties of real systems by density functional theory (DFT) [15, 16]. However, calculating the correlation energy accurately requires many-body wave function-based methods [17] such as quantum Monte Carlo (QMC) techniques [18–25]. The variational (VMC) and diffusion quantum Monte Carlo (DMC) methods [18, 26] are stochastic approaches for obtaining expectation values of quantum operators. These techniques are especially efficient for calculating the ground state energies of interacting fermions. The main object is an approximate trial wave function, whose accuracy governs the final energy and intrinsic statistical fluctuations in the simulations.

The DMC simulations of Ceperley and Alder [18] presented important data connecting the high- and low-density regimes of the correlation energy of the 3D HEG. Their data have been used in parameterizations of the correlation energy over a wide density range and are frequently used in DFT calculations. Well-known parameterizations that make use of Ceperley and Alder’s results were provided by Perdew and Zunger (PZ81) [27], Vosko, Wilk, and Nusair (VWN80) [28], and Perdew and Wang (PW92) [29], among others. The PW92 functional includes five parameters, two determined from analytic high-density constraints and three by fitting to the QMC data. A density parameter interpolation (DPI) [12], which was constructed by imposing four high-density and three low-density constraints on a seven-parameter functional form, provided a check based purely on the satisfaction of the exact constraints. Spink *et al.* [23] performed QMC calculations for spin-unpolarized and spin-polarized 3D HEGs over the high- and intermediate-density ranges, which can be regarded as the most accurate QMC data reported so far. In the present work, we provide new QMC data for the correlation energy of the paramagnetic (i.e., spin unpolarized) 3D HEG, which are lower than previously reported results. QMC energies in finite simulation cells obey the variational principle and it is reasonable to assume that the QMC energy per par-

* sam.azadi@physics.ox.ac.uk

ticle extrapolated to infinite system size is also an upper bound on the true energy per particle. Hence the fact that our energies are lower than previous works strongly suggests that our results are more accurate.

We have used the VMC and DMC methods to obtain 3D HEG correlation energies at different densities. In the VMC method, parameters in a trial wave function are optimized according to the variational principle, with energy expectation values calculated by Monte Carlo integration in the $3N$ -dimensional space of electron position vectors. In the DMC method, the imaginary-time Schrödinger equation is used to evolve a statistical ensemble of electronic configurations towards the ground state. Fermionic antisymmetry is maintained by the fixed-phase approximation, in which the complex phase of the wave function is constrained to equal that of an approximate trial wave function optimized within VMC.

The simplest fermionic wave function is a Slater determinant, which describes exchange effects but not correlation. Multideterminant wave functions and pairing (geminal) wave functions [30] can also be used. The most efficient method of going beyond the Slater wave function is to multiply it by a Jastrow factor $\exp(J)$, resulting in a Slater-Jastrow wave function [20, 21]. The Jastrow factor usually depends explicitly on the distances between particles, introducing correlation into the wave function. The Jastrow factor is positive everywhere and symmetric with respect to the exchange of indistinguishable particles, so it does not change the nodal surface defined by the rest of the wave function. By evaluating the orbitals in the Slater determinant at quasiparticle coordinates \mathbf{X} , which are functions of all the electron positions, we introduce a backflow transformation [31, 32], and the resulting wave function is referred to as a Slater-Jastrow-backflow (SJB) wave function.

II. TRIAL WAVE FUNCTION

We used a SJB trial spatial wave function $\Psi(\mathbf{R}) = e^{J(\mathbf{R})}S(\mathbf{X}(\mathbf{R}))$ for all the systems we have studied, where $\mathbf{R} = (\mathbf{r}_1, \dots, \mathbf{r}_N)$ is the $3N$ -dimensional vector of electron coordinates. The antisymmetric Slater part S is a product of determinants of single-particle orbitals for spin-up and spin-down electrons. The single-particle orbitals in S are of the free-electron form $\psi_{\mathbf{k}}(\mathbf{r}) = \exp(i\mathbf{k} \cdot \mathbf{r})$, where wavevector \mathbf{k} is a reciprocal lattice vector of the simulation cell offset by twist vector \mathbf{k}_s , where \mathbf{k}_s lies in the supercell Brillouin zone. The Jastrow exponent, which is symmetric under electron exchange, takes the form

$$\begin{aligned} J &= U + P + H \\ &= \sum_{i < j} u(r_{ij}) + \sum_{i < j} p(\mathbf{r}_{ij}) + \sum_{i < j < k} h(r_{jk}, r_{ik}, r_{ij}), \end{aligned} \quad (1)$$

where

$$u(r) = \sum_{l=0}^{N_u} \alpha_l r^l (r - L_u)^C \Theta(L_u - r), \quad (2)$$

where r is the minimum-image distance between two electrons, the cutoff length L_u is less than or equal to the radius of the largest sphere that can be inscribed in the Wigner-Seitz cell of the simulation cell, $C = 3$ specifies how smooth the function is at the cutoff length, Θ is the Heaviside step function, and $\{\alpha_l\}$ are optimizable parameters, which differ for parallel- and antiparallel-spin electrons. To satisfy the Kato cusp conditions [33, 34], we fix $\alpha_1 = \Gamma/(-L_u)^C + \alpha_0 C/L_u$, where $\Gamma = 1/2$ for opposite-spin electrons and $\Gamma = 1/4$ for same-spin electrons. We chose $N_u = 8$. The p term has the symmetry of the simulation-cell Bravais lattice and allows a description of correlation in the ‘‘corners’’ of the simulation cell. Its form is

$$p(\mathbf{r}) = \sum_A a_A \sum_{\mathbf{G} \in A^+} \cos(\mathbf{G} \cdot \mathbf{r}), \quad (3)$$

where A represents a star of symmetry-equivalent, nonzero, simulation-cell reciprocal-lattice vectors \mathbf{G} , and A^+ is a subset of A that consists of one out of each $\pm\mathbf{G}$ pair. The $\{a_A\}$ are optimizable parameters. We used 46 stars of \mathbf{G} vectors in p . The Jastrow also includes symmetric three-electron terms [35, 36]

$$\begin{aligned} h(r, r', r'') &= \sum_{l=0}^{N_h} \sum_{m=0}^{N_h} \sum_{n=0}^{N_h} c_{lmn} r^l (r')^m (r'')^n \\ &\quad \times (r - L_h)^C (r' - L_h)^C (r'' - L_h)^C \\ &\quad \times \Theta(L_h - r) \Theta(L_h - r') \Theta(L_h - r'') \end{aligned} \quad (4)$$

where L_h is a cutoff length and c_{lmn} are linear parameters. Constraints were placed on the linear parameters to ensure that h is cusplless. We chose $N_h = 4$. Different h terms, meaning different $\{c_{lmn}\}$, may be used for electron triplets involving different combinations of spins. However, in this work, the parameters in the three-electron Jastrow factors were constrained to be independent of spin.

Including a backflow transformation in the trial wave function, the Slater part of the wave function S is evaluated at transformed ‘‘quasiparticle’’ coordinates $\mathbf{X}(\mathbf{R}) = \mathbf{R} + \boldsymbol{\xi}(\mathbf{R})$, where

$$\boldsymbol{\xi}_i(\mathbf{R}) = \sum_{j \neq i} \eta(r_{ij}) \mathbf{r}_{ij} + \sum_{j \neq i} \boldsymbol{\pi}(\mathbf{r}_{ij}) \quad (5)$$

is the backflow displacement of electron i . η is a cusplless, smoothly truncated, isotropic polynomial function of minimum-image electron-electron distance r_{ij} . The polynomial coefficients are optimizable parameters, and are different for parallel- and antiparallel-spin electrons [32]. The form of $\eta(r)$ is mathematically equivalent to

that of the Jastrow $u(r)$ term [Eq. (2), with $\Gamma = 0$ for same-spin electrons and optimizable for opposite-spin electrons]. Typically we used $N_\eta = 8$ in the polynomial expansions. The π term has the form of the gradient of a Jastrow p term [Eq. (3)]:

$$\pi(\mathbf{r}) = - \sum_A c_A \sum_{\mathbf{G} \in A^+} \sin(\mathbf{G} \cdot \mathbf{r}) \mathbf{G}, \quad (6)$$

where the c_A are optimizable parameters. As the gradient of a scalar field, the π term is irrotational. We used 44 stars of \mathbf{G} vectors in π . The backflow parameters were allowed to depend on the spins of the electron pairs.

The wave functions were optimized by variance minimization [37, 38] followed by energy minimization [39]. The CASINO package was used for all our QMC calculations [40].

III. FINITE-SIZE EFFECTS

Monte Carlo-sampled canonical ensemble twist-averaged (TA) boundary conditions were used to reduce quasirandom single-particle finite-size errors in total energies due to momentum quantization effects [41–45]. The Hartree-Fock kinetic and exchange energies were used as control variates to improve the precision of the twist-averaged energy. Systematic finite-size errors due to the use of the Ewald interaction rather than $1/r$ to evaluate the interaction between each electron and its exchange-correlation hole and the neglect of long-range two-body correlations were removed by fitting $E(N) = E(\infty) + b/N$ to the TA DMC energy per particle at different system sizes [46]. Unlike the previous work of Spink *et al.* [23], we do not rely on analytic finite-size correction formulas [43, 46], but instead use the analytic results to provide the exponents used in finite-size extrapolation formulas. All our calculations were performed using face-centered cubic simulation cells, maximizing the distance between each particle and its closest periodic image.

At very high density $r_s \ll 1$, systematic finite-size effects are more challenging. In this regime, the QMC energy is close to the Hartree-Fock energy, and hence the QMC energy per particle initially shows the Hartree-Fock $O(N^{-2/3})$ scaling with system size [43], before eventually crossing over to the asymptotic $O(N^{-1})$ scaling when the finite-size error becomes small compared with the correlation energy.

IV. CORRELATION ENERGIES

We studied the paramagnetic 3D HEG at density parameters $r_s = 0.5, 0.75, 1, 2, 3, 4, 5, 7, 10,$ and 20 . For each density, QMC calculations were performed for simulation cells with $N = 130, 226,$ and 338 electrons. Our DMC energies were extrapolated linearly to zero time step, with the target walker population being varied in inverse proportion to the time step. The energies

and variances calculated using SJB wave functions for different system sizes are reported in the Supplemental Material [47]. The correlation energy is defined as the difference between the Hartree-Fock energy per electron [which is $E_{\text{HF}} = 3(9\pi/4)^{2/3}/(10r_s^2) - 3(9\pi/4)^{1/3}/(4\pi r_s)$ for the paramagnetic HEG] and the exact ground-state energy per electron, where the latter is approximated by our SJB-DMC results extrapolated to the limit of infinite system size.

Table I summarizes the contribution of each term of the trial wave function to the correlation energy per particle in a simulation cell containing $N = 54$ electrons. We considered two systems with $r_s = 0.5$ and $r_s = 20$. Figure 1 shows the improvements in the VMC and DMC correlation energies resulting from the inclusion of different terms in the Jastrow and backflow functions.

According to Spink *et al.* [23], the TA VMC energy of the spin-unpolarized 3D HEG at $r_s = 0.5$ at a system size of $N = 118$ is $3.41378(2)$ Ha/elec. Our TA VMC simulation for the same system size yields the energy as $3.412460(4)$ Ha/elec., which is ~ 36 meV/elec. lower, because of the inclusion of the π term in our work. To check the performance of our optimized wave function, we compared our VMC and DMC energies for $N = 54$ system size with full configuration interaction quantum Monte Carlo (FCIQMC) [24]. The FCIQMC energy of the spin-unpolarized 3D HEG with $N = 54$ electrons at $r_s = 0.5$ is $3.2202(2)$ Ha/elec., and our TA VMC and DMC energies are $3.40671(1)$ and $3.40658(1)$ Ha/elec., respectively. The FCIQMC energy for the same system size at $r_s = 1.0$ is $0.5300(3)$ Ha/elec., and our TA VMC and DMC energies are $0.577258(3)$ and $0.577148(5)$ Ha/elec., respectively.

TABLE I. TA VMC and DMC correlation energies for $N = 54$ system size obtained using different terms in the Jastrow exponent and using both SJB and SJ wave functions. Where Jastrow terms are not specified, U , P , and H terms were used; where the backflow terms are not specified, η and π terms were used.

Wave fn	Correlation energy (eV/elec.)			
	$r_s = 0.5$		$r_s = 20$	
	VMC	DMC	VMC	DMC
SJ(U)	-2.5112(3)	-2.5684(4)	-0.311717(2)	-0.317465(4)
SJ($U + H$)	-2.5161(3)	-2.5680(5)	-0.313798(3)	-0.317469(5)
SJ	-2.5381(3)	-2.5685(4)	-0.314464(3)	-0.317492(5)
SJB(η)	-2.6056(5)	-2.6166(5)	-0.319082(3)	-0.320579(5)
SJB	-2.6331(3)	-2.6366(4)	-0.319346(2)	-0.320756(4)

Our VMC and DMC energies extrapolated to the limit of infinite system size are listed in Table II. To obtain the best linear fit and reduce the noise in the energies as a function of the number of particles N , a larger number of twists was used for higher densities and smaller system sizes. The smallest and largest number of twists were 120 and 10^4 , respectively. Comparing our infinite-system VMC and DMC results with the DMC energies of Spink *et al.* [23] demonstrates not only the improvement of the trial wave function due to the inclusion of




FIG. 1. Twist averaged VMC and DMC correlation energies at system size $N = 54$ as a function of variance, relative to the correlation energy with a Slater-Jastrow (SJ) wave function in which the Jastrow factor only contains the isotropic two-body term U . Results are shown at density parameters $r_s = 0.5$ (top panel) and 20 (bottom panel).

long-range π backflow terms, but also the importance of removing finite-size effects by extrapolation rather than relying on analytic correction formulas.

Table III compares our VMC and DMC results for the correlation energy with the PZ81 [27], VWN80 [28], PW92 [29], and DPI [12] parameterizations, as well as the DMC data of Spink *et al.* [23]. Our correlation energies are the lowest. Even at the high density $r_s = 0.5$ our DMC correlation energy is lower than the DPI parameterization by -16 meV/elec.

TABLE II. TA VMC and DMC energies of the 3D HEG extrapolated to the thermodynamic limit from different system sizes ($N = 130, 226,$ and 338), compared with the DMC results of Spink *et al.* [23], and Ceperley and Alder [18]. Our DMC energies have been extrapolated to zero time step.

r_s	Total energy (Ha/elec.)			
	VMC	DMC		
	Pres. wk	Pres. wk	Spink <i>et al.</i>	Ceperley & Alder
0.5	3.4255(1)	3.42541(8)	3.43011(4)	...
0.75	1.28625(5)	1.28620(5)
1.0	0.58643(4)	0.58640(2)	0.58780(1)	0.5870(5)
2.0	0.00195(2)	0.001917(9)	0.002380(5)	0.002050(2)
3.0	-0.06728(1)	-0.067309(9)	-0.067075(4)	...
4.0	-0.07767(1)	-0.07771(1)
5.0	-0.07594(1)	-0.07597(1)	-0.075881(1)	-0.07560(5)
7.0	-0.066304(5)	-0.066348(2)
10.0	-0.053503(2)	-0.053527(5)	-0.0535116(5)	-0.0533750(3)
20.0	-0.031720(6)	-0.031755(3)	-0.0317686(5)	-0.0316450(1)

Following Ceperley [21], we fit

$$E_c(r_s) = \frac{\gamma}{1 + \beta_1 \sqrt{r_s} + \beta_2 r_s} \quad (7)$$

to our SJB-DMC correlation energies (Fig. 2). The fitting parameters γ , β_1 , and β_2 are $-0.151(5)$ Ha, $1.18(7)$, and $0.338(5)$, respectively. According to the all-orders perturbation theory of Gell-Mann and Brueckner [17] the correlation energy at high density is given by $E_c = A \ln(r_s) + C + O(r_s \ln(r_s))$, where $A = \frac{1}{\pi^2} [1 - \ln(2)] \approx 0.0311$ Ha and $C \approx -0.0465$ Ha. The appearance of powers of $\ln(r_s)$ in this formula shows that the correlation energy is a nonanalytic function of r_s for $r_s \rightarrow 0$ and describes the failure of the naive perturbation approach. The constant term C is the sum of the second order Onsager's exchange integral and a numerical constant caused by the sum over divergent contributions [48]. In practice, this asymptotic formula is accurate only for very small $r_s \ll 1$.

We fitted our VMC and DMC correlation energies for $r_s = 0.5, 0.75,$ and 1 to the Gell-Mann-Brueckner expression (Fig. 2). The fitting parameters are $A_{\text{VMC}} = 0.0250(5)$ Ha, $C_{\text{VMC}} = -0.06030(2)$ Ha, $A_{\text{DMC}} = 0.0250(5)$ Ha, and $C_{\text{DMC}} = -0.06040(1)$ Ha. The difference between the VMC and DMC fitting parameters is small because the random errors due to twist averaging at high density dominate. Figure 2 shows that at $r_s \leq 0.1$ the correlation energy predicted by the Gell-Mann-Brueckner formula becomes smaller than VMC and DMC.

V. CONCLUSION

In conclusion, we have performed VMC and DMC simulations using SJB trial wave functions to calculate the correlation energy of the paramagnetic 3D HEG at high and intermediate densities. We corrected finite-size errors by twist averaging and extrapolation to the thermodynamic limit. Our correlation energies, which are lower

TABLE III. Correlation energies for the spin-unpolarized 3D HEG from the PZ81 [27], VWN80 [28], PW92 [29], and DPI [12] parameters, DMC (Spink *et al.* [23]), and this work (VMC and DMC).

r_s	Correlation energy (eV/elec.)						
	PZ81	VWN80	PW92	DPI	DMC (Spink <i>et al.</i>)	VMC	DMC
0.5	-2.069	-2.097	-2.085	-2.108	-1.996	-2.121(3)	-2.124(2)
0.75	-1.829(1)	-1.831(1)
1.0	-1.623	-1.633	-1.627	-1.637	-1.605	-1.642(1)	-1.6432(5)
2.0	-1.227	-1.219	-1.218	-1.215	-1.218	-1.2301(5)	-1.2310(2)
3.0	-1.013	-1.004	-1.005	-0.996	-1.010	-1.0159(3)	-1.0166(2)
4.0	-0.8758(3)	-0.8769(3)
5.0	-0.771	-0.766	-0.768	-0.755	-0.774	-0.7756(3)	-0.7764(3)
7.0	-0.6368(1)	-0.6380(1)
10.0	-0.505	-0.485	-0.505	-0.495	-0.510	-0.5098(1)	-0.5105(2)
20.0	-0.313	-0.302	-0.314	-0.308	-0.316	-0.3149(1)	-0.3159(1)

than previously reported results, can be used to parameterize the correlation energy of the 3D HEG for use in DFT calculations.

ACKNOWLEDGMENTS

S.A. acknowledges support from the UK EPSRC Grant No. EP/P015794/1, the Royal Society, and PRACE (Partnership for Advanced Computing in Europe) for awarding us access to the High-Performance Computing Center Stuttgart, Germany, through the Project No. 2020235573.

-
- [1] S. Azadi and N. D. Drummond, *Phys. Rev. B* **105**, 245135 (2022).
- [2] G. Cassella, H. Sutterud, S. Azadi, N. D. Drummond, D. Pfau, J. S. Spencer, and W. M. C. Foulkes, arXiv:2202.05183 (2022).
- [3] P.-O. Löwdin, *Phys. Rev.* **97**, 1509 (1955).
- [4] R. O. Jones, *Rev. Mod. Phys.* **87**, 897 (2015).
- [5] R. M. Dreizler and E. K. U. Gross, *Density Functional Theory* (Springer-Verlag: Berlin, 1990).
- [6] D. Pines and P. Nozières, *The Theory of Quantum Liquids* (W.A. Benjamin, Inc., New York, 1966).
- [7] G. Giuliani and G. Vignale, *Quantum Theory of the Electron Liquid* (Cambridge University Press, Cambridge, UK, 2005).
- [8] P.-F. Loos and P. M. W. Gill, *Wiley Interdiscip. Rev.: Comput. Mol. Sci.* **6**, 410 (2016).
- [9] T. Dornheim, S. Groth, T. Sjostrom, F. D. Malone, W. M. C. Foulkes, and M. Bonitz, *Phys. Rev. Lett.* **117**, 156403 (2016).
- [10] S. Groth, T. Dornheim, T. Sjostrom, F. D. Malone, W. M. C. Foulkes, and M. Bonitz, *Phys. Rev. Lett.* **119**, 135001 (2017).
- [11] T. Dornheim, J. Vorberger, Z. A. Moldabekov, and P. Tolias, *Phys. Rev. Res.* **4**, 033018 (2022).
- [12] J. Sun, J. P. Perdew, and M. Seidl, *Phys. Rev. B* **81**, 085123 (2010).
- [13] P.-F. Loos and P. M. W. Gill, *Phys. Rev. B* **84**, 033103 (2011).
- [14] P. Bhattarai, A. Patra, C. Shahi, and J. P. Perdew, *Phys. Rev. B* **97**, 195128 (2018).
- [15] P. Hohenberg and W. Kohn, *Phys. Rev.* **136**, B864 (1964).
- [16] W. Kohn and L. J. Sham, *Phys. Rev.* **140**, A1133 (1965).
- [17] M. Gell-Mann and K. A. Brueckner, *Phys. Rev.* **106**, 364 (1957).
- [18] D. M. Ceperley and B. J. Alder, *Phys. Rev. Lett.* **45**, 566 (1980).
- [19] M. Holzmann and S. Moroni, *Phys. Rev. Lett.* **124**, 206404 (2020).
- [20] D. M. Ceperley, G. V. Chester, and M. H. Kalos, *Phys. Rev. B* **16**, 3081 (1977).
- [21] D. M. Ceperley, *Phys. Rev. B* **18**, 3126 (1978).
- [22] S. Azadi, N. D. Drummond, and W. M. C. Foulkes, *Phys. Rev. Lett.* **127**, 086401 (2021).
- [23] G. G. Spink, N. D. Drummond, and R. J. Needs, *Phys. Rev. B* **88**, 085121 (2013).
- [24] J. J. Shepherd, G. H. Booth, A. Grüneis, and A. Alavi, *Phys. Rev. B* **85**, 081103(R) (2012).
- [25] M. Ruggeri, P. Ríos López, and A. Alavi, *Phys. Rev. B* **98**, 161105 (2018).
- [26] W. M. C. Foulkes, L. Mitas, R. J. Needs, and G. Rajagopal, *Rev. Mod. Phys.* **73**, 33 (2001).
- [27] J. P. Perdew and A. Zunger, *Phys. Rev. B* **23**, 5048 (1981).
- [28] S. H. Vosko, L. Wilk, and M. Nusair, *Can. J. Phys.* **58**, 1200 (1980).
- [29] J. P. Perdew and Y. Wang, *Phys. Rev. B* **45**, 13244 (1992).
- [30] M. Marchi, S. Azadi, M. Casula, and S. Sorella, *J. Chem. Phys.* **131**, 154116 (2009).




FIG. 2. Correlation energy of the spin-unpolarized 3D HEG as a function of r_s . (Top panel) The VMC and DMC correlation energies are fitted to Eq. (7) and are compared with the PZ81 [27] parameterization. (Middle panel) VMC and DMC correlation energies at high density ($r_s = 0.5, 0.75$, and 1) are fitted to Gell-Mann and Brueckner’s asymptotic formula. The fitting parameters are presented in the main text. (Bottom panel) Difference between our DMC correlation energy (Pres. wk-DMC) and the PZ81 [27], VWN80 [28], PW92 [29], and DPI [12] DFT functionals.

- [31] Y. Kwon, D. M. Ceperley, and R. M. Martin, Phys. Rev. B **58**, 6800 (1998).
- [32] P. López Ríos, A. Ma, N. D. Drummond, M. D. Towler, and R. J. Needs, Phys. Rev. E **74**, 066701 (2006).
- [33] T. Kato, Commun. Pure Appl. Math. **10**, 151 (1957).
- [34] R. T. Pack and W. B. Brown, J. Chem. Phys. **45**, 556 (1966).
- [35] Y. Kwon, D. M. Ceperley, and R. M. Martin, Phys. Rev. B **48**, 12037 (1993).
- [36] P. López Ríos, P. Seth, N. D. Drummond, and R. J. Needs, Phys. Rev. E **86**, 036703 (2012).
- [37] C. J. Umrigar, K. G. Wilson, and J. W. Wilkins, Phys. Rev. Lett. **60**, 1719 (1988).
- [38] N. D. Drummond and R. J. Needs, Phys. Rev. B **72**, 085124 (2005).
- [39] J. Toulouse and C. J. Umrigar, J. Chem. Phys. **126**, 084102 (2007).
- [40] R. J. Needs, M. D. Towler, N. D. Drummond, P. López Ríos, and J. R. Trail, J. Chem. Phys. **152**, 154106 (2020).
- [41] C. Lin, F. H. Zong, and D. M. Ceperley, Phys. Rev. E **64**, 016702 (2001).
- [42] S. Azadi and W. M. C. Foulkes, J. Chem. Phys. **143**, 102807 (2015).
- [43] N. D. Drummond, R. J. Needs, A. Sorouri, and W. M. C. Foulkes, Phys. Rev. B **78**, 125106 (2008).
- [44] M. Holzmann, R. C. Clay, III, M. A. Morales, N. M. Tubman, D. M. Ceperley, and C. Pierleoni, Phys. Rev. B **94**, 035126 (2016).
- [45] S. Azadi and W. M. C. Foulkes, Phys. Rev. B **100**, 245142 (2019).
- [46] S. Chiesa, D. M. Ceperley, R. M. Martin, and M. Holzmann, Phys. Rev. Lett. **97**, 076404 (2006).
- [47] Raw quantum Monte Carlo data for all the system sizes studied are presented in the supplemental material.
- [48] L. Onsager, L. Mittag, and M. J. Stephen, Ann. Phys. **473**, 71 (1966).