

A Gauge Theory for Shallow Water

David Tong

*Department of Applied Mathematics and Theoretical Physics
University Cambridge, CB3 0WA, UK*

E-mail: d.tong@damtp.cam.ac.uk

ABSTRACT: The shallow water equations describe the horizontal flow of a thin layer of fluid with varying height. We show that the equations can be rewritten as a $d = 2 + 1$ dimensional gauge theory with a Chern-Simons term. The theory contains two Abelian gauge fields, corresponding to the conserved height and conserved vorticity of the fluid. In a certain linearised approximation, the shallow water equations reduce to relativistic Maxwell-Chern-Simons theory. This describes Poincaré waves. The chiral edge modes of the theory are identified as coastal Kelvin waves.

Contents

1	Introduction	1
2	A Gauge Theory for Shallow Water	3
3	The Linearised Shallow Water Equations	7
3.1	Poincaré Waves	8
3.2	A Linearised Gauge Theory	9
3.3	Coastal Kelvin Waves in Maxwell-Chern-Simons	12
A	An Alternative Derivation of the Gauge Theory	13

1 Introduction

The shallow water equations describe the dynamics of a thin layer of fluid whose height is much smaller than its horizontal extent. The basic equations, and a number of extensions, are ubiquitous in modelling of the atmosphere, oceans, rivers, and lakes.

The variables of the shallow water equations are the height $h(x, y, t)$ of the fluid, and the horizontal velocity $\mathbf{u}(x, y, t)$. Writing the components of the 2d velocity vector as u^i , with $i = 1, 2$, the equations are

$$\frac{Dh}{Dt} = -h\nabla \cdot \mathbf{u} \quad \text{and} \quad \frac{Du^i}{Dt} = f\epsilon^{ij}u^j - g\frac{\partial h}{\partial x^i} \quad (1.1)$$

Here the material time derivative, $Dh/Dt = \partial h/\partial t + \mathbf{u} \cdot \nabla h$, tracks the variation of height with the flow. Similarly, $Du^i/Dt = \partial u^i/\partial t + (\mathbf{u} \cdot \nabla)u^i$.

The shallow water equations contain two parameters: the gravitational acceleration g , and the Coriolis parameter f . On Earth, at latitude θ , the Coriolis parameter is given by $f = 4\pi \sin \theta (\text{day})^{-1}$. A derivation of the shallow water equations, together with many detailed applications can be found in the textbook [1]. A whirlwind tour of some highlights can be found in Section 4.3 of the lecture notes [2].

In recent years, a number of surprising parallels have been found between the shallow water equations, at least in their linearised form, and topological phases of matter. With a healthy dose of hindsight, the story goes back to the work of William Thomson who, in 1879, showed that the linearised shallow water equations admit chiral edge modes, now known as *coastal Kelvin waves* [3]. These waves are exponentially localised at the coast and propagate clockwise around land masses in the Northern hemisphere and anti-clockwise in the Southern hemisphere. Similar chiral modes in the shallow water equations are also found localised at the equator, now propagating from west to east [4, 5]. At a calculational level, these chiral waves are strikingly reminiscent of the chiral fermion zero modes of the Dirac equation that were discovered a century later [6].

In the quantum world, chiral edge modes are the smoking gun for a topological phase of matter [7–10]. This is also true for the classical shallow water equations, as first shown in the beautiful paper [11]. We will now give a cartoon version of the argument.

The linearised shallow water equations admit a band of solutions known as *Poincaré waves*, whose frequency ω is related to their wavevector \mathbf{k} through the dispersion relation

$$\omega^2 = c^2 \mathbf{k}^2 + f^2 \tag{1.2}$$

Here the speed is given by $c = \sqrt{gH}$, with H the average height of the fluid. (The derivation of this dispersion relation will be reviewed in Section 3.) For non-vanishing Coriolis force, $f \neq 0$, the frequency has a gap. But the gap closes at the equator where $f = 0$. Given such a gapped band of solutions, labelled by 2d momentum \mathbf{k} , one can follow lattice models and compute the Chern number by integrating a suitable Berry curvature over momentum space, á la TKNN [12] and Haldane [13]. This Chern number is non-vanishing, seemingly integer-valued, and depends on the sign of f . This means that the Chern class jumps as we cross the equator. This change in the underlying topology can be viewed as responsible for the existence of chiral equatorial waves.

The cartoon sketched above is sloppy for a number of reasons, not least the fact that momentum space in this problem is non-compact. This is in sharp contrast to the original uses of a Chern number which was for lattice models where momentum space is a compact Brillouin zone [12, 13]. This compactness is necessary for the Chern number to be an integer, and hence topological. To circumvent this issue, the original paper [11] used a slightly more subtle topological invariant, and subsequent papers have explored the possibility of regulating the momentum integral in some way [14, 15], although not without inducing further diversions [16]. Meanwhile, a topological explanation for the

original coastal Kelvin waves has also been presented, involving spatially varying $f(\mathbf{x})$ and mean height $H(\mathbf{x})$ [17]. These complications notwithstanding, the main lesson is clear: chiral modes observed in shallow water owe their existence to topology. Further studies of topological properties of the shallow water system can be found in [19–24].

The purpose of this work is to provide an effective field theory description of shallow water dynamics in the hope that it may make topological properties more manifest. We do this by writing the shallow water equations as a gauge theory. This gauge theory involves Chern-Simons terms, which are the archetypal example of a topological field theory. In particular, Chern-Simons theories describe quantum Hall states [25, 26].

In Section 2, we show that the non-linear shallow water equations have a natural rendering as a theory of two interacting Abelian gauge fields. The two magnetic fields are associated to the height and vorticity in the shallow water equations and the conservation of these quantities in the gauge theory language follows from the Bianchi identity. The two gauge fields are coupled through a Chern-Simons term. We will see that the vorticity gauge field must be formulated in a so-called Clebsch parameterisation and therein hides many of the subtleties of the system.

In Section 3, we turn to the linearised equations. We will show that the effective field theory describing Poincaré waves (1.2) is given by relativistic Maxwell-Chern-Simons theory. The coastal Kelvin waves arise as chiral edge modes of this Chern-Simons theory.

2 A Gauge Theory for Shallow Water

In this section we formulate the shallow water equations as a gauge theory in $d = 2 + 1$ dimensions. Our strategy will be to identify conserved currents in the shallow water equations and replace them with gauge fields. Rather than simply write down the Lagrangian, we will instead proceed in smaller steps, and occasional missteps, to build some intuition for the gauge theoretic formulation. The more impatient reader can find the final answer in (2.7) and (2.9).

To understand the physics, we focus on conserved quantities. The shallow water equations (1.1) enjoy a number of conservation laws, but for our immediate purposes we need just two. Both are conserved quantities in the sense that they admit a current obeying the continuity equation,

$$\partial_\mu J^\mu = \frac{\partial J^0}{\partial t} + \nabla \cdot \mathbf{J} = 0$$

Throughout, we use $\mu = 0, 1, 2$ to denote space and time indices, and $i = 1, 2$ to denote only spatial indices. Note that, unlike in relativistic systems, $x^0 = t$ with no additional factor of speed. This means that x^0 and x^i have different dimensions.

The first conserved quantity is the height h of the fluid. Indeed, the first equation in (1.1) is a continuity equation with

$$J^0 = h \quad \text{and} \quad \mathbf{J} = h\mathbf{u} \quad (2.1)$$

This is a manifestation of the conservation of mass of the fluid. The second conserved quantity is associated to the vorticity, defined in two dimensions as

$$\zeta = \frac{\partial u^2}{\partial x} - \frac{\partial u^1}{\partial y}$$

The presence of the Coriolis force means that there is a slight modification to the conserved current, which is given by

$$\tilde{J}^0 = \zeta + f \quad \text{and} \quad \tilde{\mathbf{J}} = (\zeta + f)\mathbf{u} \quad (2.2)$$

Importantly, the two conserved currents (2.1) and (2.2) are not fully independent: they obey the relation $\mathbf{J}/J^0 = \tilde{\mathbf{J}}/\tilde{J}^0 = \mathbf{u}$. Part of the challenge will be to reproduce this relation when the currents are replaced by gauge fields.

We now explain what it means to replace a conserved current with a gauge field. We denote our gauge field as A_μ , with $\mu = 0, 1, 2$. In $d = 2 + 1$ dimensions, the electric field is a vector E_i , while the magnetic field B is a (pseudo)-scalar,

$$E_i = \frac{\partial A_i}{\partial t} - \frac{\partial A_0}{\partial x^i} \quad \text{and} \quad B = \frac{\partial A_2}{\partial x^1} - \frac{\partial A_1}{\partial x^2}$$

Both are invariant under gauge transformations $A_\mu \rightarrow A_\mu + \partial_\mu \theta$.

Abelian gauge fields in $d = 2 + 1$ dimensions are special because they come with an associated conserved, global charge. This follows simply from the Bianchi identity which reads

$$\epsilon^{\mu\nu\rho} \partial_\mu \partial_\nu A_\rho = 0 \quad \Rightarrow \quad \frac{\partial B}{\partial t} - \frac{\partial E_2}{\partial x^1} + \frac{\partial E_1}{\partial x^2} = 0 \quad (2.3)$$

This has the form of the continuity equation, with the magnetic field as the charge and the components of the electric field as the current,

$$J^0 = B \quad \text{and} \quad \mathbf{J} = (-E_2, E_1) \quad \text{or} \quad J^i = -\epsilon^{ij} E_j$$

To formulate a gauge theory description of shallow water, we will think of conserved currents in terms of gauge fields. Ultimately we will introduce gauge fields associated to both the height (2.1) and the vorticity (2.2). But we will build up slowly, and we start with just a single gauge field A_μ that we identify as the height current, meaning that we take

$$B = h \quad \text{and} \quad E_i = \epsilon_{ij} h u^j \quad (2.4)$$

This single gauge field includes all degrees of freedom of the shallow water system, with the velocity given by $u^i = -\epsilon^{ij} E_j / B$. The first of the shallow water equations (1.1) is automatically obeyed by virtue of the Bianchi identity (2.3). It remains to write the second shallow water equation in the language of the gauge theory. For this, we will adopt an action principle. We will make a first attempt at writing down an action for the shallow water system and see that it fails. But the way in which it fails will be instructive.

To motivate the action, we first note that the shallow water system has a conserved energy, given by

$$\mathcal{E} = \frac{1}{2} h \mathbf{u}^2 + \frac{1}{2} g h^2 = \frac{\mathbf{E}^2}{2B} + \frac{1}{2} g B^2$$

This takes the form of kinetic energy plus potential energy, so a natural guess for the action is

$$S_{\text{fail}} = \int dt d^2x \left(\frac{\mathbf{E}^2}{2B} - \frac{1}{2} g B^2 \right) \quad (2.5)$$

The fact that there is an inverse power of B in the kinetic term should cause no more concern than an inverse power of the metric in the Einstein-Hilbert action: the dynamics should be thought of as an expansion around a non-vanishing fluid height, meaning that $B \neq 0$.

To derive the equations of motion, we vary the action with respect to A_μ . The equation of motion for A_0 is Gauss' law and reads

$$\frac{\partial}{\partial x^i} \left(\frac{E^i}{B} \right) = 0 \quad \Rightarrow \quad \zeta = 0 \quad (2.6)$$

This is the first way in which the action (2.5) fails: Gauss' law only allows irrotational flows with the vorticity ζ forced to vanish. We'll rectify this shortly. For now, we note that that vorticity is the analog of electric charge in our theory. This restriction notwithstanding, we fare slightly better with the equations of motion for A_i . These are

$$\frac{\partial}{\partial t} \left(\frac{E_i}{B} \right) + \frac{1}{2} \epsilon_{ij} \frac{\partial}{\partial x^j} \left(\frac{\mathbf{E}^2}{B^2} \right) + g \epsilon_{ij} \frac{\partial B}{\partial x^j} = 0$$

When translated into the the variables h and \mathbf{u} , the equation becomes

$$\dot{u}^i + \frac{1}{2} \frac{\partial \mathbf{u}^2}{\partial x^i} = -g \frac{\partial h}{\partial x^i} \quad \Rightarrow \quad \frac{Du^i}{Dt} + \epsilon^{ij} u^j \zeta = -g \frac{\partial h}{\partial x^i}$$

With the Gauss' law constraint $\zeta = 0$, this gives us something close to the shallow water equation (1.1). We see that, in addition to the fact the vorticity is forced to vanish, the action does not accommodate the Coriolis force. Of course, this should be no surprise as we made no attempt to include it.

With this failure behind us, we can now see how to resolve both issues. Again, our first attempt will not be a complete success. But our second will be. Clearly we need to include some additional degrees of freedom, charged under A_μ , so that Gauss' law (2.6) receives an extra contribution, breathing life into the otherwise frozen vorticity. As we've seen, the vorticity itself is also a conserved quantity which motivates us to introduce a second, dynamical gauge field \tilde{A}_μ . We couple this to the first through a Chern-Simons term. At the same time, we also introduce the Coriolis parameter f through a background charge. The resulting action is

$$S = \int dt d^2x \left(\frac{\mathbf{E}^2}{2B} - \frac{1}{2} g B^2 + f A_0 - \epsilon^{\mu\nu\rho} A_\mu \partial_\nu \tilde{A}_\rho \right) \quad (2.7)$$

The Gauss law for this revised action is

$$\frac{\partial}{\partial x^i} \left(\frac{E_i}{B} \right) + f - \tilde{B} = 0 \quad \Rightarrow \quad \tilde{B} = \zeta + f \quad (2.8)$$

This looks more promising: the new magnetic field $\tilde{B} = \partial_1 \tilde{A}_2 - \partial_2 \tilde{A}_1$ is identified with the conserved charge (2.2) associated to vorticity. But we're not yet out of the water. If we treat \tilde{A}_μ as a dynamical gauge field then we must also implement the equation of motion for \tilde{A}_0 and this would seem to fix $B = 0$ or, equivalently, $h = 0$. That's not helpful.

There is a rather charming resolution to this, again specific to gauge fields in $d = 2+1$ dimensions. Any 3d vector field can be written in a so-called *Clebsch parameterisation*. For our gauge field \tilde{A}_μ , this means that we introduce three functions $\chi(\mathbf{x}, t)$, $\beta(\mathbf{x}, t)$ and $\alpha(\mathbf{x}, t)$ and write,

$$\tilde{A}_\mu = \partial_\mu \chi + \beta \partial_\mu \alpha \quad (2.9)$$

This parameterisation is non-linear and non-unique. It does not often arise in quantum field theory but there is a long history of using this parameterisation in fluid mechanics, typically for the velocity field in three spatial dimensions. Here we will see that it will

serve our purposes nicely. We vary the action (2.7) not with respect to \tilde{A}_μ , but instead with respect to the component fields χ , β and α . The field χ is pure gauge and does not contribute to the action. Meanwhile, the equations of motion for β and α are

$$B\dot{\alpha} + \epsilon^{ij} E_i \partial_j \alpha = 0 \quad \text{and} \quad B\dot{\beta} + \epsilon^{ij} E_i \partial_j \beta = 0 \quad (2.10)$$

From this we can determine the electric field \tilde{E}_i . It is defined by

$$\tilde{E}_i = \partial_t \tilde{A}_i - \partial_i \tilde{A}_0 = \dot{\beta} \partial_i \alpha - (\partial_i \beta) \dot{\alpha}$$

From the equations of motion (2.10), we have

$$\tilde{E}_i = -\frac{E_k}{B} \epsilon^{kj} (\partial_j \beta \partial_i \alpha - \partial_i \beta \partial_j \alpha) = \frac{E_i}{B} \tilde{B} = \epsilon_{ij} u^j (\zeta + f)$$

This is what we wanted. The electric field is not independent of our other conserved quantities, but is fixed by the magnetic field \tilde{B} and the original gauge field A_μ . This is the property of the vorticity current (2.2) that we previously highlighted.

The fact that working with the Clebsch parameterisation in an action gives equations that are less restrictive than working directly with a vector field was previously noted in [27]. The observation that it allows us to introduce currents that are not fully independent is related to remarks made in [28] about the role the Clebsch parameterisation plays in the symplectic structure of the Euler equation.

It remains only to compute the equation of motion for the original spatial gauge field, A_i . This is

$$\frac{\partial}{\partial t} \left(\frac{E_i}{B} \right) + \frac{1}{2} \epsilon_{ij} \frac{\partial}{\partial x^j} \left(\frac{\mathbf{E}^2}{B^2} \right) + g \epsilon_{ij} \frac{\partial B}{\partial x^j} - \epsilon_{ij} \tilde{E}_j = 0$$

This time, when translated into the the variables h and \mathbf{u} , this equation coincides with the second shallow water equation (1.1). This confirms that the gauge theory action (2.7), with the Clebsch parameterisation (2.9), does indeed describe the shallow water system. An alternative derivation of this action is given in Appendix A.

3 The Linearised Shallow Water Equations

Much interesting physics, including aspects of topology, can be found in the linearised version of the shallow water equations. We expand about a fluid of constant, average height H , and write

$$h(\mathbf{x}, t) = H + \eta(\mathbf{x}, t)$$

Then, keeping only terms linear in η and \mathbf{u} , the shallow water equations (1.1) become

$$\frac{\partial \eta}{\partial t} + H \nabla \cdot \mathbf{u} = 0 \quad \text{and} \quad \frac{\partial u^i}{\partial t} = f \epsilon^{ij} u^j - g \frac{\partial \eta}{\partial x^i} \quad (3.1)$$

In this section we first review some properties of the solutions to (3.1) and then turn to their description in the language of gauge theory.

3.1 Poincaré Waves

It is straightforward to find solutions to the linearised equations (3.1) by looking at Fourier modes. We write

$$\eta = \eta_0 e^{i(\omega t - \mathbf{k} \cdot \mathbf{x})} \quad \text{and} \quad \mathbf{u} = \mathbf{u}_0 e^{i(\omega t - \mathbf{k} \cdot \mathbf{x})}$$

The linearised shallow water equations (3.1) can then be written in a way that looks suggestively like the Dirac equation

$$\mathcal{H} \Psi = \omega \Psi \quad \text{where} \quad \mathcal{H} = \begin{pmatrix} 0 & ck_x & ck_y \\ ck_x & 0 & -if \\ ck_y & if & 0 \end{pmatrix} \quad \text{and} \quad \Psi = \begin{pmatrix} \sqrt{g/H} \eta_0 \\ u_0 \\ v_0 \end{pmatrix} \quad (3.2)$$

Here we've written the two-component velocity as $\mathbf{u}_0 = (u_0, v_0)$ and defined $c = \sqrt{gH}$ which we recognise as the speed of surface waves in the absence of the Coriolis force. We will see that c plays a similar role in the present context.

The eigenvalue problem (3.2) is easily solved. There are two, gapped bands with frequencies

$$\omega = \pm \sqrt{c^2 \mathbf{k}^2 + f^2} \quad (3.3)$$

These are known as *Poincaré waves*. For small wavelengths, $c|\mathbf{k}| \gg f$, these waves travel with speed c . The long-wavelength modes are affected by the presence of the Coriolis force.

In addition to the Poincaré waves, there is a flat band of solutions to (3.2), with

$$\omega = 0$$

This reflects the fact that there are static solutions to the full non-linear equations (1.1) beyond the obvious $h = \text{constant}$. These solutions have a non-trivial profile in one direction, say $h = h(y)$, with the gravitational force balanced by a corresponding velocity profile $fu = -\partial h / \partial y$ generating a Coriolis force. This is known as *geostrophic balance*. These static solutions manifest themselves in the linear problem as a flat band.

There is one other feature of the linearised shallow water equations (3.1) that we will need. This follows from conservation laws. The linearised version of the conservation of height (2.1) and vorticity (2.2) becomes

$$\frac{\partial \eta}{\partial t} + H \nabla \cdot \mathbf{u} = 0 \quad \text{and} \quad \frac{\partial \zeta}{\partial t} + f \nabla \cdot \mathbf{u} = 0$$

Eliminating $\nabla \cdot \mathbf{u}$ from both of these expressions gives us the surprisingly powerful formula

$$\dot{Q} = 0 \quad \text{with} \quad Q = H\zeta - f\eta \tag{3.4}$$

This is much stronger than most conservation laws: it is telling us that there is a function $Q(\mathbf{x})$ that does not change in time. The function Q is called *potential vorticity*¹.

We can ask: what is the potential vorticity evaluated on the solutions to the eigenvalue problem (3.2)? It is simple to check that the flat band, with $\omega = 0$, has

$$Q \sim \sqrt{\frac{H}{g}} (c^2 \mathbf{k}^2 + f^2) \tag{3.5}$$

Meanwhile, the Poincaré waves with dispersion relation (3.3) have

$$Q = 0 \tag{3.6}$$

This fact will be important in the next section when we write down an effective field theory for the Poincaré waves.

3.2 A Linearised Gauge Theory

We now return to our gauge theory (2.7). We will see how the above phenomena arise in this language. We linearise the A_μ gauge fields as

$$A_\mu = \hat{A}_\mu + \delta A_\mu \quad \text{with} \quad \hat{A}_0 = 0 \quad \text{and} \quad \partial_1 \hat{A}_2 - \partial_2 \hat{A}_1 = H$$

We're going to abuse notation slightly and refer to the fluctuation δA_μ simply as A_μ . (In other words, we make the substitution, $A_\mu \rightarrow \hat{A}_\mu + A_\mu$.) This will make the subsequent equations somewhat clearer. The translation (2.4) to the fluid variables then becomes

$$B = \eta \quad \text{and} \quad E_i = H \epsilon_{ij} u^j \tag{3.7}$$

¹The existence of a function that is invariant in time is special to the linearised equations. In the full non-linear equations, potential vorticity is defined as $Q = (\zeta + f)/h$. It is not conserved in the sense that it obeys a continuity equation, but it does have vanishing material derivative: $DQ/Dt = 0$.

which ensures that the first equation in (3.1) is obeyed courtesy of the Bianchi identity. Meanwhile, we also linearise the Clebsch gauge field \tilde{A}_μ , with components

$$\beta = \hat{\beta} + p \quad \text{and} \quad \alpha = \hat{\alpha} + q \quad \text{with} \quad \partial_1 \hat{\beta} \partial_2 \hat{\alpha} - \partial_2 \hat{\beta} \partial_1 \hat{\alpha} = f \quad (3.8)$$

The fluctuations are written as $\delta\beta = p$ and $\delta\alpha = q$ to avoid an overload of $\partial\delta$'s in the expressions below. Expanding the action (2.7), we find that the terms linear in fluctuations vanish, as they must, while the terms quadratic in fluctuations read

$$S = \int dt d^2x \left(\frac{1}{2H} \mathbf{E}^2 - \frac{1}{2} g B^2 - H p \dot{q} + \epsilon^{ij} E_i \left(q \partial_j \hat{\beta} - p \partial_j \hat{\alpha} \right) \right) \quad (3.9)$$

The action is both simple and rather opaque. We will attempt to shed some light. We first look at the equations of motion. Gauss' law is

$$\partial_i E_i = H \epsilon^{ij} \left(\partial_i \hat{\beta} \partial_j q - \partial_i \hat{\alpha} \partial_j p \right) \quad (3.10)$$

while the equation of motion for A_i is

$$\dot{E}_i = -g H \epsilon_{ij} \partial_j B - H \epsilon^{ij} \left(\partial_j \hat{\beta} \dot{q} - \partial_j \hat{\alpha} \dot{p} \right) \quad (3.11)$$

Finally, the equations of motion for p and q read

$$H \dot{q} = -\epsilon^{ij} E_i \partial_j \hat{\alpha} \quad \text{and} \quad H \dot{p} = -\epsilon^{ij} E_i \partial_j \hat{\beta} \quad (3.12)$$

We can substitute (3.12) into (3.11) to find

$$\dot{E}_i = \epsilon_{ij} (f E_j - g H \partial_j B)$$

This coincides with the second linearised shallow water equation in (3.1) using the dictionary (3.7).

Things look somewhat simpler if we work in the gauge $A_0 = 0$. We treat p in (3.9) as a Lagrange multiplier, imposing the equation (3.12) for q . The linearised action (3.9) then becomes

$$S = \int dt d^2x \frac{1}{2H} \left(\dot{A}_i^2 - c^2 B^2 + f \epsilon^{ij} A_i \dot{A}_j \right) \quad (3.13)$$

where we have used the condition $\epsilon_{ij} \partial_i \hat{\beta} \partial_j \hat{\alpha} = f$. We recognise this as the action for Maxwell-Chern-Simons theory in $A_0 = 0$ gauge. However, it is not quite the same. This is because the action (3.13) should be accompanied by Gauss' law constraint (3.10). As we will now see, the presence of the supplementary variables q and p mean that this constraint is somewhat malleable, and can vary depending on the context.

The action (3.13) has two classes of solutions. We will now see that these correspond to the geostrophic flat band and the Poincaré waves.

- The flat band: The equations of motion arising from (3.13) have a class of solutions with arbitrary time-independent magnetic field $B(\mathbf{x})$, given by

$$A_i = \bar{A}_i + \frac{c^2}{f} \partial_i B t$$

where $\dot{\bar{A}}_i = 0$, and $B = \epsilon^{ij} \partial_i A_j = \epsilon^{ij} \partial_i \bar{A}_j$. This has $\ddot{A}_i = 0$ and corresponds to the flat band with vanishing frequency. The equations (3.12) for the supplementary variables q and p are solved by

$$Hq = -\epsilon^{ij} Z_i \partial_j \hat{\alpha} \quad \text{and} \quad Hp = -\epsilon^{ij} Z_i \partial_j \hat{\beta} \quad \text{with} \quad Z_i = \frac{c^2}{f} \left(\partial_i B t - \frac{1}{f} \epsilon_{ij} \partial_j B \right)$$

These solutions obey the Gauss' law constraint (3.10) which now takes the form

$$\partial_i E_i = \frac{c^2}{f} \nabla^2 B$$

While (3.13) coincides with the action for Maxwell-Chern-Simons theory in $A_0 = 0$ gauge, this is not the Gauss' law for that theory. That's why these flat band solutions are unfamiliar in the context of Chern-Simons theory.

- Poincaré waves: The other class of solutions to the equations of motion arising from (3.13) are more familiar. They are simply wave solutions of the form $A_i(\mathbf{x}, t) = e^{i(\omega t - \mathbf{k} \cdot \mathbf{x})} \hat{A}_i$. One can check that the equations of motion impose the relativistic dispersion relation (3.3). The supplementary variables are now solved by

$$Hq = -\epsilon^{ij} A_i \partial_j \hat{\alpha} \quad \text{and} \quad Hp = -\epsilon^{ij} A_i \partial_j \hat{\beta}$$

with Gauss' law (3.10) now taking the form

$$\partial_i E_i = f B \tag{3.14}$$

Translating back to the fluid variables, this tells us that the potential vorticity is $Q = H\zeta - f\eta = 0$, in agreement with expectations from Poincaré waves in (3.6). This time the form of Gauss' law (3.14) does coincide with that of relativistic Maxwell-Chern-Simons theory. Indeed, we can reinstate A_0 in the action so that its equations of motion reproduce (3.14). The result is the familiar action

$$S = \int dt d^2x \frac{1}{2H} (\mathbf{E}^2 - c^2 B^2 - f \epsilon^{\mu\nu\rho} A_\mu \partial_\nu A_\rho) \tag{3.15}$$

This, of course, is Maxwell-Chern-Simons theory. Here, it governs the dynamics of Poincaré waves. It is simple to check that the profiles of the electric and magnetic fields coincide with the velocity and height of Poincaré waves, using the dictionary (3.7).

3.3 Coastal Kelvin Waves in Maxwell-Chern-Simons

Viewed as a quantum field theory, the action (3.15) flows at low energies to a topological field theory. Indeed, it is perhaps the simplest topological field theory and actions of this type are well known to describe quantum Hall fluids [25, 26]. Here, of course, we treat (3.15) as a purely classical theory. Nonetheless, a remnant of its topological nature survives in chiral edge modes.

For concreteness, we consider the theory with a spatial boundary at $x = 0$. Varying the action then gives, in addition to the bulk term, a boundary term

$$\delta S = \delta S_{\text{bulk}} + \frac{1}{H} \int dt dy \left[-(E_1 - 2fA_2)\delta A_0 - (c^2B + 2fA_0)\delta A_2 \right] \quad (3.16)$$

The bulk term δS_{bulk} vanishes when the equations of motion are obeyed. But to have a well defined variational principle we must impose boundary conditions to ensure that the boundary term also vanishes. We will describe our boundary condition of choice shortly.

In the presence of a boundary, Chern-Simons theory comes with an additional concern. The Chern-Simons action is invariant under gauge transformations only up to a total derivative. This means that the theory risks a failure of gauge invariance in the presence of a boundary. Under a gauge transformation $A_\mu \rightarrow A_\mu + \partial_\mu \theta$, the action (3.15) transforms as

$$S \rightarrow S + \frac{f}{2H} \int dt dy \theta E_2 \quad (3.17)$$

There are various ways to deal with this lack of gauge invariance, depending on the boundary condition that we impose to ensure that (3.16) vanishes. In the context of the quantum Hall effect, one typically restricts to gauge transformations θ that vanish on the boundary. But restricting the allowed gauge transformations resurrects modes that were previously gauge redundancies [29]. These are the chiral modes on the boundary. The chiral modes that arise in this way are closely related to the anomaly inflow arguments that underlie the bulk/boundary correspondence for topological insulators.

The chiral modes that arise as coastal Kelvin waves are somewhat different because we impose different boundary conditions. This time, the boundary term in (3.16) vanishes by setting $A_0 = A_2 = 0$ on the boundary. This ensures that $E_2 = 0$ on the boundary so the gauge variation of the action (3.17) also vanishes. In electromagnetism, this is the boundary condition appropriate for a conductor. In fluid mechanics it is the relevant boundary condition because it means that $\mathbf{u} \cdot \hat{\mathbf{x}} = 0$, so that no fluid flows into the boundary. These boundary conditions in Chern-Simons theory, and their connection to anomaly inflow, were also considered recently in [30].

The bulk equations of motion of Maxwell-Chern-Simons theory are

$$\partial_i E_i = fB \quad \text{and} \quad \dot{E}_i = -c^2 \epsilon_{ij} \partial_j B + f \epsilon_{ij} E_j$$

We can look for solutions with $A_0 = A_2 = 0$ everywhere. We make the ansatz

$$A_1 = A(x) e^{i(\omega t - ky)}$$

The bulk equations tell us that

$$\dot{E}_1 = -c^2 \partial_2 B \quad \Rightarrow \quad \omega^2 = c^2 k^2 \quad \Rightarrow \quad \omega = \pm ck$$

and

$$\partial_1 E_1 = fB \quad \Rightarrow \quad \omega A' = kfA \quad \Rightarrow \quad A' = \pm \frac{f}{c} A$$

For $f > 0$, corresponding to the Northern hemisphere, a fluid restricted to lie in the region $x > 0$ has the normalisable solution only if

$$\omega = -ck \quad \Rightarrow \quad A(x) \sim e^{-fx/c}$$

This is the chiral, coastal Kelvin wave, first found in [3]. It travels in the direction of decreasing y which, in this context, means southward. Said differently, the wave moves around land masses in the clockwise direction in the Northern hemisphere. The novelty here is to see this wave emerging from the effective Maxwell-Chern-Simons description.

A An Alternative Derivation of the Gauge Theory

There is a long history of constructing variational principles for fluids in the Eulerian framework. The results do not usually look like a gauge theory. In this appendix we provide an alternative derivation of the gauge theory action (2.7) that makes contact with the more traditional approach.

For fluids in d spatial dimensions, one usually starts by introducing a map from physical spacetime to the positions of fluid parcels.

$$\alpha : \mathbb{R}^d \times \mathbb{R} \rightarrow \mathbb{R}^d$$

We write this map as $\alpha^a(\mathbf{x}, t)$ with $a = 1, \dots, d$. The map must be a diffeomorphism, with $\det(\partial\alpha^a/\partial x^i) \neq 0$. A discussion of this approach to fluids can be found in the textbook [31]. More recently, it has been revived in the context of relativistic fluids [32, 33].

Given a map $\alpha^a(\mathbf{x}, t)$, the velocity field \mathbf{u} is defined by the equation

$$\frac{D\alpha^a}{Dt} \equiv \frac{\partial\alpha^a}{\partial t} + \mathbf{u} \cdot \nabla\alpha^a = 0 \quad (\text{A.1})$$

This captures the idea that the physical label $\alpha^a(\mathbf{x}, t)$ of the point in the fluid is unchanged as the fluid moves. The velocity \mathbf{u} defined in this way is invariant under diffeomorphisms of α . Note, however, that this definition comes with a rather confusing minus sign, related to active vs passive transformations. This is seen, for example, in the simplest laminar flow $\alpha = (x - ut, y)$ which, according to (A.1), is associated to a velocity field $\mathbf{u} = (+u, 0)$.

For application to the shallow water equations, we have $d = 2$ dimensions. We also have the height variable $h(x, y, t)$, which obeys the continuity equation

$$\frac{Dh}{Dt} + h\nabla \cdot \mathbf{u} = 0 \quad (\text{A.2})$$

reflecting the incompressibility of the underlying fluid.

An action for the shallow water equations in these variables was given in [34]

$$S = \int dt d^2x \left(\frac{1}{2}h\mathbf{u}^2 - \frac{1}{2}gh^2 - h\mathbf{u} \cdot \mathbf{a}(\mathbf{x}) \right) + \phi \left(\frac{Dh}{Dt} + h\nabla \cdot \mathbf{u} \right) - h\beta_a \frac{D\alpha^a}{Dt}$$

This action should be varied with respect to h , \mathbf{u} , ϕ , α^a and β_a . Both β_a and ϕ act as Lagrange multipliers, imposing the conditions (A.1) and (A.2) respectively. We have introduced a background gauge field $\mathbf{a}(\mathbf{x})$ whose role will be to implement the Coriolis force². This background field depends only on \mathbf{x} and not on time. We will determine the form of $\mathbf{a}(\mathbf{x})$ shortly.

²In quantum Hall physics, there is a convention that lower-case gauge fields a_μ are dynamical while upper-case gauge fields A_μ are background. Annoyingly, we have chosen the opposite convention here. As we will see, this has the advantage that the field strength $f_{12} = \partial_1 a_2 - \partial_2 a_1$ coincides with the Coriolis parameter, canonically denoted in geophysics as f .

The equation of motion for \mathbf{u} reveals that the velocity has a Clebsch-like parameterisation,

$$\mathbf{u} = \nabla\phi + \beta_a \nabla\alpha^a + \mathbf{a}(\mathbf{x}) \quad (\text{A.3})$$

This differs from our previous Clebsch parameterisation for the gauge field (2.9) in that we have a pair of potentials (β_a, α_a) , with $a = 1, 2$, with α^a restricted to be a diffeomorphism. In addition, the velocity gets a contribution from the fixed background $\mathbf{a}(\mathbf{x})$.

The equation of motion for α^a reveals a conserved current, involving the Lagrange multiplier β_a ,

$$\frac{\partial(h\beta_a)}{\partial t} + \nabla \cdot (h\beta_a \mathbf{u}) = 0 \quad \Rightarrow \quad \frac{D\beta_a}{Dt} = 0 \quad (\text{A.4})$$

where the second equation follows from the conservation of the height (A.2). Finally, the equation of motion for h gives

$$\frac{D\phi}{Dt} = \frac{1}{2}\mathbf{u}^2 - gh - \mathbf{u} \cdot \mathbf{a} \quad (\text{A.5})$$

where an additional term has been set to zero courtesy of (A.1). Although it is not immediately obvious, equations (A.1), (A.3), (A.4) and (A.5) are equivalent to the second of the shallow water equations (1.1). To see this, we first use (A.3) to compute the material derivative of the velocity

$$\frac{Du_i}{Dt} = (\partial_i \dot{\phi} + u^j \partial_j \partial_i \phi) + (\dot{\beta}_a + u^j \partial_j \beta_a) \partial_i \alpha^a + \beta_a (\partial_i \dot{\alpha}^a + u^j \partial_j \partial_i \alpha^a) + u^j \partial_j a_i$$

Invoking (A.1), (A.4) and (A.5), a short calculation then reveals

$$\frac{Du_i}{Dt} = -f_{ij} u^j - g \partial_i h \quad \text{with } f_{ij} = \partial_i a_j - \partial_j a_i$$

We see that the background field strength f_{ij} acts like a Coriolis force. To reproduce the constant Coriolis force of (1.1), we simply need to find a background $\mathbf{a}(\mathbf{x})$ that satisfies $f_{ij} = -f \epsilon_{ij}$. For example, $\mathbf{a} = f(y, 0)$ or $\mathbf{a} = \frac{1}{2}f(y, -x)$ both do the job. Note that any choice necessarily breaks translational symmetry and/or rotational symmetry, even though the final shallow water equations do not. This is unavoidable in this formalism. The situation is identical to that of a charged particle in a constant background magnetic field.

We now show how this formalism is related to the gauge theory introduced in Section 2. The idea is to use a 3d duality transformation, first introduced in [35] and beloved of many a high-energy and condensed matter theorist. To proceed, it's useful to first integrate by parts, so that the Lagrangian has an overall factor of the height h . (This reflects the fact that the shallow water equations arise after integrating over the z -direction of the thin fluid.)

$$S = \int dt d^2x h \left(\frac{1}{2} \mathbf{u}^2 - \frac{1}{2} gh - \mathbf{u} \cdot \mathbf{a}(\mathbf{x}) - \frac{D\phi}{Dt} - \beta_a \frac{D\alpha^a}{Dt} \right) \quad (\text{A.6})$$

Note that only derivatives of the Lagrange multiplier ϕ appear in this action. This allows us to exchange ϕ for a vector field C_μ , by writing the action as

$$S = \int dt d^2x h \left(\frac{1}{2} \mathbf{u}^2 - \frac{1}{2} gh - \mathbf{u} \cdot \mathbf{a}(\mathbf{x}) - C_0 - u^i C_i - \beta_a \frac{D\alpha^a}{Dt} \right) + \epsilon^{\mu\nu\rho} A_\mu \partial_\nu C_\rho$$

Here we have further introduced a new Lagrange multiplier, A_μ . This can be viewed as a gauge field because the action is invariant under transformation $A_\mu \rightarrow A_\mu + \partial_\mu \theta$. The equation of motion for A_μ gives the constraint $\epsilon^{\mu\nu\rho} \partial_\nu C_\rho = 0$ which can be solved locally by $C_\mu = \partial_\mu \phi$. This takes us back to our original action (A.6). Alternatively, we may instead choose to impose the equation of motion for C_μ , which also acts as a Lagrange multiplier. This gives a constraint on the magnetic field $B = \partial_1 A_2 - \partial_2 A_1$ and electric fields $E_i = \dot{A}_i - \partial_i A_0$,

$$B = h \quad \text{and} \quad E_i = \epsilon_{ij} h u^j$$

These coincide with the relations introduced in (2.4). We may then replace h and \mathbf{u} in the action with our new gauge fields. This gives

$$S = \int dt d^2x \left(\frac{\mathbf{E}^2}{2B} - \frac{1}{2} g B^2 + \epsilon^{ij} E_i a_j(\mathbf{x}) - \epsilon^{\mu\nu\rho} \partial_\mu A_\nu \beta_a \partial_\rho \alpha^a \right)$$

In its essential details, this coincides with the action (2.7) presented in Section 2. The Coriolis term becomes the background charge $f A_0$ after an integration by parts. The form above makes it clear that this term too has its origin as a Chern-Simons term, now involving a background field. After integration by parts, the action respects both translation and rotation symmetry provided that f is constant. Meanwhile, the second gauge field arises automatically in the Clebsch parameterisation in the above presentation,

$$\tilde{A}_\mu = \partial_\mu \chi + \beta_a \partial_\mu \alpha^a$$

Here χ is pure gauge and does not contribute to the action. This differs from (2.9) only in the $a = 1, 2$ index ascribed to α and β . Diffeomorphism invariance ensures that this does not contribute any further degrees of freedom.

There are other equations in fluid mechanics that have a similar structure to the shallow water equations (1.1) and hence lend themselves to a gauge theoretic formalism. For example, a compressible fluid obeys the equations

$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{u}) = 0 \quad \text{and} \quad \rho \frac{D\mathbf{u}}{Dt} = -\nabla P$$

with an equation of state relating the pressure P to the density ρ of the form $P = C\rho^\gamma$, where C is a constant and γ is the ratio of specific heats. When the fluid lives in $d = 2$ spatial dimensions, this too may be written as gauge theory, with $B = \rho$ and $E_i = \epsilon_{ij}\rho u^j$ and with the potential term B^2 in (2.7) replaced by $B^{\gamma-1}$. Again, the slightly disconcerting fact that the power γ is not an integer is no cause for concern when it is appreciated that this theory should be expanded around the solution $B = \rho = \text{constant} \neq 0$.

Acknowledgements

I'm especially grateful to Matt Davison for introducing me to the idea of topology in fluid dynamics and for guiding me through some basic aspects of atmospheric and oceanic flows. Thanks also to Jay Armas, Sean Hartnoll and Max Metlitski for helpful discussions. This work was supported by STFC grant ST/L000385/1, the EPSRC grant EP/V047655/1 "Chiral Gauge Theories: From Strong Coupling to the Standard Model", and a Simons Investigator Award. For the purpose of open access, the author has applied a Creative Commons Attribution (CC BY) licence to any Author Accepted Manuscript version arising from this submission.

References

- [1] G. Vallis, "*Atmospheric and Oceanic Fluid Dynamics: Fundamentals and Large-scale Circulation*", Cambridge University Press (2006)
- [2] D. Tong, "*Lectures on Fluid Mechanics*", www.damtp.cam.ac.uk/user/tong/fluids.html
- [3] W. Thomson, "*On Gravitational Oscillations of Rotating Water*", Proceedings of the Royal Society of Edinburgh, Session 1878-1879, Volume 10 (1880)
- [4] T. Matsuno, "*Quasi-Geostrophic Motions in the Equatorial Area*", Journal of the Meteorological Society of Japan. Ser. II, Vol 44 (1966)
- [5] M. Longuet-Higgins, "*The Eigenfunctions of Laplace's tidal equations over a sphere*", Phil. Trans. Royal Society of London 262: 1132: 511 (1968).
- [6] R. Jackiw and C. Rebbi, "*Solitons with Fermion Number 1/2*," Phys. Rev. D **13** (1976), 3398.
- [7] R.B. Laughlin, "*Quantized Hall Conductivity in Two Dimensions*", Phys. Rev. B **23**, 5632(R)

- [8] B.I Halperin, “*Quantized Hall conductance, current-carrying edge states, and the existence of extended states in a two-dimensional disordered potential*”, Phys. Rev. B **25**, 2185.
- [9] C. G. Callan, Jr. and J. A. Harvey, “*Anomalies and Fermion Zero Modes on Strings and Domain Walls,*” Nucl. Phys. B **250** (1985), 427.
- [10] X-L Qi, Y-S Wu, and S-C Zhang, “*General theorem relating the bulk topological number to edge states in two-dimensional insulators*”, Phys. Rev. B **74**, 045125.
- [11] P. Delplace, J.B. Marston and A. Venaille, “*Topological Origin of Equatorial Waves*”, Science **358**, 1075 (2017), [arXiv:1702.07583 [cond-mat.mes-hall]]
- [12] D. J. Thouless, M. Kohmoto, M. P. Nightingale, and M. den Nijs, “*Quantized Hall Conductance in a Two-Dimensional Periodic Potential*”, Phys. Rev. Lett. **49**, 405.
- [13] F.D.M. Haldane, “*Model for a Quantum Hall Effect without Landau Levels: Condensed-Matter Realization of the “Parity Anomaly”*”. Phys. Rev. Lett. **61**, 2015 (1988).
- [14] A. Souslov, K. Dasbiswas, M Fruchart, S. Vaikuntanathan, and V. Vitelli, “*Topological waves in fluids with odd viscosity*”, Phys. Rev. Lett. **122**, 128001 (2019), [arXiv:1802.09649 [cond-mat.soft]]
- [15] C. Tauber, P. Delplace, and A. Venaille, “*A bulk-interface correspondence for equatorial waves*”, [arXiv:1812.05488 [physics.flu-dyn]]
- [16] C. Tauber, P. Delplace, and A. Venaille, “*Anomalous bulk-edge correspondence in continuous media*”, Phys. Rev. Res. **2**, 013147 (2020)[arXiv:1902.10050 [cond-mat.mes-hall]]
- [17] A. Venaille and P. Delplace, “*Wave Topology Brought to the Coast*”, Phys. Rev. Res **3**, 043002 (2021), [arXiv:2011.03440v3 [physics.flu-dyn]]
- [18] A. Souslov, B. C. van Zuiden, D. Bartolo and V. Vitelli “*Topological sound in active-liquid metamaterials*”, Nature Physics **13**, 1091 (2017) [arXiv:1610.06873 [cond-mat.soft]]
- [19] M. Perrot, P. Delplace and A. Venaille, “*Topological Transition in Stratified Fluids*”, Nat.Phys. **15**, 781 (2019) [arXiv:1810.03328 [physics.flu-dyn]]
- [20] G.M. Grad, H. Jud, and C. Tauber, “*Topology in shallow-water waves: a violation of bulk-edge correspondence*”, Comm. Math. Phys. **383**, 731 (2021) [arXiv:2001.00439 [math-ph]].
- [21] Y. Liu and Y. W. Sun, “*Topological modes in relativistic hydrodynamics,*” Phys. Rev. D **103** (2021) no.4, 044044 [arXiv:2004.13380 [hep-th]].
- [22] Y. Liu and Y. W. Sun, “*Topological hydrodynamic modes and holography,*” Phys. Rev. D **105** (2022) no.8, 086017 [arXiv:2005.02850 [hep-th]].

- [23] R. Green, J. Armas, J de Boer, and L. Giomi, “*Topological waves in passive and active fluids on curved surfaces: a unified picture*”, [arXiv:2011.12271 [cond-mat.soft]]
- [24] C. Tauber and G.C. Thiang, “*Topology in shallow-water waves: A spectral flow perspective*”, Ann. Henri Poincaré, (2022). [arXiv:2110.04097 [math-ph]]
- [25] S. C. Zhang, T. Hansson and S. Kivelson, “*Effective-Field-Theory Model for the Fractional Quantum Hall Effect*”, Phys. Rev. Lett. **62**, 82 (1989)
- [26] X.G. Wen and Q. Niu, “*Ground-state degeneracy of the fractional quantum Hall states in the presence of a random potential and on high-genus Riemann surfaces*”, Phys. Rev. B **41**, 9377 (1990).
- [27] S. Deser, R. Jackiw and A. P. Polychronakos, “*Clebsch (string) decomposition in $d=3$ field theory*,” Phys. Lett. A **279** (2001), 151 [arXiv:physics/0006056 [physics]].
- [28] R. Jackiw and A. P. Polychronakos, “*Supersymmetric fluid mechanics*,” Phys. Rev. D **62** (2000), 085019 [arXiv:hep-th/0004083 [hep-th]].
- [29] S. Elitzur, G. W. Moore, A. Schwimmer and N. Seiberg, “*Remarks on the Canonical Quantization of the Chern-Simons-Witten Theory*,” Nucl. Phys. B **326** (1989), 108.
- [30] C. T. Hsieh, Y. Tachikawa and K. Yonekura, “*Anomaly Inflow and p -Form Gauge Theories*,” Commun. Math. Phys. **391** (2022) no.2, 495-608 [arXiv:2003.11550 [hep-th]].
- [31] W. Yourgrau and S. Mandelstam, “*Variational Principles in Dynamics and Quantum Theory*”, (1955), 3rd Edition Dover (2003).
- [32] S. Dubovsky, T. Gregoire, A. Nicolis and R. Rattazzi, “*Null energy condition and superluminal propagation*,” JHEP **03** (2006), 025 [arXiv:hep-th/0512260 [hep-th]].
- [33] S. Dubovsky, L. Hui, A. Nicolis and D. T. Son, “*Effective field theory for hydrodynamics: thermodynamics, and the derivative expansion*,” Phys. Rev. D **85** (2012), 085029 [arXiv:1107.0731 [hep-th]].
- [34] R.L. Seliger, “*Variational Principles in Continuum Mechanics*”, Caltech PhD Thesis 1968.
- [35] A. M. Polyakov, “*Compact Gauge Fields and the Infrared Catastrophe*” Phys. Lett. B **59** (1975), 82