

September 23, 2022

# Neutrinoless Double Beta Decay and $\langle \eta \rangle$ Mechanism in the Left-Right Symmetric Model

Takeshi Fukuyama<sup>1</sup> and Toru Sato<sup>2</sup>

*Research Center for Nuclear Physics (RCNP), Osaka University,*

*Ibaraki, Osaka, 567-0047, Japan*

## Abstract

The neutrinoless double beta ( $0\nu\beta\beta$ ) decay is studied in the framework of left-right symmetric model. The coexistence of left and right handed currents induces rather complicated interactions in the mixing of lepton and hadrons, called  $\langle \lambda \rangle$  mechanism and  $\langle \eta \rangle$  mechanism in addition to the conventional effective neutrino mass  $\langle m_\nu \rangle$  mechanism. In this letter we indicate the possible magnification of  $\langle \eta \rangle$  mechanism and importance to survey  $0\nu\beta\beta$  decay of different nuclei for specifying New Physics beyond the Standard Model.

---

<sup>1</sup>E-mail: fukuyama@rcnp.osaka-u.ac.jp

<sup>2</sup>E-mail: tsato@rcnp.osaka-u.ac.jp

# 1 Introduction

Neutrinoless double beta ( $0\nu\beta\beta$ ) decay is one of the key probes for the new physics beyond the Standard Model. There are two conditions to realize  $0\nu\beta\beta$  decay in the context of the Standard Model with light neutrino mass ( $\nu$ SM) [1].

1.  $\nu_e$  should be the same as its anti-particle

$$\nu_e = \bar{\nu}_e \quad (1)$$

and

2. the connecting neutrinos should have the same helicity. The latter condition is satisfied if neutrinos are massive or if the right-handed ( $R$ -handed) current coexists with the left-handed ( $L$ -handed) current. The first case of 2. is described as

$$\langle m_\nu \rangle = \left| \sum_j U_{ej}^2 m_j \right|. \quad (2)$$

Here  $U_{\alpha i}$  (greek (latin) indicates flavour (mass) eigen state) is the Pontecorvo-Maki-Nakagawa-Sakata (PMNS) mixing matrix [2, 3] in  $L$ -handed current. Normal Hierarchy (NH) indicates

$$m_1 < m_2 < m_3 \quad (3)$$

and the Inverted Hierarchy (IH) does

$$m_3 < m_1 < m_2. \quad (4)$$

The central values and relative uncertainties of PMNS mixing angles and the mass square differences are [4]

$$\begin{aligned} \theta_{12} &= 0.5903, 2.3\%, \quad \theta_{23} = 0.866, 4.1\%(\text{NH}), 0.869, 4.0\%(\text{IH}), \\ \theta_{13} &= 0.150, 1.5\%(\text{NH}), 0.151, 1.5\%(\text{IH}), \\ \Delta m_{21}^2 &= 7.39 \times 10^{-5} \text{eV}^2, 2.8\%, \quad \Delta m_{32}^2 = 2.451 \times 10^{-3} \text{eV}^2, 1.3\%(\text{NH}), -2.512 \times 10^{-3} \text{eV}^2, 1.3\%(\text{IH}) \end{aligned} \quad (5)$$

and the PMNS matrix elements have the values

$$U_{11} = c_{12}c_{13}, \quad U_{12} = s_{12}c_{13}, \quad U_{13} = s_{13}e^{-i\delta} \quad (6)$$

up to the Majorana phases  $\text{diag}(e^{i\alpha}, e^{i\beta}, 1)$ . Here  $c_{ij} = \cos\theta_{ij}$  and  $s_{ij} = \sin\theta_{ij}$  as usual. Substituting these values, we obtain

$$|U_{11}|^2/|U_{13}|^2 \approx 30. \quad (7)$$

Then, IH case enhances  $\langle m_\nu \rangle$  relative to NH case. Though the final answer to the hierarchy problem is given by the observation, theoretical predictions have been given by many models.

One of the typical models is due to the predictive minimal SO(10) model [5]. Based on the SO(10) model, we fitted low energy spectra of all quark lepton masses and the CKM and the PMNS mixing angles and phases. Our results prefer the NH manifestly to the IH:  $\chi^2 \leq 1$  for the NH and  $\chi^2 > 200$  for the IH. Here the  $\chi^2$  function is defined by

$$\chi^2 = \sum_i \frac{(\chi_i - \hat{\chi}_i)^2}{\sigma_i^2}, \quad (8)$$

where  $\chi_i$ ,  $\hat{\chi}_i$ , and  $\sigma_i$  are fitted parameters, their observed values, and their standard deviations of errors, respectively (see [5] for the details). Using the fitted data, the effective neutrino mass is also predicted as

$$\langle m_\nu \rangle \approx 1 \text{ meV}. \quad (9)$$

On the other hand, the recent  $0\nu\beta\beta$  experiment in the KamLAND-Zen provides the most stringent upper limit on it, 61 – 165 meV [6, 7, 8]. So far we have discussed in the framework of  $\nu$ SM, and the half life  $T_{1/2}$  of  $0\nu\beta\beta$  decay is

$$\frac{1}{T_{1/2}} = G_{0\nu} |M_{0\nu}|^2 \left( \frac{\langle m_\nu \rangle}{m_e} \right)^2, \quad (10)$$

where  $G_{0\nu}$  is the phase-space integral and  $M_{0\nu}$  is the nuclear matrix element (NME). Unfortunately, the ambiguities in NME are still large and they are different by a factor two in each nucleus in different nuclear models. If we consider new physics beyond the  $\nu$ SM (BSM physics), we have many options furthermore. One of the most reliable ingredients is the R-handed current in L-R symmetric model [9, 10],

$$H_W = \frac{G_F \cos \theta_c}{\sqrt{2}} \left[ j_L^\mu \tilde{J}_{L\mu}^\dagger + j_R^\mu \tilde{J}_{R\mu}^\dagger \right] + H.c. \quad (11)$$

Here  $j_\mu$  ( $J_\mu$ ) indicates leptonic (hadronic) current, and the  $L$  and  $R$ -handed leptonic currents,  $j_{L\mu}$  and  $j_{R\mu}$ , are given by

$$j_{L\alpha} = \sum_{l=e,\mu,\tau} \overline{l(x)} \gamma_\alpha (1 - \gamma_5) \nu_{lL}(x) \equiv \sum \overline{l(x)} \gamma_\alpha 2P_L \nu_{lL}(x), \quad (12)$$

$$j_{R\alpha} = \sum_{l=e,\mu,\tau} \overline{l(x)} \gamma_\alpha (1 + \gamma_5) N_{lR}(x) \equiv \sum \overline{l(x)} \gamma_\alpha 2P_R N_{lR}(x), \quad (13)$$

where  $\nu_{lL}$  ( $N_{lR}$ ) are  $L$ -handed ( $R$ -handed) weak eigenstates of the neutrinos, and

$$\tilde{J}_L^\mu(\mathbf{x}) = J_L^\mu(\mathbf{x}) + \kappa J_R^\mu(\mathbf{x}), \quad (14)$$

$$\tilde{J}_R^\mu(\mathbf{x}) = \eta J_L^\mu(\mathbf{x}) + \lambda J_R^\mu(\mathbf{x}). \quad (15)$$




Figure 1: Diagrams of  $0\nu\beta\beta$  decay. (a), (b), and (c) are  $\langle m_\nu \rangle$ ,  $\langle \lambda \rangle$ , and  $\langle \eta \rangle$ -mechanisms, respectively.

The diagrams of  $0\nu\beta\beta$  decay in the  $L - R$  symmetric model are depicted in Fig.1 and its amplitude in closure approximation [15] is given as

$$R_{0\nu} = 4\sqrt{\frac{1}{2}} \left( \frac{G \cos \theta_c}{\sqrt{2}} \right)^2 \sum_i \sum_{\alpha, \beta} \int d\mathbf{x} d\mathbf{y} \int \frac{d\mathbf{k}}{(2\pi)^3} e^{i\mathbf{k}\cdot(\mathbf{y}-\mathbf{x})} H^{\nu\mu} L_{\nu\mu}, \quad (16)$$

where the lepton tensor  $L^{\nu\mu}$  is

$$L_{\nu\mu} = \bar{e}_{p_2, s'_2}(\mathbf{y}) \gamma_\nu P_\beta \frac{1}{2\omega} \left[ \frac{\omega\gamma^0 - \mathbf{k}\cdot\boldsymbol{\gamma} + m_i}{\omega + A_1} + \frac{-\omega\gamma^0 - \mathbf{k}\cdot\boldsymbol{\gamma} + m_i}{\omega + A_2} \right] P_\alpha \gamma_\mu e_{p_1, s'_1}^c(\mathbf{x}). \quad (17)$$

Here  $e_{p_i, s'_i}(\mathbf{x})$  are electron wave functions with the energy  $e_i$ , and the mixing matrices  $U$ ,  $V^*$  are omitted for simplicity. The energy denominator is given by  $A_i = e_i + \langle E_n \rangle - E_i$  and  $E_f + e_1 + e_2 = E_i$ . Here  $E_{i/f}$  and  $\langle E_n \rangle$  are energy of the initial/final nuclear state and the average energy of the intermediate nuclear state, respectively.

The nuclear tensor  $H^{\nu\mu}$  is given by the matrix element of the nuclear weak current as

$$H^{\nu\mu} = \langle F | \tilde{J}_{\beta i}^{\nu+}(\mathbf{y}) \tilde{J}_{\alpha i}^{\mu+}(\mathbf{x}) | I \rangle, \quad (18)$$

where  $\tilde{J}_{L,R}^\mu$  are given in (14) and (15).

The neutrino propagator becomes,

$$P_\alpha (\pm\omega\gamma^0 - \mathbf{k}\cdot\boldsymbol{\gamma} + m_i) P_\beta = \begin{cases} m_i P_\alpha & (\alpha = \beta) \\ (\pm\omega\gamma^0 - \mathbf{k}\cdot\boldsymbol{\gamma}) P_\beta & (\alpha \neq \beta) \end{cases}. \quad (19)$$

In the presence of the R-handed current, we have  $(\pm\omega\gamma^0 - \mathbf{k}\cdot\boldsymbol{\gamma})P_\beta$  in addition to (2). The spatial momentum exchanged between nucleon by neutrino is significantly larger than neutrino mass term,  $|\mathbf{k}| \approx 100\text{MeV} \gg E_n - E_i, m_i$ , which gives a significant effect to the decay rate.

This mechanism gives interesting interplays between particle physics and nuclear physics, whose explanation is the main theme of this letter.

The half life  $T_{1/2}$  in this case [1, 16, 17, 18, 19, 20] is given as

$$\begin{aligned} \frac{1}{T_{1/2}} = & C_{mm}^{(0)} \left( \frac{\langle m_\nu \rangle}{m_e} \right)^2 + C_{m\lambda}^{(0)} \frac{\langle m_\nu \rangle}{m_e} \langle \lambda \rangle \cos \psi + C_{m\eta}^{(0)} \frac{\langle m_\nu \rangle}{m_e} \langle \eta \rangle \cos \psi \\ & + C_{\lambda\lambda}^{(0)} \langle \lambda \rangle^2 + C_{\eta\eta}^{(0)} \langle \eta \rangle^2 + C_{\lambda\eta}^{(0)} \langle \lambda \rangle \langle \eta \rangle. \end{aligned} \quad (20)$$

Here  $C_{ab}^{(0)}$  include NME and phase space integral. The remains include BSM physics. The effective couplings  $\langle \eta \rangle$  and  $\langle \lambda \rangle$  are given as

$$\langle \lambda \rangle = \lambda \left| \sum_j 'U_{ej} V_{ej}^* \right|, \quad \langle \eta \rangle = \eta \left| \sum_j 'U_{ej} V_{ej}^* \right|. \quad (21)$$

The meanings of the mixing matrices  $U$ ,  $V$  are given in the next section.  $\psi$  is the relative phase between  $\langle m_\nu \rangle$  and  $\langle \lambda \rangle$  and  $\langle \eta \rangle$ ,

$$\psi = \arg \left[ \left( \sum_j 'm_j U_{ej}^2 \right) \left( \sum_j 'U_{ej} V_{ej}^* \right)^* \right], \quad (22)$$

where  $\sum'$  indicates the summation over only the light neutrinos. However,  $U$  and  $V$  are independent and we set  $\psi = 0$  hereafter. In the  $L - R$  symmetric model, the details of  $\lambda$  and  $\eta$  are given by (31) and (32). In the next section, we concentrate on the structure of lepton part where BSM physics resides.

## 2 Right-handed weak current

We consider  $L - R$  symmetric model [10] in this section, where Higgs sectors are composed of (3,1,2), (1,3,2) triplets ( $\Delta_L$ ,  $\Delta_R$ , respectively) and bi-doublet (2,2,0) ( $\Phi$ ) under  $SU(2)_L \otimes SU(2)_R \otimes U(1)_{B-L}$ . They have the vevs as

$$\langle \Phi \rangle_0 = \left\langle \begin{pmatrix} \phi_1^0 & \phi_2^+ \\ \phi_1^- & \phi_2^0 \end{pmatrix} \right\rangle_0 = \begin{pmatrix} v_u & 0 \\ 0 & v_d \end{pmatrix} \quad (23)$$

and

$$\langle \Delta_{L,R} \rangle_0 = \left\langle \begin{pmatrix} \frac{\delta^+}{\sqrt{2}} & \Delta^{++} \\ \Delta^0 & -\frac{\delta^+}{\sqrt{2}} \end{pmatrix}_{L,R} \right\rangle_0 = \begin{pmatrix} 0 & 0 \\ v_{L,R} & 0 \end{pmatrix}. \quad (24)$$

The neutrino mass matrix is [11, 12, 13, 14], then

$$M_\nu = \begin{pmatrix} M_L & M_D \\ M_D^T & M_R \end{pmatrix} \approx \begin{pmatrix} 0 & M_D \\ M_D^T & M_R \end{pmatrix}. \quad (25)$$

Thus we have the extended Fermi couplings (11). In (12) and (13),  $\nu_{lL}(N_{lR})$  are  $L$ -handed ( $R$ -handed) weak eigenstates of the neutrinos. Using  $3 \times 3$  blocks  $U, V, X, Y$ , the mass eigenstates  $\nu', N'$  are given as

$$\begin{pmatrix} \nu \\ (N_R)^c \end{pmatrix}_L = \begin{pmatrix} U & X \\ V & Y \end{pmatrix} \begin{pmatrix} \nu' \\ N' \end{pmatrix}_L \equiv \mathcal{U} \begin{pmatrix} \nu' \\ N' \end{pmatrix}_L. \quad (26)$$

That is,

$$(\nu_L)_\alpha = U_{\alpha i} \nu'_i + X_{\alpha I} N'_I, \quad (N_R)_\alpha^c = V_{\alpha i} \nu'_i + Y_{\alpha I} N'_I, \quad (27)$$

where  $\alpha (i)$  are the flavour (mass) eigen states.

The constants  $\lambda$  and  $\eta$  in (11) are related to the mass eigenvalues of the weak bosons in the  $L$  and  $R$ - handed gauge sectors ( $W_L, W_R$ ) as follows:

$$W_L = W_1 \cos \zeta + W_2 \sin \zeta, \quad (28)$$

$$W_R = -W_1 \sin \zeta + W_2 \cos \zeta, \quad (29)$$

$$\frac{G_F}{\sqrt{2}} = \frac{g^2}{8} \cos^2 \zeta \frac{M_{W_1}^2 \tan^2 \zeta + M_{W_2}^2}{M_{W_1}^2 M_{W_2}^2}, \quad (30)$$

$$\lambda \equiv \frac{M_{W_1}^2 + M_{W_2}^2 \tan^2 \zeta}{M_{W_1}^2 \tan^2 \zeta + M_{W_2}^2}, \quad (31)$$

$$\eta \equiv -\frac{(M_{W_2}^2 - M_{W_1}^2) \tan \zeta}{M_{W_1}^2 \tan^2 \zeta + M_{W_2}^2}. \quad (32)$$

Here  $M_{W_1}$  and  $M_{W_2}$  are the masses of the mass eigenstates  $W_1$  and  $W_2$ , respectively, and  $\zeta$  is the mixing angle which relates the mass eigenstates and the gauge eigenstates. We are considering  $L - R$  symmetric model. The gauge boson mass is generated from (23) and (24) as [21]

$$M_W^2 = \begin{pmatrix} \frac{1}{2}g^2(v_u^2 + v_d^2 + 2v_L^2) & g^2 v_u v_d \\ g^2 v_u v_d & \frac{1}{2}g^2(v_u^2 + v_d^2 + 2v_R^2) \end{pmatrix} \quad (33)$$

and the mixing angle  $\zeta$  is

$$\tan 2\zeta = \frac{2v_u v_d}{v_R^2 - v_L^2} \approx \frac{2v_u v_d}{v_R^2} = 2\xi \left( \frac{M_{WL}}{M_{WR}} \right)^2 \quad (34)$$

with

$$v_u^2 + v_d^2 = v_{ew}^2, \quad (35)$$

$$\xi = v_d/v_u = 1/\tan\beta \quad (36)$$

and

$$M_{W2} = \sqrt{2}g_R v_R \geq 5\text{TeV} \quad (37)$$

[22]. In the  $L - R$  symmetric model, we set  $g_L = g_R$ , which indicates further unification of at least rank five GUT, including SU(3) color.  $\tan\beta$  is constrained from the Yukawa coupling is renormalizable up to the GUT scale,

$$1 \leq \tan\beta \leq 60. \quad (38)$$

Furthermore, in this case, large  $\tan\beta$  induces too rapid proton decay since the proton life-time is proportional to  $1/\tan\beta^2$  and  $\tan\beta$  is limited around 10 [5]. Reflecting these relatively low mass constraint, we will discuss on the low energy seesaw mechanism soon after.

Corresponding to Figure 1, we will consider  $0\nu\beta\beta$  decay in this scheme:

- $W_L - W_L$  diagram

$$m_{eff}^{LL} = \sum_{i=1}^3 U_{ei}^2 m_i + k^2 X_{ei}^2 \frac{M_I}{k^2 - M_I^2}. \quad (39)$$

Here and in the subsequent discussions in this section, we write the subdominant terms in addition to (21), illustrating the seesaw structure.

- $W_R - W_R$  diagram

$$m_{eff}^{RR} = \sum k^2 Y_{ei}^2 \frac{M_I}{k^2 - M_I^2} \frac{g_R^4}{g_L^4} \frac{M_{WL}^4}{M_{WR}^4}, \quad (40)$$

with  $g_L = g_R$ , which was suppressed compared with the others due to  $\left(\frac{M_{WL}}{M_{WR}}\right)^4$ .

- $W_L - W_R$  diagram; the neutrino mixing ( $\lambda$ ) and  $W_L - W_R$  mixing ( $\eta$ )

$$\langle \lambda \rangle = \left( U_{ei} V_{ei}^* + X_{ei} Y_{ei}^* \frac{k^2}{k^2 - M_I^2} \right) \frac{M_{WL}^2}{M_{WR}^2}, \quad (41)$$

$$\langle \eta \rangle = \left( U_{ei} V_{ei}^* + X_{ei} Y_{ei}^* \frac{k^2}{k^2 - M_I^2} \right) (-\tan\zeta). \quad (42)$$

Here the first terms  $U_{ei} V_{ei}^*$  dominate, and let us estimate the magnitude of  $V_{ei}$ . The naive type I seesaw (25) gives tiny value for the above quantities. We are interested in TeV scale

seesaw and consider the inverse seesaw mechanism [23] hereafter. Its  $9 \times 9$  mass matrix is given by

$$M_\nu = \begin{pmatrix} 0 & M_D^T & 0 \\ M_D & 0 & M^T \\ 0 & M & \mu \end{pmatrix} \equiv \begin{pmatrix} 0_{3 \times 3} & \mathcal{M}_{D3 \times 6}^T \\ \mathcal{M}_{D6 \times 3} & \mathcal{M}_{R6 \times 6} \end{pmatrix} \quad (43)$$

Their mass scales are  $\mu \approx O(1)keV$ ,  $M_D \approx O(100)$  GeV,  $M \approx O(1)$  TeV. This matrix is diagonalized by  $9 \times 9$  mixing matrix  $\mathcal{U}$  [24, 25]

$$\mathcal{U} \approx \quad (44)$$

$$\begin{pmatrix} 1 - \frac{1}{2}\mathcal{M}_D^\dagger[\mathcal{M}_R(\mathcal{M}_R)^\dagger]^{-1}\mathcal{M}_D & \mathcal{M}_D^\dagger(\mathcal{M}_R^\dagger)^{-1} \\ -\mathcal{M}_R^{-1}\mathcal{M}_D & 1 - \frac{1}{2}\mathcal{M}_R^{-1}\mathcal{M}_D\mathcal{M}_D^\dagger(\mathcal{M}_R^\dagger)^{-1} \end{pmatrix}$$

as

$$\mathcal{U}^T M_\nu \mathcal{U} = \begin{pmatrix} m_{light} & 0_{3 \times 6} \\ 0_{6 \times 3} & M_{heavy} \ 6 \times 6 \end{pmatrix}. \quad (45)$$

Here

$$m_{light} = M_D^T M^{-1} \mu (M^T)^{-1} M_D \quad (46)$$

and

$$M_{heavy} = \mathcal{M}_R. \quad (47)$$

Thus  $\mathcal{U}$  of (26) in type I seesaw is modified in the inverse seesaw as

$$\mathcal{U} = \begin{pmatrix} U & X \\ V & Y \\ W & Z \end{pmatrix}, \quad (48)$$

where  $\{U, V, W\}$  and  $\{X, Y, Z\}$  are  $3 \times 3$  and  $3 \times 6$  matrices, respectively. All deviation from unitarity is determined by

$$\zeta = \mathcal{M}_R^{-1} \mathcal{M}_D. \quad (49)$$

It goes from (44) and (48),

$$\mathcal{M}_R^{-1} \mathcal{M}_D = \begin{pmatrix} 0 & M \\ M & \mu \end{pmatrix}^{-1} \begin{pmatrix} m_D \\ 0 \end{pmatrix} = -\frac{1}{M^2} \begin{pmatrix} -\mu m_D \\ M m_D \end{pmatrix} \quad (50)$$



Then

$$V \approx O\left(\frac{\mu m_D}{M^2}\right) \quad (51)$$

and seems to be tiny. However, this is too naive estimation. This is because the unitary matrix  $\mathcal{U}$ , as is easily checked, is unitary up to the order of  $\zeta^2$  in the context of one generation, and the estimation (51) is not reliable. Also there appears ambiguities if we consider the lepton generation. The order of the magnitude of  $V$  should be estimated from the observations.

Thus we obtain [26]

$$|\langle \lambda \rangle| \leq 5.7 \times 10^{-7} \quad (52)$$

and  $|\langle \eta \rangle|$  is suppressed by  $1/\tan \beta$  in comparison with  $|\langle \lambda \rangle|$  by virtue of eqs. (34) and (42),

$$|\langle \eta \rangle| \leq 5.7 \times 10^{-7} / \tan \beta. \quad (53)$$

### 3 Nuclear matrix elements and role of $\langle \eta \rangle$ mechanism

Possibility of non-smallness of  $\eta$  discussed in the previous section and the larger sensitivity of  $\eta$  mechanism to the decay rate than  $\lambda$  mechanism [27, 28, 29] suggests an interesting possibility to detect  $R$ -handed current from the  $0\nu\beta\beta$  decay. In this section, we revisit the mechanism that enhance the  $\eta$  mechanism but not the  $\lambda$  mechanism. A simplified derivation of the relevant nuclear operator would be useful. We then estimate the allowed region of  $\langle \eta \rangle$ .

The transition amplitude  $R_{0\nu}$  (16) can be written in terms of the nuclear transition density of  $\mathcal{O}(\mathbf{x}, \mathbf{y})$  as

$$R_{0\nu} = 4\sqrt{\frac{1}{2}} \left( \frac{G \cos \theta_c}{\sqrt{2}} \right)^2 \sum_i \sum_{\alpha, \beta} \int d\mathbf{x} d\mathbf{y} \int \frac{d\mathbf{k}}{(2\pi)^3} e^{i\mathbf{k}\cdot(\mathbf{y}-\mathbf{x})} \frac{1}{2\omega} \left[ \frac{1}{\omega + A_1} + \frac{1}{\omega + A_2} \right] \bar{e}_{p_2, s'_2}(\mathbf{y}) \mathcal{O}(\mathbf{x}, \mathbf{y}) e_{p_1, s'_1}^c(\mathbf{x}), \quad (54)$$

where  $\mathcal{O}(\mathbf{x}, \mathbf{y})$  is the matrix element of the hadronic current. We kept the  $\mathbf{k}$  of the neutrino propagator for the  $\eta$  and  $\lambda$  term.

$$\mathcal{O}(\mathbf{x}, \mathbf{y}) = -\langle F | \hat{J}_R^\dagger(\mathbf{y}) \mathbf{k} \cdot \gamma P_L \hat{J}_L^\dagger(\mathbf{x}) + \hat{J}_L^\dagger(\mathbf{y}) \mathbf{k} \cdot \gamma P_R \hat{J}_R^\dagger(\mathbf{x}) | I \rangle. \quad (55)$$

Here the Dirac matrices are for the electron spinor. Either emitted electrons or nuclear currents have to compensate the p-wave nature of neutrino propagator for the  $0^+ - 0^+$  transition. An important contribution comes from the s-wave electron and momentum dependent nuclear current due to the nuclear magnetization current.  $\mathcal{O}(\mathbf{x}, \mathbf{y})$  can be evaluated using the hadronic current of Eqs. (14) and (15). The interference term between the SM hadronic  $L$ -handed current and the BSM  $L$ -handed current  $\eta J_L^\mu$  and the  $R$ -handed current  $\lambda J_R^\mu$  contribute, while the  $\kappa J_R^\mu$  term can

be neglected. Using  $J_{L/R}^\mu = V^\mu \pm A^\mu$ , we obtain  $\mathcal{O}(\mathbf{x}, \mathbf{y})$  for the interference term between vector including magnetization current and axial vector current including Gamow-Teller operator as

$$\begin{aligned} \mathcal{O}(\mathbf{x}, \mathbf{y}) &= \langle F | \langle \lambda \rangle (\mathcal{V}^\dagger(\mathbf{y})\mathbf{k} \cdot \gamma \mathcal{A}^\dagger(\mathbf{x}) - \mathcal{A}^\dagger(\mathbf{y})\mathbf{k} \cdot \gamma \mathcal{V}^\dagger(\mathbf{x}))\gamma_5 \\ &\quad + \langle \eta \rangle (\mathcal{V}^\dagger(\mathbf{y})\mathbf{k} \cdot \gamma \mathcal{A}^\dagger(\mathbf{x}) + \mathcal{A}^\dagger(\mathbf{y})\mathbf{k} \cdot \gamma \mathcal{V}^\dagger(\mathbf{x})) | I \rangle. \end{aligned} \quad (56)$$

We focus on the spatial component of the currents and assume the s-wave electron wave function can be approximated by constant. Using  $\mathbf{V}(\mathbf{x}) = \nabla \times \boldsymbol{\mu}(\mathbf{x})$ , we obtain

$$\begin{aligned} \mathcal{O}(\mathbf{x}, \mathbf{y}) &= \langle F | \langle \lambda \rangle (\mathbf{k} \times \boldsymbol{\mu}(\mathbf{y}) \cdot \mathbf{k} \times \mathbf{A}(\mathbf{x}) - \mathbf{k} \times \mathbf{A}(\mathbf{y}) \cdot \mathbf{k} \times \boldsymbol{\mu}(\mathbf{x}))(-\gamma_0) \\ &\quad + \langle \eta \rangle (\mathbf{k} \times \boldsymbol{\mu}(\mathbf{y}) \cdot \mathbf{k} \times \mathbf{A}(\mathbf{x}) + \mathbf{k} \times \mathbf{A}(\mathbf{y}) \cdot \mathbf{k} \times \boldsymbol{\mu}(\mathbf{x}))(\gamma_5 \gamma_0) | I \rangle. \end{aligned} \quad (57)$$

In the non-relativistic and impulse approximation of nuclear current,  $\boldsymbol{\mu}(\mathbf{x})$  and  $\mathbf{A}(\mathbf{x})$  are given by the same operator  $\sim \tau^+ \boldsymbol{\sigma}$  as,

$$\mathbf{A}(\mathbf{x}) = \sum_i^A g_A(k^2) \tau_i^+ \boldsymbol{\sigma}_i \delta(\mathbf{x} - \mathbf{r}_i), \quad (58)$$

$$\boldsymbol{\mu}(\mathbf{x}) = \sum_i^A \frac{g_V(k^2) + g_M(k^2)}{2M} \tau_i^+ \boldsymbol{\sigma}_i \delta(\mathbf{x} - \mathbf{r}_i). \quad (59)$$

Here  $g_A(0) = 1.27$ ,  $g_V(0) + g_M(0) = 4.706$  and  $M$  is mass of nucleon. Within this approximation, the  $\langle \lambda \rangle$  term vanishes and only the  $\langle \eta \rangle$  term remains. Therefore the  $0\nu\beta\beta$  decay can be sensitive probe for the  $\langle \eta \rangle$  mechanism of  $R$ -handed current. Assuming the s-wave electron wave function can be approximated by constant, the amplitude  $R_{0\nu}$  is given by the NME of two-body operator  $M_{ij}$  as

$$R_{0\nu} = 4\sqrt{\frac{1}{2}} \left( \frac{G \cos \theta_c}{\sqrt{2}} \right)^2 \bar{e}_{p_2, s'_2}(\mathbf{0}) \gamma_5 \gamma_0 e_{p_1, s'_1}^c(\mathbf{0}) \langle F | \sum_{i>j=1}^A M_{ij} | I \rangle, \quad (60)$$

with

$$M_{ij} = \langle \eta \rangle \tau_i^+ \tau_j^+ \int \frac{d\mathbf{k}}{(2\pi)^3} \frac{e^{i\mathbf{k} \cdot (\mathbf{r}_i - \mathbf{r}_j)}}{\omega} \left[ \frac{1}{\omega + A_1} + \frac{1}{\omega + A_2} \right] g_A(k^2) \frac{g_V(k^2) + g_M(k^2)}{2M} \mathbf{k} \times \boldsymbol{\sigma}_i \cdot \mathbf{k} \times \boldsymbol{\sigma}_j. \quad (61)$$

The neutrino-exchange two-body operator  $M_{ij}$ , whose spin and momentum structure is similar to the  $\rho$  meson exchange nuclear potential, includes spin-spin interaction and the tensor interaction.




Figure 2: Allowed region of  $\langle \eta \rangle$  and  $\langle m_\nu \rangle$  for  $^{136}\text{Xe}$ . a,b,c are evaluated using  $C$ 's of Refs. [35],[36] and [19] (model without p-n pairing), respectively.

The spin-spin interaction includes short range the delta function for point-like nucleon. The short distance nature of the operator is well recognized and has been examined in detail. It is essential to include both the finite size of weak nucleon current and the short range correlation. The form factors of axial vector and weak magnetism of nucleon are parameterized in a dipole form as  $g_A(k^2) = g_A/(1 + \mathbf{k}^2/M_A^2)^2$  and  $g_V(k^2) + g_M(k^2) \sim (g_V + g_M)/(1 + \mathbf{k}^2/M_V^2)^2$ . The short range correlation, which is not taken into account in the model nuclear wave function, is taken into account by introducing short range correlation (SRC) function [30, 31, 32, 33]

$$F(r) = 1 - ce^{-ar^2}(1 - br^2) \quad (62)$$

with  $r = |\mathbf{r}_j - \mathbf{r}_i|$ . This correlation function vanishes at  $\lim r = 0$  when  $c = 1$ . The suppression rates relative to those without the SRC are not affirmative, depending on nuclear models and various SRC, from 5% to 30 – 40%. See the most recent result, Fig.10 of [34].

The sensitivity of the  $0\nu\beta\beta$  decay rate to  $\langle \eta \rangle$  has been studied for example in Ref. [27]. Using the typical value of the ratio  $C_{\eta\eta}^{(0)}/C_{mm}^{(0)} \sim 10^4$  to  $10^5$ , the decay rate for  $\langle m_\nu \rangle \sim 100$  meV corresponds to  $\langle \eta \rangle \sim 10^{-9}$ , which already constrains the limit given in Eq. (53). Actually, the decay rate is quadratic function of  $\langle \eta \rangle$  and  $\langle m_\nu \rangle$  for  $\langle \lambda \rangle = 0$ . Using current lower limit of  $T_{1/2} > 10.7 \times 10^{25}$  years from KamLAND-Zen [6] and  $C$ 's given in Table 27 of Ref. [20],  $\langle \eta \rangle$  and  $\langle m_\nu \rangle$  constrained from the data are inside ellipses shown in Fig. 2.

The decay rate of a single nuclear species is not possible to reveal the mechanism of  $0\nu\beta\beta$  decay including BSM physics. However, difference between the space and spin structure of the the effective nuclear operators for  $\langle m_\nu \rangle$  and  $\langle \eta \rangle$  mechanisms has a potential to generate atom (A)-dependence of the decay rate [28, 29]. A-dependence of the decay rate can be quantified by the normalized decay rate with respect to the reference  $0\nu\beta\beta$  process for two extreme cases

	<sup>48</sup> Ca	<sup>76</sup> Ge	<sup>82</sup> Se	<sup>96</sup> Zr	<sup>100</sup> Mo	<sup>116</sup> Cs	<sup>128</sup> Te	<sup>130</sup> Te
$R_A^{m\nu}$	0.75	0.51	1.2	3.0	0.47	0.39	0.095	2.1
$R_A^\eta$	0.082	0.40	0.19	0.83	0.36	0.064	0.10	2.0
$R_A = R_A^\eta/R_A^{m\nu}$	0.11	0.77	0.15	0.28	0.76	0.16	1.1	0.94

Table 1: Ratio of decay rate  $R_A^\alpha$  evaluated using  $C$ 's of [19].

$\langle m_\nu \rangle$  alone ( $\alpha = m_\nu$ ) and  $\langle \eta \rangle$  alone ( $\alpha = \eta$ ) as

$$R_A^\alpha = \frac{T_{1/2}^\alpha(A_{\text{reference}})}{T_{1/2}^\alpha(A)}, \quad (63)$$

and deviation of the ratio  $R_A = R_A^\eta/R_A^{m\nu}$  from one indicates an ability of the process to find BSM physics. Though precise A-dependence of  $R_A$  is not yet established theoretically, we estimated  $R_A$  using  $C$ 's from the Tables of [20]. In table 1,  $R_A^\alpha$  and  $R_A$  are calculated for [19](model without p-n pairing) with <sup>136</sup>Xe as  $A_{\text{reference}}$ . In this model,  $R_A$  is appreciably small for <sup>48</sup>Ca, <sup>82</sup>Se, <sup>96</sup>Zr, <sup>116</sup>Cs. The result suggests that A-dependence of the decay rate can be a key to disentangle the mechanism of  $0\nu\beta\beta$  decay and to discover the BSM signal of  $R$ -handed current.

## 4 Conclusion

We have studied  $0\nu\beta\beta$  decay in the presence of  $R$ -handed current. We have tried to clarify the arguments of hadronic side and lepton's BSM physical one. As is well known, if neutrino masses obey IH,  $\langle m_\nu \rangle$  mechanism works around 50 meV marginal to the present and near future experiments. From NME,  $\lambda$  mechanism is suppressed and  $\eta$  mechanism can dominate the decay rate even around the present or near future experimental limits. This is the case if the neutrino masses belong to NH which is much more probable than IH from theoretical reasons. Even if we get the non-null result in  $0\nu\beta\beta$  decay in a single species, though it is the great achievement, we can not limit BSM physics. It is very important to survey this process in many different nuclei to specify BSM physics.

## Acknowledgments

We would express our sincere thanks to Dr. H. Ejiri for his encouragements. This work is supported by JSPS KAKENHI Grant Numbers JP22H01237 (T.F) and JP19H05104 (TS).

## References

- [1] M. Doi, T. Kotani and E. Takasugi, Prog. Theor. Phys. Supplement **83** 1 (1985).
- [2] B. Pontecorvo, Sov. Phys. JETP, **7**, 172 (1958)
- [3] Z. Maki, M. Nakagawa, and S. Sakata, Prog. Theor. Phys. **28**, 870 (1962).
- [4] B. Abi et al. (DUNE Collaboration), Eur. Phys. J. **C80**, 978 (2020).
- [5] T. Fukuyama, K. Ichikawa, and Y. Mimura, Phys. Rev. **D94**, 075018 (2016).
- [6] A. Gando et al. (KamLAND-Zen Collaboration), Phys.Rev.Lett. **117**, 082503 (2016). [Addendum: Phys. Rev. Lett. **117**,109903 (2016)].
- [7] M. Agostini et al. (GERDA Collaboration), Science, **365**, 1445 (2019).
- [8] D. Adams et al. (CUORE Collaboration), Phys.Rev.Lett. **124**, 122501 (2020).
- [9] J.C. Pati and A. Salam, Phys.Rev. **D10**, 275 (1974).
- [10] R.N. Mohapatra and G. Senjanovic, Phys. Rev. Lett. **44**, 912 (1980).
- [11] P. Minkowski, Phys. Lett. **B67**, 421 (1977).
- [12] T. Yanagida, Conf. Proc. **C7902131**, 95 (1979).
- [13] M. Gell-Mann, P. Ramond, and R. Slansky, Conf.Proc. **C790927**, 315 (1979).
- [14] R. N. Mohapatra and G. Senjanovic, Phys. Rev. Lett. **44**, 912 (1980).
- [15] T. Tomoda, Rep.Prog.Phys. **54**, 53 (1991).
- [16] F. Simkovic, J. Vergados, and A. Faessler, Phys. Rev. **D82**, 113015 (2010).
- [17] J. Engel and J. Menendez, Rep. Prog. Phys. **80** 046301 (2017).
- [18] G. Pantis, A. Faessler, W.A. Kaminski, and J.D. Vergados, J. Phys. G: Nucl. Phys. **18**, 605 (1992).
- [19] G. Pantis, F.Simkovic, J.D. Vergados, and A. Faessler, Phys. Rev. **C53**, 695 (1996).
- [20] J. Suhonen and O. Civitarese, Phys. Rep. **300**, 123 (1998).
- [21] Y. Zhang, H. An, X. Ji, and R.N. Mohapatra, Nucl. Phys. **B802**, 247 (2008).
- [22] A.M. Sirunyan et al. (CMS Collaboration), JHEP **05**, 148 (2018).
- [23] R.N. Mohapatra and J.W.F. Valle, Phys. Rev. **D34**, 1642 (1986).

- [24] H. Hettmansperger, M. Lindner, and W. Rodejohann, *JHEP* **1104**, 123 (2011).
- [25] J.P. Pinheiro, C.A.deS. Pirez, F.S. Queiroz, and Y.S. Villamizar, *Phys. Lett.* **B823**, 136764 (2021).
- [26] J. Barry and W. Rodejohann, *JHEP* **09**, 153 (2013).
- [27] T. Tomoda and A. Faessler, *Phys. Lett.* **B199**, 475 (1987).
- [28] M. Horoi and A. Neacsu, *Phys. Rev.* **C98**, 035502 (2018).
- [29] F. Ahmed and M. Horoi, *Phys. Rev.* **C101**, 035504 (2020).
- [30] G.A. Miller and J.E. Spencer, *Ann. Physics*, **100**, 562 (1976).
- [31] F. Simkovich, A. Faessler, H. Muther, V. Rodin, and M. Stauf, *Phys.Rev.* **C79**, 055501 (2009).
- [32] H.F. Wu, H.Q. Song, T.T.S. Kuo, W.K. Cheng, and D. Strominger, *Phys. Lett.* **162B**, 227 (1988).
- [33] L.S. Song, J.M. Yao, P. Ring, and J. Meng, *Phys. Rev.* **C95**, 024305 (2017).
- [34] J.M. Yao, J. Meng, Y.Y. Niu, and P. Ring, *Prog. Part. Nucl. Phys*, **126**, 103965 (2022).
- [35] K. Muto, E. Bender, H. V. Klapdor, *Z. Phys.* **A334**, 187 (1989).
- [36] J. Suhonen, S. B. Khadkikar, A. Faessler, *Nucl. Phys.* **A535**, 509 (1991).
- [37] J. Retamosa, E. Caurier, and F. Nowacki, *Phys.Rev.* **C51**, 371 (1995).