

# Absence of entanglement transition due to feedback-induced skin effect

Yu-Peng Wang<sup>1,2</sup>, Chen Fang<sup>1,3,4</sup>, and Jie Ren<sup>1,2</sup>

<sup>1</sup>Beijing National Laboratory for Condensed Matter Physics and Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China

<sup>2</sup>University of Chinese Academy of Sciences, Beijing 100049, China

<sup>3</sup>Songshan Lake Materials Laboratory, Dongguan, Guangdong 523808, China

<sup>4</sup>Kavli Institute for Theoretical Sciences, Chinese Academy of Sciences, Beijing 100190, China

A quantum many-body system subject to unitary evolution and repeated local measurements with an increasing rate undergoes an entanglement transition from extensive (or subextensive) to area law entropy scaling. We find that certain open boundary systems under “generalized monitoring”, consisting of “projective monitoring” and conditional feedback, display an anomalous late-time particle concentration on the edge, reminiscent of the “skin effect” in non-Hermitian systems. Such feedback-induced skin effect will suppress the entanglement generation, rendering the system short-range entangled without any entanglement transition. While initially emerged in non-interacting models, such skin effect can also occur in chaotic interacting systems and Floquet quantum circuits subjected to random generalized measurements. Since the dynamics of skin-effect do not require post-selection, and can be observed at the particle number level, the phenomenon is experimentally relevant and accessible in noisy intermediate-scale quantum platforms, such as trapped ions.

## 1 Introduction

The competition between the measurement and unitary evolution produces a novel *measurement-induced phase transition* (MIPT) [14, 15, 24, 25, 40, 52–54, 63, 69, 84], where a random quantum circuit [13, 26, 62, 77, 86] interspersed by onsite measurements with an increasing rate goes from a volume-law regime to an area-law regime. Similar entanglement transitions also appear in the context of monitored fermions [1, 5, 9, 11, 12, 18, 59, 61, 73], monitored open systems [49, 75], circuits with pure measurements [41, 50], random tensor networks [45, 57, 76, 81], and quantum error correction thresholds [16, 32–34]. Measurements introduce intrinsic randomness to the otherwise deterministic dynamics, with each set of recorded measurement results corresponding to a specific trajectory [20]. Among various frameworks trying to explain the MIPT [2, 9, 11, 16, 32–34, 38, 39, 42, 44, 51, 55, 74], one approach focuses on a specific trajectory, of which the evolution is described by a non-Hermitian Hamiltonian, where the entanglement transition in some systems coincides with the spontaneous *PT* symmetry breaking [31, 46–48, 72].

One unique phenomenon in certain non-Hermitian open boundary systems is the *non-Hermitian skin effect* [6, 58, 65, 71, 80, 80, 82, 83, 85], where a finite portion of the eigenstates are spatially concentrated on the edges. Dynamically, the skin effect implies that the late-time state from the quench dynamics will have particles concentrated near the edges. The Pauli exclusion principle predicts a nearly tensor-product structure of the steady state which obeys the area-law entanglement scaling, suggesting the absence of MIPT and seemingly contradicting the putative universal entanglement transition in the monitored systems. However, this simple argument assumes that the particular non-Hermitian evolution captures the entanglement behavior of the whole ensemble

---

Chen Fang: [cfang@iphy.ac.cn](mailto:cfang@iphy.ac.cn)

Jie Ren: [jieren@iphy.ac.cn](mailto:jieren@iphy.ac.cn)

of trajectories, which is generally not guaranteed. Indeed, we will show in this work that under “projective monitoring” (the exact definition of which will be given later), on the trajectory ensemble level, there will be no skin effect.

A natural question is whether the skin effect can appear in the whole ensemble of trajectories and thus suppress the entanglement transition. In this work, we answer the question positively by considering a generalized version of monitoring, which contains conditional unitary feedback following projective monitoring. In certain generalized monitored systems, there is a dynamical skin effect induced by the feedback, which we called the feedback-induced skin effect (FISE). We introduce an extensive local order parameter, the classical entropy  $S_{\text{cl}}$ , which not only characterizes the skin effect but also imposes an upper bound on the bipartite entanglement entropy, i.e.,  $S_{\text{cl}} > 2S_A$ , where  $S_A$  is the bipartite entanglement entropy for any subsystem  $A$ . Numerical simulations of finite-size open boundary systems show a scale invariance of the steady-state classical entropy:

$$S_{\text{cl}}(\gamma, L) = L \cdot G(\gamma L^{1/\nu}), \quad G(x \rightarrow \infty) \sim c/x^\nu, \quad (1)$$

which implies the saturation of the entanglement entropy in the thermodynamics limit at arbitrarily small measurement rates. That is, for vanishing small  $\gamma$ , the scaling form in Eq. (1) implies  $S_A < c/\gamma^\nu$ , which is a large but constant value (does not scale with system size) and therefore the system is in the area-law entangled phase. Remarkably, Eq. (1) persists even when the interaction is turned on, therefore eliminating (sub)extensive entangled phases believed to appear in the weakly-monitored regime.

Besides the continuous monitoring, FISE also appears in the randomly measured Floquet circuits, where the unitary evolution has only discrete spatial-temporal translation symmetry, and the monitoring process is the conventional projective measurement followed by a feedback operation. Such quantum circuit models provide a novel type of dynamical skin effect, which is essentially a many-body phenomenon not described by any Hamiltonian. The quantum circuit model is also experimentally relevant to the noisy intermediate-scale quantum devices. As we will show in this work, the establishing of the localized steady state from a “domain-wall state” only takes constant time, and only require a moderate system size to achieve it. We propose to implement such dynamical skin effect on the trapped-ion platforms.

The structure of this paper is as follows. In Sec. 2, we analyze why projective monitoring dynamics fail to produce the desired particle accumulation and explain how the introduction of conditional feedback enables the dynamical skin effect. In Sec. 3, we investigate the phenomenologies of the feedback-induced skin effect and explore its influence on the measurement-induced entanglement transition. In Sec. 4, we extend the study of the feedback-induced skin effect to quantum circuit models with discrete spatial and temporal translation symmetry under random generalized measurements. Finally, we conclude our work in Sec. 5.

## 2 Monitoring dynamics

### 2.1 Projective monitoring

A standard quantum measurement process can be represented by a set of projectors  $P_m$ , each corresponding to a different measurement outcome. Measuring a state  $|\psi\rangle$  yields a probability distribution over the possible outcomes, given by the Born rule:  $p_m = \langle\psi|P_m|\psi\rangle$ . The state then collapses to  $P_m|\psi\rangle/\sqrt{p_m}$ . When the unitary evolution is governed by a Hamiltonian, we can consider a coarse-grained version of the dynamics known as *projective monitoring*. This approach is formulated using the stochastic Schrödinger equation (SSE) [3, 43, 79]:

$$d|\psi\rangle = -i \left[ H - \frac{i\gamma}{2} \sum_m (P_m - \langle\psi|P_m|\psi\rangle) \right] |\psi\rangle dt + \sum_m \left[ \frac{P_m|\psi\rangle}{\|P_m|\psi\rangle\|} - |\psi\rangle \right] dW_m, \quad (2)$$

where each  $dW_m$  is an independent Poisson random variable taking the values 0 or 1. In a small time interval  $\Delta t$ , the probability of observing  $dW_m = 1$  (i.e., the probe registers a quantum jump) is proportional to  $\gamma\Delta t$ . If  $dW_m = 0$  for all  $m$  (the no-click limit) [20, 79], the evolution is described

by an effective non-Hermitian Hamiltonian

$$H_{\text{eff}} = H - i\frac{\gamma}{2} \sum_m P_m. \quad (3)$$

However, the no-click limit requires exponentially many experiments to be carried out before a desired trajectory is obtained.

In this section, we first consider a spinless fermion chain described by a nearest-neighbor hopping Hamiltonian:

$$H = \sum_i (c_i^\dagger c_{i+1} + c_{i+1}^\dagger c_i). \quad (4)$$

The observable of interest in this system is the occupation number of a local quasimode created by a two-site projector  $P_i = d_i^\dagger d_i$ , where  $d_i = \frac{1}{\sqrt{2}}(c_i + ic_{i+1})$ .<sup>1</sup> The effective non-Hermitian Hamiltonian (3) for the monitored system is given by

$$H_{\text{eff}} = \sum_i \left[ \left(1 + \frac{\gamma}{4}\right) c_i^\dagger c_{i+1} + \left(1 - \frac{\gamma}{4}\right) c_{i+1}^\dagger c_i - i\frac{\gamma}{4}(n_i + n_{i+1}) \right]. \quad (5)$$

This Hamiltonian is known as the Hatano-Nelson model [35] and displays the non-Hermitian skin effect. Consider the evolution of the “domain-wall” state:

$$|\psi_0\rangle \equiv |1 \cdots 10 \cdots 0\rangle. \quad (6)$$

In the no-click limit and under open boundary conditions, the effective non-Hermitian Hamiltonian (5) results in a late-time particle accumulation on the left edge (as displayed on the left panel of Fig. 1). On the contrary, in the projective monitoring case, the detected right-moving quasimodes will balance out the momentum distribution, leaving a steady state of homogeneity (as displayed on the right panel of Fig. 1).

Besides SSE, the trajectory-averaged dynamics can also be formulated as the Lindblad master equation [8, 21, 56]:

$$\frac{d}{dt}\rho = -i[H, \rho] - \frac{\gamma}{2} \sum_m \{L_m^\dagger L_m, \rho\} + \gamma \sum_m L_m \rho L_m^\dagger, \quad (7)$$

where  $|\overline{\psi}\rangle\langle\overline{\psi}|$  is the density matrix averaged over all trajectories, and  $L_m = P_m$  is referred to as the *jump operator*. A state  $\rho_{\text{NESS}}$  is a nonequilibrium steady state (NESS) if  $\partial_t \rho_{\text{NESS}} = 0$ . Note that for the projective monitored dynamics, all jump operators  $\{P_i\}$  are Hermitian. The maximally mixed state within a given particle-number sector  $\rho = (\text{Tr } \mathbb{I}_\nu)^{-1} \mathbb{I}_\nu$  (the subscript indicates the subspace spanned by the states of filling number  $\nu$ ) is automatically a steady state since

$$\begin{aligned} \frac{d}{dt} \mathbb{I}_\nu &= -\frac{\gamma}{2} \sum_m \{P_m, \mathbb{I}_\nu\} + \gamma \sum_m P_m \mathbb{I}_\nu P_m^\dagger \\ &= -\gamma \sum_m P_m|_\nu + \gamma \sum_m P_m|_\nu = 0. \end{aligned} \quad (8)$$

In addition,  $\rho_{\text{NESS}}$  for generic Lindblad equation is nondegenerate (within each symmetry sector) [22, 28, 68, 70].<sup>2</sup> In Refs. [22, 28] (see review in Ref. [70] and application in Ref. [68], where the system is under boundary driven [4]), it was shown that a Lindblad equation has unique nonequilibrium steady state if and only if the set

$$\mathcal{O} \equiv \{H, L_1, L_1^\dagger, L_2, L_2^\dagger, \dots\} \quad (9)$$

---

<sup>1</sup>The operator  $d^\dagger$  excites a right-moving wave packet, which can be expressed as  $d^\dagger = \sum_k f(k) c_k^\dagger$  in momentum space, where  $|f(k)|^2 = \frac{1}{4}(1 + \sin k)$  peaks at  $k = \pi/2$ .

<sup>2</sup>Even if we encounter an accidental degeneracy of  $\rho_{\text{NESS}}$ , we can usually find a suitable boundary perturbation to break the degeneracy.




Figure 1: Comparison of non-Hermitian and monitoring dynamics starting from the initial state (6) and under open boundary conditions. Left panel: density evolution under non-Hermitian Hamiltonian with  $\gamma = 0.1$ , where the particles gradually concentrate on the left. Right panel: density evolution under projective monitoring with  $\gamma = 0.1$  (for a single set of measurement results). Although monitoring creates random quantum jump processes and thus introduces density fluctuation, the density distribution on a large scale is homogeneous.

generates (under multiplication and addition) the complete algebra on the Hilbert space.<sup>3</sup> We prove the uniqueness of the steady state  $\rho_{\text{NESS}}$  for the specific model considered in this work in Appendix A. In conclusion, in the projective monitoring case, we expect no skin-effect-like dynamics.

## 2.2 Generalized monitoring

It is important to note that projective monitoring is an idealized representation of an actual measurement process. In practice, detecting a quantum state requires a probe to interact with the system, which inevitably disturbs the measured states. The more general form of a quantum measurement is described by the *positive operator-valued measure* (POVM) formalism [64, 78]. A continuous version of the POVM, called the generalized monitoring, is formulated as the stochastic Schrödinger equation (SSE) in Eq.(2), with the projector  $P_m$  replaced by a general operator  $L_m$  (see Appendix B.1 for a short review). In this study, we focus on a particular form of SSE where

$$L_m = U_m P_m, \quad (10)$$

corresponding to adding a unitary feedback operator  $U_m$  to the projective monitoring process. The conditional feedback does not affect the effective non-Hermitian Hamiltonian  $H_{\text{eff}}$ , but instead operates on those trajectories that deviated from the post-selected trajectory.

Consider a conditional unitary operator

$$U_i = \exp(i\theta n_{i+1}) \quad (11)$$

---

<sup>3</sup>The general proof assumes no conserved quantity for the Lindblad equation. For the particle number conserving case, as we considered in the main text, we can focus on the Hilbert subspace  $\mathcal{H}_\nu$  spanned by states with filling number  $\nu$ . The uniqueness condition then says if  $\mathcal{O}_{|\mathcal{H}_\nu}$  generates the complete algebra on  $\mathcal{H}_\nu$ , the steady state in  $\mathcal{H}_\nu$  will be unique.




Figure 2: (a) Mean particle density evolutions of generalized monitored free fermions ( $L = 128$ ,  $\gamma = 0.1$ ) under open boundary conditions (left panel) and periodic boundary conditions (right panel). The feedback induces a strong boundary sensitivity: under OBC the particle density evolves to be evenly distributed, while under PBC the steady state still features the domain structure. (b) Mean dynamics of  $S_{cl}$  for  $\gamma = 0.05$  for different system sizes. The final saturation values of  $S_{cl}$  depend on the system size  $L$  when  $L$  is no bigger than the domain wall length  $L_{DW}$ . While for sufficiently large size ( $L \geq 256$  in this case),  $S_{cl}(t)$  shows no system size dependence. (c) Mean dynamics of  $S_{cl}$  for  $L = 512$  ( $L > L_{DW}$  for all  $\gamma$ 's we choose). The data collapse indicates that the saturated evolution  $S_{cl}(t; L > L_{DW})$  fits into the universal form Eq. (14). (d) Finite size scaling of steady-state  $S_{cl}$ , which approach the line  $S_{cl} = c/\gamma$  where  $c \approx 3.3$ . Inset: data collapse indicates the scaling form  $S_{cl} = L \cdot G(\gamma L)$ . (e) Momentum distribution of the steady state, in the periodic boundary conditions. (f) Steady-state  $S_{cl}$  for interacting monitored model, which approach the line  $S_{cl} = \tilde{c}/\gamma^\nu$  where  $\tilde{c} \approx 1.8$  and  $\nu = 0.85$ . The data collapse indicates the universal scaling form  $S_{cl} = L \cdot \tilde{G}(\gamma L^{1/\nu})$ .

acting on sites  $(i, i + 1)$  whenever the probe detect a right moving mode  $d^\dagger d$ . We temporarily fix  $\theta = \pi$  so that the feedback  $U_i$  converts the detected right-moving mode to a left-moving mode:

$$\tilde{d}^\dagger \equiv U d^\dagger U^\dagger = \frac{1}{\sqrt{2}}(c_i^\dagger + i c_{i+1}^\dagger). \quad (12)$$

Consequently, the monitored dynamics with feedback always increase left-moving particles, resulting in the FISE where particles concentrate on the left boundary.

As displayed on the left panel of Fig. 2a, starting from  $|\psi_0\rangle$  with a sharp domain wall, the late-time dynamics still feature two static domains where  $\langle n_i \rangle$  only takes the extreme value of 1 or 0. The particle exchange happens only in the vicinity of the border, where the sharp domain wall gradually blurs to a fuzzy domain-wall region. Comparing this with the dynamics with the periodic boundary conditions (right panel of Fig. 2a), where particles quickly disperse into a homogeneous state, we see the FISE features an anomalous boundary sensitivity.

### 3 Feedback-induced skin effect

#### 3.1 Dynamical scale invariance

To quantify the dynamical skin effect induced by feedback, we introduce the classical entropy:

$$S_{\text{cl}}[\{n_i\}] \equiv - \sum_i [n_i \log n_i + (1 - n_i) \log(1 - n_i)]. \quad (13)$$

where only the nontrivial density (i.e.,  $n_i \neq 0, 1$ ) contributes to  $S_{\text{cl}}$ .

In Fig. 2b, we show the time evolution of the classical entropies  $S_{\text{cl}}[\{\overline{n_i}\}]$  starting from  $|\psi_0\rangle$  under open boundary conditions. We see when the system size is small,  $S_{\text{cl}}$  will saturate to a system-size-dependent value. Whereas for large enough systems, the evolution of  $S_{\text{cl}}(t)$  will be the same. These numerical simulations support the “domain-wall blurring” picture, where the evolution of  $S_{\text{cl}}(t)$  follows a universal pattern when considering a system larger than the size of the domain-wall region (denoted as  $L_{\text{DW}}$ ).

In Fig. 2c, we display  $S_{\text{cl}}(t)$  for  $L > L_{\text{DW}}$  systems. We observed a scaling form

$$S_{\text{cl}}(\gamma, t, L > L_{\text{DW}}) = \frac{1}{\gamma} F(\gamma t), \quad (14)$$

where the system reaches the steady state with a characteristic relaxation time  $t_{\text{rlx}} \sim \gamma^{-1}$ , independent of the system size.

Furthermore, we observe that the steady-state  $S_{\text{cl}}$  has a scale invariance, as shown in Fig. 2d:

$$S_{\text{cl}}(\gamma, t > t_{\text{rlx}}, L) = L \cdot G(\gamma L), \quad (15)$$

which implies that  $L_{\text{DW}} \sim \gamma^{-1}$ . In the late-time and thermodynamics limit, the asymptotic behaviors of the scaling functions are

$$F(x \rightarrow \infty) \sim c, \quad G(x \rightarrow \infty) \sim \frac{c}{x}, \quad (16)$$

where  $c$  is a numerical constant estimated to be 3.3 in this specific case. Therefore,

$$S_{\text{cl}}(\gamma, t > t_{\text{rlx}}, L > L_{\text{DW}}) = \frac{c}{\gamma} \approx \frac{3.3}{\gamma}. \quad (17)$$

Notably, Equation (15) suggests a scale invariance for the steady-state density profile  $n(x)$  in the continuum limit.<sup>4</sup> The finite value of  $S_{\text{cl}}$  implies that the feedback-induced skin effect is present even for a vanishingly small measuring rate.

#### 3.2 Suppression of entanglement

One of the consequences of FISE is the suppression of entanglement. Specifically, we prove that  $S_{\text{cl}}$  imposes an upper bound for the (trajectory-averaged) steady-state entropy of the monitored dynamics. Consider an arbitrary subsystem  $A$  inside the monitored system. The entanglement *subadditivity* [64, 78] leads to the inequality

$$S_A \leq \sum_i S_i. \quad (18)$$

Further, consider the reduced density matrix of a local spin at site  $i$ :

$$\rho_i = \begin{bmatrix} n_i & o_i \\ o_i^* & 1 - n_i \end{bmatrix}, \quad (19)$$

---

<sup>4</sup>In contrast, when  $\gamma$  is sufficiently large ( $\gamma \sim O(c)$ ), the lattice effect becomes prominent, and the scaling behavior breaks down.

where  $o_i$  is the off-diagonal element. Mathematically, the eigenvalues of a positive  $2 \times 2$  matrix *majorize* [37] the diagonal elements, and thus have less entropy. That is,

$$S_i = -\text{Tr}[\rho_i \log \rho_i] \leq n_i \log n_i + (1 - n_i) \log(1 - n_i) = S_{\text{cl}}[n_i]. \quad (20)$$

Therefore, the entanglement entropy of subregion  $A$  is bounded by the classical entropy

$$S_A \leq \sum_{i \in A} S_{\text{cl}}[n_i] = S_{\text{cl}}[\{n_i | i \in A\}]. \quad (21)$$

Also, for a pure state, the entanglement entropy of  $A$  is equal to that of its complement  $\bar{A}$ , i.e.,  $S_A = S_{\bar{A}}$ . The classical bound on entanglement entropy also requires

$$S_A + S_{\bar{A}} \leq S_{\text{cl}}[\{n_i\}], \quad (22)$$

therefore  $S_A \leq S_{\text{cl}}/2$ .

In the context of entanglement transition in monitored systems, it is natural to consider the trajectory-averaged entanglement, denoted as  $\overline{S_A}$ , which is bounded by  $\overline{S_{\text{cl}}[\{\langle n_i \rangle\}]} / 2$ . Since the entropy function is convex,  $\overline{S_A}$  will also be bounded by:

$$\overline{S_A} \leq \frac{1}{2} \overline{S_{\text{cl}}[\{\langle n_i \rangle\}]} \leq \frac{1}{2} S_{\text{cl}}[\langle n_i \rangle]. \quad (23)$$

The asymptotic behavior, therefore, predicts the area-law entanglement scaling  $S \leq c/2\gamma$  for arbitrary  $L$ , thus proving the area-law entanglement scaling in the  $\gamma \rightarrow 0$  limit.

To complete the analysis, in Appendix C.2, we explicitly calculate the trajectory-averaged entanglement entropy (as well as the classical entropy) for the monitored dynamics. We observe that in the periodic boundary conditions, the entanglement entropy scaling indeed undergoes a logarithm-to-area-law transition, while this transition is missing in the open boundary conditions. Also, in order to show the effect of the trajectory variance, as well as find a tighter bound on the entanglement entropy, we calculate the  $\overline{S_{\text{cl}}[\{\langle n_i \rangle\}]}$ .

In the above analysis, we consider only the  $\theta = \pi$  case in the feedback operator, while similar FISE appears for arbitrary  $\theta$ . In Appendix C.3, we show the numerical results for  $\theta \neq \pi$ . The length of the domain wall, however, is minimized at  $\theta = \pi$ . When  $\theta$  approaches zero, we expect that the FISE still appears, although with a large domain wall that may exceed the numerical simulation capability. Therefore, for the projector  $P_i$ , the FISE is not a fine-tuned phenomenon and may appear for a large class of feedback operations.

### 3.3 Bulk-edge correspondence

The bulk-edge correspondence of the non-Hermitian effect has been extensively investigated [6, 65, 80, 83, 85]. In the no-click limit, where the dynamics are described by the non-interacting non-Hermitian Hamiltonian, the skin effect also manifests as a directional bulk current in the periodic boundary systems:

$$J[\{n_k\}] \equiv \int_{-\pi}^{+\pi} v_k n_k dk, \quad (24)$$

where  $n_k = \langle c_k^\dagger c_k \rangle$  is the distribution function of the  $k$ -momentum fermion modes, and  $v_k = \partial_k E(k)$  is the velocity of each quasi-mode.

The rigorous bulk-edge correspondence only happens in the no-click limit, whereas when the quantum jump processes are taken into account, the system lost the single-particle description, and thus the velocity  $v_k$  becomes ill-defined. However, we can generalize the concept of the directional bulk current  $J[\{n_k\}]$  at least to the weakly-monitored system, where the velocity  $v_k$  can be approximated by the result of the free system  $v_k \approx \partial_k E(k) = \sin(k)$ .

By simulating the system with identical parameters but under periodic boundary conditions, we show in Fig. 2e that there is an imbalance in the momentum distribution which does not vanish in the  $\gamma \rightarrow 0$  limit. Specifically, for  $\gamma = 0.01$  case, the numerical simulation shows that  $J[\{n_k\}] \approx -0.94$ . This nonzero current serves as another manifestation of the dynamical skin effect at arbitrarily small measurement rates.



### 3.4 Monitored interacted system

The monitored free-fermion systems belong to a special type of MIPT since the entanglement of the weakly monitored system is not strong, i.e., does not follow the volume-law scaling. From the quantum information point of view, the free fermion systems are not good quantum memory; the initial quantum information will be lost in polynomial time when subject to measurements [24]. It is therefore necessary to exclude the possibility that FISE only suppresses the logarithm-scaling entanglement.

For the interacting system, it is more convenient to focus on a spin-1/2 Hamiltonian

$$H = \sum_i [J_1(\sigma_i^x \sigma_{i+1}^x + \sigma_i^y \sigma_{i+1}^y) + J_z \sigma_i^z \sigma_{i+1}^z + J_2(\sigma_i^x \sigma_{i+2}^x + \sigma_i^y \sigma_{i+2}^y)]. \quad (25)$$

In Ref. [30], the non-integrable system (25) under continuous monitoring (on local  $\sigma_i^z$ ) was shown to display a volume-to-area-law entanglement transition.

In our case, however, we choose the same generalized monitoring as in the free fermion case (via Jordan-Wigner transformation):

$$P_i = \frac{1}{2}(\sigma_i^+ - i\sigma_{i+1}^+)(\sigma_i^- + i\sigma_{i+1}^-), \quad U_i = S_{i+1}^z. \quad (26)$$

In the presence of interactions, FISE also manifests in open boundary systems under the influence of this generalized monitoring. The presence of the dynamical skin effect leads to area-law entanglement entropies, allowing for efficient representation of states using matrix-product states [19, 23]. Additionally, the time evolution of large system sizes can be simulated effectively using the TEBD algorithm.

We investigate the steady-state classical entropy for different  $\gamma$  values and system sizes up to  $L = 64$ , and fix  $J_1 = 0.5$ ,  $J_z = 1.0$ ,  $J_2 = 0.1$ . As shown in Fig. 2f, the numerics indicate that for  $\gamma > 0.5$ , the classical entropy exhibits the asymptotic behavior:

$$S_{\text{cl}}(\gamma, t > t_{\text{rlx}}, L > L_{\text{DW}}) \approx \frac{\tilde{c}}{\gamma^\nu}, \quad (27)$$

where  $\tilde{c} = 1.8$  and  $\nu = 0.85$ . This suggests that a matrix product state with bond dimension  $\chi \sim O(\exp(\gamma^{-\nu}))$  is sufficient to describe the state. However, the numerical simulation becomes challenging in the small  $\gamma$  regime. While we are unable to numerically verify the scaling behavior in the small  $\gamma$  limit due to computational limitations, our finite-size simulations still demonstrate the scaling law described by (as depicted in Fig. 2f)

$$S_{\text{cl}}(\gamma, t > t_{\text{rlx}}) = L \cdot \tilde{G}(\gamma L^{1/\nu}), \quad (28)$$

where the scaling function shall obey the asymptotic behavior

$$\tilde{G}(x \rightarrow \infty) \sim \frac{\tilde{c}}{x^\nu}. \quad (29)$$

We conjecture that the asymptotic form in Eq. (28) continues to hold in the small- $\gamma$  regime.

To complete the analysis, we show that under periodic boundary conditions, the usually measurement-induced entanglement appears in this generalized monitored system. We demonstrate this transition using finite-size exact diagonalization. We remark here that the appearance of an entanglement transition can be better revealed from the mutual information  $I(A : B)$  shared by two antipodal sites  $A$  and  $B$ . This numerical technique has been used, for example, in Refs. [29, 53].

In Fig. 3a, we first demonstrate that for small  $\gamma$ , the finite size scaling of entanglement entropy  $S(L, L/2)$  shows a linear growth in the size  $L$ . In the large  $\gamma$  regime, the entanglement is greatly suppressed, suggesting steady states being area-law entangled. To better demonstrate the existence of an entanglement transition, Fig. 3b displays the mutual information  $I(A : B)$ . The peak in  $I(A : B)$  shows near  $\gamma_c \approx 1$ . Due to the finite size available to the numerical simulation, we are unable to pinpoint the critical measurement rate  $\gamma_c$ .

Therefore, we show that the FISE indeed imposes strong suppression on the entanglement which completely excludes any entanglement transition.






Figure 3: Exact diagonalization simulation of the entanglement dynamics for systems with  $J_1 = J_z = 0.5$ ,  $J_2 = 0.1$ , and under periodic boundary conditions. (a) Trajectory average of entanglement entropy for different measurement rates. (b) Trajectory average of mutual information  $I(A : B)$  between two antipodal sites  $A$  and  $B$ . The peak of  $I(A : B)$  indicates the critical measurement rate at  $\gamma_c \approx 1$ .

## 4 Floquet circuits under random measurements

Up till now, we have investigated FISE in the context of the continuously monitored Hamiltonian dynamics. Readers may wonder whether FISE is an artifact of continuous monitoring. This section serves to clarify this question by generalizing the FISE to the quantum circuits under random (generalized) measurements, which is similar to the original setup for the MIPT. The upshot is that only the discrete spatial-temporal translation symmetry is required for the skin effect.

### 4.1 Wave packet motion under Floquet evolution

The notion of discrete quantum circuits naturally appears when we use the Trotter decomposition. In general, when a local Hamiltonian can be decomposed into several groups:

$$H = \sum_{\alpha=1}^l H_{\alpha}, \quad H_{\alpha} = \sum_i o_{\alpha,i}, \quad (30)$$

where each  $o_{\alpha,i}$  is a local operator. Within each group, the local operators do not overlap. The time evolution can then be approximated by

$$U(t = N\Delta t) \approx (e^{-iH_1\Delta t} e^{-iH_2\Delta t} \dots e^{-H_l\Delta t})^N. \quad (31)$$

The decomposition approaches the real dynamics in the  $\Delta t \rightarrow 0$  limit. While for practical reasons,  $\Delta t$  is a finite value, therefore the dynamics described by Eq. (31) is a Floquet dynamics with period  $T = l\Delta t$ .

We show that the notion of wave packet motion is also valid in the context of the Floquet circuit even when the  $\Delta t$  is not vanishing small, as long as the discrete translational symmetry is preserved. For simplicity, we assume that the Floquet circuit has 2-site translational symmetry. We can group two lattice sites into a unit cell and label them as the internal degrees of freedom  $a = 1, 2$ . The plane wave is then

$$|k, a\rangle \equiv \frac{1}{\sqrt{L}} \sum_j c_{j,a}^\dagger e^{-ikj} |0\rangle. \quad (32)$$

Due to the translational invariance, the evolution  $U(T)$  is blocked-diagonal in the momentum space:

$$U_{ab} \equiv \langle k, a | U(T) | k, b \rangle = V(k) \begin{bmatrix} e^{-iE_+(k)T} & 0 \\ 0 & e^{-iE_-(k)T} \end{bmatrix} V(k)^\dagger. \quad (33)$$




Figure 4: (a) Dispersion relation for circuit model. At the maximal velocity point  $k = \pi$ , the eigenvectors are  $|\pi, -\rangle \approx 0.842|\pi, 1\rangle + (0.290 - 0.455i)|\pi, 2\rangle$  and  $|\pi, +\rangle \approx (0.290 + 0.455i)|\pi, 1\rangle - 0.842|\pi, 2\rangle$ . (b) Quantum circuit model with Floquet circuit evolution and conditional feedback. (c) Density evolution of the measured Floquet dynamics ( $L = 512$ ,  $p = 0.02$ ).

The values  $E_{\pm}(k)$  extracted from the eigenvalues of  $U_{ab}$  are the quasi-energies of the Floquet dynamics. Consider the branch with quasi-energy  $E_+$ . The wave packet with averaged momentum  $k$  and averaged position  $x$  has the form

$$|k, x; +\rangle \equiv \int \frac{dp}{2\pi} e^{-\alpha(p-k)^2 - ix(p-k)} |p, +\rangle, \quad (34)$$

where  $\alpha$  controls the variance of the momentum distribution of the wave packet. After one period  $T$ , the wave packet evolves to

$$\begin{aligned} U(T)|k, x; +\rangle &= \int \frac{dp}{2\pi} e^{-\alpha(p-k)^2 - ix(p-k) - iE_+(p)T} |p, +\rangle \\ &\propto \int \frac{dp}{2\pi} e^{-\alpha(p-k)^2 - i\left[x + \frac{E_+(p) - E_+(k)}{p-k}T\right](p-k)} |p, +\rangle. \end{aligned}$$

Specifically, when  $\alpha$  is large, most  $p$  will be close to  $k$ , and the movement of the averaged position is approximated by  $x \rightarrow x' \approx x + E'_+(k)T$ . We therefore restore the wave packet moving picture in the Hamiltonian dynamics, at least for the nearly monochromatic wave packets.

With the picture in mind, we first consider the Floquet dynamics described by the circuits

$$U(T) = \prod_j \exp \left[ -i \left( c_{2j}^\dagger c_{2j+1} + c_{2j+1}^\dagger c_{2j} \right) T \right] \times \prod_j \exp \left[ -i \left( c_{2j-1}^\dagger c_{2j} + c_{2j}^\dagger c_{2j-1} \right) T \right]. \quad (35)$$

We can choose an arbitrary finite value of  $T$ , say  $T = 0.5$ . The dispersion is then calculated and displayed in Fig. 4a. We remark that for finite  $T$ , the eigenvectors have shifted compared to the Hamiltonian case. However, the tendency remains the same. That is, the configuration  $d^\dagger|0\rangle$  still has a bigger overlap with the right-moving wave packet than with the left-moving ones. Therefore, we expect that the feedback-induced skin effect still appears in the Floquet model.

## 4.2 Skin effect in measured Floquet circuits

Besides the generalization to the Floquet circuit, we show here that the continuous monitoring setup can be replaced with randomly applying strong projective measurement followed by conditional feedback.

Consider the circuit dynamics in Fig. 4b, where the  $U_1$  and  $U_2$  come from the Floquet dynamics. Under the Jordan-Wigner transformation, they become

$$U_1 = U_2 = \exp \left( -i \frac{T}{2} \sigma_x \otimes \sigma_x \right) \exp \left( -i \frac{T}{2} \sigma_y \otimes \sigma_y \right), \quad (36)$$

and the measurements are the projective measurement of the observable

$$P_i = \frac{1}{4} \left( \sigma_i^z + \sigma_{i+1}^z + \sigma_i^x \sigma_{i+1}^y - \sigma_i^y \sigma_{i+1}^x + 2 \right), \quad (37)$$

followed by the conditional feedback  $U_i = \sigma_i^z$ . As displayed in Fig. 4c, for a small measurement rate ( $p = 0.02$ ), the feedback still induces the skin effect.

## 4.3 Experimental proposal on the trapped ions quantum computers

The two-site measurements is hard to realize in the experiments. However, we can devise a single-site measured model using a unitary transformation. To formulate the transformation in a clearer way, we temporarily reverse to the fermionic model where the quasi-mode  $d_i = c_i + ic_{i+1}$  are being measured. We now regard  $d_{2i-1}$  as the local fermion mode, which corresponds to the transformation

$$d_{2i-1} = \frac{1}{\sqrt{2}}(c_{2i-1} + ic_{2i}), \quad d_{2i} = \frac{1}{\sqrt{2}}(c_{2i} + ic_{2i-1}). \quad (38)$$

The Hamiltonian is mapped to  $H = h_1 + h_2 + h_3 + h_4$ , where  $\{h_1, h_2, h_3, h_4\}$  are hopping terms among neighboring unit cells:

$$\begin{aligned} h_1 &= \sum_i \left( d_{2i-1}^\dagger d_{2i} + H.c. \right), \\ h_2 &= \frac{1}{2} \sum_i \left( d_{2i}^\dagger d_{2i+1} + H.c. \right), \\ h_3 &= \sum_i \left( \frac{i}{2} d_{2i-1}^\dagger d_{2i+1} - \frac{i}{2} d_{2i}^\dagger d_{2i+2} + H.c. \right), \\ h_4 &= \frac{1}{2} \sum_i \left( d_{2i-1}^\dagger d_{2i+2} + H.c. \right). \end{aligned} \quad (39)$$

Under the new basis, the projector becomes an onsite measurement of particle occupation on the odd sites:

$$P_{2i-1} = d_{2i-1}^\dagger d_{2i-1}, \quad P_{2i} = d_{2i}^\dagger d_{2i}. \quad (40)$$

The feedback operator, however, is not in a neat form on this basis. However, as we remarked, the FISE is not a fine-tuned effect from the feedback, we expect FISE to happen for a wide range of conditional unitary operators. For simplicity, we choose the feedback to be the swap operator on two neighboring sites. To obtain an even simpler model, we keep only half of the measurement  $P_{2i-1}$  on the odd sites, and we choose the feedback operator to be the swap gate

$$U_{2i-1} = \text{SWAP}_{2i-1, 2i}. \quad (41)$$

Under the Jordan-Wigner transformation, the measured circuit is displayed in Fig. 4d. The unitary gate  $\{U_1, \dots, U_5\}$  in the circuits are:

$$\begin{aligned} U_1 &= \text{XX}(\theta) \otimes \text{YY}(\theta), \\ U_2 &= \text{XX}\left(\frac{\theta}{2}\right) \otimes \text{YY}\left(\frac{\theta}{2}\right), \\ U_3 &= \text{XY}\left(\frac{\theta}{2}\right) \otimes \text{YX}\left(-\frac{\theta}{2}\right), \\ U_4 &= \text{XY}\left(-\frac{\theta}{2}\right) \otimes \text{YX}\left(\frac{\theta}{2}\right), \\ U_5 &= \text{XX}\left(\frac{\theta}{2}\right) \otimes \text{YY}\left(\frac{\theta}{2}\right), \end{aligned} \quad (42)$$

where  $\theta$  is the parameter controlled by the time step of the Trotter decomposition, and the XX, XY, and YY gates are

$$\text{XX}_{i,j}(\theta) \equiv \exp(-i\theta\sigma_i^x\sigma_j^x), \quad (43)$$

$$\text{XY}_{i,j}(\theta) \equiv \exp(-i\theta\sigma_i^x\sigma_j^y), \quad (44)$$

$$\text{YY}_{i,j}(\theta) \equiv \exp(-i\theta\sigma_i^y\sigma_j^y). \quad (45)$$

The measurement operator is now simply the  $\sigma_{2i-1}^z$  on odd sites, and the conditional feedback SWAP (which does not change under Jordan-Wigner transformation) will be applied to the sites  $(2i-1)$  and  $(2i)$  when the measurement result in site  $(2i-1)$  is  $-1$ .

Experimentally, the trapped ions systems [17, 36, 60, 66] are one of the ideal platforms to realize medium-scale quantum computation and quantum simulations. Using the  $^{171}\text{Yb}^+$  ions trapped by the high-frequency electromagnetic field [66] with fined tune laser pulse, the platform is able to:

1. initiate the state as  $|0 \cdots 0\rangle$ .
2. implement any single-site rotation, which enables to initiate the state to any product state.
3. implement an ‘‘Ising’’ spin-spin interaction [67, 87]:  $\text{XX}_{i,j}(\theta) = \exp(-i\theta\sigma_i^x \otimes \sigma_j^x)$ .

4. do onsite measurement on  $\sigma^z$ , collapsing qubit to  $|0\rangle$  or  $|1\rangle$  state.

We remark that, using the combination of single-site and two-site unitary gates, it is possible to realize the YY and XY gates as:

$$\begin{aligned} XY_{i,j}(\theta) &= \exp\left(-i\frac{\pi}{2}\sigma_j^z\right) XX_{i,j}(\theta) \exp\left(i\frac{\pi}{2}\sigma_j^z\right), \\ YY_{i,j}(\theta) &= \exp\left(-i\frac{\pi}{2}\sigma_i^z\right) XY_{i,j}(\theta) \exp\left(i\frac{\pi}{2}\sigma_i^z\right). \end{aligned} \quad (46)$$

Now consider the monitored circuit model displayed in Fig. 4d, where the Floquet dynamics is the Trotterized version of Hamiltonian (39).

The measurement that needs to perform the simple  $S^z$  measurement, followed by a conditional swap operation, which is

$$U_{\text{SWAP}} \propto XX\left(\frac{\pi}{4}\right) \otimes YY\left(\frac{\pi}{4}\right). \quad (47)$$

In Fig. 4f, we simulate the dynamics on a 32-qubit chain, and choose  $\theta = \pi/4$ , with measurement probability  $p = 0.5$ . We see the late time state show a clear feedback-induced skin effect.

## 5 Conclusion

This work investigates the effect of generalized monitoring on the entanglement phase transition, with a particular focus on the emergence of the skin effect in certain open boundary monitored systems. Our analysis reveals that the skin effect qualitatively alters the entanglement structure of the nonequilibrium steady state, leading to a single area-law phase. Specifically, we show that introducing generic feedback operators can disturb the balance of particle distribution, resulting in particle accumulation. We demonstrate that the FISE is not a fine-tuned phenomenon, as the skin effect appears for different feedback parameters, and survives in the presence of interactions. The suppression of the entanglement entropy from the skin effect also enables an efficient classical simulation of the monitored interacted systems.

These results have practical implications in the context of open systems or controllable quantum devices, as the monitoring-feedback setup can enable the realization of a skin effect without the need for post-selection in non-Hermitian dynamics. For systems showing FISE, the steady states can be reached in constant steps, which is accessible for the noisy intermediate-scale quantum devices. We thus proposed a quantum circuit model displaying FISE which can be experimentally realized on the trapped ions systems.

*Note added.*— In the middle of this work, we became aware of a recent work [10], which also considers the effect of generalized monitoring in the context of MIPT. The two works are complementary to each other: in Ref. [10], the authors utilize the feedback (pre-selection) to reveal MIPT as a quantum absorbing state transition that can be directly detected, while our work shows that the presence of conditional feedback may also eliminate the MIPT.

## Acknowledgements

J.R. thanks Chenguang Liang for the valuable discussions. J.R. and Y.P.W. thank Xu Feng and Shuo Liu for alerting them of a typo in the previous version. The numerical simulations based on tensor-network use the `ITensor` package [27].

## A Uniqueness of nonequilibrium steady state

In this section, we prove the uniqueness of  $\rho_{\text{NESS}}$  for models considered in the main text, by explicitly checking the operator set

$$\mathcal{O} = \{H, L_1, L_1^\dagger, L_2, L_2^\dagger, \dots\} \quad (48)$$

generates (under multiplication and addition) the complete algebra on the Hilbert space.

## A.1 Projective monitoring

We first prove that the Hamiltonian (under open boundary conditions)

$$H = \sum_i (c_i^\dagger c_{i+1} + c_{i+1}^\dagger c_i) \quad (49)$$

and the projectors

$$P_i = \frac{1}{2}(c_i^\dagger - ic_{i+1}^\dagger)(c_i + ic_{i+1}) \quad (50)$$

generate the whole algebra. Note  $H$  and  $P_1, P_2$  together generate the following particle number operators:

$$\begin{aligned} n_1 - n_3 &= P_2 - P_1 + i[H, P_1 + P_2], \\ n_1 + n_2 &= \frac{1}{2}(P_1 + P_2 + i[H, n_1 - n_3] + n_1 - n_3), \\ n_2 + n_3 &= (n_1 + n_2) - (n_1 - n_3). \end{aligned} \quad (51)$$

Then, some straightforward algebra lead to

$$\begin{aligned} c_1^\dagger c_2 - c_2^\dagger c_1 &= i(n_1 + n_2) - iP_1, \\ c_1^\dagger c_2 + c_2^\dagger c_1 &= [c_1^\dagger c_2 - c_2^\dagger c_1, n_2 + n_3], \\ c_2^\dagger c_3 - c_3^\dagger c_2 &= i(n_2 + n_3) - iP_2, \\ c_2^\dagger c_3 + c_3^\dagger c_2 &= [n_1 + n_2, c_2^\dagger c_3 - c_3^\dagger c_2]. \end{aligned} \quad (52)$$

Upon some addition among Eqs. (52), we obtain the operator  $c_1^\dagger c_2, c_2^\dagger c_3$  and their Hermitian conjugates. The commutations of them further produce  $c_1^\dagger c_3$  and its conjugate. Also, note that

$$[c_1^\dagger c_2, c_2^\dagger c_1] = n_1 - n_2. \quad (53)$$

Together with Eqs. (51), we generate all fermion bilinear terms  $c_i^\dagger c_j$  (including  $i = j$  case) on sites  $(1, 2, 3)$ . To proceed, we subtract the hopping terms between sites  $(1, 2)$  from  $H$ . The resulting operator is equivalent to a shorter chain starting from site 2. We can then utilize the calculation above to obtain all  $c_i^\dagger c_j$  terms on sites  $(2, 3, 4)$ . We eventually obtain all fermion bilinear terms on the chain by applying the strategy iteratively. Note that fermion bilinear terms  $c_i^\dagger c_j$  generate the complete algebra within a fixed particle-number sector since any two product states in the sector can be related by applying several fermion hopping terms.

## A.2 Generalized monitoring and interactions

For the generalized monitored system described by the jump operators  $\{L_n = U_n P_n\}$ , the proof of uniqueness is essentially the same as the projective case. Note that we can generate all  $P_i$  terms by multiplying to jump operators

$$P_i = L_i^\dagger L_i. \quad (54)$$

In this way, we can generate the complete operator algebra in the same way as above.

We argue that the completeness of the operator algebra holds for generic open systems since the exact decoupling of Hilbert space is the result of symmetries or fine-tuning. For the interacting system where the Hamiltonian is

$$H = \sum_i (c_i^\dagger c_{i+1} + c_{i+1}^\dagger c_i + g n_i n_{i+1}). \quad (55)$$

The above argument means for a random value of  $g$ , we should expect the completeness of the operator algebra. We can also consider the case where the coupling constant is smoothly varying in the space. In particular, let  $g_i = 0$  for the sites near the boundary. Following the same procedure, we can generate the operator algebra of the subsystem near the boundary. Assume that we meet the first nonzero  $g$  at site  $i + 1$ . It means that we have the complete operator algebra (within

fixed filling number)  $\mathcal{A}_{[1,i]}$  of the subsystem consisting of sites  $1, \dots, i$ . We first subtract all terms within  $\mathcal{A}_{[1,i]}$  from the Hamiltonian and denote the result as  $H_{[i+1,N]}$ . Consider the commutators

$$\begin{aligned} [n_{i-1}, H_{[i+1,N]}] &= c_i^\dagger c_{i+1} - c_{i+1}^\dagger c_i, \\ [n_{i-1}, c_i^\dagger c_{i+1} - c_{i+1}^\dagger c_i] &= c_i^\dagger c_{i+1} + c_{i+1}^\dagger c_i. \end{aligned} \quad (56)$$

In this way, we generate the hopping terms  $c_i^\dagger c_{i+1}$  and  $c_{i+1}^\dagger c_i$ . Those terms together with  $\mathcal{A}_{[1,i]}$  generate the algebra  $\mathcal{A}_{[1,i+1]}$ . This procedure can proceed iteratively, therefore producing the complete algebra.

## B Stochastic Schrödinger equation

### B.1 Generalized monitoring

Microscopically, a measurement process involves a short-time interaction between the system and the probe, which are initially separable:

$$|\psi_{AB}\rangle = e^{-iH_{\text{int}}\Delta t} |\psi_A\rangle \otimes |\psi_B\rangle, \quad (57)$$

where the wave function of the measured system is denoted as  $|\psi_A\rangle$  and the probe  $|\psi_B\rangle$ . When  $\Delta t$  is much smaller than the time scale of the system, the system can be regarded as static during the measurement. Such measurement is called the *strong measurement*. The probe is thought to be a device that can convert quantum information to the classical one, which takes the form of standard projective measurement. That is, suppose the eigenbasis of the probe is  $\{|\phi_n\rangle\}$ , the probability of getting a record  $n$  is

$$p_n = \langle \phi_n | \rho_B | \phi_n \rangle, \quad \rho_B \equiv \text{Tr}_A |\psi_{AB}\rangle \langle \psi_{AB}|, \quad (58)$$

and the feedback of the measurement to the system is

$$|\tilde{\psi}_A^{(n)}\rangle = \langle \phi_n | e^{-iH_{\text{int}}\Delta t} |\psi_A\rangle \otimes |\psi_B\rangle \equiv M_n |\psi_A\rangle. \quad (59)$$

The completeness condition requires

$$\sum_n \langle \tilde{\psi}_A^{(n)} | \tilde{\psi}_A^{(n)} \rangle = 1 \implies \sum_n M_n^\dagger M_n = 1. \quad (60)$$

This is the general form of the measure. In the language of density operator, a measurement is described by a set of operators  $\{M_n\}$ . A measurement process may record a result  $n$  with probability  $p_n$  and change the state to:

$$\rho \rightarrow \frac{M_n \rho M_n^\dagger}{\|M_n \rho M_n^\dagger\|}. \quad (61)$$

If the measurement result is not known, the averaged density matrix after the measurement is

$$\rho \rightarrow \sum_n M_n \rho M_n^\dagger. \quad (62)$$

Such a map is called the quantum channel [64].

On the other hand, if the strength of system-probe coupling is comparable with the energy scale of the system, which is the case for an open quantum system, the quantum channel expression should depend on time  $\Delta t$ . This kind of measurement process is called *weak measurement*. When the system is *Markovian* (the equation of motion depends only on the near past), the course-grained dissipation process can be described by the channel:

$$M_n = L_n \sqrt{\gamma \Delta t}, \quad (63)$$

$$\begin{aligned} M_0 &= \sqrt{1 - \sum_{n>0} M_n^\dagger M_n} \\ &= 1 - \frac{\gamma}{2} \sum_{n>0} L_n^\dagger L_n \Delta t + O(\Delta t^2). \end{aligned} \quad (64)$$






Figure 5: (a) Schematic of a fermion chain under nearest neighbor interaction and generalized monitoring on each pair of neighboring sites. Unitary feedback will be applied if the probe records a quantum jump. (b) Circuits representation of the discretized Hamiltonian evolution with constant monitoring.

For the density matrix, the coarse-grained differential equation is the Lindblad equation:

$$\begin{aligned} \frac{d\rho}{dt} &= \lim_{\Delta t \rightarrow 0} \frac{1}{\Delta t} \sum_n M_n e^{-iH\Delta t} \rho e^{iH\Delta t} M_n^\dagger \\ &= -i[H, \rho] - \frac{\gamma}{2} \sum_i \{L_i^\dagger L_i, \rho\} + \gamma \sum_i L_i \rho L_i^\dagger. \end{aligned} \quad (65)$$

The joint dynamics of Hamiltonian evolution and measurement can be equivalently described by the stochastic process, as shown in Fig. 5, where for each time step  $\Delta t$ , the system first undergoes a coherent evolution  $|\psi\rangle \rightarrow e^{-iH\Delta t}|\psi\rangle$ , then the application of measurement produces a random process:

$$|\psi\rangle \rightarrow \begin{cases} M_n(\Delta t)|\psi\rangle & P_n = \langle\psi|L_n^\dagger L_n|\psi\rangle\gamma\Delta t \\ M_0(\Delta t)|\psi\rangle & P_0 = 1 - \sum_{n>0} P_n \end{cases}. \quad (66)$$

Different records of the measurement result correspond to different trajectories, and the Lindblad equation is equivalent to the trajectory averaged of such stochastic processes. In the continuum limit, the stochastic differential equation can be formulated by introducing a Poisson random variable  $dW_n$  taking the discrete values of 0 or 1. The  $dW_n = 1$  case corresponds to registering a quantum jump, otherwise,  $dW_n = 0$ . The expectation value for random  $dW_n$  is proportional to  $dt$ :

$$\overline{dW_n} = \langle\psi|L_n^\dagger L_n|\psi\rangle\gamma dt. \quad (67)$$

Different  $dW_n$ 's are independent, i.e., they satisfy the orthogonal condition

$$dW_m dW_n = \delta_{mn} dW_m. \quad (68)$$

Therefore, the random quantum jump process  $|\psi\rangle \rightarrow M_n|\psi\rangle$  is described by the expression

$$d|\psi\rangle = \left( \frac{L_n}{\sqrt{\langle L_n^\dagger L_n \rangle}} - 1 \right) |\psi\rangle dW_n, \quad (69)$$

and the null-detection case correspond to  $|\psi\rangle \rightarrow M_0|\psi\rangle$ , which is described by a non-Hermitian differential equation:

$$d|\psi\rangle = -\frac{\gamma}{2} \sum_m (L_m^\dagger L_m - \langle L_m^\dagger L_m \rangle) |\psi\rangle dt, \quad (70)$$

where the  $\langle L_m^\dagger L_m \rangle$  is introduced for the normalization purpose. Together with the coherent evolution, we obtain the stochastic Schrödinger equation in the main text:

$$d|\psi\rangle = \left[ -iH - \frac{\gamma}{2} \sum_m (L_m^\dagger L_m - \langle L_m^\dagger L_m \rangle) \right] |\psi\rangle dt + \sum_m \left( \frac{L_m}{\sqrt{\langle L_m^\dagger L_m \rangle}} - 1 \right) |\psi\rangle dW_m. \quad (71)$$

## B.2 Free fermion simulation

For numerical simulation of Eq. (71), we can first discretize the time into small interval  $\Delta t$ . The discrete evolution is then

$$|\psi(t + \Delta t)\rangle = \mathcal{M}_{\Delta t}[e^{-iH_{\text{eff}}\Delta t}|\psi(t)\rangle], \quad (72)$$

where  $\mathcal{M}_{\Delta t}$  represents the quantum jump that randomly happened in time interval  $\Delta t$ :

$$\mathcal{M}_{\Delta t}[|\psi\rangle] \propto \prod_{m \in I} L_m |\psi\rangle. \quad (73)$$

In the Eq. (73), the set  $I$  denotes the random jump processes, which can be obtained by

$$I = \{n | r_n < \gamma \langle L_n^\dagger L_n \rangle \Delta t\}, \quad (74)$$

where  $\{r_n \in (0, 1)\}$  is a set of independent random variables with evenly distributed probability.

The free fermion system can be efficiently represented by the Gaussian state [7]. For a particle number conserving system, the Gaussian state is a quasimode-occupied state, represented by a matrix  $B$ :

$$|B\rangle \equiv \prod_{j=1}^N \sum_i B_{ij} c_i^\dagger |0\rangle = \bigotimes_{j=1}^N |B_j\rangle, \quad (75)$$

where each column  $B_j$  is an occupied quasimode. Note that there is an  $\text{SU}(N)$  gauge freedom for the matrix  $B$ , i.e.,

$$|B'\rangle = |BU\rangle = |B\rangle, \quad (76)$$

where  $U$  is an arbitrary  $\text{SU}(N)$  matrix. Such gauge freedom implies that a Gaussian state is entirely specified by the linear subspace spanned by the quasimodes  $B_i$ 's.

The random Schrödinger equation can be Trotterized as Eq. (72). Using the Baker-Campbell-Hausdorff formula  $e^A B e^{-A} = e^{\text{ad}_A} B$ , the nonunitary evolution is

$$\begin{aligned} e^{-iH_{\text{eff}}\Delta t} |B_t\rangle &= \prod_{j=1}^N \sum_i B_{ij} e^{-iH_{\text{eff}}\Delta t} c_i^\dagger e^{iH_{\text{eff}}\Delta t} |0\rangle \\ &= \prod_{j=1}^N \sum_i B_{ij} e^{-i\Delta t [H_{\text{eff}}, \cdot]} c_i^\dagger |0\rangle \\ &= \prod_{j=1}^N \sum_i \sum_k B_{ij} c_k^\dagger [e^{-iH_{\text{eff}}\Delta t}]_{ki} |0\rangle \\ &= \prod_{j=1}^N \sum_k [e^{-iH_{\text{eff}}\Delta t} B_{ij}]_{kj} c_k^\dagger |0\rangle \\ &= |e^{-iH_{\text{eff}}\Delta t} B_t\rangle. \end{aligned}$$

That is, the matrix is multiplied by the exponential of the effective non-Hermitian (single-body) Hamiltonian matrix. Note that the resulting matrix is not orthogonal anymore, while the state is still well-defined by the linear space spanned by those unorthogonal vectors. In general, for a Gaussian state represented by matrix  $B$ , we can obtain a canonical form for the representing matrix using the QR decomposition  $B = Q \cdot R$ , where  $Q$  is a unitary matrix and  $R$  is upper triangular. Note that  $Q$  and  $B$  span the same linear space, so the Gaussian state can be expressed as  $|Q\rangle$ .

The supper operator  $\mathcal{M}_{\Delta t}$  in Eq. (72) corresponds to the Poisson jump process, where for each index  $i$ , we randomly decide whether a quantum jump process

$$|B\rangle \rightarrow \frac{L_i |B\rangle}{\|L_i |B\rangle\|} \quad (77)$$

happens, with the probability

$$p_i = \langle B | L_i^\dagger L_i | B \rangle \gamma \Delta t \quad (78)$$

The  $L_m$ 's we choose in the main text have the form

$$L_m = e^{ih} d^\dagger d, \quad d^\dagger = \sum_i a_i c_i^\dagger, \quad h = \sum_{ij} h_{ij} c_i^\dagger c_j, \quad (79)$$

where  $d^\dagger$  is a quasimode, and  $h$  is a fermion bilinear. The following shows that the Gaussian form is preserved by such jump operator  $L_m$ . First, the probability of the jump process is

$$\langle B | L_m^\dagger L_m | B \rangle = \langle B | d^\dagger d | B \rangle = \|d|B\rangle\|^2.$$

The action of annihilation operator  $d$  on  $|B\rangle$  is

$$d|B\rangle = \sum_k a_k^* c_k \prod_j \sum_i c_i^\dagger B_{ij} |0\rangle = \sum_j \langle a | B_j \rangle \bigotimes_{l \neq j} |B_l\rangle, \quad (80)$$

so we can obtain the probability

$$p_m = \sum_j |\langle a | B_j \rangle|^2 \gamma \Delta t. \quad (81)$$

Besides, we can utilize the gauge freedom to choose the basis such that  $\langle a | B'_j \rangle = 0$  for  $j > 1$ . The matrix  $B'$  always exists since we can always find a column  $j$  that  $\langle a | B_j \rangle \neq 0$  (otherwise, the probability of the jump is zero). We then move the column to the first and define the column as

$$|B'_j\rangle = |B_j\rangle - \frac{\langle a | B_j \rangle}{\langle a | B_1 \rangle} |B_1\rangle, \quad j > 1. \quad (82)$$

Note that such column transformation does not alter the linear space  $B$  spans, while the orthogonality and the normalization might be affected and should be renormalized afterward. Eq. (80) then simplified to:

$$d|B\rangle = \bigotimes_{j>1} |B'_j\rangle. \quad (83)$$

The result of the quantum jump is

$$L_m |B\rangle = |e^{ih} a\rangle \bigotimes_{j>1} |e^{ih} B'_j\rangle. \quad (84)$$

The representation of the outcome state is also not orthogonal. An additional QR decomposition is needed to convert it to canonical form.

## C Numerical simulations of the monitored free fermion dynamics

### C.1 Projective monitoring

We have demonstrated that a general Lindblad equation,

$$\frac{d}{dt} \rho = -i [H, \rho] + \gamma \sum_i L_i \rho L_i^\dagger - \frac{1}{2} \left\{ L_i^\dagger L_i, \rho \right\}, \quad (85)$$

where  $L_i$  are projectors, leads to a unique and homogeneous steady state (under open boundary conditions). In this section, we present numerical evidence supporting the homogeneity of late-time states for typical stochastic trajectories, described by:

$$d|\psi\rangle = \left[ -iH - \frac{\gamma}{2} \sum_m (L_m^\dagger L_m - \langle L_m^\dagger L_m \rangle) \right] |\psi\rangle dt + \sum_m \left( \frac{L_m}{\sqrt{\langle L_m^\dagger L_m \rangle}} - 1 \right) |\psi\rangle dW_m. \quad (86)$$

That is, the spatial homogeneity is not only at the density matrix level but also at the trajectory level. This suggests that the skin effect, which suppresses entanglement entropy, does not arise




Figure 6: (a) Density evolution under projective monitoring with  $\gamma = 0.1$  for a single trajectory. (b) Scaling of steady-state classical entropies. (c) Scaling of entanglement entropies. (d) Trajectory average of entanglement entropy for different measurement rates. (e) Trajectory average of mutual information  $I(A : B)$  between two  $L/8$  antipodal segments  $A$  and  $B$ . The crossing of  $I(A : B)$  indicates the critical measurement rate at  $\gamma_c \approx 0.2$ . (f) Trajectory average of classical entropy for systems in different sizes.

within the projective monitoring system. Therefore, introducing conditional feedback is essential to induce a dynamical skin effect.

In the following, we consider the model studied in the main text, of which the unitary evolution is generated by the Hamiltonian:

$$H = \sum_i (c_i^\dagger c_{i+1} + c_{i+1}^\dagger c_i) \quad (87)$$

and the monitoring is described by

$$L_i = \frac{1}{2} (c_i^\dagger - i c_{i+1}^\dagger) (c_i + i c_{i+1}). \quad (88)$$

First, Fig. 6a displays the particle density evolution of a typical trajectory. At late times, the system reaches a homogeneous state with only minor fluctuations. To further analyze the homogeneity of trajectories' densities, in Fig. 6b, we plot the trajectory-averaged classical entropy:

$$S_{\text{cl}} = - \sum_i [\langle n_i \rangle \log \langle n_i \rangle + (1 - \langle n_i \rangle) \log (1 - \langle n_i \rangle)]. \quad (89)$$

We observe a linear growth with system size, indicating a highly homogeneous density distribution. Moreover, when  $\gamma$  is small, the growth of the classical entropy approaches the saturation value

$$S_{\text{max}} = L \log 2, \quad (90)$$

which means that most trajectory has nearly homogeneous density distribution.

In contrast, Fig. 6c shows the entanglement entropy, which is considerably smaller than the classical entropy and exhibits an entanglement transition from log law to area law. These results suggest that the projective monitoring system under open boundary conditions may undergo a measurement-induced entanglement phase transition, similar to the periodic boundary condition.

## C.2 Generalized monitoring

Fig. 6d illustrates the trajectory averaging of entanglement entropy. The entanglement entropy exhibits a transition from log law to area law. We can pinpoint the critical measurement rate  $\gamma_c$  by doing the scaling analysis of the mutual information between two antipodal regions (as displayed in Fig. 6e).

In the main text, we have demonstrated trajectory averaging of classical entropy  $\overline{S_{\text{cl}}[\{\langle n_i \rangle\}]}$  provides an upper bound on entanglement entropy  $S_A$  and is itself bounded by classical entropy of averaged particle number:

$$\overline{S_{\text{cl}}[\{\langle n_i \rangle\}]} \leq S_{\text{cl}}[\{\overline{\langle n_i \rangle}\}] \quad (91)$$

In this section, we instead calculate  $\overline{S_{\text{cl}}[\{\langle n_i \rangle\}]}$  which imposes a tighter bound on the entanglement entropy. As displayed in Fig. 6f,  $\overline{S_{\text{cl}}[\{\langle n_i \rangle\}]}$  behaves similarly as  $S_{\text{cl}}[\{\overline{\langle n_i \rangle}\}]$  but with slightly different slop, indicating an asymptotic form:

$$\overline{S_{\text{cl}}[\{\langle n_i \rangle\}]} \simeq \frac{2.1}{\gamma^\nu} \quad (92)$$

where the scaling factor  $\nu \approx 1.13$  instead of 1 for  $S_{\text{cl}}[\{\overline{\langle n_i \rangle}\}]$ .

## C.3 Generalized monitoring with different conditional feedback

In this appendix, we provide more numerical results on the monitored free fermion system with Hamiltonian

$$H = \sum_i (c_i^\dagger c_{i+1} + c_{i+1}^\dagger c_i) \quad (93)$$




Figure 7: Steady-state classical entropies  $\overline{S_{\text{cl}}[\langle n_i \rangle]}$  for feedback  $U_i = \exp(i\theta n_{i+1})$ , where we chose  $\theta$  from  $\pi$  to  $0.3\pi$ . the data points all approach the asymptotic line  $\nu \log \gamma + \log S_{\text{cl}} = c'(\theta)$  where  $\nu \approx 1.13$  in the thermodynamic limit.

and the generalized monitoring characterized by the projective monitoring

$$P_i = \frac{1}{2}(c_i^\dagger - ic_{i+1}^\dagger)(c_i + ic_{i+1}) \quad (94)$$

followed by the conditional feedback  $U_i = e^{i\theta n_{i+1}}$  on site  $(i+1)$ .

In this section, we demonstrate numerically, that even under different parameters  $\theta$ 's apart from the  $\theta = \pi$ , the feedback-induced skin effect also appears. Therefore it is not a fin-tuned property of generalized monitoring. We will characterize the skin effect using the trajectory-averaged classical entropy:<sup>5</sup>

$$S_{\text{cl}}^\theta(\gamma, L) \equiv \overline{S_{\text{cl}}[\{\langle n_i \rangle\}]} \quad (95)$$

In Fig. 7, we show that The steady-state classical entropies in the thermodynamic limit  $L \rightarrow \infty$  all satisfy the general asymptotic behaviors:

$$S_{\text{cl}}^\theta(\gamma, L \rightarrow \infty) = \frac{c_\theta}{\gamma^\nu}, \quad (96)$$

where  $\nu \approx 1.13$ , while the constant  $c_\theta$  increases as  $\theta$  decreases:

$\theta/\pi$	1	0.9	0.8	0.7	0.6	0.5	0.4	0.3
$\log c_\theta$	0.75	0.85	0.95	1.1	1.4	1.9	2.6	3.4

As discussed in the main text, the constant  $c_\theta$  determines the spatial extents of the domain wall. The finite data point we obtained implies that  $c_\theta$  increases exponentially fast when  $\theta \rightarrow 0$ . In the  $\theta = 0$  case,  $c_\theta$  diverges and there is no skin effect.

The large-size domain walls impose significant overhead for numerical simulations. For small  $\theta$  ( $\leq 0.3\pi$ ), accessing the thermodynamic limit becomes practically hard.

Based on the numerical results, we speculate the scaling form (96) holds for different  $\theta$  and  $\gamma$ . Therefore we expect the skin effect to arise in a large class of generalized monitored systems.

<sup>5</sup>Note that in the main text, we use instead the classical entropy of the trajectory-averaged density. Here we use this quantity in order to find a smaller upper bound on the entanglement entropy.

## References

- [1] O. Alberton, M. Buchhold, and S. Diehl. Entanglement transition in a monitored free-fermion chain: From extended criticality to area law. *Phys. Rev. Lett.*, 126:170602, Apr 2021. DOI: [10.1103/PhysRevLett.126.170602](https://doi.org/10.1103/PhysRevLett.126.170602). URL <https://link.aps.org/doi/10.1103/PhysRevLett.126.170602>.
- [2] Yimu Bao, Soonwon Choi, and Ehud Altman. Theory of the phase transition in random unitary circuits with measurements. *Phys. Rev. B*, 101:104301, Mar 2020. DOI: [10.1103/PhysRevB.101.104301](https://doi.org/10.1103/PhysRevB.101.104301). URL <https://link.aps.org/doi/10.1103/PhysRevB.101.104301>.
- [3] Alberto Barchielli and Matteo Gregoratti. *The Stochastic Schrödinger Equation*, pages 11–49. Springer Berlin Heidelberg, Berlin, Heidelberg, 2009. ISBN 978-3-642-01298-3. DOI: [10.1007/978-3-642-01298-3\\_2](https://doi.org/10.1007/978-3-642-01298-3_2). URL [https://doi.org/10.1007/978-3-642-01298-3\\_2](https://doi.org/10.1007/978-3-642-01298-3_2).
- [4] William Berdanier, Jamir Marino, and Ehud Altman. Universal dynamics of stochastically driven quantum impurities. *Phys. Rev. Lett.*, 123:230604, Dec 2019. DOI: [10.1103/PhysRevLett.123.230604](https://doi.org/10.1103/PhysRevLett.123.230604). URL <https://link.aps.org/doi/10.1103/PhysRevLett.123.230604>.
- [5] Alberto Biella and Marco Schiró. Many-Body Quantum Zeno Effect and Measurement-Induced Subradiance Transition. *Quantum*, 5:528, August 2021. ISSN 2521-327X. DOI: [10.22331/q-2021-08-19-528](https://doi.org/10.22331/q-2021-08-19-528). URL <https://doi.org/10.22331/q-2021-08-19-528>.
- [6] Dan S. Borgnia, Alex Jura Kruchkov, and Robert-Jan Slager. Non-hermitian boundary modes and topology. *Phys. Rev. Lett.*, 124:056802, Feb 2020. DOI: [10.1103/PhysRevLett.124.056802](https://doi.org/10.1103/PhysRevLett.124.056802). URL <https://link.aps.org/doi/10.1103/PhysRevLett.124.056802>.
- [7] Sergey Bravyi. Lagrangian representation for fermionic linear optics. 2004. DOI: [10.48550/ARXIV.QUANT-PH/0404180](https://arxiv.org/abs/quant-ph/0404180). URL <https://arxiv.org/abs/quant-ph/0404180>.
- [8] Heinz-Peter Breuer and Francesco Petruccione. *The Theory of Open Quantum Systems*. Oxford University Press, 01 2007. ISBN 9780199213900. DOI: [10.1093/acprof:oso/9780199213900.001.0001](https://doi.org/10.1093/acprof:oso/9780199213900.001.0001). URL <https://doi.org/10.1093/acprof:oso/9780199213900.001.0001>.
- [9] M. Buchhold, Y. Minoguchi, A. Altland, and S. Diehl. Effective theory for the measurement-induced phase transition of dirac fermions. *Phys. Rev. X*, 11:041004, Oct 2021. DOI: [10.1103/PhysRevX.11.041004](https://doi.org/10.1103/PhysRevX.11.041004). URL <https://link.aps.org/doi/10.1103/PhysRevX.11.041004>.
- [10] M. Buchhold, T. Müller, and S. Diehl. Revealing measurement-induced phase transitions by pre-selection, 2022.
- [11] Xiangyu Cao, Antoine Tilloy, and Andrea De Luca. Entanglement in a fermion chain under continuous monitoring. *SciPost Phys.*, 7:024, 2019. DOI: [10.21468/SciPostPhys.7.2.024](https://doi.org/10.21468/SciPostPhys.7.2.024). URL <https://scipost.org/10.21468/SciPostPhys.7.2.024>.
- [12] Federico Carollo and Vincenzo Alba. Entangled multiplets and spreading of quantum correlations in a continuously monitored tight-binding chain. *Phys. Rev. B*, 106:L220304, Dec 2022. DOI: [10.1103/PhysRevB.106.L220304](https://doi.org/10.1103/PhysRevB.106.L220304). URL <https://link.aps.org/doi/10.1103/PhysRevB.106.L220304>.
- [13] Amos Chan, Andrea De Luca, and J. T. Chalker. Solution of a minimal model for many-body quantum chaos. *Phys. Rev. X*, 8:041019, Nov 2018. DOI: [10.1103/PhysRevX.8.041019](https://doi.org/10.1103/PhysRevX.8.041019). URL <https://link.aps.org/doi/10.1103/PhysRevX.8.041019>.
- [14] Amos Chan, Rahul M. Nandkishore, Michael Pretko, and Graeme Smith. Unitary-projective entanglement dynamics. *Phys. Rev. B*, 99:224307, Jun 2019. DOI: [10.1103/PhysRevB.99.224307](https://doi.org/10.1103/PhysRevB.99.224307). URL <https://link.aps.org/doi/10.1103/PhysRevB.99.224307>.
- [15] Xiao Chen, Yaodong Li, Matthew P. A. Fisher, and Andrew Lucas. Emergent conformal symmetry in nonunitary random dynamics of free fermions. *Phys. Rev. Research*, 2:033017, Jul 2020. DOI: [10.1103/PhysRevResearch.2.033017](https://doi.org/10.1103/PhysRevResearch.2.033017). URL <https://link.aps.org/doi/10.1103/PhysRevResearch.2.033017>.
- [16] Soonwon Choi, Yimu Bao, Xiao-Liang Qi, and Ehud Altman. Quantum error correction in scrambling dynamics and measurement-induced phase transition. *Phys. Rev. Lett.*, 125:030505, Jul 2020. DOI: [10.1103/PhysRevLett.125.030505](https://doi.org/10.1103/PhysRevLett.125.030505). URL <https://link.aps.org/doi/10.1103/PhysRevLett.125.030505>.



- [17] J. I. Cirac and P. Zoller. Quantum computations with cold trapped ions. *Phys. Rev. Lett.*, 74: 4091–4094, May 1995. DOI: [10.1103/PhysRevLett.74.4091](https://doi.org/10.1103/PhysRevLett.74.4091). URL <https://link.aps.org/doi/10.1103/PhysRevLett.74.4091>.
- [18] Michele Coppola, Emanuele Tirrito, Dragi Karevski, and Mario Collura. Growth of entanglement entropy under local projective measurements. *Phys. Rev. B*, 105:094303, Mar 2022. DOI: [10.1103/PhysRevB.105.094303](https://doi.org/10.1103/PhysRevB.105.094303). URL <https://link.aps.org/doi/10.1103/PhysRevB.105.094303>.
- [19] M.M. Wolf J.I. Cirac D. Perez-Garcia, F. Verstraete. Matrix product state representations, 2007.
- [20] Andrew J. Daley. Quantum trajectories and open many-body quantum systems. *Advances in Physics*, 63(2):77–149, 2014. DOI: [10.1080/00018732.2014.933502](https://doi.org/10.1080/00018732.2014.933502). URL <https://doi.org/10.1080/00018732.2014.933502>.
- [21] E. B. Davies. Markovian master equations. *Communications in Mathematical Physics*, 39(2): 91–110, 1974. DOI: [10.1007/BF01608389](https://doi.org/10.1007/BF01608389). URL <https://doi.org/10.1007/BF01608389>.
- [22] David E. Evans. Irreducible quantum dynamical semigroups. *Communications in Mathematical Physics*, 54(3):293–297, 1977. DOI: [10.1007/BF01614091](https://doi.org/10.1007/BF01614091). URL <https://doi.org/10.1007/BF01614091>.
- [23] M. Fannes, B. Nachtergaele, and R. F. Werner. Finitely correlated states on quantum spin chains. *Communications in Mathematical Physics*, 144(3):443–490, 1992. DOI: [10.1007/BF02099178](https://doi.org/10.1007/BF02099178). URL <https://doi.org/10.1007/BF02099178>.
- [24] Lukasz Fidkowski, Jeongwan Haah, and Matthew B. Hastings. How Dynamical Quantum Memories Forget. *Quantum*, 5:382, January 2021. ISSN 2521-327X. DOI: [10.22331/q-2021-01-17-382](https://doi.org/10.22331/q-2021-01-17-382). URL <https://doi.org/10.22331/q-2021-01-17-382>.
- [25] Matthew P. A. Fisher, Vedika Khemani, Adam Nahum, and Sagar Vijay. Random quantum circuits, 2022.
- [26] Matthew P. A. Fisher, Vedika Khemani, Adam Nahum, and Sagar Vijay. Random quantum circuits, 2022.
- [27] Matthew Fishman, Steven R. White, and E. Myles Stoudenmire. The ITensor software library for tensor network calculations, 2020.
- [28] Alberto Frigerio. Stationary states of quantum dynamical semigroups. *Communications in Mathematical Physics*, 63(3):269–276, 1978. DOI: [10.1007/BF01196936](https://doi.org/10.1007/BF01196936). URL <https://doi.org/10.1007/BF01196936>.
- [29] Yohei Fuji and Yuto Ashida. Measurement-induced quantum criticality under continuous monitoring. *Phys. Rev. B*, 102:054302, Aug 2020. DOI: [10.1103/PhysRevB.102.054302](https://doi.org/10.1103/PhysRevB.102.054302). URL <https://link.aps.org/doi/10.1103/PhysRevB.102.054302>.
- [30] Yohei Fuji and Yuto Ashida. Measurement-induced quantum criticality under continuous monitoring. *Phys. Rev. B*, 102:054302, Aug 2020. DOI: [10.1103/PhysRevB.102.054302](https://doi.org/10.1103/PhysRevB.102.054302). URL <https://link.aps.org/doi/10.1103/PhysRevB.102.054302>.
- [31] Sarang Gopalakrishnan and Michael J. Gullans. Entanglement and purification transitions in non-hermitian quantum mechanics. *Phys. Rev. Lett.*, 126:170503, Apr 2021. DOI: [10.1103/PhysRevLett.126.170503](https://doi.org/10.1103/PhysRevLett.126.170503). URL <https://link.aps.org/doi/10.1103/PhysRevLett.126.170503>.
- [32] Michael J. Gullans and David A. Huse. Scalable probes of measurement-induced criticality. *Phys. Rev. Lett.*, 125:070606, Aug 2020. DOI: [10.1103/PhysRevLett.125.070606](https://doi.org/10.1103/PhysRevLett.125.070606). URL <https://link.aps.org/doi/10.1103/PhysRevLett.125.070606>.
- [33] Michael J. Gullans and David A. Huse. Dynamical purification phase transition induced by quantum measurements. *Phys. Rev. X*, 10:041020, Oct 2020. DOI: [10.1103/PhysRevX.10.041020](https://doi.org/10.1103/PhysRevX.10.041020). URL <https://link.aps.org/doi/10.1103/PhysRevX.10.041020>.
- [34] Michael J. Gullans, Stefan Krastanov, David A. Huse, Liang Jiang, and Steven T. Flammia. Quantum coding with low-depth random circuits. *Phys. Rev. X*, 11:031066, Sep 2021. DOI: [10.1103/PhysRevX.11.031066](https://doi.org/10.1103/PhysRevX.11.031066). URL <https://link.aps.org/doi/10.1103/PhysRevX.11.031066>.
- [35] Naomichi Hatano and David R. Nelson. Localization transitions in non-hermitian quantum mechanics. *Phys. Rev. Lett.*, 77:570–573, Jul 1996. DOI: [10.1103/PhysRevLett.77.570](https://doi.org/10.1103/PhysRevLett.77.570). URL <https://link.aps.org/doi/10.1103/PhysRevLett.77.570>.

- [36] D. Hayes, D. N. Matsukevich, P. Maunz, D. Hucul, Q. Quraishi, S. Olmschenk, W. Campbell, J. Mizrahi, C. Senko, and C. Monroe. Entanglement of atomic qubits using an optical frequency comb. *Phys. Rev. Lett.*, 104:140501, Apr 2010. DOI: [10.1103/PhysRevLett.104.140501](https://doi.org/10.1103/PhysRevLett.104.140501). URL <https://link.aps.org/doi/10.1103/PhysRevLett.104.140501>.
- [37] Roger A. Horn and Charles R. Johnson. *Topics in Matrix Analysis*. Cambridge University Press, 1991. DOI: [10.1017/CBO9780511840371](https://doi.org/10.1017/CBO9780511840371).
- [38] Thomas Iadecola, Sriram Ganeshan, J. H. Pixley, and Justin H. Wilson. Dynamical entanglement transition in the probabilistic control of chaos, 2022.
- [39] Matteo Ippoliti and Vedika Khemani. Postselection-free entanglement dynamics via spacetime duality. *Phys. Rev. Lett.*, 126:060501, Feb 2021. DOI: [10.1103/PhysRevLett.126.060501](https://doi.org/10.1103/PhysRevLett.126.060501). URL <https://link.aps.org/doi/10.1103/PhysRevLett.126.060501>.
- [40] Matteo Ippoliti, Michael J. Gullans, Sarang Gopalakrishnan, David A. Huse, and Vedika Khemani. Entanglement phase transitions in measurement-only dynamics. *Phys. Rev. X*, 11:011030, Feb 2021. DOI: [10.1103/PhysRevX.11.011030](https://doi.org/10.1103/PhysRevX.11.011030). URL <https://link.aps.org/doi/10.1103/PhysRevX.11.011030>.
- [41] Matteo Ippoliti, Michael J. Gullans, Sarang Gopalakrishnan, David A. Huse, and Vedika Khemani. Entanglement phase transitions in measurement-only dynamics. *Phys. Rev. X*, 11:011030, Feb 2021. DOI: [10.1103/PhysRevX.11.011030](https://doi.org/10.1103/PhysRevX.11.011030). URL <https://link.aps.org/doi/10.1103/PhysRevX.11.011030>.
- [42] Matteo Ippoliti, Tibor Rakovszky, and Vedika Khemani. Fractal, logarithmic, and volume-law entangled nonthermal steady states via spacetime duality. *Phys. Rev. X*, 12:011045, Mar 2022. DOI: [10.1103/PhysRevX.12.011045](https://doi.org/10.1103/PhysRevX.12.011045). URL <https://link.aps.org/doi/10.1103/PhysRevX.12.011045>.
- [43] Kurt Jacobs and Daniel A. Steck. A straightforward introduction to continuous quantum measurement. *Contemporary Physics*, 47(5):279–303, 2006. DOI: [10.1080/00107510601101934](https://doi.org/10.1080/00107510601101934). URL <https://doi.org/10.1080/00107510601101934>.
- [44] Chao-Ming Jian, Yi-Zhuang You, Romain Vasseur, and Andreas W. W. Ludwig. Measurement-induced criticality in random quantum circuits. *Phys. Rev. B*, 101:104302, Mar 2020. DOI: [10.1103/PhysRevB.101.104302](https://doi.org/10.1103/PhysRevB.101.104302). URL <https://link.aps.org/doi/10.1103/PhysRevB.101.104302>.
- [45] Chao-Ming Jian, Bela Bauer, Anna Keselman, and Andreas W. W. Ludwig. Criticality and entanglement in non-unitary quantum circuits and tensor networks of non-interacting fermions, 2022.
- [46] Shao-Kai Jian, Zhi-Cheng Yang, Zhen Bi, and Xiao Chen. Yang-lee edge singularity triggered entanglement transition. *Phys. Rev. B*, 104:L161107, Oct 2021. DOI: [10.1103/PhysRevB.104.L161107](https://doi.org/10.1103/PhysRevB.104.L161107). URL <https://link.aps.org/doi/10.1103/PhysRevB.104.L161107>.
- [47] Kohei Kawabata, Tokiro Numasawa, and Shinsei Ryu. Entanglement phase transition induced by the non-hermitian skin effect, 2022.
- [48] Graham Kells, Dganit Meidan, and Alessandro Romito. Topological transitions in weakly monitored free fermions. *SciPost Phys.*, 14:031, 2023. DOI: [10.21468/SciPostPhys.14.3.031](https://doi.org/10.21468/SciPostPhys.14.3.031). URL <https://scipost.org/10.21468/SciPostPhys.14.3.031>.
- [49] B. Ladewig, S. Diehl, and M. Buchhold. Monitored open fermion dynamics: Exploring the interplay of measurement, decoherence, and free hamiltonian evolution. *Phys. Rev. Research*, 4:033001, Jul 2022. DOI: [10.1103/PhysRevResearch.4.033001](https://doi.org/10.1103/PhysRevResearch.4.033001). URL <https://link.aps.org/doi/10.1103/PhysRevResearch.4.033001>.
- [50] Ali Lavasani, Yahya Alavirad, and Maissam Barkeshli. Measurement-induced topological entanglement transitions in symmetric random quantum circuits. *Nature Physics*, 17(3):342–347, 2021. DOI: [10.1038/s41567-020-01112-z](https://doi.org/10.1038/s41567-020-01112-z). URL <https://doi.org/10.1038/s41567-020-01112-z>.
- [51] Yaodong Li and Matthew P. A. Fisher. Statistical mechanics of quantum error correcting codes. *Phys. Rev. B*, 103:104306, Mar 2021. DOI: [10.1103/PhysRevB.103.104306](https://doi.org/10.1103/PhysRevB.103.104306). URL <https://link.aps.org/doi/10.1103/PhysRevB.103.104306>.
- [52] Yaodong Li, Xiao Chen, and Matthew P. A. Fisher. Quantum zeno effect and the many-body entanglement transition. *Phys. Rev. B*, 98:205136, Nov 2018. DOI: [10.1103/PhysRevB.98.205136](https://doi.org/10.1103/PhysRevB.98.205136). URL <https://link.aps.org/doi/10.1103/PhysRevB.98.205136>.

- [53] Yaodong Li, Xiao Chen, and Matthew P. A. Fisher. Measurement-driven entanglement transition in hybrid quantum circuits. *Phys. Rev. B*, 100:134306, Oct 2019. DOI: [10.1103/PhysRevB.100.134306](https://doi.org/10.1103/PhysRevB.100.134306). URL <https://link.aps.org/doi/10.1103/PhysRevB.100.134306>.
- [54] Yaodong Li, Xiao Chen, Andreas W. W. Ludwig, and Matthew P. A. Fisher. Conformal invariance and quantum nonlocality in critical hybrid circuits. *Phys. Rev. B*, 104:104305, Sep 2021. DOI: [10.1103/PhysRevB.104.104305](https://doi.org/10.1103/PhysRevB.104.104305). URL <https://link.aps.org/doi/10.1103/PhysRevB.104.104305>.
- [55] Yaodong Li, Romain Vasseur, Matthew P. A. Fisher, and Andreas W. W. Ludwig. Statistical mechanics model for clifford random tensor networks and monitored quantum circuits, 2021. URL <https://arxiv.org/abs/2110.02988>.
- [56] G. Lindblad. On the generators of quantum dynamical semigroups. *Communications in Mathematical Physics*, 48(2):119–130, 1976. DOI: [10.1007/BF01608499](https://doi.org/10.1007/BF01608499). URL <https://doi.org/10.1007/BF01608499>.
- [57] Javier Lopez-Piqueres, Brayden Ware, and Romain Vasseur. Mean-field entanglement transitions in random tree tensor networks. *Phys. Rev. B*, 102:064202, Aug 2020. DOI: [10.1103/PhysRevB.102.064202](https://doi.org/10.1103/PhysRevB.102.064202). URL <https://link.aps.org/doi/10.1103/PhysRevB.102.064202>.
- [58] V. M. Martinez Alvarez, J. E. Barrios Vargas, and L. E. F. Foa Torres. Non-hermitian robust edge states in one dimension: Anomalous localization and eigenspace condensation at exceptional points. *Phys. Rev. B*, 97:121401(R), Mar 2018. DOI: [10.1103/PhysRevB.97.121401](https://doi.org/10.1103/PhysRevB.97.121401). URL <https://link.aps.org/doi/10.1103/PhysRevB.97.121401>.
- [59] Takaaki Minato, Koudai Sugimoto, Tomotaka Kuwahara, and Keiji Saito. Fate of measurement-induced phase transition in long-range interactions. *Phys. Rev. Lett.*, 128:010603, Jan 2022. DOI: [10.1103/PhysRevLett.128.010603](https://doi.org/10.1103/PhysRevLett.128.010603). URL <https://link.aps.org/doi/10.1103/PhysRevLett.128.010603>.
- [60] Klaus Mølmer and Anders Sørensen. Multiparticle entanglement of hot trapped ions. *Phys. Rev. Lett.*, 82:1835–1838, Mar 1999. DOI: [10.1103/PhysRevLett.82.1835](https://doi.org/10.1103/PhysRevLett.82.1835). URL <https://link.aps.org/doi/10.1103/PhysRevLett.82.1835>.
- [61] T. Müller, S. Diehl, and M. Buchhold. Measurement-induced dark state phase transitions in long-ranged fermion systems. *Phys. Rev. Lett.*, 128:010605, Jan 2022. DOI: [10.1103/PhysRevLett.128.010605](https://doi.org/10.1103/PhysRevLett.128.010605). URL <https://link.aps.org/doi/10.1103/PhysRevLett.128.010605>.
- [62] Adam Nahum, Sagar Vijay, and Jeongwan Haah. Operator spreading in random unitary circuits. *Phys. Rev. X*, 8:021014, Apr 2018. DOI: [10.1103/PhysRevX.8.021014](https://doi.org/10.1103/PhysRevX.8.021014). URL <https://link.aps.org/doi/10.1103/PhysRevX.8.021014>.
- [63] Adam Nahum, Sthitadhi Roy, Brian Skinner, and Jonathan Ruhman. Measurement and entanglement phase transitions in all-to-all quantum circuits, on quantum trees, and in landau-ginsburg theory. *PRX Quantum*, 2:010352, Mar 2021. DOI: [10.1103/PRXQuantum.2.010352](https://doi.org/10.1103/PRXQuantum.2.010352). URL <https://link.aps.org/doi/10.1103/PRXQuantum.2.010352>.
- [64] Michael A. Nielsen and Isaac L. Chuang. *Quantum Computation and Quantum Information: 10th Anniversary Edition*. Cambridge University Press, 2010. DOI: [10.1017/CBO9780511976667](https://doi.org/10.1017/CBO9780511976667).
- [65] Nobuyuki Okuma, Kohei Kawabata, Ken Shiozaki, and Masatoshi Sato. Topological origin of non-hermitian skin effects. *Phys. Rev. Lett.*, 124:086801, Feb 2020. DOI: [10.1103/PhysRevLett.124.086801](https://doi.org/10.1103/PhysRevLett.124.086801). URL <https://link.aps.org/doi/10.1103/PhysRevLett.124.086801>.
- [66] Wolfgang Paul. Electromagnetic traps for charged and neutral particles. *Rev. Mod. Phys.*, 62:531–540, Jul 1990. DOI: [10.1103/RevModPhys.62.531](https://doi.org/10.1103/RevModPhys.62.531). URL <https://link.aps.org/doi/10.1103/RevModPhys.62.531>.
- [67] D. Porras and J. I. Cirac. Effective quantum spin systems with trapped ions. *Phys. Rev. Lett.*, 92:207901, May 2004. DOI: [10.1103/PhysRevLett.92.207901](https://doi.org/10.1103/PhysRevLett.92.207901). URL <https://link.aps.org/doi/10.1103/PhysRevLett.92.207901>.
- [68] Tomaž Prosen. Matrix product solutions of boundary driven quantum chains. *Journal of Physics A: Mathematical and Theoretical*, 48(37):373001, aug 2015. DOI: [10.1088/1751-8113/48/37/373001](https://doi.org/10.1088/1751-8113/48/37/373001). URL <https://doi.org/10.1088/1751-8113/48/37/373001>.

- [69] Brian Skinner, Jonathan Ruhman, and Adam Nahum. Measurement-induced phase transitions in the dynamics of entanglement. *Phys. Rev. X*, 9:031009, Jul 2019. DOI: [10.1103/PhysRevX.9.031009](https://doi.org/10.1103/PhysRevX.9.031009). URL <https://link.aps.org/doi/10.1103/PhysRevX.9.031009>.
- [70] Herbert Spohn. Kinetic equations from hamiltonian dynamics: Markovian limits. *Rev. Mod. Phys.*, 52:569–615, Jul 1980. DOI: [10.1103/RevModPhys.52.569](https://doi.org/10.1103/RevModPhys.52.569). URL <https://link.aps.org/doi/10.1103/RevModPhys.52.569>.
- [71] Luis E F Foa Torres. Perspective on topological states of non-hermitian lattices. *Journal of Physics: Materials*, 3(1):014002, nov 2019. DOI: [10.1088/2515-7639/ab4092](https://doi.org/10.1088/2515-7639/ab4092). URL <https://doi.org/10.1088/2515-7639/ab4092>.
- [72] Xhek Turkeshi and Marco Schiró. Entanglement and correlation spreading in non-hermitian spin chains. *Phys. Rev. B*, 107:L020403, Jan 2023. DOI: [10.1103/PhysRevB.107.L020403](https://doi.org/10.1103/PhysRevB.107.L020403). URL <https://link.aps.org/doi/10.1103/PhysRevB.107.L020403>.
- [73] Xhek Turkeshi, Alberto Biella, Rosario Fazio, Marcello Dalmonte, and Marco Schiró. Measurement-induced entanglement transitions in the quantum ising chain: From infinite to zero clicks. *Phys. Rev. B*, 103:224210, Jun 2021. DOI: [10.1103/PhysRevB.103.224210](https://doi.org/10.1103/PhysRevB.103.224210). URL <https://link.aps.org/doi/10.1103/PhysRevB.103.224210>.
- [74] Xhek Turkeshi, Marcello Dalmonte, Rosario Fazio, and Marco Schiró. Entanglement transitions from stochastic resetting of non-hermitian quasiparticles. *Phys. Rev. B*, 105:L241114, Jun 2022. DOI: [10.1103/PhysRevB.105.L241114](https://doi.org/10.1103/PhysRevB.105.L241114). URL <https://link.aps.org/doi/10.1103/PhysRevB.105.L241114>.
- [75] Xhek Turkeshi, Lorenzo Piroli, and Marco Schiró. Enhanced entanglement negativity in boundary-driven monitored fermionic chains. *Phys. Rev. B*, 106:024304, Jul 2022. DOI: [10.1103/PhysRevB.106.024304](https://doi.org/10.1103/PhysRevB.106.024304). URL <https://link.aps.org/doi/10.1103/PhysRevB.106.024304>.
- [76] Romain Vasseur, Andrew C. Potter, Yi-Zhuang You, and Andreas W. W. Ludwig. Entanglement transitions from holographic random tensor networks. *Phys. Rev. B*, 100:134203, Oct 2019. DOI: [10.1103/PhysRevB.100.134203](https://doi.org/10.1103/PhysRevB.100.134203). URL <https://link.aps.org/doi/10.1103/PhysRevB.100.134203>.
- [77] C. W. von Keyserlingk, Tibor Rakovszky, Frank Pollmann, and S. L. Sondhi. Operator hydrodynamics, otocs, and entanglement growth in systems without conservation laws. *Phys. Rev. X*, 8:021013, Apr 2018. DOI: [10.1103/PhysRevX.8.021013](https://doi.org/10.1103/PhysRevX.8.021013). URL <https://link.aps.org/doi/10.1103/PhysRevX.8.021013>.
- [78] Mark M. Wilde. *Quantum Information Theory*. Cambridge University Press, 2013. DOI: [10.1017/CBO9781139525343](https://doi.org/10.1017/CBO9781139525343).
- [79] Howard M. Wiseman and Gerard J. Milburn. *Quantum Measurement and Control*. Cambridge University Press, 2009. DOI: [10.1017/CBO9780511813948](https://doi.org/10.1017/CBO9780511813948).
- [80] Zhesen Yang, Kai Zhang, Chen Fang, and Jiangping Hu. Non-hermitian bulk-boundary correspondence and auxiliary generalized brillouin zone theory. *Phys. Rev. Lett.*, 125:226402, Nov 2020. DOI: [10.1103/PhysRevLett.125.226402](https://doi.org/10.1103/PhysRevLett.125.226402). URL <https://link.aps.org/doi/10.1103/PhysRevLett.125.226402>.
- [81] Zhi-Cheng Yang, Yaodong Li, Matthew P. A. Fisher, and Xiao Chen. Entanglement phase transitions in random stabilizer tensor networks. *Phys. Rev. B*, 105:104306, Mar 2022. DOI: [10.1103/PhysRevB.105.104306](https://doi.org/10.1103/PhysRevB.105.104306). URL <https://link.aps.org/doi/10.1103/PhysRevB.105.104306>.
- [82] Shunyu Yao and Zhong Wang. Edge states and topological invariants of non-hermitian systems. *Phys. Rev. Lett.*, 121:086803, Aug 2018. DOI: [10.1103/PhysRevLett.121.086803](https://doi.org/10.1103/PhysRevLett.121.086803). URL <https://link.aps.org/doi/10.1103/PhysRevLett.121.086803>.
- [83] Kazuki Yokomizo and Shuichi Murakami. Non-bloch band theory of non-hermitian systems. *Phys. Rev. Lett.*, 123:066404, Aug 2019. DOI: [10.1103/PhysRevLett.123.066404](https://doi.org/10.1103/PhysRevLett.123.066404). URL <https://link.aps.org/doi/10.1103/PhysRevLett.123.066404>.
- [84] Aidan Zabalo, Michael J. Gullans, Justin H. Wilson, Sarang Gopalakrishnan, David A. Huse, and J. H. Pixley. Critical properties of the measurement-induced transition in random quantum circuits. *Phys. Rev. B*, 101:060301(R), Feb 2020. DOI: [10.1103/PhysRevB.101.060301](https://doi.org/10.1103/PhysRevB.101.060301). URL <https://link.aps.org/doi/10.1103/PhysRevB.101.060301>.
- [85] Kai Zhang, Zhesen Yang, and Chen Fang. Correspondence between winding numbers and skin modes in non-hermitian systems. *Phys. Rev. Lett.*, 125:126402, Sep

2020. DOI: [10.1103/PhysRevLett.125.126402](https://doi.org/10.1103/PhysRevLett.125.126402). URL <https://link.aps.org/doi/10.1103/PhysRevLett.125.126402>.
- [86] Tianci Zhou and Adam Nahum. Emergent statistical mechanics of entanglement in random unitary circuits. *Phys. Rev. B*, 99:174205, May 2019. DOI: [10.1103/PhysRevB.99.174205](https://doi.org/10.1103/PhysRevB.99.174205). URL <https://link.aps.org/doi/10.1103/PhysRevB.99.174205>.
- [87] Shi-Liang Zhu, C. Monroe, and L.-M. Duan. Arbitrary-speed quantum gates within large ion crystals through minimum control of laser beams. *Europhysics Letters*, 73(4):485, jan 2006. DOI: [10.1209/epl/i2005-10424-4](https://doi.org/10.1209/epl/i2005-10424-4). URL <https://dx.doi.org/10.1209/epl/i2005-10424-4>.