

Prethermalization and the local robustness of gapped systems

Chao Yin^{*} and Andrew Lucas[†]

*Department of Physics and Center for Theory of Quantum Matter,
University of Colorado, Boulder, CO 80309, USA*

(Dated: September 26, 2022)

We prove that prethermalization is a generic property of gapped local many-body quantum systems, subjected to small perturbations, in any spatial dimension. More precisely, let H_0 be a Hamiltonian, spatially local in d spatial dimensions, with a gap Δ in the many-body spectrum; let V be a spatially local Hamiltonian consisting of a sum of local terms, each of which is bounded by $\epsilon \ll \Delta$. Then, the approximation that quantum dynamics is restricted to the low-energy subspace of H_0 is accurate, in the correlation functions of local operators, for stretched exponential time scale $\tau \sim \exp[(\Delta/\epsilon)^a]$ for any $a < 1/(2d - 1)$. This result does not depend on whether the perturbation closes the gap. It significantly extends previous rigorous results on prethermalization in models where H_0 had an integer-valued spectrum. We infer the robustness of quantum simulation in low-energy subspaces, the existence of “scarring” (strongly athermal correlation functions) in gapped systems subject to generic perturbations, and the robustness of quantum information in non-frustration-free gapped phases with topological order.

Introduction.— Consider a many-body quantum Hamiltonian H_0 , assumed to be spatially local in d spatial dimensions. Suppose that we can exactly solve H_0 – or at least, learn something interesting about its spectrum or properties. Now, consider a perturbing Hamiltonian V , made out of a sum of local terms, each of bounded norm ϵ . As long as we take the thermodynamic limit *before* sending $\epsilon \rightarrow 0$, general lore states that a perturbation ($\epsilon > 0$) has drastic qualitative effects. For example, the orthogonality catastrophe shows that eigenstates are extraordinarily sensitive to perturbations [1]. A general integrable system generally exhibits a complete rearrangement of the many-body spectrum, transitioning from Poisson ($\epsilon = 0$) to Wigner-Dyson ($\epsilon \neq 0$) energy-level statistics [2, 3]. Only rare families of many-body quantum systems, such as many-body localized phases [4–9], seem to avoid the fate of many-body chaos and thermalization.

However, certain many-body quantum systems are known to undergo *prethermalization*: although at infinite time they do thermalize, the time scale associated to thermalization (as measured by physical observables, i.e. local correlation functions) may be exponentially long:

$$\tau_{\text{thermalization}} \sim \exp\left[\left(\frac{\Delta}{\epsilon}\right)^a\right] \quad (1)$$

where Δ is some intrinsic energy scale of H_0 , and $a > 0$. This was argued for in the context of the Hubbard model in the presence of large repulsive energies [10, 11]: although two particles on the same site (called a doublon) store enormous energy and should thermalize into a sea of mobile excitations by separating, there is no local perturbation that can satisfiably do this! The reason is intuitively as follows: the doublon stores energy Δ , but any

single no-doublon excitation carries at most ϵ kinetic energy. One must go to order Δ/ϵ in perturbation theory to observe the many-body resonance whereby a doublon can split apart while conserving energy: this implies (1). Only in the last few years was this intuition put on rigorous ground [12, 13].

Existing proofs of prethermalization in the Hubbard model rely fundamentally on peculiar aspects of the problem. The “unperturbed” H_0 consists exclusively of the repulsive potential energy – it is a sum of local operators which: (1) act on a single lattice site, (2) mutually commute, and (3) have an “integer spectrum”, such that the many-body spectrum of H_0 is of the form $0, \Delta, 2\Delta, \dots$. The “perturbation” V is then the kinetic (hopping) terms allowing for particle motion. While prethermalization proofs have also been extended to Floquet and other non-Hamiltonian settings [14–19] with various experimental verifications [20–25], the latter two requirements (which lead to exact solvability, among other useful features) have remained.

At the same time, one may be surprised on physical grounds by this state of affairs: the intuition for prethermalization does *not* rely on solvability of H_0 , nor even a discrete spectrum in the thermodynamic limit. In fact, it should suffice to simply say that if Δ is a many-body spectral gap of H_0 , and any local perturbation can add energy at most $\epsilon \ll \Delta$, then one has to go to order Δ/ϵ in perturbation theory to witness a many-body resonance wherein a system, prepared on one side of the gap of H_0 , can “decay” into a state on the other side. In other words, if a many-body gap protects a subspace of Hilbert space in H_0 , this should be robust against local perturbations for a long time: prethermalization, captured by (1), should hold.

Indeed, such generalizations are known in special cases. The stability of gapped phases with topological order has been proven for frustration-free Hamiltonians [26, 27], which paves the way for topological quantum computing [28, 29] and quantum memory [30, 31] at zero tempera-

^{*} chao.yin@colorado.edu

[†] andrew.j.lucas@colorado.edu

ture. Here the topological order hides information in the ground states nonlocally, so the gap does not close under small perturbations, and $\tau_{\text{thermalization}} \rightarrow \infty$. However, the gap can close rapidly in generic gapped systems, like the Hubbard model, and (1) is the optimal bound one can expect.

Main Result.— The purpose of this letter is to prove that prethermalization is a property of all spatially local gapped systems. Let H_0 and V to be local many-body Hamiltonians on a d -dimensional lattice Λ : e.g.

$$V = \sum_{S \subset \Lambda, S \text{ local}} V_S, \quad (2)$$

where V_S acts non-trivially on the degrees of freedom on sites in the geometrically local S , and trivially elsewhere, and $\|V_S\| \leq \epsilon$. H_0 has a similarly local structure, and we require the existence of a “spectral gap” of size Δ , wherein the many-body Hilbert space \mathcal{H} can be decomposed into $\mathcal{H} = \mathcal{H}_< \oplus \mathcal{H}_>$, where $\mathcal{H}_<$ contains eigenvectors of eigenvalue at most E_* , while $\mathcal{H}_>$ contains eigenvectors of eigenvalue at least $E_* + \Delta$. Here and below, precise definitions are relegated to the Supplementary Material (SM).

In the SM, we prove that for sufficiently small ϵ/Δ , there exists a unitary U , generated by finite-time evolution with a quasi-local Hamiltonian protocol with terms of strength ϵ , such that the Hamiltonian

$$U^\dagger(H_0 + V)U = H_* + V_*, \quad (3)$$

where H_* is block-diagonal in $\mathcal{H}_< \oplus \mathcal{H}_>$ (i.e. protects the low/high energy subspaces) while V_* is a sum of local terms of strength

$$\|(V_*)_S\| \lesssim \epsilon \exp \left[- \left(\frac{\Delta}{\epsilon} \right)^a \right], \quad \text{for any } a < \frac{1}{2d-1}. \quad (4)$$

(This a is likely not tight for $d > 1$: see SM.) This proves that there is prethermalization: a subspace $U\mathcal{H}_<$ of $H_0 + V$ is protected for a stretched exponentially long time scale (1). Local (few-body) operators $B \approx U^\dagger B U$, so this prethermal dynamics will be visible in physically accessible correlation functions.

Numerical Demonstration.— To showcase our result, consider the interacting $d = 1$ spin model

$$H_0 = \sum_{i=1}^{N-1} (Z_i Z_{i+1} + J_x X_i X_{i+1}) + h \sum_{i=1}^N X_i, \quad (5a)$$

$$V = Z = \epsilon \sum_{i=1}^N Z_i, \quad (5b)$$

where $h = 0.9$, $J_x = 0.37$. If $J_x = 0$, H_0 is the transverse-field Ising model with two ferromagnetic ground states, separated from the excited states by a gap $2(1-h) = 0.2$ in the thermodynamic limit [32]. The J_x term is added to break the integrability of H_0 , and we show in Fig. 1(a) that H_0 is still gapped within the ferromagnetic phase,




FIG. 1. (a) Spectrum of $H = H_0 + V$ in (5a) and (5b) at $N = 14$ (blue lines), where the lowest 60 eigenstates are shown with the ground state energy set to 0. For the lowest 3 eigenstates, data for $N = 10, 12$ are also shown by solid lines of different colors, indicating the gap closes at $\epsilon \sim 1/N$. (b) Polarization $\langle Z \rangle / N$ of evolving the initial state ψ_\uparrow with H for three values of ϵ , whose average energy E_\uparrow in H is denoted by dots in (a). The green line $\epsilon = 0$ dropped a bit, because ψ_\uparrow is superposition of the two ground states of H_0 that are not exactly degenerate for this finite size $N = 14$. (c) Weight of eigenstates of H , $|\langle E | \psi_\uparrow \rangle|^2$, as a function of energy around E_\uparrow for $N = 14, \epsilon = 0.2$. The color for each eigenstate $|E\rangle$ indicates $|\langle E | \psi_\uparrow \rangle|^2$: $|\psi_\uparrow\rangle$ is supported mainly by the three atypical states that have $\langle Z \rangle > 0$.

for system sizes accessible with exact diagonalization. However, this gap is extremely sensitive to V : the ground state $|\psi_\uparrow\rangle$ of H_0 that has polarization $\langle Z_i \rangle > 0$ quickly merges into the excitation spectrum when $\epsilon \sim N^{-1}$. As H_0 neither is commuting/frustration-free nor has integer spectrum or topological order, previous bounds could not predict prethermal dynamics in this model. Yet for finite small ϵ and $N = 14$, we see clear non-thermal behavior in $|\psi(t)\rangle = e^{-i(H_0+V)t}|\psi_\uparrow\rangle$ in Fig. 1(b). At $\epsilon = 0.1$, the polarization $\langle Z \rangle / N$ remains positive and large. While it oscillates around 0 for some time at $\epsilon = 0.2$, there is still a revival behavior $\langle Z \rangle / N \approx 0.6$ at $t \approx 180$. In sharp contrast, typical eigenstates of H around the average energy $E_\uparrow = \langle \psi_\uparrow | H | \psi_\uparrow \rangle$ (shown as the black dot in Fig. 1(a)) have $\langle Z \rangle < 0$ as shown in Fig. 1(c), since the perturbation V strongly prefers negative polarization in the lower half spectrum.

This numerical example highlights a crucial aspect of our theorem: it is valid *even if V closes the gap between $\mathcal{H}_<$ and $\mathcal{H}_>$* . Among the eigenstates shown in Fig. 1(c), there are only three atypical ones with $\langle Z \rangle > 0$, on which ψ_\uparrow has large support. Such eigenstates can be viewed as atypical “scars” in the spectrum.

Applications of our Result.— Before sketching the proof of this result, let us first highlight its implications. One immediate consequence is the generic robustness of quantum simulation of low-energy – often *constrained* – quantum dynamics in the presence of realistic experi-

mental perturbations. Namely, one often begins with a Hamiltonian $H = H_0$ which has a highly degenerate (yet non-trivially structured) ground state subspace, and then applies a small perturbation V which attempts to induce dynamics among the low-energy states of H_0 . Then it is crucial that the state remains in the low-energy subspace up to long time, even if V , or some experimental imperfections, have matrix elements acting across the gap. Such a construction has been of particular recent interest in the literature on quantum scars [33–44]. In minimal theoretical models of scarring, H_0 often is frustration-free and has an integer spectrum; our proof above demonstrates neither of these assumptions are necessary over a (stretched) exponentially long simulation time.

Moreover, our result even implies a certain *universality* to “scarring” phenomena. Consider an arbitrary Hamiltonian H_0 with a many-body spectral gap. Regardless of whether or not $H = H_0 + V$ remains gapped, our results demonstrate that on prethermally long time scales, correlation functions and entanglement entropies that probe local dynamics on one side of the gap of H_0 will largely stay decoupled from those on the other side, regardless of where they end up in the spectrum of H . This was seen explicitly in Fig. 1, where the non-thermal, atypical state $|\psi_\uparrow\rangle$ had energy deep in the continuous spectrum of H . We suggest that this phenomenon may arise more generally when adding explicit symmetry breaking V to H_0 which exhibits spontaneous symmetry breaking. But even more broadly, a notion of “scarring” is generic: one *always* finds states whose dynamics, as measured in local correlators, are extremely far from thermal under $H_0 + V$, even if $H_0 + V$ is chaotic (as measured, e.g., in level spacing). Hence, the process of thermalization (and the time scales after which eigenstate thermalization hypothesis [45–48] can hold) are guaranteed to be extraordinarily slow in all perturbations of gapped systems.

Perhaps the most commonly studied spectral gaps in many-body systems arise in gapped phases of matter, where the ground states are separated by a finite gap Δ in the spectrum from any excited state. For example, in a topological phase of matter, there are exactly degenerate ground states [49], which may serve as a robust logical qubit. Our prethermalization proof implies that such a logical qubit will always remain protected in a low-dimensional subspace (up to the local rotation by U) for extraordinarily long time scales in the presence of perturbations. This work thus provides an interesting generalization of earlier results [26, 27, 30, 50] which proved the robustness of topological order in frustration-free Hamiltonians. However, unlike the frustration-free limit, we were not able to prove the infinite-time robustness of quantum information; our work “only” holds over an stretched exponentially long time scale. In practice, the decoherence of an experimental device may be far more dangerous than the perturbation itself to a qubit.

Of course, our result does *not* guarantee that it is easy to access information in the form rotated by U with local operations. Let X_L and Z_L denote logical Paulis for a

qubit protected in the low energy subspace of H_0 . We might expect information is protected in an accessible format with a fidelity of F whenever

$$1 - F > \|X_L - U^\dagger X_L U\|, \quad (6)$$

with a similar equation for Z_L . If X_L and Z_L are spatially local, then indeed the same theorems that assure the locality of H_* and V_* ensure the right hand side of (6) is small. Then, the logicals of H_0 and $H_0 + V$ can’t act too differently in the low energy subspace (indeed, on any state). However, in most codes (e.g. the surface code) intended for near-term use, logical qubits are non-local and act on L qubits – the number of physical qubits that encode a logical cannot be too small in order to ensure it can be error-corrected. Then we anticipate the fidelity vanishes in the thermodynamic limit, as

$$\|X_L - U^\dagger X_L U\| \sim L \frac{\epsilon}{\Delta}. \quad (7)$$

Designing a logical qubit such that $L\epsilon \ll \Delta$ will ensure that, even if H_0 is not frustration-free, the logical qubit can be operated on with high fidelity for prethermally long time scales.

A somewhat similar application of our result arises in $SU(2)$ -symmetric quantum spin models, where states in the Dicke manifold (maximal S^2 subspace) can readily form squeezed states [51] of metrological value [52]. When the Dicke manifold is protected by a spectral gap (as arises in realistic models), our work demonstrates that this protection of squeezed states is robust on exponentially long time scales in the presence of inevitable perturbations (subject to an analogous caveat about whether the state is recoverable or not). Of course, many practical atomic physics experiments have long-range (power-law) interactions [53], which currently lie beyond the scope of our proof. It will be important in future work to understand whether our conclusions can be extended to this setting. Nevertheless, our work provides a crucial first step towards understanding the protection of squeezed states in more generic models.

Prethermalization in Commuting Models.— We now begin to sketch the proof of our main result (the formal proof is in the SM). Before doing so, it is useful to sketch how one proves prethermalization in the Hubbard model [13]: our proof mirrors this earlier one, albeit with substantial technical complications. In the section that follows, we assume that in the decomposition $H_0 + V$, we know that H_0 has an integer spectrum $0, \Delta, 2\Delta, \dots$ ($\Delta = U$ in the Hubbard model’s standard notation).

Suppose for the moment that V was so small that $\|V\| \ll \Delta$. In this case we would know exactly that $H_0 + V$ had a gap (it has many, but let us focus on just one of them), and moreover we could use first order perturbation theory to explicitly rotate the eigenstates $|n\rangle$ of H_0 . If V has no terms connecting degenerate states, then the rotated eigenstates would be

$$|n\rangle_1 = |n\rangle + \sum_{m \neq n} \frac{\langle m|V|n\rangle}{E_n - E_m} |m\rangle; \quad (8)$$

in particular,

$$\| |n\rangle_1 - |n\rangle \| \lesssim \frac{\epsilon}{\Delta} \quad (9)$$

Higher orders in perturbation theory are tedious but straightforward, and (9) holds for the exact all-order eigenstates $|n\rangle_{H_0+V}$. Unfortunately this series is badly behaved once we restrict to more realistic assumptions that only each local term in V is bounded by ϵ – the reason is that $\|V\| \sim \epsilon N$ now diverges with N , the number of lattice sites. Yet we have good reason to think this divergence is “artificial” – it is intuitively a consequence of the fact that (in the Hubbard model) the product state $|n\rangle$ picks up $\sim N$ local rotations by angle ϵ/Δ , each in a different direction in a high-dimensional space. Altogether, they add up to a huge unitary rotation $U \sim U_{(1)} \cdots U_{(N)}$ on states, where each $U_{(j)}$ acts only near site j . On the other hand, local operators should depend only on a small number of these rotations; if we instead rotate H_0 , the rotated $U^\dagger H_0 U$ will term-by-term look close to H_0 .

This intuition is made precise by the Schrieffer-Wolff transformations [54, 55], which proceed as follows. First, we project V onto terms acting within ($\mathbb{P}V$) and between $((1 - \mathbb{P})V)$ an eigenspace of H_0 via the transformation

$$\mathbb{P}V = \frac{\Delta}{2\pi} \int_0^{2\pi/\Delta} dt e^{iH_0 t} V e^{-iH_0 t}. \quad (10)$$

We then define

$$D_1 = \mathbb{P}V, \quad W_1 = (1 - \mathbb{P})V. \quad (11)$$

The first order unitary rotation $U_1 = e^{A_1}$ where

$$[A_1, H_0] = -W_1. \quad (12)$$

Explicit calculation shows that $e^{A_1}(H_0 + V)e^{-A_1}$ is block diagonal in the eigenspaces of H_0 up to a $O(\epsilon^2)$ piece, which we call V_2 .

This process can be iterated many times. We define $D_k = \mathbb{P}V_k$, $W_k = (1 - \mathbb{P})V_k$, $[A_k, H_0] = -W_k$, and $U_k = e^{-A_k}$. The non-trivial aspect of this iteration is to show that A_k is not too non-local (essentially, to show that the dressing of a fixed energy subspace is mild, hence there is prethermalization and not simply large rotations of an eigenspace). This is guaranteed by locality of A_k , which is proved using the Lieb-Robinson Theorem [56] (locality of operator dynamics in many-body systems). At the end of the day, one essentially discovers

$$\frac{\|V_{k+1}\|_{\text{local}}}{\|V_k\|_{\text{local}}} \sim k \frac{\epsilon}{\Delta} \quad (13)$$

where $\|A\|_{\text{local}}$ bounds the operator norm of terms that act non-trivially on any one fixed site. Hence, this iterative process stops being helpful when $k_* \sim \Delta/\epsilon$. We conclude that the quasi-local unitary $U = U_1 \cdots U_{k_*}$ rotates $H_0 + V$ to $H_* + V_*$ of the form (3), with $a = 1$.

Extending to General Models.—We now describe how to extend this proof to a generic $H_0 + V$. The first crucial feature of the prior proof is the ability to “rotate away” the off-resonant terms in the perturbation V : in this setting, terms in V that connect states across the gap in H_0 . Put another way, we need to find the projection \mathbb{P} and show that it takes local operators to local operators.

Let $|n\rangle$ formally denote the eigenstate(s) of H_0 with energy E_n , and write $V_{nm} = \langle n|V|m\rangle$. We don’t try to calculate $|n\rangle$ or V_{nm} ; nevertheless, its existence is valuable. Formally, we know the Heisenberg evolution

$$V(t) = \sum_{n,m} V_{nm} e^{i(E_n - E_m)t} |n\rangle \langle m|. \quad (14)$$

If $w(t)$ is a real-valued function whose Fourier transform is $\hat{w}(\omega)$, we can consider a projection \mathbb{P} of the form

$$\begin{aligned} \mathbb{P}V &= \int_{-\infty}^{\infty} dt w(t) V(t) \\ &= \sum_{n,m} \hat{w}(E_n - E_m) V_{nm} |n\rangle \langle m|. \end{aligned} \quad (15)$$

If we can find a function where $\hat{w}(\omega) = 0$ if $|\omega| \geq \Delta$, this transformation can project out the off-resonant terms in V . Such functions are known [57, 58], and have asymptotic decay $w(t) \sim e^{-|t|/\ln^2|t|}$ at large t . This decay is fast enough that we know, from the Lieb-Robinson bounds, that $\mathbb{P}V$ is a relatively local operator.

A bit more work allows us to show that the analogues of A_k, D_k, V_k, W_k are all quasi-local operators: terms that act non-trivially in a domain of size r have a small norm $\sim \exp[-r^\alpha]$ at large r for $\alpha = 1 - \delta$ (with $\delta \rightarrow 0$). $\delta \neq 0$ is a technical nuisance which is required due to the subexponential decay in $w(t)$. While more care is required to control the locality of the iterative Schrieffer-Wolff transformations at higher order, the procedure follows what we described before in principle. A technical remark is that we require somewhat better Lieb-Robinson bounds than the standard ones [56] to bound the growth of A_k, D_k, \dots : we use methods inspired by the equivalence class construction of [59]. After a lot of algebra, we show that at the k^{th} order of the transformation, the operators in D_k, W_k are constrained to predominantly act within balls of radius

$$R_k \lesssim (\ln k)^{1/\alpha}, \quad (16)$$

while the perturbation

$$\frac{\|V_{k+1}\|_{\text{local}}}{\|V_k\|_{\text{local}}} \lesssim \frac{\epsilon}{\Delta} k^{(2d-1)/(2\alpha-1)}. \quad (17)$$

Hence we must truncate the expansion at

$$k_* = \left(\frac{\Delta}{\epsilon} \right)^{(2\alpha-1)/(2d-1)}. \quad (18)$$

Ultimately, we obtain a rotated Hamiltonian of the form (3), whose perturbation V_* is exponentially suppressed. For any local operator B , we find that

$$\|U^\dagger e^{iHt} B e^{-iHt} U - e^{iH_*t} U^\dagger B U e^{-iH_*t}\| \leq \epsilon t^{d+1} e^{-k_*}. \quad (19)$$

Namely, there exists a mild quasi-local rotation of (sums of) local operators such that the genuine dynamics of operators (and correlation functions, etc.) appear to be restricted to the high/low-energy subspaces of H_0 for the prethermal time scale (1). This completes (the sketch of) our proof that prethermalization is a generic feature of any perturbed gapped model.

Outlook.—In this letter, we have proved that the prethermalization of doublons in the Hubbard model is but one manifestation of a universal phenomenon, whereby distinct sectors of a gapped Hamiltonian H_0 remain protected for (stretched) exponentially long times in the presence of local perturbations V . This prethermalization phenomenon, in all measurable local correla-

tion functions, is generic to any perturbation of a gapped system.

This result demonstrates that it is always reasonable to simulate quantum dynamics generated by V in constrained models, so long as one studies $H_0 + V$, where H_0 's ground state manifold is the constrained subspace of interest, and H_0 has a large spectral gap Δ . Even if $H_0 + V$ is gapless and chaotic, the (locally rotated) ground states of H_0 serve as effective “scar states” which will exhibit athermal dynamics for extraordinarily long times. We anticipate that this observation will have practical implications for the preparation of interesting entangled states on the Dicke manifold in future atomic physics experiments, and for the ease of recovering qubits under imperfect local encoding.

Acknowledgements.— This work was supported by a Research Fellowship from the Alfred P. Sloan Foundation under Grant FG-2020-13795 (AL) and by the U.S. Air Force Office of Scientific Research under Grant FA9550-21-1-0195 (CY, AL).

Supplementary Material

1. PRELIMINARIES

In this section we review a few mathematical facts, and precisely state our assumptions about the models we study.

1.1. Models of interest

We consider many-body quantum systems defined on a (finite) d -dimensional “lattice”, with vertex set Λ . Let $\mathbf{d} : \Lambda \times \Lambda \rightarrow \mathbb{Z}^+$ denote the Manhattan distance between two vertices in Λ . Note that $\mathbf{d}(i, j) = 0$ if and only if $i = j$, while two vertices are defined to be neighbors if $\mathbf{d}(i, j) = 1$. The diameter of a subset $S \subseteq \Lambda$, denoted $\text{diam}(S)$, is defined as

$$\text{diam}(S) = \max_{i, j \in S} \mathbf{d}(i, j). \quad (\text{S1})$$

Similarly, the boundary of a set S is defined precisely as

$$\partial S = \{i \in S : \text{there exists } j \notin S \text{ with } \mathbf{d}(i, j) = 1\}. \quad (\text{S2})$$

Although we will typically refer to Λ as a lattice, we do not require it to have an translation symmetry (automorphism subgroup isomorphic to \mathbb{Z}^d). Instead, we require that there exists a finite constant c_d such that for any $S \subseteq \Lambda$

$$|\partial S| \leq c_d \cdot (1 + \text{diam}S)^{d-1}, \quad \text{and} \quad |S| \leq c_d \cdot (1 + \text{diam}S)^d. \quad (\text{S3})$$

We will implicitly be interested in the regime where $|\Lambda| \rightarrow \infty$.

We associate to each vertex in Λ a q -dimensional “qudit”, such that the global Hilbert space is (on a finite lattice) $\mathcal{H} = (\mathbb{C}^q)^\Lambda$. We consider Hamiltonian

$$H = H_0 + V_1, \quad (\text{S4})$$

where H_0 and V_1 are both spatially local operators on Λ , in the sense that there exists constants $B, \kappa_0, \epsilon_0 > 0$ such that we may write

$$H_0 = B \sum_{S \subseteq \Lambda} e^{-\kappa_0 \text{diam}(S)} H_{0,S}, \quad V_1 = \epsilon_0 \sum_{S \subseteq \Lambda} e^{-\kappa_0 \text{diam}(S)} V_{1,S}, \quad (\text{S5})$$

where we assume that $\|H_{0,S}\|, \|V_{1,S}\| \leq 1$, with the operator norm here the standard infinity norm (maximal singular value), and $H_{0,S}$ and $V_{1,S}$ operators that act non-trivially only on sites in S . We do not require that $H_{0,S}$ acts non-trivially on all sites contained within S . We assume that the spectrum of H_0 has a non-trivial gap Δ , so that the many-body Hilbert space \mathcal{H} can be decomposed into $\mathcal{H} = \mathcal{H}_< \oplus \mathcal{H}_>$, where $\mathcal{H}_<$ contains eigenvectors of eigenvalue at most E_* , while $\mathcal{H}_>$ contains eigenvectors of eigenvalue at least $E_* + \Delta$.

The perturbation is weak in the sense that ϵ_0/Δ will be small – we postpone precise definition of how small to (S15). In fact, we can even slightly relax the requirements on H_0 and V_1 from above, though for practical models the above should suffice. (Models of interest not captured by the above assumptions, such as those with power-law interactions, are not within the scope of our proof.)

1.2. Superimposing a simplicial lattice

We have not specified the lattice Λ beyond requiring it being d -dimensional in (S3). However, to prove our main results, more information about Λ is needed to conveniently organize the support of operators. As a result, we fix the specific lattice by assuming that Λ is the d -dimensional simplicial lattice defined as follows (see e.g. [60]).¹ Starting from an auxiliary $(d+1)$ -dimensional hypercubic lattice with orthogonal basis $\mathbf{e}_1, \dots, \mathbf{e}_{d+1}$, define a redundant basis

$$\mathcal{E}_p = \mathbf{e}_p - \frac{\mathbf{e}_1 + \dots + \mathbf{e}_{d+1}}{d+1}, \quad p = 1, \dots, d+1, \quad (\text{S6})$$

¹ The strategy of our proof also works for other lattices, e.g., the hypercubic lattice. However, there will be more terms to keep track of when decomposing an evolved operator, so we stick with the simplicial lattice with as few terms as possible.




FIG. S1. A sketch of superimposing a simplicial lattice Λ (black solid lines) on the original lattice Λ_0 of qudits (green dots connected by blue dashed lines) in 2d. All qudits are moved to their nearest lattice site of the triangular lattice, as shown by red dashed lines. Some sites have $m > 1$ qudits, where we combine them to a “ mq -dit”. While some sites may have $m = 0$ qudits, this is not a problem since it is equivalent to having one qudit on those sites that does not interact with the rest of the system. Given the locality of the original Hamiltonian, the “ mq -dits” in Λ also only interact with their neighbors. We will consider simplices $S \subseteq \Lambda$ of fixed orientation. Namely, we say S is a simplex if it is like the magenta triangles; it can not be the cyan triangle.

that satisfies

$$\mathcal{E}_1 + \cdots + \mathcal{E}_{d+1} = 0. \quad (\text{S7})$$

All lattice points of the form $\sum_p n_p \mathbf{e}_p = \sum_p n_p \mathcal{E}_p$ with constraint $n_1 + \cdots + n_{p+1} = 0$, then lie on the d -dimensional hyperplane $\mathbf{x}_1 + \cdots + \mathbf{x}_{d+1} = 0$, and form the d -dimensional simplicial lattice. In a nutshell, each group of $d + 1$ nearest sites in the simplicial lattice, serve as the vertices of the d -dimensional regular simplex that they form. As examples, the 2d simplicial lattice is the triangular lattice, while the 3d simplicial lattice is the fcc lattice made of regular tetrahedrons.

From now on, we focus on the simplicial lattice that automatically satisfies (S3), with c_d determined by d . This is not a big restriction, since a model on an arbitrary d -dimensional lattice Λ_0 can be transformed into one on the simplicial lattice Λ as follows. One can superimpose a simplicial lattice on top of the original lattice, and move all qudits to their nearest simplicial lattice site (as measured by Euclidean distance in \mathbb{R}^d). See Fig. S1 for a sketch. A site in Λ will contain at most $O(1)$ qudits. If a site contains $m > 1$ qudits, combine them to form an “ mq -dit”: a single degree of freedom with mq -dimensional Hilbert space. Furthermore, the original Hamiltonian satisfying (S5), remains at least as local in the new simplicial lattice, since grouping sites together cannot increase Manhattan distance between (possibly now grouped) sites. Finally, all results that we prove for the new simplicial lattice, can be transformed back to the original Λ_0 . Any book-keeping factors that arise during this process will be $O(1)$ and not affect any main results.

We say a subset $S \subseteq \Lambda$ is a simplex, if there are $d + 1$ sites $i_1, \cdots, i_{d+1} \in S$, such that they are the vertices forming a d -dimensional regular simplex, and that S is exactly all sites in Λ contained in that regular simplex. Moreover, we only consider simplices S of fixed orientation, namely there are $d + 1$ fixed vectors $\mathbf{E}_1, \cdots, \mathbf{E}_{d+1}$, such that any simplex S have them as the normal vectors (pointing outwards) of its $d + 1$ faces. In Fig. S1 for example, the magenta triangles are simplices we consider, while the cyan one is not. We will use the following geometric fact: (see Fig. S4(b) as an illustration)

Proposition 1. *Let $S_0, S \subseteq \Lambda$ be two simplices with fixed direction, such that $S_0 \subseteq S$. Let f_1, \dots, f_{d+1} be the faces of S . Then*

$$\sum_{p=1}^{d+1} d(S_0, f_p) = \text{diam}S - \text{diam}S_0. \quad (\text{S8})$$

Proof. Consider the process that grows the faces f'_1, \dots, f'_{d+1} of S_0 one by one to coincide with S . First, suppose the opposite vertex of f'_1 in S_0 is $i_1 \in S_0$. We first grow f'_1 to f_1 in the sense that fixing i_1 , while enlarging all the edges of S_0 connecting i_1 to reach f_1 . Then after this first step, we get a new simplex $S_1 \subseteq S$ that has i_1 as a vertex, and its opposite face overlapping with f_1 . During this step, the edge enlarges by a length exactly $d(S_0, f_1)$:

$$d(S_0, f_1) = \text{diam}S_1 - \text{diam}S_0, \quad (\text{S9})$$

because the distance is measured by the Manhattan distance on the underlying simplicial lattice, not a Euclidean metric in \mathbb{R}^d . At the second step, we grow S_1 to S_2 in a similar way to reach f_2 , with relation

$$d(S_0, f_2) = d(S_1, f_2) = \text{diam}S_2 - \text{diam}S_1. \quad (\text{S10})$$

Iterating this, we get an equation like above at each step up to the final $(d+1)$ -th step, so that their summation produces (S8), because

$$\text{diam}S - \text{diam}S_0 = \text{diam}S_{d+1} - \text{diam}S_d + \text{diam}S_d - \text{diam}S_{d-1} + \dots + \text{diam}S_1 - \text{diam}S_0, \quad (\text{S11})$$

with $S = S_{d+1}$. \square

1.3. The κ -norm of an operator

For an extensive operator \mathcal{O} , there exist (many) local decompositions

$$\mathcal{O} = \sum_S \mathcal{O}_S, \quad (\text{S12})$$

where $S \subseteq \Lambda$ is always a simplex, and \mathcal{O}_S is supported inside S . We do not require \mathcal{O}_S acts nontrivially on the boundary of S , and the decomposition is not unique. However, there always exists an “optimal decomposition” where we assign terms in \mathcal{O} to the smallest possible simplex S . We quantify this by defining the (α, κ) -norm of \mathcal{O} as

$$\|\mathcal{O}\|_{\alpha, \kappa} := \inf_{\{\mathcal{O}_S\}} \max_{i \in \Lambda} \sum_{S \ni i} e^{\kappa(\text{diam}S)^\alpha} \|\mathcal{O}_S\|, \quad (\text{S13})$$

where $\inf_{\{\mathcal{O}_S\}}$ is the infimum over all local decompositions. The parameters α, κ are both non-negative. We will choose α as a fixed parameter that is close to 1 from below, and we will just call “ κ -norm” and use notation $\|\cdot\|_\kappa$, with α being implicit. We will always assume that we can choose a “best” decomposition $\mathcal{O} = \sum_S \mathcal{O}_S$ that realizes its κ -norm:

$$\sum_{S \ni i} e^{\kappa(\text{diam}S)^\alpha} \|\mathcal{O}_S\| = \|\mathcal{O}\|_\kappa. \quad (\text{S14})$$

Strictly speaking, this should be viewed as choosing a decomposition that is δ -close to $\|\mathcal{O}\|_\kappa$ (which is provably possible), and taking $\delta \rightarrow 0$ in the end, a mathematical annoyance that does not affect the structure of the proofs that follow.

The perturbation V_1 is weak in the sense that

$$\epsilon := \|V_1\|_{\alpha, \kappa_1} \ll \Delta, \quad (\text{S15})$$

where $\alpha < 1$, κ_1 is some order 1 constant, while Δ is a spectral gap of H_0 . While formally Δ can be any energy scale, our prethermalization bound seems most profound when it corresponds to a gap, as we will then prove that there is a notion of prethermalization – dynamics is (for long times) approximately governed by a gapped Hamiltonian, even if the true Hamiltonian is no longer gapped.

1.4. The Lieb-Robinson bound

Define the Liouvillian superoperator \mathcal{L}_0 by

$$\mathcal{L}_0 \mathcal{O} := i[H_0, \mathcal{O}], \quad \forall \mathcal{O}. \quad (\text{S16})$$

We assume H_0 contains local interactions, as defined by (S5). Then, the following Lieb-Robinson bound holds:

Proposition 2. *There exists constants μ', μ, u such that*

$$\| [e^{t\mathcal{L}_0} \mathcal{O}_S, \mathcal{O}'_{S'}] \| \leq 2 \| \mathcal{O}_S \| \| \mathcal{O}'_{S'} \| \mu' \min(|\partial S|, |\partial S'|) e^{\mu(u|t| - d(S, S'))}, \quad (\text{S17})$$

for any pair of local operators $\mathcal{O}_S, \mathcal{O}'_{S'}$ that do not overlap: $S \cap S' = \emptyset$.

This bound is slightly stronger than many commonly stated Lieb-Robinson bounds in the literature. We do not present an explicit proof here, as it can be shown following analogous methods to those we employ later in the proof of Proposition 14.

2. MAIN RESULTS

Now we summarize our main results and describe formally a few applications sketched in the main text.

2.1. Main theorem

For a given H_0 , we say an operator \mathcal{O} is Δ -diagonal, if

$$\langle E | \mathcal{O} | E' \rangle = 0, \quad (\text{S18})$$

for any pair of eigenstates $|E\rangle, |E'\rangle$ of H_0 that has energy difference $|E - E'| \geq \Delta$. We do not need to know what these eigenstates are explicitly to gain value from this definition: if we know \mathcal{O} is Δ -diagonal and H_0 has a many-body gap of size Δ in the spectrum, then \mathcal{O} is block-diagonal between $\mathcal{H}_<$ and $\mathcal{H}_>$. With this in mind, we present our main theorem:

Theorem 3. *For the Hamiltonian (S4) defined on the d -dimensional simplicial lattice, suppose H_0 has Lieb-Robinson bound (S17) (which defines parameter μ). For any $\alpha \in (0, 1)$ and*

$$\kappa_1 \leq \frac{\mu}{5}, \quad (\text{S19})$$

there exist constants c_*, c_V, c_A, c_D, c'_V determined by $\alpha, d, \mu, \mu', \kappa_1$ and the ratio u/Δ , that achieve the following:

Define

$$k_* = \left\lfloor c_* (\Delta/\epsilon)^{\frac{2\alpha-1}{2d-1}} \right\rfloor, \quad (\text{S20})$$

and

$$\kappa_k = \frac{\kappa_1}{1 + \ln k}. \quad (\text{S21})$$

For any small perturbation V_1 with

$$\|V_1\|_{\kappa_1} = \epsilon \leq c_V \Delta, \quad (\text{S22})$$

there exists a quasi-local unitary $U = e^{A_1} \dots e^{A_{k_*-1}}$ with

$$\|A_k\|_{\kappa_k} \leq c_A \epsilon 2^{-k}, \quad k = 1, \dots, k_* - 1, \quad (\text{S23})$$

that approximately block-diagonalizes $H = H_0 + V_1$:

$$U^\dagger H U = H_0 + D_* + V_*, \quad (\text{S24})$$

where D_* is Δ -diagonal with respect to H_0 . (Thus, $H_0 + D_*$ is also block-diagonal with respect to $\mathcal{H}_<$ and $\mathcal{H}_>$.) Furthermore, the local norms are bounded by

$$\|D_*\|_{\kappa_*} \leq c_D \epsilon, \quad (\text{S25a})$$

$$\|V_*\|_{\kappa_*} \leq c'_V \epsilon 2^{-k_*}, \quad (\text{S25b})$$

where $\kappa_* = \kappa_{k_*}$.

Because of (S23), the unitary transformation U rotates the Hilbert space slightly, in the sense that locality in the rotated frame is, at zeroth order of ϵ , the same as that in the original frame. Moreover, the above Theorem implies that in this locally rotated frame, local dynamics is to a high accuracy generated by a dressed Hamiltonian $H_0 + D_*$, until the prethermal time

$$t_* \sim 1/\|V_*\|_{\kappa_*} \sim \exp\left[(\Delta/\epsilon)^{\frac{2\alpha-1}{2d-1}}\right], \quad (\text{S26})$$

when V_* starts to play a role. The optimal choice of α is then $\alpha \rightarrow 1$. Before t_* , $H_0 + D_*$ preserves any gapped subspace of H_0 that has gap larger than Δ away from the complement spectrum. These heuristic arguments are formalized in the following Corollary:²

Corollary 4. *Following Theorem 3, there exist constants c_U, c_{oper} determined by $\alpha, d, \mu, \mu', \kappa_1$ and the ratio u/Δ , such that following statements hold.*

1. **Locality of U :** *For any local operator $\mathcal{O} = \mathcal{O}_S$ supported in a connected set S , $U^\dagger \mathcal{O} U$ is quasi-local and close to \mathcal{O} in the sense that*

$$U^\dagger \mathcal{O} U = \mathcal{O} + \sum_{r=0}^{\infty} \mathcal{O}_r, \quad (\text{S27})$$

where \mathcal{O}_r is supported in $B(S, r) = \{i \in \Lambda : d(i, S) \leq r\}$ [we demand each term in \mathcal{O}_r is not supported in $B(S, r-1)$], and decays rapidly with r :

$$\|\mathcal{O}_r\| \leq c_U \epsilon |S| \|\mathcal{O}\| e^{-\kappa_* r^\alpha}. \quad (\text{S28})$$

2. **Local operator dynamics is approximately generated by $H_0 + D_*$ up to an exponentially long time t_* :** *For any local operator $\mathcal{O} = \mathcal{O}_S$,*

$$\left\| e^{itH} \mathcal{O} e^{-itH} - U e^{it(H_0+D_*)} U^\dagger \mathcal{O} U e^{-it(H_0+D_*)} U^\dagger \right\| \leq c_{\text{oper}} \epsilon |S| (|t|+1)^{d+1} 2^{-k_*} \|\mathcal{O}\|. \quad (\text{S29})$$

3. **Gapped subspaces of H_0 are locally preserved up to t_* :** *Suppose the initial density matrix ρ_0 is of the form*

$$\rho_0 = U \tilde{\rho}_0 U^\dagger, \quad (\text{S30})$$

where $\tilde{\rho}_0$ is supported inside the gapped subspace $\mathcal{H}_<$ of H_0 that has gap Δ to the complement spectrum $\mathcal{H}_>$. Define the reduced density matrix on set S after time evolution

$$\rho_S(t) := \text{Tr}_{S^c} (e^{-itH} \rho_0 e^{itH}), \quad (\text{S31})$$

where partial trace is taken on S^c , the complement of S . Further define another reduced density matrix

$$\rho'_S(t) := \text{Tr}_{S^c} (U e^{-it(H_0+D_*)} U^\dagger \rho_0 U e^{-it(H_0+D_*)} U^\dagger) = \text{Tr}_{S^c} (U e^{-it(H_0+D_*)} \tilde{\rho}_0 e^{-it(H_0+D_*)} U^\dagger), \quad (\text{S32})$$

as reference, which stays in the gapped subspace $\mathcal{H}_<$ up to rotation by U . Then $\rho_S(t)$ is close to $\rho'_S(t)$ in trace norm

$$\|\rho_S(t) - \rho'_S(t)\|_1 \leq c_{\text{oper}} \epsilon |S| (|t|+1)^{d+1} 2^{-k_*}. \quad (\text{S33})$$

² This is related to Theorem 3.1-3.3 in [13].

2.2. Proof of Theorem 3

Proof of Theorem 3. We prove by iteration. Suppose at the k -th step, we have rotated H by $U_{k-1} = e^{A_1} \dots e^{A_{k-1}}$. We write

$$U_{k-1}^\dagger H U_{k-1} = H_0 + D_k + V_k, \quad (\text{S34})$$

where D_k is Δ -diagonal. For example at $k = 1$, we have $U_0 = 1$ and $D_1 = 0$. To go to step $(k + 1)$, we further rotate

$$U_k^\dagger H U_k = e^{-\mathcal{A}_k} (U_{k-1}^\dagger H U_{k-1}) = e^{-\mathcal{A}_k} (H_0 + D_k + V_k) = H_0 + D_{k+1} + V_{k+1}, \quad (\text{S35})$$

where the superoperator \mathcal{A}_k is defined similar to (S16):

$$\mathcal{A}_k = [A_k, \cdot], \quad (\text{S36})$$

so that $U_k = U_{k-1} e^{A_k}$. We choose A_k using the following Proposition, which is proved in Section 3.

Proposition 5. *For a fixed H_0 satisfying the Lieb-Robinson bound (S17), there exist superoperators \mathbb{P}, \mathbb{A} , defined by*

$$\mathbb{P}\mathcal{O} = \int_{-\infty}^{\infty} dt w(t) e^{t\mathcal{L}_0} \mathcal{O}, \quad (\text{S37})$$

$$\mathbb{A}\mathcal{O} = i \int_{-\infty}^{\infty} dt W(t) e^{t\mathcal{L}_0} \mathcal{O}, \quad (\text{S38})$$

with function $w(t), W(t)$ determined by Δ , such that for any operator \mathcal{O} , $\mathbb{P}\mathcal{O}$ is Δ -diagonal, and

$$[H_0, \mathbb{A}\mathcal{O}] + \mathcal{O} = \mathbb{P}\mathcal{O}. \quad (\text{S39})$$

Moreover, let $\kappa' < \kappa \leq \kappa_1$ with κ_1 satisfying (S19), and define $\delta\kappa = \kappa - \kappa'$. Then

$$\|\mathbb{P}\mathcal{O}\|_{\kappa'} \leq c_w [\max(1, -\ln \delta\kappa)]^{d-1} \|\mathcal{O}\|_{\kappa}, \quad (\text{S40a})$$

$$\|\mathbb{A}\mathcal{O}\|_{\kappa'} \leq \frac{c_W}{\Delta} [\max(1, -\ln \delta\kappa)]^{d-1} \|\mathcal{O}\|_{\kappa}, \quad (\text{S40b})$$

where c_w and c_W only depend on $\alpha, d, \kappa_1, \mu, \mu'$ and the ratio u/Δ .

We then choose

$$A_k = \mathbb{A}V_k, \quad (\text{S41})$$

to satisfy

$$[H_0, A_k] + V_k = \mathbb{P}V_k. \quad (\text{S42})$$

As a result, we assign

$$D_{k+1} = D_k + \mathbb{P}V_k, \quad (\text{S43})$$

that is still Δ -diagonal, and determine V_{k+1} from (S35):

$$\begin{aligned} V_{k+1} &= e^{-\mathcal{A}_k} (H_0 + D_k + V_k) - H_0 - (D_k + \mathbb{P}V_k) \\ &= (e^{-\mathcal{A}_k} - 1)H_0 + (1 - \mathbb{P})V_k + (e^{-\mathcal{A}_k} - 1)(D_k + V_k) \\ &= \int_0^1 ds (e^{-s\mathcal{A}_k} - 1) (\mathbb{P} - 1)V_k + (e^{-\mathcal{A}_k} - 1)(D_k + V_k), \end{aligned} \quad (\text{S44})$$

where we have plugged (S43) in the first line. In the third line, we have used

$$e^{-\mathcal{A}_k} - 1 = - \int_0^1 ds e^{-s\mathcal{A}_k} \mathcal{A}_k, \quad (\text{S45})$$

and (S42).

Now we calculate the local norms using the following Proposition, which generalizes Lemma 4.1 of [13]. The proof of these results is quite tedious and is postponed to Section 4.

Proposition 6. *If*

$$\mathcal{A} = [A, \cdot], \quad (\text{S46})$$

with operator A satisfying

$$\|A\|_{\kappa} \leq (\delta\kappa)^{\frac{3d+1}{\alpha}}, \quad (\text{S47})$$

with $\delta\kappa < \kappa \leq \kappa_1$, then

$$\|e^{-A}\mathcal{O} - \mathcal{O}\|_{\kappa'} \leq c_-(\delta\kappa)^{-\frac{2d-1}{\alpha}} \|A\|_{\kappa} \|\mathcal{O}\|_{\kappa}, \quad (\text{S48})$$

where $\kappa' = \kappa - \delta\kappa$, and c_- depends on α, d and κ_1 .

Define

$$a_k = \|A_k\|_{\kappa_k}, \quad d_k = \|D_k\|_{\kappa_k}, \quad v_k = \|V_k\|_{\kappa_k}, \quad \tilde{v}_k = c_w [\max(1, -\ln \delta\kappa_{k+1})]^{d-1} v_k, \quad (\text{S49})$$

where the decreasing sequence κ_k is defined in (S21), and

$$\delta\kappa_k := \kappa_{k-1} - \kappa_k = \frac{\kappa_1 \ln[1 + 1/(k-1)]}{(1 + \ln k)[1 + \ln(k-1)]} \geq \frac{\kappa_1'}{k \ln^2 k} \quad (\text{S50})$$

implies that

$$\max(1, -\ln \delta\kappa_k) \leq c_\delta \ln k, \quad (\text{S51})$$

for some constants κ_1' and c_δ determined by κ_1 . This choice (S21) of κ_k makes $\delta\kappa_k \sim 1/k \ln^2 k$ decay about as slowly as possible while keeping $\kappa_k > 0$. Using (S40a),(S40b) and (S48), the iterative definitions (S41),(S43) and (S44) lead to iterations of the local bounds

$$a_{k-1} \leq \frac{c_w \tilde{v}_{k-1}}{c_w \Delta}, \quad (\text{S52a})$$

$$d_k \leq d_{k-1} + \tilde{v}_{k-1}, \quad (\text{S52b})$$

$$v_k \leq c_-(\delta\kappa_k)^{-\frac{2d-1}{\alpha}} a_{k-1} \left[\frac{1}{2} (v_{k-1} + \tilde{v}_{k-1}) + d_{k-1} + v_{k-1} \right] \leq c_-(\delta\kappa_k)^{-\frac{2d-1}{\alpha}} a_{k-1} (d_{k-1} + \tilde{c}_w \tilde{v}_{k-1}), \quad (\text{S52c})$$

where we have shifted $k \rightarrow k-1$, and used $\tilde{v}_{k-1} \geq c_w v_{k-1}$ and $\tilde{c}_w = (c_w + 3)/(2c_w)$. Plugging (S52a) into (S52c) and combining constants using (S51), we get the iteration for \tilde{v}_k

$$\tilde{v}_k \leq \frac{\tilde{c}_w}{\Delta} (k \ln^2 k)^{\frac{2d-1}{\alpha}} (\ln k)^{d-1} \tilde{v}_{k-1} (d_{k-1} + \tilde{c}_w \tilde{v}_{k-1}) \leq \frac{c_v}{\Delta} k^{\frac{2d-1}{2\alpha-1}} \tilde{v}_{k-1} (d_{k-1} + \tilde{c}_w \tilde{v}_{k-1}), \quad (\text{S53})$$

where we have replaced a power function of $\ln k$ by a power function of k : $k^{\frac{2d-1}{2\alpha-1} - \frac{2d-1}{\alpha}}$, with the price of adjusting the prefactor. (S52b) and (S53) comprises the closed iteration for d_k and \tilde{v}_k , assuming the condition (S47) which transforms to

$$\tilde{v}_{k-1} \leq c_a (k \ln^2 k)^{\frac{3d+1}{\alpha}} \Delta, \quad (\text{S54})$$

with constant c_a . We will later verify this condition can be achieved.

For sufficiently small ϵ/Δ , d_k keeps at the order of $\epsilon = v_1$, and (S53) leads to $\tilde{v}_k \sim d_k \tilde{v}_{k-1}/\Delta \sim v_1^k/\Delta^{k-1}$. The iteration continues up to $k_* \sim (\Delta/v_1)^{\frac{2\alpha-1}{2d-1}}$ when the power of k in (S53) dominates, and the iteration terminates. In this process (S54) is guaranteed to hold, since \tilde{v}_k decays exponentially while the right hand side of (S54) decays as a power law. To make the above arguments rigorous, we assume (S54) and

$$\tilde{v}_k \leq \tilde{v}_1/2^{k-1}, \quad (\text{S55})$$

holds for all steps $k = 1, \dots, k'$ before $k' + 1$. Then (S52b) yields

$$d_{k'} \leq c_w \sum_{k=1}^{k'-1} \tilde{v}_k \leq c_w \tilde{v}_1 \frac{1}{1 - 1/2} = 2c_w \tilde{v}_1. \quad (\text{S56})$$

If (S54) also holds at $k = k' + 1$, then (S53) yields

$$\tilde{v}_{k'+1} \leq \frac{c_v}{\Delta} (k' + 1)^{\frac{2d-1}{2\alpha-1}} \tilde{v}_{k'} (2c_w + \tilde{c}_w) \tilde{v}_1. \quad (\text{S57})$$

The right hand side is bounded by $\tilde{v}_{k'}/2$, as long as

$$\frac{c_v}{\Delta} (k' + 1)^{\frac{2d-1}{2\alpha-1}} (2c_w + \tilde{c}_w) \tilde{v}_1 \leq \frac{1}{2} \quad (\text{S58})$$

which holds so long as

$$k' + 1 \leq k_* := \left\lfloor \left(\frac{\Delta}{c_v (2c_w + \tilde{c}_w) \tilde{v}_1} \right)^{\frac{2\alpha-1}{2d-1}} \right\rfloor \equiv \left\lfloor c_* \left(\frac{\Delta}{v_1} \right)^{\frac{2\alpha-1}{2d-1}} \right\rfloor, \quad (\text{S59})$$

where c_* is determined by c_v, c_w and κ_1 . Thus (S55) also holds for the next step $k' + 1$, as long as $k' + 1 \leq k_*$ and (S54) holds for $k = k' + 1$.

Finally, we verify (S54) using our assumption (S22) with a sufficiently small constant c_V : From (S55), it suffices to prove

$$\tilde{v}_1 2^{2-k'} \leq c_a (k' \ln^2 k')^{\frac{3d+1}{\alpha}} \Delta, \quad (\text{S60})$$

for all $k' \geq 2$, which indeed holds given (S22) with

$$c_V = \frac{c_a}{c_w [\max(1, -\ln \delta \kappa_2)]^{d-1}} \min_{k' \geq 2} 2^{k'-2} (k' \ln^2 k')^{\frac{3d+1}{\alpha}} > 0. \quad (\text{S61})$$

To summarize, if (S22) and (S61) hold, (S55) holds iteratively up to $k = k_*$, which further leads to (S23) by (S52a). At the final step, define $D_* = D_{k_*}$ that is Δ -diagonal and $V_* = V_{k_*}$, which then satisfy (S24) and (S25a). \square

2.3. Proof of Corollary 4

Proof of Corollary 4. We prove the three statements one by one, where latter proof relies on previous results.

1. Rewrite

$$U = \mathcal{T} \exp \left[\int_0^1 ds A(s) \right], \quad (\text{S62})$$

where \mathcal{T} is time-ordering, and

$$A(s) = \begin{cases} 0, & s < 2^{1-k_*} \\ 2^k A_k, & 2^{-k} \leq s < 2^{1-k}, \quad (k = 1, \dots, k_* - 1) \end{cases} \quad (\text{S63})$$

Then (S23) leads to

$$\|A(s)\|_{\kappa_*} \leq c_A \epsilon, \quad \forall s. \quad (\text{S64})$$

$U^\dagger \mathcal{O} U$ is the Heisenberg evolution under the time-dependent Hamiltonian $iA(s)$. However, we will use notation $U = e^A$ for simplicity, with the time-ordering being implicit. To determine the decomposition (S27), we first define

$$\mathcal{O}_0 := e^{-A|_S} \mathcal{O} e^{A|_S} - \mathcal{O} = - \int_0^1 ds e^{-sA|_S} [A|_S, \mathcal{O}] e^{sA|_S}, \quad (\text{S65})$$

which is similar to (S45). Here $A|_S := \sum_{S' \subseteq S} A_{S'}$, with the optimal decomposition $A_{S'}$ that realizes $\|A\|_{\kappa_*}$. \mathcal{O}_0 is indeed bounded by (S28):

$$\begin{aligned} \|\mathcal{O}_0\| &\leq \int_0^1 ds \left\| e^{-sA|_S} [A|_S, \mathcal{O}] e^{sA|_S} \right\| = \int_0^1 ds \|[A|_S, \mathcal{O}]\| \leq 2 \|\mathcal{O}\| \|A|_S\| \\ &\leq 2 \|\mathcal{O}\| \sum_{j \in S} \sum_{S' \ni j} \|A_{S'}\| \leq 2 \|\mathcal{O}\| |S| \|A\|_{\kappa_*} \leq 2c_A \epsilon |S| \|\mathcal{O}\|, \end{aligned} \quad (\text{S66})$$

where we used A is anti-Hermitian in the second line. In the second line of (S66), we have used the fact that each term contained in $A|_S$ must have one site $j \in S$ as its support, so that we bound by first summing over $j \in S$, and then over S' that contains j , along with $1 \leq e^{\kappa_* \text{diam}(S)^\alpha}$ to invoke the κ_* -norm for the optimal decomposition of A . Although many individual factors S' could be badly overestimated in this step for finite κ_* , the $|S|$ factor in (S66) is parametrically optimal, since it is the number of small-region factors S' with $|S'| = O(1)$ that are contained in S .

It remains to bound \mathcal{O}_r in (S27) with $r \geq 1$, using that interaction strength decays as $\|A\|_{S'} \lesssim \epsilon e^{-\kappa_* (\text{diam} S')^\alpha}$. Although such decay is too slow to have a Lieb-Robinson bound like (S17), in Proposition 14 we prove a bound $\sim \epsilon e^{-\kappa_* r^\alpha}$ in (S162) for the time-evolved commutator of two operators separated by distance r . Choosing $Z = S$ and $Z' = \Lambda \setminus B(S, r)$ as defined in Proposition 14, we may write (by the triangle inequality)

$$\|\mathcal{O}_r\| \leq \left\| \sum_{k=r}^{\infty} \mathcal{O}_k \right\| + \left\| \sum_{k=r+1}^{\infty} \mathcal{O}_k \right\| \quad (\text{S67})$$

and bound, using Eq. (12) of [61]:

$$\left\| \sum_{k=r}^{\infty} \mathcal{O}_k \right\| \leq \int_{C_r} dU \| [U, \mathcal{O}] \| \leq \sup_U \| [U, \mathcal{O}] \| \quad (\text{S68})$$

where the set C_r is over unitaries acting non-trivially on Z' and dU is the Haar measure. Proposition 14 bounds the right hand side and leads to (S29).

2. Define superoperators

$$\mathcal{L} = i[H, \cdot], \quad \tilde{\mathcal{L}} = i[U^\dagger H U, \cdot], \quad \tilde{\mathcal{L}}' = i[H_0 + D_*, \cdot], \quad \mathcal{U} = U^\dagger \cdot U. \quad (\text{S69})$$

The left hand side of (S29) is then

$$\left\| \left(e^{t\mathcal{L}} - \mathcal{U}^\dagger e^{t\tilde{\mathcal{L}}'} \mathcal{U} \right) \mathcal{O} \right\| = \left\| \left(e^{t\tilde{\mathcal{L}}} - e^{t\tilde{\mathcal{L}}'} \right) \mathcal{U} \mathcal{O} \right\|, \quad (\text{S70})$$

where we have used the fact that \mathcal{U}^\dagger does not change the operator norm. Using the Duhamel identity

$$e^{t\tilde{\mathcal{L}}} - e^{t\tilde{\mathcal{L}}'} = \int_0^t dt' e^{(t-t')\tilde{\mathcal{L}}'} (\tilde{\mathcal{L}} - \tilde{\mathcal{L}}') e^{t'\tilde{\mathcal{L}}}, \quad (\text{S71})$$

(S70) is further bounded by

$$\left\| \left(e^{t\tilde{\mathcal{L}}} - e^{t\tilde{\mathcal{L}}'} \right) \mathcal{U} \mathcal{O} \right\| \leq \int_0^t dt' \left\| (\tilde{\mathcal{L}} - \tilde{\mathcal{L}}') e^{t'\tilde{\mathcal{L}}} \mathcal{U} \mathcal{O} \right\| \leq |t| \max_{0 \leq t' \leq t} \| [UV_* U^\dagger, \mathcal{O}(t')] \|, \quad (\text{S72})$$

where $\mathcal{O}(t') = e^{t'\mathcal{L}} \mathcal{O}$, and we have used (S24). In the commutator, the first operator $UV_* U^\dagger$ is extensive, yet should be close to V_* according to statement 1 of this Corollary. Thus the local norm of $UV_* U^\dagger$ is still exponentially small:

$$\| UV_* U^\dagger \|_{\kappa'_*} \lesssim \epsilon 2^{-k_*}, \quad (\text{S73})$$

for some $0 < \kappa'_* < \kappa_*$. See Proposition 6 for details. The second operator $\mathcal{O}(t')$ in the commutator in (S72) is evolved by the Hamiltonian $H = H_0 + V_1$, where interactions $H_{S'}$ decay at least sub-exponentially with $\text{diam} S'$ according to (S17) and (S22). Although we will prove an algebraic light cone $t \sim r^\alpha$ for such Hamiltonians in Proposition 14, the light cone is actually linear. After all, sufficiently fast decaying power-law interactions is sufficient to yield linear light cones [62]. Thus $\mathcal{O}(t')$ is mostly supported in a region \tilde{S} of linear size $\text{diam} S + v_{\delta, t'} |t'|$, except for a small part of operator norm $\delta \ll 1$, according to Eq.(5) in [62]. Their last equation of section II also ensures that one can safely ignore the δ -tail of $\mathcal{O}(t')$ acting outside of \tilde{S} , because the velocity $v_{\delta, t'}$ grows very mildly when decreasing $\delta \rightarrow 0$, if the power-law interaction decays sufficiently fast. When taking commutator with $UV_* U^\dagger$, only terms in $UV_* U^\dagger$ that are within this effective support \tilde{S} will contribute. This effect is bounded by a volume factor $|\tilde{S}| \lesssim |S|(|t'| + 1)^d$. Combining the factor $|t|$ in (S72), the local norm (S73), and $\|\mathcal{O}(t')\| = \|\mathcal{O}\|$, there must exist some constant c_{oper} such that (S29) holds.

3. The trace norm is related to the operator norm by

$$\begin{aligned} \|\rho_S(t) - \rho'_S(t)\|_1 &= \max_{\mathcal{O}=\mathcal{O}_S: \|\mathcal{O}\| \leq 1} \text{Tr} [\mathcal{O} (\rho_S(t) - \rho'_S(t))] \\ &= \max_{\mathcal{O}=\mathcal{O}_S: \|\mathcal{O}\| \leq 1} \text{Tr} \left[\rho_0 \left(e^{itH} \mathcal{O} e^{-itH} - U e^{it(H_0+D_*)} U^\dagger \mathcal{O} U e^{-it(H_0+D_*)} U^\dagger \right) \right] \\ &\leq \max_{\mathcal{O}=\mathcal{O}_S: \|\mathcal{O}\| \leq 1} \left\| e^{itH} \mathcal{O} e^{-itH} - U e^{it(H_0+D_*)} U^\dagger \mathcal{O} U e^{-it(H_0+D_*)} U^\dagger \right\| \leq c_{\text{oper}} \epsilon |S| (|t| + 1)^{d+1} 2^{-k_*}, \quad (\text{S74}) \end{aligned}$$

where \mathcal{O} is an arbitrary operator supported in S . The second line comes from the definitions (S31),(S32) and rearranging orders in the trace. The last step follows from the second result of this corollary, (S29). \square

3. FILTER FUNCTION AND ITS LOCALITY WHEN ACTING ON OPERATORS

This section contains the proof of Proposition 5, which requires the existence of a function $w(t)$ (as sketched in the main text) that can be used to build a projector \mathbb{P} onto Δ -diagonal operators.

3.1. Defining the filter function

In the proof above we frequently want to project out, for some operator, “off-resonant” matrix elements that connects pairs of eigenstates of H_0 that have energy difference $|E - E'| > \Delta$. This can be achieved as follows. Define the Δ -dependent filter function $w \in L^1(\mathbb{R})$ [57, 58]

$$w(t) := c_\Delta \Delta \prod_{n=1}^{\infty} \left(\frac{\sin a_n t}{a_n t} \right)^2, \quad \text{where } a_1 = c_1 \Delta, \quad a_n = \frac{a_1}{n \ln^2 n}, \quad \forall n \geq 2. \quad (\text{S75})$$

Here $c_1 \approx 0.161$ is chosen such that

$$\sum_{n=1}^{\infty} a_n = \frac{\Delta}{2}, \quad (\text{S76})$$

and $c_\Delta \in (1/(2\pi), 1/\pi)$ is a pure number chosen so that the function is normalized:

$$\int_{-\infty}^{\infty} dt w(t) = 1. \quad (\text{S77})$$

We define a similarly related odd function $W(t)$ by

$$W(t) = -W(-t) = \int_t^{\infty} ds w(s), \quad (t > 0). \quad (\text{S78})$$

These two functions have useful properties summarized in the following Proposition, which is proved in [57, 58].

Proposition 7. *w(t) satisfies the following:*

1. It is even in t, with bound $0 \leq w(t) \leq c_\Delta \Delta$.
2. The Fourier transform is compact

$$\hat{w}(\omega) = 0, \quad \forall |\omega| \geq \Delta, \quad (\text{S79})$$

and bounded $|\hat{w}(\omega)| \leq \int w(t) dt = 1 = \hat{w}(0)$.

3. subexponential decay:

$$w(t) \leq 2(e\Delta)^2 t e^{-\frac{2}{7} \frac{\Delta t}{\ln^2 \Delta t}}, \quad \text{if } t \geq e^{1/\sqrt{2}} \Delta^{-1}. \quad (\text{S80})$$

$W(t)$ satisfies

1. $W(t)$ is bounded as

$$|W(t)| \leq W(0+) = -W(0-) = 1/2. \quad (\text{S81})$$

2. For any bounded function $f(t)$,

$$\int_{-\infty}^{\infty} W(t) \frac{df(t)}{dt} dt = -f(0) + \int_{-\infty}^{\infty} w(t) f(t) dt. \quad (\text{S82})$$

3. $W(t)$ has weakly subexponential decay:

$$|W(t)| \leq 2e^2 \int_{\Delta|t|}^{\infty} s e^{-\frac{2}{7} \frac{s}{\ln^2 s}} ds, \quad \text{if } |t| \geq e^{1/\sqrt{2}} \Delta^{-1}. \quad (\text{S83})$$

3.2. Lieb-Robinson bound for the κ -norm

In this section, we establish a technical lemma for proving Proposition 5, which can be skipped in a first reading. In a nutshell, we wish to bound the growth in the κ -norm with the Lieb-Robinson bounds. This approach uses established, albeit tedious, methods.

Since the κ -norm takes infimum over all possible local decompositions of the form (S12), it suffices to prove bound on a particular local decomposition. Moreover, we will frequently use the κ -norm probed at vertex i :

$$\|\mathcal{O}\|_{\kappa,i} := \sum_{S \ni i} e^{\kappa(\text{diam}S)^\alpha} \|\mathcal{O}_S\|, \quad (\text{S84})$$

where a local decomposition is implicitly chosen. Then at the final step of the proof, we will use $\|\mathcal{O}\|_\kappa = \inf_{\{\mathcal{O}_S\}} \max_{i \in \Lambda} \|\mathcal{O}\|_{\kappa,i}$.

Any evolved local operator $e^{t\mathcal{L}_0}\mathcal{O}_{S_0}$, can be decomposed by

$$e^{t\mathcal{L}_0}\mathcal{O}_{S_0} = \sum_{r=0}^{\infty} \mathbb{Q}_r (e^{t\mathcal{L}_0}\mathcal{O}_{S_0}), \quad (\text{S85})$$

where the projector $\mathbb{Q}_r := \mathbb{Q}'_r - \mathbb{Q}'_{r-1}$ and

$$\mathbb{Q}'_r := \int_{\text{Haar outside } S(S_0,r)} dU U^\dagger \mathcal{O} U, \quad (\text{S86})$$

where $S(S_0, r)$ is the simplex $S \supset S_0$ whose faces are all of distance r to the parallel faces of S_0 . The measure ‘‘Haar outside $S(S_0, r)$ ’’ denotes the Haar measure on all unitary operators supported outside the set $S(S_0, r)$. Put simply, \mathbb{Q}_r is a projection onto all operators whose farthest support from S_0 is a ‘‘distance’’ r away from S_0 . Note that

$$\|\mathbb{Q}'_r \mathcal{O}\| \leq \|\mathcal{O}\|, \quad \|\mathbb{Q}_r \mathcal{O}\| \leq 2\|\mathcal{O}\|, \quad (\text{S87})$$

where the second inequality comes from (S86) and the triangle inequality applied to the definition of \mathbb{Q}_r .

Using (S85), we define a local decomposition for $e^{t\mathcal{L}_0}\mathcal{O}$ by simple extension:

$$e^{t\mathcal{L}_0}\mathcal{O} = \sum_{S_0} \sum_{r=0}^{\infty} \mathbb{Q}_{r,S_0} (e^{t\mathcal{L}_0}\mathcal{O}_{S_0}), \quad (\text{S88})$$

where we have shown the explicit dependence of \mathbb{Q}_r on S_0 , and each \mathbb{Q}_{r,S_0} term is viewed as supported in simplex $S(S_0, r)$. The decomposition $\mathcal{O} = \sum_{S_0} \mathcal{O}_{S_0}$ is optimal and is chosen to be (arbitrarily close to) minimizing the κ -norm $\|\mathcal{O}\|_\kappa$. The advantage of this decomposition is to invoke the Lieb-Robinson bound (S17), which implies for any r ,

$$\|\mathbb{Q}_r (e^{t\mathcal{L}_0}\mathcal{O}_{S_0})\| \leq 2\|\mathcal{O}_{S_0}\| \min\left(1, c_d \mu' (1 + \text{diam}S_0)^{d-1} e^{\mu(u|t|-r)}\right), \quad (\text{S89})$$

where we have used (S3). Here the first argument 1 in min is from (S87).

Lemma 8. *If $\kappa' < \kappa \leq \kappa_1$ and H_0 satisfies the Lieb-Robinson bound (S17) with*

$$\mu \geq 5\kappa_1, \quad (\text{S90})$$

then

$$\|e^{t\mathcal{L}_0}\mathcal{O}\|_{\kappa'} \leq c_0 [\max(1, -\ln \delta\kappa)]^{d-1} \|\mathcal{O}\|_\kappa (4u|t| + c_\mu)^d e^{\kappa'(4u|t|)^\alpha}, \quad (\text{S91})$$

where $\delta\kappa = \kappa - \kappa'$, c_0 and c_μ only depends on d, κ_1, μ, μ' , and α .

Proof. Assume $t \geq 0$ without loss of generality (alternatively set $|t| \rightarrow t$). For the decomposition (S88), suppose that a given vertex i in (S84) is the one responsible for the κ' -norm of $e^{t\mathcal{L}_0}\mathcal{O}$: in what follows, we will assume this i is fixed, as in (S84), but the result holds for any i and thus eventually for (S91) as well. Further fix a set S_0 . Then the initial operator \mathcal{O}_{S_0} can contribute an amount

$$K_{\kappa',i}(\mathcal{O}_{S_0}) \leq \|\mathbb{Q}_* (e^{t\mathcal{L}_0}\mathcal{O}_{S_0})\| e^{\kappa'(s-1+2r_0)^\alpha} \mathbb{I}(r_0 \geq \mathbf{d}(i, S_0)) + \sum_{r=r_0+1, r \geq \mathbf{d}(i, S_0)}^{\infty} \|\mathbb{Q}_r (e^{t\mathcal{L}_0}\mathcal{O}_{S_0})\| e^{\kappa'(s-1+2r)^\alpha}, \quad (\text{S92})$$

according to decomposition (S85), where we write $s = 1 + \text{diam}S_0$ for convenience, and $\mathbb{I}(\cdot)$ is the indicator function that returns 1 for input True and 0 for False. We have combined the leading terms

$$\mathbb{Q}_0 + \cdots + \mathbb{Q}_{r_0} = \mathbb{Q}_* \quad (\text{S93})$$

into a single operator, which has support in $S(S_0, r_0)$, with r_0 a constant chosen shortly to distinguish between pieces of the operator with “large” and “small” support.

Intuitively, the contribution $K_{\kappa',i}(\mathcal{O}_{S_0})$ is small if i is far from the initial support S_0 , compared to the distance ut that an operator can expand during time t . To formalize, observe that (S89) decays exponentially with r at sufficiently large r . This exponential decay is assured to kick in once $r > r'_0$, where we define r'_0 as the solution to

$$1 = c_d \mu' s^{d-1} e^{\mu(ut-r'_0)}, \quad \Rightarrow \quad r'_0 = ut + \frac{1}{\mu} \ln(c_d \mu' s^{d-1}). \quad (\text{S94})$$

We now choose $r_0 \approx 2r'_0$, noting that r_0 depends on t and s :

$$r_0 = 2 \left\lfloor ut + \frac{1}{\mu} \ln(c_d \mu' s^{d-1}) \right\rfloor + 1. \quad (\text{S95})$$

Here $\lfloor a \rfloor$ is the largest integer below a . Note that we are free to choose a large enough μ' so that r_0 is always positive. As a result, (S89) transforms to

$$\|\mathbb{Q}_*(e^{t\mathcal{L}_0} \mathcal{O}_{S_0})\| \leq 2 \|\mathcal{O}_{S_0}\|, \quad (\text{S96a})$$

$$\|\mathbb{Q}_r(e^{t\mathcal{L}_0} \mathcal{O}_{S_0})\| \leq 2c_d \mu' \|\mathcal{O}_{S_0}\| s^{d-1} e^{\mu(ut-r)} \leq 2c_d \mu' \|\mathcal{O}_{S_0}\| s^{d-1} e^{\mu(ut - \frac{r_0+1}{2} - \frac{r}{2})} \leq 2 \|\mathcal{O}_{S_0}\| e^{-\mu r/2}, \quad r > r_0. \quad (\text{S96b})$$

We can now bound $K_{\kappa',i}(\mathcal{O}_{S_0})$. We need to consider whether r_0 is larger or smaller than $\mathbf{d}(i, S_0)$. Let us start with the possibility that $r_0 < \mathbf{d}(i, S_0)$. Then (S96b) yields

$$\begin{aligned} K_{\kappa',i}(\mathcal{O}_{S_0}) &= 2 \|\mathcal{O}_{S_0}\| \sum_{r=\mathbf{d}(i,S_0)}^{\infty} e^{-\mu r/2} e^{\kappa'(s-1+2r)\alpha} \leq 2 \|\mathcal{O}_{S_0}\| \sum_{r=\mathbf{d}(i,S_0)}^{\infty} e^{-\mu r/2} e^{\kappa'[(s-1)\alpha+2r]} \\ &\leq 2 \|\mathcal{O}_{S_0}\| e^{-(\mu/2-2\kappa')\mathbf{d}(i,S_0)} e^{\kappa'(s-1)\alpha} \left(1 - e^{-(\mu/2-2\kappa')}\right)^{-1} \\ &\leq \frac{2}{1 - e^{-\mu/10}} \|\mathcal{O}_{S_0}\| e^{-\mu\mathbf{d}(i,S_0)/10} e^{\kappa'(s-1)\alpha}, \quad r_0 < \mathbf{d}(i, S_0). \end{aligned} \quad (\text{S97})$$

In the first line, we have used $\alpha < 1$ and

$$(a+b)^\alpha \leq a^\alpha + b^\alpha, \quad \forall a, b \geq 0. \quad (\text{S98})$$

In the second line we have summed the geometric series and used (S90). For the second case, $r_0 \geq \mathbf{d}(i, S_0)$, the $r > r_0$ part of summation is done exactly as (S97), with $\mathbf{d}(i, S_0)$ replaced by $r_0 + 1$. Thus combining with (S96a) yields

$$K_{\kappa',i}(\mathcal{O}_{S_0}) \leq 2 \|\mathcal{O}_{S_0}\| \left[e^{\kappa'(s-1+2r_0)\alpha} + \frac{1}{e^{\mu/10} - 1} e^{-\mu r_0/10} e^{\kappa'(s-1)\alpha} \right] \leq \left(2 + \frac{20}{\mu}\right) \|\mathcal{O}_{S_0}\| e^{\kappa'(s-1+2r_0)\alpha}, \quad r_0 \geq \mathbf{d}(i, S_0), \quad (\text{S99})$$

where we have simplified the prefactor using

$$e^a - 1 \geq a. \quad (\text{S100})$$

The κ' -norm at i , denoted by $\|e^{t\mathcal{L}_0} \mathcal{O}\|_{\kappa',i}$, is the sum $K_{\kappa',i}(\mathcal{O}_{S_0})$ over all \mathcal{O}_{S_0} . For each \mathcal{O}_{S_0} , let $x = \mathbf{d}(i, S_0)$: by definition, there is at least one site $j \in S_0$ such that $\mathbf{d}(i, j) = \mathbf{d}(i, S_0) \equiv x$. We can sum over S_0 by grouping the sums according to the x and j : the outermost sum will be over x , then we will sum over j at a fixed distance x , and then sum over sets S_0 with $j \in S_0$. Note that there can be multiple valid j for each S_0 , so this sum will overestimate the bound:

$$\|e^{t\mathcal{L}_0} \mathcal{O}\|_{\kappa',i} \leq \sum_{x=0}^{\infty} \sum_{j:\mathbf{d}(i,j)=x} \sum_{\mathcal{O}_{S_0}}^{j,x} K_{\kappa',i}(\mathcal{O}_{S_0}) = \sum_{x=0}^{\infty} \sum_{j:\mathbf{d}(i,j)=x} \left[\sum_{\mathcal{O}_{S_0}:s < f_0(x)}^{j,x} K_{\kappa',i}(\mathcal{O}_{S_0}) + \sum_{\mathcal{O}_{S_0}:s \geq f_0(x)}^{j,x} K_{\kappa',i}(\mathcal{O}_{S_0}) \right], \quad (\text{S101})$$

where $\sum_{\mathcal{O}_{S_0}}^{j,x}$ means the restriction that $S_0 \ni j$ and $\mathbf{d}(i, S_0) = x$ (as described above in words). In the latter equality, we separated the sum according to whether $r_0(s) < \mathbf{d}(i, S_0) = x$, which is equivalent to whether $1 + \text{diam}S_0 = s < f_0(x)$, where

$$x = 2 \left[ut + \frac{1}{\mu} \ln (c_d \mu' f_0(x)^{d-1}) \right] + 1. \quad (\text{S102})$$

Note that for $d = 1$ there is no solution for $f_0(x)$ since r_0 is a fixed number independent of s , and hence we do not need to sum over $s \geq f_0(x)$ in $d = 1$: (S101) simply vanishes for all $x > r_0$. Thus for $d = 1$ we can simply set $f_0(x) = 1$ for $x \leq r_0$ and $f_0(x) = +\infty$ otherwise, so that (S101) and the following equations still make sense.

We first bound the first term in (S101) using (S97):

$$\begin{aligned} \sum_{x=0}^{\infty} \sum_{j:\mathbf{d}(i,j)=x} \sum_{\mathcal{O}_{S_0}:s < f_0(x)}^{j,x} K_{\kappa',i}(\mathcal{O}_{S_0}) &\leq \frac{2}{1 - e^{-\mu/10}} \sum_{x=0}^{\infty} \sum_{j:\mathbf{d}(i,j)=x} \sum_{\mathcal{O}_{S_0}:S_0 \ni j} \|\mathcal{O}_{S_0}\| e^{-\mu x/10} e^{\kappa'(s-1)^\alpha} \\ &\leq \frac{2}{1 - e^{-\mu/10}} \sum_{x=0}^{\infty} \sum_{j:\mathbf{d}(i,j)=x} e^{-\mu x/10} \|\mathcal{O}\|_{\kappa'} \\ &\leq \frac{2c_d}{1 - e^{-\mu/10}} \|\mathcal{O}\|_{\kappa'} \sum_{x=0}^{\infty} (1 + 2x)^{d-1} e^{-\mu x/10} = c_{<} \|\mathcal{O}\|_{\kappa}, \end{aligned} \quad (\text{S103})$$

which is well upper bounded by the right hand side of (S91). Here we have used (S3) with $S = \{j : \mathbf{d}(i, j) \leq x\}$ to get

$$\sum_{j:\mathbf{d}(i,j)=x} \leq c_d (1 + 2x)^{d-1}, \quad (\text{S104})$$

because $\partial S = \{j : \mathbf{d}(i, j) = x\}$ and $\text{diam}S \leq 2x$ from triangle inequality. The constant $c_{<}$ only depends on d and μ . We also used $\{\mathcal{O}_{S_0}\}$ is the optimal decomposition of \mathcal{O} that realizes its κ' -norm.

We now evaluate the second term in (S101). First, we use that if $s \geq f_0(x)$, we may as well use (S99) to bound

$$\begin{aligned} K_{\kappa',i}(\mathcal{O}_{S_0}) &\leq \left(2 + \frac{20}{\mu}\right) \|\mathcal{O}_{S_0}\| e^{\kappa'(s-1+2r_0)^\alpha} \\ &\leq \left(2 + \frac{20}{\mu}\right) \|\mathcal{O}_{S_0}\| \exp \left\{ \kappa' \left[(4ut)^\alpha + \left(s + 1 + \frac{4}{\mu} \ln (c_d \mu' s^{d-1}) \right)^\alpha \right] \right\} \\ &\leq \left(2 + \frac{20}{\mu}\right) e^{\kappa'(4ut)^\alpha} \|\mathcal{O}_{S_0}\| \exp \{ \kappa' [(s-1)^\alpha + c_{\text{ln}}] \}. \end{aligned} \quad (\text{S105})$$

In the second line we have used (S98). In the third line we have used the fact that the function $(s + a \ln s + b)^\alpha - s^\alpha$ is upper bounded for any a, b (given $\alpha < 1$), and the resulting constant c_{ln} in (S105) is determined by α, d, μ, μ' . Now, combining (S101), (S103), and (S105), we find

$$\begin{aligned} \|e^{t\mathcal{L}_0} \mathcal{O}\|_{\kappa',i} &\leq c_{<} \|\mathcal{O}\|_{\kappa} + \left(2 + \frac{20}{\mu}\right) e^{\kappa'(4ut)^\alpha} \sum_{x=0}^{\infty} \sum_{j:\mathbf{d}(i,j)=x} \sum_{\mathcal{O}_{S_0}:s \geq f_0(x)}^{j,x} \|\mathcal{O}_{S_0}\| \exp \{ \kappa' [(s-1)^\alpha + c_{\text{ln}}] \} \\ &\leq c_{<} \|\mathcal{O}\|_{\kappa} + \left(2 + \frac{20}{\mu}\right) e^{\kappa'[c_{\text{ln}} + (4ut)^\alpha]} \sum_{x=0}^{\infty} \sum_{j:\mathbf{d}(i,j)=x} e^{-\delta\kappa(f_0(x)-1)^\alpha} \|\mathcal{O}\|_{\kappa} \\ &\leq c_{<} \|\mathcal{O}\|_{\kappa} + \left(2 + \frac{20}{\mu}\right) e^{\kappa'[c_{\text{ln}} + (4ut)^\alpha]} \|\mathcal{O}\|_{\kappa} \sum_{x=0}^{\infty} c_d (2x+1)^{d-1} e^{-\delta\kappa(f_0(x)-1)^\alpha}. \end{aligned} \quad (\text{S106})$$

In the second line of (S106) we have used the Markov inequality

$$\begin{aligned} \sum_{\mathcal{O}_{S_0}:s \geq f_0(x)}^{j,x} \|\mathcal{O}_{S_0}\| e^{\kappa'(s-1)^\alpha} &= \sum_{\mathcal{O}_{S_0}:s \geq f_0(x)}^{j,x} \|\mathcal{O}_{S_0}\| e^{(\kappa - \delta\kappa)(s-1)^\alpha} \\ &\leq \sum_{\mathcal{O}_{S_0}:s \geq f_0(x)}^{j,x} \|\mathcal{O}_{S_0}\| e^{\kappa(s-1)^\alpha} e^{-\delta\kappa(f_0(x)-1)^\alpha} \leq \|\mathcal{O}\|_{\kappa} e^{-\delta\kappa(f_0(x)-1)^\alpha}. \end{aligned} \quad (\text{S107})$$

In the last line of (S107) we have overestimated the final sum over S_0 . Returning to the last line of (S106), we have used (S104).

If $d = 1$, the sum over x in (S106) is finite: $x \leq r_0 \approx 2ut$, and (S91) follows easily.

For $d > 1$, note that there exists a constant c'_μ that depends on d, μ, μ' , such that when $x \geq 4ut + c'_\mu$, (S102) yields

$$f_0(x) \geq (c_d \mu')^{-\frac{1}{d-1}} e^{\frac{\mu}{d-1}(\frac{x-1}{2}-ut-1)} \geq e^{\frac{\mu x}{4(d-1)}} \equiv y, \quad (x \geq 4ut + c'_\mu). \quad (\text{S108})$$

Changing variable from x to y defined above, (S106) becomes

$$\begin{aligned} \|e^{t\mathcal{L}_0} \mathcal{O}\|_{\kappa', i} &\leq c_{<} \|\mathcal{O}\|_\kappa + \frac{(2\mu + 20)c_d}{\mu} e^{\kappa' \{c_{\text{in}} + (4ut)^\alpha\}} \|\mathcal{O}\|_\kappa \left[(4ut + c'_\mu + 1)^d + \left(\frac{4(d-1)}{\mu}\right)^d c_{\text{int}} \int_1^\infty \frac{dy}{y} (\ln y)^{d-1} e^{-\delta\kappa(y-1)^\alpha} \right] \\ &\leq c_{<} \|\mathcal{O}\|_\kappa + \frac{(2\mu + 20)c_d}{\mu} e^{\kappa' \{c_{\text{in}} + (4ut)^\alpha\}} \|\mathcal{O}\|_\kappa \left[(4ut + c'_\mu + 1)^d + \left(\frac{4(d-1)}{\mu}\right)^d c'_{\text{in}} [\max(1, -\ln \delta\kappa)]^{d-1} \right] \\ &\leq c_{<} \|\mathcal{O}\|_\kappa + \frac{(2\mu + 20)c_d c'_{\text{in}}}{\mu} e^{\kappa' \{c_{\text{in}} + (4ut)^\alpha\}} \|\mathcal{O}\|_\kappa [\max(1, -\ln \delta\kappa)]^{d-1} \left(4ut + c'_\mu + \frac{4(d-1)}{\mu}\right)^d. \end{aligned} \quad (\text{S109})$$

Here in the first line we have summed over $x < 4ut + c'_\mu$, and c_{int} is a constant due to replacing the sum in (S106) by an integral. In the second line we have used the fast decay of the exponential function: if $\delta\kappa = \Omega(1)$, then the integral is $O(1)$; otherwise if $\delta\kappa \ll 1$, then rescaling $\delta\kappa y^\alpha = \tilde{y}^\alpha$ will pull out an overall factor $(\ln \frac{1}{\delta\kappa})^{d-1}$. In the last line, we combined the two terms using $a^d + b^d \leq (a+b)^d$. Finally, as our result does not depend at all on i , we are free to re-label our final $O(1)$ constants sitting out in front; in doing so, we arrive at (S91). \square

3.3. Proof of Proposition 5

Proof of Proposition 5. (S39) comes from

$$[H_0, \mathbb{A}\mathcal{O}] = \int_{-\infty}^\infty W(t) \frac{d}{dt} (e^{t\mathcal{L}_0} \mathcal{O}) dt = (-1 + \mathbb{P})\mathcal{O}, \quad (\text{S110})$$

following (S82). To prove (S40a), note that the condition (S90) of Lemma 8 is just (S19). Thus we use (S91) to get

$$\begin{aligned} \|\mathbb{P}\mathcal{O}\|_{\kappa'} &\leq \int_{-\infty}^\infty dt w(t) \|e^{t\mathcal{L}_0} \mathcal{O}\|_{\kappa'} \leq 2c_0 [\max(1, -\ln \delta\kappa)]^{d-1} \|\mathcal{O}\|_\kappa \int_0^\infty dt w(t) (4ut + c_\mu)^d e^{\kappa' (4ut)^\alpha} \\ &\leq 2c_0 [\max(1, -\ln \delta\kappa)]^{d-1} \|\mathcal{O}\|_\kappa \left[\left(c_\mu + e^{\frac{1}{\sqrt{2}} \frac{4u}{\Delta}}\right)^d e^{\kappa \left(e^{\frac{1}{\sqrt{2}} \frac{4u}{\Delta}}\right)^\alpha} + 2e^2 \int_{e^{\frac{1}{\sqrt{2}}}}^\infty d\tilde{t} e^{\kappa \left(e^{\frac{1}{\sqrt{2}} \tilde{t} \frac{4u}{\Delta}}\right)^\alpha - \frac{2\tilde{t}}{7 \ln^2 \tilde{t}}} \left(c_\mu + e^{\frac{1}{\sqrt{2}} \tilde{t} \frac{4u}{\Delta}}\right)^d \right]. \end{aligned} \quad (\text{S111})$$

In the second line, we have first used (S80) to bound the large t tails of the integral; for the small t limit we have simply bounded the integral by using the maximum of each term in the integrand separately in the domain $t < e^{1/\sqrt{2}} \Delta^{-1}$, together with (S77), which reduces to the form of (S40a) since the integral converges for any $\alpha < 1$. (S40b) comes from (S83) using almost identical manipulations. \square

4. PROOF OF PROPOSITION 6

4.1. Motivation

It remains to prove Proposition 6, which is the main difficulty of generalizing [13]. Since this Proposition is a self-contained bound on local dynamics beyond conventional Lieb-Robinson bound, we restate it below replacing A with itH . This notation will be used for this whole section. Therefore, in this section, the Hamiltonian H is *not* the same one as (S4): we only require it to be local in the sense of (S114).

Proposition 6 (restatement). *Suppose $\kappa' < \kappa \leq \kappa_1$ with*

$$\delta\kappa = \kappa - \kappa'. \quad (\text{S112})$$

If

$$\mathcal{L} = i[H, \cdot], \quad (\text{S113})$$

with Hamiltonian H on a d -dimensional simplicial lattice, satisfying

$$|t| \|H\|_{\kappa} \leq (\delta\kappa)^{\frac{3d+1}{\alpha}}, \quad (\text{S114})$$

then

$$\|e^{t\mathcal{L}}\mathcal{O} - \mathcal{O}\|_{\kappa'} \leq c_-(\delta\kappa)^{-\frac{2d-1}{\alpha}} |t| \|H\|_{\kappa} \|\mathcal{O}\|_{\kappa}, \quad (\text{S115})$$

where $0 < c_- < \infty$ depends on d, α and κ_1 .³

The difference between Proposition 6 and its counterpart, Lemma 4.1 in [13], is that here H is less local:

$$\|H_S\| \lesssim e^{-\kappa(\text{diam}S)^\alpha}, \quad \alpha < 1. \quad (\text{S116})$$

In contrast, $\|H_S\| \lesssim e^{-\kappa|S|}$ in [13]. As a result, a simple combinatorial expansion of $e^{t\mathcal{L}}\mathcal{O}$ over t , which was done in [13], no longer converges for $d > 1$. Essentially, the issue is that while H is sub-exponentially localized⁴, the growth in the κ -norm can be dominated by tiny terms in H with unusually large polygons S in which they are supported. In $d > 1$, there are an increasingly large number of ways for such large polygons to intersect with \mathcal{O} , so they must be summed up with some care to not overcount.

To sketch the proof that follows, we first observe that evolution generated by a Hamiltonian H of form (S116) still has a Lieb-Robinson bound that we will prove in Proposition 14 using established methods (see, e.g., [61]), which resums the divergence of the simple Taylor expansion mentioned above. The idea is illustrated in Fig. S2 for 1d, where we focus on a single local operator \mathcal{O}_{S_0} , since the conclusion for the extensive \mathcal{O} follows simply by superposition. The rectangle at each layer represents commuting the operator with a local term H_X in the Hamiltonian $H = \sum_X H_X$. Suppose at some step, the operator has support on S . Then there are $\sim |S|$ terms of H_X that act nontrivially on the operator. However, only $\sim |\partial S|$ of them (red rectangles in Fig. S2) grow the operator to a strictly larger support, while the ‘‘bulk’’ ones (yellow rectangles) yield unitary rotations inside the support. Using Lieb-Robinson techniques, one can essentially ignore these internal rotations, and bound the operator growth only by the ‘‘boundary’’ ones, leading to a convergent series.

Unfortunately there is an important technical difference between a standard Lieb-Robinson bound, which bounds $\|[A_S(t), B_R]\|$ for fixed sets S and R , and a bound on $\|\mathcal{O}(t)\|_{\kappa}$. In the former, we need only keep track of (loosely speaking) terms that grow set S towards set R . In the latter, we need to keep track of all terms which grow the operator in any direction on the lattice – for $d > 1$, increasingly large operators have a large perimeter with many possible ways to grow. In particular, suppose the Hamiltonian terms are all of size $\text{diam}S \sim r$. Then the typical size of a grown operator is $\geq 2r$, because each end of the operator can be attached by a H_S of size r that just touches the end. Even in $d = 1$, a Lieb-Robinson bound can affirm that it took time ut for (terms of high weight) in the operator to expand a distance r to the right; during the same time it also will likely expand a distance r to the left. When $r \gg |S|$, this implies that $|S|$ grows twice as fast as the Lieb-Robinson velocity. So in every iteration, the typical size of an operator in H_k will at least double. Returning to the overarching sketch of our proof, this would make $k_* \sim \log \frac{\Delta}{v_1}$. So we need to use the extra fact that, by attaching more H_S to the initial operator, the amplitude is suppressed by more powers of $\|H\|_{\kappa}$. In other words, we need to differentiate cases where 1 or 2 ends are attached by H_S , which is not considered in conventional Lieb-Robinson bounds.

Our strategy can be intuitively described first in $d = 1$, and we will do so now. Let operator \mathcal{O}_S have support on connected subset $S \subseteq \Lambda$: namely S is an interval. \mathcal{O}_S can only grow at the two ends (left and right) of the support interval. For example in Fig. S2, we are studying a term in the expansion of the time-evolved operator of the form

$$e^{\mathcal{L}t}\mathcal{O}_{S_0} \subseteq \frac{t^5}{5!} \mathcal{L}_{X_5} \mathcal{L}_{X_4} \mathcal{L}_{X_3} \mathcal{L}_{X_2} \mathcal{L}_{X_1} \mathcal{O}_{S_0}.$$

Intuitively the operator grows as follows: the left end first moves from site 4 to 3 by X_1 , and then from 3 to 2 by X_3 ; similarly on the right end, X_4 moves us from 6 to 8. So, in order to grow the initial domain $S_0 = \{4, 5, 6\}$ by two

³ We believe the exponent of $\delta\kappa$ can be improved, for example, to $(\delta\kappa)^{-\frac{1}{\alpha} \max(d, 2d-2)}$, using complicated geometrical facts about simplices. Such improvement will lead to a larger $a > \frac{1}{2d-1}$ in the prethermal time scale $t_* \sim \exp\left(\frac{\Delta}{\epsilon}\right)^a$.

⁴ As Hamiltonians with algebraic tails have rigorous Lieb-Robinson bounds [62–66], this remains an extremely strong condition. Still, it requires some care to re-sum and find a bound on $\|e^{\mathcal{L}t}\mathcal{O}\|_{\kappa}$, which is not the object usually bounded by a Lieb-Robinson Theorem!




FIG. S2. A sketch of the Heisenberg evolution for an operator, which is equivalent to step-by-step taking its commutator with local terms H_X of the Hamiltonian H . Starting from the initial operator supported on $S_0 = \{4, 5, 6\}$, some H_{X_n} s truly grow the support (red rectangles) to finally reach $S = \{2, 3, \dots, 8\}$, while others do not grow the support and serve as internal rotations (yellow rectangles). Only the former, which intersect with the boundary of the operator at each step, contributes to enlarging the κ -norm of the final operator.

sites on each end, we must traverse:

$$\begin{cases} \text{left : } 4 \xrightarrow{X_1} 3 \xrightarrow{X_3} 2 \\ \text{right : } 6 \xrightarrow{X_3} 7 \xrightarrow{X_4} 8 \end{cases}.$$

This pattern only depends on the boundary terms H_X (red rectangles in Fig. S2). However, observe that in general the Taylor expansion of $e^{\mathcal{L}t}\mathcal{O}_{S_0}$ will contain many additional terms which act entirely inside of S_0 . We do not want to count these terms, since they cannot grow the support of the operator at all. The key observation, first made in [?], is that one can elegantly classify all of the possible orderings for the “red” terms in H (those that grow the operator), in such a way that all possible intermediate sequences of yellow terms can be re-exponentiated to form a unitary operation (which leaves operator norms invariant)! The practical consequence of this observation is that we only need to bound the contributions of red terms (and the number of possible patterns of red terms) when building a Lieb-Robinson bound for the κ -norm. We emphasize that it is crucial that we track both the left and right end: the main technical issue addressed in this section is how to find such a “direction-resolved” Lieb-Robinson bound in $d > 1$.

4.2. Irreducible skeleton representation

We first focus on a single initial operator $\mathcal{O} = \mathcal{O}_{S_0}$, locally supported in a fixed d -dimensional simplex S_0 . Recall that we work with the simplicial lattice, and the Hamiltonian $H = \sum_X H_X$ is also expanded in the simplices X on which operators are supported. All simplices are regular and have the same orientation, as shown by the magenta triangles in Fig. S1.

To describe how the Hamiltonian couples different lattice sites, we use the factor graph theory approach developed in [59]. A factor graph $G = (\Lambda, F, E)$ is defined as follows: Λ contains all of the lattice sites (as before), while F is the set of all factors X that appear in $H = \sum_X H_X$. Their union $\Lambda \cup F$ serves as the vertices of the factor graph G , and we connect each factor $X \subseteq F$ to all its contained sites $i \in X$. Hence, the edges connecting F and Λ form the edge set E of the factor graph G . As an example, in Fig. S3(a) the green circles are lattice sites and the red rectangles are factors. Rectangles representing nearest-neighbor three-site interaction are connected to their supporting sites by black lines, while the connection for the six-site interactions are explicitly shown only for one of its supporting sites.

We expand $e^{t\mathcal{L}}\mathcal{O}$ in powers of t :

$$e^{t\mathcal{L}}\mathcal{O} = \sum_{n=0}^{\infty} \frac{t^n}{n!} \mathcal{L}^n \mathcal{O}. \quad (\text{S117})$$

Further expanding \mathcal{L} into factors using $\mathcal{L} = \sum_X \mathcal{L}_X$, each term is of the form $\mathcal{L}_{X_n} \cdots \mathcal{L}_{X_1} \mathcal{O}$, where $\mathcal{L}_X := i\text{Adj}_{H_X}$. Crucially, observe that such a term is nonzero only if the sequence $T = (S_0, X_1, \cdots, X_n)$ obeys a *casual structure*: each X_m intersects with the previous support $S_0 \cup X_1 \cup \cdots \cup X_{m-1}$.⁵ If this condition holds, we say T is a causal tree. The tree structure, embedded in the factor graph G , is thoroughly defined in [59]; in what follows we will borrow closely from their formalism. We see that (S117) only contains contributions from the sequences $\mathcal{L}_{X_n} \cdots \mathcal{L}_{X_1}$ arising from causal trees T whose root is S_0 . Defining \mathcal{T}_{S_0} to be the set of all causal trees that have a root in S_0 , we write:

$$e^{t\mathcal{L}}\mathcal{O} = \sum_{T \in \mathcal{T}_{S_0}} \frac{t^n}{n!} \mathcal{L}_{X_n} \cdots \mathcal{L}_{X_1} \mathcal{O}. \quad (\text{S118})$$

For a given causal tree T , let us denote with $S(T)$ the unique (smallest) circumscribed simplex of T . For example, the circumscribed simplex for the subset $\{2, 3, 4, 5, 9\} \subseteq \Lambda$ in Fig. S3(a) is the triangle $S = \{1, 2, \cdots, 10\}$, since the subset touches all three faces of S . Then (S118) gives rise to a decomposition

$$e^{t\mathcal{L}}\mathcal{O} = \sum_{S \supset S_0} \sum_{T \in \mathcal{T}_{S_0}: S(T)=S} \frac{t^n}{n!} \mathcal{L}^n \mathcal{O} := \sum_{S \supset S_0} (e^{t\mathcal{L}}\mathcal{O})_S, \quad (\text{S119})$$

of the evolved operator to simplices S . Note that some local terms of the evolved operator are assigned to a looser support S , if the operator accidentally becomes the identity operator at sites in ∂S . For example, if $\mathcal{O} = Z_1 Z_2$ and $X_1 = Z_1 X_2$, then $\mathcal{L}_{X_1} \mathcal{O} \propto I_1 Y_2$ acts trivially on site 1, which we however assign a support of $S = \{1, 2\}$ in the decomposition (S117).

The next crucial step is to realize that the causal trees T can be re-summed in an elegant way by sorting them into equivalence classes based on *irreducible* subsequences of T . Given T and its circumscribed simplex $S(T)$, for any of the $d+1$ faces $f_1, f_2, \cdots, f_{d+1}$ of $S(T)$, there is a unique self-avoiding path $\Gamma_p \subseteq T$ that starts from the initial support S_0 and ends at the face f_p [59]. For the casual tree example in Fig. S3(a), the path Γ_1 is trivial containing only the root S_0 , while $\Gamma_2 = S_0 \ni 8 \rightarrow X_1 \rightarrow 5 \rightarrow X_3 \rightarrow 3 \in f_2$ (or equivalently, arriving at $6 \in f_2$). The $d+1$ paths Γ_p form a tree Γ , of the form in Fig. S3(b) for 2d, which we call the irreducible skeleton $\Gamma(T)$ for the causal tree T . Here “irreducible” means self-avoiding. Note that some of the branches of the irreducible skeleton may be absent, for example Γ looks like Fig. S3(c) in 2d, if S_0 already shares one face of S . For the casual tree example in Fig. S3(a), the irreducible skeleton is

$$S_0 \ni 8 - X_1 \begin{cases} 5 - X_3 - 3 \in f_2 \\ 9 - X_5 - 14 \in f_3 \end{cases} \quad (\text{S120})$$

$\Gamma(T)$ is a well-defined function of T , provided that

Proposition 9. *For each causal tree T , there is a unique irreducible skeleton Γ .*

Proof. Γ is unique because it is composed by the unique irreducible paths Γ_p from S_0 to f_p . For the uniqueness of Γ_p , see Proposition 2 of [59]. Intuitively, for a given face f_p , one can first uniquely determine the first factor \tilde{X}_1 in T that hits it. Next, there is a unique factor \tilde{X}_2 in T that first hits the support of \tilde{X}_1 . Further tracing back along the casual tree T , the unique Γ_p is then $S_0 \rightarrow \cdots \rightarrow \tilde{X}_2 \rightarrow \tilde{X}_1$. \square

As a result, we can define $[\Gamma] \subseteq \mathcal{T}_{S_0}$ as the equivalence class of causal trees that have Γ as its irreducible skeleton, and define $\mathbf{\Gamma}(S)$ as the set of all Γ that have S as their circumscribed equilateral triangle, with each face of S touched only once by Γ , and only by its $d+1$ end points.

Each irreducible skeleton Γ corresponds to factors $\{X_1^\Gamma, \cdots, X_\ell^\Gamma\}$, where $\ell = \ell(\Gamma)$ is the “length” of Γ . However, their sequence acting on the operator can vary because different branches of Γ can grow “independently”: the only constraint is causality, that each factor in the irreducible path Γ_p to a fixed face must appear in that same order in Γ . Thus we define $\gamma = (X_1^\gamma, \cdots, X_\ell^\gamma)$ as the ordered irreducible skeleton, and use $\gamma \in \mathbf{\Gamma}$ to mean γ is one ordered skeleton of Γ . Each γ thus corresponds to a finer equivalence class $[\gamma] \subseteq \mathcal{T}_{S_0}$ of casual trees T . Now we are prepared to give our re-summation lemma.

⁵ If this does not happen, then the expression must vanish as it contains within it a commutator of operators supported on disjoint sets.




FIG. S3. (a) Example of the factor graph and the irreducible skeleton construction at $d = 2$. The lattice sites are numbered green circles composing a triangular lattice. We consider two kinds of factors here represented by red squares: three-site interactions in H supported on the smallest triangles, and six-site interactions on the second smallest triangles. In the factor graph G , each factor is connected to all of the lattice sites that support it. In the figure, however, three-site square factors are connected to all of their lattices sites by black lines, while only one connection edge is explicitly shown for each six-site square factor. Having constructed the factor graph, we consider operator growth from $S_0 = \{4, 7, 8\}$ (the cyan triangle) to the whole triangle $S = \{1, 2, \dots, 15\}$. In particular, we consider a causal tree $T = (S_0, X_1, \dots, X_6)$ that starts from S_0 and reaches the three faces of S , i.e., $S(T) = S$. Since S_0 already intersects with the face $f_1 = \{1, 2, 4, 7, 11\}$, one only needs to figure out the unique irreducible paths from S_0 to $f_2 = \{1, 3, 6, 10, 15\}$ and $f_3 = \{11, 12, \dots, 15\}$. For example, the irreducible path from S_0 to f_2 is $\Gamma_2 = 8 \rightarrow X_1 \rightarrow 5 \rightarrow X_3 \rightarrow 3$ (or equivalently, 6), shown in blue solid lines. The reason is that f_2 is first reached by X_3 , and because X_3 grows from 5, one needs to figure out how 5 is first reached: it is reached by X_1 that acts nontrivially on $8 \in S_0$. Similarly, the irreducible path from S_0 to f_3 is the orange path $\Gamma_3 = 8 \rightarrow X_1 \rightarrow 9 \rightarrow X_5 \rightarrow 14(13, 15)$. Merging these two paths, we find the ordered irreducible skeleton $\gamma = (X_1^\gamma, X_3^\gamma, X_5^\gamma) = (X_1, X_3, X_5)$ that T belongs to. The length is $\ell = 3$. The corresponding non-ordered irreducible skeleton is thus $\Gamma = \{X_1, X_3, X_5\}$, which coincidentally, only has γ as its ordered counterpart. The other factors $\{X_2, X_4, X_6\}$ in the casual tree T , are linked to the skeleton by causality, as shown by brown dashed lines. They are allowed factors in $\mathcal{L}_1^\gamma, \mathcal{L}_2^\gamma, \mathcal{L}_3^\gamma$ respectively, defined in (S123), which are resummed to unitaries inserted between steps of the irreducible skeleton γ , in (S121). In Subsection 4.4, we make further definitions. Γ has two bifurcation factors: $Y_0 = S_0$, and $Y_1 = X_1$ (pink triangle) where the two paths Γ_2 and Γ_3 merge. As for the irreducible paths, \mathcal{P}_1 is trivial (length $l_1 = 0$) because S_0 intersects with f_1 . \mathcal{P}_2 and \mathcal{P}_3 , which start from Y_1 , are $5 \rightarrow X_3 \rightarrow 3(6)$ and $9 \rightarrow X_5 \rightarrow 14(13, 15)$ respectively. The paths $\mathcal{P}'_1, \mathcal{P}'_2$ connecting Y_p s are both trivial. (b-d) Topological structure of irreducible skeletons. (b-c) are for $2d$, with (c) being the special case that $l_1 = 0$, as the example in (a). (d) A more complicated example for $5d$.

Lemma 10. *The following identities hold:*

$$\begin{aligned}
 (e^{t\mathcal{L}}\mathcal{O})_S &= \sum_{\Gamma \in \mathbf{\Gamma}(S)} (e^{t\mathcal{L}}\mathcal{O})_\Gamma, \quad \text{where} \\
 (e^{t\mathcal{L}}\mathcal{O})_\Gamma &= \sum_{\gamma \in \Gamma_{\mathbb{T}^\ell(t)}} \int dt_1 \cdots dt_\ell e^{(t-t_\ell)\mathcal{L}_\ell^\gamma} \mathcal{L}_{X_\ell^\gamma} \cdots e^{(t_2-t_1)\mathcal{L}_1^\gamma} \mathcal{L}_{X_1^\gamma} e^{t_1\mathcal{L}_0^\gamma} \mathcal{O},
 \end{aligned} \tag{S121}$$

with $\ell = \ell(\Gamma) = \ell(\gamma)$, and

$$\mathbb{T}^\ell(t) = \{(t_1, \dots, t_\ell) \in [0, t]^\ell : t_1 \leq t_2 \leq \dots \leq t_\ell\}. \tag{S122}$$

The set of allowed factors between steps k and $k+1$ are included in

$$\mathcal{L}_k^\gamma = \mathcal{L}|_{R_k^\gamma} - \sum_{Y \subseteq R_k^\gamma: Y \cap W_k^\gamma \neq \emptyset} \mathcal{L}_Y, \quad (\text{S123})$$

where

$$W_k^\gamma = \bigcup_{m=k+2: X_m^\gamma \cap \gamma_k = \emptyset}^{\ell(\gamma)} X_m^\gamma, \quad \text{where } \gamma_k := S_0 \cup X_1^\gamma \cup \dots \cup X_k^\gamma, \quad (\text{S124})$$

and

$$R_k^\gamma = \begin{cases} S, & k = \ell \\ S - \bigcup_{p'=p}^{d+1} f_{p'}^\gamma, & \ell_{p-1} \leq k = \ell_{p-1} < \ell_p, \quad (p = 1, \dots, d+1) \end{cases} \quad (\text{S125})$$

where γ first reaches the faces in order $(f_1^\gamma, f_2^\gamma, \dots, f_{d+1}^\gamma)$, by factors $(X_{\ell_1}^\gamma, X_{\ell_2}^\gamma, \dots, X_{\ell_{d+1}}^\gamma)$ respectively. Here $\ell_0 = 0, \ell_{d+1} = \ell$, and R_k^γ is always a simplex.

Proof. Define

$$(e^{t\mathcal{L}}\mathcal{O})_\Gamma := \sum_{T \in [\Gamma]} \frac{t^n}{n!} \mathcal{L}_{X_n} \cdots \mathcal{L}_{X_1} \mathcal{O} = \sum_{\gamma \in \Gamma} \sum_{T \in [\gamma]} \frac{t^n}{n!} \mathcal{L}_{X_n} \cdots \mathcal{L}_{X_1} \mathcal{O}. \quad (\text{S126})$$

The first equality in (S121) holds because each sequence T is a causal tree and by Proposition 9 each T belongs to a unique equivalence class. Now we need to verify the second equality of (S121). To do so, we first prove

$$\sum_{T \in [\gamma]} \frac{t^n}{n!} \mathcal{L}_{X_n} \cdots \mathcal{L}_{X_1} \mathcal{O} = \sum_{m_0, \dots, m_\ell = 0}^{\infty} \frac{t^{\ell + m_0 + \dots + m_\ell}}{(\ell + m_0 + \dots + m_\ell)!} (\mathcal{L}_\ell^\gamma)^{m_\ell} \mathcal{L}_{X_\ell^\gamma} \cdots (\mathcal{L}_1^\gamma)^{m_1} \mathcal{L}_{X_1^\gamma} (\mathcal{L}_0^\gamma)^{m_0} \mathcal{O}, \quad (\text{S127})$$

by identifying terms on the two sides, as a generalization of Lemma 4 in [59]. Observe that when expanding \mathcal{L}_k^γ into factors using (S123), each term on the right hand side forms a casual tree T that has γ as its ordered irreducible skeleton. Indeed, before acting with $\mathcal{L}_{X_{k+1}^\gamma}$, all previous factors included in $\mathcal{L}_0^\gamma, \dots, \mathcal{L}_k^\gamma$ do not touch W_k^γ , the future skeleton factors that have not been touched by the previous skeleton factors $S_0, X_1^\gamma, \dots, X_k^\gamma$. Thus when tracing back through T , the irreducible skeleton has to pass the skeleton factors X_k^γ , not any of the factors contained in \mathcal{L}_k^γ . To summarize, each term on the right hand side, which is a casual tree $T \in [\gamma]$, appears on the left hand side with an identical prefactor due to $n = \ell + m_0 + \dots + m_\ell$, and appears only once.

(S127) then holds if the left hand side contains no more terms, i.e., each $T \in [\gamma]$ can be expressed as one term on the right hand side (when expanding \mathcal{L}_k^γ to factors). To prove this statement, consider all possible ‘‘candidates’’ that may have γ as its ordered irreducible skeleton. They must belong to terms in

$$(\mathcal{L})^{m_\ell} \mathcal{L}_{X_\ell^\gamma} \cdots (\mathcal{L})^{m_1} \mathcal{L}_{X_1^\gamma} (\mathcal{L})^{m_0} \mathcal{O}, \quad (\text{S128})$$

with m_0, \dots, m_ℓ being non-negative integers. However, not all terms in each \mathcal{L} are allowed, because they may modify the irreducible skeleton to another one rather than γ . For each $(\mathcal{L})^{m_k}$ before acting with $\mathcal{L}_{X_{k+1}^\gamma}$, there are three kinds of them. First, any factor not contained in S will make the irreducible skeleton different from γ , because it will not have S as its circumscribed simplex. Second, any factor not contained in R_k^γ defined in (S125) will also modify the irreducible skeleton, because it touches one of the faces f_p, \dots, f_{d+1} of S that ought to be touched at the first time by later skeleton factors $X_{k+1}^\gamma, \dots, X_\ell^\gamma$. Third, any factor Y that touches W_k^γ in (S124) is also not allowed. Suppose Y touches X_{k+3}^γ that does not overlap with γ_k : $X_{k+3}^\gamma \cap \gamma_k = \emptyset$. Then when tracing backwards the casual tree to find the irreducible skeleton, X_{k+3}^γ should be traced back to Y instead of some factor in $S_0, X_1^\gamma, \dots, X_{k+2}^\gamma$, making the irreducible skeleton different than γ . Indeed, these are the only three cases that are avoided in (S123), so any $T \in [\gamma]$ must appear on the right hand side of (S127).

As a result, we have constructed an one-to-one correspondence between the terms on the two sides of (S123), so the two sides are equal given that the prefactors always agree. We then ‘‘exponentiate’’ the right hand side using Lemma 5 in [59], to get

$$\sum_{T \in [\gamma]} \frac{t^n}{n!} \mathcal{L}_{X_n} \cdots \mathcal{L}_{X_1} \mathcal{O} = \int_{\mathbb{T}^\ell(t)} dt_1 \cdots dt_\ell e^{(t-t_\ell)\mathcal{L}_\ell^\gamma} \mathcal{L}_{X_\ell^\gamma} \cdots e^{(t_2-t_1)\mathcal{L}_1^\gamma} \mathcal{L}_{X_1^\gamma} e^{t_1\mathcal{L}_0^\gamma} \mathcal{O}. \quad (\text{S129})$$

(S121) then follows from (S126) and (S129). \square

Taking the operator norm of (S121) using $\|\mathcal{L}_X \mathcal{O}\| \leq 2 \|H_X\| \|\mathcal{O}\|$, we obtain:

Corollary 11. *The operator norm of (S121) is bounded by*

$$\|(e^{t\mathcal{L}}\mathcal{O})_S\| \leq \|\mathcal{O}\| \sum_{\Gamma \in \Gamma(S)} \frac{(2|t|)^\ell}{\ell!} N(\Gamma) \prod_{k=1}^{\ell} \|H_{X_k^\Gamma}\|, \quad (\text{S130})$$

where $N(\Gamma)$ is the number of ordered γ that corresponds to Γ .

4.3. Proof of Proposition 6 (restatement)

Proof of Proposition 6 (restatement). We assume $t \geq 0$ without loss of generality in the remainder of this section. To bound $\|e^{t\mathcal{L}}\mathcal{O} - \mathcal{O}\|_{\kappa'}$, we choose the local decomposition for $e^{t\mathcal{L}}\mathcal{O}$ in the previous section, namely

$$(e^{t\mathcal{L}}\mathcal{O})_S = \sum_{S_0 \subseteq S} (e^{t\mathcal{L}}\mathcal{O}_{S_0})_S, \quad (\text{S131})$$

where each term can be further expanded to irreducible skeletons (S121). The decomposition $\mathcal{O} = \sum_{S_0} \mathcal{O}_{S_0}$ is chosen as the optimal one that realizes $\|\mathcal{O}\|_{\kappa}$. The bound for this decomposition would immediately lead to a bound on the infimum (S13) over all possible decompositions of $\|e^{t\mathcal{L}}\mathcal{O} - \mathcal{O}\|_{\kappa'}$. We divide the terms in (S131) into two cases: $S_0 = S$ for case 1 and $S_0 \neq S$ for case 2. We will further divide case 2 to 2A and 2B. We prove this Proposition 6 by showing that the contribution to $\|e^{t\mathcal{L}}\mathcal{O} - \mathcal{O}\|_{\kappa',i}$ from each of the three cases, is bounded by the form of the right hand side of (S115) that does not depend on i . Recall that $\|e^{t\mathcal{L}}\mathcal{O} - \mathcal{O}\|_{\kappa',i}$ defined in (S84), probes the κ' -norm of $e^{t\mathcal{L}}\mathcal{O} - \mathcal{O}$ at site $i \in \Lambda$. The bound (S115) for $\|e^{t\mathcal{L}}\mathcal{O} - \mathcal{O}\|_{\kappa'}$ then follows by a trivial maximization over i . We begin with the simplest case 1.

Case 1: If $S_0 = S$, we have

$$(e^{t\mathcal{L}}\mathcal{O}_S)_S = e^{t\mathcal{L}|_S} \mathcal{O}_S, \quad (\text{S132})$$

where

$$\mathcal{L}|_S := \sum_{X \subseteq S} \mathcal{L}_X. \quad (\text{S133})$$

The corresponding local terms H_X are chosen as the optimal decomposition that realizes $\|H\|_{\kappa}$. (S132) cancels \mathcal{O}_S at zeroth order of t :

$$\|(e^{t\mathcal{L}}\mathcal{O}_S)_S - \mathcal{O}_S\| = \|e^{t\mathcal{L}|_S} \mathcal{O}_S - \mathcal{O}_S\| \leq 2t \|H|_S\| \|\mathcal{O}_S\| \leq 2t \|\mathcal{O}_S\| \sum_{j \in S} \sum_{X \ni j} \|H_X\| \leq 2t \|\mathcal{O}_S\| |S| \|H\|_{\kappa}, \quad (\text{S134})$$

which follows the treatment in (S66).

An operator $(e^{t\mathcal{L}}\mathcal{O}_S)_S - \mathcal{O}_S$ can contribute to $\|e^{t\mathcal{L}}\mathcal{O} - \mathcal{O}\|_{\kappa',i}$ only if $S \ni i$, so the total contribution of such ‘‘local rotations’’ (case 1 terms in (S131)) to $\|e^{t\mathcal{L}}\mathcal{O} - \mathcal{O}\|_{\kappa',i}$ is

$$\begin{aligned} \sum_{S \ni i} \|(e^{t\mathcal{L}}\mathcal{O}_S)_S - \mathcal{O}_S\| e^{\kappa'(\text{diam}S)^\alpha} &\leq 2t \|H\|_{\kappa} \sum_{S \ni i} |S| \|\mathcal{O}_S\| e^{\kappa'(\text{diam}S)^\alpha} \\ &\leq 2c_{\text{vol}} t \|H\|_{\kappa} \sum_{S \ni i} (\delta\kappa)^{-d/\alpha} e^{\delta\kappa(\text{diam}S)^\alpha} \|\mathcal{O}_S\| e^{\kappa'(\text{diam}S)^\alpha} \\ &= 2c_{\text{vol}} (\delta\kappa)^{-d/\alpha} t \|H\|_{\kappa} \|\mathcal{O}\|_{\kappa}. \end{aligned} \quad (\text{S135})$$

Here we have used (S134) and (S3):

$$|S| \leq c_d (1 + \text{diam}S)^d \leq c_{\text{vol}} (\delta\kappa)^{-d/\alpha} e^{\delta\kappa(\text{diam}S)^\alpha}, \quad (\text{S136})$$

for some constant c_{vol} that only depends on d and α . We also used the optimality of decomposition \mathcal{O}_S in the final step. Therefore, the case 1 contribution (S135) to $\|e^{t\mathcal{L}}\mathcal{O} - \mathcal{O}\|_{\kappa',i}$ is bounded by the right hand side of (S115), since $\delta\kappa$ is bounded from above by an $O(1)$ constant, so we can always make the power of $\delta\kappa$ more negative:

$$c_{\text{vol}} (\delta\kappa)^{-d/\alpha} \leq c_{\text{vol}} \left(\frac{\kappa_1}{\delta\kappa}\right)^{\frac{d-1}{\alpha}} (\delta\kappa)^{-d/\alpha} = c'_{\text{vol}} (\delta\kappa)^{-\frac{2d-1}{\alpha}}. \quad (\text{S137})$$

Separate case 2 to 2A and 2B: For a given \mathcal{O}_{S_0} , define its contribution $K_{\kappa',i}(\mathcal{O}_{S_0})$ to $\|e^{t\mathcal{L}}\mathcal{O} - \mathcal{O}\|_{\kappa',i}$ that is beyond the local rotation above, as

$$\begin{aligned} K_{\kappa',i}(\mathcal{O}_{S_0}) &:= \sum_{S \ni i: S_0 \subsetneq S} \|(e^{t\mathcal{L}}\mathcal{O}_{S_0})_S\| e^{\kappa'(\text{diam}S)^\alpha} = \sum_{S \ni i: S_0 \subsetneq S} \|(e^{t\mathcal{L}}\mathcal{O}_{S_0})_S\| e^{(\kappa'' - \frac{\delta\kappa}{2})(\text{diam}S)^\alpha} \\ &\leq e^{-\frac{\delta\kappa}{2}d^\alpha(i,S_0)} \sum_{S: S_0 \subsetneq S} \|(e^{t\mathcal{L}}\mathcal{O}_{S_0})_S\| e^{\kappa''(\text{diam}S)^\alpha}, \end{aligned} \quad (\text{S138})$$

where we have defined

$$\kappa'' = \kappa' + \frac{\delta\kappa}{2} = \kappa - \frac{\delta\kappa}{2}, \quad (\text{S139})$$

and used Markov inequality like (S107) with $\text{diam}S \geq d(i, S_0)$, since $\{i\} \cup S_0 \subseteq S$. We have also relaxed the restriction that S must contain i , since the prefactor $e^{-\frac{\delta\kappa}{2}d^\alpha(i,S_0)}$ already suppresses contributions from S_0 that are faraway from i . According to the irreducible skeleton expansion (S121), (S138) becomes

$$e^{\frac{\delta\kappa}{2}d^\alpha(i,S_0)} K_{\kappa',i}(\mathcal{O}_{S_0}) \leq \sum_{\ell=1}^{\infty} \sum_{\Gamma: \ell(\Gamma)=\ell} \|(e^{t\mathcal{L}}\mathcal{O}_{S_0})_\Gamma\| e^{\kappa''(\text{diam}S(\Gamma))^\alpha} \equiv \sum_{\ell=1}^{\infty} K_\ell(\mathcal{O}_{S_0}), \quad (\text{S140})$$

where $K_\ell(\mathcal{O}_{S_0})$ contains contributions from irreducible skeletons of length ℓ . Note that $K_\ell(\mathcal{O}_{S_0})$ no longer cares about whether i is reached or not. We further separate (S140) to the sum of $K_1(\mathcal{O}_{S_0})$ and $\sum_{\ell=2}^{\infty} K_\ell(\mathcal{O}_{S_0})$, whose contributions to $\|e^{t\mathcal{L}}\mathcal{O} - \mathcal{O}\|_{\kappa',i}$ are called case 2A and 2B respectively.

Case 2A: We compute $K_1(\mathcal{O}_{S_0})$ using (S130), where each irreducible skeleton Γ corresponds to a single factor X that grows S_0 to a strictly larger simplex $S(X)$ so that $N(\Gamma) = 1$.

$$\begin{aligned} K_1(\mathcal{O}_{S_0}) &= \sum_{\Gamma: \ell(\Gamma)=1} \|(e^{t\mathcal{L}}\mathcal{O}_{S_0})_\Gamma\| e^{\kappa''(\text{diam}S(\Gamma))^\alpha} \\ &\leq 2t \|\mathcal{O}_{S_0}\| \sum_{X: X \cap S_0 \neq \emptyset, X \cap S_0^c \neq \emptyset} \|H_X\| e^{\kappa''(\text{diam}S(X))^\alpha} \\ &\leq 2t \|\mathcal{O}_{S_0}\| \sum_{X: X \cap S_0 \neq \emptyset, X \cap S_0^c \neq \emptyset} \|H_X\| e^{\kappa''[(\text{diam}S_0)^\alpha + (\text{diam}X)^\alpha]} \\ &\leq 2t \|\mathcal{O}_{S_0}\| \sum_{j \in \partial S_0} \sum_{X \ni j} \|H_X\| e^{\kappa''[(\text{diam}S_0)^\alpha + (\text{diam}X)^\alpha]} \\ &\leq 2t \|\mathcal{O}_{S_0}\| \|\partial S_0\| \|H\|_\kappa e^{\kappa''(\text{diam}S_0)^\alpha}. \end{aligned} \quad (\text{S141})$$

Here in the third line, we have used (S98) with $\text{diam}S(X) \leq \text{diam}S_0 + \text{diam}X$. In the fourth line, we have used the fact that every X that grows S_0 must contain at least one vertex j in the boundary ∂S_0 .

We then bound the contribution from all $K_1(\mathcal{O}_{S_0})$ to $\|e^{t\mathcal{L}}\mathcal{O} - \mathcal{O}\|_{\kappa',i}$, following the same strategy of (S101). Namely, all \mathcal{O}_{S_0} are grouped by their distance x to the site i , and the site $j \in S_0$ that has distance x to i . Using (S141), the contribution is

$$\begin{aligned} \sum_{\mathcal{O}_{S_0}} e^{-\frac{\delta\kappa}{2}d^\alpha(i,S_0)} K_1(\mathcal{O}_{S_0}) &\leq \sum_{x=0}^{\infty} e^{-\frac{\delta\kappa}{2}x^\alpha} \sum_{j: d(i,j)=x} \sum_{\mathcal{O}_{S_0}} 2t \|\mathcal{O}_{S_0}\| \|\partial S_0\| \|H\|_\kappa e^{\kappa''(\text{diam}S_0)^\alpha} \\ &\leq 2c_1 t (\delta\kappa)^{-\frac{d-1}{\alpha}} \|H\|_\kappa \sum_{x=0}^{\infty} e^{-\frac{\delta\kappa}{2}x^\alpha} \sum_{j: d(i,j)=x} \sum_{S_0 \ni j} \|\mathcal{O}_{S_0}\| e^{\frac{\delta\kappa}{2}(\text{diam}S_0)^\alpha} e^{\kappa''(\text{diam}S_0)^\alpha} \\ &\leq 2c_1 t (\delta\kappa)^{-\frac{d-1}{\alpha}} \|H\|_\kappa \sum_{x=0}^{\infty} e^{-\frac{\delta\kappa}{2}x^\alpha} \sum_{j: d(i,j)=x} \|\mathcal{O}\|_\kappa \\ &\leq 2c_1 t (\delta\kappa)^{-\frac{d-1}{\alpha}} \|H\|_\kappa \|\mathcal{O}\|_\kappa \sum_{x=0}^{\infty} c_d (2x+1)^{d-1} e^{-\frac{\delta\kappa}{2}x^\alpha} \\ &\leq c_{2A} t (\delta\kappa)^{-\frac{2d-1}{\alpha}} \|H\|_\kappa \|\mathcal{O}\|_\kappa, \end{aligned} \quad (\text{S142})$$

bounded by the form of the right hand side of (S115). Here in the second line, we have used the surface analog of (S136) with numerical factors adjusted:

$$|\partial S_0| \leq c_d(1 + \text{diam}S_0)^{d-1} \leq c_1(\delta\kappa)^{-\frac{d-1}{\alpha}} e^{\frac{\delta\kappa}{2}(\text{diam}S_0)^\alpha}. \quad (\text{S143})$$

In the third line we have used (S139) and the κ -norm of \mathcal{O} . (S104) is used in the fourth line. The final sum over x in (S142) is bounded by transforming to an integral, and rescaling $x \rightarrow (\delta\kappa)^{-1/\alpha}x$, so that

$$\sum_{x=0}^{\infty} (2x+1)^{d-1} e^{-\frac{\delta\kappa}{2}x^\alpha} \leq \frac{c_{2A}}{2c_1c_d} (\delta\kappa)^{-d/\alpha}, \quad (\text{S144})$$

for a constant c_{2A} determined by d, α and κ_1 .

Case 2B: It is more tedious to find higher orders $K_\ell(\mathcal{O}_{S_0})$, because each irreducible skeleton has many branches. Fortunately, the assumption (S114) simplifies the analysis by cancelling powers of $\delta\kappa$ from geometrical factors like $|S|$ and $|\partial S|$. This is achieved in the following Lemma, whose proof is the goal of later subsections.

Lemma 12. *If (S114) holds, then there exists a constant c_K depending on d, α and κ_1 , such that*

$$\sum_{\ell=2}^{\infty} K_\ell(\mathcal{O}_{S_0}) := \sum_{\ell=2}^{\infty} \sum_{\Gamma: \ell(\Gamma)=\ell} \|(e^{t\mathcal{L}}\mathcal{O}_{S_0})_\Gamma\| e^{\kappa''(\text{diam}S(\Gamma))^\alpha} \leq c_K t \|\mathcal{O}_{S_0}\| \|H\|_\kappa e^{\kappa(\text{diam}S_0)^\alpha}. \quad (\text{S145})$$

Comparing to (S141), (S145) no longer has the surface factor $|\partial S_0|$. As a result, the contribution to $\|e^{t\mathcal{L}}\mathcal{O} - \mathcal{O}\|_{\kappa',i}$ in this 2B case is bounded by the same procedure of (S142):

$$\sum_{\mathcal{O}_{S_0}} e^{-\frac{\delta\kappa}{2}d^\alpha(i, S_0)} \sum_{\ell=2}^{\infty} K_\ell(\mathcal{O}_{S_0}) \leq c_{2B} t (\delta\kappa)^{-\frac{d}{\alpha}} \|H\|_\kappa \|\mathcal{O}\|_\kappa, \quad (\text{S146})$$

which is smaller than (S142) by a factor $(\delta\kappa)^{\frac{d-1}{\alpha}}$. In fact, the exponent in (S114) is deliberately chosen such that (S146) is of the same order as (S135) for case 1, so that case 2A dominates at small $\delta\kappa$.

Summarizing the three cases 1, 2A and 2B, (S135), (S142) and (S146) combine to produce (S115), with the constant c_- depending only on d, α and κ_1 . \square

4.4. Bounding the contributions of irreducible skeletons

The rest of the subsections serve to prove Lemma 12, where we need to sum over all irreducible skeletons Γ , starting from a fixed local operator $\mathcal{O} = \mathcal{O}_{S_0}$. In this subsection, we transform this sum into more tractable sums over factors and paths. Recall that each skeleton Γ is topologically a tree that have branches to reach all faces of the final simplex $S = S(\Gamma)$. Thus, we can identify the unique set of bifurcation factors

$$Y = \{Y_0 \equiv S_0, Y_1, \dots, Y_{l(Y)}\} \subseteq \{S_0, X_1^f, \dots, X_\ell^f\}, \quad (\text{S147})$$

as the internal vertices (including the root) of the tree Γ , while the faces f_1, \dots, f_{d+1} are terminal vertices. In Fig. S3 for example, $Y = \{Y_0, Y_1\}$ where $Y_1 = X_1$. $l(Y) = |Y| - 1 \leq d$ is the number of distinct factors in the sequence at which the path to another face bifurcates (see Figure S4 for an illustration); if $l(Y) = 0$ it means that the paths never share any common Y_n s. We do not care about the labeling order in Y_n , but keep track of their connection. I.e., for each Y_n , define $\mathfrak{p}(Y_n)$ as its parent vertex in the tree, which is some bifurcation factor $Y_{n'}$. Similarly define $\mathfrak{p}(f_p)$ as the parent bifurcation factor of face f_p . In Γ , f_p is then reached by an irreducible path \mathcal{P}_p that starts from $\mathfrak{p}(f_p)$. Each such path \mathcal{P}_p is a single branch of the skeleton Γ . Each bifurcation factor Y_n is also connected to its parent $\mathfrak{p}(Y_n)$ by an irreducible path \mathcal{P}'_n . See Fig. S4(a) as a sketch in 2d. We let $\mathbf{Y}(S)$ be all possible sets Y of bifurcation factors that support a $\Gamma \in \mathbf{\Gamma}(S)$.

Furthermore, we define the sum over paths that bounds operator growth between two given sets $Z, Z' \subseteq \Lambda$ (not necessarily simplices). There are irreducible paths \mathcal{P} in the factor graph G that connects them, which we label by $\mathcal{P} : Z \rightarrow Z'$. We define the sum over such irreducible paths

$$C_l(Z, Z') := \sum_{\mathcal{P}: Z \rightarrow Z', \ell(\mathcal{P})=l} \frac{(2t)^\ell}{\ell!} \prod_{k=1}^{\ell} \|H_{X_k^{\mathcal{P}}}\|, \quad (\text{S148})$$




FIG. S4. (a) A 2d sketch of bifurcation factors Y_p and irreducible paths $\mathcal{P}_p, \mathcal{P}'_n$ that comprise an irreducible skeleton Γ . The irreducible paths Γ_p from S_0 to the three faces are also denoted by lines of three different colors. (b) A 2d sketch of why the inequality (S189) holds. In particular, here the $l(Y) = 1$ case takes the equality, guided by the red dashed lines.

as a bound for the relative amplitude that an operator in Z can grow to Z' , where $(X_1^{\mathcal{P}}, \dots, X_\ell^{\mathcal{P}})$ are the factors in \mathcal{P} . Note that we also use the parameter l to control the length of a path that can contribute. Depending on whether Z intersects with Z' , we have

$$C_l(Z, Z') \begin{cases} = \delta_{l0}, & Z \cap Z' \neq \emptyset \\ \propto (1 - \delta_{l0}), & Z \cap Z' = \emptyset \end{cases} \quad (\text{S149})$$

With the definitions above and the short hand notation

$$J(S) := \sum_{\Gamma \in \mathbf{\Gamma}(S): l(\Gamma) > 1,} \|(e^{t\mathcal{L}}\mathcal{O})_{\Gamma}\|, \quad (\text{S150})$$

we prove the following Lemma. Namely, the sum over skeletons in (S121) is bounded by first summing over the bifurcation factors, and then summing over the irreducible paths that connect them to the faces of S and to themselves, with the constraint that the length ℓ of the skeleton should agree with the number of factors contained in the bifurcation factors and the paths.

Lemma 13. *The following identities hold:*

$$J(S) \leq (d+1)! \|\mathcal{O}\| \sum_{Y \in \mathbf{Y}(S)} (2t)^{l(Y)} \left(\prod_{n=1}^{l(Y)} \|H_{Y_n}\| \right) \mathcal{G}(Y), \quad \text{where}$$

$$\mathcal{G}(Y) := \sum_{\substack{l_1, \dots, l_{d+1}, l'_1, \dots, l'_{l(Y)} \geq 0: \\ l(Y) + l_1 + \dots + l_{d+1} + l'_1 + \dots + l'_{l(Y)} > 1}} \left[\prod_{n=1}^{l(Y)} C_{l'_n}(Y_n, \mathbf{p}(Y_n)) \right] \left[\prod_{p=1}^{d+1} C_{l_p}(f_p, \mathbf{p}(f_p)) \right]. \quad (\text{S151})$$

Here $\{f_1, \dots, f_{d+1}\}$ are the faces of S , and $Y = \{Y_0, \dots, Y_{l(Y)}\}$ is the set of bifurcation factors in (S147).

Proof. Suppose $\|\mathcal{O}\| = 1$. According to (S150) and (S130),

$$\begin{aligned}
J(S) &\leq \sum_{\Gamma \in \Gamma(S): \ell(\Gamma) \equiv \ell > 1} N(\Gamma) \frac{(2t)^\ell}{\ell!} \prod_{k=1}^{\ell} \|H_{X_k^\Gamma}\| \\
&\leq (d+1)! \sum_{Y \in \mathbf{Y}(S)} \sum_{\mathcal{P}_1: \mathbf{p}(f_1) \rightarrow f_1} \cdots \sum_{\mathcal{P}_{d+1}: \mathbf{p}(f_{d+1}) \rightarrow f_{d+1}} \sum_{\mathcal{P}'_1: \mathbf{p}(Y_1) \rightarrow Y_1} \cdots \sum_{\mathcal{P}'_{l(Y)}: \mathbf{p}(Y_{l(Y)}) \rightarrow Y_{l(Y)}} N(\Gamma) \frac{(2t)^\ell}{\ell!} \prod_{k=1}^{\ell} \|H_{X_k^\Gamma}\| \mathbb{I}(\ell > 1) \\
&= (d+1)! \sum_{Y \in \mathbf{Y}(S)} (2t)^{l(Y)} \left(\prod_{n=1}^{l(Y)} \|H_{Y_n}\| \right) \\
&\quad \left(\sum_{\mathcal{P}_1: \mathbf{p}(f_1) \rightarrow f_1} (2t)^{l_1} \prod_{k=1}^{l_1} \|H_{X_k^{\mathcal{P}_1}}\| \right) \cdots \left(\sum_{\mathcal{P}_{d+1}: \mathbf{p}(f_{d+1}) \rightarrow f_{d+1}} (2t)^{l_{d+1}} \prod_{k=1}^{l_{d+1}} \|H_{X_k^{\mathcal{P}_{d+1}}}\| \right) \\
&\quad \left(\sum_{\mathcal{P}'_1: \mathbf{p}(Y_1) \rightarrow Y_1} (2t)^{l'_1} \prod_{k=1}^{l'_1} \|H_{X_k^{\mathcal{P}'_1}}\| \right) \cdots \left(\sum_{\mathcal{P}'_{l(Y)}: \mathbf{p}(Y_{l(Y)}) \rightarrow Y_{l(Y)}} (2t)^{l'_{l(Y)}} \prod_{k=1}^{l'_{l(Y)}} \|H_{X_k^{\mathcal{P}'_{l(Y)}}}\| \right) \frac{N(\Gamma)}{\ell!} \mathbb{I}(\ell > 1). \tag{S152}
\end{aligned}$$

In the second line, we have disintegrated each skeleton Γ as its bifurcation factors Y_n and branch irreducible paths $\mathcal{P}_p, \mathcal{P}'_n$. $\mathbb{I}(\cdot)$ is the indicator function that returns 1 if the input is True and 0 if False, so that only Γ with $\ell = \ell(\Gamma) > 1$ is included in the sum. The prefactor $(d+1)!$ comes from the number of sequences of faces $\{f_1, \dots, f_{d+1}\}$ that a skeleton reaches in order. Note that the inequality is loose in general, because certain pairs of irreducible paths like \mathcal{P}'_n and \mathcal{P}_p that starts from Y_n , are allowed to intersect with each other here, while they are not allowed to intersect as two consecutive branches of the irreducible skeleton Γ . In the third line of (S152), we have moved factors around (each factor X_k^Γ belongs to either an irreducible path \mathcal{P}_p or \mathcal{P}'_n , or the bifurcation factors Y), and used

$$\ell = l(Y) + \sum_{p=1}^{d+1} l_p + \sum_{n=1}^{l(Y)} l'_n, \tag{S153}$$

where l_p (l'_n) is the length of \mathcal{P}_p (\mathcal{P}'_n). Now compare (S152) with the goal (S151). The indicator function $\mathbb{I}(\ell > 1)$ matches exactly the sum rule $l(Y) + l_1 + \dots + l_{d+1} + l'_1 + \dots + l'_{l(Y)} > 1$ in (S151), so (S152) implies (S151) as long as

$$\frac{N(\Gamma)}{\ell!} \leq \left(\prod_{p=1}^{d+1} \frac{1}{l_p!} \right) \left(\prod_{n=1}^{l(Y)} \frac{1}{l'_n!} \right), \tag{S154}$$

because then the $\frac{1}{l_p!}$ ($\frac{1}{l'_n!}$) factor can be moved into the sum over \mathcal{P}_p (\mathcal{P}'_n), and each sum over paths independently yields a C_l in (S151).

In the remaining proof, we verify (S154) by the recursion structure of Γ as a tree. Any tree $\tilde{\Gamma}$ can be viewed as the root connecting to some subtrees $\tilde{\Gamma}_m$. For example, the tree Γ in Fig. S3(d) has three subtrees: a subtree $\tilde{\Gamma}_1$ with root Y_1 , a trivial tree $\tilde{\Gamma}_2 = f_2$, and a subtree $\tilde{\Gamma}_3$ with root Y_2 . They are all connected to Y_0 , the root of Γ . $\tilde{\Gamma}_1$ has a further subtree $\tilde{\Gamma}_4$ with root Y_3 . Each subtree $\tilde{\Gamma}_m$ also corresponds to an irreducible skeleton (although it does not necessarily start from S_0), so it has length $\ell(\tilde{\Gamma}_m)$, and $N(\tilde{\Gamma}_m)$ as the number of corresponding ordered irreducible skeletons. We first prove that, if a tree $\tilde{\Gamma}_0$ with root \tilde{Y}_0 is composed by subtrees $\tilde{\Gamma}_m : m = 1, \dots, m_0$, whose roots \tilde{Y}_m are directly connected to \tilde{Y}_0 by irreducible paths $\tilde{\mathcal{P}}_m$ of length $\tilde{\ell}_m$, then

$$\frac{N(\tilde{\Gamma}_0)}{\ell(\tilde{\Gamma}_0)!} \leq \frac{1}{\ell(\tilde{\Gamma}_0)!} \frac{\left(\sum_{m=1}^{m_0} \tilde{\ell}_m + \ell(\tilde{\Gamma}_m) \right)!}{\prod_{m=1}^{m_0} \left(\tilde{\ell}_m + \ell(\tilde{\Gamma}_m) \right)!} \prod_{m=1}^{m_0} N(\tilde{\Gamma}_m) \leq \left(\prod_{m=1}^{m_0} \frac{1}{\tilde{\ell}_m!} \right) \prod_{m=1}^{m_0} \frac{N(\tilde{\Gamma}_m)}{\ell(\tilde{\Gamma}_m)!}. \tag{S155}$$

The second inequality comes from $\ell(\tilde{\Gamma}_0) \geq \sum_{m=1}^{m_0} \tilde{\ell}_m + \ell(\tilde{\Gamma}_m)$, and

$$(a+b)! \geq a! b!, \quad \forall a, b > 0. \tag{S156}$$

The first inequality of (S155) comes from the following. Starting from \tilde{Y}_0 there are m_0 branches, where factors of different branches can stagger in an ordered skeleton. Thus assuming the order within each branch is fixed, there are

at most

$$\frac{\left(\sum_{m=1}^{m_0} \tilde{\ell}'_m\right)!}{\prod_{m=1}^{m_0} \tilde{\ell}'_m!}, \quad (\text{S157})$$

number of ways to stagger the m_0 branches, where $\tilde{\ell}'_m = \tilde{\ell}_m + \ell(\tilde{T}_m)$ is the total number of factors in the m -th branch. The multinomial coefficient can be understood as the number of orders to place m_0 groups of balls in a line, where each group labeled by m has $\tilde{\ell}'_m$ identical balls, while balls in different groups are distinguishable. The number of inter-branch orders may be strictly smaller than (S157), because some ordering may produce a γ that is not a skeleton. In Fig. S3(a) for example, $X_5 \in \mathcal{P}_3$ must act after $X_3 \in \mathcal{P}_2$, since otherwise f_2 is already reached by X_5 and X_3 is no longer needed for a skeleton.

Now we let the relative order within each branch to vary. The intra-branch order for the m -th branch has exactly $N(\tilde{T}_m)$ possibilities, since the irreducible path $\tilde{\mathcal{P}}_m$ must act in a fixed order, followed by factors in \tilde{T}_m that have $N(\tilde{T}_m)$ number of different orders. Because the intra-branch order is independent of the inter-branch order counted by (S157), the total number of ordering is the product of them, and (S155) is established.

Finally, we show (S155) yields (S154). Set $\tilde{T}_0 = \Gamma$, we have

$$\begin{aligned} \frac{N(\Gamma)}{\ell(\Gamma)!} &\leq \left(\prod_{m_1 \in \text{st}(\Gamma)} \frac{1}{\tilde{\ell}_{m_1}!} \right) \prod_{m_1 \in \text{st}(\Gamma)} \frac{N(\tilde{T}_{m_1})}{\ell(\tilde{T}_{m_1})!} \\ &\leq \left(\prod_{m_1 \in \text{st}(\Gamma)} \frac{1}{\tilde{\ell}_{m_1}!} \right) \prod_{m_1 \in \text{st}(\Gamma)} \left[\left(\prod_{m_2 \in \text{st}(\tilde{T}_{m_1})} \frac{1}{\tilde{\ell}_{m_2}!} \right) \prod_{m_2 \in \text{st}(\tilde{T}_{m_1})} \frac{N(\tilde{T}_{m_2})}{\ell(\tilde{T}_{m_2})!} \right] \leq \dots \\ &\leq \left(\prod_{p=1}^{d+1} \frac{1}{l_p!} \right) \left(\prod_{n=1}^{l(Y)} \frac{1}{l'_n!} \right), \end{aligned} \quad (\text{S158})$$

where $m \in \text{st}(\tilde{T})$ represents all subtrees \tilde{T}_m connecting to the root of \tilde{T} , and we have used (S155) recursively to smaller and smaller subtrees. The iteration terminates when all subtrees \tilde{T} become trivial: $\frac{N(\tilde{T})}{\ell(\tilde{T})!} = 1$, and what remain are exactly one factor of either $\frac{1}{l_p!}$ or $\frac{1}{l'_n!}$ for each of the edges of the tree Γ . \square

4.5. Sum over irreducible paths: Lieb-Robinson bound with sub-exponential tails

Next, we prove a Lieb-Robinson bound for $C_l(Z, Z')$ with a Hamiltonian $H = \sum H_S$, where $\|H_S\| \sim e^{-r^\alpha}$ has subexponential decay. Amusingly, this goal is rather elegant to achieve, as it turns out that the sub-exponential decay $e^{-\kappa r^\alpha}$ with $\alpha < 1$ is reproducing (a technical property explained below),⁶ while the faster decay with $\alpha = 1$ is not.

Proposition 14 (Lieb-Robinson bound for interactions decaying with a stretched exponential). *The commutator of two local operators, one time-evolved, is bounded by the irreducible path summation defined in (S148):*

$$\|[\mathcal{O}_Z(t), \mathcal{O}'_{Z'}]\| \leq 2 \|\mathcal{O}_Z\| \|\mathcal{O}'_{Z'}\| \sum_{l \geq 0} C_l(Z, Z'). \quad (\text{S159})$$

Suppose the interaction decays as a stretched exponential, which is guaranteed by

$$\|H\|_{\alpha, \kappa} < \infty. \quad (\text{S160})$$

There exists

$$\lambda = c_\lambda \kappa^{-\frac{d+1}{\alpha}}, \quad (\text{S161})$$

with c_λ determined by d and α , such that

$$\sum_{l \geq 1} C_l(Z, Z') \leq \min(|\partial Z|, |\partial Z'|) e^{-\kappa d^\alpha(Z, Z')} \lambda^{-1} \left(e^{2\lambda t \|H\|_\kappa} - 1 \right), \quad (\text{S162})$$

⁶ This appears related to the fact that power-law decaying interactions are reproducing.

and

$$\sum_{l \geq 2} C_l(Z, Z') \leq \min(|\partial Z|, |\partial Z'|) e^{-\kappa d^\alpha(Z, Z')} \lambda^{-1} \left(e^{2\lambda t \|H\|_\kappa} - 1 - 2\lambda t \|H\|_\kappa \right). \quad (\text{S163})$$

Proof. (S159) is proven in Theorem 3 of [59], so we focus on (S162) and (S163). (S148) can be bounded explicitly as

$$C_l(Z, Z') \leq \frac{(2t)^l}{l!} \sum_{Z_1: Z_1 \text{ grows } Z} \|H_{Z_1}\| \sum_{Z_2: Z_2 \text{ grows } Z_1} \|H_{Z_2}\| \cdots \sum_{Z_l: Z_l \text{ grows } Z_{l-1}, Z_l \cap Z' \neq \emptyset} \|H_{Z_l}\|, \quad (\text{S164})$$

where “ X grows Y ” means the condition $X \cap Y \neq \emptyset$ and $X \cap Y^c \neq \emptyset$ with Y^c being the complement of Y as a subset of Λ . We require $X \cap Y^c \neq \emptyset$ because otherwise X is a factor of local rotation that does not appear in irreducible paths. Nevertheless, the path (Z_1, \dots, Z_l) in (S164) is not necessarily irreducible because Z_3 may touch Z_1 for example, which makes (S164) a loose bound in general. Now we follow closely [61], starting at their Eq. (27). To simplify the sums in (S164) we may replace

$$\sum_{X: X \text{ grows } Y} \leq \sum_{i \in \partial Y} \sum_{X \ni i}, \quad (\text{S165})$$

because X must intersect with ∂Y in order to grow Y nontrivially.⁷ Now consider, the last term in (S164), which ends in:

$$\begin{aligned} & \sum_{i \in \partial Z_{l-2}} \sum_{Z_{l-1} \ni i: Z_{l-1} \cap Z_l \neq \emptyset} \sum_{j \in \partial Z_{l-1}} \sum_{Z_l \ni j: Z_l \cap Z' \neq \emptyset} \|H_{Z_{l-1}}\| \|H_{Z_l}\| \\ & \leq \sum_{i \in \partial Z_{l-2}} \sum_{Z_{l-1} \ni i: Z_{l-1} \cap Z_l \neq \emptyset} \sum_{j \in \partial Z_{l-1}} \sum_{Z_l \ni j} \|H_{Z_{l-1}}\| \|H_{Z_l}\| e^{-\kappa d^\alpha(j, Z') + \kappa \text{diam}(Z_l)^\alpha} \\ & \leq \sum_{i \in \partial Z_{l-2}} \sum_{Z_{l-1} \ni i: Z_{l-1} \cap Z_l \neq \emptyset} \sum_{j \in \partial Z_{l-1}} \|H_{Z_{l-1}}\| e^{-\kappa d^\alpha(i, j) + \kappa \text{diam}(Z_{l-1})^\alpha} \times e^{-\kappa d^\alpha(j, Z')} \|H\|_\kappa, \end{aligned} \quad (\text{S166})$$

where in the second line we used the identity $d(j, Z') \leq \text{diam}(Z_l)$, and subsequently relaxed the restriction that $Z_l \cap Z'$ was non-empty. In the third line, we summed over Z_l , and *also* began the process anew by including a similar identity $d(i, j) \leq \text{diam}(Z_{l-1})$, since both $i, j \in Z_{l-1}$. At this point, we will need to use the *reproducibility* ansatz (which we will prove a little later):

$$\sum_{m \in \Lambda} \mathcal{K}(d(i, m)) \mathcal{K}(d(m, j)) \leq \lambda \mathcal{K}(d(i, j)), \quad (\text{S167})$$

for some constant λ . Combining (S166) and (S167):

$$\begin{aligned} & \sum_{i \in \partial Z_{l-2}} \sum_{Z_{l-1} \ni i: Z_{l-1} \cap Z_l \neq \emptyset} \sum_{j \in \partial Z_{l-1}} \sum_{Z_l \ni j: Z_l \cap Z' \neq \emptyset} \|H_{Z_{l-1}}\| \|H_{Z_l}\| \\ & \leq \sum_{i \in \partial Z_{l-2}} \sum_{Z_{l-1} \ni i: Z_{l-1} \cap Z_l \neq \emptyset} \lambda e^{-\kappa d^\alpha(i, Z') + \kappa \text{diam}(Z_{l-1})^\alpha} \|H_{Z_{l-1}}\| \|H\|_\kappa \\ & \leq \sum_{i \in \partial Z_{l-2}} \lambda e^{-\kappa d^\alpha(i, Z')} \|H\|_\kappa^2. \end{aligned} \quad (\text{S168})$$

Hence the process iterates l times and we obtain (S162) by resumming a simple exponential and subtracting off the constant term. The surface factor $\min(|\partial Z|, |\partial Z'|)$ comes from our improvement (S165), and the fact that irreducible paths from Z to Z' are exactly those from Z' to Z , so we can pick whichever as the starting point. Furthermore, (S163) also follows by deleting the linear in t -term as well.

To complete the proof, it only remains to verify the reproducing property (S167). For any pair $i, j \in \Lambda$, we use a d -dimensional “prolate spheroidal coordinate” suited for the lattice Λ . Namely, for any $m \in \Lambda$, define its distance to the two “focal points” i and j as r_i and r_j , respectively. Further define

$$\sigma = r_i + r_j, \quad (\text{S169a})$$

$$r = \min(r_i, r_j) \leq \frac{\sigma}{2}. \quad (\text{S169b})$$

⁷ Recall that X is mandated to be a simplex, which is convex. Thus this intersection must be non-vanishing.

Suppose there are $M(\sigma, r)$ vertices m corresponding to a given (σ, r) . Then the left hand side of (S167) becomes

$$\sum_{m \in \Lambda} e^{-\kappa(r_i^\alpha + r_j^\alpha)} = \sum_{r=0}^{\infty} \sum_{\sigma=\max(r_0, 2r)}^{2r+r_0} M(\sigma, r) e^{-\kappa[r^\alpha + (\sigma-r)^\alpha]}, \quad (\text{S170})$$

where $r_0 = d(i, j)$, and we have used the triangle inequalities $\sigma = r_i + r_j \geq r_0$ and $\sigma - 2r = |r_i - r_j| \leq r_0$ to constrain the sum. To proceed, observe that

$$r^\alpha + (\sigma - r)^\alpha \geq \sigma^\alpha + (2 - 2^\alpha)r^\alpha. \quad (\text{S171})$$

This equation comes from bounding the function

$$g(\rho) := 1 + (\rho - 1)^\alpha - \rho^\alpha \geq g(2) = 2 - 2^\alpha, \quad \rho \equiv \frac{\sigma}{r} \geq 2, \quad (\text{S172})$$

as one can verify by taking the derivative to confirm $g(\rho)$ is monotonic increasing.

Combining (S170) with (S171),

$$\begin{aligned} \sum_{m \in \Lambda} e^{-\kappa(r_i^\alpha + r_j^\alpha)} &\leq \sum_{r=0}^{\infty} \sum_{\sigma=\max(r_0, 2r)}^{2r+r_0} M(\sigma, r) e^{-\kappa[\sigma^\alpha + (2-2^\alpha)r^\alpha]} \\ &\leq \sum_{r=0}^{\infty} \sum_{\sigma=\max(r_0, 2r)}^{2r+r_0} 2c_d(r+1)^{d-1} e^{-\kappa[\sigma^\alpha + (2-2^\alpha)r^\alpha]} \\ &\leq 2c_d \sum_{r=0}^{\infty} (2r+1)(r+1)^{d-1} e^{-\kappa[r_0^\alpha + (2-2^\alpha)r^\alpha]} \\ &\leq c_\lambda \kappa^{-\frac{d+1}{\alpha}} e^{-\kappa r_0^\alpha}. \end{aligned} \quad (\text{S173})$$

Here in the second line we have used (S104) that there are at most $c_d(r+1)^{d-1}$ sites that are distance r to a given site i or j (each of which must be counted leading to an additional multiplicative factor of 2). In the third line we have used $\sigma \geq r_0$ in the exponent, and that there are at most $2r+1$ values for σ to choose. We arrive at the last line analogously to (S144), where c_λ depends only on d and α .

To conclude, we have verified $\mathcal{K}(r) = e^{-\kappa r^\alpha}$ is reproducing, with λ given by (S161). (S162) and (S163) then follow from (S165) and (S167). \square

4.6. Proof of Lemma 12

Proof of Lemma 12. First, observe that a sufficient condition to prove (S145) is if, for any $\mathcal{O} = \mathcal{O}_{S_0}$ with $\|\mathcal{O}\| = 1$,

$$J(S) \leq c_J (\delta\kappa)^{-\frac{2d}{\alpha}} t^2 \|H\|_\kappa^2 e^{\frac{\delta\kappa}{2} s_0^\alpha} e^{-\tilde{\kappa}(s-s_0)^\alpha}, \quad (\text{S174})$$

where $s = \text{diam}S$, $s_0 = \text{diam}S_0$,

$$\tilde{\kappa} = \kappa - \frac{\delta\kappa}{4} = \kappa'' + \frac{\delta\kappa}{4}, \quad (\text{S175})$$

and c_J is a constant determined by d, α and κ_1 . The reason is that, using (S114),

$$\begin{aligned} \sum_{\ell \geq 2} K_\ell(\mathcal{O}) &= \sum_{S \supset S_0} e^{\kappa'' s^\alpha} J(S) \leq c_J (\delta\kappa)^{-\frac{2d}{\alpha}} t^2 \|H\|_\kappa^2 e^{\frac{\delta\kappa}{2} s_0^\alpha} \sum_{S \supset S_0} e^{\kappa'' s^\alpha} e^{-\tilde{\kappa}(s-s_0)^\alpha} \\ &\leq c_J (\delta\kappa)^{-\frac{2d}{\alpha}} t^2 \|H\|_\kappa^2 e^{\frac{\delta\kappa}{2} s_0^\alpha} \sum_{s > s_0} \tilde{c}_\Lambda (s-s_0)^d e^{\kappa'' [s_0^\alpha + (s-s_0)^\alpha]} e^{-\tilde{\kappa}(s-s_0)^\alpha} \\ &\leq c_J \tilde{c}'_\Lambda (\delta\kappa)^{-\frac{2d}{\alpha}} t^2 \|H\|_\kappa^2 e^{\kappa s_0^\alpha} (\delta\kappa)^{-\frac{d+1}{\alpha}} \leq c'_J t \|H\|_\kappa e^{\kappa s_0^\alpha}, \end{aligned} \quad (\text{S176})$$

which implies (S145) for \mathcal{O} with a general normalization $\|\mathcal{O}\|$. In the second line, we have used the fact that for a given S_0 and s , there are at most $\tilde{c}_\Lambda (s-s_0)^d$ simplices $S \supset S_0$ that has diameter s . (S98) is also used. In the third

line we have summed over $s - s_0$, and used (S114) to cancel extra powers of $(\delta\kappa)^{-1}$: observe that the form of (S114) was chosen precisely so that this identity might hold. The final prefactor c'_j then only depends on d, α and κ_1 .

Thus, it remains to prove the condition (S174). According to (S151), we separate the sum over $Y \in \mathbf{Y}(S)$ into two cases: $l(Y) = 0$ for case 1, where the $d + 1$ paths to faces originate independently from S_0 , and $l(Y) \geq 1$ for case 2, where $l(Y)$ is the number of nontrivial bifurcation factors defined in (S147). For each case, we prove below that the contribution to $J(S)$ is bounded by the right hand side of (S174), albeit with a case-dependent constant prefactor. We start with the simpler $l(Y) = 0$ case, where some technical results are also used for the later case.

Case 1: If $l(Y) = 0$, or equivalently $Y = \{S_0\}$, it suffices to bound $\mathcal{G}(Y)$ in the second line of (S151), which describes the $d + 1$ faces of S_0 growing independently to the corresponding parallel faces of S , via irreducible paths $\mathcal{P}_p : S_0 \rightarrow f_p$. Here $\mathbf{p}(f_p) = S_0$ because S_0 is the parent of all faces.

If S_0 shares a simplex vertex with S , such that only one face f_{d+1} of S_0 grows nontrivially, then $\mathcal{G}(Y)$ is bounded by (S163):

$$\begin{aligned} \mathcal{G}(Y) &= \sum_{l_{d+1} > 1} C_{l_{d+1}}(S_0, f_{d+1}) \leq |\partial S_0| e^{-\kappa(s-s_0)^\alpha} \lambda^{-1} \left(e^{2\lambda t \|H\|_\kappa} - 1 - 2\lambda t \|H\|_\kappa \right) \\ &\leq c_1 (\delta\kappa)^{-\frac{d-1}{\alpha}} e^{\frac{\delta\kappa}{2} s_0^\alpha} e^{-\kappa(s-s_0)^\alpha} \lambda^{-1} \left(e^{2\lambda t \|H\|_\kappa} - 1 - 2\lambda t \|H\|_\kappa \right) \\ &\leq c_1 (\delta\kappa)^{-\frac{d-1}{\alpha}} e^{\frac{\delta\kappa}{2} s_0^\alpha} e^{-\kappa(s-s_0)^\alpha} c_\lambda'' c_\lambda \kappa^{-\frac{d+1}{\alpha}} t^2 \|H\|_\kappa^2 \\ &\leq c_1 c_\lambda'' c_\lambda (\delta\kappa)^{-\frac{2d}{\alpha}} t^2 \|H\|_\kappa^2 e^{\frac{\delta\kappa}{2} s_0^\alpha} e^{-\kappa(s-s_0)^\alpha}. \end{aligned} \quad (\text{S177})$$

Here (S143) is used in the second line. In the third line, we have used that the argument of the exponential function is bounded by $O(1)$:

$$2\lambda t \|H\|_\kappa \leq 2c_\lambda \kappa^{\frac{3d+1-(d+1)}{\alpha}} \leq 2c_\lambda \kappa_1^{\frac{2d}{\alpha}}, \quad (\text{S178})$$

which holds by combining (S161), (S114) and

$$\delta\kappa < \kappa \leq \kappa_1. \quad (\text{S179})$$

The constant

$$c_\lambda'' = \max_{\xi \leq 2c_\lambda \kappa_1^{2d/\alpha}} \frac{e^{2\xi} - 1 - 2\xi}{\xi^2} = \frac{e^{2\xi} - 1 - 2\xi}{\xi^2} \Big|_{\xi=2c_\lambda \kappa_1^{2d/\alpha}} \quad (\text{S180})$$

is completely determined by d, α and κ_1 . We have also plugged in the definition of λ from (S161). In the last line of (S177), we have used (S179). Note that (S177) actually has a stronger exponential decay exponent (κ) than the $\tilde{\kappa}$ advertised in (S174).

If S_0 does not share a simplex vertex with S instead, there are $1 < q \leq d + 1$ faces that grow nontrivially. The $\sum_p l_p > 1$ summation condition in (S174) can be relaxed, so that the sum over l_p are independent for different p . For each of the q faces labeled by $p \in \{d + 2 - q, \dots, d + 1\}$, we invoke (S162) to get

$$\sum_{l \geq 1} C_l(S_0, f_p) \leq |\partial S_0| e^{-\kappa d^\alpha(S_0, f_p)} \lambda^{-1} \left(e^{2\lambda t \|H\|_\kappa} - 1 \right) \leq |\partial S_0| e^{-\kappa d^\alpha(S_0, f_p)} c'_\lambda t \|H\|_\kappa, \quad (\text{S181})$$

where we again used (S178) with c'_λ determined by d, α and κ_1 in a similar way to (S180). Combining all the p faces,

$$\begin{aligned} \mathcal{G}(Y) &\leq (c'_\lambda |\partial S_0| t \|H\|_\kappa)^q \exp \left[-\kappa \sum_{p=d+2-q}^{d+1} d^\alpha(S_0, f_p) \right] \\ &\leq c_q (\delta\kappa)^{-q \frac{d-1}{\alpha}} e^{\frac{\delta\kappa}{2} s_0^\alpha} (c'_\lambda t \|H\|_\kappa)^q \exp(-\kappa(s-s_0)^\alpha) \\ &\leq c_q (c'_\lambda)^q (\delta\kappa)^{-2 \frac{d-1}{\alpha}} t^2 \|H\|_\kappa^2 e^{\frac{\delta\kappa}{2} s_0^\alpha} e^{-\kappa(s-s_0)^\alpha} (\delta\kappa)^{(q-2) \frac{2d+2}{\alpha}}. \end{aligned} \quad (\text{S182})$$

Here in the second line we have used

$$|\partial S_0|^q \leq c_q (\delta\kappa)^{-q \frac{d-1}{\alpha}} e^{\frac{\delta\kappa}{2} s_0^\alpha}, \quad (\text{S183})$$

similar to (S143). We have also used

$$\sum_{p=d+2-q}^{d+1} d^\alpha(S_0, f_p) \geq (s-s_0)^\alpha, \quad (\text{S184})$$

which follows from Proposition 1. In the third line of (S182) we have used (S114).

Summarizing Case 1, either (S177) for $q = 1$, or (S182) for $q \geq 2$, is bounded by the form of (S174), considering $\delta\kappa < \kappa \leq \kappa_1$.

Case 2: If $l(Y) \geq 1$, recall that besides paths \mathcal{P}_p from bifurcation factor $\mathfrak{p}(f_p)$ to f_p , we also have paths \mathcal{P}'_n connecting $\mathfrak{p}(Y_n)$ to Y_n , where $\mathfrak{p}(Y_n)$ is the parent bifurcation factor of Y_n . Among them, at least one path must be nontrivial according to the sum rule in (S174) ($l'_p > 0$ and/or $l_p > 0$) i.e., the path connects two sets Z and Z' that do not overlap. For each nontrivial path from $Z \rightarrow Z'$, the bound (S162) becomes

$$\sum_{l \geq 1} C_l(Z, Z') \leq |\partial S| e^{-\kappa d^\alpha(Z, Z')} c'_\lambda t \|H\|_\kappa, \quad (\text{S185})$$

following (S181). For a trivial path with $Z \cap Z' \neq \emptyset$ instead, we have factor $\sum_{l \geq 0} C_l(Z, Z') = 1 = e^{-\kappa d^\alpha(Z, Z')}$ from (S149). Thus we only need to keep track of the number of paths $1 \leq q(Y) \leq 2d + 1$ that are nontrivial, and do not need to track of which individual path is trivial or not, as if $d(\mathfrak{p}(f), f) = 0$ (e.g.) it just contributes trivially to

$$\mathcal{G}(Y) \leq (|\partial S| c'_\lambda t \|H\|_\kappa)^{q(Y)} \exp \left\{ -\kappa \left[\sum_{p=1}^{d+1} d^\alpha(\mathfrak{p}(f_p), f_p) + \sum_{n=1}^{l(Y)} d^\alpha(\mathfrak{p}(Y_n), Y_n) \right] \right\}. \quad (\text{S186})$$

To proceed, we bound the exponent of (S186) by

$$\sum_{p=1}^{d+1} d^\alpha(\mathfrak{p}(f_p), f_p) + \sum_{n=1}^{l(Y)} d^\alpha(\mathfrak{p}(Y_n), Y_n) \geq (s - s_0)^\alpha - \sum_{n=1}^{l(Y)} s_n^\alpha, \quad (\text{S187})$$

where

$$s_n := \text{diam} Y_n. \quad (\text{S188})$$

(S187) comes from applying (S98) repeatedly to

$$\sum_{p=1}^{d+1} d(\mathfrak{p}(f_p), f_p) + \sum_{n=1}^{l(Y)} d(\mathfrak{p}(Y_n), Y_n) \geq s - \sum_{n=0}^{l(Y)} s_n, \quad (\text{S189})$$

which is derived as follows. First, we introduce some notations. Let $\text{ch}(Y_n)$ be the set of all children faces of Y_n in the tree Γ ,

$$\text{ch}(Y_n) = \{f_p : \mathfrak{p}(f_p) = Y_n\}, \quad (\text{S190})$$

and let $\text{de}(Y_n)$ be the set of all descendant faces of Y_n . Namely, any $f_p \in \text{de}(Y_n)$ is reached by an irreducible path Γ_p that passes Y_n . Similarly, define $\text{ch}'(Y_n)$ to be the set of all children bifurcation factors of Y_n :

$$\text{ch}'(Y_n) = \{Y_{n'} \in Y : \mathfrak{p}(Y_{n'}) = Y_n\}. \quad (\text{S191})$$

Observe the following geometric fact

$$d(\mathfrak{p}(Y_n), Y_n) + s_n + \sum_{f_p \in \text{de}(Y_n)} d(Y_n, f_p) \geq \sum_{f_p \in \text{de}(Y_n)} d(\mathfrak{p}(Y_n), f_p). \quad (\text{S192})$$

This is trivial for $\text{de}(Y_n) = \{f_1, \dots, f_{d+1}\}$, because then according to Proposition 1, the right hand side is not larger than s , while the left hand side is not smaller than s , which equals the sum of the last two terms. Otherwise if $\text{de}(Y_n) \neq \{f_1, \dots, f_{d+1}\}$, (S192) comes from triangle inequality

$$d(\mathfrak{p}(Y_n), Y_n) + s_n + d(Y_n, W_n) \geq d(\mathfrak{p}(Y_n), W_n), \quad (\text{S193})$$

with $W_n := \bigcap_{f_p \in \text{de}(Y_n)} f_p \neq \emptyset$, and

$$d(Y_{n'}, W_n) = \sum_{f_p \in \text{de}(Y_n)} d(Y_{n'}, f_p), \quad (\text{S194})$$

due to the Manhattan metric of the simplicial lattice.

(S192) gives a relation on the distance from a parent $p(Y_n)$ to one of its children factors Y_n . If Y_n contains further children factors $Y_{n'}$ – i.e. $\text{de}(Y_n) \neq \text{ch}(Y_n)$ – then (S192) also gives a relation that connects the distances from Y_n to $f_p \notin \text{ch}(Y_n)$, with the distances from $Y_{n'}$ s to these faces. Thus (S192) can be used recursively to eliminate all “indirect” distances $d(Y_n, f_p)$ where $f_p \in \text{de}(Y_n) \setminus \text{ch}(Y_n)$, with the “direct” ones $f_p \in \text{ch}(Y_n)$. To be specific, we start from the root $Y_0 = S_0$,

$$\begin{aligned}
s - s_0 &= \sum_{f_p \in \text{de}(Y_0)} d(Y_0, f_p) = \sum_{f_p \in \text{ch}(Y_0)} d(Y_0, f_p) + \sum_{Y_{n_1} \in \text{ch}'(Y_0)} \sum_{f_p \in \text{de}(Y_{n_1})} d(Y_0, f_p) \\
&\leq \sum_{f_p \in \text{ch}(Y_0)} d(Y_0, f_p) + \sum_{Y_{n_1} \in \text{ch}'(Y_0)} \left[d(Y_0, Y_{n_1}) + s_{n_1} + \sum_{f_p \in \text{de}(Y_{n_1})} d(Y_{n_1}, f_p) \right] \\
&= \sum_{f_p \in \text{ch}(Y_0)} d(p(f_p), f_p) + \sum_{Y_{n_1} \in \text{ch}'(Y_0)} [d(p(Y_{n_1}), Y_{n_1}) + s_{n_1}] + \sum_{Y_{n_1} \in \text{ch}'(Y_0)} \sum_{f_p \in \text{de}(Y_{n_1})} d(Y_{n_1}, f_p) \\
&\leq \sum_{f_p \in \text{ch}(Y_0)} d(p(f_p), f_p) + \sum_{Y_{n_1} \in \text{ch}'(Y_0)} [d(p(Y_{n_1}), Y_{n_1}) + s_{n_1}] \\
&\quad + \sum_{Y_{n_1} \in \text{ch}'(Y_0)} \left\{ \sum_{f_p \in \text{ch}(Y_{n_1})} d(Y_{n_1}, f_p) + \sum_{Y_{n_2} \in \text{ch}'(Y_{n_1})} \left[d(Y_{n_1}, Y_{n_2}) + s_{n_2} + \sum_{f_p \in \text{de}(Y_{n_2})} d(Y_{n_2}, f_p) \right] \right\} \leq \dots \\
&\leq \sum_{p=1}^{d+1} d(p(f_p), f_p) + \sum_{n=1}^{l(Y)} [d(p(Y_n), Y_n) + s_n]. \tag{S195}
\end{aligned}$$

Here in the first line, we have used (S8) together with the fact that any f_p belongs to either $\text{ch}(Y_0)$, or one of $\text{de}(Y_{n_1})$ with Y_{n_1} being one child of Y_0 . I.e., $\{Y_{n_1}\}$ are the bifurcation factors at the second layer of the tree Γ (the first layer being the root Y_0). In the second line we invoked (S192). In the third line we identified Y_0 with nothing but the parent of its children. In the fourth line, we repeated the treatment in the first line for each Y_{n_1} , and used (S192) for each of the third-layer factors Y_{n_2} . Traversing down the tree Γ recursively, we cover the distances corresponding to all edges in the tree exactly once, together with all the s_n s. As a result, we arrived at the final line and established (S189), which further yields (S187).

Plugging (S187) into (S186),

$$\begin{aligned}
\mathcal{G}(Y) &\leq |S|^{-l(Y)} |S|^{l(Y)} (|\partial S| c'_\lambda t \|H\|_\kappa)^{q(Y)} \exp \left\{ -\kappa \left[(s - s_0)^\alpha - \sum_{n=1}^{l(Y)} s_n^\alpha \right] \right\} \\
&\leq |S|^{-l(Y)} c_{l(Y), q(Y)} (c'_\lambda)^{q(Y)} (\delta\kappa)^{-\frac{dl(Y) + (d-1)q(Y)}{\alpha}} t \|H\|_\kappa (\delta\kappa)^{[q(Y)-1] \frac{3d-1}{\alpha}} e^{\frac{\delta\kappa}{2} s_0^\alpha} e^{-\tilde{\kappa}(s-s_0)^\alpha} e^{\kappa \sum_{n=1}^{l(Y)} s_n^\alpha} \\
&\leq c_{max} |S|^{-l(Y)} (\delta\kappa)^{-\frac{d-1+dl(Y)}{\alpha}} t \|H\|_\kappa e^{\frac{\delta\kappa}{2} s_0^\alpha} e^{-\tilde{\kappa}(s-s_0)^\alpha} e^{\kappa \sum_{n=1}^{l(Y)} s_n^\alpha}, \tag{S196}
\end{aligned}$$

In the second line we have used a generalization of (S183):

$$|S|^l |\partial S_0|^q \leq c_{l,q} (\delta\kappa)^{-\frac{ld+q(d-1)}{\alpha}} e^{\frac{\delta\kappa}{4} (s^\alpha + s_0^\alpha)} \leq c_{l,q} (\delta\kappa)^{-\frac{ld+q(d-1)}{\alpha}} e^{\frac{\delta\kappa}{4} [(s-s_0)^\alpha + 2s_0^\alpha]}.$$
 \tag{S197}

and (S114). In the third line we have maximized the prefactor over $1 \leq q(Y) \leq 2d + 1$, using $\delta\kappa < \kappa \leq \kappa_1$.

For a fixed vertex number $l + 1$, there are at most $l!$ different trees. Thus for a given $l(Y) = l \leq d$, plugging (S196) into (S174) yields

$$\begin{aligned}
J(S) &\leq (d+1)! l! \sum_{Y_1, \dots, Y_l \subseteq S} (2t)^l \prod_{n=1}^l \|H_{Y_n}\| \mathcal{G}(Y) \\
&\leq (d+1)! l! c_{max} (\delta\kappa)^{-\frac{d-1+dl}{\alpha}} (2t)^l t \|H\|_\kappa e^{\frac{\delta\kappa}{2} s_0^\alpha} e^{-\tilde{\kappa}(s-s_0)^\alpha} \sum_{i_1, \dots, i_l \in S} |S|^{-l} \prod_{n=1}^l \left(\sum_{Y_n \ni i_n} \|H_{Y_n}\| e^{\kappa s_n^\alpha} \right) \\
&\leq (d+1)! l! c_{max} (\delta\kappa)^{-\frac{2d-1+d(l-1)}{\alpha}} (2t)^l t \|H\|_\kappa e^{\frac{\delta\kappa}{2} s_0^\alpha} e^{-\tilde{\kappa}(s-s_0)^\alpha} \|H\|_\kappa^l \\
&\leq c'_{max} (\delta\kappa)^{-\frac{2d-1}{\alpha}} t^2 \|H\|_\kappa^2 e^{\frac{\delta\kappa}{2} s_0^\alpha} e^{-\tilde{\kappa}(s-s_0)^\alpha}. \tag{S198}
\end{aligned}$$

Here in the first line, we sum over $Y \in \mathbf{Y}(S)$ by first summing over the topological tree structure that gives the prefactor $l!$, and then summing over each $Y_n \in Y$. In the second line, we overcount by replacing each $\sum_{Y_n \subseteq S}$ by $\sum_{i_n \in S} \sum_{Y_n \ni i_n}$. While this counts the same Y_n , $|Y_n|$ times, we can use the κ -norm of H to sum over Y_n efficiently. Furthermore, the sum over sites is canceled by $|S|^{-l}$ from $\mathcal{G}(Y)$. In the last line, we have used $(2t \|H\|_\kappa)^{l-1}$ to kill $(\delta\kappa)^{-\frac{d(l-1)}{\alpha}}$ according to (S114), and finally arrived at the form of (S174). \square

-
- [1] P. W. Anderson, “Infrared catastrophe in fermi gases with local scattering potentials,” *Phys. Rev. Lett.* **18**, 1049–1051 (1967).
- [2] D Poilblanc, T Ziman, J Bellissard, F Mila, and G Montambaux, “Poisson vs. GOE statistics in integrable and non-integrable quantum hamiltonians,” *Europhysics Letters (EPL)* **22**, 537–542 (1993).
- [3] D. A. Rabson, B. N. Narozhny, and A. J. Millis, “Crossover from poisson to wigner-dyson level statistics in spin chains with integrability breaking,” *Phys. Rev. B* **69**, 054403 (2004).
- [4] D.M. Basko, I.L. Aleiner, and B.L. Altshuler, “Metal–insulator transition in a weakly interacting many-electron system with localized single-particle states,” *Annals of Physics* **321**, 1126–1205 (2006).
- [5] Vadim Oganesyan and David A. Huse, “Localization of interacting fermions at high temperature,” *Phys. Rev. B* **75**, 155111 (2007).
- [6] David A. Huse, Rahul Nandkishore, and Vadim Oganesyan, “Phenomenology of fully many-body-localized systems,” *Phys. Rev. B* **90**, 174202 (2014).
- [7] Rahul Nandkishore and David A. Huse, “Many-body localization and thermalization in quantum statistical mechanics,” *Annual Review of Condensed Matter Physics* **6**, 15–38 (2015).
- [8] John Z Imbrie, “On many-body localization for quantum spin chains,” *Journal of Statistical Physics* **163**, 998–1048 (2016).
- [9] Dmitry A. Abanin, Ehud Altman, Immanuel Bloch, and Maksym Serbyn, “Colloquium: Many-body localization, thermalization, and entanglement,” *Rev. Mod. Phys.* **91**, 021001 (2019).
- [10] Rajdeep Sensarma, David Pekker, Ehud Altman, Eugene Demler, Niels Strohmaier, Daniel Greif, Robert Jördens, Leticia Tarruell, Henning Moritz, and Tilman Esslinger, “Lifetime of double occupancies in the fermi-hubbard model,” *Phys. Rev. B* **82**, 224302 (2010).
- [11] A. L. Chudnovskiy, D. M. Gangardt, and A. Kamenev, “Doublon relaxation in the bose-hubbard model,” *Phys. Rev. Lett.* **108**, 085302 (2012).
- [12] Dmitry A. Abanin, Wojciech De Roeck, and Fran çois Huveneers, “Exponentially slow heating in periodically driven many-body systems,” *Phys. Rev. Lett.* **115**, 256803 (2015).
- [13] Dmitry Abanin, Wojciech De Roeck, Wen Wei Ho, and Fran çois Huveneers, “A rigorous theory of many-body prethermalization for periodically driven and closed quantum systems,” *Communications in Mathematical Physics* **354**, 809–827 (2017).
- [14] Dmitry A. Abanin, Wojciech De Roeck, Wen Wei Ho, and Fran çois Huveneers, “Effective hamiltonians, prethermalization, and slow energy absorption in periodically driven many-body systems,” *Phys. Rev. B* **95**, 014112 (2017).
- [15] Tomotaka Kuwahara, Takashi Mori, and Keiji Saito, “Floquet–magnus theory and generic transient dynamics in periodically driven many-body quantum systems,” *Annals of Physics* **367**, 96–124 (2016).
- [16] Takashi Mori, Tomotaka Kuwahara, and Keiji Saito, “Rigorous bound on energy absorption and generic relaxation in periodically driven quantum systems,” *Phys. Rev. Lett.* **116**, 120401 (2016).
- [17] Dominic V. Else, Bela Bauer, and Chetan Nayak, “Prethermal phases of matter protected by time-translation symmetry,” *Phys. Rev. X* **7**, 011026 (2017).
- [18] Francisco Machado, Dominic V. Else, Gregory D. Kahanamoku-Meyer, Chetan Nayak, and Norman Y. Yao, “Long-range prethermal phases of nonequilibrium matter,” *Phys. Rev. X* **10**, 011043 (2020).
- [19] Dominic V. Else, Wen Wei Ho, and Philipp T. Dumitrescu, “Long-lived interacting phases of matter protected by multiple time-translation symmetries in quasiperiodically driven systems,” *Phys. Rev. X* **10**, 021032 (2020).
- [20] Ken Xuan Wei, Pai Peng, Oles Shtanko, Iman Marvian, Seth Lloyd, Chandrasekhar Ramanathan, and Paola Cappellaro, “Emergent prethermalization signatures in out-of-time ordered correlations,” *Phys. Rev. Lett.* **123**, 090605 (2019).
- [21] Pai Peng, Chao Yin, Xiaoyang Huang, Chandrasekhar Ramanathan, and Paola Cappellaro, “Floquet prethermalization in dipolar spin chains,” *Nature Physics* **17**, 444–447 (2021).
- [22] Antonio Rubio-Abadal, Matteo Ippoliti, Simon Hollerith, David Wei, Jun Rui, S. L. Sondhi, Vedika Khemani, Christian Gross, and Immanuel Bloch, “Floquet prethermalization in a bose-hubbard system,” *Phys. Rev. X* **10**, 021044 (2020).
- [23] William Beatrix, Otto Janes, Amala Akkijaru, Arjun Pillai, Alexander Oddo, Paul Reshetikhin, Emanuel Druga, Maxwell McAllister, Mark Elo, Benjamin Gilbert, Dieter Suter, and Ashok Ajoy, “Floquet prethermalization with lifetime exceeding 90 s in a bulk hyperpolarized solid,” *Phys. Rev. Lett.* **127**, 170603 (2021).
- [24] A. Kyprianidis, F. Machado, W. Morong, P. Becker, K. S. Collins, D. V. Else, L. Feng, P. W. Hess, C. Nayak, G. Pagano, N. Y. Yao, and C. Monroe, “Observation of a prethermal discrete time crystal,” *Science* **372**, 1192–1196 (2021).
- [25] Constantine Shkredov, Meny Menashes, Gal Ness, Anastasiya Vainbaum, Ehud Altman, and Yoav Sagi, “Absence of heating in a uniform fermi gas created by periodic driving,” *Phys. Rev. X* **12**, 011041 (2022).

- [26] Spyridon Michalakis and Justyna P Zwolak, “Stability of frustration-free hamiltonians,” *Communications in Mathematical Physics* **322**, 277–302 (2013).
- [27] Bruno Nachtergaele, Robert Sims, and Amanda Young, “Quasi-Locality Bounds for Quantum Lattice Systems. Part II. Perturbations of Frustration-Free Spin Models with Gapped Ground States,” *Annales Henri Poincare* **23**, 393–511 (2022), [arXiv:2010.15337 \[math-ph\]](https://arxiv.org/abs/2010.15337).
- [28] A.Yu. Kitaev, “Fault-tolerant quantum computation by anyons,” *Annals of Physics* **303**, 2–30 (2003).
- [29] Chetan Nayak, Steven H. Simon, Ady Stern, Michael Freedman, and Sankar Das Sarma, “Non-abelian anyons and topological quantum computation,” *Rev. Mod. Phys.* **80**, 1083–1159 (2008).
- [30] Sergey Bravyi, Matthew B. Hastings, and Spyridon Michalakis, “Topological quantum order: Stability under local perturbations,” *Journal of Mathematical Physics* **51**, 093512 (2010), <https://doi.org/10.1063/1.3490195>.
- [31] Benjamin J. Brown, Daniel Loss, Jiannis K. Pachos, Chris N. Self, and James R. Wootton, “Quantum memories at finite temperature,” *Rev. Mod. Phys.* **88**, 045005 (2016).
- [32] Subir Sachdev, *Quantum Phase Transitions*, 2nd ed. (Cambridge University Press, 2011).
- [33] Naoto Shiraishi and Takashi Mori, “Systematic construction of counterexamples to the eigenstate thermalization hypothesis,” *Phys. Rev. Lett.* **119**, 030601 (2017).
- [34] Hannes Bernien, Sylvain Schwartz, Alexander Keesling, Harry Levine, Ahmed Omran, Hannes Pichler, Soonwon Choi, Alexander S Zibrov, Manuel Endres, Markus Greiner, *et al.*, “Probing many-body dynamics on a 51-atom quantum simulator,” *Nature* **551**, 579–584 (2017).
- [35] Christopher J Turner, Alexios A Michailidis, Dmitry A Abanin, Maksym Serbyn, and Zlatko Papić, “Weak ergodicity breaking from quantum many-body scars,” *Nature Physics* **14**, 745–749 (2018).
- [36] Pablo Sala, Tibor Rakovszky, Ruben Verresen, Michael Knap, and Frank Pollmann, “Ergodicity breaking arising from hilbert space fragmentation in dipole-conserving hamiltonians,” *Phys. Rev. X* **10**, 011047 (2020).
- [37] Vedika Khemani, Michael Hermele, and Rahul Nandkishore, “Localization from hilbert space shattering: From theory to physical realizations,” *Phys. Rev. B* **101**, 174204 (2020).
- [38] Maksym Serbyn, Dmitry A. Abanin, and Zlatko Papić, “Quantum many-body scars and weak breaking of ergodicity,” *Nature Phys.* **17**, 675–685 (2021), [arXiv:2011.09486 \[quant-ph\]](https://arxiv.org/abs/2011.09486).
- [39] Zhi-Cheng Yang, Fangli Liu, Alexey V. Gorshkov, and Thomas Iadecola, “Hilbert-space fragmentation from strict confinement,” *Phys. Rev. Lett.* **124**, 207602 (2020).
- [40] Atsuki Yoshinaga, Hideaki Hakoshima, Takashi Imoto, Yuichiro Matsuzaki, and Ryusuke Hamazaki, “Emergence of hilbert space fragmentation in ising models with a weak transverse field,” *Phys. Rev. Lett.* **129**, 090602 (2022).
- [41] Sanjay Moudgalya and Olexei I. Motrunich, “Hilbert space fragmentation and commutant algebras,” *Phys. Rev. X* **12**, 011050 (2022).
- [42] Julia Wildeboer, Thomas Iadecola, and Dominic J. Williamson, “Symmetry-protected infinite-temperature quantum memory from subsystem codes,” *PRX Quantum* **3**, 020330 (2022).
- [43] David T. Stephen, Oliver Hart, and Rahul M. Nandkishore, “Ergodicity breaking provably robust to arbitrary perturbations,” (2022), [arXiv:2209.03966 \[cond-mat.stat-mech\]](https://arxiv.org/abs/2209.03966).
- [44] Anushya Chandran, Thomas Iadecola, Vedika Khemani, and Roderich Moessner, “Quantum Many-Body Scars: A Quasi-particle Perspective,” (2022), [arXiv:2206.11528 \[cond-mat.str-el\]](https://arxiv.org/abs/2206.11528).
- [45] J. M. Deutsch, “Quantum statistical mechanics in a closed system,” *Phys. Rev. A* **43**, 2046–2049 (1991).
- [46] Mark Srednicki, “Chaos and quantum thermalization,” *Phys. Rev. E* **50**, 888–901 (1994).
- [47] Luca D’Alessio, Yariv Kafri, Anatoli Polkovnikov, and Marcos Rigol, “From quantum chaos and eigenstate thermalization to statistical mechanics and thermodynamics,” *Advances in Physics* **65**, 239–362 (2016), <https://doi.org/10.1080/00018732.2016.1198134>.
- [48] Takashi Mori, Tatsuhiko N Ikeda, Eriko Kaminishi, and Masahito Ueda, “Thermalization and prethermalization in isolated quantum systems: a theoretical overview,” *Journal of Physics B: Atomic, Molecular and Optical Physics* **51**, 112001 (2018).
- [49] Bei Zeng, Xie Chen, Duan-Lu Zhou, Xiao-Gang Wen, *et al.*, *Quantum information meets quantum matter* (Springer, 2019).
- [50] Sergey Bravyi and Matthew B Hastings, “A short proof of stability of topological order under local perturbations,” *Communications in mathematical physics* **307**, 609–627 (2011).
- [51] Jian Ma, Xiaoguang Wang, C.P. Sun, and Franco Nori, “Quantum spin squeezing,” *Physics Reports* **509**, 89–165 (2011).
- [52] Vittorio Giovannetti, Seth Lloyd, and Lorenzo Maccone, “Quantum metrology,” *Phys. Rev. Lett.* **96**, 010401 (2006).
- [53] Michael A. Perlin, Chunlei Qu, and Ana Maria Rey, “Spin squeezing with short-range spin-exchange interactions,” *Phys. Rev. Lett.* **125**, 223401 (2020).
- [54] J. R. Schrieffer and P. A. Wolff, “Relation between the anderson and kondo hamiltonians,” *Phys. Rev.* **149**, 491–492 (1966).
- [55] Sergey Bravyi, David P. DiVincenzo, and Daniel Loss, “Schrieffer–wolff transformation for quantum many-body systems,” *Annals of Physics* **326**, 2793–2826 (2011).
- [56] Elliott H. Lieb and Derek W. Robinson, “The finite group velocity of quantum spin systems,” *Commun. Math. Phys.* **28**, 251–257 (1972).
- [57] MB Hastings, “Quasi-adiabatic continuation for disordered systems: Applications to correlations, lieb-schultz-mattis, and hall conductance,” [arXiv:1001.5280 \[math-ph\]](https://arxiv.org/abs/1001.5280).
- [58] Sven Bachmann, Spyridon Michalakis, Bruno Nachtergaele, and Robert Sims, “Automorphic equivalence within gapped phases of quantum lattice systems,” *Communications in Mathematical Physics* **309**, 835–871 (2012).
- [59] Chi-Fang Chen and Andrew Lucas, “Operator Growth Bounds from Graph Theory,” *Commun. Math. Phys.* **385**, 1273–1323 (2021), [arXiv:1905.03682 \[math-ph\]](https://arxiv.org/abs/1905.03682).
- [60] J.M. Drouffe and K.J.M. Moriarty, “Gauge theories on a simplicial lattice,” *Nuclear Physics B* **220**, 253–268 (1983).

- [61] M. B. Hastings, “Locality in quantum systems,” (2010), [arXiv:1008.5137 \[math-ph\]](#).
- [62] Tomotaka Kuwahara and Keiji Saito, “Strictly linear light cones in long-range interacting systems of arbitrary dimensions,” *Phys. Rev. X* **10**, 031010 (2020), [arXiv:1910.14477 \[quant-ph\]](#).
- [63] Michael Foss-Feig, Zhe-Xuan Gong, Charles W. Clark, and Alexey V. Gorshkov, “Nearly linear light cones in long-range interacting quantum systems,” *Phys. Rev. Lett.* **114**, 157201 (2015).
- [64] Chi-Fang Chen and Andrew Lucas, “Finite speed of quantum scrambling with long range interactions,” *Phys. Rev. Lett.* **123**, 250605 (2019), [arXiv:1907.07637 \[quant-ph\]](#).
- [65] Minh C. Tran, Chi-Fang Chen, Adam Ehrenberg, Andrew Y. Guo, Abhinav Deshpande, Yifan Hong, Zhe-Xuan Gong, Alexey V. Gorshkov, and Andrew Lucas, “Hierarchy of Linear Light Cones with Long-Range Interactions,” *Phys. Rev. X* **10**, 031009 (2020), [arXiv:2001.11509 \[quant-ph\]](#).
- [66] Minh C. Tran, Andrew Y. Guo, Christopher L. Baldwin, Adam Ehrenberg, Alexey V. Gorshkov, and Andrew Lucas, “Lieb-Robinson Light Cone for Power-Law Interactions,” *Phys. Rev. Lett.* **127**, 160401 (2021), [arXiv:2103.15828 \[quant-ph\]](#).