

On the W mass anomaly in models with right-handed currents

Henry Diaz,^{1,*} V. Pleitez,^{2,†} and O. P. Ravinez^{1,‡}

¹*Facultad de Ciencias, Universidad Nacional de Ingeniería (UNI)*

av. Tupac Amaru s/n, Lima - Rímac 15333, Perú

²*Instituto de Física Teórica–Universidade Estadual Paulista (UNESP)*

R. Dr. Bento Teobaldo Ferraz 271, Barra Funda

São Paulo - SP, 01140-070, Brazil

(Dated: 11/22/2022)

Abstract

We show that, in electroweak models with right-handed currents and gauge symmetry $SU(2)_L \otimes SU(2)_R \otimes U(1)_{B-L}$, it is possible to accommodate, at the tree level, the anomaly of the W mass obtained by the CDF-II. The solution to that anomaly is possible independently if a generalized parity is broken explicitly or spontaneously. A change in the mass of Z is also predicted. This seems to be a characteristic of any of this type of models.

PACS numbers: 12.60.Fr; 12.15.-y

* hdiaz@uni.edu.pe

† v.pleitez@unesp.br

‡ opereyra@uni.edu.pe

I. INTRODUCTION

Recently the CDF collaboration has obtained a W mass which does not agree with the value predicted by the standard model (SM) [1] at the 7σ level:

$$M_W|_{CDF} = 80,433.5 \pm 9.4 \text{ MeV}, \quad (1)$$

that if compared with the value in the context of the SM (with the Higgs boson mass measured at the LHC used as input and using radiative corrections) $M_W|_{SM} = 80,357 \pm 0.0056 \text{ MeV}$ [2], it means a disagreement at 7σ level with the theoretically prediction. The CDF value is also in disagreement with other experimental data [3]. Then, if confirmed, the CDF's result implies that new physics has been revealed.

In the literature some explanations for this fact have been proposed. For instance, it may be new physics whose effects appear through radiative corrections [4–6], or also at tree level [7–9]. Any solution to this problem requires that new physics would manifest itself in other observables since it is difficult for its effects to occur only in the mass of the W .

A characteristic that we will explore here is the fact that in models with many scalar fields some of them do not have couplings with the known fermions. For instance, if a symmetry, or the representation content, of the model prohibits a scalar to interact with the fermions but it contributes to the mass of W , the respective VEV would be, in principle, a free parameter and it would be in charge of the extra contribution to the mass from W . An example of this mechanism is possible in the supersymmetric models in which the supersymmetry forbids the interaction of a scalar with fermions but its VEV contributes to the vector boson masses [10, 11].

Another example that we will consider here, are the well motivated models with electroweak gauge symmetry [12–15]

$$SU(2)_L \otimes SU(2)_R \otimes U(1)_{B-L} \quad (2)$$

with, or without, a conserved (at a given energy scale) generalized parity \mathcal{P} .

In this sort of models left- and right-handed fermions transform as doublets of $SU(2)_L$ and $SU(2)_R$, respectively. The minimal scalar sector consist of one bidoublet $\Phi \sim (\mathbf{2}_L, \mathbf{2}_R, 0)$, two doublets $\chi_L \sim (\mathbf{2}_L, \mathbf{1}_R, +1)$ and $\chi_R \sim (\mathbf{1}_L, \mathbf{2}_R, +1)$. Instead of doublets these scalars may be complex triplets, say $\chi_L \sim (\mathbf{3}_L, \mathbf{1}_R, +2)$ and $\chi_R \sim (\mathbf{1}_L, \mathbf{3}_R, +2)$, if we want to have Majorana neutrino. However, we will not consider the latter possibility here because our idea does not depend on this. Only the bi-doublets, and the complex triplets, if they are added, can generate all fermions masses.

In fact, in order to allow the doublet χ_L to interact with quarks, new quarks singlets must be introduced as in Ref. [16]. If $\chi_{L,R}$ are triplets, they may couple with left- and right-handed leptons, unless a discrete symmetry, say \mathbb{Z}_2 , forbids the coupling with the triplet χ_L . Below we consider $\chi_{L,R}$ being doublets of the respective $SU(2)_{L,R}$ factor.

Since the vacuum expectation value of $\langle \chi_L \rangle = v_L/\sqrt{2}$ contributes only to the mass of the $W_L \approx W$ (where W is the SM charged vector boson) and for this reason it has an arbitrary value, it may be even zero [14]. Here we explore this possibility in order to explain the W mass anomaly in the context of these models. Of course, $\langle \chi_R \rangle = v_R/\sqrt{2} \neq 0$ because it is in charge of the mass of the $W_R \approx W'$ vector boson. We denote k and k' the VEVs of the bi-doublets.

In these models the mass square matrix for the charged vector bosons has the following structure

$$M_{CB}^2 = \frac{g^2 v_R^2}{4} \begin{pmatrix} x+y & -2z\varepsilon \\ & 1+x \end{pmatrix}, \quad (3)$$

where we have defined $x = K^2/v_R^2$, with $K^2 = k^2 + k'^2$, $y = v_L^2/v_R^2$, and $z = kk'/v_R^2$, $\varepsilon = g_R^2/g_L^2$. Below we call the mass eigenstates of the matrix in (3) just W and W' , similarly with Z and Z' , but we omit the mass matrix of the neutral vector bosons.

The outline of this paper is as follows: In Sec. II we consider the model in which there is a left-right symmetry which is spontaneously broken; in Sec. III we consider a model with the same representation content of the first one but now with explicit parity breaking. Our conclusion appears in Sec. IV.

II. MODEL WITH LEFT-RIGHT SYMMETRY

In the first model there is also the symmetry under a generalized parity, \mathcal{P} ,

$$\begin{aligned} g_L &\leftrightarrow g_R, \quad W_{L\mu} \leftrightarrow W_R^\mu, \quad f_L \leftrightarrow f_R, \\ \chi_L &\leftrightarrow \chi_R, \quad \Phi_i \leftrightarrow \Phi_i^\dagger, \quad \tilde{\Phi}_i \leftrightarrow \tilde{\Phi}_i^\dagger, \end{aligned} \quad (4)$$

$\tilde{\Phi}_i = \tau_2 \Phi_i^* \tau_2$, which is broken spontaneously [12–14].

As we said above, the scalar doublet, χ_R , under $SU(2)_R$ has a VEV whose value is determined by the LR symmetry breaking scale, but the χ_L which is introduced just to maintain the generalized parity symmetry has a free VEV, that would be zero, or having an appropriate value to solve the case of W mass as we will do here. In this case, we have the parameters x, y and z defined as above

but with $\varepsilon = 1$, and the eigenvalues of the mass matrix in Eq. (3) are

$$\begin{aligned} M_W^2 &= \frac{g^2 v_R^2}{4} \left(x + \frac{1+y}{2} - \sqrt{\Delta_1} \right), \\ M_{W'}^2 &= \frac{g^2 v_R^2}{4} \left(x + \frac{1+y}{2} + \sqrt{\Delta_1} \right), \\ \Delta_1 &= 4z^2 + \frac{1}{4}(1-y)^2. \end{aligned} \quad (5)$$

Expanding in x we have

$$\begin{aligned} M_W^2 &\approx \frac{g^2 K^2}{4} [1 + \xi - x(1+y+y^2) + x^3], \\ M_{W'}^2 &\approx \frac{g^2}{4} v_R^2, \quad \xi = v_L^2 / K^2 \end{aligned} \quad (6)$$

Notice that $M_{W'}^2 \gg M_W^2$, and we identify the W^\pm of the standard model with W_1^\pm . We can see that the extra mass contributions comes from the ν_L , and (6) for $x \ll 1$ we have

$$\delta M_W^2 = \frac{g^2}{4} v_L^2, \quad (7)$$

and we see that an additional contribution to the mass of the W boson is possible since ν_L is a free parameter.

However, we use the exact expression in Eq. (5) for calculating M_W as a function of ν_L , using several values for ν_R , say from 1 TeV to ∞ (in fact $\nu_R \gg k, k'$), using $K^2 = (246 \text{ GeV})^2$, and for the sake of simplicity we assume $k = k'$. Moreover, we use $g = 0.6539$ and $s_\theta = 0.2312$ [3]. In fact, although the model has two mixing angles in the neutral vector bosons [15, 17], in the limit of the large mass of W' ($\nu_R \rightarrow \infty$) we can identify $\sin\theta = \sin\theta_W$. The result are shown in Fig. 1, the colored strip indicates the value of the CDF at the level of 1σ and we show the result for several values of ν_R . We have verified that with $2.4 \leq \nu_L \leq 60.6 \text{ GeV}$, the lower bound refers to $\nu_R \rightarrow \infty$ and the larger one to $\nu_R = 1 \text{ TeV}$, we get the mass of the W obtained by the CDF [1]. Below we will show that the solution with $\nu_R = 24 \text{ TeV}$, and $\nu_L = 3.47 \text{ GeV}$ is an interesting possibility.

However, there are also contributions to the Z and Z' masses (we omit the mass square matrix in the neutral vector bosons) given by

$$M_Z^2 = \frac{g^2 v_R^2}{4} \left(x + \frac{1}{2}(1+y)(1+\delta') - \frac{1}{2}\sqrt{\Xi_1} \right) \quad (8)$$

$$M_{Z'}^2 = \frac{g^2 v_R^2}{4} \left(x + \frac{1}{2}(1+y)(1+\delta') + \frac{1}{2}\sqrt{\Xi_1} \right) \quad (9)$$




FIG. 1. W mass as a function of v_L in a model with left-right symmetry.

where:

$$\Xi_1 = (1+y)^2(1+\delta')^2 - 4\delta'x(1+y) - 4(1+2\delta')y + 4x^2$$

$\delta' = g'^2/g^2 = \tan^2\theta = 03987$, expanding in x

$$M_Z^2 \approx \frac{g^2 K^2}{4\cos^2\theta}(1+\xi), \quad \xi = v_L^2/K^2 \quad (10)$$

however the usual relation

$$\delta M_Z^2 \approx \frac{\delta M_W^2}{\cos^2\theta}, \quad (11)$$

is still valid, as expected, since $\chi_{L,R}$ are a doublet of $SU(2)_L$, then $\rho = M_W^2/M_Z^2 \cos^2\theta = 1$ at tree level. If χ_L is a complex triplet, i.e., carrying hypercharge, the relation in Eq. (11) implies a upper bound in its VEV, denoted by v_T , in order to keep $\rho = 1$, within the experimental error, then $v_T \lesssim 4$ GeV which is also consistent with the CDF II result of the W mass. This happens in any model with this sort of scalar [18].

Since $\delta M_W|_{CDF} = 0.0094$ GeV then $\delta M_Z = 0.0107$ GeV. Notice that it is a prediction of the model that the mass of the Z is also different than its value in the context of the SM. This make the model falsifiable.

III. MODEL WITH EXPLICIT PARITY BREAKING

Next, let us now consider a model with the gauge symmetry as in Eq. (2) but with \mathcal{P} being explicitly broken [19], i.e., in which $g_L \neq g_R$ and the transformations in Eq. (4) are not valid.

Hence, the bosons and all parameters in the $SU(2)_L$ are not related with those of the $SU(2)_R$ factor, including the mixing matrices in each quiral fermion sector. In this case the mass matrix of the charged vector bosons is as in Eq. (3) with x, y and z are defined as in the previous model but now $\varepsilon = g_R^2/g_L^2 \neq 1$. Then we obtain for the charged vector bosons the masses:

$$\begin{aligned} M_W^2 &= \frac{g_L^2 v_R^2}{8} \left[(1 + \varepsilon)x + y + \varepsilon - \sqrt{\Delta_2} \right] \\ M_{W'}^2 &= \frac{g_L^2 v_R^2}{8} \left[(1 + \varepsilon)x + y + \varepsilon + \sqrt{\Delta_2} \right], \\ \Delta_2 &= 4\varepsilon x^2 + [\varepsilon - y - (1 - \varepsilon)x]^2, \end{aligned} \quad (12)$$

from these equations we see that

$$\begin{aligned} \delta M_W^2 &\approx \frac{g_L^2}{4} K^2 \left[\xi - \left(1 + \frac{y}{\varepsilon}\right)x + \left(1 - \frac{y}{\varepsilon}\right)x^2 \right] \\ &= \delta M_W^2|_{\varepsilon=1} + \delta M_W^2|_{\varepsilon \neq 1} \\ M_{W'}^2 &\approx \frac{g_L^2}{4} \varepsilon v_R^2 (1+x) = \frac{g_R^2}{4} v_R^2 (1+x), \end{aligned} \quad (13)$$

where $\delta M_W^2|_{\varepsilon=1}$ is as in Eq. (7) but now with $g \rightarrow g_L$, and

$$\delta M_W^2|_{\varepsilon \neq 1} = -\frac{g_L^2}{4} K^2 \left[\left(1 + \frac{y}{\varepsilon}\right)x - \left(1 - \frac{y}{\varepsilon}\right)x^2 \right], \quad (14)$$

and $\delta M_W^2|_{\varepsilon \neq 1} \rightarrow 0$ when $x \rightarrow 0$. Hence the δM_W is again as given in Eq. (7).

Notice that, in order to maintain the δM_W mass at a given value, the greater the v_R ($x \ll 1$) smaller the $\varepsilon \ll 1$ ($g_R \ll g_L$) for any energy. This also maintain the mass of the W' at a given value. In Fig. 2(a) we show the W mass in terms of ε for several values v_L and $v_R = 1$ TeV. We see that even with $v_L = 0$ it is possible to have solutions for the W mass anomaly. In Fig. 2(b) we show the mass of W in terms of v_L for several values of ε and also $v_R = 1$ TeV. For the neutral vector bosons we have in this case

$$\begin{aligned} M_Z^2 &= \frac{1}{8} g_L^2 v_R^2 \left(x(1 + \varepsilon) + y(1 + \delta_L) + \varepsilon + \delta_L - \sqrt{\Xi_2} \right) \\ M_{Z'}^2 &= \frac{1}{8} g_L^2 v_R^2 \left(x(1 + \varepsilon) + y(1 + \delta_L) + \varepsilon + \delta_L + \sqrt{\Xi_2} \right), \\ \Xi_2 &= y^2(1 + \delta_L)^2 + x^2(\varepsilon + 1)^2 + (\delta_L + \varepsilon)^2 + 4xy(1 - \varepsilon(1 + \delta_L) - \delta_L), \end{aligned} \quad (15)$$

where $\delta_L = g'^2/g_L^2 = 0.3987$. Again when $x, y \rightarrow 0$ ($v_R \rightarrow \infty$) the relation in Eq. (11) is obtained, independently of the value of ε .

From the exact expression in Eq. (12) we can calculate the mass of the W' gauge boson. We recall that most searches for W' , and also for Z' , assume that the couplings of these bosons to

quarks and leptons are taken to be identical with those of SM's W and Z and also that $g_L = g_R$ [3]. This is not the case in the present model which has explicit parity breaking and in this case the limits are considerably weaker than for the special case of manifest left-right symmetry [20]. Moreover, since we have considered only Dirac neutrinos and no scalar triplets (no doubly charged scalars), the results obtained in Ref. [21] do not apply straitforward, moreover in the model with explicit parity breaking does not exist any relation between the left- and right- parameters, g_L , g_R and the respective mixing matrices at any energy scale.

Notice, in particular, that for $\nu_R = 1$ TeV and $\nu_L = 60$ GeV, we have $0.8 \lesssim \epsilon \lesssim 1.2$. We have found but do not show in Fig. 2, that when $\nu_L = 0$ the solution still exist but only at the 5σ level with ϵ rather small, $g_R \sim 10^{-6} g_L$.

IV. CONCLUSIONS

Here we have shown that the well motivated minimal version of electroweak models with right-handed currents and $SU(2)_L \otimes SU(2) \otimes U(1)_{B-l}$ gauge symemtry, with or whitout left-right symmetry, are able to explain the W mass anomaly while keeping the usual phenomeology of the models. It is interesting that if these models are the solution to the W mass anomaly, there must also be an anomaly in the mass of Z .

Another possibility, which we have not considered here, is the case in which the mass of a new charged vector, say V^- , does exist and has a mass near the mass of W^- , so that the measures before the CDF [3] have obtained an average value of the mass of both charged vectors, but CDF II with more precise data has measured the mass of one of them (may be the mass of the lightest one). Finally, it is possible that loop effects are important also and even if $\nu_L = 0$; in this case the doublet χ_L would be an example of an inert doublet [22].




FIG. 2. In the upper picture we show the W mass as a function of ϵ and two values of $\nu_{L,R}$. In the lower picture it appears the W mass as a function of ϵ for several values of ν_L and $\nu_R = 1$ TeV in a model with explicit parity breakdown.

Acknowledgments

V. P. is thankful for the support of FAPESP funding Grant 2014/19164-6.

-
- [1] T. Aaltonen *et al.* [CDF], High-precision measurement of the W boson mass with the CDF II detector, *Science* **376**, no.6589, 170-176 (2022).
 - [2] J. de Blas, M. Pierini, L. Reina and L. Silvestrini, Impact of the recent measurements of the top-quark and W -boson masses on electroweak precision fits, [arXiv:2204.04204 [hep-ph]].
 - [3] R. L. Workman *et al.* [Particle Data Group], Review of Particle Physics, *PTEP* **2022**, 083C01 (2022)

doi:10.1093/ptep/ptac097

- [4] A. Strumia, Interpreting electroweak precision data including the W-mass CDF anomaly, JHEP **08**, 248 (2022); [arXiv:2204.04191 [hep-ph]].
- [5] K. Sakurai, F. Takahashi and W. Yin, *Singlet extensions and W boson mass in light of the CDF II result*, Phys. Lett. B **833**, 137324 (2022); [arXiv:2204.04770 [hep-ph]].
- [6] K. S. Babu, S. Jana and V. P. K., Correlating W-Boson Mass Shift with Muon g-2 in the Two Higgs Doublet Model, Phys. Rev. Lett. **129**, no.12, 12 (2022); [arXiv:2204.05303 [hep-ph]].
- [7] G. Lazarides, R. Maji, R. Roshan and Q. Shafi, Heavier W boson, dark matter, and gravitational waves from strings in an SO(10) axion model, Phys. Rev. D **106**, no.5, 5 (2022); [arXiv:2205.04824 [hep-ph]].
- [8] J. L. Evans, T. T. Yanagida and N. Yokozaki, W boson mass anomaly and grand unification, Phys. Lett. B **833**, 137306 (2022); [arXiv:2205.03877 [hep-ph]].
- [9] G. Senjanović and M. Zantedeschi, *SU(5) grand unification and W-boson mass*, [arXiv:2205.05022 [hep-ph]].
- [10] M. C. Rodriguez, The W-boson mass anomaly and Supersymmetric $SU(3)_C \otimes SU(3)_L \otimes U(1)_N$ Model, [arXiv:2205.09109 [hep-ph]].
- [11] M. C. Rodriguez, Explain the W-boson mass anomaly in the context of Supersymmetric $U(1)_{Y'} \otimes U(1)_{B-L}$ with three identical neutrinos, [arXiv:2209.04653 [hep-ph]].
- [12] G. Senjanovic and R. N. Mohapatra, Exact Left-Right Symmetry and Spontaneous Violation of Parity, Phys. Rev. D **12**, 1502 (1975).
- [13] R. N. Mohapatra and G. Senjanovic, Neutrino Mass and Spontaneous Parity Nonconservation, Phys. Rev. Lett. **44**, 912 (1980).
- [14] G. Senjanovic, Spontaneous Breakdown of Parity in a Class of Gauge Theories, Nucl. Phys. B **153**, 334-364 (1979).
- [15] H. Diaz, V. Pleitez and O. P. Ravinez, Dirac neutrinos in an SU(2) left-right symmetric model, Phys. Rev. D **102**, no.7, 075006 (2020); doi:10.1103/PhysRevD.102.075006, [arXiv:1908.02828 [hep-ph]], and references therein.
- [16] A. E. C. Hernández and I. Schmidt, A renormalizable left-right symmetric model with low scale seesaw mechanisms, Nucl. Phys. B **976**, 115696 (2022); [arXiv:2101.02718 [hep-ph]].
- [17] A. Kokado and T. Saito, New Mixing Angles in the Left-Right Symmetric Model, Phys. Rev. D **92**, 125008 (2015); [arXiv:1508.07406 [hep-ph]].

- [18] J. C. Montero, C. A. de Sousa Pires and V. Pleitez, Spontaneous breaking of a global symmetry in a 331 model, *Phys. Rev. D* **60**, 115003 (1999); doi:10.1103/PhysRevD.60.115003, [arXiv:hep-ph/9903251 [hep-ph]].
- [19] H. Diaz, E. Castillo-Ruiz, O. Pereyra Ravinez and V. Pleitez, Explicit parity violation in $SU(2)_L \otimes SU(2)_R \otimes U(1)_{B-L}$ models, *J. Phys. G* **48**, no.8, 085010 (2021); doi:10.1088/1361-6471/abf9f2, [arXiv:2002.03524 [hep-ph]].
- [20] P. Langacker and S. U. Sankar, Bounds on the Mass of $W(R)$ and the $W(L)$ - $W(R)$ Mixing Angle ξ in General $SU(2)_L \times SU(2)_R \times U(1)$ Models, *Phys. Rev. D* **40**, 1569-1585 (1989).
- [21] M. Frank, Ö. Özdal and P. Poulose, Relaxing LHC constraints on the W_R mass, *Phys. Rev. D* **99**, no.3, 035001 (2019); [arXiv:1812.05681 [hep-ph]].
- [22] N. G. Deshpande and E. Ma, Pattern of Symmetry Breaking with Two Higgs Doublets, *Phys. Rev. D* **18**, 2574 (1978).