

# Generalized parton distributions in spin-3/2 particles

Dongyan Fu<sup>a,b,\*</sup>, Bao-Dong Sun<sup>c,d,e,†</sup>, and Yubing Dong<sup>a,b,‡</sup>

Institute of High Energy Physics, Chinese Academy of Sciences, Beijing 100049, China<sup>a</sup>

School of Physical Sciences, University of Chinese Academy of Sciences, Beijing 101408, China<sup>b</sup>

Guangdong Provincial Key Laboratory of Nuclear Science, Institute of Quantum Matter,

South China Normal University, Guangzhou 510006, China<sup>c</sup>

Guangdong-Hong Kong Joint Laboratory of Quantum Matter,

Southern Nuclear Science Computing Center,

South China Normal University, Guangzhou 510006, China<sup>d</sup>

Helmholtz Institut für Strahlen- und Kernphysik and Bethe

Center for Theoretical Physics, Universität Bonn, D-53115 Bonn, Germany<sup>e</sup>

September 30, 2022

## Abstract

Generalized parton distribution functions (GPDs) of spin-3/2 particles are defined for the first time in this paper. Eight unpolarized and eight polarized GPDs are found. The sum rules between the GPDs and the electromagnetic and gravitational form factors are explicitly displayed. In particular, the relations between GPDs and the helicity amplitudes of the system are derived, based on which several sum rules for the structure functions are obtained.

## 1 Introduction

The electromagnetic form factors (EMFFs) of delta isobar (spin-3/2) have been explored extensively both experimentally and theoretically for a long history [1–6]. It shows the richer information of charge and magnetic structures carried by spin-3/2 particles w.r.t. the lower spin hadrons like nucleon [7–10], deuteron [11–16] and rho meson [17–21] etc. In parallel, the gravitational form factors (GFFs) which are defined by factorizing the matrix element of energy momentum tensor (EMT) characterize the mechanical properties such as the mass, spin densities, as well as the pressure and shear force distributions inside the particles [22–27]. Both the EMFFs and GFFs can be related through the GPDs which is firstly introduced in describing the DVCS process for spin-1/2 particles [28–30] and then for spin-1 particles [31]. The GPDs can also produce PDFs and structure functions in the forward limit. Though the crossing symmetry, GPDs are connected with generalized distribution amplitudes (GDAs) which is also a powerful tool to access GFFs [32]. It makes the concept of GPDs very important to understand the abundant experimental and theoretical (by model-(in)dependent studies, Lattice QCD calculations etc.) information of hadron structures in a unified theoretical framework [33–36]. Thus, the GPDs and the related quantities have been receiving many interest for particles of spin-0 (e.g. pion and kaon mesons [37–42]), spin-1/2 (e.g. nucleon [43–46], and for a nuclei, like <sup>3</sup>He [47, 48]) and spin-1 (e.g. rho meson [49–52] and deuteron [47, 48, 53–55]), etc..

In contrast to the increasing interest on the EMFFs and GFFs of delta isobar (spin-3/2), the definition of GPDs for spin-3/2 hadrons is still missing, to the best of our knowledge. One could expect that spin-3/2 GPDs

---

\*fudongyan@ihep.ac.cn

†bao-dong.sun@m.scnu.edu.cn

‡dongyb@ihep.ac.cn

would expand and deepen our understanding on the quantities such as PDFs [56], structure functions, EMFFs and GFFs etc. for higher spin ( $\geq 3/2$ ) particles in a similar method as lower spin cases. Therefore it is of great interest to give the definition and properties of spin-3/2 GPDs and it is the main purpose of this work. Now the most promising "measurement" is the Lattice QCD calculation, e.g. the recent Lattice calculation on spin-3/2 gluonic GFFs [57]. In the future electron-ion collision experiments, EIC [58] (under construction) and EicC [59] (planned), the candidate targets may possess higher spins, including spin-3/2 nuclei like  ${}^7_3\text{Li}$ ,  ${}^9_4\text{Be}$ , and  ${}^{11}_4\text{B}$ , and even spin-5/2 ones like  ${}^{17}_8\text{O}$ ,  ${}^{25}_{12}\text{Mg}$ , and  ${}^{27}_{13}\text{Al}$  [59]. It would be possible to access the spin-3/2 GPDs and helicity amplitudes experimentally [53] in the near future. Nevertheless, spin-3/2 GPDs by itself is important enough in the theoretical aspect.

This work is organized as follows: In Section 2, we give the definitions and the properties of GPDs of the spin-3/2 system. Section 3 displays the sum rules between EMFFs, GFFs and GPDs, and the explicit helicity amplitudes and their relations to the GPDs. Furtherly, the connections of structure functions to GPDs, and two sum rules of the structure functions are derived for the spin-3/2 system. Section 4 will be devoted for a short summary and discussion.

## 2 Generalized parton distributions of a spin-3/2 particle

In this section, we will firstly give the conventions used in this work and then the GPDs for the spin-3/2 state. The four-vector  $v$  in light-cone coordinates is given as

$$v = (v^+, v^-, \mathbf{v}), \quad \text{with} \quad v^\pm = v^0 \pm v^3 \quad \text{and} \quad \mathbf{v} = (v^1, v^2), \quad (1)$$

and the light-like four vector  $n = (0, 2, \mathbf{0})$  with  $n^2 = 0$ . The scalar product of two four-vectors is  $u \cdot v = \frac{1}{2}u^+v^- + \frac{1}{2}u^-v^+ - \mathbf{u} \cdot \mathbf{v}$ . The convention for the momenta is

$$P = \frac{p+p'}{2}, \quad q = p' - p, \quad t = q^2, \quad (2)$$

where  $p$  and  $p'$  are the initial and final momenta, respectively. The conventions of variables skewness  $\xi$  and  $x$  in the representations of GPDs are

$$\xi = -\frac{q \cdot n}{2P \cdot n} = -\frac{q^+}{2P^+} \quad (|\xi| \leq 1), \quad \text{and} \quad x = \frac{k \cdot n}{P \cdot n} = \frac{k^+}{P^+} \quad (-1 \leq x \leq 1). \quad (3)$$

The partonic four-momentum is  $k - q/2$  and the fraction of the momentum carried by parton over that of the total system in the light cone direction is  $x_q = (x + \xi)/(1 + \xi)$  [60].

The GPDs are defined through the nondiagonal matrix elements of quark and gluon nonlocal current operators at a light-like separation [61]. The general decompositions for quarks can be written as

$$\begin{aligned} V_{\lambda'\lambda} &= \frac{1}{2} \int \frac{dz^-}{2\pi} e^{ix(P \cdot z)} \langle p', \lambda' | \bar{\psi}(-z/2) \not{n} \psi(z/2) | p, \lambda \rangle \Big|_{z^+=0, \mathbf{z}=\mathbf{0}} \\ &= -\bar{u}_{\alpha'}(p', \lambda') \mathcal{H}^{\alpha'\alpha}(x, \xi, t) u_\alpha(p, \lambda), \end{aligned} \quad (4)$$

for unpolarized case, and

$$\begin{aligned} A_{\lambda'\lambda} &= \frac{1}{2} \int \frac{dz^-}{2\pi} e^{ix(P \cdot z)} \langle p', \lambda' | \bar{\psi}(-z/2) \not{n} \gamma_5 \psi(z/2) | p, \lambda \rangle \Big|_{z^+=0, \mathbf{z}=\mathbf{0}} \\ &= -\bar{u}_{\alpha'}(p', \lambda') \tilde{\mathcal{H}}^{\alpha'\alpha}(x, \xi, t) u_\alpha(p, \lambda), \end{aligned} \quad (5)$$

for polarized case, where  $\lambda(\lambda')$  are the helicities of incoming (outgoing) spin-3/2 particles and  $\alpha(\alpha')$  are reserved indices for the initial (final) states in this work. The tensors  $\mathcal{H}^{\alpha'\alpha}$  and  $\tilde{\mathcal{H}}^{\alpha'\alpha}$  define the GPDs as will

be shown in the later context. The Rarita-Schwinger spinor  $u_\alpha(p, \lambda)$  shown in [Appendix A](#) is normalized to  $\bar{u}_\alpha(p, \lambda')u^\alpha(p, \lambda) = -2M\delta_{\lambda'\lambda}$ .

To count the number of independent GPDs, one can define the helicity amplitudes for the scattering of a quark on a spin-3/2 particle as [\[33\]](#)

$$\mathcal{A}_{\lambda'\mu', \lambda\mu} = \int \frac{dz^-}{2\pi} e^{ix(P \cdot z)} \langle p', \lambda' | \mathcal{O}_{\mu'\mu}(z) | p, \lambda \rangle_{z^+=0, \mathbf{z}=\mathbf{0}}. \quad (6)$$

The operators  $\mathcal{O}_{\mu'\mu}$  describe a quark transferring from helicity  $\mu$  to  $\mu'$ . The two helicity-conserved operators are

$$\begin{aligned} \mathcal{O}_{++}(z) &= \frac{1}{4} \bar{\psi}(-z/2) \gamma^+ (1 + \gamma_5) \psi(z/2), \\ \mathcal{O}_{--}(z) &= \frac{1}{4} \bar{\psi}(-z/2) \gamma^+ (1 - \gamma_5) \psi(z/2). \end{aligned} \quad (7)$$

With those definitions, one has

$$\mathcal{A}_{\lambda'\pm, \lambda\pm} = \frac{1}{2} (V_{\lambda'\lambda} \pm A_{\lambda'\lambda}), \quad (8)$$

where  $\pm$  represents the quark helicities. The constraints

$$\begin{aligned} \mathcal{A}_{-\lambda'-\mu, -\lambda-\mu} &= (-1)^{\lambda'-\lambda} \mathcal{A}_{\lambda'\mu, \lambda\mu}^*, \\ \mathcal{A}(x, \xi, t)_{\lambda'\mu, \lambda\mu} &= (-1)^{\lambda'-\lambda} \mathcal{A}^*(x, -\xi, t)_{\lambda\mu, \lambda'\mu}, \end{aligned} \quad (9)$$

can be obtained from the parity and time reversal invariances. And the analogous forms have been found for spin 1/2 and 1 cases in Refs. [\[31, 33\]](#). With these constraints, we finally obtain up to eight independent unpolarized GPDs and eight independent polarized GPDs for the spin-3/2 system.

The building blocks to construct the Lorentz structures accompanying the spin-3/2 GPDs are:  $P^\mu$ ,  $q^\mu$ ,  $n^\mu$ ,  $\gamma^\mu$ ,  $\gamma^5$  (or Levi-Civita tensors  $\epsilon^{\mu\nu\alpha\beta}$ ),  $g^{\mu\nu}$  and  $\sigma^{\mu\nu}$ . With help of the on-shell identities given in Refs. [\[6, 23\]](#) and the properties of the Rarita-Schwinger spinors (see [Appendix A](#)), one can obtain eight independent tensors for the unpolarized case and eight for the polarized case, which agrees with the number-counting from the helicity amplitudes. Equivalently, one can build the tensors accompanying the GPDs of the spin-3/2 particle through the direct product between the unpolarized spin-1/2 and unpolarized spin-1 structures. It is known that tensors accompanying the unpolarized GPDs of a spin-1/2 particle [\[62, 63\]](#), have two linear independent Lorentz structures

$$1, \quad \not{q}, \quad (10)$$

and tensors for the spin-1 case are [\[31, 49\]](#)

$$g^{\alpha'\alpha}, \quad P^{\alpha'} P^\alpha, \quad n^{[\alpha' P^\alpha]}, \quad n^{\{\alpha' P^\alpha\}}, \quad n^{\alpha'} n^\alpha, \quad (11)$$

with  $a^{[\mu b^\nu]} = a^\mu b^\nu - a^\nu b^\mu$  and  $a^{\{\mu b^\nu\}} = a^\mu b^\nu + a^\nu b^\mu$ . The direct product gives,

$$\begin{aligned} g^{\alpha'\alpha}, \quad P^{\alpha'} P^\alpha, \quad n^{[\alpha' P^\alpha]}, \quad n^{\{\alpha' P^\alpha\}}, \quad n^{\alpha'} n^\alpha, \\ g^{\alpha'\alpha} \not{q}, \quad P^{\alpha'} P^\alpha \not{q}, \quad n^{[\alpha' P^\alpha]} \not{q}, \quad n^{\{\alpha' P^\alpha\}} \not{q}, \quad n^{\alpha'} n^\alpha \not{q}. \end{aligned} \quad (12)$$

We use two more on-shell identities, Eqs. [\(B 1\)](#) and [\(B 2\)](#) (see [Appendix B](#)), that can be used to reduce two tensors,  $n^{\{\alpha' P^\alpha\}}$  and  $n^{\{\alpha' P^\alpha\}} \not{q}$  in terms of the rest eight independent ones. Therefore there are eight independent unpolarized GPDs for the spin-3/2 case which can be defined as

$$\begin{aligned} \mathcal{H}^{\alpha'\alpha} &= H_1 g^{\alpha'\alpha} + H_2 \frac{P^{\alpha'} P^\alpha}{M^2} + H_3 \frac{n^{[\alpha' P^\alpha]}}{P \cdot n} + H_4 \frac{M^2 n^{\alpha'} n^\alpha}{(P \cdot n)^2} + H_5 \frac{M g^{\alpha'\alpha} \not{q}}{P \cdot n} \\ &+ H_6 \frac{P^{\alpha'} P^\alpha \not{q}}{M P \cdot n} + H_7 \frac{M n^{[\alpha' P^\alpha]} \not{q}}{(P \cdot n)^2} + H_8 \frac{M^3 n^{\alpha'} n^\alpha \not{q}}{(P \cdot n)^3}. \end{aligned} \quad (13)$$

Similarly, the two independent Lorentz structures of the polarized spin-1/2 GPDs are [61]

$$\gamma^5, \not{n}\gamma^5, \quad (14)$$

and the four independent tensor structures of the spin-1 polarized GPDs are [31, 49]

$$i\epsilon^{nP\alpha'\alpha}, \quad i\epsilon^{nPq\{\alpha P\alpha'\}}, \quad i\epsilon^{nPq[\alpha P\alpha']}, \quad i\epsilon^{nPq\{\alpha n\alpha'\}}. \quad (15)$$

Analogy to the unpolarized case, one can expect that the Lorentz structures of the polarized GPDs in spin-3/2 case can also be expressed as the direct product between the polarized and unpolarized structures for the spin-1/2 and spin-1 cases. Actually, the Lorentz structures of the polarized GPDs can be written in two equivalent ways: with polarization comes from either spin-1/2 part ( $\gamma^5$  terms) or spin-1 part (the Levi-Civita tensors). With the constraints by on-shell identities (Eqs. (B 3) and (B 4)), there are eight independent polarized GPDs in spin-3/2 case:

$$\begin{aligned} \tilde{\mathcal{H}}^{\alpha'\alpha} = & \tilde{H}_1 g^{\alpha'\alpha} \gamma^5 + \tilde{H}_2 \frac{P^{\alpha'} P^\alpha}{M^2} \gamma^5 + \tilde{H}_3 \frac{n^{\{\alpha' P \alpha\}}}{P \cdot n} \gamma^5 + \tilde{H}_4 \frac{M^2 n^{\alpha'} n^\alpha}{(P \cdot n)^2} \gamma^5 + \tilde{H}_5 \frac{M g^{\alpha'\alpha}}{P \cdot n} \not{n} \gamma^5 \\ & + \tilde{H}_6 \frac{P^{\alpha'} P^\alpha}{M (P \cdot n)} \not{n} \gamma^5 + \tilde{H}_7 \frac{M n^{\{\alpha' P \alpha\}}}{(P \cdot n)^2} \not{n} \gamma^5 + \tilde{H}_8 \frac{M^3 n^{\alpha'} n^\alpha}{(P \cdot n)^3} \not{n} \gamma^5. \end{aligned} \quad (16)$$

The time reversal doesn't provide further limits on the number of GPDs but determines their behavior under sign change of the skewedness parameter  $\xi$ ,

$$\begin{aligned} H_i(x, \xi, t) &= H_i(x, -\xi, t) \quad \text{with } i = 1, 2, 4, 5, 6, 8, \\ H_i(x, \xi, t) &= -H_i(x, -\xi, t) \quad \text{with } i = 3, 7, \\ \tilde{H}_j(x, \xi, t) &= -\tilde{H}_j(x, -\xi, t) \quad \text{with } j = 1, 2, 3, 4, \\ \tilde{H}_j(x, \xi, t) &= \tilde{H}_j(x, -\xi, t) \quad \text{with } j = 5, 6, 7, 8. \end{aligned} \quad (17)$$

In the forward limit,  $\xi = 0$ ,  $H_{3,7}(x, 0, t) = 0$ ,  $\tilde{H}_{1,2,3,4}(x, 0, t) = 0$ .

In addition, for the gluon distributions in the spin-3/2 system, instead of the matrix elements in (4) and (5) we now have

$$\begin{aligned} & \frac{n_{\beta'} n_\beta}{P \cdot n} \int \frac{dz^-}{2\pi} e^{ix(P \cdot z)} \left\langle p', \lambda' \left| F^{\beta'\mu}(-z/2) F_\mu^\beta(z/2) \right| p, \lambda \right\rangle \Big|_{z^+=0, \mathbf{z}=0} \\ &= -\bar{u}_{\alpha'}(p', \lambda') \mathcal{H}_g^{\alpha'\alpha}(x, \xi, t) u_\alpha(p, \lambda), \\ & -i \frac{n_{\beta'} n_\beta}{P \cdot n} \int \frac{dz^-}{2\pi} e^{ix(P \cdot z)} \left\langle p', \lambda' \left| F^{\beta'\mu}(-z/2) \tilde{F}_\mu^\beta(z/2) \right| p, \lambda \right\rangle \Big|_{z^+=0, \mathbf{z}=0} \\ &= -\bar{u}_{\alpha'}(p', \lambda') \tilde{\mathcal{H}}_g^{\alpha'\alpha}(x, \xi, t) u_\alpha(p, \lambda), \end{aligned} \quad (18)$$

with  $\tilde{F}^{\alpha\beta} = \frac{1}{2} \epsilon^{\alpha\beta\gamma\delta} F_{\gamma\delta}$  which represents the gluon tensor. The tensors  $\mathcal{H}_g^{\alpha'\alpha}$ ,  $\tilde{\mathcal{H}}_g^{\alpha'\alpha}$  have same structures as those for quark distributions, given in Eqs. (13) and (16).

### 3 Sum rules to form factors and PDFs

#### 3.1 Sum rules

As shown in Ref. [33], the  $(a+1)$ th Mellin moments (in  $x$ ) of the operator defining the GPDs of the system (4) lead to derivative operators between the two fields,

$$\begin{aligned} & (P \cdot n)^{a+1} \int dx x^a \int \frac{dz^-}{2\pi} e^{ixP^+z^-} [\bar{\psi}(-z/2) \not{n} \psi(z/2)] \Big|_{z^+=0, \mathbf{z}=0} \\ &= \left( i \frac{d}{dz^-} \right)^a [\bar{\psi}(-z/2) \not{n} \psi(z/2)] \Big|_{z=0} = \bar{\psi}(0) \not{n} (i \overleftrightarrow{\partial}^+)^a \psi(0). \end{aligned} \quad (19)$$

This relation at operator level connects the GPDs to EMFFs ( $a=0$ ), to GFFs ( $a=1$ ), and other FFs from higher rank current operators.

The decompositions of the matrix elements of the vector [3] and axial [64, 65] currents of the spin-3/2 case should be:

$$\begin{aligned} \langle p', \lambda' | \bar{\psi}(0) \gamma^\mu \psi(0) | p, \lambda \rangle &= -2\bar{u}_{\alpha'}(p', \lambda') \left[ g^{\alpha'\alpha} \left( G_1(t) \frac{P^\mu}{M} + G_5(t) \gamma^\mu \right) \right. \\ &\quad \left. + \frac{P^{\alpha'} P^\alpha}{M^2} \left( G_2(t) \frac{P^\mu}{M} + G_6(t) \gamma^\mu \right) \right] u_\alpha(p, \lambda), \end{aligned} \quad (20)$$

$$\begin{aligned} \langle p', \lambda' | \bar{\psi}(0) \gamma^\mu \gamma^5 \psi(0) | p, \lambda \rangle &= -2\bar{u}_{\alpha'}(p', \lambda') \left[ g^{\alpha'\alpha} \left( -\tilde{G}_1(t) \frac{q^\mu}{2M} + \tilde{G}_5(t) \gamma^\mu \right) \right. \\ &\quad \left. + \frac{P^{\alpha'} P^\alpha}{M^2} \left( -\tilde{G}_2(t) \frac{q^\mu}{2M} + \tilde{G}_6(t) \gamma^\mu \right) \right] \gamma^5 u_\alpha(p, \lambda), \end{aligned} \quad (21)$$

where the different parameters are used for the later convenience of exhibiting the relations between helicity amplitudes and GPDs in a more compact way, and its relation to the traditional notation is  $2(G_1, G_2, G_5, G_6) = (-a_2, c_2, -a_1, c_1)$  [3],  $2(\tilde{G}_1, \tilde{G}_2, \tilde{G}_5, \tilde{G}_6) = (-g_3, h_3, g_1, -h_1)$  [65]. Note that the matrix elements (as well as tensors  $\mathcal{H}^{\alpha'\alpha}$ ,  $\tilde{\mathcal{H}}^{\alpha'\alpha}$  and GPDs) are defined flavor by flavor and one should multiply the electric or weak charges and sum over flavors to get the conventional form factors. The isospin symmetry is not specified in this work.

The GFFs for the spin-3/2 particle are defined as [23, 25]

$$\begin{aligned} & \langle p', \lambda' | \hat{T}^{\mu\nu}(0) | p, \lambda \rangle \\ &= -\bar{u}_{\alpha'}(p', \lambda') \left[ \frac{P^\mu P^\nu}{M} \left( g^{\alpha'\alpha} F_{1,0}^T(t) + \frac{2P^{\alpha'} P^\alpha}{M^2} F_{1,1}^T(t) \right) \right. \\ &\quad \left. + \frac{(q^\mu q^\nu - g^{\mu\nu} q^2)}{4M} \left( g^{\alpha'\alpha} F_{2,0}^T(t) + \frac{2P^{\alpha'} P^\alpha}{M^2} F_{2,1}^T(t) \right) \right. \\ &\quad \left. + Mg^{\mu\nu} \left( g^{\alpha'\alpha} F_{3,0}^T(t) + \frac{2P^{\alpha'} P^\alpha}{M^2} F_{3,1}^T(t) \right) + \frac{P^{\{\mu} i \sigma^{\nu\} q}}{2M} \left( g^{\alpha'\alpha} F_{4,0}^T(t) + \frac{2P^{\alpha'} P^\alpha}{M^2} F_{4,1}^T(t) \right) \right. \\ &\quad \left. - \frac{1}{M} \left( 2q^{\{\mu} g^{\nu\} [\alpha' P^\alpha]} + 8g^{\mu\nu} P^{\alpha'} P^\alpha - g^{\alpha'\{\mu} g^{\nu\}\alpha} q^2 \right) F_{5,0}^T(t) + Mg^{\alpha'\{\mu} g^{\nu\}\alpha} F_{6,0}^T(t) \right] u_\alpha(p, \lambda). \end{aligned} \quad (22)$$

With the relation in Eq. (19) taking  $a = 0$ , the sum rules between GPDs and EMFFs are obtained as

$$\begin{aligned}
M \int_{-1}^1 dx H_i(x, \xi, t) &= G_i(t) \quad \text{with } i = 1, 2, 5, 6, \\
M \int_{-1}^1 dx \tilde{H}_i(x, \xi, t) &= \xi \tilde{G}_i(t) \quad \text{with } i = 1, 2, \\
M \int_{-1}^1 dx \tilde{H}_i(x, \xi, t) &= \tilde{G}_i(t) \quad \text{with } i = 5, 6, \\
M \int_{-1}^1 dx H_j(x, \xi, t) &= M \int_{-1}^1 dx \tilde{H}_j(x, \xi, t) = 0 \quad \text{with } j = 3, 4, 7, 8.
\end{aligned} \tag{23}$$

As one can see from Eqs. (13), (17), (20), and Eqs. (16), (17), (21), the first moments of  $H_3$  and  $\tilde{H}_3$  vanish because of the time reversal. Similarly, the first moments of  $H_i$  ( $i = 4, 7, 8$ ) and  $\tilde{H}_i$  ( $i = 4, 7, 8$ ) disappear due to that they carry the tensor structure  $n^\mu n^\nu / (P \cdot n)^2$  or  $n^\mu n^\nu n^\rho / (P \cdot n)^3$  which has more indices and therefore has no correspondences in the factorization of the matrix element of local current.

Similar procedure can be done for the case of GFFs. With the relation in Eq. (19) taking  $a = 1$ , the sum rules between GPDs and GFFs are obtained as

$$\begin{aligned}
M \int_{-1}^1 dx x H_1(x, \xi, t) &= F_{1,0}^T(t) + \xi^2 F_{2,0}^T(t) - 2F_{4,0}^T(t), \\
M \int_{-1}^1 dx x H_2(x, \xi, t) &= 2F_{1,1}^T(t) + 2\xi^2 F_{2,1}^T(t) - 4F_{4,1}^T(t), \\
M \int_{-1}^1 dx x H_3(x, \xi, t) &= 8\xi F_{5,0}^T(t), \\
M \int_{-1}^1 dx x H_4(x, \xi, t) &= \frac{2t}{M^2} F_{5,0}^T(t) + 2F_{6,0}^T(t), \\
M \int_{-1}^1 dx x H_5(x, \xi, t) &= 2F_{4,0}^T(t), \\
M \int_{-1}^1 dx x H_6(x, \xi, t) &= 4F_{4,1}^T(t), \\
M \int_{-1}^1 dx x H_i(x, \xi, t) &= 0, \quad \text{with } i = 7, 8.
\end{aligned} \tag{24}$$

The second Mellin moment of  $H_7$  vanishes because of the constraint by time reversal invariance and the second Mellin moment of  $H_8$  disappears because of the tensor structure  $n^\mu n^\nu n^\rho / (P \cdot n)^3$  has no correspondences in the parametrization of GFFs in Eq. (22). Analogously, it also holds that the second Mellin moment of  $H_3$  vanishes in the forward limit because of the constraint by time reversal invariance.

### 3.2 Helicity amplitudes

Here we derive the explicit expressions of helicity amplitudes in terms of the obtained GPDs. First of all, we introduce some notations. Let's define  $p_1 \pm ip_2 \equiv |p_\perp|e^{\pm i\phi}$  and  $p'_1 \pm ip'_2 \equiv |p'_\perp|e^{\pm i\phi'}$ , and

$$\begin{aligned}
C &\equiv \sqrt{\frac{1-\xi}{1+\xi}}|p_\perp|e^{-i\phi} - \sqrt{\frac{1+\xi}{1-\xi}}|p'_\perp|e^{-i\phi'}, \\
D &\equiv -\frac{t}{4M^2} - \frac{\xi^2}{1-\xi^2}, \\
K_{\pm i} &\equiv A_1 H_i \pm \xi A_1 \tilde{H}_i + A_2 \tilde{K}_{\pm(i+4)} \quad \text{with } i = 1 \sim 4, \\
\tilde{K}_{\pm j} &\equiv H_j \pm \tilde{H}_j \quad \text{with } j = 1 \sim 8,
\end{aligned} \tag{25}$$

where

$$A_1 \equiv \frac{2}{\sqrt{1-\xi^2}}, \quad A_2 \equiv 2\sqrt{1-\xi^2}. \tag{26}$$

With the above notations, the helicity amplitudes have the following forms,

$$2\mathcal{A}_{\frac{3}{2}+, \frac{3}{2}+} = M \left( K_{+1} + \frac{|C|^2}{8M^2} K_{+2} \right), \tag{27}$$

$$\begin{aligned}
2\mathcal{A}_{\frac{3}{2}+, \frac{1}{2}+} &= -\sqrt{\frac{1+\xi}{1-\xi}} \frac{C}{\sqrt{3}} \left( K_{+1} - \frac{1+\xi}{2} K_{-3} \right) - \frac{C}{\sqrt{3}} \left( \tilde{K}_{-1} + \frac{|C|^2}{8M^2} \tilde{K}_{-2} \right) \\
&\quad - \sqrt{\frac{1+\xi}{1-\xi}} \frac{[D(1-\xi^2) + \xi] C}{2\sqrt{3}(1-\xi^2)} K_{+2},
\end{aligned} \tag{28}$$

$$2\mathcal{A}_{\frac{3}{2}+, (-\frac{1}{2})+} = \sqrt{\frac{1+\xi}{1-\xi}} \frac{C^2}{\sqrt{3}M} \left( \tilde{K}_{-1} - \frac{1+\xi}{2} \tilde{K}_{+3} \right) - \frac{C^2}{8\sqrt{3}M} K_{+2} + \sqrt{\frac{1+\xi}{1-\xi}} \frac{[D(1-\xi^2) + \xi] C^2}{2\sqrt{3}M(1-\xi^2)} \tilde{K}_{-2}, \tag{29}$$

$$2\mathcal{A}_{\frac{3}{2}+, (-\frac{3}{2})+} = \frac{C^3}{8M^2} \tilde{K}_{-2}, \tag{30}$$

$$\begin{aligned}
2\mathcal{A}_{\frac{1}{2}+, \frac{3}{2}+} &= \sqrt{\frac{1-\xi}{1+\xi}} \frac{C^*}{\sqrt{3}} \left( K_{+1} + \frac{1-\xi}{2} K_{+3} \right) + \frac{C^*}{\sqrt{3}} \left( \tilde{K}_{+1} + \frac{|C|^2}{8M^2} \tilde{K}_{+2} \right) \\
&\quad + \sqrt{\frac{1-\xi}{1+\xi}} \frac{[D(1-\xi^2) - \xi] C^*}{2\sqrt{3}(1-\xi^2)} K_{+2},
\end{aligned} \tag{31}$$

$$\begin{aligned}
2\mathcal{A}_{\frac{1}{2}+, \frac{1}{2}+} &= -\frac{2M}{3} \left[ K_{+1} - \frac{1}{2} \left( K_{-1} + \frac{|C|^2}{8M^2} K_{-2} \right) + K_{+3} - K_{-3} + (1-\xi^2) K_{+4} \right] \\
&\quad - \frac{|C|^2}{3M} \left[ \sqrt{\frac{1+\xi}{1-\xi}} \tilde{K}_{+1} + \sqrt{\frac{1-\xi}{1+\xi}} \tilde{K}_{-1} \right] + \frac{|C|^2}{6M\sqrt{1-\xi^2}} (\tilde{K}_{+2} + \tilde{K}_{-2} + 4\xi \tilde{K}_{-3}) \\
&\quad + \frac{2M[D(\xi^2-1)+1]}{3} \left( \frac{2K_{+1}}{(1-\xi^2)} + \frac{K_{+3}}{1+\xi} - \frac{K_{-3}}{1-\xi} \right) - \frac{2M[D^2(1-\xi^2)^2 - \xi^2]}{3(1-\xi^2)^2} K_{+2} \\
&\quad + \frac{|C|^2[D(\xi^2-1)-1]}{6M(1-\xi^2)} \left( \sqrt{\frac{1+\xi}{1-\xi}} \tilde{K}_{+2} + \sqrt{\frac{1-\xi}{1+\xi}} \tilde{K}_{-2} \right),
\end{aligned} \tag{32}$$

$$\begin{aligned}
2\mathcal{A}_{\frac{1}{2}+,(-\frac{1}{2})+} &= \frac{2C}{3} \left[ \tilde{K}_{-1} - \frac{|C|^2}{16M^2} \tilde{K}_{+2} - \tilde{K}_{+3} + \tilde{K}_{-3} + (1-\xi^2)\tilde{K}_{-4} \right] \\
&\quad - \frac{C}{3} \left( \sqrt{\frac{1+\xi}{1-\xi}} K_{-1} + \sqrt{\frac{1-\xi}{1+\xi}} K_{+1} \right) + \frac{C}{6\sqrt{1-\xi^2}} (K_{+2} + K_{-2} + 4\xi K_{+3}) \\
&\quad - \frac{2[D(\xi^2-1)+1]C}{3} \left( \frac{2\tilde{K}_{-1}}{(1-\xi^2)} - \frac{\tilde{K}_{+3}}{1-\xi} + \frac{\tilde{K}_{-3}}{1+\xi} \right) + \frac{2[D^2(\xi^2-1)^2 - \xi^2]C}{3(1-\xi^2)^2} \tilde{K}_{-2} \\
&\quad + \frac{[D(\xi^2-1)-1]C}{6(1-\xi^2)} \left( \sqrt{\frac{1+\xi}{1-\xi}} K_{-2} + \sqrt{\frac{1-\xi}{1+\xi}} K_{+2} \right), \tag{33}
\end{aligned}$$

$$2\mathcal{A}_{\frac{1}{2}+,(-\frac{3}{2})+} = \sqrt{\frac{1-\xi}{1+\xi}} \frac{C^2}{\sqrt{3}M} \left( \tilde{K}_{-1} + \frac{1-\xi}{2} \tilde{K}_{-3} \right) - \frac{C^2}{8\sqrt{3}M} K_{-2} - \sqrt{\frac{1-\xi}{1+\xi}} \frac{[D(\xi^2-1)+\xi]C^2}{2\sqrt{3}M(1-\xi^2)} \tilde{K}_{-2}. \tag{34}$$

The remaining helicity amplitudes can be given by Eqs. (8) and (9). Obviously,  $\mathcal{A}_{\frac{3}{2}+, \frac{3}{2}+}$  and  $\mathcal{A}_{\frac{1}{2}+, \frac{1}{2}+}$  in Eqs. (27) and (32) are unchanged after the time reversal. All the helicity amplitudes satisfy the time reversal constraints in Eq.(9). Noted that  $C$  is complex number and the helicity amplitudes depend on the transverse momentum of both initial and final states. It is known that the forward limit of the GPDs gives the usual parton distributions. In the parton model, for spin-3/2 sector, there are four independent structure functions in deep inelastic scattering at leading twist and leading order in  $\alpha_s$ . They are  $F_1$ ,  $b_1$ ,  $g_1$ ,  $g_2$  whose probabilistic interpretation in terms of quark densities reads [66]

$$\begin{aligned}
F_1(x) &= \frac{1}{2} \sum_q e_q^2 \frac{q_{\uparrow}^{\frac{3}{2}}(x) + q_{\uparrow}^{-\frac{3}{2}}(x) + q_{\uparrow}^{\frac{1}{2}}(x) + q_{\uparrow}^{-\frac{1}{2}}(x)}{2} + \{q \rightarrow \bar{q}\}, \\
b_1(x) &= \frac{1}{2} \sum_q e_q^2 \frac{\left( q_{\uparrow}^{\frac{3}{2}}(x) + q_{\uparrow}^{-\frac{3}{2}}(x) \right) - \left( q_{\uparrow}^{\frac{1}{2}}(x) + q_{\uparrow}^{-\frac{1}{2}}(x) \right)}{2} + \{q \rightarrow \bar{q}\}, \\
g_1(x) &= \frac{1}{2} \sum_q e_q^2 \frac{3 \left( q_{\uparrow}^{\frac{3}{2}}(x) - q_{\uparrow}^{-\frac{3}{2}}(x) \right) + \left( q_{\uparrow}^{\frac{1}{2}}(x) - q_{\uparrow}^{-\frac{1}{2}}(x) \right)}{\sqrt{20}} + \{q \rightarrow \bar{q}\}, \\
g_2(x) &= \frac{1}{2} \sum_q e_q^2 \frac{\left( q_{\uparrow}^{\frac{3}{2}}(x) - q_{\uparrow}^{-\frac{3}{2}}(x) \right) - 3 \left( q_{\uparrow}^{\frac{1}{2}}(x) - q_{\uparrow}^{-\frac{1}{2}}(x) \right)}{\sqrt{20}} + \{q \rightarrow \bar{q}\}. \tag{35}
\end{aligned}$$

In addition, one has  $q_{\uparrow}^{\lambda}(x) = q_{\downarrow}^{-\lambda}(x)$  from parity and  $g_2$  is the new structure function as the spin goes from 1 up to 3/2. Moreover, in the forward limit we have  $\mathcal{A}(x, 0, 0)_{\lambda+, \lambda+} = q_{\uparrow}^{\lambda}(x)$ . Using the results for the helicity amplitudes shown in Eqs. (27-34), one gets

$$\begin{aligned}
2H_1(x) - \frac{2}{3}H_4(x) + 2H_5(x) - \frac{2}{3}H_8(x) &= \frac{q_{\uparrow}^{\frac{3}{2}}(x) + q_{\uparrow}^{-\frac{3}{2}}(x) + q_{\uparrow}^{\frac{1}{2}}(x) + q_{\uparrow}^{-\frac{1}{2}}(x)}{2M}, \\
\frac{2}{3}H_4(x) + \frac{2}{3}H_8(x) &= \frac{\left( q_{\uparrow}^{\frac{3}{2}}(x) + q_{\uparrow}^{-\frac{3}{2}}(x) \right) - \left( q_{\uparrow}^{\frac{1}{2}}(x) + q_{\uparrow}^{-\frac{1}{2}}(x) \right)}{2M}, \\
\frac{2\sqrt{5}}{3}\tilde{H}_5(x) - \frac{2}{3\sqrt{5}}\tilde{H}_8(x) &= \frac{3 \left( q_{\uparrow}^{\frac{3}{2}}(x) - q_{\uparrow}^{-\frac{3}{2}}(x) \right) + \left( q_{\uparrow}^{\frac{1}{2}}(x) - q_{\uparrow}^{-\frac{1}{2}}(x) \right)}{\sqrt{20}M}, \\
\frac{2}{\sqrt{5}}\tilde{H}_8(x) &= \frac{\left( q_{\uparrow}^{\frac{3}{2}}(x) - q_{\uparrow}^{-\frac{3}{2}}(x) \right) - 3 \left( q_{\uparrow}^{\frac{1}{2}}(x) - q_{\uparrow}^{-\frac{1}{2}}(x) \right)}{\sqrt{20}M}. \tag{36}
\end{aligned}$$



With the sum rules (23) for the GPDs, the structure functions have the following sum rules,

$$\int_0^1 dx b_1(x) = 0, \quad \int_0^1 dx g_2(x) = 0, \quad (37)$$

if the quark sea  $q - \bar{q}$  does not contribute to the integral [67]. These two equalities in Eq. (37) are consistent with the sum rules derived from the rotation properties of the structure functions [66].

## 4 Summary and discussions

In this work, the GPDs of spin-3/2 particle are given for the first time. There are eight independent unpolarized GPDs and eight polarized ones. The independent tensors that accompany the distributions can be constructed through the direct product between the tensors accompanying the spin-1/2 unpolarized GPDs and that of the spin-1 unpolarized or polarized GPDs. The sum rules between EMFFs, GFFs and GPDs are obtained through the Mellin moments. In the last section, the helicity amplitudes of the spin-3/2 particle are given in term of GPDs we defined and they depend on the transverse momentums of the initial and final states. The parity and time reversal invariances are satisfied throughoutly. Finally, the structure functions can be expressed in terms of the GPDs in the forward limit. Hopefully, it is expected to be tested in the coming electron ion collision experiments at future EIC and EicC with spin-3/2 targets, such as  ${}^7_3\text{Li}$ . Other measurements of the GPDs of the spin-3/2 particle, like of the  $\Omega$  baryon, may also be possible in the heavy ion collision, where the outgoing  $\Omega$  baryon is rich. Finally, the numerical results for the unpolarized and polarized GPDs of a spin-3/2 particle, take  $\Delta$  isobar as an example, will be given in our forthcoming paper.

## Acknowledgments

The authours are grateful to Wim Cosyn for the helpful discussion on the forms of spinor and polarization vector. This work is supported by the National Natural Science Foundation of China under Grants No. 11975245, No. 11947224, No. 11947228, and No. 12035007. This work is also supported by the Sino-German CRC 110 ‘‘Symmetries and the Emergence of Structure in QCD’’ project by NSFC under Grant No. 12070131001, the Key Research Program of Frontier Sciences, CAS, under Grant No. Y7292610K1, and the National Key Research and Development Program of China under Contracts No. 2020YFA0406300, and Guangdong Provincial funding with Grant No. 2019QN01X172, Guangdong Major Project of Basic and Applied Basic Research No. 2020B0301030008, and the Department of Science and Technology of Guangdong Province with Grant No. 2022A0505030010.

## Appendix A The Rarita-Schwinger spinor

The explicit form of the spinor of spin-3/2 particles, Rarita-Schwinger spinor employed used in our work is [68]

$$u^\alpha(p, \lambda) = \sum_{\rho, \sigma} C_{1\rho, \frac{1}{2}\sigma}^{\frac{3}{2}\lambda} \epsilon^\alpha(p, \rho) u(p, \sigma), \quad (\text{A } 1)$$

where the coefficient in (A 1) is the Clebsch-Gordan coefficient. The explicit light-front form expressions of the

polarization vectors are derived from Ref. [69] by boosting from the rest momentum frame to the moving frame

$$\begin{aligned}
\epsilon^\alpha(p, 0) &= \frac{1}{M} \left( p^+, p^- - \frac{2M^2}{p^+}, \epsilon_\perp(p, 0) \right)^T \quad \text{with} \quad \epsilon_\perp(p, 0) = (p_1, p_2), \\
\epsilon^\alpha(p, +1) &= - \left( 0, \frac{\sqrt{2}(p_1 + ip_2)}{p^+}, \epsilon_\perp(p, +1) \right)^T \quad \text{with} \quad \epsilon_\perp(p, +1) = \left( \frac{1}{\sqrt{2}}, \frac{i}{\sqrt{2}} \right), \\
\epsilon^\alpha(p, -1) &= \left( 0, \frac{\sqrt{2}(p_1 - ip_2)}{p^+}, \epsilon_\perp(p, -1) \right)^T \quad \text{with} \quad \epsilon_\perp(p, -1) = \left( \frac{1}{\sqrt{2}}, \frac{-i}{\sqrt{2}} \right).
\end{aligned} \tag{A 2}$$

The massive positive energy Dirac spinor can be written as [70]

$$u(p, \sigma) = \frac{(\not{p} + M)}{\sqrt{2p \cdot n}} \not{n} \chi_\sigma, \tag{A 3}$$

where  $\chi_\sigma$  is the rest frame spinor. Note that the spinor of Eq. (A 1) satisfies the Rarita-Schwinger equation, as well as the subsidiary constraint equation

$$(\not{p} - M) u^\alpha(p, \lambda) = 0, \quad \gamma_\alpha u^\alpha(p, \lambda) = 0, \quad \partial_\alpha u^\alpha(p, \lambda) = 0. \tag{A 4}$$

## Appendix B On-shell identities

There are some useful on-shell identities which can be derived in Refs. [6, 23]:

$$P^{\{\alpha' n^\alpha\}} i\sigma^{nq} \doteq P \cdot n g^{\alpha' \alpha} i\sigma^{nq} + 2P \cdot n P^{\{\alpha' n^\alpha\}} + \frac{1}{2}(q \cdot n)^2 g^{\alpha' \alpha} + 2q \cdot n P^{\{\alpha' n^\alpha\}} + t n^{\alpha'} n^\alpha, \tag{B 1}$$

$$n^{\{\alpha' P^\alpha\}} \doteq -M \left( 1 - \frac{t}{4M^2} \right) g^{\alpha' \alpha} \left( \frac{P \cdot n}{M} + \frac{i\sigma^{nq}}{2M} \right) + g^{\alpha' \alpha} P \cdot n + \frac{2}{M} P^{\alpha'} P^\alpha \left( \frac{P \cdot n}{M} + \frac{i\sigma^{nq}}{2M} \right), \tag{B 2}$$

in which  $\doteq$  represents the relation in the similar manner as Gordon identity. Analogously, for the case of polarized GPDs, one will need these on-shell identities:

$$\begin{aligned}
4M n^{\{\alpha' P^\alpha\}} \not{n} \gamma^5 &\doteq [4(P \cdot n)^2 - (q \cdot n)^2] g^{\alpha' \alpha} \gamma^5 - 8(P \cdot n) n^{\{\alpha' P^\alpha\}} \gamma^5 + 4(q \cdot n) n^{\{\alpha' P^\alpha\}} \gamma^5 \\
&\quad + 8P^2 n^{\alpha'} n^\alpha \gamma^5 + 2M(q \cdot n) g^{\alpha' \alpha} \not{n} \gamma^5,
\end{aligned} \tag{B 3}$$

$$n^{\{\alpha' P^\alpha\}} \gamma^5 \doteq -\frac{1}{2} g^{\alpha' \alpha} (q \cdot n) \gamma^5 + M g^{\alpha' \alpha} \not{n} \gamma^5 - \frac{P^2}{M} g^{\alpha' \alpha} \not{n} \gamma^5 + \frac{2}{M} P^{\alpha'} P^\alpha \not{n} \gamma^5. \tag{B 4}$$

## References

- [1] H. J. Weber and H. Arenhovel. Isobar Configurations in Nuclei. Phys. Rept., 36:277–348, 1978.
- [2] M. Benmerrouche, R. M. Davidson, and N. C. Mukhopadhyay. Problems of Describing Spin 3/2 Baryon Resonances in the Effective Lagrangian Theory. Phys. Rev. C, 39:2339–2348, 1989.
- [3] S. Nozawa and D. B. Leinweber. Electromagnetic form-factors of spin 3/2 baryons. Phys. Rev. D, 42:3567–3571, 1990.
- [4] V. Pascalutsa. Quantization of an interacting spin - 3 / 2 field and the Delta isobar. Phys. Rev. D, 58:096002, 1998.
- [5] Mauro Napsuciale, Mariana Kirchbach, and Simon Rodriguez. Spin 3/2 Beyond the Rarita-Schwinger Framework. Eur. Phys. J. A, 29:289–306, 2006.

- [6] Dongyan Fu, Bao-Dong Sun, and Yubing Dong. Electromagnetic and gravitational form factors of  $\Delta$  resonance in a covariant quark-diquark approach. Phys. Rev. D, 105(9):096002, 2022.
- [7] Franz Gross, G. Ramalho, and M. T. Pena. A Pure S-wave covariant model for the nucleon. Phys. Rev. C, 77:015202, 2008.
- [8] J. P. B. C. de Melo, T. Frederico, E. Pace, S. Pisano, and G. Salme. Time- and Spacelike Nucleon Electromagnetic Form Factors beyond Relativistic Constituent Quark Models. Phys. Lett. B, 671:153–157, 2009.
- [9] Ian C. Cloët, Wolfgang Bentz, and Anthony W. Thomas. Role of diquark correlations and the pion cloud in nucleon elastic form factors. Phys. Rev. C, 90:045202, 2014.
- [10] W. R. B. de Araújo, J. P. B. C. de Melo, and K. Tsushima. Study of the in-medium nucleon electromagnetic form factors using a light-front nucleon wave function combined with the quark-meson coupling model. Nucl. Phys. A, 970:325–352, 2018.
- [11] Franz Gross. Electromagnetic structure of the deuteron: Review of recent theoretical and experimental results. Eur. Phys. J. A, 17:407–413, 2003.
- [12] Franz Gross and Ronald A. Gilman. The Deuteron: A Mini review. AIP Conf. Proc., 603(1):55–68, 2001.
- [13] Ronald A. Gilman and Franz Gross. Electromagnetic structure of the deuteron. J. Phys. G, 28:R37–R116, 2002.
- [14] M. Garçon and J. W. Van Orden. The Deuteron: Structure and form-factors. Adv. Nucl. Phys., 26:293, 2001.
- [15] Bao-dong Sun and Yu-bing Dong. Deuteron electromagnetic form factors with the light-front approach. Chin. Phys. C, 41(1):013102, 2017.
- [16] Yu-bing Dong, Amand Faessler, Thomas Gutsche, and Valery E. Lyubovitskij. Phenomenological Lagrangian approach to the electromagnetic deuteron form factors. Phys. Rev. C, 78:035205, 2008.
- [17] A. F. Krutov, R. G. Polezhaev, and V. E. Troitsky. Magnetic moment of the  $\rho$  meson in instant-form relativistic quantum mechanics. Phys. Rev. D, 97(3):033007, 2018.
- [18] A. F. Krutov, R. G. Polezhaev, and V. E. Troitsky. The radius of the rho meson determined from its decay constant. Phys. Rev. D, 93(3):036007, 2016.
- [19] Ho-Meoyng Choi and Chueng-Ryong Ji. Electromagnetic structure of the rho meson in the light front quark model. Phys. Rev. D, 70:053015, 2004.
- [20] J. P. B. C de Melo and T. Frederico. Covariant and light front approaches to the rho meson electromagnetic form-factors. Phys. Rev. C, 55:2043, 1997.
- [21] D. Garcia Gudino and G. Toledo Sanchez. Finite width induced modification to the electromagnetic form factors of spin-1 particles. Phys. Rev. D, 81:073006, 2010.
- [22] Maxim V. Polyakov and Peter Schweitzer. Forces inside hadrons: pressure, surface tension, mechanical radius, and all that. Int. J. Mod. Phys. A, 33(26):1830025, 2018.
- [23] Sabrina Cotogno, Cédric Lorcé, Peter Lowdon, and Manuel Morales. Covariant multipole expansion of local currents for massive states of any spin. Phys. Rev. D, 101(5):056016, 2020.
- [24] Bao-Dong Sun and Yu-Bing Dong. Gravitational form factors of  $\rho$  meson with a light-cone constituent quark model. Phys. Rev. D, 101(9):096008, 2020.
- [25] June-Young Kim and Bao-Dong Sun. Gravitational form factors of a baryon with spin-3/2. Eur. Phys. J. C, 81(1):85, 2021.

- [26] Xiangdong Ji and Yizhuang Liu. Momentum-Current Gravitational Multipoles of Hadrons. Phys. Rev. D, 106(3):034028, 2022.
- [27] H. Alharazin, E. Epelbaum, J. Gegelia, U. G. Meißner, and B. D. Sun. Gravitational form factors of the delta resonance in chiral EFT. 9 2022. arXiv: 2209.01233.
- [28] Dieter Müller, D. Robaschik, B. Geyer, F. M. Dittes, and J. Hořejši. Wave functions, evolution equations and evolution kernels from light ray operators of QCD. Fortsch. Phys., 42:101–141, 1994.
- [29] A. V. Radyushkin. Scaling limit of deeply virtual Compton scattering. Phys. Lett. B, 380:417–425, 1996.
- [30] Xiang-Dong Ji. Deeply virtual Compton scattering. Phys. Rev. D, 55:7114–7125, 1997.
- [31] Edgar R. Berger, F. Cano, M. Diehl, and B. Pire. Generalized parton distributions in the deuteron. Phys. Rev. Lett., 87:142302, 2001.
- [32] S. Kumano, Qin-Tao Song, and O. V. Teryaev. Hadron tomography by generalized distribution amplitudes in pion-pair production process  $\gamma^*\gamma \rightarrow \pi^0\pi^0$  and gravitational form factors for pion. Phys. Rev. D, 97(1):014020, 2018.
- [33] M. Diehl. Generalized parton distributions. Phys. Rept., 388:41–277, 2003.
- [34] A. V. Belitsky and A. V. Radyushkin. Unraveling hadron structure with generalized parton distributions. Phys. Rept., 418:1–387, 2005.
- [35] Markus Diehl. Introduction to GPDs and TMDs. Eur. Phys. J. A, 52(6):149, 2016.
- [36] Adam Freese and Gerald A. Miller. Unified formalism for electromagnetic and gravitational probes: Densities. Phys. Rev. D, 105(1):014003, 2022.
- [37] Cristiano Fanelli, Emanuele Pace, Giovanni Romanelli, Giovanni Salme, and Marco Salmistraro. Pion Generalized Parton Distributions within a fully covariant constituent quark model. Eur. Phys. J. C, 76(5):253, 2016.
- [38] Jin-Li Zhang and Jia-Lun Ping. Kaon generalized parton distributions and light-front wave functions in the Nambu–Jona-Lasinio model. Eur. Phys. J. C, 81(9):814, 2021.
- [39] Jin-Li Zhang, Khépani Raya, Lei Chang, Zhu-Fang Cui, José Manuel Morgado, Craig D Roberts, and Jose Rodríguez-Quintero. Measures of pion and kaon structure from generalised parton distributions. Phys. Lett. B, 815:136158, 2021.
- [40] Khepani Raya, Zhu-Fang Cui, Lei Chang, Jose-Manuel Morgado, Craig D. Roberts, and Jose Rodriguez-Quintero. Revealing pion and kaon structure via generalised parton distributions \*. Chin. Phys. C, 46(1):013105, 2022.
- [41] Ho-Meoyng Choi, Chueng-Ryong Ji, and L. S. Kisslinger. Skewed quark distribution of the pion in the light front quark model. Phys. Rev. D, 64:093006, 2001.
- [42] Wojciech Broniowski, Alexander E. Dorokhov, and Enrique Ruiz Arriola. Transversity Form Factors and Generalized Parton Distributions of the Pion in Chiral Quark Models. Few Body Syst., 52:295–300, 2012.
- [43] Neetika Sharma. Momentum transfer dependence of generalized parton distributions. Eur. Phys. J. A, 52(11):338, 2016.
- [44] Markus Diehl and Peter Kroll. Nucleon form factors, generalized parton distributions and quark angular momentum. Eur. Phys. J. C, 73(4):2397, 2013.
- [45] O. V. Selyugin. Nucleon structure and the high energy interactions. Phys. Rev. D, 91(11):113003, 2015. [Erratum: Phys.Rev.D 92, 099901 (2015)].

- [46] Thomas Gutsche, Valery E. Lyubovitskij, and Ivan Schmidt. Nucleon parton distributions in a light-front quark model. Eur. Phys. J. C, 77(2):86, 2017.
- [47] A. Kirchner and Dieter Mueller. Deeply virtual Compton scattering off nuclei. Eur. Phys. J. C, 32:347–375, 2003.
- [48] F. Cano and B. Pire. Deep electroproduction of photons and mesons on the deuteron. Eur. Phys. J. A, 19:423–438, 2004.
- [49] Bao-Dong Sun and Yu-Bing Dong.  $\rho$  meson unpolarized generalized parton distributions with a light-front constituent quark model. Phys. Rev. D, 96(3):036019, 2017.
- [50] Bao-Dong Sun and Yu-Bing Dong. Polarized generalized parton distributions and structure functions of the  $\rho$  meson. Phys. Rev. D, 99(1):016023, 2019.
- [51] Narinder Kumar. Transverse densities and generalized parton distributions of the  $\rho$  meson in the light front quark model. Phys. Rev. D, 99(1):014039, 2019.
- [52] Jin-Li Zhang, Guang-Zhen Kang, and Jia-Lun Ping.  $\rho$  meson generalized parton distributions in the Nambu–Jona-Lasinio model. Phys. Rev. D, 105(9):094015, 2022.
- [53] A. Airapetian et al. Measurement of azimuthal asymmetries associated with deeply virtual Compton scattering on an unpolarized deuterium target. Nucl. Phys. B, 829:1–27, 2010.
- [54] Yubing Dong and Cuiying Liang. Generalized parton distribution functions of a deuteron in a phenomenological Lagrangian approach. J. Phys. G, 40:025001, 2013.
- [55] Chandan Mondal, Dipankar Chakrabarti, and Xingbo Zhao. Deuteron transverse densities in holographic QCD. Eur. Phys. J. A, 53(5):106, 2017.
- [56] Yahui Chai et al. Parton distribution functions of  $\Delta^+$  on the lattice. Phys. Rev. D, 102(1):014508, 2020.
- [57] Dimitra A. Pefkou, Daniel C. Hackett, and Phiala E. Shanahan. Gluon gravitational structure of hadrons of different spin. Phys. Rev. D, 105(5):054509, 2022.
- [58] Proceedings, Probing Nucleons and Nuclei in High Energy Collisions: Dedicated to the Physics of the Electron Ion Collider: Seattle (WA), United States, October 1 - November 16, 2018. WSP, 2 2020.
- [59] Daniele P. Anderle et al. Electron-ion collider in China. Front. Phys. (Beijing), 16(6):64701, 2021.
- [60] T. Frederico, E. Pace, B. Pasquini, and G. Salme. Pion Generalized Parton Distributions with covariant and Light-front constituent quark models. Phys. Rev. D, 80:054021, 2009.
- [61] Xiang-Dong Ji. Gauge-Invariant Decomposition of Nucleon Spin. Phys. Rev. Lett., 78:610–613, 1997.
- [62] Pervez Hoodbhoy and Xiang-Dong Ji. Helicity flip off forward parton distributions of the nucleon. Phys. Rev. D, 58:054006, 1998.
- [63] M. Diehl. Generalized parton distributions with helicity flip. Eur. Phys. J. C, 19:485–492, 2001.
- [64] Constantia Alexandrou, Eric B. Gregory, Tomasz Korzec, Giannis Koutsou, John Negele, Toru Sato, and Antonios Tsapalis. Axial and pseudoscalar form-factors of the  $\Delta^+(1232)$ . PoS, LATTICE2010:141, 2010.
- [65] Yu-Son Jun, Jung-Min Suh, and Hyun-Chul Kim. Axial-vector form factors of the baryon decuplet with flavor SU(3) symmetry breaking. Phys. Rev. D, 102(5):054011, 2020.
- [66] R. L. Jaffe and Aneesh Manohar. Deep Inelastic Scattering from Arbitrary Spin Targets. Nucl. Phys. B, 321:343, 1989.

- [67] S. Kumano. Tensor-polarized quark and antiquark distribution functions in a spin-one hadron. Phys. Rev. D, 82:017501, 2010.
- [68] William Rarita and Julian Schwinger. On a theory of particles with half integral spin. Phys. Rev., 60:61, 1941.
- [69] B. D. Keister and W. N. Polyzou. Relativistic Hamiltonian dynamics in nuclear and particle physics. Adv. Nucl. Phys., 20:225–479, 1991.
- [70] Cédric Lorcé. New explicit expressions for Dirac bilinears. Phys. Rev. D, 97(1):016005, 2018.