

Kaon spectrum revisited

U. Taboada-Nieto,^{1,*} P. G. Ortega,^{1,2,†} D. R. Entem,^{1,2,‡} F. Fernández,^{1,2,§} and J. Segovia^{3,¶}

¹*Departamento de Física Fundamental, Universidad de Salamanca, E-37008 Salamanca, Spain*

²*Instituto Universitario de Física Fundamental y Matemáticas (IUFFyM),*

Universidad de Salamanca, E-37008 Salamanca, Spain

³*Departamento de Sistemas Físicos, Químicos y Naturales,
Universidad Pablo de Olavide, E-41013 Sevilla, Spain*

(Dated: September 27, 2022)

The European Organization for Nuclear Research (CERN) has recently approved a world-unique QCD facility in which an updated version of the external M2 beam line of the CERN SPS in conjunction with a universal spectrometer of the COMPASS experiment is used. One of its main goals is to use highly intense and energetic kaon beams to map out the complete spectrum of excited kaons with an unprecedented precision; having a broad impact not only on low-energy QCD phenomenology, but also on many high-energy particle processes where excited kaons appear, such as the study of CP violation in heavy-meson decays studied at LHCb and Belle II. In support of the experimental effort, the kaon spectrum is computed herein using a constituent quark model which has been successfully applied to a wide range of hadronic observables, from light to heavy quark sectors, and thus the model parameters are completely constrained. The model's prediction can be used as a template against which to compare the already collected data and future experimental findings, in order to distinguish between conventional and exotic kaon states. We also compare our results with those available in the literature in order to provide some general statements, common to all calculations.

I. INTRODUCTION

Most of the data on strange mesons are based on experiments that were performed around 30 years ago. Nevertheless, strange mesons appear in many high-energy particle processes as searches for CP violation in multi-body heavy-meson decays like $B^\pm \rightarrow D^0 K^\pm \rightarrow (K_S^0 \pi^+ \pi^-) K^\pm$ performed by the BaBar [1], Belle [2] and LHCb [3] collaborations. Nowadays, the Particle Data Group (PDG) lists 25 strange mesons that have been detected within the energy region [0.5,3.1] GeV [4]. Among the 25 states, only 14 are well established and included in the summary tables whereas the remaining 11 states still need further confirmation; with two of them having their spin-parity quantum numbers not yet determined.

A completely new horizon in kaon physics has been opened with the CERN's approval of upgrading the external M2 beam line of the CERN SPS in order to provide radio-frequency separated high-intensity and high-energy kaon and antiproton beams. The kaon beam, in combination with a universal spectrometer of the COMPASS experiment, will allow to map out the complete spectrum of excited kaons with unprecedented precision, using novel analysis methods.

There are also proposals and plans for future measurements of strange mesons using either τ or D -meson decays at other facilities such as Belle II, BES III and

LHCb. These studies will be limited by the mass of the initial state which is around 2 GeV in both cases. Besides, a new kaon beam line will be built at J-PARC in the near future [5]; however, the expected low momenta would make it difficult to separate beam and target excitations, which might lead to large systematic uncertainties in the determination of strange-meson spectrum.

In support of the experimental effort, we revisit herein the kaon spectrum using a constituent quark model [6, 7] in which the quark-antiquark interaction is based on the non-perturbative phenomena of dynamical chiral symmetry breaking and color confinement, plus the perturbative one-gluon exchange (OGE) force. As it is well known, the quark model parameters are crucial; we have used values that have been fitted before through hadron [8–12], hadron-hadron [13–16] and multiquark [17–21] phenomenology, from light to heavy quark sectors, and thus the model parameters are completely constrained. The model can be used as a template against which to compare the already collected data of kaons in the PDG, and the future experimental findings, in order to distinguish between conventional quark-antiquark bound states and those whose nature could be in conflict with standard quarkonium interpretations.

The bound-state problem is solved using the Rayleigh-Ritz variational method in which the genuine state's wave function is expanded by means of Gaussian trial functions whose ranges are in geometric progression [22]. This provides enough accuracy and it makes the subsequent evaluation of the decay amplitude matrix elements easier. Moreover, the geometric progression is dense at short distances, so that it allows the description of the dynamics mediated by short range potentials. The fast damping of the gaussian tail is not a problem, since we can choose

* id00746851@usal.es

† pgortega@usal.es

‡ entem@usal.es

§ fdz@usal.es

¶ jsegovia@upo.es

the maximal range much longer than the hadronic size.

The manuscript is arranged as follows. After this introduction, the theoretical framework is presented in Sec. II. Section III is mostly devoted to the analysis and discussion of our theoretical results; we end this section by comparing our spectrum with those produced by other theoretical frameworks. Finally, we summarize and give some prospects in Sec. IV.

II. THEORETICAL FRAMEWORK

We describe in this section the basic phenomenological properties of QCD and how they are implemented into the constituent quark model. There is a detailed description of all different terms of the interacting potential and the variational method used to solve the Schrödinger equation. This section also contains a discussion related with the convenience of using a non-relativistic approach to the kaon sector.

A. CONSTITUENT QUARK MODEL

A consequence of the spontaneous chiral symmetry breaking is that the nearly massless current light quarks that appear in the QCD Lagrangian acquire a constituent quark mass, $M(p)$, which is momentum dependent. To preserve chiral invariance of the QCD Lagrangian new interaction terms appear between constituent quarks which are given by Goldstone boson exchanges.

A simplistic Lagrangian invariant under chiral transformations can be derived as [23]

$$\mathcal{L} = \bar{\psi}(i\gamma^\mu \partial_\mu - MU^{\gamma_5})\psi, \quad (1)$$

where $U^{\gamma_5} = \exp(i\pi^a \lambda^a \gamma_5 / f_\pi)$ with π^a denoting the pseudoscalar fields ($\vec{\pi}, K, \eta_8$) and M the constituent quark mass. The momentum dependent constituent quark mass acts as a natural cut-off of the theory. The chiral quark-quark interaction can be written in the following way

$$V_{qq}(\vec{r}_{ij}) = V_{qq}^C(\vec{r}_{ij}) + V_{qq}^T(\vec{r}_{ij}) + V_{qq}^{SO}(\vec{r}_{ij}), \quad (2)$$

where C , T and SO stand for central, tensor and spin-orbit potentials. The central contribution has four different terms,

$$V_{qq}^C(\vec{r}_{ij}) = V_\pi^C(\vec{r}_{ij}) + V_\sigma^C(\vec{r}_{ij}) + V_K^C(\vec{r}_{ij}) + V_\eta^C(\vec{r}_{ij}), \quad (3)$$

which are given by the following expressions

$$V_\pi^C(\vec{r}_{ij}) = \frac{g_{ch}^2}{4\pi} \frac{m_\pi^2}{12m_i m_j} \frac{\Lambda_\pi^2}{\Lambda_\pi^2 - m_\pi^2} m_\pi \left[Y(m_\pi r_{ij}) - \frac{\Lambda_\pi^3}{m_\pi^3} Y(\Lambda_\pi r_{ij}) \right] \times (\vec{\sigma}_i \cdot \vec{\sigma}_j) \sum_{a=1}^3 (\lambda_i^a \cdot \lambda_j^a), \quad (4a)$$

$$V_\sigma^C(\vec{r}_{ij}) = -\frac{g_{ch}^2}{4\pi} \frac{\Lambda_\sigma^2}{\Lambda_\sigma^2 - m_\sigma^2} m_\sigma \left[Y(m_\sigma r_{ij}) - \frac{\Lambda_\sigma}{m_\sigma} Y(\Lambda_\sigma r_{ij}) \right], \quad (4b)$$

$$V_K^C(\vec{r}_{ij}) = \frac{g_{ch}^2}{4\pi} \frac{m_K^2}{12m_i m_j} \frac{\Lambda_K^2}{\Lambda_K^2 - m_K^2} m_K \left[Y(m_K r_{ij}) - \frac{\Lambda_K^3}{m_K^3} Y(\Lambda_K r_{ij}) \right] \times (\vec{\sigma}_i \cdot \vec{\sigma}_j) \sum_{a=4}^7 (\lambda_i^a \cdot \lambda_j^a), \quad (4c)$$

$$V_\eta^C(\vec{r}_{ij}) = \frac{g_{ch}^2}{4\pi} \frac{m_\eta^2}{12m_i m_j} \frac{\Lambda_\eta^2}{\Lambda_\eta^2 - m_\eta^2} m_\eta \left[Y(m_\eta r_{ij}) - \frac{\Lambda_\eta^3}{m_\eta^3} Y(\Lambda_\eta r_{ij}) \right] \times (\vec{\sigma}_i \cdot \vec{\sigma}_j) [\cos \theta_p (\lambda_i^8 \cdot \lambda_j^8) - \sin \theta_p]. \quad (4d)$$

We consider the physical η -meson instead of the octet one and thus the angle θ_p is introduced. The standard Yukawa function is defined by $Y(x) = e^{-x}/x$. The λ^a are the SU(3) flavour Gell-Mann matrices. The quark masses are m_i and m_π , m_K and m_η are the masses of the SU(3) Goldstone bosons, taken from experimental values. The value of m_σ is given by the partially conserved axial current relation $m_\sigma^2 \simeq m_\pi^2 + 4m_{u,d}^2$ [24]. Note, however, that better determinations of the mass of the σ -meson have been reported since then [25, 26] – see also the recent review [27]; one should simply consider the value used here as a model parameter. Finally, the chiral coupling

constant, g_{ch} , is determined from the πNN coupling constant through

$$\frac{g_{ch}^2}{4\pi} = \frac{9}{25} \frac{g_{\pi NN}^2}{4\pi} \frac{m_{u,d}^2}{m_N^2}, \quad (5)$$

which assumes that flavour SU(3) is an exact symmetry only broken by the different mass of the strange quark.

There are three different contributions to the tensor potential:

$$V_{qq}^T(\vec{r}_{ij}) = V_\pi^T(\vec{r}_{ij}) + V_K^T(\vec{r}_{ij}) + V_\eta^T(\vec{r}_{ij}), \quad (6)$$

given by

$$V_{\pi}^T(\vec{r}_{ij}) = \frac{g_{ch}^2}{4\pi} \frac{m_{\pi}^2}{12m_i m_j} \frac{\Lambda_{\pi}^2}{\Lambda_{\pi}^2 - m_{\pi}^2} m_{\pi} \left[H(m_{\pi} r_{ij}) - \frac{\Lambda_{\pi}^3}{m_{\pi}^3} H(\Lambda_{\pi} r_{ij}) \right] \times S_{ij} \sum_{a=1}^3 (\lambda_i^a \cdot \lambda_j^a), \quad (7a)$$

$$V_K^T(\vec{r}_{ij}) = \frac{g_{ch}^2}{4\pi} \frac{m_K^2}{12m_i m_j} \frac{\Lambda_K^2}{\Lambda_K^2 - m_K^2} m_K \left[H(m_K r_{ij}) - \frac{\Lambda_K^3}{m_K^3} H(\Lambda_K r_{ij}) \right] \times S_{ij} \sum_{a=4}^7 (\lambda_i^a \cdot \lambda_j^a), \quad (7b)$$

$$V_{\eta}^T(\vec{r}_{ij}) = \frac{g_{ch}^2}{4\pi} \frac{m_{\eta}^2}{12m_i m_j} \frac{\Lambda_{\eta}^2}{\Lambda_{\eta}^2 - m_{\eta}^2} m_{\eta} \left[H(m_{\eta} r_{ij}) - \frac{\Lambda_{\eta}^3}{m_{\eta}^3} H(\Lambda_{\eta} r_{ij}) \right] \times S_{ij} [\cos \theta_p (\lambda_i^8 \cdot \lambda_j^8) - \sin \theta_p]. \quad (7c)$$

The quark tensor operator is $S_{ij} = 3(\vec{\sigma}_i \cdot \hat{r}_{ij})(\vec{\sigma}_j \cdot \hat{r}_{ij}) - \vec{\sigma}_i \cdot \vec{\sigma}_j$ and $H(x) = (1 + 3/x + 3/x^2)Y(x)$.

Finally, the spin-orbit potential only presents a contribution coming from the scalar part of the interaction

$$V_{qq}^{SO}(\vec{r}_{ij}) = V_{\sigma}^{SO}(\vec{r}_{ij}) = -\frac{g_{ch}^2}{4\pi} \frac{m_{\sigma}^3}{2m_i m_j} \frac{\Lambda_{\sigma}^2}{\Lambda_{\sigma}^2 - m_{\sigma}^2} \times \left[G(m_{\sigma} r_{ij}) - \frac{\Lambda_{\sigma}^3}{m_{\sigma}^3} G(\Lambda_{\sigma} r_{ij}) \right] (\vec{L} \cdot \vec{S}). \quad (8)$$

In the last equation $G(x)$ is the function $(1+1/x)Y(x)/x$. Beyond the scale of chiral symmetry breaking, the dy-

namics should be governed by QCD perturbative effects. They are taken into account by one-gluon fluctuations around the instanton vacuum. The effective vertex Lagrangian that define such effect is given by

$$\mathcal{L}_{qqg} = i\sqrt{4\pi}\alpha_s \bar{\psi} \gamma_{\mu} G_c^{\mu} \lambda^c \psi, \quad (9)$$

with λ^c being the $SU(3)$ color matrices and G_c^{μ} the gluon field.

The different terms of the potential derived from the Lagrangian above contain central, tensor and spin-orbit contributions. They can be written in the following way

$$V_{\text{OGE}}^{\text{C}}(\vec{r}_{ij}) = \frac{1}{4}\alpha_s (\vec{\lambda}_i^c \cdot \vec{\lambda}_j^c) \left[\frac{1}{r_{ij}} - \frac{1}{6m_i m_j} (\vec{\sigma}_i \cdot \vec{\sigma}_j) \frac{e^{-r_{ij}/r_0(\mu)}}{r_{ij} r_0^2(\mu)} \right], \quad (10a)$$

$$V_{\text{OGE}}^{\text{T}}(\vec{r}_{ij}) = -\frac{1}{16} \frac{\alpha_s}{m_i m_j} (\vec{\lambda}_i^c \cdot \vec{\lambda}_j^c) \left[\frac{1}{r_{ij}^3} - \frac{e^{-r_{ij}/r_g(\mu)}}{r_{ij}} \left(\frac{1}{r_{ij}^2} + \frac{1}{3r_g^2(\mu)} + \frac{1}{r_{ij} r_g(\mu)} \right) \right] S_{ij}, \quad (10b)$$

$$V_{\text{OGE}}^{\text{SO}}(\vec{r}_{ij}) = -\frac{1}{16} \frac{\alpha_s}{m_i^2 m_j^2} (\vec{\lambda}_i^c \cdot \vec{\lambda}_j^c) \left[\frac{1}{r_{ij}^3} - \frac{e^{-r_{ij}/r_g(\mu)}}{r_{ij}^3} \left(1 + \frac{r_{ij}}{r_g(\mu)} \right) \right] \times \left[((m_i + m_j)^2 + 2m_i m_j) (\vec{S}_+ \cdot \vec{L}) + (m_i^2 - m_j^2) (\vec{S}_- \cdot \vec{L}) \right], \quad (10c)$$

where $\vec{S}_{\pm} = \frac{1}{2}(\vec{\sigma}_i \pm \vec{\sigma}_j)$ are the so-called symmetric and anti-symmetric spin-orbit operators, respectively. Besides, $r_0(\mu) = \hat{r}_0 \frac{\mu_{nn}}{\mu_{ij}}$ and $r_g(\mu) = \hat{r}_g \frac{\mu_{nn}}{\mu_{ij}}$ are space-regulators which depend on μ_{ij} , the reduced mass of the quark-antiquark pair. The contact term of the central potential has been regularized as

$$\delta(r_{ij}) \sim \frac{1}{4\pi r_0^2} \frac{e^{-r_{ij}/r_0}}{r_{ij}} \quad (11)$$

The wide energy range needed to provide a consistent description of light, strange and heavy mesons requires an effective scale-dependent strong coupling constant. We

use the frozen coupling constant of Ref. [6]:

$$\alpha_s(\mu_{ij}) = \frac{\alpha_0}{\ln \left(\frac{\mu_{ij}^2 + \mu_0^2}{\Lambda_0^2} \right)}, \quad (12)$$

in α_0 , μ_0 and Λ_0 are parameters of the model determined by a global fit to the meson spectra.

Color confinement is an empirical fact in which quarks, antiquarks and gluons appear always inside colorless structures named hadrons. It is well known that multi-gluon exchanges produce an attractive linearly rising potential proportional to the distance between infinite-heavy quarks. However, sea quarks are also important ingredients of the strong interaction dynamics that con-

tribute to the screening of the rising potential at low momenta and eventually to the breaking of the quark-

antiquark binding string [28]. All these facts have been taken into account in our model by including the terms

$$V_{\text{CON}}^{\text{C}}(\vec{r}_{ij}) = [-a_c(1 - e^{-\mu_c r_{ij}}) + \Delta] (\vec{\lambda}_i^c \cdot \vec{\lambda}_j^c), \quad (13a)$$

$$V_{\text{CON}}^{\text{SO}}(\vec{r}_{ij}) = - \left(\vec{\lambda}_i^c \cdot \vec{\lambda}_j^c \right) \frac{a_c \mu_c e^{-\mu_c r_{ij}}}{4m_i^2 m_j^2 r_{ij}} \left[((m_i^2 + m_j^2)(1 - 2a_s) + 4m_i m_j(1 - a_s))(\vec{S}_+ \cdot \vec{L}) + (m_j^2 - m_i^2)(1 - 2a_s)(\vec{S}_- \cdot \vec{L}) \right], \quad (13b)$$

where a_s controls the mixture between the scalar and vector Lorentz structures of the confinement. At short distances this potential presents a linear behavior with an effective confinement strength $\sigma = -a_c \mu_c (\vec{\lambda}_i^c \cdot \vec{\lambda}_j^c)$ and becomes constant at large distances with a threshold defined by

$$V_{\text{thr}} = \{-a_c + \Delta\} (\vec{\lambda}_i^c \cdot \vec{\lambda}_j^c). \quad (14)$$

No bound states can be found for energies higher than this threshold. The system suffers a transition from a color string configuration between two static color sources into a pair of static mesons due to the breaking of the color string and the most favored decay into hadrons.

Among the different methods to solve the Schrödinger equation in order to find the quark-antiquark bound states, we use the Gaussian Expansion Method [22] because it provides enough accuracy and it makes the subsequent evaluation of the decay amplitude matrix elements easier.

This procedure provides the radial wave function solution of the Schrödinger equation as an expansion in terms of basis functions

$$R_\alpha(r) = \sum_{n=1}^{n_{\text{max}}} c_n^\alpha \phi_{nl}^G(r), \quad (15)$$

where α refers to the channel quantum numbers. The coefficients, c_n^α , and the eigenvalue, E , are determined from the Rayleigh-Ritz variational principle

$$\sum_{n=1}^{n_{\text{max}}} \left[(T_{n'n}^\alpha - EN_{n'n}^\alpha) c_n^\alpha + \sum_{\alpha'} V_{n'n}^{\alpha\alpha'} c_n^{\alpha'} = 0 \right], \quad (16)$$

where $T_{n'n}^\alpha$, $N_{n'n}^\alpha$ and $V_{n'n}^{\alpha\alpha'}$ are the matrix elements of the kinetic energy, the normalization and the potential, respectively. $T_{n'n}^\alpha$ and $N_{n'n}^\alpha$ are diagonal whereas the mixing between different channels is given by $V_{n'n}^{\alpha\alpha'}$.

Following Ref. [22], we employ Gaussian trial functions whose ranges are in geometric progression. This enables the optimization of ranges employing a small number of free parameters. Moreover, the geometric progression is dense at short distances, so that it allows the description

TABLE I. Quark model parameters.

Quark masses	m_n (MeV)	313
	m_s (MeV)	555
Goldstone Bosons	m_π (fm ⁻¹)	0.70
	m_σ (fm ⁻¹)	3.42
	m_K (fm ⁻¹)	2.51
	m_η (fm ⁻¹)	2.77
	Λ_π (fm ⁻¹)	4.20
	Λ_σ (fm ⁻¹)	4.20
	Λ_K (fm ⁻¹)	4.20
	Λ_η (fm ⁻¹)	4.20
OGE	$g_{ch}^2/4\pi$	0.54
	θ_p (°)	-15
	α_0	2.118
	Λ_0 (fm ⁻¹)	0.113
	μ_0 (MeV)	36.976
	\hat{r}_0 (fm)	0.181
Confinement	\hat{r}_g (fm)	0.259
	a_c (MeV)	507.4
	μ_c (fm ⁻¹)	0.576
	Δ (MeV)	184.432
	a_s	0.81

of the dynamics mediated by short range potentials. The fast damping of the gaussian tail is not a problem, since we can choose the maximal range much longer than the hadronic size.

The model parameters fitted over all meson spectra and relevant for this work are shown in Table I.

B. Relativity and model-independence

Model estimates of the mean momentum, $\langle p \rangle$, of a light constituent quark, with mass M , inside a meson typically yield $\langle p \rangle \sim M$. It might therefore be argued that bound-state calculations involving light quark systems should only be undertaken within models that, at some level, incorporate relativity. This potential weakness of

the nonrelativistic quark model has long been considered. For example, Ref. [29] remarks that a non-relativistic treatment of quark motion is inaccurate. However, using scales that are internally consistent, it is not ultra-relativistic. Therefore, the non-relativistic approximation must be useful. The point is also canvassed in Ref. [30], which opens with the question “Why does the non-relativistic quark model work?” and proceeds to provide a range of plausible answers. These discussions are complemented by Ref. [31], which devotes itself to “The significance of the treatment of relativistically moving constituents by an effective non-relativistic Schrödinger equation [...]”. The conclusion of these discourses and many others is simple: the non-relativistic model has proved very useful, unifying a wide range of observables within a single framework.

This last observation provides our rationale for employing a non-relativistic model for the analysis herein. Namely, we take a pragmatic view: the non-relativistic quark model is a useful tool. The practical reason for its success is simple: the model has many parameters; they are fitted to a body of data; and, consequently, on this domain, the model cannot be wrong numerically. If one adds relativistic effects in one way or another, there are similar parameters in the new potential. They, too, are fitted to data; and hence the resulting model cannot produce results that are very different from the original non-relativistic version. The values of the parameters in the potential are modified, but the potential is not observable, so nothing substantive is altered.

III. RESULTS

Beyond the renewed experimental interest in studying the spectrum of kaons, its theoretical study is interesting for several reasons. For instance, the strange mesons exhibit explicit flavor, and thus C -parity cannot be a good quantum number, resulting in the fact that J^{PC} -exotic quark-gluon hybrid kaons should appear as an overpopulation of states in the conventional excited kaon spectrum. An example of this is the $J^P = 1^-$ channel that should exhibit a rich number of kaon-like states, with some of them not fitting the quark model expectations, because the $J^{PC} = 1^{-+}$ exotics are included beside the conventional 1^{--} states. Another important feature of kaons, also related with the fact that they exhibit explicit flavor, is the non-presence of quark-antiquark annihilation which makes its spectrum an ideal ground to compare with the isoscalar light-meson sector, where $q\bar{q}$ states can mix with the so-called glueballs.

Tables II to XIII show the masses, in MeV, of the strange mesons predicted by our constituent quark model for spin-parity $J^P = 0^-$ to 5^- , respectively. Moreover, they compare our spectra with those reported by other theoretical approaches. Figure 1 shows a global picture of our kaon spectrum, comparing it with the states collected in PDG. If the kaon state is well-established and

TABLE II. Masses, in MeV, predicted by our constituent quark model for kaon states with quantum numbers $J^P = 0^-$. First column shows the name of the corresponding meson, second column refers to the spin-parity quantum numbers, third column is dedicated to the $n^{2S+1}L_J$ dominant partial wave in the kaon’s wave function, fourth column shows our predicted mass and the successive columns collect the masses reported by PDG first and by other theoretical approaches later.

Meson	J^P	Dom.	The.	Exp.	[33]	[34]	[35]
K	0^-	1^1S_0	481	495	498	462	482
		2^1S_0	1512	1482 ± 16	1457	1454	1538
		3^1S_0	2018	...	1924	2065	2065
		4^1S_0	2318	...	2248
		5^1S_0	2488
		6^1S_0	2567

thus it appears in the PDG’s summary table, the corresponding box is colored in blue, otherwise it is drawn in red. It is worth mentioning here that our spectrum is finite because the energy of the quark-antiquark bound-state cannot be larger than $m_q + m_{\bar{q}} + V_{\text{thr}} = 2591$ MeV. There is only one kaon state listed in the PDG whose mass is larger than such a threshold. Particularly, the $K(3100)$, which was measured by only one collaboration in 1993 [32], is omitted from the summary table of PDG and collected with the warning of needing confirmation.

We proceed now to describe in detail our theoretical findings.

Besides the ground state kaon $K(495)$, the excited states with equal spin-parity $J^P = 0^-$ collected by the PDG are $K(1460)$, $K(1630)$, $K(1830)$ and $K(3100)$. From those, only the first one is confirmed whereas the remaining three are omitted from the summary table, with $K(1630)$ and $K(3100)$ even having unknown quantum numbers. One can see in Table II that our constituent quark model predicts correctly the masses of the well-established $K(495)$ and $K(1460)$ mesons. The deviation with respect to the experimental values is less than 25 MeV, which is within the expected theoretical uncertainty. Moreover, our results for these two states are in reasonable agreement with those predicted by other theoretical frameworks, and this gives us confidence to contrast the rest of excited states. They also compare well and thus, based on our discussion, we do not believe that the observed $K(1630)$, $K(1830)$ and $K(3100)$ fit the quantum numbers $J^P = 0^-$.

The identification of scalar mesons is problematic because some of them have very large decay widths causing a strong overlap between resonances and background. Furthermore, in the mass range of interest, one should expect non- $q\bar{q}$ scalar systems such as glueballs, multi-quark states and even kinematic phenomena like cusp effects due to the presence of close thresholds (the interested reader is referred to, for instance, Refs. [27, 36–39]). The $K_0^*(700)$, $K_0^*(1430)$ and $K_0^*(1950)$ are the three

TABLE III. Masses, in MeV, predicted by our constituent quark model for kaon states with quantum numbers $J^P = 0^+$. Note that we use the same column description as in Table II.

Meson	J^P	Dom.	The.	Exp.	[33]	[34]	[35]
K_0^*	0^+	1^3P_0	1305	1425 ± 50	1257	1234	1362
		2^3P_0	1894	1944 ± 18	1829	1890	1791
		3^3P_0	2242	...	2176	2160	2160
		4^3P_0	2447	...	2424
		5^3P_0	2552
		6^3P_0	2587

TABLE IV. Masses, in MeV, predicted by our constituent quark model for kaon states with quantum numbers $J^P = 1^+$. Note that we use the same column description as in Table II.

Meson	J^P	Dom.	The.	Exp.	[33]	[34]	[35]
K_1	1^+	1^1P_1	1370	1253 ± 7	1364	1352	1294
		1^3P_1	1455	1403 ± 7	1377	1366	1412
		2^1P_1	1925	1650 ± 50	1840	1897	1757
		2^3P_1	1971	...	1861	1928	1893
		3^1P_1	2260	...	2177	2164	2164
		3^3P_1	2286	...	2192	2200	2200
		4^1P_1	2458	...	2422
		4^3P_1	2471	...	2434
		5^1P_1	2556
		5^3P_1	2561
		6^1P_1	2589
6^3P_1	2590		

states collected in the PDG with scalar nature. All theoretical studies of πK scattering that include constraints from chiral symmetry at low energies naturally find the $K_0^*(700)$ as a pole, with a mass less than 800 MeV and a very large width of about 500 MeV. Its nature, however, does not seem to point only to a quark-antiquark bound-state, rather to a dynamically generated 4-quark resonance or a mixture of 2- and 4-quark structures. As other theoretical approaches, we have considered that our constituent quark model does not predict the $K_0^*(700)$ as a naive quark-antiquark bound state and thus the $K_0^*(1430)$ and $K_0^*(1950)$ mesons are assigned to the 1^3P_0 and 2^3P_0 states, respectively. Looking at Table III, our theoretical mass values are in reasonable agreement with those predicted by other approaches but suffer of discrepancies with respect to the experimental data; therefore, every formalism fails on describing quantitatively the scalar strange meson sector although ours seems to deliver a better agreement, but the nature of all K_0^* candidates is still full of controversies.

The mesons with quantum numbers $(I, J^P) = (1/2, 1^+)$ that are summarized in the PDG are $K_1(1270)$, $K_1(1400)$ and $K_1(1650)$. Their experimental masses are somewhat lower than those predicted by our constituent

TABLE V. Masses, in MeV, predicted by our constituent quark model for kaon states with quantum numbers $J^P = 1^-$. Note that we use the same column description as in Table II.

Meson	J^P	Dom.	The.	Exp.	[33]	[34]	[35]
K^*	1^-	1^3S_1	900	896	896	903	897
		2^3S_1	1676	...	1548	1579	1675
		1^3D_1	1787	1718 ± 18	1766	1776	1699
		3^3S_1	2112	...	1983	1950	2156
		2^3D_1	2173	...	2127	2251	2063
		4^3S_1	2372	...	2287
		3^3D_1	2408	...	2385
		5^3S_1	2516
		4^3D_1	2537	...	2573
		6^3S_1	2576
		5^3D_1	2585

quark model. However, the disagreement is alleviated as we advance in excited states taking into account that the so-called $K_1(1650)$ signal is observed experimentally as various peaks in $K^+\phi$ and $K\pi\pi$ invariant mass distributions in the $[1.6 - 1.9]$ GeV energy region. Concerning other theoretical tools, the predicted spectrum is better than ours in this channel; however, it should be remembered that our model parameters are completely constrained and the kaon spectrum is a pure prediction in our case, whereas the other theoretical approaches have performed a fit to the experimental data before providing the full spectrum. An example, it is the mixing between spin-singlet, n^1P_1 , and spin-triplet, n^3P_1 , states that is driven in our model by the so-called anti-symmetric spin-orbit interaction, given unequal quark and antiquark masses; and fixed in a phenomenological way by other approaches. The masses of the axial kaons are quite sensible to this mixing and we predict states with approximately 50% of both 1^1P_1 and 3^1P_1 partial-waves in their wave function.

The strange mesons reported by PDG with spin-parity $J^P = 1^-$ are $K^*(896)$, $K^*(1410)$ and $K^*(1680)$. One can see in Table V that there is agreement between theory and experiment for the masses of the $K^*(896)$ and $K^*(1680)$ mesons; being the last one traditionally assigned to the 1^3D_1 state, but perfectly compatible with our prediction for the 2^3S_1 case, and that reported in Ref. [35]. Note now that the mass of $K^*(1410)$ does not fit any theoretical result, with an experimental value $(150 - 300)$ MeV lower than the reported theoretical ones. Despite the PDG lists $K^*(1410)$ meson as a plausible candidate of the 2^3S_1 state, one must face the following puzzles: (i) the mass of the $K^*(1410)$ is smaller than that of the experimentally assigned 2^1S_0 state, and (ii) if $K^*(1410)$ is a member of the 2^3S_1 nonet, together with $\rho(1450)$, $\omega(1420)$ and $\phi(1680)$, its mass is unexpectedly low from a theoretical point of view. All this makes us to be cautious with respect to the nature of the $K^*(1410)$

TABLE VI. Masses, in MeV, predicted by our constituent quark model for kaon states with quantum numbers $J^P = 2^-$. Note that we use the same column description as in Table II.

Meson	J^P	Dom.	The.	Exp.	[33]	[34]	[35]
K_2	2^-	1^1D_2	1760	1773 ± 8	1778	1791	1709
		1^3D_2	1854	1819 ± 12	1789	1804	1824
		2^1D_2	2160	...	2121	2238	2066
		2^3D_2	2214	2247 ± 17	2131	2254	2163
		3^1D_2	2403	...	2380
		3^3D_2	2432	...	2388
		4^1D_2	2535	...	2570
		4^3D_2	2547	...	2575
		5^1D_2	2585
		5^3D_2	2588

meson and thus we leave it without a conventional quark-antiquark assignment.

The PDG reports four candidates to be strange mesons with spin-parity $J^P = 2^-$; they are $K_2(1580)$, $K_2(1770)$, $K_2(1820)$ and $K_2(2250)$. Among them, the first and last ones are omitted from the PDG's summary table because distinct reasons. The $K_2(1580)$ meson was observed in 1979 when performing a partial-wave analysis of the $K^-\pi^+\pi^-$ system, and needs confirmation. As one can see in Table VI, none of the theoretical approaches locates the ground state in such mass region; in fact, all agree that the ground state should have a mass of around 1.75 GeV, which match the mass of the $K_2(1770)$ meson, making it the most plausible ground state of the strange mesons with $J^P = 2^-$. The last mass measurement of the $K(1820)$ candidate was reported by the LHCb collaboration in 2017 with an experimental value of $(1853 \pm 27^{+18}_{-35})$ MeV [40], which coincides quite nicely with our model prediction; the PDG's average value is a bit smaller because a measurement of 1993 with a mass value 1816 ± 13 [41] is also taken into account. The $K_2(2250)$ signal contains various peaks in strange meson systems within the invariant mass energy region 2.15 – 2.26 GeV, as well as enhancements seen in the antihyperon-nucleon system, either in the mass spectra or in the $J^P = 2^-$ wave. From a theoretical point of view, this multiplicity can be explained because the 2^1D_2 and 2^3D_2 states are close in mass, and within the energy range reported experimentally. Our constituent quark model predicts the following masses 2160 MeV and 2214 MeV for 2^1D_2 and 2^3D_2 , respectively. They are also in the energy range predicted by other theoretical formalisms; in fact, the excited spectra are quite similar when comparison is possible.

The $(I, J^P) = (1/2, 2^+)$ channel is populated until now by two candidates, the $K_2^*(1430)$ and $K_2^*(1980)$ mesons reported by PDG. They have masses (1427 ± 2) MeV and (1994^{+60}_{-50}) MeV, respectively, and match quite well our quark model predictions for the 1^3P_2 and 2^3P_2 states,

TABLE VII. Masses, in MeV, predicted by our constituent quark model for kaon states with quantum numbers $J^P = 2^+$. Note that we use the same column description as in Table II.

Meson	J^P	Dom.	The.	Exp.	[33]	[34]	[35]
K_2^*	2^+	1^3P_2	1454	1427 ± 2	1431	1428	1424
		2^3P_2	1975	1994^{+60}_{-50}	1870	1938	1896
		1^3F_2	2095	...	2093	2151	1964
		3^3P_2	2290	...	2198	2206	2206
		2^3F_2	2364	...	2356	2551	...
		4^3P_2	2474	...	2438
		3^3F_2	2519	...	2560
		5^3P_2	2563
		4^3F_2	2584	...	2695

TABLE VIII. Masses, in MeV, predicted by our constituent quark model for kaon states with quantum numbers $J^P = 3^+$. Note that we use the same column description as in Table II.

Meson	J^P	Dom.	The.	Exp.	[33]	[34]	[35]
K_3	3^+	1^1F_3	2047	...	2075	2131	2009
		1^3F_3	2132	...	2084	2143	2080
		2^1F_3	2340	2324 ± 24	2340	2524	2348
		2^3F_3	2387	...	2347	2536	...
		3^1F_3	2507	...	2550
		3^3F_3	2529	...	2546
		4^1F_3	2581	...	2688
		4^3F_3	2587	...	2691

as can be seen in Table VII. It is worth to note here that the $K_2^*(1980)$ needs confirmation, but the agreement between the different theoretical approaches in the $J^P = 2^+$ medium-energy spectrum makes us confident that $K_2^*(1980)$ is assigned correctly. As one can notice in Table VII, the very high energy spectra differs between the two approaches that are able to deliver results. This is mostly due to the fact that the linear-screening confining interaction is fixed differently, in our case a global fitting from the light to the heavy quark sectors is pursued while in Ref. [33] only the kaon sector is considered.

The strange mesons with spin-parity $J^P = 3^+$ are characterized in its wave function by the presence of the 1^1F_3 and 3^3F_3 partial waves. Their mixture is still quite large, *i.e.* it is around 50%, with a slightly dominance of one of them, exchanging as we go up in energy. Such a mixture for states with higher orbital angular momenta indicates that it is not a short-range interaction, producing pairs of states quite degenerate in mass even for $J = 3$ (and higher), as can be verified in Table VIII. All theoretical computations predict ground states at (2.05 – 2.14) GeV and first excited ones at (2.32 – 2.54) GeV, and the two cases able to provide very high excited states seem also to agree. The only meson collected in the PDG with quantum numbers $(I, J^P) = (1/2, 3^+)$ is the $K_3(2320)$.

TABLE IX. Masses, in MeV, predicted by our constituent quark model for kaon states with quantum numbers $J^P = 3^-$. Note that we use the same column description as in Table II.

Meson	J^P	Dom.	The.	Exp.	[33]	[34]	[35]
K_3^*	3^-	1^3D_3	1810	1779 ± 8	1781	1794	1789
		2^3D_3	2191	...	2121	2237	2182
		1^3G_3	2316	...	2336	2458	2207
		3^3D_3	2421	...	2382
		2^3G_3	2499	...	2540	2814	...
		4^3D_3	2543	...	2571
		3^3G_3	2584	...	2687
		5^3D_3	2587
		4^3G_3	2790

TABLE X. Masses, in MeV, predicted by our constituent quark model for kaon states with quantum numbers $J^P = 4^-$. Note that we use the same column description as in Table II.

Meson	J^P	Dom.	The.	Exp.	[33]	[34]	[35]
K_4	4^-	1^1G_4	2270	...	2309	2422	2255
		1^3G_4	2337	...	2317	2433	2285
		2^1G_4	2476	2490 ± 20	2520	2779	2575
		2^3G_4	2510	...	2526	2789	...
		3^1G_4	2576	...	2673
		3^3G_4	2587	...	2677
		4^1G_4	2782
		4^3G_4	2785

It has been seen in the $J^P = 3^+$ wave of the antihyperon-nucleon system but needs confirmation and thus it has been omitted from the summary table. In any case, the mass of the $K_3(2320)$ is (2324 ± 24) MeV, which fits well the energy range corresponding to the first excited multiplet that contains the 2^1F_3 and 2^3F_3 states.

With respect to the $J^P = 3^+$ strange mesons, a complete different picture is found for the $(I, J^P) = (1/2, 3^-)$ channel. Two partial waves also coexist in this case, the triplet-spin D and G waves, but its mixture is negligible, given by the tensor terms of the quark-antiquark potential. When comparison with other theoretical approaches is possible, our constituent quark model spectrum coincides quite well. For instance, considering all theoretical works, the ground state of the $(I, J^P) = (1/2, 3^-)$ channel is predicted to have a mass between 1.78 GeV and 1.81 GeV, see Table IX. Moreover, the PDG provides a unique $J^P = 3^-$ meson, $K_3^*(1780)$, whose mass (1779 ± 8) MeV makes it a perfect candidate to be the lowest-lying state. Analyzing the medium and high energy spectrum of our quark model and that predicted by Ref. [33], *i.e.* the most completed spectra, one can conclude that both are quite similar with more apparent discrepancies in the very high energy region.

Concerning the $J^P = 4^-$ channel, the $K_4(2500)$ state

TABLE XI. Masses, in MeV, predicted by our constituent quark model for kaon states with quantum numbers $J^P = 4^+$. Note that we use the same column description as in Table II.

Meson	J^P	Dom.	The.	Exp.	[33]	[34]	[35]
K_4^*	4^+	1^3F_4	2080	2048_{-9}^{+8}	2058	2108	2096
		2^3F_4	2359	...	2328	2504	2436
		1^3H_4	2477
		3^3F_4	2517	...	2533
		2^3H_4	2584
		4^3F_4	2585	...	2683

TABLE XII. Masses, in MeV, predicted by our constituent quark model for kaon states with quantum numbers $J^P = 5^+$. Note that we use the same column description as in Table II.

Meson	J^P	Dom.	The.	Exp.	[33]	[34]	[35]
K_5	5^+	1^1H_5	2442
		1^3H_5	2489
		2^1H_5	2572
		2^3H_5	2589

is the only one listed by the PDG but omitted from the summary table because it needs confirmation. Table X shows that the $K_4(2500)$ mass, (2490 ± 20) MeV, is clearly higher than the one predicted by any theoretical approach for the ground state, either 1^1G_4 or 1^3G_4 . Moreover, there is reasonable agreement between the different approaches with respect to the spectrum of the $(I, J^P) = (1/2, 4^-)$ channel. Therefore, we assign the $K_4(2500)$ meson to the 2^1G_4 state, without ruling out the possibility of being the 2^3G_4 state, due to the fact that these two states are quite close in mass. Again, one can see that the spectrum predicted by Ref. [33] differs with ours at very large energies because the way of fitting the model parameters related with the linear-screening potential, *i.e.* our kaon spectrum is a pure prediction where the model parameters have been fitted to the full meson spectra whereas that of Ref. [33] is obtained from a fit of some lowest-lying kaon states.

Table XI shows the $J^P = 4^+$ strange meson spectrum predicted by our constituent quark model and compares it with those predicted by other theoretical approximations. There is good agreement for the ground-state mass, lying around $[2.06 - 2.11]$ GeV. The $K_4^*(2045)$ is the only state collected in the PDG and we are pleased to see that its mass lies a little below the theoretical interval. There is agreement between our constituent quark model and Ref. [33] for the mass of the first excited state whereas Refs. [34, 35] predict such state 100-150 MeV higher. Concerning the medium and high energy mass range, our model and that of Ref. [33] provide similar spectra except for the latest state, probably due to the different way of fitting the confining interaction at large inter-quark distances.

TABLE XIII. Masses, in MeV, predicted by our constituent quark model for kaon states with quantum numbers $J^P = 5^-$. Note that we use the same column description as in Table II.

Meson	J^P	Dom.	The.	Exp.	[33]	[34]	[35]
K_5^*	5^-	1^3G_5	2291	2382 ± 24	2286	2388	2356
		2^3G_5	2488	...	2504	2749	2656
		1^3I_5	2581
		3^3G_5	2588	...	2662
		2^3I_5
		4^3G_5	2776

Unfortunately, there is no experimental data neither theoretical computations to compare our spectrum of strange mesons with spin-parity $J^P = 5^+$. This is mostly due to the fact that these states have very high orbital angular momentum with either spin zero or one. As shown in Table XII, we predict four states, which are very close in mass, until reaching threshold. The ground-state mass is 2442 MeV and the remaining three excited states have masses of 2489 MeV, 2572 MeV and 2589 MeV, respectively. That is to say, there is only 150 MeV between the lowest and highest masses.

The $K_5^*(2380)$ is the only state reported by PDG with quantum numbers $(I, J^P) = (1/2, 5^+)$. It was discovered in the 1986 and not confirmed by any other experimental collaboration. This is the reason because it has been omitted from the summary table. Its mass is (2382 ± 24) MeV lies just in the middle of the masses predicted by our constituent quark model for the 1^3G_5 and 2^3G_5 states. The same situation is found in the latest work related with kaon spectrum [33]; however, it coincides in mass with the ground states reported by Refs. [34, 35]. This feature could be related with the slope of the linear confining potential but it is difficult to make a strong statement without having more data related with this state.

Finally, we show the complete kaon spectrum predicted by our constituent quark model in Figure 1. It also compares the different states with those listed in the PDG with the following legend of colors: the states collected in the summary table are colored in blue whereas those listed by the PDG but omitted from the summary table are shown in red. The states are grouped by their spin-parity J^P quantum numbers, and named accordingly. The dashed line in the figure represents the energy threshold of quark-antiquark bound-states. It is clearly shown in the figure, and common to any theoretical approach, that the kaon spectrum is rich with many quark model states not yet seen experimentally. Moreover, exotic states in this sector are expected and must be added to this quark-antiquark picture. From an experimental point of view, there are 25 strange mesons that have been detected; however, only 14 are well established and included in the summary tables whereas 11 states still need further confirmation. Among the

14 well-established kaons, our quark model picture faces fitting problems with only two of them: $K^*(1410)$ and $K_1(1650)$, which usually present issues to fit in other theoretical frameworks and they are candidates for exotic states. Only three of the eleven states that are omitted from the summary table do not present natural assignments within our quark model. They are $K_0^*(700)$, $K(1630)$ and $K_2(1580)$. The first one shows a nature not compatible with a quark-antiquark state, the second one suffers from even not fixed its spin-parity quantum numbers, and the third one is an isolated measure performed in 1979.

IV. SUMMARY

A completely new horizon in kaon physics has been opened with the CERN's approval of upgrading the external M2 beam line of the CERN SPS in order to provide radio-frequency separated high-intensity and high-energy kaon and antiproton beams. The kaon beam, in combination with a universal spectrometer of the COMPASS experiment, will allow to map out the complete spectrum of excited kaons with unprecedented precision, using novel analysis methods.

In support of the experimental effort, we have revisited herein the kaon spectrum using a constituent quark model in which the quark-antiquark interaction (based on the non-perturbative phenomena of dynamical chiral symmetry breaking and color confinement, plus the perturbative one-gluon exchange force) has been fitted elsewhere through hadron, hadron-hadron and multi-quark phenomenology, from light to heavy quark sectors, and thus the model parameters are completely constrained. The bound-state problem is solved using the Rayleigh-Ritz variational method in which the genuine state's wave function is expanded by means of Gaussian trial functions whose ranges are in geometric progression.

We have compare our results with those available in the literature in order to provide some general statements, common to all calculations. In addition, we have faced our spectrum to the states listed by the PDG. Among the results we have described, the following are of particular interest: First, it is common to any theoretical approach that the kaon spectrum is rich with many quark model states not yet seen experimentally. Second, our quark model picture faces fitting problems with only two of the fourteen well-established kaons. The $K^*(1410)$ and $K_1(1650)$ are usually difficult to describe by any theoretical computation and thus they are candidates for exotic states. Third, only three of the eleven states that are omitted from the summary table do not present natural assignments within our quark model. They are $K_0^*(700)$, $K(1630)$ and $K_2(1580)$. The first one shows a nature not compatible with a quark-antiquark state, the second one suffers from even not fixed its spin-parity quantum numbers, and the third one is an isolated measure performed in 1979. And fourth, when comparison with other theo-




FIG. 1. Spectrum of strange mesons predicted by our constituent quark model (black solid lines). It is compared with that collected in PDG [4]. States included in the PDG summary table are shown in blue, the remaining one are shown in red. The states are grouped by their spin-parity J^P quantum numbers, and named accordingly. It is worth noting that our spectrum is finite because the energy of the quark-antiquark bound-state cannot be larger than $m_q + m_{\bar{q}} + V_{\text{thr}} = 2591$ MeV, which is marked by a dashed line.

retical approaches is possible, most of the states in the low- and medium-energy region compare nicely, whereas the (very) high-energy region is characterized differently because assumed confining interaction.

It is important to remark that our predicted *naïve* quark-antiquark bound states can be influenced by nearby meson-meson thresholds and thus their composition, binding energy and decay properties can be modified. The expansion of the Fock's space including, together with the quark-antiquark degrees of freedom, states of four (anti-)quarks has been done within our approach in the heavy quark sectors and it is expected to

carry out kindred studies in the kaon spectrum, where such effects appear to be even more important, because the ubiquitous Goldstone-boson exchange interactions, and could explain the discrepancies found herein between theory and experiment. This task, however, is beyond the scope of the present manuscript.

ACKNOWLEDGMENTS

This work has been partially funded by EU Horizon 2020 research and innovation program, STRONG-2020

project, under grant agreement no. 824093; Ministerio Español de Ciencia e Innovación, grant no. PID2019-

107844GB-C22 and PID2019-105439GB-C22; and Junta de Andalucía, contract nos. P18-FR-5057 and Operativo FEDER Andalucía 2014-2020 UHU-1264517.

-
- [1] B. Aubert *et al.* (BaBar), *Phys. Rev. D* **78**, 034023 (2008), arXiv:0804.2089 [hep-ex].
- [2] A. Poluektov *et al.* (Belle), *Phys. Rev. D* **81**, 112002 (2010), arXiv:1003.3360 [hep-ex].
- [3] R. Aaij *et al.* (LHCb), *Nucl. Phys. B* **888**, 169 (2014), arXiv:1407.6211 [hep-ex].
- [4] P. A. Zyla *et al.* (Particle Data Group), *PTEP* **2020**, 083C01 (2020).
- [5] K. Aoki *et al.*, (2021), arXiv:2110.04462 [nucl-ex].
- [6] J. Vijande, F. Fernandez, and A. Valcarce, *J. Phys. G* **31**, 481 (2005), arXiv:hep-ph/0411299.
- [7] J. Segovia, D. R. Entem, F. Fernandez, and E. Hernandez, *Int. J. Mod. Phys. E* **22**, 1330026 (2013), arXiv:1309.6926 [hep-ph].
- [8] A. Valcarce, F. Fernandez, P. Gonzalez, and V. Vento, *Phys. Lett. B* **367**, 35 (1996), arXiv:nucl-th/9509009.
- [9] A. Valcarce, H. Garcilazo, and J. Vijande, *Phys. Rev. C* **72**, 025206 (2005), arXiv:hep-ph/0507297.
- [10] J. Segovia, D. R. Entem, and F. Fernandez, *Phys. Rev. D* **83**, 114018 (2011).
- [11] J. Segovia, D. R. Entem, and F. Fernandez, *Phys. Rev. D* **91**, 094020 (2015), arXiv:1502.03827 [hep-ph].
- [12] G. Yang, J. Ping, P. G. Ortega, and J. Segovia, *Chin. Phys. C* **44**, 023102 (2020), arXiv:1904.10166 [hep-ph].
- [13] D. R. Entem, F. Fernandez, and A. Valcarce, *Phys. Rev. C* **62**, 034002 (2000).
- [14] A. Valcarce, H. Garcilazo, F. Fernandez, and P. Gonzalez, *Rept. Prog. Phys.* **68**, 965 (2005), arXiv:hep-ph/0502173.
- [15] P. G. Ortega, J. Segovia, D. R. Entem, and F. Fernández, *Phys. Rev. D* **95**, 034010 (2017), arXiv:1612.04826 [hep-ph].
- [16] P. G. Ortega, J. Segovia, D. R. Entem, and F. Fernández, *Eur. Phys. J. C* **79**, 78 (2019), arXiv:1808.00914 [hep-ph].
- [17] J. Vijande, F. Fernandez, A. Valcarce, and B. Silvestre-Brac, *Eur. Phys. J. A* **19**, 383 (2004), arXiv:hep-ph/0310007.
- [18] G. Yang, J. Ping, and J. Segovia, *Phys. Rev. D* **99**, 014035 (2019), arXiv:1809.06193 [hep-ph].
- [19] G. Yang, J. Ping, and J. Segovia, *Phys. Rev. D* **101**, 074030 (2020), arXiv:2003.05253 [hep-ph].
- [20] P. G. Ortega, J. Segovia, and F. Fernandez, *Phys. Rev. D* **104**, 094004 (2021), arXiv:2107.02544 [hep-ph].
- [21] G. Yang, J. Ping, and J. Segovia, *Phys. Rev. D* **104**, 094035 (2021), arXiv:2109.04311 [hep-ph].
- [22] E. Hiyama, Y. Kino, and M. Kamimura, *Prog. Part. Nucl. Phys.* **51**, 223 (2003).
- [23] D. Diakonov, *Prog. Part. Nucl. Phys.* **51**, 173 (2003), arXiv:hep-ph/0212026.
- [24] M. D. Scadron, *Phys. Rev. D* **26**, 239 (1982).
- [25] R. Garcia-Martin, R. Kaminski, J. R. Pelaez, and J. Ruiz de Elvira, *Phys. Rev. Lett.* **107**, 072001 (2011), arXiv:1107.1635 [hep-ph].
- [26] M. Albaladejo and J. A. Oller, *Phys. Rev. D* **86**, 034003 (2012), arXiv:1205.6606 [hep-ph].
- [27] J. R. Pelaez, *Phys. Rept.* **658**, 1 (2016), arXiv:1510.00653 [hep-ph].
- [28] G. S. Bali, H. Neff, T. Duessel, T. Lippert, and K. Schilling (SESAM), *Phys. Rev. D* **71**, 114513 (2005), arXiv:hep-lat/0505012.
- [29] S. Capstick and N. Isgur, *AIP Conf. Proc.* **132**, 267 (1985).
- [30] A. Manohar and H. Georgi, *Nucl. Phys. B* **234**, 189 (1984).
- [31] W. Lucha, F. F. Schoberl, and D. Gromes, *Phys. Rept.* **200**, 127 (1991).
- [32] A. N. Aleev *et al.* (EXCHARM), *Phys. Atom. Nucl.* **56**, 1358 (1993).
- [33] C.-Q. Pang, J.-Z. Wang, X. Liu, and T. Matsuki, *Eur. Phys. J. C* **77**, 861 (2017), arXiv:1705.03144 [hep-ph].
- [34] S. Godfrey and N. Isgur, *Phys. Rev. D* **32**, 189 (1985).
- [35] D. Ebert, R. N. Faustov, and V. O. Galkin, *Phys. Rev. D* **79**, 114029 (2009), arXiv:0903.5183 [hep-ph].
- [36] F. E. Close and N. A. Tornqvist, *J. Phys. G* **28**, R249 (2002), arXiv:hep-ph/0204205.
- [37] C. Amsler and N. A. Törnqvist, *Physics Reports* **389**, 61 (2004).
- [38] D. V. Bugg, *Phys. Rept.* **397**, 257 (2004), arXiv:hep-ex/0412045.
- [39] E. Klempt and A. Zaitsev, *Phys. Rept.* **454**, 1 (2007), arXiv:0708.4016 [hep-ph].
- [40] R. Aaij *et al.* (LHCb), *Phys. Rev. Lett.* **118**, 022003 (2017), arXiv:1606.07895 [hep-ex].
- [41] D. Aston, T. Bienz, F. Bird, W. Dunwoodie, W. Johnson, P. Kunz, Y. Kwon, D. Leith, L. Levinson, B. Ratcliff, P. Rensing, D. Schultz, S. Shapiro, P. Sinervo, G. Tarnopolsky, N. Toge, A. Waite, S. Williams, N. Awaji, K. Fujii, H. Hayashii, S. Iwata, R. Kajikawa, T. Matsui, A. Miyamoto, H. Ozaki, C. Pak, T. Shimomura, A. Sugiyama, S. Suzuki, T. Tauchi, J. D'Amore, R. Endorf, J. Martinez, B. Meadows, M. Nussbaum, and K. Ukai, *Physics Letters B* **308**, 186 (1993).