

Bottom hadro-chemistry in high-energy hadronic collisions

Min He¹ and Ralf Rapp²

¹*Department of Applied Physics, Nanjing University of Science and Technology, Nanjing 210094, China*

²*Cyclotron Institute and Department of Physics and Astronomy,
Texas A&M University, College Station, Texas 77843-3366, U.S.A.*

(Dated: September 28, 2022)

The hadro-chemistry of bottom quarks (b) produced in hadronic collisions encodes valuable information on the mechanism of color-neutralization in these reactions. Since the b -quark mass is much larger than the typical hadronic scale of ~ 1 GeV, the $b\bar{b}$ pair production is expected to be well separated from their subsequent hadronization process. In particular, a significantly larger fraction of b baryons has been observed in proton-proton (pp) and proton-antiproton ($p\bar{p}$) reactions relative to that measured in e^+e^- collisions, posing a challenge to theoretical descriptions. We propose to resolve this problem by employing a statistical hadronization approach with a largely augmented set of b -hadron states beyond the currently measured spectrum as listed by the particle data group, by using guidance from the relativistic quark model and lattice-QCD computations. Assuming *relative* chemical equilibrium between the yields of different kinds of b hadrons, their thermo-statistical densities are used as fragmentation weights of a b -quarks into the various hadron species. Aided by quark model estimates of the decay systematics of excited states, the fragmentation fractions of weakly-decaying b hadrons are computed and found to agree well with measurements in $p\bar{p}$ collisions at the Tevatron. Employing transverse-momentum (p_T) distributions of b -quarks from perturbative QCD and combining thermo-statistical fragmentation weights with independent fragmentation toward high p_T , we deduce the total $b\bar{b}$ cross section based on the computed hadro-chemistry. At the same time a fair description of the \bar{B}_s^0/B^- and Λ_b^0/B^- ratios measured in pp collisions at the LHC is obtained. In particular, the observed enhancement of Λ_b^0 production is attributed to the feeddown from thus far unobserved excited b baryons. Our study thus corroborates the statistical nature of b -quark hadronization in hadronic collisions at collider energies.

PACS numbers: 25.75.Dw, 12.38.Mh, 25.75.Nq

Keywords: Bottom Quarks, Proton-Proton Collisions, Statistical Hadronization

I. INTRODUCTION

The masses of charm (c) and especially bottom (b) quarks are much greater than the nonperturbative scale of Quantum Chromodynamics (QCD), Λ_{QCD} , and therefore their production in scattering experiments offers valuable tests of perturbative-QCD dynamics [1, 2]. However, the heavy-quark (HQ) conversion into heavy-flavor (HF) hadrons is an intrinsically soft process that usually requires phenomenological modeling of nonperturbative fragmentation functions (FFs) to describe the production yields and momentum spectra of the observed hadrons [1, 3–5]. Fragmentation fractions of heavy quarks into weakly-decaying heavy hadrons, which include feeddown from excited states via strong or electromagnetic decays, provide a critical test of hadronization mechanisms. Taking bottom (b) as an example, these fragmentation fractions are denoted as f_u , f_d , f_s and f_{baryon} , representing the probabilities of a b quark hadronizing into a B^- , \bar{B}^0 , \bar{B}_s^0 meson or a b baryon (or their charge-conjugate counterparts), respectively. Precise knowledge of these fractions is also important to improve the sensitivity of searches for physics beyond the standard model via rare decays of b hadrons [6].

Traditionally, b -quark fragmentation was assumed to be universal across different colliding systems based on the idea that hadronization occurs nonperturbatively at the scale of Λ_{QCD} [7]. This appears to be supported

within uncertainties by consistent values of the f_s/f_d ratio as first measured in e^+e^- collisions at the Z^0 resonance in LEP experiments [8, 9] and recently in pp collisions at the LHC [10–17]. However, a substantially larger value of f_{Λ_b}/f_d has been measured in high-energy b jets produced in $p\bar{p}$ [18] and pp [14, 19, 20] compared to b jets from Z^0 decays [9, 21, 22], thus challenging the universality assumption. Similar discrepancies have been reported in the charm (c) sector [23–25].

In practice FFs for b or c hadrons are usually inferred from e^+e^- annihilation data. Employing these empirical FFs gives a satisfactory description of p_T -differential cross sections for b and c mesons in hadronic collisions as well, within various calculational schemes, such as Fixed Order Next-to-Leading Log (FONLL) [26–28], k_T -factorization [29–31] or general-mass variable flavor number scheme (GM-VFNS) [32–34]. However, the application of empirical FFs for c baryons substantially underestimates Λ_c production, especially at low p_T [31, 34], as recently measured in pp collisions at the LHC [23, 24], further questioning universality assumptions. An extraction of the FFs for b baryons is currently lacking [35].

It has been argued that HQ hadronization depends on the partonic environment of the color-neutralization process, even in elementary reactions [36–42]. In hadronic collisions, HQ hadronization may be affected by the presence of valence quarks in the initial state [36–40], or multi-parton interactions in the final state [41], rendering

recombination as a plausible non-fragmentation mechanism in a quark-rich environment. This has been pursued in the form of statistical coalescence production of c hadrons [43, 44] in pp collisions at LHC energies. In the present work, we generalize the treatment to the bottom sector and compute the b hadro-chemistry in the statistical hadronization approach that determines the hadron yields by the thermo-statistical weights governed by their masses at a universal hadronization temperature [45–47]. A critical ingredient is an enlarged set of b -hadron states, well beyond the currently measured spectrum [48], using guidance from relativistic-quark model (RQM) and lattice-QCD (lQCD) computations [49–52]. Aided by a quark model estimate of the decay systematics of excited b hadrons, we are able to predict the full set of fragmentation fractions of weakly-decaying b hadrons. We also carry out an evaluation of the p_T dependence of the hadro-chemistry via a combined recombination/fragmentation scheme, which will further allow to make predictions for the total $b\bar{b}$ cross section as well as for the \bar{B}_s^0/B^- and Λ_b^0/B^- ratios.

II. BOTTOM-HADRON SPECTRUM AND STRONG DECAY SYSTEMATICS

The experimental effort to search for missing resonances in the HF sector has been ongoing for several decades [53–55]. The current particle data group (PDG) listings are rather scarce [48], especially for b baryons. Many additional b hadrons are predicted by quark model studies [49, 50, 56], in good agreement with lQCD results for the excitation spectrum [51, 52]. We therefore employ a statistical hadronization model (SHM) using two different sets of b -hadrons as input: (a) PDG-only states [48] and (b) a RQM [49, 50] which additionally includes 18 B 's, 16 B_s 's, 27 Λ_b 's, 45 Σ_b 's, 71 Ξ_b 's, and 41 Ω_b 's, not counting spin/isospin degeneracies, amounting to meson (baryon) masses up to 6.5 (7) GeV (including more massive states has little influence on our final results). Since we are mostly concerned with the *relative* production yields of b hadrons, we use the grand-canonical version of SHM, which works well for bulk hadron production in minimum-bias pp collisions at the LHC energies [57, 58] (the smallness of the total number of b hadrons requires a canonical treatment of the b number when computing the *absolute* yields, but the induced canonical suppression factor is common for each b hadron containing a single b quark and thus cancels out in hadron ratios [59]; for the same reason the b fugacity factor, which is to be fixed by the total $b\bar{b}$ cross section, is dropped). The thermal density of a given b hadron of mass m_i and spin-isospin degeneracy d_i and containing N_s^i strange or antistrange quarks is then evaluated at the hadronization temperature T_H as

$$n_i^{\text{primary}} = \frac{d_i}{2\pi^2} \gamma_s^{N_s^i} m_i^2 T_H K_2\left(\frac{m_i}{T_H}\right), \quad (1)$$




FIG. 1: The (a) meson and (b) baryon decay process $A \rightarrow B + C$ in the 3P_0 model.

where K_2 is the modified Bessel function of second order and $\gamma_s \sim 0.6$ [44, 60] the strangeness suppression factor in elementary reactions. While the SHM analysis of light-hadron yields in heavy-ion collisions at the LHC [47] indicates a hadronization temperature very comparable to the pseudo-critical chiral transition temperature $T_c^x \sim 155$ MeV determined in lattice QCD [61, 62], a higher hadronization temperature $T_H \sim 170$ MeV appears to be more appropriate for HF hadrons in elementary reactions [43]. A flavor hierarchy in the effective hadronization temperature has also been suggested based on lattice calculations of flavor-specific susceptibilities [63]. From theoretical arguments, it has long been expected that the bound states of heavy quarks are more robust in the medium than their light-quark counterparts, with higher melting temperatures, cf. *e.g.*, Ref. [64] for recent calculations using the in-medium T -matrix. In the following, we therefore use $T_H = 170$ MeV as our default value and $T_H = 160$ MeV as part of our error estimate.

Information about decays of excited b hadrons into their weakly-decaying ground states is very limited, even for observed states [48]. Instead of taking rather incomplete results available in the literature, we estimate the branching ratios (BR s) of the OZI-allowed strong decays for all employed b hadrons within the 3P_0 pair creation model [65]. As schematically shown in Fig. 1, a pair of quarks with $J^{PC} = 0^{++}$ is created from the vacuum and regroups with the quarks within the initial b hadron into the outgoing meson or baryon. For example, in the case of a baryon decay, there are three ways of quark regrouping [66], leading to a b -baryon plus a light meson or a b -meson plus a light baryon in the final state, cf. Fig. 1(b). We do not attempt to perform full calculations using realistic hadron wavefunctions [67–69], but obtain the needed BR s via counting all possible diagrams of the types in Fig. 1 once a decay channel opens up, by assuming the number of diagrams for the channel to be proportional to its BR . The probability of creating a quark pair is assumed to be $\propto e^{-2m_q/T_H}$; therefore a diagram involving creating a $s\bar{s}$ is weighted by $e^{-2m_s/T_H} / e^{-2m_{u/d}/T_H} \simeq 1/3$ taking the current-quark masses $m_{u/d} \simeq 8$ MeV and $m_s \simeq 100$ MeV (cf. Ref. [70] for a logarithmic dependence of the pair creation strength in the 3P_0 model). We account for direct three-body decays of b -baryons by counting the regroupings resulting from creation of two $u\bar{u}$ or $d\bar{d}$ pairs and find signif-

n_α ($\cdot 10^{-12} \text{ fm}^{-3}$)	B^-	\bar{B}^0	\bar{B}_s^0	Λ_b^0	$\Xi_b^{0,-}$	Ω_b^-
PDG(170)	1.0094	1.0089	0.29308	0.31591	0.10097	0.002341
PDG(160)	0.12655	0.12649	0.036622	0.034241	0.010520	0.00023076
RQM(170)	1.2045	1.2041	0.32513	0.61702	0.19548	0.0063204
RQM(160)	0.14567	0.14561	0.039664	0.061914	0.018819	0.00061087

TABLE I: Thermal densities of weakly-decaying ground-state b hadrons for hadronization temperatures of $T_H=170$ and 160 MeV in the PDG and RQM scenarios.

icant BR s for the $\Lambda_b^0 + n\pi$ channels of the excited Λ_b 's or Σ_b 's decays, which is supported by the measured information on $\Lambda_b(5912)^0$ and $\Lambda_b(5920)^0$ (reminiscent of $\Lambda_c(2595)^+$ and $\Lambda_c(2625)^+$ in the c sector) [48]. While low-lying excited states all end up in their corresponding ground states (including the pure electromagnetic decay of B^* and B_s^*), higher states have significant cross-feeddowns, *e.g.*, $BR \sim 14\%$ for high B 's to $\bar{B}_s^0 + K$, $\sim 85\text{-}100\%$ for high B_s 's to $B^-/\bar{B}^0 + K$, $\sim 20\text{-}30\%$ for high Λ_b 's (Σ_b 's) to $B^-/\bar{B}^0 + N$, $\sim 20\%$ (20%) for high Ξ_b 's to $\Lambda_b^0 + \pi + K$ ($B^-/\bar{B}^0 + \Sigma$), and $\sim 75\%$ ($\sim 25\%$) for high Ω_b 's to $\Xi_b^{0,-} + K$ ($B^-/\bar{B}^0 + \Xi$). These estimates are overall consistent with available results from a relativized quark model [68, 69].

III. BOTTOM-HADRON FRACTIONS AND RATIOS

With the BR s as estimated above, the total densities of the weakly-decaying ground states are obtained via

$$n_\alpha = n_\alpha^{\text{primary}} + \sum_i n_i^{\text{primary}} \cdot BR(i \rightarrow \alpha), \quad (2)$$

which are summarized in Table I, where PDG and RQM scenarios are compared at hadronization temperatures $T_H=170$ MeV and 160 MeV. These densities are converted into fractions of the total b content in Table II, under the constraint of $f_u + f_d + f_s + f_{\Lambda_b^0} + f_{\Xi_b^{0,-}} + f_{\Omega_b^-} = 1$ when neglecting the tiny fractions of states made of two or more heavy quarks (*e.g.*, B_c mesons, doubly-bottom baryons or bottomonia). When going from the PDG to the RQM scenario, a marked transfer of the b content from the meson to the baryon sector occurs, comparable to the experimentally observed b -hadron fractions in e^+e^- vs. $p\bar{p}$ collisions [22] (and reminiscent of the charm sector [25]). Specifically, the fraction of B^- (\bar{B}_s^0) is reduced by $\sim 10(15)\%$, but the fractions of Λ_b^0 and $\Xi_b^{0,-}$ are both enhanced by $\sim 50\%$ upon inclusion of additional baryons in the RQM at $T_H=170$ MeV, relative to the PDG scenario (similarly also at $T_H=160$ MeV). The weakly-decaying b -hadron fractions obtained in the RQM for both $T_H=170$ and 160 MeV turn out to agree with the measurements in $p\bar{p}$ collisions at the Tevatron within uncertainties, $f_u = f_d = 0.340 \pm 0.021$, $f_s = 0.101 \pm 0.015$ and $f_{\text{baryon}} = 0.220 \pm 0.048$ as averaged in [22].

f_α	B^-	\bar{B}^0	\bar{B}_s^0	Λ_b^0	$\Xi_b^{0,-}$
PDG(170)	0.3697	0.3695	0.1073	0.1157	0.03698
PDG(160)	0.3782	0.3780	0.1094	0.1023	0.03144
RQM(170)	0.3391	0.3389	0.09152	0.1737	0.05503
RQM(160)	0.3533	0.3532	0.09620	0.1502	0.04565

TABLE II: Fractions of weakly-decaying ground-state b hadrons (relative to total $b\bar{b}$) from the SHM with $T_H=170$ and 160 MeV in the PDG and RQM scenarios. $\Xi_b^{0,-}$ denotes the sum of two isospin states. The Ω_b^- fraction is $\sim 0.1\%$ (not shown here).

r_α	\bar{B}^0/B^-	\bar{B}_s^0/B^-	Λ_b^0/B^-	$\Xi_b^{0,-}/B^-$
PDG(170)	0.9995	0.2904	0.3129	0.1000
PDG(160)	0.9995	0.2894	0.2706	0.08313
RQM(170)	0.9994	0.2699	0.5122	0.1623
RQM(160)	0.9996	0.2723	0.4250	0.1292

TABLE III: Ratios of \bar{B}^0 , \bar{B}_s^0 , Λ_b^0 and $\Xi_b^{0,-}$ to B^- at $T_H=170$ and 160 MeV in the PDG and RQM scenarios.

The calculated ratios of \bar{B}^0 , \bar{B}_s^0 , Λ_b^0 and $\Xi_b^{0,-}$ to B^- are summarized in Table III. The mesonic ratios are rather stable, but baryonic ratios are more sensitive to variations in the hadronization temperature. While an equal production of B^- and \bar{B}^0 always holds due to isospin symmetry, the \bar{B}_s^0/B^- ratio is reduced by $\sim 7\%$ upon inclusion of additional states in the RQM scenario. The most pronounced effect is caused by the inclusion of missing baryons, enhancing the baryonic ratios by $\sim 60\%$ relative to the PDG scenario, leading to $\Lambda_b^0/B^- \sim 0.51$ with $T_H=170$ MeV, rather comparable to $f_{\Lambda_b^0}/(f_u + f_d) = 0.259 \pm 0.018$ as measured by LHCb in 13 TeV $p\bar{p}$ collisions [14].

IV. BOTTOM-HADRON p_T -SPECTRA FROM FRAGMENTATION AND DECAY SIMULATIONS

To compute the p_T differential cross sections of weakly-decaying ground-state b hadrons, we simulate the fragmentation and decay processes using the b -quark p_t




FIG. 2: Upper panel: p_T -differential cross sections of ground-state B^- (including strong and electromagnetic feeddown) in $\sqrt{s}=7$ and 13 TeV pp collisions for the RQM (solid lines) and PDG (dashed lines) scenarios at $T_H=170$ MeV, in comparison with LHCb data at $2 < y < 2.5$ [71]. Lower panel: The same for B^- , \bar{B}_s^0 , Λ_b^0 and $\Xi_b^{0,-}$ in $\sqrt{s}=7$ TeV pp collisions.

spectrum in $\sqrt{s} = 7$ or 13 TeV pp collisions from FONLL [26, 27]. A b quark sampled from this spectrum is fragmented into b mesons and baryons via the same FF [3] as implemented in FONLL

$$D_{b \rightarrow H_b}(z) \propto z^\alpha (1-z), \quad (3)$$

where $z = p_T/p_t$ is the fraction of the b -hadron's (H_b) momentum, p_T , over the b -quark momentum, p_t . The fragmentation weight of H_b is determined by its thermal density $n_{H_b}^{\text{primary}}$, Eq. (1), normalized by the sum $\sum_{H_b} n_{H_b}^{\text{primary}}$. Each H_b produced from fragmentation is then decayed into the ground-state particles with a constant matrix element, *i.e.* decay kinematics solely determined by phase space and BRs estimated in Sec. II.

The parameter α in Eq. (3) is tuned to fit the slope of the p_T spectra of ground state b hadrons. Within the RQM scenario at $T_H = 170$ MeV, we find that with $\alpha_B = 45$ (for simplicity taken the same for all B mesons), the measured $B^+ + B^-$ p_T -differential cross section at $2 < y < 2.5$ [71] can be described with an underlying total $b\bar{b}$ cross section of $d\sigma^{b\bar{b}}/dy = 34.55 \mu\text{b}$ ($68.87 \mu\text{b}$) in $\sqrt{s} = 7$ TeV (13 TeV) pp collisions, cf. Fig. 2(a). The




FIG. 3: p_T -dependent ratios of \bar{B}_s^0/B^- (upper panel), Λ_b^0/B^- (middle panel) and $\Xi_b^{0,-}/B^-$ (lower panel) for RQM (solid lines) and PDG (dashed lines) scenarios, in comparison to LHCb data [14, 15, 19] as available in $\sqrt{s} = 7$ and 13 TeV pp collisions. The horizontal arrow in the middle panel indicates the LEP average for f_{Λ_b}/f_d [22] from Z^0 decays at an average b -quark transverse momentum of $\langle p_T(b) \rangle \sim 40$ GeV.

latter emerges as a prediction based on the computed hadro-chemistry content and is consistent with the LHCb measurements via semileptonic b decays [72]. Within the PDG scenario, the $B^+ + B^-$ data turn out to be equally well described but with a $\sim 10\%$ smaller total $b\bar{b}$ cross section. The decrease in the latter is due to the reduction of b content in the baryon sector, as demonstrated by the significantly smaller p_T -differential cross sections

of Λ_b^0 and $\Xi_b^{0,-}$ compared to their RQM counterparts, cf. Fig. 2(b) (the \bar{B}_s^0 spectrum slightly increases in accordance with its fraction change as seen in Table II), where we have used $\alpha_{B_s} = 25$ for all B_s mesons and $\alpha_{\text{baryon}} = 8$ for all b baryons.

We now turn to the p_T -dependent b -hadron ratios. For \bar{B}_s^0/B^- , shown in Fig. 3(a), the additional states in the RQM scenario bring down the PDG results significantly, by over 10% at low p_T , providing a better description of the LHCb data for f_s/f_d which is approximately constant with p_T . While the data indicate a slightly larger \bar{B}_s^0/B^- at $\sqrt{s} = 13$ TeV than at 7 TeV, our results have essentially no collision energy dependence (we refrain from varying the strangeness suppression factor γ_s with collision energy). For Λ_b^0/B^- , the RQM scenario is clearly favored by the LHCb data, cf. Fig. 3(b). The substantial gap between data and the results of the PDG scenario is largely overcome by the feeddown of the large set of “missing” baryons included in the RQM calculation, leading to a fair description of the data, including its increasing trend toward low p_T . For comparison, the LEP average of f_{Λ_b}/f_d [22] in Z decays is indicated as a horizontal arrow. Finally, our predictions for the $\Xi_b^{0,-}/B^-$ ratio from RQM and PDG scenarios are compared in Fig. 3(c), exhibiting very similar features as in the case of Λ_b^0/B^- .

V. SUMMARY

Employing the statistical hadronization model we have evaluated the hadro-chemistry of b hadrons in pp collisions at collider energies. The input spectrum of b -hadrons has been taken from theoretical predictions of the relativistic quark model which is largely supported by lattice-QCD computations of the vacuum spectroscopy.

Many of the RQM states, especially in the baryon sector, are not yet observed and therefore go well beyond the current listings of the particle data group. With strong and electromagnetic feeddown as a key ingredient, we have performed quark model estimates of the decay systematics of excited b hadrons which enabled a comprehensive prediction of fragmentation fractions of ground-state (*i.e.*, weakly-decaying) b hadrons for the first time; pertinent ratios turn out to agree with measurements in $p\bar{p}$ collisions at the Tevatron. We have further calculated p_T differential cross sections of the ground-state b hadrons using fragmentation weights of an underlying (perturbative) b -quark spectrum determined by the SHM. The resulting p_T -dependent \bar{B}_s^0/B^- and Λ_b^0/B^- were also found to agree with LHCb data. All our results critically depend on the excited states that the RQM predicts beyond the PDG listings.

Our findings further support the importance of statistical hadronization in hadronic systems, here for the first time systematically elaborated in the bottom sector. They also have significant ramifications for the investigation of the quark-gluon plasma using bottom probes. The latter are among the central pillars of experimental efforts at both RHIC [73] and the LHC in the near future [74–76]. Our analysis provides a controlled reference for transport calculations of b -hadron yields and spectra in the hot QCD medium. In addition, the total $b\bar{b}$ cross section as determined here is a fundamental input to bottomonium transport calculations in heavy-ion collisions, governing their regeneration reactions [77].

Acknowledgments

This work was supported by NSFC grant 12075122 and the U.S. NSF under grant no. PHY-1913286.

-
- [1] E. Norrbin and T. Sjostrand, *Eur. Phys. J. C* **17**, 137-161 (2000).
 - [2] M. L. Mangano, P. Nason and G. Ridolfi, *Nucl. Phys. B* **373**, 295-345 (1992).
 - [3] V. G. Kartvelishvili, A. K. Likhoded and V. A. Petrov, *Phys. Lett. B* **78**, 615-617 (1978).
 - [4] C. Peterson, D. Schlatter, I. Schmitt and P. M. Zerwas, *Phys. Rev. D* **27**, 105 (1983).
 - [5] E. Braaten, K. m. Cheung, S. Fleming and T. C. Yuan, *Phys. Rev. D* **51**, 4819-4829 (1995).
 - [6] V. Khachatryan *et al.* [CMS and LHCb], *Nature* **522**, 68-72 (2015).
 - [7] M. Lisovyi, A. Verbytskyi and O. Zenaiev, *Eur. Phys. J. C* **76**, no.7, 397 (2016).
 - [8] D. Buskulic *et al.* [ALEPH], *Phys. Lett. B* **361**, 221-233 (1995).
 - [9] J. Abdallah *et al.* [DELPHI], *Phys. Lett. B* **576**, 29-42 (2003).
 - [10] R. Aaij *et al.* [LHCb], *Phys. Rev. Lett.* **107**, 211801 (2011).
 - [11] R. Aaij *et al.* [LHCb], *Phys. Rev. D* **85**, 032008 (2012).
 - [12] R. Aaij *et al.* [LHCb], *JHEP* **04**, 001 (2013).
 - [13] G. Aad *et al.* [ATLAS], *Phys. Rev. Lett.* **115**, no.26, 262001 (2015).
 - [14] R. Aaij *et al.* [LHCb], *Phys. Rev. D* **100**, no.3, 031102 (2019).
 - [15] R. Aaij *et al.* [LHCb], *Phys. Rev. Lett.* **124**, no.12, 122002 (2020).
 - [16] R. Aaij *et al.* [LHCb], *Phys. Rev. D* **104**, no.3, 032005 (2021).
 - [17] S. Acharya *et al.* [ALICE], *JHEP* **05**, 220 (2021).
 - [18] T. Aaltonen *et al.* [CDF], *Phys. Rev. D* **77**, 072003 (2008).
 - [19] R. Aaij *et al.* [LHCb], *JHEP* **08**, 143 (2014).
 - [20] R. Aaij *et al.* [LHCb], *Chin. Phys. C* **40**, no.1, 011001 (2016).
 - [21] R. Barate *et al.* [ALEPH], *Eur. Phys. J. C* **5**, 205-227 (1998).
 - [22] Y. S. Amhis *et al.* [HFLAV], *Eur. Phys. J. C* **81**, no.3, 226 (2021).

- [23] S. Acharya *et al.* [ALICE], JHEP **04**, 108 (2018).
- [24] S. Acharya *et al.* [ALICE], Phys. Rev. Lett. **127**, no.20, 202301 (2021).
- [25] S. Acharya *et al.* [ALICE], Phys. Rev. D **105**, no.1, L011103 (2022).
- [26] S. Frixione, P. Nason and G. Ridolfi, JHEP **09**, 126 (2007).
- [27] M. Cacciari, S. Frixione, N. Houdeau, M. L. Mangano, P. Nason and G. Ridolfi, JHEP **10**, 137 (2012).
- [28] S. Catani, S. Devoto, M. Grazzini, S. Kallweit and J. Mazitelli, JHEP **03**, 029 (2021).
- [29] S. Catani, M. Ciafaloni and F. Hautmann, Nucl. Phys. B **366**, 135-188 (1991).
- [30] J. C. Collins and R. K. Ellis, Nucl. Phys. B **360**, 3-30 (1991).
- [31] R. Maciula and A. Szczurek, Phys. Rev. D **98**, no.1, 014016 (2018).
- [32] B. A. Kniehl, G. Kramer, I. Schienbein and H. Spiesberger, Eur. Phys. J. C **41**, 199-212 (2005).
- [33] B. A. Kniehl, G. Kramer, I. Schienbein and H. Spiesberger, Phys. Rev. D **84**, 094026 (2011).
- [34] B. A. Kniehl, G. Kramer, I. Schienbein and H. Spiesberger, Phys. Rev. D **101**, no.11, 114021 (2020).
- [35] G. Kramer and H. Spiesberger, Chin. Phys. C **42**, no.8, 083102 (2018).
- [36] R. C. Hwa, Phys. Rev. D **51**, 85-89 (1995).
- [37] E. Cuautle, G. Herrera and J. Magnin, Eur. Phys. J. C **2**, 473-476 (1998).
- [38] E. Braaten, Y. Jia and T. Mehen, Phys. Rev. Lett. **89**, 122002 (2002).
- [39] R. Rapp and E. V. Shuryak, Phys. Rev. D **67**, 074036 (2003).
- [40] A. V. Berezhnoy and A. K. Likhoded, Phys. Atom. Nucl. **78**, no.2, 292-300 (2015).
- [41] J. R. Christiansen and P. Z. Skands, JHEP **08**, 003 (2015).
- [42] M. He and R. Rapp, Phys. Rev. Lett. **124**, no.4, 042301 (2020).
- [43] A. Andronic, F. Beutler, P. Braun-Munzinger, K. Redlich and J. Stachel, Phys. Lett. B **678**, 350-354 (2009).
- [44] M. He and R. Rapp, Phys. Lett. B **795**, 117-121 (2019).
- [45] P. Braun-Munzinger, K. Redlich and J. Stachel, arXiv:nucl-th/0304013 [nucl-th].
- [46] F. Becattini, arXiv:0901.3643 [hep-ph].
- [47] A. Andronic, P. Braun-Munzinger, K. Redlich and J. Stachel, Nature **561**, no.7723, 321-330 (2018).
- [48] P. A. Zyla *et al.* [Particle Data Group], PTEP **2020**, no.8, 083C01 (2020).
- [49] D. Ebert, R. N. Faustov and V. O. Galkin, Eur. Phys. J. C **66**, 197-206 (2010).
- [50] D. Ebert, R. N. Faustov and V. O. Galkin, Phys. Rev. D **84**, 014025 (2011).
- [51] R. Lewis and R. M. Woloshyn, Phys. Rev. D **79**, 014502 (2009).
- [52] Z. S. Brown, W. Detmold, S. Meinel and K. Orginos, Phys. Rev. D **90**, no.9, 094507 (2014).
- [53] V. Crede and W. Roberts, Rept. Prog. Phys. **76**, 076301 (2013).
- [54] H. X. Chen, W. Chen, X. Liu, Y. R. Liu and S. L. Zhu, Rept. Prog. Phys. **80**, no.7, 076201 (2017).
- [55] M. Charles, tel-01340092.
- [56] W. Roberts and M. Pervin, Int. J. Mod. Phys. A **23**, 2817-2860 (2008).
- [57] N. Sharma, J. Cleymans, B. Hippolyte and M. Paradz, Phys. Rev. C **99**, no.4, 044914 (2019).
- [58] S. Das, D. Mishra, S. Chatterjee and B. Mohanty, Phys. Rev. C **95**, no.1, 014912 (2017).
- [59] Y. Chen and M. He, Phys. Lett. B **815**, 136144 (2021).
- [60] B. I. Abelev *et al.* [STAR], Phys. Rev. C **79**, 034909 (2009).
- [61] A. Bazavov *et al.* [HotQCD], Phys. Rev. D **90**, 094503 (2014).
- [62] S. Borsanyi, Z. Fodor, C. Hoelbling, S. D. Katz, S. Krieg and K. K. Szabo, Phys. Lett. B **730**, 99-104 (2014).
- [63] R. Bellwied, S. Borsanyi, Z. Fodor, S. D. Katz and C. Ratti, Phys. Rev. Lett. **111**, 202302 (2013).
- [64] S. Y. F. Liu and R. Rapp, Phys. Rev. C **97**, no.3, 034918 (2018).
- [65] S. Capstick and W. Roberts, Prog. Part. Nucl. Phys. **45**, S241-S331 (2000).
- [66] C. Chen, X. L. Chen, X. Liu, W. Z. Deng and S. L. Zhu, Phys. Rev. D **75**, 094017 (2007).
- [67] W. Roberts and B. Silvestre-Brac, Few Body Syst. **11**, no.4, 171-193 (1992).
- [68] J. Ferretti and E. Santopinto, Phys. Rev. D **97**, no.11, 114020 (2018).
- [69] G. L. Yu, Z. Y. Li, Z. G. Wang, J. Lu and M. Yan, arXiv:2206.08128 [hep-ph].
- [70] J. Segovia, D. R. Entem and F. Fernández, Phys. Lett. B **715**, 322-327 (2012).
- [71] R. Aaij *et al.* [LHCb], JHEP **12**, 026 (2017).
- [72] R. Aaij *et al.* [LHCb], Phys. Rev. Lett. **118**, no.5, 052002 (2017) [erratum: Phys. Rev. Lett. **119**, no.16, 169901 (2017)].
- [73] C. Dean [Sphenix], PoS **ICHEP2020**, 731 (2021).
- [74] Z. Citron, A. Dainese, J. F. Grosse-Oetringhaus, J. M. Jowett, Y. J. Lee, U. A. Wiedemann, M. Winn, A. Andronic, F. Bellini and E. Bruna, *et al.* CERN Yellow Rep. Monogr. **7**, 1159-1410 (2019).
- [75] Letter of intent for ALICE 3: A next generation heavy-ion experiment at the LHC, <https://cds.cern.ch/record/2803563?ln=en> (2022).
- [76] [LHCb], LHCb-PUB-2022-012.
- [77] X. Du, R. Rapp and M. He, Phys. Rev. C **96**, no.5, 054901 (2017).