

$\Lambda(1520)$ resonance production with respect to transverse sphericity using EPOS3

Nasir Mehdi Malik* and Sanjeev Singh Sambyal

University of Jammu

* nasir.mehdi.malik@cern.ch

September 29, 2022



51st International Symposium on Multiparticle Dynamics (ISMD2022)

Pitlochry, Scottish Highlands, 1-5 August 2022

doi:[10.21468/SciPostPhysProc.7](https://doi.org/10.21468/SciPostPhysProc.7)

Abstract

Resonances are sensitive to the properties of the medium created in heavy ion collision. They also provide insight into the properties of the hadronic phase. Studying the dependence of the yield of resonances on transverse sphericity and multiplicity allows us to understand the resonance production mechanism with event topology and system size, respectively. The results reported pertains to $\Lambda(1520)$. The data from EPOS3 is used for the present analysis.

Contents

1	Introduction	1
2	Distribution plots	2
3	p_T spectra of $\Lambda(1520)$	2
4	Conclusion	3
	References	4

1 Introduction

In collider experiment, people are trying to understand production/interaction of resonances in a hot dense medium because they have life-time comparable to that of hadronic phases [1]. Resonance particle $\Lambda(1520)$ having a life-time between $K^{*0\pm}(980)$ and $\phi(1020)$ make it a good candidate to understand the hadronic phase [2] [3]. Here we are analysing production of $\Lambda(1520)$ with respect to transverse sphericity S_0 using event-generator EPOS3 [4] [5] [6] with UrQMD as

afterburner.

Transverse sphericity is an event shape variable which help to distinguish isotropic and jetty event. Its value ranges form 0 to 1 and it is given by

$$S_0 = \frac{\pi^2}{4} \min_{\vec{n}=(n_x, n_y, 0)} \left(\frac{\sum_i |p_{T_i} \times \vec{n}|}{N_{particles>5}} \right)^2$$

Where,

$$S_0 = \begin{cases} 0, & \text{"pencil-like" limit (hard events)} \\ 1, & \text{"isotropic" limit (soft events)} \end{cases}$$

and \hat{n} is a two dimensional unit vector in the transverse plane, chosen in a way so that S_0 is minimised. By restricting it to the transverse plane, transverse sphericity becomes infrared and collinear safe. Also the jetty events are usually hard events while the isotropic ones are the result of soft processes [7] [8] [9] [10].

2 Distribution plots

For the present study, events having charge particles greater than 5 are considered. Moreover, to study in various multiplicity interval and transverse sphericity classes, First we took the quantile of multiplicity distribution in two regions 0-10%(high multiplicity) and 30-60%(low multiplicity). To disentangle the sphericity classes i.e jetty and isotropic events, we have applied sphericity cuts on sphericity distribution. The values, cuts and definition of various multiplicity percentile and sphericity classes can be found in Table 1a and 1b. We observed that the sphericity distributions are shifted toward 1 from low multiplicity percentile to high multiplicity. Figure 1 displays charged particle multiplicity, transverse sphericity and sphericity as a function of multiplicity.

No. of charged particle	Percentile (lower)	Percentile (upper)
28 - 110	90-100%	0-10%
18-28	70-90%	10-30%
11 - 18	40-70%	30-60%
0-110	0-100%	0-100%

(a) Percentile of multiplicity distribution

Multiplicity(%)	Jetty 0-10% (lower)	Isotropic 0-10% (upper)
0-100	0.00-0.381	0.834-1.00
0-10	0.00-0.644	0.896-1.00
30-60	0.00-0.442	0.824-1.00

(b) Quantile of sphericity distribution

Table 1

3 pT spectra of $\Lambda(1520)$

Figure 2(2a,2b) shows the pT-spectra of $\Lambda(1520)$ for 0-10% and 30-60% multiplicity classes with top and bottom 10% sphericity values responsible for selecting jetty and isotropic events respectively. The bottom panels show the ratio with respect to the sphericity integrated class.




Figure 1: Shows distribution of :- (a) Charge particle multiplicities. (b) Transverse sphericity. (c): Transverse sphericity vs multiplicity.




Figure 2: (a) p_T spectra of $\Lambda(1520)$ evaluated at 0-10% multiplicity for Isotropic(blue), Jetty (red) and Integrated(black). (b) p_T spectra of $\Lambda(1520)$ evaluated at 30-60% multiplicity for Isotropic(blue), Jetty (red) and Integrated(black).

4 Conclusion

A study of the transverse sphericity of $\Lambda(1520)$ in p-p collisions at $\sqrt{s} = 13\text{TeV}$ using EPOS3 generator with URQMD as afterburner was reported. The observables are measured using primary charge particles and reported as a function of charged particle multiplicity for events with different scales defined by the transverse sphericity. As it can be seen in Figure(1c), that jetty events are dominating in low multiplicities. When comparing 0-10% and 30-60% multiplicity class, but we

see that production of $\Lambda(1520)$ in isotropic event shape is dominating in both cases.

Acknowledgements

We thank Vikash Sumberia, Randhir Singh and Dr. Ranbir Singh for inspiring us to do this work and providing also valuable advice. Thanks are also due to Johannes Jahan Ph.D. student in Subatech, Nantes -France for helping us to understand technical term in EPOS3. This work was supported by Council of Scientific and Industrial Research (CSIR), India. The authors acknowledge the HEP Lab. Department of Physics, University of Jammu for providing infrasture for computing etc.

References

- [1] M. Bleicher and J. Aichelin, *Strange resonance production: probing chemical and thermal freeze-out in relativistic heavy ion collisions*, Physics Letters B **530**(1), 81 (2002), doi:[https://doi.org/10.1016/S0370-2693\(02\)01334-5](https://doi.org/10.1016/S0370-2693(02)01334-5).
- [2] A. G. Knospe, C. Markert, K. Werner, J. Steinheimer and M. Bleicher, *Hadronic resonance production and interaction in partonic and hadronic matter in the epos3 model with and without the hadronic afterburner urqmd*, Phys. Rev. C **93**, 014911 (2016), doi:[10.1103/PhysRevC.93.014911](https://doi.org/10.1103/PhysRevC.93.014911).
- [3] M. Tanabashi, K. Hagiwara, K. Hikasa, K. Nakamura, Y. Sumino, F. Takahashi, J. Tanaka, K. Agashe, G. Aielli, C. AMSLER, M. Antonelli, D. M. Asner *et al.*, *Review of particle physics*, Phys. Rev. D **98**, 030001 (2018), doi:[10.1103/PhysRevD.98.030001](https://doi.org/10.1103/PhysRevD.98.030001).
- [4] H. Drescher, M. Hladik, S. Ostapchenko, T. Pierog and K. Werner, *Parton-based Gribov-regge theory*, Physics Reports **350**(2), 93 (2001), doi:[https://doi.org/10.1016/S0370-1573\(00\)00122-8](https://doi.org/10.1016/S0370-1573(00)00122-8).
- [5] K. Werner, B. Guiot, I. Karpenko and T. Pierog, *Analyzing radial flow features in p-pb and p-p collisions at several tev by studying identified-particle production with the event generator epos3*, Phys. Rev. C **89**, 064903 (2014), doi:[10.1103/PhysRevC.89.064903](https://doi.org/10.1103/PhysRevC.89.064903).
- [6] K. Werner, *Core-corona separation in ultrarelativistic heavy ion collisions*, Phys. Rev. Lett. **98**, 152301 (2007), doi:[10.1103/PhysRevLett.98.152301](https://doi.org/10.1103/PhysRevLett.98.152301).
- [7] N. Mallick, R. Sahoo, S. Tripathy and A. Ortiz, *Study of transverse spherocity and azimuthal anisotropy in pb-pb collisions at sNN=5.02 TeV using a multi-phase transport model*, Journal of Physics G: Nuclear and Particle Physics **48**(4), 045104 (2021), doi:[10.1088/1361-6471/abeb59](https://doi.org/10.1088/1361-6471/abeb59).
- [8] A. Ortiz, E. Cuautle and G. Paić, *Mid-rapidity charged hadron transverse spherocity in pp collisions simulated with pythia*, Nuclear Physics A **941**, 78 (2015), doi:<https://doi.org/10.1016/j.nuclphysa.2015.05.010>.

- [9] S. Prasad, N. Mallick, D. Behera, R. Sahoo and S. Tripathy, *Event topology and global observables in heavy-ion collisions at the large hadron collider*, Scientific Reports **12**(1), 3917 (2022), doi:[10.1038/s41598-022-07547-z](https://doi.org/10.1038/s41598-022-07547-z).
- [10] ALICE Collaboration, S. Acharya, D. Adamová, S. P. Adhya, A. Adler, J. Adolfsson, M. M. Aggarwal, G. A. Rinella, M. Agnello, N. Agrawal, Z. Ahammed, S. Ahmad *et al.*, *Charged-particle production as a function of multiplicity and transverse sphericity in pp collisions at $\sqrt{s} = 5.02$ and 13 TeV*, The European Physical Journal C **79**(10), 857 (2019), doi:[10.1140/epjc/s10052-019-7350-y](https://doi.org/10.1140/epjc/s10052-019-7350-y).