

WELL-POSEDNESS OF A VISCOELASTIC RESISTIVE FORCE THEORY AND APPLICATIONS TO SWIMMING

LAUREL OHM

ABSTRACT. We propose and analyze a simple model for the evolution of an immersed, inextensible filament which incorporates linear viscoelastic effects of the surrounding fluid. The model is a closed-form system of equations along the curve only which includes a ‘memory’ term due to viscoelasticity. For a planar filament, given a forcing in the form of a preferred curvature, we prove well-posedness of the fiber evolution as well as the existence of a unique time-periodic solution in the case of time-periodic forcing. Moreover, we obtain an expression for the swimming speed of the filament in terms of the preferred curvature. The swimming speed depends in a complicated way on the viscoelastic parameters corresponding to the fluid relaxation time and additional polymeric viscosity. We study this expression in detail, accompanied by numerical simulations, and show that this simple model can capture complex effects of viscoelasticity on swimming. In particular, the viscoelastic swimmer is faster than its Newtonian counterpart in some situations and slower in others, although the differences are often quite small. Most strikingly, we can predict a situation where viscoelastic effects lead to a reversal in swimming direction from the Newtonian setting.

CONTENTS

1	Introduction	1
1.1	The model	3
1.2	Analytical setup	5
1.3	Statement of results	7
2	Applications and numerical results	10
3	Well-posedness and periodic solutions	16
3.1	Semigroup properties	16
3.2	Tension equation	19
3.3	Evolution equation	21
3.4	Existence of a unique periodic solution	24
3.5	Small relaxation time limit	27
4	Viscoelastic swimming	29
	Appendix A Numerical method	32
	References	34

1. INTRODUCTION

The effect of fluid viscoelasticity on swimming microorganisms is a subject of great interest in the biofluids community. A major focus of studies on swimming in viscoelastic media is on the complexity of the impacts that viscoelasticity can have on swimming speeds, depending on the situation. In experimental settings and in computational models, viscoelastic effects have been shown to hinder swimming [34], enhance swimming [35, 13, 4, 31, 32], or both hinder and enhance swimming depending on factors such as the kinematics of the swimmer [9, 3, 1, 37, 19]. Many studies emphasize the non-monotonic dependence of swimming speed on parameters relating to fluid viscoelasticity [21, 36, 20, 38, 33]. Much of this prior work is either experimental, e.g. [34], or

based on computational models which couple an equation for curve evolution with bulk viscoelastic fluid equations (such as Oldroyd-B) via, e.g. the immersed boundary method [17, 18, 37, 38].

Here we present a simple model for the evolution of an immersed, inextensible curve which incorporates linear viscoelastic effects of the surrounding fluid in a closed-form system of equations along the curve only. The model is derived from the linear viscoelastic resistive force theory described in [6, 38], but requires some additional interpretation to yield a well-posed curve evolution. The resulting fiber evolution equations look like classical resistive force theory plus Euler beam theory [11, 2, 12, 39], but incorporate the evolution of an additional variable corresponding to a memory of the fiber curvature at previous times. This system of PDEs satisfies a very natural energy identity: in the absence of forcing, the bending energy of the filament plus an energy corresponding to the memory term are non-increasing in time. We prescribe a time-periodic forcing along the filament in the form of a preferred curvature and consider swimming as an emergent property. We show that this simple model, which is not coupled to any equations in the bulk, can capture the complexity of viscoelastic effects on swimming, including slowdowns and speedups relative to a Newtonian swimmer, depending on the form of the preferred curvature and the size of two viscoelastic parameters.

Although the viscoelastic effect is linear, the system of PDEs we obtain is nonlinear due to the fiber inextensibility constraint. For a planar filament, we prove well-posedness of the fiber evolution problem, including global existence and uniqueness for small data and local existence and uniqueness for large data. The model inherits many of the features of the Newtonian problem, studied in detail from a PDE perspective in our previous work [26], but is fundamentally different in that the analysis now includes an additional ODE for the memory variable. As in [26], we prove that given a (small) time-periodic forcing along the fiber in the form of a preferred curvature, there exists a unique periodic solution to the filament evolution equations. Furthermore, we show that the periodic solution to the viscoelastic PDE converges to the unique periodic solution in the Newtonian setting as a parameter corresponding to the fluid relaxation time goes to zero. These analysis questions are interesting in their own right and continue to develop the PDE theory of the hydrodynamics of slender filaments initiated in [27, 28, 24, 25, 29].

Finally, we calculate an expression for the fiber swimming speed in terms of the prescribed preferred curvature. The expression depends in a complicated way on the viscoelastic parameters corresponding to the fluid relaxation time and the additional (polymer) viscosity of the fluid. Nevertheless, we are able to make a few predictions about the swimming speed, which we test via numerical simulations. The numerical method that we use is a natural extension of the method we proposed in [26], which is based on a combination of the methods used in [23, 22]. We show that varying two viscoelastic parameters corresponding to the fluid relaxation time and the additional polymeric stress of the fluid can have complex effects on the fiber swimming speed. Most strikingly, we find a scenario in which viscoelastic effects cause the swimmer to reverse direction. Our results are for a small set of parameter values and two choices of time-periodic preferred curvature, meaning that much remains to be explored.

We note that prior asymptotic calculations have indicated that linear viscoelasticity does not affect the swimming speed of a filament to leading order in small amplitude deformations, and nonlinear viscoelastic effects are needed to see changes in the swimming speed from a Newtonian swimmer [5, 7, 8, 15]. These results rely on an expression for the swimming speed in terms of the actual fiber shape rather than a given forcing. Here we show that prescribing the same preferred curvature along the filament results in differences in the emerging shape and hence in the swimming speed.

We also note that our analysis accounts for effects of boundary conditions on finite fibers and applies to fibers with small curvatures rather than small amplitude deformations.

We further note that the numerical results reported here are qualitative rather than quantitative in the sense that the scaling of the model as presented here is not physically realistic. In particular, our choice of timescale for the fiber evolution removes the dependence of the dynamics on the swimmer slenderness and its bending stiffness. This makes the analysis and comparison to the Newtonian setting in [26] more convenient, but makes direct comparison with experiments or biological models less convenient. These physical considerations are expected to play a role: indeed, the important role of fiber flexibility on swimming has been emphasized in, e.g., [33, 38]. From an applications perspective, our main aim is to show that we can indeed get complex swimming behaviors from the simple model presented here. A more physical rescaling of this model may even amplify these differences.

1.1. The model. Let $\mathbf{X} : [0, 1] \times [0, T] \rightarrow \mathbb{R}^3$ denote the centerline of an inextensible elastic filament. Throughout, we will use the notation $I = [0, 1]$ to denote the unit interval, $s \in I$ to denote the arclength parameter along \mathbf{X} , and subscript $(\cdot)_s$ to denote $\frac{\partial}{\partial s}$. At each $s \in I$, the unit tangent vector to \mathbf{X} is given by $\mathbf{e}_t(s, t) = \mathbf{X}_s / |\mathbf{X}_s| = \mathbf{X}_s$ due to inextensibility. Here we will consider a fiber undergoing planar deformations only; in particular, at each point $s \in I$ we may define an in-plane unit normal vector $\mathbf{e}_n \perp \mathbf{e}_t$ to the filament.

The motion of the filament is driven by a prescribed active forcing in the form of a preferred curvature $\kappa_0(s, t)$ along the fiber. Given κ_0 , the filament evolves according to

$$\frac{\partial \mathbf{X}}{\partial t} = -(1 + \mu)(\mathbf{I} + \gamma \mathbf{X}_s \mathbf{X}_s^T)(\mathbf{X}_{sss} - \tau \mathbf{X}_s - (\kappa_0)_s \mathbf{e}_n - \frac{\mu}{1 + \mu} \xi_s \mathbf{e}_n)_s \quad (1)$$

$$\delta \frac{\partial \xi}{\partial t} = -\xi + \kappa - \kappa_0 \quad (2)$$

$$|\mathbf{X}_s|^2 = 1, \quad (3)$$

with boundary conditions

$$(\mathbf{X}_{ss} - \kappa_0 \mathbf{e}_n)|_{s=0,1} = 0, \quad (\mathbf{X}_{sss} - \tau \mathbf{X}_s - (\kappa_0)_s \mathbf{e}_n)|_{s=0,1} = 0, \quad \xi|_{s=0,1} = \xi_s|_{s=0,1} = 0. \quad (4)$$

Here the matrix $(\mathbf{I} + \gamma \mathbf{X}_s \mathbf{X}_s^T)$ in (1) is the resistive force theory approximation relating the hydrodynamic force along a slender filament in a Stokes (Newtonian) fluid to its velocity [11, 30, 16]. The parameter γ is a shape factor which depends on the aspect ratio of the filament; for a very slender filament, $\gamma \approx 1$.

The first three components $(\mathbf{X}_{sss} - \tau \mathbf{X}_s - (\kappa_0)_s \mathbf{e}_n)_s$ of the forcing term in (1) are identical to the Newtonian setting. The term \mathbf{X}_{sss} is the elastic response of the filament to deformations and may be rewritten as $(\kappa_s \mathbf{e}_n - \kappa^2 \mathbf{e}_t)_s$, where $\kappa(s, t)$ is the filament curvature. The function $\tau(s, t)$ plays the role of the (unknown) filament tension and enforces the inextensibility constraint (3). As noted, κ_0 is the prescribed preferred curvature of the filament and serves as an active forcing along the fiber.

The leading order effects of the filament response to a viscoelastic fluid are encoded in the variable ξ . As seen from equation (1), ξ modifies the evolution of \mathbf{X} in the same way as the preferred curvature κ_0 , and, from equation (2), may be interpreted as the “memory” of the curvature difference $\kappa - \kappa_0$ at previous times. The parameters $\delta \geq 0$ and $\mu \geq 0$ are associated with viscoelasticity: δ is the additional relaxation time of the filament due to viscoelastic effects of the fluid, and μ relates to the additional (polymer) viscosity of the viscoelastic medium. Note that if $\mu = 0$ or if $\delta = 0$, we

recover the classical Newtonian formulation (see [2, 12, 40, 41, 39, 26])

$$\frac{\partial \mathbf{X}}{\partial t} = -(\mathbf{I} + \gamma \mathbf{X}_s \mathbf{X}_s^T)(\mathbf{X}_{sss} - \tau \mathbf{X}_s - (\kappa_0)_s \mathbf{e}_n)_s \quad (5)$$

(up to a rescaling of the unknown tension in the $\delta = 0$ case). However, showing convergence of solutions of (1)-(2) to solutions of (5) as $\delta \rightarrow 0$ is more subtle than simply verifying that $\delta = 0$ yields the Newtonian formulation, since $\delta > 0$ is a singular perturbation. We address the $\delta \rightarrow 0$ limit in the time-periodic setting in Theorem 1.2.

The model (1)-(4) is inspired by the following (perhaps more familiar) framing of linear viscoelastic effects on filament evolution. We consider

$$\frac{\partial \mathbf{X}}{\partial t} = -(\mathbf{I} + \gamma \mathbf{X}_s \mathbf{X}_s^T)(\boldsymbol{\sigma}^{\text{vis}})_s \quad (6)$$

$$(1 + \mu)\delta \frac{\partial \boldsymbol{\sigma}^{\text{ve}}}{\partial t} + \boldsymbol{\sigma}^{\text{ve}} = \delta \frac{\partial \boldsymbol{\sigma}^{\text{vis}}}{\partial t} + \boldsymbol{\sigma}^{\text{vis}} \quad (7)$$

$$(\boldsymbol{\sigma}^{\text{ve}})_s = (\mathbf{X}_{sss} - \tau \mathbf{X}_s - (\kappa_0)_s \mathbf{e}_n)_s \quad (8)$$

$$|\mathbf{X}_s|^2 = 1. \quad (9)$$

Here $\boldsymbol{\sigma}^{\text{ve}}(s, t)$ and $\boldsymbol{\sigma}^{\text{vis}}(s, t)$ are both vectors along the filament \mathbf{X} . Equation (6) relates the filament velocity $\frac{\partial \mathbf{X}}{\partial t}$ to the viscous drag $\mathbf{f}^{\text{vis}} = (\boldsymbol{\sigma}^{\text{vis}})_s$ along the fiber via resistive force theory. Equation (7) has the form of a linearized Oldroyd-B model [6, 38], which relates the viscoelastic stresses in the fluid to the viscous strain rate. Equation (7) is restricted to the filament only; $\boldsymbol{\sigma}^{\text{ve}}(s, t)$ and $\boldsymbol{\sigma}^{\text{vis}}(s, t)$ are both vectors along the fiber. When s and t derivatives commute (i.e. for a straight filament), equation (7) agrees with the linear viscoelastic resistive force theory derived in [6, 38] in terms of \mathbf{f}^{ve} and \mathbf{f}^{vis} instead. Here again δ is the fluid relaxation time and $1 + \mu$ is the total viscosity of the medium. Note that rescaling $\boldsymbol{\sigma}^{\text{vis}}$ by $\frac{1}{1+\mu}$ and δ by $1 + \mu$, we may rewrite (7) in the (perhaps more usual) form $\delta \frac{\partial \boldsymbol{\sigma}^{\text{ve}}}{\partial t} + \boldsymbol{\sigma}^{\text{ve}} = \delta \frac{\partial \boldsymbol{\sigma}^{\text{vis}}}{\partial t} + (1 + \mu)\boldsymbol{\sigma}^{\text{vis}}$, but we will use the form (7) for analysis.

Equation (8) is the force balance between the viscoelastic forces in the fluid $\mathbf{f}^{\text{ve}} = (\boldsymbol{\sigma}^{\text{ve}})_s$ and the elastic forces $(\mathbf{X}_{sss} - \tau \mathbf{X}_s - (\kappa_0)_s \mathbf{e}_n)_s$ along the rod, which are subject to the inextensibility constraint (9). See [2, 38, 26] for a variational derivation of the elastic forces along the fiber; note that the boundary conditions (4) come from this variational derivation.

Let $\sigma_n^{\text{vis}} = \boldsymbol{\sigma}^{\text{vis}} \cdot \mathbf{e}_n$, $\sigma_t^{\text{vis}} = \boldsymbol{\sigma}^{\text{vis}} \cdot \mathbf{e}_t$, and likewise for σ_n^{ve} , σ_t^{ve} . Now, because the filament is inextensible, the stresses in the tangential direction along the fiber are unknown and are grouped into the filament tension (see (8)). Up to a redefinition of the (unknown) filament tension, we will use the same form of stress in the tangential direction as in the Newtonian setting, i.e. $\sigma_t^{\text{vis}} = \tau \mathbf{X}_s$ for some unknown function τ . We will thus consider equation (7) as an equation holding along the normal direction of the fiber only:

$$(1 + \mu)\delta \frac{\partial \sigma_n^{\text{ve}}}{\partial t} + \sigma_n^{\text{ve}} = \delta \frac{\partial \sigma_n^{\text{vis}}}{\partial t} + \sigma_n^{\text{vis}}.$$

We can then solve for σ_n^{vis} in terms of σ_n^{ve} :

$$\begin{aligned} \sigma_n^{\text{vis}} &= (1 + \mu)\sigma_n^{\text{ve}} - \mu\delta^{-1} \int_0^t e^{-(t-t')/\delta} \sigma_n^{\text{ve}}|_{t=t'} dt' + e^{-t/\delta} (\sigma_n^{\text{vis}, \text{in}} - (1 + \mu)\sigma_n^{\text{ve}, \text{in}}) \\ &= (1 + \mu)\sigma_n^{\text{ve}} - \mu\delta^{-1} \int_0^t e^{-(t-t')/\delta} (\kappa - \kappa_0)_s dt' + e^{-t/\delta} (\sigma_n^{\text{vis}, \text{in}} - (1 + \mu)\sigma_n^{\text{ve}, \text{in}}), \end{aligned}$$

where $\sigma_n^{\text{vis},\text{in}} = \sigma_n^{\text{vis}}|_{t=0}$, $\sigma_n^{\text{ve},\text{in}} = \sigma_n^{\text{ve}}|_{t=0}$. Then $\xi = \delta^{-1} \int_0^t e^{-(t-t')/\delta} (\kappa - \kappa_0)_s dt' + \mu^{-1} e^{-t/\delta} (\sigma_n^{\text{vis},\text{in}} - (1 + \mu)\sigma_n^{\text{ve},\text{in}})$ satisfies $\delta \dot{\xi} = -\xi + \kappa - \kappa_0$, yielding the system (1)-(4).

The model (1)-(4) has advantages over other potential ways of incorporating linear viscoelastic effects of the surrounding fluid due to both its simplicity and because it has an associated energy. In particular, taking $\kappa_0 = 0$ (no internal forcing), equations (1) and (2) reduce to

$$\begin{aligned} \frac{\partial \mathbf{X}}{\partial t} &= -(1 + \mu)(\mathbf{I} + \gamma \mathbf{X}_s \mathbf{X}_s^T)(\mathbf{X}_{sss} - \tau \mathbf{X}_s - \frac{\mu}{1 + \mu} \xi_s \mathbf{e}_n)_s \\ \delta \frac{\partial \xi}{\partial t} &= -\xi + \kappa. \end{aligned}$$

Multiplying both sides of the first equation by $(\mathbf{X}_{sss} - \tau \mathbf{X}_s - \frac{\mu}{1 + \mu} \xi_s \mathbf{e}_n)_s$ and integrating in s , on the right hand side we obtain the negative quantity

$$-R^2(t) := -(1 + \mu) \int_0^1 \left(\left| (\mathbf{X}_{sss} - \tau \mathbf{X}_s - \frac{\mu}{1 + \mu} \xi_s \mathbf{e}_n)_s \right|^2 + \gamma \left(\mathbf{X}_s \cdot (\mathbf{X}_{sss} - \tau \mathbf{X}_s - \frac{\mu}{1 + \mu} \xi_s \mathbf{e}_n)_s \right)^2 \right) ds.$$

On the left hand side, using that $\mathbf{X}_s = \mathbf{e}_t$ and $\mathbf{X}_{ss} = \kappa \mathbf{e}_n$, we have

$$\begin{aligned} \int_0^1 \frac{\partial \mathbf{X}}{\partial t} \cdot (\mathbf{X}_{sss} - \tau \mathbf{X}_s - \frac{\mu}{1 + \mu} \xi_s \mathbf{e}_n)_s ds &= - \int_0^1 \frac{\partial \mathbf{X}_s}{\partial t} \cdot (\mathbf{X}_{sss} - \tau \mathbf{X}_s - \frac{\mu}{1 + \mu} \xi_s \mathbf{e}_n) ds \\ &= \int_0^1 \dot{\kappa} \left(\kappa - \frac{\mu}{1 + \mu} \xi \right) ds, \end{aligned}$$

where we use $\dot{\kappa}$ to denote $\frac{\partial \kappa}{\partial t}$. Now, $\dot{\kappa} \xi$ may be rewritten as

$$\dot{\kappa} \xi = \partial_t(\kappa \xi) - \kappa \dot{\xi} = \partial_t(\kappa \xi) - \dot{\xi}(\xi + \delta \dot{\xi}) = \partial_t(\kappa \xi) - \frac{1}{2} \partial_t(\xi^2) - \delta \dot{\xi}^2,$$

so the left hand side becomes

$$\int_0^1 \dot{\kappa} \left(\kappa - \frac{\mu}{1 + \mu} \xi \right) ds = \int_0^1 \left(\frac{1}{2} \partial_t(\kappa^2) - \frac{\mu}{1 + \mu} (\partial_t(\kappa \xi) - \frac{1}{2} \partial_t(\xi^2) - \delta \dot{\xi}^2) \right) ds.$$

We thus obtain the energy equality

$$\frac{1}{2} \partial_t \int_0^1 \left(\kappa^2 + \mu(\kappa - \xi)^2 \right) ds = -\delta \mu \int_0^1 \dot{\xi}^2 ds - (1 + \mu) R^2(t); \quad (10)$$

in particular, $\|\kappa\|_{L^2}^2 + \mu \|\kappa - \xi\|_{L^2}^2$ is a monotone quantity. Notice that $\kappa - \xi = \delta \dot{\xi}$, so this quantity may be rewritten as $\|\kappa\|_{L^2}^2 + \mu \delta \|\dot{\xi}\|_{L^2}^2$.

1.2. Analytical setup. Rather than working directly with the formulation (13)-(16), we will use the inextensibility of the filament and the planarity of its deformation to write the tangent vector along the filament as

$$\mathbf{X}_s = \mathbf{e}_t = \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix}, \quad (11)$$

where $\theta(s, t)$ is the angle between $\mathbf{e}_t(s, t)$ and $\mathbf{e}_t(0, 0)$. Differentiating (1) in s , we may then obtain a system of three equations: two evolution equations for θ and ξ , and one elliptic equation for the

tension τ , given by

$$\begin{aligned}
\dot{\theta} &= (1 + \mu) \left(-\theta_{ssss} + (2 + \gamma)(\theta_s^3)_s + (2 + \gamma)\tau_s\theta_s + \tau\theta_{ss} + (\kappa_0)_{sss} \right. \\
&\quad \left. - (1 + \gamma)\theta_s^2(\kappa_0)_s \right) + \mu \left(\xi_{sss} - (1 + \gamma)\theta_s^2\xi_s \right) \\
\delta\dot{\xi} &= -\xi + \theta_s - \kappa_0 \\
(1 + \gamma)\tau_{ss} &= (\theta_s)^2\tau + (\theta_s)^4 + \theta_{ss}^2 - (4 + 3\gamma)(\theta_{ss}\theta_s)_s + (2 + \gamma)(\kappa_0)_{ss}\theta_s + (1 + \gamma)\theta_{ss}(\kappa_0)_s \\
&\quad + \frac{\mu}{1 + \mu} \left((2 + \gamma)(\theta_s\xi_s)_s - \theta_{ss}\xi_s \right) \\
(\theta_s - \kappa_0)|_{s=0,1} &= 0, \quad (\theta_{ss} - (\kappa_0)_s)|_{s=0,1} = 0, \quad (\tau + \kappa_0^2)|_{s=0,1} = 0 \quad \xi|_{s=0,1} = \xi_s|_{s=0,1} = 0.
\end{aligned} \tag{12}$$

However, it will be more useful to consider the filament evolution in term of the fiber curvature $\kappa = \theta_s$ rather than θ , as is done in [10, 38, 26]. Furthermore, due to the boundary conditions in (12), it will be most useful to recast the system (12) in terms of $\bar{\kappa} = \kappa - \kappa_0$ and $\bar{\tau} = \tau + \kappa_0^2$. The equations (12) may be written as

$$\dot{\bar{\kappa}} = -(1 + \mu)\bar{\kappa}_{ssss} + \mu\xi_{ssss} - \dot{\kappa}_0 + (1 + \mu)(\mathcal{N}[\bar{\kappa}, \kappa_0])_s - \mu(1 + \gamma)((\bar{\kappa} + \kappa_0)^2\xi_s)_s \tag{13}$$

$$\delta\dot{\xi} = \bar{\kappa} - \xi \tag{14}$$

$$(1 + \gamma)\bar{\tau}_{ss} = (\bar{\kappa} + \kappa_0)^2\bar{\tau} + \mathcal{T}[\bar{\kappa}, \kappa_0] + \frac{\mu}{1 + \mu} \left((2 + \gamma)((\bar{\kappa} + \kappa_0)\xi_s)_s - (\bar{\kappa} + \kappa_0)_s\xi_s \right) \tag{15}$$

$$\bar{\kappa}|_{s=0,1} = \bar{\kappa}_s|_{s=0,1} = 0, \quad \xi|_{s=0,1} = \xi_s|_{s=0,1} = 0, \quad \bar{\tau}|_{s=0,1} = 0. \tag{16}$$

This may be compared with the curvature formulation in the Newtonian case [26], where the evolution is given by

$$\begin{aligned}
\dot{\bar{\kappa}}^{\text{nw}} &= -\bar{\kappa}_{ssss}^{\text{nw}} - \dot{\kappa}_0 + (\mathcal{N}[\bar{\kappa}^{\text{nw}}, \kappa_0])_s \\
(1 + \gamma)\bar{\tau}_{ss}^{\text{nw}} &= (\bar{\kappa}^{\text{nw}} + \kappa_0)^2\bar{\tau}^{\text{nw}} + \mathcal{T}[\bar{\kappa}^{\text{nw}}, \kappa_0].
\end{aligned} \tag{17}$$

Here we use the superscript $(\cdot)^{\text{nw}}$ to distinguish the solution to the Newtonian PDE (17) from the viscoelastic $\bar{\kappa}$. The nonlinear terms \mathcal{N} and \mathcal{T} have the same form in both the viscoelastic and Newtonian cases and are given by

$$\begin{aligned}
\mathcal{N}[\bar{\kappa}, \kappa_0] &:= 3(2 + \gamma)\bar{\kappa}(\bar{\kappa} + 2\kappa_0)\bar{\kappa}_s + (5 + 3\gamma)\kappa_0^2\bar{\kappa}_s + (5 + 2\gamma)\bar{\kappa}^2(\kappa_0)_s \\
&\quad + 2(3 + \gamma)\bar{\kappa}\kappa_0(\kappa_0)_s + (2 + \gamma)\bar{\tau}_s(\bar{\kappa} + \kappa_0) + \bar{\tau}(\bar{\kappa} + \kappa_0)_s \\
\mathcal{T}[\bar{\kappa}, \kappa_0] &:= \bar{\kappa}(\bar{\kappa} + \kappa_0)^2(\bar{\kappa} + 2\kappa_0) + (\bar{\kappa} + \kappa_0)_s\bar{\kappa}_s \\
&\quad - (1 + \gamma)(\bar{\kappa}(\bar{\kappa} + 2\kappa_0))_{ss} - (2 + \gamma)(\bar{\kappa}_s(\bar{\kappa} + \kappa_0))_s.
\end{aligned}$$

Note that $\bar{\tau}$ appears in \mathcal{N} , but since $\bar{\tau} = \bar{\tau}(\bar{\kappa}, \kappa_0)$, we will not denote this $\bar{\tau}$ dependence in our notation. The formulation (13)-(16) will serve as the basis for our analysis.

Given $\theta|_{t=0}$ and $(\bar{\kappa}, \xi)$ solving (18), we may recover $\theta(s, t)$ via

$$\dot{\theta} = -(1 + \mu)\bar{\kappa}_{ss} - \dot{\kappa}_0 + (1 + \mu)\mathcal{N}[\bar{\kappa}, \kappa_0] + \mu\xi_{ss} - \mu(1 + \gamma)(\bar{\kappa} + \kappa_0)^2\xi_s, \tag{18}$$

and the evolution of the fiber frame $(\mathbf{e}_t, \mathbf{e}_n)$ via

$$\dot{\mathbf{e}}_t(s, t) = \dot{\theta}(s, t)\mathbf{e}_n(s, t), \quad \dot{\mathbf{e}}_n(s, t) = -\dot{\theta}(s, t)\mathbf{e}_t(s, t). \tag{19}$$

Using (19), we may then obtain the full fiber evolution by

$$\begin{aligned} \frac{\partial \mathbf{X}}{\partial t}(s, t) = & \left(-(1 + \mu)\bar{\kappa}_{ss} + \mu\xi_{ss} + (1 + \mu)(\bar{\kappa} + \kappa_0)(\bar{\tau} - \bar{\kappa}(\bar{\kappa} + 2\kappa_0)) \right) \mathbf{e}_n \\ & + (1 + \gamma) \left((\bar{\kappa} + \kappa_0)((1 + \mu)\bar{\kappa}_s - \mu\xi_s) + (1 + \mu)(\bar{\tau}_s - (\bar{\kappa}(\bar{\kappa} + 2\kappa_0))_s) \right) \mathbf{e}_t. \end{aligned} \quad (20)$$

To analyze the system (13)-(16), as in [26], we will begin by defining the linear operator \mathcal{L} by

$$\mathcal{L}[\psi] := \partial_{ssss}\psi, \quad \psi(0) = \psi(1) = 0, \quad \psi_s(0) = \psi_s(1) = 0. \quad (21)$$

From [14, 40, 41], we have that the eigenfunctions ψ_k and eigenvalues λ_k of the operator \mathcal{L} (21) are given by

$$\begin{aligned} \psi_k(s) &= \frac{\widehat{\psi}_k(s)}{\|\widehat{\psi}_k\|_{L^2}}, \quad \lambda_k = \alpha_k^4, \quad k = 1, 2, \dots \\ \text{where } \cos(\alpha_k) \cosh(\alpha_k) &= 1, \quad \alpha_0 = 0 \\ \text{and } \widehat{\psi}_k(s) &= (\cos(\alpha_k) - \cosh(\alpha_k))(\cos(\alpha_k s) - \cosh(\alpha_k s)) \\ &\quad + (\sin(\alpha_k) + \sinh(\alpha_k))(\sin(\alpha_k s) - \sinh(\alpha_k s)). \end{aligned} \quad (22)$$

We note that $\alpha_k \rightarrow \frac{(2k+1)\pi}{2}$ as $k \rightarrow \infty$, and that the smallest eigenvalue of \mathcal{L} is given by $\lambda_1 \approx (4.73)^4 \approx 500$. We further note that $\psi_k(s)$ is even about $s = \frac{1}{2}$ for odd k , and odd about $s = \frac{1}{2}$ for even k .

We may consider the expansion of any $u \in L^2(I)$ in eigenfunctions of \mathcal{L} :

$$u = \sum_{k=1}^{\infty} \tilde{u}_k \psi_k, \quad \tilde{u}_k = \int_0^1 u(s) \psi_k(s) ds. \quad (23)$$

The domain of \mathcal{L}^r , $0 \leq r \leq 1$, may then be defined by

$$D(\mathcal{L}^r) = \left\{ u \in L^2(I) : \sum_{k=1}^{\infty} \lambda_k^{2r} \tilde{u}_k^2 < \infty \right\}. \quad (24)$$

Note that $D(\mathcal{L}^r) \subseteq H^{4r}(I)$ for $0 \leq r \leq 1$, and $D(\mathcal{L}^0) = L^2(I)$.

With \mathcal{L} as defined in (21), we may define a mild solution $(\bar{\kappa}, \xi)$ to the system (13)-(16) by the Duhamel formula

$$\begin{aligned} \begin{pmatrix} \bar{\kappa} \\ \xi \end{pmatrix} &= e^{\mathcal{A}t} \begin{pmatrix} \bar{\kappa}^{\text{in}} \\ \xi^{\text{in}} \end{pmatrix} - \int_0^t e^{\mathcal{A}(t-t')} \begin{pmatrix} \dot{\kappa}_0 \\ 0 \end{pmatrix} dt' + (1 + \mu) \int_0^t e^{\mathcal{A}(t-t')} \begin{pmatrix} (\mathcal{N}[\bar{\kappa}, \kappa_0])_s \\ 0 \end{pmatrix} dt' \\ &\quad - \mu(1 + \gamma) \int_0^t e^{\mathcal{A}(t-t')} \begin{pmatrix} ((\bar{\kappa} + \kappa_0)^2 \xi_s)_s \\ 0 \end{pmatrix} dt', \end{aligned} \quad (25)$$

where \mathcal{A} denotes the operator

$$\mathcal{A} = \begin{pmatrix} -(1 + \mu)\mathcal{L} & \mu\mathcal{L} \\ \delta^{-1} & -\delta^{-1} \end{pmatrix}. \quad (26)$$

1.3. Statement of results. Our first result is well-posedness for the system (13)-(16) in the case of small κ_0 and either short time or small initial data $\bar{\kappa}^{\text{in}}$. We note that the viscoelastic system inherits all of the subtleties of the Newtonian case (17) that make global well-posedness difficult for large initial data. In particular, the behavior of the filament tension $\bar{\tau}$, particularly its dependence on powers of $\bar{\kappa}_s$, limits what we can show in terms of well-posedness. See [26] for a deeper discussion of these issues in the Newtonian setting.

Here and throughout, we use the notation

$$\left\| \begin{pmatrix} u \\ \phi \end{pmatrix} \right\|_{\dot{H}^m \times \dot{H}^m} := \|u\|_{\dot{H}^m} + \|\phi\|_{\dot{H}^m}.$$

The well-posedness results for the system (13)-(16) may be stated as follows.

Theorem 1.1 (Well-posedness). *There exist constants $\varepsilon > 0, \varepsilon_1 \geq 0, \varepsilon_2 \geq 0$ such that, given $\kappa_0 \in C^1([0, T]; H^1(I))$ satisfying*

$$\sup_{t \in [0, T]} \|\kappa_0\|_{H^1(I)} = \varepsilon_1 \leq \varepsilon, \quad \sup_{t \in [0, T]} \|\dot{\kappa}_0\|_{L^2(I)} = \varepsilon_2 \leq \varepsilon,$$

there exist

- (1) *A time $T_\varepsilon > 0$ depending on $(\bar{\kappa}^{\text{in}}, \xi^{\text{in}})$ such that the system (13)-(16) admits a unique mild solution $(\bar{\kappa}, \xi) \in C([0, T_\varepsilon]; L^2(I) \times L^2(I)) \cap C((0, T_\varepsilon]; \dot{H}^1(I) \times \dot{H}^1(I))$.*
- (2) *A constant $\varepsilon_3 > 0$ such that if*

$$\left\| \begin{pmatrix} \bar{\kappa}^{\text{in}} \\ \xi^{\text{in}} \end{pmatrix} \right\|_{L^2(I) \times L^2(I)} = \varepsilon_3 \leq \varepsilon,$$

then, for any $T > 0$, the system (13)-(16) admits a unique mild solution $(\bar{\kappa}, \xi) \in C([0, T]; L^2(I) \times L^2(I)) \cap C((0, T]; \dot{H}^1(I) \times \dot{H}^1(I))$ satisfying

$$\sup_{t \in [0, T]} \left(\left\| \begin{pmatrix} \bar{\kappa} \\ \xi \end{pmatrix} \right\|_{L^2 \times L^2} + \min\{t^{1/4}, 1\} \left\| \begin{pmatrix} \bar{\kappa} \\ \xi \end{pmatrix} \right\|_{\dot{H}^1 \times \dot{H}^1} \right) \leq c(\varepsilon_1 + \varepsilon_2 + \varepsilon_3). \quad (27)$$

In case (2), in the absence of an internal forcing ($\kappa_0 \equiv 0$), we may obtain the bound

$$\left\| \begin{pmatrix} \kappa \\ \xi \end{pmatrix} \right\|_{L^2(I) \times L^2(I)} + \min\{t^{1/4}, 1\} \left\| \begin{pmatrix} \kappa \\ \xi \end{pmatrix} \right\|_{\dot{H}^1(I) \times \dot{H}^1(I)} \leq c e^{-t\Lambda} \left\| \begin{pmatrix} \kappa^{\text{in}} \\ \xi^{\text{in}} \end{pmatrix} \right\|_{L^2(I)}, \quad (28)$$

where $\Lambda = \min\{\lambda_1, \frac{1}{\delta(1+\mu)}\}$ for λ_1 as in (22).

As in the Newtonian case, in the absence of internal fiber forcing, the straight filament is nonlinearly stable to small perturbations. However, the fiber relaxation time Λ in (28) depends on the viscoelastic parameters δ and μ . In particular, if either δ or μ is large (even if, e.g., $\delta = \mu = 1$), the decay rate of perturbations to the straight filament is much slower than in the Newtonian setting (where $\Lambda = \lambda_1 \approx 500$). Finally, similar to the Newtonian setting, the quantity $\|(\bar{\kappa}^{\text{in}}, \xi^{\text{in}})\|_{L^2(I) \times L^2(I)}$ in case (2) corresponds to the initial viscoelastic bending energy (10) of the filament; in particular, a small initial energy leads to global existence.

For applications to undulatory swimming, we are most interested in prescribing a time-periodic preferred curvature κ_0 and understanding properties of the resulting time-periodic solution. Given a T -periodic κ_0 , we prove the existence of a unique T -periodic solution $(\bar{\kappa}, \xi)$ to (13)-(16). Moreover, we show that since the prescribed κ_0 is small, the unique periodic solution $(\bar{\kappa}, \xi)$ is close to the solution to the linearized version of (13)-(16). This will be useful for computing an expression for the fiber swimming speed. Finally, we show that as the viscoelastic relaxation time $\delta \rightarrow 0$, the unique periodic solution converges to the unique periodic solution of the Newtonian PDE (17), studied in detail in [26]. Recall that $\delta > 0$ is a singular perturbation of the $\delta = 0$ case (see (14)).

Theorem 1.2 (Periodic solutions and properties). *There exists a constant $\varepsilon > 0$ such that, given a T -periodic $\kappa_0 \in C^1([0, T]; H^1(I))$ satisfying*

$$\sup_{t \in [0, T]} \|\dot{\kappa}_0\|_{L^2(I)} = \varepsilon_1 \leq \varepsilon, \quad \sup_{t \in [0, T]} \|\kappa_0\|_{H^1(I)} = \varepsilon_2 \leq \varepsilon, \quad (29)$$

(a) *There exists a unique T -periodic solution $(\bar{\kappa}, \xi)$ to the system (13)-(16) satisfying*

$$\sup_{t \in [0, T]} \left\| \begin{pmatrix} \bar{\kappa} \\ \xi \end{pmatrix} \right\|_{H^1(I) \times H^1(I)} \leq c(\varepsilon_1 + \varepsilon_2). \quad (30)$$

(b) *Defining $(\bar{\kappa}^{\text{lin}}, \xi^{\text{lin}})$ to be the unique periodic solution to the linear PDE*

$$\begin{aligned} \dot{\bar{\kappa}}^{\text{lin}} &= -(1 + \mu)\bar{\kappa}_{ssss}^{\text{lin}} + \mu\xi_{ssss}^{\text{lin}} - \dot{\kappa}_0 \\ \delta \dot{\xi}^{\text{lin}} &= -\xi^{\text{lin}} + \bar{\kappa}^{\text{lin}} \\ \bar{\kappa}^{\text{lin}}|_{s=0,1} &= \xi^{\text{lin}}|_{s=0,1} = 0, \quad \bar{\kappa}_s^{\text{lin}}|_{s=0,1} = \xi_s^{\text{lin}}|_{s=0,1} = 0, \end{aligned} \quad (31)$$

the periodic solution $(\bar{\kappa}, \xi)$ of (30) satisfies

$$\sup_{t \in [0, T]} \left\| \begin{pmatrix} \bar{\kappa} - \bar{\kappa}^{\text{lin}} \\ \xi - \xi^{\text{lin}} \end{pmatrix} \right\|_{H^1(I) \times H^1(I)} \leq c\varepsilon^3. \quad (32)$$

(c) *In the limit $\delta \rightarrow 0$, the periodic solution $(\bar{\kappa}, \xi)$ satisfies*

$$\|\bar{\kappa} - \xi\|_{H^1(I)} \leq c\delta^{1/2} \left(\sup_{t \in [0, T]} \|\dot{\kappa}_0\|_{L^2} + \sup_{t \in [0, T]} \|\kappa_0\|_{H^1} \right). \quad (33)$$

In particular, as $\delta \rightarrow 0$, $(\bar{\kappa}, \xi)$ converges to the solution to the Newtonian PDE (17) with the same forcing κ_0 .

The proof of Theorem 1.2 appears in two parts: parts (a) and (b) are shown in section 3.4, and part (c) is shown in section 3.5.

Given a T -periodic κ_0 and the corresponding T -periodic solution guaranteed by Theorem 1.2, we may now study the actual swimming speed of the filament. We first calculate an expression for the swimming velocity $\mathbf{V}(t) = \int_0^1 \frac{\partial \mathbf{X}}{\partial t} ds$. The expression involves κ_0 , $\bar{\kappa}$, and ξ . To better understand the swimming speed of the filament, especially in relation to the Newtonian setting, we need an expression in terms of κ_0 only. We use the closeness of $(\bar{\kappa}, \xi)$ to $(\bar{\kappa}^{\text{lin}}, \xi^{\text{lin}})$ from Theorem 1.2 (b) to obtain an expression for the average filament swimming speed at leading order in terms of κ_0 only. Here and throughout, for $u = u(t)$, we will denote the time average over one period by

$$\langle u \rangle := \frac{1}{T} \int_0^T u dt. \quad (34)$$

The fiber swimming velocity $\mathbf{V}(t)$ and average speed in direction $\mathbf{e}_t(0, 0)$ are given as follows.

Theorem 1.3. *For ε as in Theorem 1.2, given a T -periodic $\kappa_0 \in C^1([0, T]; H^3(I))$ satisfying*

$$\sup_{t \in [0, T]} \|\dot{\kappa}_0\|_{L^2} = \varepsilon_1 \leq \varepsilon, \quad \sup_{t \in [0, T]} \|\kappa_0\|_{H^1} = \varepsilon_2 \leq \varepsilon, \quad (35)$$

a filament satisfying equations (13)-(16) swims with velocity

$$\mathbf{V}(t) = U(t)\mathbf{e}_t(0, 0) + \mathbf{r}_v(t), \quad (36)$$

where $\sup_{t \in [0, T]} |U| \leq c\varepsilon^2$, $\sup_{t \in [0, T]} |\mathbf{r}_v| \leq c\varepsilon^3$, and

$$U(t) = -\gamma \int_0^1 (\kappa_0)_s \bar{\kappa} ds - \gamma\mu \int_0^1 (\bar{\kappa} - \xi)(\bar{\kappa} + \kappa_0)_s ds. \quad (37)$$

Moreover, expanding κ_0 as $\kappa_0(s, t) = \sum_{m,k=1}^{\infty} (a_{m,k} \cos(\omega mt) - b_{m,k} \sin(\omega mt)) \psi_k(s)$, where ψ_k are the eigenfunctions (22) of the operator \mathcal{L} and $\omega = \frac{2\pi}{T}$, the average swimming speed $\langle U \rangle$ over the

course of one time period is given by

$$\begin{aligned} \langle U \rangle = \frac{\gamma}{2} \sum_{m,k,\ell=1}^{\infty} & \left(W_{1,m\ell k} (a_{m,k} b_{m,\ell} - a_{m,\ell} b_{m,k}) + W_{2,m\ell k} (a_{m,k} a_{m,\ell} + b_{m,k} b_{m,\ell}) \right. \\ & \left. + W_{3,m\ell k} (a_{m,k} a_{m,\ell} - b_{m,k} b_{m,\ell}) + W_{4,m\ell k} (a_{m,\ell} b_{m,k} + b_{m,\ell} a_{m,k}) \right) \int_0^1 \psi_k(\psi_\ell)_s ds + r_u. \end{aligned} \quad (38)$$

Here coefficients $W_{j,m\ell k}$ are explicitly computable (see equation (39)) and the remainder term r_u satisfies $|r_u| \leq c\varepsilon^4$.

The proof of Theorem 1.3 is given in section 4, and various observations and applications of the swimming expression (38) are explored immediately in the following section 2.

2. APPLICATIONS AND NUMERICAL RESULTS

Here we detail some of the main takeaways and applications of Theorem 1.3, interspersed with numerical simulations. For the numerical simulations, we will rely on a reformulation of (1)-(4) which avoids the need to solve for the fiber tension τ . We propose a natural extension of the numerical method in [26], which is itself based on a combination of the formulations of [23] and [22]. The method is described in detail in Appendix A.

We begin with some observations about the viscoelastic swimming speed expression (38). We first note the forms of the coefficients $W_{j,m\ell k}$, which are calculated in section 4. We have

$$\begin{aligned} W_{1,m\ell k} &= Q_{m,k} - \frac{\mu\delta\omega m}{1 + (\delta\omega m)^2} \left(-\delta\omega m(1 - H_{m,\ell})Q_{m,k} + Q_{m,\ell}Q_{m,k} \right) \\ W_{2,m\ell k} &= H_{m,k} - \frac{\mu\delta\omega m}{1 + (\delta\omega m)^2} \left(\delta\omega m Q_{m,\ell}Q_{m,k} + (1 - H_{m,\ell})Q_{m,k} \right) \\ W_{3,m\ell k} &= -\frac{\mu\delta\omega m}{1 + (\delta\omega m)^2} \left(\delta\omega m(1 - H_{m,\ell})H_{m,k} + Q_{m,\ell}H_{m,k} \right) \\ W_{4,m\ell k} &= -\frac{\mu\delta\omega m}{1 + (\delta\omega m)^2} \left(\delta\omega m Q_{m,\ell}H_{m,k} - (1 - H_{m,\ell})H_{m,k} \right), \end{aligned} \quad (39)$$

where

$$\begin{aligned} Q_{m,k} &= \frac{\lambda_k \omega m (1 + (1 + \mu)(\delta\omega m)^2)}{\lambda_k^2 (1 + (1 + \mu)^2 (\delta\omega m)^2) + \omega^2 m^2 (2\mu\delta\lambda_k + 1 + (\delta\omega m)^2)}, \\ H_{m,k} &= \frac{\omega^2 m^2 (\mu\delta\lambda_k + 1 + (\delta\omega m)^2)}{\lambda_k^2 (1 + (1 + \mu)^2 (\delta\omega m)^2) + \omega^2 m^2 (2\mu\delta\lambda_k + 1 + (\delta\omega m)^2)}. \end{aligned} \quad (40)$$

We will be comparing the coefficients (39) with the Newtonian swimmer for the same preferred curvature κ_0 . In [26], the Newtonian swimmer satisfying (17) was shown to swim with speed U^{nw} of the form

$$U^{\text{nw}}(t) = -\gamma \int_0^1 (\kappa_0)_s \bar{\kappa}^{\text{nw}} ds, \quad (41)$$

which, again averaging over one period, may be written to leading order in ε as

$$\begin{aligned} \langle U^{\text{nw}} \rangle = \frac{\gamma}{2} \sum_{m,k,\ell=1}^{\infty} \frac{\omega^2 m^2}{\omega^2 m^2 + \lambda_k^2} & \left(\frac{\lambda_k}{\omega m} (a_{m,k} b_{m,\ell} - b_{m,k} a_{m,\ell}) \right. \\ & \left. + a_{m,k} a_{m,\ell} + b_{m,k} b_{m,\ell} \right) \int_0^1 \psi_k(\psi_\ell)_s ds. \end{aligned} \quad (42)$$

Some things that we can immediately note in comparing (36) and (38) to (41) and (42) include:

0. The viscoelastic swimming speed has a complicated dependence on the parameters μ , δ , and ω and it is not immediately clear how it compares to the Newtonian swimming speed.
1. The velocity expression (36) has the form of the Newtonian swimming speed (41) plus a correction term proportional to $\mu(\bar{\kappa} - \xi)$, but note that $\bar{\kappa} \neq \bar{\kappa}^{\text{nw}}$ in general.
2. If either $\mu = 0$ or $\delta = 0$, the viscoelastic expression (38) reduces to the Newtonian expression (42). In particular, we obtain $W_{1,m\ell k} = Q_{m,k}$ where $Q_{m,k} = \frac{\lambda_k \omega m}{\lambda_k^2 + \omega^2 m^2}$, $W_{2,m\ell k} = H_{m,k}$ where $H_{m,k} = \frac{\omega^2 m^2}{\lambda_k^2 + \omega^2 m^2}$, and $W_{3,m\ell k} = W_{4,m\ell k} = 0$.
3. Due to the form of the eigenfunctions ψ_k of the operator \mathcal{L} (see (22)), in both the Newtonian and viscoelastic cases, if the preferred curvature $\kappa_0(s, t)$ is always odd or always even about the fiber midpoint $s = \frac{1}{2}$, the swimming speed will vanish. This is because $\psi_{2k}(s)$ is odd about $s = \frac{1}{2}$ and $\psi_{2k-1}(s)$ is even for each $k = 1, 2, \dots$, and thus $\int_0^1 \psi_{2k}(\psi_{2\ell})_s ds = 0$ and $\int_0^1 \psi_{2k-1}(\psi_{2\ell-1})_s ds = 0$ for each k, ℓ . In particular, if $\kappa_0(s, t)$ can be written purely in terms of either ψ_{2k} or ψ_{2k-1} , the swimming speed (38) will vanish.

Otherwise, owing to the complicated nature of the expression (38) and particularly of the coefficients (39), it is difficult to say much in general about the viscoelastic swimmer, but we can make some predictions in certain scenarios. For the following, we will consider κ_0 of the form

$$\kappa_0(s, t) = F_1(s) \cos(\omega t) + F_2(s) \sin(\omega t), \quad (43)$$

i.e. we will force only a single mode in time. As such, we will drop all dependence on the temporal mode m in our notation.

We will begin by considering the case $F_1 = F_2$, in which case the swimming speed expression (38) reduces to

$$\langle U \rangle = \gamma \sum_{k, \ell=1}^{\infty} (W_{2, \ell k} + W_{4, \ell k}) a_k a_\ell \int_0^1 \psi_k(\psi_\ell)_s ds. \quad (44)$$

We referred to these swimmers as ‘bad swimmers’ in the Newtonian setting [26] because the coefficient $W_{2, \ell k} + W_{4, \ell k}$ is given by $\omega^2/(\lambda_k^2 + \omega^2)$ and thus decays very rapidly as k increases. Even $\lambda_1^2 \approx 500^2$ is extremely large, and thus to see noticeable displacement for the swimmer over one period, we need to take ω very large as well. For numerical simulations, we will take $\omega = 32\pi$.

Since λ_k is so large, we consider the leading order behavior of the coefficient $W_{2, \ell k} + W_{4, \ell k}$ in $1/\lambda_k$, given by

$$W_{2, \ell k} + W_{4, \ell k} \approx H_k - \frac{\mu \delta \omega}{1 + (\delta \omega)^2} (Q_k + H_k). \quad (45)$$

We immediately note that, for large ω and fixed $\mu > 0$, the expression (45) behaves differently (and in fact, non-monotonically) depending on the size of δ with respect to ω . If $\delta = 0$ or if $\delta \approx 1$ or larger, then (45) reduces to just H_k . However, if $\delta \approx 1/\omega$, the second term plays a more prominent role and we should expect to see cancellations and possibly even a reversal of swimming direction.

We study the behavior of H_k (40) in more detail. We first rewrite H_k as

$$H_k = \frac{\omega^2}{\lambda_k^2 H_{1,k} + \omega^2 H_{2,k}} \quad \text{where} \quad H_{1,k} = \frac{(1 + (1 + \mu)^2 (\delta \omega)^2)}{(\mu \delta \lambda_k + 1 + (\delta \omega)^2)}, \quad H_{2,k} = \frac{(2\mu \delta \lambda_k + 1 + (\delta \omega)^2)}{(\mu \delta \lambda_k + 1 + (\delta \omega)^2)}.$$

In the Newtonian setting, we have $H_{1,k} = H_{2,k} = 1$. We still have $1 \leq H_{2,k} \leq 2$ for all $\delta, \mu > 0$, so we do not expect much of a contribution from this term for different δ and μ . However, we note that, for fixed $\delta > 0$, $H_{1,k}$ is strictly increasing in μ , and therefore H_k is strictly decreasing in μ . Furthermore, since we are taking ω to be large, if $\delta \gtrsim 1$, we should not see much of an effect from varying δ .

We therefore predict the following.

- a. For fixed $\delta \gtrsim 1$, we will see a slowdown in the swimmer as μ is increased. For fixed $\mu > 0$, we will see very little change in the swimming speed as $\delta \gtrsim 1$ is increased.
- b. For small $\delta = \frac{1}{\omega}$, however, we expect that the swimmer will be much slower than both the Newtonian ($\delta = 0$) or $\delta \gtrsim 1$ swimmers, and can potentially reverse direction.

To test these predictions, we choose a preferred curvature of the form (43) with $F_1 = (s-1)^2$ and $F_2 = F_1$ (note that we normalize such that $\|F_1\|_{L^2} = \|F_2\|_{L^2} = 1$). We take $\omega = 32\pi$ and simulate the fiber motion until $t = 2$ beginning from a straight line from $x = 0$ to $x = 1$ along the x -axis. We record the fiber's displacement $\int_0^1 \mathbf{X}(s, t_2) ds - \int_0^1 \mathbf{X}(s, t_1) ds$ between times $t_1 = 1$ and $t_2 = 2$ to ensure that the periodic solution has been reached.




FIGURE 1. (a) Location of the fiber at time $t = 2$ for fixed $\delta = 1$ and five different values of μ . All swimmers swim poorly, but the Newtonian swimmer (blue) is noticeably faster. (b) Comparison of swimmer shapes at ten different snapshots in time for the Newtonian (blue) and $\mu = \delta = 1$ (orange) swimmers.

To test prediction (a), we first fix $\delta = 1$ and compare the swimming displacement for 5 different values of μ (see Figure 1a). We compare $\mu = 0$ (Newtonian) against $\mu = 1, 2, 4, 8$. From Figure 1a, we can see that the Newtonian swimmer swims the farthest, although none of the swimmers swim very well. Between $t = 1$ and $t = 2$, the Newtonian swimmer's displacement is -0.036 . For $\mu = 1, 2, 4, 8$, respectively, the displacement is -0.018 , -0.012 , -0.0069 , -0.0035 . As predicted, the distance decreases with increasing μ . Furthermore, when μ is fixed at $\mu = 1$ and $\delta \gtrsim 1$ is varied, there is very little difference in the swimming displacement versus the $\delta = 1$ swimmer. When $\delta = 1, 2, 4, 8$, respectively, the swimming displacement is still -0.018 .

In Figure 1b, snapshots of the location of the swimmer with $\mu = \delta = 1$ at ten different points in time between $t = 0$ and $t = 2$ are plotted against the same points in time for the Newtonian swimmer.

To test prediction (b), we fix $\delta = 1/\omega$ and simulate the swimmer until $t = 2$ using different values of μ . We again calculate the swimmer's displacement between $t = 1$ and $t = 2$. When $\mu = 1$, the displacement is -0.019 , i.e. roughly the same as when $\delta = 1$. However, when $\mu = 2$, the swimmer's displacement is $+0.0062$, in particular, we do indeed find that the swimmer moves in the opposite direction (see Figure 2). This is the combination of μ and δ for which the H_k terms in




FIGURE 2. (a) Location of the fiber at time $t = 2$ for fixed $\delta = 1/\omega$ and five different values of μ . Notice that each of the viscoelastic swimmers appear to have moved to the right initially, but except for the $\mu = 2$ case, after an initial adjustment the swimmers do move leftward. (b) Comparison of swimmer shapes at ten different snapshots in time for the Newtonian (blue) and $\mu = 2, \delta = 1/\omega$ (yellow) swimmers. Note that the yellow swimmer moves backward.

(45) cancel, reducing the leading order coefficient to $-Q_k$. The behavior of the swimmer becomes more complicated as μ increases: for $\mu = 4$ and $\mu = 8$, we observe a displacement of -0.0030 and -0.0040 , respectively. The swimmer now moves in the same direction as for large δ , but instead of losing speed as μ increases, it appears to gain a bit of speed. For $\mu = 8$, the $\delta = 1/\omega$ swimmer even swims a bit further than the $\delta = 1$ swimmer. This may be due to the behavior of the coefficient Q_k in (45), which we consider in a bit more detail later.

In summary, for the simple case of a preferred curvature (43) satisfying $F_1 = F_2$, we already see a complex, non-monotonic response to varying δ and μ . The Newtonian swimmer $\delta = 0$ is by far the fastest. For $\delta \gtrsim 1$, the swimmer's displacement decreases with increasing μ . When $\delta = 1/\omega$, for some values of μ the swimmer's displacement is in the opposite direction, and as μ increases, the swimmers's displacement does not monotonically decrease. Importantly, most of this behavior can be predicted from the swimming speed expression (38).

We next consider the more complicated scenario of $F_1 \neq F_2$ in the preferred curvature equation (43). Now all terms are present in the swimming expression (38), and it is more difficult to make detailed predictions. However, we can still make some predictions for large $\delta \gtrsim 1$ and test the behavior of small $\delta \sim 1/\omega$ numerically.

To leading order in $\frac{1}{\lambda_k}$, the coefficients (39) of the swimming expression (38) are given by

$$\begin{aligned} W_{1,\ell k} &= Q_k + \frac{\mu(\delta\omega)^2}{1 + (\delta\omega)^2} Q_k, & W_{2,\ell k} &= H_k - \frac{\mu\delta\omega}{1 + (\delta\omega)^2} Q_k, \\ W_{3,\ell k} &= -\frac{\mu(\delta\omega)^2}{1 + (\delta\omega)^2} H_k, & W_{4,\ell k} &= \frac{\mu\delta\omega}{1 + (\delta\omega)^2} H_k. \end{aligned} \quad (46)$$

Again we will consider ω large. Due to the differing signs of coefficients in the general swimming expression (38), it is not so clear what behavior we should expect for $\delta \sim 1/\omega$, but we can still run

numerical simulations to see what happens.

For large $\delta \gtrsim 1$, however, we can further reduce the expressions (46). We will also use that $Q_k \sim \omega/\lambda_k$ while $H_k \sim \omega^2/\lambda_k^2$; in particular, the coefficients Q_k do not decay as quickly in k and indeed appear to play a more important role in numerics. We thus approximate the swimming speed for a general swimmer by

$$\langle U \rangle \approx \frac{\gamma}{2} \sum_{k,\ell=1}^{\infty} \frac{1 + (1+\mu)(\delta\omega)^2}{1 + (\delta\omega)^2} Q_k (a_k b_\ell - a_\ell b_k) \int_0^1 \psi_k(\psi_\ell)_s ds. \quad (47)$$

We now need to study the behavior of Q_k for different values of μ and $\delta \gtrsim 1$. We rewrite the expression (40) as

$$Q_k = \frac{\lambda_k \omega}{\lambda_k^2 Q_{1,k} + \omega^2 Q_{2,k}} \quad \text{where} \quad Q_{1,k} = \frac{1 + (1+\mu)^2(\delta\omega)^2}{1 + (1+\mu)(\delta\omega)^2}, Q_{2,k} = \frac{2\mu\delta\lambda_k + 1 + (\delta\omega)^2}{1 + (1+\mu)(\delta\omega)^2}.$$

Again, in the Newtonian setting, $Q_{1,k} = Q_{2,k} = 1$. We note that we have $1 \leq Q_{1,k} \leq 1 + \mu$, while the behavior of $Q_{2,k}$ is a bit more complicated but perhaps plays a smaller role than $Q_{1,k}$ due to the leading coefficient of ω^2 rather than λ_k^2 . When ω is large and $\delta \gtrsim 1$, the coefficient in (47) behaves approximately as

$$\frac{1 + (1+\mu)(\delta\omega)^2}{1 + (\delta\omega)^2} Q_k \approx \frac{(1 + (1+\mu)(\delta\omega)^2)^2}{(1 + (\delta\omega)^2)(1 + (1+\mu)^2(\delta\omega)^2)} \frac{\omega}{\lambda_k}.$$

In particular, for fixed δ , increasing μ may very slightly increase the swimming speed, and for fixed μ , increasing $\delta \gtrsim 1$ again does not appear to have a strong effect. This is not a very exciting prediction, but we can again test it numerically. We can also test what happens for $\delta = 1/\omega$, where predictions are harder to make.




FIGURE 3. The form of F_1 and F_2 in the preferred curvature (43) used in numerical tests. This F_1 and F_2 came from a small numerical optimization of the Newtonian swimming speed (42) in [26].

To test our predictions for large $\delta \gtrsim 1$ and to see what happens for small $\delta = 1/\omega$, we use the preferred curvature components F_1 and F_2 pictured in Figure 3. These F_1 and F_2 were computed in [26] as one of two ‘optimal’ preferred curvatures κ_0 of the form (43) resulting in the greatest average swimming speed (42) in the Newtonian setting. The optimization of (42) was performed over the first 12 spatial modes k of κ_0 and thus may not exactly represent the true optimal κ_0 for the Newtonian swimmer. However, we note that in the Newtonian setting, the combination of F_1 and F_2

plotted in Figure 3 does outperform the classical traveling wave forcing $F_1 = \sin(\omega s)$, $F_2 = \cos(\omega s)$.

As before, we take $\omega = 32\pi$ and simulate the swimmer until $t = 2$. The swimmer begins as a straight line along the x -axis from $x = 0$ to $x = 1$. Again, we keep track of the displacement of the swimmer $\int_0^1 \mathbf{X}(s, t_2) ds - \int_0^1 \mathbf{X}(s, t_1) ds$ between times $t_1 = 1$ and $t_2 = 2$.




FIGURE 4. (a) Location of the fiber at time $t = 2$ for fixed $\delta = 1$ and five different values of μ . All swimmers swim roughly the same distance, but note that the Newtonian swimmer (blue) is slightly slower. This is opposite from the viscoelastic effects pictured in Figure 1. (b) Comparison of swimmer shapes at ten different snapshots in time for the Newtonian (blue), $\mu = \delta = 1$ (orange), and $\mu = 8, \delta = 1$ (green) swimmers.

For the case $\delta \gtrsim 1$, we again start by fixing $\delta = 1$ and compare the fiber displacement for 5 values of μ (see Figure 4a). In both the viscoelastic and Newtonian settings, the swimmers swim much further than in the case $F_1 = F_2$ above, and the differences among their displacements is much smaller. However, we note that the Newtonian swimmer ($\mu = 0$) has the smallest displacement between $t = 1$ and $t = 2$ of -0.380 . This may be compared with each of the $\mu = 1, 2, 4, 8$ swimmers, which have a displacement of -0.390 , -0.391 , -0.392 , and -0.390 , respectively. Besides the jump in swimming speed between the Newtonian swimmer ($\mu = 0$) and $\mu = 1$, there is not much difference in displacement among different values of μ . A similar result holds when $\mu = 1$ is fixed and $\delta = 1, 2, 4, 8$ is varied. The displacement in each of these cases is -0.390 .

The effect of varying μ seems to be a bit more interesting at small δ . Fixing $\delta = 1/\omega$, we consider $\mu = 0, 1, 2, 4, 8$. Recalling that the displacement of the Newtonian swimmer from $t = 1$ to $t = 2$ was -0.380 , we note that for $\mu = 1, 2, 4, 8$, the swimmer's displacement was -0.373 , -0.375 , -0.380 , and -0.384 , respectively (see Figure 5). The effect of varying μ is still relatively small, but already more complex than at large δ , varying from a slight enhancement of the swimming speed at larger μ to a slight inhibition of the swimming speed at smaller μ .

We note that the numerical tests performed in this section cover a very small portion of the possible parameter space, and indeed an even smaller portion of the possible forcing functions κ_0 . We hope, however, that the tests included here serve to emphasize the complexity of possible behaviors in this model over just a small range of the possible options. We believe this justifies studying the




FIGURE 5. (a) Location of the fiber at time $t = 2$ for fixed $\delta = 1/\omega$ and five different values of μ . Again, swimmers swim roughly the same distance, but now the Newtonian swimmer (blue) is faster than the $\mu = 1, 2$ viscoelastic swimmers and slower than the $\mu = 4, 8$ swimmers. (b) Comparison of swimmer shapes at ten different snapshots in time for the Newtonian (blue), $\mu = \delta = 1$ (orange), and $\mu = 8, \delta = 1$ (green) swimmers.

model (1)-(4) in more detail, and hopefully provides convincing evidence that linear viscoelasticity can have an interesting effect on small-amplitude undulatory swimming.

The remainder of this paper is devoted to proving Theorems 1.1, 1.2, and 1.3 regarding the PDE behavior of the model (1)-(4).

3. WELL-POSEDNESS AND PERIODIC SOLUTIONS

In this section we prove Theorems 1.1 and 1.2. We start by showing some preliminary bounds in sections 3.1 and 3.2, and then proceed to the proof of Theorem 1.1 in section 3.3. Sections 3.4 and 3.5 contain the proof of Theorem 1.2.

3.1. Semigroup properties. We begin by deriving the following estimates for the semigroup generated by the linear operator \mathcal{A} , given by (26).

Lemma 3.1. *For any $(u, \phi)^T \in L^2(I) \times L^2(I)$, for $0 \leq m + j \leq 4$, we have*

$$\left\| e^{t\mathcal{A}} \begin{pmatrix} \partial_s^j u \\ \partial_s^j \phi \end{pmatrix} \right\|_{\dot{H}^m(I) \times \dot{H}^m(I)} \leq c \max\{t^{-(m+j)/4}, 1\} e^{-t\Lambda} \left\| \begin{pmatrix} u \\ \phi \end{pmatrix} \right\|_{L^2(I) \times L^2(I)}, \quad (48)$$

where $\Lambda = \min\{\lambda_1, \frac{1}{\delta(1+\mu)}\}$ for λ_1 as in (22), and the constant c is independent of δ .

As a consequence of Lemma 3.1, we may also show the following small time estimate, which relies on approximating $(u, \phi)^T \in L^2(I) \times L^2(I)$ by functions in $D(\mathcal{L}^r) \times D(\mathcal{L}^r)$, $0 < r \leq 1$.

Lemma 3.2. *Fix $(u, \phi)^T \in L^2(I) \times L^2(I)$ and let $0 < r \leq 1$ and $\varepsilon > 0$. There exists $T_\varepsilon > 0$ depending on u and ϕ such that*

$$\sup_{t \in [0, T_\varepsilon]} \min\{t^r, 1\} e^{t\Lambda} \left\| e^{t\mathcal{A}} \begin{pmatrix} u \\ \phi \end{pmatrix} \right\|_{\dot{H}^{4r}(I) \times \dot{H}^{4r}(I)} \leq \varepsilon. \quad (49)$$

Proof of Lemma 3.1. For $(w, \varphi)^T \in D(\mathcal{L}) \times D(\mathcal{L})$, the eigenfunction expansion of $\mathcal{A} \begin{pmatrix} w \\ \varphi \end{pmatrix}$ (see (23)) may be written as $\sum_{k=1}^{\infty} \tilde{\mathcal{A}}_k \begin{pmatrix} \tilde{w}_k \\ \tilde{\varphi}_k \end{pmatrix} \psi_k$, where

$$\tilde{\mathcal{A}}_k = \begin{pmatrix} -(1+\mu)\lambda_k & \mu\lambda_k \\ \delta^{-1} & -\delta^{-1} \end{pmatrix}. \quad (50)$$

We study the properties of $\tilde{\mathcal{A}}_k$. The eigenvalues of $\tilde{\mathcal{A}}_k$ are given by

$$\nu_k^{\pm} = \frac{1}{2\delta} \left(- (1 + (1+\mu)\delta\lambda_k) \pm \sqrt{(1 + (1+\mu)\delta\lambda_k)^2 - 4\delta\lambda_k} \right), \quad (51)$$

with corresponding eigenvectors

$$\mathbf{v}_k^- = \begin{pmatrix} 1 \\ \frac{1}{1+\delta\nu_k^-} \end{pmatrix}, \quad \mathbf{v}_k^+ = \begin{pmatrix} 1 + \delta\nu_k^+ \\ 1 \end{pmatrix}. \quad (52)$$

We note in particular that $\delta\nu_k^+ \rightarrow -\frac{1}{1+\mu}$ monotonically as $k \rightarrow \infty$, while $\lambda_k^{-1}\delta\nu_k^- \rightarrow -\delta(1+\mu)$ as $k \rightarrow \infty$. We may decompose the unit vectors $(1, 0)^T$ and $(0, 1)^T$ in terms of the eigenvectors \mathbf{v}_k^{\pm} of $\tilde{\mathcal{A}}_k$ as:

$$\begin{aligned} \begin{pmatrix} 1 \\ 0 \end{pmatrix} &= a_k^- \mathbf{v}_k^- + a_k^+ \mathbf{v}_k^+, & a_k^- &= -\frac{1 + \delta\nu_k^-}{\delta(\nu_k^+ - \nu_k^-)}; & a_k^+ &= \frac{1}{\delta(\nu_k^+ - \nu_k^-)}, \\ \begin{pmatrix} 0 \\ 1 \end{pmatrix} &= b_k^- \mathbf{v}_k^- + b_k^+ \mathbf{v}_k^+, & b_k^- &= \frac{(1 + \delta\nu_k^+)(1 + \delta\nu_k^-)}{\delta(\nu_k^+ - \nu_k^-)}; & b_k^+ &= -\frac{1 + \delta\nu_k^-}{\delta(\nu_k^+ - \nu_k^-)}. \end{aligned} \quad (53)$$

Noting that

$$\delta(\nu_k^+ - \nu_k^-) = \sqrt{(1 + (1+\mu)\delta\lambda_k)^2 - 4\delta\lambda_k},$$

we have that there exist constants c independent of both k and δ such that

$$|a_k^- \mathbf{v}_k^-| \leq c, \quad |b_k^- \mathbf{v}_k^-| \leq c, \quad \text{and} \quad (\delta\lambda_k)^r |a_k^+ \mathbf{v}_k^+| \leq c, \quad (\delta\lambda_k)^r |b_k^+ \mathbf{v}_k^+| \leq c, \quad 0 \leq r \leq 1. \quad (54)$$

Using the decomposition (53) and the bounds (54), for any $0 \leq r \leq 1$, we may estimate

$$\begin{aligned} \left\| \mathcal{L}^r e^{t\mathcal{A}} \begin{pmatrix} u \\ 0 \end{pmatrix} \right\|_{L^2 \times L^2} &= \left\| \sum_{k=1}^{\infty} \lambda_k^r e^{t\tilde{\mathcal{A}}_k} \begin{pmatrix} \tilde{u}_k \psi_k \\ 0 \end{pmatrix} \right\|_{L^2 \times L^2} = \left\| \sum_{k=1}^{\infty} \lambda_k^r \left(a_k^- \mathbf{v}_k^- e^{t\nu_k^-} + a_k^+ \mathbf{v}_k^+ e^{t\nu_k^+} \right) \tilde{u}_k \psi_k \right\|_{L^2 \times L^2} \\ &\leq c \sup_k \left(\lambda_k^r e^{t\nu_k^-} + \delta^{-r} e^{t\nu_k^+} \right) \left\| \sum_{k=1}^{\infty} \tilde{u}_k \psi_k \right\|_{L^2} \\ &\leq c \left(\sup_k \left(\lambda_k^r e^{-t(\lambda_k - \lambda_1)} \right) e^{-t\lambda_1} + \delta^{-r} e^{-t/(\delta(1+\mu))} \right) \|u\|_{L^2} \\ &\leq c \max\{t^{-r}, 1\} \left(e^{-t\lambda_1} + e^{-t/(\delta(1+\mu))} \right) \|u\|_{L^2}. \end{aligned}$$

By an analogous series of estimates, we may also show

$$\begin{aligned} \left\| \mathcal{L}^r e^{t\mathcal{A}} \begin{pmatrix} 0 \\ \phi \end{pmatrix} \right\|_{L^2 \times L^2} &= \left\| \sum_{k=1}^{\infty} \lambda_k^r e^{t\tilde{\mathcal{A}}_k} \begin{pmatrix} 0 \\ \tilde{\phi}_k \psi_k \end{pmatrix} \right\|_{L^2 \times L^2} = \left\| \sum_{k=1}^{\infty} \left(b_k^- \mathbf{v}_k^- e^{\nu_k^- t} + b_k^+ \mathbf{v}_k^+ e^{\nu_k^+ t} \right) \lambda_k^r \tilde{\phi}_k \psi_k \right\|_{L^2 \times L^2} \\ &\leq c \max\{t^{-r}, 1\} \left(e^{-t\lambda_1} + e^{-t/(\delta(1+\mu))} \right) \|\phi\|_{L^2}. \end{aligned}$$

Recalling that $D(\mathcal{L}^r) \subseteq H^{4r}$ for each $0 \leq r \leq 1$ by (24), we obtain estimate (48) for $j = 0$.

For $0 < j \leq 4$, we proceed by a duality argument as in [26]. In particular, for $(u, \phi), (w, \varphi) \in C_c^\infty(I) \times C_c^\infty(I)$, we have

$$\begin{aligned} \sup_{\|(u, \phi)\|_{L^2 \times L^2} = 1} \left\| e^{t\mathcal{A}} \begin{pmatrix} \partial_s^j u \\ \partial_s^j \phi \end{pmatrix} \right\|_{L^2 \times L^2} &= \sup_{\|(u, \phi)\|_{L^2 \times L^2} = \|(w, \varphi)\|_{L^2 \times L^2} = 1} \left(\begin{pmatrix} w \\ \varphi \end{pmatrix}, e^{t\mathcal{A}} \begin{pmatrix} \partial_s^j u \\ \partial_s^j \phi \end{pmatrix} \right)_{L^2 \times L^2} \\ &= \sup_{\|(u, \phi)\|_{L^2 \times L^2} = \|(w, \varphi)\|_{L^2 \times L^2} = 1} \left(\partial_s^j e^{t\mathcal{A}^*} \begin{pmatrix} w \\ \varphi \end{pmatrix}, \begin{pmatrix} u \\ \phi \end{pmatrix} \right)_{L^2 \times L^2} \\ &\leq \sup_{\|(w, \varphi)\|_{L^2 \times L^2} = 1} \left\| \partial_s^j e^{t\mathcal{A}^*} \begin{pmatrix} w \\ \varphi \end{pmatrix} \right\|_{L^2 \times L^2}, \end{aligned}$$

where we are using the notation

$$\left(\begin{pmatrix} w \\ \varphi \end{pmatrix}, \begin{pmatrix} u \\ \phi \end{pmatrix} \right)_{L^2 \times L^2} := (w, u)_{L^2} + (\varphi, \phi)_{L^2}.$$

Now, the adjoint operator \mathcal{A}^* satisfies

$$\widetilde{\mathcal{A}}^*_k = (\widetilde{\mathcal{A}}_k)^T = \begin{pmatrix} -(1+\mu)\lambda_k & \delta^{-1} \\ \mu\lambda_k & -\delta^{-1} \end{pmatrix},$$

with the same eigenvalues (51) as $\widetilde{\mathcal{A}}_k$ but with eigenvectors given by

$$\mathbf{v}_k^{*+} = \begin{pmatrix} \frac{1+\delta\nu_k^+}{\mu\delta\lambda_k} \\ 1 \end{pmatrix}, \quad \mathbf{v}_k^{*-} = \begin{pmatrix} 1 \\ \frac{\mu\delta\lambda_k}{1+\delta\nu_k^-} \end{pmatrix}. \quad (55)$$

We may again decompose the unit vectors $(1, 0)^T$ and $(0, 1)^T$ in terms of the eigenvectors (55) of $\widetilde{\mathcal{A}}^*_k$ as

$$\begin{aligned} \begin{pmatrix} 1 \\ 0 \end{pmatrix} &= a_k^{*-} \mathbf{v}_k^{*-} + a_k^{*+} \mathbf{v}_k^{*+}, \quad a_k^{*-} = -\frac{1+\delta\nu_k^-}{\delta(\nu_k^+ - \nu_k^-)}; \quad a_k^{*+} = \frac{\mu\delta\lambda_k}{\delta(\nu_k^+ - \nu_k^-)}, \\ \begin{pmatrix} 0 \\ 1 \end{pmatrix} &= b_k^{*-} \mathbf{v}_k^{*-} + b_k^{*+} \mathbf{v}_k^{*+}, \quad b_k^{*-} = \frac{(1+\delta\nu_k^+)(1+\delta\nu_k^-)}{\mu\delta\lambda_k \delta(\nu_k^+ - \nu_k^-)}; \quad b_k^{*+} = -\frac{1+\delta\nu_k^-}{\delta(\nu_k^+ - \nu_k^-)}. \end{aligned} \quad (56)$$

Note that

$$\begin{aligned} 1 + \delta\nu_k^+ &= \frac{1}{2} - \frac{1}{2} \sqrt{(1 + (1+\mu)\delta\lambda_k)^2 - 4\delta\lambda_k} - \frac{1}{2}(1+\mu)\delta\lambda_k \\ &= -\frac{\delta\lambda_k}{2} \left(\frac{2(1+\mu) + (1+\mu)^2\delta\lambda_k + 4}{1 + \sqrt{(1 + (1+\mu)\delta\lambda_k)^2 - 4\delta\lambda_k}} + (1+\mu) \right); \end{aligned}$$

in particular, we may bound

$$\left| \frac{1 + \delta\nu_k^+}{\delta\lambda_k} \right| \leq c$$

for some c independent of k and δ . Then, as in (54), we have that the components in (56) satisfy

$$|a_k^{*-} \mathbf{v}_k^{*-}| \leq c, \quad |b_k^{*-} \mathbf{v}_k^{*-}| \leq c, \quad (\delta\lambda_k)^r |a_k^{*+} \mathbf{v}_k^{*+}| \leq c, \quad (\delta\lambda_k)^r |b_k^{*+} \mathbf{v}_k^{*+}| \leq c, \quad 0 \leq r \leq 1. \quad (57)$$

Using the decomposition (56) and the bounds (57) in the same way as above, we have that

$$\begin{aligned} \left\| \mathcal{L}^{j/4} e^{t\mathcal{A}^*} \begin{pmatrix} w \\ 0 \end{pmatrix} \right\|_{L^2 \times L^2} &= \left\| \sum_{k=1}^{\infty} \lambda_k^{j/4} \left(a_k^{*-} \mathbf{v}_k^{*-} e^{t\nu_k^-} + a_k^{*+} \mathbf{v}_k^{*+} e^{t\nu_k^+} \right) \widetilde{w}_k \psi_k \right\|_{L^2 \times L^2} \\ &\leq c \max\{t^{-j/4}, 1\} \left(e^{-t\lambda_1} + e^{-t/(\delta(1+\mu))} \right) \|w\|_{L^2} \end{aligned}$$

and

$$\begin{aligned} \left\| \mathcal{L}^{j/4} e^{t\mathcal{A}^*} \begin{pmatrix} 0 \\ \varphi \end{pmatrix} \right\|_{L^2 \times L^2} &= \left\| \sum_{k=1}^{\infty} \lambda_k^{j/4} \left(b_k^{*-} \mathbf{v}_k^{*-} e^{t\nu_k^-} + b_k^{*+} \mathbf{v}_k^{*+} e^{t\nu_k^+} \right) \tilde{\varphi}_k \psi_k \right\|_{L^2 \times L^2} \\ &\leq c \max\{t^{-j/4}, 1\} \left(e^{-t\lambda_1} + e^{-t/(\delta(1+\mu))} \right) \|\varphi\|_{L^2}. \end{aligned}$$

In particular, since $D(\mathcal{L}^{j/4}) \subseteq H^j$ for $0 \leq j \leq 4$, we have

$$\begin{aligned} \sup_{\|(u, \phi)\|_{L^2 \times L^2} = 1} \left\| e^{t\mathcal{A}} \begin{pmatrix} \partial_s^j u \\ \partial_s^j \phi \end{pmatrix} \right\|_{L^2 \times L^2} &= \sup_{\|(w, \varphi)\|_{L^2 \times L^2} = 1} \left\| \partial_s^j e^{t\mathcal{A}^*} \begin{pmatrix} w \\ \varphi \end{pmatrix} \right\|_{L^2 \times L^2} \\ &\leq c \max\{t^{-j/4}, 1\} \left(e^{-t\lambda_1} + e^{-t/(\delta(1+\mu))} \right). \end{aligned}$$

The desired estimate then holds for $(u, \phi) \in L^2 \times L^2$ by density. \square

Proof of Lemma 3.2. Let $u_n = \sum_{k=1}^n \tilde{u}_k \psi_k$ and $\phi_n = \sum_{k=1}^n \tilde{\phi}_k \psi_k$. Since these sums are finite, we have $u_n, \phi_n \in D(\mathcal{L}^r)$ (see (24)) and we may estimate

$$\left\| e^{t\mathcal{A}} \begin{pmatrix} u_n \\ \phi_n \end{pmatrix} \right\|_{\dot{H}^{4r} \times \dot{H}^{4r}} \leq \left\| \mathcal{L}^r e^{t\mathcal{A}} \begin{pmatrix} u_n \\ \phi_n \end{pmatrix} \right\|_{L^2 \times L^2} \leq c e^{-t\Lambda} \left\| \sum_{k=1}^n \lambda_k^r \psi_k \begin{pmatrix} \tilde{u}_k \\ \tilde{\phi}_k \end{pmatrix} \right\|_{L^2 \times L^2} \leq c_n e^{-t\Lambda}.$$

By Lemma 3.1, we then have

$$\begin{aligned} e^{t\Lambda} \min\{t^r, 1\} \left\| e^{t\mathcal{A}} \begin{pmatrix} u \\ \phi \end{pmatrix} \right\|_{\dot{H}^{4r} \times \dot{H}^{4r}} &\leq e^{t\Lambda} \min\{t^r, 1\} \left(\left\| e^{t\mathcal{A}} \begin{pmatrix} u - u_n \\ \phi - \phi_n \end{pmatrix} \right\|_{\dot{H}^{4r} \times \dot{H}^{4r}} + \left\| e^{t\mathcal{A}} \begin{pmatrix} u_n \\ \phi_n \end{pmatrix} \right\|_{\dot{H}^{4r} \times \dot{H}^{4r}} \right) \\ &\leq c \left\| \begin{pmatrix} u - u_n \\ \phi - \phi_n \end{pmatrix} \right\|_{L^2 \times L^2} + \min\{t^r, 1\} c_n. \end{aligned}$$

Taking n sufficiently large and t sufficiently small (depending on n), we obtain Lemma 3.2. \square

3.2. Tension equation. We next prove the following lemma regarding the elliptic equation (15) for the tension $\bar{\tau}$.

Lemma 3.3. *Given $(\bar{\kappa}, \xi, \kappa_0) \in H^1(I) \times H^1(I) \times H^1(I)$, there exists a unique weak solution $\bar{\tau} \in H_0^1(I)$ to (15) satisfying*

$$\|\bar{\tau}\|_{H^1(I)} \leq c \left(\|\bar{\kappa}\|_{\dot{H}^1}^2 (\|\bar{\kappa}\|_{L^2} + 1) + \|\xi\|_{\dot{H}^1} (\|\bar{\kappa}\|_{\dot{H}^1} + \|\kappa_0\|_{H^1}) + \|\bar{\kappa}\|_{\dot{H}^1} \|\kappa_0\|_{H^1} (\|\kappa_0\|_{L^2} + 1) \right). \quad (58)$$

Furthermore, given $(\bar{\kappa}_a, \xi_a), (\bar{\kappa}_b, \xi_b) \in H^1(I) \times H^1(I)$, define $\bar{\tau}_a, \bar{\tau}_b \in H_0^1(I)$ to be the corresponding unique weak solutions to (15). The difference $\bar{\tau}_a - \bar{\tau}_b$ then satisfies

$$\begin{aligned} \|\bar{\tau}_a - \bar{\tau}_b\|_{H^1} &\leq c \|\bar{\kappa}_a - \bar{\kappa}_b\|_{L^2} \left((\|\xi_a\|_{\dot{H}^1} + \|\xi_b\|_{\dot{H}^1}) (\|\bar{\kappa}_a\|_{\dot{H}^1} + \|\bar{\kappa}_b\|_{\dot{H}^1} + \|\kappa_0\|_{H^1}) \right. \\ &\quad \left. + (\|\bar{\kappa}_a\|_{\dot{H}^1}^2 + \|\bar{\kappa}_b\|_{\dot{H}^1}^2 + \|\kappa_0\|_{H^1}^2) \right) (\|\bar{\kappa}_a\|_{L^2} + \|\bar{\kappa}_b\|_{L^2} + \|\kappa_0\|_{L^2} + 1)^2 \\ &\quad + c \|\bar{\kappa}_a - \bar{\kappa}_b\|_{\dot{H}^1} (\|\bar{\kappa}_a\|_{\dot{H}^1} + \|\bar{\kappa}_b\|_{\dot{H}^1} + \|\xi_a\|_{\dot{H}^1} + \|\kappa_0\|_{H^1}) \\ &\quad + c \|\xi_a - \xi_b\|_{\dot{H}^1} (\|\bar{\kappa}_b\|_{\dot{H}^1} + \|\kappa_0\|_{H^1}). \end{aligned} \quad (59)$$

Proof. We begin by decomposing $\bar{\tau}$ into $\bar{\tau} = \bar{\tau}^{\text{nw}} + \bar{\tau}^{\text{ve}}$, where

$$\bar{\tau}_{ss}^{\text{nw}} - \frac{(\bar{\kappa} + \kappa_0)^2}{1 + \gamma} \bar{\tau}^{\text{nw}} = \frac{1}{1 + \gamma} \mathcal{T}[\bar{\kappa}, \kappa_0] \quad (60)$$

$$\bar{\tau}_{ss}^{\text{ve}} - \frac{(\bar{\kappa} + \kappa_0)^2}{1 + \gamma} \bar{\tau}^{\text{ve}} = \frac{1}{1 + \gamma} \frac{\mu}{1 + \mu} \left((2 + \gamma)((\bar{\kappa} + \kappa_0)\xi_s)_s - (\bar{\kappa} + \kappa_0)_s \xi_s \right). \quad (61)$$

From [26], we have that there exists a unique $\bar{\tau}^{\text{nw}} \in H_0^1(I)$ satisfying (60) in a weak sense, with

$$\|\bar{\tau}^{\text{nw}}\|_{H^1} \leq c \left(\|\bar{\kappa}\|_{\dot{H}^1}^2 (\|\bar{\kappa}\|_{L^2} + 1) + \|\bar{\kappa}\|_{\dot{H}^1} \|\kappa_0\|_{H^1} (\|\kappa_0\|_{L^2} + 1) \right). \quad (62)$$

It thus remains to consider (61). As in the Newtonian setting, we define the bilinear form

$$\mathcal{B}(\bar{\tau}, \phi) := \int_0^1 \left(\bar{\tau}_s \phi_s + \frac{(\bar{\kappa} + \kappa_0)^2}{1 + \gamma} \bar{\tau} \phi \right) ds,$$

which is bounded and coercive on $H_0^1(I)$. A weak solution to (61) may then be defined as $\bar{\tau}^{\text{ve}} \in H_0^1$ satisfying

$$\mathcal{B}(\bar{\tau}^{\text{ve}}, \phi) = \frac{1}{1 + \gamma} \frac{\mu}{1 + \mu} \int_0^1 \left((2 + \gamma)(\bar{\kappa} + \kappa_0)\xi_s \phi_s + (\bar{\kappa} + \kappa_0)_s \xi_s \phi \right) ds$$

for all $\phi \in H_0^1(I)$, and existence and uniqueness follow from the Lax-Milgram lemma. Furthermore, we may estimate

$$\mathcal{B}(\bar{\tau}^{\text{ve}}, \bar{\tau}^{\text{ve}}) \leq c(\|\bar{\kappa}\|_{\dot{H}^1} + \|\kappa_0\|_{H^1}) \|\xi\|_{\dot{H}^1} \|\bar{\tau}^{\text{ve}}\|_{H^1},$$

and using that $\|\bar{\tau}^{\text{ve}}\|_{H^1}^2 \leq c\mathcal{B}(\bar{\tau}^{\text{ve}}, \bar{\tau}^{\text{ve}})$ along with Young's inequality, we obtain

$$\|\bar{\tau}^{\text{ve}}\|_{H^1} \leq c(\|\bar{\kappa}\|_{\dot{H}^1} + \|\kappa_0\|_{H^1}) \|\xi\|_{\dot{H}^1}. \quad (63)$$

Combining estimates (62) and (63), we obtain (58).

To show the Lipschitz estimate (59), we first recall that, from [26], we have

$$\begin{aligned} \|\bar{\tau}_a^{\text{nw}} - \bar{\tau}_b^{\text{nw}}\|_{H^1} &\leq c \|\bar{\kappa}_a - \bar{\kappa}_b\|_{L^2} \left(\|\bar{\kappa}_a\|_{\dot{H}^1}^2 + \|\bar{\kappa}_b\|_{\dot{H}^1}^2 + \|\kappa_0\|_{H^1}^2 \right) (\|\bar{\kappa}_a\|_{L^2} + \|\bar{\kappa}_b\|_{L^2} + \|\kappa_0\|_{L^2} + 1)^2 \\ &\quad + c \|\bar{\kappa}_a - \bar{\kappa}_b\|_{\dot{H}^1} \left(\|\bar{\kappa}_a\|_{\dot{H}^1} + \|\bar{\kappa}_b\|_{\dot{H}^1} + \|\kappa_0\|_{H^1} \right). \end{aligned} \quad (64)$$

It remains to estimate the viscoelastic contribution, which (weakly) satisfies

$$\begin{aligned} (\bar{\tau}_a^{\text{ve}} - \bar{\tau}_b^{\text{ve}})_{ss} - \frac{(\bar{\kappa}_a + \kappa_0)^2 + (\bar{\kappa}_b + \kappa_0)^2}{2(1 + \gamma)} (\bar{\tau}_a^{\text{ve}} - \bar{\tau}_b^{\text{ve}}) \\ = \frac{1}{1 + \gamma} \left(\frac{1}{2} (\bar{\kappa}_a - \bar{\kappa}_b)(\bar{\kappa}_a + \bar{\kappa}_b + 2\kappa_0)(\bar{\tau}_a^{\text{ve}} + \bar{\tau}_b^{\text{ve}}) \right. \\ \left. + \frac{\mu}{1 + \mu} \left((2 + \gamma)((\bar{\kappa}_a - \bar{\kappa}_b)(\xi_a)_s + (\bar{\kappa}_b + \kappa_0)(\xi_a - \xi_b)_s)_s \right. \right. \\ \left. \left. - (\bar{\kappa}_a - \bar{\kappa}_b)_s (\xi_a)_s + (\bar{\kappa}_b + \kappa_0)_s (\xi_a - \xi_b)_s \right) \right). \end{aligned}$$

In particular, we have

$$\begin{aligned} \|\bar{\tau}_a^{\text{ve}} - \bar{\tau}_b^{\text{ve}}\|_{H^1}^2 &\leq c \int_0^1 \left((\bar{\tau}_a^{\text{ve}} - \bar{\tau}_b^{\text{ve}})_s^2 + \frac{(\bar{\kappa}_a + \kappa_0)^2 + (\bar{\kappa}_b + \kappa_0)^2}{2(1 + \gamma)} (\bar{\tau}_a^{\text{ve}} - \bar{\tau}_b^{\text{ve}})^2 \right) ds \\ &\leq c \left(\|\bar{\kappa}_a - \bar{\kappa}_b\|_{L^2} (\|\bar{\kappa}_a\|_{L^2} + \|\bar{\kappa}_b\|_{L^2} + \|\kappa_0\|_{L^2}) (\|\bar{\kappa}_a\|_{\dot{H}^1} \|\xi_a\|_{\dot{H}^1} + \|\bar{\kappa}_b\|_{\dot{H}^1} \|\xi_b\|_{\dot{H}^1} \right. \\ &\quad \left. + (\|\xi_a\|_{\dot{H}^1} + \|\xi_b\|_{\dot{H}^1}) \|\kappa_0\|_{H^1} \right) \|\bar{\tau}_a^{\text{ve}} - \bar{\tau}_b^{\text{ve}}\|_{L^2} \end{aligned}$$

$$+ \left(\|\bar{\kappa}_a - \bar{\kappa}_b\|_{\dot{H}^1} \|\xi_a\|_{\dot{H}^1} + (\|\bar{\kappa}_b\|_{\dot{H}^1} + \|\kappa_0\|_{H^1}) \|\xi_a - \xi_b\|_{\dot{H}^1} \right) \|\bar{\tau}_a^{\text{ve}} - \bar{\tau}_b^{\text{ve}}\|_{H^1} \Bigg).$$

Applying Young's inequality, we obtain

$$\begin{aligned} \|\bar{\tau}_a^{\text{ve}} - \bar{\tau}_b^{\text{ve}}\|_{H^1} &\leq c \left(\|\bar{\kappa}_a - \bar{\kappa}_b\|_{L^2} (\|\bar{\kappa}_a\|_{L^2} + \|\bar{\kappa}_b\|_{L^2} + \|\kappa_0\|_{L^2}) (\|\bar{\kappa}_a\|_{\dot{H}^1} + \|\bar{\kappa}_b\|_{\dot{H}^1} \right. \\ &\quad \left. + \|\kappa_0\|_{H^1}) (\|\xi_a\|_{\dot{H}^1} + \|\xi_b\|_{\dot{H}^1}) \right. \\ &\quad \left. + \|\bar{\kappa}_a - \bar{\kappa}_b\|_{\dot{H}^1} \|\xi_a\|_{\dot{H}^1} + (\|\bar{\kappa}_b\|_{\dot{H}^1} + \|\kappa_0\|_{H^1}) \|\xi_a - \xi_b\|_{\dot{H}^1} \right). \end{aligned} \quad (65)$$

Combining (64) and (65) yields (59). \square

3.3. Evolution equation. We now proceed to the proof of Theorem 1.1.

Proof of Theorem 1.1. We will consider the map

$$\begin{aligned} \Psi \left[\begin{pmatrix} \bar{\kappa} \\ \xi \end{pmatrix} \right] &= e^{\mathcal{A}t} \begin{pmatrix} \bar{\kappa}^{\text{in}} \\ \xi^{\text{in}} \end{pmatrix} - \int_0^t e^{\mathcal{A}(t-t')} \begin{pmatrix} \dot{\kappa}_0 \\ 0 \end{pmatrix} dt' + (1 + \mu) \int_0^t e^{\mathcal{A}(t-t')} \begin{pmatrix} (\mathcal{N}[\bar{\kappa}, \kappa_0])_s \\ 0 \end{pmatrix} dt' \\ &\quad - \mu(1 + \gamma) \int_0^t e^{\mathcal{A}(t-t')} \begin{pmatrix} ((\bar{\kappa} + \kappa_0)^2 \xi_s)_s \\ 0 \end{pmatrix} dt' \end{aligned} \quad (66)$$

and show that Ψ admits a unique fixed point in a suitable function space. To construct such a function space, we first define the spaces

$$\begin{aligned} \mathcal{Y}_0 &= \{u \in C([0, T]; L^2(I)) : \|u\|_{\mathcal{Y}_0} < \infty\}, \quad \|\cdot\|_{\mathcal{Y}_0} := \sup_{t \in [0, T]} \|\cdot\|_{L^2(I)} \\ \mathcal{Y}_1 &= \{u \in C((0, T]; \dot{H}^1(I)) : \|u\|_{\mathcal{Y}_1} < \infty\}, \quad \|\cdot\|_{\mathcal{Y}_1} := \sup_{t \in [0, T]} \min\{t^{1/4}, 1\} \|\cdot\|_{\dot{H}^1(I)}. \end{aligned} \quad (67)$$

We close our contraction mapping argument for $(\bar{\kappa}, \xi)$ in $(\mathcal{Y}_0 \times \mathcal{Y}_0) \cap (\mathcal{Y}_1 \times \mathcal{Y}_1)$.

Given a function space $\mathcal{X} \times \mathcal{X}$, we will use the notation $B_M(\mathcal{X} \times \mathcal{X})$ to denote the closed ball in $\mathcal{X} \times \mathcal{X}$ of radius M , i.e.

$$B_M(\mathcal{X} \times \mathcal{X}) = \left\{ \begin{pmatrix} u \\ \phi \end{pmatrix} \in \mathcal{X} \times \mathcal{X} : \left\| \begin{pmatrix} u \\ \phi \end{pmatrix} \right\|_{\mathcal{X} \times \mathcal{X}} \leq M \right\}. \quad (68)$$

We first show that Ψ maps $B_{M_0}(\mathcal{Y}_0 \times \mathcal{Y}_0) \cap B_{M_1}(\mathcal{Y}_1 \times \mathcal{Y}_1)$ into itself for some $M_1, M_0 > 0$.

Since the nonlinear terms \mathcal{N} have the same form as in the Newtonian setting, from [26], we have

$$\begin{aligned} \|\mathcal{N}[\bar{\kappa}, \kappa_0]\|_{L^2(I)} &\leq c (\|\bar{\kappa}\|_{\dot{H}^1}^2 + \|\kappa_0\|_{\dot{H}^1}^2 + \|\bar{\tau}\|_{H^1}) (\|\bar{\kappa}\|_{\dot{H}^1} + \|\kappa_0\|_{\dot{H}^1}) \\ &\leq c \left((\|\bar{\kappa}\|_{\dot{H}^1}^3 + \|\kappa_0\|_{\dot{H}^1}^3) (\|\bar{\kappa}\|_{L^2} + \|\kappa_0\|_{L^2} + 1) + \|\xi\|_{\dot{H}^1} (\|\bar{\kappa}\|_{\dot{H}^1}^2 + \|\kappa_0\|_{\dot{H}^1}^2) \right). \end{aligned} \quad (69)$$

Here we have used the new viscoelastic tension estimate (Lemma 3.3) in the second line. Then, using Lemma 3.1, for $(\bar{\kappa}, \xi) \in B_{M_0}(\mathcal{Y}_0 \times \mathcal{Y}_0) \cap B_{M_1}(\mathcal{Y}_1 \times \mathcal{Y}_1)$, we may estimate the second forcing

term of (66) in $\dot{H}^m(I) \times \dot{H}^m(I)$, $m = 0, 1$, as

$$\begin{aligned}
& \left\| \int_0^t e^{\mathcal{A}(t-t')} \begin{pmatrix} (\mathcal{N}[\bar{\kappa}, \kappa_0])_s \\ 0 \end{pmatrix} dt' \right\|_{\dot{H}^m \times \dot{H}^m} \\
& \leq c \int_0^t \max\{(t-t')^{-(m+1)/4}, 1\} e^{-(t-t')\Lambda} \|\mathcal{N}[\bar{\kappa}, \kappa_0]\|_{L^2} dt' \\
& \leq c \int_0^t \max\{(t-t')^{-(m+1)/4}, 1\} \max\{(t')^{-3/4}, 1\} e^{-(t-t')\Lambda} dt' M_1^3 (M_0 + M_1 + 1) \\
& \leq c \max\{t^{-m/4}, 1\} M_1^3 (M_0 + M_1 + 1).
\end{aligned} \tag{70}$$

Here we have also taken $\sup_{t \in [0, T]} \|\kappa_0\|_{H^1(I)} \leq cM_1$. Furthermore, using Lemma 3.1, we may estimate the third forcing term of (66) in $\dot{H}^m(I) \times \dot{H}^m(I)$, $m = 0, 1$, by

$$\begin{aligned}
& \left\| \int_0^t e^{\mathcal{A}(t-t')} \begin{pmatrix} ((\bar{\kappa} + \kappa_0)^2 \xi_s)_s \\ 0 \end{pmatrix} dt' \right\|_{\dot{H}^m \times \dot{H}^m} \\
& \leq c \int_0^t \max\{(t-t')^{-(m+1)/4}, 1\} e^{-(t-t')\Lambda} \|\xi\|_{\dot{H}^1} (\|\bar{\kappa}\|_{L^2}^2 + \|\kappa_0\|_{L^2}^2) dt' \\
& \leq c \int_0^t \max\{(t-t')^{-(m+1)/4}, 1\} \max\{(t')^{-1/4}, 1\} e^{-(t-t')\Lambda} dt' M_1 (M_0^2 + M_1^2) \\
& \leq c M_1 (M_0^2 + M_1^2).
\end{aligned} \tag{71}$$

Finally, the forcing term involving $\dot{\kappa}_0$ may be estimated in $\dot{H}^m(I) \times \dot{H}^m(I)$, $m = 0, 1$, as

$$\begin{aligned}
& \left\| \int_0^t e^{\mathcal{A}(t-t')} \begin{pmatrix} \dot{\kappa}_0 \\ 0 \end{pmatrix} dt' \right\|_{\dot{H}^m \times \dot{H}^m} \leq c \int_0^t \max\{(t-t')^{-m/4}, 1\} e^{-(t-t')\Lambda} \|\dot{\kappa}_0\|_{L^2(I)} dt' \\
& \leq c \left(\sup_{t \in [0, T]} \|\dot{\kappa}_0\|_{L^2(I)} \right).
\end{aligned}$$

Combining the above three estimates and using Lemma 3.1 to estimate the initial data, we obtain the following $\mathcal{Y}_0 \times \mathcal{Y}_0$ bound:

$$\begin{aligned}
& \left\| \Psi \begin{bmatrix} \bar{\kappa} \\ \xi \end{bmatrix} \right\|_{\mathcal{Y}_0 \times \mathcal{Y}_0} \leq c \left(\left\| \begin{pmatrix} \bar{\kappa}^{\text{in}} \\ \xi^{\text{in}} \end{pmatrix} \right\|_{L^2 \times L^2} + M_1^3 (M_0 + M_1 + 1) \right. \\
& \quad \left. + M_1 (M_0^2 + M_1^2) + \sup_{t \in [0, T]} \|\dot{\kappa}_0\|_{L^2} \right) \leq M_0,
\end{aligned} \tag{72}$$

provided that we choose $M_0 = c_0 \left\| \begin{pmatrix} \bar{\kappa}^{\text{in}} \\ \xi^{\text{in}} \end{pmatrix} \right\|_{L^2 \times L^2}$ for c_0 small enough, $c \sup_{t \in [0, T]} \|\dot{\kappa}_0\|_{L^2} \leq M_0/4$, and M_1 small enough that $c(M_1^3(M_0 + M_1 + 1) + M_1(M_0^2 + M_1^2)) \leq M_0/4$.

We may also obtain the following $\mathcal{Y}_1 \times \mathcal{Y}_1$ bound for Ψ :

$$\begin{aligned}
& \left\| \Psi \begin{bmatrix} \bar{\kappa} \\ \xi \end{bmatrix} \right\|_{\mathcal{Y}_1 \times \mathcal{Y}_1} \leq \sup_{t \in [0, T]} \min\{t^{1/4}, 1\} \left\| e^{t\mathcal{A}} \begin{pmatrix} \bar{\kappa}^{\text{in}} \\ \xi^{\text{in}} \end{pmatrix} \right\|_{\dot{H}^1 \times \dot{H}^1} + c \left(M_1^3 (M_0 + M_1 + 1) \right. \\
& \quad \left. + M_1 (M_0^2 + M_1^2) + \sup_{t \in [0, T]} \|\dot{\kappa}_0\|_{L^2} \right) \\
& \leq \sup_{t \in [0, T]} \min\{t^{1/4}, 1\} \left\| e^{t\mathcal{A}} \begin{pmatrix} \bar{\kappa}^{\text{in}} \\ \xi^{\text{in}} \end{pmatrix} \right\|_{\dot{H}^1 \times \dot{H}^1} + \frac{M_1}{2},
\end{aligned} \tag{73}$$

provided that $c \sup_{t \in [0, T]} \|\dot{\kappa}_0\|_{L^2} \leq M_1/4$, and M_0 and M_1 are small enough that $c(M_1^3(M_0 + M_1 + 1) + M_1(M_0^2 + M_1^2)) \leq M_1/4$.

It remains to show that

$$\sup_{t \in [0, T]} \min\{t^{1/4}, 1\} \left\| e^{t\mathcal{A}} \begin{pmatrix} \bar{\kappa}^{\text{in}} \\ \xi^{\text{in}} \end{pmatrix} \right\|_{\dot{H}^1 \times \dot{H}^1} \leq \frac{M_1}{2}, \quad (74)$$

which we may achieve by either choosing a small time interval T or small initial data. For small time, we may use Lemma 3.2 to find $T_{M_1} > 0$ such that (74) holds. For small initial data, we may use Lemma 3.1 to obtain

$$\sup_{t \in [0, T]} \min\{t^{1/4}, 1\} \left\| e^{t\mathcal{A}} \begin{pmatrix} \bar{\kappa}^{\text{in}} \\ \xi^{\text{in}} \end{pmatrix} \right\|_{\dot{H}^1 \times \dot{H}^1} \leq c \left\| \begin{pmatrix} \bar{\kappa}^{\text{in}} \\ \xi^{\text{in}} \end{pmatrix} \right\|_{L^2 \times L^2},$$

and for sufficiently small $(\bar{\kappa}^{\text{in}}, \xi^{\text{in}})$, we may take $M_1 = c_1 \left\| \begin{pmatrix} \bar{\kappa}^{\text{in}} \\ \xi^{\text{in}} \end{pmatrix} \right\|_{L^2 \times L^2}$ to obtain the bound (74).

We next show that the map Ψ is a contraction on $B_{M_0}(\mathcal{Y}_0 \times \mathcal{Y}_0) \cap B_{M_1}(\mathcal{Y}_1 \times \mathcal{Y}_1)$. Given two pairs $(\bar{\kappa}_a, \xi_a), (\bar{\kappa}_b, \xi_b)$, we seek an estimate for $\Psi \left[\begin{pmatrix} \bar{\kappa}_a \\ \xi_a \end{pmatrix} \right] - \Psi \left[\begin{pmatrix} \bar{\kappa}_b \\ \xi_b \end{pmatrix} \right]$.

First, from [26], we may borrow the estimate

$$\begin{aligned} & \|\mathcal{N}[\bar{\kappa}_a(\cdot, t)] - \mathcal{N}[\bar{\kappa}_b(\cdot, t)]\|_{L^2(I)} \\ & \leq c \left(\|\bar{\kappa}_a - \bar{\kappa}_b\|_{\dot{H}^1} (\|\bar{\kappa}_a\|_{\dot{H}^1}^2 + \|\bar{\kappa}_b\|_{\dot{H}^1}^2 + \|\kappa_0\|_{\dot{H}^1}^2) \right. \\ & \quad \left. + \|\bar{\tau}_a - \bar{\tau}_b\|_{H^1} (\|\bar{\kappa}_a\|_{\dot{H}^1} + \|\kappa_0\|_{H^1}) + \|\bar{\kappa}_a - \bar{\kappa}_b\|_{\dot{H}^1} \|\bar{\tau}_b\|_{H^1} \right) \\ & \leq c \left(\|\bar{\kappa}_a - \bar{\kappa}_b\|_{\dot{H}^1} + \|\bar{\kappa}_a - \bar{\kappa}_b\|_{L^2} (\|\bar{\kappa}_a\|_{\dot{H}^1} + \|\kappa_0\|_{H^1}) \right) \left(\|\bar{\kappa}_a\|_{\dot{H}^1}^2 + \|\bar{\kappa}_b\|_{\dot{H}^1}^2 + \|\kappa_0\|_{\dot{H}^1}^2 \right. \\ & \quad \left. + (\|\xi_a\|_{\dot{H}^1} + \|\xi_b\|_{\dot{H}^1}) (\|\bar{\kappa}_a\|_{\dot{H}^1} + \|\bar{\kappa}_b\|_{\dot{H}^1} + \|\kappa_0\|_{H^1}) \right) (\|\bar{\kappa}_a\|_{L^2} + \|\bar{\kappa}_b\|_{L^2} + \|\kappa_0\|_{L^2} + 1)^2 \\ & \quad + c \|\xi_a - \xi_b\|_{\dot{H}^1} (\|\bar{\kappa}_a\|_{\dot{H}^1} + \|\kappa_0\|_{H^1}) (\|\bar{\kappa}_b\|_{\dot{H}^1} + \|\kappa_0\|_{H^1}). \end{aligned} \quad (75)$$

Here we have again used the new viscoelastic estimates of Lemma 3.3 to bound the tension in the second inequality. Furthermore, we have the following Lipschitz bound for the new viscoelastic nonlinear term:

$$\begin{aligned} & \|(\bar{\kappa}_a + \kappa_0)^2(\xi_a)_s - (\bar{\kappa}_b + \kappa_0)^2(\xi_b)_s\|_{L^2(I)} \\ & \leq \|\xi_a - \xi_b\|_{\dot{H}^1} (\|\bar{\kappa}_a\|_{\dot{H}^1}^2 + \|\kappa_0\|_{\dot{H}^1}^2) + \|\bar{\kappa}_a - \bar{\kappa}_b\|_{\dot{H}^1} \|\xi_b\|_{\dot{H}^1} (\|\bar{\kappa}_a\|_{\dot{H}^1} + \|\bar{\kappa}_b\|_{\dot{H}^1} + \|\kappa_0\|_{H^1}). \end{aligned} \quad (76)$$

Together, we may then obtain the Lipschitz estimate

$$\begin{aligned} & \left\| \Psi \left[\begin{pmatrix} \bar{\kappa}_a \\ \xi_a \end{pmatrix} \right] - \Psi \left[\begin{pmatrix} \bar{\kappa}_b \\ \xi_b \end{pmatrix} \right] \right\|_{\dot{H}^m \times \dot{H}^m} \\ & \leq c \int_0^t \max\{(t-t')^{-(m+1)/4}, 1\} e^{-(t-t')\Lambda} (\|\mathcal{N}[\bar{\kappa}_a(\cdot, t)] - \mathcal{N}[\bar{\kappa}_b(\cdot, t)]\|_{L^2(I)} \\ & \quad + \|(\bar{\kappa}_a + \kappa_0)^2(\xi_a)_s - (\bar{\kappa}_b + \kappa_0)^2(\xi_b)_s\|_{L^2(I)}) dt' \\ & \leq c \int_0^t \max\{(t-t')^{-(m+1)/4}, 1\} e^{-(t-t')\Lambda} \left(\|\bar{\kappa}_a - \bar{\kappa}_b\|_{\dot{H}^1} \right. \end{aligned}$$

$$\begin{aligned}
& + \|\bar{\kappa}_a - \bar{\kappa}_b\|_{L^2} (\|\bar{\kappa}_a\|_{\dot{H}^1} + \|\kappa_0\|_{H^1}) \Big) \Big((\|\xi_a\|_{\dot{H}^1} + \|\xi_b\|_{\dot{H}^1}) (\|\bar{\kappa}_a\|_{\dot{H}^1} + \|\bar{\kappa}_b\|_{\dot{H}^1} + \|\kappa_0\|_{H^1}) \\
& + \|\bar{\kappa}_a\|_{\dot{H}^1}^2 + \|\bar{\kappa}_b\|_{\dot{H}^1}^2 + \|\kappa_0\|_{H^1}^2 \Big) (\|\bar{\kappa}_a\|_{L^2} + \|\bar{\kappa}_b\|_{L^2} + \|\kappa_0\|_{L^2} + 1)^2 dt' \\
& + c \int_0^t \max\{(t-t')^{-(m+1)/4}, 1\} e^{-(t-t')\Lambda} \|\xi_a - \xi_b\|_{\dot{H}^1} (\|\bar{\kappa}_a\|_{\dot{H}^1}^2 + \|\bar{\kappa}_b\|_{\dot{H}^1}^2 + \|\kappa_0\|_{H^1}^2) dt' \\
& \leq c \int_0^t \max\{(t-t')^{-(m+1)/4}, 1\} \max\{(t')^{-3/4}, 1\} e^{-(t-t')\Lambda} dt' M_1^2 \Big(\|\xi_a - \xi_b\|_{\mathcal{Y}_1} \\
& + M_0^2 \|\bar{\kappa}_a - \bar{\kappa}_b\|_{\mathcal{Y}_1} + M_1 M_0^2 \|\bar{\kappa}_a - \bar{\kappa}_b\|_{\mathcal{Y}_0} \Big) \\
& \leq c \max\{t^{-m/4}, 1\} M_1^2 \Big(\|\xi_a - \xi_b\|_{\mathcal{Y}_1} + M_0^2 \|\bar{\kappa}_a - \bar{\kappa}_b\|_{\mathcal{Y}_1} + M_1 M_0^2 \|\bar{\kappa}_a - \bar{\kappa}_b\|_{\mathcal{Y}_0} \Big).
\end{aligned}$$

We thus have

$$\begin{aligned}
& \left\| \Psi \left[\begin{pmatrix} \bar{\kappa}_a \\ \xi_a \end{pmatrix} \right] - \Psi \left[\begin{pmatrix} \bar{\kappa}_b \\ \xi_b \end{pmatrix} \right] \right\|_{\mathcal{Y}_0 \times \mathcal{Y}_0} \\
& \leq c M_1^2 \left(\|\xi_a - \xi_b\|_{\mathcal{Y}_1} + M_0^2 \|\bar{\kappa}_a - \bar{\kappa}_b\|_{\mathcal{Y}_1} + M_1 M_0^2 \|\bar{\kappa}_a - \bar{\kappa}_b\|_{\mathcal{Y}_0} \right), \\
& \left\| \Psi \left[\begin{pmatrix} \bar{\kappa}_a \\ \xi_a \end{pmatrix} \right] - \Psi \left[\begin{pmatrix} \bar{\kappa}_b \\ \xi_b \end{pmatrix} \right] \right\|_{\mathcal{Y}_1 \times \mathcal{Y}_1} \\
& \leq c M_1^2 \left(\|\xi_a - \xi_b\|_{\mathcal{Y}_1} + M_0^2 \|\bar{\kappa}_a - \bar{\kappa}_b\|_{\mathcal{Y}_1} + M_1 M_0^2 \|\bar{\kappa}_a - \bar{\kappa}_b\|_{\mathcal{Y}_0} \right).
\end{aligned}$$

For sufficiently small $M_0, M_1 < 1$, we obtain a contraction on $B_{M_0}(\mathcal{Y}_0 \times \mathcal{Y}_0) \cap B_{M_1}(\mathcal{Y}_1 \times \mathcal{Y}_1)$, thus proving Theorem 1.1 for $\kappa_0 \not\equiv 0$.

If $\kappa_0 \equiv 0$, we may replace the norms in the definition (67) of \mathcal{Y}_0 and \mathcal{Y}_1 with the exponentially weighted norms

$$\|\cdot\|_{\widehat{\mathcal{Y}}_0} := \sup_{t \in [0, T]} e^{-t\Lambda} \|\cdot\|_{L^2(I)}, \quad \|\cdot\|_{\widehat{\mathcal{Y}}_1} := \sup_{t \in [0, T]} \min\{t^{1/4}, 1\} e^{-t\Lambda} \|\cdot\|_{\dot{H}^1(I)}, \quad (77)$$

where Λ is given by Lemma 3.1. We obtain analogous estimates to (72) and (73) in $\widehat{\mathcal{Y}}_0 \times \widehat{\mathcal{Y}}_0$ and $\widehat{\mathcal{Y}}_1 \times \widehat{\mathcal{Y}}_1$, except, crucially, no term depending on κ_0 , allowing for the desired time decay. \square

3.4. Existence of a unique periodic solution. We next consider solutions to the system (13)-(16) when the internal fiber forcing κ_0 is T -periodic in time. We prove Theorem 1.2 in two parts: in this section, we prove parts (a) and (b) on the existence of a unique periodic solution $(\bar{\kappa}, \xi)$, and in section 3.5 we show part (c) concerning the limiting behavior of this periodic solution as $\delta \rightarrow 0$.

To prove parts (a) and (b) of Theorem 1.2, we begin by considering the following linear PDE, where κ_0 and g are both given, T -periodic functions:

$$\begin{aligned}
\dot{\bar{\kappa}} &= -(1 + \mu)\bar{\kappa}_{ssss} + \mu\xi_{ssss} - \dot{\kappa}_0 + g_s \\
\delta\dot{\xi} &= -\xi + \bar{\kappa},
\end{aligned} \quad (78)$$

with boundary conditions as in (16). For the system (78), we show the following lemma.

Lemma 3.4. *There exists a constant $\varepsilon > 0$ such that, given a T -periodic $\kappa_0 \in C^1([0, T]; L^2(I))$ satisfying*

$$\sup_{t \in [0, T]} \|\dot{\kappa}_0\|_{L^2} = \varepsilon_1 \leq \varepsilon \quad (79)$$

and a T -periodic $g(s, t) \in C([0, T]; L^2(I))$ satisfying

$$\sup_{t \in [0, T]} \|g\|_{L^2} = \varepsilon_2 \leq \varepsilon, \quad (80)$$

there exists a unique T -periodic solution to (78) satisfying

$$\sup_{t \in [0, T]} \left\| \begin{pmatrix} \bar{\kappa} \\ \xi \end{pmatrix} \right\|_{H^1 \times H^1} \leq c \left(\sup_{t \in [0, T]} \|\dot{\kappa}_0\|_{L^2} + \sup_{t \in [0, T]} \|g\|_{L^2} \right). \quad (81)$$

Proof. We consider the map Ψ^T taking the initial data $(\bar{\kappa}^{\text{in}}, \xi^{\text{in}})$ to the solution to (78) at time T , which may be written as

$$\Psi^T \left[\begin{pmatrix} \bar{\kappa}^{\text{in}} \\ \xi^{\text{in}} \end{pmatrix} \right] = e^{T\mathcal{A}} \begin{pmatrix} \bar{\kappa}^{\text{in}} \\ \xi^{\text{in}} \end{pmatrix} - \int_0^T e^{(T-t')\mathcal{A}} \begin{pmatrix} \dot{\kappa}_0 \\ 0 \end{pmatrix} dt' + \int_0^T e^{(T-t')\mathcal{A}} \begin{pmatrix} g_s \\ 0 \end{pmatrix} dt'.$$

We show that Ψ^T maps $B_M(H^1 \times H^1)$ to itself, where $B_M(H^1 \times H^1)$ is as in (68). Using Lemma 3.1, for $m = 0, 1$, we have

$$\begin{aligned} \left\| \Psi^T \left[\begin{pmatrix} \bar{\kappa}^{\text{in}} \\ \xi^{\text{in}} \end{pmatrix} \right] \right\|_{\dot{H}^m \times \dot{H}^m} &\leq c e^{-T\Lambda} \left\| \begin{pmatrix} \bar{\kappa}^{\text{in}} \\ \xi^{\text{in}} \end{pmatrix} \right\|_{\dot{H}^m \times \dot{H}^m} + \int_0^T \left\| e^{(T-t')\mathcal{A}} \begin{pmatrix} \dot{\kappa}_0 \\ 0 \end{pmatrix} \right\|_{\dot{H}^m \times \dot{H}^m} dt' \\ &\quad + \int_0^T \left\| e^{(T-t')\mathcal{A}} \begin{pmatrix} g_s \\ 0 \end{pmatrix} \right\|_{\dot{H}^m \times \dot{H}^m} dt' \\ &\leq c e^{-T\Lambda} \left\| \begin{pmatrix} \bar{\kappa}^{\text{in}} \\ \xi^{\text{in}} \end{pmatrix} \right\|_{\dot{H}^m \times \dot{H}^m} \\ &\quad + c \int_0^T \max\{(T-t')^{-m/4}, 1\} e^{-(T-t')\Lambda} \|\dot{\kappa}_0\|_{L^2} dt' \\ &\quad + \int_0^T \max\{(T-t')^{-(m+1)/4}, 1\} e^{-(T-t')\Lambda} \|g\|_{L^2} dt' \\ &\leq c \left(e^{-T\Lambda} \left\| \begin{pmatrix} \bar{\kappa}^{\text{in}} \\ \xi^{\text{in}} \end{pmatrix} \right\|_{\dot{H}^m \times \dot{H}^m} + \sup_{t \in [0, T]} \|\dot{\kappa}_0\|_{L^2} + \sup_{t \in [0, T]} \|g\|_{L^2} \right). \end{aligned} \quad (82)$$

In particular, provided that the period T is large enough that $c e^{-T\Lambda} \leq \frac{1}{3}$, we may choose κ_0 and g such that $c \sup_{t \in [0, T]} \|\dot{\kappa}_0\|_{L^2} \leq \frac{M}{3}$ and $c \sup_{t \in [0, T]} \|g\|_{L^2} \leq \frac{M}{3}$ to obtain

$$\left\| \Psi^T \left[\begin{pmatrix} \bar{\kappa}^{\text{in}} \\ \xi^{\text{in}} \end{pmatrix} \right] \right\|_{H^1 \times H^1} \leq \left(\frac{M}{3} + \frac{M}{3} + \frac{M}{3} \right) \leq M. \quad (83)$$

Furthermore, again using Lemma 3.1, we may obtain the following Lipschitz estimate:

$$\begin{aligned} \left\| \Psi^T \left[\begin{pmatrix} \bar{\kappa}_a^{\text{in}} \\ \xi_a^{\text{in}} \end{pmatrix} \right] - \Psi^T \left[\begin{pmatrix} \bar{\kappa}_b^{\text{in}} \\ \xi_b^{\text{in}} \end{pmatrix} \right] \right\|_{H^1 \times H^1} &= \left\| e^{T\mathcal{A}} \begin{pmatrix} \bar{\kappa}_a^{\text{in}} - \bar{\kappa}_b^{\text{in}} \\ \xi_a^{\text{in}} - \xi_b^{\text{in}} \end{pmatrix} \right\|_{H^1 \times H^1} \\ &\leq c e^{-T\Lambda} \left\| \begin{pmatrix} \bar{\kappa}_a^{\text{in}} - \bar{\kappa}_b^{\text{in}} \\ \xi_a^{\text{in}} - \xi_b^{\text{in}} \end{pmatrix} \right\|_{H^1 \times H^1} \leq \frac{1}{4} \left\| \begin{pmatrix} \bar{\kappa}_a^{\text{in}} - \bar{\kappa}_b^{\text{in}} \\ \xi_a^{\text{in}} - \xi_b^{\text{in}} \end{pmatrix} \right\|_{H^1 \times H^1}, \end{aligned}$$

as long as the period T is sufficiently large. By the contraction mapping theorem, there exists a unique fixed point of the map Ψ^T , i.e. $\Psi^T \left[\begin{pmatrix} \bar{\kappa}^{\text{in}} \\ \xi^{\text{in}} \end{pmatrix} \right] = \begin{pmatrix} \bar{\kappa}^{\text{in}} \\ \xi^{\text{in}} \end{pmatrix}$, corresponding to a unique T -periodic solution $(\bar{\kappa}, \xi)$ to (78).

In addition, using (82) and (83), the T -periodic solution $(\bar{\kappa}, \xi)$ satisfies

$$\left\| \begin{pmatrix} \bar{\kappa}^{\text{in}} \\ \xi^{\text{in}} \end{pmatrix} \right\|_{H^1 \times H^1} \leq c \left(\sup_{t \in [0, T]} \|\dot{\kappa}_0\|_{L^2} + \sup_{t \in [0, T]} \|g\|_{L^2} \right). \quad (84)$$

To obtain the bound (81), we may use Duhamel's formula to write $(\bar{\kappa}, \xi)$ as

$$\begin{pmatrix} \bar{\kappa} \\ \xi \end{pmatrix} = e^{At} \begin{pmatrix} \bar{\kappa}^{\text{in}} \\ \xi^{\text{in}} \end{pmatrix} - \int_0^t e^{\mathcal{A}(t-t')} \begin{pmatrix} \dot{\kappa}_0 \\ 0 \end{pmatrix} dt' + \int_0^t e^{\mathcal{A}(t-t')} \begin{pmatrix} g_s \\ 0 \end{pmatrix} dt'.$$

Then, as for the time- T map (82), but now for any $t \in [0, T]$ and $m = 0, 1$, we have

$$\begin{aligned} \left\| \begin{pmatrix} \bar{\kappa} \\ \xi \end{pmatrix} \right\|_{\dot{H}^m \times \dot{H}^m} &\leq c e^{-t\Lambda} \left\| \begin{pmatrix} \bar{\kappa}^{\text{in}} \\ \xi^{\text{in}} \end{pmatrix} \right\|_{\dot{H}^m \times \dot{H}^m} + c \int_0^t \max\{(t-t')^{-m/4}, 1\} e^{-(t-t')\Lambda} \|\dot{\kappa}_0\|_{L^2} dt' \\ &\quad + \int_0^t \max\{(t-t')^{-(m+1)/4}, 1\} e^{-(t-t')\Lambda} \|g\|_{L^2} dt' \\ &\leq c \left(e^{-t\Lambda} \left\| \begin{pmatrix} \bar{\kappa}^{\text{in}} \\ \xi^{\text{in}} \end{pmatrix} \right\|_{\dot{H}^m \times \dot{H}^m} + \sup_{t \in [0, T]} \|\dot{\kappa}_0\|_{L^2} + \sup_{t \in [0, T]} \|g\|_{L^2} \right). \end{aligned}$$

Using (84), we obtain (81). \square

We now show parts (a) and (b) of Theorem 1.2.

Proof of Theorem 1.2, parts (a) & (b). We will use Lemma 3.4. Let $\mathcal{A}_T^{\text{per}}$ denote the solution operator mapping $(-\dot{\kappa}_0 + g_s, 0)^T$ to the unique periodic $(\bar{\kappa}, \xi)$:

$$\begin{pmatrix} \bar{\kappa} \\ \xi \end{pmatrix} = \mathcal{A}_T^{\text{per}} \left[\begin{pmatrix} -\dot{\kappa}_0 + g_s \\ 0 \end{pmatrix} \right].$$

We will consider $g(\bar{\kappa}, \xi, \kappa_0) = (1 + \mu)\mathcal{N}[\bar{\kappa}, \kappa_0] - \mu(1 + \gamma)(\bar{\kappa} + \kappa_0)^2 \xi_s$, i.e. the nonlinear terms from (13), and show that, given κ_0 , the operator $\mathcal{A}_T^{\text{per}}$ admits a unique fixed point in the space $\mathcal{X}_T \times \mathcal{X}_T$, where

$$\mathcal{X}_T := \{u \in C([0, T]; H^1(I)) : u \text{ is } T\text{-periodic}\}, \quad \|\cdot\|_{\mathcal{X}_T} := \sup_{t \in [0, T]} \|u\|_{H^1(I)}.$$

We show that $\mathcal{A}_T^{\text{per}}$ maps the ball $B_M(\mathcal{X}_T, \mathcal{X}_T)$ (68) to itself for some $M > 0$. For g as above, we first note that, by (69), we have

$$\|g\|_{L^2(I)} \leq c \left((\|\bar{\kappa}\|_{H^1}^3 + \|\kappa_0\|_{H^1}^3)(\|\bar{\kappa}\|_{L^2} + \|\kappa_0\|_{L^2} + 1) + \|\xi\|_{H^1} (\|\bar{\kappa}\|_{H^1}^2 + \|\kappa_0\|_{H^1}^2) \right). \quad (85)$$

Then, using Lemma 3.4, taking κ_0 such that $\|\kappa_0\|_{\mathcal{X}_T} = c_1 M$, for $(\bar{\kappa}, \xi) \in B_M(\mathcal{X}_T, \mathcal{X}_T)$ we have

$$\begin{aligned} \left\| \mathcal{A}_T^{\text{per}} \left[\begin{pmatrix} -\dot{\kappa}_0 + g_s \\ 0 \end{pmatrix} \right] \right\|_{\mathcal{X}_T \times \mathcal{X}_T} &\leq c \left(\sup_{t \in [0, T]} \|\dot{\kappa}_0\|_{L^2} + \|\bar{\kappa}\|_{\mathcal{X}_T}^4 + \|\kappa_0\|_{\mathcal{X}_T}^4 + \|\bar{\kappa}\|_{\mathcal{X}_T}^3 + \|\kappa_0\|_{\mathcal{X}_T}^3 \right. \\ &\quad \left. + \|\xi\|_{\mathcal{X}_T} (\|\bar{\kappa}\|_{\mathcal{X}_T}^2 + \|\kappa_0\|_{\mathcal{X}_T}^2) \right) \\ &\leq \left(\frac{M}{2} + c(M^4 + M^3) \right) \leq M, \end{aligned} \quad (86)$$

where we have taken $c \sup_{t \in [0, T]} \|\dot{\kappa}_0\|_{L^2} = c_2 M$ for $c_2 \leq \frac{1}{2}$ and M sufficiently small.

To show that $\mathcal{A}_T^{\text{per}}$ is a contraction on $B_M(\mathcal{X}_T, \mathcal{X}_T)$, we note that from (75) and (76), given two pairs $(\bar{\kappa}_a, \xi_a)$, $(\bar{\kappa}_b, \xi_b)$ and defining $g_a = g(\bar{\kappa}_a, \xi_a, \kappa_0)$, $g_b = g(\bar{\kappa}_b, \xi_b, \kappa_0)$, we have

$$\|g_a - g_b\|_{L^2} \leq c \left(\|\bar{\kappa}_a - \bar{\kappa}_b\|_{H^1} + \|\bar{\kappa}_a - \bar{\kappa}_b\|_{L^2} (\|\bar{\kappa}_a\|_{H^1} + \|\kappa_0\|_{H^1}) \right) \left(\|\bar{\kappa}_a\|_{H^1}^2 + \|\bar{\kappa}_b\|_{H^1}^2 + \|\kappa_0\|_{H^1}^2 \right)$$

$$\begin{aligned}
& + (\|\xi_a\|_{H^1} + \|\xi_b\|_{H^1}) (\|\bar{\kappa}_a\|_{H^1} + \|\bar{\kappa}_b\|_{H^1} + \|\kappa_0\|_{H^1}) \Big) (\|\bar{\kappa}_a\|_{L^2} + \|\bar{\kappa}_b\|_{L^2} + \|\kappa_0\|_{L^2} + 1)^2 \\
& + c \|\xi_a - \xi_b\|_{H^1} (\|\bar{\kappa}_a\|_{H^1}^2 + \|\bar{\kappa}_b\|_{H^1}^2 + \|\kappa_0\|_{H^1}^2).
\end{aligned}$$

We then have

$$\begin{aligned}
\left\| \mathcal{A}_T^{\text{per}} \left[\begin{pmatrix} (g_a - g_b)_s \\ 0 \end{pmatrix} \right] \right\|_{\mathcal{X}_T \times \mathcal{X}_T} & \leq c \left(M^2 (M^3 + 1) \|\bar{\kappa}_a - \bar{\kappa}_b\|_{\mathcal{X}_T} + M^2 \|\xi_a - \xi_b\|_{\mathcal{X}_T} \right) \\
& \leq \frac{1}{4} \left\| \begin{pmatrix} \bar{\kappa}_a - \bar{\kappa}_b \\ \xi_a - \xi_b \end{pmatrix} \right\|_{\mathcal{X}_T \times \mathcal{X}_T}
\end{aligned}$$

for M sufficiently small, yielding a contraction on $B_M(\mathcal{X}_T, \mathcal{X}_T)$.

For part (b) of Theorem 1.2, we note that $(\bar{\kappa}^{\text{lin}}, \xi^{\text{lin}})$ is the solution to (78) with $g = 0$. In particular, using the bound (85) on $g(\bar{\kappa}, \xi, \kappa_0) = (1 + \mu)\mathcal{N}[\bar{\kappa}, \kappa_0] - \mu(1 + \gamma)(\bar{\kappa} + \kappa_0)^2 \xi_s$ and the estimate (86), for the periodic solution $(\bar{\kappa}, \xi)$ of part (a), we have

$$\begin{aligned}
\left\| \begin{pmatrix} \bar{\kappa} - \bar{\kappa}^{\text{lin}} \\ \xi - \xi^{\text{lin}} \end{pmatrix} \right\|_{\mathcal{X}_T \times \mathcal{X}_T} & = \left\| \mathcal{A}_T^{\text{per}} \left[\begin{pmatrix} -\dot{\kappa}_0 + g_s \\ 0 \end{pmatrix} \right] - \mathcal{A}_T^{\text{per}} \left[\begin{pmatrix} -\dot{\kappa}_0 \\ 0 \end{pmatrix} \right] \right\|_{\mathcal{X}_T \times \mathcal{X}_T} \\
& \leq c \left(\|\bar{\kappa}\|_{\mathcal{X}_T}^4 + \|\kappa_0\|_{\mathcal{X}_T}^4 + \|\bar{\kappa}\|_{\mathcal{X}_T}^3 + \|\kappa_0\|_{\mathcal{X}_T}^3 + \|\xi\|_{\mathcal{X}_T} (\|\bar{\kappa}\|_{\mathcal{X}_T}^2 + \|\kappa_0\|_{\mathcal{X}_T}^2) \right) \\
& \leq c \varepsilon^3.
\end{aligned}$$

□

3.5. Small relaxation time limit. We next show part (c) of Theorem 1.2 concerning the behavior of the periodic solution $(\bar{\kappa}, \xi)$ of part (a) as the relaxation time $\delta \rightarrow 0$. To show that $(\bar{\kappa}, \xi)$ satisfies the estimate (33), we need the following lemma.

Lemma 3.5. *For $f \in L^2(I)$, let*

$$\begin{pmatrix} u_j \\ \phi_j \end{pmatrix} = e^{tA} \begin{pmatrix} \partial_s^j f \\ 0 \end{pmatrix} \quad j = 0, 1. \quad (87)$$

Then for $0 \leq m \leq 4 - j$ and $t \in (0, T]$, we have

$$\|u_j - \phi_j\|_{\dot{H}^m} \leq c \delta^{1-(j+m)/4} \sup_k \left(|\nu_k^-| e^{t\nu_k^-} + |\nu_k^+| e^{t\nu_k^+} \right) \|f\|_{L^2(I)} \quad (88)$$

where

$$\nu_k^\pm = \frac{1}{2\delta} \left(-(\delta(1 + \mu)\lambda_k + 1) \pm \sqrt{(\delta(1 + \mu)\lambda_k + 1)^2 - 4\delta\lambda_k} \right) \quad (89)$$

are the eigenvalues of the matrix \tilde{A}_k defined in (50).

Proof. It suffices to show that

$$\begin{pmatrix} w_j \\ \varphi_j \end{pmatrix} = \mathcal{L}^{j/4} e^{tA} \begin{pmatrix} f \\ 0 \end{pmatrix}$$

satisfies

$$\|w_j - \varphi_j\|_{\dot{H}^m} \leq c \delta^{1-(j+m)/4} \sup_k \left(|\nu_k^-| e^{t\nu_k^-} + |\nu_k^+| e^{t\nu_k^+} \right) \|f\|_{L^2(I)} ;$$

Lemma 3.5 then follows by a duality argument as in the proof of Lemma 3.1.

We begin by recalling the decomposition (53) in terms of eigenvectors of $\tilde{\mathcal{A}}_k$; in particular

$$\begin{pmatrix} 1 \\ 0 \end{pmatrix} = a_k^- \mathbf{v}_k^- + a_k^+ \mathbf{v}_k^+, \quad a_k^- \mathbf{v}_k^- = -\frac{1}{\delta(\nu_k^+ - \nu_k^-)} \begin{pmatrix} 1 + \delta\nu_k^- \\ 1 \end{pmatrix}, \quad a_k^+ \mathbf{v}_k^+ = \frac{1}{\delta(\nu_k^+ - \nu_k^-)} \begin{pmatrix} 1 + \delta\nu_k^+ \\ 1 \end{pmatrix},$$

where

$$\delta(\nu_k^+ - \nu_k^-) = \sqrt{(\delta(1 + \mu)\lambda_k + 1)^2 - 4\delta\lambda_k}.$$

Note that for $0 \leq r \leq 1$, we have

$$\frac{(\delta\lambda_k)^r}{\delta(\nu_k^+ - \nu_k^-)} = \frac{(\delta\lambda_k)^r}{\sqrt{(1 - \delta\lambda_k)^2 + 2\mu\delta\lambda_k + (2\mu + \mu^2)\delta^2\lambda_k^2}} \leq c$$

for c independent of both δ and λ_k . Letting $a_k^\pm v_{k,(1)}^\pm, a_k^\pm v_{k,(2)}^\pm$ denote the first and second component, respectively, of the vectors $a_k^\pm \mathbf{v}_k^\pm$, for $0 \leq r \leq 1$, we then have

$$\begin{aligned} \left| a_k^- (v_{k,(1)}^- - v_{k,(2)}^-) \right| \lambda_k^r &= \left| \frac{\delta\nu_k^-}{\delta(\nu_k^+ - \nu_k^-)} \right| \lambda_k^r \leq c \delta^{1-r} |\nu_k^-|, \\ \left| a_k^+ (v_{k,(1)}^+ - v_{k,(2)}^+) \right| \lambda_k^r &= \left| \frac{\delta\nu_k^+}{\delta(\nu_k^+ - \nu_k^-)} \right| \lambda_k^r \leq c \delta^{1-r} |\nu_k^+|. \end{aligned}$$

Using the decomposition (53), we then have

$$\begin{aligned} \|w_j - \varphi_j\|_{\dot{H}^m} &\leq \left\| \sum_{k=1}^{\infty} \left(a_k^- (v_{k,(1)}^- - v_{k,(2)}^-) e^{t\nu_k^-} + a_k^+ (v_{k,(1)}^+ - v_{k,(2)}^+) e^{t\nu_k^+} \right) \lambda_k^{(j+m)/4} \tilde{f}_k \psi_k \right\|_{L^2} \\ &\leq c \delta^{1-(j+m)/4} \sup_k \left(|\nu_k^-| e^{t\nu_k^-} + |\nu_k^+| e^{t\nu_k^+} \right) \left\| \sum_{k=1}^{\infty} \tilde{f}_k \psi_k \right\|_{L^2}. \end{aligned}$$

□

Using Lemma 3.5, we may now show part (c) of Theorem 1.2.

Proof of Theorem 1.2, part (c). We consider the T -periodic solution $(\bar{\kappa}, \xi)$ of part (a), which satisfies the bound (30). Note that c in (30) is bounded independent of δ as $\delta \rightarrow 0$, due to the δ -independence of the constant c in Lemma 3.1.

By T -periodicity, we have that $(\bar{\kappa}, \xi)$ at time $t \in [0, T]$ may be written

$$\begin{pmatrix} \bar{\kappa} \\ \xi \end{pmatrix} = e^{(t+NT)\mathcal{A}} \begin{pmatrix} \bar{\kappa}^{\text{in}} \\ \xi^{\text{in}} \end{pmatrix} + \begin{pmatrix} \bar{\kappa}^{\text{f}} \\ \xi^{\text{f}} \end{pmatrix}, \quad \begin{pmatrix} \bar{\kappa}^{\text{f}} \\ \xi^{\text{f}} \end{pmatrix} := \int_0^{t+NT} e^{(t+NT-t')\mathcal{A}} \begin{pmatrix} -\dot{\kappa}_0 + g_s \\ 0 \end{pmatrix} dt'$$

for any $N \in \mathbb{N}$, where $g(\bar{\kappa}, \xi, \kappa_0) = (1 + \mu)\mathcal{N}[\bar{\kappa}, \kappa_0] - \mu(1 + \gamma)(\bar{\kappa} + \kappa_0)^2 \xi_s$. Recall that by (85) and the estimate (30) on $(\bar{\kappa}, \xi)$, we have that $\sup_{t \in [0, T]} \|g\|_{L^2(I)} \leq c(\sup_{t \in [0, T]} \|\dot{\kappa}_0\|_{L^2} + \sup_{t \in [0, T]} \|\kappa_0\|_{H^1})$.

By Lemma 3.1, for $m = 0, 1$ we have

$$\left\| e^{(t+NT)\mathcal{A}} \begin{pmatrix} \bar{\kappa}^{\text{in}} \\ \xi^{\text{in}} \end{pmatrix} \right\|_{\dot{H}^m \times \dot{H}^m} \leq c e^{-(t+NT)\Lambda} \left\| \begin{pmatrix} \bar{\kappa}^{\text{in}} \\ \xi^{\text{in}} \end{pmatrix} \right\|_{\dot{H}^m \times \dot{H}^m}$$

for c independent of δ and $\Lambda = \min\{\lambda_1, \frac{1}{\delta(1+\mu)}\}$. In particular, for sufficiently small δ , we have $\Lambda = \lambda_1$. Note that since $(\bar{\kappa}, \xi) \in H^1 \times H^1$, we use $\dot{H}^m \times \dot{H}^m$, $m = 0, 1$ on the right hand side. For any (small) δ , we may choose $N = N_\delta$ large enough that

$$\left\| e^{(t+N_\delta T)\mathcal{A}} \begin{pmatrix} \bar{\kappa}^{\text{in}} \\ \xi^{\text{in}} \end{pmatrix} \right\|_{H^1 \times H^1} \leq \delta. \quad (90)$$

Furthermore, for $N = N_\delta$ as above, by Lemma 3.5 we may estimate the difference $\bar{\kappa}^f - \xi^f$ as

$$\begin{aligned} \left\| \bar{\kappa}^f - \xi^f \right\|_{\dot{H}^m(I)} &\leq c \delta^{1-m/4} \int_0^{t+N_\delta T} \sup_k \left(|\nu_k^-| e^{t\nu_k^-} + |\nu_k^+| e^{t\nu_k^+} \right) \|\dot{\kappa}_0\|_{L^2(I)} dt' \\ &\quad + c \delta^{1-(1+m)/4} \int_0^{t+N_\delta T} \sup_k \left(|\nu_k^-| e^{t\nu_k^-} + |\nu_k^+| e^{t\nu_k^+} \right) \|g\|_{L^2(I)} dt' \\ &\leq c \delta^{1-(1+m)/4} \left((1 + \delta^{1/4}) \sup_{t \in [0, T]} \|\dot{\kappa}_0\|_{L^2} + \sup_{t \in [0, T]} \|\kappa_0\|_{H^1} \right). \end{aligned} \quad (91)$$

Here we have integrated in time and used the T -periodicity of both κ_0 and g to take the supremum only over time $t \in [0, T]$. Combining (90) and (91), as $\delta \rightarrow 0$ we obtain

$$\|\bar{\kappa} - \xi\|_{H^1(I)} \leq \delta + \left\| \bar{\kappa}^f - \xi^f \right\|_{H^1(I)} \leq \delta + c \delta^{1/2} \left(\sup_{t \in [0, T]} \|\dot{\kappa}_0\|_{L^2} + \sup_{t \in [0, T]} \|\kappa_0\|_{H^1} \right).$$

□

4. VISCOELASTIC SWIMMING

In this section we give a proof of the fiber swimming expressions in Theorem 1.3. We will first need a brief lemma. A version of this lemma also appears in the Newtonian case [26] and states that, given some additional regularity on our (small) κ_0 , we can ensure that the fiber frame $(\mathbf{e}_t, \mathbf{e}_n)$ is not varying much over time.

Lemma 4.1. *Suppose that $\kappa_0 \in C^1([0, T]; H^3(I))$ is T -periodic and satisfies*

$$\sup_{t \in [0, T]} \|\dot{\kappa}_0\|_{L^2} = \varepsilon_1 \leq \varepsilon, \quad \sup_{t \in [0, T]} \|\kappa_0\|_{H^1} = \varepsilon_2 \leq \varepsilon,$$

for some $0 < \varepsilon < 1$, and let $(\bar{\kappa}, \xi)$ be the corresponding T -periodic solution to (13)-(16). The evolution of the fiber tangent vector \mathbf{e}_t (19) then satisfies

$$\sup_{t \in [0, T]} \|\mathbf{e}_t(\cdot, t) - \mathbf{e}_t(0, 0)\|_{L^2(I)} \leq c\varepsilon. \quad (92)$$

Proof. Since $\kappa_0 \in H^3(I)$, we may use estimates (70) and (71) for the Duhamel formula (25) for $(\bar{\kappa}, \xi)$ along with Lemma 3.1 to show

$$\sup_{t \in [0, T]} \min\{t^{m/4}, 1\} \left\| \begin{pmatrix} \bar{\kappa} \\ \xi \end{pmatrix} \right\|_{\dot{H}^m \times \dot{H}^m} \leq c \left\| \begin{pmatrix} \bar{\kappa}^{\text{in}} \\ \xi^{\text{in}} \end{pmatrix} \right\|_{L^2 \times L^2}, \quad 0 \leq m \leq 3.$$

Due to the T -periodicity of κ_0 , we in fact have

$$\sup_{t \in [0, T]} \left\| \begin{pmatrix} \bar{\kappa} \\ \xi \end{pmatrix} \right\|_{\dot{H}^m \times \dot{H}^m} \leq c \left\| \begin{pmatrix} \bar{\kappa}^{\text{in}} \\ \xi^{\text{in}} \end{pmatrix} \right\|_{L^2 \times L^2} \leq c\varepsilon, \quad 0 \leq m \leq 3.$$

Using the equation (18) for $\dot{\theta}$ in the frame evolution (19), we thus have

$$\begin{aligned} \sup_{t \in [0, T]} \|\mathbf{e}_t(\cdot, t) - \mathbf{e}_t(\cdot, 0)\|_{L^2(I)} &\leq c \sup_{t \in [0, T]} \|\dot{\theta}\|_{L^2(I)} \\ &\leq c \sup_{t \in [0, T]} \left(\|\bar{\kappa}_{ss}\|_{L^2(I)} + \|\dot{\kappa}_0\|_{L^2(I)} + \|\xi_{sss}\|_{L^2(I)} + \varepsilon^2 \right) \\ &\leq c\varepsilon. \end{aligned}$$

In addition, since $(\mathbf{e}_t)_s = \kappa \mathbf{e}_n$, we have

$$\|\mathbf{e}_t(\cdot, 0) - \mathbf{e}_t(0, 0)\|_{L^2(I)} \leq c \|\kappa\|_{L^2(I)} \leq c\varepsilon.$$

Together, these two estimates give Lemma 4.1. □

Equipped with Lemma 4.1, we may now prove Theorem 1.3.

Proof of Theorem 1.3. It will be convenient to define the difference $z := \bar{\kappa} - \xi$ and work in terms of $\bar{\kappa}$ and z rather than $\bar{\kappa}$ and ξ . Using the definition of z and the equation (1) for $\frac{\partial \mathbf{X}}{\partial t}$, we may calculate the velocity of the swimming fiber as

$$\mathbf{V}(t) = \int_0^1 \frac{\partial \mathbf{X}}{\partial t}(s, t) ds = \mathbf{V}_{\text{vis}}(t) + \mathbf{V}_{\text{ve}}(t),$$

where \mathbf{V}_{vis} and \mathbf{V}_{ve} are given by

$$\begin{aligned} \mathbf{V}_{\text{vis}}(t) &:= - \int_0^1 (\mathbf{I} + \gamma \mathbf{e}_t \mathbf{e}_t^T) \left(-(\bar{\kappa}^2 + 2\bar{\kappa}\kappa_0) \mathbf{e}_t + \bar{\kappa}_s \mathbf{e}_n - \bar{\tau} \mathbf{e}_t \right)_s ds \\ &= \gamma \int_0^1 (3\bar{\kappa}\bar{\kappa}_s + 3\bar{\kappa}_s \kappa_0 + 2\bar{\kappa}(\kappa_0)_s + \bar{\tau}_s) \mathbf{e}_t ds; \\ \mathbf{V}_{\text{ve}}(t) &:= -\mu \int_0^1 (\mathbf{I} + \gamma \mathbf{e}_t \mathbf{e}_t^T) \left(-(\bar{\kappa}^2 + 2\bar{\kappa}\kappa_0) \mathbf{e}_t - \bar{\tau} \mathbf{e}_t + z_s \mathbf{e}_n \right)_s ds \\ &= \gamma\mu \int_0^1 ((\bar{\kappa}^2 + 2\bar{\kappa}\kappa_0)_s + \bar{\tau}_s + z_s \bar{\kappa} + z_s \kappa_0) \mathbf{e}_t ds. \end{aligned}$$

Then, using Lemma 4.1 along with the vanishing boundary conditions for $\bar{\kappa}$, $\bar{\tau}$, and z , for small κ_0 we may write

$$\begin{aligned} \mathbf{V}_{\text{vis}}(t) &= \gamma \int_0^1 \kappa_0 \bar{\kappa}_s \mathbf{e}_t(0, 0) ds + \mathbf{r}_{\text{vis}}(t) = -\gamma \int_0^1 (\kappa_0)_s \bar{\kappa} ds \mathbf{e}_t(0, 0) + \mathbf{r}_{\text{vis}}(t), \\ \mathbf{V}_{\text{ve}}(t) &= -\gamma\mu \int_0^1 z(\bar{\kappa} + \kappa_0)_s ds \mathbf{e}_t(0, 0) + \mathbf{r}_{\text{ve}}(t), \end{aligned}$$

where both

$$|\mathbf{r}_{\text{vis}}(t)| \leq c\varepsilon^3, \quad |\mathbf{r}_{\text{ve}}(t)| \leq c\varepsilon^3.$$

We thus have

$$\mathbf{V}(t) = \left(-\gamma \int_0^1 (\kappa_0)_s \bar{\kappa} ds - \gamma\mu \int_0^1 z(\bar{\kappa} + \kappa_0)_s ds \right) \mathbf{e}_t(0, 0) + \mathbf{r}_{\text{vis}}(t) + \mathbf{r}_{\text{ve}}(t),$$

from which we obtain the first swimming expression (36)-(37).

To obtain the second expression (38), we first note that by Theorem 1.2, part (b), we may approximate $\bar{\kappa}$ and $z = \bar{\kappa} - \xi$ by $\bar{\kappa}^{\text{lin}}$ and $z^{\text{lin}} = \bar{\kappa}^{\text{lin}} - \xi^{\text{lin}}$, the T -periodic solutions to the linear equations

$$\begin{aligned} \dot{\bar{\kappa}}^{\text{lin}} &= -\mathcal{L}\bar{\kappa}^{\text{lin}} - \mu\mathcal{L}z^{\text{lin}} - \dot{\kappa}_0 \\ \dot{z}^{\text{lin}} &= \dot{\bar{\kappa}}^{\text{lin}} - \delta^{-1}z^{\text{lin}}. \end{aligned} \tag{93}$$

More specifically, defining

$$U^{\text{lin}} = -\gamma \int_0^1 (\kappa_0)_s \bar{\kappa}^{\text{lin}} ds - \gamma\mu \int_0^1 z^{\text{lin}}(\bar{\kappa}^{\text{lin}} + \kappa_0)_s ds, \tag{94}$$

we have that

$$\begin{aligned} |U - U^{\text{lin}}| &\leq c \int_0^1 |(\kappa_0)_s| |\bar{\kappa} - \bar{\kappa}^{\text{lin}}| ds + \int_0^1 |z - z^{\text{lin}}| (|\bar{\kappa}_s| + |(\kappa_0)_s|) ds \\ &\quad + \int_0^1 |z^{\text{lin}}| |(\bar{\kappa} - \bar{\kappa}^{\text{lin}})_s| ds \\ &\leq c \|\kappa_0\|_{H^1} \|\bar{\kappa} - \bar{\kappa}^{\text{lin}}\|_{L^2} + \|z - z^{\text{lin}}\|_{L^2} (\|\bar{\kappa}\|_{H^1} + \|\kappa_0\|_{H^1}) + \|z^{\text{lin}}\|_{L^2} \|\bar{\kappa} - \bar{\kappa}^{\text{lin}}\|_{H^1} \\ &\leq c\varepsilon^4. \end{aligned}$$

Therefore, it suffices to use $\bar{\kappa}^{\text{lin}}$ and z^{lin} to compute a more detailed expression for the time-averaged swimming speed $\langle U \rangle$. We begin by solving for $\bar{\kappa}^{\text{lin}}$ and z^{lin} in terms of κ_0 . Defining $\omega = \frac{2\pi}{T}$, we expand each of κ_0 , $\bar{\kappa}^{\text{lin}}$, and z^{lin} as a Fourier series in time:

$$\begin{aligned}\kappa_0 &= \sum_{m=1}^{\infty} A_m(s) \cos(\omega m t) - B_m(s) \sin(\omega m t), & \bar{\kappa}^{\text{lin}} &= \sum_{m=1}^{\infty} C_m(s) \cos(\omega m t) - D_m(s) \sin(\omega m t), \\ z^{\text{lin}} &= \sum_{m=1}^{\infty} E_m(s) \cos(\omega m t) - F_m(s) \sin(\omega m t).\end{aligned}$$

Using (93), the coefficients of this expansion then satisfy the following system of equations:

$$\begin{aligned}-\omega m C_m &= \mathcal{L} D_m + \mu \mathcal{L} F_m + \omega m A_m, & -\omega m E_m &= -\omega m C_m + \delta^{-1} F_m, \\ -\omega m D_m &= -\mathcal{L} C_m - \mu \mathcal{L} E_m + \omega m B_m, & -\omega m F_m &= -\omega m D_m - \delta^{-1} E_m.\end{aligned}$$

Further expanding each of these coefficients in eigenfunctions (22) of the operator \mathcal{L} , i.e.

$$\begin{aligned}A_m &= \sum_{k=1}^{\infty} a_{m,k} \psi_k, & B_m &= \sum_{k=1}^{\infty} b_{m,k} \psi_k, & C_m &= \sum_{k=1}^{\infty} c_{m,k} \psi_k, \\ D_m &= \sum_{k=1}^{\infty} d_{m,k} \psi_k, & E_m &= \sum_{k=1}^{\infty} e_{m,k} \psi_k, & F_m &= \sum_{k=1}^{\infty} f_{m,k} \psi_k,\end{aligned}$$

we may solve for the coefficients $c_{m,k}$ and $d_{m,k}$ as

$$c_{m,k} = Q_{m,k} b_{m,k} - H_{m,k} a_{m,k}, \quad d_{m,k} = -Q_{m,k} a_{m,k} - H_{m,k} b_{m,k},$$

where

$$\begin{aligned}Q_{m,k} &= \frac{\lambda_k \omega m (1 + (1 + \mu)(\delta \omega m)^2)}{\lambda_k^2 (1 + (1 + \mu)^2 (\delta \omega m)^2) + \omega^2 m^2 (2\mu \delta \lambda_k + 1 + (\delta \omega m)^2)}, \\ H_{m,k} &= \frac{\omega^2 m^2 (\mu \delta \lambda_k + 1 + (\delta \omega m)^2)}{\lambda_k^2 (1 + (1 + \mu)^2 (\delta \omega m)^2) + \omega^2 m^2 (2\mu \delta \lambda_k + 1 + (\delta \omega m)^2)}.\end{aligned}$$

Additionally we may solve for $e_{m,k}$ and $f_{m,k}$ as

$$\begin{aligned}e_{m,k} &= \frac{\delta \omega m}{1 + (\delta \omega m)^2} \left(Q_{m,k} (a_{m,k} + \delta \omega m b_{m,k}) - H_{m,k} (b_{m,k} - \delta \omega m a_{m,k}) \right) \\ f_{m,k} &= \frac{\delta \omega m}{1 + (\delta \omega m)^2} \left(Q_{m,k} (b_{m,k} - \delta \omega m a_{m,k}) + H_{m,k} (-a_{m,k} - \delta \omega m b_{m,k}) \right).\end{aligned}$$

We now need to use the above expansions in the expression (94) to calculate the average swimming speed $\langle U^{\text{lin}} \rangle$. We first calculate

$$\begin{aligned}-\gamma \int_0^1 \langle (\kappa_0)_s \bar{\kappa}^{\text{lin}} \rangle ds &= -\gamma \sum_{m,k,\ell=1}^{\infty} \left(\frac{1}{2} a_{m,\ell} c_{m,k} + \frac{1}{2} b_{m,\ell} d_{m,k} \right) \int_0^1 \psi_k(\psi_\ell)_s ds \\ &= \frac{\gamma}{2} \sum_{m,k,\ell=1}^{\infty} \left(Q_{m,k} (a_{m,k} b_{m,\ell} - a_{m,\ell} b_{m,k}) + H_{m,k} (a_{m,k} a_{m,\ell} + b_{m,k} b_{m,\ell}) \right) \int_0^1 \psi_k(\psi_\ell)_s ds.\end{aligned}$$

Furthermore, noting that

$$(\bar{\kappa}^{\text{lin}} + \kappa_0)_s = \sum_{m,\ell} (\psi_\ell)_s ((a_{m,\ell} + c_{m,\ell}) \cos(\omega m t) - (b_{m,\ell} + d_{m,\ell}) \sin(\omega m t)),$$

we may also calculate

$$\begin{aligned}
-\gamma\mu \int_0^1 \langle z^{\text{lin}}(\bar{\kappa}^{\text{lin}} + \kappa_0)_s \rangle ds &= -\frac{\gamma\mu}{2} \sum_{m,k,\ell} \left((a_{m,\ell} + c_{m,\ell})e_{m,k} + (b_{m,\ell} + d_{m,\ell})f_{m,k} \right) \int_0^1 \psi_k(\psi_\ell)_s ds \\
&= -\frac{\gamma\mu}{2} \sum_{m,k,\ell} \frac{\delta\omega m}{1 + (\delta\omega m)^2} \left((Q_{m,\ell}b_{m,\ell} + (1 - H_{m,\ell})a_{m,\ell}) \left(Q_{m,k}(a_{m,k} + \delta\omega mb_{m,k}) \right. \right. \\
&\quad \left. \left. - H_{m,k}(b_{m,k} - \delta\omega ma_{m,k}) \right) + (b_{m,\ell} - Q_{m,\ell}a_{m,\ell} - H_{m,\ell}b_{m,\ell}) \left(Q_{m,k}(b_{m,k} - \delta\omega ma_{m,k}) \right. \right. \\
&\quad \left. \left. - H_{m,k}(a_{m,k} + \delta\omega mb_{m,k}) \right) \right) \int_0^1 \psi_k(\psi_\ell)_s ds.
\end{aligned}$$

Rearranging the above expression and combining the two components of $\langle U^{\text{lin}} \rangle$, we obtain the form of the swimming speed reported in (38) and (39). \square

APPENDIX A. NUMERICAL METHOD

For the numerical simulations of section 2, we use the formulation introduced in [26], which readily adapts to the viscoelastic setting. The formulation is based on a combination of works by Moreau, et al. [23] and Maxian, et al. [22]. For convenience, we recall the original formulation (1)-(4) of the viscoelastic resistive force theory equations:

$$\begin{aligned}
\frac{\partial \mathbf{X}}{\partial t}(s, t) &= -(1 + \mu)(\mathbf{I} + \gamma \mathbf{X}_s \mathbf{X}_s^T)(\mathbf{X}_{sss} - \tau \mathbf{X}_s - (\kappa_0)_s \mathbf{e}_n - \frac{\mu}{1 + \mu} \xi_s \mathbf{e}_n)_s \\
\delta \dot{\xi} &= -\xi + \kappa - \kappa_0 \\
|\mathbf{X}_s|^2 &= 1 \\
(\mathbf{X}_{ss} - \kappa_0 \mathbf{e}_n)|_{s=0,1} &= 0, \quad (\mathbf{X}_{sss} - \tau \mathbf{X}_s - (\kappa_0)_s \mathbf{e}_n)|_{s=0,1} = 0, \quad \xi|_{s=0,1} = \xi_s|_{s=0,1} = 0.
\end{aligned} \tag{95}$$

The formulation used in numerical simulations will be derived from (95). We begin by parameterizing the filament using the tangent angle description (11). In particular, we write

$$\mathbf{X}(s, t) = \mathbf{X}_0(t) + \int_0^s \mathbf{e}_t(s', t) ds', \quad \mathbf{e}_t = \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix}, \quad \mathbf{e}_n = \begin{pmatrix} -\sin \theta \\ \cos \theta \end{pmatrix} \tag{96}$$

As in [26, 23, 22], we exploit that, due to the inextensibility constraint, only the normal components of the hydrodynamic force along the filament actually contribute to the fiber motion. In particular, using the parameterization (96), we may rewrite the first three equations of (95) as a closed system:

$$\dot{\mathbf{X}}_0 + \int_0^s \dot{\mathbf{e}}_t(s') ds' = -(1 + \mu)(\mathbf{I} + \gamma \mathbf{e}_t \mathbf{e}_t^T) \mathbf{h}(s) \tag{97}$$

$$(\mathbf{I} - \mathbf{e}_t(s) \mathbf{e}_t(s)^T) \int_0^s \mathbf{h}(s') ds' = (\mathbf{I} - \mathbf{e}_t(s) \mathbf{e}_t(s)^T) \left(\mathbf{X}_{sss} - (\kappa_0)_s \mathbf{e}_n - \frac{\mu}{1 + \mu} \xi_s \mathbf{e}_n \right) \tag{98}$$

$$\delta \dot{\xi} = -\xi + \kappa - \kappa_0. \tag{99}$$

Note that in (98), by projecting away from the tangential direction along the filament, we have eliminated the need to solve for the unknown fiber tension. Instead, inextensibility is enforced directly via the parameterization (96).

Solving (97) directly for \mathbf{h} and inserting this expression in (98), we obtain the system

$$\mathbf{e}_n(s, t) \cdot \int_0^s \left(\mathbf{I} - \frac{\gamma}{1 + \gamma} \mathbf{e}_t \mathbf{e}_t^T \right) \left(\dot{\mathbf{X}}_0 + \int_0^{s'} \dot{\mathbf{e}}_t(\bar{s}) d\bar{s} \right) ds' = -(1 + \mu) \theta_{ss} + (1 + \mu)(\kappa_0)_s + \mu \xi_s \tag{100}$$

$$\delta \dot{\xi} = -\xi + \theta_s - \kappa_0 \tag{101}$$

for unknowns $\mathbf{X}_0(t)$, $\theta(s, t)$, and $\xi(s, t)$. Equations (100) and (101) serve as the basis for our numerical method. The boundary conditions $(\theta_s - \kappa_0)|_{s=0,1} = 0$ are enforced directly in the discretization of θ_{ss} on the right hand side of (100), while $\xi|_{s=0,1} = \xi_s|_{s=0,1} = 0$ is enforced in the discretization of ξ in (101). To enforce the boundary condition $(-\theta_{ss} + (\kappa_0)_s)|_{s=1} = 0$, we will also need to require

$$\int_0^1 \left(\mathbf{I} - \frac{\gamma}{1+\gamma} \mathbf{e}_t \mathbf{e}_t^T \right) \left(\dot{\mathbf{X}}_0 + \int_0^s \dot{\mathbf{e}}_t(s') ds' \right) ds = 0. \quad (102)$$

The analogous condition at $s = 0$ is then satisfied automatically via the formulation (100).

We discretize the arclength coordinate $s \in [0, 1]$ into $N + 1$ equally spaced points s_i , $i = 0, \dots, N$ and define $\mathbf{X}_i = \mathbf{X}(s_i)$. We consider the fiber as N straight segments between each \mathbf{X}_i , and define θ_i , $i = 1, \dots, N$, to be the angle between segment i and the x -axis.

The evolution equation (100) is enforced at the midpoint of each segment $\mathbf{X}_{i-\frac{1}{2}} := \frac{\mathbf{X}_{i-1} + \mathbf{X}_i}{2}$, $i = 1, \dots, N$. In particular, we parameterize the evolution $\dot{\mathbf{X}}_{i-\frac{1}{2}}$ of each fiber segment as

$$\dot{\mathbf{X}}_{i-\frac{1}{2}} = \begin{pmatrix} \dot{x}_0 \\ \dot{y}_0 \end{pmatrix} + \frac{1}{2N} \begin{pmatrix} -\sin \theta_i \\ \cos \theta_i \end{pmatrix} \dot{\theta}_i + \frac{1}{N} \sum_{k=1}^i \begin{pmatrix} -\sin \theta_k \\ \cos \theta_k \end{pmatrix} \dot{\theta}_k, \quad i = 1, \dots, N.$$

We also define $\kappa_{0,i} = \kappa_0(s_{i-\frac{1}{2}})$ where $s_{i-\frac{1}{2}} = \frac{s_{i-1} + s_i}{2}$, $i = 1, \dots, N$, and we define ξ_i similarly.

For the middle segments $j = 2, \dots, N - 1$, we obtain $2(N - 2)$ equations from the discretization of (100) and (101):

$$\frac{1}{N} \begin{pmatrix} -\sin \theta_j \\ \cos \theta_j \end{pmatrix} \cdot \sum_{i=1}^j \mathbf{M}_{\text{RFT}}(\theta_i) \dot{\mathbf{X}}_{i-\frac{1}{2}} = -N^2(1 + \mu)(\theta_{j-1} - 2\theta_j + \theta_{j+1}) + (1 + \mu)(\kappa_0)_{s,j} \quad (103)$$

$$+ \mu \frac{N}{2} (\xi_{j+1} - \xi_{j-1}), \quad j = 2, \dots, N - 1$$

$$\delta \dot{\xi}_j = -\xi_j + 2N(\theta_{j+1} - \theta_{j-1}) - \kappa_{0,j}, \quad j = 2, \dots, N - 1. \quad (104)$$

Here the $2N \times 2N$ matrix $\mathbf{M}_{\text{RFT}}(\theta_i)$ is given by

$$\mathbf{M}_{\text{RFT}}(\theta_i) = \begin{pmatrix} 1 - \frac{\gamma}{1+\gamma} \cos^2 \theta_i & -\frac{\gamma}{1+\gamma} \cos \theta_i \sin \theta_i \\ -\frac{\gamma}{1+\gamma} \cos \theta_i \sin \theta_i & 1 - \frac{\gamma}{1+\gamma} \sin^2 \theta_i \end{pmatrix}.$$

At the fiber endpoints, we set $\xi_1 = \xi_N = 0$ and enforce the boundary conditions $(-\theta_s + \kappa_0)_s|_{s=0,1} = 0$ via the following two equations:

$$\frac{1}{N} \begin{pmatrix} -\sin \theta_1 \\ \cos \theta_1 \end{pmatrix} \cdot \mathbf{M}_{\text{RFT}}(\theta_1) \dot{\mathbf{X}}_{\frac{1}{2}} = -N^2(2\theta_2 - 2\theta_1) + 2N\kappa_{0,1} \quad (105)$$

$$\frac{1}{N} \sum_{i=1}^{N-1} \mathbf{M}_{\text{RFT}}(\theta_i) \dot{\mathbf{X}}_{i-\frac{1}{2}} = 0, \quad (106)$$

To enforce $(\theta_s - \kappa_0)|_{s=0,1} = 0$, we discretize θ_{ss} near the fiber endpoints as

$$\theta_{ss}|_{s=0} \approx N^2(2\theta_2 - 2\theta_1) - 2N\kappa_{0,1}, \quad \theta_{ss}|_{s=1} \approx N^2(2\theta_{N-1} - 2\theta_N) + 2N\kappa_{0,N}. \quad (107)$$

At the $s = 1$ endpoint, equations (100) and (102) coincide to give the boundary condition $\theta_{ss}|_{s=1} = (\kappa_0)_s|_{s=1}$, which, using (107), becomes an equation for θ_N :

$$\theta_N = \theta_{N-1} + \frac{1}{N} \kappa_{0,N} - \frac{1}{2N^2} (\kappa_0)_{s,N}. \quad (108)$$

Counting equations, we have $2(N - 2)$ equations from (103) and (104), 1 equation from (105), 2 equations from (106), and 1 equation from (108) for a total of $2N$ equations. These uniquely determine the $2N$ unknowns $x_0, y_0, \theta_1, \dots, \theta_N, \xi_2, \dots, \xi_{N-1}$.

The equations (103)-(108) are evolved in time using a built-in ODE solver in Matlab.

Acknowledgments. L.O. acknowledges support from NSF Postdoctoral Fellowship DMS-2001959 and thanks Becca Thomases, Yoichiro Mori, and Dallas Albritton for helpful discussion.

REFERENCES

- [1] V. Angeles, F. A. Godínez, J. A. Puente-Velazquez, R. Mendez-Rojano, E. Lauga, and R. Zenit. Front-back asymmetry controls the impact of viscoelasticity on helical swimming. *Physical Review Fluids*, 6(4):043102, 2021.
- [2] S. Camalet and F. Jülicher. Generic aspects of axonemal beating. *New Journal of Physics*, 2(1):24, 2000.
- [3] G. J. Elfring and G. Goyal. The effect of gait on swimming in viscoelastic fluids. *Journal of Non-Newtonian Fluid Mechanics*, 234:8–14, 2016.
- [4] J. Espinosa-Garcia, E. Lauga, and R. Zenit. Fluid elasticity increases the locomotion of flexible swimmers. *Physics of Fluids*, 25(3):031701, 2013.
- [5] H. C. Fu, T. R. Powers, and C. W. Wolgemuth. Theory of swimming filaments in viscoelastic media. *Physical review letters*, 99(25):258101, 2007.
- [6] H. C. Fu, C. W. Wolgemuth, and T. R. Powers. Beating patterns of filaments in viscoelastic fluids. *Physical Review E*, 78(4):041913, 2008.
- [7] H. C. Fu, C. W. Wolgemuth, and T. R. Powers. Swimming speeds of filaments in nonlinearly viscoelastic fluids. *Physics of Fluids*, 21(3):033102, 2009.
- [8] G. R. Fulford, D. F. Katz, and R. L. Powell. Swimming of spermatozoa in a linear viscoelastic fluid. *Biorheology*, 35(4-5):295–309, 1998.
- [9] F. A. Godínez, L. Koens, T. D. Montenegro-Johnson, R. Zenit, and E. Lauga. Complex fluids affect low-reynolds number locomotion in a kinematic-dependent manner. *Experiments in Fluids*, 56(5):1–10, 2015.
- [10] R. E. Goldstein and S. A. Langer. Nonlinear dynamics of stiff polymers. *Physical review letters*, 75(6):1094, 1995.
- [11] J. Gray and G. Hancock. The propulsion of sea-urchin spermatozoa. *Journal of Experimental Biology*, 32(4):802–814, 1955.
- [12] M. Hines and J. Blum. Bend propagation in flagella. i. derivation of equations of motion and their simulation. *Biophysical Journal*, 23(1):41–57, 1978.
- [13] N. C. Keim, M. Garcia, and P. E. Arratia. Fluid elasticity can enable propulsion at low reynolds number. *Physics of Fluids*, 24(8):081703, 2012.
- [14] L. Landau and E. Lifschitz. Theory of elasticity—course of theoretical physics, vol. 7 oxford. UK: *Butterworth-Heinemann*, 1986.
- [15] E. Lauga. Propulsion in a viscoelastic fluid. *Physics of Fluids*, 19(8):083104, 2007.
- [16] E. Lauga. *The fluid dynamics of cell motility*, volume 62. Cambridge University Press, 2020.
- [17] C. Li, B. Qin, A. Gopinath, P. E. Arratia, B. Thomases, and R. D. Guy. Flagellar swimming in viscoelastic fluids: role of fluid elastic stress revealed by simulations based on experimental data. *Journal of The Royal Society Interface*, 14(135):20170289, 2017.
- [18] C. Li, B. Thomases, and R. D. Guy. Orientation dependent elastic stress concentration at tips of slender objects translating in viscoelastic fluids. *Physical Review Fluids*, 4(3):031301, 2019.
- [19] G. Li, E. Lauga, and A. M. Ardekani. Microswimming in viscoelastic fluids. *Journal of Non-Newtonian Fluid Mechanics*, 297:104655, 2021.
- [20] B. Liu, T. R. Powers, and K. S. Breuer. Force-free swimming of a model helical flagellum in viscoelastic fluids. *Proceedings of the National Academy of Sciences*, 108(49):19516–19520, 2011.
- [21] V. A. Martinez, J. Schwarz-Linek, M. Reufer, L. G. Wilson, A. N. Morozov, and W. C. Poon. Flagellated bacterial motility in polymer solutions. *Proceedings of the National Academy of Sciences*, 111(50):17771–17776, 2014.
- [22] O. Maxian, A. Mogilner, and A. Donev. Integral-based spectral method for inextensible slender fibers in stokes flow. *Physical Review Fluids*, 6(1):014102, 2021.
- [23] C. Moreau, L. Giralardi, and H. Gadêlha. The asymptotic coarse-graining formulation of slender-rods, bio-filaments and flagella. *Journal of the Royal Society Interface*, 15(144):20180235, 2018.

- [24] Y. Mori and L. Ohm. An error bound for the slender body approximation of a thin, rigid fiber sedimenting in stokes flow. *Research in the Mathematical Sciences*, 7(2):1–27, 2020.
- [25] Y. Mori and L. Ohm. Accuracy of slender body theory in approximating force exerted by thin fiber on viscous fluid. *Studies in Applied Mathematics*, 2021.
- [26] Y. Mori and L. Ohm. Well-posedness and applications of classical elastohydrodynamics for a swimming filament. *arXiv preprint arXiv:2208.03350*, 2022.
- [27] Y. Mori, L. Ohm, and D. Spirn. Theoretical justification and error analysis for slender body theory. *Communications on Pure and Applied Mathematics*, 73(6):1245–1314, 2020.
- [28] Y. Mori, L. Ohm, and D. Spirn. Theoretical justification and error analysis for slender body theory with free ends. *Archive for Rational Mechanics and Analysis*, 235(3):1905–1978, 2020.
- [29] L. Ohm. Remarks on regularized stokeslets in slender body theory. *Fluids*, 6(8):283, 2021.
- [30] O. Pironneau and D. Katz. Optimal swimming of flagellated micro-organisms. *Journal of Fluid Mechanics*, 66(2):391–415, 1974.
- [31] E. E. Riley and E. Lauga. Enhanced active swimming in viscoelastic fluids. *EPL (Europhysics Letters)*, 108(3):34003, 2014.
- [32] E. E. Riley and E. Lauga. Small-amplitude swimmers can self-propel faster in viscoelastic fluids. *Journal of theoretical biology*, 382:345–355, 2015.
- [33] D. Salazar, A. M. Roma, and H. D. Ceniceros. Numerical study of an inextensible, finite swimmer in stokesian viscoelastic flow. *Physics of Fluids*, 28(6):063101, 2016.
- [34] X. Shen and P. E. Arratia. Undulatory swimming in viscoelastic fluids. *Physical review letters*, 106(20):208101, 2011.
- [35] S. E. Spagnolie, B. Liu, and T. R. Powers. Locomotion of helical bodies in viscoelastic fluids: enhanced swimming at large helical amplitudes. *Physical review letters*, 111(6):068101, 2013.
- [36] J. Teran, L. Fauci, and M. Shelley. Viscoelastic fluid response can increase the speed and efficiency of a free swimmer. *Physical review letters*, 104(3):038101, 2010.
- [37] B. Thomases and R. D. Guy. Mechanisms of elastic enhancement and hindrance for finite-length undulatory swimmers in viscoelastic fluids. *Physical review letters*, 113(9):098102, 2014.
- [38] B. Thomases and R. D. Guy. The role of body flexibility in stroke enhancements for finite-length undulatory swimmers in viscoelastic fluids. *Journal of Fluid Mechanics*, 825:109–132, 2017.
- [39] A.-K. Tornberg and M. J. Shelley. Simulating the dynamics and interactions of flexible fibers in stokes flows. *Journal of Computational Physics*, 196(1):8–40, 2004.
- [40] C. H. Wiggins and R. E. Goldstein. Flexive and propulsive dynamics of elastica at low reynolds number. *Physical Review Letters*, 80(17):3879, 1998.
- [41] C. H. Wiggins, D. Riveline, A. Ott, and R. E. Goldstein. Trapping and wiggling: elastohydrodynamics of driven microfilaments. *Biophysical Journal*, 74(2):1043–1060, 1998.

(L. Ohm) DEPARTMENT OF MATHEMATICS, PRINCETON UNIVERSITY, PRINCETON, NJ 08540
 Email address: laurel.ohm@princeton.edu