

On Quantum Complexity

Mohsen Alishahiha*

*School of Physics, Institute for Research in Fundamental Sciences (IPM),
P.O. Box 19395-5531, Tehran, Iran*

The ETH ansatz for matrix elements of a given operator in the energy eigenstate basis results in a notion of thermalization for a chaotic system. In this context for a certain quantity - to be found for a given model - one may impose a particular condition on its matrix elements in the energy eigenstate basis so that the corresponding quantity exhibit linear growth at late times. The condition is to do with a possible pole structure the corresponding matrix elements may have. Based on the general expectation of complexity one may want to think of this quantity as a possible candidate for the quantum complexity. We note, however, that for the explicit examples we have considered in this paper, there are infinitely many quantities exhibiting similar behavior.

I. INTRODUCTION

For chaotic systems with a finite entropy S , complexity is expected to grow for exponentially large times in the entropy, long after thermal equilibrium has been reached [1, 2]. For such systems the notion of thermalization may be described by the eigenstate thermalization hypothesis (ETH) which gives an understanding of how an observable thermalizes to its thermal equilibrium value [3, 4] (for review see [5]).

To be concrete let us consider a Hamiltonian, H , whose eigenvalues and eigenstates are denoted by E and $|E\rangle$, respectively. Given a general state $|\psi\rangle$, the quantum expectation value of an operator, \mathcal{O} , at given time is

$$\langle \mathcal{O}(t) \rangle = \langle \psi | e^{itH} \mathcal{O} e^{-itH} | \psi \rangle = \int_0^\infty dE_1 dE_2 e^{it(E_1 - E_2)} \langle \psi | E_1 \rangle \langle E_1 | \mathcal{O} | E_2 \rangle \langle E_2 | \psi \rangle. \quad (1)$$

In the context of the thermalization of a quantum chaotic system one is typically interested in the equal time averages of observables. More precisely, one would like to find the time average of \mathcal{O} over an interval of time, which eventually it will be sent to infinity.

According to the ETH, thermalization occurs at the level of individual eigenstates of the Hamiltonian. In fact setting

$$\varepsilon = \frac{E_1 + E_2}{2}, \quad \omega = E_1 - E_2, \quad (2)$$

the ETH states that the matrix elements of observables in the basis of the eigenstates of the Hamiltonian obey the following ansatz [6]

$$\langle E_1 | \mathcal{O} | E_2 \rangle = \bar{\mathcal{O}}(\varepsilon) \delta_{E_1, E_2} + e^{-S} f(\varepsilon, \omega) \mathcal{R}_{E_1 E_2} \quad (3)$$

where $\bar{\mathcal{O}}(\varepsilon)$ is the micro canonical average of the corresponding operator, S is thermodynamical entropy of the system, $f(\varepsilon, \omega)$ is a smooth function of its arguments and \mathcal{R} is unit variance random function with zero mean.

The above ETH ansatz has an immediate application in understanding of the thermalization which indicates that the quantum expectation value of an observable must approach its thermal average for long enough times. We note, however, that this ansatz does not tell us how long the process of thermalization is.

Of course our main concern in this note is not to explore the thermalization of the system. Actually the aim of the present letter is to understand the late time behavior of a certain observable when the system is in the thermal equilibrium. Indeed, within the context of the ETH assuming to have a system in the thermal equilibrium we are interested in finding, if any, an observable that is still growing with time.

In the next section we would like to study such a possibility. More precisely, we will explore a possible condition one may impose on the matrix elements of an observable so that the corresponding expectation value exhibits time growth even though the system has been already reached the thermal equilibrium.

II. QUANTUM COMPLEXITY

To formulate our question and its possible answer it is useful to utilize the wave function formalism as we are interested in the late time behavior of physical quantities which may not even be associated with local operators. In this formalism for a given quantum system we will define the Hartle-Hawking wave function by which one could study expectation values of different observables.

To proceed, let us start with a Hamiltonian H , whose eigenvalues and eigenstate wave functions are denoted by E and $\psi_E(x)$ ¹, respectively. We assume that the “non-normalized” eigenstate wave functions satisfy the following orthogonality condition (with an appropriate integral

* alishah@ipm.ir

¹ In what follows we will consider a chaotic system with non-degenerate continuous spectrum.

measure)

$$\int_0^\infty dx \psi_{E_1}^*(x) \psi_{E_2}(x) = \frac{\delta(E_1 - E_2)}{N(E_1)}, \quad (4)$$

where N is a function of the energy by which the eigenstates may be normalized.

For the system at finite temperature given by β^{-1} , let us define the *time shifted Hartle-Hawking wave function* $\phi(\beta, t, x)$ as follows

$$\phi(\beta, t, x) = \int_0^\infty dE e^{-(\frac{\beta}{2} + it)E} \sqrt{\eta(E)} N(E) \psi_E(x), \quad (5)$$

so that the partition function of the system is

$$Z(\beta) = \int_0^\infty dx \phi^*(\beta, t, x) \phi(\beta, t, x). \quad (6)$$

Here $\eta(E)$ is an arbitrary function which for more cases we will be considering in this paper we will set it to one.

Using the orthogonality condition for the eigenstate wave functions one gets the standard expression for partition function if one identifies the density of states as $\rho(E) = \eta(E)N(E)$

$$Z \equiv Z(\beta) = \int dE e^{-\beta E} \rho(E). \quad (7)$$

Given a function f of the coordinate x one may define a *quantum object* \mathcal{A} associated with it as follows²

$$\mathcal{A}_f(\beta, t) = \frac{1}{Z} \int_0^\infty dx dx' \phi^*(\beta, t, x) \phi(\beta, t, x') f(x, x'), \quad (8)$$

which may be simplified to get³

$$\mathcal{A}_f(\beta, t) = \frac{1}{Z} \int_0^\infty dE_1 dE_2 e^{-\frac{\beta}{2}(E_1 + E_2)} e^{it(E_1 - E_2)} \times \rho(E_1) \rho(E_2) A(E_1, E_2), \quad (9)$$

where

$$A(E_1, E_2) = \int_0^\infty dx dx' \psi_{E_1}^*(x) \psi_{E_2}(x') f(x, x'). \quad (10)$$

Indeed this is a generalization of (1) in which the corresponding object may not necessarily represent an expectation value or correlation functions of local operators, though for appropriate choices of the function f it reduces to that for quantum correlation functions of local operators. In this case the function $f(x, x')$ may be thought of as the matrix elements of the corresponding operators in coordinate basis. More precisely for a given

local operator \mathcal{O} , one has $f(x, x') = \langle x | \mathcal{O} | x' \rangle$, that results in $A(E_1, E_2) = \langle E_1 | \mathcal{O} | E_2 \rangle$. Clearly if the function f only depends on coordinates the corresponding matrix is diagonal $f(x, x') = \delta(x - x') f(x)$. Of course, as already mentioned, we would like to emphasize that in general the function f could be any function which may not be given in terms of the matrix elements of certain local operators.

As far as the time dependence of the corresponding quantum object is concerned, as we will see, the main role is played by the function A (10) (A -function) associated with the given function f .

For the cases in which the quantum object \mathcal{A}_f reduces to quantum expectation value of local operators, one would expect that the A -function follows the ETH ansatz and therefore the long time average of \mathcal{A}_f approaches its value in thermal equilibrium.

In our case, however, since we are interested in the late time behavior of the quantum object, we will not perform the long time average and instead will look for possible conditions from which the late time behavior of \mathcal{A}_f may be read.

More precisely, we would like to see whether there is a condition under which the corresponding quantum quantity keeps growing with time even though the whole system is reached thermal equilibrium.

To address this equation, following the ETH idea, it is clear from the equation (9) that the corresponding information should be encoded in the behavior of A -function (10). To proceed, since we are interested in the late time behavior it useful to rewrite the expression (9) in terms of variables defined in (2)

$$\mathcal{A}_f(\beta, t) = \frac{1}{Z} \int_0^\infty d\varepsilon e^{-\beta\varepsilon} \int_{-\infty}^\infty d\omega e^{i\omega t} \rho(\varepsilon + \frac{\omega}{2}) \rho(\varepsilon - \frac{\omega}{2}) \times A(\varepsilon, \omega) \quad (11)$$

and then study the behavior of $A(\varepsilon, \omega)$ in the limit of $\omega \rightarrow 0$.

Actually, as it is evident from the above expression, the time dependence of \mathcal{A}_f should be read from the ω -integral. Indeed, due to the simple factor of $e^{i\omega t}$ in the integrand, using the Cauchy's residue theorem with the assumption that the density of state $\rho(\varepsilon \pm \omega/2)$ is a smooth function in the limit of $\omega \rightarrow 0$, in order to get a non-trivial time dependence, the A -function must have a pole structure of order of n for $n \geq 2$. In particular, for the case of a double pole structure where the A -function has the following limiting behavior

$$A(\varepsilon, \omega) = -\frac{a(\varepsilon)}{\omega^2} + \text{local terms}, \quad \text{for } \omega \rightarrow 0, \quad (12)$$

with a positive smooth function $a(\varepsilon)$, one finds that the quantum object \mathcal{A} exhibits a linear growth at late times⁴

$$\mathcal{A}_f(\beta, t) = C_0 + \frac{1}{Z} \int_0^\infty d\varepsilon e^{-\beta\varepsilon} \rho^2(\varepsilon) a(\varepsilon) (2\pi t), \quad (13)$$

² Here we have used the partition function to normalize \mathcal{A}_f , so that for $f = \delta(x - x')$ one has $\mathcal{A} = 1$.

³ In what follows we have set $\eta(E) = 1$ so that the density of state is given by $\rho(E) = N(E)$.

⁴ It is worth noting that this linear growth must not be confused

where C_0 is a time independent constant that is a function of β . It is worth noting that for poles of higher order, one generically gets power low time dependent behavior. We will back to this point latter.

Having found a quantum object that has linear growth at late times, it is tempting to identify the corresponding quantum object, \mathcal{A}_f , as the quantum complexity. To be precise, we would like to define the complexity as follows.

For a chaotic quantum system the quantum complexity is defined by (8) for a particular function f -to be found for a given system- so that the associated A -function exhibits a double pole structure in the limit of $E_1 \rightarrow E_2$

$$A(E_1, E_2) \approx -\frac{a(E_1, E_2)}{(E_1 - E_2)^2} + \text{local terms} \quad (14)$$

where $a(E_1, E_2)$ is a smooth positive function.

Of course for a given quantum system and for a given state, a priori, it is not obvious how to find the function f that results in the desired double pole structure for A -function. Moreover, in general the corresponding quantity may not be given in terms of local operators.

To further explore this observation in the next section we will present an explicit example in which one could identify a proper function f , that results in a linear growth for \mathcal{A}_f .

III. EXPLICIT EXAMPLE

Let us consider a quantum system with the following Hamiltonian

$$H = \frac{P^2}{2} + 2\mu e^{-x} + 2e^{-2x}. \quad (15)$$

Then the corresponding Schrödinger equation is

$$\left(-\frac{d^2}{dx^2} + 4\mu e^{-x} + 4e^{-2x}\right) \psi(x) = 2E\psi(x). \quad (16)$$

The eigenstate wave functions of the above equation are given in terms of the Whittaker function of the second kind with imaginary order

$$\psi_{\mu,E}(x) = e^{x/2} W_{-\mu, i\sqrt{2E}}(4e^{-x}). \quad (17)$$

Actually this Hamiltonian is used to study different aspects of two-dimensional JT gravity (see *e.g.* [7–10]). For general $\mu \neq 0$ it corresponds to JT gravity with end of the world brane whose tension is given by μ . For the particular value of $\mu = \frac{1}{2}$ it may also be considered as supersymmetric version of JT-gravity [11]⁵. It may also be

thought of as a Liouville Quantum Mechanics describing Sachdev-Ye-Kitaev Model [12].

Using this Hamiltonian the complexity of JT gravity has also been studied in [13–15]. Of course in what follows the relevance of this quantum system to the two-dimensional JT gravity is not important for us, and we will consider it as a one dimensional quantum system.

The orthogonality condition for the eigenstates $\psi_{\mu,E}(x)$ is [16]

$$\int_0^\infty dx \psi_{\mu,E_1}(x) \psi_{\mu,E_2}(x) = \frac{\delta(E_1 - E_2)}{N_\mu(E_1)}, \quad (18)$$

where

$$N_\mu(E) = \left| \Gamma\left(\frac{1}{2} + \mu + i\sqrt{2E}\right) \right|^2 \frac{\sinh 2\pi\sqrt{2E}}{4\pi^2}, \quad (19)$$

which is essentially the density of state of the system, $\rho_\mu(E) = N_\mu(E)$ (here we set $\eta = 1$). Therefore the time shifted Hartle-Hawking wave function is

$$\phi(\beta, t, x) = \int_0^\infty dE e^{-(\frac{\beta}{2} + it)E} \rho_\mu(E) \psi_{\mu,E}(x). \quad (20)$$

Motivated by the result of [14] we will consider $f(x, x') = \delta(x - x')x$ by which the associated A -function reads

$$A(E_1, E_2) = \int_0^\infty dx \psi_{\mu,E_1}(x) \psi_{\mu,E_2}(x) x. \quad (21)$$

Actually if one recalls the fact that the function f may be interpreted as matrix elements in the coordinate basis, the above choice corresponds to the matrix elements of position operator that is obviously diagonal leading to a delta function. On the other hand since the wave function satisfies the Schrödinger equation, essentially in this case what we are evaluating is the average of position operator (distance).

It is then straightforward to study the pole structure of the A -function. Indeed, using the variables defined in (2) and in the limit of $E_1 \rightarrow E_2$ one finds [14]

$$A(\varepsilon, \omega) = -\frac{\sqrt{2\varepsilon}}{\pi\rho_\mu(\varepsilon)} \frac{1}{\omega^2} + \text{local terms}. \quad (22)$$

Therefore from the equation (9) one can find the late time behavior as follows

$$\mathcal{A}(\beta, t) = C_0 + \frac{1}{Z} \int_0^\infty d\varepsilon e^{-\beta\varepsilon} \rho_\mu(\varepsilon) \sqrt{2\varepsilon} (2t) \quad (23)$$

that is the linear growth, as expected.

As another example we note that in the context of random matrix model and its connection with chaos we are typically dealing with matter two point functions whose matrix elements in energy eigenstates have the following general form [17]

$$\mathcal{O}_{E_1, E_2} = \frac{|\Gamma(\Delta + i(\sqrt{E_1} - \sqrt{E_2}))\Gamma(\Delta + i(\sqrt{E_1} + \sqrt{E_2}))|^2}{\Gamma(2\Delta)}. \quad (24)$$

with that of the ramp phase in *e.g.* the spectrum form factor where the linear growth was the consequence of subleading connected part of the density-density correlation. Here we have a linear growth at leading disconnected level. We will back to this point later.

⁵ Note that in this case one has $\eta(E) = 1/E$.

where Δ is the dimension of the corresponding matter field. From this expression one may define an A -function as follows

$$A(E_1, E_2) = - \lim_{\Delta \rightarrow 0} \frac{d}{d\Delta} \mathcal{O}_{E_1, E_2} \quad (25)$$

$$= -2|\Gamma(i(\sqrt{E_1} - \sqrt{E_2}))\Gamma(i(\sqrt{E_1} + \sqrt{E_2}))|^2.$$

It is then easy to see that in the limit of $E_1 \rightarrow E_2$ this A -function exhibits a double pole structure

$$A(E_1, E_2) = -\frac{4\pi\sqrt{\varepsilon}}{\sinh(2\pi\sqrt{\varepsilon})\omega^2} + \text{local terms}. \quad (26)$$

Actually recalling the relation between random matrix model and two-dimensional JT-gravity, the above expression corresponds to the case of $\mu = 0$ in (22).

It is also interesting to look at the rate of complexity growth

$$\frac{d\mathcal{A}_f(t)}{dt} = \frac{1}{Z} \int_0^\infty d\varepsilon e^{-\beta\varepsilon} \rho_\mu(\varepsilon) \sqrt{2\varepsilon}, \quad (27)$$

which might be compared with the Lloyd's bound [18]. Actually we note that in the context of holographic complexity where the complexity may be computed using CA conjecture [19] the rate of complexity turns out to be twice of the energy of the system, saturating the Lloyd's bound [18]. Here, instead, we get rather an expression that is not that of Lloyd's bound. We note, however, that at low energies one has $\rho_\mu(\varepsilon) \sim \sqrt{2\varepsilon}$ and thus the above expression may be thought of as the average of energy in a canonical ensemble.

If one works with a non-normalized situation by dropping $1/Z$ factor, one then could present the rate of complexity growth in the macro canonical ensemble by making use of the inverse Laplace transformation. In this case the corresponding rate is given by $\rho(E_0)\sqrt{2E_0}$ that at low energies results in $\sim 2E_0$, reminiscing of Lloyd's bound. Here E_0 is the energy of the macro canonical ensemble.

In general for $\mu = 0$ the integral may be performed exactly to find

$$\frac{d\mathcal{A}_f(t)}{dt} = \frac{2e^{-\frac{2\pi^2}{\beta}}}{\sqrt{2\pi\beta}} + \left(\frac{1}{2\pi} + \frac{4\pi^2}{\beta}\right) \text{Erf}\left(\frac{\sqrt{2\pi}}{\sqrt{\beta}}\right), \quad (28)$$

which at low temperatures goes as $\sim \beta^{-1/2}$ while at high temperatures it is $\sim \beta^{-1}$. Although for general μ the full expression for the rate of complexity growth may not be written explicitly, asymptotic behaviors for low and high temperatures are the same as that of $\mu = 0$.

IV. DISCUSSIONS

In this letter we have defined a quantum object associated with a given function in a chaotic system. We

have demonstrated that under certain condition the corresponding quantum object exhibits linear growth at late times much longer than the system reaches the thermal equilibrium.

Given the function f , we will impose a condition on its matrix elements in the energy basis. We have shown that if for a given function f -to be found- the matrix elements in energy basis exhibit a double pole structure at late times (14), the corresponding quantum object given by the equation (9) will have late times linear growth which could be interpreted as the quantum complexity. Of course for given state in a given system, a priori, it is not clear how to find the function f with the above desired property.

In the context of thermalization of quantum system when the function is given by the expectation value of local operators, one generally assumes that matrix elements follow the ETH ansatz.

Actually in order to get a non-trivial time dependence at late times, the A -function should have poles of order of n with $n \geq 2$. For general $n > 2$ one generally gets power law growth at late times, though for $n = 2$ one has a linear growth. Since having a linear growth at late times might be a signature of the complexity [1] that is expected to be the fastest growth, one may propose a hypothesis that the double pole structure is the highest pole structure the A -function could have.

It is worth mentioning that for a give chaotic model there could be several f 's that give double pole structure for A -function which in turns results in the late time linear growth. For the explicit example we have presented in the previous section it is straightforward to see that for any functions in the form of $f(x, x') = \delta(x - x')x^m$, with integer m , one finds double pole structure. Moreover from the matrix elements (24) it is easy to construct several A -functions with the desired property. They can be obtained by taking $\Delta \rightarrow 0$ limit of m^{th} Δ -derivative of the matrix elements (24).

This is very similar to the observation made in [20] in the context of holographic complexity where it was shown that there are infinite class of gravitational observables in asymptotically Anti-de Sitter space which living on codimension one slices of the geometry, that exhibit universal features as that in complexity. Namely they grow linearly in time at late times.

To conclude this part we should emphasis that having known the late time behavior is not enough to define complexity and we need more information to fix it. Actually it seems that this is also the case even for a rather well studied candidate for the complexity in the context of Krylov complexity⁶ (for discussions about Krylov complexity closely related to our point of view see [21–25]). It would be interesting to explore this point better [28].

⁶ Yet another interesting notion of complexity which is related to what we have studied in this paper, spread complexity, has been introduced in [26, 27].

Another interesting feature of complexity is that it saturates at the very late times given by exponential of the entropy of the system. It is then natural to see how the saturation could occur in this context.

To address this question we note that the quantum object \mathcal{A} associated with the function f may be written as follows

$$\mathcal{A}_f(\beta, t) = \int_0^\infty dE_1 dE_2 e^{-\frac{\beta}{2}(E_1+E_2)} e^{it(E_1-E_2)} \times \rho(E_1, E_2) A(E_1, E_2), \quad (29)$$

where $\rho(E_1, E_2)$ is a function of E_1 and E_2 which has the following general form

$$\rho(E_1, E_2) = \rho(E_1)\rho(E_2) + \rho_c(E_1, E_2). \quad (30)$$

Here ρ_c represents the connected term meaning that it cannot be written in a factorized form of $g_1(E_1)g_2(E_2)$ with $g_{1,2}$ being arbitrary functions. Clearly for the first factorized term the above function reduces to that of (9). The connected terms could have either perturbative or non-perturbative origins which may generally have non-trivial pole structure that could result in the saturation phase for very late times.

Actually this is a well known structure which has been seen in the literature for spectrum form factor of chaotic models such as JT-gravity in which the pole structure of $\rho(E_1, E_2)$ results in the ramp phase. Of course for spectrum form factor there is no A -function and whole time dependence controls by the density-density correlator. In the context of holographic complexity for JT gravity in which we also have the A -function the connected part of $\rho(E_1, E_2)$ which has non-trivial pole structure at late times is, indeed, responsible for the saturation phase [13–15].

We note, however, that in the present case, where we are dealing with a general formalism which is not directly related to the holography picture, it is not clear how the full expression of the connected term could be computed.

Nonetheless as far as the late time behavior is concerned, one would expect that the main contribution comes from the short range correlation which is given by the universal sine kernel term [29]⁷

$$\rho_c(\varepsilon, \omega) \approx -\frac{\sin^2(D\omega\rho(\varepsilon))}{(D\omega)^2}, \quad \text{for } \omega \ll 1. \quad (31)$$

Here D is the dimension of Hilbert space which is given by the exponential of the entropy of the system. Therefore the whole late time behavior of the quantum object \mathcal{A}_f is described as follows: the double point structure of the A -function leads to linear growth at the leading disconnected part of the density-density correlation, while there is the saturation phase which can be described by subleading connected term given in the universal sine-kernel term multiplied by the double pole structure of A -function. It is then easy to see that the saturation occurs at $t \sim D$.

As a final comment we note that the structure we have presented in this letter, naively, has the similar feature as that of the Krylov complexity [21–24]. In particular for the explicit example we have considered the complexity may be thought of as the average of position. It would also be interesting to understand this connection better [28].

ACKNOWLEDGMENTS

I would like to thank Alexandre Belin, Kyriakos Papadodimas and Julian Sonner for discussions. The author would also like to thank Sonner’s group in Geneva University for discussions and comments. I would also like to thank Souvik Banerjee for comments and discussions on different aspects of complexity as well as collaborating on several past and ongoing projects. The author would also like to thank Department of Theoretical Physics of CERN for very warm hospitality

-
- [1] L. Susskind, “Computational Complexity and Black Hole Horizons,” *Fortsch. Phys.* **64** (2016), 24-43 doi:10.1002/prop.201500092 [arXiv:1403.5695 [hep-th]].
 - [2] L. Susskind, “Entanglement is not enough,” *Fortsch. Phys.* **64** (2016), 49-71 doi:10.1002/prop.201500095 [arXiv:1411.0690 [hep-th]].
 - [3] J. M. Deutsch, “Quantum statistical mechanics in a closed system,” *Phys Rev A* **43** 2046 (1991), doi:10.1103/PhysRevA.43.2046.
 - [4] M. Srednicki, “Chaos and Quantum Thermalization,” doi:10.1103/PhysRevE.50.888 [arXiv:cond-mat/9403051 [cond-mat]].

- [5] L. D’Alessio, Y. Kafri, A. Polkovnikov and M. Rigol, “From quantum chaos and eigenstate thermalization to statistical mechanics and thermodynamics,” *Adv. Phys.* **65** (2016) no.3, 239-362 doi:10.1080/00018732.2016.1198134 [arXiv:1509.06411 [cond-mat.stat-mech]].
- [6] M. Srednicki, “The approach to thermal equilibrium in quantized chaotic systems,” *J. Phys. A* **32** (1999) 1163, doi:10.1088/0305-4470/32/7/007.
- [7] D. Harlow and D. Jafferis, “The Factorization Problem in Jackiw-Teitelboim Gravity,” *JHEP* **02** (2020), 177 doi:10.1007/JHEP02(2020)177 [arXiv:1804.01081 [hep-th]].
- [8] Z. Yang, “The Quantum Gravity Dynamics of Near Extremal Black Holes,” *JHEP* **05** (2019), 205 doi:10.1007/JHEP05(2019)205 [arXiv:1809.08647 [hep-th]].

⁷ I would like to thank Julian Sonner for pointing out this to me.

- [9] P. Saad, “Late Time Correlation Functions, Baby Universes, and ETH in JT Gravity,” [arXiv:1910.10311 [hep-th]].
- [10] P. Gao, D. L. Jafferis and D. K. Kolchmeyer, “An effective matrix model for dynamical end of the world branes in Jackiw-Teitelboim gravity,” JHEP **01** (2022), 038 doi:10.1007/JHEP01(2022)038 [arXiv:2104.01184 [hep-th]].
- [11] M. R. Douglas, I. R. Klebanov, D. Kutasov, J. M. Maldacena, E. J. Martinec and N. Seiberg, “A New hat for the $c=1$ matrix model,” [arXiv:hep-th/0307195 [hep-th]].
- [12] D. Bagrets, A. Altland and A. Kamenev, “Sachdev–Ye–Kitaev model as Liouville quantum mechanics,” Nucl. Phys. B **911** (2016), 191–205 doi:10.1016/j.nuclphysb.2016.08.002 [arXiv:1607.00694 [cond-mat.str-el]].
- [13] L. V. Iliesiu, M. Mezei and G. Sárosi, “The volume of the black hole interior at late times,” JHEP **07**, 073 (2022) doi:10.1007/JHEP07(2022)073 [arXiv:2107.06286 [hep-th]].
- [14] M. Alishahiha, S. Banerjee and J. Kames-King, “Complexity via replica trick,” JHEP **08** (2022), 181 doi:10.1007/JHEP08(2022)181 [arXiv:2205.01150 [hep-th]].
- [15] M. Alishahiha and S. Banerjee, “On the saturation of late-time growth of complexity in supersymmetric JT gravity,” [arXiv:2209.02441 [hep-th]].
- [16] R. Szmytkowski and S. Bielski, “An orthogonality relation for the Whittaker functions of the second kind of imaginary order,” arXiv:0910.1492 [math.CA]
- [17] D. L. Jafferis, D. K. Kolchmeyer, B. Mukhametzhanov and J. Sonner, “JT gravity with matter, generalized ETH, and Random Matrices,” [arXiv:2209.02131 [hep-th]].
- [18] S. Lloyd, “Ultimate Physical limits to computation,” Nature **406** (2000) 1047, quant-ph/9908043.
- [19] A. R. Brown, D. A. Roberts, L. Susskind, B. Swingle and Y. Zhao, “Holographic Complexity Equals Bulk Action?,” Phys. Rev. Lett. **116** (2016) no.19, 191301 doi:10.1103/PhysRevLett.116.191301 [arXiv:1509.07876 [hep-th]].
- [20] A. Belin, R. C. Myers, S. M. Ruan, G. Sárosi and A. J. Speranza, “Does Complexity Equal Anything?,” Phys. Rev. Lett. **128** (2022) no.8, 081602 doi:10.1103/PhysRevLett.128.081602 [arXiv:2111.02429 [hep-th]].
- [21] D. E. Parker, X. Cao, A. Avdoshkin, T. Scaffidi and E. Altman, “A Universal Operator Growth Hypothesis,” Phys. Rev. X **9** (2019) no.4, 041017 doi:10.1103/PhysRevX.9.041017 [arXiv:1812.08657 [cond-mat.stat-mech]].
- [22] J. L. F. Barbón, E. Rabinovici, R. Shir and R. Sinha, “On The Evolution Of Operator Complexity Beyond Scrambling,” JHEP **10** (2019), 264 doi:10.1007/JHEP10(2019)264 [arXiv:1907.05393 [hep-th]].
- [23] S. K. Jian, B. Swingle and Z. Y. Xian, “Complexity growth of operators in the SYK model and in JT gravity,” JHEP **03** (2021), 014 doi:10.1007/JHEP03(2021)014 [arXiv:2008.12274 [hep-th]].
- [24] E. Rabinovici, A. Sánchez-Garrido, R. Shir and J. Sonner, “Operator complexity: a journey to the edge of Krylov space,” JHEP **06** (2021), 062 doi:10.1007/JHEP06(2021)062 [arXiv:2009.01862 [hep-th]].
- [25] A. Kar, L. Lamprou, M. Rozali and J. Sully, “Random matrix theory for complexity growth and black hole interiors,” JHEP **01** (2022), 016 doi:10.1007/JHEP01(2022)016 [arXiv:2106.02046 [hep-th]].
- [26] V. Balasubramanian, P. Caputa, J. M. Magan and Q. Wu, “Quantum chaos and the complexity of spread of states,” Phys. Rev. D **106** (2022) no.4, 046007 doi:10.1103/PhysRevD.106.046007 [arXiv:2202.06957 [hep-th]].
- [27] V. Balasubramanian, J. M. Magan and Q. Wu, “A Tale of Two Hungarians: Tridiagonalizing Random Matrices,” [arXiv:2208.08452 [hep-th]].
- [28] M. Alishahiha, S. Banerjee, Work in progress
- [29] M. L. Mehta, “Random Matrices.” (Elsevier, San Diego, 2004) Third edition