KÄHLER–EINSTEIN METRICS ON SMOOTH FANO TOROIDAL SYMMETRIC VARIETIES OF TYPE AIII

KYUSIK HONG, DONGSEON HWANG, AND KYEONG-DONG PARK

ABSTRACT. The wonderful compactification X_m of a symmetric homogeneous space of type AIII(2, m) for each $m \ge 4$ is Fano, and its blowup Y_m along the unique closed orbit is Fano if $m \ge 5$ and Calabi–Yau if m = 4. Using a combinatorial criterion for K-polystability of smooth Fano spherical varieties obtained by Delcroix, we prove that X_m admits a Kähler–Einstein metric for each $m \ge 4$ and Y_m admits a Kähler– Einstein metric if and only if m = 4, 5.

1. INTRODUCTION

A Kähler metric on a complex manifold is said to be Kähler-Einstein if it satisfies the Einstein condition

 $\operatorname{Ric}(\omega) = \lambda \,\omega,$

where $\lambda \in \mathbb{R}$ and $\operatorname{Ric}(\omega)$ denotes the Ricci curvature form of the associated Kähler form ω . While Calabi–Yau manifolds and Kähler manifolds with ample canonical bundle always admit a Kähler–Einstein metric by the works of Aubin [Aub78] and Yau [Yau78], Fano manifolds do not necessarily admit a Kähler–Einstein metric in general. By the Yau–Tian–Donaldson conjecture solved by Chen–Donaldson–Sun [CDS15a, CDS15b, CDS15c] and Tian [Tia15], the existence of a Kähler–Einstein metric on a Fano manifold has turned out to be equivalent to the *K*-polystability which is an algebro-geometric condition introduced by Tian [Tia94, Tia97] and Donaldson [Don02].

However, it is a difficult problem to determine K-polystability of a given Fano manifold. Most known results are concentrated on del Pezzo surfaces and Fano 3-folds by focusing on each individual manifold or a general member of a finitely many families of Fano manifolds ([TY87], [Tia90], and [ACCFKMGSSV]) except for the recent breakthrough [AZ22] and [Zhu21].

Fortunately, more manageable K-polystability criterion have been found for Fano varieties equipped with algebraic group actions. For instance, the works of Mabuchi [Mab87] together with that of Wang and Zhu [WZ04] imply that a toric Fano manifold is K-polystable if and only if the barycenter of the moment polytope is the origin. This gave us a powerful combinatorial criterion for the K-polystability of a toric Fano manifold.

This is largely generalized to smooth Fano spherical varieties in terms of its moment polytope and spherical data by Delcroix ([Del20a]). More precisely, a smooth Fano spherical variety X is K-polystable if and only if the barycenter of the moment polytope of X with respect to the Duistermaat–Heckman measure belongs to certain cone. See also [Del22] for some examples including the blow-up of a quadric along a linear subquadric. Here, a normal variety is called *spherical* if it admits an action of a reductive algebraic group whose Borel subgroup acts with an open orbit on the variety.

If a Fano spherical manifold is homogeneous, it is K-polystable by [Mat72, Section 5]. There are two additional well-known classes of spherical varieties: symmetric varieties and horospherical varieties. It is known that, under the assumption that it is a non-homogeneous Fano manifold of Picard number one, the former is K-polystable by [LPY21] and the latter is not since the automorphism group is non-reductive by [Pas09, Theorem 0.1]. Thus one might be interested in determining K-polystability of non-homogeneous Fano symmetric manifolds of Picard number two or higher.

Recall that symmetric varieties are normal equivariant open embeddings of (algebraic) symmetric homogeneous spaces (see Section 2.3 for details). From [DCP83, Theorem 3.1], we know that a symmetric homogeneous space $G/N_G(G^{\theta})$ admits a maximal smooth equivariant compactification having a unique

²⁰¹⁰ Mathematics Subject Classification. Primary: 14M27, 32Q20, Secondary: 32M12, 32Q26, 53C55.

Key words and phrases. symmetric variety, wonderful compactification, Kähler–Einstein metric, K-stability, moment polytope, spherical variety, greatest Ricci lower bound.

closed *G*-orbit, which is called the *wonderful compactification* of $G/N_G(G^{\theta})$. A symmetric variety without colors is said to be *toroidal*, and a toroidal symmetric variety of $G/N_G(G^{\theta})$ dominates the wonderful compactification of $G/N_G(G^{\theta})$ by [DCP83, Section 2.3]. If a Fano symmetric variety is obtainable from a wonderful compactification *S* by a sequence of blow-ups along closed orbits, then it is either *S* or the blow-up of *S* along the unique closed orbit by [Ruz12, Theorem B]. In summary, one can ask the following question:

Question 1.1. Is the wonderful compactification of a symmetric homogeneous space K-polystable? Which of the blow-ups of wonderful compactifications of symmetric homogeneous spaces along the (unique) closed orbit are K-polystable?

In this paper, we answer the question in the case of the wonderful compactifications X_m of symmetric homogeneous spaces of type AIII(2, m) for $m \ge 4$ and their blow-ups Y_m along the closed orbit. It is known that X_m is Fano for each $m \ge 4$ and Y_m is Fano if $m \ge 5$ and Calabi–Yau if m = 4. Recall that X_m is of Picard number two and Y_m is of Picard number three. Geometrically, for $m \ge 5$, the wonderful compactification of symmetric homogeneous space

$$\operatorname{SL}_m(\mathbb{C})/S(\operatorname{GL}_2(\mathbb{C}) \times \operatorname{GL}_{m-2}(\mathbb{C}))$$

of type AIII(2, m) can be described as the blow-up of the product $\operatorname{Gr}(2,m) \times \operatorname{Gr}(m-2,m)$ of Grassmannians along the closed orbit $\mathcal{F}\ell(2,m-2;m)$. In the case of m = 4, the symmetric homogeneous space $\operatorname{SL}_4(\mathbb{C})/N_{\operatorname{SL}_4(\mathbb{C})}(S(\operatorname{GL}_2(\mathbb{C}) \times \operatorname{GL}_2(\mathbb{C})))$ admits the wonderful compactification.

Theorem 1.2. All wonderful compactifications X_m of symmetric homogeneous spaces of type AIII(2, m) for $m \ge 4$ are K-polystable.

For the intersection Z of the two G-stable divisors Y_1 and Y_2 in X_m , it turns out that the blow-up $B\ell_Z(X_m)$ of X_m along the closed orbit Z admits a Kähler–Einstein metric if and only if m = 4, 5. Note that when $m = 4, B\ell_Z(X_m)$ is Calabi–Yau, so it admits a Kähler–Einstein metric.

Theorem 1.3. Let X_m be the wonderful compactification of symmetric homogeneous space of type AIII(2, m) for $m \ge 5$. The blow-up $B\ell_Z(X_m)$ of X_m along the closed orbit Z is K-polystable if m = 5 and K-unstable if $m \ge 6$.

In fact, we show that, in Theorems 3.12 and 4.4, both X_m for $m \ge 4$ and $B\ell_Z(X_m)$ of X_m for m = 5 are $SL_m(\mathbb{C})$ -equivariantly uniformly K-stable, not just K-polystable, which implies Theorems 1.2 and 1.3 by Proposition 2.12. Here, we emphasize that Theorems 1.2 and 1.3 are one of the few cases, including [Del22] and [LL22], that the K-stability or K-unstability of a reasonable class consisting of infinitely many Fano manifolds of different dimensions is verified. In practice, the estimate of the barycenter becomes extremely hard when considering *infinite* series of the moment polytope of Fano spherical manifolds.

The paper is organized as follows. In Section 2, we recall the theory of spherical varieties by focusing on symmetric varieties, and we state a combinatorial criterion for *G*-equivariant K-stability of smooth Fano spherical varieties obtained by Delcroix in terms of algebraic moment polytopes. In Section 3, we study the wonderful compactification X_m of symmetric homogeneous space of type AIII(2, m) for $m \ge 4$, and prove Theorem 1.2 by computing the barycenter of the moment polytope of X_m with respect to the Duistermaat– Heckman measure. In Section 4, we prove Theorem 1.3 and compute the greatest Ricci lower bound of the blow-up of X_6 along the closed orbit.

Acknowledgments. The authors would like to thank JongHae Keum for his interest and encouragement. They would like to express their gratitude to the anonymous referees for carefully reading the manuscript and providing excellent suggestions for improvement.

Kyusik Hong was supported by the National Research Foundation of Korea(NRF) grant funded by the Korea government(MSIT) (NRF-2019R1A2C3010487, NRF-2021R1F1A1059506). DongSeon Hwang was supported by the Samsung Science and Technology Foundation under Project SSTF-BA1602-03, the National Research Foundation of Korea(NRF) grant funded by the Korea government(MSIT) (NRF-2021R1A2C1093787), and the Institute for Basic Science(IBS-R032-D1). Kyeong-Dong Park was supported by the National Research Foundation of Korea(NRF) grant funded by the Korea government(MSIT) (NRF-2019R1A2C3010487, NRF-2021R1C1C2092610), and by the fund of research promotion program, Gyeongsang National University, 2022. He was also supported by Learning & Academic research institution for Master's PhD students, and Postdocs(LAMP) Program of the National Research Foundation of Korea(NRF) grant funded by the Ministry of Education(RS-2023-00301974).

2. Spherical varieties and symmetric varieties

Let G be a connected reductive algebraic group over \mathbb{C} .

2.1. Spherical varieties and algebraic moment polytopes. We review general notions and results about spherical varieties. We refer [Kno91], [Tim11] and [Gan18] as references for spherical varieties.

Definition 2.1. A normal variety X equipped with an action of G is called *spherical* if a Borel subgroup B of G acts on X with an open orbit.

Let G/H be an open dense G-orbit of a spherical variety X and T a maximal torus of B. By definition, the spherical weight lattice \mathcal{M} of G/H is a subgroup of characters $\chi \in \mathfrak{X}(B) = \mathfrak{X}(T)$ of (nonzero) B-semiinvariant functions in the function field $\mathbb{C}(G/H) = \mathbb{C}(X)$, that is,

$$\mathcal{M} = \{ \chi \in \mathfrak{X}(T) : \mathbb{C}(G/H)_{\chi}^{(B)} \neq 0 \},\$$

where $\mathbb{C}(G/H)_{\chi}^{(B)} = \{f \in \mathbb{C}(G/H) : b \cdot f = \chi(b)f \text{ for all } b \in B\}$. Note that every function f_{χ} in $\mathbb{C}(G/H)^{(B)}$ is determined by its weight χ up to constant because $\mathbb{C}(G/H)^B = \mathbb{C}$, that is, any *B*-invariant rational function on *X* is constant. The spherical weight lattice \mathcal{M} is a free abelian group of finite rank. We define the rank of G/H as the rank of the lattice \mathcal{M} . Let $\mathcal{N} = \text{Hom}(\mathcal{M}, \mathbb{Z})$ denote its dual lattice together with the natural pairing $\langle \cdot, \cdot \rangle : \mathcal{N} \times \mathcal{M} \to \mathbb{Z}$.

Let L be a G-linearized ample line bundle on a spherical G-variety X. By the multiplicity-free property of spherical varieties, the algebraic moment polytope $\Delta(X, L)$ encodes the structure of representation of G in the spaces of multi-sections of tensor powers of L.

Definition 2.2. The algebraic moment polytope $\Delta(X, L)$ of L with respect to B is defined as the closure of $\bigcup_{k \in \mathbb{N}} \Delta_k / k$ in $\mathfrak{X}(T) \otimes \mathbb{R}$, where Δ_k is a finite set consisting of (dominant) weights λ such that

$$H^0(X, L^{\otimes k}) = \bigoplus_{\lambda \in \Delta_k} V_G(\lambda).$$

Here, $V_G(\lambda)$ means the irreducible representation of G with highest weight λ .

The algebraic moment polytope $\Delta(X, L)$ for a polarized (spherical) *G*-variety *X* was introduced by Brion in [Bri87] as a purely algebraic version of the Kirwan polytope. This is indeed the convex hull of finitely many points in $\mathfrak{X}(T) \otimes \mathbb{R}$ (see [Bri87]).

2.2. Colors of spherical varieties and types of colors. As the open B-orbit of a spherical variety X is an affine variety, its complement has pure codimension one and is a finite union of B-stable prime divisors.

Definition 2.3. For a spherical homogeneous space G/H, *B*-stable prime divisors in G/H are called *colors* of G/H. We denote by $\mathfrak{D} = \{D_1, \dots, D_k\}$ the set of colors of G/H.

As a *B*-semi-invariant function f_{χ} in $\mathbb{C}(G/H)_{\chi}^{(B)}$ is unique up to constant, we define the *color map* $\rho: \mathfrak{D} \to \mathcal{N}$ by $\langle \rho(D), \chi \rangle = \nu_D(f_{\chi})$ for $\chi \in \mathcal{M}$, where ν_D is the discrete valuation associated to a divisor D, that is, $\nu_D(f)$ is the vanishing order of f along D. Unfortunately, the color map is generally not injective. In addition, every discrete \mathbb{Q} -valued valuation ν of the function field $\mathbb{C}(G/H)$ induces a homomorphism $\hat{\rho}(\nu): \mathcal{M} \to \mathbb{Q}$ defined by $\langle \hat{\rho}(\nu), \chi \rangle = \nu(f_{\chi})$, so that we get a map $\hat{\rho}$: {discrete \mathbb{Q} -valued valuations on G/H} $\to \mathcal{N} \otimes \mathbb{Q}$. Luna and Vust [LV83] showed that the restriction of $\hat{\rho}$ to the set of G-invariant discrete valuations on G/H is injective. From now on, we will regard a G-invariant discrete valuation on G/H as an element of $\mathcal{N} \otimes \mathbb{Q}$ via the map $\hat{\rho}$, and in order to simplify the notation $\hat{\rho}(\nu_E)$ will be written as $\hat{\rho}(E)$ for a G-stable divisor E in X.

Let \mathcal{V} be the set of *G*-invariant discrete \mathbb{Q} -valued valuations on G/H. Since the map $\hat{\rho}$ is injective, we may consider \mathcal{V} as a subset of $\mathcal{N} \otimes \mathbb{Q}$ and it is known that \mathcal{V} is a full-dimensional cosimplicial cone, called the *valuation cone* of G/H. Denote by Σ the set of primitive generators in \mathcal{M} of the extremal rays of the negative of the dual of the valuation cone \mathcal{V} .

Remark 2.4. The normal equivariant embeddings of a given spherical homogeneous space are classified by combinatorial objects called *colored fans*, which generalize the fans appearing in the classification of toric varieties. In a brief way, a colored fan is a finite collection of colored cones, which is a pair $(\mathcal{C}, \mathfrak{R})$ consisting

of $\mathfrak{R} \subset \mathfrak{D}$ and a strictly convex cone $\mathcal{C} \subset \mathcal{N} \otimes \mathbb{Q}$ generated by $\rho(\mathfrak{R})$ and finitely many elements in the valuation cone \mathcal{V} (see [Kno91] for details).

For a simple root α , we denote by P_{α} the corresponding minimal parabolic subgroup containing B. We define the set $\mathfrak{D}(\alpha)$ of colors moved by the minimal parabolic subgroup P_{α} :

$$\mathfrak{D}(\alpha) = \{ D_i \in \mathfrak{D} : P_\alpha \cdot D_i \neq D_i \}.$$

Note that every color is moved by at least one minimal parabolic subgroup since the colors are not G-stable. Now, we recall the types of colors introduced by Brion [Bri97] and Luna [Lun97].

Definition 2.5. Suppose that a color D_i is moved by a minimal parabolic subgroup P_{α} , that is, $D_i \in \mathfrak{D}(\alpha)$.

- If $\alpha \in \Sigma$, we say that D_i is of type **a**.
- If $2\alpha \in \Sigma$, we say that D_i is of type 2a.
- Otherwise, we say that D_i is of type **b**.

Remark 2.6. The type of a color D_i does not depend on the choice of a simple root α such that $D_i \in \mathfrak{D}(\alpha)$.

From [Bri97, Theorem 4.2] and [Lun97, Section 3.6], we can get an explicit expression of the anticanonical divisor $-K_X$ for a spherical variety X.

Proposition 2.7. Let X be an embedding of a spherical homogeneous space G/H. There exists a B-semiinvariant global section $s \in \Gamma(X, K_X^{-1})$ of the anticanonical bundle K_X^{-1} such that

$$\operatorname{div}(s) = \sum_{i=1}^{k} m_i D_i + \sum_{j=1}^{\ell} E_j$$

for colors D_i and G-stable divisors E_j in X. Denoting by $\xi \in \mathfrak{X}(B)$ the B-weight of the section s, the positive integers m_i are obtained in terms of ξ and the types of colors, more explicitly,

$$m_i = \begin{cases} 1 & \text{if } D_i \text{ is of type a or } 2a, \\ \langle \alpha^{\vee}, \xi \rangle & \text{if } D_i \text{ is of type b and } D_i \in \mathfrak{D}(\alpha). \end{cases}$$

Let *P* be the stabilizer of the open orbit of *B* in a spherical *G*-variety *X*. Then *P* is a parabolic subgroup of *G* containing *B*, and admits a unique Levi subgroup *L* containing *T*. We denote by Φ_{P^u} the set of roots of the unipotent radical P^u of *P*, which are the roots of *P* that are not roots of *L*. By [GH15, Proposition 4.2], we have $\xi = \sum_{\alpha \in \Phi_{P^u}} \alpha = \sum_{\alpha \in \Phi^+ \setminus \Phi_L} \alpha$. Alternatively, $\xi = 2\rho_G - 2\rho_L$, where ρ_L is the sum of the fundamental

weights of the root system of L.

Based on the works of Brion [Bri89, Bri97], Gagliardi and Hofscheier [GH15, Section 9] described the (algebraic) moment polytope of the anticanonical line bundle on a Gorenstein Fano spherical variety.

Proposition 2.8. Let X be a Gorenstein Fano embedding of a spherical homogeneous space G/H. If a B-stable Weil divisor $-K_X = \sum_{i=1}^k m_i D_i + \sum_{j=1}^{\ell} E_j$ represents the anticanonical line bundle K_X^{-1} for colors D_i and G-stable divisors E_j in X, the moment polytope $\Delta(X, K_X^{-1})$ is $\xi + Q_X^*$, where the polytope Q_X is the convex hull of the set

$$\left\{\frac{\rho(D_i)}{m_i}: i = 1, \cdots, k\right\} \cup \left\{\hat{\rho}(E_j): j = 1, \cdots, \ell\right\}$$

in $\mathcal{N} \otimes \mathbb{R}$ and its dual polytope Q_X^* is defined as $\{m \in \mathcal{M} \otimes \mathbb{R} : \langle n, m \rangle \geq -1 \text{ for every } n \in Q_X\}$.

2.3. Criterion for existence of Kähler–Einstein metrics on symmetric varieties. For an algebraic group involution θ of a connected reductive algebraic group G, let $G^{\theta} = \{g \in G : \theta(g) = g\}$ be the subgroup consisting of elements fixed by θ . If H is a closed subgroup of G such that the identity component of H coincides with the identity component of G^{θ} , then the homogeneous space G/H is called a *symmetric homogeneous space*. By taking a universal cover of G, we can always assume that G is simply connected. When G is simply connected, G^{θ} is connected (see [Ste68, Section 8.1]) and H is a closed subgroup between G^{θ} and its normalizer $N_G(G^{\theta})$ in G, that is, $G^{\theta} \subset H \subset N_G(G^{\theta})$.

For an algebraic group involution θ of G, a torus T in G is *split* if $\theta(t) = t^{-1}$ for any $t \in T$. A torus T is *maximally split* if T is a θ -stable maximal torus in G which contains a split torus T_s of maximal dimension

among split tori. Then θ descends to an involution of $\mathfrak{X}(T)$ for a maximally split torus T, and the rank of a symmetric homogeneous space G/H is equal to the dimension of a maximal split subtorus T_s of T.

Let $\Phi = \Phi(G, T)$ be the root system of G with respect to a maximally split torus T. By [DCP83, Lemma 1.2], we can take a set of positive roots Φ^+ such that either $\theta(\alpha) = \alpha$ or $\theta(\alpha)$ is a negative root for all $\alpha \in \Phi^+$; then we denote $2\rho_{\theta} = \sum_{\alpha \in \Phi^+ \setminus \Phi^{\theta}} \alpha$, where $\Phi^{\theta} = \{\alpha \in \Phi : \theta(\alpha) = \alpha\}$. The set

$$\Phi_{\theta} = \{ \alpha - \theta(\alpha) : \alpha \in \Phi \setminus \Phi^{\theta} \}$$

is a (possibly non-reduced) root system, which is called the *restricted root system*. Let C^+_{θ} denote the cone generated by positive restricted roots in $\Phi^+_{\theta} = \{\alpha - \theta(\alpha) : \alpha \in \Phi^+ \setminus \Phi^{\theta}\}.$

Vust proved that a symmetric homogeneous space G/H is spherical as a consequence of the Iwasawa decomposition (see [Vus74, Theorem 1 in Section 1.3]). The spherical data of G/H can be described in terms of the restricted root system Φ_{θ} (see [Vus90] and [Tim11, Section 26]).

Proposition 2.9. For a symmetric homogeneous space G/H,

- the spherical weight lattice \mathcal{M} is the lattice $\mathfrak{X}(T/T \cap H)$,
- the valuation cone \mathcal{V} is the negative restricted Weyl chamber determined by $-(\Phi_{\theta}^{+})^{\vee}$ in $\mathcal{N} \otimes \mathbb{Q}$, and
- the image $\rho(\mathfrak{D}(G/H))$ under the color map ρ is the set of simple restricted coroots $\bar{\alpha}_i^{\vee}$.

Definition 2.10. A normal G-variety X together with an equivariant open embedding $G/H \hookrightarrow X$ of a symmetric homogeneous space G/H is called a symmetric variety.

Combining Propositions 2.7, 2.8, 2.9, we can easily compute (algebraic) moment polytopes of the anticanonical line bundles on Fano symmetric varieties.

Corollary 2.11. Let X be a Fano embedding of a symmetric homogeneous space G/H. If all colors in G/H are of type **b** and $\langle \rho(D_i), 2\rho_\theta \rangle = \langle \alpha^{\vee}, 2\rho_\theta \rangle$ for any $D_i \in \mathfrak{D}(\alpha)$, then the moment polytope $\Delta(X_m, K_{X_m}^{-1})$ is the intersection of the positive restricted Weyl chamber and the half-spaces $\{m \in \mathcal{M} \otimes \mathbb{R} : \langle \hat{\rho}(E_j), m - 2\rho_\theta \rangle \ge -1\}$ for all G-stable divisors E_j in X.

Proof. Suppose that a Weil divisor $-K_X = \sum_{i=1}^k m_i D_i + \sum_{j=1}^\ell E_j$ represents the anticanonical line bundle K_X^{-1} for colors D_i and G-stable divisors E_j in X. If a color D_i is of type **b** and $D_i \in \mathfrak{D}(\alpha)$ for a simple root α , then we have $m_i = \langle \alpha^{\vee}, \xi \rangle$ by Proposition 2.7. Note that when X is a symmetric variety associated to an involution θ of G, the B-weight ξ of a general global section in $\Gamma(X, K_X^{-1})^{(B)}$ is equal to $2\rho_\theta = \sum_{\alpha \in \Phi^+ \setminus \Phi^\theta} \alpha$. As $\{\rho(D_1), \cdots, \rho(D_k)\} = \{\bar{\alpha}_1^{\vee}, \cdots, \bar{\alpha}_r^{\vee}\}$ from Proposition 2.9, where r denotes the rank of \mathcal{M} and $k \geq r$,

As $\{\rho(D_1), \dots, \rho(D_k)\} = \{\alpha_1, \dots, \alpha_r\}$ from Proposition 2.9, where r denotes the rank o $\rho(D_i) = \bar{\alpha}_p^{\vee}$ gives an inequality

$$\left\langle \frac{\bar{\alpha}_p^{\vee}}{\langle \alpha^{\vee}, 2\rho_{\theta} \rangle}, m - 2\rho_{\theta} \right\rangle \ge -1 \quad \iff \quad \left\langle \bar{\alpha}_p^{\vee}, m \right\rangle \ge 0 \quad \text{for all } 1 \le p \le r.$$

Thus, the images of all colors D_1, \dots, D_k determine the positive restricted Weyl chamber, and the result follows from Proposition 2.8.

Note that, for smooth Fano symmetric varieties, the linear part of the valuation cone \mathcal{V} is trivial. Thus, Delcroix's criterion for *G*-equivariant K-stability of smooth Fano symmetric varieties in [Del20a, Corollary 5.9] can be slightly strengthened as follows due to the observation in [Gol20, Proposition 5.10].

Proposition 2.12. Let X be a smooth Fano embedding of a symmetric homogeneous space G/H associated to an involution θ of G. Then the following are equivalent:

- (1) X admits a Kähler-Einstein metric.
- (2) X is G-equivariantly uniformly K-stable.
- (3) the barycenter of the moment polytope $\Delta(X, K_X^{-1})$ with respect to the Duistermaat-Heckman measure

$$\prod_{\alpha \in \Phi^+ \setminus \Phi^\theta} \kappa(\alpha, p) \, dp$$

is in the relative interior of the translated cone $2\rho_{\theta} + C_{\theta}^+$, where κ denotes the Cartan-Killing form on the Lie algebra \mathfrak{g} of G. 2.4. Greatest Ricci lower bounds of smooth Fano symmetric varieties. For a Fano manifold X, the greatest Ricci lower bound R(X) of X is defined as the supremum of all $0 \le t \le 1$ such that there exists a Kähler form ω in the first Chern class $c_1(X)$ with $\operatorname{Ric}(\omega) \ge t\omega$. This invariant was first studied by Tian [Tia92], and was explicitely defined by Rubinstein [Rub08, Rub09], where it was called Tian's β -invariant. It was further studied by Székelyhidi [Szé11]; Song and Wang [SW16]. It is shown to be the same as the maximum existence time of Aubin and Yau's continuity path for finding a Kähler–Einstein metric. Given a Kähler form $\omega \in c_1(X)$, the continuity method involves introducing a family of equations

$$\operatorname{Ric}(\omega_t) = t \,\omega_t + (1-t)\omega \quad \text{for } t \in [0,1]$$

depending on a parameter t and finding a Kähler form ω_t to solve the equation, which for t = 1 gives the Einstein equation we want to solve. Hence, in the absence of a Kähler–Einstein metric R(X) may be regarded as a sort of numerical measure of how a Fano manifold X fails to be Kähler–Einstein. Furthermore, the greatest Ricci lower bound R(X) of a Fano manifold X is closely related with the δ -invariant defined by Fujita and Odaka [FO18] using log canonical thresholds of basis type divisors. In fact, the basis log canonical threshold $\delta(X, -K_X)$ and R(X) are related by the formula $R(X) = \min\{1, \delta(X, -K_X)\}$ by [BBJ21, Corollary 7.6] and [CRZ19, Theorem 5.7] (see also [Gol20, Section 3.3]).

By definition, if X admits a Kähler–Einstein metric, then the greatest Ricci lower bound R(X) is equal to 1. However, the converse does not hold. For example, Tian [Tia97] constructed a small deformation of the Mukai–Umemura 3-fold whose general members satisfy R(X) = 1 but are not K-polystable (see [Szé11, Section 3]).

For any toric Fano manifold X, Li [Li11] found an explicit formula for the greatest Ricci lower bound R(X) purely in terms of the moment polytope associated to X. In the case of smooth Fano equivariant compactifications of complex Lie groups, Delcroix [Del17] obtained a formula for the greatest Ricci lower bound where the barycenter of the moment polytope with respect to the Lebesgue measure is replaced by the barycenter with respect to the Duistermaat–Heckman measure. Furthermore, the recent result of Delcroix and Hultgren [DH21] extends the formula to the case of horosymmetric manifolds introduced in [Del20b], which is a class of spherical varieties including smooth horospherical varieties and smooth symmetric varieties.

Now, we state an explicit expression of the greatest Ricci lower bound for smooth symmetric varieties using the following notion.

Definition 2.13. Let X be a smooth Fano embedding of a symmetric homogeneous space G/H. A toric polytope Δ^{tor} of X is defined as the convex hull of the images by the restricted Weyl group \overline{W} of the moment polytope $\Delta(X, K_X^{-1})$.

The following proposition follows from [DH21, Corollary 1.3].

Proposition 2.14. Let X be a smooth Fano embedding of a symmetric homogeneous space G/H associated to an involution θ of G. The greatest Ricci lower bound R(X) of X is equal to

$$\sup\left\{t\in(0,1):2\rho_{\theta}+\frac{t}{1-t}(2\rho_{\theta}-\boldsymbol{bar}_{DH}(\Delta))\in\operatorname{Relint}(\Delta^{tor}-\mathcal{C}_{\theta}^{+})\right\},\$$

where $bar_{DH}(\Delta)$ is the barycenter of the moment polytope $\Delta(X, K_X^{-1})$ with respect to the Duistermaat-Heckman measure and $\operatorname{Relint}(\Delta^{tor} - \mathcal{C}_{\theta}^+)$ means the relative interior of the Minkowski difference $\Delta^{tor} - \mathcal{C}_{\theta}^+$.

We immediately get an elementary geometric expression for the greatest Ricci lower bound from the above result.

Corollary 2.15. Suppose that a smooth Fano embedding X of a symmetric homogeneous space associated to an involution θ of G does not admit a Kähler–Einstein metric. Let A be the point corresponding to $2\rho_{\theta}$ in $\mathfrak{X}(T)$ and C be $bar_{DH}(\Delta)$. If Q is the point at which the half-line starting from the barycenter C in the direction of A intersects the boundary of $\Delta^{tor} - C^+_{\theta}$, then the greatest Ricci lower bound of X is equal to

$$R(X) = \frac{AQ}{\overline{CQ}}.$$



FIGURE 1. Positive restricted roots of type AIII(2, m) for $m \ge 5$.



FIGURE 2. Positive restricted roots of type AIII(2, 4).

3. Wonderful compactifications of symmetric spaces of type AIII

Let us recall the involution on $G = \operatorname{SL}_m(\mathbb{C})$ of type $\operatorname{AIII}(r, m)$ for $m \ge 2r$. Let θ be the involution of G defined by $g \mapsto J_r g J_r$, where $J_r = \begin{pmatrix} O & O & S_r \\ O & I_{m-2r} & O \\ S_r & O & O \end{pmatrix}$ for the $r \times r$ matrix $(S_r)_{i,j} = \delta_{i+j,r+1}$. The subgroup

 G^{θ} fixed by θ is conjugate to the subgroup $S(\operatorname{GL}_r(\mathbb{C}) \times \operatorname{GL}_{m-r}(\mathbb{C}))$. If m > 2r, then the normalizer $N_G(G^{\theta})$ is equal to G^{θ} . However, if m = 2r, then $N_G(G^{\theta})$ is different from G^{θ} and $N_G(G^{\theta})/G^{\theta} \cong \mathbb{Z}_2$.

3.1. Symmetric homogeneous spaces of type AIII(2, m). The homogeneous space G/G^{θ} is considered as the variety parametrizing pairs (V_1, V_2) of linear subspaces in an m-dimensional complex vector space such that dim $V_1 = r$, dim $V_2 = m - r$, and $V_1 \cap V_2 = \{\mathbf{0}\}$. Note that the dimension of the homogeneous space $\mathrm{SL}_m(\mathbb{C})/S(\mathrm{GL}_r(\mathbb{C}) \times \mathrm{GL}_{m-r}(\mathbb{C}))$ is equal to $(m^2 - 1) - \{r^2 + (m - r)^2 - 1\} = 2r(m - r)$. Since the Grassmannians $\mathrm{Gr}(r,m)$ and $\mathrm{Gr}(m - r,m)$ have a natural action of $\mathrm{SL}_m(\mathbb{C})$ as homogeneous spaces, the symmetric space $G/G^{\theta} = \mathrm{SL}_m(\mathbb{C})/S(\mathrm{GL}_r(\mathbb{C}) \times \mathrm{GL}_{m-r}(\mathbb{C}))$ is an open orbit for the diagonal action of $\mathrm{SL}_m(\mathbb{C})$ on the product $\mathrm{Gr}(r,m) \times \mathrm{Gr}(m - r,m)$.

For the root system A_{m-1} of $SL_m(\mathbb{C})$ with respect to the maximal torus T of diagonal matrices, we know that the roots of A_{m-1} consist of algebraic group homomorphisms $\alpha_{i,j}: T \to \mathbb{C}^*$ defined by

$$\alpha_{i,j}(\operatorname{diag}(a_1, a_2, \cdots, a_{m-1}, a_m)) = \frac{a_i}{a_j}$$

for $1 \leq i \neq j \leq m$. For simplicity, the simple roots $\alpha_{i,i+1}$ will be denoted by α_i for $1 \leq i \leq m-1$, and we use the same notations for the corresponding roots of the Lie algebra $\mathfrak{sl}_m(\mathbb{C})$.

Lemma 3.1. Let θ be the involution on $SL_m(\mathbb{C})$ of type AIII(2,m) for $m \ge 4$. The restricted root system Φ_{θ} is as follow:

$$\Phi_{\theta} = \begin{cases} \{\pm \alpha_{1,m}, \pm \alpha_{2,m-1}, \pm 2\alpha_{1,m}, \pm 2\alpha_{2,m-1}, \pm (\alpha_{1,m} - \alpha_{2,m-1}), \pm (\alpha_{1,m} + \alpha_{2,m-1})\} \text{ of type } \mathsf{BC}_2 & \text{when } m \ge 5 \\ \{\pm 2\alpha_{1,4}, \pm 2\alpha_{2,3}, \pm (\alpha_{1,4} - \alpha_{2,3}), \pm (\alpha_{1,4} + \alpha_{2,3})\} \text{ of type } \mathsf{C}_2 & \text{when } m = 4 \end{cases}$$

Moreover, positive restricted roots $\alpha_{1,m}, \alpha_{2,m-1}, \alpha_{1,m} - \alpha_{2,m-1}, \alpha_{1,m} + \alpha_{2,m-1}$ have multiplicities 2(m-4), 2(m-4), 2, 2, respectively.

Proof. Since the involution θ of type AIII(2, m) acts on diagonal matrices as

 $\theta(\operatorname{diag}(a_1, a_2, a_3, \cdots, a_{m-2}, a_{m-1}, a_m)) = \operatorname{diag}(a_m, a_{m-1}, a_3, \cdots, a_{m-2}, a_2, a_1),$

the roots $\alpha_{i,j}$ for $3 \leq i,j \leq m-2$ are fixed by the action of θ . In particular, the cardinality of the set $\Phi_{\theta}^+ = \Phi^+ \setminus \Phi^{\theta}$ is equal to $\binom{m}{2} - \binom{m-4}{2} = 2(2m-5)$.

(i) $\alpha_{1,2} - \theta(\alpha_{1,2}) = \alpha_{1,2} - \alpha_{m,m-1} = \alpha_{1,m} - \alpha_{2,m-1} = \alpha_{m-1,m} - \alpha_{2,1} = \alpha_{m-1,m} - \theta(\alpha_{m-1,m})$ and $\alpha_{1,m-1} - \theta(\alpha_{1,m-1}) = \alpha_{1,m-1} - \alpha_{m,2} = \alpha_{1,m} + \alpha_{2,m-1} = \alpha_{2,m} - \alpha_{m-1,1} = \alpha_{2,m} - \theta(\alpha_{2,m}).$

(ii) For $3 \le k \le m-2$, $\alpha_{1,k} - \theta(\alpha_{1,k}) = \alpha_{1,k} - \alpha_{m,k} = \alpha_{1,m} = \alpha_{k,m} - \alpha_{k,1} = \alpha_{k,m} - \theta(\alpha_{k,m})$ and $\alpha_{2,k} - \theta(\alpha_{2,k}) = \alpha_{2,k} - \alpha_{m-1,k} = \alpha_{2,m-1} = \alpha_{k,m-1} - \alpha_{k,2} = \alpha_{k,m-1} - \theta(\alpha_{k,m-1})$.

(iii)
$$\alpha_{1,m} - \theta(\alpha_{1,m}) = \alpha_{1,m} - \alpha_{m,1} = 2\alpha_{1,m}$$
 and $\alpha_{2,m-1} - \theta(\alpha_{2,m-1}) = \alpha_{2,m-1} - \alpha_{m-1,2} = 2\alpha_{2,m-1}$.

Lemma 3.2. For the involution θ on $SL_m(\mathbb{C})$ of type AIII(2,m), we have $2\rho_\theta = (m-1)\alpha_{1,m} + (m-3)\alpha_{2,m-1}$.

Proof. From the proof of Lemma 3.1, if $m \ge 5$ then we have

$$2\rho_{\theta} = \sum_{\alpha \in \Phi^+ \setminus \Phi^{\theta}} \alpha = \sum_{i=1}^{m-1} \sum_{j=1}^{i} \alpha_j + \sum_{i=2}^{m-1} \sum_{j=2}^{i} \alpha_j + \sum_{i=1}^{m-3} \sum_{j=1}^{i} \alpha_{m-j} + \sum_{i=2}^{m-3} \sum_{j=2}^{i} \alpha_{m-j}$$
$$= (m-1)\alpha_{1,m} + (m-3)\alpha_{2,m-1}$$

by the straightforward computations. When m = 4, as $\Phi^{\theta} = \emptyset$ we can easily check that

 $2\rho_{\theta} = \alpha_{1,4} + \alpha_{2,3} + (\alpha_{1,2} + \alpha_{2,4}) + (\alpha_{1,3} + \alpha_{3,4}) = 3\alpha_{1,4} + \alpha_{2,3}.$

Alternatively, it is possible to calculate $2\rho_{\theta}$ from the description of the restricted root system Φ_{θ} in Lemma 3.1 because the sum of the positive restricted roots with multiplicities is equal to $2 \times 2\rho_{\theta}$.

From the detailed descriptions in [Vus90, Section 6.1], if m > 2r then we know that the symmetric homogeneous space $\operatorname{SL}_m(\mathbb{C})/S(\operatorname{GL}_r(\mathbb{C}) \times \operatorname{GL}_{m-r}(\mathbb{C}))$ of type $\operatorname{AIII}(r,m)$ has r+1 colors $D_1, \cdots D_{r-1}, D^+, D^$ such that

$$\mathcal{D}(\alpha_1) = \mathcal{D}(\alpha_{m-1}) = \{D_1\}, \cdots, \mathcal{D}(\alpha_{r-1}) = \mathcal{D}(\alpha_{m-r+1}) = \{D_{r-1}\}, \mathcal{D}(\alpha_r) = \{D^+\} \text{ and } \mathcal{D}(\alpha_{m-r}) = \{D^-\}.$$

On the other hand, if m = 2r, that is, m - r = r, then $\alpha_{m-r} = \alpha_r$ so that $D^+ = D^-$; hence the symmetric homogeneous space $\operatorname{SL}_{2r}(\mathbb{C})/S(\operatorname{GL}_r(\mathbb{C}) \times \operatorname{GL}_r(\mathbb{C}))$ has r colors $D_1, \cdots D_{r-1}, D^+ = D^-$.

Lemma 3.3. For colors D_1 , D^+ , D^- in the symmetric space $\operatorname{SL}_m(\mathbb{C})/S(\operatorname{GL}_2(\mathbb{C}) \times \operatorname{GL}_{m-2}(\mathbb{C}))$ of type $\operatorname{AIII}(2,m)$, we have $\mathcal{D}(\alpha_1) = \mathcal{D}(\alpha_{m-1}) = \{D_1\}$, $\mathcal{D}(\alpha_2) = \{D^+\}$ and $\mathcal{D}(\alpha_{m-2}) = \{D^-\}$. Moreover, all colors D_1 , D^+ , D^- are of type b.

Proof. The first statement follows from [Vus90, Section 6.1]. As the valuation cone \mathcal{V} of a symmetric homogeneous space is the negative restricted Weyl chamber determined by $-(\Phi_{\theta}^+)^{\vee}$ in $\mathcal{N} \otimes \mathbb{Q}$ by Proposition 2.9, \mathcal{V} is the cone generated by two restricted coroots $\bar{\alpha}_{1,m}^{\vee}$ and $\bar{\alpha}_{1,m-1}^{\vee}$. Thus, the primitive generators in \mathcal{M} of the extremal rays of $-\mathcal{V}^{\vee}$ are $\bar{\alpha}_{1,2} = \alpha_{1,m} - \alpha_{2,m-1}$ and $\bar{\alpha}_{2,3} = \alpha_{2,m-1}$, that is, $\Sigma = \{\alpha_{1,m} - \alpha_{2,m-1}, \alpha_{2,m-1}\}$. For any simple root α , neither α nor 2α belongs to Σ , which means that all colors are of type **b**.

3.2. Wonderful compactifications of symmetric homogeneous spaces of type AIII(2, m). Wonderful varieties are *G*-varieties where the orbit structure is particularly nice. These varieties first appeared as compactifications with remarkable properties of symmetric homogeneous spaces in the work of De Concini and Procesi [DCP83], and were generalized by Luna [Lun96].

Definition 3.4. An irreducible *G*-variety is called *wonderful* of rank r if it is smooth, complete, and it has exactly r prime *G*-stable divisors such that they intersect transversally and their intersection is a (non-empty) single *G*-orbit.

By definition, a wonderful G-variety of rank r has 2^r G-orbits; among them exactly one is open, and exactly one is closed. Note that any wonderful variety is spherical from [Lun96]. In fact, wonderful varieties play a crucial role in the theory of spherical varieties; see [Lun01, BP16, Pez18] for details.

Example 3.5. (1) For a maximal torus $T \subset SL_2(\mathbb{C})$, the wonderful compactification of the symmetric space $SL_2(\mathbb{C})/T$ is exactly $\mathbb{P}^1 \times \mathbb{P}^1$.

(2) The projective plane $\mathbb{P}^2 = \mathbb{P}(\text{Sym}^2(\mathbb{C}^2))$ under the action of $\text{SL}_2(\mathbb{C})$ is a wonderful variety of rank 1, that is, the wonderful compactification of the symmetric space $\text{SL}_2(\mathbb{C})/N_{\text{SL}_2}(T)$.

(3) A classical example studied by De Concini and Procesi [DCP83] is the variety of complete conics $X = \{([A], [B]) \in \mathbb{P}(Mat_{3\times3}) \times \mathbb{P}(Mat_{3\times3}) : AB \in \mathbb{C}I_{3\times3}\}$. Then X is the wonderful compactification of the homogeneous space $SL_3(\mathbb{C})/SO_3(\mathbb{C})Z(SL_3(\mathbb{C}))$, which is the space of smooth conics in \mathbb{P}^2 . Since X has the two SL_3 -stable prime divisors consisting of non-invertible matrices, X is wonderful of rank 2.

For $m \geq 5$, the wonderful compactification of the symmetric space $\operatorname{SL}_m(\mathbb{C})/S(\operatorname{GL}_2(\mathbb{C}) \times \operatorname{GL}_{m-2}(\mathbb{C}))$ is the blow-up of the product of Grassmannians $\operatorname{Gr}(2,m) \times \operatorname{Gr}(m-2,m)$ along the closed orbit $\mathcal{F}\ell(2,m-2;m)$. In the case of m = 4, the symmetric homogeneous space $\operatorname{SL}_4(\mathbb{C})/N_{\operatorname{SL}_4(\mathbb{C})}(S(\operatorname{GL}_2(\mathbb{C}) \times \operatorname{GL}_2(\mathbb{C})))$ admits the wonderful compactification. As the rank of the symmetric space is two, its wonderful compactification has the two *G*-stable divisors Y_1, Y_2 .

Proposition 3.6. Let X_m be the wonderful compactification of the symmetric homogeneous space of type AIII(2, m) for $m \ge 4$. If $m \ge 5$, then there is a B-semi-invariant section of the anticanonical line bundle $K_{X_m}^{-1}$ with Weil divisor $-K_{X_m} = Y_1 + Y_2 + 2D_1 + (m-3)D^+ + (m-3)D^-$, and the B-weight of this section is equal to $(m-1)\alpha_{1,m} + (m-3)\alpha_{2,m-1}$. Moreover, for the wonderful compactification X_4 of the symmetric homogeneous space of type AIII(2, 4), we have $-K_{X_4} = Y_1 + Y_2 + 2D_1 + D^+$.

Proof. First, let us consider the case $m \ge 5$. By [GH15, Theorem 1.2], there is a *B*-semi-invariant section *s* of the anticanonical line bundle $K_{X_m}^{-1}$ with Weil divisor div $(s) = Y_1 + Y_2 + m_1D_1 + m^+D^+ + m^-D^-$ and its *B*-weight is equal to $2\rho_{\theta} = (m-1)\alpha_{1,m} + (m-3)\alpha_{2,m-1}$ from Lemma 3.2. As each color is of type b from Lemma 3.3, we obtain the coefficients

$$\begin{split} m_1 &= \langle \alpha_1^{\vee}, 2\rho_{\theta} \rangle = \langle \alpha_{m-1}^{\vee}, 2\rho_{\theta} \rangle = 2\\ m^+ &= \langle \alpha_2^{\vee}, 2\rho_{\theta} \rangle = m - 3,\\ m^- &= \langle \alpha_{m-2}^{\vee}, 2\rho_{\theta} \rangle = m - 3 \end{split}$$

by [GH15, Theorem 1.5]. Thus, we get $-K_{X_m} = Y_1 + Y_2 + 2D_1 + (m-3)D^+ + (m-3)D^-$. Similarly, as $-K_{X_4} = Y_1 + Y_2 + m_1D_1 + m^+D^+$, Lemmas 3.2 and 3.3 imply the result.

Two real roots $\alpha_{1,m}$ and $\alpha_{2,m-1}$ generate the spherical weight lattice \mathcal{M} .

Proposition 3.7. Let X_m be the wonderful compactification of the symmetric homogeneous space of type AIII(2, m) for $m \ge 4$. The moment polytope $\Delta_m = \Delta(X_m, K_{X_m}^{-1})$ of the anticanonical line bundle $K_{X_m}^{-1}$ is the convex hull of four points 0, $m\alpha_{1,m}, m\alpha_{1,m} + (m-3)\alpha_{2,m-1}, (m-\frac{3}{2})\alpha_{1,m} + (m-\frac{3}{2})\alpha_{2,m-1}$ in $\mathcal{M} \otimes \mathbb{R}$.

Proof. From the description in [Vus90, Section 6.1], we know that the three colors D_1, D^+, D^- and the G-stable divisors Y_1, Y_2 in X_m have the images

$$\rho(D_1) = \bar{\alpha}_{1,2}^{\vee} = \frac{1}{2} \alpha_{1,m}^{\vee} - \frac{1}{2} \alpha_{2,m-1}^{\vee},$$

$$\rho(D^+) = \rho(D^-) = \bar{\alpha}_{2,m-1}^{\vee} = \frac{1}{2} \alpha_{2,m-1}^{\vee},$$

$$\hat{\rho}(Y_1) = -\bar{\alpha}_{1,m-1}^{\vee} = -\frac{1}{2} \alpha_{1,m}^{\vee},$$

$$\hat{\rho}(Y_2) = -\bar{\alpha}_{1,m}^{\vee} = -\frac{1}{2} \alpha_{1,m}^{\vee} - \frac{1}{2} \alpha_{2,m-1}^{\vee}$$

in \mathcal{N} , respectively.

Using Corollary 2.11 and Proposition 3.6, $\frac{1}{2}\rho(D_1)$, $\frac{1}{m-3}\rho(D^+) = \frac{1}{m-3}\rho(D^-)$, $\hat{\rho}(Y_1)$ and $\hat{\rho}(Y_2)$ are used as inward-pointing facet normal vectors of the moment polytope $\Delta(X_m, K_{X_m}^{-1})$. First, $\frac{1}{m-3}\rho(D^+) = \frac{1}{m-3}\rho(D^-) = \frac{1}{m-3}\rho(D^-)$

 $\frac{1}{2(m-3)}\alpha_{2,m-1}^{\vee}$ gives an inequality

$$\left\langle \frac{1}{2(m-3)} \alpha_{2,m-1}^{\vee}, x \alpha_{1,m} + y \alpha_{2,m-1} - 2\rho_{\theta} \right\rangle = \frac{1}{m-3} \{y - (m-3)\} \ge -1$$

because $2\rho_{\theta} = (m-1)\alpha_{1,m} + (m-3)\alpha_{2,m-1}$. Similarly, as $\frac{1}{2}\rho(D_1) = \frac{1}{4}\alpha_{1,m}^{\vee} - \frac{1}{4}\alpha_{2,m-1}^{\vee}$ gives an inequality

$$\left\langle \frac{1}{4}\alpha_{1,m}^{\vee} - \frac{1}{4}\alpha_{2,m-1}^{\vee}, x\alpha_{1,m} + y\alpha_{2,m-1} - 2\rho_{\theta} \right\rangle = \frac{1}{2}[x - (m-1) - \{y - (m-3)\}] \ge -1,$$

from which we get a domain $\{x\alpha_{1,m} + y\alpha_{2,m-1} \in \mathcal{M} \otimes \mathbb{R} : x \geq y\}$. Thus, the images of two colors D_1, D_2 determine the positive restricted Weyl chamber. Next, for two *G*-stable divisors $\hat{\rho}(Y_1) = -\frac{1}{2}\alpha_{1,m}^{\vee}$ and $\hat{\rho}(Y_2) = -\frac{1}{2}\alpha_{1,m}^{\vee} - \frac{1}{2}\alpha_{2,m-1}^{\vee}$ give inequalities

$$\left\langle -\frac{1}{2}\alpha_{1,m}^{\vee}, x\alpha_{1,m} + y\alpha_{2,m-1} - 2\rho_{\theta} \right\rangle = -\{x - (m-1)\} \ge -1$$

and

$$\left\langle -\frac{1}{2}\alpha_{1,m}^{\vee} - \frac{1}{2}\alpha_{2,m-1}^{\vee}, x\alpha_{1,m} + y\alpha_{2,m-1} - 2\rho_{\theta} \right\rangle = -\{x - (m-1)\} - \{y - (m-3)\} \ge -1.$$

Therefore, the moment polytope $\Delta(X_m, K_{X_m}^{-1})$ is the intersection of the positive restricted Weyl chamber and two half-spaces $\{x\alpha_{1,m} + y\alpha_{2,m-1} \in \mathcal{M} \otimes \mathbb{R} : x \leq m\}, \{x\alpha_{1,m} + y\alpha_{2,m-1} \in \mathcal{M} \otimes \mathbb{R} : x + y \leq 2m - 3\}$. \Box

For $p = x\alpha_{1,m} + y\alpha_{2,m-1} \in \mathcal{M} \otimes \mathbb{R}$, the Duistermaat–Heckman measure on $\mathcal{M} \otimes \mathbb{R}$ is given as $\prod_{\alpha \in \Phi^+ \setminus \Phi^\theta} \kappa(\alpha, p) \, dp = x^{2m-8} (2x) y^{2m-8} (2y) (x+y)^2 (x-y)^2 \, dx dy = 2^2 x^{2m-7} y^{2m-7} (x+y)^2 (x-y)^2 \, dx dy$

$$=: P_{DH}(x, y) dx dy$$

up to a multiplicative constant.

Example 3.8. In the case of m = 5, we compute the volume of the moment polytope Δ_5

$$\begin{aligned} \operatorname{Vol}_{DH}(\Delta_5) &= \int_0^{\frac{7}{2}} \int_0^x 4x^3 y^3 (x+y)^2 (x-y)^2 \, dy dx + \int_{\frac{7}{2}}^5 \int_0^{7-x} 4x^3 y^3 (x+y)^2 (x-y)^2 \, dy dx \\ &= \frac{1}{72} \left(\frac{7}{2}\right)^{12} + \left[\frac{1}{72} x^{12} - \frac{5488}{27} x^9 + 4802 x^8 - 48020 x^7 + \frac{2235331}{9} x^6 - \frac{3294172}{5} x^5 + \frac{5764801}{8} x^4\right]_{\frac{7}{2}}^5 \\ &= \frac{391880669}{360} \end{aligned}$$

and the barycenter of Δ_5 with respect to the Duistermaat–Heckman measure

$$\begin{aligned} \mathbf{bar}_{DH}(\Delta_5) &= (\bar{x}, \bar{y}) = \frac{1}{\mathrm{Vol}_{DH}(\Delta_5)} \left(\int_{\Delta_5} x \prod_{\alpha \in \Phi^+ \setminus \Phi^\theta} \kappa(\alpha, p) \, dp, \int_{\Delta_5} y \prod_{\alpha \in \Phi^+ \setminus \Phi^\theta} \kappa(\alpha, p) \, dp \right) \\ &= \left(\frac{1426329931935}{326044716608}, \frac{4418316612263}{2282313016256} \right) \approx (4.37, 1.94). \end{aligned}$$

Since $\bar{x} > 4$ and $\bar{x} + \bar{y} > 6$, $\mathbf{bar}_{DH}(\Delta_5)$ is in the relative interior of the translated cone $2\rho_{\theta} + \mathcal{C}^+_{\theta}$ (see Figure 3). Therefore, the wonderful compactification X_5 of the symmetric homogeneous space $\mathrm{SL}_5(\mathbb{C})/S(\mathrm{GL}_2(\mathbb{C}) \times \mathrm{GL}_3(\mathbb{C}))$ admits a Kähler–Einstein metric by Proposition 2.12.

Similarly, we obtain the barycenter of the moment polytope Δ_6 with respect to the Duistermaat–Heckman measure

$$\mathbf{bar}_{DH}(\Delta_6) = (\bar{x}, \bar{y}) = \left(\frac{421619272419}{78063549568}, \frac{226450079005}{78063549568}\right) \approx (5.40, 2.90)$$

Since $\bar{x} > 5$ and $\bar{x} + \bar{y} > 8$, $\mathbf{bar}_{DH}(\Delta_6)$ is in the relative interior of the translated cone $2\rho_{\theta} + \mathcal{C}^+_{\theta}$ (see Figure 4). Therefore, the wonderful compactification X_6 of the symmetric homogeneous space $\mathrm{SL}_6(\mathbb{C})/S(\mathrm{GL}_2(\mathbb{C}) \times \mathrm{GL}_4(\mathbb{C}))$ admits a Kähler–Einstein metric by Proposition 2.12.



FIGURE 4. $\Delta_6 = \Delta(X_6, K_{X_6}^{-1})$

Proposition 3.9. Let (\bar{x}, \bar{y}) be the barycenter $\operatorname{bar}_{DH}(\Delta_m)$ of the moment polytope $\Delta(X_m, K_{X_m}^{-1})$ with respect to the Duistermaat–Heckman measure. Then we have $\bar{x} > m - 1$ and $\bar{x} + \bar{y} > 2m - 4$ for all $m \ge 4$.

Proof. From Proposition 3.7, we can divide the moment polytope Δ_m into two regions Ω_1 and Ω_2 :

$$\Omega_1 := \{ s(m,t) : 0 \le s \le 1, \ 0 \le t \le m-3 \},$$

$$\Omega_2 := \{ s(t,2m-3-t) : 0 \le s \le 1, \ m-\frac{3}{2} \le t \le m \}.$$

Denoting $(\bar{x}_i, \bar{y}_i) = \mathbf{bar}_{DH}(\Omega_i)$ for each i = 1, 2, in order to prove $\bar{x} > m - 1$ it suffices to show that $\bar{x}_i > m - 1$ for each i. Indeed, $\bar{x}_1 > m - 1$ and $\bar{x}_2 > m - 1$ imply that $\bar{x} > m - 1$ since

$$\bar{x} = \frac{1}{\operatorname{Vol}_{DH}(\Delta_m)} \int_{\Delta_m} x P_{DH}(x, y) \, dx dy = \frac{1}{\operatorname{Vol}_{DH}(\Omega_1 \cup \Omega_2)} \int_{\Omega_1 \cup \Omega_2} x P_{DH}(x, y) \, dx dy$$
$$= \frac{\bar{x}_1 \operatorname{Vol}_{DH}(\Omega_1) + \bar{x}_2 \operatorname{Vol}_{DH}(\Omega_2)}{\operatorname{Vol}_{DH}(\Omega_1) + \operatorname{Vol}_{DH}(\Omega_2)}.$$

(i) Using the parametrization x = ms, y = st of Ω_1 , we get the volume form

$$dx \wedge dy = d(ms) \wedge d(st) = m \, ds \wedge s \, dt = ms \, ds \wedge dt$$

Then we compute the volume of Ω_1 with respect to the Duistermaat–Heckman measure:

$$\begin{aligned} \operatorname{Vol}_{DH}(\Omega_1) &= \int_{\Omega_1} P_{DH}(x, y) \, dx dy = \int_{\Omega_1} 2^2 x^{2m-7} y^{2m-7} (x+y)^2 (x-y)^2 \, dx dy \\ &= \int_0^1 \int_0^{m-3} 4(ms)^{2m-7} (st)^{2m-7} \{s(m+t)\}^2 \{s(m-t)\}^2 \, ms \, dt ds \\ &= \int_0^1 \int_0^{m-3} 4m^{2m-6} s^{4m-9} t^{2m-7} \{(m+t)(m-t)\}^2 \, dt ds \\ &= 4m^{2m-6} \int_0^1 s^{4m-9} \, ds \int_0^{m-3} t^{2m-7} \{(m+t)(m-t)\}^2 \, dt \\ &= \frac{4m^{2m-6}}{4m-8} \int_0^{m-3} t^{2m-7} (m+t)^2 (m-t)^2 \, dt. \end{aligned}$$

Hence, when $m \ge 4$ we see that

$$\bar{x}_{1} = \frac{1}{\operatorname{Vol}_{DH}(\Omega_{1})} \int_{\Omega_{1}} x P_{DH}(x, y) \, dx \, dy$$

= $\frac{1}{\operatorname{Vol}_{DH}(\Omega_{1})} \int_{0}^{1} \int_{0}^{m-3} 4m^{2m-5} s^{4m-8} t^{2m-7} \{(m+t)(m-t)\}^{2} \, dt \, ds$
= $\frac{m(4m-8)}{4m-7} > m-1.$

(ii) Using the parametrization x = st, y = s(2m - 3 - t) of Ω_2 , we get the volume form

$$dx \wedge dy = d(st) \wedge d(s(2m - 3 - t)) = (s \, dt + t \, ds) \wedge [(2m - 3 - t)ds + s(-dt)]$$

= $s(2m - 3 - t) \, dt \wedge ds - ts \, ds \wedge dt = s(2m - 3) \, dt \wedge ds.$

Then we compute the volume of Ω_2 with respect to the Duistermaat–Heckman measure:

$$\begin{aligned} \operatorname{Vol}_{DH}(\Omega_2) &= \int_{\Omega_2} P_{DH}(x,y) \, dx dy = \int_{\Omega_2} 2^2 x^{2m-7} y^{2m-7} (x+y)^2 (x-y)^2 \, dx dy \\ &= \int_0^1 \int_{m-\frac{3}{2}}^m 4(st)^{2m-7} \{s(2m-3-t)\}^{2m-7} \{s(2m-3)\}^2 \{s(2m-3-2t)\}^2 \, s(2m-3) \, dt ds \\ &= \int_0^1 \int_{m-\frac{3}{2}}^m 4(2m-3)^3 s^{4m-9} t^{2m-7} (2m-3-t)^{2m-7} (2m-3-2t)^2 \, dt ds \\ &= 4(2m-3)^3 \int_0^1 s^{4m-9} \, ds \int_{m-\frac{3}{2}}^m \{t(2m-3-t)\}^{2m-7} (2m-3-2t)^2 \, dt \\ &= \frac{4(2m-3)^3}{4m-8} \int_{m-\frac{3}{2}}^m \{t(2m-3-t)\}^{2m-7} (2m-3-2t)^2 \, dt. \end{aligned}$$

As x + y = s(2m - 3) on Ω_2 , we can verify the inequality

$$\bar{x}_{2} + \bar{y}_{2} = \frac{1}{\operatorname{Vol}_{DH}(\Omega_{2})} \int_{\Omega_{2}} (x+y) P_{DH}(x,y) \, dx \, dy$$

$$= \frac{1}{\operatorname{Vol}_{DH}(\Omega_{2})} \cdot 4(2m-3)^{4} \int_{0}^{1} s^{4m-8} \, ds \int_{m-\frac{3}{2}}^{m} \{t(2m-3-t)\}^{2m-7} (2m-3-2t)^{2} \, dt.$$

$$= (2m-3) \cdot \frac{4m-8}{4m-7} = \frac{4m-6}{4m-7} \cdot (2m-4) > 2m-4.$$

Since

$$\bar{x}_{2} = \frac{1}{\operatorname{Vol}_{DH}(\Omega_{2})} \int_{\Omega_{2}} x P_{DH}(x, y) \, dx \, dy$$
$$= \frac{4m - 8}{4m - 7} \cdot \frac{\int_{m - \frac{3}{2}}^{m} t\{t(2m - 3 - t)\}^{2m - 7}(2m - 3 - 2t)^{2} \, dt}{\int_{m - \frac{3}{2}}^{m} \{t(2m - 3 - t)\}^{2m - 7}(2m - 3 - 2t)^{2} \, dt}$$

we conclude that $\bar{x}_2 > m - 1$ by Lemma 3.10.

(iii) To show $\bar{x} + \bar{y} > 2m - 4$, for $m \ge 4$ we will verify the inequality

$$\bar{x} + \bar{y} = \frac{\int_{\Omega_1} (x+y) P_{DH}(x,y) dxdy + \int_{\Omega_2} (x+y) P_{DH}(x,y) dxdy}{\operatorname{Vol}_{DH}(\Omega_1) + \operatorname{Vol}_{DH}(\Omega_2)} \\ = \frac{\frac{4m^{2m-6}}{4m-7} \int_0^{m-3} t^{2m-7} (m+t)^3 (m-t)^2 dt + \frac{4(2m-3)^4}{4m-7} \int_{m-\frac{3}{2}}^m \{t(2m-3-t)\}^{2m-7} (2m-3-2t)^2 dt + \frac{4(2m-3)^3}{4m-8} \int_0^m t^{2m-7} (m+t)^2 (m-t)^2 dt + \frac{4(2m-3)^3}{4m-8} \int_{m-\frac{3}{2}}^m \{t(2m-3-t)\}^{2m-7} (2m-3-2t)^2 dt + 2m-4 dt + 2m-4$$

To do this, it suffices to show the following inequality

$$m^{2m-6} \int_0^{m-3} t^{2m-7} (m+t)^2 (m-t)^2 (2t-2m+7) dt + (2m-3)^3 \int_{m-\frac{3}{2}}^m \{t(2m-3-t)\}^{2m-7} (2m-3-2t)^2 dt > 0.$$

From Lemma 3.11, we get the result as claimed.

From Lemma 3.11, we get the result as claimed.

Now, we complete the proof by showing the remaining two lemmas.

Lemma 3.10. For $m \ge 4$, the inequality

$$\int_{m-\frac{3}{2}}^{m} \left(\frac{4m-8}{4m-7}t - (m-1)\right) \left\{t(2m-3-t)\right\}^{2m-7} (2m-3-2t)^2 dt > 0$$

holds.

Proof. For a fixed m, since t(2m-3-t) is a decreasing function on the interval $[m-\frac{3}{2},m]$, we have t(2m-3-t) > m(m-3) for $m - \frac{3}{2} < t < m$. Thus,

$$\begin{split} I(m) &:= \int_{m-\frac{3}{2}}^{m} \left(\frac{4m-8}{4m-7}t - (m-1)\right) \{t(2m-3-t)\}^{2m-7}(2m-3-2t)^2 \, dt \\ &> \int_{m-\frac{3}{2}}^{m} \left(\frac{4m-8}{4m-7}t - (m-1)\right) \{m(m-3)\}^{2m-7}(2m-3-2t)^2 \, dt \\ &= \{m(m-3)\}^{2m-7} \int_{m-\frac{3}{2}}^{m} \left(\frac{4m-8}{4m-7}t - (m-1)\right) (2m-3-2t)^2 \, dt \\ &= \{m(m-3)\}^{2m-7} \cdot \frac{4m-8}{4m-7} \int_{0}^{\frac{3}{2}} \left(u + m - \frac{3}{2} - (m-1)\frac{4m-7}{4m-8}\right) (-2u)^2 \, du \qquad (\text{let } u := t - (m-\frac{3}{2})). \end{split}$$

Putting $\varphi(m) = (m-1)\frac{4m-7}{4m-8} - m + \frac{3}{2} = \frac{3}{4} + \frac{1}{4(m-2)}$, it suffices to show that $\int_0^2 u^2(u-\varphi(m)) \, du > 0$. As $\frac{3}{4} < \varphi(m) \le \frac{7}{8}$ for $m \ge 4$, we obtain that $\int_0^{\frac{3}{2}} u^2(u-\varphi(m)) \, du = \left(\frac{3}{2}\right)^3 \left(\frac{3}{8} - \frac{\varphi(m)}{3}\right) > 0$.

Lemma 3.11. For $m \ge 4$, the inequality

$$m^{2m-6} \int_{0}^{m-3} t^{2m-7} (m+t)^2 (m-t)^2 (2t-2m+7) dt + (2m-3)^3 \int_{m-\frac{3}{2}}^{m} \{t(2m-3-t)\}^{2m-7} (2m-3-2t)^2 dt > 0$$
 holds.

Proof. From the direct computation, we get

$$\begin{split} &\int_{0}^{m-3} t^{2m-7} (m+t)^2 (m-t)^2 (2t-2m+7) \, dt \\ &= \int_{0}^{m-3} t^{2m-7} \{ 2t^5 - (2m-7)t^4 - 4m^2t^3 + 2(2m-7)m^2t^2 + 2m^4t - (2m-7)m^4 \} \, dt \\ &= (m-3)^{2m-6} \cdot \frac{-512m^7 + 9024m^6 - 62848m^5 + 233160m^4 - 507384m^3 + 656100m^2 - 471420m + 145800}{(2m-1)(2m-2)(2m-3)(2m-4)(2m-5)(2m-6)} \\ &=: (m-3)^{2m-6}q(m) \end{split}$$

As t(2m - 3 - t) > m(m - 3) for $m - \frac{3}{2} < t < m$, we have

$$\begin{split} &m^{2m-6} \int_{0}^{m-3} t^{2m-7} (m+t)^2 (m-t)^2 (2t-2m+7) \, dt + (2m-3)^3 \int_{m-\frac{3}{2}}^{m} \{t(2m-3-t)\}^{2m-7} (2m-3-2t)^2 \, dt \\ &> m^{2m-6} \int_{0}^{m-3} t^{2m-7} (m+t)^2 (m-t)^2 (2t-2m+7) \, dt + (2m-3)^3 \int_{m-\frac{3}{2}}^{m} \{m(m-3)\}^{2m-7} (2m-3-2t)^2 \, dt \\ &= m^{2m-6} (m-3)^{2m-6} q(m) + \frac{9}{2} \{m(m-3)\}^{2m-7} (2m-3)^3 \\ &= m^{2m-7} (m-3)^{2m-7} \left\{ m(m-3)q(m) + \frac{9}{2} (2m-3)^3 \right\}. \end{split}$$

We can check that a polynomial

$$p(m) := (2m-1)(2m-2)(2m-3)(2m-4)(2m-5)(2m-6)\left\{m(m-3)q(m) + \frac{9}{2}(2m-3)^3\right\}$$

= $m(m-3)(-512m^7 + 9024m^6 - 62848m^5 + 233160m^4 - 507384m^3 + 656100m^2 - 471420m + 145800)$
+ $\frac{9}{2}(2m-1)(2m-2)(2m-3)(2m-4)(2m-5)(2m-6)(2m-3)^3$

of degree 9 is increasing on $[4, \infty)$, and p(4) > 0. Hence, we conclude the result.

Theorem 3.12. All wonderful compactifications X_m of symmetric homogeneous spaces of type AIII(2, m) for $m \ge 4$ are $SL_m(\mathbb{C})$ -equivariantly uniformly K-stable.

Proof. Recall that $2\rho_{\theta} = (m-1)\alpha_{1,m} + (m-3)\alpha_{2,m-1} = (m-1,m-3)$ from Lemma 3.2 and the cone \mathcal{C}^+_{θ} is generated by the vectors (1,-1) and (0,1). As $\bar{x} > m-1$ and $\bar{x} + \bar{y} > 2m-4$ by Proposition 3.9, the barycenter $\mathbf{bar}_{DH}(\Delta_m)$ is in the relative interior of the translated cone $2\rho_{\theta} + \mathcal{C}^+_{\theta}$. Therefore, X_m is $\mathrm{SL}_m(\mathbb{C})$ -equivariantly uniformly K-stable by Proposition 2.12.

4. Blow-ups of the wonderful compactifications along the closed orbit

Let $Z = Y_1 \cap Y_2$ be the intersection of two *G*-stable divisors Y_1, Y_2 in the wonderful compactification X_m of the symmetric homogeneous space of type AIII(2, m), which is the unique closed *G*-orbit in X_m . In this section, we study the existence of a Kähler–Einstein metric on the blow-up $B\ell_Z(X_m)$ of X_m along the closed orbit Z.

4.1. Moment polytopes of the blow-ups of X_m along the closed orbit. In the case of m = 4, the blow-up $B\ell_Z(X_4)$ of the wonderful compactification X_4 of symmetric homogeneous space

$$\operatorname{SL}_4(\mathbb{C})/N_{\operatorname{SL}_4(\mathbb{C})}(S(\operatorname{GL}_2(\mathbb{C}) \times \operatorname{GL}_2(\mathbb{C})))$$

of type AIII(2, 4) along the unique closed orbit Z is not Fano but Calabi–Yau from [Ruz12, Proposition 2.1].

Proposition 4.1. For $m \geq 5$, the moment polytope $\Delta_m^{B\ell} = \Delta(B\ell_Z(X_m), K_{B\ell_Z(X_m)}^{-1})$ of the anticanonical line bundle $K_{B\ell_Z(X_m)}^{-1}$ is the convex hull of five points 0, $m\alpha_{1,m}$, $m\alpha_{1,m} + (m-4)\alpha_{2,m-1}$, $(m-1)\alpha_{1,m} + (m-2)\alpha_{2,m-1}$, $(m-\frac{3}{2})\alpha_{1,m} + (m-\frac{3}{2})\alpha_{2,m-1}$ in $\mathcal{M} \otimes \mathbb{R}$.



FIGURE 6. $\Delta_6^{B\ell} = \Delta(B\ell_Z(X_6), K_{B\ell_Z(X_6)}^{-1})$

Proof. Since Z is the intersection of two G-stable divisors Y_1, Y_2 in the wonderful compactification X_m , the exceptional divisor E for the blow-up $\pi : B\ell_Z(X_m) \to X_m$ is linearly equivalent to $Y_1 + Y_2$. As the G-stable divisors Y_1, Y_2 in X_m have the images $\hat{\rho}(Y_1) = -\bar{\alpha}_{1,m-1}^{\vee} = -\frac{1}{2}\alpha_{1,m}^{\vee}$ and $\hat{\rho}(Y_2) = -\bar{\alpha}_{1,m}^{\vee} = -\frac{1}{2}\alpha_{1,m}^{\vee} - \frac{1}{2}\alpha_{2,m-1}^{\vee}$ in \mathcal{N} , respectively, the exceptional divisor E for the blow-up $\pi : B\ell_Z(X_m) \to X_m$ has the image $\hat{\rho}(E) = -\bar{\alpha}_{1,m-1}^{\vee} - \bar{\alpha}_{1,m}^{\vee} \in \mathcal{N}$ (see [Ruz12, Table 7]). The same argument as in Proposition 3.7 implies the result. \Box

Example 4.2. In the case of m = 5, we compute the volume of the moment polytope $\Delta_5^{B\ell}$

$$\begin{aligned} \operatorname{Vol}_{DH}(\Delta_5^{B\ell}) &= \int_0^{\frac{7}{2}} \int_0^x 4x^3 y^3 (x+y)^2 (x-y)^2 \, dy dx + \int_{\frac{7}{2}}^4 \int_0^{7-x} 4x^3 y^3 (x+y)^2 (x-y)^2 \, dy dx \\ &+ \int_4^5 \int_0^{11-2x} 4x^3 y^3 (x+y)^2 (x-y)^2 \, dy dx = \frac{1553111579}{2520} \end{aligned}$$

and the barycenter of $\Delta_5^{B\ell}$ with respect to the Duistermaat–Heckman measure

$$\begin{aligned} \mathbf{bar}_{DH}(\Delta_5^{B\ell}) &= (\bar{x}, \bar{y}) = \frac{1}{\mathrm{Vol}_{DH}(\Delta_5^{B\ell})} \left(\int_{\Delta_5^{B\ell}} x \prod_{\alpha \in \Phi^+ \setminus \Phi^\theta} \kappa(\alpha, p) \, dp, \int_{\Delta_5^{B\ell}} y \prod_{\alpha \in \Phi^+ \setminus \Phi^\theta} \kappa(\alpha, p) \, dp \right) \\ &= \left(\frac{5341911643737}{1292188833728}, \frac{2416468747943}{1292188833728} \right) \approx (4.134, 1.870). \end{aligned}$$

Since $\bar{x} > 4$ and $\bar{x} + \bar{y} > 6$, $\mathbf{bar}_{DH}(\Delta_5^{B\ell})$ is in the relative interior of the translated cone $2\rho_{\theta} + \mathcal{C}_{\theta}^+$ (see Figure 5). Therefore, the blow-up $B\ell_Z(X_5)$ of X_5 along the closed orbit Z admits a Kähler–Einstein metric by Proposition 2.12.

Similarly, we obtain the barycenter of the moment polytope $\Delta_6^{B\ell}$ with respect to the Duistermaat– Heckman measure

$$\mathbf{bar}_{DH}(\Delta_6^{B\ell}) = (\bar{x}, \bar{y}) = \left(\frac{5817870364882097045}{1125028875118233728}, \frac{15784597990157403671}{5625144375591168640}\right) \approx (5.171, 2.806)$$

Since $\bar{x} + \bar{y} < 8$, $\mathbf{bar}_{DH}(\Delta_6^{B\ell})$ is not in the relative interior of the translated cone $2\rho_{\theta} + C_{\theta}^+$ (see Figure 6). Therefore, the blow-up $B\ell_Z(X_6)$ of X_6 along the closed orbit Z cannot admit any Kähler–Einstein metric by Proposition 2.12.

4.2. Barycenters of moment polytopes $\Delta_m^{B\ell}$. By Example 4.2, it is enough to prove Theorem 1.3 when $m \geq 7$.

Proposition 4.3. Let (\bar{x}, \bar{y}) be the barycenter $\mathbf{bar}_{DH}(\Delta_m^{B\ell})$ of the moment polytope $\Delta_m^{B\ell}$ with respect to the Duistermaat–Heckman measure. Then we have $\bar{x} + \bar{y} < 2m - 4$ for $m \ge 6$.

Proof. From Proposition 4.1, we can divide the moment polytope $\Delta_m^{B\ell}$ into three regions Ω_1 , Ω_2 and Ω_3 :

$$\begin{split} \Omega_1 &:= \{s(m,t): 0 \le s \le 1, \ 0 \le t \le m-4\},\\ \Omega_2 &:= \{s(t,3m-4-2t): 0 \le s \le 1, \ m-1 \le t \le m\},\\ \Omega_3 &:= \{s(t,2m-3-t): 0 \le s \le 1, \ m-\frac{3}{2} \le t \le m-1\}. \end{split}$$

Using the parametrization x = st, y = s(3m - 4 - 2t) of Ω_2 , we get the volume form

$$dx \wedge dy = d(st) \wedge d(s(3m - 4 - 2t)) = (s \, dt + t \, ds) \wedge [(3m - 4 - 2t)ds + s(-2dt)]$$

= $s(3m - 4 - 2t) \, dt \wedge ds - 2ts \, ds \wedge dt = s(3m - 4) \, dt \wedge ds.$

Then we compute the volume of Ω_2 with respect to the Duistermaat-Heckman measure:

$$\begin{aligned} \operatorname{Vol}_{DH}(\Omega_2) &= \int_{\Omega_2} P_{DH}(x,y) \, dx dy = \int_{\Omega_2} 2^2 x^{2m-7} y^{2m-7} (x+y)^2 (x-y)^2 \, dx dy \\ &= \int_0^1 \int_{m-1}^m 4(st)^{2m-7} \{s(3m-4-2t)\}^{2m-7} \{s(3m-4-t)\}^2 \{s(3m-4-3t)\}^2 \, s(3m-4) \, dt ds \\ &= \int_0^1 \int_{m-1}^m 4(3m-4) s^{4m-9} t^{2m-7} (3m-4-2t)^{2m-7} (3m-4-t)^2 (3m-4-3t)^2 \, dt ds \\ &= 4(3m-4) \int_0^1 s^{4m-9} \, ds \int_{m-1}^m \{t(3m-4-2t)\}^{2m-7} (3m-4-t)^2 (3m-4-3t)^2 \, dt \\ &= \frac{4(3m-4)}{4m-8} \int_{m-1}^m \{t(3m-4-2t)\}^{2m-7} (3m-4-t)^2 (3m-4-3t)^2 \, dt. \end{aligned}$$

As x + y = s(3m - 4 - t) on Ω_2 , we get

$$\bar{x}_{2} + \bar{y}_{2} = \frac{1}{\operatorname{Vol}_{DH}(\Omega_{2})} \int_{\Omega_{2}} (x+y) P_{DH}(x,y) dxdy$$

$$= \frac{1}{\operatorname{Vol}_{DH}(\Omega_{2})} \cdot 4(3m-4) \int_{0}^{1} s^{4m-8} ds \int_{m-1}^{m} \{t(3m-4-2t)\}^{2m-7} (3m-4-t)^{3} (3m-4-3t)^{2} dt$$

$$= \frac{1}{\operatorname{Vol}_{DH}(\Omega_{2})} \cdot \frac{4(3m-4)}{4m-7} \int_{m-1}^{m} \{t(3m-4-2t)\}^{2m-7} (3m-4-t)^{3} (3m-4-3t)^{2} dt.$$

Using the results in the proof of Proposition 3.9, we have

$$\begin{aligned} \operatorname{Vol}_{DH}(\Delta_m^{B\ell}) &= \frac{4m^{2m-6}}{4m-8} \int_0^{m-4} t^{2m-7} (m+t)^2 (m-t)^2 \, dt \\ &+ \frac{4(3m-4)}{4m-8} \int_{m-1}^m \{t(3m-4-2t)\}^{2m-7} (3m-4-t)^2 (3m-4-3t)^2 \, dt \\ &+ \frac{4(2m-3)^3}{4m-8} \int_{m-\frac{3}{2}}^{m-1} \{t(2m-3-t)\}^{2m-7} (2m-3-2t)^2 \, dt, \end{aligned}$$

and

$$\begin{aligned} (\bar{x}+\bar{y})\mathrm{Vol}_{DH}(\Delta_m^{B\ell}) &= \frac{4m^{2m-6}}{4m-7} \int_0^{m-4} t^{2m-7} (m+t)^3 (m-t)^2 \, dt \\ &+ \frac{4(3m-4)}{4m-7} \int_{m-1}^m \{t(3m-4-2t)\}^{2m-7} (3m-4-t)^3 (3m-4-3t)^2 \, dt \\ &+ \frac{4(2m-3)^4}{4m-7} \int_{m-\frac{3}{2}}^{m-1} \{t(2m-3-t)\}^{2m-7} (2m-3-2t)^2 \, dt. \end{aligned}$$

We can check $\bar{x}+\bar{y} < 2m-4$ for $6 \le m \le 41$ with the aid of a computer program SAGE. To prove $\bar{x}+\bar{y} < 2m-4$ for $m \ge 42$, it suffices to show the following inequality

$$\begin{split} &m^{2m-6} \int_{0}^{m-4} t^{2m-7} (m+t)^2 (m-t)^2 (2t-2m+7) \, dt \\ &+ (3m-4) \int_{m-1}^{m} \{t(3m-4-2t)\}^{2m-7} (3m-4-t)^2 (3m-4-3t)^2 (2m-2t-1) \, dt \\ &+ (2m-3)^3 \int_{m-\frac{3}{2}}^{m-1} \{t(2m-3-t)\}^{2m-7} (2m-3-2t)^2 \, dt < 0. \end{split}$$

We first derive explicit expressions for each integrals.

$$\begin{split} I_1 &= m^{2m-6} \int_0^{m-4} t^{2m-7} (m+t)^2 (m-t)^2 (2t-2m+7) \, dt \\ &= m^{2m-6} (m-4)^{2m-6} \\ &\times \frac{-5888m^7 + 82112m^6 - 491008m^5 + 1637752m^4 - 3300288m^3 + 4031616m^2 - 2774016m + 829440}{(2m-1)(2m-2)(2m-3)(2m-4)(2m-5)(2m-6)} \\ &=: m^{2m-6} (m-4)^{2m-8} \cdot (m-4)^2 r(m). \end{split}$$

Since 2t - 2m + 7 is negative for 0 < t < m - 4, we know $I_1 < 0$ so that r(m) < 0. By a well-known inequality $\left(1 - \frac{1}{n}\right)^n < e^{-1} < \left(1 - \frac{1}{n}\right)^{n-1}$ for all n > 1, we obtain $m^{2m-6}(m-4)^{2m-8} = m^{4m-14} \left(1 - \frac{4}{m}\right)^{2m-8} > m^{4m-14}(e^{-1})^{\frac{4}{m-4}(2m-8)} = m^{4m-14}e^{-8}.$

17

Hence, $I_1 < m^{4m-14}e^{-8}(m-4)^2r(m)$. For the second integral, we note that the term 2m - 2t - 1 is negative for $m - \frac{1}{2} < t < m$, which implies that we have to separate two cases for the sake of estimate.

$$\begin{split} I_2 &= (3m-4) \int_{m-1}^m \{t(3m-4-2t)\}^{2m-7} (3m-4-t)^2 (3m-4-3t)^2 (2m-2t-1) \, dt \\ &< (3m-4) \Big[\{(m-1)(m-2)\}^{2m-7} \int_{m-1}^{m-\frac{1}{2}} (3m-4-t)^2 (3m-4-3t)^2 (2m-2t-1) \, dt \\ &+ \{m(m-4)\}^{2m-7} \int_{m-\frac{1}{2}}^m (3m-4-t)^2 (3m-4-3t)^2 (2m-2t-1) \, dt \Big] \\ &= (3m-4) \Big[\{(m-1)(m-2)\}^{2m-7} \left(\frac{19}{8}m^2 - \frac{1837}{240}m + \frac{5929}{960} \right) + \{m(m-4)\}^{2m-7} \left(-\frac{99}{8}m^2 + \frac{11453}{240}m - \frac{44201}{960} \right) \Big] \end{split}$$

Using inequalities

$$\{(m-1)(m-2)\}^{2m-7} = m^{4m-14} \left(1 - \frac{1}{m}\right)^{2m-7} \left(1 - \frac{2}{m}\right)^{2m-7} < m^{4m-14} (e^{-1})^{2 - \frac{7}{m}} (e^{-1})^{4 - \frac{14}{m}} \le m^{4m-14} e^{-6 + \frac{1}{2}}$$

and $\{m(m-4)\}^{2m-7} = m^{4m-14} \left(1 - \frac{4}{m}\right)^{2m-7} > m^{4m-14} (e^{-1})^{8 + \frac{4}{m-4}} \ge m^{4m-14} e^{-8 - \frac{2}{19}}$ for $m \ge 42$, we have

$$I_2 < m^{4m-14}(3m-4) \left\{ e^{-6+\frac{1}{2}} \left(\frac{19}{8}m^2 - \frac{1837}{240}m + \frac{5929}{960} \right) + e^{-8-\frac{2}{19}} \left(-\frac{99}{8}m^2 + \frac{11453}{240}m - \frac{44201}{960} \right) \right\}.$$

For the third term, we use the same method:

$$I_{3} = (2m-3)^{3} \int_{m-\frac{3}{2}}^{m-1} \{t(2m-3-t)\}^{2m-7} (2m-3-2t)^{2} dt$$

$$< (2m-3)^{3} \left(m-\frac{3}{2}\right)^{4m-14} \int_{m-\frac{3}{2}}^{m-1} (2m-3-2t)^{2} dt = \frac{1}{6} (2m-3)^{3} \left(m-\frac{3}{2}\right)^{4m-14}$$

$$< \frac{1}{6} (2m-3)^{3} \cdot m^{4m-14} e^{\frac{21}{m}-6} \le \frac{1}{6} (2m-3)^{3} m^{4m-14} e^{-6+\frac{1}{2}}$$

for $m \ge 42$. Combining the calculations, we get the inequality

$$I_{1} + I_{2} + I_{3} < m^{4m-14}e^{-8} \Big[(m-4)^{2}r(m) + (3m-4) \left\{ e^{\frac{5}{2}} \left(\frac{19}{8}m^{2} - \frac{1837}{240}m + \frac{5929}{960} \right) + e^{-\frac{2}{19}} \left(-\frac{99}{8}m^{2} + \frac{11453}{240}m - \frac{44201}{960} \right) \right\} + \frac{e^{\frac{5}{2}}}{6} (2m-3)^{3} \Big]$$

Consequently, it suffices to show that a polynomial

 $R(m) := (m-4)^2 (-5888m^7 + 82112m^6 - 491008m^5 + 1637752m^4 - 3300288m^3 + 4031616m^2 - 2774016m + 829440) + (2m-1)(2m-2)(2m-3)(2m-4)(2m-5)(2m-6)$ $\times \left[(2m-4) \int_{-5}^{5} \left(\frac{19}{2m^2} - \frac{1837}{2m} + \frac{5929}{2} \right) + e^{-\frac{2}{27}} \left(-\frac{99}{2m^2} + \frac{11453}{2m} - \frac{44201}{2m} \right) \right] + e^{\frac{5}{2}} (2m-2)^3 \right]$

$$\times \left[(3m-4) \left\{ e^{\frac{5}{2}} \left(\frac{19}{8}m^2 - \frac{1837}{240}m + \frac{5929}{960} \right) + e^{-\frac{2}{19}} \left(-\frac{99}{8}m^2 + \frac{11453}{240}m - \frac{44201}{960} \right) \right\} + \frac{e^{\frac{5}{2}}}{6} (2m-3)^3 \right]$$

of degree 9 is always negative for $m \ge 42$. We can check that the polynomial R(m) is decreasing on $[42, \infty)$, and R(42) < 0. Therefore, we conclude the result.

Theorem 4.4. Let X_m be the wonderful compactification of symmetric homogeneous space of type AIII(2, m) for $m \ge 5$. The blow-up $B\ell_Z(X_m)$ of X_m along the closed orbit Z is $SL_m(\mathbb{C})$ -equivariantly uniformly K-stable if m = 5 and K-unstable if $m \ge 6$.

Proof. For $\mathbf{bar}_{DH}(\Delta_m^{B\ell}) = (\bar{x}, \bar{y})$, if $m \ge 6$ then $\mathbf{bar}_{DH}(\Delta_m^{B\ell})$ is not in the relative interior of the translated cone $2\rho_{\theta} + \mathcal{C}^+_{\theta}$ as $\bar{x} + \bar{y} < 2m - 4$ by Proposition 4.3. Thus, we get the result from Proposition 2.12 and Example 4.2.

4.3. Greatest Ricci lower bounds of the blow-ups $B\ell_Z(X_m)$. From Corollary 2.15, we can compute the greatest Ricci lower bound of the blow-up of X_m along the closed orbit for $m \ge 6$.

Example 4.5. We already know that the blow-up $B\ell_Z(X_6)$ of X_6 along the closed orbit Z cannot admit any Kähler–Einstein metric. Let Q be the point at which the half-line starting from the barycenter C = $\mathbf{bar}_{DH}(\Delta_6^{B\ell})$ in the direction of A = (5,3) intersects the boundary of $(\Delta_6^{B\ell})^{\mathrm{tor}} - \mathcal{C}_{\theta}^+$. Considering the line x + y = 9 giving a part of $\partial((\Delta_6^{B\ell})^{\mathrm{tor}} - \mathcal{C}_{\theta}^+)$ and the half-line \overrightarrow{CA} , from the results in Example 4.2 we can compute

$$Q = \left(-\frac{327603995647340905}{127205190161460224}, \frac{1472450707100482921}{127205190161460224}\right) \approx (-2.575, 11.575).$$

Thus, we obtain the greatest Ricci lower bound $R(B\ell_Z(X_6)) = \frac{AQ}{\overline{CQ}} \approx 0.978$ by Corollary 2.15.

References

- [AZ22] Hamid Abban and Ziquan Zhuang, K-stability of Fano varieties via admissible flags, Forum Math. Pi 10 (2022), Art. No. e15.
- [ACCFKMGSSV] Carolina Araujo, Ana-Maria Castravet, Ivan Cheltsov, Kento Fujita, Anne-Sophie Kaloghiros, Jesus Martinez-Garcia, Constantin Shramov, Hendrik Süß, and Nivedita Viswanathan, The Calabi problem for Fano threefolds, London Math. Soc. Lect. Note Ser., vol. 485, Cambridge Univ. Press, Cambridge, 2023.
- [Aub78] Thierry Aubin, Équations du type Monge-Ampère sur les variétés kählériennes compactes, Bull. Sci. Math.(2) 102 (1978), 63–95.
- [BBJ21] Robert Berman, Sébastien Boucksom, and Mattias Jonsson, A variational approach to the Yau-Tian-Donaldson conjecture, J. Amer. Math. Soc. **34** (2021), no. 3, 605–652.
- [BP16] Paolo Bravi and Guido Pezzini, Primitive wonderful varieties, Math. Z. 282 (2016), no. 3-4, 1067–1096.
- [Bri87] Michel Brion, Sur l'image de l'application moment, Séminaire d'algèbre Paul Dubreil et Marie-Paule Malliavin (Paris, 1986), Lecture Notes in Math., vol. 1296, Springer, Berlin, 1987, pp. 177–192.
- [Bri89] _____, Groupe de Picard et nombres caractéristiques des variétés sphériques, Duke Math. J. 58 (1989), no. 2, 397–424.
- [Bri97] _____, Curves and divisors in spherical varieties, Algebraic groups and Lie groups, Austral. Math. Soc. Lect. Ser., vol. 9, Cambridge Univ. Press, Cambridge, 1997, pp. 21–34.
- [CRZ19] Ivan Cheltsov, Yanir Rubinstein, and Kewei Zhang, Basis log canonical thresholds, local intersection estimates, and asymptotically log del Pezzo surfaces, Selecta Math. 25 (2019), Article number 34.
- [CDS15a] Xiuxiong Chen, Simon Donaldson, and Song Sun, Kähler-Einstein metrics on Fano manifolds. I: Approximation of metrics with cone singularities, J. Amer. Math. Soc. 28 (2015), no. 1, 183–197.
- [CDS15b] _____, Kähler-Einstein metrics on Fano manifolds. II: Limits with cone angle less than 2π, J. Amer. Math. Soc. 28 (2015), no. 1, 199–234.
- [CDS15c] _____, Kähler–Einstein metrics on Fano manifolds. III: Limits as cone angle approaches 2π and completion of the main proof, J. Amer. Math. Soc. 28 (2015), no. 1, 235–278.
- [DCP83] Corrado De Concini and Claudio Procesi, Complete symmetric varieties, Invariant theory (Montecatini, 1982), Lecture Notes in Math., Vol. 996, Springer, Berlin, 1983, pp. 1–44.
- [Del17] Thibaut Delcroix, Kähler–Einstein metrics on group compactifications, Geom. Funct. Anal. 27 (2017), no. 1, 78–129.
- [Del20a] _____, K-stability of Fano spherical varieties, Ann. Sci. Éc. Norm. Supér. (4) 53 (2020), no. 3, 615–662.
- [Del20b] _____, Kähler geometry of horosymmetric varieties, and application to Mabuchi's K-energy functional, J. Reine Angew. Math. **763** (2020), 129–199.
- [Del22] _____, Examples of K-unstable Fano manifolds, Ann. Inst. Fourier **72** (2022), no. 5, 2079–2108.
- [DH21] Thibaut Delcroix and Jakob Hultgren, Coupled complex Monge-Ampère equations on Fano horosymmetric manifolds, J. Math. Pures Appl. 153 (2021), 281–315.
- [Don02] Simon K. Donaldson, Scalar curvature and stability of toric varieties, J. Differential Geom. 62 (2002), no. 2, 289–349.
- [FO18] Kento Fujita and Yuji Odaka, On the K-stability of Fano varieties and anticanonical divisors, Tohoku Math. J. 70 (2018), no. 4, 511–521.
- [Gan18] Jacopo Gandini, Embeddings of spherical homogeneous spaces, Acta Math. Sin. (Engl. Ser.) 34 (2018), no. 3, 299–340.
 [GH15] Giuliano Gagliardi and Johannes Hofscheier, Gorenstein spherical Fano varieties, Geom. Dedicata 178 (2015), 111–133.
- [Gol20] Aleksei Golota, Delta-invariants for Fano varieties with large automorphism groups, Internat. J. Math. 31 (2020), no. 10, 2050077.
- [Kno91] Friedrich Knop, The Luna-Vust theory of spherical embeddings, Proceedings of the Hyderabad Conference on Algebraic Groups (Hyderabad, 1989), Manoj Prakashan, Madras, 1991, pp. 225–249.
- [LPY21] Jae-Hyouk Lee, Kyeong-Dong Park, and Sungmin Yoo, Kähler-Einstein metrics on smooth Fano symmetric varieties with Picard number one, Mathematics 9 (2021), no. 1, 102.

[Li11] Chi Li, Greatest lower bounds on Ricci curvature for toric Fano manifolds, Adv. Math. 226 (2011), no. 6, 4921–4932.

[LL22] Yan Li and Zhenye Li, Finiteness of Q-Fano compactifications of semisimple groups with Kähler-Einstein metrics, Int. Math. Res. Not. IMRN 2022, no. 15, 11776–11795.

- [Lun96] Domingo Luna, Toute variété magnifique est sphérique, Transform. Groups, 1 (1996), no. 3, 249–258.
- [Lun97] _____, Grosses cellules pour les variétés sphériques, Algebraic groups and Lie groups, Austral. Math. Soc. Lect. Ser., vol. 9, Cambridge Univ. Press, Cambridge, 1997, pp. 267–280.
- [Lun01] _____, Variétés sphériques de type A, Inst. Hautes 'Etudes Sci. Publ. Math., 94 (2001), 161–226.
- [LV83] Domingo Luna and Thierry Vust, Plongements d'espaces homogènes, Comment. Math. Helv. 58 (1983), no. 2, 186– 245.
- [Mab87] Toshiki Mabuchi, Einstein-Kähler forms, Futaki invariants and convex geometry on toric Fano varieties, Osaka J. Math. 24 (1987), no. 4, 705–737.
- [Mat72] Yozô Matsushima, Remarks on Kähler-Einstein manifolds, Nagoya Math. J. 46 (1972), 161–173.
- [Pas09] Boris Pasquier, On some smooth projective two-orbit varieties of Picard number 1, Math. Ann. 344 (2009), no. 4, 963–987.
- [Pez18] Guido Pezzini, Lectures on wonderful varieties, Acta Math. Sin. (Engl. Ser.) 34 (2018), no. 3, 417-438.
- [Rub08] Yanir A. Rubinstein, Some discretizations of geometric evolution equations and the Ricci iteration on the space of Kähler metrics, Adv. Math. 218 (2008), no. 5, 1526–1565.
- [Rub09] _____, On the construction of Nadel multiplier ideal sheaves and the limiting behavior of the Ricci flow, Trans. Amer. Math. Soc. **361** (2009), no. 11, 5839–5850.
- [Ruz12] Alessandro Ruzzi, Fano symmetric varieties with low rank, Publ. Res. Inst. Math. Sci. 48 (2012), no. 2, 235–278.
- [SW16] Jian Song and Xiaowei Wang, The greatest Ricci lower bound, conical Einstein metrics and Chern number inequality, Geom. Topol. 20 (2016), no. 1, 49–102.
- [Ste68] Robert Steinberg, Endomorphisms of linear algebraic groups, Memoirs of the American Mathematical Society, No. 80, American Mathematical Society, Providence, RI, 1968.
- [Szé11] Gábor Székelyhidi, Greatest lower bounds on the Ricci curvature of Fano manifolds, Compos. Math. 147, no. 1, 319–331.
- [Tia90] Gang Tian, On Calabi's conjecture for complex surfaces with positive first Chern class, Invent. Math. 101 (1990), 101–172.
- [Tia92] _____, On stability of the tangent bundles of Fano varieties, Internat. J. Math. 3 (1992), no. 3, 401–413.
- [Tia94] _____, The K-energy on hypersurfaces and stability, Comm. Anal. Geom. 2 (1994), no. 2, 239–265.
- [Tia97] _____, Kähler-Einstein metrics with positive scalar curvature, Invent. Math. 130 (1997), no. 1, 1–37.
- [Tia15] _____, K-stability and Kähler-Einstein metrics, Commun. Pure Appl. Math. 68 (2015), no. 7, 1085–1156.
- [TY87] Gang Tian and Shing-Tung Yau, Kähler—Einstein metrics on complex surfaces with $C_1 > 0$, Commun. Math. Phys. **112**, (1987), 175–203.
- [Tim11] Dmitry A. Timashev, Homogeneous spaces and equivariant embeddings, Encyclopaedia of Mathematical Sciences, vol. 138, Springer, Heidelberg, 2011, Invariant Theory and Algebraic Transformation Groups, 8.
- [Vus74] Thierry Vust, Opération de groupes réductifs dans un type de cônes presque homogènes, Bull. Soc. Math. France 102 (1974), 317–333.
- [Vus90] _____, Plongements d'espaces symétriques algébriques: une classification, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) 17 (1990), no. 2, 165–195.
- [WZ04] Xu-Jia Wang and Xiaohua Zhu, Kähler-Ricci solitons on toric manifolds with positive first Chern class, Adv. Math. 188 (2004), no. 1, 87–103.
- [Yau78] Shing-Tung Yau, On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation, I, Comm. Pure Appl. Math. 31 (1978), no. 3, 339–411.
- [Zhu21] Ziquan Zhuang, Optimal destabilizing centers and equivariant K-stability, Invent. Math. 226 (2021), no. 1, 195–223.

DEPARTMENT OF MATHEMATICS EDUCATION, JEONJU UNIVERSITY, JEONJU 55069, REPUBLIC OF KOREA *Email address*: kszooj@jj.ac.kr

CENTER FOR COMPLEX GEOMETRY, INSTITUTE FOR BASIC SCIENCE (IBS), DAEJEON 34126, REPUBLIC OF KOREA *Email address:* dshwang@ibs.re.kr

DEPARTMENT OF MATHEMATICS AND RESEARCH INSTITUTE OF NATURAL SCIENCE/RESEARCH INSTITUTE OF MOLECULAR Alchemy, Gyeongsang National University, Jinju 52828, Republic of Korea

Email address: kdpark@gnu.ac.kr

20