

Robust Unsupervised Multi-task and Transfer Learning on Gaussian Mixture Models

Ye Tian¹, Haolei Weng², and Yang Feng³

¹Department of Statistics, Columbia University

²Department of Statistics and Probability, Michigan State University

³Department of Biostatistics, School of Global Public Health, New York University

Abstract

Unsupervised learning has been widely used in many real-world applications. One of the simplest and most important unsupervised learning models is the Gaussian mixture model (GMM). In this work, we study the multi-task learning problem on GMMs, which aims to leverage potentially similar GMM parameter structures among tasks to obtain improved learning performance compared to single-task learning. We propose a multi-task GMM learning procedure based on the EM algorithm that not only can effectively utilize unknown similarity between related tasks but is also robust against a fraction of outlier tasks from arbitrary distributions. The proposed procedure is shown to achieve minimax optimal rate of convergence for both parameter estimation error and the excess mis-clustering error, in a wide range of regimes. Moreover, we generalize our approach to tackle the problem of transfer learning for GMMs, where similar theoretical results are derived. Finally, we demonstrate the effectiveness of our methods through simulations and real data examples. To the best of our knowledge, this is the first work studying multi-task and transfer learning on GMMs with theoretical guarantees.

Keywords: Multi-task learning, transfer learning, unsupervised learning, Gaussian mixture models, robustness, minimax rate, EM algorithm

1 Introduction

1.1 Gaussian mixture models (GMMs)

Unsupervised learning that learns patterns from unlabeled data is a prevalent problem in statistics and machine learning. Clustering is one of the most important problems in unsupervised learning, where the goal is to group the observations based on some metrics of similarity. Researchers have developed numerous clustering methods including k -means (Forgy, 1965), k -medians (Jain and Dubes, 1988), spectral clustering (Ng et al., 2001), and hierarchical clustering (Murtagh and Contreras, 2012), among others. On the other hand, clustering problems have been analyzed from the perspective of the mixture of several probability distributions (Scott and Symons, 1971). The mixture of Gaussian distributions is one of the simplest models in this category and has been widely applied in many real applications (Yang and Ahuja, 1998; Lee et al., 2012).

In the binary Gaussian mixture models (GMMs) with common covariances, each observation $Z \in \mathbb{R}^p$ comes from the following mixture of two Gaussian distributions:

$$Y = \begin{cases} 1, & \text{with probability } 1 - w, \\ 2, & \text{with probability } w, \end{cases}$$
$$Z|Y = r \sim \mathcal{N}(\boldsymbol{\mu}_r, \boldsymbol{\Sigma}), \quad r = 1, 2,$$

where $w \in (0, 1)$, $\boldsymbol{\mu}_1 \in \mathbb{R}^p$, $\boldsymbol{\mu}_2 \in \mathbb{R}^p$ and $\boldsymbol{\Sigma} \succ 0$ are parameters. This is the same setting as the linear discriminant analysis (LDA) problem in classification (Hastie et al., 2009), except that the label Y is unknown in the clustering problem, while it is observed in the classification case. It has been shown that the Bayes classifier for the LDA problem is

$$\mathcal{C}(\mathbf{z}) = \begin{cases} 1, & \text{if } \boldsymbol{\beta}^T \mathbf{z} - \delta \leq \log\left(\frac{1-w}{w}\right); \\ 2, & \text{otherwise,} \end{cases} \quad (1)$$

where $\boldsymbol{\beta} = \boldsymbol{\Sigma}^{-1}(\boldsymbol{\mu}_2 - \boldsymbol{\mu}_1) \in \mathbb{R}^p$ and $\delta = \boldsymbol{\beta}^T(\boldsymbol{\mu}_1 + \boldsymbol{\mu}_2)/2$. Note that $\boldsymbol{\beta}$ is usually referred to as the discriminant coefficient (Anderson, 1958; Efron, 1975). Naturally, this classifier is useful in clustering too. In clustering, after learning w , $\boldsymbol{\mu}_1$, $\boldsymbol{\mu}_2$ and $\boldsymbol{\beta}$, we can plug their estimators into (1) to group any new observation Z^{new} . Generally, we define the mis-clustering error rate of any given clustering method \mathcal{C} as

$$R(\mathcal{C}) = \min_{\pi: \{1,2\} \rightarrow \{1,2\}} \mathbb{P}(\mathcal{C}(Z^{\text{new}}) \neq \pi(Y^{\text{new}})),$$

where π is a permutation function, Y^{new} is the label of a future observation Z^{new} , and the probability is taken w.r.t. the joint distribution of $(Z^{\text{new}}, Y^{\text{new}})$ based on parameters w , $\boldsymbol{\mu}_1$, $\boldsymbol{\mu}_2$ and $\boldsymbol{\Sigma}$. Here the error is calculated up to a permutation due to the lack of label information. It is clear that in the ideal case where the parameters are known, $\mathcal{C}(\cdot)$ in (1) achieves the optimal mis-clustering error. Multi-cluster Gaussian mixture models with $R \geq 3$ components can be described in a similar way. We provide the details in Section 3.

There is a large volume of published studies on learning a GMM. The vast majority of approaches can be roughly divided into three categories. The first category is the method of moments, where the parameters are estimated through several moment equations (Pearson, 1894; Kalai et al., 2010; Hsu and Kakade, 2013; Ge et al., 2015). The second category is the spectral method, where the estimation is based on the spectral decomposition (Vempala and Wang, 2004; Hsu and Kakade, 2013; Jin et al., 2017). The last category is the likelihood-based method including the popular expectation-maximization (EM) algorithm as a canonical example. The general form of EM algorithm was formalized by Dempster et al. (1977) in the context of incomplete data, though earlier works (Hartley, 1958; Hasselblad, 1966; Baum et al., 1970; Sundberg, 1974) have studied EM-style algorithms in various concrete settings. Classical convergence results on the EM algorithm (Wu, 1983; Redner and Walker, 1984; Meng and Rubin, 1994; McLachlan and Krishnan, 2007) guarantee local convergence of the algorithm to fixed points of the sample likelihood. Recent advances in the analysis of EM algorithm and its variants provide stronger guarantees by establishing geometric convergence rates of the algorithm to the underlying true parameters under mild initialization conditions. See, for example, Dasgupta and Schulman (2013); Wang et al. (2014); Xu et al. (2016); Balakrishnan et al. (2017); Yan et al. (2017); Cai et al. (2019); Kwon and Caramanis (2020); Zhao et al. (2020) for GMM related works. In this paper, we propose modified versions of the EM algorithm with similarly strong guarantees to learn GMMs, under the new context of multi-task and transfer learning.

1.2 Multi-task learning and transfer learning

Multi-tasking is an ability that helps people pay attention to more than one task simultaneously. Moreover, we often find that the knowledge learned from one task can also be useful in other tasks. Multi-task learning (MTL) is a learning paradigm inspired by the human learning ability, which aims to learn multiple tasks and improve performance by utilizing the similarity between these tasks (Zhang and Yang, 2021). There have been numerous research on MTL, which can be classified into five categories (Zhang and Yang, 2021): feature learning approach (Argyriou et al., 2008; Obozinski et al., 2006), low-rank approach (Ando et al., 2005), task clustering approach (Thrun and O’Sullivan, 1996), task relation learning approach (Evgeniou and Pontil, 2004) and decomposition approach (Jalali et al., 2010). The majority of existing works focus on the use of MTL in supervised learning problems, while the application of MTL in unsupervised learning, such as clustering, has received less attention. Zhang and Zhang (2011) developed an MTL clustering method based on a penalization framework, where the objective function consists of a local loss function and a pairwise task regularization term, both of which are related to the Bregman divergence. In Gu et al. (2011), a reproducing kernel Hilbert space (RKHS) was first established, then a multi-task kernel k-means clustering was applied based on that RKHS. Yang et al. (2014) proposed a spectral MTL clustering method with a novel $\ell_{2,p}$ -norm, which can also produce a linear regression function to predict labels for out-of-sample data. Zhang et al. (2018) suggested a new method based on the similarity matrix of samples in each task, which can learn the within-task clustering structure as well as the task relatedness simultaneously.

Marfoq et al. (2021) established a new federated multi-task EM algorithm to learn the mixture of distributions and provided some theory on the convergence guarantee, but the statistical properties of the estimators were not fully understood. In general, there are very few theoretical results about unsupervised MTL.

Transfer learning (TL) is another learning paradigm similar to multi-task learning but has different objectives. While MTL aims to learn all the tasks well with no priority for any specific task, the goal of TL is to improve the performance on the *target* task using the information from the *source* tasks (Zhang and Yang, 2021). According to Pan and Yang (2009), most of TL approaches can be classified into four categories: instance-based transfer (Dai et al., 2007), feature representation transfer (Dai et al., 2008), parameter transfer (Lawrence and Platt, 2004) and relational-knowledge transfer (Mihalkova et al., 2007). Similar to MTL, most of the TL methods focus on supervised learning. Some TL approaches are also developed for the semi-supervised learning setting (Chattopadhyay et al., 2012; Li et al., 2013), where only part of target or source data is labeled. There are much less discussions on the unsupervised TL approaches¹, which focus on the cases where both target and source data are unlabeled. Dai et al. (2008) developed a co-clustering approach to transfer information from a single source to the target, which relies on the loss in mutual information and requires the features to be discrete. Wang et al. (2008) proposed a TL discriminant analysis method, where the target data is allowed to be unlabeled, but some labeled source data is necessary. In Wang et al. (2021), a TL approach was developed to learn Gaussian mixture models with only one source by weighting the target and source likelihood functions. Zuo et al. (2018) proposed a TL method based on infinite Gaussian mixture models and active learning, but their approach needs sufficient labeled source data and a few labeled target samples.

There are some recent studies on TL and MTL under various statistical settings, including high-dimensional linear regression (Xu and Bastani, 2021; Li et al., 2022b; Zhang et al., 2022; Li et al., 2022a), high-dimensional generalized linear models (Li et al., 2021a; Bastani, 2021; Tian and Feng, 2022), functional linear regression (Lin and Reimherr, 2022), high-dimensional graphical models (Li et al., 2022b), reinforcement learning (Chen et al., 2022), among others. The recent work Duan and Wang (2022) developed an adaptive and robust MTL framework with sharp statistical guarantees for a broad class of models. We discuss its connection to our work in Section 2.

1.3 Our contributions and paper structure

Our main contributions in this work can be summarized in the following:

- (i) We develop efficient polynomial-time iterative procedures to learn GMMs in both MTL and TL settings. These procedures can be viewed as adaptations of the standard EM algorithm for MTL and TL problems.

¹There are different definitions for unsupervised TL. Sometimes people call the semi-supervised TL an unsupervised TL as well. We follow the definition in Pan and Yang (2009) here.

- (ii) The developed procedures come with provable statistical guarantees. Specifically, we derive the upper bounds of their estimation and excess mis-clustering error rates under mild conditions. For MTL, it is shown that when the tasks are close to each other, our method can achieve better upper bounds than those from the single-task learning; when the tasks are substantially different from each other, our method can still obtain competitive convergence rates compared to single-task learning. Similarly for TL, our method can achieve better upper bounds than those from fitting GMM only to target data when the target and sources are similar, and remains competitive otherwise. In addition, the derived upper bounds reveal the robustness of our methods against a fraction of outlier tasks (for MTL) or outlier sources (for TL) from arbitrary distributions. These guarantees certify our procedures as adaptive (to the unknown task relatedness) and robust (to contaminated data) learning approaches.
- (iii) We derive the minimax lower bounds for parameter estimation and excess mis-clustering errors. In various regimes, the upper bounds from our methods match the lower bounds (up to small order terms), showing that the proposed methods are (nearly) minimax rate optimal.
- (iv) Our MTL and TL approaches require the initial estimates for different tasks to be “well-aligned”, due to the non-identifiability of GMM. We propose two pre-processing alignment algorithms to provably resolve the alignment problem. Similar problems arise in many unsupervised MTL settings. However, to our knowledge, there is no formal discussion for the alignment issue in existing literatures of unsupervised MTL (Gu et al., 2011; Zhang and Zhang, 2011; Yang et al., 2014; Zhang et al., 2018; Dieuleveut et al., 2021; Marfoq et al., 2021). Therefore, our rigorous treatment on the alignment problem is an important step forward in this field.

The rest of the paper is organized as follows. In Section 2, we first discuss the multi-task learning problem for binary GMMs, by introducing the problem setting, our method, and the associated theory. The above-mentioned alignment problem is discussed in Section 2.4. We then extend our methods and theory to multi-cluster GMMs in Section 3. We present the results of numerical experiments, including simulations and real-data examples, in Section 4. Finally, in Section 5, we point out some interesting future research directions. Due to the space limit, additional numerical results on multi-task learning, a full treatment of the transfer learning problem, and all the technical proofs are delegated to Appendix.

We summarize the notations used throughout the paper here for convenience. We use bold capital letters (e.g. $\mathbf{\Sigma}$) to denote matrices, and use bold small letters (e.g. \mathbf{x} , \mathbf{y}) to denote vectors. For a matrix $\mathbf{A} = [a_{ij}]_{p \times q} \in \mathbb{R}^{p \times q}$, its 2-norm or spectral norm is defined as $\|\mathbf{A}\|_2 = \max_{\mathbf{x}: \|\mathbf{x}\|_2=1} \|\mathbf{A}\mathbf{x}\|_2$. If $q = 1$, \mathbf{A} becomes a p -dimensional vector and $\|\mathbf{A}\|_2$ equals its Euclidean norm. For symmetric \mathbf{A} , we define $\lambda_{\max}(\mathbf{A})$ and $\lambda_{\min}(\mathbf{A})$ as the maximum and minimum eigenvalues of \mathbf{A} , respectively. For two non-zero real sequences $\{a_n\}_{n=1}^{\infty}$ and $\{b_n\}_{n=1}^{\infty}$, we use $a_n \ll b_n$, $b_n \gg a_n$ or $a_n = o(b_n)$ to represent $|a_n/b_n| \rightarrow 0$ as $n \rightarrow \infty$. And $a_n \lesssim b_n$, $b_n \gtrsim a_n$ or $a_n = \mathcal{O}(b_n)$ means $\sup_n |a_n/b_n| < \infty$. For two random variable sequences $\{x_n\}_{n=1}^{\infty}$ and $\{y_n\}_{n=1}^{\infty}$, notation $x_n \lesssim_p y_n$ or $x_n = \mathcal{O}_p(y_n)$ means that for any

$\epsilon > 0$, there exists a positive constant M such that $\sup_n \mathbb{P}(|x_n/y_n| > M) \leq \epsilon$. For two real numbers a and b , $a \vee b$ and $a \wedge b$ represent $\max(a, b)$ and $\min(a, b)$, respectively. For any positive integer K , both $1 : K$ and $[K]$ stand for the set $\{1, 2, \dots, K\}$. For any set $S \subseteq [K]$, $|S|$ denotes its cardinality and S^c denotes its complement. Without further notice, c, C, C_1, C_2, \dots represent some positive constants and can change from line to line.

2 Multi-task Learning

2.1 Problem setting

Suppose there are K tasks, for which we have n_k observations $\{\mathbf{z}_i^{(k)}\}_{i=1}^{n_k}$ from the k -th task. Suppose there exists an *unknown* subset $S \subseteq 1 : K$, such that observations from each task in S independently follow a GMM, while samples from tasks outside S can be arbitrarily distributed. This means,

$$y_i^{(k)} = \begin{cases} 1, & \text{with probability } 1 - w^{(k)*}; \\ 2, & \text{with probability } w^{(k)*}; \end{cases} \quad \mathbf{z}_i^{(k)} | y_i^{(k)} = r \sim \mathcal{N}(\boldsymbol{\mu}_r^{(k)*}, \boldsymbol{\Sigma}^{(k)*}), \quad r = 1, 2,$$

for all $k \in S$, $i = 1 : n_k$, and

$$\{\mathbf{z}_i^{(k)}\}_{i,k \in S^c} \sim \mathbb{Q}_S,$$

where \mathbb{Q}_S is some probability measure on $(\mathbb{R}^p)^{\otimes n_{S^c}}$ and $n_{S^c} = \sum_{k \in S^c} n_k$. In unsupervised learning, we have no access to the true labels $\{y_i^{(k)}\}_{i,k}$. To formalize the multi-task learning problem, we first introduce the parameter space for a single GMM:

$$\begin{aligned} \bar{\Theta} = \{\bar{\boldsymbol{\theta}} = (w, \boldsymbol{\mu}_1, \boldsymbol{\mu}_2, \boldsymbol{\Sigma}) : \|\boldsymbol{\mu}_1\|_2 \vee \|\boldsymbol{\mu}_2\|_2 \leq M, w \in (c_w, 1 - c_w), \\ c_{\boldsymbol{\Sigma}}^{-1} \leq \lambda_{\min}(\boldsymbol{\Sigma}) \leq \lambda_{\max}(\boldsymbol{\Sigma}) \leq c_{\boldsymbol{\Sigma}}\}, \end{aligned} \quad (2)$$

where $M, c_w \in (0, 1/2]$ and $c_{\boldsymbol{\Sigma}}$ are some fixed positive constants. For simplicity, throughout the main text we assume these constants are fixed, hence we have suppressed the dependency on them in the notation $\bar{\Theta}$. The parameter space $\bar{\Theta}$ is a standard formulation. Similar parameter spaces have been considered, for example, in [Cai et al. \(2019\)](#).

Our goal of multi-task learning is to leverage the potential similarity shared by different tasks in S to collectively learn them all. The tasks outside S can be arbitrarily distributed and they can be potentially outlier tasks. This motivates us to define a joint parameter space for GMMs in S

$$\bar{\Theta}_S(h) = \left\{ \{\bar{\boldsymbol{\theta}}^{(k)}\}_{k \in S} = \{(w^{(k)}, \boldsymbol{\mu}_1^{(k)}, \boldsymbol{\mu}_2^{(k)}, \boldsymbol{\Sigma}^{(k)})\}_{k \in S} : \bar{\boldsymbol{\theta}}^{(k)} \in \bar{\Theta}, \inf_{\bar{\boldsymbol{\beta}}} \max_{k \in S} \|\boldsymbol{\beta}^{(k)} - \bar{\boldsymbol{\beta}}\|_2 \leq h \right\}, \quad (3)$$

where $\boldsymbol{\beta}^{(k)} = (\boldsymbol{\Sigma}^{(k)})^{-1}(\boldsymbol{\mu}_2^{(k)} - \boldsymbol{\mu}_1^{(k)})$ is called the discriminant coefficient in the k -th task (recall Section 1.1). For convenience, we define $\delta^{(k)} = (\boldsymbol{\beta}^{(k)})^T(\boldsymbol{\mu}_1^{(k)} + \boldsymbol{\mu}_2^{(k)})/2$, which together

with $\log((1 - w^{(k)})/w^{(k)})$ is part of the decision boundary. Note that this parameter space is defined only for GMMs of tasks in S . To model potentially corrupted or contaminated data, we do not impose any distributional constraints for tasks in S^c . Such a modeling framework is reminiscent of Huber’s ϵ -contamination model (Huber, 1964). Similar formulations have been adopted in recent multi-task learning research such as Konstantinov et al. (2020) and Duan and Wang (2022).

For GMMs in $\bar{\Theta}_S(h)$, we assume that they share similar discriminant coefficients. The similarity is formalized by assuming that all the discriminant coefficients in S are within Euclidean distance h from a “center”. Given that the discriminant coefficient has a major impact on the clustering performance (see the discriminant rule in (1)), the parameter space $\bar{\Theta}_S(h)$ is tailored to characterize the task relatedness from the clustering perspective. A similar viewpoint that focuses on modeling the discriminant coefficient has appeared in the study of high-dimensional GMM clustering (Cai et al., 2019) and sparse linear discriminant analysis (Cai and Liu, 2011; Mai et al., 2012). With both S and h being unknown in practice (similar to that in Konstantinov et al. (2020) and Duan and Wang (2022)), we aim to develop a multi-task learning procedure that is robust to outlier tasks in S^c , and achieves improved performance for tasks in S (compared to the single-task learning), in terms of discriminant coefficient estimation and clustering, whenever h is small.

The parameter space does not require the mean vectors $\{\boldsymbol{\mu}_1^{(k)}, \boldsymbol{\mu}_2^{(k)}\}_{k \in S}$ or the covariance matrices $\{\boldsymbol{\Sigma}^{(k)}\}_{k \in S}$ to be similar, although they are not free parameters due to the constraint on $\{\boldsymbol{\beta}^{(k)}\}_{k \in S}$. And the mixture proportions $\{w^{(k)}\}_{k \in S}$ do not need to be similar either. We thus avoid imposing restrictive conditions on those parameters. On the other hand, it implies that estimation of the mixture proportions, mean vectors, and covariance matrices in multi-task learning may not be generally improvable over that in the single-task learning. This is verified by the theoretical results in Section 2.3. While the current treatment in the paper does not consider similarity structure among $\{\boldsymbol{\mu}_1^{(k)}, \boldsymbol{\mu}_2^{(k)}\}_{k \in S}$, $\{\boldsymbol{\Sigma}^{(k)}\}_{k \in S}$ or $\{w^{(k)}\}_{k \in S}$, our methods and theory can be readily adapted to handle such scenarios, if desired.

There are two main reasons why this MTL problem can be challenging. First, commonly used strategies like data pooling are fragile with respect to outlier tasks and can lead to arbitrarily inaccurate outcomes in the presence of even a small number of outliers. Also, since the distribution of data from outlier tasks can be adversarial to the learner, the idea of outlier task detection in the recent literature (Li et al., 2021b; Tian and Feng, 2022) may not be applicable. Second, to address the nonconvexity of the likelihood, we propose to explore the similarity among tasks via a generalization of the EM algorithm. However, a clear theoretical understanding of such an iterative procedure requires a delicate analysis of the whole iterative process. In particular, as in the analysis of EM algorithms (Cai et al., 2019; Kwon and Caramanis, 2020), the estimates of similar discriminant vectors $\{\boldsymbol{\beta}^{(k)*}\}_{k \in S}$ and other potentially dissimilar parameters are entangled in the iterations. It is highly non-trivial to separate the impact of estimating $\{\boldsymbol{\beta}^{(k)*}\}_{k \in S}$ and other parameters to derive the desired statistical error rates. We manage to address this challenge through a localization technique by carefully shrinking the analysis radius of estimators as the iteration proceeds.

2.2 Method

We aim to tackle the problem of GMM estimation under the context of multi-task learning. The EM algorithm is commonly used to address the non-convexity of the log-likelihood function arising from the latent labels. In the standard EM algorithm, we “classify” the observations (update the posterior) in E-step and update the parameter estimations in M-step (Redner and Walker, 1984). For multi-task and transfer learning problems, the penalization framework is very popular, where we solve an optimization problem based on a new objective function. This objective function consists of a local loss function and a penalty term, forcing the estimators of similar tasks to be close to each other. See Zhang and Zhang (2011); Zhang et al. (2015); Bastani (2021); Xu and Bastani (2021); Li et al. (2021b,a); Tian and Feng (2022); Duan and Wang (2022); Lin and Reimherr (2022). Thusly motivated, our method seeks a combination of the EM algorithm and the penalization framework.

In particular, we adapt the penalization framework of Duan and Wang (2022) and modify the updating formulas in M-step accordingly. The proposed procedure is summarized in Algorithm 1. For simplicity, in Algorithm 1 we have used the notation

$$\gamma_{\theta}(\mathbf{z}) = \frac{w \exp(\boldsymbol{\beta}^T \mathbf{z} - \delta)}{1 - w + w \exp(\boldsymbol{\beta}^T \mathbf{z} - \delta)}, \quad \text{for } \boldsymbol{\theta} = (w, \boldsymbol{\beta}, \delta).$$

Note that $\gamma_{\theta}(\mathbf{z})$ is the posterior probability $\mathbb{P}(Y = 2|Z = \mathbf{z})$ given the observation \mathbf{z} . The estimated posterior probability is calculated in every E-step given the updated parameter estimates.

Recall that the parameter space $\bar{\Theta}_S(h)$ introduced in (3) does not encode similarity for the mixture proportions $\{w^{(k)*}\}_{k \in S}$, mean vectors $\{\boldsymbol{\mu}_1^{(k)*}, \boldsymbol{\mu}_2^{(k)*}\}_{k \in S}$, or covariance matrices $\{\boldsymbol{\Sigma}^{(k)*}\}_{k \in S}$. Hence, the updates of them in Steps 5-7 are kept the same as in the standard EM algorithm. Regarding the update for discriminant coefficients in Step 9, the quadratic loss function is motivated by the direct estimation of the discriminant coefficient in high-dimensional GMM (Cai et al., 2019) and high-dimensional LDA literature (Cai and Liu, 2011; Witten and Tibshirani, 2011; Fan et al., 2012; Mai et al., 2012, 2019). The penalty term in Step 9 penalizes the contrasts of $\boldsymbol{\beta}^{(k)}$ ’s to exploit the similarity structure among tasks. Having the “center” parameter $\bar{\boldsymbol{\beta}}$ in the penalization induces robustness against outlier tasks. We refer to Duan and Wang (2022) for a systematic treatment of this penalization framework. It is straightforward to verify that when the tuning parameters $\{\lambda^{[t]}\}_{t=1}^T$ are set to zero, Algorithm 1 reduces to the standard EM algorithm performed separately on the K tasks. That is, for each $k = 1 : K$, given the parameter estimate from the previous step $\hat{\boldsymbol{\theta}}^{(k)[t-1]} = (\hat{w}^{(k)[t-1]}, \hat{\boldsymbol{\beta}}^{(k)[t-1]}, \hat{\delta}^{(k)[t-1]})$, we update $\hat{w}^{(k)[t]}$, $\hat{\boldsymbol{\mu}}_1^{(k)[t]}$, $\hat{\boldsymbol{\mu}}_2^{(k)[t]}$, $\hat{\boldsymbol{\Sigma}}^{(k)[t]}$, and $\hat{\delta}^{(k)[t]}$ as in Algorithm 1, and update $\hat{\boldsymbol{\beta}}^{(k)[t]}$ via

$$\hat{\boldsymbol{\beta}}^{(k)[t]} = (\hat{\boldsymbol{\Sigma}}^{(k)[t]})^{-1}(\hat{\boldsymbol{\mu}}_2^{(k)[t]} - \hat{\boldsymbol{\mu}}_1^{(k)[t]}).$$

For the maximum iteration round number T , our theory will show that $T \gtrsim \log(\max_{k=1:K} n_k)$ is sufficient to reach the desired statistical error rates. In practice, we can terminate the it-

- (i) $n_S = \sum_{k \in S} n_k \geq C_1 |S| \max_{k=1:K} n_k$ with a constant $C_1 > 0$;
- (ii) $\min_{k \in S} n_k \geq C_2(p + \log K)$ with a large constant C_2 ;
- (iii) Either of the following two conditions holds with a sufficiently small constant C_3 :
 - (a) $\max_{k \in S} (\|\widehat{\boldsymbol{\beta}}^{(k)[0]} - \boldsymbol{\beta}^{(k)*}\|_2 \vee \|\widehat{\boldsymbol{\mu}}_1^{(k)[0]} - \boldsymbol{\mu}_1^{(k)*}\|_2 \vee \|\widehat{\boldsymbol{\mu}}_2^{(k)[0]} - \boldsymbol{\mu}_2^{(k)*}\|_2) \leq C_3 \min_{k \in S} \Delta^{(k)},$
 $\max_{k \in S} |\widehat{w}^{(k)[0]} - w^{(k)*}| \leq c_w/2$;
 - (b) $\max_{k \in S} (\|\widehat{\boldsymbol{\beta}}^{(k)[0]} + \boldsymbol{\beta}^{(k)*}\|_2 \vee \|\widehat{\boldsymbol{\mu}}_1^{(k)[0]} - \boldsymbol{\mu}_2^{(k)*}\|_2 \vee \|\widehat{\boldsymbol{\mu}}_2^{(k)[0]} - \boldsymbol{\mu}_1^{(k)*}\|_2) \leq C_3 \min_{k \in S} \Delta^{(k)},$
 $\max_{k \in S} |1 - \widehat{w}^{(k)[0]} - w^{(k)*}| \leq c_w/2$.
- (iv) $\min_{k \in S} \Delta^{(k)} \geq C_4 > 0$ with a large constant C_4 ;

Remark 1. These are common and mild conditions related to the sample size, initialization, and signal-to-noise ratio of GMMs. Condition (i) requires the maximum sample size of all tasks not to be much larger than the average sample size of tasks in S . Similar conditions can be found in [Duan and Wang \(2022\)](#). Condition (ii) is the requirement of the sample size of tasks in S . The usual condition for low-dimensional single-task learning is $n_k \gtrsim p$ ([Cai et al., 2019](#)). The additional $\log K$ term arises from the simultaneous control of performance on all tasks in S , where S can be as large as $1 : K$. Condition (iii) requires that the initialization should not be too far away from the truth, which is commonly assumed in either the analysis of EM algorithm ([Redner and Walker, 1984](#); [Balakrishnan et al., 2017](#); [Cai et al., 2019](#)) or other iterative algorithms like the local estimation used in semi-parametric models ([Carroll et al., 1997](#); [Li and Liang, 2008](#)) and adaptive Lasso ([Zou, 2006](#)). The two possible forms considered in this condition is due to the fact that binary GMM is only identifiable up to label permutation. Condition (iv) requires that the signal strength of GMM (in terms of Mahalanobis distance) is strong enough, which is usually assumed in literature about the likelihood-based methods of GMMs ([Dasgupta and Schulman, 2013](#); [Azizyan et al., 2013](#); [Balakrishnan et al., 2017](#); [Cai et al., 2019](#)).

We first establish the rate of convergence for the estimation. Recalling the parameter space $\overline{\Theta}_S(h)$ in (3), let us denote the true parameter by

$$\{\overline{\boldsymbol{\theta}}^{(k)*}\}_{k \in S} = \{(w^{(k)*}, \boldsymbol{\mu}_1^{(k)*}, \boldsymbol{\mu}_2^{(k)*}, \boldsymbol{\Sigma}^{(k)*})\}_{k \in S} \in \overline{\Theta}_S(h).$$

To better present the results for parameters related to the optimal discriminant rule (1), we further denote

$$\boldsymbol{\theta}^{(k)*} = (w^{(k)*}, \boldsymbol{\beta}^{(k)*}, \delta^{(k)*}), \quad \forall k \in S,$$

where $\boldsymbol{\beta}^{(k)*} = (\boldsymbol{\Sigma}^{(k)*})^{-1}(\boldsymbol{\mu}_2^{(k)*} - \boldsymbol{\mu}_1^{(k)*})$, $\delta^{(k)*} = \frac{1}{2}(\boldsymbol{\beta}^{(k)*})^T(\boldsymbol{\mu}_1^{(k)*} + \boldsymbol{\mu}_2^{(k)*})$. Note that $\boldsymbol{\theta}^{(k)*}$ is a function of $\overline{\boldsymbol{\theta}}^{(k)*}$. For the estimators returned by MTL-GMM (see Algorithm 1), we are particularly interested in the following two error metrics:

$$d(\widehat{\boldsymbol{\theta}}^{(k)[T]}, \boldsymbol{\theta}^{(k)*}) := \min\{|\widehat{w}^{(k)[T]} - w^{(k)*}| \vee \|\widehat{\boldsymbol{\beta}}^{(k)[T]} - \boldsymbol{\beta}^{(k)*}\|_2 \vee |\widehat{\delta}^{(k)[T]} - \delta^{(k)*}|\},$$

$$|1 - \widehat{w}^{(k)[T]} - w^{(k)*}| \vee \|\widehat{\boldsymbol{\beta}}^{(k)[T]} + \boldsymbol{\beta}^{(k)*}\|_2 \vee |\widehat{\delta}^{(k)[T]} + \delta^{(k)*}|, \quad (4)$$

$$\left(\min_{\pi: [2] \rightarrow [2]} \max_{r=1:2} \|\widehat{\boldsymbol{\mu}}_r^{(k)[T]} - \boldsymbol{\mu}_{\pi(r)}^{(k)*}\|_2 \right) \vee \|\widehat{\boldsymbol{\Sigma}}^{(k)[T]} - \boldsymbol{\Sigma}^{(k)*}\|_2, \quad (5)$$

where $\pi: [2] \rightarrow [2]$ is a permutation on $\{1, 2\}$. Again, we take the minimum above because binary GMM is identifiable up to label permutation. The first error metric $d(\widehat{\boldsymbol{\theta}}^{(k)[T]}, \boldsymbol{\theta}^{(k)*})$ involves the error for discriminant coefficients and is closely related to the clustering performance. It reveals how well our method utilizes similarity structure in multi-task learning. The second error metric is about the mean vectors and covariance matrix. As discussed in Section 2.1, we shall not expect it to be improved compared to that in single-task learning, as these parameters are not necessarily similar.

We are ready to present upper bounds for the estimation error of MTL-GMM. We recall that $\bar{\Theta}_S(h)$ and \mathbb{Q}_S are the parameter space and probability measure that we use in Section 2.1 to describe the data distributions for tasks in S and S^c , respectively.

Theorem 1. *(Upper bounds of the estimation error of GMM parameters for MTL-GMM) Suppose Assumption 1 holds for some S with $|S| \geq s$ and $\epsilon := \frac{K-s}{K} < a$ small constant c . Let $\lambda^{[0]} \geq C_1 \max_{k=1:K} \sqrt{n_k}$, $C_\lambda \geq C_1$ and $\kappa > C_2$ with some constants $C_1 > 0, C_2 \in (0, 1)$ ². Then there exist constants $C_3, C_4 > 0$, such that for any $\{\bar{\boldsymbol{\theta}}^{(k)*}\}_{k \in S} = \{(w^{(k)*}, \boldsymbol{\mu}_1^{(k)*}, \boldsymbol{\mu}_2^{(k)*}, \boldsymbol{\Sigma}^{(k)*})\}_{k \in S} \in \bar{\Theta}_S(h)$ and any probability measure \mathbb{Q}_S on $(\mathbb{R}^p)^{\otimes n_{S^c}}$, with probability $1 - C_3(K^{-1} + e^{-C_4 p})$, the following hold for all $k \in S$:*

$$d(\widehat{\boldsymbol{\theta}}^{(k)[T]}, \boldsymbol{\theta}^{(k)*}) \lesssim \sqrt{\frac{p}{n_S}} + \sqrt{\frac{\log K}{n_k}} + h \wedge \sqrt{\frac{p + \log K}{n_k}} + \epsilon \sqrt{\frac{p + \log K}{\max_{k=1:K} n_k}} + T^2(\kappa')^T,$$

$$\left(\min_{\pi: [2] \rightarrow [2]} \max_{r=1:2} \|\widehat{\boldsymbol{\mu}}_r^{(k)[T]} - \boldsymbol{\mu}_{\pi(r)}^{(k)*}\|_2 \right) \vee \|\widehat{\boldsymbol{\Sigma}}^{(k)[T]} - \boldsymbol{\Sigma}^{(k)*}\|_2 \lesssim \sqrt{\frac{p + \log K}{n_k}} + T^2(\kappa')^T,$$

where $\kappa' \in (0, 1)$ is some constant and $n_S = \sum_{k \in S} n_k$. When $T \geq C \log(\max_{k=1:K} n_k)$ with some large constant $C > 0$, the last term on the right-hand side will be dominated by other terms in both inequalities.

The upper bound of $\left(\min_{\pi: [2] \rightarrow [2]} \max_{r=1:2} \|\widehat{\boldsymbol{\mu}}_r^{(k)[T]} - \boldsymbol{\mu}_{\pi(r)}^{(k)*}\|_2 \right) \vee \|\widehat{\boldsymbol{\Sigma}}^{(k)[T]} - \boldsymbol{\Sigma}^{(k)*}\|_2$ contains two parts. The first part is comparable to the single-task learning rate (Cai et al., 2019) (up to a $\sqrt{\log K}$ term due to the simultaneous control over all tasks in S), and the second part characterizes the geometric convergence of iterates. As expected, since $\boldsymbol{\mu}_1^{(k)*}, \boldsymbol{\mu}_2^{(k)*}, \boldsymbol{\Sigma}^{(k)*}$ in S are not necessarily similar, an improved error rate over single-task learning is generally impossible. The upper bound for $d(\widehat{\boldsymbol{\theta}}^{(k)[T]}, \boldsymbol{\theta}^{(k)*})$ is directly related to the clustering performance of our method. Thus we will provide a detailed discussion about it after presenting the clustering result in the next theorem.

² C_1 and C_2 depend on the constants M, c_w , and c_Σ etc.

As introduced in Section 1.1, using the estimate $\hat{\boldsymbol{\theta}}^{(k)[T]} = (\hat{w}^{(k)[T]}, \hat{\boldsymbol{\beta}}^{(k)[T]}, \hat{\delta}^{(k)[T]})$ from Algorithm 1, we can construct a classifier for task k as

$$\hat{\mathcal{C}}^{(k)[T]}(\mathbf{z}) = \begin{cases} 1, & \text{if } (\hat{\boldsymbol{\beta}}^{(k)[T]})^T \mathbf{z} - \hat{\delta}^{(k)[T]} \leq \log \left(\frac{1 - \hat{w}^{(k)[T]}}{\hat{w}^{(k)[T]}} \right); \\ 2, & \text{otherwise.} \end{cases} \quad (6)$$

Recall that for a clustering method $\mathcal{C} : \mathbb{R}^p \rightarrow \{1, 2\}$, its mis-clustering error rate under GMM with parameter $\bar{\boldsymbol{\theta}} = (w, \boldsymbol{\mu}_1, \boldsymbol{\mu}_2, \boldsymbol{\Sigma})$ is

$$R_{\bar{\boldsymbol{\theta}}}(\mathcal{C}) = \min_{\pi: [2] \rightarrow [2]} \mathbb{P}_{\bar{\boldsymbol{\theta}}}(\mathcal{C}(Z^{\text{new}}) \neq \pi(Y^{\text{new}})), \quad (7)$$

where $Z^{\text{new}} \sim (1 - w)\mathcal{N}(\boldsymbol{\mu}_1, \boldsymbol{\Sigma}) + w\mathcal{N}(\boldsymbol{\mu}_2, \boldsymbol{\Sigma})$ is a future observation associated with the label Y^{new} , independent from \mathcal{C} ; the probability $\mathbb{P}_{\bar{\boldsymbol{\theta}}}$ is w.r.t. $(Z^{\text{new}}, Y^{\text{new}})$, and the minimum is taken over two permutation functions on $\{1, 2\}$. Denote $\mathcal{C}_{\bar{\boldsymbol{\theta}}}$ as the Bayes classifier that minimizes $R_{\bar{\boldsymbol{\theta}}}(\mathcal{C})$. In the following theorem, we obtain the upper bound of the excess mis-clustering error of $\hat{\mathcal{C}}^{(k)[T]}$ for $k \in S$.

Theorem 2. (*Upper bound of the excess mis-clustering error for MTL-GMM*) Suppose the same conditions in Theorem 1 hold. Then there exist constants C_1 and C_2 such that for any $\{\bar{\boldsymbol{\theta}}^{(k)*}\}_{k \in S} \in \bar{\Theta}_S(h)$ and any probability measure \mathbb{Q}_S on $(\mathbb{R}^p)^{\otimes n_{S^c}}$, with probability $1 - C_1(K^{-1} + e^{-C_2 p})$, the following holds for all $k \in S$:

$$\begin{aligned} R_{\bar{\boldsymbol{\theta}}^{(k)*}}(\hat{\mathcal{C}}^{(k)[T]}) - R_{\bar{\boldsymbol{\theta}}^{(k)*}}(\mathcal{C}_{\bar{\boldsymbol{\theta}}^{(k)*}}) &\lesssim d^2(\hat{\boldsymbol{\theta}}^{(k)[T]}, \boldsymbol{\theta}^{(k)*}) \\ &\lesssim \underbrace{\frac{p}{n_S}}_{\text{(I)}} + \underbrace{\frac{\log K}{n_k}}_{\text{(II)}} + \underbrace{h^2 \wedge \frac{p + \log K}{n_k}}_{\text{(III)}} + \underbrace{\epsilon^2 \frac{p + \log K}{\max_{k=1:K} n_k}}_{\text{(IV)}} + \underbrace{T^4 (\kappa')^{2T}}_{\text{(V)}}, \end{aligned}$$

with some $\kappa' \in (0, 1)$. When $T \geq C \log(\max_{k=1:K} n_k)$ with a large constant $C > 0$, the last term on the right-hand side will be dominated by the second term.

The upper bounds of $d(\hat{\boldsymbol{\theta}}^{(k)[T]}, \boldsymbol{\theta}^{(k)*})$ in Theorem 1 and $R_{\bar{\boldsymbol{\theta}}^{(k)*}}(\hat{\mathcal{C}}^{(k)[T]}) - R_{\bar{\boldsymbol{\theta}}^{(k)*}}(\mathcal{C}_{\bar{\boldsymbol{\theta}}^{(k)*}})$ in Theorem 2 consist of five parts with one-to-one correspondence. It is sufficient to discuss the bound of $R_{\bar{\boldsymbol{\theta}}^{(k)*}}(\hat{\mathcal{C}}^{(k)[T]}) - R_{\bar{\boldsymbol{\theta}}^{(k)*}}(\mathcal{C}_{\bar{\boldsymbol{\theta}}^{(k)*}})$. Part (I) represents the ‘‘oracle rate’’, which can be achieved when all tasks in S are the same. This is the best rate to possibly achieve. Part (II) is a dimension-free error caused by estimating scalar parameters $\delta^{(k)*}$ and $w^{(k)*}$ that appears in the optimal discriminant rule. Part (III) includes h that measures the degree of similarity among the tasks in S . When these tasks are very similar, h will be small, contributing a small term to the upper bound. Nicely, even when h is large, the term becomes $\frac{p + \log K}{n_k}$ and it is still comparable to the minimax error rate of single-task learning $\mathcal{O}_p(p/n_k)$ (e.g., Theorems 4.1 and 4.2 in Cai et al. (2019)). We have the extra $\log K$ term here due to the simultaneous control over all tasks in S . Part (IV) quantifies the influence from the outlier tasks in S^c . When there are more outlier tasks, ϵ increases and the

bound becomes worse. On the other hand, as long as ϵ is small enough to make this term dominated by any other part, the error rate induced by outlier tasks becomes negligible. Given that data from outlier tasks can be arbitrarily contaminated, we can conclude that our method is robust against a fraction of outlier tasks from arbitrary sources. The term in Part (V) decreases geometrically in the iteration number T , implying that the iterates in Algorithm 1 converge geometrically to a ball of radius determined by the errors from Parts (I)-(IV).

After explaining each part of the upper bound, we now compare it with the convergence rate $\mathcal{O}_p(\frac{p+\log K}{n_k})$ (including $\log K$ here since we consider all the tasks simultaneously) in the single-task learning and reveal how our method performs. With a quick inspection we can conclude the following:

- The rate of the upper bound is never larger than $\frac{p+\log K}{n_k}$. So in terms of rate of convergence, our method MTL-GMM performs at least as good as single-task learning, regardless of the similarity level h and outlier task fraction ϵ .
- When $n_S \gg n_k$ (large total sample size), p increases with n_k (diverging dimension), $h \ll \sqrt{\frac{p+\log K}{n_k}}$ (sufficient similarity between tasks in S), and $\epsilon \ll \sqrt{(\max_{k=1:K} n_k)/n_k}$ (small fraction of outlier tasks), MTL-GMM attains a faster excess mis-clustering error rate and improves over single-task learning.

The preceding discussions on the upper bounds have demonstrated the superiority of our method. But can we do better? To further evaluate the upper bounds of our method, we next derive complementary minimax lower bounds for both estimation error and excess mis-clustering error. We will show that our method is (nearly) minimax rate optimal in a broad range of regimes.

Theorem 3. (Lower bounds of the estimation error of GMM parameters in multi-task learning) Suppose $\epsilon = \frac{K-s}{K}$ is bounded by a small constant. Suppose there exists a subset S with $|S| \geq s$ such that $\min_{k \in S} n_k \geq C_1(p + \log K)$ with a large constant C_1 and $\min_{k \in S} \Delta^{(k)} \geq$ some constant C_2 . Then

$$\inf_{\{\widehat{\boldsymbol{\theta}}^{(k)}\}_{k=1}^K} \sup_{S: |S| \geq s} \sup_{\substack{\{\boldsymbol{\theta}^{(k)*}\}_{k \in S} \in \overline{\Theta}_S(h) \\ \mathbb{Q}_S}} \mathbb{P} \left(\bigcup_{k \in S} \left\{ d(\widehat{\boldsymbol{\theta}}^{(k)}, \boldsymbol{\theta}^{(k)*}) \gtrsim \sqrt{\frac{p}{n_S}} + \sqrt{\frac{\log K}{n_k}} \right. \right. \\ \left. \left. + h \wedge \sqrt{\frac{p + \log K}{n_k}} + \frac{\epsilon}{\sqrt{\max_{k=1:K} n_k}} \right\} \right) \geq \frac{1}{10},$$

$$\inf_{\substack{\{\widehat{\boldsymbol{\mu}}_1^{(k)}, \widehat{\boldsymbol{\mu}}_2^{(k)}\}_{k=1}^K \\ \{\widehat{\boldsymbol{\Sigma}}^{(k)}\}_{k=1}^K}} \sup_{S: |S| \geq s} \sup_{\substack{\{\boldsymbol{\theta}^{(k)*}\}_{k \in S} \in \overline{\Theta}_S(h) \\ \mathbb{Q}_S}} \mathbb{P} \left(\bigcup_{k \in S} \left\{ \left(\min_{\pi: [2] \rightarrow [2]} \max_{r=1:2} \|\widehat{\boldsymbol{\mu}}_r^{(k)} - \boldsymbol{\mu}_{\pi(r)}^{(k)*}\|_2 \right) \vee \|\widehat{\boldsymbol{\Sigma}}^{(k)} - \boldsymbol{\Sigma}^{(k)*}\|_2 \right\} \right)$$

$$\gtrsim \sqrt{\frac{p + \log K}{n_k}} \Bigg\} \Bigg) \geq \frac{1}{10}.$$

Theorem 4. (Lower bound of the excess mis-clustering error in multi-task learning) Suppose the same conditions in Theorem 3 hold. Then

$$\inf_{\{\widehat{\mathcal{C}}^{(k)}\}_{k=1}^K} \sup_{S: |S| \geq s} \sup_{\{\boldsymbol{\theta}^{(k)*}\}_{k \in S} \in \overline{\Theta}_S(h)} \mathbb{P} \left(\bigcup_{k \in S} \left\{ R_{\boldsymbol{\theta}^{(k)*}}(\widehat{\mathcal{C}}^{(k)}) - R_{\boldsymbol{\theta}^{(k)*}}(\mathcal{C}_{\boldsymbol{\theta}^{(k)*}}) \gtrsim \frac{p}{n_S} + \frac{\log K}{n_k} \right. \right. \\ \left. \left. + h^2 \wedge \frac{p + \log K}{n_k} + \frac{\epsilon^2}{\max_{k=1:K} n_k} \right\} \right) \geq \frac{1}{10}.$$

Comparing the upper and lower bounds in Theorems 1-4, we make several remarks:

- Regarding the estimation of mean vectors $\{\boldsymbol{\mu}_1^{(k)*}, \boldsymbol{\mu}_2^{(k)*}\}_{k \in S}$ and covariance matrices $\{\boldsymbol{\Sigma}^{(k)*}\}_{k \in S}$, the upper and lower bounds match, hence our method is minimax rate optimal.
- For the estimation error $d(\widehat{\boldsymbol{\theta}}^{(k)}, \boldsymbol{\theta}^{(k)*})$ and excess mis-clustering error $R_{\widehat{\boldsymbol{\theta}}^{(k)*}}(\widehat{\mathcal{C}}^{(k)[T]}) - R_{\boldsymbol{\theta}^{(k)*}}(\mathcal{C}_{\boldsymbol{\theta}^{(k)*}})$ with $k \in S$, the first three terms in the upper and lower bounds match. Only the term involving ϵ in the lower bound differs from that in the upper bound by a factor $\sqrt{p + \log K}$ or $p + \log K$. As a result, in the classical low-dimensional regime where p is bounded, the upper and lower bounds match (up to a logarithmic factor). Therefore, our method is (nearly) minimax rate optimal for estimating $\{\boldsymbol{\theta}^{(k)*}\}_{k \in S}$ and clustering in such a classical regime.
- When the dimension p diverges, there might exist a non-negligible gap between the upper and lower bounds for $d(\widehat{\boldsymbol{\theta}}^{(k)}, \boldsymbol{\theta}^{(k)*})$ and $R_{\widehat{\boldsymbol{\theta}}^{(k)*}}(\widehat{\mathcal{C}}^{(k)[T]}) - R_{\boldsymbol{\theta}^{(k)*}}(\mathcal{C}_{\boldsymbol{\theta}^{(k)*}})$ with $k \in S$. Nevertheless, this only occurs when the fraction of outlier task ϵ is above the threshold $\sqrt{\frac{\max_{k=1:K} n_k}{p + \log K}} (h^2 \vee \frac{\log K}{n_k} \vee \frac{p}{n_S})$. Below the threshold, our method remains minimax rate optimal even though p is unbounded.
- Does the gap, when exist, arise from the upper bound or the lower bound? We believe that it is the upper bound that sometimes becomes not sharp. As can be seen from the proof of Theorems 1, the term $\epsilon \sqrt{\frac{p + \log K}{\max_{k=1:K} n_k}}$ is due to the estimation of those “center” parameters in Algorithm 1. Recent advances in robust statistics (Chen et al., 2018) have shown that estimators based on statistical depth functions such as Tukey’s depth function (Tukey, 1975) can achieve optimal minimax rate under Huber’s ϵ -contamination model for location and covariance estimation. It might be possible to utilize depth functions to estimate “center” parameters in our problem and kill the factor \sqrt{p} in the upper bound. We leave a rigorous development of optimal robustness as an interesting future research. On the other hand, such a statistical improvement may come with expensive computation, as depth function based estimation typically requires solving a challenging non-convex optimization problem.

2.4 Initialization and cluster alignment

As specified by Condition (iii) in Assumption 1, our proposed learning procedure requires that for each task in S , initial values of the GMM parameter estimates lie within a distance of SNR-order from the ground truth. This can be satisfied by the method of moments proposed in Ge et al. (2015). In practice, a natural initialization method is to run the standard EM algorithm or other common clustering methods like k -means on each task and use the corresponding estimate as the initial values. We adopted the standard EM algorithm in our numerical experiments, and the numerical results in Section 4 and Appendix showed that this practical initialization works quite well. However, in the context of multi-task learning, Condition (iii) further require a correct alignment of those good initializations from each task, owing to the non-identifiability of GMMs. We discuss in detail the alignment issue in Section 2.4.1 and propose two algorithms to resolve this issue in Section 2.4.2.

2.4.1 The alignment issue

Recall that Section 2.1 introduces the binary GMM with parameters $(w^{(k)*}, \boldsymbol{\mu}_1^{(k)*}, \boldsymbol{\mu}_2^{(k)*}, \boldsymbol{\Sigma}^{(k)*})$ for each task $k \in S$. Because the two sets of parameter values $\{(w, \mathbf{u}, \mathbf{v}, \boldsymbol{\Sigma}), (1-w, \mathbf{v}, \mathbf{u}, \boldsymbol{\Sigma})\}$ for $(w^{(k)*}, \boldsymbol{\mu}_1^{(k)*}, \boldsymbol{\mu}_2^{(k)*}, \boldsymbol{\Sigma}^{(k)*})$ index the same distribution, a good initialization close to the truth is up to a permutation of the two cluster labels. The permutations in the initialization of different tasks could be different. Therefore, in light of the joint parameter space $\bar{\Theta}_S(h)$ defined in (3) and Condition (iii) in Assumption 1, for given initializations from different tasks, we may need to permute their cluster labels to feed the well-aligned initialization into Algorithm 1.

We further elaborate on the alignment issue using Algorithm 1. The penalization in Step 9 aims to push the estimators $\hat{\boldsymbol{\beta}}^{(k)[t]}$'s with different k towards each other, which is expected to improve the performance thanks to the similarity among underlying true parameters $\{\boldsymbol{\beta}^{(k)*}\}_{k \in S}$. However, due to the potential permutation of two cluster labels, the vanilla single-task initializations (without alignment) cannot guarantee that the estimators $\{\hat{\boldsymbol{\beta}}^{(k)[t]}\}_{k \in S}$ at each iteration are all estimating the corresponding $\boldsymbol{\beta}^{(k)*}$'s (some may estimate $-\boldsymbol{\beta}^{(k)*}$'s).

Figure 1 illustrates the alignment issue in the case of two tasks. The left-hand-side situation is ideal where $\hat{\boldsymbol{\beta}}^{(1)[0]}, \hat{\boldsymbol{\beta}}^{(2)[0]}$ are estimates of $\boldsymbol{\beta}^{(1)*}, \boldsymbol{\beta}^{(2)*}$ (which are similar). The right-hand-side situation is problematic because $\hat{\boldsymbol{\beta}}^{(1)[0]}, \hat{\boldsymbol{\beta}}^{(2)[0]}$ are estimates of $\boldsymbol{\beta}^{(1)*}, -\boldsymbol{\beta}^{(2)*}$ (which are not similar). Therefore, in practice, after obtaining the initializations from each task, it is necessary to align their cluster labels to ensure that estimators of similar parameters are correctly put together in the penalization framework in Algorithm 1. We formalize the problem and provide two solutions in the next subsection.

2.4.2 Two alignment algorithms

Suppose $\{\hat{\boldsymbol{\beta}}^{(k)[0]}\}_{k=1}^K$ are the initial estimates of discriminant coefficients with potentially bad alignment for $k \in S$. Note that a good initialization and alignment is not required (in




Figure 1: Examples of well-aligned (left) and badly-aligned (right) initializations.

fact it is not even well defined) for the outlier tasks in S^c , because they can be from arbitrary distributions. However, since S is unknown, we will have to address the alignment issue for tasks in S based on initial estimates from all the tasks. For binary GMMs, each alignment of $\{\hat{\beta}^{(k)[0]}\}_{k=1}^K$ can be represented by a K -dimensional Rademacher vector $\mathbf{r} \in \{\pm 1\}^K$. Define the ideal alignment as $r_k^* = \arg \min_{r_k = \pm 1} \|r_k \hat{\beta}^{(k)[0]} - \beta^{(k)*}\|_2, k \in S$. The goal is to recover the well-aligned initializers $\{r_k^* \hat{\beta}^{(k)[0]}\}_{k \in S}$ from the initial estimates $\{\hat{\beta}^{(k)[0]}\}_{k=1}^K$ (equivalently, to recover $\{r_k^*\}_{k \in S}$), which can then be fed into Algorithm 1. Once $\{\hat{\beta}^{(k)[0]}\}_{k \in S}$ are well aligned, other initial estimates in Algorithm 1 will be automatically well aligned.

In the following, we will introduce two alignment algorithms. The first one is the “exhaustive search” method (Algorithm 2), where we search among all possible alignments to find the best one. The second one is the “greedy search” method (Algorithm 3), where we flip the sign of $\hat{\beta}^{(k)[0]}$ in a greedy way to recover $\{r_k^* \hat{\beta}^{(k)[0]}\}_{k \in S}$. Both methods are proved to recover $\{r_k^*\}_{k \in S}$ under mild conditions. The conditions required by the “exhaustive search” method are slightly weaker than those required by the “greedy search” method. As for computational complexity, the latter enjoys a linear time complexity $\mathcal{O}(K)$, while the former suffers from an exponential time complexity $\mathcal{O}(2^K)$ due to optimization over all possible 2^K alignments.

To this end, for a given alignment $\mathbf{r} = \{r_k\}_{k=1}^K \in \{\pm 1\}^K$ with the correspondingly aligned estimates $\{r_k \hat{\beta}^{(k)[0]}\}_{k=1}^K$, define its alignment score as

$$\text{score}(\mathbf{r}) = \sum_{1 \leq k_1 \neq k_2 \leq K} \|r_{k_1} \hat{\beta}^{(k_1)[0]} - r_{k_2} \hat{\beta}^{(k_2)[0]}\|_2.$$

The intuition is that as long as the initializations $\{\hat{\beta}^{(k)[0]}\}_{k \in S}$ are close to the ground

truth, a smaller score indicates less difference among $\{r_k \hat{\beta}^{(k)[0]}\}_{k \in S}$, which implies a better alignment. The score can be thus used to evaluate the quality of an alignment. Note that the score is defined in a symmetric way, that is, $\text{score}(\mathbf{r}) = \text{score}(-\mathbf{r})$. The exhaustive search algorithm is presented in Algorithm 2, where scores of all alignments are calculated, and the alignment that minimizes the score is output. Since the score is symmetric, there are at least two alignments with the minimum score. The algorithm can arbitrarily choose and output one of them.

Algorithm 2: Exhaustive search for the alignment

Input: Initialization $\{\hat{\beta}^{(k)[0]}\}_{k=1}^K$
1 $\hat{\mathbf{r}} \leftarrow \arg \min_{\mathbf{r} \in \{\pm 1\}^K} \text{score}(\mathbf{r})$
Output: $\hat{\mathbf{r}}$

The following theorem reveals that the exhaustive search algorithm can successfully find the ideal alignment under mild conditions.

Theorem 5 (Alignment correctness for Algorithm 2). *Assume that*

$$(i) \quad \epsilon < \frac{1}{3};$$

$$(ii) \quad \min_{k \in S} \|\beta^{(k)*}\|_2 \geq \frac{4(1-\epsilon)}{1-3\epsilon} h + \frac{2(2-\epsilon)}{1-3\epsilon} \max_{k \in S} (\|\hat{\beta}^{(k)[0]} - \beta^{(k)*}\|_2 \wedge \|\hat{\beta}^{(k)[0]} + \beta^{(k)*}\|_2),$$

where $\epsilon = \frac{K-|S|}{K}$ is the outlier task proportion introduced in Theorem 1, and h is the similarity level of discriminant coefficient defined in (3). Then the output of Algorithm 2 satisfies

$$\hat{r}_k = r_k^* \text{ for all } k \in S \quad \text{or} \quad \hat{r}_k = -r_k^* \text{ for all } k \in S$$

Remark 2. The conditions imposed in Theorem 5 are **no stronger** than conditions required by Theorem 1. First of all, Condition (i) is also required in Theorem 1. Moreover, from the definition of h in (3), it is bounded by a constant. This together with Conditions (iii) and (iv) in Assumption 1 implies Condition (ii) in Theorem 5.

Remark 3. With Theorem 5, we can relax the original Condition (iii) in Assumption 1 to the following condition:

For all $k \in S$, either of the following two conditions holds with a sufficiently small constant C_3 :

$$(a) \quad \|\hat{\beta}^{(k)[0]} - \beta^{(k)*}\|_2 \vee \|\hat{\mu}_1^{(k)[0]} - \mu_1^{(k)*}\|_2 \vee \|\hat{\mu}_2^{(k)[0]} - \mu_2^{(k)*}\|_2 \leq C_3 \min_{k \in S} \Delta^{(k)}, \quad |\hat{w}^{(k)[0]} - w^{(k)*}| \leq c_w/2;$$

$$(b) \quad \|\hat{\beta}^{(k)[0]} + \beta^{(k)*}\|_2 \vee \|\hat{\mu}_1^{(k)[0]} - \mu_2^{(k)*}\|_2 \vee \|\hat{\mu}_2^{(k)[0]} - \mu_1^{(k)*}\|_2 \leq C_3 \min_{k \in S} \Delta^{(k)}, \quad |1 - \hat{w}^{(k)[0]} - w^{(k)*}| \leq c_w/2.$$

In the relaxed version, the initialization for each task only needs to be good up to an **arbitrary** permutation, while in the original version, the initialization for each task needs to be good under **the same permutation**.

Next, we would like to introduce the second alignment algorithm, the “greedy search” method, summarized in Algorithm 3. The main idea is to flip the sign of the discriminant coefficient estimates (equivalently, swap the two cluster labels) from K tasks in a sequential fashion to check whether the alignment score decreases or not. If yes, we keep the alignment after the flip and proceed with the next task. Otherwise, we keep the alignment before the flip and proceed with the next task. A surprising fact of Algorithm 3 is that it is sufficient to iterate this procedure for all tasks just **once** to recover the ideal alignment, making the algorithm computationally efficient.

Algorithm 3: Greedy search for the alignment

Input: Initialization $\{\widehat{\beta}^{(k)[0]}\}_{k=1}^K$
 $\mathbf{1} \ \widehat{\mathbf{r}} = (1, \dots, 1) \in \{\pm 1\}^K$
 $\mathbf{2} \ \text{for } k = 1 \text{ to } K \text{ do}$
 $\mathbf{3} \quad \widetilde{\mathbf{r}} \leftarrow \text{flip the sign of } \widehat{r}_k \text{ in } \widehat{\mathbf{r}}$
 $\mathbf{4} \quad \text{if } \text{score}(\widehat{\mathbf{r}}) > \text{score}(\widetilde{\mathbf{r}}) \text{ then}$
 $\mathbf{5} \quad \quad \widehat{\mathbf{r}} \leftarrow \widetilde{\mathbf{r}}$
 $\mathbf{6} \quad \text{end}$
 $\mathbf{7} \ \text{end}$
Output: $\widehat{\mathbf{r}}$

To help state the theory of the greedy search algorithm, we define the “mismatch proportion” of $\{\widehat{\beta}^{(k)[0]}\}_{k \in S}$ as

$$p_a = \min\{\#\{k \in S : r_k^* = 1\}, \#\{k \in S : r_k^* = -1\}\} / |S|$$

Intuitively, p_a represents the level of mismatch between the initial alignment and the ideal one. It's straightforward to verify that $p_a \in [0, 1/2]$; $p_a = 0$ means the initial alignment equals the ideal one, while $p_a = 1/2$ (or $\frac{|S|-1}{2|S|}$ when $|S|$ is odd) represents the “worst” alignment, where almost half of tasks are badly-aligned. The smaller p_a is, the better alignment $\{\widehat{\beta}^{(k)[0]}\}_{k \in S}$ is. Note that we only care about the alignment of tasks in S .

The following theorem shows that the greedy search algorithm can succeed in finding the ideal alignment under mild conditions.

Theorem 6 (Alignment correctness for Algorithm 3). *Assume that*

- (i) $\epsilon < \frac{1}{2}$;
- (ii) $\min_{k \in S} \|\beta^{(k)*}\|_2 \geq \frac{2(1-\epsilon)}{2(1-\epsilon)(1-p_a)-1} h + \frac{2-\epsilon}{2(1-\epsilon)(1-p_a)-1} \max_{k \in S} (\|\widehat{\beta}^{(k)[0]} - \beta^{(k)*}\|_2 \wedge \|\widehat{\beta}^{(k)[0]} + \beta^{(k)*}\|_2)$;
- (iii) $p_a < \frac{1-2\epsilon}{2(1-\epsilon)}$,

where ϵ and h are the same as in Theorem 5. Then the output of Algorithm 3 satisfies

$$r_k = r_k^* \text{ for all } k \in S \quad \text{or} \quad r_k = -r_k^* \text{ for all } k \in S$$

Remark 4. Conditions (i) and (ii) required by Theorem 6 are similar to the requirements in Theorem 5, which have been shown to be **no stronger** than conditions in Assumption 1 and Theorem 1 (See Remark 2). However, Condition (iii) is an additional requirement for the success of the greedy label swapping algorithm. The intuition is that in the exhaustive search algorithm, we compare the scores of all alignments and only need to ensure the ideal alignment can defeat the badly-aligned ones in terms of the alignment score. In contrast, the success of the greedy search algorithm relies on the correct move at each step. We need to guarantee that the “better” alignment after the swap (which may still be badly aligned) can outperform the “worse” one before the swap. This is more difficult to satisfy, hence more conditions are needed for the success of Algorithm 3. Condition (iii) is one such condition to provide a reasonably good initial alignment to start the greedy search process. More details of the analysis can be found in the proofs of Theorems 5 and 6 in Appendix.

Remark 5. In practice, Condition (iii) can fail to hold with a non-zero probability. One solution is to start with random alignments, run the greedy search algorithm multiple times and use the alignment that appears most frequently in the output. Nevertheless, this will increase the computational burden. In our numerical studies, Algorithm 3 without multiple random alignments works well.

One appealing feature of the two alignment algorithms is that they are robust against a fraction of outlier tasks from arbitrary distributions. According to the definition of the alignment score, this may appear impossible at the first glance because the score depends on the estimators from all tasks. However, it turns out that the impact of outliers when comparing the scores in Algorithm 2 and 3 can be bounded by parameters and constants that are unrelated to outlier tasks via the triangle inequality of Euclidean norms. The key idea is that the alignment of outlier tasks in S^c does not matter in Theorems 5 and 6. More details can be found in the proof of Theorems 5 and 6 in Appendix.

In contrast with supervised MTL, the alignment issue commonly exists in unsupervised MTL. It generally occurs when aggregating information (up to latent label permutation) across different tasks. Alignment pre-processing is thus necessary and important. However, to our knowledge, there is no formal discussion regarding alignment in existing literatures of unsupervised MTL (Gu et al., 2011; Zhang and Zhang, 2011; Yang et al., 2014; Zhang et al., 2018; Dieuleveut et al., 2021; Marfoq et al., 2021). Our treatment of alignment in Section 2.4 is an important step forward in this field. Our algorithms can be potentially extended to other unsupervised MTL scenarios and we leave it for future studies.

3 Extension to Multi-cluster GMMs

In previous sections, we have discussed the MTL problem for binary GMMs. In this section, we extend our methods and theory to Gaussian mixtures with R clusters ($R \geq 3$).

We first generalize the problem setting introduced in Sections 1.1 and 2.1. There are K tasks where we have n_k observations $\{\mathbf{z}_i^{(k)}\}_{i=1}^{n_k}$ from the k -th task. An *unknown* subset

$S \subseteq 1 : K$ denotes tasks whose samples follow multi-cluster GMMs, and S^c refers to outlier tasks that can have arbitrary distributions. Specifically, for all $k \in S, i = 1 : n_k$,

$$y_i^{(k)} = r \text{ with probability } w_r^{(k)*}, \quad \mathbf{z}_i^{(k)} | y_i^{(k)} = r \sim \mathcal{N}(\boldsymbol{\mu}_r^{(k)*}, \boldsymbol{\Sigma}^{(k)*}), \quad r = 1 : R,$$

with $\sum_{r=1}^R w_r^{(k)*} = 1$, and

$$\{\mathbf{z}_i^{(k)}\}_{i,k \in S^c} \sim \mathbb{Q}_S,$$

where \mathbb{Q}_S is some probability measure on $(\mathbb{R}^p)^{\otimes n_{S^c}}$ and $n_{S^c} = \sum_{k \in S^c} n_k$. We focus on the following joint parameter space

$$\bar{\Theta}_S(h) = \left\{ \{\bar{\boldsymbol{\theta}}^{(k)}\}_{k \in S} = \{(\{w_r^{(k)}\}_{r=1}^R, \{\boldsymbol{\mu}_r^{(k)}\}_{r=1}^R, \boldsymbol{\Sigma}^{(k)})\}_{k \in S} : \bar{\boldsymbol{\theta}}^{(k)} \in \bar{\Theta}, \max_{r=2:R} \inf_{\boldsymbol{\beta}} \max_{k \in S} \|\boldsymbol{\beta}_r^{(k)} - \boldsymbol{\beta}\|_2 \leq h \right\}, \quad (8)$$

where $\boldsymbol{\beta}_r^{(k)} = (\boldsymbol{\Sigma}^{(k)})^{-1}(\boldsymbol{\mu}_r^{(k)} - \boldsymbol{\mu}_1^{(k)})$ is the r -th discriminant coefficient in the k -th task, and $\bar{\Theta}$ is the parameter space for a single multi-cluster GMM,

$$\bar{\Theta} = \left\{ \bar{\boldsymbol{\theta}} = (\{w_r\}_{r=1}^R, \{\boldsymbol{\mu}_r\}_{r=1}^R, \boldsymbol{\Sigma}) : \max_{r=1:R} \|\boldsymbol{\mu}_r\|_2 \leq M, c_w \leq \min_{r=1:R} w_r \leq \max_{r=1:R} w_r \leq 1 - c_w, \sum_{r=1}^R w_r = 1, c_{\boldsymbol{\Sigma}}^{-1} \leq \lambda_{\min}(\boldsymbol{\Sigma}) \leq \lambda_{\max}(\boldsymbol{\Sigma}) \leq c_{\boldsymbol{\Sigma}} \right\}. \quad (9)$$

Note that (8) and (9) are natural generalizations of (3) and (2), respectively.

Under a multi-cluster GMM with parameter $\bar{\boldsymbol{\theta}} = (\{w_r\}_{r=1}^R, \{\boldsymbol{\mu}_r\}_{r=1}^R, \boldsymbol{\Sigma})$, compared with (1), the optimal discriminant rule now becomes

$$\mathcal{C}_{\bar{\boldsymbol{\theta}}}(\mathbf{z}) = \arg \max_{r=1:R} \left\{ \left(\mathbf{z} - \frac{\boldsymbol{\mu}_1 + \boldsymbol{\mu}_r}{2} \right)^T \boldsymbol{\beta}_r + \log \left(\frac{w_r}{w_1} \right) \right\}, \quad (10)$$

where $\boldsymbol{\beta}_r = \boldsymbol{\Sigma}^{-1}(\boldsymbol{\mu}_r - \boldsymbol{\mu}_1)$. Once we have the parameter estimators, we plug them into the above rule to obtain the plug-in clustering method. Recall that for a clustering method $\mathcal{C} : \mathbb{R}^p \rightarrow [R]$, its mis-clustering error is

$$R_{\bar{\boldsymbol{\theta}}}(\mathcal{C}) = \min_{\pi : [R] \rightarrow [R]} \mathbb{P}_{\bar{\boldsymbol{\theta}}}(\mathcal{C}(Z^{\text{new}}) \neq \pi(Y^{\text{new}})).$$

Here, $Z^{\text{new}} \sim \sum_{r=1}^R w_r \mathcal{N}(\boldsymbol{\mu}_r, \boldsymbol{\Sigma})$ is an independent future observation associated with the label Y^{new} , the probability $\mathbb{P}_{\bar{\boldsymbol{\theta}}}$ is w.r.t. $(Z^{\text{new}}, Y^{\text{new}})$, and the minimum is taken over $R!$ permutation functions on $[R]$. Since (10) is the optimal clustering method that minimizes $R_{\bar{\boldsymbol{\theta}}}(\mathcal{C})$, the excess mis-clustering error for a given clustering \mathcal{C} is $R_{\bar{\boldsymbol{\theta}}}(\mathcal{C}) - R_{\bar{\boldsymbol{\theta}}}(\mathcal{C}_{\bar{\boldsymbol{\theta}}})$. The rest of the section aims to extend the EM-stylized multi-task learning procedure and the two alignment algorithms in Section 2 to the general multi-cluster GMM setting, and provide similar statistical guarantees in terms of estimation and excess mis-clustering errors. For simplicity, throughout this section, we assume the number of clusters R to be bounded and known. We leave the case of diverging R as a future work.

Since both the EM algorithm and the penalization framework work beyond binary GMM, our methodological idea described in Section 2.2 can be directly adapted to extend Algorithm 1 to the multi-cluster case. We summarize the general procedure in Algorithm 4. Like in Algorithm 1, we have adopted the following notation for posterior probability in Algorithm 4,

$$\gamma_{\boldsymbol{\theta}}^{(r)}(\mathbf{z}) = \frac{w_r \exp(\boldsymbol{\beta}_r^T \mathbf{z} - \delta_r)}{\sum_{r=1}^R w_r \exp(\boldsymbol{\beta}_r^T \mathbf{z} - \delta_r)}, \quad \text{for } \boldsymbol{\theta} = (\{w_r\}_{r=2}^R, \{\boldsymbol{\beta}_r\}_{r=2}^R, \{\delta_r\}_{r=2}^R),$$

where $w_1 = 1 - \sum_{r=2}^R w_r$, $\boldsymbol{\beta}_1 := \mathbf{0}$, and $\delta_1 := 0$. Specifically, $\gamma_{\boldsymbol{\theta}}^{(r)}(\mathbf{z})$ is the posterior probability $\mathbb{P}(Y = r | Z = \mathbf{z})$ given the observation \mathbf{z} , when the true parameter of a multi-cluster GMM $(\{w_r^*\}_{r=1}^R, \{\boldsymbol{\mu}_r^*\}_{r=1}^R, \boldsymbol{\Sigma}^*)$ satisfies $w_r = w_r^*$, $\boldsymbol{\beta}_r = (\boldsymbol{\Sigma}^*)^{-1}(\boldsymbol{\mu}_r^* - \boldsymbol{\mu}_1^*)$, $\delta_r = \frac{1}{2}\boldsymbol{\beta}_r^T(\boldsymbol{\mu}_1^* + \boldsymbol{\mu}_r^*)$, for $r = 1 : R$.

Having the estimates $(\{\hat{w}_r^{(k)[T]}\}_{r=1}^R, \{\hat{\boldsymbol{\beta}}_r^{(k)[T]}\}_{r=2}^R, \{\hat{\boldsymbol{\mu}}_r^{(k)[T]}\}_{r=1}^R)$ from Algorithm 4, we can plug them into (10) to construct the clustering method, denoted by $\hat{\mathcal{C}}^{(k)[T]}(\mathbf{z})$. Equivalently,

$$\hat{\mathcal{C}}^{(k)[T]}(\mathbf{z}) = \arg \max_{r=1:R} \gamma_{\hat{\boldsymbol{\theta}}^{(k)[T]}}^{(r)}(\mathbf{z}). \quad (11)$$

3.1 Theory

We need the following assumption before stating the theory.

Assumption 2. Denote $\Delta_{rj}^{(k)} = \sqrt{(\boldsymbol{\mu}_r^{(k)*} - \boldsymbol{\mu}_j^{(k)*})^T (\boldsymbol{\Sigma}^{(k)*})^{-1} (\boldsymbol{\mu}_r^{(k)*} - \boldsymbol{\mu}_j^{(k)*})}$ for $k \in S$. Suppose the following conditions hold:

- (i) $n_S = \sum_{k \in S} n_k \geq C_1 |S| \max_{k=1:K} n_k$ with a constant $C_1 > 0$;
- (ii) $\min_{k \in S} n_k \geq C_2(p + \log K)$ with a large constant C_2 ;
- (iii) There exists a permutation $\pi : [R] \rightarrow [R]$ such that
 - (a) $\max_{k \in S} \{ [\max_{r=2:R} \|\hat{\boldsymbol{\beta}}_r^{(k)[0]} - (\boldsymbol{\Sigma}^{(k)*})^{-1}(\boldsymbol{\mu}_{\pi(r)}^{(k)*} - \boldsymbol{\mu}_{\pi(1)}^{(k)*})\|_2] \vee (\max_{r=1:R} \|\hat{\boldsymbol{\mu}}_r^{(k)[0]} - \boldsymbol{\mu}_{\pi(r)}^{(k)*}\|_2) \} \leq C_3 \min_{k \in S} \min_{r \neq j} \Delta_{rj}^{(k)}$, with a sufficiently small constant C_3 ;
 - (b) $\max_{k \in S} \max_{r=2:R} |\hat{w}_r^{(k)[0]} - w_{\pi(r)}^{(k)*}| \leq c_w/2$.
- (iv) $\min_{k \in S} \min_{r \neq j} \Delta_{rj}^{(k)} \geq C_4 > 0$ with a large constant C_4 ;

Remark 6. The above set of conditions are analogues of those in Assumption 1. We refer to Remark 1 for a detailed explanation of each condition.

We first present the result for parameter estimation. We adopt similar error metrics as the ones in (4) and (5). Specifically, denote the true parameter by $\{\bar{\boldsymbol{\theta}}^{(k)*}\}_{k \in S} = \{(\{w_r^{(k)*}\}_{r=2}^R, \{\boldsymbol{\mu}_r^{(k)*}\}_{r=1}^R, \boldsymbol{\Sigma}^{(k)*})\}_{k \in S}$ which belongs to the parameter space $\bar{\Theta}_S(h)$ in (8).

Algorithm 4: MTL-GMM (Multi-cluster)

Input: Initialization $\{(\{\widehat{w}_r^{(k)[0]}\}_{r=1}^R, \{\widehat{\beta}_r^{(k)[0]}\}_{r=2}^R, \{\widehat{\mu}_r^{(k)[0]}\}_{r=1}^R)\}_{k=1}^K$, round number T , initial penalty parameter $\lambda^{[0]}$, tuning parameters $C_\lambda > 0$, $\kappa \in (0, 1)$

- 1 $\widehat{\theta}^{(k)[0]} = (\{\widehat{w}_r^{(k)[0]}\}_{r=1}^R, \{\widehat{\beta}_r^{(k)[0]}\}_{r=2}^R, \{\widehat{\delta}_r^{(k)[0]}\}_{r=2}^R)$, where
 $\widehat{\delta}_r^{(k)[0]} = \frac{1}{2}(\widehat{\beta}_r^{(k)[0]})^T(\widehat{\mu}_1^{(k)[0]} + \widehat{\mu}_r^{(k)[0]})$, for $k = 1 : K$
- 2 **for** $t = 1$ **to** T **do**
- 3 $\lambda^{[t]} = \kappa \lambda^{[t-1]} + C_\lambda \sqrt{p + \log K}$ // Update the penalty parameter
- 4 **for** $k = 1$ **to** K **do** // Local update for each task
- 5 **for** $r = 1$ **to** R **do**
- 6 $\widehat{w}_r^{(k)[t]} = \frac{1}{n_k} \sum_{i=1}^{n_k} \gamma_{\widehat{\theta}^{(k)[t-1]}}^{(r)}(\mathbf{z}_i^{(k)})$
- 7 $\widehat{\mu}_r^{(k)[t]} = \frac{\sum_{i=1}^{n_k} \gamma_{\widehat{\theta}^{(k)[t-1]}}^{(r)}(\mathbf{z}_i^{(k)}) \mathbf{z}_i^{(k)}}{n_k \widehat{w}_r^{(k)[t]}}$
- 8 **end**
- 9 $\widehat{\Sigma}^{(k)[t]} = \frac{1}{n_k} \sum_{i=1}^{n_k} \sum_{r=1}^R \gamma_{\widehat{\theta}^{(k)[t-1]}}^{(r)}(\mathbf{z}_i^{(k)}) \cdot (\mathbf{z}_i^{(k)} - \widehat{\mu}_r^{(k)[t]})(\mathbf{z}_i^{(k)} - \widehat{\mu}_r^{(k)[t]})^T$
- 10 **end**
- 11 **for** $r = 2$ **to** R **do**
- 12 $\{\widehat{\beta}_r^{(k)[t]}\}_{k=1}^K, \overline{\beta}_r^{[t]} = \arg \min_{\beta^{(1)}, \dots, \beta^{(K)}, \overline{\beta}} \left\{ \sum_{k=1}^K n_k \left[\frac{1}{2} (\beta^{(k)})^T \widehat{\Sigma}^{(k)[t]} \beta^{(k)} - (\beta^{(k)})^T (\widehat{\mu}_r^{(k)[t]} - \widehat{\mu}_1^{(k)[t]}) \right] + \sum_{k=1}^K \sqrt{n_k} \lambda^{[t]} \cdot \|\beta^{(k)} - \overline{\beta}\|_2 \right\}$ // Aggregation
- 13 **end**
- 14 **for** $k = 1$ **to** K **do** // Local update for each task
- 15 **for** $r = 2$ **to** R **do**
- 16 $\widehat{\delta}_r^{(k)[t]} = \frac{1}{2}(\widehat{\beta}_r^{(k)[t]})^T(\widehat{\mu}_1^{(k)[t]} + \widehat{\mu}_r^{(k)[t]})$
- 17 **end**
- 18 Let $\widehat{\theta}^{(k)[t]} = (\{\widehat{w}_r^{(k)[t]}\}_{r=1}^R, \{\widehat{\beta}_r^{(k)[t]}\}_{r=2}^R, \{\widehat{\delta}_r^{(k)[t]}\}_{r=2}^R)$
- 19 **end**
- 20 **end**

Output: $\{(\widehat{\theta}^{(k)[T]}, \{\widehat{\mu}_r^{(k)[T]}\}_{r=1}^R, \widehat{\Sigma}^{(k)[T]})\}_{k=1}^K$ with $\widehat{\theta}^{(k)[T]} = (\{\widehat{w}_r^{(k)[T]}\}_{r=1}^R, \{\widehat{\beta}_r^{(k)[T]}\}_{r=2}^R, \{\widehat{\delta}_r^{(k)[T]}\}_{r=2}^R)$

For each $k \in S$, define the functional $\theta^{(k)*} = (\{w_r^{(k)*}\}_{r=2}^R, \{\beta_r^{(k)*}\}_{r=2}^R, \{\delta_r^{(k)*}\}_{r=2}^R)$, where $\beta_r^{(k)*} = (\Sigma^{(k)*})^{-1}(\mu_r^{(k)*} - \mu_1^{(k)*})$, $\delta_r^{(k)*} = \frac{1}{2}(\beta_r^{(k)*})^T(\mu_1^{(k)*} + \mu_r^{(k)*})$. For the estimators returned by Algorithm 4, we are interested in the error metrics ³:

$$d(\widehat{\theta}^{(k)[T]}, \theta^{(k)*}) = \min_{\pi: [R] \rightarrow [R]} \max_{r=2:R} \left\{ |\widehat{w}_r^{(k)[T]} - w_{\pi(r)}^{(k)*}| \vee \|\widehat{\beta}_r^{(k)[T]} - (\Sigma^{(k)*})^{-1}(\mu_{\pi(r)}^{(k)*} - \mu_{\pi(1)}^{(k)*})\|_2 \right\}$$

³Similar to the binary case, the minimum is taken due to the non-identifiability in multi-cluster GMMs.

$$\vee |\widehat{\delta}_r^{(k)[T]} - (\boldsymbol{\mu}_{\pi(r)}^{(k)*} + \boldsymbol{\mu}_{\pi(1)}^{(k)*})^T (\boldsymbol{\Sigma}^{(k)*})^{-1} (\boldsymbol{\mu}_{\pi(r)}^{(k)*} - \boldsymbol{\mu}_{\pi(1)}^{(k)*})/2| \Big\},$$

$$\left(\min_{\pi: [R] \rightarrow [R]} \max_{r=1:R} \|\widehat{\boldsymbol{\mu}}_r^{(k)[T]} - \boldsymbol{\mu}_{\pi(r)}^{(k)*}\|_2 \right) \vee \|\widehat{\boldsymbol{\Sigma}}^{(k)[T]} - \boldsymbol{\Sigma}^{(k)*}\|_2.$$

Theorem 7. (Upper bounds of the estimation error of GMM parameters for multi-cluster MTL-GMM) Suppose Assumption 2 holds, $|S| \geq s$, and $\epsilon = \frac{K-s}{K}$ is bounded by a small constant. Let $\lambda^{[0]} \geq C_1 \max_{k=1:K} \sqrt{n_k}$, $C_\lambda \geq C_1$ and $\kappa > C_2$ with some constants $C_1 > 0, C_2 \in (0, 1)$ ⁴ Then there exist constants $C_3, C_4, > 0$, such that for any $\{\bar{\boldsymbol{\theta}}^{(k)*}\}_{k \in S} \in \bar{\Theta}_S(h)$ and any probability measure \mathbb{Q}_S on $(\mathbb{R}^p)^{\otimes n_{sc}}$, with probability $1 - C_3(K^{-1} + e^{-C_4 p})$, the following hold for all $k \in S$:

$$d(\widehat{\boldsymbol{\theta}}^{(k)[T]}, \boldsymbol{\theta}^{(k)*}) \lesssim \sqrt{\frac{p}{n_S}} + \sqrt{\frac{\log K}{n_k}} + h \wedge \sqrt{\frac{p + \log K}{n_k}} + \epsilon \sqrt{\frac{p + \log K}{\max_{k=1:K} n_k}} + T^2(\kappa')^T,$$

$$\left(\min_{\pi: [R] \rightarrow [R]} \max_{r=1:R} \|\widehat{\boldsymbol{\mu}}_r^{(k)[T]} - \boldsymbol{\mu}_{\pi(r)}^{(k)*}\|_2 \right) \vee \|\widehat{\boldsymbol{\Sigma}}^{(k)[T]} - \boldsymbol{\Sigma}^{(k)*}\|_2 \lesssim \sqrt{\frac{p + \log K}{n_k}} + T^2(\kappa')^T,$$

where $\kappa' \in (0, 1)$ is some constant and $n_S = \sum_{k \in S} n_k$. When $T \geq C \log(\max_{k=1:K} n_k)$ with some large constant $C > 0$, the last term on the right-hand side will be dominated by other terms in both inequalities.

Recall the clustering method $\widehat{\mathcal{C}}^{(k)[T]}$ defined in (11). The next theorem obtains the upper bound of the excess mis-clustering error of $\widehat{\mathcal{C}}^{(k)[T]}$ for $k \in S$.

Theorem 8. (Upper bound of the excess mis-clustering error for multi-cluster MTL-GMM) Suppose the same conditions in Theorem 7 hold. Then there exist constants C_1 and C_2 such that for any $\{\bar{\boldsymbol{\theta}}^{(k)*}\}_{k \in S} \in \bar{\Theta}_S(h)$ and any probability measure \mathbb{Q}_S on $(\mathbb{R}^p)^{\otimes n_{sc}}$, with probability at least $1 - C_1(K^{-1} + e^{-C_2 p})$, the following holds for all $k \in S$:

$$\begin{aligned} R_{\bar{\boldsymbol{\theta}}^{(k)*}}(\widehat{\mathcal{C}}^{(k)[T]}) - R_{\bar{\boldsymbol{\theta}}^{(k)*}}(\mathcal{C}_{\bar{\boldsymbol{\theta}}^{(k)*}}) &\lesssim d^2(\widehat{\boldsymbol{\theta}}^{(k)[T]}, \boldsymbol{\theta}^{(k)*}) \cdot \log d^{-1}(\widehat{\boldsymbol{\theta}}^{(k)[T]}, \boldsymbol{\theta}^{(k)*}) \\ &\lesssim \left[\frac{p}{n_S} + \frac{\log K}{n_k} + h^2 \wedge \frac{p + \log K}{n_k} + \epsilon^2 \frac{p + \log K}{\max_{k=1:K} n_k} + T^4(\kappa')^{2T} \right] \cdot \log \left(\frac{n_S}{p} \wedge \frac{n_k}{\log K} \right), \end{aligned}$$

where $\kappa' \in (0, 1)$ is some constant. When $T \geq C \log(\max_{k=1:K} n_k)$ with a large constant $C > 0$, the term involving T on the right-hand side will be dominated by other terms.

Comparing the upper bounds in Theorems 7 and 8 with those in Theorems 1 and 2, the only difference is an extra logarithmic term $\log \left(\frac{n_S}{p} \wedge \frac{n_k}{\log K} \right)$ in Theorem 8, which we believe is a proof artifact. Similar logarithmic terms appear in other multi-cluster GMM literatures as well, see for example, Yan et al. (2017) and Zhao et al. (2020). To understand

⁴ C_1 and C_2 depend on the constants M , c_w , and c_Σ etc.

the upper bounds in Theorems 7 and 8, we can follow the discussion after Theorems 1 and 2. We do not repeat it here.

The following lower bounds together with the derived upper bounds will show that our method is (nearly) minimax optimal in a wide range of regimes.

Theorem 9. (Lower bounds of the estimation error of GMM parameters in multi-task learning) Suppose $\epsilon = \frac{K-s}{K}$ is bounded by a small constant. When there exists a subset S with $|S| \geq s$ such that $\min_{k \in S} n_k \geq C_1(p + \log K)$ with a large constant C_1 and $\min_{k \in S} \min_{r,j} \Delta_{rj}^{(k)}$ is larger than some constant C_2 , we have

$$\inf_{\{\hat{\boldsymbol{\theta}}^{(k)}\}_{k=1}^K} \sup_{S: |S| \geq s} \sup_{\substack{\{\bar{\boldsymbol{\theta}}^{(k)*}\}_{k \in S} \in \bar{\Theta}_S(h) \\ \mathbb{Q}_S}} \mathbb{P} \left(\bigcup_{k \in S} \left\{ d(\hat{\boldsymbol{\theta}}^{(k)}, \boldsymbol{\theta}^{(k)*}) \gtrsim \sqrt{\frac{p}{n_S}} + \sqrt{\frac{\log K}{n_k}} \right. \right. \\ \left. \left. + h \wedge \sqrt{\frac{p + \log K}{n_k}} + \frac{\epsilon}{\sqrt{\max_{k=1:K} n_k}} \right\} \right) \geq \frac{1}{10},$$

$$\inf_{\substack{\{\hat{\boldsymbol{\mu}}_r^{(k)}\}_{k=1:K, r=1:R} \\ \{\hat{\boldsymbol{\Sigma}}^{(k)}\}_{k=1}^K}} \sup_{S: |S| \geq s} \sup_{\substack{\{\bar{\boldsymbol{\theta}}^{(k)*}\}_{k \in S} \in \bar{\Theta}_S(h) \\ \mathbb{Q}_S}} \mathbb{P} \left(\bigcup_{k \in S} \left\{ \left(\min_{\pi: [R] \rightarrow [R]} \max_{r=1:R} \|\hat{\boldsymbol{\mu}}_r^{(k)} - \boldsymbol{\mu}_{\pi(r)}^{(k)*}\|_2 \vee \right. \right. \right. \\ \left. \left. \left. \|\hat{\boldsymbol{\Sigma}}^{(k)} - \boldsymbol{\Sigma}^{(k)*}\|_2 \gtrsim \sqrt{\frac{p + \log K}{n_k}} \right\} \right) \geq \frac{1}{10}.$$

Theorem 10. (Lower bound of the excess mis-clustering error in multi-task learning) Suppose the same conditions in Theorem 9 hold. Then

$$\inf_{\{\hat{\mathcal{C}}^{(k)}\}_{k=1}^K} \sup_{S: |S| \geq s} \sup_{\substack{\{\bar{\boldsymbol{\theta}}^{(k)*}\}_{k \in S} \in \bar{\Theta}_S(h) \\ \mathbb{Q}_S}} \mathbb{P} \left(\bigcup_{k \in S} \left\{ R_{\bar{\boldsymbol{\theta}}^{(k)*}}(\hat{\mathcal{C}}^{(k)}) - R_{\bar{\boldsymbol{\theta}}^{(k)*}}(\mathcal{C}_{\bar{\boldsymbol{\theta}}^{(k)*}}) \gtrsim \frac{p}{n_S} + \frac{\log K}{n_k} \right. \right. \\ \left. \left. + h^2 \wedge \frac{p + \log K}{n_k} + \frac{\epsilon^2}{\max_{k=1:K} n_k} \right\} \right) \geq \frac{1}{10}.$$

The lower bounds in Theorems 9 and 10 are the same as those in Theorems 3 and 4. Therefore, the remarks on the comparison of upper and lower bounds presented after Theorem 4 carry over to the multi-cluster setting (up to the logarithmic term from Theorem 8). We do not repeat the details here.

3.2 Alignment

The alignment issue discussed in Section 2.4 continues to exist in multi-cluster GMMs. In fact, it requires a more careful treatment because of an increased number of latent labels. Due to the space limit, we leave the details to Section B in Appendix.

4 Numerical Results

In this section, we present a thorough experimental study of our multi-task learning procedure MTL-GMM, i.e., Algorithms 1 and 4. The tuning parameter $\kappa \in (0, 1)$ is set as $1/3$, and the value of C_λ is determined by a 10-fold cross validation based on the log-likelihood of the final fitted model. The candidates of C_λ are chosen in a data-driven way, which is described in detail in Section C.1.5 of Appendix. All the experiments in this section are implemented in R. Function `Mcluster` in R package `mclust` is called to fit a single GMM. Due to page limit, we move one simulation example to Appendix C.1.3.

4.1 Simulations

4.1.1 Simulation 1

We first consider a binary GMM setting. There are $K = 10$ tasks of which each has sample size $n_k = 100$ and dimension $p = 15$. When $k \in S$, we generate each $w^{(k)*}$ from $\text{Unif}(0.1, 0.9)$ and $\mu_1^{(k)*}$ from $(2, 2, \mathbf{0}_{p-2})^T + h/2 \cdot (\Sigma^{(k)*})^{-1} \mathbf{u}$, where $\mathbf{u} \sim \text{Unif}(\{\mathbf{u} \in \mathbb{R}^p : \|\mathbf{u}\|_2 = 1\})$, $\Sigma^{(k)*} = (0.2^{|i-j|})_{p \times p}$, and let $\mu_2^{(k)*} = -\mu_1^{(k)*}$. When $k \notin S$, the distributions still follow GMM, but we generate each $w^{(k)*}$ from $\text{Unif}(0.2, 0.4)$ and $\mu_1^{(k)*}$ from $\text{Unif}(\{\mathbf{u} \in \mathbb{R}^p : \|\mathbf{u}\|_2 = 0.5\})$, and let $\mu_2^{(k)*} = -\mu_1^{(k)*}$, $\Sigma^{(k)*} = (0.5^{|i-j|})_{p \times p}$. In this setup, it is clear that h quantifies the similarity among tasks in S , and tasks in S^c have very distinct distributions and can be viewed as outlier tasks. For a given $\epsilon \in [0, 1)$, the outlier task index set S^c in each replication is uniformly sampled from all subsets of $1 : K$ with cardinality $K\epsilon$. We consider two cases:

- (i) No outlier tasks ($\epsilon = 0$), and h changes from 0 to 10 with increment 1;
- (ii) 2 outlier tasks ($\epsilon = 0.2$), and h changes from 0 to 10 with increment 1;

We fit Single-task-GMM on each separate task, Pooled-GMM on the merged data of all tasks, and our MTL-GMM in Algorithm 1 coupled with the exhaustive search for the alignment in Algorithm 2. The performance of all three methods are evaluated by the estimation error of $w^{(k)*}$, $\mu_1^{(k)*}$, $\mu_2^{(k)*}$, $\beta^{(k)*}$, $\delta^{(k)*}$, and $\Sigma^{(k)*}$, as well as the empirical mis-clustering error calculated on a test data set of size 500, for tasks in S . Due to page limit, we only present the estimation error of $\beta^{(k)*}$ and mis-clustering error here, and leave the others to Section C.1.1 in Appendix. These two errors are the maximum errors over tasks in S . For each setting, the simulation is replicated 200 times, and the average of the maximum errors together with the standard deviation are reported in Figure 2.

When there are no outlier tasks, it can be seen that MTL-GMM and Pooled-GMM are competitive when h is small (i.e. the tasks are similar), and they outperform Single-task-GMM. As h increases (i.e. tasks become more heterogenous), MTL-GMM starts to outperform Pooled-GMM by a large margin. Moreover, MTL-GMM is significantly better than Single-task-GMM in terms of both estimation and mis-clustering errors over a wide range of h . These comparisons demonstrate that MTL-GMM not only effectively utilizes




Figure 2: The performance of different methods in Simulation 1 under different outlier proportions. The upper panel shows the performance without outlier tasks ($\epsilon = 0$), and the lower panel shows the performance with 2 outlier tasks ($\epsilon = 0.2$). h changes from 0 to 10 with increment 1. Estimation error of $\{\beta^{(k)*}\}_{k \in S}$ stands for $\max_{k \in S} (\|\hat{\beta}^{(k)[T]} - \beta^{(k)*}\|_2 \wedge \|\hat{\beta}^{(k)[T]} + \beta^{(k)*}\|_2)$ and maximum mis-clustering error represents the maximum empirical mis-clustering error rate calculated on the test set of tasks in S .

the unknown similarity structure among tasks, but also adapts to it. When the outlier tasks exist, even when h is very small, MTL-GMM still performs better than Pooled-GMM, showing the robustness of MTL-GMM against a fraction of outlier tasks.

4.1.2 Simulation 2

The second simulation is a multi-cluster example, which is built based on Simulation 1. Consider a multi-task learning problem with $K = 10$ tasks, where each task has sample size $n_k = 100$ and dimension $p = 15$, and follows a GMM with $R = 4$ clusters. For all $k \in [K]$, we generate $(w_1^{(k)*}, \dots, w_R^{(k)*})$ independently from $\text{Dirichlet}(\alpha)$ with $\alpha = 5 \cdot \mathbf{1}_R$. When $k \in S$, we generate $\mu_r^{(k)*}$ from $(2 \cdot \mathbf{0}_{2r-2}, 2, 2, \mathbf{0}_{p-2r})^T + h/2 \cdot (\Sigma^{(k)*})^{-1} \mathbf{u}$, where $\mathbf{u} \sim \text{Unif}(\{\mathbf{u} \in \mathbb{R}^p : \|\mathbf{u}\|_2 = 1\})$, $\Sigma^{(k)*} = (0.2^{|i-j|})_{p \times p}$. When $k \notin S$, we generate each $w^{(k)*}$ from the same Dirichlet distribution and set $\Sigma^{(k)*} = (0.5^{|i-j|})_{p \times p}$ and $\mu_r^{(k)*}$ from $\text{Unif}(\{\mathbf{u} \in \mathbb{R}^p : \|\mathbf{u}\|_2 = 0.5\})$ for $r = 1 : R$. For a given $\epsilon \in [0, 1]$, in each replication the outlier task index set S^c is uniformly sampled from all subsets of $1 : K$ with cardinality $K\epsilon$. We consider two cases:

- (i) No outlier tasks ($\epsilon = 0$), and h changes from 0 to 10 with increment 1;
- (ii) 2 outlier tasks ($\epsilon = 0.2$), and h changes from 0 to 10 with increment 1.




Figure 3: The performance of different methods in Simulation 2 under different outlier proportions. The upper panel shows the performance without outlier tasks ($\epsilon = 0$), and the lower panel shows the performance with 2 outlier tasks ($\epsilon = 0.2$). h changes from 0 to 10 with increment 1. Estimation error of $\{\beta_r^{(k)*}\}_{r \in [R], k \in S}$ stands for $\max_{k \in S} \min_{\pi: [R] \rightarrow [R]} \max_{r=1:R} \|\hat{\beta}_r^{(k)[T]} - (\Sigma^{(k)*})^{-1}(\mu_{\pi(r)}^{(k)*} - \mu_{\pi(1)}^{(k)*})\|_2$ and maximum mis-clustering error represents the maximum empirical mis-clustering error rate calculated on the test set of tasks in S .

Algorithm 4 is run with the alignment Algorithm 8 (in Appendix B). The results of it and other benchmarks are reported in Figures 3. The main message is the same as in Simulation 1: Pooled-GMM is sensitive to outlier tasks and suffers from negative transfer when h is large, while MTL-GMM is robust to outliers and can adapt to the unknown similarity level h . Note that in this example, $\{\mu_r^{(k)*}\}_{k \in S}$ are similar and $\{\Sigma^{(k)*}\}_{k \in S}$ are the same, therefore running the EM algorithm by pooling all the data when h is small without outliers may be more effective than our MTL algorithm. This could explain why MTL-GMM performs slightly worse than Pooled-GMM in terms of maximum mis-clustering error when h is small and $\epsilon = 0$.

4.2 Real-data analysis

4.2.1 Human activity recognition

Human Activity Recognition (HAR) Using Smartphones Data Set contains the data collected from 30 volunteers when they performed six activities (walking, walking upstairs, walking downstairs, sitting, standing, and laying) wearing a smartphone (Anguita et al., 2013). Each observation has 561 time and frequency domain variables. Each volunteer can

be viewed as a task, and the sample size of each task varies from 281 to 409. The original data set is available at UCI Machine Learning Repository: <https://archive.ics.uci.edu/ml/datasets/human+activity+recognition+using+smartphones>.

Here we first focus on two activities: standing and laying, and perform clustering without the label information, to test our method in the binary case. This is a binary MTL clustering problem with 30 tasks. The sample size of each task varies from 95 to 179. For each task, in each replication, we use 90% of the samples as training data and hold 10% of the samples as test data.

We first run a principal component analysis (PCA) on the training data of each task and project both the training and test data onto the first 15 principal components. PCA has often been used for dimension reduction in pre-processing the HAR data set (Zeng et al., 2014; Walse et al., 2016; Aljarrah and Ali, 2019; Duan and Wang, 2022). We fit Single-task-GMM on each task separately, Pooled-GMM on merged data from 30 tasks, and our MTL-GMM with the greedy label swapping alignment algorithm. The performance of the three methods is evaluated by the mis-clustering error rate on the test data of all 30 tasks. The maximum and average mis-clustering errors among the 30 tasks are calculated in each replication. The mean and standard deviation of these two errors over 200 replications are reported on the left side of Table 1. To better display the clustering performance on each task, we further generate the box plot of mis-clustering errors of 30 tasks (averaged over 200 replications) for each method in the left plot of Figure 4. It is clear that MTL-GMM outperforms both Pooled-GMM and Single-task-GMM.

Method	Binary			Multi-cluster		
	Single-task	Pooled	MTL	Single-task	Pooled	MTL
Max. error	0.49 (0.02)	0.40 (0.12)	0.37 (0.09)	0.51 (0.04)	0.50 (0.04)	0.51 (0.04)
Avg. error	0.28 (0.02)	0.18 (0.18)	0.04 (0.04)	0.25 (0.01)	0.35 (0.03)	0.25 (0.01)

Table 1: Maximum and average mis-clustering errors and standard deviations (numbers in the parentheses) in binary and multi-cluster HAR data sets.

Next, we consider all six activities and compare the performance of the three approaches using the same sample-splitting strategy, to test our method in a multi-cluster scenario. Now the sample size of each task varies from 281 to 409. The maximum and average mis-clustering error rates and standard deviations over 200 replications are reported on the right side of Table 1. We can see that Pooled-GMM might suffer from negative transfer with a worse performance than the other two methods, while MTL-GMM and Single-task-GMM have similar performances. The right plot in Figure 4 reveal the same comparison results.

In summary, the HAR data set exhibits different levels of similarity in binary and multi-cluster cases: tasks in the binary data are sufficiently similar so that Pooled-GMM achieves a large margin of improvement over Single-task GMM, while tasks in the multi-cluster data become much more heterogeneous, resulting in the degraded performance of Pooled-




Figure 4: Box plots of mis-clustering errors of 30 tasks for each method for HAR data set. (Left: binary case; Right: multi-cluster case)

GMM compared to Single-task GMM. Nevertheless, our method MTL-GMM performs either competitively or better than the best of the two, regardless of the similarity level. These results lend further support to the effectiveness of our method.

4.2.2 Pen-based recognition of handwritten digits (PRHD)

Pen-based Recognition of Handwritten Digits (PRHD) data set contains 250 samples from each of 44 writers. Each of these writers was asked to write digits 0-9 on a pressure sensitive tablet with an integrated LCD display and a cordless stylus. The x and y tablet coordinates and pressure level values of the pen were recorded. After some transformations, each observation has 16 features. The data set and more information about it are available at UCI Machine Learning Repository: <https://archive.ics.uci.edu/dataset/81/pen+based+recognition+of+handwritten+digits>.

Similar to the previous real-data example, we first focus on a binary clustering problem by clustering observations of digits 8 and 9. The number of observations varies between 47 and 48 among the 44 tasks, showing that this is a more balanced data set with a smaller sample size (per dimension) than the HAR data. For each task, in each replication, we use 90% of the samples as training data and hold 10% of the samples as test data. The maximum and average mis-clustering error rates and standard deviations over 200 replications are reported on the left side of Table 2, and the box plots of mis-clustering errors of 44 tasks are shown in Figure 5. We can see that Pooled-GMM and MTL-GMM perform similarly and are much better than Single-task-GMM.

Next, we consider the observations of digits 5-9, i.e. a 5-class clustering problem. The maximum and average mis-clustering error rates and standard deviations over 200 replications are reported on the right side of Table 2, and the box plots of mis-clustering errors of 44 tasks are shown in Figure 5. In this multi-cluster case, MTL-GMM and Single-task-GMM have similar performance which is better than that of Pooled-GMM. Like in

Method	Binary			Multi-cluster		
	Single-task	Pooled	MTL	Single-task	Pooled	MTL
Max. error	0.32 (0.10)	0.03 (0.07)	0.03 (0.09)	0.26 (0.07)	0.37 (0.06)	0.27 (0.07)
Avg. error	0.02 (0.01)	0.00 (0.00)	0.00 (0.02)	0.03 (0.01)	0.12 (0.01)	0.03 (0.01)

Table 2: Maximum and average mis-clustering errors and standard deviations (numbers in the parentheses) in binary and multi-cluster PRHD data sets.




Figure 5: Box plots of mis-clustering errors of 44 tasks for each method for PRHD data set. (Left: binary case; Right: multi-cluster case)

the first real-data example, our method MTL-GMM adapts to the unknown similarity and is competitive with the best of the other two methods.

5 Discussions

We would like to highlight several interesting open problems for potential future work:

- What if only some clusters are similar among different tasks? This may be a more realistic situation in particular when there are more than 2 clusters in each task. Our current proposed algorithms may not work well because they do not take into account this extra layer of heterogeneity. Furthermore, in this situation, different tasks may have a different number of Gaussian clusters. Such a setting with various numbers of clusters has been considered in some literature on general unsupervised multi-task learning (Zhang and Zhang, 2011; Yang et al., 2014; Zhang et al., 2018). It would be of great interest to develop multi-task and transfer learning methods with provable guarantees for GMMs under these more complicated settings.
- How to accommodate heterogeneous covariance matrices for different Gaussian clus-

ters within each task? This is related to the quadratic discriminant analysis (QDA) in supervised learning where the Bayes classifier has a leading quadratic term. It may require more delicate analysis for methodological and theoretical development. Some recent QDA literature might be helpful (Li and Shao, 2015; Fan et al., 2015; Hao et al., 2018; Jiang et al., 2018).

- In this paper, we only considered the ℓ_2 -norm between the discriminant coefficients as the similarity metric. When dimension p is fixed, all norms in \mathbb{R}^p are equivalent. When p is diverging, other metrics can also be considered. For example, in the regression setting, under the context of MTL or TL, ℓ_1 -norm (Li et al., 2021b; Tian and Feng, 2022) and ℓ_0 -“norm” (Lounici et al., 2011; Liu et al., 2021; Maity et al., 2022) are also commonly used in literature. And some works describe the similarity through the latent structure (Du et al., 2021; Tripuraneni et al., 2021; Tian et al., 2023). By changing the penalty function used in M-step, we may accommodate our EM algorithm to these similarity metrics, which we leave for future study.
- In this paper, we have been focused on the pure unsupervised learning problem, where all the samples are unlabeled. It would be interesting to consider the semi-supervised learning setting, where labels in some tasks (or sources) are known. Li et al. (2022a) discusses a similar problem under the linear regression setting, but how the labeled data can help the estimation and clustering in the context of GMMs remains unknown.

References

- Aljarrah, A. A. and Ali, A. H. (2019). Human activity recognition using pca and bilstm recurrent neural networks. In *2019 2nd International Conference on Engineering Technology and its Applications (IICETA)*, pages 156–160. IEEE.
- Anderson, T. W. (1958). An introduction to multivariate statistical analysis: Wiley series in probability and mathematical statistics: Probability and mathematical statistics.
- Ando, R. K., Zhang, T., and Bartlett, P. (2005). A framework for learning predictive structures from multiple tasks and unlabeled data. *Journal of Machine Learning Research*, 6(11).
- Anguita, D., Ghio, A., Oneto, L., Parra Perez, X., and Reyes Ortiz, J. L. (2013). A public domain dataset for human activity recognition using smartphones. In *Proceedings of the 21th international European symposium on artificial neural networks, computational intelligence and machine learning*, pages 437–442.
- Argyriou, A., Evgeniou, T., and Pontil, M. (2008). Convex multi-task feature learning. *Machine learning*, 73(3):243–272.

- Audibert, J.-Y. and Tsybakov, A. B. (2007). Fast learning rates for plug-in classifiers. *Annals of statistics*, 35(2):608–633.
- Azizyan, M., Singh, A., and Wasserman, L. (2013). Minimax theory for high-dimensional gaussian mixtures with sparse mean separation. *Advances in Neural Information Processing Systems*, 26.
- Balakrishnan, S., Wainwright, M. J., and Yu, B. (2017). Statistical guarantees for the em algorithm: From population to sample-based analysis. *The Annals of Statistics*, 45(1):77–120.
- Bastani, H. (2021). Predicting with proxies: Transfer learning in high dimension. *Management Science*, 67(5):2964–2984.
- Baum, L. E., Petrie, T., Soules, G., and Weiss, N. (1970). A maximization technique occurring in the statistical analysis of probabilistic functions of markov chains. *The annals of mathematical statistics*, 41(1):164–171.
- Boucheron, S., Lugosi, G., and Massart, P. (2013). *Concentration inequalities: A nonasymptotic theory of independence*. Oxford university press.
- Cai, T. and Liu, W. (2011). A direct estimation approach to sparse linear discriminant analysis. *Journal of the American statistical association*, 106(496):1566–1577.
- Cai, T. T., Ma, J., and Zhang, L. (2019). Chime: Clustering of high-dimensional gaussian mixtures with em algorithm and its optimality. *The Annals of Statistics*, 47(3):1234–1267.
- Cai, T. T. and Zhou, H. H. (2012). Optimal rates of convergence for sparse covariance matrix estimation. *The Annals of Statistics*, 40(5):2389–2420.
- Carroll, R. J., Fan, J., Gijbels, I., and Wand, M. P. (1997). Generalized partially linear single-index models. *Journal of the American Statistical Association*, 92(438):477–489.
- Chattopadhyay, R., Sun, Q., Fan, W., Davidson, I., Panchanathan, S., and Ye, J. (2012). Multisource domain adaptation and its application to early detection of fatigue. *ACM Transactions on Knowledge Discovery from Data (TKDD)*, 6(4):1–26.
- Chen, D.-R. and Sun, T. (2006). Consistency of multiclass empirical risk minimization methods based on convex loss. *The Journal of Machine Learning Research*, 7:2435–2447.
- Chen, E. Y., Jordan, M. I., and Li, S. (2022). Transferred q-learning. *arXiv preprint arXiv:2202.04709*.
- Chen, M., Gao, C., and Ren, Z. (2018). Robust covariance and scatter matrix estimation under huber’s contamination model. *The Annals of Statistics*, 46(5):1932–1960.

- Dai, W., Yang, Q., Xue, G., and Yu, Y. (2007). Boosting for transfer learning. In *ACM International Conference Proceeding Series*, volume 227, page 193.
- Dai, W., Yang, Q., Xue, G.-R., and Yu, Y. (2008). Self-taught clustering. In *Proceedings of the 25th international conference on Machine learning*, pages 200–207.
- Dasgupta, S. and Schulman, L. (2013). A two-round variant of em for gaussian mixtures. *arXiv preprint arXiv:1301.3850*.
- Dempster, A. P., Laird, N. M., and Rubin, D. B. (1977). Maximum likelihood from incomplete data via the em algorithm. *Journal of the Royal Statistical Society: Series B (Methodological)*, 39(1):1–22.
- Dieuleveut, A., Fort, G., Moulines, E., and Robin, G. (2021). Federated-em with heterogeneity mitigation and variance reduction. *Advances in Neural Information Processing Systems*, 34:29553–29566.
- Du, S. S., Hu, W., Kakade, S. M., Lee, J. D., and Lei, Q. (2021). Few-shot learning via learning the representation, provably. In *9th International Conference on Learning Representations, ICLR 2021*.
- Duan, Y. and Wang, K. (2022). Adaptive and robust multi-task learning. *arXiv preprint arXiv:2202.05250*.
- Efron, B. (1975). The efficiency of logistic regression compared to normal discriminant analysis. *Journal of the American Statistical Association*, 70(352):892–898.
- Evgeniou, T. and Pontil, M. (2004). Regularized multi-task learning. In *Proceedings of the tenth ACM SIGKDD international conference on Knowledge discovery and data mining*, pages 109–117.
- Fan, J., Feng, Y., and Tong, X. (2012). A road to classification in high dimensional space: the regularized optimal affine discriminant. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 74(4):745–771.
- Fan, Y., Kong, Y., Li, D., and Zheng, Z. (2015). Innovated interaction screening for high-dimensional nonlinear classification. *The Annals of Statistics*, 43(3):1243–1272.
- Forgy, E. W. (1965). Cluster analysis of multivariate data: efficiency versus interpretability of classifications. *biometrics*, 21:768–769.
- Ge, R., Huang, Q., and Kakade, S. M. (2015). Learning mixtures of gaussians in high dimensions. In *Proceedings of the forty-seventh annual ACM symposium on Theory of computing*, pages 761–770.
- Gu, Q., Li, Z., and Han, J. (2011). Learning a kernel for multi-task clustering. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 25, pages 368–373.

- Hao, N., Feng, Y., and Zhang, H. H. (2018). Model selection for high-dimensional quadratic regression via regularization. *Journal of the American Statistical Association*, 113(522):615–625.
- Hartley, H. O. (1958). Maximum likelihood estimation from incomplete data. *Biometrics*, 14(2):174–194.
- Hasselblad, V. (1966). Estimation of parameters for a mixture of normal distributions. *Technometrics*, 8(3):431–444.
- Hastie, T., Tibshirani, R., Friedman, J. H., and Friedman, J. H. (2009). *The elements of statistical learning: data mining, inference, and prediction*, volume 2. Springer.
- Hsu, D. and Kakade, S. M. (2013). Learning mixtures of spherical gaussians: moment methods and spectral decompositions. In *Proceedings of the 4th conference on Innovations in Theoretical Computer Science*, pages 11–20.
- Huber, P. J. (1964). Robust estimation of a location parameter. *The Annals of Mathematical Statistics*, pages 73–101.
- Jain, A. K. and Dubes, R. C. (1988). *Algorithms for clustering data*. Prentice-Hall, Inc.
- Jalali, A., Sanghavi, S., Ruan, C., and Ravikumar, P. (2010). A dirty model for multi-task learning. *Advances in neural information processing systems*, 23.
- Jiang, B., Wang, X., and Leng, C. (2018). A direct approach for sparse quadratic discriminant analysis. *The Journal of Machine Learning Research*, 19(1):1098–1134.
- Jin, J., Ke, Z. T., and Wang, W. (2017). Phase transitions for high dimensional clustering and related problems. *The Annals of Statistics*, 45(5):2151–2189.
- Kalai, A. T., Moitra, A., and Valiant, G. (2010). Efficiently learning mixtures of two gaussians. In *Proceedings of the forty-second ACM symposium on Theory of computing*, pages 553–562.
- Konstantinov, N., Frantar, E., Alistarh, D., and Lampert, C. (2020). On the sample complexity of adversarial multi-source pac learning. In *International Conference on Machine Learning*, pages 5416–5425. PMLR.
- Kwon, J. and Caramanis, C. (2020). The em algorithm gives sample-optimality for learning mixtures of well-separated gaussians. In *Conference on Learning Theory*, pages 2425–2487. PMLR.
- Lawrence, N. D. and Platt, J. C. (2004). Learning to learn with the informative vector machine. In *Proceedings of the twenty-first international conference on Machine learning*, page 65.

- Lee, K., Guillemot, L., Yue, Y., Kramer, M., and Champion, D. (2012). Application of the gaussian mixture model in pulsar astronomy-pulsar classification and candidates ranking for the fermi 2fgl catalogue. *Monthly Notices of the Royal Astronomical Society*, 424(4):2832–2840.
- Li, Q. and Shao, J. (2015). Sparse quadratic discriminant analysis for high dimensional data. *Statistica Sinica*, pages 457–473.
- Li, R. and Liang, H. (2008). Variable selection in semiparametric regression modeling. *The Annals of Statistics*, 36(1):261–286.
- Li, S., Cai, T., and Duan, R. (2021a). Targeting underrepresented populations in precision medicine: A federated transfer learning approach. *arXiv preprint arXiv:2108.12112*.
- Li, S., Cai, T. T., and Li, H. (2021b). Transfer learning for high-dimensional linear regression: Prediction, estimation and minimax optimality. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, pages 1–25.
- Li, S., Cai, T. T., and Li, H. (2022a). Estimation and inference with proxy data and its genetic applications. *arXiv preprint arXiv:2201.03727*.
- Li, S., Cai, T. T., and Li, H. (2022b). Transfer learning in large-scale gaussian graphical models with false discovery rate control. *Journal of the American Statistical Association*, pages 1–13.
- Li, W., Duan, L., Xu, D., and Tsang, I. W. (2013). Learning with augmented features for supervised and semi-supervised heterogeneous domain adaptation. *IEEE transactions on pattern analysis and machine intelligence*, 36(6):1134–1148.
- Lin, H. and Reimherr, M. (2022). On transfer learning in functional linear regression. *arXiv preprint arXiv:2206.04277*.
- Liu, M., Xia, Y., Cho, K., and Cai, T. (2021). Integrative high dimensional multiple testing with heterogeneity under data sharing constraints. *The Journal of Machine Learning Research*, 22(1):5607–5632.
- Lounici, K., Pontil, M., van de Geer, S., and Tsybakov, A. B. (2011). Oracle inequalities and optimal inference under group sparsity. *The Annals of Statistics*, pages 2164–2204.
- Mai, Q., Yang, Y., and Zou, H. (2019). Multiclass sparse discriminant analysis. *Statistica Sinica*, 29(1):97–111.
- Mai, Q., Zou, H., and Yuan, M. (2012). A direct approach to sparse discriminant analysis in ultra-high dimensions. *Biometrika*, 99(1):29–42.
- Maity, S., Sun, Y., and Banerjee, M. (2022). Meta-analysis of heterogeneous data: integrative sparse regression in high-dimensions. *The Journal of Machine Learning Research*, 23(1):8975–9024.

- Marfoq, O., Neglia, G., Bellet, A., Kameni, L., and Vidal, R. (2021). Federated multi-task learning under a mixture of distributions. *Advances in Neural Information Processing Systems*, 34:15434–15447.
- Maurer, A. (2016). A vector-contraction inequality for rademacher complexities. In *Algorithmic Learning Theory: 27th International Conference, ALT 2016, Bari, Italy, October 19-21, 2016, Proceedings 27*, pages 3–17. Springer.
- Maurer, A. and Pontil, M. (2021). Concentration inequalities under sub-gaussian and sub-exponential conditions. *Advances in Neural Information Processing Systems*, 34:7588–7597.
- McLachlan, G. J. and Krishnan, T. (2007). *The EM algorithm and extensions*. John Wiley & Sons.
- Meng, X.-L. and Rubin, D. B. (1994). On the global and componentwise rates of convergence of the em algorithm. *Linear Algebra and its Applications*, 199:413–425.
- Mihalkova, L., Huynh, T., and Mooney, R. J. (2007). Mapping and revising markov logic networks for transfer learning. In *Proceedings of the 22nd national conference on Artificial intelligence-Volume 1*, pages 608–614.
- Murtagh, F. and Contreras, P. (2012). Algorithms for hierarchical clustering: an overview. *Wiley Interdisciplinary Reviews: Data Mining and Knowledge Discovery*, 2(1):86–97.
- Ng, A., Jordan, M., and Weiss, Y. (2001). On spectral clustering: Analysis and an algorithm. *Advances in neural information processing systems*, 14.
- Obozinski, G., Taskar, B., and Jordan, M. (2006). Multi-task feature selection. *Statistics Department, UC Berkeley, Tech. Rep*, 2(2.2):2.
- Pan, S. J. and Yang, Q. (2009). A survey on transfer learning. *IEEE Transactions on knowledge and data engineering*, 22(10):1345–1359.
- Pearson, K. (1894). Contributions to the mathematical theory of evolution. *Philosophical Transactions of the Royal Society of London. A*, 185:71–110.
- Redner, R. A. and Walker, H. F. (1984). Mixture densities, maximum likelihood and the em algorithm. *SIAM review*, 26(2):195–239.
- Scott, A. J. and Symons, M. J. (1971). Clustering methods based on likelihood ratio criteria. *Biometrics*, pages 387–397.
- Sundberg, R. (1974). Maximum likelihood theory for incomplete data from an exponential family. *Scandinavian Journal of Statistics*, pages 49–58.
- Thrun, S. and O’Sullivan, J. (1996). Discovering structure in multiple learning tasks: The tc algorithm. In *ICML*, volume 96, pages 489–497.

- Tian, Y. and Feng, Y. (2022). Transfer learning under high-dimensional generalized linear models. *Journal of the American Statistical Association*, (just-accepted):1–30.
- Tian, Y., Gu, Y., and Feng, Y. (2023). Learning from similar linear representations: Adaptivity, minimaxity, and robustness. *arXiv preprint arXiv:2303.17765*.
- Tripuraneni, N., Jin, C., and Jordan, M. (2021). Provable meta-learning of linear representations. In *International Conference on Machine Learning*, pages 10434–10443. PMLR.
- Tsybakov, A. B. (2009). *Introduction to nonparametric estimation*. Springer, New York.
- Tukey, J. W. (1975). Mathematics and the picturing of data. In *Proceedings of the International Congress of Mathematicians, Vancouver, 1975*, volume 2, pages 523–531.
- Vempala, S. and Wang, G. (2004). A spectral algorithm for learning mixture models. *Journal of Computer and System Sciences*, 68(4):841–860.
- Vigogna, S., Meanti, G., De Vito, E., and Rosasco, L. (2022). Multiclass learning with margin: exponential rates with no bias-variance trade-off. In *International Conference on Machine Learning*, pages 22260–22269. PMLR.
- Wainwright, M. J. (2019). *High-dimensional statistics: A non-asymptotic viewpoint*, volume 48. Cambridge University Press.
- Walse, K. H., Dharaskar, R. V., and Thakare, V. M. (2016). Pca based optimal ann classifiers for human activity recognition using mobile sensors data. In *Proceedings of First International Conference on Information and Communication Technology for Intelligent Systems: Volume 1*, pages 429–436. Springer.
- Wang, R., Zhou, J., Jiang, H., Han, S., Wang, L., Wang, D., and Chen, Y. (2021). A general transfer learning-based gaussian mixture model for clustering. *International Journal of Fuzzy Systems*, 23(3):776–793.
- Wang, Z., Gu, Q., Ning, Y., and Liu, H. (2014). High dimensional expectation-maximization algorithm: Statistical optimization and asymptotic normality. *arXiv preprint arXiv:1412.8729*.
- Wang, Z., Song, Y., and Zhang, C. (2008). Transferred dimensionality reduction. In *Joint European conference on machine learning and knowledge discovery in databases*, pages 550–565. Springer.
- Witten, D. M. and Tibshirani, R. (2011). Penalized classification using fisher’s linear discriminant. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 73(5):753–772.
- Wu, C. J. (1983). On the convergence properties of the em algorithm. *The Annals of statistics*, pages 95–103.

- Xu, J., Hsu, D. J., and Maleki, A. (2016). Global analysis of expectation maximization for mixtures of two gaussians. *Advances in Neural Information Processing Systems*, 29.
- Xu, K. and Bastani, H. (2021). Learning across bandits in high dimension via robust statistics. *arXiv preprint arXiv:2112.14233*.
- Yan, B., Yin, M., and Sarkar, P. (2017). Convergence of gradient em on multi-component mixture of gaussians. *Advances in Neural Information Processing Systems*, 30.
- Yang, M.-H. and Ahuja, N. (1998). Gaussian mixture model for human skin color and its applications in image and video databases. In *Storage and retrieval for image and video databases VII*, volume 3656, pages 458–466. SPIE.
- Yang, Y., Ma, Z., Yang, Y., Nie, F., and Shen, H. T. (2014). Multitask spectral clustering by exploring intertask correlation. *IEEE transactions on cybernetics*, 45(5):1083–1094.
- Zeng, M., Nguyen, L. T., Yu, B., Mengshoel, O. J., Zhu, J., Wu, P., and Zhang, J. (2014). Convolutional neural networks for human activity recognition using mobile sensors. In *6th international conference on mobile computing, applications and services*, pages 197–205. IEEE.
- Zhang, J. and Zhang, C. (2011). Multitask bregman clustering. *Neurocomputing*, 74(10):1720–1734.
- Zhang, X., Blanchet, J., Ghosh, S., and Squillante, M. S. (2022). A class of geometric structures in transfer learning: Minimax bounds and optimality. In *International Conference on Artificial Intelligence and Statistics*, pages 3794–3820. PMLR.
- Zhang, X., Zhang, X., and Liu, H. (2015). Smart multitask bregman clustering and multi-task kernel clustering. *ACM Transactions on Knowledge Discovery from Data (TKDD)*, 10(1):1–29.
- Zhang, X., Zhang, X., Liu, H., and Luo, J. (2018). Multi-task clustering with model relation learning. In *Proceedings of the 27th International Joint Conference on Artificial Intelligence*, pages 3132–3140.
- Zhang, Y. and Yang, Q. (2021). A survey on multi-task learning. *IEEE Transactions on Knowledge and Data Engineering*.
- Zhao, R., Li, Y., and Sun, Y. (2020). Statistical convergence of the em algorithm on gaussian mixture models. *Electronic Journal of Statistics*, 14:632–660.
- Zou, H. (2006). The adaptive lasso and its oracle properties. *Journal of the American statistical association*, 101(476):1418–1429.
- Zuo, H., Lu, J., Zhang, G., and Liu, F. (2018). Fuzzy transfer learning using an infinite gaussian mixture model and active learning. *IEEE Transactions on Fuzzy Systems*, 27(2):291–303.

Appendix

Contents

A	Transfer Learning	40
A.1	Problem setting	40
A.2	Method	41
A.3	Theory	43
A.4	Label alignment	46
B	Alignment for Multi-cluster GMMs in MTL	47
C	Additional Numerical Studies	50
C.1	Simulations	50
C.1.1	Simulation 1 of MTL	50
C.1.2	Simulation 2 of MTL	50
C.1.3	Simulation 3 of MTL	50
C.1.4	Simulation of TL	57
C.1.5	Tuning parameters C_λ and C_{λ_0} in Algorithms 1, 4, and 5	57
C.1.6	Tuning parameter κ and κ_0 in Algorithms 1, 4, and 5	60
D	Technical Lemmas	60
D.1	General lemmas	60
E	Proofs of Theorems in Main Text	64
E.1	Proofs of general lemmas	64
E.1.1	Proof of Lemma 2	64
E.1.2	Proof of Lemma 4	64
E.1.3	Proof of Lemma 5	65
E.1.4	Proof of Lemma 6	65
E.2	Proof of Theorem 1	65
E.2.1	Lemmas	66
E.2.2	Main proof of of Theorem 1	68
E.2.3	Proofs of lemmas	81
E.3	Proof of Theorem 3	88
E.3.1	Lemmas	88
E.3.2	Main proof of Theorem 3	89
E.3.3	Proofs of lemmas	90
E.4	Proof of Theorem 2	98
E.5	Proof of Theorem 4	100
E.5.1	Lemmas	100
E.5.2	Main proof of Theorem 4	101

E.5.3	Proof of lemmas	101
E.6	Proof of Theorem 5	107
E.7	Proof of Theorem 6	108
E.8	Proof of Theorem 11	110
E.8.1	Lemmas	110
E.8.2	Main proof of Theorem 11	111
E.8.3	Proof of lemmas	113
E.9	Proof of Theorem 13	114
E.9.1	Lemmas	114
E.9.2	Main proof of Theorem 13	115
E.9.3	Proof of lemmas	115
E.10	Proof of Theorem 12	118
E.11	Proof of Theorem 14	118
E.11.1	Lemmas	118
E.11.2	Main proof of Theorem 14	118
E.11.3	Proof of lemmas	119
E.12	Proof of Theorem 15	122
E.13	Proof of Theorem 7	123
E.13.1	Lemmas	123
E.13.2	Main proof of Theorem 7	124
E.13.3	Proof of lemmas	127
E.14	Proof of Theorem 8	132
E.15	Proof of Theorem 9	134
E.16	Proof of Theorem 10	135
E.16.1	Lemmas	135
E.16.2	Main proof of Theorem 10	135
E.16.3	Proof of lemmas	135
E.17	Proof of Theorem 16	137
E.18	Proof of Theorem 17	143

A Transfer Learning

A.1 Problem setting

In the main paper, we discussed GMMs under the context of *multi-task learning*, where the goal is to learn all tasks jointly by utilizing the potential similarities shared by different tasks. In this section, we will study binary GMMs in the *transfer learning* context where the focus is on the improvement of learning in one target task through the transfer of knowledge from related source tasks. Multi-cluster results can be obtained similarly as in the MTL case, and we omit the details given the extensive length of the paper.

Suppose that there are $(K + 1)$ tasks in total, where the first task is called the target and the K remaining ones are called K sources. Like in the multi-task learning, we assume

that there exists an unknown subset $S \subseteq 1 : K$, such that samples from sources in S follow an independent GMM, while samples from sources outside S can be arbitrarily distributed. This means,

$$y_i^{(k)} = \begin{cases} 1, & \text{with probability } 1 - w^{(k)*}; \\ 2, & \text{with probability } w^{(k)*}; \end{cases}$$

$$\mathbf{z}_i^{(k)} | y_i^{(k)} = j \sim \mathcal{N}(\boldsymbol{\mu}_j^{(k)*}, \boldsymbol{\Sigma}^{(k)*}), \quad j = 1, 2,$$

for all $k \in S$, $i = 1 : n_k$, and

$$\{\mathbf{z}_i^{(k)}\}_{i,k \in S^c} \sim \mathbb{Q}_S,$$

where \mathbb{Q}_S is some probability measure on $(\mathbb{R}^p)^{\otimes n_{S^c}}$ and $n_{S^c} = \sum_{k \in S^c} n_k$.

For the target task, we observe sample $\{\mathbf{z}_i^{(0)}\}_{i=1}^{n_0}$ independently sampled from the following GMM:

$$y_i^{(0)} = \begin{cases} 1, & \text{with probability } 1 - w^{(0)*}; \\ 2, & \text{with probability } w^{(0)*}; \end{cases}$$

$$\mathbf{z}_i^{(0)} | y_i^{(0)} = j \sim \mathcal{N}(\boldsymbol{\mu}_j^{(0)*}, \boldsymbol{\Sigma}^{(0)*}), \quad j = 1, 2.$$

The objective of transfer learning is to use source data to help improve GMM learning in the *target* task. As for multi-task learning, we measure the learning performance by both parameter estimation error and the excess mis-clustering error, but only on the *target* GMM. Toward this end, we define the joint parameter space for GMM parameters of the target and sources in S :

$$\overline{\Theta}'_S(h) = \left\{ \{\overline{\boldsymbol{\theta}}^{(k)}\}_{k \in \{0\} \cup S} = \{(w^{(k)}, \boldsymbol{\mu}_1^{(k)}, \boldsymbol{\mu}_2^{(k)}, \boldsymbol{\Sigma}^{(k)})\}_{k \in \{0\} \cup S} : \overline{\boldsymbol{\theta}}^{(k)} \in \overline{\Theta}, \max_{k \in S} \|\boldsymbol{\beta}^{(k)} - \boldsymbol{\beta}^{(0)}\|_2 \leq h \right\}, \quad (\text{A.12})$$

where $\overline{\Theta}$ is the single GMM parameter space introduced in (2), and $\boldsymbol{\beta}^{(k)} = (\boldsymbol{\Sigma}^{(k)})^{-1}(\boldsymbol{\mu}_2^{(k)} - \boldsymbol{\mu}_1^{(k)})$, $k \in \{0\} \cup S$. Comparing $\overline{\Theta}'_S(h)$ with the parameter space $\overline{\Theta}_S(h)$ from multi-task learning in (3), here the target discriminant coefficient $\boldsymbol{\beta}^{(0)}$ serves as the “center” of discriminant coefficients of sources in S . The quantity h characterizes the closeness between sources in S and the target.

A.2 Method

Like the MTL-GMM procedure developed in Section 2.2, we combine the EM algorithm and the penalization framework to develop a variant of EM algorithm for transfer learning. The key idea is to first apply MTL-GMM to all the sources to obtain estimates of discriminant coefficient “center” as good summary statistics of the K source data sets, and then shrink target discriminant coefficient towards those center estimates in the EM iterations to explore the relatedness between sources and the target. See Section 3.3 of Duan and

Wang (2022) for more general discussions on this idea. Our proposed transfer learning procedure TL-GMM is summarized in Algorithm 5.

While the steps of TL-GMM look very similar to those of MTL-GMM, there exist two major differences between them. First, for each optimization problem in TL-GMM, the first part of the objective function only involves the target data $\{\mathbf{z}_i^{(0)}\}_{i=1}^{n_0}$, while in MTL-GMM it is a weighted average of all tasks. Second, in TL-GMM, the penalty is imposed on the distance between a discriminant coefficient estimator and a given center estimator produced by MTL-GMM from the source data. In contrast, the center is estimated simultaneously with other parameters through the penalization in MTL-GMM. In light of existing transfer learning approaches in literature, TL-GMM can be considered as the “debiasing” step described in [Li et al. \(2021b\)](#) and [Tian and Feng \(2022\)](#), which corrects potential bias of the center estimate using the target data.

The tuning parameters $\{\lambda_0^{[t]}\}_{t=1}^{T_0}$ in Algorithm 5 control the amount of knowledge to be transferred from sources. Setting tuning parameters large enough pushes parameter estimates for target task to be exactly equal to the center learned from sources, while letting them be zero makes TL-GMM reduce to the standard EM algorithm on the target data.

Algorithm 5: TL-GMM

Input: Initialization $\widehat{\boldsymbol{\theta}}^{(0)[0]} = (\widehat{w}^{(0)[0]}, \widehat{\boldsymbol{\beta}}^{(0)[0]}, \widehat{\delta}^{(0)[0]})$, output $\overline{\boldsymbol{\beta}}^{[T]}$ from Algorithm 1, round number T_0 , initial penalty parameter $\lambda_0^{[0]}$, tuning parameters $C_{\lambda_0} > 0$ and $\kappa_0 \in (0, 1)$

```

1 for  $t = 1$  to  $T_0$  do
2    $\lambda_0^{[t]} = \kappa_0 \lambda_0^{[t-1]} + C_{\lambda_0} \sqrt{p + \log K}$  // Update the penalty parameter
3    $\hat{w}^{(0)[t]} = \frac{1}{n_0} \sum_{i=1}^{n_0} \gamma_{\hat{\theta}^{(0)[t-1]}}(\mathbf{z}_i^{(0)})$ 
4    $\hat{\mu}_1^{(0)[t]} = \frac{\sum_{i=1}^{n_0} [1 - \gamma_{\hat{\theta}^{(0)[t-1]}}(\mathbf{z}_i^{(0)})] \mathbf{z}_i^{(0)}}{n_0(1 - \hat{w}^{(0)[t]})}$ ,  $\hat{\mu}_2^{(0)[t]} = \frac{\sum_{i=1}^{n_0} \gamma_{\hat{\theta}^{(0)[t-1]}}(\mathbf{z}_i^{(0)}) \mathbf{z}_i^{(0)}}{n_0 \hat{w}^{(0)[t]}}$ 
5    $\hat{\Sigma}^{(0)[t]} = \frac{1}{n_0} \sum_{i=1}^{n_0} \left\{ [1 - \gamma_{\hat{\theta}^{(0)[t-1]}}(\mathbf{z}_i^{(0)})] \cdot (\mathbf{z}_i^{(0)} - \hat{\mu}_1^{(0)[t]})(\mathbf{z}_i^{(0)} - \hat{\mu}_1^{(0)[t]})^T \right.$ 
6      $\left. + \gamma_{\hat{\theta}^{(0)[t-1]}}(\mathbf{z}_i^{(0)}) \cdot (\mathbf{z}_i^{(0)} - \hat{\mu}_2^{(0)[t]})(\mathbf{z}_i^{(0)} - \hat{\mu}_2^{(0)[t]})^T \right\}$ 
7    $\hat{\beta}^{(0)[t]} = \arg \min_{\beta^{(0)}} \left\{ \left[ \frac{1}{2} (\beta^{(0)})^T \hat{\Sigma}^{(0)[t]} \beta^{(0)} - (\beta^{(0)})^T (\hat{\mu}_2^{(0)[t]} - \hat{\mu}_1^{(0)[t]}) \right] + \frac{\lambda_0^{[t]}}{\sqrt{n_0}} \|\beta^{(0)} - \bar{\beta}^{[T]}\|_2 \right\}$ 
8    $\hat{\delta}^{(0)[t]} = \frac{1}{2} (\hat{\beta}^{(0)[t]})^T (\hat{\mu}_1^{(0)[t]} + \hat{\mu}_2^{(0)[t]})$ 
9   Let  $\hat{\theta}^{(0)[t]} = (\hat{w}^{(0)[t]}, \hat{\beta}^{(0)[t]}, \hat{\delta}^{(0)[t]})$ 
10 end

```

Output: $(\hat{\theta}^{(0)[T_0]}, \hat{\mu}_1^{(0)[T_0]}, \hat{\mu}_2^{(0)[T_0]}, \hat{\Sigma}^{(0)[T_0]})$ with $\hat{\theta}^{(0)[T_0]} = (\hat{w}^{(0)[T_0]}, \hat{\beta}^{(0)[T_0]}, \hat{\delta}^{(0)[T_0]})$

A.3 Theory

In this section, we will establish the upper and lower bounds for the GMM parameter estimation error and the excess mis-clustering error on the *target* task. First, we impose the following assumption set.

Assumption 3. Denote $\Delta^{(0)} = \sqrt{(\boldsymbol{\mu}_1^{(0)*} - \boldsymbol{\mu}_2^{(0)*})^T (\boldsymbol{\Sigma}^{(0)*})^{-1} (\boldsymbol{\mu}_1^{(0)*} - \boldsymbol{\mu}_2^{(0)*})}$. Assume the following conditions hold:

$$(i) \quad C_1 \left[\frac{\log K}{p} + \epsilon^2 \left(1 + \frac{\log K}{p} \right) \right] \leq \frac{\max_{k \in S} n_k}{n_0} \leq C_2 \left(1 + \frac{\log K}{p} \right) \text{ with constants } C_1 \text{ and } C_2, \text{ where } \epsilon = \frac{K-s}{K}.$$

$$(ii) \quad n_0 \geq C_3 p \text{ with a large constant } C_3;$$

(iii) Either of the following two conditions holds with a sufficiently small constant C_4 :

$$(a) \quad \|\widehat{\boldsymbol{\beta}}^{(0)[0]} - \boldsymbol{\beta}^{(0)*}\|_2 \vee |\widehat{\delta}^{(0)[0]} - \delta^{(0)*}| \leq C_4 \Delta^{(0)}, \quad |\widehat{w}^{(0)[0]} - w^{(0)*}| \leq c_w/2;$$

$$(b) \quad \|\widehat{\boldsymbol{\beta}}^{(0)[0]} + \boldsymbol{\beta}^{(0)*}\|_2 \vee |\widehat{\delta}^{(0)[0]} + \delta^{(0)*}| \leq C_4 \Delta^{(0)}, \quad |1 - \widehat{w}^{(0)[0]} - w^{(0)*}| \leq c_w/2.$$

$$(iv) \quad \Delta^{(0)} \geq C_5 > 0 \text{ with a large constant } C_5;$$

Remark 7. Condition (i) requires the target sample size not to be much smaller than the maximum source sample size, which appears due to technical reasons in the proof. Conditions (ii)-(iv) can be seen as the counterpart of Conditions (ii)-(iv) in Assumption 1 for the target GMM.

We are in the position to present the upper bounds of the estimation error of GMM parameters for TL-GMM.

Theorem 11. (Upper bounds of the estimation error of GMM parameters for TL-GMM) Suppose the conditions in Theorem 1 and Assumption 3 hold. Let $\lambda_0^{[0]} \geq C_1 \max_{k=1:K} \sqrt{n_k}$, $C_{\lambda_0} \geq C_1$, $\kappa_0 > C_2$ with some specific constants $C_1 > 0, C_2 \in (0, 1)$. Then there exist constants C_3 and C_4 , such that for any $\{\boldsymbol{\theta}^{(k)*}\}_{k \in \{0\} \cup S} \in \overline{\Theta}'_S(h)$ and any probability measure \mathbb{Q}_S on $(\mathbb{R}^p)^{\otimes n_{Sc}}$, we have

$$\begin{aligned} d(\widehat{\boldsymbol{\theta}}^{(0)[T_0]}, \boldsymbol{\theta}^{(0)*}) &\lesssim \sqrt{\frac{p}{n_S + n_0}} + \sqrt{\frac{1}{n_0}} + h \wedge \sqrt{\frac{p}{n_0}} + \left(\epsilon \sqrt{\frac{p + \log K}{\max_{k=1:K} n_k}} \right) \wedge \sqrt{\frac{p}{n_0}} \\ &\quad + \sqrt{\frac{\log K}{\max_{k=1:K} n_k}} + T_0(\kappa'_0)^{T_0}, \end{aligned}$$

$$\min_{\pi: [2] \rightarrow [2]} \max_{r=1:2} \|\widehat{\boldsymbol{\mu}}_{\pi(r)}^{(0)[T_0]} - \boldsymbol{\mu}_r^{(0)*}\|_2 \vee \|\widehat{\boldsymbol{\Sigma}}^{(0)[T_0]} - \boldsymbol{\Sigma}^{(0)*}\|_2 \lesssim \sqrt{\frac{p}{n_0}} + T_0(\kappa'_0)^{T_0},$$

with probability at least $1 - C_3(K^{-1} + e^{-C_4 p})$, where $\kappa'_0 \in (0, 1)$ and $n_S = \sum_{k \in S} n_k$. When $T_0 \geq C \log n_0$ with a large constant $C > 0$, in both inequalities, the last term on the right-hand side will be dominated by other terms.

Next, we present the upper bound of the excess mis-clustering error on target task for TL-GMM. Having the estimator $\hat{\boldsymbol{\theta}}^{(0)[T_0]}$ and the truth $\boldsymbol{\theta}^{(0)*}$, the clustering method $\hat{\mathcal{C}}^{(0)[T_0]}$ and its mis-clustering error $R_{\hat{\boldsymbol{\theta}}^{(0)*}}(\hat{\mathcal{C}}^{(0)[T_0]})$ are defined in the same way as in (6) and (7).

Theorem 12. (*Upper bound of the target excess mis-clustering error for TL-GMM*) Suppose the same conditions in Theorem 11 hold. Then there exist constants C_1 and C_2 such that for any $\{\bar{\boldsymbol{\theta}}^{(k)*}\}_{k \in \{0\} \cup S} \in \bar{\Theta}'_S(h)$ and any probability measure \mathbb{Q}_S on $(\mathbb{R}^p)^{\otimes n_{Sc}}$, with probability at least $1 - C_1(K^{-1} + e^{-C_2 p})$ the following holds:

$$\begin{aligned} R_{\hat{\boldsymbol{\theta}}^{(0)*}}(\hat{\mathcal{C}}^{(0)[T_0]}) - R_{\hat{\boldsymbol{\theta}}^{(0)*}}(\mathcal{C}_{\hat{\boldsymbol{\theta}}^{(0)*}}) &\lesssim \underbrace{\frac{p}{n_S + n_0}}_{\text{(I)}} + \underbrace{\frac{1}{n_0}}_{\text{(II)}} + \underbrace{h^2 \wedge \frac{p}{n_0}}_{\text{(III)}} + \underbrace{\epsilon^2 \frac{p + \log K}{\max_{k=1:K} n_k} \wedge \frac{p}{n_0}}_{\text{(IV)}} \\ &\quad + \underbrace{\frac{\log K}{\max_{k=1:K} n_k}}_{\text{(V)}} + \underbrace{T_0^2 (\kappa'_0)^{2T_0}}_{\text{(VI)}}, \end{aligned}$$

with some constant $\kappa'_0 \in (0, 1)$. When $T_0 \geq C \log n_0$ with some large constant $C > 0$, the last term in the upper bound will be dominated by the second term.

Similar to the upper bounds of $d(\hat{\boldsymbol{\theta}}^{(k)[T]}, \boldsymbol{\theta}^{(k)*})$ and $R_{\hat{\boldsymbol{\theta}}^{(k)*}}(\hat{\mathcal{C}}^{(k)[T]}) - R_{\hat{\boldsymbol{\theta}}^{(k)*}}(\mathcal{C}_{\hat{\boldsymbol{\theta}}^{(k)*}})$ in Theorems 1 and 2, the upper bounds for $d(\hat{\boldsymbol{\theta}}^{(0)[T_0]}, \boldsymbol{\theta}^{(0)*})$ and $R_{\hat{\boldsymbol{\theta}}^{(0)*}}(\hat{\mathcal{C}}^{(0)[T_0]}) - R_{\hat{\boldsymbol{\theta}}^{(0)*}}(\mathcal{C}_{\hat{\boldsymbol{\theta}}^{(0)*}})$ consist of multiple parts with one-to-one correspondence. We take the bound of $R_{\hat{\boldsymbol{\theta}}^{(0)*}}(\hat{\mathcal{C}}^{(0)[T_0]}) - R_{\hat{\boldsymbol{\theta}}^{(0)*}}(\mathcal{C}_{\hat{\boldsymbol{\theta}}^{(0)*}})$ in Theorem 12 as an example. Part (I) is the oracle rate $\mathcal{O}_p(\frac{p}{n_S + n_0})$. Part (II) is the error caused by estimating scalar parameters $\delta^{(0)*}$ and $w^{(0)*}$ in the decision boundary, which thus do not depend on dimension p . Part (III) quantifies the contribution of related sources to the target task learning. The more related the sources in S to the target (i.e. the smaller h is), the smaller Part (III) becomes. Part (IV) captures the impact of outlier sources on the estimation error. As ϵ increases (i.e. the proportion of outlier sources increases), Part (IV) first increases and then flats out. It never exceeds the minimax rate $\mathcal{O}_p(p/n_0)$ of the single task learning on *target* task (Balakrishnan et al., 2017; Cai et al., 2019). Therefore, our method is robust against a fraction of outlier sources with arbitrary contaminated data. Part (V) is an extra term caused by estimating the center in MTL-GMM, which by Assumption 3.(i) is smaller than the single-task learning rate $\mathcal{O}_p(p/n_0)$. Part (VI) decreases geometrically in the iteration number T_0 of Algorithm 5, which becomes negligible by setting the iteration numbers T_0 large enough.

Consider the general scenario $T_0 \gtrsim \log n_0$. Then the upper bound of excess mis-clustering error rate $R_{\hat{\boldsymbol{\theta}}^{(0)*}}(\hat{\mathcal{C}}^{(0)[T_0]}) - R_{\hat{\boldsymbol{\theta}}^{(0)*}}(\mathcal{C}_{\hat{\boldsymbol{\theta}}^{(0)*}})$ is guaranteed to be no worse than the optimal single-task learning rate $\mathcal{O}_p(p/n_0)$. More importantly, in the general regime where $\epsilon \ll \sqrt{\frac{p \max_{k=1:K} n_k}{(p + \log K) n_0}}$ (small number of outlier sources), $h \ll \sqrt{p/n_0}$ (enough similarity between sources and target), $n_S \gg n_0$ (large total source sample size), and $\max_{k \in S} n_k/n_0 \gg \log K/p$ (large maximum source sample size), TL-GMM improves the GMM learning on the target task by achieving a better estimation error rate. As for the upper bound of

$\min_{\pi:[2]\rightarrow[2]} \max_{r=1:2} \|\widehat{\boldsymbol{\mu}}_{\pi(r)}^{(0)[T_0]} - \boldsymbol{\mu}_r^{(0)*}\|_2$ and $\|\widehat{\boldsymbol{\Sigma}}^{(0)[T_0]} - \boldsymbol{\Sigma}^{(0)*}\|_2$, when $T_0 \gtrsim \log n_0$, it has the single-task learning rate $\mathcal{O}_p(\sqrt{p/n_0})$. This is expected since the mean vectors and covariance matrices from sources are not necessarily similar to the one from target in the parameter space $\overline{\Theta}'_S(h)$.

The following result of minimax lower bounds shows that the upper bounds in Theorems 11 and 12 are optimal in a broad range of regimes.

Theorem 13. *(Lower bounds of the estimation error of GMM parameters in transfer learning) Suppose $\epsilon = \frac{K-s}{K} < a$ small constant c . Suppose there exists a subset S with $|S| \geq s$ such that $\min_{k \in S} n_k \geq C_1(p + \log K)$, $n_0 \geq C_1 p$ with a large constant C_1 and $\min_{k \in \{0\} \cup S} \Delta^{(k)} \geq C_2$ with some constant C_2 . Then we have*

$$\inf_{\widehat{\boldsymbol{\theta}}^{(0)}} \sup_{S: |S| \geq s} \sup_{\{\widehat{\boldsymbol{\theta}}^{(k)*}\}_{k \in \{0\} \cup S} \in \overline{\Theta}'_S(h)} \mathbb{P} \left(d(\widehat{\boldsymbol{\theta}}^{(0)}, \boldsymbol{\theta}^{(0)*}) \gtrsim \sqrt{\frac{p}{n_S + n_0}} + \sqrt{\frac{1}{n_0}} + h \wedge \sqrt{\frac{p}{n_0}} \right. \\ \left. + \frac{\epsilon}{\sqrt{\max_{k=1:K} n_k}} \wedge \sqrt{\frac{p}{n_0}} \right) \geq \frac{1}{10},$$

$$\inf_{\substack{\widehat{\boldsymbol{\mu}}_1^{(0)}, \widehat{\boldsymbol{\mu}}_2^{(0)} \\ \widehat{\boldsymbol{\Sigma}}^{(0)}}} \sup_{S: |S| \geq s} \sup_{\{\widehat{\boldsymbol{\theta}}^{(k)*}\}_{k \in \{0\} \cup S} \in \overline{\Theta}'_S(h)} \mathbb{P} \left(\min_{\pi:[2]\rightarrow[2]} \max_{r=1:2} \|\widehat{\boldsymbol{\mu}}_{\pi(r)}^{(0)} - \boldsymbol{\mu}_r^{(0)*}\|_2 \vee \|\widehat{\boldsymbol{\Sigma}}^{(0)} - \boldsymbol{\Sigma}^{(0)*}\|_2 \gtrsim \sqrt{\frac{p}{n_0}} \right) \geq \frac{1}{10}.$$

Theorem 14. *(Lower bound of the target excess mis-clustering error in transfer learning) Suppose the same conditions in Theorem 13 hold. Then we have*

$$\inf_{\widehat{\mathcal{C}}^{(0)}} \sup_{S: |S| \geq s} \sup_{\{\widehat{\boldsymbol{\theta}}^{(k)*}\}_{k \in \{0\} \cup S} \in \overline{\Theta}'_S(h)} \mathbb{P} \left(R_{\widehat{\boldsymbol{\theta}}^{(0)*}}(\widehat{\mathcal{C}}^{(0)}) - R_{\widehat{\boldsymbol{\theta}}^{(0)*}}(\mathcal{C}_{\widehat{\boldsymbol{\theta}}^{(0)*}}) \gtrsim \frac{p}{n_S + n_0} + \frac{1}{n_0} + h^2 \wedge \frac{p}{n_0} \right. \\ \left. + \frac{\epsilon^2}{\max_{k=1:K} n_k} \wedge \frac{1}{n_0} \right) \geq \frac{1}{10}.$$

Comparing the upper and lower bounds in Theorems 11- 14, several remarks are in order:

- With $T_0 \gtrsim \log n_0$, our estimators $\widehat{\boldsymbol{\mu}}_1^{(0)[T_0]}$, $\widehat{\boldsymbol{\mu}}_2^{(0)[T_0]}$, $\widehat{\boldsymbol{\Sigma}}^{(0)[T_0]}$ achieve the minimax optimal rate for estimating the mean vectors $\boldsymbol{\mu}_1^{(0)*}$, $\boldsymbol{\mu}_2^{(0)*}$ and the covariance matrix $\boldsymbol{\Sigma}^{(0)*}$.
- Regarding the target excess mis-clustering error, with the choices $T_0 \gtrsim \log n_0$, Part (VI) in the upper bound becomes negligible. We thus compare the other five terms in the upper bound with the corresponding terms in the lower bound.

1. Part (IV) in the upper bound differs from the one in the lower bound by a factor p (up to $\log K$). Hence the gap can arise when the dimension p diverges. The reason is similar to the one in multi-task learning setting and using statistical depth function based “center” estimates might be able to close the gap. We refer to the paragraph after Theorem 4 for more details.
 2. Part (V) in the upper bound does not appear in the lower bound. This term is due to the center estimate from the upper bound in MTL-GMM. When $\max_{k \in S} n_k/n_0 \gtrsim \log K$, this term is dominated by Part (II).
 3. The other three terms from the upper bound match with the ones in the lower bound.
- Based on the above comparisons, we can conclude that under the mild condition $\max_{k \in S} n_k/n_0 \gtrsim \log K$, our method is minimax rate optimal for the estimation of $\boldsymbol{\theta}^{(0)*}$ in the classical low-dimensional regime $p = O(1)$. Even when p is unbounded, the gap between the upper and lower bounds appears only when the fourth or fifth term is the dominating term in the upper bound. Like the discussions after Theorem 4, similar restricted regimes where our method might become sub-optimal can be derived.

A.4 Label alignment

As in multi-task learning, the alignment issue exists in transfer learning as well. Referring to the parameter space $\overline{\Theta}'(h)$ and the conditions of initialization in Assumptions 1 and 3, the success of Algorithm 5 requires correct alignments in two places. First, the center estimate $\widehat{\boldsymbol{\beta}}^{[T]}$ used as input of Algorithm 5 are obtained from Algorithm 1 which involves the alignment of initial estimates for sources. This alignment problem can be readily solved by Algorithm 2 or 3. Second, the initialization of the target problem $\widehat{\boldsymbol{\beta}}^{(0)[0]}$ needs to be correctly aligned with the aforementioned center estimates. This is easy to address using the alignment score described in Section 2.4.2 as there are only two different alignment options. We summarize the steps in Algorithm 6.

Like Algorithms 2 and 3, Algorithm 6 is able to find the correct alignments under mild conditions. Suppose $\{\widehat{\boldsymbol{\beta}}^{(k)[0]}\}_{k=0}^K$ are the initialization values with potentially wrong alignment. Define the correct alignment as $\mathbf{r}^* = (r_0^*, r_1^*, \dots, r_K^*)$ with $r_k^* = \arg \min_{r_k = \pm 1} \|r_k \widehat{\boldsymbol{\beta}}^{(k)[0]} - \boldsymbol{\beta}^{(k)*}\|_2$. For any $\mathbf{r} = \{r_k\}_{k=0}^K \in \{\pm 1\}^{K+1}$ which is a permutation order of $\{\widehat{\boldsymbol{\beta}}^{(k)[0]}\}_{k=0}^K$ and its corresponding alignment $\{r_k \widehat{\boldsymbol{\beta}}^{(k)[0]}\}_{k=0}^K$, define its alignment score as

$$\text{score}(\mathbf{r}) = \sum_{0 \leq k_1 \neq k_2 \leq K} \|r_{k_1} \widehat{\boldsymbol{\beta}}^{(k_1)[0]} - r_{k_2} \widehat{\boldsymbol{\beta}}^{(k_2)[0]}\|_2.$$

As expected, under the conditions from Algorithms 2 or 3 for sources together with some similar conditions on the target, Algorithm 6 will output the ideal alignment $\widehat{\mathbf{r}}'$ (equivalently, the good initialization $\widehat{\mathbf{r}}'_0 \widehat{\boldsymbol{\beta}}^{(0)[0]}$ for Algorithm 5).

Algorithm 6: Alignment for transfer learning

Input: Initialization $\{(\hat{\beta}^{(k)[0]})\}_{k=0}^K$, and $\hat{\mathbf{r}}$ from Algorithm 2 or 3
1 **if** $\text{score}((-1, \hat{\mathbf{r}})) > \text{score}((1, \hat{\mathbf{r}}))$ **then**
2 $\hat{\mathbf{r}}' = (1, \hat{\mathbf{r}})$
3 **else**
4 $\hat{\mathbf{r}}' = (-1, \hat{\mathbf{r}})$
5 **end**
Output: $\hat{\mathbf{r}}'$

Theorem 15 (Alignment correctness for Algorithm 6). *Assume that*

$$(i) \quad \epsilon < \frac{1}{2};$$

$$(ii) \quad \|\beta^{(0)*}\|_2 > \frac{2(1-\epsilon)}{1-2\epsilon}h + \frac{2-\epsilon}{1-2\epsilon} \max_{k \in \{0\} \cup S} (\|\hat{\beta}^{(k)[0]} - \beta^{(k)*}\|_2 \wedge \|\hat{\beta}^{(k)[0]} + \beta^{(k)*}\|_2),$$

where $\epsilon = \frac{K-s}{K}$ is the outlier source task proportion, and h is the degree of discriminant coefficient relatedness defined in (A.12).

For $\hat{\mathbf{r}}$ in Algorithm 6: if it is from Algorithm 2, assume the conditions of Theorem 5 hold; if it is from Algorithm 3, assume the conditions of Theorem 6 hold. Then the output of Algorithm 6 satisfies

$$\hat{\mathbf{r}}'_k = r_k^* \text{ for all } k \in \{0\} \cup S \quad \text{or} \quad \hat{\mathbf{r}}'_k = -r_k^* \text{ for all } k \in \{0\} \cup S.$$

B Alignment for Multi-cluster GMMs in MTL

Similar to the binary case, we have the alignment issues in multi-cluster case as well. In this section, we propose two alignment algorithms as the extension to the Algorithms 2 and 3.

In the multi-cluster case, the alignment of each task can be represented as a permutation of $[R]$. Consider a series of permutations $\boldsymbol{\pi} = \{\pi_k\}_{k=1}^K$, where each π_k is a permutation function on $[R]$. Define a score of $\boldsymbol{\pi}$ as

$$\text{score}(\boldsymbol{\pi}) = \sum_{r=2}^R \sum_{k \neq k'} \left\| (\hat{\Sigma}^{(k)[0]})^{-1} (\hat{\boldsymbol{\mu}}_{\pi_k(r)}^{(k)[0]} - \hat{\boldsymbol{\mu}}_{\pi_k(1)}^{(k)[0]}) - (\hat{\Sigma}^{(k')[0]})^{-1} (\hat{\boldsymbol{\mu}}_{\pi_{k'}(r)}^{(k')[0]} - \hat{\boldsymbol{\mu}}_{\pi_{k'}(1)}^{(k')[0]}) \right\|_2.$$

We want to recover the correct alignment $\pi_k^* = \arg \min_{\pi_k: [R] \rightarrow [R]} \sum_{r=2}^R \left\| (\hat{\Sigma}^{(k)[0]})^{-1} (\hat{\boldsymbol{\mu}}_{\pi_k(r)}^{(k)[0]} - \hat{\boldsymbol{\mu}}_{\pi_k(1)}^{(k)[0]}) - \beta_r^{(k)*} \right\|_2$.

We proposed an exhaustive search algorithm, which is summarized in Algorithm 7.

The following theorem shows that under certain conditions, the output from Algorithm 7 recovers the correct alignment up to a permutation.

Theorem 16 (Alignment correctness for Algorithm 7). *Assume that*

Algorithm 7: Exhaustive search for the alignment in multi-cluster GMMs

Input: Initialization $\{\{\hat{\mu}_r^{(k)[0]}\}_{r=1}^R, \hat{\Sigma}^{(k)[0]}\}_{k=1}^K$

1 $\hat{\pi} = \{\hat{\pi}_k\}_{k=1}^K \leftarrow \arg \min_{\pi} \text{score}(\pi)$

Output: $\hat{\pi} = \{\hat{\pi}_k\}_{k=1}^K$

(i) $\max_{k \in S} \max_{r \neq j} \Delta_{rj}^{(k)} / \min_{k \in S} \min_{r \neq j} \Delta_{rj}^{(k)} \leq D$ with $D \geq 1$.

(ii) $\epsilon < \frac{1}{24Dc_{\Sigma}+1}$;

(iii) $\min_{k \in S} \min_{r \neq j} \Delta_{rj}^{(k)} \geq \left[\frac{4(1-\epsilon)c_{\Sigma}^{1/2}}{1-(24Dc_{\Sigma}+1)\epsilon} h + \frac{(4+20\epsilon)c_{\Sigma}^{1/2}}{1-(24Dc_{\Sigma}+1)\epsilon} \xi \right] \vee \left[\frac{13(1-\epsilon)c_{\Sigma}^{1/2}}{1-(9Dc_{\Sigma}+1)\epsilon} h + \frac{(13-4\epsilon)c_{\Sigma}^{1/2}}{1-(9Dc_{\Sigma}+1)\epsilon} \xi \right],$

where $\epsilon = \frac{K-s}{K}$ is the outlier task proportion introduced in Theorem 7, h is degree of discriminant coefficient similarity defined in (8), and

$$\begin{aligned} \xi &= \max_{k \in S} \min_{\pi: [R] \rightarrow [R]} \max_{r=1:R} \left\| (\hat{\Sigma}^{(k)[0]})^{-1} (\hat{\mu}_{\pi(r)}^{(k)[0]} - \hat{\mu}_{\pi(1)}^{(k)[0]}) - \beta_r^{(k)*} \right\|_2 \\ &= \max_{k \in S} \min_{\pi: [R] \rightarrow [R]} \max_{r=1:R} \left\| (\hat{\Sigma}^{(k)[0]})^{-1} (\hat{\mu}_{\pi(r)}^{(k)[0]} - \hat{\mu}_{\pi(1)}^{(k)[0]}) - (\Sigma^{(k)*})^{-1} (\mu_r^{(k)*} - \mu_1^{(k)*}) \right\|_2. \end{aligned}$$

Then there exists a permutation $\iota: [R] \rightarrow [R]$, such that the output of Algorithm 7 satisfies

$$\hat{\pi}_k = \iota \circ \pi_k^*,$$

for all $k \in S$.

The biggest issue of Algorithm 7 is the computational time. It is easy to see that the time complexity of it is $\mathcal{O}((R!)^K \cdot RK^2)$, because it needs to search over all permutations. This is not practically feasible when R and K are large. Therefore, we propose the following greedy search algorithm to reduce the computational cost, which is summarized in Algorithm 8. Note that its main idea is similar to Algorithm 3 for the binary GMM, but the procedure is different. We define the score of alignments $\{\pi_{k'}\}_{k'=1}^k$ of tasks 1- k' as

$$\begin{aligned} &\text{score}(\{\pi_{k'}\}_{k'=1}^k | \{\{\hat{\mu}_r^{(k')[0]}\}_{r=1}^R\}_{k'=1}^k, \{\hat{\Sigma}^{(k')[0]}\}_{k'=1}^k) \\ &= \sum_{r=2}^R \sum_{\tilde{k}, k' \leq k} \left\| (\hat{\Sigma}^{(\tilde{k})[0]})^{-1} (\hat{\mu}_{\pi_{\tilde{k}}(r)}^{(\tilde{k})[0]} - \hat{\mu}_{\pi_{\tilde{k}}(1)}^{(\tilde{k})[0]}) - (\hat{\Sigma}^{(k')[0]})^{-1} (\hat{\mu}_{\pi_{k'}(r)}^{(k')[0]} - \hat{\mu}_{\pi_{k'}(1)}^{(k')[0]}) \right\|_2. \end{aligned}$$

The subsequent theorem demonstrates that, with slightly stronger assumptions than those required by Algorithm 7, the greedy search algorithm can recover the correct alignment up to a permutation with high probability. Importantly, this approach significantly alleviates the computational cost from $\mathcal{O}((R!)^K \cdot RK^2)$ to $\mathcal{O}(R!K \cdot RK^2)$.

Theorem 17. Assume there are no outlier tasks in the first K_0 tasks, and

Algorithm 8: Greedy search for the alignment in multi-cluster GMMs

Input: Initialization $\{\{\hat{\mu}_r^{(k)[0]}\}_{r=1}^R, \hat{\Sigma}^{(k)[0]}\}_{k=1}^K$

1 **for** $k = 1$ **to** K **do**

2 With $\{\hat{\pi}_{k'}\}_{k'=1}^{k-1}$ fixed (\emptyset when $k = 1$), set

$$\hat{\pi}_k = \arg \min_{\pi: [R] \rightarrow [R]} \text{score}(\{\hat{\pi}_{k'}\}_{k'=1}^{k-1} \cup \pi | \{\{\hat{\mu}_r^{(k')[0]}\}_{r=1}^R\}_{k'=1}^k, \{\hat{\Sigma}^{(k')[0]}\}_{k'=1}^k)$$

3 **end**

Output: $\hat{\pi} = \{\hat{\pi}_k\}_{k=1}^K$

$$(i) \max_{k \in S} \max_{r \neq j} \Delta_{rj}^{(k)} / \min_{k \in S} \min_{r \neq j} \Delta_{rj}^{(k)} \leq D \text{ with } D \geq 1.$$

$$(ii) \epsilon < \frac{1}{2Dc_{\Sigma}+1};$$

$$(iii) K_0 > 2Dc_{\Sigma}K\epsilon;$$

$$(iv) \min_{k \in S} \min_{r \neq j} \Delta_{rj}^{(k)} \geq \left[\frac{4K_0c_{\Sigma}^{1/2}}{K_0 - Dc_{\Sigma}K\epsilon}h + \frac{(4K_0+K\epsilon)c_{\Sigma}^{1/2}}{K_0 - Dc_{\Sigma}K\epsilon}\xi \right] \vee \left[\frac{2K_0c_{\Sigma}^{1/2}}{K_0 - 2Dc_{\Sigma}K\epsilon}h + \frac{(2K_0+2K\epsilon)c_{\Sigma}^{1/2}}{K_0 - 2Dc_{\Sigma}K\epsilon}\xi \right],$$

where $\epsilon = \frac{K-s}{K}$ is the outlier task proportion and c_{Σ} appears in the condition that $c_{\Sigma}^{-1} \leq \min_{k \in S} \lambda_{\min}(\Sigma^{(k)*}) \leq \max_{k \in S} \lambda_{\max}(\Sigma^{(k)*}) \leq c_{\Sigma}$. Then there exists a permutation $\iota : [R] \rightarrow [R]$, such that the output of Algorithm 8 satisfies

$$\hat{\pi}_k = \iota \circ \pi_k^*,$$

for all $k \in S$.

Remark 8. Conditions (ii)-(iv) are similar to the conditions in Theorem 6. The inclusion of Condition (i) aims to facilitate the analysis in the proof, and we conjecture that the obtained results persist even if this condition is omitted.

When R is very large, the computational burden becomes prohibitive, rendering even the $\mathcal{O}(R!K \cdot RK^2)$ time complexity impractical. Addressing this computational challenge requires the development of more efficient alignment algorithms, a pursuit that we defer to future investigations. In addition, one caveat of the greedy search algorithm is that we need to know K_0 non-outlier tasks in priori, which may not be unrealistic in practice. In our empirical examinations, we enhance the algorithm's performance by introducing a random shuffle of the K tasks in each iteration. Specifically, we execute Algorithm 8 200 times, yielding 200 alignment candidates. The final alignment is then determined by selecting the configuration that attains the minimum score among the candidates.

C Additional Numerical Studies

C.1 Simulations

C.1.1 Simulation 1 of MTL

In this subsection, we provide some additional performance evaluation of the three methods (MTL-GMM, Pooled-GMM, and Single-task-GMM) in Simulation 1. The results are presented in Figures 6 and 7.

Referring to Figure 6 for the case without outlier tasks, MTL-GMM outperforms Pooled-GMM in estimating $w^{(k)*}$ all the time. This makes sense because Pooled-GMM does not take the heterogeneity of $w^{(k)*}$'s into account. For the estimation of other parameters (except $\delta^{(k)*}$ ⁵) and clustering, MTL-GMM and Pooled-GMM are competitive when h is small (i.e. the tasks are similar). As h increases (i.e. tasks become more heterogenous), MTL-GMM starts to outperform Pooled-GMM by a large margin. Moreover, MTL-GMM is significantly better than Single-task-GMM in terms of both estimation and mis-clustering errors over a wide range of h . They only become comparable when h is very large. These comparisons demonstrate that MTL-GMM not only effectively utilizes the unknown similarity structure among tasks, but also adapts to it.

The results for the case with two outlier tasks are shown in Figure 7. It is clear that the comparison between MTL-GMM and Single-task-GMM is similar to the one in Figure 6. What is new here is that even when h is very small, MTL-GMM still performs much better than Pooled-GMM, showing the robustness of MTL-GMM against a fraction of outlier tasks. Note that in this simulation, $\delta^{(k)*} = 0$ for all $k \in [K]$, which might explain the phenomenon where Pooled-GMM outperforms MTL-GMM in estimating $\delta^{(k)*}$'s.

C.1.2 Simulation 2 of MTL

In this subsection, we provide some additional performance evaluation of three methods in Simulation 2. The results are presented in Figures 8 and 9.

The results are reported in Figures 8 and 9. The main message is the same as in the previous simulation example: Pooled-GMM is sensitive to outlier tasks and suffers from negative transfer when h is large, while MTL-GMM is robust to outliers and can adapt to the unknown similarity level h . The results verify the theoretical findings in the multi-cluster case.

C.1.3 Simulation 3 of MTL

In the third simulation of MTL, we consider a different similarity structure among tasks in S and a different type of outlier tasks. For a multi-task learning problem with $K = 10$ tasks, set the sample size of each task equal to 100. Let $\beta^{(1)*} = (2.5, 0, 0, 0, 0)$, $\Sigma^{(1)*} =$

⁵Actually it is not suprising to see Pooled-GMM estimates $\mu_1^{(k)*}$, $\mu_2^{(k)*}$, $\delta^{(k)*}$, and $\Sigma^{(k)*}$ better than MTL-GMM when h is small in this example. The reason is that these parameters are similar to each other (although MTL-GMM does not rely on this similarity) which makes pooling the data a good approach.




Figure 6: The performance of different methods in Simulation 1.(i) of multi-task learning, with no outlier tasks ($\epsilon = 0$), and h changing from 0 to 10 with increment 1. Estimation error of $\{w^{(k)*}\}_{k \in S}$ stands for $\max_{k \in S} (|\hat{w}^{(k)[T]} - w^{(k)*}| \wedge |1 - \hat{w}^{(k)[T]} - w^{(k)*}|)$. Estimation error of $\{\mu_1^{(k)*}\}_{k \in S}$ and $\{\mu_2^{(k)*}\}_{k \in S}$ stands for $\max_{k \in S} \min_{\pi: [2] \rightarrow [2]} (\|\hat{\mu}_1^{(k)[T]} - \mu_{\pi(1)}^{(k)*}\|_2 \vee \|\hat{\mu}_2^{(k)[T]} - \mu_{\pi(2)}^{(k)*}\|_2)$. Estimation error of $\{\Sigma^{(k)*}\}_{k \in S}$ stands for $\max_{k \in S} \|\hat{\Sigma}^{(k)[T]} - \Sigma^{(k)*}\|_2$. Estimation error of $\{\delta^{(k)*}\}_{k \in S}$ stands for $\max_{k \in S} (|\hat{\delta}^{(k)[T]} - \delta^{(k)*}| \wedge |\hat{\delta}^{(k)[T]} + \delta^{(k)*}|)$. Average mis-clustering error represents the average empirical mis-clustering error rate calculated on the test set of tasks in S .

$(0.5^{|i-j|})_{5 \times 5}$, and $1 \in S$, i.e., the first task is not an outlier task. We generate each $w^{(k)*}$ from $\text{Unif}(0.1, 0.9)$ for all $k \in S$. For $k \in S \setminus \{1\}$, we generate $\Sigma^{(k)*}$ as

$$\Sigma^{(k)*} = \begin{cases} (0.5^{|i-j|})_{5 \times 5}, & \text{with probability } 1/2, \\ (a^{|i-j|})_{5 \times 5}, & \text{with probability } 1/2, \end{cases} \quad (\text{C.13})$$




Figure 7: The performance of different methods in Simulation 1.(ii) of multi-task learning, with 2 outlier tasks ($\epsilon = 0.2$), and h changing from 0 to 10 with increment 1. The meaning of each subfigure's title is the same as in Figure 6.

and set $\beta^{(k)*} = (\Sigma^{(k)*})^{-1} \Sigma^{(1)*} \beta^{(1)*}$. Here, the value of a is determined by $\max\{a \in [0.5, 1) : \|\beta^{(k)*} - \beta^{(1)*}\|_2 \leq h\}$ for a given h . Let $\mu_2^{(k)*} = \Sigma^{(k)*} \beta^{(k)*}$ and $\mu_1^{(k)*} = \mathbf{0}$, $\forall k \in S$. In this generation process, $\mu_2^{(k)*} = \mu_2^{(1)*} = \Sigma^{(1)*} \beta^{(1)*} = (5/2, 5/4, 5/8, 5/16, 5/32)^T$ for all $k \in S$. The covariance matrix of tasks in S can differ. When $k \notin S$, we generate the data of task k from two clusters with probability $1 - w^{(k)*}$ and $w^{(k)*}$, where $w^{(k)*} \sim \text{Unif}(0.1, 0.9)$. Samples from the second cluster follow $N(\mu_2^{(k)*}, \Sigma^{(k)*})$, with $\Sigma^{(k)*}$ coming from (C.13), $\beta^{(k)*} = (-2.5, -2.5, -2.5, -2.5, -2.5)^T$, and $\mu_2^{(k)*} = \Sigma^{(k)*} \beta^{(k)*}$. For each sample from the first cluster, each component is independently generated from a t -distribution with degrees of freedom 4. In each replication, for given ϵ , the outlier task index set S^c is uniformly sampled from all subsets of $2 : K$ with cardinality $K\epsilon$ (since task 1 has been fixed in S).




Figure 8: The performance of different methods in Simulation 2.(i) of multi-task learning, with no outlier tasks ($\epsilon = 0$), and h changing from 0 to 10 with increment 1. Estimation error of $\{w_r^{(k)*}\}_{r \in [R], k \in S}$ stands for $\max_{k \in S} \min_{\pi: [R] \rightarrow [R]} \max_{r \in [R]} |\hat{w}_r^{(k)[T]} - w_{\pi(r)}^{(k)*}|$. Estimation error of $\{\mu_r^{(k)*}\}_{r \in [R], k \in S}$ stands for $\max_{k \in S} \min_{\pi: [R] \rightarrow [R]} \max_{r \in [R]} \|\hat{\mu}_r^{(k)[T]} - \mu_{\pi(r)}^{(k)*}\|_2$. Estimation error of $\{\Sigma^{(k)*}\}_{k \in S}$ stands for $\max_{k \in S} \|\hat{\Sigma}^{(k)[T]} - \Sigma^{(k)*}\|_2$. Estimation error of $\{\delta_r^{(k)*}\}_{r \in [R], k \in S}$ stands for $\max_{k \in S} \min_{\pi: [R] \rightarrow [R]} \max_{r \in [R]} |\hat{\delta}_r^{(k)[T]} - (\mu_{\pi(r)}^{(k)*} + \mu_{\pi(1)}^{(k)*})^T (\Sigma^{(k)*})^{-1} (\mu_{\pi(r)}^{(k)*} - \mu_{\pi(1)}^{(k)*})/2|$. Average mis-clustering error represents the average empirical mis-clustering error rate calculated on the test set of tasks in S .

We consider two cases:




Figure 9: The performance of different methods in Simulation 2.(ii) of multi-task learning, with 2 outlier tasks ($\epsilon = 0.2$), and h changing from 0 to 10 with increment 1. The meaning of each subfigure's title is the same as in Figure 8.

- (i) No outlier tasks ($\epsilon = 0$), and h changes from 0 to 10 with increment 1;
- (ii) 2 outlier tasks ($\epsilon = 0.2$), and h changes from 0 to 10 with increment 1.

We implement the same three methods as in Simulation 1 and the results are reported in Figures 10 and 11. When there are no outlier tasks, both MTL-GMM and Pooled-GMM significantly outperform Single-task-GMM. Note that in this simulation, $\mu_1^{(k)*} = \mu_1^{(k')*}$ and $\mu_2^{(k)*} = \mu_2^{(k')*}$ for all $k \neq k' \in [K]$, which might explain the phenomenon where Pooled-




Figure 10: The performance of different methods in Simulation 3.(i) of multi-task learning, with no outlier tasks ($\epsilon = 0$), and h changing from 0 to 10 with increment 1. Estimation error of $\{\beta^{(k)*}\}_{k \in S}$ stands for $\max_{k \in S} (\|\hat{\beta}^{(k)[T]} - \beta^{(k)*}\|_2 \wedge \|\hat{\beta}^{(k)[T]} + \beta^{(k)*}\|_2)$. Maximum mis-clustering error represents maximum empirical mis-clustering error rate calculated on test set of tasks in S . The meaning of other subfigures' titles is the same as in Figure 6.




Figure 11: The performance of different methods in Simulation 3.(ii) of multi-task learning, with 2 outlier tasks ($\epsilon = 0.2$), and h changing from 0 to 10 with increment 1. The meaning of each subfigure's title is the same as in Figure 10.

GMM outperforms MTL-GMM in estimating $\mu_1^{(k)*}$ and $\mu_2^{(k)*}$'s. When there are two outlier tasks, Figure 11 shows that Pooled-GMM performs much worse than Single-task-GMM on

most of the estimation errors of GMM parameters as well as the mis-clustering error rate. In contrast, MTL-GMM greatly improves the performance of Single-task-GMM, showing the advantage of MTL-GMM when dealing with outlier tasks and heterogeneous covariance matrices.

C.1.4 Simulation of TL

Consider a transfer learning problem with $K = 10$ source data sets, where all sources are from the same GMM. The setting is modified based on Simulation 1 of MTL. The source and target sample sizes are equal to 100. For each of the source and target task, $w^{(k)*} \sim \text{Unif}(0.1, 0.9)$ and $\boldsymbol{\mu}_1^{(k)*} = -\boldsymbol{\mu}_2^{(k)*} = (2, 2, \mathbf{0}_{p-2})^T + h/2 \cdot (\boldsymbol{\Sigma}^{(k)*})^{-1} \mathbf{u}$, where $p = 15$, $\boldsymbol{\Sigma}^{(k)*} = (0.2^{|i-j|})_{5 \times 5}$, and $\mathbf{u} \sim \text{Unif}(\{\mathbf{u} \in \mathbb{R}^p : \|\mathbf{u}\|_2 = 1\})$. We consider the case that h changes from 0 to 10 with increment 1.

We compare five different methods, including Target-GMM fitted on target data only, MTL-GMM fitted on all the data, MTL-GMM-center which fits MTL-GMM on source data and outputs the estimated “center” $\bar{\boldsymbol{\beta}}^{[T]}$ as the target estimate⁶, Pooled-GMM which fits a merged GMM on all the data, and our TL-GMM. The performance is evaluated by the estimation errors of $w^{(0)*}$, $\boldsymbol{\mu}_1^{(0)*}$, $\boldsymbol{\mu}_2^{(0)*}$, $\boldsymbol{\beta}^{(0)*}$, $\delta^{(0)*}$, and $\boldsymbol{\Sigma}^{(0)*}$ as well as the mis-clustering error rate calculated on an independent test target data of size 500. Results are presented in Figure 12.

Figure 12 shows that when h is small, the performances of MTL-GMM, MTL-GMM-center, Pooled-GMM, and TL-GMM are comparable, and all of them are much better than Target-GMM. This is expected, because the sources are very similar to the target and can be easily used to improve the target task learning. As h keeps increasing, the target and sources become increasingly different. This is the phase where the knowledge of sources need be carefully transferred for the possible learning improvement on the target task. As is clear from Figure 12, MTL-GMM, MTL-GMM-center, and Pooled-GMM do not handle heterogeneous resources well, thus outperformed by Target-GMM. By contrast, TL-GMM remains effective in transferring source knowledge to improve over Target-GMM; when h is very large so that sources are not useful anymore, TL-GMM is robust enough to still have competitive performance compared to Target-GMM.

Figure 13 shows the results when there are two outlier tasks ($\epsilon = 0.2$). It can be seen that MTL-GMM is robust to outliers.

C.1.5 Tuning parameters C_λ and C_{λ_0} in Algorithms 1, 4, and 5

The candidates of C_λ and C_{λ_0} values used in the 10-fold cross-validation are chosen through a data-driven way. For C_λ in Algorithm 1, we first determine the smallest C_λ value which makes all $\boldsymbol{\beta}^{(k)}$ estimators identical, which is denoted as C_{\max} . Then the C_λ candidates are set to be the sequence from $C_{\max}/50$ and $2C_{\max}$ with equal logarithm distance. For C_{λ_0} in Algorithm 5, we first determine the smallest C_{λ_0} value which makes the $\boldsymbol{\beta}^{(0)}$ estimator

⁶MTL-GMM-center only appears in the comparison of estimation error of $\boldsymbol{\beta}^{(k)*}$ s.




Figure 12: The performance of different methods in the simulation of transfer learning, with no outlier tasks ($\epsilon = 0$) and h changing from 0 to 10 with increment 1. Estimation error of $w^{(0)*}$ stands for $|\hat{w}^{(0)[T_0]} - w^{(0)*}| \wedge |1 - \hat{w}^{(0)[T_0]} - w^{(0)*}|$. Estimation error of $\mu_1^{(0)*}$ and $\mu_2^{(0)*}$ stands for $\min_{\pi: [2] \rightarrow [2]} \max_{r \in [2]} \|\hat{\mu}_r^{(0)[T_0]} - \mu_{\pi(r)}^{(0)*}\|_2$. Estimation error of $\beta^{(0)*}$ stands for $\|\hat{\beta}^{(0)[T_0]} - \beta^{(0)*}\|_2 \wedge \|\hat{\beta}^{(0)[T_0]} + \beta^{(0)*}\|_2$. Estimation error of $\Sigma^{(k)*}$ stands for $\|\hat{\Sigma}^{(0)[T_0]} - \Sigma^{(0)*}\|_2$. Estimation error of $\delta_r^{(0)*}$ stands for $|\hat{\delta}^{(0)[T_0]} - \delta^{(0)*}| \wedge |\hat{\delta}^{(0)[T_0]} + \delta^{(0)*}|$. Mis-clustering error represents the empirical mis-clustering error rate calculated on the test set of the target data.




Figure 13: The performance of different methods in the simulation of transfer learning, with 2 outlier tasks ($\epsilon = 0.2$) and h changing from 0 to 10 with increment 1. The meaning of each subfigure's title is the same as in Figure 12.

equal to $\bar{\beta}^{[T]}$, which is denoted as C'_{\max} . Then the C_{λ_0} candidates are set to be the sequence

from $C''_{\max}/50$ and $2C''_{\max}$ with equal logarithm distance.

We also run MTL-GMM with different C_λ values in Simulation 1 to test the impact of penalty parameter. The results are presented in Figure 14. The values 1.29, 2.15, 3.59, 5.99, and 10 are the last 5 elements in sequence from 0.1 to 10 with equal logarithm distance. It can be seen that with small C_λ values like 1.29 and 2.15, the performance of MTL-GMM is similar to that of Single-task-GMM, although MTL-GMM-2.15 improves Single-task-GMM a lot when h is small. With large C_λ values like 5.99 and 10, MTL-GMM performs similarly to Pooled-GMM when h is small, while suffers from negative transfer when h is large. However, as h continues to increase, the performance of MTL-GMM with large C_λ values start to improve and finally become similar to Single-task-GMM. This phenomenon is in accordance with the theory, as the theory predicts that MTL-GMM achieves the same rate as Single-task-GMM for large h . The negative transfer effect of MTL-GMM with large C_λ could be caused by large unknown constants in the upper bound. Comparing Figure 14 with figures in Sections 4.1.1 and C.1.1, we can see that cross-validation enhances the performance of MTL-GMM.

C.1.6 Tuning parameter κ and κ_0 in Algorithms 1, 4, and 5

We set $\kappa = \kappa_0 = 1/3$ in Algorithms 1, 4, and 5. We run MTL-GMM with different κ values in Simulation 1 to test the impact of κ on the performance. The results are presented in Figure 15. We tried $\kappa = 0.1, 0.3, 0.5, 0.7, 0.9$ in Algorithms 1. It can be seen that the lines representing MTL-GMM with different κ values highly overlap with each other, which shows that the performance of MTL-GMM is very robust to the choice of κ . In practice, we take $\kappa = 1/3$ for convenience.

D Technical Lemmas

D.1 General lemmas

Denote the unit ball $\mathcal{B}^p = \{\mathbf{u} \in \mathbb{R}^p : \|\mathbf{u}\|_2 \leq 1\}$ and the unit sphere $\mathcal{S}^{p-1} = \{\mathbf{u} \in \mathbb{R}^p : \|\mathbf{u}\|_2 = 1\}$.

Lemma 1 (Covering number of the unit ball under Euclidean norm, Example 5.8 in Wainwright (2019)). *Denote the ϵ -covering number of a unit ball \mathcal{B}^p in \mathbb{R}^p under Euclidean norm as $N(\epsilon, \mathcal{B}^p, \|\cdot\|_2)$, where the centers of covering balls are required to be on the sphere. We have $(1/\epsilon)^p \leq N(\epsilon, \mathcal{B}^p, \|\cdot\|_2) \leq (1 + 2/\epsilon)^p$.*

Lemma 2 (Packing number of the unit sphere under Euclidean norm). *Denote the ϵ -packing number of the unit sphere \mathcal{S}^{p-1} in \mathbb{R}^p under Euclidean norm as $M(\epsilon, \mathcal{S}^{p-1}, \|\cdot\|_2)$. When $p \geq 2$, we have $M(\epsilon, \mathcal{S}^{p-1}, \|\cdot\|_2) \geq N(\epsilon, \mathcal{B}^{p-1}, \|\cdot\|_2) \geq (1/\epsilon)^{p-1}$.*

Lemma 3 (Fano's lemma, see Chapter 2 of Tsybakov (2009), Chapter 15 of Wainwright (2019)). *Suppose (Θ, d) is a metric space and each θ in this space is associated with a*




Figure 14: The performance of different methods in Simulation 1 of multi-task learning, with no outlier tasks ($\epsilon = 0$) and h changing from 0 to 10 with increment 1. The meaning of each subfigure's title is the same as in Figures 2 and 6.

probability measure \mathbb{P}_θ . If $\{\theta_j\}_{j=1}^N$ is an s -separated set (i.e. $d(\theta_j, \theta_k) \geq s$ for any $j \neq k$),




Figure 15: The performance of different methods in Simulation 1 of multi-task learning, with no outlier tasks ($\epsilon = 0$) and h changing from 0 to 10 with increment 1. The meaning of each subfigure's title is the same as in Figures 2 and 6.

and $\text{KL}(\mathbb{P}_{\theta_j}, \mathbb{P}_{\theta_k}) \leq \alpha \log N$, then

$$\inf_{\hat{\theta}} \sup_{\theta \in \Theta} \mathbb{P}_{\theta}(d(\hat{\theta}, \theta) \geq s/2) \geq \inf_{\psi} \sup_{j=1:N} \mathbb{P}_{\theta_j}(\psi \neq j) \geq 1 - \alpha - \frac{\log 2}{\log N},$$

where $\psi : X \mapsto \psi(X) \in \{1, \dots, N\}$.

Lemma 4 (Packing number of the unit sphere in a quadrant under Euclidean norm). *In \mathbb{R}^p , we can use a vector $\mathbf{v} \in \{\pm 1\}^{\otimes p}$ to indicate each quadrant $\mathcal{Q}_{\mathbf{v}} = \{[0, +\infty) \cdot \mathbf{1}(v_j = +1) + (-\infty, 0) \cdot \mathbf{1}(v_j = -1)\}^{\otimes p}$. Then when $p \geq 2$, there exists a quadrant $\mathcal{Q}_{\mathbf{v}_0}$ such that $M(\epsilon, \mathcal{S}^{p-1} \cap \mathcal{Q}_{\mathbf{v}_0}, \|\cdot\|_2) \geq (\frac{1}{2})^p (\frac{1}{\epsilon})^{p-1}$.*

Lemma 5. *For one-dimensional Gaussian mixture variable $Z \sim (1-w)\mathcal{N}(\mu_1, \sigma^2) + w\mathcal{N}(\mu_2, \sigma^2)$ with $(1-w)\mu_1 + w\mu_2 = 0$, it is a $\sqrt{\sigma^2 + \frac{1}{4}|\mu_1 - \mu_2|^2}$ -subGaussian variable. That means,*

$$\mathbb{E}e^{\lambda Z} \leq \exp \left\{ \frac{1}{2} \lambda \left(\sigma^2 + \frac{1}{4} |\mu_1 - \mu_2|^2 \right) \right\}.$$

Lemma 6 (Duan and Wang (2022)). *Let*

$$(\{\hat{\boldsymbol{\theta}}_j\}_{j=1}^K, \hat{\boldsymbol{\beta}}) = \arg \min_{\boldsymbol{\theta}_k, \boldsymbol{\beta} \in \mathbb{R}^p} \left\{ \sum_{k=1}^K \omega_k f^{(k)}(\boldsymbol{\theta}_k) + \sqrt{\omega_k} \lambda \|\boldsymbol{\beta} - \boldsymbol{\theta}_k\|_2 \right\}.$$

Suppose there exists $S \subseteq 1 : K$ such that the following conditions are satisfied:

(i) *For any $k \in S$, f_k is $(\boldsymbol{\theta}_k^*, M, \rho, L)$ -regular, that is*

- f_k is convex and twice differentiable;
- $\rho \mathbf{I} \preceq \nabla^2 f_k(\boldsymbol{\theta}) \preceq L \mathbf{I}$ for all $\boldsymbol{\theta} \in \mathcal{B}(\boldsymbol{\theta}_k^*, M)$;
- $\|\nabla f_k(\boldsymbol{\theta}_k^*)\|_2 \leq \rho M/2$.

(ii) $\min_{\boldsymbol{\theta} \in \mathbb{R}^d} \max_{k \in S} \{\|\boldsymbol{\theta}_k^* - \boldsymbol{\theta}\|_2\} \leq h$ and $\sum_{k \in S^c} \sqrt{\omega_k} \leq \epsilon \sum_{k \in S} \omega_k / \max_{k \in S} \sqrt{\omega_k}$.

Then we have the following conclusions:

(i) $\|\hat{\boldsymbol{\theta}}_k - \boldsymbol{\theta}_k^*\|_2 \leq \frac{1}{\rho} (\|\nabla f^{(k)}(\boldsymbol{\theta}_k^*)\|_2 + \lambda / \sqrt{\omega_k})$ for all $k \in S$.

(ii) *If*

$$\frac{5\rho\kappa_w \max_{k \in S} \{\sqrt{\omega_k} \|\nabla f^{(k)}(\boldsymbol{\theta}_k^*)\|_2\}}{1 - \rho\epsilon} < \lambda < \frac{\rho M}{2} \min_{k \in S} \sqrt{\omega_k},$$

where $\rho = L/\rho$, $\rho\epsilon < 1$, and $\max_{k \in S} \sqrt{n_k} \cdot \frac{\sum_{k \in S} \sqrt{n_k}}{n_S} \leq \sqrt{\frac{|S| \max_{k \in S} n_k}{n_S}} := \kappa_w$, then

$$\|\hat{\boldsymbol{\theta}}_k - \boldsymbol{\theta}_k^*\|_2 \leq \frac{\|\sum_{k \in S} \omega_k \nabla f^{(k)}(\boldsymbol{\theta}_k^*)\|_2}{\rho \sum_{k \in S} \omega_k} + \frac{6}{1 - \rho\epsilon} \min \left\{ 3\rho^2 \kappa_w h, \frac{2\lambda}{5\rho \sqrt{\omega_k}} \right\} + \frac{\lambda\epsilon}{\rho \max_{k \in S} \sqrt{\omega_k}}.$$

*Furthermore, if we **also** have*

$$\lambda \geq \frac{15\rho\kappa_w L \max_{k \in S} \sqrt{\omega_k} h}{1 - \rho\epsilon},$$

then $\hat{\boldsymbol{\theta}}_k = \hat{\boldsymbol{\beta}}$ for all $k \in S$, and

$$\sup_{k \in S} \|\hat{\boldsymbol{\theta}}_k - \boldsymbol{\theta}_k^*\|_2 \leq \frac{\|\sum_{k \in S} \omega_k \nabla f^{(k)}(\boldsymbol{\theta}_k^*)\|_2}{\rho \sum_{k \in S} \omega_k} + 2\rho\kappa_w h + \frac{\lambda\epsilon}{\rho \max_{k \in S} \sqrt{\omega_k}}.$$

Lemma 7. *Suppose*

$$\hat{\boldsymbol{\theta}} = \arg \min_{\boldsymbol{\theta}} \left\{ f^{(0)}(\boldsymbol{\theta}) + \frac{\lambda}{\sqrt{n_0}} \|\hat{\boldsymbol{\theta}} - \bar{\boldsymbol{\theta}}\|_2 \right\}$$

with some $\bar{\boldsymbol{\theta}} \in \mathbb{R}^p$. Assume $f^{(0)}$ is convex and twice differentiable, and $\rho \mathbf{I}_p \leq \nabla^2 f^{(0)}(\boldsymbol{\theta}) \leq L \mathbf{I}_p$ for any $\boldsymbol{\theta} \in \mathbb{R}^p$. Then

$$(i) \quad \|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|_2 \leq \frac{\nabla f^{(0)}(\boldsymbol{\theta})}{\rho} + \frac{\lambda}{\rho \sqrt{n_0}}, \text{ for any } \boldsymbol{\theta}^* \in \mathbb{R}^p \text{ and } \lambda \geq 0;$$

$$(ii) \quad \hat{\boldsymbol{\theta}} = \bar{\boldsymbol{\theta}}, \text{ if } \lambda \geq 2 \|\nabla f^{(0)}(\boldsymbol{\theta}^*)\|_2 \sqrt{n_0} \text{ and } \|\bar{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|_2 \leq (\lambda/\sqrt{n_0} - \|\nabla f^{(0)}(\boldsymbol{\theta}^*)\|_2)/L.$$

E Proofs of Theorems in Main Text

E.1 Proofs of general lemmas

E.1.1 Proof of Lemma 2

The second half inequality is due to Lemma 1. It suffices to show the first half inequality. For any $\mathbf{x} = (x_1, \dots, x_{p-1})^T \in \mathcal{B}^{p-1}$, define $x_p = \sqrt{1 - \sum_{j=1}^{p-1} x_j^2}$. Then we can define a mapping

$$\mathbf{x} \in \mathcal{B}^{p-1} \mapsto \tilde{\mathbf{x}} = (\tilde{x}_1, \dots, \tilde{x}_{p-1}, \tilde{x}_p) \in \mathcal{S}^p,$$

with $\tilde{x}_j = x_j$ for $j \leq p-1$ and $\tilde{x}_p = \pm x_p$. And it's easy to see that for any $\mathbf{x}, \mathbf{y} \in \mathcal{B}^{p-1}$, we have $\|\mathbf{x} - \mathbf{y}\|_2 \leq \|\tilde{\mathbf{x}} - \tilde{\mathbf{y}}\|_2$. Therefore, if $\{\tilde{\mathbf{x}}_j\}_{j=1}^N$ is an ϵ -cover of \mathcal{S}^{p-1} under Euclidean norm, then it $\{\mathbf{x}_j\}_{j=1}^N$ must be an ϵ -cover of \mathcal{B}^{p-1} under Euclidean norm. Then

$$N(\epsilon, \mathcal{B}^{p-1}, \|\cdot\|_2) \leq N(\epsilon, \mathcal{S}^{p-1}, \|\cdot\|_2) \leq M(\epsilon, \mathcal{S}^{p-1}, \|\cdot\|_2).$$

E.1.2 Proof of Lemma 4

If $\{\mathbf{u}_j\}_{j=1}^N$ is an ϵ -packing of \mathcal{S}^{p-1} under Euclidean norm, then $\{\mathbf{u}_j\}_{j=1}^N \cap \mathcal{Q}_v$ must be an ϵ -packing of $\mathcal{S}^p \cap \mathcal{Q}_v$ under Euclidean norm. Then by Lemma 2,

$$\begin{aligned} 2^p \max_{v \in \{\pm 1\}^{\otimes p}} M(\epsilon, \mathcal{S}^{p-1} \cap \mathcal{Q}_v, \|\cdot\|_2) &\geq \sum_{v \in \{\pm 1\}^{\otimes p}} M(\epsilon, \mathcal{S}^{p-1} \cap \mathcal{Q}_v, \|\cdot\|_2) \\ &\geq M(\epsilon, \mathcal{S}^{p-1}, \|\cdot\|_2) \\ &\geq \left(\frac{1}{\epsilon}\right)^{p-1}, \end{aligned}$$

implying

$$\max_{v \in \{\pm 1\}^{\otimes p}} M(\epsilon, \mathcal{S}^p \cap \mathcal{Q}_v, \|\cdot\|_2) \geq \left(\frac{1}{2}\right)^p \left(\frac{1}{\epsilon}\right)^{p-1}.$$

E.1.3 Proof of Lemma 5

Suppose $Z_1 \sim \mathcal{N}(\mu_1, \sigma^2) \perp Z_2 \sim \mathcal{N}(\mu_2, \sigma^2)$, then we can write $Z = (1 - I)Z_1 + IZ_2 = (1 - I)(Z_1 - \mu_1) + I(Z_2 - \mu_2) + [\mu_1(1 - I) + \mu_2 I]$, where $I \sim \text{Bernoulli}(w)$ that is independent with Z_1 and Z_2 . Then

$$\begin{aligned} \mathbb{E}e^{\lambda Z} &\leq \mathbb{E}e^{\lambda(1-I)(Z_1 - \mu_1) + \lambda I(Z_2 - \mu_2)} \cdot \mathbb{E}e^{\lambda[\mu_1(1-I) + \mu_2 I]} \\ &\leq \mathbb{E}_I \left[(1 - I)\mathbb{E}e^{\lambda Z_1} + I\mathbb{E}e^{\lambda Z_2} \right] \cdot \mathbb{E}e^{\lambda[\mu_1(1-I) + \mu_2 I]} \\ &\leq \exp \left\{ \frac{1}{2}\sigma^2\lambda^2 + \frac{1}{8}(\mu_2 - \mu_1)^2\lambda^2 \right\}, \end{aligned}$$

where the last second inequality comes from Jensen's inequality and the independence between Z_1 , Z_2 and I . This completes the proof.

E.1.4 Proof of Lemma 6

The result follows from Theorem A.2, Lemma B.1 and Claim B.1 in [Duan and Wang \(2022\)](#).

E.2 Proof of Theorem 1

Define the contraction basin of one GMM as

$$B_{\text{con}}(\boldsymbol{\theta}^{(k)*}) = \left\{ \boldsymbol{\theta} = \{w, \boldsymbol{\beta}, \delta\} : w_r \in [c_w/2, 1 - c_w/2], \|\boldsymbol{\beta} - \boldsymbol{\beta}^{(k)*}\|_2 \leq C_b\Delta, \delta = \frac{1}{2}\boldsymbol{\beta}^T(\boldsymbol{\mu}_1 + \boldsymbol{\mu}_2) \right. \\ \left. \max_{r=1:2} \|\boldsymbol{\mu}_r - \boldsymbol{\mu}_r^*\|_2 \leq C_b\Delta \right\},$$

for which we may shorthand as B_{con} in the following. And given the index set S , two joint contraction basins are defined as

$$\begin{aligned} B_{\text{con}}^{J,1}(\{\boldsymbol{\theta}^{(k)*}\}_{k \in S}) &= \left\{ \{\boldsymbol{\theta}^{(k)}\}_{k \in S} = \{(w^{(k)}, \boldsymbol{\beta}^{(k)}, \delta^{(k)})\}_{k \in S} : \boldsymbol{\theta}^{(k)} \in B_{\text{con}}(\boldsymbol{\theta}^{(k)*}) \right\}, \\ B_{\text{con}}^{J,2}(\{\boldsymbol{\theta}^{(k)*}\}_{k \in S}) &= \left\{ \{\boldsymbol{\theta}^{(k)}\}_{k \in S} = \{(w^{(k)}, \boldsymbol{\beta}^{(k)}, \delta^{(k)})\}_{k \in S} : \boldsymbol{\theta}^{(k)} \in B_{\text{con}}(\boldsymbol{\theta}^{(k)*}), \boldsymbol{\beta}^{(k)} \equiv \bar{\boldsymbol{\beta}} \text{ for all } k \right\}. \end{aligned}$$

For simplicity, at some places, we will write them as $B_{\text{con}}^{J,1}$ and $B_{\text{con}}^{J,2}$, respectively.

For $\boldsymbol{\theta} = (w, \boldsymbol{\beta}, \delta)$ and $\boldsymbol{\theta}' = (w', \boldsymbol{\beta}', \delta')$, define

$$d(\boldsymbol{\theta}, \boldsymbol{\theta}') = |w - w'| \vee \|\boldsymbol{\beta} - \boldsymbol{\beta}'\|_2 \vee |\delta - \delta'|.$$

And denote the minimum SNR $\Delta = \min_{k \in S} \Delta^{(k)}$.

E.2.1 Lemmas

For GMM $\mathbf{z} \sim (1 - w^*)\mathcal{N}(\boldsymbol{\mu}_1^*, \boldsymbol{\Sigma}^*) + w^*\mathcal{N}(\boldsymbol{\mu}_2^*, \boldsymbol{\Sigma}^*)$ and any $\boldsymbol{\theta} = (w, \boldsymbol{\beta}, \delta)$, define

$$\begin{aligned}\gamma_{\boldsymbol{\theta}}(\mathbf{z}) &= \frac{w \exp\{\boldsymbol{\beta}^T \mathbf{z} - \delta\}}{1 - w + w \exp\{\boldsymbol{\beta}^T \mathbf{z} - \delta_r\}}, \quad w(\boldsymbol{\theta}) = \mathbb{E}[\gamma_{\boldsymbol{\theta}}(\mathbf{z})], \\ \boldsymbol{\mu}_1(\boldsymbol{\theta}) &= \frac{\mathbb{E}[(1 - \gamma_{\boldsymbol{\theta}}(\mathbf{z}))\mathbf{z}]}{\mathbb{E}[1 - \gamma_{\boldsymbol{\theta}}(\mathbf{z})]}, \quad \boldsymbol{\mu}_2(\boldsymbol{\theta}) = \frac{\mathbb{E}[\gamma_{\boldsymbol{\theta}}(\mathbf{z})\mathbf{z}]}{\mathbb{E}[\gamma_{\boldsymbol{\theta}}(\mathbf{z})]}.\end{aligned}$$

Lemma 8 (Contraction of binary GMMs, a special case of Lemma 37 when $R = 2$). *When $C_b \leq cc_{\boldsymbol{\Sigma}}^{-1/2}$ with a small constant $c > 0$ and $\Delta \geq CC_b M$ with a large constant $C > 0$, there exist positive constants $C' > 0$ and $C'' > 0$, for any $\boldsymbol{\theta} \in B_{\text{con}}(\boldsymbol{\theta}^{(k)*})$,*

$$|w_r(\boldsymbol{\theta}) - w_r^*| \leq C' \Delta \exp\{-C'' \Delta^2\} \cdot d(\boldsymbol{\theta}, \boldsymbol{\theta}^*), \quad \|\boldsymbol{\mu}_r(\boldsymbol{\theta}) - \boldsymbol{\mu}_r^*\|_2 \leq C' \Delta \exp\{-C'' \Delta^2\} \cdot d(\boldsymbol{\theta}, \boldsymbol{\theta}^*).$$

Hence when $\Delta > \text{some constant } C''$, we have $C \Delta \exp\{-C' \Delta^2\} \leq \kappa_0 < 1$, such that $C \Delta \exp\{-C' \Delta^2\}$ can be replaced by κ_0 in the inequalities above.

Lemma 9. *When $h \leq C_b \Delta$, $B_{\text{con}}^{J,2}(\{\boldsymbol{\theta}^{(k)*}\}_{k \in S}) \neq \emptyset$.*

Lemma 10 (Theorem 3 in Maurer and Pontil (2021)). *Let $f : \mathcal{X}^n \rightarrow \mathbb{R}$ and $X = (X_1, \dots, X_n)$ be a vector of independent random variables with values in a space \mathcal{X} . Then for any $t > 0$ we have*

$$\mathbb{P}(f(X) - \mathbb{E}f(X) > t) \leq \exp \left\{ -\frac{t^2}{32e \left\| \sum_{i=1}^n \|f_i(X)\|_{\psi_2}^2 \right\|_{\infty}} \right\},$$

where $f_i(X)$ as a random function of x is defined to be $(f_i(X))(x) := f(x_1, \dots, x_{i-1}, X_i, x_{i+1}, \dots, X_n) - \mathbb{E}_{X_i}[f(x_1, \dots, x_{i-1}, X_i, x_{i+1}, \dots, X_n)]$, the sub-Gaussian norm $\|Z\|_{\psi_2} := \sup_{d \geq 1} \{\|Z\|_d / \sqrt{d}\}$, and $\|Z\|_d = (\mathbb{E}|Z|^d)^{1/d}$.

Lemma 11. *Suppose Assumption 1 holds.*

(i) *With probability at least $1 - C'K^{-2}$,*

$$\sup_{\substack{\boldsymbol{\theta}^{(k)} \in B_{\text{con}} \\ \|\boldsymbol{\beta}^{(k)} - \boldsymbol{\beta}^{(k)*}\|_2 \leq \xi^{(k)}}} \left| \frac{1}{n_k} \sum_{i=1}^{n_k} \gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}_i^{(k)}) - \mathbb{E}[\gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}^{(k)})] \right| \lesssim \xi^{(k)} \sqrt{\frac{p}{n_k}} + \sqrt{\frac{\log K}{n_k}},$$

for all $k \in S$.

(ii) *With probability at least $1 - C'K^{-1}e^{-C''p}$,*

$$\sup_{\{\boldsymbol{\theta}^{(k)}\}_{k \in S} \in B_{\text{con}}^{J,2}} \sup_{|\tilde{w}_k| \leq 1} \frac{1}{n_S} \left| \sum_{k \in S} \tilde{w}_k \sum_{i=1}^{n_k} [\gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}_i^{(k)}) - \mathbb{E}[\gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}^{(k)})]] \right| \lesssim \sqrt{\frac{p + K}{n_S}}.$$

Lemma 12. Suppose Assumption 1 holds.

(i) With probability at least $1 - C'(K^{-2} + K^{-1}e^{-C''p})$,

$$\sup_{\substack{\boldsymbol{\theta}^{(k)} \in B_{con} \\ \|\boldsymbol{\beta}^{(k)} - \boldsymbol{\beta}^{(k)*}\|_2 \leq \xi^{(k)}}} \left| \frac{1}{n_k} \sum_{i=1}^{n_k} [1 - \gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}_i^{(k)})] (\mathbf{z}_i^{(k)})^T \boldsymbol{\beta}^{(k)*} - \mathbb{E}[[1 - \gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}^{(k)})] (\mathbf{z}^{(k)})^T \boldsymbol{\beta}^{(k)*}] \right|$$

$$\lesssim \xi^{(k)} \sqrt{\frac{p}{n_k}} + \sqrt{\frac{\log K}{n_k}},$$

for all $k \in S$.

(ii) With probability at least $1 - C'K^{-1}e^{-C''p}$,

$$\sup_{\{\boldsymbol{\theta}^{(k)}\}_{k \in S} \in B_{con}^{J,2}} \sup_{|\tilde{w}_k| \leq 1} \frac{1}{n_S} \left| \sum_{k \in S} \tilde{w}_k \sum_{i=1}^{n_k} \left[[1 - \gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}_i^{(k)})] (\mathbf{z}_i^{(k)})^T \boldsymbol{\beta}^{(k)*} - \mathbb{E}[[1 - \gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}^{(k)})] (\mathbf{z}^{(k)})^T \boldsymbol{\beta}^{(k)*}] \right] \right|$$

$$\lesssim \sqrt{\frac{p + K}{n_S}}.$$

Lemma 13. Suppose Assumption 1 holds.

(i) With probability at least $1 - C'(K^{-2} + K^{-1}e^{-C''p})$,

$$\sup_{\boldsymbol{\theta}^{(k)} \in B_{con}} \left\| \frac{1}{n_k} \sum_{i=1}^{n_k} \gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}_i^{(k)}) \mathbf{z}_i^{(k)} - \mathbb{E}[\gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}^{(k)}) \mathbf{z}^{(k)}] \right\|_2 \lesssim \sqrt{\frac{p + \log K}{n_k}},$$

for all $k \in S$.

(ii) With probability at least $1 - C'K^{-1}e^{-C''p}$,

$$\sup_{\{\boldsymbol{\theta}^{(k)}\}_{k \in S} \in B_{con}^{J,2}} \sup_{|\tilde{w}_k| \leq 1} \frac{1}{n_S} \left\| \sum_{k \in S} \tilde{w}_k \sum_{i=1}^{n_k} \left[\gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}_i^{(k)}) \mathbf{z}_i^{(k)} - \mathbb{E}[\gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}^{(k)}) \mathbf{z}^{(k)}] \right] \right\|_2 \lesssim \sqrt{\frac{p + K}{n_S}}.$$

(iii) With probability at least $1 - C'K^{-1}e^{-C''p}$,

$$\sup_{\{\boldsymbol{\theta}^{(k)}\}_{k \in S} \in B_{con}^{J,2}} \sup_{|\tilde{w}_k| \leq 1} \frac{1}{n_S} \left\| \sum_{k \in S} \tilde{w}_k \sum_{i=1}^{n_k} \left[\gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}_i^{(k)}) - \mathbb{E}[\gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}^{(k)})] \right] \boldsymbol{\mu}_1^{(k)*} \right\|_2 \lesssim \sqrt{\frac{p + K}{n_S}}.$$

Lemma 14. Suppose Assumption 1 holds.

(i) With probability at least $1 - C'(K^{-2} + K^{-1}e^{-C''p})$,

$$\left\| \frac{1}{n_k} \sum_{i=1}^{n_k} [\mathbf{z}_i^{(k)} (\mathbf{z}_i^{(k)})^T - \mathbb{E}[\mathbf{z}_i^{(k)} (\mathbf{z}_i^{(k)})^T]] \boldsymbol{\beta}^{(k)*} \right\|_2 \lesssim \sqrt{\frac{p + \log K}{n_k}},$$

for all $k \in S$.

(ii) With probability at least $1 - C'K^{-1}e^{-C''p}$,

$$\left\| \frac{1}{n_S} \sum_{k \in S} \sum_{i=1}^{n_k} [\mathbf{z}_i^{(k)} (\mathbf{z}_i^{(k)})^T - \mathbb{E}[\mathbf{z}_i^{(k)} (\mathbf{z}_i^{(k)})^T]] \boldsymbol{\beta}^{(k)*} \right\|_2 \lesssim \sqrt{\frac{p}{n_S}}.$$

E.2.2 Main proof of Theorem 1

WLOG, in Assumptions 1.(iii) and 1.(iv), we assume

- $\max_{k \in S} (\|\widehat{\boldsymbol{\beta}}^{(k)[0]} - \boldsymbol{\beta}^{(k)*}\|_2 \vee |\widehat{\delta}^{(k)[0]} - \delta^{(k)*}|) \leq C_3 \min_{k \in S} \Delta^{(k)}$, with a sufficiently small constant C_3 ;
- $\max_{k \in S} |\widehat{w}^{(k)[0]} - w^{(k)*}| \leq c_w/2$.

(I) Case 1: Let us consider the case that $h \geq C\sqrt{\frac{p + \log K}{\max_{k \in S} n_k}}$. Consider an event \mathcal{E} defined to be the intersection of the events in Lemmas 11.(i), 12.(i), 13.(i), and 14.(i), with $\xi^{(k)} =$ a large constant C , which satisfies $\mathbb{P}(\mathcal{E}) \geq 1 - C'(K^{-1} + e^{-C''p})$. Throughout the analysis in Case 1, we condition on \mathcal{E} , therefore all the arguments hold with probability at least $1 - C'(K^{-1} + e^{-C''p})$.

Consider the case $t = 1$. Lemma 6 tells us that when $\lambda^{[t]} \geq C \max_{k \in S} \{\sqrt{n_k} \|\widehat{\boldsymbol{\Sigma}}^{(k)[t]} \boldsymbol{\beta}^{(k)*} - (\widehat{\boldsymbol{\mu}}_2^{(k)[t]} - \widehat{\boldsymbol{\mu}}_1^{(k)[t]})\|_2\}$, we have

$$\|\widehat{\boldsymbol{\beta}}^{(k)[t]} - \boldsymbol{\beta}^{(k)*}\|_2 \lesssim \left\| \sum_{k \in S} \frac{n_k}{n_S} [\widehat{\boldsymbol{\Sigma}}^{(k)[t]} \boldsymbol{\beta}^{(k)*} - (\widehat{\boldsymbol{\mu}}_2^{(k)[t]} - \widehat{\boldsymbol{\mu}}_1^{(k)[t]})] \right\|_2 + h \wedge \frac{\lambda^{[t]}}{\sqrt{n_k}} + \epsilon \frac{\lambda^{[t]}}{\sqrt{n_k}}. \quad (\text{E.14})$$

And if further $\lambda^{[t]} \geq C \max_{k \in S} \sqrt{n_k} h$, we have (E.14) holds with $\widehat{\boldsymbol{\beta}}^{(k)[t]} = \overline{\boldsymbol{\beta}}^{[t]}$ for all $k \in S$. Note that

$$\|\widehat{\boldsymbol{\Sigma}}^{(k)[t]} \boldsymbol{\beta}^{(k)*} - (\widehat{\boldsymbol{\mu}}_2^{(k)[t]} - \widehat{\boldsymbol{\mu}}_1^{(k)[t]})\|_2 \leq \|(\widehat{\boldsymbol{\Sigma}}^{(k)[t]} - \boldsymbol{\Sigma}^{(k)*}) \boldsymbol{\beta}^{(k)*}\|_2 + \|\widehat{\boldsymbol{\mu}}_2^{(k)[t]} - \widehat{\boldsymbol{\mu}}_1^{(k)[t]} - \boldsymbol{\mu}_2^{(k)*} + \boldsymbol{\mu}_1^{(k)*}\|_2. \quad (\text{E.15})$$

And the first term on the RHS can be controlled as

$$\begin{aligned} & \|(\widehat{\boldsymbol{\Sigma}}^{(k)[t]} - \boldsymbol{\Sigma}^{(k)*}) \boldsymbol{\beta}^{(k)*}\|_2 \\ & \leq \underbrace{\left\| \frac{1}{n_k} \sum_{i=1}^{n_k} [\mathbf{z}_i^{(k)} (\mathbf{z}_i^{(k)})^T - \mathbb{E}[\mathbf{z}_i^{(k)} (\mathbf{z}_i^{(k)})^T]] \boldsymbol{\beta}^{(k)*} \right\|_2}_{\textcircled{1}} \end{aligned}$$

$$\begin{aligned}
& + \underbrace{\left\| \left[(1 - \widehat{w}^{(k)[t]}) \widehat{\boldsymbol{\mu}}_1^{(k)[t]} (\widehat{\boldsymbol{\mu}}_1^{(k)[t]})^T - (1 - w^{(k)*}) \boldsymbol{\mu}_1^{(k)*} (\boldsymbol{\mu}_1^{(k)*})^T \right] \boldsymbol{\beta}^{(k)*} \right\|_2}_{\textcircled{2}} \\
& + \underbrace{\left\| \left[\widehat{w}^{(k)[t]} \widehat{\boldsymbol{\mu}}_2^{(k)[t]} (\widehat{\boldsymbol{\mu}}_2^{(k)[t]})^T - w^{(k)*} \boldsymbol{\mu}_2^{(k)*} (\boldsymbol{\mu}_2^{(k)*})^T \right] \boldsymbol{\beta}^{(k)*} \right\|_2}_{\textcircled{3}}. \tag{E.16}
\end{aligned}$$

Conditioned on \mathcal{E} , we have

$$\textcircled{1} \lesssim \sqrt{\frac{p + \log K}{n_k}},$$

And

$$\begin{aligned}
\textcircled{2} & \leq \underbrace{\left\| (1 - \widehat{w}^{(k)[t]}) (\widehat{\boldsymbol{\mu}}_1^{(k)[t]} - \boldsymbol{\mu}_1^{(k)*}) \cdot (\widehat{\boldsymbol{\mu}}_1^{(k)[t]})^T \boldsymbol{\beta}^{(k)*} \right\|_2}_{\textcircled{2.1}} \\
& + \underbrace{\left\| \left[(1 - \widehat{w}^{(k)[t]}) \boldsymbol{\mu}_1^{(k)*} (\widehat{\boldsymbol{\mu}}_1^{(k)[t]})^T - (1 - w^{(k)*}) \boldsymbol{\mu}_1^{(k)*} (\boldsymbol{\mu}_1^{(k)*})^T \right] \boldsymbol{\beta}^{(k)*} \right\|_2}_{\textcircled{2.2}},
\end{aligned}$$

where

$$\textcircled{2.2} \leq \left\| (\widehat{w}^{(k)[t]} - w^{(k)*}) \boldsymbol{\mu}_1^{(k)*} (\widehat{\boldsymbol{\mu}}_1^{(k)[t]})^T \boldsymbol{\beta}^{(k)*} \right\|_2 + \left\| (1 - \widehat{w}^{(k)[t]}) \boldsymbol{\mu}_1^{(k)*} (\widehat{\boldsymbol{\mu}}_1^{(k)[t]} - \boldsymbol{\mu}_1^{(k)*})^T \boldsymbol{\beta}^{(k)*} \right\|_2. \tag{E.17}$$

Before we discuss how to control the terms on the RHS, let us first try to control $|\widehat{w}^{(k)[t]} - w^{(k)*}|$ as it will be used to bound the existing terms. Note that

$$\begin{aligned}
|\widehat{w}^{(k)[t]} - w^{(k)*}| & \leq |w^{(k)}(\widehat{\boldsymbol{\theta}}^{(k)[t-1]}) - w^{(k)*}| + |\widehat{w}^{(k)[t]} - w^{(k)}(\widehat{\boldsymbol{\theta}}^{(k)[t-1]})| \\
& \leq \kappa_0 d(\widehat{\boldsymbol{\theta}}^{(k)[t-1]}, \boldsymbol{\theta}^{(k)*}) + \left| \frac{1}{n_k} \sum_{i=1}^{n_k} \gamma_{\widehat{\boldsymbol{\theta}}^{(k)[t-1]}}(\mathbf{z}_i^{(k)}) - \mathbb{E}[\gamma_{\widehat{\boldsymbol{\theta}}^{(k)[t-1]}}(\mathbf{z}^{(k)})] \right| \\
& \lesssim \kappa_0 d(\widehat{\boldsymbol{\theta}}^{(k)[t-1]}, \boldsymbol{\theta}^{(k)*}) + \sqrt{\frac{p + \log K}{n_k}} \\
& \leq c, \tag{E.18}
\end{aligned}$$

where c is a small constant. And

$$\begin{aligned}
\|\widehat{\boldsymbol{\mu}}_1^{(k)[t]} - \boldsymbol{\mu}_1^{(k)*}\|_2 & = \left\| \frac{\frac{1}{n_k} \sum_{i=1}^{n_k} (1 - \gamma_{\widehat{\boldsymbol{\theta}}^{(k)[t-1]}}(\mathbf{z}_i^{(k)})) \mathbf{z}_i^{(k)}}{1 - \widehat{w}^{(k)[t]}} - \frac{\mathbb{E}[(1 - \gamma_{\widehat{\boldsymbol{\theta}}^{(k)[t-1]}}(\mathbf{z}^{(k)})) \mathbf{z}^{(k)}]}{1 - w^{(k)}(\widehat{\boldsymbol{\theta}}^{(k)[t-1]})} \right\|_2 \\
& \leq \left\| \frac{\frac{1}{n_k} \sum_{i=1}^{n_k} (1 - \gamma_{\widehat{\boldsymbol{\theta}}^{(k)[t-1]}}(\mathbf{z}_i^{(k)})) \mathbf{z}_i^{(k)} - \mathbb{E}[(1 - \gamma_{\widehat{\boldsymbol{\theta}}^{(k)[t-1]}}(\mathbf{z}^{(k)})) \mathbf{z}^{(k)}]}{1 - \widehat{w}^{(k)[t]}} \right\|_2 \\
& + \left\| \frac{\mathbb{E}[(1 - \gamma_{\widehat{\boldsymbol{\theta}}^{(k)[t-1]}}(\mathbf{z}^{(k)})) \mathbf{z}^{(k)}]}{(1 - \widehat{w}^{(k)[t]})(1 - w^{(k)}(\widehat{\boldsymbol{\theta}}^{(k)[t-1]}))} (\widehat{w}^{(k)[t]} - w^{(k)}(\widehat{\boldsymbol{\theta}}^{(k)[t-1]})) \right\|_2
\end{aligned}$$

$$\begin{aligned}
&\lesssim \left\| \frac{1}{n_k} \sum_{i=1}^{n_k} (1 - \gamma_{\hat{\boldsymbol{\theta}}^{(k)}[t-1]}(\mathbf{z}_i^{(k)})) \mathbf{z}_i^{(k)} - \mathbb{E}[(1 - \gamma_{\hat{\boldsymbol{\theta}}^{(k)}[t-1]}(\mathbf{z}^{(k)})) \mathbf{z}^{(k)}] \right\|_2 \\
&\quad + |\hat{w}^{(k)[t]} - w^{(k)*}| + \kappa_0 d(\hat{\boldsymbol{\theta}}^{(k)[t-1]}, \boldsymbol{\theta}^{(k)*}) \\
&\lesssim \kappa_0 d(\hat{\boldsymbol{\theta}}^{(k)[t-1]}, \boldsymbol{\theta}^{(k)*}) + \sqrt{\frac{p + \log K}{n_k}} \\
&\leq C_b \Delta.
\end{aligned}$$

Therefore, we can bound the RHS of (E.17) as

$$\textcircled{2}.2 \lesssim |\hat{w}^{(k)[t]} - w^{(k)*}| + \|\hat{\boldsymbol{\mu}}_1^{(k)[t]} - \boldsymbol{\mu}_1^{(k)*}\|_2 \lesssim \kappa_0 d(\hat{\boldsymbol{\theta}}^{(k)[t-1]}, \boldsymbol{\theta}^{(k)*}) + \sqrt{\frac{p + \log K}{n_k}}. \quad (\text{E.19})$$

Similarly, we have

$$\textcircled{2}.1 \lesssim \|\hat{\boldsymbol{\mu}}_1^{(k)[t]} - \boldsymbol{\mu}_1^{(k)*}\|_2 \lesssim \kappa_0 d(\hat{\boldsymbol{\theta}}^{(k)[t-1]}, \boldsymbol{\theta}^{(k)*}) + \sqrt{\frac{p + \log K}{n_k}}, \quad (\text{E.20})$$

Combining (E.19) and (E.20), we have

$$\textcircled{2} \lesssim \kappa_0 d(\hat{\boldsymbol{\theta}}^{(k)[t-1]}, \boldsymbol{\theta}^{(k)*}) + \sqrt{\frac{p + \log K}{n_k}},$$

Similarly, we can bound $\textcircled{3}$ in the same way, and get

$$\textcircled{3} \lesssim \kappa_0 d(\hat{\boldsymbol{\theta}}^{(k)[t-1]}, \boldsymbol{\theta}^{(k)*}) + \sqrt{\frac{p + \log K}{n_k}},$$

Hence

$$\|(\hat{\boldsymbol{\Sigma}}^{(k)[t]} - \boldsymbol{\Sigma}^{(k)*}) \boldsymbol{\beta}^{(k)*}\|_2 \lesssim \kappa_0 d(\hat{\boldsymbol{\theta}}^{(k)[t-1]}, \boldsymbol{\theta}^{(k)*}) + \sqrt{\frac{p + \log K}{n_k}},$$

And the second term on the RHS of (E.15) satisfies

$$\|\hat{\boldsymbol{\mu}}_2^{(k)[t]} - \hat{\boldsymbol{\mu}}_1^{(k)[t]} - \boldsymbol{\mu}_2^{(k)*} + \boldsymbol{\mu}_1^{(k)*}\|_2 \leq \|\hat{\boldsymbol{\mu}}_2^{(k)[t]} - \boldsymbol{\mu}_2^{(k)*}\|_2 \vee \|\hat{\boldsymbol{\mu}}_1^{(k)[t]} - \boldsymbol{\mu}_1^{(k)*}\|_2 \lesssim \kappa_0 d(\hat{\boldsymbol{\theta}}^{(k)[t-1]}, \boldsymbol{\theta}^{(k)*}) + \sqrt{\frac{p + \log K}{n_k}}. \quad (\text{E.21})$$

All together, we have

$$\|\hat{\boldsymbol{\Sigma}}^{(k)[t]} \boldsymbol{\beta}^{(k)*} - (\hat{\boldsymbol{\mu}}_2^{(k)[t]} - \hat{\boldsymbol{\mu}}_1^{(k)[t]})\|_2 \lesssim \kappa_0 d(\hat{\boldsymbol{\theta}}^{(k)[t-1]}, \boldsymbol{\theta}^{(k)*}) + \sqrt{\frac{p + \log K}{n_k}}. \quad (\text{E.22})$$

This implies that $\lambda^{[t]} = C_\lambda \sqrt{p + \log K} + \kappa \lambda^{[0]} \geq C \max_{k \in S} \{\sqrt{n_k} \|\hat{\boldsymbol{\Sigma}}^{(k)[t]} \boldsymbol{\beta}^{(k)*} - (\hat{\boldsymbol{\mu}}_2^{(k)[t]} - \hat{\boldsymbol{\mu}}_1^{(k)[t]})\|_2\}$, therefore by (E.14),

$$\|\hat{\boldsymbol{\beta}}^{(k)[t]} - \boldsymbol{\beta}^{(k)*}\|_2 \lesssim \kappa_0 \sum_{k \in S} \frac{n_k}{n_S} d(\hat{\boldsymbol{\theta}}^{(k)[t-1]}, \boldsymbol{\theta}^{(k)*}) + \sqrt{\frac{K(p + \log K)}{n_S}} + \lambda^{[t]}, \quad (\text{E.23})$$

And by (E.18),

$$\sum_{k \in S} \frac{n_k}{n_S} |\widehat{w}^{(k)[t]} - w^{(k)*}| \lesssim \kappa_0 \sum_{k \in S} \frac{n_k}{n_S} d(\widehat{\boldsymbol{\theta}}^{(k)[t-1]}, \boldsymbol{\theta}^{(k)*}) + \sqrt{\frac{K(p + \log K)}{n_S}}. \quad (\text{E.24})$$

Also,

$$\begin{aligned} |\widehat{\delta}^{(k)[t]} - \delta^{(k)*}| &= \frac{1}{2} \left\| (\widehat{\boldsymbol{\beta}}^{(k)[t]})^T (\widehat{\boldsymbol{\mu}}_1^{(k)[t]} + \widehat{\boldsymbol{\mu}}_2^{(k)[t]}) - (\boldsymbol{\beta}^{(k)*})^T (\boldsymbol{\mu}_1^{(k)*} + \boldsymbol{\mu}_2^{(k)*}) \right\|_2 \\ &\lesssim \|\widehat{\boldsymbol{\beta}}^{(k)[t]} - \boldsymbol{\beta}^{(k)*}\|_2 + \|\widehat{\boldsymbol{\mu}}_1^{(k)[t]} - \boldsymbol{\mu}_1^{(k)*}\|_2 + \|\widehat{\boldsymbol{\mu}}_2^{(k)[t]} - \boldsymbol{\mu}_2^{(k)*}\|_2 \\ &\lesssim \kappa_0 d(\widehat{\boldsymbol{\theta}}^{(k)[t-1]}, \boldsymbol{\theta}^{(k)*}) + \sqrt{\frac{p + \log K}{n_k}}, \end{aligned} \quad (\text{E.25})$$

which entails that

$$\sum_{k \in S} \frac{n_k}{n_S} |\widehat{\delta}^{(k)[t]} - \delta^{(k)*}| \lesssim \kappa_0 \sum_{k \in S} \frac{n_k}{n_S} d(\widehat{\boldsymbol{\theta}}^{(k)[t-1]}, \boldsymbol{\theta}^{(k)*}) + \sqrt{\frac{K(p + \log K)}{n_S}}. \quad (\text{E.26})$$

Combining (E.23), (E.24), and (E.26), we have

$$\sum_{k \in S} \frac{n_k}{n_S} d(\widehat{\boldsymbol{\theta}}^{(k)[t]}, \boldsymbol{\theta}^{(k)*}) \lesssim \kappa_0 \sum_{k \in S} \frac{n_k}{n_S} d(\widehat{\boldsymbol{\theta}}^{(k)[t-1]}, \boldsymbol{\theta}^{(k)*}) + \sqrt{\frac{K(p + \log K)}{n_S}} + \sqrt{\frac{K}{n_S}} \lambda^{[t]}. \quad (\text{E.27})$$

Also,

$$\max_{k \in S} \{\sqrt{n_k} d(\widehat{\boldsymbol{\theta}}^{(k)[t]}, \boldsymbol{\theta}^{(k)*})\} \lesssim \kappa_0 \max_{k \in S} \{\sqrt{n_k} d(\widehat{\boldsymbol{\theta}}^{(k)[t-1]}, \boldsymbol{\theta}^{(k)*})\} + \lambda^{[t]}. \quad (\text{E.28})$$

When we assume (E.23), (E.27), (E.28) hold for all $t = 1 : t'$, via the same analysis we will have (E.22) hold again for $t = t' + 1$. Hence

$$\max_{k \in S} \{\sqrt{n_k} \|\widehat{\boldsymbol{\Sigma}}^{(k)[t'+1]} \boldsymbol{\beta}^{(k)*} - (\widehat{\boldsymbol{\mu}}_2^{(k)[t'+1]} - \widehat{\boldsymbol{\mu}}_1^{(k)[t'+1]})\|_2\} \lesssim \kappa_0 \max_{k \in S} \{\sqrt{n_k} d(\widehat{\boldsymbol{\theta}}^{(k)[t']}, \boldsymbol{\theta}^{(k)*})\} + \sqrt{p + \log K}.$$

Then by (E.28) when $t = t'$,

$$\begin{aligned} \kappa_0 \max_{k \in S} \{\sqrt{n_k} d(\widehat{\boldsymbol{\theta}}^{(k)[t']}, \boldsymbol{\theta}^{(k)*})\} + \sqrt{p + \log K} &\lesssim \kappa_0^2 \max_{k \in S} \{\sqrt{n_k} d(\widehat{\boldsymbol{\theta}}^{(k)[t'-1]}, \boldsymbol{\theta}^{(k)*})\} + \sqrt{p + \log K} + \kappa_0 \lambda^{[t']} \\ &\leq \kappa_0 \lambda^{[t']} + \sqrt{p + \log K} \\ &\leq \lambda^{[t'+1]}, \end{aligned}$$

where we need $\kappa \geq C \kappa_0$ with a large constant $C > 0$. Recall that $\kappa \in (0, 1)$ is one of the tuning parameters in the update formula of $\lambda^{[t]}$. Therefore we can follow the same arguments as above to obtain (E.18), (E.21), (E.23), (E.25), (E.27), (E.28) for $t = t' + 1$.

So far, we have shown that (E.18), (E.21), (E.23), (E.25), (E.27), (E.28) hold for any t . By the update formula of $\lambda^{[t]}$, when $t \geq 1$, we have

$$\lambda^{[t]} = \frac{1 - \kappa^t}{1 - \kappa} C_\lambda \sqrt{p + \log K} + \kappa^{t-1} \lambda^{[0]}.$$

Therefore by (E.27),

$$\begin{aligned} \sum_{k \in S} \frac{n_k}{n_S} d(\widehat{\boldsymbol{\theta}}^{(k)[t]}, \boldsymbol{\theta}^{(k)*}) &\leq C\kappa_0 \sum_{k \in S} \frac{n_k}{n_S} d(\widehat{\boldsymbol{\theta}}^{(k)[t-1]}, \boldsymbol{\theta}^{(k)*}) + C' \sqrt{\frac{K(p + \log K)}{n_S}} + C' \sqrt{\frac{K}{n_S}} \lambda^{[t]} \\ &\leq (C\kappa_0)^t \sum_{k \in S} \frac{n_k}{n_S} d(\widehat{\boldsymbol{\theta}}^{(k)[0]}, \boldsymbol{\theta}^{(k)*}) + C' \sqrt{\frac{K(p + \log K)}{n_S}} + C' \sqrt{\frac{K}{n_S}} \sum_{t'=1}^t \lambda^{[t']} \cdot (C\kappa_0)^{t-t'} \\ &\leq (C\kappa_0)^t \sum_{k \in S} \frac{n_k}{n_S} d(\widehat{\boldsymbol{\theta}}^{(k)[0]}, \boldsymbol{\theta}^{(k)*}) + C' \sqrt{\frac{K(p + \log K)}{n_S}} + C' \sqrt{\frac{K}{n_S}} \sum_{t'=1}^t \lambda^{[t']} \cdot \kappa^{t-t'} \\ &\leq C'' t \kappa^t + C'' \sqrt{\frac{K(p + \log K)}{n_S}}. \end{aligned} \tag{E.29}$$

Consider an new event \mathcal{E}' defined to be the intersection of the events in Lemmas 11.(i), 12.(i), 13.(i), and 14.(i), with $\xi^{(k)} = C \sqrt{\frac{n_k}{\max_{k \in S} n_k}} + C \sqrt{\frac{\log K}{p}}$, which satisfies $\mathbb{P}(\mathcal{E}') \geq 1 - C'(K^{-1} + e^{-C''p})$. Throughout the following analysis in Case 1, we condition on $\mathcal{E} \cap \mathcal{E}'$, therefore all the arguments hold with probability at least $1 - C'(K^{-1} + e^{-C''p})$. When $h \geq C \sqrt{\frac{p + \log K}{\max_{k \in S} n_k}}$, since $n_S \gtrsim K \max_{k \in S} n_k$, we have $\sqrt{\frac{K(p + \log K)}{n_S}} \lesssim \sqrt{\frac{p + \log K}{\max_{k \in S} n_k}} \lesssim h \wedge \sqrt{\frac{p + \log K}{n_k}}$. Furthermore, when $t \geq C' \log \left(\frac{\max_{k \in S} n_k}{\min_{k \in S} n_k} \right) \wedge \log \left(\frac{p}{\log K} \vee 1 \right)$, we have $\xi^{(k)} \sqrt{\frac{p}{n_k}} \lesssim h \wedge \sqrt{\frac{p + \log K}{n_k}} + \epsilon \sqrt{\frac{p + \log K}{n_k}} + \sqrt{\frac{\log K}{n_k}}$ and $t \kappa^t + \sqrt{\frac{p + \log K}{n_k}} \leq \xi^{(k)}$.

Plugging (E.29) back into (E.23), we have

$$\begin{aligned} \|\widehat{\boldsymbol{\beta}}^{(k)[t]} - \boldsymbol{\beta}^{(k)*}\|_2 &\leq C'' t \kappa^t + C'' \sqrt{\frac{K(p + \log K)}{n_S}} + C'' \xi^{(k)} \sqrt{\frac{p}{n_k}} + C'' \sqrt{\frac{\log K}{n_k}} \\ &\leq C t \kappa^t + C \sqrt{\frac{K(p + \log K)}{n_S}} + C \cdot h \wedge \sqrt{\frac{p + \log K}{n_k}} + C \epsilon \sqrt{\frac{p + \log K}{n_k}} + C \sqrt{\frac{\log K}{n_k}} \\ &\leq C t \kappa^t + C \cdot h \wedge \sqrt{\frac{p + \log K}{n_k}} + C \epsilon \sqrt{\frac{p + \log K}{n_k}} + C \sqrt{\frac{\log K}{n_k}} \end{aligned} \tag{E.30}$$

Then by (E.18),

$$|\widehat{w}^{(k)[t]} - w^{(k)*}| \lesssim \kappa_0 d(\widehat{\boldsymbol{\theta}}^{(k)[t-1]}, \boldsymbol{\theta}^{(k)*}) + \xi^{(k)} \sqrt{\frac{p}{n_k}} + \sqrt{\frac{\log K}{n_k}}$$

$$\begin{aligned}
&\lesssim \kappa_0 \|\widehat{\boldsymbol{\beta}}^{(k)[t-1]} - \boldsymbol{\beta}^{(k)*}\|_2 + \kappa_0 |\widehat{w}^{(k)[t-1]} - w^{(k)*}| \vee |\widehat{\delta}^{(k)[t-1]} - \delta^{(k)*}| + \xi^{(k)} \sqrt{\frac{p}{n_k}} + \sqrt{\frac{\log K}{n_k}} \\
&\lesssim Ct\kappa^t + \kappa_0 |\widehat{w}^{(k)[t-1]} - w^{(k)*}| \vee |\widehat{\delta}^{(k)[t-1]} - \delta^{(k)*}| + h \wedge \sqrt{\frac{p + \log K}{n_k}} \\
&\quad + \epsilon \sqrt{\frac{p + \log K}{n_k}} + \sqrt{\frac{\log K}{n_k}}.
\end{aligned}$$

Similarly, by (E.25),

$$|\widehat{\delta}^{(k)[t-1]} - \delta^{(k)*}| \lesssim Ct\kappa^t + \kappa_0 |\widehat{w}^{(k)[t-1]} - w^{(k)*}| \vee |\widehat{\delta}^{(k)[t-1]} - \delta^{(k)*}| + h \wedge \sqrt{\frac{p + \log K}{n_k}} + \epsilon \sqrt{\frac{p + \log K}{n_k}} + \sqrt{\frac{\log K}{n_k}}.$$

Therefore,

$$\begin{aligned}
|\widehat{w}^{(k)[t]} - w^{(k)*}| \vee |\widehat{\delta}^{(k)[t]} - \delta^{(k)*}| &\leq Ct\kappa^t + C\kappa_0 |\widehat{w}^{(k)[t-1]} - w^{(k)*}| \vee |\widehat{\delta}^{(k)[t-1]} - \delta^{(k)*}| \\
&\quad + h \wedge \sqrt{\frac{p + \log K}{n_k}} + \epsilon \sqrt{\frac{p + \log K}{n_k}} + \sqrt{\frac{\log K}{n_k}} \\
&\leq C''t^2\kappa^t + C'' \cdot h \wedge \sqrt{\frac{p + \log K}{n_k}} + C''\epsilon \sqrt{\frac{p + \log K}{n_k}} + C'' \sqrt{\frac{\log K}{n_k}}.
\end{aligned}$$

Combine it with (E.30), we obtain that

$$d(\widehat{\boldsymbol{\theta}}^{(k)[t]}, \boldsymbol{\theta}^{(k)*}) \leq C''t^2\kappa^t + C'' \cdot h \wedge \sqrt{\frac{p + \log K}{n_k}} + C''\epsilon \sqrt{\frac{p + \log K}{n_k}} + C'' \sqrt{\frac{\log K}{n_k}}.$$

Plugging this back into (E.20), we get

$$\|\widehat{\boldsymbol{\mu}}_1^{(k)[t]} - \boldsymbol{\mu}_1^{(k)*}\|_2 \leq C''t^2\kappa^t + C'' \sqrt{\frac{p + \log K}{n_k}}.$$

And the same bound holds for $\|\widehat{\boldsymbol{\mu}}_2^{(k)[t]} - \boldsymbol{\mu}_2^{(k)*}\|_2$ as well. The same bound for $\|\widehat{\boldsymbol{\Sigma}}^{(k)[t]} - \boldsymbol{\Sigma}^{(k)*}\|_2$ can be obtained in the same spirit as in (E.16).

(II) Case 2: We now focus on the case that $h \leq C \sqrt{\frac{p + \log K}{\max_{k \in S} n_k}}$. Similar to Case 1, we consider an event \mathcal{E} defined to be the intersection of the events in Lemmas 11, 12, 13, and 14, with $\xi^{(k)} =$ a large constant C , which satisfies $\mathbb{P}(\mathcal{E}) \geq 1 - C'(K^{-1} + e^{-C''p})$. Throughout the analysis in Case 2, we condition on \mathcal{E} , therefore all the arguments hold with probability at least $1 - C'(K^{-1} + e^{-C''p})$.

Consider t_0 that satisfies $t_0^2\kappa^{t_0} \asymp \sqrt{\frac{K(p + \log K)}{n_S}}$. When $t = 1 : t_0$, we can go through the same analysis as in Case 1, and show that conditioned on \mathcal{E} ,

$$\sum_{k \in S} \frac{n_k}{n_S} d(\widehat{\boldsymbol{\theta}}^{(k)[t]}, \boldsymbol{\theta}^{(k)*}) \lesssim C''t\kappa^t + C'' \sqrt{\frac{K(p + \log K)}{n_S}},$$

and

$$d(\widehat{\boldsymbol{\theta}}^{(k)[t]}, \boldsymbol{\theta}^{(k)*}) \leq C'' t^2 \kappa^t + C'' \sqrt{\frac{K(p + \log K)}{n_S}} + C'' \cdot h \wedge \sqrt{\frac{p + \log K}{n_k}} + C'' \epsilon \sqrt{\frac{p + \log K}{n_k}} + C'' \sqrt{\frac{\log K}{n_k}},$$

$$\|\widehat{\boldsymbol{\mu}}_1^{(k)[t]} - \boldsymbol{\mu}_1^{(k)*}\|_2 \vee \|\widehat{\boldsymbol{\mu}}_2^{(k)[t]} - \boldsymbol{\mu}_2^{(k)*}\|_2 \vee \|\widehat{\boldsymbol{\Sigma}}^{(k)[t]} - \boldsymbol{\Sigma}^{(k)*}\|_2 \leq C'' t^2 \kappa^t + C'' \sqrt{\frac{p + \log K}{n_k}}.$$

Since $t_0^2 \kappa^{t_0} \asymp \sqrt{\frac{K(p + \log K)}{n_S}}$, the rates above are the desired rates. In the following, we will derive the results for the case $t \geq t_0 + 1$.

Define

$$\begin{aligned} \xi_{t_0}^{(k)} &= C'' \sqrt{\frac{K(p + \log K)}{n_S}} + C'' \cdot h \wedge \sqrt{\frac{p + \log K}{n_k}} + C'' \epsilon \sqrt{\frac{p + \log K}{n_k}} + C'' \sqrt{\frac{\log K}{n_k}}, \\ \bar{\xi}_{t_0} &= \sum_{k \in S} \frac{n_k}{n_S} \xi_{t_0}^{(k)} \leq C'' \sqrt{\frac{K(p + \log K)}{n_S}} + C'' \cdot h \wedge \sqrt{\frac{K(p + \log K)}{n_S}} + C'' \epsilon \sqrt{\frac{K(p + \log K)}{n_S}} \\ &\quad + C'' \sqrt{\frac{K \log K}{n_S}}. \end{aligned}$$

Consider an event \mathcal{E}_{t_0} defined to be the intersection of the events in Lemmas 11, 12, 13, and 14, with $\xi^{(k)} = \xi_{t_0}^{(k)}$, which satisfies $\mathbb{P}(\mathcal{E}_{t_0}) \geq 1 - C''(K^{-1} + e^{-C''p})$. In the following, we condition on $\mathcal{E} \cap \mathcal{E}_{t_0}$, therefore all the arguments hold with probability at least $1 - C''(K^{-1} + e^{-C''p})$.

Let $t = t_0 + 1$. Since $\lambda^{[t-1]} \geq C\sqrt{p + \log K} \geq C \max_{k \in S} \sqrt{n_k} h$, by Lemma 6, we also have $\widehat{\boldsymbol{\beta}}^{(k)[t-1]} = \bar{\boldsymbol{\beta}}^{[t-1]}$ for all $k \in S$. Similar to (E.18),

$$\begin{aligned} |\widehat{w}^{(k)[t]} - w^{(k)*}| &\leq |w^{(k)}(\widehat{\boldsymbol{\theta}}^{(k)[t-1]}) - w^{(k)*}| + |\widehat{w}^{(k)[t]} - w^{(k)}(\widehat{\boldsymbol{\theta}}^{(k)[t-1]})| \\ &\leq \kappa_0 d(\widehat{\boldsymbol{\theta}}^{(k)[t-1]}, \boldsymbol{\theta}^{(k)*}) + \left| \frac{1}{n_k} \sum_{i=1}^{n_k} \gamma_{\widehat{\boldsymbol{\theta}}^{(k)[t-1]}}(\mathbf{z}_i^{(k)}) - \mathbb{E}[\gamma_{\widehat{\boldsymbol{\theta}}^{(k)[t-1]}}(\mathbf{z}^{(k)})] \right| \\ &\leq \kappa_0 d(\widehat{\boldsymbol{\theta}}^{(k)[t-1]}, \boldsymbol{\theta}^{(k)*}) + C'' \xi_{t-1}^{(k)} \sqrt{\frac{p}{n_k}} + C'' \sqrt{\frac{\log K}{n_k}} \\ &\leq \kappa_0 d(\widehat{\boldsymbol{\theta}}^{(k)[t-1]}, \boldsymbol{\theta}^{(k)*}) + \kappa_0 \xi_{t-1}^{(k)} + C'' \sqrt{\frac{\log K}{n_k}}. \end{aligned}$$

This implies that

$$\sum_{k \in S} \frac{n_k}{n_S} |\widehat{w}^{(k)[t]} - w^{(k)*}| \leq \kappa_0 \sum_{k \in S} \frac{n_k}{n_S} d(\widehat{\boldsymbol{\theta}}^{(k)[t-1]}, \boldsymbol{\theta}^{(k)*}) + \kappa_0 \sum_{k \in S} \frac{n_k}{n_S} \xi_{t-1}^{(k)} + C'' \sqrt{\frac{K \log K}{n_S}}.$$

And by Lemma 6,

$$\begin{aligned}
\|\widehat{\boldsymbol{\beta}}^{(k)[t]} - \boldsymbol{\beta}^{(k)*}\|_2 &\lesssim \left\| \sum_{k \in S} \frac{n_k}{n_S} [\widehat{\boldsymbol{\Sigma}}^{(k)[t]} \boldsymbol{\beta}^{(k)*} - (\widehat{\boldsymbol{\mu}}_2^{(k)[t]} - \widehat{\boldsymbol{\mu}}_1^{(k)[t]})] \right\|_2 + h \wedge \frac{\lambda^{[t]}}{\sqrt{n_k}} + \epsilon \frac{\lambda^{[t]}}{\sqrt{n_k}} \\
&\lesssim \left\| \sum_{k \in S} \frac{n_k}{n_S} [\widehat{\boldsymbol{\Sigma}}^{(k)[t]} \boldsymbol{\beta}^{(k)*} - (\widehat{\boldsymbol{\mu}}_2^{(k)[t]} - \widehat{\boldsymbol{\mu}}_1^{(k)[t]})] \right\|_2 + h \wedge \sqrt{\frac{p + \log K}{n_k}} + \epsilon \sqrt{\frac{p + \log K}{n_k}},
\end{aligned} \tag{E.31}$$

where

$$\begin{aligned}
\left\| \sum_{k \in S} \frac{n_k}{n_S} [\widehat{\boldsymbol{\Sigma}}^{(k)[t]} \boldsymbol{\beta}^{(k)*} - (\widehat{\boldsymbol{\mu}}_2^{(k)[t]} - \widehat{\boldsymbol{\mu}}_1^{(k)[t]})] \right\|_2 &\leq \left\| \sum_{k \in S} \frac{n_k}{n_S} (\widehat{\boldsymbol{\Sigma}}^{(k)[t]} - \boldsymbol{\Sigma}^{(k)*}) \boldsymbol{\beta}^{(k)*} \right\|_2 \\
&\quad + \left\| \sum_{k \in S} \frac{n_k}{n_S} (\widehat{\boldsymbol{\mu}}_2^{(k)[t]} - \widehat{\boldsymbol{\mu}}_1^{(k)[t]} - \boldsymbol{\mu}_2^{(k)*} + \boldsymbol{\mu}_1^{(k)*}) \right\|_2.
\end{aligned}$$

And the first term on the RHS can be controlled as

$$\begin{aligned}
&\left\| \sum_{k \in S} \frac{n_k}{n_S} (\widehat{\boldsymbol{\Sigma}}^{(k)[t]} - \boldsymbol{\Sigma}^{(k)*}) \boldsymbol{\beta}^{(k)*} \right\|_2 \\
&\leq \underbrace{\left\| \frac{1}{n_S} \sum_{k \in S} \sum_{i=1}^{n_k} [\mathbf{z}_i^{(k)} (\mathbf{z}_i^{(k)})^T - \mathbb{E}[\mathbf{z}_i^{(k)} (\mathbf{z}_i^{(k)})^T]] \boldsymbol{\beta}^{(k)*} \right\|_2}_{\textcircled{4}} \\
&\quad + \underbrace{\left\| \sum_{k \in S} \frac{n_k}{n_S} [(1 - \widehat{w}^{(k)[t]}) \widehat{\boldsymbol{\mu}}_1^{(k)[t]} (\widehat{\boldsymbol{\mu}}_1^{(k)[t]})^T - (1 - w^{(k)*}) \boldsymbol{\mu}_1^{(k)*} (\boldsymbol{\mu}_1^{(k)*})^T] \boldsymbol{\beta}^{(k)*} \right\|_2}_{\textcircled{5}} \\
&\quad + \underbrace{\left\| \sum_{k \in S} \frac{n_k}{n_S} [\widehat{w}^{(k)[t]} \widehat{\boldsymbol{\mu}}_2^{(k)[t]} (\widehat{\boldsymbol{\mu}}_2^{(k)[t]})^T - w^{(k)*} \boldsymbol{\mu}_2^{(k)*} (\boldsymbol{\mu}_2^{(k)*})^T] \boldsymbol{\beta}^{(k)*} \right\|_2}_{\textcircled{6}}.
\end{aligned}$$

Conditioned on \mathcal{E} , we have

$$\textcircled{4} \lesssim \sqrt{\frac{p}{n_S}}.$$

And

$$\textcircled{5} \leq \underbrace{\left\| \sum_{k \in S} \frac{n_k}{n_S} (1 - \widehat{w}^{(k)[t]}) (\widehat{\boldsymbol{\mu}}_1^{(k)[t]} - \boldsymbol{\mu}_1^{(k)*}) \cdot (\widehat{\boldsymbol{\mu}}_1^{(k)[t]})^T \boldsymbol{\beta}^{(k)*} \right\|_2}_{\textcircled{5.1}}$$

$$+ \underbrace{\left\| \sum_{k \in S} \frac{n_k}{n_S} [(1 - \hat{w}^{(k)[t]}) \boldsymbol{\mu}_1^{(k)*} (\hat{\boldsymbol{\mu}}_1^{(k)[t]})^T - (1 - w^{(k)*}) \boldsymbol{\mu}_1^{(k)*} (\boldsymbol{\mu}_1^{(k)*})^T] \boldsymbol{\beta}^{(k)*} \right\|_2}_{\textcircled{5}.2},$$

where

$$\textcircled{5}.2 \leq \left\| \sum_{k \in S} \frac{n_k}{n_S} (\hat{w}^{(k)[t]} - w^{(k)*}) \boldsymbol{\mu}_1^{(k)*} (\hat{\boldsymbol{\mu}}_1^{(k)[t]})^T \boldsymbol{\beta}^{(k)*} \right\|_2 + \left\| \sum_{k \in S} \frac{n_k}{n_S} (1 - \hat{w}^{(k)[t]}) \boldsymbol{\mu}_1^{(k)*} (\hat{\boldsymbol{\mu}}_1^{(k)[t]} - \boldsymbol{\mu}_1^{(k)*})^T \boldsymbol{\beta}^{(k)*} \right\|_2.$$

For the first term, we have

$$\begin{aligned} & \left\| \sum_{k \in S} \frac{n_k}{n_S} (\hat{w}^{(k)[t]} - w^{(k)*}) \boldsymbol{\mu}_1^{(k)*} (\hat{\boldsymbol{\mu}}_1^{(k)[t]})^T \boldsymbol{\beta}^{(k)*} \right\|_2 \\ & \lesssim \sum_{k \in S} \frac{n_k}{n_S} |w^{(k)}(\hat{\boldsymbol{\theta}}^{(k)[t-1]}) - w^{(k)*}| \\ & \quad + \frac{1}{n_S} \left\| \sum_{k \in S} \sum_{i=1}^{n_k} \left\{ \gamma_{\hat{\boldsymbol{\theta}}^{(k)[t-1]}}(\mathbf{z}_i^{(k)}) \boldsymbol{\mu}_1^{(k)*} (\hat{\boldsymbol{\mu}}_1^{(k)[t]})^T \boldsymbol{\beta}^{(k)*} - \mathbb{E}[\gamma_{\hat{\boldsymbol{\theta}}^{(k)[t-1]}}(\mathbf{z}^{(k)}) \boldsymbol{\mu}_1^{(k)*} (\hat{\boldsymbol{\mu}}_1^{(k)[t]})^T \boldsymbol{\beta}^{(k)*}] \right\} \right\|_2 \\ & \lesssim \kappa_0 \sum_{k \in S} \frac{n_k}{n_S} d(\hat{\boldsymbol{\theta}}^{(k)[t-1]}, \boldsymbol{\theta}^{(k)*}) \\ & \quad + \frac{1}{n_S} \sup_{|\tilde{w}_k| \leq U} \left\| \sum_{k \in S} \sum_{i=1}^{n_k} \tilde{w}_k \left\{ \gamma_{\hat{\boldsymbol{\theta}}^{(k)[t-1]}}(\mathbf{z}_i^{(k)}) \boldsymbol{\mu}_1^{(k)*} - \mathbb{E}[\gamma_{\hat{\boldsymbol{\theta}}^{(k)[t-1]}}(\mathbf{z}^{(k)}) \boldsymbol{\mu}_1^{(k)*}] \right\} \right\|_2, \end{aligned}$$

where $U > 0$ is some constant. By Lemma 13.(iii) and the definition of \mathcal{E}_{t_0} , the second term can be bounded as

$$\frac{1}{n_S} \sup_{|\tilde{w}_k| \leq U} \left\| \sum_{k \in S} \sum_{i=1}^{n_k} \tilde{w}_k \left\{ \gamma_{\hat{\boldsymbol{\theta}}^{(k)[t-1]}}(\mathbf{z}_i^{(k)}) \boldsymbol{\mu}_1^{(k)*} - \mathbb{E}[\gamma_{\hat{\boldsymbol{\theta}}^{(k)[t-1]}}(\mathbf{z}^{(k)}) \boldsymbol{\mu}_1^{(k)*}] \right\} \right\|_2 \lesssim \sqrt{\frac{p+K}{n_S}}.$$

Therefore,

$$\left\| \sum_{k \in S} \frac{n_k}{n_S} (\hat{w}^{(k)[t]} - w^{(k)*}) \boldsymbol{\mu}_1^{(k)*} (\hat{\boldsymbol{\mu}}_1^{(k)[t]})^T \boldsymbol{\beta}^{(k)*} \right\|_2 \lesssim \kappa_0 \sum_{k \in S} \frac{n_k}{n_S} d(\hat{\boldsymbol{\theta}}^{(k)[t-1]}, \boldsymbol{\theta}^{(k)*}) + \sqrt{\frac{p+K}{n_S}}.$$

On the other hand, by simple calculations, we have

$$\begin{aligned} & \left\| \sum_{k \in S} \frac{n_k}{n_S} (1 - \hat{w}^{(k)[t]}) \boldsymbol{\mu}_1^{(k)*} (\hat{\boldsymbol{\mu}}_1^{(k)[t]} - \boldsymbol{\mu}_1^{(k)*})^T \boldsymbol{\beta}^{(k)*} \right\|_2 \\ & \lesssim \kappa_0 \sum_{k \in S} \frac{n_k}{n_S} d(\hat{\boldsymbol{\theta}}^{(k)[t-1]}, \boldsymbol{\theta}^{(k)*}) + \sup_{|\tilde{w}_k| \leq U} \left\| \sum_{k \in S} \frac{n_k}{n_S} \tilde{w}_k [w^{(k)}(\hat{\boldsymbol{\theta}}^{(k)[t-1]}) - \hat{w}^{(k)[t]}] \boldsymbol{\mu}_1^{(k)*} \right\|_2 \end{aligned}$$

$$\begin{aligned}
& + \frac{1}{n_S} \sup_{|\tilde{w}_k| \leq U'} \left\| \sum_{k \in S} \tilde{w}_k \sum_{i=1}^{n_k} \{ \gamma_{\hat{\boldsymbol{\theta}}^{(k)}[t-1]}(\mathbf{z}_i^{(k)}) (\mathbf{z}_i^{(k)})^T \boldsymbol{\beta}^{(k)*} - \mathbb{E}[\gamma_{\hat{\boldsymbol{\theta}}^{(k)}[t-1]}(\mathbf{z}^{(k)}) (\mathbf{z}^{(k)})^T \boldsymbol{\beta}^{(k)*}] \} \boldsymbol{\mu}_1^{(k)*} \right\|_2 \\
& \lesssim \kappa_0 \sum_{k \in S} \frac{n_k}{n_S} d(\hat{\boldsymbol{\theta}}^{(k)}[t-1], \boldsymbol{\theta}^{(k)*}) + \sup_{|\tilde{w}_k| \leq U} \left\| \sum_{k \in S} \frac{n_k}{n_S} \tilde{w}_k [w^{(k)}(\hat{\boldsymbol{\theta}}^{(k)}[t-1]) - \hat{w}^{(k)}[t]] \boldsymbol{\mu}_1^{(k)*} \right\|_2 \\
& + \frac{1}{n_S} \sup_{|\tilde{w}_k| \leq U'} \left\| \sum_{k \in S} \tilde{w}_k \sum_{i=1}^{n_k} \{ \gamma_{\hat{\boldsymbol{\theta}}^{(k)}[t-1]}(\mathbf{z}_i^{(k)}) (\mathbf{z}_i^{(k)})^T \boldsymbol{\beta}^{(k)*} - \mathbb{E}[\gamma_{\hat{\boldsymbol{\theta}}^{(k)}[t-1]}(\mathbf{z}^{(k)}) (\mathbf{z}^{(k)})^T \boldsymbol{\beta}^{(k)*}] \} \boldsymbol{\mu}_1^{(k)*} \right\|_2 \\
& \lesssim \kappa_0 \sum_{k \in S} \frac{n_k}{n_S} d(\hat{\boldsymbol{\theta}}^{(k)}[t-1], \boldsymbol{\theta}^{(k)*}) + \frac{1}{n_S} \sup_{|\tilde{w}_k| \leq U} \left\| \sum_{k \in S} \tilde{w}_k \sum_{i=1}^{n_k} [\gamma_{\hat{\boldsymbol{\theta}}^{(k)}[t-1]}(\mathbf{z}_i^{(k)}) - \mathbb{E}[\gamma_{\hat{\boldsymbol{\theta}}^{(k)}[t-1]}(\mathbf{z}^{(k)})]] \boldsymbol{\mu}_1^{(k)*} \right\|_2 \\
& + \frac{1}{n_S} \sup_{|\tilde{w}_k| \leq U'} \sup_{\|u\|_2 \leq 1} \left| \sum_{k \in S} \tilde{w}_k \sum_{i=1}^{n_k} \{ \gamma_{\hat{\boldsymbol{\theta}}^{(k)}[t-1]}(\mathbf{z}_i^{(k)}) (\mathbf{z}_i^{(k)})^T \boldsymbol{\beta}^{(k)*} - \mathbb{E}[\gamma_{\hat{\boldsymbol{\theta}}^{(k)}[t-1]}(\mathbf{z}^{(k)}) (\mathbf{z}^{(k)})^T \boldsymbol{\beta}^{(k)*}] \} (\boldsymbol{\mu}_1^{(k)*})^T \mathbf{u} \right| \\
& \lesssim \kappa_0 \sum_{k \in S} \frac{n_k}{n_S} d(\hat{\boldsymbol{\theta}}^{(k)}[t-1], \boldsymbol{\theta}^{(k)*}) + \sqrt{\frac{p+K}{n_S}} \\
& + \frac{1}{n_S} \sup_{|\tilde{w}_k| \leq U'} \left| \sum_{k \in S} \tilde{w}_k \sum_{i=1}^{n_k} \{ \gamma_{\hat{\boldsymbol{\theta}}^{(k)}[t-1]}(\mathbf{z}_i^{(k)}) (\mathbf{z}_i^{(k)})^T \boldsymbol{\beta}^{(k)*} - \mathbb{E}[\gamma_{\hat{\boldsymbol{\theta}}^{(k)}[t-1]}(\mathbf{z}^{(k)}) (\mathbf{z}^{(k)})^T \boldsymbol{\beta}^{(k)*}] \} \right| \\
& \lesssim \kappa_0 \sum_{k \in S} \frac{n_k}{n_S} d(\hat{\boldsymbol{\theta}}^{(k)}[t-1], \boldsymbol{\theta}^{(k)*}) + \sqrt{\frac{p+K}{n_S}}.
\end{aligned}$$

Hence

$$\textcircled{5}.2 \lesssim \kappa_0 \sum_{k \in S} \frac{n_k}{n_S} d(\hat{\boldsymbol{\theta}}^{(k)}[t-1], \boldsymbol{\theta}^{(k)*}) + \sqrt{\frac{p+K}{n_S}}.$$

A similar discussion leads to the same bound for $\textcircled{5}.1$. Therefore,

$$\textcircled{5} \lesssim \kappa_0 \sum_{k \in S} \frac{n_k}{n_S} d(\hat{\boldsymbol{\theta}}^{(k)}[t-1], \boldsymbol{\theta}^{(k)*}) + \sqrt{\frac{p+K}{n_S}}.$$

And the same bound holds for $\textcircled{6}$, which can be shown in the same spirit. Putting all the pieces together,

$$\left\| \sum_{k \in S} \frac{n_k}{n_S} (\hat{\boldsymbol{\Sigma}}^{(k)}[t] - \boldsymbol{\Sigma}^{(k)*}) \boldsymbol{\beta}^{(k)*} \right\|_2 \lesssim \kappa_0 \sum_{k \in S} \frac{n_k}{n_S} d(\hat{\boldsymbol{\theta}}^{(k)}[t-1], \boldsymbol{\theta}^{(k)*}) + \sqrt{\frac{p+K}{n_S}}.$$

Therefore by Lemma 6 and (E.31), we have $\widehat{\beta}^{(k)[t-1]} = \bar{\beta}^{[t-1]}$ for all $k \in S$, and

$$\begin{aligned}
\|\widehat{\beta}^{(k)[t]} - \beta^{(k)*}\|_2 &\lesssim \kappa_0 \sum_{k \in S} \frac{n_k}{n_S} d(\widehat{\theta}^{(k)[t-1]}, \theta^{(k)*}) + \sqrt{\frac{p+K}{n_S}} + h \wedge \sqrt{\frac{p+\log K}{n_k}} + \epsilon \sqrt{\frac{p+\log K}{n_k}} \\
&\leq C\kappa_0 \cdot \sum_{k \in S} \frac{n_k}{n_S} \xi_{t-1}^{(k)} + \sqrt{\frac{p+K}{n_S}} + h \wedge \sqrt{\frac{p+\log K}{n_k}} + \epsilon \sqrt{\frac{p+\log K}{n_k}} \\
&\leq \kappa'_0 \bar{\xi}_{t-1} + C' \sqrt{\frac{p}{n_S}} + C' \cdot h \wedge \sqrt{\frac{p+\log K}{n_k}} + C' \epsilon \sqrt{\frac{p+\log K}{n_k}} + C' \sqrt{\frac{K \log K}{n_S}} \\
&:= \xi_t^{(k)}.
\end{aligned} \tag{E.32}$$

This entails that

$$\bar{\xi}_t = \sum_{k \in S} \frac{n_k}{n_S} \xi_t^{(k)} = \kappa'_0 \bar{\xi}_{t-1} + C' \sqrt{\frac{p}{n_S}} + C' \cdot h \wedge \sqrt{\frac{K(p+\log K)}{n_S}} + C' \epsilon \sqrt{\frac{K(p+\log K)}{n_S}} + C' \sqrt{\frac{K \log K}{n_S}} \tag{E.33}$$

This implies that

$$\sum_{k \in S} \frac{n_k}{n_S} \|\widehat{\beta}^{(k)[t]} - \beta^{(k)*}\|_2 \lesssim \kappa_0 \sum_{k \in S} \frac{n_k}{n_S} d(\widehat{\theta}^{(k)[t-1]}, \theta^{(k)*}) + \sqrt{\frac{p+K}{n_S}} + h \wedge \sqrt{\frac{K(p+\log K)}{n_S}} + \epsilon \sqrt{\frac{K(p+\log K)}{n_S}}.$$

Also,

$$\begin{aligned}
|\widehat{\delta}^{(k)[t]} - \delta^{(k)*}| &= \frac{1}{2} \left\| (\widehat{\beta}^{(k)[t]})^T (\widehat{\mu}_1^{(k)[t]} + \widehat{\mu}_2^{(k)[t]}) - (\beta^{(k)*})^T (\mu_1^{(k)*} + \mu_2^{(k)*}) \right\|_2 \\
&\lesssim \|\widehat{\beta}^{(k)[t]} - \beta^{(k)*}\|_2 + \|(\beta^{(k)*})^T (\widehat{\mu}_1^{(k)[t]} - \mu_1^{(k)*})\|_2 + \|(\beta^{(k)*})^T (\widehat{\mu}_2^{(k)[t]} - \mu_2^{(k)*})\|_2 \\
&\lesssim \kappa_0 \sum_{k \in S} \frac{n_k}{n_S} d(\widehat{\theta}^{(k)[t-1]}, \theta^{(k)*}) + \sqrt{\frac{p+K}{n_S}} + h \wedge \sqrt{\frac{p+\log K}{n_k}} + \epsilon \sqrt{\frac{p+\log K}{n_k}} \\
&\quad + \left\| \frac{1}{n_k} \sum_{i=1}^{n_k} \gamma_{\widehat{\theta}^{(k)[t-1]}}(\mathbf{z}_i^{(k)}) (\beta^{(k)*})^T \mathbf{z}_i^{(k)} - \mathbb{E}[\gamma_{\widehat{\theta}^{(k)[t-1]}}(\mathbf{z}^{(k)}) (\beta^{(k)*})^T \mathbf{z}^{(k)}] \right\|_2 \\
&\lesssim \kappa_0 \sum_{k \in S} \frac{n_k}{n_S} d(\widehat{\theta}^{(k)[t-1]}, \theta^{(k)*}) + \sqrt{\frac{p+K}{n_S}} + h \wedge \sqrt{\frac{p+\log K}{n_k}} + \epsilon \sqrt{\frac{p+\log K}{n_k}} \\
&\quad + \xi_{t-1}^{(k)} \sqrt{\frac{p}{n_k}} + \sqrt{\frac{\log K}{n_k}}
\end{aligned}$$

$$\begin{aligned}
&\lesssim \kappa_0 \sum_{k \in S} \frac{n_k}{n_S} d(\widehat{\boldsymbol{\theta}}^{(k)[t-1]}, \boldsymbol{\theta}^{(k)*}) + \sqrt{\frac{p}{n_S}} + h \wedge \sqrt{\frac{p + \log K}{n_k}} + \epsilon \sqrt{\frac{p + \log K}{n_k}} \\
&\quad + \kappa_0 \xi_{t-1}^{(k)} + \sqrt{\frac{\log K}{n_k}}
\end{aligned}$$

Therefore,

$$\begin{aligned}
\sum_{k \in S} \frac{n_k}{n_S} |\widehat{\delta}^{(k)[t]} - \delta^{(k)*}| &\lesssim \kappa_0 \sum_{k \in S} \frac{n_k}{n_S} d(\widehat{\boldsymbol{\theta}}^{(k)[t-1]}, \boldsymbol{\theta}^{(k)*}) + \sqrt{\frac{p}{n_S}} + h \wedge \sqrt{\frac{K(p + \log K)}{n_S}} \\
&\quad + \epsilon \sqrt{\frac{K(p + \log K)}{n_S}} + \kappa_0 \sum_{k \in S} \frac{n_k}{n_S} \xi_{t-1}^{(k)} + \sqrt{\frac{K \log K}{n_S}}.
\end{aligned}$$

Hence

$$\begin{aligned}
\sum_{k \in S} \frac{n_k}{n_S} d(\widehat{\boldsymbol{\theta}}^{(k)[t]}, \boldsymbol{\theta}^{(k)*}) &\leq C \kappa_0 \sum_{k \in S} \frac{n_k}{n_S} d(\widehat{\boldsymbol{\theta}}^{(k)[t-1]}, \boldsymbol{\theta}^{(k)*}) + C \sqrt{\frac{p}{n_S}} + C \cdot h \wedge \sqrt{\frac{K(p + \log K)}{n_S}} \\
&\quad + C \epsilon \sqrt{\frac{K(p + \log K)}{n_S}} + C \kappa_0 \bar{\xi}_{t-1} + C \sqrt{\frac{K \log K}{n_S}} \\
&\leq \kappa'_0 \bar{\xi}_{t-1} + C \sqrt{\frac{p}{n_S}} + C \cdot h \wedge \sqrt{\frac{K(p + \log K)}{n_S}} \\
&\quad + C \epsilon \sqrt{\frac{K(p + \log K)}{n_S}} + C' \sqrt{\frac{K \log K}{n_S}},
\end{aligned}$$

and

$$\begin{aligned}
d(\widehat{\boldsymbol{\theta}}^{(k)[t]}, \boldsymbol{\theta}^{(k)*}) &\leq C \kappa_0 \sum_{k \in S} \frac{n_k}{n_S} d(\widehat{\boldsymbol{\theta}}^{(k)[t-1]}, \boldsymbol{\theta}^{(k)*}) + C \sqrt{\frac{p}{n_S}} + C h \wedge \sqrt{\frac{p + \log K}{n_k}} + C \epsilon \sqrt{\frac{p + \log K}{n_k}} \\
&\quad + C \kappa_0 \xi_{t-1}^{(k)} + C \sqrt{\frac{\log K}{n_k}} \\
&\leq \frac{1}{2} \kappa'_0 \sum_{k \in S} \frac{n_k}{n_S} d(\widehat{\boldsymbol{\theta}}^{(k)[t-1]}, \boldsymbol{\theta}^{(k)*}) + C \sqrt{\frac{p}{n_S}} + C h \wedge \sqrt{\frac{p + \log K}{n_k}} + C \epsilon \sqrt{\frac{p + \log K}{n_k}} \\
&\quad + \frac{1}{2} \kappa'_0 \xi_{t-1}^{(k)} + C \sqrt{\frac{\log K}{n_k}} \\
&\leq \kappa'_0 \xi_{t-1}^{(k)} + C \sqrt{\frac{p}{n_S}} + C h \wedge \sqrt{\frac{p + \log K}{n_k}} + C \epsilon \sqrt{\frac{p + \log K}{n_k}} + C \sqrt{\frac{\log K}{n_k}} \text{ (E.34)}
\end{aligned}$$

where $\kappa'_0 = 2C\kappa_0 \in (0, 1)$.

Therefore, for $t = (t_0 + 1) : (t_0 + t'_0)$, where $(\kappa'_0)^{t'_0} \asymp K^{-1/2}$ hence $t'_0 \asymp \log K$, we have updating formulas (E.32) and (E.33) for $\xi_t^{(k)}$. We can get

$$\begin{aligned} \bar{\xi}_{t_0+t'} &= (\kappa'_0)^{t'} \bar{\xi}_{t_0} + C' \sqrt{\frac{p}{n_S}} + C' \cdot h \wedge \sqrt{\frac{K(p + \log K)}{n_S}} + C' \epsilon \sqrt{\frac{K(p + \log K)}{n_S}} + C' \sqrt{\frac{K \log K}{n_S}} \\ &\leq C(\kappa'_0)^{t'} \sqrt{\frac{K(p + \log K)}{n_S}} + C' \sqrt{\frac{p}{n_S}} + C' \cdot h \wedge \sqrt{\frac{K(p + \log K)}{n_S}} + C' \epsilon \sqrt{\frac{K(p + \log K)}{n_S}} \\ &\quad + C' \sqrt{\frac{K \log K}{n_S}}, \end{aligned}$$

and

$$\begin{aligned} \xi_{t_0+t'}^{(k)} &= \kappa'_0 \bar{\xi}_{t_0+t'-1} + C' \sqrt{\frac{p}{n_S}} + C' \cdot h \wedge \sqrt{\frac{p + \log K}{n_k}} + C' \epsilon \sqrt{\frac{p + \log K}{n_k}} + C' \sqrt{\frac{K \log K}{n_S}} \\ &\leq C(\kappa'_0)^{t'} \sqrt{\frac{K(p + \log K)}{n_S}} + C' \sqrt{\frac{p}{n_S}} + C' \cdot h \wedge \sqrt{\frac{p + \log K}{n_k}} + C' \epsilon \sqrt{\frac{p + \log K}{n_k}} + C' \sqrt{\frac{K \log K}{n_S}}, \end{aligned}$$

with

$$\xi_{t_0+t'_0}^{(k)} \leq C' \sqrt{\frac{p}{n_S}} + C' \cdot h \wedge \sqrt{\frac{p + \log K}{n_k}} + C' \epsilon \sqrt{\frac{p + \log K}{n_k}} + C' \sqrt{\frac{K \log K}{n_S}}, \quad (\text{E.35})$$

where the last inequality is due to $(\kappa'_0)^{t'_0} \sqrt{\frac{K(p + \log K)}{n_S}} \asymp \sqrt{\frac{p + \log K}{n_S}}$.

Consider an event series $\{\mathcal{E}_t\}_{t=t_0}^{t_0+t'_0}$ each of which is defined to be the intersection of the events in Lemmas 11, 12, 13, and 14, with $\xi_t^{(k)} = \xi_t^{(k)}$, which satisfies $\mathbb{P}(\mathcal{E}_t) \geq 1 - C'(K^{-2} + K^{-1}e^{-C''p})$ hence $\mathbb{P}(\bigcap_{t=t_0}^{t_0+t'_0} \mathcal{E}_t) \geq 1 - C't'_0(K^{-2} + K^{-2}e^{-C''p}) \geq 1 - C''(K^{-2} + K^{-2}e^{-C''p}) \log K \geq 1 - C'(K^{-1} + e^{-C''p})$. In the following, we condition on $\mathcal{E} \cap (\bigcap_{t=t_0}^{t_0+t'_0} \mathcal{E}_t)$, therefore all the arguments hold with probability at least $1 - C'(K^{-1} + e^{-C''p})$. Therefore, for $t = (t_0 + 1) : (t_0 + t'_0)$, we have (E.34) hold, which leads to

$$\begin{aligned} d(\hat{\boldsymbol{\theta}}^{(k)[t_0+t']}, \boldsymbol{\theta}^{(k)*}) &\leq \kappa'_0 \xi_{t_0+t'-1}^{(k)} + C \sqrt{\frac{p}{n_S}} + Ch \wedge \sqrt{\frac{p + \log K}{n_k}} + C \epsilon \sqrt{\frac{p + \log K}{n_k}} + C \sqrt{\frac{\log K}{n_k}} \\ &\leq C(\kappa'_0)^{t'} \sqrt{\frac{K(p + \log K)}{n_S}} + C \sqrt{\frac{p}{n_S}} + Ch \wedge \sqrt{\frac{p + \log K}{n_k}} + C \epsilon \sqrt{\frac{p + \log K}{n_k}} \end{aligned}$$

$$\begin{aligned}
& + C\sqrt{\frac{\log K}{n_k}} \\
& \leq C'(\kappa'_0)^{t'} \cdot t_0^2 \kappa^{t_0} + C\sqrt{\frac{p}{n_S}} + Ch \wedge \sqrt{\frac{p + \log K}{n_k}} + C\epsilon\sqrt{\frac{p + \log K}{n_k}} \\
& + C\sqrt{\frac{\log K}{n_k}} \\
& \leq (t_0 + t')^2 (\kappa \vee \kappa'_0)^{t_0+t'} + C\sqrt{\frac{p}{n_S}} + Ch \wedge \sqrt{\frac{p + \log K}{n_k}} + C\epsilon\sqrt{\frac{p + \log K}{n_k}} \\
& + C\sqrt{\frac{\log K}{n_k}},
\end{aligned}$$

where $t' = 1, \dots, t'_0$, which provides the desired rate for $t = (t_0 + 1) : (t_0 + t'_0)$. When $t' \geq t'_0$, by (E.35), we have

$$\begin{aligned}
d(\widehat{\boldsymbol{\theta}}^{(k)[t_0+t']}, \boldsymbol{\theta}^{(k)*}) & \leq (\kappa'_0)^{t'-t'_0} \xi_{t_0+t'_0}^{(k)} + C\sqrt{\frac{p}{n_S}} + Ch \wedge \sqrt{\frac{p + \log K}{n_k}} + C\epsilon\sqrt{\frac{p + \log K}{n_k}} + C\sqrt{\frac{\log K}{n_k}} \\
& \leq C\sqrt{\frac{p}{n_S}} + Ch \wedge \sqrt{\frac{p + \log K}{n_k}} + C\epsilon\sqrt{\frac{p + \log K}{n_k}} + C\sqrt{\frac{\log K}{n_k}},
\end{aligned}$$

which is the desired rate. We complete the proof for Theorem 1.

E.2.3 Proofs of lemmas

Proof of Lemma 11. We prove part (i) first.

Denote $W = \sup_{\substack{\boldsymbol{\theta}^{(k)} \in B_{\text{con}} \\ \|\boldsymbol{\beta}^{(k)} - \boldsymbol{\beta}^{(k)*}\|_2 \leq \xi^{(k)}}} \left| \frac{1}{n_k} \sum_{i=1}^{n_k} \gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}_i^{(k)}) - \mathbb{E}[\gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}^{(k)})] \right|$. By bounded difference inequality,

$$W \leq \mathbb{E}W + C\sqrt{\frac{\log K}{n_k}}, \quad (\text{E.36})$$

with probability at least $1 - C'K^{-2}$. By the generalized symmetrization inequality (Proposition 4.11 in Wainwright (2019)), with i.i.d. Rademacher variables $\{\epsilon_i^{(k)}\}_{i=1}^{n_k}$,

$$\begin{aligned}
\mathbb{E}W & \leq \frac{2}{n_k} \mathbb{E}_{\mathbf{z}} \mathbb{E}_{\boldsymbol{\epsilon}} \left[\sup_{\substack{\boldsymbol{\theta}^{(k)} \in B_{\text{con}} \\ \|\boldsymbol{\beta}^{(k)} - \boldsymbol{\beta}^{(k)*}\|_2 \leq \xi^{(k)}}} \left| \sum_{i=1}^{n_k} \gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}_i^{(k)}) \epsilon_i^{(k)} \right| \right] \\
& \leq \frac{2}{n_k} \mathbb{E}_{\mathbf{z}} \mathbb{E}_{\boldsymbol{\epsilon}} \left[\sup_{\substack{\boldsymbol{\theta}^{(k)} \in B_{\text{con}} \\ \|\boldsymbol{\beta}^{(k)} - \boldsymbol{\beta}^{(k)*}\|_2 \leq \xi^{(k)}}} \left| \sum_{i=1}^{n_k} \frac{w^{(k)} \exp\{C_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}_i^{(k)})\}}{1 - w^{(k)} + w^{(k)} \exp\{C_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}_i^{(k)})\}} \epsilon_i^{(k)} \right| \right]
\end{aligned}$$

$$\leq \frac{2}{n_k} \mathbb{E}_{\mathbf{z}} \mathbb{E}_{\epsilon} \left[\sup_{\substack{\boldsymbol{\theta}^{(k)} \in B_{\text{con}} \\ \|\boldsymbol{\beta}^{(k)} - \boldsymbol{\beta}^{(k)*}\|_2 \leq \xi^{(k)}}} \left| \sum_{i=1}^{n_k} \frac{1}{1 + \exp\{C_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}_i^{(k)}) - \log((w^{(k)})^{-1} - 1)\}} \epsilon_i^{(k)} \right| \right], \quad (\text{E.37})$$

where $C_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}_i^{(k)}) = (\boldsymbol{\beta}^{(k)})^T \mathbf{z}_i^{(k)} - \delta^{(k)}$. Denote $\boldsymbol{\mu}^{(k)*} = (1 - w^{(k)*}) \boldsymbol{\mu}_1^{(k)*} + w^{(k)*} \boldsymbol{\mu}_2^{(k)*} = \mathbb{E}[\mathbf{z}_i^{(k)}]$. By the contraction inequality for Rademacher variables (Theorem 11.6 in [Boucheron et al. \(2013\)](#)),

$$\begin{aligned} \text{RHS of (E.37)} &\leq \frac{C}{n_k} \mathbb{E}_{\mathbf{z}} \mathbb{E}_{\epsilon} \left[\sup_{\substack{\boldsymbol{\theta}^{(k)} \in B_{\text{con}} \\ \|\boldsymbol{\beta}^{(k)} - \boldsymbol{\beta}^{(k)*}\|_2 \leq \xi^{(k)}}} \left| \sum_{i=1}^{n_k} [C_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}_i^{(k)}) - \log((w^{(k)})^{-1} - 1)] \epsilon_i^{(k)} \right| \right] \\ &\leq \frac{C}{n_k} \mathbb{E}_{\mathbf{z}} \mathbb{E}_{\epsilon} \left[\sup_{\substack{\boldsymbol{\theta}^{(k)} \in B_{\text{con}} \\ \|\boldsymbol{\beta}^{(k)} - \boldsymbol{\beta}^{(k)*}\|_2 \leq \xi^{(k)}}} \left| \sum_{i=1}^{n_k} (\boldsymbol{\beta}^{(k)})^T (\mathbf{z}_i^{(k)} - \boldsymbol{\mu}^{(k)*}) \cdot \epsilon_i^{(k)} \right| \right] \\ &\quad + \frac{C}{n_k} \mathbb{E}_{\mathbf{z}} \mathbb{E}_{\epsilon} \left[\sup_{\substack{\boldsymbol{\theta}^{(k)} \in B_{\text{con}} \\ \|\boldsymbol{\beta}^{(k)} - \boldsymbol{\beta}^{(k)*}\|_2 \leq \xi^{(k)}}} \left| \sum_{i=1}^{n_k} (\boldsymbol{\beta}^{(k)})^T \boldsymbol{\mu}^{(k)*} \cdot \epsilon_i^{(k)} \right| \right] \\ &\quad + \frac{C}{n_k} \mathbb{E}_{\epsilon} \left[\sup_{\substack{\boldsymbol{\theta}^{(k)} \in B_{\text{con}} \\ \|\boldsymbol{\beta}^{(k)} - \boldsymbol{\beta}^{(k)*}\|_2 \leq \xi^{(k)}}} |\log((w^{(k)})^{-1} - 1)| \cdot \left| \sum_{i=1}^{n_k} \epsilon_i^{(k)} \right| \right] \\ &\leq \frac{C}{n_k} \mathbb{E}_{\mathbf{z}} \mathbb{E}_{\epsilon} \left[\sup_{\|\boldsymbol{\beta}^{(k)} - \boldsymbol{\beta}^{(k)*}\|_2 \leq \xi^{(k)}} \left| \sum_{i=1}^{n_k} (\boldsymbol{\beta}^{(k)} - \boldsymbol{\beta}^{(k)*})^T (\mathbf{z}_i^{(k)} - \boldsymbol{\mu}^{(k)*}) \cdot \epsilon_i^{(k)} \right| \right] \\ &\quad + \frac{C}{n_k} \mathbb{E}_{\mathbf{z}} \mathbb{E}_{\epsilon} \left| \sum_{i=1}^{n_k} (\boldsymbol{\beta}^{(k)*})^T (\mathbf{z}_i^{(k)} - \boldsymbol{\mu}^{(k)*}) \cdot \epsilon_i^{(k)} \right| \\ &\quad + \frac{C}{n_k} \mathbb{E}_{\epsilon} \left| \sum_{i=1}^{n_k} \epsilon_i^{(k)} \right| + \frac{C}{n_k} \mathbb{E}_{\mathbf{z}} \mathbb{E}_{\epsilon} \left[\sup_{\substack{\boldsymbol{\theta}^{(k)} \in B_{\text{con}} \\ \|\boldsymbol{\beta}^{(k)} - \boldsymbol{\beta}^{(k)*}\|_2 \leq \xi^{(k)}}} \left| \sum_{i=1}^{n_k} (\boldsymbol{\beta}^{(k)})^T \boldsymbol{\mu}^{(k)*} \cdot \epsilon_i^{(k)} \right| \right]. \end{aligned}$$

Since $\{\epsilon_i^{(k)}\}_{i=1}^{n_k}$ and $\{(\boldsymbol{\beta}^{(k)} - \boldsymbol{\beta}^{(k)*})^T (\mathbf{z}_i^{(k)} - \boldsymbol{\mu}^{(k)*}) \epsilon_i^{(k)}\}_{i=1}^{n_k}$ are i.i.d. sub-Gaussian variables, we know that

$$\frac{C}{n_k} \mathbb{E}_{\mathbf{z}} \mathbb{E}_{\epsilon} \left| \sum_{i=1}^{n_k} (\boldsymbol{\beta}^{(k)*})^T (\mathbf{z}_i^{(k)} - \boldsymbol{\mu}^{(k)*}) \cdot \epsilon_i^{(k)} \right| + \frac{C}{n_k} \mathbb{E}_{\epsilon} \left| \sum_{i=1}^{n_k} \epsilon_i^{(k)} \right| \lesssim \sqrt{\frac{1}{n_k}}.$$

Suppose $\{\mathbf{u}_j\}_{j=1}^N$ is a $1/2$ -cover of $\mathcal{B}^p := \{\mathbf{u} \in \mathbb{R}^p : \|\mathbf{u}\|_2 \leq 1\}$ with $N = 5^p$ (see Example

5.8 in [Wainwright \(2019\)](#)). Hence by standard arguments,

$$\begin{aligned} & \frac{C}{n_k} \mathbb{E}_{\mathbf{z}} \mathbb{E}_{\epsilon} \left[\sup_{\|\beta^{(k)} - \beta^{(k)*}\|_2 \leq \xi^{(k)}} \left| \sum_{i=1}^{n_k} (\beta^{(k)} - \beta^{(k)*})^T (\mathbf{z}_i^{(k)} - \boldsymbol{\mu}^{(k)*}) \cdot \epsilon_i^{(k)} \right| \right] \\ & \lesssim \frac{\xi^{(k)}}{n_k} \mathbb{E}_{\mathbf{z}} \mathbb{E}_{\epsilon} \left[\sup_{j=1:N} \left| \sum_{i=1}^{n_k} \mathbf{u}_j^T (\mathbf{z}_i^{(k)} - \boldsymbol{\mu}^{(k)*}) \cdot \epsilon_i^{(k)} \right| \right]. \end{aligned}$$

Again, since $\{\mathbf{u}_j^T (\mathbf{z}_i^{(k)} - \boldsymbol{\mu}^{(k)*}) \epsilon_i^{(k)}\}_{i=1}^{n_k}$ are i.i.d. sub-Gaussian variables,

$$\frac{1}{n_k} \mathbb{E}_{\mathbf{z}} \mathbb{E}_{\epsilon} \left[\sup_{j=1:N} \left| \sum_{i=1}^{n_k} \mathbf{u}_j^T (\mathbf{z}_i^{(k)} - \boldsymbol{\mu}^{(k)*}) \cdot \epsilon_i^{(k)} \right| \right] \lesssim \sqrt{\frac{\log N}{n_k}} = \sqrt{\frac{p}{n_k}}.$$

Putting all pieces together,

$$\mathbb{E}W \lesssim \xi^{(k)} \sqrt{\frac{p}{n_k}} + \sqrt{\frac{1}{n_k}}. \quad (\text{E.38})$$

Combining (E.37) and (E.38), we get the result in (i).

Next, we derive part (ii) using a similar analysis.

Denote $W' = \sup_{\{\boldsymbol{\theta}^{(k)}\}_{k \in S} \in B_{\text{con}}^{J,2}} \sup_{|\tilde{w}_k| \leq 1} \frac{1}{n_S} \left| \sum_{k \in S} \tilde{w}_k \sum_{i=1}^{n_k} [\gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}_i^{(k)}) - \mathbb{E}[\gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}^{(k)})]] \right|$.

By a similar standard symmetrization and contraction arguments we used in part (i), with i.i.d. Rademacher variables $\{\epsilon_i^{(k)}\}_{i=1}^{n_k}$, for any $\lambda \in \mathbb{R}$, we have

$$\begin{aligned} & \mathbb{E} \exp\{\lambda W'\} \\ & \leq C \mathbb{E}_{\mathbf{z}} \mathbb{E}_{\epsilon} \exp \left\{ \frac{2\lambda}{n_S} \sup_{\{\boldsymbol{\theta}^{(k)}\}_{k \in S} \in B_{\text{con}}^{J,2}} \sup_{|\tilde{w}_k| \leq 1} \left| \sum_{k \in S} \tilde{w}_k \sum_{i=1}^{n_k} \gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}_i^{(k)}) \epsilon_i^{(k)} \right| \right\} \\ & \leq C \mathbb{E}_{\mathbf{z}} \mathbb{E}_{\epsilon} \exp \left\{ \frac{4\lambda}{n_S} \sup_{\{\boldsymbol{\theta}^{(k)}\}_{k \in S} \in B_{\text{con}}^{J,2}} \sup_{\tilde{w}_k = \pm 1/2} \left| \sum_{k \in S} \tilde{w}_k \sum_{i=1}^{n_k} \gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}_i^{(k)}) \epsilon_i^{(k)} \right| \right\} \\ & \leq C \sum_{\tilde{w}_k = \pm 1/2} \mathbb{E}_{\mathbf{z}} \mathbb{E}_{\epsilon} \exp \left\{ \frac{4\lambda}{n_S} \sup_{\{\boldsymbol{\theta}^{(k)}\}_{k \in S} \in B_{\text{con}}^{J,2}} \left| \sum_{k \in S} \tilde{w}_k \sum_{i=1}^{n_k} \gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}_i^{(k)}) \epsilon_i^{(k)} \right| \right\} \\ & \leq C \sum_{\tilde{w}_k = \pm 1/2} \mathbb{E}_{\mathbf{z}} \mathbb{E}_{\epsilon} \exp \left\{ \frac{4\lambda}{n_S} \sup_{\{\boldsymbol{\theta}^{(k)}\}_{k \in S} \in B_{\text{con}}^{J,2}} \left| \sum_{k \in S} \tilde{w}_k \sum_{i=1}^{n_k} \frac{1}{1 + \exp\{C_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}_i^{(k)}) - \log((w^{(k)})^{-1} - 1)\}} \epsilon_i^{(k)} \right| \right\} \\ & \leq C \sum_{\tilde{w}_k = \pm 1/2} \mathbb{E}_{\mathbf{z}} \mathbb{E}_{\epsilon} \exp \left\{ \frac{4\lambda}{n_S} \sup_{\{\boldsymbol{\theta}^{(k)}\}_{k \in S} \in B_{\text{con}}^{J,2}} \left| \sum_{k \in S} \tilde{w}_k \sum_{i=1}^{n_k} [C_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}_i^{(k)}) - \log((w^{(k)})^{-1} - 1)] \epsilon_i^{(k)} \right| \right\}, \end{aligned}$$

where $C_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}_i^{(k)}) = (\beta^{(k)})^T \mathbf{z}_i^{(k)} - \delta^{(k)}$. Denote $\boldsymbol{\mu}^{(k)*} = (1 - w^{(k)*}) \boldsymbol{\mu}_1^{(k)*} + w^{(k)*} \boldsymbol{\mu}_2^{(k)*} = \mathbb{E}[\mathbf{z}_i^{(k)}]$. Suppose $\{\mathbf{u}_j\}_{j=1}^N$ is a $1/2$ -cover of $\mathcal{B}^p := \{\mathbf{u} \in \mathbb{R}^p : \|\mathbf{u}\|_2 \leq 1\}$ with $N = 5^p$. Then by

Cauchy-Schwarz inequality and standard arguments,

$$\begin{aligned}
& \mathbb{E}_{\mathbf{z}, \epsilon} \exp \left\{ \frac{4\lambda}{n_S} \sup_{\{\boldsymbol{\theta}^{(k)}\}_{k \in S} \in B_{\text{con}}^{J,2}} \left| \sum_{k \in S} \tilde{w}_k \sum_{i=1}^{n_k} [C_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}_i^{(k)}) - \log((w^{(k)})^{-1} - 1)] \epsilon_i^{(k)} \right| \right\} \\
& \lesssim \left[\mathbb{E}_{\mathbf{z}, \epsilon} \exp \left\{ \frac{C\lambda}{n_S} \sup_{\|\boldsymbol{\beta}\|_2 \leq U} \left| \sum_{k \in S} \sum_{i=1}^{n_k} \tilde{w}_k \boldsymbol{\beta}^T (\mathbf{z}_i^{(k)} - \boldsymbol{\mu}^{(k)*}) \epsilon_i^{(k)} \right| \right\} \right]^{1/3} \\
& + \left[\mathbb{E}_{\epsilon} \exp \left\{ \frac{C\lambda}{n_S} \sup_{\|\boldsymbol{\beta}\|_2 \leq U} \left| \sum_{k \in S} \sum_{i=1}^{n_k} \tilde{w}_k \boldsymbol{\beta}^T \boldsymbol{\mu}^{(k)*} \epsilon_i^{(k)} \right| \right\} \right]^{1/3} \\
& + \left[\mathbb{E}_{\epsilon} \exp \left\{ \frac{C\lambda}{n_S} \sup_{c_w/2 \leq w^{(k)} \leq 1-c_w/2} \left| \sum_{k \in S} \sum_{i=1}^{n_k} \tilde{w}_k \log((w^{(k)})^{-1} - 1) \epsilon_i^{(k)} \right| \right\} \right]^{1/3} \\
& \lesssim \left[\mathbb{E}_{\mathbf{z}, \epsilon} \exp \left\{ \frac{C\lambda}{n_S} \sup_{j=1:N} \left| \sum_{k \in S} \sum_{i=1}^{n_k} \tilde{w}_k \mathbf{u}_j^T (\mathbf{z}_i^{(k)} - \boldsymbol{\mu}^{(k)*}) \epsilon_i^{(k)} \right| \right\} \right]^{1/3} \\
& + \left[\mathbb{E}_{\epsilon} \exp \left\{ \frac{C\lambda}{n_S} \sup_{j=1:N} \left| \sum_{k \in S} \sum_{i=1}^{n_k} \tilde{w}_k \mathbf{u}_j^T \boldsymbol{\mu}^{(k)*} \epsilon_i^{(k)} \right| \right\} \right]^{1/3} \\
& + \left[\mathbb{E}_{\epsilon} \exp \left\{ \frac{C\lambda}{n_S} \sup_{c_w/2 \leq w^{(k)} \leq 1-c_w/2} \left| \sum_{k \in S} \sum_{i=1}^{n_k} \tilde{w}_k \log((w^{(k)})^{-1} - 1) \epsilon_i^{(k)} \right| \right\} \right]^{1/3} \\
& \lesssim \underbrace{\sum_{j=1}^N \left[\mathbb{E}_{\mathbf{z}, \epsilon} \exp \left\{ \frac{C\lambda}{n_S} \left| \sum_{k \in S} \sum_{i=1}^{n_k} \tilde{w}_k \mathbf{u}_j^T (\mathbf{z}_i^{(k)} - \boldsymbol{\mu}^{(k)*}) \epsilon_i^{(k)} \right| \right\} \right]}_{[1]} \\
& + \underbrace{\sum_{j=1}^N \left[\mathbb{E}_{\epsilon} \exp \left\{ \frac{C\lambda}{n_S} \left| \sum_{k \in S} \sum_{i=1}^{n_k} \tilde{w}_k \mathbf{u}_j^T \boldsymbol{\mu}^{(k)*} \epsilon_i^{(k)} \right| \right\} \right]}_{[2]} \\
& + \underbrace{\left[\mathbb{E}_{\epsilon} \exp \left\{ \frac{C\lambda}{n_S} \sum_{k \in S} \left| \sum_{i=1}^{n_k} \tilde{w}_k \epsilon_i^{(k)} \right| \right\} \right]}_{[3]}
\end{aligned}$$

Since $\{\tilde{w}_k \mathbf{u}_j^T (\mathbf{z}_i^{(k)} - \boldsymbol{\mu}^{(k)*}) \epsilon_i^{(k)}\}_{i,j}$, $\{\tilde{w}_k (\boldsymbol{\beta}^{(k)})^T \boldsymbol{\mu}^{(k)*} \epsilon_i^{(k)}\}_{i,k}$, and $\{\tilde{w}_k \epsilon_i^{(k)}\}_{i=1}^{n_k}$ are independent sub-Gaussian variables, we can bound the three terms on the RHS as

$$\begin{aligned}
[1] & \lesssim 5^p \cdot \exp \left\{ \frac{C\lambda^2}{n_S} \right\}, \\
[2] & \lesssim 5^p \cdot \exp \left\{ \frac{C\lambda^2}{n_S} \right\},
\end{aligned}$$

$$[3] \leq \left[\prod_{k \in S} \mathbb{E}_\epsilon \exp \left\{ \frac{C\lambda}{n_S} \left| \sum_{i=1}^{n_k} \tilde{w}_k \epsilon_i^{(k)} \right| \right\} \right]^{1/3} \lesssim \left[\prod_{k \in S} \exp \left\{ C \frac{\lambda^2}{n_S^2} n_k \right\} \right]^{1/3} \lesssim \exp \left\{ \frac{C\lambda^2}{n_S} \right\}.$$

Putting all pieces together,

$$\mathbb{E} \exp\{\lambda W'\} \leq C' 2^K 5^p \cdot \exp \left\{ \frac{C\lambda^2}{n_S} \right\} = \exp \left\{ C \frac{\lambda^2}{n_S} + C''(K+p) \right\}.$$

Therefore, for any $\delta > 0$

$$\mathbb{P}(W' \geq \delta) \leq e^{-\lambda\delta} \mathbb{E} \exp\{\lambda W'\} \leq \exp \left\{ C \frac{\lambda^2}{n_S} + C''(K+p) - \lambda\delta \right\}.$$

Let $\lambda = \frac{n_S}{2C}\delta$ and $\delta = 4\sqrt{\frac{CC''(K+p)}{n_S}}$, we have

$$\mathbb{P}(W' \geq \delta) \leq \exp \left\{ -\frac{n_S}{4C}\delta^2 + C''(K+p) \right\} = \exp\{-3C''(K+p)\} \leq C' K^{-1} \exp\{-3C''p\},$$

which completes the proof. \square

Proof of Lemma 12. The proof of part (ii) is the same as the proof of part (ii) for Lemma 11, so we omit it. The only difference between the proofs of part (i) for two lemmas is that here the bounded difference inequality is not available. Denote

$$W = \sup_{\substack{\boldsymbol{\theta}^{(k)} \in B_{\text{con}} \\ \|\boldsymbol{\beta}^{(k)} - \boldsymbol{\beta}^{(k)*}\|_2 \leq \xi^{(k)}}} \left| \frac{1}{n_k} \sum_{i=1}^{n_k} [1 - \gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}_i^{(k)})] (\mathbf{z}_i^{(k)})^T \boldsymbol{\beta}^{(k)*} - \mathbb{E}[1 - \gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}^{(k)})] (\mathbf{z}^{(k)})^T \boldsymbol{\beta}^{(k)*}] \right|.$$

We need to use Lemma 10 to upper bound $W - \mathbb{E}W$. Prior to that, we first verify the conditions required by the lemma. Fix $\mathbf{z}_1^{(k)}, \dots, \mathbf{z}_{i-1}^{(k)}, \mathbf{z}_{i+1}^{(k)}, \dots, \mathbf{z}_{n_k}^{(k)}$, and define

$$g_i^{(k)}(\mathbf{z}_i^{(k)}) = W - \mathbb{E}[W | \mathbf{z}_1^{(k)}, \dots, \mathbf{z}_{i-1}^{(k)}, \mathbf{z}_{i+1}^{(k)}, \dots, \mathbf{z}_{n_k}^{(k)}].$$

By triangle inequality,

$$\begin{aligned} & \left| g_i^{(k)}(\mathbf{z}_i^{(k)}) \right| \\ &= \left| \sup_{\substack{\boldsymbol{\theta}^{(k)} \in B_{\text{con}} \\ \|\boldsymbol{\beta}^{(k)} - \boldsymbol{\beta}^{(k)*}\|_2 \leq \xi^{(k)}}} \left| \frac{1}{n_k} \sum_{i=1}^{n_k} \gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}_i^{(k)}) (\mathbf{z}_i^{(k)})^T \boldsymbol{\beta}^{(k)*} - \mathbb{E}[\gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}_i^{(k)}) (\mathbf{z}_i^{(k)})^T \boldsymbol{\beta}^{(k)*}] \right| \right. \\ & \quad \left. - \mathbb{E} \left[\sup_{\substack{\boldsymbol{\theta}^{(k)} \in B_{\text{con}} \\ \|\boldsymbol{\beta}^{(k)} - \boldsymbol{\beta}^{(k)*}\|_2 \leq \xi^{(k)}}} \left| \frac{1}{n_k} \sum_{i=1}^{n_k} \gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}_i^{(k)}) (\mathbf{z}_i^{(k)})^T \boldsymbol{\beta}^{(k)*} - \mathbb{E}[\gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}_i^{(k)}) (\mathbf{z}_i^{(k)})^T \boldsymbol{\beta}^{(k)*}] \right| \left| \{\mathbf{z}_{i'}^{(k)}\}_{i' \neq i} \right| \right] \right| \end{aligned}$$

$$\leq \underbrace{\frac{1}{n_k} \sup_{\substack{\boldsymbol{\theta}^{(k)} \in B_{\text{con}} \\ \|\boldsymbol{\beta}^{(k)} - \boldsymbol{\beta}^{(k)*}\|_2 \leq \xi^{(k)}}} \left| \gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}_i^{(k)}) (\mathbf{z}_i^{(k)})^T \boldsymbol{\beta}^{(k)*} \right|}_{W_1} + \underbrace{\frac{2}{n_k} \mathbb{E} \left[\sup_{\substack{\boldsymbol{\theta}^{(k)} \in B_{\text{con}} \\ \|\boldsymbol{\beta}^{(k)} - \boldsymbol{\beta}^{(k)*}\|_2 \leq \xi^{(k)}}} \gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}_i^{(k)}) (\mathbf{z}_i^{(k)})^T \boldsymbol{\beta}^{(k)*} \right]}_{W_2}.$$

Note that $[\mathbb{E}(W_1 + W_2)^d]^{1/d} \leq (\mathbb{E}W_1^d)^{1/d} + (\mathbb{E}W_2^d)^{1/d}$, where

$$\begin{aligned} (\mathbb{E}W_1^d)^{1/d} &\leq \frac{1}{n_k} \left[\mathbb{E} \sup_{\boldsymbol{\theta}^{(k)} \in B_{\text{con}}} [\gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}_i^{(k)})]^{2d} \right]^{1/2d} \left[\mathbb{E} |(\mathbf{z}_i^{(k)})^T \boldsymbol{\beta}^{(k)*}|^{2d} \right]^{1/2d} \\ &\leq \frac{1}{n_k} \cdot C \cdot \sqrt{d} \\ (\mathbb{E}W_2^d)^{1/d} &= \mathbb{E}W_1 \leq (\mathbb{E}W_1^d)^{1/d} \leq \frac{1}{n_k} \cdot C \cdot \sqrt{d}. \end{aligned}$$

Therefore $[\mathbb{E}(W_1 + W_2)^d]^{1/d} \leq \frac{C}{n_k} \sqrt{d}$. Hence by applying Lemma 10, we have

$$W \leq \mathbb{E}W + C \sqrt{\frac{\log K}{n_k}},$$

with probability at least $1 - C'K^{-2}$. □

Proof of Lemma 13. For part (i), denote

$$W = \sup_{\boldsymbol{\theta}^{(k)} \in B_{\text{con}}} \left\| \frac{1}{n_k} \sum_{i=1}^{n_k} \gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}_i^{(k)}) \mathbf{z}_i^{(k)} - \mathbb{E}[\gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}^{(k)}) \mathbf{z}^{(k)}] \right\|_2.$$

Suppose $\{\mathbf{u}_j\}_{j=1}^N$ is a $1/2$ -cover of $\mathcal{B}^p := \{\mathbf{u} \in \mathbb{R}^p : \|\mathbf{u}\|_2 \leq 1\}$ with $N = 5^p$. Define $\boldsymbol{\mu}^{(k)*} = (1 - w^{(k)*})\boldsymbol{\mu}_1^{(k)*} + w^{(k)*}\boldsymbol{\mu}_2^{(k)*}$. Then by the generalized symmetrization inequality (Proposition 4.11 in Wainwright (2019)), with i.i.d. Rademacher variables $\{\epsilon_i^{(k)}\}_{i=1}^{n_k}$, for any $\lambda \in \mathbb{R}$,

$$\begin{aligned} &\mathbb{E} \exp\{\lambda W\} \\ &\lesssim \mathbb{E}_{\mathbf{z}, \epsilon} \exp \left\{ \frac{C\lambda}{n_k} \sup_{\boldsymbol{\theta}^{(k)} \in B_{\text{con}}} \left\| \sum_{i=1}^{n_k} \gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}_i^{(k)}) \mathbf{z}_i^{(k)} \epsilon_i^{(k)} \right\|_2 \right\} \\ &\lesssim \mathbb{E}_{\mathbf{z}, \epsilon} \exp \left\{ \frac{C\lambda}{n_k} \sup_{j=1:N} \sup_{\boldsymbol{\theta}^{(k)} \in B_{\text{con}}} \left| \sum_{i=1}^{n_k} \gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}_i^{(k)}) (\mathbf{z}_i^{(k)})^T \mathbf{u}_j \cdot \epsilon_i^{(k)} \right| \right\} \\ &\lesssim \sum_{j=1}^N \mathbb{E}_{\mathbf{z}, \epsilon} \exp \left\{ \frac{C\lambda}{n_k} \sup_{\boldsymbol{\theta}^{(k)} \in B_{\text{con}}} \left| \sum_{i=1}^{n_k} \gamma_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}_i^{(k)}) (\mathbf{z}_i^{(k)})^T \mathbf{u}_j \cdot \epsilon_i^{(k)} \right| \right\} \end{aligned}$$

$$\begin{aligned}
&\lesssim \sum_{j=1}^N \mathbb{E}_{\mathbf{z}, \epsilon} \exp \left\{ \frac{C\lambda}{n_k} \sup_{\boldsymbol{\theta}^{(k)} \in B_{\text{con}}} \left| \sum_{i=1}^{n_k} [C_{\boldsymbol{\theta}^{(k)}}(\mathbf{z}_i^{(k)}) - \log((w^{(k)})^{-1} - 1)] (\mathbf{z}_i^{(k)})^T \mathbf{u}_j \cdot \epsilon_i^{(k)} \right| \right\} \\
&\lesssim \sum_{j=1}^N \mathbb{E}_{\mathbf{z}, \epsilon} \exp \left\{ \frac{C\lambda}{n_k} \sup_{\|\boldsymbol{\beta}^{(k)}\|_2 \leq U} \left| \sum_{i=1}^{n_k} (\boldsymbol{\beta}^{(k)})^T (\mathbf{z}_i^{(k)} - \boldsymbol{\mu}^{(k)*}) (\mathbf{z}_i^{(k)})^T \mathbf{u}_j \cdot \epsilon_i^{(k)} \right| \right\} \\
&\quad + \sum_{j=1}^N \mathbb{E}_{\mathbf{z}, \epsilon} \exp \left\{ \frac{C\lambda}{n_k} \sup_{\|\boldsymbol{\beta}^{(k)}\|_2 \leq U} \left| \sum_{i=1}^{n_k} [(\boldsymbol{\beta}^{(k)})^T \boldsymbol{\mu}^{(k)*} - \delta^{(k)}] (\mathbf{z}_i^{(k)})^T \mathbf{u}_j \cdot \epsilon_i^{(k)} \right| \right\} \\
&\quad + \sum_{j=1}^N \mathbb{E}_{\mathbf{z}, \epsilon} \exp \left\{ \frac{C\lambda}{n_k} \sup_{c_w/2 \leq w^{(k)} \leq 1 - c_w/2} \left| \sum_{i=1}^{n_k} \log((w^{(k)})^{-1} - 1) (\mathbf{z}_i^{(k)})^T \mathbf{u}_j \cdot \epsilon_i^{(k)} \right| \right\} \\
&\lesssim \sum_{j=1}^N \sum_{j'=1}^N \mathbb{E}_{\mathbf{z}, \epsilon} \exp \left\{ \frac{C\lambda}{n_k} \left| \sum_{i=1}^{n_k} \mathbf{u}_{j'}^T (\mathbf{z}_i^{(k)} - \boldsymbol{\mu}^{(k)*}) (\mathbf{z}_i^{(k)})^T \mathbf{u}_j \cdot \epsilon_i^{(k)} \right| \right\} \\
&\quad + \sum_{j=1}^N \mathbb{E}_{\mathbf{z}, \epsilon} \exp \left\{ \frac{C\lambda}{n_k} \left| \sum_{i=1}^{n_k} (\mathbf{z}_i^{(k)})^T \mathbf{u}_j \cdot \epsilon_i^{(k)} \right| \right\}.
\end{aligned}$$

Note that since $\{\mathbf{u}_{j'}^T (\mathbf{z}_i^{(k)} - \boldsymbol{\mu}^{(k)*}) (\mathbf{z}_i^{(k)})^T \mathbf{u}_j \cdot \epsilon_i^{(k)}\}_{i=1}^{n_k}$ are i.i.d. sub-exponential variables and $\{(\mathbf{z}_i^{(k)})^T \mathbf{u}_j \cdot \epsilon_i^{(k)}\}_{i=1}^{n_k}$ are i.i.d. sub-Gaussian variables, we have

$$\begin{aligned}
\mathbb{E}_{\mathbf{z}, \epsilon} \exp \left\{ \frac{C\lambda}{n_k} \left| \sum_{i=1}^{n_k} \mathbf{u}_{j'}^T (\mathbf{z}_i^{(k)} - \boldsymbol{\mu}^{(k)*}) (\mathbf{z}_i^{(k)})^T \mathbf{u}_j \cdot \epsilon_i^{(k)} \right| \right\} &\lesssim \exp \left\{ C \frac{\lambda^2}{n_k} \right\}, \\
\mathbb{E}_{\mathbf{z}, \epsilon} \exp \left\{ \frac{C\lambda}{n_k} \left| \sum_{i=1}^{n_k} (\mathbf{z}_i^{(k)})^T \mathbf{u}_j \cdot \epsilon_i^{(k)} \right| \right\} &\lesssim \exp \left\{ C \frac{\lambda^2}{n_k} \right\},
\end{aligned}$$

where the first inequality holds when $\lambda \leq C'' n_k$ where C'' is small. Therefore,

$$\mathbb{E} \exp\{\lambda W\} \lesssim \exp \left\{ C \frac{\lambda^2}{n_k} + C' p \right\},$$

when $\lambda \leq C'' n_k$. The desired result follows from Chernoff's bound.

The proofs of parts (ii) and (iii) are almost the same as the proofs of part (ii) of Lemma 11, so we do not repeat them here. \square

Proof of Lemma 14. Note that

$$\left\| \frac{1}{n_k} \sum_{i=1}^{n_k} [\mathbf{z}_i^{(k)} (\mathbf{z}_i^{(k)})^T - \mathbb{E}[\mathbf{z}_i^{(k)} (\mathbf{z}_i^{(k)})^T]] \boldsymbol{\beta}^{(k)*} \right\|_2 \lesssim \left\| \frac{1}{n_k} \sum_{i=1}^{n_k} [\mathbf{z}_i^{(k)} (\mathbf{z}_i^{(k)})^T - \mathbb{E}[\mathbf{z}_i^{(k)} (\mathbf{z}_i^{(k)})^T]] \right\|_2.$$

The bound of the RHS come from Theorem 6.5 in Wainwright (2019). And the bound in part (ii) can be proved in the same way. \square

E.3 Proof of Theorem 3

E.3.1 Lemmas

Recall the parameter space

$$\overline{\Theta}_S(h) = \left\{ \{\overline{\boldsymbol{\theta}}^{(k)}\}_{k \in S} = \{(w^{(k)}, \boldsymbol{\mu}_1^{(k)}, \boldsymbol{\mu}_2^{(k)}, \boldsymbol{\Sigma}^{(k)})\}_{k \in S} : \overline{\boldsymbol{\theta}}^{(k)} \in \overline{\Theta}, \inf_{\boldsymbol{\beta}} \max_{k \in S} \|\boldsymbol{\beta}^{(k)} - \overline{\boldsymbol{\beta}}\|_2 \leq h \right\},$$

where $\boldsymbol{\beta}^{(k)} = (\boldsymbol{\Sigma}^{(k)})^{-1}(\boldsymbol{\mu}_2^{(k)} - \boldsymbol{\mu}_1^{(k)})$ and $\delta^{(k)} = \frac{1}{2}(\boldsymbol{\beta}^{(k)})^T(\boldsymbol{\mu}_1^{(k)} + \boldsymbol{\mu}_2^{(k)})$.

Lemma 15 (Lemma 8.4 in Cai et al. (2019)). *For any $\boldsymbol{\mu}, \tilde{\boldsymbol{\mu}} \in \mathbb{R}^p$ and $w \in (0, 1)$, denote $\mathbb{P}_{\boldsymbol{\mu}} = (1 - w)\mathcal{N}(\boldsymbol{\mu}, \mathbf{I}_p) + \mathcal{N}(-\boldsymbol{\mu}, \mathbf{I}_p)$ and $\mathbb{P}_{\tilde{\boldsymbol{\mu}}} = (1 - w)\mathcal{N}(\tilde{\boldsymbol{\mu}}, \mathbf{I}_p) + \mathcal{N}(-\tilde{\boldsymbol{\mu}}, \mathbf{I}_p)$. Then*

$$\text{KL}(\mathbb{P}_{\boldsymbol{\mu}} \|\mathbb{P}_{\tilde{\boldsymbol{\mu}}}) \leq \left(4\|\boldsymbol{\mu}\|_2^2 + \frac{1}{2} \log \left(\frac{w}{1-w} \right) \right) \cdot 2\|\boldsymbol{\mu} - \tilde{\boldsymbol{\mu}}\|_2^2.$$

Lemma 16. *For any $\boldsymbol{\mu}, \boldsymbol{\mu}', \tilde{\boldsymbol{\mu}}, \tilde{\boldsymbol{\mu}}' \in \mathbb{R}^p$ and $w \in (0, 1)$, denote $\mathbb{P}_{\boldsymbol{\mu}, \tilde{\boldsymbol{\mu}}} = (1 - w)\mathcal{N}(\boldsymbol{\mu}, \mathbf{I}_p) + w\mathcal{N}(\tilde{\boldsymbol{\mu}}, \mathbf{I}_p)$ and $\mathbb{P}_{\boldsymbol{\mu}', \tilde{\boldsymbol{\mu}}'} = (1 - w)\mathcal{N}(\boldsymbol{\mu}', \mathbf{I}_p) + w\mathcal{N}(\tilde{\boldsymbol{\mu}}', \mathbf{I}_p)$. Then*

$$\text{KL}(\mathbb{P}_{\boldsymbol{\mu}, \tilde{\boldsymbol{\mu}}} \|\mathbb{P}_{\boldsymbol{\mu}', \tilde{\boldsymbol{\mu}}'}) \leq (1 - w)\|\boldsymbol{\mu} - \boldsymbol{\mu}'\|_2^2 + w\|\tilde{\boldsymbol{\mu}} - \tilde{\boldsymbol{\mu}}'\|_2^2.$$

Lemma 17. *Denote distribution $(1 - w)\mathcal{N}(\boldsymbol{\mu}, \mathbf{I}_p) + w\mathcal{N}(-\boldsymbol{\mu}, \mathbf{I}_p)$ as \mathbb{P}_w for any $w \in (c_w, 1 - c_w)$, where $\boldsymbol{\mu} \in \mathbb{R}^p$. Then*

$$\text{KL}(\mathbb{P}_w \|\mathbb{P}_{w'}) \leq \frac{1}{2c_w^2}(w - w')^2.$$

Lemma 18. *Denote distribution $\frac{1}{2}\mathcal{N}((-1/2, \mathbf{0}_{p-1}^T)^T, \mathbf{I}_p) + \frac{1}{2}\mathcal{N}((1/2 + \tilde{u}, \mathbf{0}_{p-1}^T)^T, \mathbf{I}_p)$ as $\mathbb{P}_{\tilde{u}}$ for any $\tilde{u} \in [-1, 1]$. Then*

$$\text{KL}(\mathbb{P}_{\tilde{u}} \|\mathbb{P}_{\tilde{u}'}) \leq \frac{1}{2}(\tilde{u} - \tilde{u}')^2.$$

Lemma 19. *When there exists an subset S such that $\min_{k \in S} n_k \geq C(p + \log K)$ with some constant $C > 0$, we have*

$$\inf_{\{\hat{\boldsymbol{\theta}}^{(k)}\}_{k=1}^K} \sup_{S: |S| \geq s} \sup_{\substack{\{\boldsymbol{\theta}^{(k)*}\}_{k \in S} \in \overline{\Theta}_S \\ \mathbb{Q}_S}} \mathbb{P} \left(\bigcup_{k \in S} \left\{ d(\hat{\boldsymbol{\theta}}^{(k)}, \boldsymbol{\theta}^{(k)*}) \geq C_1 \sqrt{\frac{p}{n_S}} + C_2 \sqrt{\frac{\log K}{n_k}} \right. \right. \\ \left. \left. + C_3 h \wedge \sqrt{\frac{p + \log K}{n_k}} \right\} \right) \geq \frac{1}{4}.$$

Lemma 20. *Denote $\tilde{\epsilon} = \frac{K-s}{s}$. Then*

$$\inf_{\{\hat{\boldsymbol{\theta}}^{(k)}\}_{k=1}^K} \sup_{S: |S| \geq s} \sup_{\substack{\{\boldsymbol{\theta}^{(k)*}\}_{k \in S} \in \overline{\Theta}_S \\ \mathbb{Q}_S}} \mathbb{P} \left(\max_{k \in S} d(\hat{\boldsymbol{\theta}}^{(k)}, \boldsymbol{\theta}^{(k)*}) \geq C_1 \tilde{\epsilon} \sqrt{\frac{1}{\max_{k=1:K} n_k}} \right) \geq \frac{1}{10}.$$

Lemma 21 (The first variant of Theorem 5.1 in [Chen et al. \(2018\)](#)). *Given a series of distributions $\{\{\mathbb{P}_\theta^{(k)}\}_{k=1}^K : \theta \in \Theta\}$, each of which is indexed by the same parameter $\theta \in \Theta$. Consider $\mathbf{x}^{(k)} \sim (1-\tilde{\epsilon})\mathbb{P}_\theta^{(k)} + \tilde{\epsilon}\mathbb{Q}^{(k)}$ independently for $k = 1 : K$. Denote the joint distribution of $\{\mathbf{x}^{(k)}\}_{k=1}^K$ as $\mathbb{P}_{(\tilde{\epsilon}, \theta, \{\mathbb{Q}^{(k)}\}_{k=1}^K)}$. Then*

$$\inf_{\hat{\theta}} \sup_{\substack{\theta \in \Theta \\ \{\mathbb{Q}^{(k)}\}_{k=1}^K}} \mathbb{P}_{(\tilde{\epsilon}, \theta, \{\mathbb{Q}^{(k)}\}_{k=1}^K)} \left(\|\hat{\theta} - \theta\| \geq C\varpi(\tilde{\epsilon}, \Theta) \right) \geq \frac{1}{2},$$

where $\varpi(\tilde{\epsilon}, \Theta) := \sup \{ \|\theta_1 - \theta_2\| : \max_{k=1:K} d_{\text{TV}}(\mathbb{P}_{\theta_1}^{(k)}, \mathbb{P}_{\theta_2}^{(k)}) \leq \tilde{\epsilon}/(1-\tilde{\epsilon}) \}$.

Lemma 22. *Suppose $K - s \geq 1$. Consider two data generating mechanisms:*

- (i) $\mathbf{x}^{(k)} \sim (1 - \tilde{\epsilon}')\mathbb{P}_\theta^{(k)} + \tilde{\epsilon}'\mathbb{Q}^{(k)}$ independently for $k = 1 : K$, where $\tilde{\epsilon}' = \frac{K-s}{K}$;
- (ii) With a preserved set $S \subseteq 1 : K$, generate $\{\mathbf{x}^{(k)}\}_{k \in S^c} \sim \mathbb{Q}_S$ and $\mathbf{x}^{(k)} \sim \mathbb{P}_\theta^{(k)}$ independently for $k \in S$.

Denote the joint distributions of $\{\mathbf{x}^{(k)}\}_{k=1}^K$ in (i) and (ii) as $\mathbb{P}_{(\tilde{\epsilon}, \theta, \{\mathbb{Q}^{(k)}\}_{k=1}^K)}$ and $\mathbb{P}_{(S, \theta, \mathbb{Q})}$, respectively. We claim that if

$$\inf_{\hat{\theta}} \sup_{\substack{\theta \in \Theta \\ \{\mathbb{Q}^{(k)}\}_{k=1}^K}} \mathbb{P}_{(\frac{K-s}{50K}, \theta, \{\mathbb{Q}^{(k)}\}_{k=1}^K)} \left(\|\hat{\theta} - \theta\| \geq C\varpi\left(\frac{K-s}{50K}, \Theta\right) \right) \geq \frac{1}{2},$$

then

$$\inf_{\hat{\theta}} \sup_{S: |S| \geq s} \sup_{\substack{\theta \in \Theta \\ \mathbb{Q}_S}} \mathbb{P}_{(S, \theta, \mathbb{Q}_S)} \left(\|\hat{\theta} - \theta\| \geq C\varpi\left(\frac{K-s}{50K}, \Theta\right) \right) \geq \frac{1}{10}, \quad (\text{E.39})$$

where $\varpi(\tilde{\epsilon}, \Theta) := \sup \{ \|\theta_1 - \theta_2\| : \max_{k=1:K} \text{KL}(\mathbb{P}_{\theta_1}^{(k)} \| \mathbb{P}_{\theta_2}^{(k)}) \leq [\tilde{\epsilon}/(1-\tilde{\epsilon})]^2 \}$ for any $\tilde{\epsilon} \in (0, 1)$.

Lemma 23. *When there exists an subset S such that $\min_{k \in S} n_k \geq C(p \vee \log K)$ with some constant $C > 0$, we have*

$$\inf_{\{\hat{\Sigma}^{(k)}\}_{k=1}^K} \sup_{S: |S| \geq s} \sup_{\substack{\{\hat{\theta}^{(k)*}\}_{k \in S} \in \bar{\Theta}_S \\ \mathbb{Q}_S}} \mathbb{P} \left(\bigcup_{k \in S} \left\{ \min \{ \|\hat{\mu}_1^{(k)} - \mu_1^{(k)*}\|_2 \vee \|\hat{\mu}_2^{(k)} - \mu_2^{(k)*}\|_2, \right. \right. \\ \left. \left. \|\hat{\mu}_1^{(k)} - \mu_2^{(k)*}\|_2 \vee \|\hat{\mu}_2^{(k)} - \mu_1^{(k)*}\|_2 \right\} \vee \|\hat{\Sigma}^{(k)} - \Sigma^{(k)*}\|_2 \geq C \sqrt{\frac{p + \log K}{n_k}} \right) \geq \frac{1}{10}.$$

E.3.2 Main proof of Theorem 3

Combine conclusions of Lemmas 19 and 20 to get the first lower bound. Lemma 23 implies the second one.

E.3.3 Proofs of lemmas

Proof of Lemma 17. Denote $g(w; \tilde{z}) = \log[(1-w)\tilde{z} + w]$, $g'(w; \tilde{z}) = \frac{1-\tilde{z}}{(1-w)\tilde{z} + w}$, $g''(w; \tilde{z}) = -\frac{(1-\tilde{z})^2}{[(1-w)\tilde{z} + w]^2}$ and $f(w; \mathbf{z}, \boldsymbol{\mu}) = \frac{1-w}{(2\pi)^{p/2}} \exp\{-\frac{1}{2}\|\mathbf{z} - \boldsymbol{\mu}\|_2^2\} + \frac{w}{(2\pi)^{p/2}} \exp\{-\frac{1}{2}\|\mathbf{z} + \boldsymbol{\mu}\|_2^2\}$.

By Taylor expansion,

$$\log \left[\frac{f(w'; \mathbf{z}, \boldsymbol{\mu})}{f(w; \mathbf{z}, \boldsymbol{\mu})} \right] = \frac{\partial \log f(w; \mathbf{z}, \boldsymbol{\mu})}{\partial w} \Big|_w \cdot (w' - w) + \frac{1}{2} \frac{\partial^2 \log f(w; \mathbf{z}, \boldsymbol{\mu})}{\partial w^2} \Big|_{w_0} \cdot (w' - w)^2,$$

where $w_0 = w_0(\mathbf{z}, \boldsymbol{\mu})$ is between w and w' . By the property of score function,

$$\int \frac{\partial \log f(w; \mathbf{z}, \boldsymbol{\mu})}{\partial w} d\mathbb{P}_w = 0.$$

Besides,

$$\frac{\partial^2 \log f(w; \mathbf{z}, \boldsymbol{\mu})}{\partial w^2} = \frac{\partial^2 \log [f(w; \mathbf{z}, \boldsymbol{\mu}) / ((2\pi)^{-p/2} \exp\{-\frac{1}{2}\|\mathbf{z} + \boldsymbol{\mu}\|_2^2\})]}{\partial w^2} = g''(w; \tilde{z}),$$

where $\tilde{z} = e^{-\boldsymbol{\mu}^T \mathbf{z}}$. Note that

$$-g''(w; \tilde{z}) = \frac{1}{(1-w)^2} \cdot \frac{(\tilde{z} - 1)^2}{(\tilde{z} + w/(1-w))^2} \leq \frac{1}{c_w^2},$$

for any $\tilde{z} > 0$. Therefore,

$$\begin{aligned} \text{KL}(\mathbb{P}_w \| \mathbb{P}_{w'}) &= - \int \log \left[\frac{f(w'; \mathbf{z}, \boldsymbol{\mu})}{f(w; \mathbf{z}, \boldsymbol{\mu})} \right] d\mathbb{P}_w \\ &= -\frac{1}{2} (w' - w)^2 \cdot \int g''(w_0(\mathbf{z}, \boldsymbol{\mu}); \tilde{z}) d\mathbb{P}_w \\ &\leq \frac{1}{2c_w^2} (w' - w)^2, \end{aligned}$$

which completes the proof. \square

Proof of Lemma 18. Recall that we denote distribution $\frac{1}{2}\mathcal{N}((-1/2, \mathbf{0}_{p-1}^T)^T, \mathbf{I}_p) + \frac{1}{2}\mathcal{N}((1/2 + \tilde{u}, \mathbf{0}_{p-1}^T)^T, \mathbf{I}_p)$ as $\mathbb{P}_{\tilde{u}}$ for any $\tilde{u} \in [-1, 1]$. By the bi-convexity of KL divergence, we have

$$\begin{aligned} \text{KL}(\mathbb{P}_{\tilde{u}} \| \mathbb{P}_{\tilde{u}'} &\leq \frac{1}{2} \text{KL}(\mathcal{N}((1/2 + \tilde{u}, \mathbf{0}_{p-1}^T)^T, \mathbf{I}_p) \| \mathcal{N}((1/2 + \tilde{u}', \mathbf{0}_{p-1}^T)^T, \mathbf{I}_p)) \\ &= \frac{1}{2} \text{KL}(\mathcal{N}(1/2 + \tilde{u}, 1) \| \mathcal{N}(1/2 + \tilde{u}', 1)) \\ &= \frac{1}{2} (\tilde{u} - \tilde{u}')^2, \end{aligned}$$

which completes the proof. \square

Proof of Lemma 19. WLOG, suppose $\Delta \geq 1$. It's easy to see that given any S , $\overline{\Theta}_S \supseteq \overline{\Theta}_{S,w} \cup \overline{\Theta}_{S,\beta} \cup \overline{\Theta}_{S,\delta}$, where

$$\overline{\Theta}_{S,w} = \left\{ \{\overline{\boldsymbol{\theta}}^{(k)}\}_{k \in S} : \boldsymbol{\mu}_1^{(k)} = \mathbf{1}_p / \sqrt{p}, \boldsymbol{\mu}_2^{(k)} = -\boldsymbol{\mu}_1^{(k)} = \tilde{\boldsymbol{\mu}}, \boldsymbol{\Sigma}^{(k)} = \mathbf{I}_p, w^{(k)} \in (c_w, 1 - c_w) \right\},$$

$$\overline{\Theta}_{S,\beta} = \left\{ \{\overline{\boldsymbol{\theta}}^{(k)}\}_{k \in S} : \boldsymbol{\Sigma}^{(k)} = \mathbf{I}_p, w^{(k)} = \frac{1}{2}, \|\boldsymbol{\mu}_1^{(k)}\|_2 \vee \|\boldsymbol{\mu}_2^{(k)}\|_2 \leq M, \min_{\boldsymbol{\beta}} \max_{k \in S} \|\boldsymbol{\beta}^{(k)} - \boldsymbol{\beta}\|_2 \leq h \right\},$$

$$\overline{\Theta}_{S,\delta} = \left\{ \{\overline{\boldsymbol{\theta}}^{(k)}\}_{k \in S} : \boldsymbol{\Sigma}^{(k)} = \mathbf{I}_p, w^{(k)} = \frac{1}{2}, \|\boldsymbol{\mu}_1^{(k)}\|_2 \vee \|\boldsymbol{\mu}_2^{(k)}\|_2 \leq M, \boldsymbol{\mu}_1^{(k)} = -\frac{1}{2}\boldsymbol{\mu}_0, \boldsymbol{\mu}_2^{(k)} = \frac{1}{2}\boldsymbol{\mu}_0 + \mathbf{u}, \right. \\ \left. \|\mathbf{u}\|_2 \leq 1 \right\}.$$

(i) By fixing an S and a \mathbb{Q}_S , we want to show

$$\inf_{\{\hat{\boldsymbol{\beta}}^{(k)}\}_{k=1}^K} \sup_{\{\overline{\boldsymbol{\theta}}^{(k)*}\}_{k \in S} \in \overline{\Theta}_{S,\beta}} \mathbb{P} \left(\bigcup_{k \in S} \left\{ \|\hat{\boldsymbol{\beta}}^{(k)} - \boldsymbol{\beta}^{(k)*}\|_2 \wedge \|\hat{\boldsymbol{\beta}}^{(k)} + \boldsymbol{\beta}^{(k)*}\|_2 \geq C \sqrt{\frac{p}{n_S}} \right\} \right) \geq \frac{1}{4}$$

By Lemma 4, \exists a quadrant \mathcal{Q}_v of \mathbb{R}^p and a $r/8$ -packing of $(r\mathcal{S}^p) \cap \mathcal{Q}_v$ under Euclidean norm: $\{\tilde{\boldsymbol{\mu}}_j\}_{j=1}^N$, where $r = (c\sqrt{p/n_S}) \wedge M \leq 1$ with a small constant $c > 0$ and $N \geq (\frac{1}{2})^p 8^{p-1} = \frac{1}{2} \times 4^{p-1} \geq 2^{p-1}$ when $p \geq 2$. For any $\boldsymbol{\mu} \in \mathbb{R}^p$, denote distribution $\frac{1}{2}\mathcal{N}(\boldsymbol{\mu}_0 + \boldsymbol{\mu}, \mathbf{I}_p) + \frac{1}{2}\mathcal{N}(-\boldsymbol{\mu}_0 + \boldsymbol{\mu}, \mathbf{I}_p)$ as $\mathbb{P}_{\boldsymbol{\mu}}$, where $\boldsymbol{\mu}_0$ can be any vector in \mathbb{R}^p with $\|\boldsymbol{\mu}_0\|_2 \geq 1$. Then

$$\begin{aligned} \text{LHS} &\geq \inf_{\hat{\boldsymbol{\mu}}} \sup_{\boldsymbol{\mu} \in (r\mathcal{S}^p) \cap \mathcal{Q}_v} \mathbb{P} \left(\|\hat{\boldsymbol{\mu}} - \boldsymbol{\mu}\|_2 \wedge \|\hat{\boldsymbol{\mu}} + \boldsymbol{\mu}\|_2 \geq C \sqrt{\frac{p}{n_S}} \right) \\ &\geq \inf_{\hat{\boldsymbol{\mu}}} \sup_{\boldsymbol{\mu} \in (r\mathcal{S}^p) \cap \mathcal{Q}_v} \mathbb{P} \left(\|\hat{\boldsymbol{\mu}} - \boldsymbol{\mu}\|_2 \geq C \sqrt{\frac{p}{n_S}} \right), \end{aligned} \quad (\text{E.40})$$

where the last inequality holds because it suffices to consider estimator $\hat{\boldsymbol{\mu}}$ satisfying $\hat{\boldsymbol{\mu}}(X) \in (r\mathcal{S}^p) \cap \mathcal{Q}_v$ almost surely. In addition, for any $\mathbf{x}, \mathbf{y} \in \mathcal{Q}_v$, $\|\mathbf{x} - \mathbf{y}\|_2 \leq \|\mathbf{x} + \mathbf{y}\|_2$.

By Lemma 15,

$$\begin{aligned} \text{KL} \left(\prod_{k \in S} \mathbb{P}_{\tilde{\boldsymbol{\mu}}_j}^{\otimes n_k} \cdot \mathbb{Q}_S \middle\| \prod_{k \in S} \mathbb{P}_{\tilde{\boldsymbol{\mu}}_{j'}}^{\otimes n_k} \cdot \mathbb{Q}_S \right) &= \sum_{k \in S} n_k \text{KL}(\mathbb{P}_{\tilde{\boldsymbol{\mu}}_j} \| \mathbb{P}_{\tilde{\boldsymbol{\mu}}_{j'}}) \\ &\leq \sum_{k \in S} n_k \cdot 8 \|\tilde{\boldsymbol{\mu}}_j\|_2^2 \|\tilde{\boldsymbol{\mu}}_j - \tilde{\boldsymbol{\mu}}_{j'}\|_2^2 \\ &\leq 32 n_S r^2 \\ &\leq 32 n_S c^2 \cdot \frac{2(p-1)}{n_S} \\ &\leq \frac{64c^2}{\log 2} \log N. \end{aligned}$$

By Lemma 3,

$$\text{LHS of (E.40)} \geq 1 - \frac{\log 2}{\log N} - \frac{64c^2}{\log 2} \geq 1 - \frac{1}{p-1} - \frac{1}{4} \geq \frac{1}{4},$$

when $C = c/2$, $p \geq 3$ and $c = \sqrt{\log 2}/16$.

(ii) By fixing an S and a \mathbb{Q}_S , we want to show

$$\inf_{\{\hat{\beta}^{(k)}\}_{k=1}^K} \sup_{\{\bar{\theta}^{(k)*}\}_{k \in S \in \bar{\Theta}_S}} \mathbb{P} \left(\bigcup_{k \in S} \left\{ \|\hat{\beta}^{(k)} - \beta^{(k)*}\|_2 \wedge \|\hat{\beta}^{(k)} + \beta^{(k)*}\|_2 \geq C \left[h \wedge \left(c \sqrt{\frac{p}{n_k}} \right) \right] \right\} \right) \geq \frac{1}{4}.$$

WLOG, suppose $1 \in S$. We have

$$\inf_{\hat{\beta}^{(1)}} \sup_{\{\bar{\theta}^{(k)*}\}_{k \in S \in \bar{\Theta}_S}} \mathbb{P} \left(\|\hat{\beta}^{(1)} - \beta^{(1)*}\|_2 \wedge \|\hat{\beta}^{(1)} + \beta^{(1)*}\|_2 \geq C \left[h \wedge \left(c \sqrt{\frac{p}{n_1}} \right) \right] \right) \geq \frac{1}{4}, \quad (\text{E.41})$$

By Lemma 4, \exists a quadrant \mathcal{Q}_v of \mathbb{R}^p and a $r/8$ -packing of $(r\mathcal{S}^{p-1}) \cap \mathcal{Q}_v$ under Euclidean norm: $\{\tilde{\boldsymbol{\vartheta}}_j\}_{j=1}^N$, where $r = h_\beta \wedge (c\sqrt{p/n_1}) \wedge M \leq 1$ with a small constant $c > 0$ and $N \geq (\frac{1}{2})^{p-1} 8^{p-2} = \frac{1}{2} \times 4^{p-2} \geq 2^{p-2}$ when $p \geq 3$. WLOG, assume $M \geq 2$. Denote $\tilde{\boldsymbol{\mu}}_j = (1, \tilde{\boldsymbol{\vartheta}}_j^T)^T \in \mathbb{R}^p$. Let $\boldsymbol{\mu}_1^{(k)*} = \tilde{\boldsymbol{\mu}} = (1, \mathbf{0}_{p-1})^T$ for all $k \in S \setminus \{1\}$. And let $\boldsymbol{\mu}_1^{(0)*} = \boldsymbol{\mu} = (1, \boldsymbol{\vartheta})$ with $\boldsymbol{\vartheta} \in (r\mathcal{S}^{p-1}) \cap \mathcal{Q}_v$. For any $\boldsymbol{\mu} \in \mathbb{R}^p$, denote distribution $\frac{1}{2}\mathcal{N}(\boldsymbol{\mu}, \mathbf{I}_p) + \frac{1}{2}\mathcal{N}(-\boldsymbol{\mu}, \mathbf{I}_p)$ as \mathbb{P}_μ . Then similar to the arguments in (i),

$$\begin{aligned} \text{LHS} &\geq \inf_{\hat{\boldsymbol{\mu}}} \sup_{\substack{\boldsymbol{\vartheta} \in (r\mathcal{S}^{p-1}) \cap \mathcal{Q}_v \\ \boldsymbol{\mu} = (1, \boldsymbol{\vartheta})^T}} \mathbb{P} \left(\|\hat{\boldsymbol{\mu}} - \boldsymbol{\mu}\|_2 \wedge \|\hat{\boldsymbol{\mu}} + \boldsymbol{\mu}\|_2 \geq C \left[h \wedge \left(c \sqrt{\frac{p}{n_1}} \right) \right] \right) \\ &\geq \inf_{\hat{\boldsymbol{\mu}}} \sup_{\substack{\boldsymbol{\vartheta} \in (r\mathcal{S}^{p-1}) \cap \mathcal{Q}_v \\ \boldsymbol{\mu} = (1, \boldsymbol{\vartheta})^T}} \mathbb{P} \left(\|\hat{\boldsymbol{\mu}} - \boldsymbol{\mu}\|_2 \geq C \left[h \wedge \left(c \sqrt{\frac{p}{n_1}} \right) \right] \right). \end{aligned}$$

Then by Lemma 15,

$$\begin{aligned} \text{KL} \left(\prod_{k \in S \setminus \{1\}} \mathbb{P}_{\tilde{\boldsymbol{\mu}}_k}^{\otimes n_k} \cdot \mathbb{P}_{\tilde{\boldsymbol{\mu}}_j}^{\otimes n_1} \cdot \mathbb{Q}_S \left\| \prod_{k \in S \setminus \{1\}} \mathbb{P}_{\tilde{\boldsymbol{\mu}}_k}^{\otimes n_k} \cdot \mathbb{P}_{\tilde{\boldsymbol{\mu}}_{j'}}^{\otimes n_1} \cdot \mathbb{Q}_S \right. \right) &= n_1 \text{KL}(\mathbb{P}_{\tilde{\boldsymbol{\mu}}_j} \| \mathbb{P}_{\tilde{\boldsymbol{\mu}}_{j'}}) \\ &\leq n_1 \cdot 8 \|\tilde{\boldsymbol{\mu}}_j\|_2^2 \|\tilde{\boldsymbol{\mu}}_j - \tilde{\boldsymbol{\mu}}_{j'}\|_2^2 \\ &\leq 32n_1 r^2 \\ &\leq 32n_1 c^2 \cdot \frac{3(p-2)}{n_1} \\ &\leq \frac{96c^2}{\log 2} \log N, \end{aligned}$$

when $n_1 \geq (c^2 \vee M^{-2})p$ and $p \geq 3$. By Fano's lemma (See Corollary 2.6 in [Tsybakov \(2009\)](#)),

$$\text{LHS of (E.41)} \geq 1 - \frac{\log 2}{\log N} - \frac{96c^2}{\log 2} \geq 1 - \frac{1}{p-2} - \frac{1}{4} \geq \frac{1}{4},$$

when $C = 1/2$, $p \geq 4$ and $c = \sqrt{(\log 2)/384}$.

(iii) By fixing an S and a \mathbb{Q}_S , we want to show

$$\inf_{\{\hat{\boldsymbol{\theta}}^{(k)}\}_{k=1}^K} \sup_{\{\bar{\boldsymbol{\theta}}^{(k)*}\}_{k \in S} \in \bar{\Theta}_S} \mathbb{P} \left(\bigcup_{k \in S} \left\{ \|\hat{\boldsymbol{\beta}}^{(k)} - \boldsymbol{\beta}^{(k)*}\|_2 \wedge \|\hat{\boldsymbol{\beta}}^{(k)} + \boldsymbol{\beta}^{(k)*}\|_2 \geq C \left[h \wedge \left(c \sqrt{\frac{\log K}{n_k}} \right) \right] \right\} \right) \geq \frac{1}{4}.$$

Suppose $\mathbf{v} = \mathbf{1}_p$ and denote the associated quadrant $\mathcal{Q}_{\mathbf{v}} = \mathbb{R}_+^p$, $\Upsilon_S = \{\{\boldsymbol{\mu}^{(k)}\}_{k \in S} : \boldsymbol{\mu}^{(k)} \in \mathbb{R}_+^p, \min_{\boldsymbol{\mu}} \max_{k \in S} \|\boldsymbol{\mu}^{(k)} - \boldsymbol{\mu}\|_2 \leq h, \|\boldsymbol{\mu}^{(k)}\|_2 \leq M\}$. Let $r_k = h \wedge (c \sqrt{\log K / n_k}) \wedge M$ with a small constant $c > 0$ for $k \in S$. For any $\mathbf{M} = \{\boldsymbol{\mu}^{(k)}\}_{k \in S}$, where $\boldsymbol{\mu}^{(k)} \in \mathbb{R}^p$, denote distribution $\prod_{k \in S} [\frac{1}{2}\mathcal{N}(\boldsymbol{\mu}^{(k)}, \mathbf{I}_p) + \frac{1}{2}\mathcal{N}(-\boldsymbol{\mu}^{(k)}, \mathbf{I}_p)]^{\otimes n_k}$ as $\mathbb{P}_{\mathbf{M}}$, and the joint distribution of $\mathbb{P}_{\mathbf{M}}$ and \mathbb{Q}_S as $\mathbb{P}_{\mathbf{M}} \cdot \mathbb{Q}_S$. And denote distribution $(1 - \bar{w})\mathcal{N}(\boldsymbol{\mu}, \mathbf{I}_p) + \bar{w}\mathcal{N}(-\boldsymbol{\mu}, \mathbf{I}_p)$ as $\mathbb{P}_{\boldsymbol{\mu}}$ for any $\boldsymbol{\mu} \in \mathbb{R}^p$. Similar to the arguments in (i), since it suffices to consider the estimators $\{\hat{\boldsymbol{\mu}}^{(k)}\}_{k \in S}$ satisfying $\{\hat{\boldsymbol{\mu}}^{(k)}\}_{k \in S} \in \Upsilon_S$ almost surely and $\|\mathbf{x} - \mathbf{y}\|_2 \leq \|\mathbf{x} + \mathbf{y}\|_2$ for any $\mathbf{x}, \mathbf{y} \in \mathbb{R}_+^p$, we have

$$\begin{aligned} \text{LHS} &\geq \inf_{\{\hat{\boldsymbol{\mu}}^{(k)}\}_{k \in S}} \sup_{\{\boldsymbol{\mu}^{(k)}\}_{k \in S} \in \Upsilon_S} \mathbb{P}_{\{\boldsymbol{\mu}^{(k)}\}_{k \in S}} \cdot \mathbb{Q}_S \left(\bigcup_{k \in S} \left\{ \|\hat{\boldsymbol{\mu}}^{(k)} - \boldsymbol{\mu}^{(k)}\|_2 \wedge \|\hat{\boldsymbol{\mu}}^{(k)} + \boldsymbol{\mu}^{(k)}\|_2 \right. \right. \\ &\quad \left. \left. \geq C \left[h \wedge \left(c \sqrt{\frac{\log K}{n_k}} \right) \right] \right\} \right) \\ &\geq \inf_{\{\hat{\boldsymbol{\mu}}^{(k)}\}_{k \in S}} \sup_{\{\boldsymbol{\mu}^{(k)}\}_{k \in S} \in \Upsilon_S} \mathbb{P}_{\{\boldsymbol{\mu}^{(k)}\}_{k \in S}} \cdot \mathbb{Q}_S \left(\bigcup_{k \in S} \left\{ \|\hat{\boldsymbol{\mu}}^{(k)} - \boldsymbol{\mu}^{(k)}\|_2 \geq C \left[h \wedge \left(c \sqrt{\frac{\log K}{n_k}} \right) \right] \right\} \right), \end{aligned}$$

Consider $\mathbf{M}^{(k)} = \{\boldsymbol{\mu}^{(j)}\}_{j \in S}$ where $\boldsymbol{\mu}^{(j)} = \frac{r_j}{\sqrt{p-3/4}} \cdot \mathbf{1}_p + \boldsymbol{\mu}_0$ for $j \neq k$ and $\boldsymbol{\mu}^{(k)} = \frac{r_k}{2\sqrt{p-3/4}} \cdot \mathbf{1}_p + \boldsymbol{\mu}_0$, where $\boldsymbol{\mu}_0 = (1, \mathbf{0}_{p-1}^T)^T$. Define two new “distances” (which are not rigorously distances because triangle inequalities and the definiteness do not hold) between $\mathbf{M} = \{\boldsymbol{\mu}^{(k)}\}_{k \in S}$ and as $\mathbf{M}' = \{\boldsymbol{\mu}'^{(k)}\}_{k \in S}$

$$\begin{aligned} \tilde{d}(\mathbf{M}, \mathbf{M}') &:= \sum_{k \in S} \mathbb{1} \left(\|\boldsymbol{\mu}^{(k)} - \boldsymbol{\mu}'^{(k)}\|_2 \geq \frac{r_k}{2\sqrt{p-3/4}} \right), \\ \tilde{d}'(\mathbf{M}, \mathbf{M}') &:= \sum_{k \in S} \mathbb{1} \left(\|\boldsymbol{\mu}^{(k)} - \boldsymbol{\mu}'^{(k)}\|_2 \geq \frac{r_k}{4\sqrt{p-3/4}} \right). \end{aligned}$$

Therefore $\tilde{d}(\mathbf{M}^{(k)}, \mathbf{M}^{(k')}) = 2$ when $k \neq k'$. For $\{\hat{\boldsymbol{\mu}}^{(k)}\}_{k \in S}$, define $\psi^* = \arg \min_{k \in S} \tilde{d}'(\{\hat{\boldsymbol{\mu}}^{(k)}\}_{k \in S}, \mathbf{M}^{(k)})$. Because $\tilde{d}(\mathbf{M}_1, \mathbf{M}_2) \leq \tilde{d}'(\mathbf{M}_1, \mathbf{M}_2) + \tilde{d}'(\mathbf{M}_2, \mathbf{M}_3)$ for any $\mathbf{M}_1, \mathbf{M}_2$ and \mathbf{M}_3 , it's

easy to see that

$$\begin{aligned}
& \inf_{\{\hat{\boldsymbol{\mu}}^{(k)}\}_{k \in S}} \sup_{\{\boldsymbol{\mu}^{(k)}\}_{k \in S} \in \Upsilon_S} \mathbb{P}_{\{\boldsymbol{\mu}^{(k)}\}_{k \in S}} \cdot \mathbb{Q}_S \left(\bigcup_{k \in S} \left\{ \|\hat{\boldsymbol{\mu}}^{(k)} - \boldsymbol{\mu}^{(k)}\|_2 \geq \frac{r_k}{4\sqrt{p-3/4}} \right\} \right) \\
& \geq \inf_{\{\hat{\boldsymbol{\mu}}^{(k)}\}_{k \in S}} \sup_{k \in S} \mathbb{P}_{\mathbf{M}^{(k)}} \left(\tilde{d}'(\{\hat{\boldsymbol{\mu}}_1^{(k)}\}_{k \in S}, \mathbf{M}^{(k)}) \geq 1 \right) \\
& \geq \inf_{\{\hat{\boldsymbol{\mu}}^{(k)}\}_{k \in S}} \sup_{k \in S} \mathbb{P}_{\mathbf{M}^{(k)}} (\psi^* \neq k) \\
& \geq \inf_{\psi} \sup_{k \in S} \mathbb{P}_{\mathbf{M}^{(k)}} (\psi \neq k). \tag{E.42}
\end{aligned}$$

By Lemma 15,

$$\begin{aligned}
\text{KL}(\mathbb{P}_{\mathbf{M}^{(k)}} \cdot \mathbb{Q}_S \| \mathbb{P}_{\mathbf{M}^{(k')}} \cdot \mathbb{Q}_S) &= n_k \text{KL} \left(\mathbb{P}^{\otimes n_k}_{\frac{r_k}{\sqrt{p-3/4}} \mathbf{1}_p + \boldsymbol{\mu}_0} \| \mathbb{P}^{\otimes n_k}_{\frac{r_k}{2\sqrt{p-3/4}} \mathbf{1}_p + \boldsymbol{\mu}_0} \right) \\
&\quad + n_{k'} \text{KL} \left(\mathbb{P}^{\otimes n_{k'}}_{\frac{r_{k'}}{\sqrt{p-3/4}} \mathbf{1}_p + \boldsymbol{\mu}_0} \| \mathbb{P}^{\otimes n_{k'}}_{\frac{r_{k'}}{2\sqrt{p-3/4}} \mathbf{1}_p + \boldsymbol{\mu}_0} \right) \\
&\leq n_k \cdot 8 \left\| \frac{r_k}{\sqrt{p-3/4}} \mathbf{1}_p + \boldsymbol{\mu}_0 \right\|_2^2 \left\| \frac{r_k}{2\sqrt{p-3/4}} \mathbf{1}_p \right\|_2^2 \\
&\quad + n_{k'} \cdot 8 \left\| \frac{r_{k'}}{\sqrt{p-3/4}} \mathbf{1}_p + \boldsymbol{\mu}_0 \right\|_2^2 \left\| \frac{r_{k'}}{2\sqrt{p-3/4}} \mathbf{1}_p \right\|_2^2 \\
&\leq n_k \cdot 8 \cdot 2 \cdot (2r_k^2 + 1) \cdot \frac{1}{4} \cdot 2r_k^2 + n_{k'} \cdot 8 \cdot 2 \cdot (2r_{k'}^2 + 1) \cdot \frac{1}{4} \cdot 2r_{k'}^2 \\
&\leq 16c^2 \log K,
\end{aligned}$$

when $p \geq 3$. By Fano's lemma (See Corollary 2.6 in [Tsybakov \(2009\)](#)),

$$\text{RHS of (E.42)} \geq 1 - \frac{\log 2}{\log K} - 16c^2 \geq \frac{1}{4},$$

when $K \geq 3$, $c = \sqrt{1/160}$, and $\min_{k \in S} n_k \geq (c^2 \vee M^{-2}) \log K$.

(iv) We want to show

$$\begin{aligned}
& \inf_{\{\hat{\boldsymbol{\theta}}^{(k)}\}_{k=1}^K} \sup_{S: |S| \geq s} \sup_{\{\bar{\boldsymbol{\theta}}^{(k)*}\}_{k \in S} \in \bar{\Theta}_{S,w}} \mathbb{P} \left(\bigcup_{k \in S} \left\{ |\hat{w}^{(k)} - w^{(k)*}| \wedge |1 - \hat{w}^{(k)} - w^{(k)*}| \right. \right. \\
& \quad \left. \left. \geq C \sqrt{\frac{\log K}{n_k}} \right\} \right) \geq \frac{1}{4}.
\end{aligned}$$

The argument is similar to part (iii). The only two differences here are that the dimension of interested parameter w equals 1, and Lemma 15 is replaced by Lemma 17.

(v) We want to show

$$\inf_{\{\widehat{\boldsymbol{\mu}}^{(k)}\}_{k=1}^K} \sup_{S: |S| \geq s} \sup_{\{\boldsymbol{\theta}^{(k)*}\}_{k \in S} \in \overline{\Theta}_{S, \delta} \atop \mathbb{Q}_S} \mathbb{P} \left(\bigcup_{k \in S} \left\{ |\widehat{\delta}^{(k)} - \delta^{(k)*}| \wedge |\widehat{\delta}^{(k)} + \delta^{(k)*}| \right. \right. \\ \left. \left. \geq C \sqrt{\frac{\log K}{n_k}} \right\} \right) \geq \frac{1}{4},$$

The argument is similar to (iii). The only two differences here are that the dimension of interested parameter δ equals 1, and Lemma 15 is replaced by Lemma 18.

Finally, we get the desired conclusion by combining (i)-(v). \square

Proof of Lemma 20. Let $\tilde{\epsilon} = \frac{K-s}{s}$ and $\tilde{\epsilon}' = \frac{K-s}{K}$. Since $s/K \geq c > 0$, $\tilde{\epsilon} \lesssim \tilde{\epsilon}'$. Denote $\Upsilon_S = \{\{\boldsymbol{\mu}^{(k)}\}_{k \in S} : \boldsymbol{\mu}^{(k)} \in \mathbb{R}_+^p, \min_{\boldsymbol{\mu}} \max_{k \in S} \|\boldsymbol{\mu}^{(k)} - \boldsymbol{\mu}\|_2 \leq h_\beta/2, \|\boldsymbol{\mu}^{(k)}\|_2 \leq M\}$. For any $\boldsymbol{\mu} \in \mathbb{R}$, denote distribution $\frac{1}{2}\mathcal{N}(\boldsymbol{\mu}, \mathbf{I}_p) + \frac{1}{2}\mathcal{N}(-\boldsymbol{\mu}, \mathbf{I}_p)$ as $\mathbb{P}_{\boldsymbol{\mu}}$, and denote $\prod_{k \in S} \mathbb{P}_{\boldsymbol{\mu}^{(k)}}^{\otimes n_k}$ as $\mathbb{P}_{\{\boldsymbol{\mu}^{(k)}\}_{k \in S}}$. Note that $\boldsymbol{\beta}^{(k)} = 2\boldsymbol{\mu}^{(k)}$ for $\mathbb{P}_{\boldsymbol{\mu}^{(k)}}$ with $\{\boldsymbol{\mu}^{(k)}\}_{k \in S} \in \Upsilon_S$. Then it suffices to show

$$\inf_{\{\widehat{\boldsymbol{\mu}}^{(k)}\}_{k=1}^K} \sup_{S: |S| \geq s} \sup_{\{\boldsymbol{\mu}^{(k)}\}_{k \in S} \in \Upsilon_S \atop \mathbb{Q}_S} \mathbb{P} \left(\max_{k \in S} \|\widehat{\boldsymbol{\mu}}^{(k)} - \boldsymbol{\mu}^{(k)}\|_2 \geq C_1 \tilde{\epsilon}' \sqrt{\frac{1}{\max_{k=1:K} n_k}} \right) \geq \frac{1}{10}, \quad (\text{E.43})$$

where $\mathbb{P} = \mathbb{P}_{\{\boldsymbol{\mu}^{(k)}\}_{k \in S}} \cdot \mathbb{Q}_S$. WLOG, assume $M \geq 1$. For any $\tilde{\boldsymbol{\mu}}_1, \tilde{\boldsymbol{\mu}}_2 \in \mathbb{R}^p$ with $\|\tilde{\boldsymbol{\mu}}_1\|_2 = \|\tilde{\boldsymbol{\mu}}_2\|_2 = 1$, by Lemma 15,

$$\max_{k=1:K} \text{KL}(\mathbb{P}_{\tilde{\boldsymbol{\mu}}_1}^{\otimes n_k} \|\mathbb{P}_{\tilde{\boldsymbol{\mu}}_2}^{\otimes n_k}) \leq \max_{k=1:K} n_k \cdot 8 \|\tilde{\boldsymbol{\mu}}_1 - \tilde{\boldsymbol{\mu}}_2\|_2^2.$$

Let $8 \max_{k=1:K} n_k \cdot \|\tilde{\boldsymbol{\mu}}_1 - \tilde{\boldsymbol{\mu}}_2\|_2^2 \leq (\frac{\tilde{\epsilon}'}{1-\tilde{\epsilon}'})^2$, then $\|\tilde{\boldsymbol{\mu}}_1 - \tilde{\boldsymbol{\mu}}_2\|_2 \leq C \sqrt{\frac{1}{\max_{k=1:K} n_k}} \tilde{\epsilon}'$ for some constant $C > 0$. Then (E.43) follows by Lemma 22. \square

Proof of Lemma 21. The proof is similar to the proof of Theorem 5.1 in Chen et al. (2018), so we omit it here. \square

Proof of Lemma 22. It's easy to see that

$$\text{LHS of (E.39)} \geq \inf_{\hat{\boldsymbol{\theta}}} \sup_{\substack{\boldsymbol{\theta} \in \Theta \\ \mathbb{Q}_S}} \mathbb{E}_{S \sim \mathbb{P}_s} \left[\mathbb{P}_{(S, \boldsymbol{\theta}, \mathbb{Q}_S)} \left(\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}\| \geq C \varpi \left(\frac{K-s}{50K}, \Theta \right) \right) \right],$$

where \mathbb{P}_s can be any probability measure on all subsets of $1 : K$ with $\mathbb{P}_s(|S| \geq s) = 1$.

Consider a special distribution $\tilde{\mathbb{P}}_s$ as \mathbb{P}_s :

$$\tilde{\mathbb{P}}_s(S = S') = \frac{\mathbb{P}_{|S^c| \sim \text{Bin}(K, \frac{K-s}{50K})}(|S| = |S'|)}{\mathbb{P}_{|S^c| \sim \text{Bin}(K, \frac{K-s}{50K})}(|S^c| \leq \frac{41(K-s)}{50})} \cdot \frac{1}{\binom{K}{|S'|}},$$

for any S' with $|(S')^c| \leq \frac{41(K-s)}{50}$. Given S , consider the distribution of $\{\mathbf{x}^{(k)}\}_{k=1}^K$ as

$$\mathbb{P}^S = \prod_{k \in S} \mathbb{P}_\theta^{(k)} \cdot \prod_{k \notin S} \mathbb{Q}^{(k)}$$

Then consider the distribution of $\{\mathbf{x}^{(k)}\}_{k=1}^K$ as

$$\mathbb{P}' = \sum_{S: |S^c| \leq \frac{41(K-s)}{50}} \tilde{\mathbb{P}}_s(S) \cdot \mathbb{P}^S.$$

It's easy to see that \mathbb{P}' is the same as $\mathbb{P}_{(\frac{K-s}{50K}, \theta, \{\mathbb{Q}^{(k)}\}_{k=1}^K)}$ conditioning on the event $\{S : |S^c| \leq \frac{41(K-s)}{50}\}$. Therefore,

$$\begin{aligned} & \inf_{\hat{\theta}} \sup_{\substack{\theta \in \Theta \\ \mathbb{Q}_S}} \mathbb{E}_{S \sim \tilde{\mathbb{P}}_s} \left[\mathbb{P}_{(S, \theta, \mathbb{Q}_S)} \left(\|\hat{\theta} - \theta\| \geq C\varpi \left(\frac{K-s}{50K}, \Theta \right) \right) \right] \\ & \geq \inf_{\hat{\theta}} \sup_{\substack{\theta \in \Theta \\ \{\mathbb{Q}^{(k)}\}_{k=1}^K}} \mathbb{E}_{S \sim \tilde{\mathbb{P}}_s} \left[\mathbb{P}^S \left(\|\hat{\theta} - \theta\| \geq C\varpi \left(\frac{K-s}{50K}, \Theta \right) \right) \right] \\ & = \inf_{\hat{\theta}} \sup_{\substack{\theta \in \Theta \\ \{\mathbb{Q}^{(k)}\}_{k=1}^K}} \mathbb{P}' \left(\|\hat{\theta} - \theta\| \geq C\varpi \left(\frac{K-s}{50K}, \Theta \right) \right) \\ & \geq \inf_{\hat{\theta}} \sup_{\substack{\theta \in \Theta \\ \{\mathbb{Q}^{(k)}\}_{k=1}^K}} \mathbb{P}_{(\frac{K-s}{50K}, \theta, \{\mathbb{Q}^{(k)}\}_{k=1}^K)} \left(\|\hat{\theta} - \theta\| \geq C\varpi \left(\frac{K-s}{50K}, \Theta \right) \middle| |S^c| \leq \frac{41(K-s)}{50} \right) \\ & \geq \inf_{\hat{\theta}} \sup_{\substack{\theta \in \Theta \\ \{\mathbb{Q}^{(k)}\}_{k=1}^K}} \mathbb{P}_{(\frac{K-s}{50K}, \theta, \{\mathbb{Q}^{(k)}\}_{k=1}^K)} \left(\|\hat{\theta} - \theta\| \geq C\varpi \left(\frac{K-s}{50K}, \Theta \right) \right) - \mathbb{P}_{|S^c| \sim \text{Bin}(K, \frac{K-s}{50K})} \left(|S^c| > \frac{41(K-s)}{50} \right) \\ & \geq \frac{1}{2} - \exp \left\{ -\frac{\frac{1}{2}[\frac{4}{5}(K-s)]^2}{K \cdot \frac{K-s}{50K} \left(1 - \frac{K-s}{50K}\right) + \frac{1}{3} \cdot \frac{4}{5}(K-s)} \right\} \\ & \geq \frac{1}{2} - \exp \left\{ -\frac{\frac{1}{2} \cdot \left(\frac{4}{5}\right)^2}{\frac{1}{50} + \frac{4}{15}} \right\} \\ & \geq \frac{1}{10}, \end{aligned}$$

where the last third inequality comes from Bernstein's inequality, application of Lemma 17 and the fact that $d_{\text{TV}}^2(\mathbb{P}_{\theta_1}, \mathbb{P}_{\theta_2}) \leq \text{KL}(\mathbb{P}_{\theta_1} \parallel \mathbb{P}_{\theta_2})$. \square

Proof of Lemma 23. (i) We want to show

$$\inf_{\{\hat{\Sigma}^{(k)}\}_{k=1}^K} \sup_{S: |S| \geq s} \sup_{\substack{\{\theta^{(k)*}\}_{k \in S} \in \bar{\Theta}_S \\ \mathbb{Q}_S}} \mathbb{P} \left(\bigcup_{k \in S} \left\{ \|\hat{\Sigma}^{(k)} - \Sigma^{(k)*}\|_2 \geq C\sqrt{\frac{p}{n_k}} \right\} \right) \geq \frac{1}{10}.$$

Fix S and some \mathbb{Q}_S . WLOG, assume $1 \in S$. Then it suffices to show

$$\inf_{\widehat{\Sigma}^{(k)}} \sup_{\{\bar{\theta}^{(k)*}\}_{k \in S} \in \bar{\Theta}_S} \mathbb{P} \left(\|\widehat{\Sigma}^{(k)} - \Sigma^{(k)*}\|_2 \geq C \sqrt{\frac{p}{n_k}} \right) \geq \frac{1}{10}.$$

Consider a special subset of $\bar{\Theta}_S$ as

$$\bar{\Theta}_{S, \Sigma} = \{\bar{\theta} : w = 1/2, \boldsymbol{\mu}_1 = \boldsymbol{\mu}_2 = 0, \Sigma = \Sigma(\gamma), \gamma \in \{0, 1\}^p\},$$

where

$$\Sigma(\gamma) = \begin{pmatrix} \gamma_1 \mathbf{e}_1^T \\ \vdots \\ \gamma_p \mathbf{e}_1^T \end{pmatrix} \cdot \tau + \mathbf{I}_p,$$

and $\tau > 0$ is a small constant which we will specify later. For any $\gamma \in \{0, 1\}^p$, denote $\mathcal{N}(\mathbf{0}, \Sigma(\gamma))$ as \mathbb{P}_γ . Therefore it suffices to show

$$\inf_{\widehat{\Sigma}^{(1)}} \sup_{\gamma \in \{0, 1\}^p} \mathbb{P} \left(\|\widehat{\Sigma}^{(1)} - \Sigma^{(1)*}\|_2 \geq C \sqrt{\frac{p}{n_1}} \right) \geq \frac{1}{10}. \quad (\text{E.44})$$

Note that for any $\widehat{\Sigma}^{(1)}$, we can define $\widehat{\gamma} = \arg \min_{\gamma \in \{0, 1\}^p} \|\widehat{\Sigma}^{(1)} - \Sigma(\gamma)\|_2$. Then by triangle inequality and definition of $\widehat{\gamma}$, $\|\widehat{\Sigma}^{(1)} - \Sigma(\gamma)\|_2 \geq \|\Sigma(\widehat{\gamma}) - \Sigma(\gamma)\|_2/2$. Therefore

$$\text{LHS of (E.44)} \geq \inf_{\widehat{\gamma} \in \{0, 1\}^p} \sup_{\gamma \in \{0, 1\}^p} \mathbb{P} \left(\|\Sigma(\widehat{\gamma}) - \Sigma(\gamma)\|_2 \geq 2C \sqrt{\frac{p}{n_1}} \right). \quad (\text{E.45})$$

Let $\tau = c\sqrt{1/n_1}$ where $c > 0$ is a small constant. Since $\|\Sigma(\widehat{\gamma}) - \Sigma(\gamma)\|_2 \leq \tau$ for any $\widehat{\gamma}$ and $\gamma \in \{0, 1\}^p$, by Lemma D.2 in [Duan and Wang \(2022\)](#),

$$\text{LHS of (E.45)} \geq \frac{\inf_{\widehat{\gamma} \in \{0, 1\}^p} \sup_{\gamma \in \{0, 1\}^p} \mathbb{E} \|\Sigma(\widehat{\gamma}) - \Sigma(\gamma)\|_2^2 - 4C^2 \cdot \frac{p}{n_1}}{(c^2 - 4C^2) \frac{p}{n_1}}. \quad (\text{E.46})$$

Applying Assouad's lemma (Theorem 2.12 in [Tsybakov \(2009\)](#) or Lemma 2 in [Cai and Zhou \(2012\)](#)), we get

$$\begin{aligned} \inf_{\widehat{\gamma} \in \{0, 1\}^p} \sup_{\gamma \in \{0, 1\}^p} \mathbb{E} \|\Sigma(\widehat{\gamma}) - \Sigma(\gamma)\|_2^2 &\geq \frac{p}{8} \min_{\rho_H(\gamma, \gamma') \geq 1} \left[\frac{\|\Sigma(\gamma) - \Sigma(\gamma')\|_2^2}{\rho_H(\gamma, \gamma')} \right] \\ &\cdot \left[1 - \max_{\rho_H(\gamma, \gamma')=1} (\text{KL}(\mathbb{P}_\gamma^{\otimes n_1} \|\mathbb{P}_{\gamma'}^{\otimes n_1}))^{1/2} \right], \quad (\text{E.47}) \end{aligned}$$

where ρ_H is the Hamming distance. For the first term on the RHS, it's easy to see that

$$\|\Sigma(\gamma) - \Sigma(\gamma')\|_2^2 = \tau^2 \rho_H(\gamma, \gamma'),$$

for any γ and $\gamma' \in \{0, 1\}^p$. For the second term, by the density form of Gaussian distribution, we can show that if $\rho_H(\gamma, \gamma') = 1$, then

$$\begin{aligned} \text{KL}(\mathbb{P}_\gamma^{\otimes n_1} \parallel \mathbb{P}_{\gamma'}^{\otimes n_1}) &= n_1 \text{KL}(\mathbb{P}_\gamma \parallel \mathbb{P}_{\gamma'}) \\ &\leq n_1 \cdot \frac{1}{2} \left\{ \log(|\Sigma(\gamma')|/|\Sigma(\gamma)|) - \text{Tr} [(\Sigma(\gamma)^{-1} - \Sigma(\gamma')^{-1})\Sigma(\gamma)] \right\} \\ &\leq n_1 \cdot \frac{1}{4} \tau^2 \\ &\leq \frac{c^2}{4}. \end{aligned}$$

Plugging this back into (E.47), combining with (E.46), we have

$$\text{LHS of (E.45)} \geq \frac{c^2 \cdot \frac{p}{8n_1} (1 - \frac{c}{2}) - 4C^2 \cdot \frac{p}{n_1}}{(c^2 - 4C^2) \frac{p}{n_1}} \geq \frac{1}{10},$$

when $c = 2/9$ and $C \leq c/\sqrt{324}$.

(ii) We want to show

$$\inf_{\{\widehat{\Sigma}^{(k)}\}_{k=1}^K} \sup_{S: |S| \geq s} \sup_{\{\bar{\theta}^{(k)*}\}_{k \in S} \in \bar{\Theta}_S} \mathbb{P} \left(\bigcup_{k \in S} \left\{ \|\widehat{\Sigma}^{(k)} - \Sigma^{(k)*}\|_2 \geq C \sqrt{\frac{\log K}{n_k}} \right\} \right) \geq \frac{1}{10}.$$

The proof idea is similar to part (iii) of the proof of Lemma 19, so we omit the details here. It suffices to consider $\mathbf{M}^{(k)} = \{\Sigma^{(j)}\}_{j=1}^K$ where $\Sigma^{(j)} = \mathbf{I}_p$ when $j \neq k$ and $\Sigma^{(k)} = \mathbf{I}_p + \sqrt{\log K/n_k} \cdot \mathbf{e}_1 \mathbf{e}_1^T$.

□

E.4 Proof of Theorem 2

We claim that with probability at least $1 - C \exp\{-C'p\}$,

$$R_{\bar{\theta}^{(k)*}}(\widehat{\mathcal{C}}^{(k)[t]}) - R_{\bar{\theta}^{(k)*}}(\mathcal{C}_{\bar{\theta}^{(k)*}}) \lesssim d^2(\widehat{\theta}^{(k)[t]}, \theta^{(k)*}). \quad (\text{E.48})$$

Then the conclusion immediately follows from Theorem 1. Hence it suffices to verify the claim. For convenience, we write $\widehat{\mathcal{C}}^{(k)[t]} = \mathcal{C}_{\widehat{\theta}^{(k)[t]}}$ simply as $\mathcal{C}_{\widehat{\theta}^{(k)}}$ and $\widehat{\theta}^{(k)[t]}$ as $\widehat{\theta}^{(k)}$.

By simple calculations, we have

$$\begin{aligned} R_{\bar{\theta}^{(k)*}}(\mathcal{C}_{\widehat{\theta}^{(k)}}) &= (1 - w^{(k)*}) \Phi \left(\frac{-\log(\frac{1-\widehat{w}^{(k)}}{\widehat{w}^{(k)}}) - \widehat{\delta}^{(k)} + (\widehat{\beta}^{(k)})^T \boldsymbol{\mu}_1^{(k)*}}{\sqrt{(\widehat{\beta}^{(k)})^T \Sigma^{(k)*} \widehat{\beta}^{(k)}}} \right) \\ &\quad + w^{(k)*} \Phi \left(\frac{\log(\frac{1-\widehat{w}^{(k)}}{\widehat{w}^{(k)}}) + \widehat{\delta}^{(k)} - (\widehat{\beta}^{(k)})^T \boldsymbol{\mu}_2^{(k)*}}{\sqrt{(\widehat{\beta}^{(k)})^T \Sigma^{(k)*} \widehat{\beta}^{(k)}}} \right), \end{aligned}$$

$$R_{\hat{\theta}^{(k)*}}(\mathcal{C}_{\theta^{(k)*}}) = (1 - w^{(k)*})\Phi \left(\frac{-\log(\frac{1-w^{(k)*}}{w^{(k)*}}) - \delta^{(k)*} + (\beta^{(k)*})^T \mu_1^{(k)*}}{\sqrt{(\beta^{(k)*})^T \Sigma^{(k)*} \beta^{(k)*}}} \right) \\ + w^{(k)*}\Phi \left(\frac{\log(\frac{1-w^{(k)*}}{w^{(k)*}}) + \delta^{(k)*} - (\beta^{(k)*})^T \mu_2^{(k)*}}{\sqrt{(\beta^{(k)*})^T \Sigma^{(k)*} \beta^{(k)*}}} \right).$$

Then by Taylor expansion,

$$R_{\hat{\theta}^{(k)*}}(\mathcal{C}_{\hat{\theta}^{(k)*}}) - R_{\hat{\theta}^{(k)*}}(\mathcal{C}_{\theta^{(k)*}}) \leq (1 - w^{(k)*})\Phi' \left(\frac{-\log(\frac{1-w^{(k)*}}{w^{(k)*}}) - \delta^{(k)*} + (\beta^{(k)*})^T \mu_1^{(k)*}}{\sqrt{(\beta^{(k)*})^T \Sigma^{(k)*} \beta^{(k)*}}} \right) \\ \cdot \left[\frac{-\log(\frac{1-\hat{w}^{(k)*}}{\hat{w}^{(k)*}}) - \hat{\delta}^{(k)*} + (\hat{\beta}^{(k)*})^T \mu_1^{(k)*}}{\sqrt{(\hat{\beta}^{(k)*})^T \Sigma^{(k)*} \hat{\beta}^{(k)*}}} - \frac{-\log(\frac{1-w^{(k)*}}{w^{(k)*}}) - \delta^{(k)*} + (\beta^{(k)*})^T \mu_1^{(k)*}}{\sqrt{(\beta^{(k)*})^T \Sigma^{(k)*} \beta^{(k)*}}} \right] \\ + w^{(k)*}\Phi' \left(\frac{\log(\frac{1-w^{(k)*}}{w^{(k)*}}) + \delta^{(k)*} - (\beta^{(k)*})^T \mu_2^{(k)*}}{\sqrt{(\beta^{(k)*})^T \Sigma^{(k)*} \beta^{(k)*}}} \right) \\ \cdot \left[\frac{\log(\frac{1-\hat{w}^{(k)*}}{\hat{w}^{(k)*}}) + \hat{\delta}^{(k)*} - (\hat{\beta}^{(k)*})^T \mu_2^{(k)*}}{\sqrt{(\hat{\beta}^{(k)*})^T \Sigma^{(k)*} \hat{\beta}^{(k)*}}} - \frac{\log(\frac{1-w^{(k)*}}{w^{(k)*}}) + \delta^{(k)*} - (\beta^{(k)*})^T \mu_2^{(k)*}}{\sqrt{(\beta^{(k)*})^T \Sigma^{(k)*} \beta^{(k)*}}} \right] \\ + C \left[\frac{-\log(\frac{1-\hat{w}^{(k)*}}{\hat{w}^{(k)*}}) - \hat{\delta}^{(k)*} + (\hat{\beta}^{(k)*})^T \mu_1^{(k)*}}{\sqrt{(\hat{\beta}^{(k)*})^T \Sigma^{(k)*} \hat{\beta}^{(k)*}}} - \frac{-\log(\frac{1-w^{(k)*}}{w^{(k)*}}) - \delta^{(k)*} + (\beta^{(k)*})^T \mu_1^{(k)*}}{\sqrt{(\beta^{(k)*})^T \Sigma^{(k)*} \beta^{(k)*}}} \right]^2 \\ + C \left[\frac{\log(\frac{1-\hat{w}^{(k)*}}{\hat{w}^{(k)*}}) + \hat{\delta}^{(k)*} - (\hat{\beta}^{(k)*})^T \mu_2^{(k)*}}{\sqrt{(\hat{\beta}^{(k)*})^T \Sigma^{(k)*} \hat{\beta}^{(k)*}}} - \frac{\log(\frac{1-w^{(k)*}}{w^{(k)*}}) + \delta^{(k)*} - (\beta^{(k)*})^T \mu_2^{(k)*}}{\sqrt{(\beta^{(k)*})^T \Sigma^{(k)*} \beta^{(k)*}}} \right]^2.$$

Denote $\mathcal{A} = (1 - w^{(k)*})\Phi' \left(\frac{-\log(\frac{1-w^{(k)*}}{w^{(k)*}}) - \delta^{(k)*} + (\beta^{(k)*})^T \mu_1^{(k)*}}{\sqrt{(\beta^{(k)*})^T \Sigma^{(k)*} \beta^{(k)*}}} \right) \cdot \left[\frac{-\log(\frac{1-\hat{w}^{(k)*}}{\hat{w}^{(k)*}}) - \hat{\delta}^{(k)*} + (\hat{\beta}^{(k)*})^T \mu_1^{(k)*}}{\sqrt{(\hat{\beta}^{(k)*})^T \Sigma^{(k)*} \hat{\beta}^{(k)*}}} - \frac{-\log(\frac{1-w^{(k)*}}{w^{(k)*}}) - \delta^{(k)*} + (\beta^{(k)*})^T \mu_1^{(k)*}}{\sqrt{(\beta^{(k)*})^T \Sigma^{(k)*} \beta^{(k)*}}} \right] \\ + w^{(k)*}\Phi' \left(\frac{\log(\frac{1-w^{(k)*}}{w^{(k)*}}) + \delta^{(k)*} - (\beta^{(k)*})^T \mu_2^{(k)*}}{\sqrt{(\beta^{(k)*})^T \Sigma^{(k)*} \beta^{(k)*}}} \right) \cdot \left[\frac{\log(\frac{1-\hat{w}^{(k)*}}{\hat{w}^{(k)*}}) + \hat{\delta}^{(k)*} - (\hat{\beta}^{(k)*})^T \mu_2^{(k)*}}{\sqrt{(\hat{\beta}^{(k)*})^T \Sigma^{(k)*} \hat{\beta}^{(k)*}}} - \frac{\log(\frac{1-w^{(k)*}}{w^{(k)*}}) + \delta^{(k)*} - (\beta^{(k)*})^T \mu_2^{(k)*}}{\sqrt{(\beta^{(k)*})^T \Sigma^{(k)*} \beta^{(k)*}}} \right] \\ \text{and } \mathcal{B} = C \left[\frac{-\log(\frac{1-\hat{w}^{(k)*}}{\hat{w}^{(k)*}}) - \hat{\delta}^{(k)*} + (\hat{\beta}^{(k)*})^T \mu_1^{(k)*}}{\sqrt{(\hat{\beta}^{(k)*})^T \Sigma^{(k)*} \hat{\beta}^{(k)*}}} - \frac{-\log(\frac{1-w^{(k)*}}{w^{(k)*}}) - \delta^{(k)*} + (\beta^{(k)*})^T \mu_1^{(k)*}}{\sqrt{(\beta^{(k)*})^T \Sigma^{(k)*} \beta^{(k)*}}} \right]^2 \\ + C \left[\frac{\log(\frac{1-\hat{w}^{(k)*}}{\hat{w}^{(k)*}}) + \hat{\delta}^{(k)*} - (\hat{\beta}^{(k)*})^T \mu_2^{(k)*}}{\sqrt{(\hat{\beta}^{(k)*})^T \Sigma^{(k)*} \hat{\beta}^{(k)*}}} - \frac{\log(\frac{1-w^{(k)*}}{w^{(k)*}}) + \delta^{(k)*} - (\beta^{(k)*})^T \mu_2^{(k)*}}{\sqrt{(\beta^{(k)*})^T \Sigma^{(k)*} \beta^{(k)*}}} \right]^2. \text{ By plugging in the density formula of standard Gaussian distribution, it is easy to see that}$

$$\mathcal{A} \lesssim \sqrt{(1 - w^{(k)*})w^{(k)*}} \cdot \exp \left\{ -\frac{[\log(\frac{1-w^{(k)*}}{w^{(k)*}}) + \frac{1}{2}(\beta^{(k)*})^T \Sigma^{(k)*} \beta^{(k)*}]^2}{2(\beta^{(k)*})^T \Sigma^{(k)*} \beta^{(k)*}} + \frac{1}{2} \log \left(\frac{1 - w^{(k)*}}{w^{(k)*}} \right) \right\}$$

$$\begin{aligned}
& \cdot \left[\frac{-\log(\frac{1-\widehat{w}^{(k)}}{\widehat{w}^{(k)}}) - \widehat{\delta}^{(k)} + (\widehat{\boldsymbol{\beta}}^{(k)})^T \boldsymbol{\mu}_1^{(k)*}}{\sqrt{(\widehat{\boldsymbol{\beta}}^{(k)})^T \boldsymbol{\Sigma}^{(k)*} \widehat{\boldsymbol{\beta}}^{(k)}}} - \frac{-\log(\frac{1-w^{(k)*}}{w^{(k)*}}) - \delta^{(k)*} + (\boldsymbol{\beta}^{(k)*})^T \boldsymbol{\mu}_1^{(k)*}}{\sqrt{(\boldsymbol{\beta}^{(k)*})^T \boldsymbol{\Sigma}^{(k)*} \boldsymbol{\beta}^{(k)*}}} \right] \\
& + \sqrt{(1-w^{(k)*})w^{(k)*}} \cdot \exp \left\{ -\frac{[\log(\frac{1-w^{(k)*}}{w^{(k)*}}) - \frac{1}{2}(\boldsymbol{\beta}^{(k)*})^T \boldsymbol{\Sigma}^{(k)*} \boldsymbol{\beta}^{(k)*}]^2}{2(\boldsymbol{\beta}^{(k)*})^T \boldsymbol{\Sigma}^{(k)*} \boldsymbol{\beta}^{(k)*}} + \frac{1}{2} \log \left(\frac{w^{(k)*}}{1-w^{(k)*}} \right) \right\} \\
& \cdot \left[\frac{\log(\frac{1-\widehat{w}^{(k)}}{\widehat{w}^{(k)}}) + \widehat{\delta}^{(k)} - (\widehat{\boldsymbol{\beta}}^{(k)})^T \boldsymbol{\mu}_2^{(k)*}}{\sqrt{(\widehat{\boldsymbol{\beta}}^{(k)})^T \boldsymbol{\Sigma}^{(k)*} \widehat{\boldsymbol{\beta}}^{(k)}}} - \frac{\log(\frac{1-w^{(k)*}}{w^{(k)*}}) + \delta^{(k)*} - (\boldsymbol{\beta}^{(k)*})^T \boldsymbol{\mu}_2^{(k)*}}{\sqrt{(\boldsymbol{\beta}^{(k)*})^T \boldsymbol{\Sigma}^{(k)*} \boldsymbol{\beta}^{(k)*}}} \right] \\
& = \sqrt{(1-w^{(k)*})w^{(k)*}} \cdot \exp \left\{ -\frac{1}{8}[(\boldsymbol{\beta}^{(k)*})^T \boldsymbol{\Sigma}^{(k)*} \boldsymbol{\beta}^{(k)*}]^2 - \frac{1}{2} \cdot \frac{\log^2(\frac{1-w^{(k)*}}{w^{(k)*}})}{(\boldsymbol{\beta}^{(k)*})^T \boldsymbol{\Sigma}^{(k)*} \boldsymbol{\beta}^{(k)*}} \right\} \\
& \cdot \left| \frac{(\widehat{\boldsymbol{\beta}}^{(k)})^T (\boldsymbol{\mu}_1^{(k)*} - \boldsymbol{\mu}_2^{(k)*})}{\sqrt{(\widehat{\boldsymbol{\beta}}^{(k)})^T \boldsymbol{\Sigma}^{(k)*} \widehat{\boldsymbol{\beta}}^{(k)}}} - \frac{(\boldsymbol{\beta}^{(k)*})^T (\boldsymbol{\mu}_1^{(k)*} - \boldsymbol{\mu}_2^{(k)*})}{\sqrt{(\boldsymbol{\beta}^{(k)*})^T \boldsymbol{\Sigma}^{(k)*} \boldsymbol{\beta}^{(k)*}}} \right| \\
& \lesssim \left| \frac{(\widehat{\boldsymbol{\xi}}^{(k)})^T \boldsymbol{\xi}^{(k)*}}{\|\widehat{\boldsymbol{\xi}}^{(k)}\|_2} - \|\boldsymbol{\xi}^{(k)*}\|_2 \right| \\
& \lesssim \|\widehat{\boldsymbol{\xi}}^{(k)} - \boldsymbol{\xi}^{(k)*}\|_2^2, \tag{E.49}
\end{aligned}$$

with probability at least $1 - C' \exp\{-C''p\}$, where $\widehat{\boldsymbol{\xi}}^{(k)} = (\boldsymbol{\Sigma}^{(k)*})^{1/2} \widehat{\boldsymbol{\beta}}^{(k)}$ and $\boldsymbol{\xi}^{(k)*} = (\boldsymbol{\Sigma}^{(k)*})^{1/2} \boldsymbol{\beta}^{(k)*}$, so $\|\widehat{\boldsymbol{\xi}}^{(k)} - \boldsymbol{\xi}^{(k)*}\|_2^2 \lesssim \|\widehat{\boldsymbol{\beta}}^{(k)} - \boldsymbol{\beta}^{(k)*}\|_2^2$. Only the last inequality in (E.49) holds with high probability and the others are deterministic. It comes from Lemma 8.1 in Cai et al. (2019) and the fact that $\|\widehat{\boldsymbol{\beta}}^{(k)} - \boldsymbol{\xi}^{(k)*}\|_2 \leq c \leq \|\boldsymbol{\xi}^{(k)*}\|_2$ for some $c > 0$ with probability at least $1 - C' \exp\{-C''p\}$. On the other hand, it is easy to see that $\mathcal{B} \lesssim d^2(\widehat{\boldsymbol{\theta}}^{(k)}, \boldsymbol{\theta}^{(k)*})$. Combining these two facts leads to (E.48).

E.5 Proof of Theorem 4

E.5.1 Lemmas

Recall that for GMM associated with parameter set $\bar{\boldsymbol{\theta}} = (w, \boldsymbol{\mu}_1, \boldsymbol{\mu}_2, \boldsymbol{\Sigma})$, we define the mis-clustering error rate of any classifier \mathcal{C} as $R_{\bar{\boldsymbol{\theta}}}(\mathcal{C}) = \min_{\pi: \{1,2\} \rightarrow \{1,2\}} \mathbb{P}_{\bar{\boldsymbol{\theta}}}(\mathcal{C}(Z^{\text{new}}) \neq \pi(Y^{\text{new}}))$, where $\mathbb{P}_{\bar{\boldsymbol{\theta}}}$ represents the distribution of $(Z^{\text{new}}, Y^{\text{new}})$, i.e. $(1-w)\mathcal{N}(\boldsymbol{\mu}_1, \boldsymbol{\Sigma}) + w\mathcal{N}(\boldsymbol{\mu}_2, \boldsymbol{\Sigma})$. Denote $\mathcal{C}_{\bar{\boldsymbol{\theta}}}$ as the Bayes classifier corresponding to $\bar{\boldsymbol{\theta}}$. Define a surrogate loss $L_{\bar{\boldsymbol{\theta}}}(\mathcal{C}) = \min_{\pi: \{1,2\} \rightarrow \{1,2\}} \mathbb{P}_{\bar{\boldsymbol{\theta}}}(\mathcal{C}(Z^{\text{new}}) \neq \pi(\mathcal{C}_{\bar{\boldsymbol{\theta}}}(Y^{\text{new}})))$.

Lemma 24. *Assume there exists an subset S such that $\min_{k \in S} n_k \geq C(p \vee \log K)$ and $\min_{k \in S} \Delta^{(k)} \geq \sigma^2 > 0$ with some constants $C > 0$. We have*

$$\inf_{\{\widehat{\mathcal{C}}^{(k)}\}_{k=1}^K} \sup_{S: |S| \geq s} \sup_{\{\boldsymbol{\theta}^{(k)*}\}_{k \in S} \in \bar{\Theta}_S} \mathbb{P} \left(\bigcup_{k \in S} \left\{ R_{\bar{\boldsymbol{\theta}}^{(k)*}}(\widehat{\mathcal{C}}^{(k)}) - R_{\bar{\boldsymbol{\theta}}^{(k)*}}(\mathcal{C}_{\bar{\boldsymbol{\theta}}^{(k)*}}) \geq C_1 \frac{p}{n_S} + C_2 \frac{\log K}{n_k} \right\} \right)$$

$$+ C_3 h^2 \wedge \frac{p + \log K}{n_k} + C_4 \frac{\epsilon^2}{\max_{k \in S} n_k} \Bigg\} \geq \frac{1}{10}.$$

Lemma 25. Suppose $\bar{\theta} = (w, \boldsymbol{\mu}_1, \boldsymbol{\mu}_2, \boldsymbol{\beta}, \boldsymbol{\Sigma})$ satisfies $\Delta^2 := (\boldsymbol{\mu}_1 - \boldsymbol{\mu}_2)^T \boldsymbol{\Sigma}^{-1} (\boldsymbol{\mu}_1 - \boldsymbol{\mu}_2) \geq \sigma^2 > 0$ with some constant $\sigma^2 > 0$ and $w, w' \in (c_w, 1 - c_w)$. Then $\exists c > 0$ such that

$$cL_{\bar{\theta}}^2(\mathcal{C}) \leq R_{\bar{\theta}}(\mathcal{C}) - R_{\bar{\theta}}(\mathcal{C}_{\bar{\theta}}),$$

for any classifier \mathcal{C} , where $R_{\bar{\theta}}(\mathcal{C}) := \min_{\pi: \{1,2\} \rightarrow \{1,2\}} \mathbb{P}_{\bar{\theta}}(\mathcal{C}(\mathbf{z}) \neq \pi(y))$, $L_{\bar{\theta}}(\mathcal{C}) := \min_{\pi: \{0,1\} \rightarrow \{0,1\}} \mathbb{P}_{\bar{\theta}}(\mathcal{C}(\mathbf{z}) \neq \pi(\mathcal{C}_{\bar{\theta}}(\mathbf{z})))$, and $\mathcal{C}_{\bar{\theta}}$ is the corresponding Bayes classifier.

Lemma 26. Consider $\bar{\theta} = (w, \boldsymbol{\mu}_1, \boldsymbol{\mu}_2, \boldsymbol{\beta}, \boldsymbol{\Sigma})$ and $\bar{\theta}' = (w', \boldsymbol{\mu}_1, \boldsymbol{\mu}_2, \boldsymbol{\beta}, \boldsymbol{\Sigma})$ satisfies $\Delta^2 := (\boldsymbol{\mu}_1 - \boldsymbol{\mu}_2)^T \boldsymbol{\Sigma}^{-1} (\boldsymbol{\mu}_1 - \boldsymbol{\mu}_2) \geq \sigma^2 > 0$ with some constant $\sigma^2 > 0$. We have

$$c|w - w'| \leq L_{\bar{\theta}}(\mathcal{C}_{\bar{\theta}'}), \leq c'|w - w'|,$$

for some constants $c, c' > 0$.

Lemma 27. Consider $\bar{\theta} = (w, \boldsymbol{\mu}_1, \boldsymbol{\mu}_2, \boldsymbol{\Sigma})$ and $\bar{\theta}' = (w, \boldsymbol{\mu}'_1, \boldsymbol{\mu}'_2, \boldsymbol{\Sigma})$ satisfies $w = 1/2$, $\boldsymbol{\mu}_1 = -\boldsymbol{\mu}_0/2 + \mathbf{u}$, $\boldsymbol{\mu}_2 = \boldsymbol{\mu}_0/2 + \mathbf{u}$, $\boldsymbol{\mu}'_1 = -\boldsymbol{\mu}_0/2 + \mathbf{u}'$, $\boldsymbol{\mu}'_2 = \boldsymbol{\mu}_0/2 + \mathbf{u}'$, $\boldsymbol{\Sigma} = \mathbf{I}_p$, $\boldsymbol{\mu}_0 = (1, \mathbf{0}_{p-1}^T)^T$, $\mathbf{u} = (\tilde{u}, \mathbf{0}_{p-1}^T)^T$, and $\mathbf{u}' = (\tilde{u}', \mathbf{0}_{p-1}^T)^T$. We have

$$c|\tilde{u} - \tilde{u}'| \leq L_{\bar{\theta}}(\mathcal{C}_{\bar{\theta}'}), \leq c'|\tilde{u} - \tilde{u}'|,$$

for some constants $c, c' > 0$.

Lemma 28. Denote $\tilde{\epsilon} = \frac{K-s}{s}$. We have

$$\inf_{\{\hat{\mathcal{C}}^{(k)}\}_{k=1}^K} \sup_{S: |S| \geq s} \sup_{\substack{\{\bar{\theta}^{(k)*}\}_{k \in S} \in \bar{\Theta}_S \\ \mathbb{Q}_S}} \mathbb{P} \left(\max_{k \in S} \left[R_{\bar{\theta}^{(k)*}}(\hat{\mathcal{C}}^{(k)}) - R_{\bar{\theta}^{(k)*}}(\mathcal{C}_{\bar{\theta}^{(k)*}}) \right] \geq C_1 \frac{\tilde{\epsilon}^2}{\max_{k=1:K} n_k} \right) \geq \frac{1}{10}.$$

E.5.2 Main proof of Theorem 4

Combine conclusions of Lemmas 24 and 28 to get the lower bound.

E.5.3 Proof of lemmas

Proof of Lemma 24. Recall the definitions and proof idea of Lemma 19. We have $\bar{\Theta}_S \supseteq \bar{\Theta}_{|S|,w} \cup \bar{\Theta}_{|S|,\delta} \cup \bar{\Theta}_{|S|,\beta}$, where

$$\begin{aligned} \bar{\Theta}_{S,w} &= \left\{ \{\bar{\theta}^{(k)}\}_{k \in S} : \boldsymbol{\mu}_1^{(k)} = \mathbf{1}_p / \sqrt{p}, \boldsymbol{\mu}_2^{(k)} = -\boldsymbol{\mu}_1^{(k)} = \tilde{\boldsymbol{\mu}}, \boldsymbol{\Sigma}^{(k)} = \mathbf{I}_p, w^{(k)} \in (c_w, 1 - c_w) \right\}, \\ \bar{\Theta}_{S,\beta} &= \left\{ \{\bar{\theta}^{(k)}\}_{k \in S} : \boldsymbol{\Sigma}^{(k)} = \mathbf{I}_p, w^{(k)} = \frac{1}{2}, \|\boldsymbol{\mu}_1^{(k)}\|_2 \vee \|\boldsymbol{\mu}_2^{(k)}\|_2 \leq M, \boldsymbol{\mu}_2^{(k)} = -\boldsymbol{\mu}_1^{(k)} \right\}, \end{aligned}$$

$$\min_{\beta} \max_{k \in S} \|\beta^{(k)} - \beta\|_2 \leq h\},$$

$$\overline{\Theta}_{S,\delta} = \left\{ \{\bar{\theta}^{(k)}\}_{k \in S} : \Sigma^{(k)} = \mathbf{I}_p, w^{(k)} = \frac{1}{2}, \|\mu_1^{(k)}\|_2 \vee \|\mu_2^{(k)}\|_2 \leq M, \mu_1^{(k)} = -\frac{1}{2}\mu_0, \mu_2^{(k)} = \frac{1}{2}\mu_0 + \mathbf{u}, \right.$$

$$\left. \|\mathbf{u}\|_2 \leq 1 \right\}.$$

Recall the mis-clustering error for GMM associated with parameter set $\bar{\theta}$ of any classifier \mathcal{C} is $R_{\bar{\theta}}(\mathcal{C}) = \min_{\pi: \{1,2\} \rightarrow \{1,2\}} \mathbb{P}_{\bar{\theta}}(\mathcal{C}(Z) \neq \pi(Y))$. To help the analysis, following [Azizyan et al. \(2013\)](#) and [Cai et al. \(2019\)](#), we define a surrogate loss $L_{\bar{\theta}}(\mathcal{C}) = \min_{\pi: \{1,2\} \rightarrow \{1,2\}} \mathbb{P}_{\bar{\theta}}(\mathcal{C}(Z) \neq \pi(\mathcal{C}_{\bar{\theta}}(Z)))$, where $\mathcal{C}_{\bar{\theta}}$ is the Bayes classifier. Suppose $\sigma = \sqrt{0.005}$.

(i) We want to show

$$\inf_{\{\hat{\mathcal{C}}^{(k)}\}_{k=1}^K} \sup_{S: |S| \geq s} \sup_{\{\bar{\theta}^{(k)*}\}_{k \in S} \in \overline{\Theta}_S} \mathbb{P} \left(\bigcup_{k \in S} \left\{ R_{\bar{\theta}^{(k)*}}(\hat{\mathcal{C}}^{(k)}) - R_{\bar{\theta}^{(k)*}}(\mathcal{C}_{\bar{\theta}^{(k)*}}) \geq C \sqrt{\frac{p}{n_S}} \right\} \right) \geq \frac{1}{4}. \quad (\text{E.50})$$

Consider $S = 1 : K$ and space $\overline{\Theta}_0 = \{\{\bar{\theta}^{(k)}\}_{k=1}^K : \Sigma^{(k)} = \mathbf{I}_p, w^{(k)} = 1/2, \mu_1^{(k)} = \mu_1, \mu_2^{(k)} = \mu_2, \|\mu_1\|_2 \vee \|\mu_2\|_2 \leq M\}$. And

$$\text{LHS of (E.50)} \geq \inf_{\hat{\mathcal{C}}^{(1)}} \sup_{\{\bar{\theta}^{(k)*}\}_{k=1}^K \in \overline{\Theta}_0} \mathbb{P} \left(R_{\bar{\theta}^{(1)*}}(\hat{\mathcal{C}}^{(1)}) - R_{\bar{\theta}^{(1)*}}(\mathcal{C}_{\bar{\theta}^{(1)*}}) \geq C \sqrt{\frac{p}{n_S}} \right).$$

Let $r = c\sqrt{p/n_S} \leq 0.001$ with some small constant $c > 0$. For any $\mu \in \mathbb{R}^p$, denote distribution $\frac{1}{2}\mathcal{N}(\mu, \mathbf{I}_p) + \frac{1}{2}\mathcal{N}(-\mu, \mathbf{I}_p)$ as \mathbb{P}_{μ} . Consider a $r/4$ -packing of $r\mathcal{S}^{p-1}$: $\{\tilde{\mathbf{v}}_j\}_{j=1}^N$. By Lemma 2, $N \geq 4^{p-1}$. Denote $\tilde{\mu}_j = (\sigma, \tilde{\mathbf{v}}_j^T)^T \in \mathbb{R}^p$, where $\sigma = \sqrt{0.005}$. Then by definition of KL divergence and Lemma 8.4 in [Cai et al. \(2019\)](#),

$$\begin{aligned} \text{KL} \left(\prod_{k \in S} \mathbb{P}_{\tilde{\mu}_j}^{\otimes n_k} \cdot \mathbb{Q}_S \left\| \prod_{k \in S} \mathbb{P}_{\tilde{\mu}_{j'}}^{\otimes n_k} \cdot \mathbb{Q}_S \right. \right) &= \sum_{k \in S} n_k \text{KL}(\mathbb{P}_{\tilde{\mu}_j} \|\mathbb{P}_{\tilde{\mu}_{j'}}) \\ &\leq n_S \cdot 8(1 + \sigma^2) \|\tilde{\mu}_j - \tilde{\mu}_{j'}\|_2^2 \\ &\leq 32(1 + \sigma^2) n_S r^2 \\ &\leq 32(1 + \sigma^2) n_S \cdot c^2 \frac{2(p-1)}{n_S} \\ &\leq \frac{32(1 + \sigma^2) c^2}{\log 2} \log N. \end{aligned}$$

For simplicity, we write $L_{\bar{\theta}}$ with $\bar{\theta} \in \overline{\Theta}_0$ and $\mu_1 = -\mu_2 = \mu$ as L_{μ} . By Lemma 8.5 in [Cai et al. \(2019\)](#),

$$L_{\tilde{\mu}_i}(\mathcal{C}_{\tilde{\mu}_j}) \geq \frac{1}{\sqrt{2}} g \left(\frac{\sqrt{\sigma^2 + r^2}}{2} \right) \frac{\|\tilde{\mu}_i - \tilde{\mu}_j\|_2}{\|\tilde{\mu}_i\|_2} \geq \frac{1}{\sqrt{2}} \cdot 0.15 \cdot \frac{r/4}{\sqrt{\sigma^2 + r^2}} \geq 2r,$$

where $g(x) = \phi(x)[\phi(x) - x\Phi(x)]$. The last inequality holds because $\sqrt{\sigma^2 + r^2} \geq \sqrt{2}\sigma$ and $g(\sqrt{\sigma^2 + r^2}/2) \geq 0.15$ when $r^2 \leq \sigma^2 = 0.001$. Then by Lemma 3.5 in [Cai et al. \(2019\)](#) (Proposition 2 in [Azizyan et al. \(2013\)](#)), for any classifier \mathcal{C} , and $i \neq j$,

$$L_{\tilde{\mu}_i}(\mathcal{C}) + L_{\tilde{\mu}_j}(\mathcal{C}) \geq L_{\tilde{\mu}_i}(\mathcal{C}_{\tilde{\mu}_j}) - \sqrt{\text{KL}(\mathbb{P}_{\tilde{\mu}_i} \|\mathbb{P}_{\tilde{\mu}_j})/2} \geq 2r - r = c\sqrt{\frac{p}{n_S}}. \quad (\text{E.51})$$

For any $\hat{\mathcal{C}}^{(1)}$, consider a test $\psi^* = \arg \min_{j=1:N} L_{\tilde{\mu}_j}(\hat{\mathcal{C}}^{(1)})$. Therefore if there exists j_0 such that $L_{\tilde{\mu}_{j_0}}(\hat{\mathcal{C}}^{(1)}) < \frac{c}{2}\sqrt{\frac{p}{n_S}}$, then by (E.51), we must have $\psi^* = j_0$. Let $C_1 \leq c/2$, then by Fano's lemma (Corollary 6 in [Tsybakov \(2009\)](#))

$$\begin{aligned} \inf_{\hat{\mathcal{C}}^{(1)}} \sup_{\{\bar{\theta}^{(k)*}\}_{k=1}^K \in \bar{\Theta}_0} \mathbb{P} \left(L_{\bar{\theta}^{(1)*}}(\hat{\mathcal{C}}^{(1)}) \geq C_1 \sqrt{\frac{p}{n_S}} \right) &\geq \inf_{\hat{\mathcal{C}}^{(1)}} \sup_{j=1:N} \mathbb{P} \left(L_{\tilde{\mu}^{(j)}}(\hat{\mathcal{C}}^{(1)}) \geq C_1 \sqrt{\frac{p}{n_S}} \right) \\ &\geq \inf_{\hat{\mathcal{C}}^{(1)}} \sup_{j=1:N} \mathbb{P} \left(\psi^* \neq j \right) \\ &\geq \inf_{\psi} \sup_{j=1:N} \mathbb{P} \left(\psi \neq j \right) \\ &\geq 1 - \frac{\log 2}{\log N} - \frac{32(1 + \sigma^2)c^2}{\log 2} \\ &\geq \frac{1}{4}, \end{aligned}$$

when $p \geq 2$ and $c = \sqrt{\frac{\log 2}{128(1 + \sigma^2)}}$. Then apply Lemma 25 to get the (E.50).

(ii) We want to show

$$\inf_{\{\hat{\mathcal{C}}^{(k)}\}_{k=1}^K} \sup_{S: |S| \geq s} \sup_{\substack{\{\bar{\theta}^{(k)*}\}_{k \in S} \in \bar{\Theta}_{S, \beta} \\ \mathbb{Q}_S}} \mathbb{P} \left(\bigcup_{k \in S} \left\{ R_{\bar{\theta}^{(k)*}}(\hat{\mathcal{C}}^{(k)}) - R_{\bar{\theta}^{(k)*}}(\mathcal{C}_{\bar{\theta}^{(k)*}}) \geq Ch \wedge \sqrt{\frac{p}{n_k}} \right\} \right) \geq \frac{1}{4}. \quad (\text{E.52})$$

Fixing an S and a \mathbb{Q}_S . Suppose $1 \in S$. We have

$$\text{LHS of (E.52)} \geq \inf_{\hat{\mathcal{C}}^{(1)}} \sup_{\substack{\{\bar{\theta}^{(k)*}\}_{k \in S} \in \bar{\Theta}_{S, \beta} \\ \mathbb{Q}_S}} \mathbb{P} \left(R_{\bar{\theta}^{(1)*}}(\hat{\mathcal{C}}^{(1)}) - R_{\bar{\theta}^{(1)*}}(\mathcal{C}_{\bar{\theta}^{(1)*}}) \geq Ch \wedge \sqrt{\frac{p}{n_k}} \right). \quad (\text{E.53})$$

Let $r = h \wedge (c\sqrt{p/n_1}) \wedge M$ with a small constant $c > 0$. For any $\mu \in \mathbb{R}^p$, denote distribution $\frac{1}{2}\mathcal{N}(\mu, \mathbf{I}_p) + \frac{1}{2}\mathcal{N}(-\mu, \mathbf{I}_p)$ as \mathbb{P}_μ . Consider a $r/4$ -packing of $r\mathcal{S}^{p-1}$. By Lemma 2, $N \geq 4^{p-1}$. Denote $\tilde{\mu}_j = (\sigma, \tilde{\mathbf{v}}_j^T)^T \in \mathbb{R}^p$. WLOG, assume $M \geq 2$. Let $\mu_1^{(k)*} = \tilde{\mu} = (\sigma, \mathbf{0}_{p-1})^T$ for all $k \in S \setminus \{1\}$. Then by following the same arguments in (i) and part (ii) of the proof of Lemma 19, we can show that the RHS of (E.53) is larger than or equal to $1/4$ when $p \geq 3$.

(iii) We want to show

$$\inf_{\{\widehat{\mathcal{C}}^{(k)}\}_{k=1}^K} \sup_{S:|S|\geq s} \sup_{\substack{\{\bar{\theta}^{(k)*}\}_{k\in S}\in\bar{\Theta}_{S,\beta} \\ \mathbb{Q}_S}} \mathbb{P}\left(\bigcup_{k\in S} \left\{R_{\bar{\theta}^{(k)*}}(\widehat{\mathcal{C}}^{(k)}) - R_{\bar{\theta}^{(k)*}}(\mathcal{C}_{\bar{\theta}^{(k)*}}) \geq Ch \wedge \sqrt{\frac{\log K}{n_k}}\right\}\right) \geq \frac{1}{4}.$$

This can be proved by following similar ideas used in step (iii) of the proof of Lemma 19, so we omit the proof here.

(iv) We want to show

$$\inf_{\{\widehat{\mathcal{C}}^{(k)}\}_{k=1}^K} \sup_{S:|S|\geq s} \sup_{\substack{\{\bar{\theta}^{(k)*}\}_{k\in S}\in\bar{\Theta}_{S,w} \\ \mathbb{Q}_S}} \mathbb{P}\left(\bigcup_{k\in S} \left\{R_{\bar{\theta}^{(k)*}}(\widehat{\mathcal{C}}^{(k)}) - R_{\bar{\theta}^{(k)*}}(\mathcal{C}_{\bar{\theta}^{(k)*}}) \geq C\sqrt{\frac{\log K}{n_k}}\right\}\right) \geq \frac{1}{4}.$$

This can be similarly proved by following the arguments in part (i) with Lemmas 25 and 26.

(v) We want to show

$$\inf_{\{\widehat{\mathcal{C}}^{(k)}\}_{k=1}^K} \sup_{S:|S|\geq s} \sup_{\substack{\{\bar{\theta}^{(k)*}\}_{k\in S}\in\bar{\Theta}_{S,\delta} \\ \mathbb{Q}_S}} \mathbb{P}\left(\bigcup_{k\in S} \left\{R_{\bar{\theta}^{(k)*}}(\widehat{\mathcal{C}}^{(k)}) - R_{\bar{\theta}^{(k)*}}(\mathcal{C}_{\bar{\theta}^{(k)*}}) \geq C\sqrt{\frac{\log K}{n_k}}\right\}\right) \geq \frac{1}{4}.$$

This can be similarly proved by following the arguments in part (i) with Lemmas 25 and 27.

Finally, we get the desired conclusion by combining (i)-(v). \square

Proof of Lemma 25. We follow a similar proof idea used in the proof of Lemma 3.4 in Cai et al. (2019). Let ϕ_1 and ϕ_2 be the density of $\mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ and $\mathcal{N}(-\boldsymbol{\mu}, \boldsymbol{\Sigma})$, respectively. Denote $\eta_{\bar{\theta}}(\mathbf{z}) = \frac{(1-w)\phi_1(\mathbf{z})}{(1-w)\phi_1(\mathbf{z})+w\phi_2(\mathbf{z})}$ and $S_{\mathcal{C}} = \{\mathbf{z} \in \mathbb{R}^p : \mathcal{C}(\mathbf{z}) = 1\}$ for any classifier \mathcal{C} . Note that $S_{\mathcal{C}_{\bar{\theta}}} = \{\mathbf{z} \in \mathbb{R}^p : (1-w)\phi_1(\mathbf{z}) \geq w\phi_2(\mathbf{z})\}$. The permutation actually doesn't matter in the proof. WLOG, we drop the permutations in the definition of misclassification error and surrogate loss by assuming π to be the identity function. If π is not identity in the definition of $R_{\bar{\theta}}(\mathcal{C})$, for example, we can define $S_{\mathcal{C}} = \{\mathbf{z} \in \mathbb{R}^p : \mathcal{C}(\mathbf{z}) = 2\}$ instead and all the following steps still follow.

By definition,

$$\begin{aligned} \mathbb{P}_{\bar{\theta}}(\mathcal{C}(\mathbf{z}) \neq y) &= (1-w) \int_{S_{\mathcal{C}}^c} \phi_1 d\mathbf{z} + w \int_{S_{\mathcal{C}}} \phi_2 d\mathbf{z}, \\ \mathbb{P}_{\bar{\theta}}(\mathcal{C}_{\bar{\theta}}(\mathbf{z}) \neq y) &= (1-w) \int_{S_{\mathcal{C}_{\bar{\theta}}}^c} \phi_1 d\mathbf{z} + w \int_{S_{\mathcal{C}_{\bar{\theta}}}} \phi_2 d\mathbf{z}, \end{aligned}$$

which leads to

$$\mathbb{P}_{\bar{\theta}}(\mathcal{C}(\mathbf{z}) \neq y) - \mathbb{P}_{\bar{\theta}}(\mathcal{C}_{\bar{\theta}}(\mathbf{z}) \neq y) = \int_{S_{\mathcal{C}_{\bar{\theta}}} \setminus S_{\mathcal{C}}} [(1-w)\phi_1 - w\phi_2] d\mathbf{z} + \int_{S_{\mathcal{C}_{\bar{\theta}}}^c \setminus S_{\mathcal{C}}^c} [w\phi_2 - (1-w)\phi_1] d\mathbf{z}$$

$$\begin{aligned}
&= \int_{S_{C_{\bar{\theta}}} \triangle S_C} |(1-w)\phi_1 - w\phi_2| dz \\
&= \mathbb{E}_{\mathbf{z} \sim (1-w)\phi_1 + w\phi_2} [|2\eta_{\bar{\theta}}(\mathbf{z}) - 1| \mathbf{1}(S_{C_{\bar{\theta}}} \triangle S_C)] \\
&\geq 2t \cdot \mathbb{P}_{\bar{\theta}}(S_{C_{\bar{\theta}}} \triangle S_C, |2\eta_{\bar{\theta}}(\mathbf{z}) - 1| > 2t) \\
&= 2t [\mathbb{P}_{\bar{\theta}}(S_{C_{\bar{\theta}}} \triangle S_C) - \mathbb{P}_{\bar{\theta}}(|2\eta_{\bar{\theta}}(\mathbf{z}) - 1| \leq 2t)] \\
&\geq 2t [\mathbb{P}_{\bar{\theta}}(S_{C_{\bar{\theta}}} \triangle S_C) - ct] \\
&\geq \frac{1}{2c} \mathbb{P}_{\bar{\theta}}^2(S_{C_{\bar{\theta}}} \triangle S_C),
\end{aligned}$$

where we let $t = \frac{1}{2c} \mathbb{P}_{\boldsymbol{\theta}}(S_{\mathcal{C}_{\boldsymbol{\theta}}} \triangle S_{\mathcal{C}})$ with $c = 1 + \frac{8}{\sqrt{2\pi}\sigma}$. This completes the proof. The last second inequality depends on the fact that

$$\mathbb{P}_{\bar{\theta}}(|\eta_{\bar{\theta}}(\mathbf{z}) - 1/2| \leq t) \leq ct.$$

holds for all $t \leq 1/(2c)$. This is because

$$\begin{aligned}
& \mathbb{P}_{\bar{\boldsymbol{\theta}}}(|\eta_{\bar{\boldsymbol{\theta}}}(\mathbf{z}) - 1/2| \leq t) \\
&= \mathbb{P}_{\bar{\boldsymbol{\theta}}} \left(\log \left(\frac{w}{1-w} \right) + \log \left(\frac{1-2t}{1+2t} \right) \leq \log \left(\frac{\phi_1}{\phi_2}(\mathbf{z}) \right) \leq \log \left(\frac{w}{1-w} \right) + \log \left(\frac{1+2t}{1-2t} \right) \right) \\
&= \mathbb{P}_{\bar{\boldsymbol{\theta}}} \left(\log \left(\frac{w}{1-w} \right) + \log \left(\frac{1-2t}{1+2t} \right) \leq (\boldsymbol{\mu}_1 - \boldsymbol{\mu}_2)^T \boldsymbol{\Sigma}^{-1} \left(\mathbf{z} - \frac{\boldsymbol{\mu}_1 + \boldsymbol{\mu}_2}{2} \right) \right. \\
&\quad \left. \leq \log \left(\frac{w}{1-w} \right) + \log \left(\frac{1+2t}{1-2t} \right) \right) \\
&= \frac{1}{2} \mathbb{P}_{\bar{\boldsymbol{\theta}}} \left(\log \left(\frac{w}{1-w} \right) + \log \left(\frac{1-2t}{1+2t} \right) \leq \mathcal{N}(\Delta^2/2, \Delta^2) \leq \log \left(\frac{w}{1-w} \right) + \log \left(\frac{1+2t}{1-2t} \right) \right) \\
&\quad + \frac{1}{2} \mathbb{P}_{\bar{\boldsymbol{\theta}}} \left(\log \left(\frac{w}{1-w} \right) + \log \left(\frac{1-2t}{1+2t} \right) \leq \mathcal{N}(-\Delta^2/2, \Delta^2) \leq \log \left(\frac{w}{1-w} \right) + \log \left(\frac{1+2t}{1-2t} \right) \right) \\
&\hspace{10cm} \text{(E.54)} \\
&\leq \frac{1}{\sqrt{2\pi}\sigma} \cdot \left[\log \left(\frac{1+2t}{1-2t} \right) - \log \left(\frac{1-2t}{1+2t} \right) \right] \\
&\leq \frac{1}{\sqrt{2\pi}\sigma} \cdot \frac{8t}{1-2t} \\
&\leq ct, \hspace{10cm} \text{(E.55)}
\end{aligned}$$

when $t \leq 1/(2c)$. Note that (E.55) implies that a binary GMM under the separation assumption $\Delta \gtrsim 1$ has Tsybakov’s margin with margin parameter 1. For the notion of Tsybakov’s margin, see [Audibert and Tsybakov \(2007\)](#). We will prove a more general result showing that a multi-cluster GMM under the separation assumption also has Tsybakov’s margin with margin parameter 1. This turns out to be useful in proving the upper and lower bounds of misclassification error. \square

Proof of Lemma 26. WLOG, suppose $w \geq w'$. Similar to (E.54), it's easy to see that

$$\begin{aligned}
L_{\bar{\theta}}(\mathcal{C}_{\bar{\theta}'}) &= (1-w)\mathbb{P}_{\bar{\theta}}(\mathcal{C}_{\bar{\theta}}(\mathbf{z}) \neq \mathcal{C}_{\bar{\theta}'}(\mathbf{z}) | \mathbf{z} = 1) + w\mathbb{P}_{\bar{\theta}}(\mathcal{C}_{\bar{\theta}}(\mathbf{z}) \neq \mathcal{C}_{\bar{\theta}'}(\mathbf{z}) | \mathbf{z} = 2) \\
&= (1-w)\mathbb{P}\left(\log\left(\frac{w'}{1-w'}\right) \leq \mathcal{N}(\Delta^2/2, \Delta^2) \leq \log\left(\frac{w}{1-w}\right)\right) \\
&\quad + w\mathbb{P}\left(\log\left(\frac{w'}{1-w'}\right) \leq \mathcal{N}(-\Delta^2/2, \Delta^2) \leq \log\left(\frac{w}{1-w}\right)\right) \\
&\leq \frac{1}{\sqrt{2\pi}\sigma} \left[\log\left(\frac{w}{1-w}\right) - \log\left(\frac{w'}{1-w'}\right) \right] \\
&= \frac{1}{\sqrt{2\pi}c_w(1-c_w)\sigma} \cdot |w - w'|.
\end{aligned} \tag{E.56}$$

On the other hand,

$$\begin{aligned}
(\text{E.56}) &\geq \frac{1}{\sqrt{2\pi}Mc_{\Sigma}} \cdot \exp\left\{-\frac{1}{2\sigma^2} \left[\log\left(\frac{1-c_w}{c_w}\right) + \frac{1}{2}M^2c_{\Sigma} \right]^2\right\} \left[\log\left(\frac{w}{1-w}\right) - \log\left(\frac{w'}{1-w'}\right) \right] \\
&\geq \frac{1}{\sqrt{2\pi}Mc_{\Sigma}c_w(1-c_w)} \cdot \exp\left\{-\frac{1}{2\sigma^2} \left[\log\left(\frac{1-c_w}{c_w}\right) + \frac{1}{2}M^2c_{\Sigma} \right]^2\right\} |w - w'|,
\end{aligned}$$

which completes the proof. \square

Proof of Lemma 28. By Lemma 25, it suffices to prove

$$\inf_{\{\hat{\mathcal{C}}^{(k)}\}_{k=1}^K} \sup_{S: |S| \geq s} \sup_{\{\bar{\theta}^{(k)*}\}_{k \in S} \in \bar{\Theta}_S} \mathbb{P}\left(\max_{k \in S} L_{\bar{\theta}^{(k)*}}(\hat{\mathcal{C}}^{(k)}) \geq C_1 \frac{\tilde{\epsilon}^2}{\max_{k=1:K} n_k}\right) \geq \frac{1}{10}.$$

For any $\boldsymbol{\mu} \in \mathbb{R}^p$, denote distribution $\frac{1}{2}\mathcal{N}(\boldsymbol{\mu}, \mathbf{I}_p) + \frac{1}{2}\mathcal{N}(-\boldsymbol{\mu}, \mathbf{I}_p)$ as $\mathbb{P}_{\boldsymbol{\mu}}$. For simplicity, we write $L_{\bar{\theta}}$ with $\bar{\theta}$ satisfying $\boldsymbol{\mu}_1 = -\boldsymbol{\mu}_2 = \boldsymbol{\mu}$, $w = 1/2$ and $\Sigma = \mathbf{I}_p$ as $L_{\boldsymbol{\mu}}$. Consider $L_{\boldsymbol{\mu}}(\mathcal{C}_{\boldsymbol{\mu}'})$ as a loss function between $\boldsymbol{\mu}$ and $\boldsymbol{\mu}'$ in Lemmas 21 and 22. Considering $\|\boldsymbol{\mu}\|_2 = \|\boldsymbol{\mu}'\|_2 = 1$, by Lemma 15, note that

$$\max_{k=1:K} \text{KL}(\mathbb{P}_{\boldsymbol{\mu}}^{\otimes n_k} \| \mathbb{P}_{\boldsymbol{\mu}'}^{\otimes n_k}) \leq 8 \max_{k=1:K} n_k \cdot \|\boldsymbol{\mu} - \boldsymbol{\mu}'\|_2^2.$$

By Lemma 8.5 in Cai et al. (2019), this implies for some constants $c, C > 0$

$$\begin{aligned}
&\sup \left\{ L_{\boldsymbol{\mu}}(\mathcal{C}_{\boldsymbol{\mu}'}) : \max_{k=1:K} \text{KL}(\mathbb{P}_{\boldsymbol{\mu}}^{\otimes n_k} \| \mathbb{P}_{\boldsymbol{\mu}'}^{\otimes n_k}) \leq (\tilde{\epsilon}'/(1-\tilde{\epsilon}))^2 \right\} \\
&\geq \sup \left\{ c\|\boldsymbol{\mu} - \boldsymbol{\mu}'\|_2 : \max_{k=1:K} \text{KL}(\mathbb{P}_{\boldsymbol{\mu}}^{\otimes n_k} \| \mathbb{P}_{\boldsymbol{\mu}'}^{\otimes n_k}) \leq (\tilde{\epsilon}'/(1-\tilde{\epsilon}))^2 \right\} \\
&\geq \sup \left\{ c\|\boldsymbol{\mu} - \boldsymbol{\mu}'\|_2 : 8 \max_{k=1:K} n_k \cdot \|\boldsymbol{\mu} - \boldsymbol{\mu}'\|_2^2 \leq (\tilde{\epsilon}'/(1-\tilde{\epsilon}))^2 \right\} \\
&= C \cdot \frac{\tilde{\epsilon}'}{\sqrt{\max_{k=1:K} n_k}}.
\end{aligned}$$

Then apply Lemmas 21 and 22 to get the desired bound. \square

E.6 Proof of Theorem 5

Denote $\xi = \max_{k \in S} \min_{r_k = \pm 1} \|r_k \widehat{\beta}^{(k)[0]} - \beta^{(k)*}\|_2 = \max_{k \in S} (\|\widehat{\beta}^{(k)[0]} - \beta^{(k)*}\|_2 \wedge \|\widehat{\beta}^{(k)[0]} + \beta^{(k)*}\|_2)$. WLOG, assume $S = \{1, \dots, s\}$ and $r_k^* = 1$ for all $k \in S$. Hence $\xi = \max_{k \in S} \|\widehat{\beta}^{(k)[0]} - \beta^{(k)*}\|_2$. For any $k' = 1, \dots, s$, define

$$\begin{aligned} \mathbf{r} &= (\underbrace{r_1, \dots, r_{k'}}_{=-1}, \underbrace{r_{k'+1}, \dots, r_s}_{=1}, \underbrace{r_{s+1}, \dots, r_K}_{\text{outlier tasks}}), \\ \mathbf{r}' &= (1, 1, \dots, 1, 1, 1, \dots, 1, 1, \underbrace{r_{s+1}, \dots, r_K}_{\text{outlier tasks}}), \\ \mathbf{r}'' &= (-1, \dots, -1, -1, \dots, -1, \underbrace{r_{s+1}, \dots, r_K}_{\text{outlier tasks}}). \end{aligned}$$

WLOG, it suffices to prove that

$$\text{score}(\mathbf{r}) - \text{score}(\mathbf{r}') > 0 \quad \text{when } k' \leq \lfloor s/2 \rfloor, \quad (\text{E.57})$$

$$\text{score}(\mathbf{r}) - \text{score}(\mathbf{r}'') > 0 \quad \text{when } k' > \lfloor s/2 \rfloor. \quad (\text{E.58})$$

In fact, if this holds, then we must have

$$\widehat{r}_k = 1 \text{ for all } k \in S \quad \text{or} \quad \widehat{r}_k = -1 \text{ for all } k \in S.$$

Otherwise, according to (E.57), if $\#\{k \in S : \widehat{r}_k = -1\} \leq \lfloor s/2 \rfloor$, by replacing the first s entries of $\widehat{\mathbf{r}}$ with 1, we get a different alignment whose score is smaller than the score of $\widehat{\mathbf{r}}$, which is contradicted with the definition of $\widehat{\mathbf{r}}$. If $\#\{k \in S : \widehat{r}_k = -1\} > \lfloor s/2 \rfloor$, based on (E.58), by replacing the first s entries of $\widehat{\mathbf{r}}$ with -1 , we get a different alignment whose score is smaller than the score of $\widehat{\mathbf{r}}$, which is again contradicted with the definition of $\widehat{\mathbf{r}}$.

In the following, we prove (E.57). The proof of (E.58) is almost the same, so we do not repeat it. Under the conditions we assume, it can be shown that

$$\begin{aligned} \text{score}(\mathbf{r}) - \text{score}(\mathbf{r}') &= \sum_{k_1=1}^{k'} \sum_{k_2=1}^{k'} \|\widehat{\beta}^{(k_1)[0]} - \widehat{\beta}^{(k_2)[0]}\|_2 + 2 \sum_{k_1=1}^{k'} \sum_{k_2=k'+1}^s \|\widehat{\beta}^{(k_1)[0]} + \widehat{\beta}^{(k_2)[0]}\|_2 \\ &\quad + 2 \sum_{k_1=1}^{k'} \sum_{k_2=s+1}^K \|\widehat{\beta}^{(k_1)[0]} + r_{k_2} \widehat{\beta}^{(k_2)[0]}\|_2 \\ &\quad - \sum_{k_1=1}^{k'} \sum_{k_2=1}^{k'} \|\widehat{\beta}^{(k_1)[0]} - \widehat{\beta}^{(k_2)[0]}\|_2 - 2 \sum_{k_1=1}^{k'} \sum_{k_2=k'+1}^s \|\widehat{\beta}^{(k_1)[0]} - \widehat{\beta}^{(k_2)[0]}\|_2 \\ &\quad - 2 \sum_{k_1=1}^{k'} \sum_{k_2=s+1}^K \|\widehat{\beta}^{(k_1)[0]} + r_{k_2} \widehat{\beta}^{(k_2)[0]}\|_2 \\ &= 2 \underbrace{\sum_{k_1=1}^{k'} \sum_{k_2=k'+1}^s \|\widehat{\beta}^{(k_1)[0]} + \widehat{\beta}^{(k_2)[0]}\|_2}_{(1)} + 2 \underbrace{\sum_{k_1=1}^{k'} \sum_{k_2=s+1}^K \|\widehat{\beta}^{(k_1)[0]} + r_{k_2} \widehat{\beta}^{(k_2)[0]}\|_2}_{(2)} \end{aligned}$$

$$\begin{aligned}
& -2 \underbrace{\sum_{k_1=1}^{k'} \sum_{k_2=k'+1}^s \|\widehat{\beta}^{(k_1)[0]} - \widehat{\beta}^{(k_2)[0]}\|_2}_{(1)'} - 2 \underbrace{\sum_{k_1=1}^{k'} \sum_{k_2=s+1}^K \|\widehat{\beta}^{(k_1)[0]} + r_{k_2} \widehat{\beta}^{(k_2)[0]}\|_2}_{(2)'} . \\
& \hspace{15em} \text{(E.59)}
\end{aligned}$$

And

$$\begin{aligned}
(1) - (1)' &= \sum_{k_1=1}^{k'} \sum_{k_2=k'+1}^s (\|\widehat{\beta}^{(k_1)[0]} + \widehat{\beta}^{(k_2)[0]}\|_2 - \|\widehat{\beta}^{(k_1)[0]} - \widehat{\beta}^{(k_2)[0]}\|_2) \\
&\geq \sum_{k_1=1}^{k'} \sum_{k_2=k'+1}^s (\|\beta^{(k_1)*} + \beta^{(k_2)*}\|_2 - \|\beta^{(k_1)*} - \beta^{(k_2)*}\|_2 - 4\xi) \\
&\geq \sum_{k_1=1}^{k'} \sum_{k_2=k'+1}^s (2\|\beta^{(k_1)*}\|_2 - 2\|\beta^{(k_1)*} - \beta^{(k_2)*}\|_2 - 4\xi) \\
&\geq 2(s - k') \sum_{k_1=1}^{k'} \|\beta^{(k_1)*}\|_2 - 4k'(s - k')h_{\beta} - 4k'(s - k')\xi, \\
(2) - (2)' &\geq - \sum_{k_1=1}^{k'} \sum_{k_2=s+1}^K 2\|\widehat{\beta}^{(k_1)[0]}\|_2 \geq -2(K - s) \sum_{k_1=1}^{k'} \|\beta^{(k_1)*}\|_2 - 2k'(K - s)\xi.
\end{aligned}$$

Combining all these pieces,

$$\begin{aligned}
& \text{score}(\mathbf{r}) - \text{score}(\mathbf{r}') \\
& \geq 2(2s - k' - K) \sum_{k_1=1}^{k'} \|\beta^{(k_1)*}\|_2 - 4k'(s - k')h_{\mu} - 2k'(K - s)\xi - 4k'(s - k')\xi \\
& \geq 2k' \left[(2s - k' - K) \min_{k \in S} \|\beta^{(k)*}\|_2 - 2(s - k')h_{\mu} - (K - s)\xi - 2(s - k')\xi \right] \\
& > 2k' \left[\left(\frac{3}{2}s - K \right) \min_{k \in S} \|\beta^{(k)*}\|_2 - 2sh_{\mu} - (K - s)\xi - 2s\xi \right] \tag{E.60} \\
& \geq 0, \tag{E.61}
\end{aligned}$$

where (E.60) holds because $1 \leq k' \leq \lfloor s/2 \rfloor$ and (E.61) is due to the condition (ii).

E.7 Proof of Theorem 6

Denote $\xi = \max_{k \in S} \min_{r_k = \pm 1} \|r_k \widehat{\beta}^{(k)[0]} - \beta^{(k)*}\|_2 = \max_{k \in S} (\|\widehat{\beta}^{(k)[0]} - \beta^{(k)*}\|_2 \wedge \|\widehat{\beta}^{(k)[0]} + \beta^{(k)*}\|_2)$. WLOG, assume $S = \{1, \dots, s\}$ and $r_k^* = 1$ for all $k \in S$. Hence $\xi = \max_{k \in S} \|\widehat{\beta}^{(k)[0]} - \beta^{(k)*}\|_2$. For any $k' = 1, \dots, sp_a$, define

$$\mathbf{r} = (\underbrace{r_1, \dots, r_{k'}}_{=-1}, \underbrace{r_{k'+1}, \dots, r_s}_{=1}, \underbrace{r_{s+1}, \dots, r_K}_{\text{outlier tasks}}),$$

$$\mathbf{r}' = \underbrace{(r_1, \dots, r_{k'-1})}_{=-1}, \underbrace{(\mathbf{r}_{k'}', r_{k'+1}, \dots, r_s)}_{=1}, \underbrace{(r_{s+1}, \dots, r_K)}_{\text{outlier tasks}}.$$

By the definition of p_a , we must have $\#\{k \in S : \hat{r}_k = -1\} = sp_a$ or $\#\{k \in S : \hat{r}_k = 1\} = sp_a$. If $\#\{k \in S : \hat{r}_k = -1\} = sp_a$ and we have

$$\text{score}(\mathbf{r}) - \text{score}(\mathbf{r}') > 0, \quad (\text{E.62})$$

then for each $k \in S$ in the for loop of Algorithm 3, the algorithm will flip the sign of $\hat{r}_{k'}$ to decrease the mis-alignment proportion p_a . Then after the for loop, the mis-alignment proportion p_a will become zero, which means the correct alignment is achieved. The case that $\#\{k \in S : \hat{r}_k = 1\} = sp_a$ can be similarly discussed.

Now we derive (E.62). Similar to the decomposition in (E.59), we have

$$\begin{aligned} \text{score}(\mathbf{r}) - \text{score}(\mathbf{r}') &= 2 \underbrace{\sum_{k=k'+1}^s \|\hat{\boldsymbol{\beta}}^{(k')}[0] + \hat{\boldsymbol{\beta}}^{(k)}[0]\|_2}_{(1)} + 2 \underbrace{\sum_{k=1}^{k'-1} \|\hat{\boldsymbol{\beta}}^{(k')}[0] - \hat{\boldsymbol{\beta}}^{(k)}[0]\|_2}_{(2)} \\ &\quad + 2 \underbrace{\sum_{k=s+1}^K \|\hat{\boldsymbol{\beta}}^{(k')}[0] - r_k \hat{\boldsymbol{\beta}}^{(k)}[0]\|_2}_{(3)} \\ &\quad - 2 \underbrace{\sum_{k=k'+1}^s \|\hat{\boldsymbol{\beta}}^{(k')}[0] - \hat{\boldsymbol{\beta}}^{(k)}[0]\|_2}_{(1)'} - 2 \underbrace{\sum_{k=1}^{k'-1} \|\hat{\boldsymbol{\beta}}^{(k')}[0] + \hat{\boldsymbol{\beta}}^{(k)}[0]\|_2}_{(2)'} \\ &\quad - 2 \underbrace{\sum_{k=s+1}^K \|\hat{\boldsymbol{\beta}}^{(k')}[0] - r_k \hat{\boldsymbol{\beta}}^{(k)}[0]\|_2}_{(3)'} . \end{aligned}$$

Note that

$$\begin{aligned} (1) - (1)' &= \sum_{k=k'+1}^s (\|\hat{\boldsymbol{\beta}}^{(k')}[0] + \hat{\boldsymbol{\beta}}^{(k)}[0]\|_2 - \|\hat{\boldsymbol{\beta}}^{(k')}[0] - \hat{\boldsymbol{\beta}}^{(k)}[0]\|_2) \\ &\geq \sum_{k=k'+1}^s (\|\boldsymbol{\beta}^{(k')*} + \boldsymbol{\beta}^{(k)*}\|_2 - \|\boldsymbol{\beta}^{(k')*} - \boldsymbol{\beta}^{(k)*}\|_2 - 4\xi) \\ &\geq \sum_{k=k'+1}^s (2\|\boldsymbol{\beta}^{(k')*}\|_2 - 2\|\boldsymbol{\beta}^{(k')*} - \boldsymbol{\beta}^{(k)*}\|_2 - 4\xi) \\ &\geq (s - k')(2\|\boldsymbol{\beta}^{(k')*}\|_2 - 4h_{\boldsymbol{\beta}} - 4\xi), \\ (2) - (2)' &\geq - \sum_{k=1}^{k'-1} 2\|\hat{\boldsymbol{\beta}}^{(k')}[0]\|_2 \geq -2(k' - 1)\|\boldsymbol{\beta}^{(k')*}\|_2 - 2(k' - 1)\xi, \end{aligned}$$

$$(3) - (3)' \geq - \sum_{k=s+1}^K 2\|\widehat{\boldsymbol{\beta}}^{(k')^{[0]}}\|_2 \geq -2(K-s)\|\boldsymbol{\beta}^{(k')^*}\|_2 - 2(K-s)\xi.$$

Putting all pieces together,

$$\begin{aligned} & \text{score}(\mathbf{r}) - \text{score}(\mathbf{r}') \\ & \geq 2 \left[(2s - 2k' - K + 1)\|\boldsymbol{\beta}^{(k')^*}\|_2 - 2(s - k')h_{\boldsymbol{\beta}} - (s - k' + K - 1)\xi \right] \\ & > 2 \left[(2s - 2sp_a - K)\|\boldsymbol{\beta}^{(k')^*}\|_2 - 2sh_{\boldsymbol{\beta}} - 2(s + K)\xi \right] \tag{E.63} \\ & \geq 0. \tag{E.64} \end{aligned}$$

where (E.63) holds because $1 \leq k' \leq sp_a$ and (E.64) is due to the condition (iii).

E.8 Proof of Theorem 11

E.8.1 Lemmas

Define the contraction basin of one GMM as

$$B_{\text{con}}(\boldsymbol{\theta}^{(k)*}) = \{\boldsymbol{\theta} = \{w, \boldsymbol{\beta}, \delta\} : w_r \in [c_w/2, 1 - c_w/2], \|\boldsymbol{\beta} - \boldsymbol{\beta}^{(k)*}\|_2 \leq C_b\Delta, |\delta - \delta^{(k)*}| \leq C_b\Delta\},$$

for which we may shorthand as B_{con} in the following.

For GMM $\mathbf{z} \sim (1 - w^*)\mathcal{N}(\boldsymbol{\mu}_1^*, \boldsymbol{\Sigma}^*) + w^*\mathcal{N}(\boldsymbol{\mu}_2^*, \boldsymbol{\Sigma}^*)$ and any $\boldsymbol{\theta} = (w, \boldsymbol{\beta}, \delta)$, define

$$\begin{aligned} \gamma_{\boldsymbol{\theta}}(\mathbf{z}) &= \frac{w \exp\{\boldsymbol{\beta}^T \mathbf{z} - \delta\}}{1 - w + w \exp\{\boldsymbol{\beta}^T \mathbf{z} - \delta_r\}}, \quad w(\boldsymbol{\theta}) = \mathbb{E}[\gamma_{\boldsymbol{\theta}}(\mathbf{z})], \\ \boldsymbol{\mu}_1(\boldsymbol{\theta}) &= \frac{\mathbb{E}[(1 - \gamma_{\boldsymbol{\theta}}(\mathbf{z}))\mathbf{z}]}{\mathbb{E}[1 - \gamma_{\boldsymbol{\theta}}(\mathbf{z})]}, \quad \boldsymbol{\mu}_2(\boldsymbol{\theta}) = \frac{\mathbb{E}[\gamma_{\boldsymbol{\theta}}(\mathbf{z})\mathbf{z}]}{\mathbb{E}[\gamma_{\boldsymbol{\theta}}(\mathbf{z})]}. \end{aligned}$$

Lemma 29. *Suppose Assumption 3 holds.*

(i) *With probability at least $1 - \tau$,*

$$\sup_{\substack{\boldsymbol{\theta}^{(0)} \in B_{\text{con}} \\ \|\boldsymbol{\beta}^{(0)} - \boldsymbol{\beta}^{(0)*}\|_2 \leq \xi^{(0)}}} \left| \frac{1}{n_0} \sum_{i=1}^{n_0} \gamma_{\boldsymbol{\theta}^{(0)}}(\mathbf{z}_i^{(0)}) - \mathbb{E}[\gamma_{\boldsymbol{\theta}^{(0)}}(\mathbf{z}^{(0)})] \right| \lesssim \xi^{(0)} \sqrt{\frac{p}{n_0}} + \sqrt{\frac{\log(1/\tau)}{n_0}}.$$

(ii) *With probability at least $1 - \tau$,*

$$\sup_{\substack{\boldsymbol{\theta}^{(0)} \in B_{\text{con}} \\ \|\boldsymbol{\beta}^{(0)} - \boldsymbol{\beta}^{(0)*}\|_2 \leq \xi^{(0)}}} \left| \frac{1}{n_0} \sum_{i=1}^{n_0} \gamma_{\boldsymbol{\theta}^{(0)}}(\mathbf{z}_i^{(0)}) (\mathbf{z}_i^{(0)})^T \boldsymbol{\beta}^{(0)*} - \mathbb{E}[\gamma_{\boldsymbol{\theta}^{(0)}}(\mathbf{z}^{(0)}) (\mathbf{z}_i^{(0)})^T \boldsymbol{\beta}^{(0)*}] \right| \lesssim \xi^{(0)} \sqrt{\frac{p}{n_0}} + \sqrt{\frac{\log(1/\tau)}{n_0}}.$$

(iii) *With probability at least $1 - \tau$,*

$$\sup_{\substack{\boldsymbol{\theta}^{(0)} \in B_{\text{con}} \\ \|\boldsymbol{\beta}^{(0)} - \boldsymbol{\beta}^{(0)*}\|_2 \leq \xi^{(0)}}} \left\| \frac{1}{n_0} \sum_{i=1}^{n_0} \gamma_{\boldsymbol{\theta}^{(0)}}(\mathbf{z}_i^{(0)}) \mathbf{z}_i^{(0)} - \mathbb{E}[\gamma_{\boldsymbol{\theta}^{(0)}}(\mathbf{z}^{(0)}) \mathbf{z}_i^{(0)}] \right\|_2 \lesssim \xi^{(0)} \sqrt{\frac{p}{n_0}} + \sqrt{\frac{\log(1/\tau)}{n_0}}.$$

E.8.2 Main proof of Theorem 11

WLOG, in Assumptions 3.(iii) and 3.(iv), we assume

- $\|\widehat{\beta}^{(0)[0]} - \beta^{(0)*}\|_2 \vee |\widehat{\delta}^{(0)[0]} - \delta^{(0)*}| \leq C_4 \Delta^{(0)}$, with a sufficiently small constant C_4 ;
- $|\widehat{w}^{(0)[0]} - w^{(0)*}| \leq c_w/2$.

(I) Case 1: We first consider the case that $h \geq C\sqrt{\frac{p}{n_0}}$. Consider an event \mathcal{E} defined to be the intersection of the events in Lemma 29, with $\xi^{(k)} =$ a large constant C , which satisfies $\mathbb{P}(\mathcal{E}) \geq 1 - \tau$. Throughout the analysis in Case 1, we condition on \mathcal{E} , therefore all the arguments hold with probability at least $1 - \tau$.

Similar to our analysis in the proof of Theorem 1, conditioned on \mathcal{E} , we have

$$\begin{aligned} |\widehat{w}^{(0)[t]} - w^{(0)*}| &\lesssim \kappa_0'' d(\widehat{\theta}^{(0)[t-1]}, \theta^{(0)*}) + \sqrt{\frac{p}{n_0}}, \\ \max_{r=1:2} \|\widehat{\mu}_r^{(0)[t]} - \mu_r^{(0)*}\|_2 &\lesssim \kappa_0'' d(\widehat{\theta}^{(0)[t-1]}, \theta^{(0)*}) + \sqrt{\frac{p}{n_0}}, \end{aligned} \quad (\text{E.65})$$

$$\|(\widehat{\Sigma}^{(0)[t]} - \Sigma^{(0)*})\beta^{(0)*}\|_2 \lesssim \kappa_0'' d(\widehat{\theta}^{(0)[t-1]}, \theta^{(0)*}) + \sqrt{\frac{p}{n_0}}. \quad (\text{E.66})$$

Hence

$$\begin{aligned} \|(\widehat{\Sigma}^{(0)[t]})\beta^{(0)*} - (\widehat{\mu}_2^{(0)[t-1]} - \widehat{\mu}_1^{(0)[t-1]})\|_2 &\lesssim \|(\widehat{\Sigma}^{(0)[t]} - \Sigma^{(0)*})\beta^{(0)*}\|_2 + \max_{r=1:2} \|\widehat{\mu}_r^{(0)[t]} - \mu_r^{(0)*}\|_2 \\ &\lesssim \kappa_0'' d(\widehat{\theta}^{(0)[t-1]}, \theta^{(0)*}) + \sqrt{\frac{p}{n_0}}. \end{aligned}$$

By Lemma 7, we have

$$\begin{aligned} \|\widehat{\beta}^{(0)[t]} - \beta^{(0)*}\|_2 &\lesssim \|(\widehat{\Sigma}^{(0)[t]})\beta^{(0)*} - (\widehat{\mu}_2^{(0)[t-1]} - \widehat{\mu}_1^{(0)[t-1]})\|_2 + \frac{\lambda_0^{[t]}}{\sqrt{n_0}} \\ &\lesssim \kappa_0'' d(\widehat{\theta}^{(0)[t-1]}, \theta^{(0)*}) + \sqrt{\frac{p}{n_0}} + \frac{\lambda_0^{[t]}}{\sqrt{n_0}}. \end{aligned}$$

Combining these results, we have

$$d(\widehat{\theta}^{(0)[t]}, \theta^{(0)*}) \leq C\kappa_0'' d(\widehat{\theta}^{(0)[t-1]}, \theta^{(0)*}) + C'\sqrt{\frac{p}{n_0}} + C'\frac{\lambda_0^{[t]}}{\sqrt{n_0}}.$$

By the construction of $\lambda_0^{[t]}$, we know that

$$\lambda_0^{[t]} = \frac{1 - \kappa_0^t}{1 - \kappa_0} C_{\lambda_0} \sqrt{p} + \kappa_0^t \lambda_0^{[0]},$$

implies that

$$\begin{aligned}
d(\widehat{\boldsymbol{\theta}}^{(0)[t]}, \boldsymbol{\theta}^{(0)*}) &\leq (C\kappa_0'')^t d(\widehat{\boldsymbol{\theta}}^{(0)[0]}, \boldsymbol{\theta}^{(0)*}) + C'' \sqrt{\frac{p}{n_0}} + C' \sum_{t'=1}^t \frac{\lambda_0^{[t']}}{\sqrt{n_0}} (C\kappa_0'')^{t-t'} \\
&\leq (\kappa_0')^t d(\widehat{\boldsymbol{\theta}}^{(0)[0]}, \boldsymbol{\theta}^{(0)*}) + C''' \sqrt{\frac{p}{n_0}} + C''' t (\kappa_0')^t \\
&\leq Ct(\kappa_0')^t + C''' \sqrt{\frac{p}{n_0}},
\end{aligned}$$

which is the desired rate. The bound of $\max_{r=1:2} \|\widehat{\boldsymbol{\mu}}_r^{(0)[t]} - \boldsymbol{\mu}_r^{(0)*}\|_2$ and $\|\widehat{\boldsymbol{\Sigma}}^{(0)[t]} - \boldsymbol{\Sigma}^{(0)*}\|_2$ can be derived similar to (E.65) and (E.66).

(II) Case 2: Next, we consider the case that $h \leq C\sqrt{\frac{p}{n_0}}$. According to Assumption 3, we have $\sqrt{\frac{p}{n_0}} \lesssim \sqrt{\frac{p+\log K}{\max_{k \in S} n_k}}$. It is easy to see that the analysis in part (I) does not depend on the condition $h \geq C\sqrt{\frac{p}{n_0}}$. Hence we have proved the desired bounds of $\max_{r=1:2} \|\widehat{\boldsymbol{\mu}}_r^{(0)[t]} - \boldsymbol{\mu}_r^{(0)*}\|_2$ and $\|\widehat{\boldsymbol{\Sigma}}^{(0)[t]} - \boldsymbol{\Sigma}^{(0)*}\|_2$. Denote t_0 as an integer such that $t_0 \kappa_0^{t_0} \asymp \sqrt{\frac{p}{n_0}}$. When $1 \leq t \leq t_0$, the bound in part (I) is the desired bound since the term $t\kappa_0^t$ dominates the other terms. Let us consider the case $t = t_0 + 1$.

Consider an event \mathcal{E}' defined to be the event of

$$\|\bar{\boldsymbol{\beta}}^{[T]} - \boldsymbol{\beta}^{(k')*}\|_2 \lesssim h + \sqrt{\frac{\log K}{n_{k'}}} + \sqrt{\frac{p}{n_S}} + \epsilon \sqrt{\frac{p + \log K}{\max_{k=1:K} n_k}}, \quad k' \in \arg \min_{k \in S} n_k.$$

Note that since $h \leq C\sqrt{\frac{p+\log K}{\max_{k \in S} n_k}}$, by part (II) of the proof of Theorem 1, we know that $\mathbb{P}(\mathcal{E}') \geq 1 - C(K^{-1} + \exp\{-C'p\})$. And \mathcal{E}' implies that

$$\|\bar{\boldsymbol{\beta}}^{[T]} - \boldsymbol{\beta}^{(0)*}\|_2 \lesssim h + \sqrt{\frac{\log K}{\max_{k \in S} n_k}} + \sqrt{\frac{p}{n_S}} + \epsilon \sqrt{\frac{p + \log K}{\max_{k=1:K} n_k}} \lesssim \sqrt{\frac{p}{n_0}},$$

where the second inequality comes from Assumption 3.

Also consider another event \mathcal{E}'' defined to be the intersection of the events in Lemma 29, with $\xi = C\left[h + \sqrt{\frac{\log K}{\max_{k \in S} n_k}} + \sqrt{\frac{p}{n_S}} + \epsilon \sqrt{\frac{p+\log K}{\max_{k=1:K} n_k}}\right]$, which satisfies $\mathbb{P}(\mathcal{E}'') \geq 1 - \tau$. Throughout the analysis in Case 1, we condition on $\mathcal{E} \cap \mathcal{E}' \cap \mathcal{E}''$, therefore all the arguments hold with probability at least $1 - \tau - C(K^{-1} + \exp\{-C'p\})$.

Note that $\lambda_0^{[t]} \geq C\sqrt{p} \geq C'\|\bar{\boldsymbol{\beta}}^{[T]} - \boldsymbol{\beta}^{(0)*}\|_2$ and $\lambda_0^{[t]} \geq C\sqrt{p} \geq C'\sqrt{n_0}(\widehat{\boldsymbol{\Sigma}}^{(0)[t]})\boldsymbol{\beta}^{(0)*} - (\widehat{\boldsymbol{\mu}}_2^{(0)[t-1]} - \widehat{\boldsymbol{\mu}}_1^{(0)[t-1]})\|_2$. Hence by Lemma 7, we have $\widehat{\boldsymbol{\beta}}^{(0)[t]} = \bar{\boldsymbol{\beta}}^{[T]}$ thus

$$\|\widehat{\boldsymbol{\beta}}^{(0)[t]} - \boldsymbol{\beta}^{(0)*}\|_2 \lesssim h + \sqrt{\frac{\log K}{\max_{k \in S} n_k}} + \sqrt{\frac{p}{n_S}} + \epsilon \sqrt{\frac{p + \log K}{\max_{k=1:K} n_k}}.$$

Similar to the analysis in part (II) in the proof of Theorem 1, we have

$$\begin{aligned}
|\widehat{w}^{(0)[t]} - w^{(0)*}| &\lesssim \kappa_0'' d(\widehat{\boldsymbol{\theta}}^{(0)[t-1]}, \boldsymbol{\theta}^{(0)*}) + \xi \sqrt{\frac{p}{n_0}} + \sqrt{\frac{1}{n_0}} \\
&\lesssim \kappa_0'' d(\widehat{\boldsymbol{\theta}}^{(0)[t-1]}, \boldsymbol{\theta}^{(0)*}) + \xi + \sqrt{\frac{1}{n_0}}, \\
|\widehat{\delta}^{(0)[t]} - \delta^{(0)*}| &\lesssim \kappa_0'' d(\widehat{\boldsymbol{\theta}}^{(0)[t-1]}, \boldsymbol{\theta}^{(0)*}) + \xi + \xi \sqrt{\frac{p}{n_0}} + \sqrt{\frac{1}{n_0}}, \\
&\lesssim \kappa_0'' d(\widehat{\boldsymbol{\theta}}^{(0)[t-1]}, \boldsymbol{\theta}^{(0)*}) + \xi + \sqrt{\frac{1}{n_0}}.
\end{aligned}$$

Putting all pieces together,

$$d(\widehat{\boldsymbol{\theta}}^{(0)[t]}, \boldsymbol{\theta}^{(0)*}) \leq C\kappa_0'' d(\widehat{\boldsymbol{\theta}}^{(0)[t-1]}, \boldsymbol{\theta}^{(0)*}) + h + \sqrt{\frac{\log K}{\max_{k \in S} n_k}} + \sqrt{\frac{p}{n_S}} + \epsilon \sqrt{\frac{p + \log K}{\max_{k=1:K} n_k}} + \sqrt{\frac{1}{n_0}}.$$

We can continue this analysis from $t = t_0 + 1$ to $t_0 + 2$, and so on. Hence for any $t' \geq 1$, we have

$$\begin{aligned}
d(\widehat{\boldsymbol{\theta}}^{(0)[t_0+t']}, \boldsymbol{\theta}^{(0)*}) &\leq (C\kappa_0'')^{t'} d(\widehat{\boldsymbol{\theta}}^{(0)[t_0]}, \boldsymbol{\theta}^{(0)*}) + C'h + C' \sqrt{\frac{\log K}{\max_{k \in S} n_k}} + C' \sqrt{\frac{p}{n_S}} \\
&\quad + C' \epsilon \sqrt{\frac{p + \log K}{\max_{k=1:K} n_k}} + C' \sqrt{\frac{1}{n_0}} \\
&\leq (\kappa_0')^{t'} d(\widehat{\boldsymbol{\theta}}^{(0)[t_0]}, \boldsymbol{\theta}^{(0)*}) + C'h + C' \sqrt{\frac{\log K}{\max_{k \in S} n_k}} + C' \sqrt{\frac{p}{n_S}} \\
&\quad + C' \epsilon \sqrt{\frac{p + \log K}{\max_{k=1:K} n_k}} + C' \sqrt{\frac{1}{n_0}} \\
&\leq (t' + t_0)(\kappa_0')^{t'+t_0} + C'h + C' \sqrt{\frac{\log K}{\max_{k \in S} n_k}} + C' \sqrt{\frac{p}{n_S}} \\
&\quad + C' \epsilon \sqrt{\frac{p + \log K}{\max_{k=1:K} n_k}} + C' \sqrt{\frac{1}{n_0}},
\end{aligned}$$

where the last inequality holds because t_0 is chosen to be the integer satisfying $t_0(\kappa_0)^{t_0} \asymp \sqrt{\frac{p}{n_0}} \gtrsim d(\widehat{\boldsymbol{\theta}}^{(0)[t_0]}, \boldsymbol{\theta}^{(0)*})$.

E.8.3 Proof of lemmas

Proof of Lemma 29. The proof is almost the same as the proofs of lemmas in Theorem 1, so we do not repeat it here. \square

E.9 Proof of Theorem 13

E.9.1 Lemmas

Recall

$$\overline{\Theta}'_S(h) = \left\{ \{\overline{\boldsymbol{\theta}}^{(k)}\}_{k \in \{0\} \cup S} = \{(w^{(k)}, \boldsymbol{\mu}_1^{(k)}, \boldsymbol{\mu}_2^{(k)}, \boldsymbol{\Sigma}^{(k)})\}_{k \in \{0\} \cup S} : \boldsymbol{\theta}^{(k)} \in \overline{\Theta}, \max_{k \in S} \|\boldsymbol{\beta}^{(k)} - \boldsymbol{\beta}^{(0)}\|_2 \leq h \right\}.$$

Lemma 30. *When $n_0 \geq Cp$ with some constant $C > 0$, we have*

$$\inf_{\widehat{\boldsymbol{\theta}}^{(0)}} \sup_{S: |S| \geq s} \sup_{\substack{\{\overline{\boldsymbol{\theta}}^{(k)*}\}_{k \in \{0\} \cup S} \in \overline{\Theta}'_S \\ \mathbb{Q}_S}} \mathbb{P} \left(d(\widehat{\boldsymbol{\theta}}^{(0)}, \boldsymbol{\theta}^{(0)*}) \geq C_1 \sqrt{\frac{p}{n_S + n_0}} + C_1 h \wedge \sqrt{\frac{p}{n_0}} + C_1 \sqrt{\frac{1}{n_0}} \right) \geq \frac{1}{10}.$$

Lemma 31. *Denote $\tilde{\epsilon} = \frac{K-s}{s}$. Then*

$$\inf_{\widehat{\boldsymbol{\theta}}^{(0)}} \sup_{S: |S| \geq s} \sup_{\substack{\{\overline{\boldsymbol{\theta}}^{(k)*}\}_{k \in \{0\} \cup S} \in \overline{\Theta}'_S \\ \mathbb{Q}_S}} \mathbb{P} \left(d(\widehat{\boldsymbol{\theta}}^{(0)}, \boldsymbol{\theta}^{(0)*}) \geq \left(C_1 \frac{\tilde{\epsilon}}{\sqrt{\max_{k=1:K} n_k}} \right) \wedge \left(C_2 \sqrt{\frac{1}{n_0}} \right) \right) \geq \frac{1}{10}.$$

Lemma 32 (The second variant of Theorem 5.1 in [Chen et al. \(2018\)](#)). *Given a series of distributions $\{\{\mathbb{P}_\theta^{(k)}\}_{k=0}^K : \theta \in \Theta\}$, each of which is indexed by the same parameter $\theta \in \Theta$. Consider $\mathbf{x}^{(k)} \sim (1 - \tilde{\epsilon})\mathbb{P}_\theta^{(k)} + \tilde{\epsilon}\mathbb{Q}^{(k)}$ independently for $k = 1 : K$ and $\mathbf{x}^{(0)} \sim \mathbb{P}_\theta^{(0)}$. Denote the joint distribution of $\{\mathbf{x}^{(k)}\}_{k=0}^K$ as $\mathbb{P}_{(\tilde{\epsilon}, \theta, \{\mathbb{Q}^{(k)}\}_{k=1}^K)}$. Then*

$$\inf_{\widehat{\theta}} \sup_{\substack{\theta \in \Theta \\ \{\mathbb{Q}^{(k)}\}_{k=1}^K}} \mathbb{P}_{(\tilde{\epsilon}, \theta, \{\mathbb{Q}^{(k)}\}_{k=1}^K)} \left(\|\widehat{\theta} - \theta\| \geq C\varpi'(\tilde{\epsilon}, \Theta) \right) \geq \frac{9}{20},$$

where $\varpi'(\tilde{\epsilon}, \Theta) := \sup \{ \|\theta_1 - \theta_2\| : \max_{k=1:K} d_{\text{TV}}(\mathbb{P}_{\theta_1}^{(k)}, \mathbb{P}_{\theta_2}^{(k)}) \leq \tilde{\epsilon}/(1 - \tilde{\epsilon}), d_{\text{TV}}(\mathbb{P}_{\theta_1}^{(0)}, \mathbb{P}_{\theta_2}^{(0)}) \leq 1/20 \}$.

Lemma 33. *Consider two data generating mechanisms:*

- (i) $\mathbf{x}^{(k)} \sim (1 - \tilde{\epsilon}')\mathbb{P}_\theta^{(k)} + \tilde{\epsilon}'\mathbb{Q}^{(k)}$ independently for $k = 1 : K$ and $\mathbf{x}^{(0)} \sim \mathbb{P}_\theta^{(0)}$, where $\tilde{\epsilon}' = \frac{K-s}{K}$;
- (ii) With a preserved set $S \subseteq 1 : K$, generate $\{\mathbf{x}^{(k)}\}_{k \in S^c} \sim \mathbb{Q}_S$ and $\mathbf{x}^{(k)} \sim \mathbb{P}_\theta^{(k)}$ independently for $k \in S$. And $\mathbf{x}^{(0)} \sim \mathbb{P}_\theta^{(0)}$.

Denote the joint distributions of $\{\mathbf{x}^{(k)}\}_{k=0}^K$ in (i) and (ii) as $\mathbb{P}_{(\tilde{\epsilon}, \theta, \{\mathbb{Q}^{(k)}\}_{k=1}^K)}$ and $\mathbb{P}_{(S, \theta, \mathbb{Q})}$, respectively. We claim that if

$$\inf_{\widehat{\theta}} \sup_{\substack{\theta \in \Theta \\ \{\mathbb{Q}^{(k)}\}_{k=1}^K}} \mathbb{P}_{(\frac{K-s}{50K}, \theta, \{\mathbb{Q}^{(k)}\}_{k=1}^K)} \left(\|\widehat{\theta} - \theta\| \geq C\varpi' \left(\frac{K-s}{50K}, \Theta \right) \right) \geq \frac{9}{20},$$

then

$$\inf_{\hat{\theta}} \sup_{S: |S| \geq s} \sup_{\substack{\theta \in \Theta \\ \mathbb{Q}_S}} \mathbb{P}_{(S, \theta, \mathbb{Q}_S)} \left(\|\hat{\theta} - \theta\| \geq C \varpi' \left(\frac{K-s}{50K}, \Theta \right) \right) \geq \frac{1}{10},$$

where $\varpi'(\tilde{\epsilon}, \Theta) := \sup \{ \|\theta_1 - \theta_2\| : \max_{k=1:K} \text{KL}(\mathbb{P}_{\theta_1}^{(k)} \| \mathbb{P}_{\theta_2}^{(k)}) \leq [\tilde{\epsilon}/(1-\tilde{\epsilon})]^2, \text{KL}(\mathbb{P}_{\theta_1}^{(0)} \| \mathbb{P}_{\theta_2}^{(0)}) \leq 1/400 \}$ for any $\tilde{\epsilon} \in (0, 1)$.

Lemma 34. When $n_0 \geq Cp$ with some constant $C > 0$, we have

$$\inf_{\hat{\Sigma}^{(0)}} \sup_{S: |S| \geq s} \sup_{\substack{\{\bar{\theta}^{(k)*}\}_{k \in \{0\} \cup S} \in \bar{\Theta}'_S \\ \mathbb{Q}_S}} \mathbb{P} \left(\|\hat{\Sigma}^{(0)} - \Sigma^{(0)*}\|_2 \geq C \sqrt{\frac{p}{n_0}} \right) \geq \frac{1}{10}.$$

E.9.2 Main proof of Theorem 13

E.9.3 Proof of lemmas

Proof of Lemma 30. It's easy to see that $\bar{\Theta}'_S \supseteq \bar{\Theta}'_{S,w} \cup \bar{\Theta}'_{S,\beta} \cup \bar{\Theta}'_{S,\delta}$, where

$$\begin{aligned} \bar{\Theta}'_{S,w} &= \left\{ \{\bar{\theta}^{(k)}\}_{k \in \{0\} \cup S} : \mu_1^{(k)} = \mathbf{1}_p / \sqrt{p}, \mu_2^{(k)} = -\mu_1^{(k)}, \Sigma^{(k)} = \mathbf{I}_p, w^{(k)} \in (c_w, 1 - c_w) \right\}, \\ \bar{\Theta}'_{S,\beta} &= \left\{ \{\bar{\theta}^{(k)}\}_{k \in \{0\} \cup S} : \Sigma^{(k)} = \mathbf{I}_p, w^{(k)} = \frac{1}{2}, \|\mu_1^{(k)}\|_2 \vee \|\mu_2^{(k)}\|_2 \leq M, \max_{k \in S} \|\beta^{(k)} - \beta^{(0)}\|_2 \leq h \right\}, \\ \bar{\Theta}'_{S,\delta} &= \left\{ \{\bar{\theta}^{(k)}\}_{k \in \{0\} \cup S} : \Sigma^{(k)} = \mathbf{I}_p, w^{(k)} = \frac{1}{2}, \|\mu_1^{(k)}\|_2 \vee \|\mu_2^{(k)}\|_2 \leq M, \mu_1^{(k)} = -\frac{1}{2}\mu_0, \right. \\ &\quad \left. \mu_2^{(k)} = \frac{1}{2}\mu_0 + \mathbf{u}, \|\mathbf{u}\|_2 \leq 1 \right\}. \end{aligned}$$

(i) By fixing an S and a \mathbb{Q}_S , we want to show

$$\inf_{\hat{\theta}^{(0)}} \sup_{\{\bar{\theta}^{(k)*}\}_{k \in \{0\} \cup S} \in \bar{\Theta}'_{S,\beta}} \mathbb{P} \left(\|\hat{\beta}^{(0)} - \beta^{(0)*}\|_2 \vee \|\hat{\beta}^{(0)} + \beta^{(0)*}\|_2 \geq C \sqrt{\frac{p}{n_S + n_0}} \right) \geq \frac{1}{4}.$$

By Lemma 4, \exists a quadrant \mathcal{Q}_v of \mathbb{R}^p and a $r/8$ -packing of $(r\mathcal{S}^p) \cap \mathcal{Q}_v$ under Euclidean norm: $\{\tilde{\mu}_j\}_{j=1}^N$, where $r = (c\sqrt{p/(n_S + n_0)}) \wedge M \leq 1$ with a small constant $c > 0$ and $N \geq (\frac{1}{2})^p 8^{p-1} = \frac{1}{2} \times 4^{p-1} \geq 2^{p-1}$ when $p \geq 2$. For any $\mu \in \mathbb{R}^p$, denote distribution $\frac{1}{2}\mathcal{N}(\mu, \mathbf{I}_p) + \frac{1}{2}\mathcal{N}(-\mu, \mathbf{I}_p)$ as \mathbb{P}_μ . Then

$$\begin{aligned} \text{LHS} &\geq \inf_{\hat{\mu}} \sup_{\mu \in (r\mathcal{S}^p) \cap \mathcal{Q}_v} \mathbb{P} \left(\|\hat{\mu} - \mu\|_2 \wedge \|\hat{\mu} + \mu\|_2 \geq C \sqrt{\frac{p}{n_S}} \right) \\ &\geq \inf_{\hat{\mu}} \sup_{\mu \in (r\mathcal{S}^p) \cap \mathcal{Q}_v} \mathbb{P} \left(\|\hat{\mu} - \mu\|_2 \geq C \sqrt{\frac{p}{n_S}} \right), \end{aligned} \tag{E.67}$$

where the last inequality holds because it suffices to consider estimator $\hat{\boldsymbol{\mu}}$ satisfying $\hat{\boldsymbol{\mu}}(X) \in (r\mathcal{S}^p) \cap \mathcal{Q}_v$ almost surely. In addition, for any $\mathbf{x}, \mathbf{y} \in \mathcal{Q}_v$, $\|\mathbf{x} - \mathbf{y}\|_2 \leq \|\mathbf{x} + \mathbf{y}\|_2$.

By Lemma 15,

$$\begin{aligned} \text{KL} \left(\prod_{k \in \{0\} \cup S} \mathbb{P}_{\tilde{\boldsymbol{\mu}}_j}^{\otimes n_k} \cdot \mathbb{Q}_S \left\| \prod_{k \in \{0\} \cup S} \mathbb{P}_{\tilde{\boldsymbol{\mu}}_{j'}}^{\otimes n_k} \cdot \mathbb{Q}_S \right. \right) &= \sum_{k \in \{0\} \cup S} n_k \text{KL}(\mathbb{P}_{\tilde{\boldsymbol{\mu}}_j} \|\mathbb{P}_{\tilde{\boldsymbol{\mu}}_{j'}}) \\ &\leq \sum_{k \in \{0\} \cup S} n_k \cdot 8 \|\tilde{\boldsymbol{\mu}}_j\|_2^2 \|\tilde{\boldsymbol{\mu}}_j - \tilde{\boldsymbol{\mu}}_{j'}\|_2^2 \\ &\leq 32(n_S + n_0)r^2 \\ &\leq 32n_S c^2 \cdot \frac{2(p-1)}{n_S + n_0} \\ &\leq \frac{64c^2}{\log 2} \log N. \end{aligned}$$

By Lemma 3,

$$\text{LHS of (E.67)} \geq 1 - \frac{\log 2}{\log N} - \frac{64c^2}{\log 2} \geq 1 - \frac{1}{p-1} - \frac{1}{4} \geq \frac{1}{4},$$

when $C = c/2$, $p \geq 3$ and $c = \sqrt{\log 2}/16$.

(ii) By fixing an S and a \mathbb{Q}_S , we want to show

$$\inf_{\hat{\boldsymbol{\theta}}^{(0)}} \sup_{\{\bar{\boldsymbol{\theta}}^{(k)*}\}_{k \in \{0\} \cup S} \in \bar{\Theta}'_S} \mathbb{P} \left(\|\hat{\boldsymbol{\beta}}^{(0)} - \boldsymbol{\beta}^{(0)*}\|_2 \vee \|\hat{\boldsymbol{\beta}}^{(0)} + \boldsymbol{\beta}^{(0)*}\|_2 \geq C \left[h \wedge \left(c \sqrt{\frac{p}{n_0}} \right) \right] \right) \geq \frac{1}{4}. \quad (\text{E.68})$$

By Lemma 4, \exists a quadrant \mathcal{Q}_v of \mathbb{R}^p and a $r/8$ -packing of $(r\mathcal{S}^{p-1}) \cap \mathcal{Q}_v$ under Euclidean norm: $\{\tilde{\boldsymbol{\theta}}_j\}_{j=1}^N$, where $r = h \wedge (c\sqrt{p/n_0}) \wedge M \leq 1$ with a small constant $c > 0$ and $N \geq (\frac{1}{2})^{p-1} 8^{p-2} = \frac{1}{2} \times 4^{p-2} \geq 2^{p-2}$ when $p \geq 3$. WLOG, assume $M \geq 2$. Denote $\tilde{\boldsymbol{\mu}}_j = (1, \tilde{\boldsymbol{\theta}}_j^T)^T \in \mathbb{R}^p$. Let $\boldsymbol{\mu}_1^{(k)*} = \tilde{\boldsymbol{\mu}} = (1, \mathbf{0}_{p-1})^T$ for all $k \in S$ and $\boldsymbol{\mu}_1^{(0)*} = \boldsymbol{\mu} = (1, \boldsymbol{\vartheta})$ with $\boldsymbol{\vartheta} \in (r\mathcal{S}^{p-1}) \cap \mathcal{Q}_v$. For any $\boldsymbol{\mu} \in \mathbb{R}^p$, denote distribution $\frac{1}{2}\mathcal{N}(\boldsymbol{\mu}, \mathbf{I}_p) + \frac{1}{2}\mathcal{N}(-\boldsymbol{\mu}, \mathbf{I}_p)$ as $\mathbb{P}_{\boldsymbol{\mu}}$. Then similar to the arguments in (i),

$$\begin{aligned} \text{LHS} &\geq \inf_{\hat{\boldsymbol{\mu}}} \sup_{\substack{\boldsymbol{\vartheta} \in (r\mathcal{S}^{p-1}) \cap \mathcal{Q}_v \\ \boldsymbol{\mu} = (1, \boldsymbol{\vartheta})^T}} \mathbb{P} \left(\|\hat{\boldsymbol{\mu}} - \boldsymbol{\mu}\|_2 \wedge \|\hat{\boldsymbol{\mu}} + \boldsymbol{\mu}\|_2 \geq C \left[h \wedge \left(c \sqrt{\frac{p}{n_0}} \right) \right] \right) \\ &\geq \inf_{\hat{\boldsymbol{\mu}}} \sup_{\substack{\boldsymbol{\vartheta} \in (r\mathcal{S}^{p-1}) \cap \mathcal{Q}_v \\ \boldsymbol{\mu} = (1, \boldsymbol{\vartheta})^T}} \mathbb{P} \left(\|\hat{\boldsymbol{\mu}} - \boldsymbol{\mu}\|_2 \geq C \left[h \wedge \left(c \sqrt{\frac{p}{n_0}} \right) \right] \right). \end{aligned}$$

Then by Lemma 15,

$$\text{KL} \left(\mathbb{P}_{\tilde{\boldsymbol{\mu}}_j}^{\otimes n_0} \cdot \prod_{k \in S} \mathbb{P}_{\tilde{\boldsymbol{\mu}}}^{\otimes n_k} \cdot \mathbb{Q}_S \left\| \mathbb{P}_{\tilde{\boldsymbol{\mu}}_{j'}}^{\otimes n_0} \cdot \prod_{k \in S} \mathbb{P}_{\tilde{\boldsymbol{\mu}}}^{\otimes n_k} \cdot \mathbb{Q}_S \right. \right) = n_0 \text{KL}(\mathbb{P}_{\tilde{\boldsymbol{\mu}}_j} \|\mathbb{P}_{\tilde{\boldsymbol{\mu}}_{j'}})$$

$$\begin{aligned}
&\leq n_0 \cdot 8 \|\tilde{\boldsymbol{\mu}}_j\|_2^2 \|\tilde{\boldsymbol{\mu}}_j - \tilde{\boldsymbol{\mu}}_{j'}\|_2^2 \\
&\leq 32n_0 r^2 \\
&\leq 32n_0 c^2 \cdot \frac{3(p-2)}{n_0} \\
&\leq \frac{96c^2}{\log 2} \log N,
\end{aligned}$$

when $n_0 \geq (c^2 \vee M^{-2})p$ and $p \geq 3$. By Fano's lemma (See Corollary 2.6 in [Tsybakov \(2009\)](#)),

$$\text{LHS of (E.68)} \geq 1 - \frac{\log 2}{\log N} - \frac{96c^2}{\log 2} \geq 1 - \frac{1}{p-2} - \frac{1}{4} \geq \frac{1}{4},$$

when $C = 1/2$, $p \geq 4$ and $c = \sqrt{(\log 2)/384}$.

(iii) We want to show

$$\inf_{\hat{\boldsymbol{\theta}}^{(0)}} \sup_{S: |S| \geq s} \sup_{\{\bar{\boldsymbol{\theta}}^{(k)*}\}_{k \in \{0\} \cup S} \in \bar{\Theta}'_{S,w}} \mathbb{P} \left(|\hat{w}^{(0)} - w^{(0)*}| \geq C \sqrt{\frac{1}{n_0}} \right) \geq \frac{1}{4}.$$

The argument is similar to (ii). The only two differences here are that the dimension of interested parameter w equals 1, and Lemma 15 is replaced by Lemma 17.

(iv) We want to show

$$\inf_{\hat{\boldsymbol{\theta}}^{(0)}} \sup_{S: |S| \geq s} \sup_{\{\bar{\boldsymbol{\theta}}^{(k)*}\}_{k \in \{0\} \cup S} \in \bar{\Theta}'_{S,\delta}} \mathbb{P} \left(|\hat{\delta}^{(0)} - \delta^{(0)*}| \geq C \sqrt{\frac{1}{n_0}} \right) \geq \frac{1}{4}$$

The argument is similar to (ii).

Finally, we get the desired conclusion by combining (i)-(iv). \square

Proof of Lemma 31. Let $\tilde{\epsilon} = \frac{K-s}{s}$ and $\tilde{\epsilon}' = \frac{K-s}{K}$. Since $s/K \geq c > 0$, $\tilde{\epsilon} \lesssim \tilde{\epsilon}'$. Denote $\Upsilon_S = \{\{\boldsymbol{\mu}^{(k)}\}_{k \in \{0\} \cup S} : \boldsymbol{\mu}^{(k)} \in \mathbb{R}_+^p, \max_{k \in S} \|\boldsymbol{\mu}^{(k)} - \boldsymbol{\mu}^{(0)}\|_2 \leq h/2, \|\boldsymbol{\mu}^{(k)}\|_2 \leq M\}$. For any $\boldsymbol{\mu} \in \mathbb{R}$, denote distribution $\frac{1}{2}\mathcal{N}(\boldsymbol{\mu}, \mathbf{I}_p) + \frac{1}{2}\mathcal{N}(-\boldsymbol{\mu}, \mathbf{I}_p)$ as $\mathbb{P}_{\boldsymbol{\mu}}$, and denote $\prod_{k \in S} \mathbb{P}_{\boldsymbol{\mu}^{(k)}}^{\otimes n_k}$ as $\mathbb{P}_{\{\boldsymbol{\mu}^{(k)}\}_{k \in S}}$. It suffices to show

$$\inf_{\hat{\boldsymbol{\mu}}^{(0)}} \sup_{S: |S| \geq s} \sup_{\{\bar{\boldsymbol{\mu}}^{(k)*}\}_{k \in \{0\} \cup S} \in \bar{\Theta}_S} \mathbb{P} \left(\|\hat{\boldsymbol{\mu}}^{(0)} - \boldsymbol{\mu}^{(0)*}\|_2 \geq \left(C_1 \tilde{\epsilon}' \sqrt{\frac{1}{\max_{k=1:K} n_k}} \right) \wedge \left(C_2 \sqrt{\frac{1}{n_0}} \right) \right) \geq \frac{1}{10}. \quad (\text{E.69})$$

where $\mathbb{P} = \mathbb{P}_{\boldsymbol{\mu}^{(0)}}^{\otimes n_0} \cdot \mathbb{P}_{\{\boldsymbol{\mu}^{(k)}\}_{k \in S}} \cdot \mathbb{Q}_S$.

For any $\boldsymbol{\mu} \in \mathbb{R}$, denote distribution $\frac{1}{2}\mathcal{N}(\boldsymbol{\mu}, \mathbf{I}_p) + \frac{1}{2}\mathcal{N}(-\boldsymbol{\mu}, \mathbf{I}_p)$ as $\mathbb{P}_{\boldsymbol{\mu}}$. WLOG, assume $M \geq 1$. For any $\tilde{\boldsymbol{\mu}}_1, \tilde{\boldsymbol{\mu}}_2 \in \mathbb{R}^p$ with $\|\tilde{\boldsymbol{\mu}}_1\|_2 = \|\tilde{\boldsymbol{\mu}}_2\|_2 = 1$, by Lemma 15,

$$\max_{k=1:K} \text{KL}(\mathbb{P}_{\tilde{\boldsymbol{\mu}}_1}^{\otimes n_k} \|\mathbb{P}_{\tilde{\boldsymbol{\mu}}_2}^{\otimes n_k}) \leq \max_{k=1:K} n_k \cdot 8\|\tilde{\boldsymbol{\mu}}_1 - \tilde{\boldsymbol{\mu}}_2\|_2^2.$$

for any $k = 1 : K$. Let $8 \max_{k=1:K} n_k \cdot \|\tilde{\boldsymbol{\mu}}_1 - \tilde{\boldsymbol{\mu}}_2\|_2^2 \leq (\frac{\tilde{\epsilon}'}{1-\tilde{\epsilon}'})^2$, then $\|\tilde{\boldsymbol{\mu}}_1 - \tilde{\boldsymbol{\mu}}_2\|_2 \leq C \sqrt{\frac{1}{\max_{k=1:K} n_k}} \tilde{\epsilon}'$ for some constant $C > 0$. On the other hand, let $\text{KL}(\mathbb{P}_{\tilde{\boldsymbol{\mu}}_1}^{\otimes n_0} \|\mathbb{P}_{\tilde{\boldsymbol{\mu}}_2}^{\otimes n_0}) = 8n_0 \cdot \|\tilde{\boldsymbol{\mu}}_1 - \tilde{\boldsymbol{\mu}}_2\|_2^2 \leq 1/100$, then $\|\tilde{\boldsymbol{\mu}}_1 - \tilde{\boldsymbol{\mu}}_2\|_2 \leq \sqrt{\frac{1}{800}} \sqrt{\frac{1}{n_0}}$ for some constant $C > 0$. Then (E.69) follows by Lemma 33. \square

Proof of Lemma 32. The proof is similar to the proof of Theorem 5.1 in Chen et al. (2018), so we omit it here. \square

Proof of Lemma 34. This can be similarly shown by Assouad's Lemma as in the proof of Lemma 23. We omit the proof here. \square

E.10 Proof of Theorem 12

The result follows from (E.48) and Theorem 11.

E.11 Proof of Theorem 14

E.11.1 Lemmas

Lemma 35. Assume $n_0 \geq Cp$ and $\Delta^{(0)} \geq C' > 0$ with some constants $C, C' > 0$. We have

$$\inf_{\hat{\mathcal{C}}^{(0)}} \sup_{S: |S| \geq s} \sup_{\{\tilde{\boldsymbol{\theta}}^{(k)*}\}_{k \in \{0\} \cup S} \in \bar{\Theta}'_S \atop \mathbb{Q}_S} \mathbb{P} \left(R_{\tilde{\boldsymbol{\theta}}^{(0)*}}(\hat{\mathcal{C}}^{(0)}) - R_{\tilde{\boldsymbol{\theta}}^{(0)*}}(\mathcal{C}_{\tilde{\boldsymbol{\theta}}^{(0)*}}) \geq C_1 \frac{p}{n_S + n_0} + C_2 h^2 \wedge \frac{p}{n_0} + \frac{1}{n_0} \right) \geq \frac{1}{10}.$$

Lemma 36. Denote $\tilde{\epsilon} = \frac{K-s}{s}$. We have

$$\inf_{\hat{\mathcal{C}}^{(0)}} \sup_{S: |S| \geq s} \sup_{\{\tilde{\boldsymbol{\theta}}^{(k)*}\}_{k \in \{0\} \cup S} \in \bar{\Theta}_S^{(T)} \atop \mathbb{Q}_S} \mathbb{P} \left(R_{\tilde{\boldsymbol{\theta}}^{(0)*}}(\hat{\mathcal{C}}^{(0)}) - R_{\tilde{\boldsymbol{\theta}}^{(0)*}}(\mathcal{C}_{\tilde{\boldsymbol{\theta}}^{(0)*}}) \geq \left(C_1 \frac{\tilde{\epsilon}^2}{\max_{k=1:K} n_k} \right) \wedge \left(C_2 \sqrt{\frac{1}{n_0}} \right) \right) \geq \frac{1}{10}.$$

E.11.2 Main proof of Theorem 14

Combine Lemmas 35 and 36 to finish the proof.

E.11.3 Proof of lemmas

Proof of Lemma 35. We proceed with similar proof ideas used in the proof of Lemma 35. Recall the definitions and proof idea of Lemma 30. We have $\bar{\Theta}'_S \supseteq \bar{\Theta}'_{S,w} \cup \bar{\Theta}'_{S,\beta} \cup \bar{\Theta}'_{S,\delta}$, where

$$\begin{aligned}\bar{\Theta}'_{S,w} &= \left\{ \{\bar{\boldsymbol{\theta}}^{(k)}\}_{k \in \{0\} \cup S} : \boldsymbol{\mu}_1^{(k)} = \mathbf{1}_p / \sqrt{p}, \boldsymbol{\mu}_2^{(k)} = -\boldsymbol{\mu}_1^{(k)}, \boldsymbol{\Sigma}^{(k)} = \mathbf{I}_p, w^{(k)} \in (c_w, 1 - c_w) \right\}, \\ \bar{\Theta}'_{S,\beta} &= \left\{ \{\bar{\boldsymbol{\theta}}^{(k)}\}_{k \in \{0\} \cup S} : \boldsymbol{\Sigma}^{(k)} = \mathbf{I}_p, w^{(k)} = \frac{1}{2}, \|\boldsymbol{\mu}_1^{(k)}\|_2 \vee \|\boldsymbol{\mu}_2^{(k)}\|_2 \leq M, \max_{k \in S} \|\boldsymbol{\beta}^{(k)} - \boldsymbol{\beta}^{(0)}\|_2 \leq h \right\}, \\ \bar{\Theta}'_{S,\delta} &= \left\{ \{\bar{\boldsymbol{\theta}}^{(k)}\}_{k \in \{0\} \cup S} : \boldsymbol{\Sigma}^{(k)} = \mathbf{I}_p, w^{(k)} = \frac{1}{2}, \|\boldsymbol{\mu}_1^{(k)}\|_2 \vee \|\boldsymbol{\mu}_2^{(k)}\|_2 \leq M, \boldsymbol{\mu}_1^{(k)} = -\frac{1}{2}\boldsymbol{\mu}_0, \right. \\ &\quad \left. \boldsymbol{\mu}_2^{(k)} = \frac{1}{2}\boldsymbol{\mu}_0 + \mathbf{u}, \|\mathbf{u}\|_2 \leq 1 \right\}.\end{aligned}$$

Recall the mis-clustering error for GMM associated with parameter set $\bar{\boldsymbol{\theta}}$ of any classifier \mathcal{C} is $R_{\bar{\boldsymbol{\theta}}}(\mathcal{C}) = \min_{\pi: \{1,2\} \rightarrow \{1,2\}} \mathbb{P}_{\bar{\boldsymbol{\theta}}}(\mathcal{C}(Z) \neq \pi(Y))$. To help the analysis, following Azizyan et al. (2013) and Cai et al. (2019), we define a surrogate loss $L_{\bar{\boldsymbol{\theta}}}(\mathcal{C}) = \min_{\pi: \{1,2\} \rightarrow \{1,2\}} \mathbb{P}_{\bar{\boldsymbol{\theta}}}(\mathcal{C}(Z) \neq \pi(\mathcal{C}_{\bar{\boldsymbol{\theta}}}(Z)))$, where $\mathcal{C}_{\bar{\boldsymbol{\theta}}}$ is the Bayes classifier. Suppose $\sigma = \sqrt{0.005}$.

(i) We want to show

$$\inf_{\hat{\mathcal{C}}^{(0)}} \sup_{S: |S| \geq s} \sup_{\substack{\{\bar{\boldsymbol{\theta}}^{(k)*}\}_{k \in \{0\} \cup S} \in \bar{\Theta}'_S \\ \mathbb{Q}_S}} \mathbb{P} \left(R_{\bar{\boldsymbol{\theta}}^{(0)*}}(\hat{\mathcal{C}}^{(0)}) - R_{\bar{\boldsymbol{\theta}}^{(0)*}}(\mathcal{C}_{\bar{\boldsymbol{\theta}}^{(0)*}}) \geq C \sqrt{\frac{p}{n_S + n_0}} \right) \geq \frac{1}{4}. \quad (\text{E.70})$$

Consider $S = 1 : K$ and space $\bar{\Theta}'_0 = \{\{\bar{\boldsymbol{\theta}}^{(k)}\}_{k=0}^K : \boldsymbol{\Sigma}^{(k)} = \mathbf{I}_p, w^{(k)} = 1/2, \boldsymbol{\mu}_1^{(k)} = \boldsymbol{\mu}_1, \boldsymbol{\mu}_2^{(k)} = \boldsymbol{\mu}_2, \|\boldsymbol{\mu}_1\|_2 \vee \|\boldsymbol{\mu}_2\|_2 \leq M\}$. And

$$\text{LHS of (E.50)} \geq \inf_{\hat{\mathcal{C}}^{(0)}} \sup_{\{\bar{\boldsymbol{\theta}}^{(k)*}\}_{k=0}^K \in \bar{\Theta}'_0} \mathbb{P} \left(R_{\bar{\boldsymbol{\theta}}^{(0)*}}(\hat{\mathcal{C}}^{(0)}) - R_{\bar{\boldsymbol{\theta}}^{(0)*}}(\mathcal{C}_{\bar{\boldsymbol{\theta}}^{(0)*}}) \geq C \sqrt{\frac{p}{n_S + n_0}} \right).$$

Let $r = c\sqrt{p/(n_S + n_0)} \leq 0.001$ with some small constant $c > 0$. For any $\boldsymbol{\mu} \in \mathbb{R}^p$, denote distribution $\frac{1}{2}\mathcal{N}(\boldsymbol{\mu}, \mathbf{I}_p) + \frac{1}{2}\mathcal{N}(-\boldsymbol{\mu}, \mathbf{I}_p)$ as $\mathbb{P}_{\boldsymbol{\mu}}$. Consider a $r/4$ -packing of $r\mathcal{S}^{p-1}$: $\{\tilde{\mathbf{v}}_j\}_{j=1}^N$. By Lemma 2, $N \geq 4^{p-1}$. Denote $\tilde{\boldsymbol{\mu}}_j = (\sigma, \tilde{\mathbf{v}}_j^T)^T \in \mathbb{R}^p$, where $\sigma = \sqrt{0.005}$. Then by definition of KL divergence and Lemma 8.4 in Cai et al. (2019),

$$\begin{aligned}\text{KL} \left(\prod_{k \in \{0\} \cup S} \mathbb{P}_{\tilde{\boldsymbol{\mu}}_j}^{\otimes n_k} \cdot \mathbb{Q}_S \left\| \prod_{k \in \{0\} \cup S} \mathbb{P}_{\tilde{\boldsymbol{\mu}}_{j'}}^{\otimes n_k} \cdot \mathbb{Q}_S \right. \right) &= \sum_{k \in \{0\} \cup S} n_k \text{KL}(\mathbb{P}_{\tilde{\boldsymbol{\mu}}_j} \| \mathbb{P}_{\tilde{\boldsymbol{\mu}}_{j'}}) \\ &\leq (n_S + n_0) \cdot 8(1 + \sigma^2) \|\tilde{\boldsymbol{\mu}}_j - \tilde{\boldsymbol{\mu}}_{j'}\|_2^2 \\ &\leq 32(1 + \sigma^2)(n_S + n_0)r^2 \\ &\leq 32(1 + \sigma^2)(n_S + n_0) \cdot c^2 \frac{2(p-1)}{n_S + n_0}\end{aligned}$$

$$\leq \frac{32(1 + \sigma^2)c^2}{\log 2} \log N.$$

For simplicity, we write $L_{\bar{\theta}}$ with $\bar{\theta} \in \bar{\Theta}_0$ and $\mu_1 = -\mu_2 = \mu$ as L_{μ} . By Lemma 8.5 in [Cai et al. \(2019\)](#),

$$L_{\tilde{\mu}_i}(\mathcal{C}_{\tilde{\mu}_j}) \geq \frac{1}{\sqrt{2}} g\left(\frac{\sqrt{\sigma^2 + r^2}}{2}\right) \frac{\|\tilde{\mu}_i - \tilde{\mu}_j\|_2}{\|\tilde{\mu}_i\|_2} \geq \frac{1}{\sqrt{2}} \cdot 0.15 \cdot \frac{r/4}{\sqrt{\sigma^2 + r^2}} \geq 2r,$$

where $g(x) = \phi(x)[\phi(x) - x\Phi(x)]$. The last inequality holds because $\sqrt{\sigma^2 + r^2} \geq \sqrt{2}\sigma$ and $g(\sqrt{\sigma^2 + r^2}/2) \geq 0.15$ when $r^2 \leq \sigma^2 = 0.001$. Then by Lemma 3.5 in [Cai et al. \(2019\)](#) (Proposition 2 in [Azizyan et al. \(2013\)](#)), for any classifier \mathcal{C} , and $i \neq j$,

$$L_{\tilde{\mu}_i}(\mathcal{C}) + L_{\tilde{\mu}_j}(\mathcal{C}) \geq L_{\tilde{\mu}_i}(\mathcal{C}_{\tilde{\mu}_j}) - \sqrt{\text{KL}(\mathbb{P}_{\tilde{\mu}_i} \|\mathbb{P}_{\tilde{\mu}_j})/2} \geq 2r - r = c\sqrt{\frac{p}{n_S + n_0}}. \quad (\text{E.71})$$

For any $\hat{\mathcal{C}}^{(0)}$, consider a test $\psi^* = \arg \min_{j=1:N} L_{\tilde{\mu}_j}(\hat{\mathcal{C}}^{(0)})$. Therefore if there exists j_0 such that $L_{\tilde{\mu}_{j_0}}(\hat{\mathcal{C}}^{(0)}) < \frac{c}{2} \sqrt{\frac{p}{n_S + n_0}}$, then by (E.71), we must have $\psi^* = j_0$. Let $C_1 \leq c/2$, then by Fano's lemma (Corollary 6 in [Tsybakov \(2009\)](#))

$$\begin{aligned} \inf_{\hat{\mathcal{C}}^{(0)}} \sup_{\{\bar{\theta}^{(k)*}\}_{k=0}^K \in \bar{\Theta}'_0} \mathbb{P}\left(L_{\bar{\theta}^{(0)*}}(\hat{\mathcal{C}}^{(0)}) \geq C_1 \sqrt{\frac{p}{n_S + n_0}}\right) &\geq \inf_{\hat{\mathcal{C}}^{(0)}} \sup_{j=1:N} \mathbb{P}\left(L_{\tilde{\mu}^{(j)}}(\hat{\mathcal{C}}^{(0)}) \geq C_1 \sqrt{\frac{p}{n_S + n_0}}\right) \\ &\geq \inf_{\hat{\mathcal{C}}^{(0)}} \sup_{j=1:N} \mathbb{P}\left(\psi^* \neq j\right) \\ &\geq \inf_{\psi} \sup_{j=1:N} \mathbb{P}\left(\psi \neq j\right) \\ &\geq 1 - \frac{\log 2}{\log N} - \frac{32(1 + \sigma^2)c^2}{\log 2} \\ &\geq \frac{1}{4}, \end{aligned}$$

when $p \geq 2$ and $c = \sqrt{\frac{\log 2}{128(1 + \sigma^2)}}$. Then apply Lemma 25 to get the (E.70).

(ii) We want to show

$$\inf_{\hat{\mathcal{C}}^{(0)}} \sup_{S: |S| \geq s} \sup_{\{\bar{\theta}^{(k)*}\}_{k \in \{0\} \cup S} \in \bar{\Theta}'_{S, \beta}} \mathbb{P}\left(R_{\bar{\theta}^{(0)*}}(\hat{\mathcal{C}}^{(0)}) - R_{\bar{\theta}^{(0)*}}(\mathcal{C}_{\bar{\theta}^{(0)*}}) \geq C \left(h \wedge \sqrt{\frac{p}{n_0}}\right)\right) \geq \frac{1}{4}. \quad (\text{E.72})$$

Fixing an S and a \mathbb{Q}_S . Suppose $1 \in S$. We have

$$\text{LHS of (E.72)} \geq \inf_{\widehat{\mathcal{C}}^{(0)}} \sup_{\{\bar{\boldsymbol{\theta}}^{(k)*}\}_{k \in \{0\} \cup S} \in \bar{\Theta}'_{S, \beta}} \mathbb{P} \left(R_{\bar{\boldsymbol{\theta}}^{(0)*}}(\widehat{\mathcal{C}}^{(0)}) - R_{\bar{\boldsymbol{\theta}}^{(0)*}}(\mathcal{C}_{\bar{\boldsymbol{\theta}}^{(0)*}}) \geq C \left(h \wedge \sqrt{\frac{p}{n_0}} \right) \right). \quad (\text{E.73})$$

Let $r = h \wedge (c\sqrt{p/n_0}) \wedge M$ with a small constant $c > 0$. For any $\boldsymbol{\mu} \in \mathbb{R}^p$, denote distribution $\frac{1}{2}\mathcal{N}(\boldsymbol{\mu}, \mathbf{I}_p) + \frac{1}{2}\mathcal{N}(-\boldsymbol{\mu}, \mathbf{I}_p)$ as $\mathbb{P}_{\boldsymbol{\mu}}$. Consider a $r/4$ -packing of $r\mathcal{S}^{p-1}$. By Lemma 2, $N \geq 4^{p-1}$. Denote $\tilde{\boldsymbol{\mu}}_j = (\sigma, \tilde{\mathbf{v}}_j^T)^T \in \mathbb{R}^p$. WLOG, assume $M \geq 2$. Let $\boldsymbol{\mu}_1^{(k)*} = \tilde{\boldsymbol{\mu}} = (\sigma, \mathbf{0}_{p-1})^T$ for all $k \in S$ and $\boldsymbol{\mu}_1^{(0)*} = \boldsymbol{\mu} = (1, \boldsymbol{\vartheta})$ with $\boldsymbol{\vartheta} \in (r\mathcal{S}^{p-1}) \cap \mathcal{Q}_v$. Then by following the same arguments in part (ii) of the proof of Lemma 30, we can show that the RHS of (E.73) is larger than or equal to $1/4$ when $p \geq 3$.

(iii) We want to show

$$\inf_{\widehat{\mathcal{C}}^{(0)}} \sup_{S: |S| \geq s} \sup_{\{\bar{\boldsymbol{\theta}}^{(k)*}\}_{k \in \{0\} \cup S} \in \bar{\Theta}'_S} \mathbb{P} \left(R_{\bar{\boldsymbol{\theta}}^{(0)*}}(\widehat{\mathcal{C}}^{(0)}) - R_{\bar{\boldsymbol{\theta}}^{(0)*}}(\mathcal{C}_{\bar{\boldsymbol{\theta}}^{(0)*}}) \geq Ch_w^2 \wedge \frac{1}{n_0} \right) \geq \frac{1}{4}.$$

This can be similarly proved by following the arguments in part (i) with Lemmas 25 and 26.

(iv) We want to show

$$\inf_{\widehat{\mathcal{C}}^{(0)}} \sup_{S: |S| \geq s} \sup_{\{\bar{\boldsymbol{\theta}}^{(k)*}\}_{k \in \{0\} \cup S} \in \bar{\Theta}'_S} \mathbb{P} \left(R_{\bar{\boldsymbol{\theta}}^{(0)*}}(\widehat{\mathcal{C}}^{(0)}) - R_{\bar{\boldsymbol{\theta}}^{(k)*}}(\mathcal{C}_{\bar{\boldsymbol{\theta}}^{(0)*}}) \geq Ch_{\beta}^2 \wedge \frac{p}{n_0} \right) \geq \frac{1}{4}.$$

The conclusion can be obtained immediately from (ii), by noticing that $\bar{\Theta}'_{S, \beta} \supseteq \bar{\Theta}'_{S, \boldsymbol{\mu}}$. Finally, we get the desired conclusion by combining (i)-(iv). \square

Proof of Lemma 36. By Lemma 25, it suffices to prove

$$\inf_{\widehat{\mathcal{C}}^{(0)}} \sup_{S: |S| \geq s} \sup_{\{\bar{\boldsymbol{\theta}}^{(k)*}\}_{k \in \{0\} \cup S} \in \bar{\Theta}'_S} \mathbb{P} \left(L_{\bar{\boldsymbol{\theta}}^{(0)*}}(\widehat{\mathcal{C}}^{(0)}) \geq C_1 \frac{\tilde{\epsilon}^2}{\max_{k=1:K} n_k} \wedge \frac{1}{n_0} \right) \geq \frac{1}{10}.$$

For any $\boldsymbol{\mu} \in \mathbb{R}^p$, denote distribution $\frac{1}{2}\mathcal{N}(\boldsymbol{\mu}, \mathbf{I}_p) + \frac{1}{2}\mathcal{N}(-\boldsymbol{\mu}, \mathbf{I}_p)$ as $\mathbb{P}_{\boldsymbol{\mu}}$. For simplicity, we write $L_{\bar{\boldsymbol{\theta}}}$ with $\bar{\boldsymbol{\theta}}$ satisfying $\boldsymbol{\mu}_1 = -\boldsymbol{\mu}_2 = \boldsymbol{\mu}$, $w = 1/2$ and $\boldsymbol{\Sigma} = \mathbf{I}_p$ as $L_{\boldsymbol{\mu}}$. Consider $L_{\boldsymbol{\mu}}(\mathcal{C}_{\boldsymbol{\mu}'})$ as a loss function between $\boldsymbol{\mu}$ and $\boldsymbol{\mu}'$ in Lemmas 32 and 33. Considering $\|\boldsymbol{\mu}\|_2 = \|\boldsymbol{\mu}'\|_2 = 1$, by Lemma 15, note that

$$\begin{aligned} \max_{k=1:K} \text{KL}(\mathbb{P}_{\boldsymbol{\mu}}^{\otimes n_k} \parallel \mathbb{P}_{\boldsymbol{\mu}'}^{\otimes n_k}) &\leq 8 \max_{k=1:K} n_k \cdot \|\boldsymbol{\mu} - \boldsymbol{\mu}'\|_2^2, \\ \text{KL}(\mathbb{P}_{\boldsymbol{\mu}}^{\otimes n_0} \parallel \mathbb{P}_{\boldsymbol{\mu}'}^{\otimes n_0}) &\leq 8n_0 \cdot \|\boldsymbol{\mu} - \boldsymbol{\mu}'\|_2^2. \end{aligned}$$

By Lemma 8.5 in [Cai et al. \(2019\)](#), this implies for some constants $c, C > 0$

$$\begin{aligned}
& \sup \left\{ L_{\boldsymbol{\mu}}(\mathcal{C}_{\boldsymbol{\mu}'}) : \max_{k=1:K} \text{KL}(\mathbb{P}_{\boldsymbol{\mu}}^{\otimes n_k} \| \mathbb{P}_{\boldsymbol{\mu}'}^{\otimes n_k}) \leq (\tilde{\epsilon}'/(1-\tilde{\epsilon}))^2, \text{KL}(\mathbb{P}_{\boldsymbol{\mu}}^{\otimes n_0} \| \mathbb{P}_{\boldsymbol{\mu}'}^{\otimes n_0}) \leq 1/100 \right\} \\
& \geq \sup \left\{ c \|\boldsymbol{\mu} - \boldsymbol{\mu}'\|_2 : \max_{k=1:K} \text{KL}(\mathbb{P}_{\boldsymbol{\mu}}^{\otimes n_k} \| \mathbb{P}_{\boldsymbol{\mu}'}^{\otimes n_k}) \leq (\tilde{\epsilon}'/(1-\tilde{\epsilon}))^2, \text{KL}(\mathbb{P}_{\boldsymbol{\mu}}^{\otimes n_0} \| \mathbb{P}_{\boldsymbol{\mu}'}^{\otimes n_0}) \leq 1/100 \right\} \\
& \geq \sup \left\{ c \|\boldsymbol{\mu} - \boldsymbol{\mu}'\|_2 : 8 \max_{k=1:K} n_k \cdot \|\boldsymbol{\mu} - \boldsymbol{\mu}'\|_2^2 \leq (\tilde{\epsilon}'/(1-\tilde{\epsilon}))^2, 8n_0 \cdot \|\boldsymbol{\mu} - \boldsymbol{\mu}'\|_2^2 \leq 1/800 \right\} \\
& = C \cdot \frac{\tilde{\epsilon}'}{\sqrt{\max_{k=1:K} n_k}} \wedge \sqrt{\frac{1}{n_0}}.
\end{aligned}$$

Then apply Lemmas 32 and 33 to get the desired bound. \square

E.12 Proof of Theorem 15

Denote $\xi = \max_{k \in \{0\} \cup S} \min_{r_k = \pm 1} \|r_k \widehat{\boldsymbol{\beta}}^{(k)[0]} - \boldsymbol{\beta}^{(k)*}\|_2 = \max_{k \in \{0\} \cup S} (\|\widehat{\boldsymbol{\beta}}^{(k)[0]} - \boldsymbol{\beta}^{(k)*}\|_2 \wedge \|\widehat{\boldsymbol{\beta}}^{(k)[0]} + \boldsymbol{\beta}^{(k)*}\|_2)$. WLOG, assume $S = \{1, \dots, s\}$ and $r_k^* = 1$ for all $k \in \{0\} \cup S$. Hence $\xi = \max_{k \in \{0\} \cup S} \|\widehat{\boldsymbol{\beta}}^{(k)[0]} - \boldsymbol{\beta}^{(k)*}\|_2$. WLOG, consider $\widehat{r}_k = 1$ for all $k \in S$ (i.e., the tasks in S are already well-aligned). Consider

$$\begin{aligned}
(1, \widehat{\mathbf{r}}) &= (\underbrace{1}_{\text{target}}, \underbrace{1, \dots, 1}_S, \underbrace{r_{s+1}, \dots, r_K}_{\text{outlier tasks}}), \\
(-1, \widehat{\mathbf{r}}) &= (\underbrace{-1}_{\text{target}}, \underbrace{1, \dots, 1}_S, \underbrace{r_{s+1}, \dots, r_K}_{\text{outlier tasks}}).
\end{aligned}$$

It suffices to prove that

$$\text{score}((-1, \widehat{\mathbf{r}})) - \text{score}((1, \widehat{\mathbf{r}})) > 0.$$

In fact,

$$\begin{aligned}
\text{score}((-1, \widehat{\mathbf{r}})) - \text{score}((1, \widehat{\mathbf{r}})) &= 2 \underbrace{\sum_{k=1}^s \|\widehat{\boldsymbol{\beta}}^{(0)[0]} + \widehat{\boldsymbol{\beta}}^{(k)[0]}\|_2}_{[1]} + 2 \underbrace{\sum_{k=s+1}^K \|\widehat{\boldsymbol{\beta}}^{(0)[0]} + r_k \widehat{\boldsymbol{\beta}}^{(k)[0]}\|_2}_{[2]} \\
&\quad - 2 \underbrace{\sum_{k=1}^s \|\widehat{\boldsymbol{\beta}}^{(0)[0]} - \widehat{\boldsymbol{\beta}}^{(k)[0]}\|_2}_{[1]'} - 2 \underbrace{\sum_{k=s+1}^K \|\widehat{\boldsymbol{\beta}}^{(0)[0]} + r_k \widehat{\boldsymbol{\beta}}^{(k)[0]}\|_2}_{[2]},
\end{aligned}$$

where

$$[1] - [1]' \geq \sum_{k=1}^s (\|\boldsymbol{\beta}^{(0)*} + \boldsymbol{\beta}^{(k)*}\|_2 - \|\boldsymbol{\beta}^{(0)*} - \boldsymbol{\beta}^{(k)*}\|_2 - 4\xi)$$

$$\begin{aligned}
&\geq \sum_{k=1}^s (2\|\boldsymbol{\beta}^{(0)*}\|_2 - 2\|\boldsymbol{\beta}^{(0)*} - \boldsymbol{\beta}^{(k)*}\|_2 - 4\xi) \\
&\geq s(2\|\boldsymbol{\beta}^{(0)*}\|_2 - 4h - 4\xi),
\end{aligned}$$

and

$$[2] - [2]' \geq -4 \sum_{k=s+1}^K \|\widehat{\boldsymbol{\beta}}^{(0)[0]}\|_2 \geq -4(K-s)(\|\boldsymbol{\beta}^{(0)*}\|_2 + \xi).$$

Hence

$$\begin{aligned}
\text{score}((-1, \widehat{\mathbf{r}})) - \text{score}((1, \widehat{\mathbf{r}})) &= 2([1] - [1]') + 2([2] - [2]') \\
&\geq 4[(2s - K)\|\boldsymbol{\beta}^{(0)*}\|_2 - 2sh - (K + s)\xi] \\
&> 0,
\end{aligned}$$

when $\|\boldsymbol{\beta}^{(0)*}\|_2 > \frac{2(1-\epsilon)}{1-2\epsilon}h + \frac{2-\epsilon}{1-2\epsilon}\xi$, which completes our proof.

E.13 Proof of Theorem 7

Define the contraction basin of one GMM as

$$\begin{aligned}
B_{\text{con}}(\boldsymbol{\theta}^{(k)*}) &= \{\boldsymbol{\theta} = \{\{w_r\}_{r=2}^R, \{\boldsymbol{\beta}_r\}_{r=2}^R, \{\delta_r\}_{r=2}^R\} : w_r^* \in (c_w/2, 1 - c_w/2), \\
&\quad \|\boldsymbol{\beta}_r - \boldsymbol{\beta}_r^*\|_2 \leq C_b\Delta, |\delta_r - \delta_r^*| \leq C_b\Delta\}.
\end{aligned}$$

And the joint contraction basin is defined as $B_{\text{con}}(\{\boldsymbol{\theta}^{(k)*}\}_{k \in S}) = \bigcap_{r=1}^R B_{\text{con}}(\boldsymbol{\theta}^{(k)*})$.

For $\boldsymbol{\theta} = (\{w_r\}_{r=2}^R, \{\boldsymbol{\beta}_r\}_{r=2}^R, \{\delta_r\}_{r=2}^R)$ and $\boldsymbol{\theta}' = (\{w'_r\}_{r=2}^R, \{\boldsymbol{\beta}'_r\}_{r=2}^R, \{\delta'_r\}_{r=2}^R)$, define

$$d(\boldsymbol{\theta}, \boldsymbol{\theta}') = \max_{r=2:R} \{|w_r - w'_r| \vee \|\boldsymbol{\beta}_r - \boldsymbol{\beta}'_r\|_2 \vee |\delta_r - \delta'_r|\}.$$

E.13.1 Lemmas

For GMM $\mathbf{z} \sim \sum_{r=1}^R w_r^* \mathcal{N}(\boldsymbol{\mu}_r^*, \boldsymbol{\Sigma}^*)$ and any $\boldsymbol{\theta}$, define

$$\gamma_{\boldsymbol{\theta}}^{(r)}(\mathbf{z}) = \frac{w_r \exp\{\boldsymbol{\beta}_r^T \mathbf{z} - \delta_r\}}{w_1 + \sum_{r=2}^R w_r \exp\{\boldsymbol{\beta}_r^T \mathbf{z} - \delta_r\}}, \quad r = 2 : R, \quad \gamma_{\boldsymbol{\theta}}^{(1)}(\mathbf{z}) = \frac{w_1}{w_1 + \sum_{r=2}^R w_r \exp\{\boldsymbol{\beta}_r^T \mathbf{z} - \delta_r\}}.$$

Denote $w_r(\boldsymbol{\theta}) = \mathbb{E}[\gamma_{\boldsymbol{\theta}}^{(r)}(\mathbf{z})]$ and $\boldsymbol{\mu}_r(\boldsymbol{\theta}) = \frac{\mathbb{E}[\gamma_{\boldsymbol{\theta}}^{(r)}(\mathbf{z})\mathbf{z}]}{\mathbb{E}[\gamma_{\boldsymbol{\theta}}^{(r)}(\mathbf{z})]}$.

Lemma 37 (Contraction of multi-cluster GMM). *When $C_b \leq cc_{\boldsymbol{\Sigma}}^{-1/2}$ with a small constant $c > 0$ and $\Delta \geq C \log(cc_{\boldsymbol{\Sigma}} M c_w^{-1})$ with a large constant $C > 0$, there exist positive constants $C' > 0$ and $C'' > 0$, for any $\boldsymbol{\theta} \in B_{\text{con}}(\boldsymbol{\theta}^{(k)*})$,*

$$|w_r(\boldsymbol{\theta}) - w_r^*| \leq C' \exp\{-C''\Delta^2\} \cdot d(\boldsymbol{\theta}, \boldsymbol{\theta}^*), \quad \|\boldsymbol{\mu}_r(\boldsymbol{\theta}) - \boldsymbol{\mu}_r^*\|_2 \leq C' \exp\{-C''\Delta^2\} \cdot d(\boldsymbol{\theta}, \boldsymbol{\theta}^*),$$

where $C' \exp\{-C''\Delta^2\} \leq \kappa_0 < 1$ with a constant κ_0 .

Lemma 38 (Vectorized contraction of Rademacher complexity, Corollary 1 in [Maurer \(2016\)](#)). Suppose $\{\epsilon_{ir}\}_{i \in [n], r \in [R]}$ and $\{\epsilon_i\}_{i=1}^n$ are independent Rademacher variables. Let \mathcal{F} be a class of functions $f : \mathbb{R}^d \rightarrow \mathcal{S} \subseteq \mathbb{R}^R$ and $h : \mathcal{S} \rightarrow \mathbb{R}$ is L -Lipschitz under ℓ_2 -norm, i.e., $|h(\mathbf{y}) - h(\mathbf{y}')| \leq L\|\mathbf{y} - \mathbf{y}'\|_2$, where $\mathbf{y} = (y_1, \dots, y_R)^T$, $\mathbf{y}' = (y'_1, \dots, y'_R)^T \in \mathcal{S}$. Then

$$\mathbb{E} \sup_{f \in \mathcal{F}} \sum_{i=1}^n \epsilon_i h(f(x_i)) \leq \sqrt{2} L \mathbb{E} \sup_{f \in \mathcal{F}} \sum_{i=1}^n \sum_{r=1}^R \epsilon_{ir} f_r(x_i),$$

where $f_r(x_i)$ is the r -th component of $f(x_i) \in \mathcal{S} \subseteq \mathbb{R}^R$.

E.13.2 Main proof of Theorem 7

The proof idea is almost the same as the idea used in the proof of Theorem 1. We still need to show similar results presented in the lemmas associated with Theorem 1, then go through the same arguments in the proof of Theorem 7. We only sketch the key steps and the differences here.

The biggest difference appear in the proofs of the lemmas associated with Theorem 1 under the context of multi-cluster GMM. The original arguments in the proofs of Lemmas 11-14 involve the contraction inequality for Rademacher variables and univariate Lipschitz functions, which is not available any more. We replace this part by an argument through a vectorized Rademacher contraction inequality ([Maurer, 2016](#)).

First, we will show that

$$\sup_{\substack{\boldsymbol{\theta}^{(k)} \in B_{\text{con}} \\ \|\boldsymbol{\beta}_r^{(k)} - \boldsymbol{\beta}_r^{(k)*}\|_2 \leq \xi^{(k)}}} \left| \frac{1}{n_k} \sum_{i=1}^{n_k} \gamma_{\boldsymbol{\theta}^{(k)}}^{(r)}(\mathbf{z}_i^{(k)}) - \mathbb{E}[\gamma_{\boldsymbol{\theta}^{(k)}}^{(r)}(\mathbf{z}^{(k)})] \right| \lesssim \xi^{(k)} \sqrt{\frac{p}{n_k}} + \sqrt{\frac{\log K}{n_k}}, \quad (\text{E.74})$$

for all $k \in S$ and $r \in 1 : R$, with probability at least $1 - CK^{-2}$. Denote the LHS as W . By changing one observation $\mathbf{z}_i^{(k)}$, denote the new W as W' . Since $\gamma_{\boldsymbol{\theta}}^{(r)}(\mathbf{z})$ is bounded for all $\mathbf{z} \in \mathbb{R}^p$, we know that $|W - W'| \leq 1/n_k$. Then by bounded difference inequality, we have

$$W \leq \mathbb{E}W + C\sqrt{\frac{\log K}{n_k}},$$

with probability at least $1 - C'K^{-2}$. On the other hand, by symmetrization,

$$\mathbb{E}W \leq \frac{2}{n_k} \mathbb{E}_{\mathbf{z}} \mathbb{E}_{\epsilon} \sup_{\substack{\boldsymbol{\theta}^{(k)} \in B_{\text{con}} \\ \|\boldsymbol{\beta}_r^{(k)} - \boldsymbol{\beta}_r^{(k)*}\|_2 \leq \xi^{(k)}}} \left| \sum_{i=1}^{n_k} \epsilon_i^{(k)} \gamma_{\boldsymbol{\theta}^{(k)}}^{(r)}(\mathbf{z}_i^{(k)}) \right|.$$

Note that $\gamma_{\boldsymbol{\theta}^{(k)}}^{(r)}(\mathbf{z}) = \frac{w_r^{(k)} \cdot \exp\{(\boldsymbol{\beta}_r^{(k)})^T \mathbf{z} - \delta_r^{(k)}\}}{w_1^{(k)} + \sum_{r=2}^R w_r^{(k)} \exp\{(\boldsymbol{\beta}_r^{(k)})^T \mathbf{z} - \delta_r^{(k)}\}} = \frac{\exp\{(\boldsymbol{\beta}_r^{(k)})^T \mathbf{z} - \delta_r^{(k)} + \log w_r^{(k)} - \log w_1^{(k)}\}}{1 + \sum_{r=2}^R \exp\{(\boldsymbol{\beta}_r^{(k)})^T \mathbf{z} - \delta_r^{(k)} + \log w_r^{(k)} - \log w_1^{(k)}\}} = \varphi(\{(\boldsymbol{\beta}_r^{(k)})^T \mathbf{z} - \delta_r^{(k)} + \log w_r^{(k)} - \log w_1^{(k)}\}_{r=2}^R)$, where $\varphi(\mathbf{x}) = \frac{\exp\{x_r\}}{1 + \sum_{r=2}^R \exp\{x_r\}}$ is a 1-Lipschitz

function (w.r.t. ℓ_2 -norm). By Lemma 38,

$$\begin{aligned} \frac{2}{n_k} \mathbb{E}_{\mathbf{z}} \mathbb{E}_{\epsilon} \sup_{\substack{\boldsymbol{\theta}^{(k)} \in B_{\text{con}} \\ \|\boldsymbol{\beta}_r^{(k)} - \boldsymbol{\beta}_r^{(k)*}\|_2 \leq \xi^{(k)}}} \left| \sum_{i=1}^{n_k} \epsilon_i^{(k)} \gamma_{\boldsymbol{\theta}^{(k)}}^{(r)}(\mathbf{z}_i^{(k)}) \right| &\lesssim \frac{1}{n_k} \mathbb{E}_{\mathbf{z}} \mathbb{E}_{\epsilon} \sup_{\substack{\boldsymbol{\theta}^{(k)} \in B_{\text{con}} \\ \|\boldsymbol{\beta}_r^{(k)} - \boldsymbol{\beta}_r^{(k)*}\|_2 \leq \xi^{(k)}}} \left| \sum_{i=1}^{n_k} \sum_{r=2}^R \epsilon_{ir}^{(k)} g_{ir}^{(k)} \right| \\ &\lesssim \frac{1}{n_k} \sum_{r=2}^R \mathbb{E}_{\mathbf{z}} \mathbb{E}_{\epsilon} \sup_{\substack{\boldsymbol{\theta}^{(k)} \in B_{\text{con}} \\ \|\boldsymbol{\beta}_r^{(k)} - \boldsymbol{\beta}_r^{(k)*}\|_2 \leq \xi^{(k)}}} \left| \sum_{i=1}^{n_k} \epsilon_{ir}^{(k)} g_{ir}^{(k)} \right|, \end{aligned}$$

where $g_{ir}^{(k)} := (\boldsymbol{\beta}_r^{(k)})^T \mathbf{z}_i^{(k)} - \delta_r^{(k)} + \log w_r^{(k)} - \log w_1^{(k)}$. It follows that

$$\begin{aligned} &\frac{1}{n_k} \mathbb{E}_{\mathbf{z}} \mathbb{E}_{\epsilon} \sup_{\substack{\boldsymbol{\theta}^{(k)} \in B_{\text{con}} \\ \|\boldsymbol{\beta}_r^{(k)} - \boldsymbol{\beta}_r^{(k)*}\|_2 \leq \xi^{(k)}}} \left| \sum_{i=1}^{n_k} \epsilon_{ir}^{(k)} g_{ir}^{(k)} \right| \\ &\leq \frac{1}{n_k} \mathbb{E}_{\mathbf{z}, \epsilon} \sup_{\substack{\boldsymbol{\theta}^{(k)} \in B_{\text{con}} \\ \|\boldsymbol{\beta}_r^{(k)} - \boldsymbol{\beta}_r^{(k)*}\|_2 \leq \xi^{(k)}}} \left| \sum_{i=1}^{n_k} \epsilon_{ir}^{(k)} (\boldsymbol{\beta}_r^{(k)})^T \mathbf{z}_i^{(k)} \right| \\ &\quad + \frac{1}{n_k} \mathbb{E}_{\epsilon} \sup_{\substack{\boldsymbol{\theta}^{(k)} \in B_{\text{con}} \\ \|\boldsymbol{\beta}_r^{(k)} - \boldsymbol{\beta}_r^{(k)*}\|_2 \leq \xi^{(k)}}} \left| \sum_{i=1}^{n_k} \epsilon_{ir}^{(k)} (\delta_r^{(k)} - \log w_r^{(k)} + \log w_1^{(k)}) \right| \\ &\leq \frac{1}{n_k} \mathbb{E}_{\mathbf{z}, \epsilon} \sup_{\substack{\boldsymbol{\theta}^{(k)} \in B_{\text{con}} \\ \|\boldsymbol{\beta}_r^{(k)} - \boldsymbol{\beta}_r^{(k)*}\|_2 \leq \xi^{(k)}}} \left| \sum_{i=1}^{n_k} \epsilon_{ir}^{(k)} (\boldsymbol{\beta}_r^{(k)})^T (\mathbf{z}_i^{(k)} - \boldsymbol{\mu}^{(k)*}) \right| \\ &\quad + \frac{1}{n_k} \mathbb{E}_{\mathbf{z}, \epsilon} \sup_{\substack{\boldsymbol{\theta}^{(k)} \in B_{\text{con}} \\ \|\boldsymbol{\beta}_r^{(k)} - \boldsymbol{\beta}_r^{(k)*}\|_2 \leq \xi^{(k)}}} \left| \sum_{i=1}^{n_k} \epsilon_{ir}^{(k)} (\boldsymbol{\beta}_r^{(k)})^T \boldsymbol{\mu}^{(k)*} \right| \\ &\quad + \frac{1}{n_k} \mathbb{E}_{\epsilon} \sup_{\substack{|\delta_r^{(k)}| \leq U \\ c_w/2 \leq w_r^{(k)} \leq 1 - c_w/2}} \left| \sum_{i=1}^{n_k} \epsilon_{ir}^{(k)} (\delta_r^{(k)} - \log w_r^{(k)} + \log w_1^{(k)}) \right| \\ &\leq \frac{\xi^{(k)}}{n_k} \mathbb{E}_{\mathbf{z}, \epsilon} \sup_{j=1:N} \left| \sum_{i=1}^{n_k} \epsilon_{ir}^{(k)} \mathbf{u}_j^T (\mathbf{z}_i^{(k)} - \boldsymbol{\mu}^{(k)*}) \right| + \frac{1}{n_k} \mathbb{E}_{\mathbf{z}, \epsilon} \left| \sum_{i=1}^{n_k} \epsilon_{ir}^{(k)} (\boldsymbol{\beta}_r^{(k)*})^T (\mathbf{z}_i^{(k)} - \boldsymbol{\mu}^{(k)*}) \right| \\ &\quad + \frac{C}{n_k} \mathbb{E}_{\epsilon} \left| \sum_{i=1}^{n_k} \epsilon_{ir}^{(k)} \right| \tag{E.75} \end{aligned}$$

where $\boldsymbol{\mu}^{(k)*} := \sum_{r=1}^R w_r^{(k)*} \boldsymbol{\mu}_r^{(k)*}$, $\{\mathbf{u}_j\}_{j=1}^N$ is a $1/2$ -cover of \mathcal{S}^{d-1} with $N = 5^p$, and $\{\epsilon_{ir}^{(k)} \mathbf{u}_j^T (\mathbf{z}_i^{(k)} - \boldsymbol{\mu}^{(k)*})\}_{i=1}^{n_k}$, $\{\epsilon_{ir}^{(k)} (\boldsymbol{\beta}_r^{(k)*})^T (\mathbf{z}_i^{(k)} - \boldsymbol{\mu}^{(k)*})\}_{i=1}^{n_k}$, and $\{\epsilon_{ir}^{(k)}\}_{i=1}^{n_k}$ are all sub-Gaussian processes. Then by the property of sub-Gaussian variables,

$$\text{RHS of (E.75)} \lesssim \xi^{(k)} \sqrt{\frac{p}{n_k}} + \sqrt{\frac{\log K}{n_k}}.$$

Putting all the pieces together, we obtain $W \lesssim \xi^{(k)} \sqrt{\frac{p}{n_k}} + \sqrt{\frac{\log K}{n_k}}$ with probability at least $1 - CK^{-2}$.

The second bound we want to show is

$$\sup_{\{\boldsymbol{\theta}^{(k)}\}_{k \in S} \in B_{\text{con}}^{J,2}} \sup_{|\tilde{w}_k| \leq 1} \frac{1}{n_S} \left| \sum_{k \in S} \tilde{w}_k \sum_{i=1}^{n_k} \left[\gamma_{\boldsymbol{\theta}^{(k)}}^{(r)}(\mathbf{z}_i^{(k)}) - \mathbb{E}[\gamma_{\boldsymbol{\theta}^{(k)}}^{(r)}(\mathbf{z}^{(k)})] \right] \right| \lesssim \sqrt{\frac{p+K}{n_S}}.$$

Denote the LHS as W' . Again by bounded difference inequality,

$$W' \leq \mathbb{E}W' + C \sqrt{\frac{p}{n_S}},$$

with probability at least $1 - C' \exp\{-C''p\}$. It remains to control $\mathbb{E}W'$. By symmetrization,

$$\mathbb{E}W' \leq \frac{2}{n_S} \mathbb{E} \sup_{\{\boldsymbol{\theta}^{(k)}\}_{k \in S} \in B_{\text{con}}^{J,2}} \sup_{|\tilde{w}_k| \leq 1} \left| \sum_{k \in S} \sum_{i=1}^{n_k} \tilde{w}_k \gamma_{\boldsymbol{\theta}^{(k)}}^{(r)}(\mathbf{z}_i^{(k)}) \right|.$$

Denote

$$\begin{aligned} & \varphi(\tilde{w}, \{(\boldsymbol{\beta}_r^{(k)})^T \mathbf{z} - \delta_r^{(k)} + \log w_r^{(k)} - \log w_1^{(k)}\}_{r=2}^R) \\ &= \tilde{w}_k \gamma_{\boldsymbol{\theta}^{(k)}}^{(r)}(\mathbf{z}_i^{(k)}) \\ &= \tilde{w}_k \cdot \frac{\exp\{(\boldsymbol{\beta}_r^{(k)})^T \mathbf{z} - \delta_r^{(k)} + \log w_r^{(k)} - \log w_1^{(k)}\}}{1 + \sum_{r=2}^R \exp\{(\boldsymbol{\beta}_r^{(k)})^T \mathbf{z} - \delta_r^{(k)} + \log w_r^{(k)} - \log w_1^{(k)}\}}, \end{aligned}$$

which is C-Lipschitz w.r.t. $(\tilde{w}, \{(\boldsymbol{\beta}_r^{(k)})^T \mathbf{z} - \delta_r^{(k)} + \log w_r^{(k)} - \log w_1^{(k)}\}_{r=2}^R)$ as a R -dimensional vector with a constant C . Denote $g_{ir}^{(k)} = (\boldsymbol{\beta}_r^{(k)})^T \mathbf{z}_i^{(k)} - \delta_r^{(k)} + \log w_r^{(k)} - \log w_1^{(k)}$. A direct application of Lemma 38 implies that

$$\begin{aligned} & \frac{1}{n_S} \mathbb{E} \sup_{\{\boldsymbol{\theta}^{(k)}\}_{k \in S} \in B_{\text{con}}^{J,2}} \sup_{|\tilde{w}_k| \leq 1} \left| \sum_{k \in S} \sum_{i=1}^{n_k} \tilde{w}_k \gamma_{\boldsymbol{\theta}^{(k)}}^{(r)}(\mathbf{z}_i^{(k)}) \right| \\ & \lesssim \frac{1}{n_S} \mathbb{E} \sup_{|\tilde{w}_k| \leq 1} \left| \sum_{k \in S} \sum_{i=1}^{n_k} \tilde{w}_k \epsilon_{i1}^{(k)} \right| + \frac{1}{n_S} \sum_{r=2}^R \mathbb{E} \sup_{\{\boldsymbol{\theta}^{(k)}\}_{k \in S} \in B_{\text{con}}^{J,2}} \left| \sum_{k \in S} \sum_{i=1}^{n_k} g_{ir}^{(k)} \epsilon_{ir}^{(k)} \right|. \end{aligned}$$

By a similar argument involving covering number as before, we can show that

$$\frac{1}{n_S} \mathbb{E} \sup_{|\tilde{w}_k| \leq 1} \left| \sum_{k \in S} \sum_{i=1}^{n_k} \tilde{w}_k \epsilon_{i1}^{(k)} \right| + \frac{1}{n_S} \sum_{r=2}^R \mathbb{E} \sup_{\{\boldsymbol{\theta}^{(k)}\}_{k \in S} \in B_{\text{con}}^{J,2}} \left| \sum_{k \in S} \sum_{i=1}^{n_k} g_{ir}^{(k)} \epsilon_{ir}^{(k)} \right| \lesssim \sqrt{\frac{p+K}{n_S}}.$$

Therefore $W \lesssim \sqrt{\frac{p+K}{n_S}}$ with probability at least $1 - C' \exp\{-C''p\}$.

The third bound we want to show is

$$\begin{aligned} & \sup_{\substack{\boldsymbol{\theta}^{(k)} \in B_{\text{con}} \\ \|\boldsymbol{\beta}^{(k)} - \boldsymbol{\beta}^{(k)*}\|_2 \leq \xi^{(k)}}} \left| \frac{1}{n_k} \sum_{i=1}^{n_k} [1 - \gamma_{\boldsymbol{\theta}^{(k)}}^{(r)}(\mathbf{z}_i^{(k)})] (\mathbf{z}_i^{(k)})^T \boldsymbol{\beta}^{(k)*} - \mathbb{E}[1 - \gamma_{\boldsymbol{\theta}^{(k)}}^{(r)}(\mathbf{z}^{(k)})] (\mathbf{z}^{(k)})^T \boldsymbol{\beta}^{(k)*} \right| \\ & \lesssim \xi^{(k)} \sqrt{\frac{p}{n_k}} + \sqrt{\frac{\log K}{n_k}}, \end{aligned}$$

for all $k \in S$ and $r = 1 : R$, with probability at least $1 - C'(K^{-2} + K^{-1}e^{-C''p})$. Denote the LHS as W'' . Similar to the previous two proofs, we derive an upper bound for W'' by controlling $W'' - \mathbb{E}W''$ and $\mathbb{E}W''$, separately. The first part involving $W'' - \mathbb{E}W''$ is similar to the proof of part (i) in Lemma 12 and the second part involving $\mathbb{E}W''$ is similar to the proof of (E.74), so we omit the details.

The arguments to derive these three bounds can be used to derive other results similar to the lemmas used in the proof of Theorem 1. With these lemmas in hand, the remaining proof is almost the same as the proof of Theorem 1.

E.13.3 Proof of lemmas

Proof of Lemma 37. We will prove the contraction of w_r first, and only sketch the different part for the proof of contraction of $\boldsymbol{\mu}_r$ because the proofs are quite similar.

Part 1: Contraction of $|w_r(\boldsymbol{\theta}) - w_r^*|$:

First, note that $w_r(\boldsymbol{\theta}^*) = w_r^*$ and $\boldsymbol{\mu}_r(\boldsymbol{\theta}^*) = \boldsymbol{\mu}_r^*$. Therefore,

$$\begin{aligned} |w_r(\boldsymbol{\theta}) - w_r^*| &= \left| \mathbb{E}[\gamma_{\boldsymbol{\theta}}^{(r)}(\mathbf{z}) - \gamma_{\boldsymbol{\theta}^*}^{(r)}(\mathbf{z})] \right| \\ &\leq \sum_{\tilde{r}=1}^R w_r^{(k)*} \left| \mathbb{E}[\gamma_{\boldsymbol{\theta}}^{(r)}(\mathbf{z}) - \gamma_{\boldsymbol{\theta}^*}^{(r)}(\mathbf{z}) | y = \tilde{r}] \right| \\ &\leq \sum_{\tilde{r}=1}^R w_r^{(k)*} \sum_{r'=2}^R \left| \mathbb{E} \left[\frac{\partial \gamma_{\boldsymbol{\theta}}^{(r)}(\mathbf{z})}{\partial w_{r'}} \bigg|_{\boldsymbol{\theta}=\tilde{\boldsymbol{\theta}}_t} \bigg| y = \tilde{r} \right] \right| \cdot |w_{r'} - w_r^*| \\ &\quad + \sum_{\tilde{r}=1}^R w_r^{(k)*} \sum_{r'=2}^R \left| \mathbb{E} \left[\frac{\partial \gamma_{\boldsymbol{\theta}}^{(r)}(\mathbf{z})}{\partial \delta_{r'}} \bigg|_{\boldsymbol{\theta}=\tilde{\boldsymbol{\theta}}_t} \bigg| y = \tilde{r} \right] \right| \cdot |\delta_{r'} - \delta_r^*| \\ &\quad + \sum_{\tilde{r}=1}^R w_r^{(k)*} \sum_{r'=2}^R \left| \mathbb{E} \left[\frac{\partial \gamma_{\boldsymbol{\theta}}^{(r)}(\mathbf{z})}{\partial \boldsymbol{\beta}_{r'}} \bigg|_{\boldsymbol{\theta}=\tilde{\boldsymbol{\theta}}_t} \bigg| y = \tilde{r} \right]^T (\boldsymbol{\beta}_{r'} - \boldsymbol{\beta}_r^*) \right|, \end{aligned}$$

We only show how to bound $\left| \mathbb{E}[\gamma_{\boldsymbol{\theta}}^{(r)}(\mathbf{z}) - \gamma_{\boldsymbol{\theta}^*}^{(r)}(\mathbf{z}) | y = 1] \right|$, i.e. the case when $\tilde{r} = 1$. For the other $\tilde{r} = 2 : R$, the proof is the same by changing the reference level from $y = 1$ to $y = \tilde{r}$. Note that

$$\left| \mathbb{E}[\gamma_{\boldsymbol{\theta}}^{(r)}(\mathbf{z}) - \gamma_{\boldsymbol{\theta}^*}^{(r)}(\mathbf{z}) | y = 1] \right| \leq \sum_{r'=2}^R \left| \mathbb{E} \left[\frac{\partial \gamma_{\boldsymbol{\theta}}^{(r)}(\mathbf{z})}{\partial w_{r'}} \bigg|_{\boldsymbol{\theta}=\tilde{\boldsymbol{\theta}}_t} \bigg| y = 1 \right] \right| \cdot |w_{r'} - w_r^*|$$

$$\begin{aligned}
& + \sum_{r'=2}^R \left| \mathbb{E} \left[\frac{\partial \gamma_{\boldsymbol{\theta}}^{(r)}(\mathbf{z})}{\partial \delta_{r'}} \bigg|_{\boldsymbol{\theta}=\tilde{\boldsymbol{\theta}}_t} \bigg| y=1 \right] \right| \cdot |\delta_{r'} - \delta_r^*| \\
& + \sum_{r'=2}^R \left| \mathbb{E} \left[\frac{\partial \gamma_{\boldsymbol{\theta}}^{(r)}(\mathbf{z})}{\partial \boldsymbol{\beta}_{r'}} \bigg|_{\boldsymbol{\theta}=\tilde{\boldsymbol{\theta}}_t} \bigg| y=1 \right]^T (\boldsymbol{\beta}_{r'} - \boldsymbol{\beta}_r^*) \right|.
\end{aligned}$$

where $\tilde{\boldsymbol{\theta}}_t = (\{\tilde{w}_r\}_{r=2}^R, \{\tilde{\boldsymbol{\beta}}_r\}_{r=2}^R, \{\tilde{\delta}_r\}_{r=2}^R)$ with $\tilde{w}_r = tw_r + (1-t)w_r^*$, $\tilde{\boldsymbol{\beta}}_r = t\boldsymbol{\beta}_r + (1-t)\boldsymbol{\beta}_r^*$, $\tilde{\delta}_r = t\delta_r + (1-t)\delta_r^*$, and $\delta_r = \frac{1}{2}\boldsymbol{\beta}_r^T(\boldsymbol{\mu}_r + \boldsymbol{\mu}_1)$. We will bound the three terms on the RHS separately. Note that when $\boldsymbol{\theta} \in B_{\text{con}}(\boldsymbol{\theta}^*)$, we have $w_r \in [c_w/2, 1 - c_w]$, $\|\boldsymbol{\beta}_r - \boldsymbol{\beta}_r^*\|_2 \leq C_b\Delta$, and $\max_{r=1:R} \|\boldsymbol{\mu}_r - \boldsymbol{\mu}_r^*\|_2 \leq C_b\Delta$, hence $\tilde{w}_r \in [c_w/2, 1 - c_w]$, $\|\tilde{\boldsymbol{\beta}}_r - \boldsymbol{\beta}_r^*\|_2 \leq tC_b\Delta$.

(i) Bounding $|\mathbb{E}[\frac{\partial \gamma_{\boldsymbol{\theta}}^{(r)}(\mathbf{z})}{\partial w_{r'}} \big|_{\boldsymbol{\theta}=\tilde{\boldsymbol{\theta}}_t} | y=1]|$: Note that

$$\begin{aligned}
\frac{\partial \gamma_{\boldsymbol{\theta}}^{(r)}(\mathbf{z})}{\partial w_{r'}} &= \frac{\exp\{\tilde{\boldsymbol{\beta}}_r^T \mathbf{z} - \delta_r\}}{\tilde{w}_1 + \sum_{r=2}^R \tilde{w}_r \exp\{\tilde{\boldsymbol{\beta}}_r^T \mathbf{z} - \delta_r\}} - \frac{\tilde{w}_r \exp\{\tilde{\boldsymbol{\beta}}_r^T \mathbf{z} - \delta_r\} (\exp\{\tilde{\boldsymbol{\beta}}_r^T \mathbf{z} - \delta_r\} - 1)}{(\tilde{w}_1 + \sum_{r=2}^R \tilde{w}_r \exp\{\tilde{\boldsymbol{\beta}}_r^T \mathbf{z} - \delta_r\})^2} \\
&= \begin{cases} \frac{\exp\{\tilde{\boldsymbol{\beta}}_r^T \mathbf{z} - \delta_r\} (\tilde{w}_1 + \tilde{w}_r + \sum_{r' \neq r} \tilde{w}_{r'} \exp\{\tilde{\boldsymbol{\beta}}_{r'}^T \mathbf{z} - \delta_{r'}\})}{(\tilde{w}_1 + \sum_{r=2}^R \tilde{w}_r \exp\{\tilde{\boldsymbol{\beta}}_r^T \mathbf{z} - \delta_r\})^2}, & r' = r, \\ -\frac{\tilde{w}_r \cdot \exp\{\tilde{\boldsymbol{\beta}}_r^T \mathbf{z} - \delta_r\} (\exp\{\tilde{\boldsymbol{\beta}}_{r'}^T \mathbf{z} - \delta_{r'}\} - 1)}{(\tilde{w}_1 + \sum_{r=2}^R \tilde{w}_r \exp\{\tilde{\boldsymbol{\beta}}_r^T \mathbf{z} - \delta_r\})^2}, & r' \neq r. \end{cases}
\end{aligned}$$

Hence

$$\mathbb{E} \left[\frac{\partial \gamma_{\boldsymbol{\theta}}^{(r)}(\mathbf{z})}{\partial w_r} \bigg| y=1 \right] = \mathbb{E} \left[\underbrace{\frac{\exp\{\tilde{\boldsymbol{\beta}}_r^T \mathbf{z}^{(1)} - \delta_r\} (\tilde{w}_1 + \tilde{w}_r + \sum_{r' \neq r} \tilde{w}_{r'} \exp\{\tilde{\boldsymbol{\beta}}_{r'}^T \mathbf{z}^{(1)} - \delta_{r'}\})}{(\tilde{w}_1 + \sum_{r=2}^R \tilde{w}_r \exp\{\tilde{\boldsymbol{\beta}}_r^T \mathbf{z}^{(1)} - \delta_r\})^2}}_{(*)} \right].$$

Let $\tilde{z}_{r'} = \tilde{\boldsymbol{\beta}}_{r'}^T(\mathbf{z}^{(1)} - \boldsymbol{\mu}_1^*) \sim \mathcal{N}(0, \tilde{\boldsymbol{\beta}}_{r'}^T \boldsymbol{\Sigma}^* \tilde{\boldsymbol{\beta}}_{r'})$. And notice that

$$\tilde{\boldsymbol{\beta}}_{r'}^T \boldsymbol{\mu}_1^* - \tilde{\delta}_{r'} = t(\boldsymbol{\beta}_{r'}^T \boldsymbol{\mu}_1^* - \delta_{r'}) + (1-t)[(\boldsymbol{\beta}_{r'}^*)^T \boldsymbol{\mu}_1^* - \delta_{r'}^*],$$

where

$$\begin{aligned}
\boldsymbol{\beta}_{r'}^T \boldsymbol{\mu}_1^* - \delta_{r'} &= [\boldsymbol{\beta}_{r'}^* + (\boldsymbol{\beta}_{r'} - \boldsymbol{\beta}_{r'}^*)]^T \left[\frac{1}{2}(\boldsymbol{\mu}_1^* - \boldsymbol{\mu}_{r'}^*) + \frac{1}{2}(\boldsymbol{\mu}_{r'}^* - \boldsymbol{\mu}_{r'}) + \frac{1}{2}(\boldsymbol{\mu}_1^* - \boldsymbol{\mu}_1) \right] \\
&= -\frac{1}{2} \underbrace{(\boldsymbol{\mu}_{r'}^* - \boldsymbol{\mu}_1^*)^T (\boldsymbol{\Sigma}^*)^{-1} (\boldsymbol{\mu}_{r'}^* - \boldsymbol{\mu}_1^*)}_{A_{r'}^2} + \frac{1}{2} (\boldsymbol{\beta}_{r'}^*)^T (\boldsymbol{\Sigma}^*)^{1/2} (\boldsymbol{\Sigma}^*)^{-1/2} [(\boldsymbol{\mu}_{r'}^* - \boldsymbol{\mu}_{r'}) + (\boldsymbol{\mu}_1^* - \boldsymbol{\mu}_1)] \\
&\quad + \frac{1}{2} (\boldsymbol{\beta}_{r'} - \boldsymbol{\beta}_{r'}^*)^T (\boldsymbol{\Sigma}^*)^{1/2} (\boldsymbol{\Sigma}^*)^{-1/2} (\boldsymbol{\mu}_1^* - \boldsymbol{\mu}_{r'}^*) \\
&\quad + \frac{1}{2} (\boldsymbol{\beta}_{r'} - \boldsymbol{\beta}_{r'}^*)^T [(\boldsymbol{\mu}_{r'}^* - \boldsymbol{\mu}_{r'}) + (\boldsymbol{\mu}_1^* - \boldsymbol{\mu}_1)], \\
(\boldsymbol{\beta}_{r'}^*)^T \boldsymbol{\mu}_1^* - \delta_{r'}^* &= -\frac{1}{2} A_{r'}^2,
\end{aligned}$$

and $A_{r'} = \sqrt{(\boldsymbol{\mu}_{r'}^* - \boldsymbol{\mu}_1^*)^T (\boldsymbol{\Sigma}^*)^{-1} (\boldsymbol{\mu}_{r'}^* - \boldsymbol{\mu}_1^*)} = \sqrt{(\boldsymbol{\beta}_r^*)^T \boldsymbol{\Sigma}^* \boldsymbol{\beta}_r^*}$. By the fact that $\max_{r=1:R} \|\boldsymbol{\mu}_r - \boldsymbol{\mu}_r^*\|_2 \leq C_b \Delta$ and $\max_{r=1:R} \|\boldsymbol{\beta}_r - \boldsymbol{\beta}_r^*\|_2 \leq C_b \Delta$, we have

$$-\frac{1}{2}A_{r'}^2 - 2c_{\boldsymbol{\Sigma}}^{1/2}C_b\Delta A_{r'} - C_b^2\Delta^2 \leq \boldsymbol{\beta}_{r'}^T \boldsymbol{\mu}_1^* - \delta_{r'} \leq -\frac{1}{2}A_{r'}^2 + 2c_{\boldsymbol{\Sigma}}^{1/2}C_b\Delta A_{r'} + C_b^2\Delta^2,$$

implying that

$$-\frac{1}{2}A_{r'}^2 - 2c_{\boldsymbol{\Sigma}}^{1/2}C_b\Delta A_{r'} - C_b^2\Delta^2 \leq \tilde{\boldsymbol{\beta}}_{r'}^T \boldsymbol{\mu}_1^* - \tilde{\delta}_{r'} \leq -\frac{1}{2}A_{r'}^2 + 2c_{\boldsymbol{\Sigma}}^{1/2}C_b\Delta A_{r'} + C_b^2\Delta^2.$$

By Gaussian tail, we have

$$\mathbb{P}\left(\bigcap_{r'=2}^R \left\{|\tilde{z}_{r'}| \leq \frac{1}{4}\tilde{\boldsymbol{\beta}}_{r'}^T \boldsymbol{\Sigma}^* \tilde{\boldsymbol{\beta}}_{r'}\right\}\right) \geq 1 - CR \exp\left\{-\frac{1}{32}\tilde{\boldsymbol{\beta}}_{r'}^T \boldsymbol{\Sigma}^* \tilde{\boldsymbol{\beta}}_{r'}\right\}.$$

Denote event $\mathcal{E} = \bigcap_{r'=1}^R \left\{|\tilde{z}_{r'}| \leq \frac{1}{4}c_{\boldsymbol{\Sigma}}C_b^2\Delta^2 + \frac{1}{2}C_b c_{\boldsymbol{\Sigma}}^{1/2}\Delta A_{r'} + \frac{1}{4}A_{r'}^2\right\}$. Since

$$\begin{aligned} \frac{1}{4}\tilde{\boldsymbol{\beta}}_{r'}^T \boldsymbol{\Sigma}^* \tilde{\boldsymbol{\beta}}_{r'} &= \frac{1}{4}(\tilde{\boldsymbol{\beta}}_{r'} - \boldsymbol{\beta}_{r'}^*)^T \boldsymbol{\Sigma}^* (\tilde{\boldsymbol{\beta}}_{r'} - \boldsymbol{\beta}_{r'}^*) + \frac{1}{2}(\tilde{\boldsymbol{\beta}}_{r'} - \boldsymbol{\beta}_{r'}^*)^T (\boldsymbol{\Sigma}^*)^{1/2} (\boldsymbol{\Sigma}^*)^{1/2} \boldsymbol{\beta}_{r'}^* + \frac{1}{4}(\boldsymbol{\beta}_{r'}^*)^T \boldsymbol{\Sigma}^* \boldsymbol{\beta}_{r'}^* \\ &\leq \frac{1}{4}c_{\boldsymbol{\Sigma}}C_b^2\Delta^2 + \frac{1}{2}C_b c_{\boldsymbol{\Sigma}}^{1/2}\Delta A_{r'} + \frac{1}{4}A_{r'}^2, \end{aligned}$$

and

$$\tilde{\boldsymbol{\beta}}_{r'}^T \boldsymbol{\Sigma}^* \tilde{\boldsymbol{\beta}}_{r'} \geq A_{r'}^2 - c_{\boldsymbol{\Sigma}}C_b^2\Delta^2 - 2C_b c_{\boldsymbol{\Sigma}}^{1/2}\Delta A_{r'} \geq (1 - c_{\boldsymbol{\Sigma}}C_b^2 - 2C_b c_{\boldsymbol{\Sigma}}^{1/2})\Delta^2 \geq \frac{1}{2}\Delta^2,$$

we have

$$\mathbb{P}(\mathcal{E}) \geq 1 - CR \exp\left\{-\frac{1}{64}\Delta^2\right\}.$$

Then since $\min_{r'=1:R} A_{r'} \geq \Delta \geq 5c_{\boldsymbol{\Sigma}}^{1/2}C_b\Delta$ and $C_b \leq \frac{c_{\boldsymbol{\Sigma}}^{-1/2}}{40} \wedge (2c_{\boldsymbol{\Sigma}} + 8)^{-1/2}$, we have

$$\begin{aligned} (*) &\leq \mathbb{E}\left[\frac{\exp\{-\frac{1}{4}A_r^2 + \frac{5}{2}c_{\boldsymbol{\Sigma}}^{1/2}C_b\Delta A_r + (\frac{1}{4}c_{\boldsymbol{\Sigma}} + 1)C_b^2\Delta^2\}}{\tilde{w}_1^2}\right. \\ &\quad \cdot \left.\left(\tilde{w}_1 + \tilde{w}_r + \sum_{r' \neq r} \tilde{w}_{r'} \exp\left\{-\frac{1}{4}A_{r'}^2 + \frac{5}{2}c_{\boldsymbol{\Sigma}}^{1/2}C_b\Delta A_{r'} + \left(\frac{1}{4}c_{\boldsymbol{\Sigma}} + 1\right)C_b^2\Delta^2\right\}\right)\right] + \mathbb{P}(\mathcal{E}^c) \\ &\lesssim c_w^{-2} \exp\{-C\Delta^2\}. \end{aligned}$$

Hence,

$$\left|\mathbb{E}\left[\frac{\partial \gamma_{\boldsymbol{\theta}}^{(r)}(\mathbf{z})}{\partial w_r} \Big|_{\boldsymbol{\theta}=\tilde{\boldsymbol{\theta}}_t} \Big| y=1\right]\right| \lesssim c_w^{-2} \exp\{-C\Delta^2\}.$$

Similarly, it can be shown that

$$\left| \mathbb{E} \left[\frac{\partial \gamma_{\boldsymbol{\theta}}^{(r)}(\mathbf{z})}{\partial w_{r'}} \Big|_{\boldsymbol{\theta}=\tilde{\boldsymbol{\theta}}_t} \Big| y=1 \right] \right| \lesssim c_w^{-2} \exp\{-C\Delta^2\}.$$

for any $r' = 2 : R$.

(ii) Bounding $|\mathbb{E}[\frac{\partial \gamma_{\boldsymbol{\theta}}^{(r)}(\mathbf{z})}{\partial \delta_{r'}} \Big|_{\boldsymbol{\theta}=\tilde{\boldsymbol{\theta}}_t} | y=1]|$: Note that

$$\frac{\partial \gamma_{\boldsymbol{\theta}}^{(r)}(\mathbf{z})}{\partial \delta_{r'}} = \begin{cases} \frac{-w_r \cdot \exp\{\boldsymbol{\beta}_r^T \mathbf{z} - \delta_r\} \cdot \sum_{r' \neq r} w_{r'} \exp\{\boldsymbol{\beta}_{r'}^T \mathbf{z} - \delta_{r'}\}}{(w_1 + \sum_{r=2}^R w_r \exp\{\boldsymbol{\beta}_r^T \mathbf{z} - \delta_r\})^2}, & r' = r, \\ -\frac{w_r \cdot \exp\{\boldsymbol{\beta}_r^T \mathbf{z} - \delta_r\} \cdot w_{r'} \cdot \exp\{\boldsymbol{\beta}_{r'}^T \mathbf{z} - \delta_{r'}\}}{(w_1 + \sum_{r=2}^R w_r \exp\{\boldsymbol{\beta}_r^T \mathbf{z} - \delta_r\})^2}, & r' \neq r. \end{cases}$$

The analysis is almost the same as in (i), which leads to

$$\left| \mathbb{E} \left[\frac{\partial \gamma_{\boldsymbol{\theta}}^{(r)}(\mathbf{z})}{\partial \delta_{r'}} \Big|_{\boldsymbol{\theta}=\boldsymbol{\theta}} \right] \right| \lesssim c_w^{-2} \exp\{-C\Delta^2\},$$

for any $r' = 2 : R$. We omit the proof here.

(iii) Bounding $|\mathbb{E}[\frac{\partial \gamma_{\boldsymbol{\theta}}^{(r)}(\mathbf{z})}{\partial \boldsymbol{\beta}_{r'}} \Big|_{\boldsymbol{\theta}=\tilde{\boldsymbol{\theta}}_t} | y=1]^T (\boldsymbol{\beta}_{r'} - \boldsymbol{\beta}_r^*)|$: Note that

$$\frac{\partial \gamma_{\boldsymbol{\theta}}^{(r)}(\mathbf{z})}{\partial \boldsymbol{\beta}_{r'}} = \begin{cases} \frac{-w_r \cdot \exp\{\boldsymbol{\beta}_r^T \mathbf{z} - \delta_r\} \cdot \sum_{r' \neq r} w_{r'} \exp\{\boldsymbol{\beta}_{r'}^T \mathbf{z} - \delta_{r'}\} \mathbf{z}}{(w_1 + \sum_{r=2}^R w_r \exp\{\boldsymbol{\beta}_r^T \mathbf{z} - \delta_r\})^2}, & r' = r, \\ -\frac{w_r \cdot \exp\{\boldsymbol{\beta}_r^T \mathbf{z} - \delta_r\} \cdot w_{r'} \cdot \exp\{\boldsymbol{\beta}_{r'}^T \mathbf{z} - \delta_{r'}\} \mathbf{z}}{(w_1 + \sum_{r=2}^R w_r \exp\{\boldsymbol{\beta}_r^T \mathbf{z} - \delta_r\})^2}, & r' \neq r. \end{cases}$$

• When $r' = r$:

$$\begin{aligned} & \mathbb{E} \left[\left(\frac{\partial \gamma_{\boldsymbol{\theta}}^{(r)}(\mathbf{z})}{\partial \boldsymbol{\beta}_{r'}} \Big|_{\boldsymbol{\theta}=\tilde{\boldsymbol{\theta}}_t} \right)^T (\boldsymbol{\beta}_r - \boldsymbol{\beta}_r^*) \Big| y=1 \right] \\ &= \mathbb{E} \left[\frac{\tilde{w}_r \exp\{\tilde{\boldsymbol{\beta}}_r^T \mathbf{z}^{(1)} - \tilde{\delta}_r\} \sum_{r' \neq r} \tilde{w}_{r'} \cdot \exp\{\tilde{\boldsymbol{\beta}}_{r'}^T \mathbf{z}^{(1)} - \tilde{\delta}_{r'}\} \cdot (\mathbf{z}^{(1)})^T (\boldsymbol{\beta}_r - \boldsymbol{\beta}_r^*)}{(\tilde{w}_1 + \sum_{r=2}^R \tilde{w}_r \exp\{\tilde{\boldsymbol{\beta}}_r^T \mathbf{z}^{(1)} - \tilde{\delta}_r\})^2} \right] \\ &\leq \underbrace{\sqrt{\mathbb{E} \left[\frac{\tilde{w}_r \exp\{\tilde{\boldsymbol{\beta}}_r^T \mathbf{z}^{(1)} - \tilde{\delta}_r\} \sum_{r' \neq r} \tilde{w}_{r'} \cdot \exp\{\tilde{\boldsymbol{\beta}}_{r'}^T \mathbf{z}^{(1)} - \tilde{\delta}_{r'}\}}{(\tilde{w}_1 + \sum_{r=2}^R \tilde{w}_r \exp\{\tilde{\boldsymbol{\beta}}_r^T \mathbf{z}^{(1)} - \tilde{\delta}_r\})^2} \right]^2}}_{(1)} \cdot \underbrace{\sqrt{\mathbb{E}[(\mathbf{z}^{(1)})^T (\boldsymbol{\beta}_r - \boldsymbol{\beta}_r^*)]^2}}_{(2)}. \end{aligned}$$

Similar to the previous argument in (i), let $\tilde{z}_{r'} = \tilde{\boldsymbol{\beta}}_{r'}^T (\mathbf{z}^{(\tilde{r})} - \boldsymbol{\mu}_1^*) \sim \mathcal{N}(0, \tilde{\boldsymbol{\beta}}_{r'}^T \boldsymbol{\Sigma}^* \tilde{\boldsymbol{\beta}}_{r'})$ and event $\mathcal{E} = \bigcap_{r'=1}^R \{|\tilde{z}_{r'}| \leq \frac{1}{4} c_{\boldsymbol{\Sigma}} C_b^2 \Delta^2 + \frac{1}{2} C_b c_{\boldsymbol{\Sigma}}^{1/2} \Delta A_{r'} + \frac{1}{4} A_{r'}^2\}$, then

$$\mathbb{P}(\mathcal{E}) \geq 1 - CR \exp\left\{-\frac{1}{64} \Delta^2\right\}.$$

Similar to (i), we have

$$(1) \lesssim \sqrt{\mathbb{E} \left[\left(\frac{\tilde{w}_r \exp\{\tilde{\beta}_r^T \mathbf{z}^{(1)} - \tilde{\delta}_r\} \sum_{r' \neq r} \tilde{w}_{r'} \cdot \exp\{\tilde{\beta}_{r'}^T \mathbf{z}^{(1)} - \tilde{\delta}_{r'}\}}{(\tilde{w}_1 + \sum_{r=2}^R \tilde{w}_r \exp\{\tilde{\beta}_r^T \mathbf{z}^{(1)} - \tilde{\delta}_r\})^2} \right)^2 \middle| \mathcal{E} \right]} + \mathbb{P}(\mathcal{E}^c) \lesssim \exp\{-C\Delta^2\}.$$

Moreover, $(\mathbf{z}^{(1)})^T(\beta_r - \beta_r^*) = (\mathbf{z}^{(1)} - \mu_1^*)^T(\beta_r - \beta_r^*) + (\mu_1^*)^T(\beta_r - \beta_r^*)$, where $(\mathbf{z}^{(1)} - \mu_1^*)^T(\beta_r - \beta_r^*) \sim \mathcal{N}(0, (\beta_r - \beta_r^*)^T \Sigma^* (\beta_r - \beta_r^*))$ and $|(\mu_1^*)^T(\beta_r - \beta_r^*)| \leq M \|\beta_r - \beta_r^*\|_2$, hence

$$(2) \lesssim \sqrt{(\beta_r - \beta_r^*)^T \Sigma^* (\beta_r - \beta_r^*)} + M \|\beta_r - \beta_r^*\|_2 \lesssim (c_\Sigma^{1/2} + M) \|\beta_r - \beta_r^*\|_2 \lesssim (c_\Sigma^{1/2} + M) C_b \Delta.$$

Therefore, since $\Delta \leq 2M c_\Sigma^{1/2}$ and $\Delta \gtrsim \log^{1/2}(c_\Sigma M c_w^{-1})$,

$$\mathbb{E} \left[\left(\frac{\partial \gamma_\theta^{(r)}(\mathbf{z})}{\partial \beta_{r'}} \right)^T (\beta_r - \beta_r^*) \middle| y = 1 \right] \lesssim c_w^{-2} \exp\{-C' \Delta^2\} \cdot (c_\Sigma^{1/2} + M) M c_\Sigma^{1/2} \lesssim \exp\{-C' \Delta^2\}.$$

• When $r' \neq r$: we can obtain

$$\mathbb{E} \left[\left(\frac{\partial \gamma_\theta^{(r)}(\mathbf{z})}{\partial \beta_{r'}} \right)^T (\beta_r - \beta_r^*) \middle| y = 1 \right] \lesssim \exp\{-C' \Delta^2\}.$$

similarly.

Combining (i)-(iii), we have

$$|w_r(\theta) - w_r^*| \lesssim \exp\{-C'' \Delta^2\} \cdot \sum_{r=2}^R (|w_r - w_r^*| + |\delta_r - \delta_r^*| + \|\beta_r - \beta_r^*\|_2).$$

Part 2: Contraction of $\|\mu_r(\theta) - \mu_r^*\|_2$:

By definition,

$$\|\mu_r(\theta) - \mu_r^*\|_2 \leq \frac{\|\mathbb{E}[\gamma_\theta^{(r)}(\mathbf{z}) \mathbf{z}]\|_2}{w_r(\theta) w_r^*} \cdot |w_r^* - w_r(\theta)| + \frac{\|\mathbb{E}[(\gamma_\theta^{(r)}(\mathbf{z}) - \gamma_{\theta^*}^{(r)}(\mathbf{z})) \mathbf{z}]\|_2}{w_r^*}, \quad (\text{E.76})$$

implying that

$$\begin{aligned} \|\mathbb{E}[(\gamma_\theta^{(r)}(\mathbf{z}) - \gamma_{\theta^*}^{(r)}(\mathbf{z})) \mathbf{z}]\|_2 &\leq \sum_{r'=2}^R \left\| \mathbb{E} \left[\frac{\partial \gamma_\theta^{(r)}(\mathbf{z})}{\partial w_{r'}} \middle|_{\theta=\tilde{\theta}_t} \mathbf{z} \right] \right\|_2 \cdot |w_{r'} - w_{r'}^*| \\ &\quad + \sum_{r=2}^R \left\| \mathbb{E} \left[\frac{\partial \gamma_\theta^{(r)}(\mathbf{z})}{\partial \delta_{r'}} \middle|_{\theta=\tilde{\theta}_t} \mathbf{z} \right] \right\|_2 \cdot |\delta_{r'} - \delta_{r'}^*| \\ &\quad + \sum_{r=2}^R \left\| \mathbb{E} \left[\frac{\partial \gamma_\theta^{(r)}(\mathbf{z})}{\partial \beta_{r'}} \middle|_{\theta=\tilde{\theta}_t} \mathbf{z} \right] \right\|_2 \cdot \|\beta_{r'} - \beta_{r'}^*\|_2, \end{aligned}$$

where $\tilde{\boldsymbol{\theta}}_t = (\{\tilde{w}_r\}_{r=2}^R, \{\tilde{\boldsymbol{\beta}}_r\}_{r=2}^R, \{\tilde{\delta}_r\}_{r=2}^R)$ with $\tilde{w}_r = tw_r + (1-t)w_r^*$, $\tilde{\boldsymbol{\beta}}_r = t\boldsymbol{\beta}_r + (1-t)\boldsymbol{\beta}_r^*$, and $\tilde{\delta}_r = t\delta_r + (1-t)\delta_r^*$. We will bound the three terms on the RHS separately. Note that when $\boldsymbol{\theta} \in B_{\text{con}}(\boldsymbol{\theta}^*)$, we have $\tilde{w}_r \in (c_w/2, 1 - c_w)$, $\|\tilde{\boldsymbol{\beta}}_r - \boldsymbol{\beta}_r^*\|_2 \leq C_b\Delta$, and $|\tilde{\delta}_r - \delta_r| \leq C_b\Delta$.

For any $\mathbf{u} \in \mathbb{R}^p$ with $\|\mathbf{u}\|_2 \leq 1$ and any $\tilde{r} \in 1 : R$, similar to our previous arguments, we have

$$\left| \mathbb{E} \left[\frac{\partial \gamma_{\boldsymbol{\theta}}^{(r)}(\mathbf{z})}{\partial w_{r'}} \middle|_{\boldsymbol{\theta}=\tilde{\boldsymbol{\theta}}_t} \mathbf{z}^T \mathbf{u} \middle| y = \tilde{r} \right] \right| \leq \sqrt{\mathbb{E} \left[\frac{\partial \gamma_{\boldsymbol{\theta}}^{(r)}(\mathbf{z})}{\partial w_{r'}} \right]^2} \cdot \sqrt{\mathbb{E}[(\mathbf{z}^{(\tilde{r})})^T \mathbf{u}]^2} \lesssim \exp\{-C''\Delta^2\},$$

which leads to

$$\left\| \mathbb{E} \left[\frac{\partial \gamma_{\boldsymbol{\theta}}^{(r)}(\mathbf{z})}{\partial w_{r'}} \mathbf{z} \right] \right\|_2 \lesssim \exp\{-C''\Delta^2\},$$

for any $r' \in 2 : R$. Similarly, we have

$$\left\| \mathbb{E} \left[\frac{\partial \gamma_{\boldsymbol{\theta}}^{(r)}(\mathbf{z})}{\partial \delta_{r'}} \mathbf{z} \right] \right\|_2, \left\| \mathbb{E} \left[\frac{\partial \gamma_{\boldsymbol{\theta}}^{(r)}(\mathbf{z})}{\partial \boldsymbol{\beta}_{r'}} \mathbf{z} \right] \right\|_2 \lesssim \exp\{-C''\Delta^2\}.$$

for any $r' \in 2 : R$. Therefore, $\|\mathbb{E}[(\gamma_{\boldsymbol{\theta}}^{(r)}(\mathbf{z}) - \gamma_{\boldsymbol{\theta}^*}^{(r)}(\mathbf{z}))\mathbf{z}]\|_2 \lesssim \exp\{-C''\Delta^2\}$. By part 1, we have $\frac{\|\mathbb{E}[\gamma_{\boldsymbol{\theta}}^{(r)}(\mathbf{z})\mathbf{z}]\|_2}{w_r(\boldsymbol{\theta})w_r^*} \cdot |w_r^* - w_r(\boldsymbol{\theta})| \lesssim \exp\{-C''\Delta^2\} \cdot d(\boldsymbol{\theta}, \boldsymbol{\theta}^*)$. Hence by (E.76), we have $\|\boldsymbol{\mu}_r(\boldsymbol{\theta}) - \boldsymbol{\mu}_r^*\|_2 \lesssim \exp\{-C''\Delta^2\} \cdot d(\boldsymbol{\theta}, \boldsymbol{\theta}^*)$.

Combining part 1 and part 2, we complete the proof. \square

E.14 Proof of Theorem 8

Note that the excess risk

$$\begin{aligned} & R_{\hat{\boldsymbol{\theta}}^{(k)*}}(\mathcal{C}_{\hat{\boldsymbol{\theta}}^{(k)}}) - R_{\boldsymbol{\theta}^{(k)*}}(\mathcal{C}_{\boldsymbol{\theta}^{(k)*}}) \\ &= \mathbb{P}(y^{(k)} \neq \mathcal{C}_{\hat{\boldsymbol{\theta}}^{(k)}}(\mathbf{z}^{(k)})) - \mathbb{P}(y^{(k)} \neq \mathcal{C}_{\boldsymbol{\theta}^{(k)*}}(\mathbf{z}^{(k)})) \\ &= \int_{\mathcal{C}_{\hat{\boldsymbol{\theta}}^{(k)}} \neq \mathcal{C}_{\boldsymbol{\theta}^{(k)*}}} [\mathbb{P}(y^{(k)} = \mathcal{C}_{\boldsymbol{\theta}^{(k)*}}(\mathbf{z}) | \mathbf{z}^{(k)} = \mathbf{z}) - \mathbb{P}(y^{(k)} = \mathcal{C}_{\hat{\boldsymbol{\theta}}^{(k)}}(\mathbf{z}) | \mathbf{z}^{(k)} = \mathbf{z})] d\mathbb{P}_{\boldsymbol{\theta}^{(k)*}}(\mathbf{z}) \\ &= \int_{\mathcal{C}_{\hat{\boldsymbol{\theta}}^{(k)}} \neq \mathcal{C}_{\boldsymbol{\theta}^{(k)*}}} \left[\max_{r=1:R} \mathbb{P}(y^{(k)} = r | \mathbf{z}^{(k)} = \mathbf{z}) - \mathbb{P}(y^{(k)} = \mathcal{C}_{\hat{\boldsymbol{\theta}}^{(k)}}(\mathbf{z}) | \mathbf{z}^{(k)} = \mathbf{z}) \right] d\mathbb{P}_{\boldsymbol{\theta}^{(k)*}}(\mathbf{z}) \end{aligned} \quad (\text{E.77})$$

Let event $\mathcal{E} = \{\mathbf{z} : \max_r \mathbb{P}(y^{(k)} = r | \mathbf{z}^{(k)} = \mathbf{z}) - \max_j \{\mathbb{P}(y^{(k)} = j | \mathbf{z}^{(k)} = \mathbf{z}) : \mathbb{P}(y^{(k)} = j | \mathbf{z}^{(k)} = \mathbf{z}) < \max_r \mathbb{P}(y^{(k)} = r | \mathbf{z}^{(k)} = \mathbf{z})\} \leq t\}$. We claim that the margin condition $\mathbb{P}(\mathcal{E}) \lesssim t$ holds for any $t \leq$ a small constant c (to be verified). If this is the case, then denote $\tilde{\mathcal{E}} = \{\max_r |\eta_{\hat{\boldsymbol{\theta}}^{(k)}}^{(r)}(\mathbf{z}^{(k)}) - \eta_{\boldsymbol{\theta}^{(k)*}}^{(r)}(\mathbf{z}^{(k)})| \leq t/2\}$.

$$r^* = \arg \max_r \eta_{\boldsymbol{\theta}^{(k)*}}^{(r)}(\mathbf{z}),$$

$$\hat{r} = \arg \max_r \eta_{\hat{\theta}^{(k)}}^{(r)}(\mathbf{z}).$$

We have

RHS of (E.77)

$$\begin{aligned} &\leq \int_{\substack{r^* \neq \hat{r} \\ \mathcal{E}, \tilde{\mathcal{E}}}} [\eta_{\theta^{(k)*}}^{(r^*)}(\mathbf{z}) - \eta_{\theta^{(k)*}}^{(\hat{r})}(\mathbf{z})] d\mathbb{P}_{\theta^{(k)*}}(\mathbf{z}) + \int_{\substack{r^* \neq \hat{r} \\ \mathcal{E}^c, \tilde{\mathcal{E}}}} [\eta_{\theta^{(k)*}}^{(r^*)}(\mathbf{z}) - \eta_{\hat{\theta}^{(k)}}^{(\hat{r})}(\mathbf{z})] d\mathbb{P}_{\theta^{(k)*}}(\mathbf{z}) + \mathbb{P}(\tilde{\mathcal{E}}^c) \\ &\leq t\mathbb{P}(\mathcal{E}) + \mathbb{P}(\tilde{\mathcal{E}}^c), \end{aligned} \tag{E.78}$$

where the last inequality comes from the fact that when $r^* \neq \hat{r}$, $\eta_{\theta^{(k)*}}^{(r^*)}(\mathbf{z}) - \eta_{\theta^{(k)*}}^{(\hat{r})}(\mathbf{z}) \leq t$ on \mathcal{E} . And notice that on $\mathcal{E}^c \cap \tilde{\mathcal{E}}$, we must have $\hat{r} = r^*$ because if $\hat{r} \neq r^*$, then

$$\eta_{\hat{\theta}^{(k)}}^{(\hat{r})}(\mathbf{z}) - \eta_{\theta^{(k)*}}^{(r^*)}(\mathbf{z}) \leq \eta_{\theta^{(k)*}}^{(\hat{r})}(\mathbf{z}) - \eta_{\theta^{(k)*}}^{(r^*)}(\mathbf{z}) + \frac{t}{2} + \frac{t}{2} \leq -t + t < 0,$$

which is a contradiction with the definition of \hat{r} . Hence $\{r^* \neq \hat{r}\} \cap \mathcal{E} \cap \tilde{\mathcal{E}}$ is empty. Therefore $\int_{\substack{r^* \neq \hat{r} \\ \mathcal{E}^c, \tilde{\mathcal{E}}}} [\eta_{\theta^{(k)*}}^{(r^*)}(\mathbf{z}) - \eta_{\hat{\theta}^{(k)}}^{(\hat{r})}(\mathbf{z})] d\mathbb{P}_{\theta^{(k)*}}(\mathbf{z}) = 0$. Finally, by Lipschitzness,

$$\begin{aligned} \mathbb{P}(\tilde{\mathcal{E}}^c) &= \mathbb{P}\left(\max_r |\eta_{\hat{\theta}^{(k)}}^{(r)}(\mathbf{z}^{(k)}) - \eta_{\theta^{(k)*}}^{(r)}(\mathbf{z}^{(k)})| > t/2\right) \\ &\leq \sum_{r=1}^R \mathbb{P}\left(|\eta_{\hat{\theta}^{(k)}}^{(r)}(\mathbf{z}^{(k)}) - \eta_{\theta^{(k)*}}^{(r)}(\mathbf{z}^{(k)})| > t/2\right) \\ &\lesssim \mathbb{P}(|(\hat{\beta}^{(k)} - \beta^{(k)*})^T \mathbf{z}^{(k)} - \hat{\delta}^{(k)} + \delta^{(k)*} - \log \hat{w}_r^{(k)} + \log \hat{w}_r^{(1)} + \log w_r^{(k)*} - \log w_1^{(k)*}| > Ct) \\ &\lesssim \mathbb{P}(|(\hat{\beta}^{(k)} - \beta^{(k)*})^T (\mathbf{z}^{(k)} - \boldsymbol{\mu}^{(k)})| > C't) \\ &\lesssim \exp\left\{-\frac{Ct^2}{\|\hat{\beta}^{(k)} - \beta^{(k)*}\|_2^2}\right\}. \end{aligned}$$

Plugging back into (E.78), we have

$$R_{\hat{\theta}^{(k)*}}(\mathcal{C}_{\hat{\theta}^{(k)}}) - R_{\hat{\theta}^{(k)*}}(\mathcal{C}_{\theta^{(k)*}}) \lesssim t^2 + \exp\left\{-\frac{Ct^2}{\|\hat{\beta}^{(k)} - \beta^{(k)*}\|_2^2}\right\}.$$

Let $t \asymp d(\hat{\theta}^{(k)}, \theta^{(k)*}) \sqrt{\log d^{-1}(\hat{\theta}^{(k)}, \theta^{(k)*})}$:

$$R_{\hat{\theta}^{(k)*}}(\mathcal{C}_{\hat{\theta}^{(k)}}) - R_{\hat{\theta}^{(k)*}}(\mathcal{C}_{\theta^{(k)*}}) \lesssim d^2(\hat{\theta}^{(k)}, \theta^{(k)*}) \log d^{-1}(\hat{\theta}^{(k)}, \theta^{(k)*}) \lesssim d^2(\hat{\theta}^{(k)}, \theta^{(k)*}) \log\left(\frac{n_S}{p + \log n_S}\right).$$

Then plugging in the upper bound of $d(\hat{\theta}^{(k)}, \theta^{(k)*})$ in Theorem 7 completes the proof.

It remains to verify the margin condition $\mathbb{P}(\mathcal{E}) \lesssim t$ for any $t \leq$ a small constant c . In fact,

$$\begin{aligned}
\mathbb{P}(\mathcal{E}) &= \sum_{r=1}^R \sum_{j \neq r} \mathbb{P} \left(\arg \max_{r'} \mathbb{P}(y = r' | \mathbf{z}^{(k)}) = r, \arg \max_{r' \neq r} \mathbb{P}(y = r' | \mathbf{z}^{(k)}) = j, \right. \\
&\quad \left. \mathbb{P}(y = r | \mathbf{z}^{(k)}) - \mathbb{P}(y = j | \mathbf{z}^{(k)}) \leq t \right) \\
&\leq \sum_{r=1}^R \sum_{j \neq r} \mathbb{P} \left(\arg \max_{r'} \mathbb{P}(y = r' | \mathbf{z}^{(k)}) = r, \arg \max_{r' \neq r} \mathbb{P}(y = r' | \mathbf{z}^{(k)}) = j, \right. \\
&\quad \left. 1 - \frac{\mathbb{P}(y = j | \mathbf{z}^{(k)})}{\mathbb{P}(y = r | \mathbf{z}^{(k)})} \leq \frac{t}{\mathbb{P}(y = r | \mathbf{z}^{(k)})} \leq Rt \right) \\
&\leq \sum_{r=1}^R \sum_{j \neq r} \mathbb{P} \left(1 - Rt \leq \frac{\mathbb{P}(y = j | \mathbf{z}^{(k)})}{\mathbb{P}(y = r | \mathbf{z}^{(k)})} \right. \\
&\quad \left. = \exp \{ (\boldsymbol{\beta}_j^{(k)*} - \boldsymbol{\beta}_r^{(k)*})^T \mathbf{z}^{(k)} - \delta_j^{(k)*} + \delta_r^{(k)*} + \log w_j^{(k)*} - \log w_r^{(k)*} \} \right) \\
&\lesssim \sum_{r=1}^R \sum_{j \neq r} \sum_{r'=1}^R \mathbb{P} \left(\log(1 - Rt) \leq \mathcal{N}((\boldsymbol{\beta}_j^{(k)*} - \boldsymbol{\beta}_r^{(k)*})^T \boldsymbol{\mu}_r^{(k)*} - \delta_j^{(k)*} + \delta_r^{(k)*} + \log w_j^{(k)*} - \log w_r^{(k)*}, \right. \\
&\quad \left. (\boldsymbol{\beta}_j^{(k)*} - \boldsymbol{\beta}_r^{(k)*})^T \boldsymbol{\Sigma}^{(k)*} (\boldsymbol{\beta}_j^{(k)*} - \boldsymbol{\beta}_r^{(k)*})) \leq 0 \right) \\
&\lesssim -\log(1 - Rt) \\
&\lesssim t,
\end{aligned}$$

when $t > 0$ is less than some constant $c > 0$. Note that we used the fact that $(\boldsymbol{\beta}_j^{(k)*} - \boldsymbol{\beta}_r^{(k)*})^T \boldsymbol{\Sigma}^{(k)*} (\boldsymbol{\beta}_j^{(k)*} - \boldsymbol{\beta}_r^{(k)*}) \geq \Delta^2 \geq$ some constant C , which implies that the Gaussian density is upper bounded by a constant. Hence the marginal condition is true.

We want to point out that this multi-class extension of margin condition in binary case has been widely used in literature of multi-class classification. For example, see [Chen and Sun \(2006\)](#) and [Vigogna et al. \(2022\)](#).

E.15 Proof of Theorem 9

The proof is almost the same as the proof of Theorem 3, by noticing that we can make the GMM parameters the same across r -th task with $r \geq 3$ to reduce the problem to the case $R = 2$, so we do not repeat it here.

E.16 Proof of Theorem 10

E.16.1 Lemmas

Lemma 39. Consider $\bar{\theta} = \{\{w_r\}_{r=2}^R, \{\beta_r\}_{r=2}^R, \{\delta_r\}_{r=2}^R, \{\mu_r\}_{r=1}^R, \Sigma\}$ and $\bar{\theta}' = \{\{w'_r\}_{r=2}^R, \{\beta'_r\}_{r=2}^R, \{\delta'_r\}_{r=2}^R, \{\mu'_r\}_{r=1}^R, \Sigma'\}$ with $w_r = w'_r$ for $r \geq 3$, $\mu_r = \mu'_r = \mathbf{e}_r$ for $r \geq 2$, $\mu_1 = \mu'_1 = \mathbf{0}$, and $\Sigma = \Sigma' = \mathbf{I}_p$. Then $\beta_r = \beta'_r = \mathbf{e}_r$, $\delta_r = \delta'_r = \beta_r^T(\frac{\mu_1 + \mu_2}{2}) = \frac{1}{2}$ for $r \geq 2$, and

$$\mathbb{P}_{\bar{\theta}}(\mathcal{C}_{\bar{\theta}} \neq \mathcal{C}_{\bar{\theta}'}) \gtrsim |w_2 - w'_2|.$$

Lemma 40. Consider $\bar{\theta} = \{\{w_r\}_{r=2}^R, \{\beta_r\}_{r=2}^R, \{\delta_r\}_{r=2}^R, \{\mu_r\}_{r=1}^R, \Sigma\}$ and $\bar{\theta}' = \{\{w'_r\}_{r=2}^R, \{\beta'_r\}_{r=2}^R, \{\delta'_r\}_{r=2}^R, \{\mu'_r\}_{r=1}^R, \Sigma'\}$ with $w_r = w'_r = \frac{1}{R}$ for $r = 1 : R$, $\mu_r = (u+1)\mathbf{e}_r$, $\mu'_r = \mathbf{e}_r$ for $r \geq 2$, $\mu_1 = u\mathbf{e}_r$, $\mu'_1 = \mathbf{0}$, and $\Sigma = \Sigma' = \mathbf{I}_p$, where $0 < u \leq C$. Then $\beta_r = \beta'_r = \mathbf{e}_r$, $\delta_r = \mathbf{e}_r^T(\frac{\mu_1 + \mu_2}{2}) = u + \frac{1}{2}$, $\delta'_r = \mathbf{e}_r^T(\frac{\mu'_1 + \mu'_2}{2}) = \frac{1}{2}$ for $r \geq 2$, and

$$\mathbb{P}_{\bar{\theta}}(\mathcal{C}_{\bar{\theta}} \neq \mathcal{C}_{\bar{\theta}'}) \gtrsim u.$$

Lemma 41. Consider $\bar{\theta} = \{\{w_r\}_{r=2}^R, \{\beta_r\}_{r=2}^R, \{\delta_r\}_{r=2}^R, \{\mu_r\}_{r=1}^R, \Sigma\}$ and $\bar{\theta}' = \{\{w'_r\}_{r=2}^R, \{\beta'_r\}_{r=2}^R, \{\delta'_r\}_{r=2}^R, \{\mu'_r\}_{r=1}^R, \Sigma'\}$ with $w_r = w'_r = \frac{1}{R}$ for $r = 1 : R$, $\mu_r = \mu'_r = \mathbf{e}_r$ for $r \geq 3$, $\mu_1 = \mu'_1 = \mathbf{0}$, and $\Sigma = \Sigma' = \mathbf{I}_p$. Suppose μ_2 and μ'_2 satisfy $(\mu_2)_{3:R} = (\mu'_2)_{3:R} = \mathbf{0}$. Then $\beta_r = \beta'_r = \mathbf{e}_r$ for $r \geq 3$, $\delta_r = \delta'_r$ for $r \geq 3$, $\beta_2 = \mu_2$, $\beta'_2 = \mu'_2$, $\delta_2 = \frac{1}{2}\|\mu_2\|_2^2$, $\delta'_2 = \frac{1}{2}\|\mu'_2\|_2^2$ where $\|\mu_2\|_2 = \|\mu'_2\|_2 = 1$ with $\mu_2^T \mu'_2 > \frac{\sqrt{2}}{2}$, and

$$\mathbb{P}_{\bar{\theta}}(\mathcal{C}_{\bar{\theta}} \neq \mathcal{C}_{\bar{\theta}'}) \gtrsim \|\mu_2 - \mu'_2\|_2.$$

E.16.2 Main proof of Theorem 10

Given the three lemmas we presented, the proof is almost the same as the proof of Theorem 4. We do not repeat it here.

E.16.3 Proof of lemmas

Proof of Lemma 39. Note that z_j 's are independent given $y = 3$. We have

$$\begin{aligned} \mathbb{P}_{\bar{\theta}}(\mathcal{C}_{\bar{\theta}} \neq \mathcal{C}_{\bar{\theta}'}) &\geq \mathbb{P}_{\bar{\theta}}(\mathcal{C}_{\bar{\theta}}(\mathbf{z}) = 2, \mathcal{C}_{\bar{\theta}'}(\mathbf{z}) \neq 2) \\ &\geq \mathbb{P}_{\bar{\theta}}\left(z_2 - \frac{1}{2} - \log w_1 + \log w_2 \geq 0, z_2 - \frac{1}{2} - \log w'_1 + \log w'_2 \leq 0 \right. \\ &\quad \left. z_r - \frac{1}{2} - \log w_1 + \log w_r \leq 0 \text{ for all } r \geq 3\right) \\ &\geq w_3 \mathbb{P}_{\bar{\theta}}\left(z_2 - \frac{1}{2} - \log w_1 + \log w_2 \geq 0, z_2 - \frac{1}{2} - \log w'_1 + \log w'_2 \leq 0 \right. \\ &\quad \left. z_r - \frac{1}{2} - \log w_1 + \log w_r \leq 0 \text{ for all } r \geq 3 \middle| y = 3\right) \\ &\gtrsim \mathbb{P}_{\bar{\theta}}\left(\frac{1}{2} + \log w_1 - \log w_2 \leq z_2 \leq \frac{1}{2} + \log w'_1 - \log w'_2 \middle| y = 3\right) \end{aligned}$$

$$\begin{aligned}
& \cdot \prod_{r=3}^R \mathbb{P}\left(z_r - \frac{1}{2} - \log w_1 + \log w_r \leq 0\right) \\
& \gtrsim |\log w_1 - \log w_2 - \log w'_1 + \log w'_2| \\
& = |\log(1 - w_2) - \log w_2 - \log(1 - w'_2) + \log w'_2| \\
& \gtrsim |w_2 - w'_2|.
\end{aligned}$$

□

Proof of Lemma 40. Note that z_j 's are independent given $y = 3$. We have

$$\begin{aligned}
\mathbb{P}_{\bar{\theta}}(\mathcal{C}_{\bar{\theta}} \neq \mathcal{C}_{\bar{\theta}'}) & \geq \mathbb{P}_{\bar{\theta}}(\mathcal{C}_{\bar{\theta}}(\mathbf{z}) = 2, \mathcal{C}_{\bar{\theta}'}(\mathbf{z}) \neq 2) \\
& \geq \mathbb{P}_{\bar{\theta}}\left(z_2 - \frac{1}{2} - u \geq 0, z_2 - \frac{1}{2} > 0, z_r - \frac{1}{2} - u \leq 0, z_r - \frac{1}{2} \leq 0 \text{ for all } r \geq 3\right) \\
& \geq \mathbb{P}_{\bar{\theta}}\left(\frac{1}{2} \leq z_2 \leq \frac{1}{2} + u, z_r \leq \frac{1}{2} \text{ for all } r \geq 3\right) \\
& \gtrsim w_3 \mathbb{P}_{\bar{\theta}}\left(\frac{1}{2} \leq z_2 \leq \frac{1}{2} + u, z_r \leq \frac{1}{2} \text{ for all } r \geq 3 \middle| y = 3\right) \\
& \gtrsim w_3 \mathbb{P}_{\bar{\theta}}\left(\frac{1}{2} \leq z_2 \leq \frac{1}{2} + u \middle| y = 3\right) \cdot \prod_{r=3}^R \mathbb{P}\left(z_r \leq \frac{1}{2}\right) \\
& \gtrsim u,
\end{aligned}$$

where we used the fact that $\mathbb{P}(z_r \leq \frac{1}{2} | y = 3) \geq$ some constant C .

□

Proof of Lemma 40. Note that z_j 's are independent given $y = 3$. We have

$$\begin{aligned}
\mathbb{P}_{\bar{\theta}}(\mathcal{C}_{\bar{\theta}} \neq \mathcal{C}_{\bar{\theta}'}) & \geq \mathbb{P}_{\bar{\theta}}(\mathcal{C}_{\bar{\theta}}(\mathbf{z}) = 2, \mathcal{C}_{\bar{\theta}'}(\mathbf{z}) \neq 2) \\
& \geq \mathbb{P}_{\bar{\theta}}\left(\boldsymbol{\mu}_2^T \mathbf{z} - \frac{1}{2} \|\boldsymbol{\mu}_2\|_2^2 \geq 0, (\boldsymbol{\mu}'_2)^T z_2 - \frac{1}{2} \|\boldsymbol{\mu}'_2\|_2^2 \leq 0, z_r \leq \frac{1}{2} \text{ for all } r \geq 3\right) \\
& \geq w_3 \mathbb{P}_{\bar{\theta}}\left(\boldsymbol{\mu}_2^T \mathbf{z} - \frac{1}{2} \geq 0, (\boldsymbol{\mu}'_2)^T z_2 - \frac{1}{2} \leq 0, z_r \leq \frac{1}{2} \text{ for all } r \geq 3 \middle| y = 3\right) \\
& \gtrsim w_3 \mathbb{P}_{\bar{\theta}}\left(\boldsymbol{\mu}_2^T \mathbf{z} \geq \frac{1}{2}, (\boldsymbol{\mu}'_2)^T \mathbf{z} \leq \frac{1}{2} \middle| y = 3\right) \cdot \prod_{r=3}^R \mathbb{P}\left(z_r \leq \frac{1}{2} \middle| y = 3\right) \\
& \gtrsim \sqrt{1 - \frac{|\boldsymbol{\mu}_2^T \boldsymbol{\mu}'_2|^2}{\|\boldsymbol{\mu}_2\|_2^4}} \cdot \frac{|\boldsymbol{\mu}_2^T \boldsymbol{\mu}'_2|}{\|\boldsymbol{\mu}_2\|_2^2} \\
& \gtrsim \|\boldsymbol{\mu}_2 - \boldsymbol{\mu}'_2\|_2
\end{aligned}$$

The second last inequality is due to the fact that $\mathbb{P}(z_r \leq \frac{1}{2} | y = 3) \geq$ some constant C and Proposition 23 in [Azizyan et al. \(2013\)](#). The last inequality comes from Lemma 8.1 in [Cai et al. \(2019\)](#).

□

E.17 Proof of Theorem 16

WLOG, suppose that π_k^* satisfies $\pi_k^*(r) =$ the “majority class” \tilde{r} , if $\#\{k \in S : \pi_k(r) = \tilde{r}\} \geq \frac{1}{2}|S|$. Note that we can make this assumption because it suffices to recover $\{\iota(\pi_k^*)\}_{k \in S}$ with a permutation ι . And WLOG, suppose $\pi_k^*(r) = r$ for all $k \in S$. Let us consider any $\pi = \{\pi_k\}_{k=1}^K$ with $\pi_k(r) = \pi_k^*(r)$ for all $k \in S^c$ and $\pi \neq \pi^*$. It suffices to prove that $\text{score}(\pi) > \text{score}(\pi^*)$.

Recall that $\xi = \max_{k \in S} \min_{\pi} \max_{r \in [R]} \|(\hat{\Sigma}^{(k)})^{-1}(\hat{\mu}_{\pi_k(r)}^{(k)[0]} - \hat{\mu}_{\pi_k(1)}^{(k)[0]} - \beta_r^{(k)*})\|_2$. Note that

$$\begin{aligned}
\text{score}(\pi) - \text{score}(\pi^*) &= \underbrace{\sum_{\substack{k \neq k' \in S \\ \pi_k(1) = \pi_{k'}(1)}} \sum_{r=2}^R \|(\hat{\Sigma}^{(k)[0]})^{-1}(\hat{\mu}_{\pi_k(r)}^{(k)[0]} - \hat{\mu}_{\pi_k(1)}^{(k)[0]}) - (\hat{\Sigma}^{(k')[0]})^{-1}(\hat{\mu}_{\pi_{k'}(r)}^{(k')[0]} - \hat{\mu}_{\pi_{k'}(1)}^{(k')[0]})\|_2}_{[1]} \\
&+ \underbrace{\sum_{\substack{k \neq k' \in S \\ \pi_k(1) \neq \pi_{k'}(1)}} \sum_{r=2}^R \|(\hat{\Sigma}^{(k)[0]})^{-1}(\hat{\mu}_{\pi_k(r)}^{(k)[0]} - \hat{\mu}_{\pi_k(1)}^{(k)[0]}) - (\hat{\Sigma}^{(k')[0]})^{-1}(\hat{\mu}_{\pi_{k'}(r)}^{(k')[0]} - \hat{\mu}_{\pi_{k'}(1)}^{(k')[0]})\|_2}_{[2]} \\
&+ 2 \underbrace{\sum_{\substack{k \in S \\ k' \in S^c}} \sum_{r=2}^R \|(\hat{\Sigma}^{(k)[0]})^{-1}(\hat{\mu}_{\pi_k(r)}^{(k)[0]} - \hat{\mu}_{\pi_k(1)}^{(k)[0]}) - (\hat{\Sigma}^{(k')[0]})^{-1}(\hat{\mu}_{\pi_{k'}(r)}^{(k')[0]} - \hat{\mu}_{\pi_{k'}(1)}^{(k')[0]})\|_2}_{[3]} \\
&- \underbrace{\sum_{\substack{k \neq k' \in S \\ \pi_k(1) = \pi_{k'}(1)}} \sum_{r=2}^R \|(\hat{\Sigma}^{(k)[0]})^{-1}(\hat{\mu}_r^{(k)[0]} - \hat{\mu}_1^{(k)[0]}) - (\hat{\Sigma}^{(k')[0]})^{-1}(\hat{\mu}_r^{(k')[0]} - \hat{\mu}_1^{(k')[0]})\|_2}_{[1]'} \\
&- \underbrace{\sum_{\substack{k \neq k' \in S \\ \pi_k(1) \neq \pi_{k'}(1)}} \sum_{r=2}^R \|(\hat{\Sigma}^{(k)[0]})^{-1}(\hat{\mu}_r^{(k)[0]} - \hat{\mu}_1^{(k)[0]}) - (\hat{\Sigma}^{(k')[0]})^{-1}(\hat{\mu}_r^{(k')[0]} - \hat{\mu}_1^{(k')[0]})\|_2}_{[2]'}
\end{aligned}$$

$$- 2 \underbrace{\sum_{\substack{k \in S \\ k' \in S^c}} \sum_{r=2}^R \|(\widehat{\Sigma}^{(k)[0]})^{-1}(\widehat{\mu}_r^{(k)[0]} - \widehat{\mu}_1^{(k)[0]}) - (\widehat{\Sigma}^{(k')[0]})^{-1}(\widehat{\mu}_{\pi_{k'}(r)}^{(k')[0]} - \widehat{\mu}_{\pi_{k'}(1)}^{(k')[0]})\|_2}_{[3]'}$$

We have

$$\begin{aligned} [2] &\geq \sum_{\substack{k \neq k' \in S \\ \pi_k(1) \neq \pi_{k'}(1)}} \sum_{r=2}^R \left(\|(\widehat{\Sigma}^{(k)[0]})^{-1}(\widehat{\mu}_{\pi_k(r)}^{(k)[0]} - \widehat{\mu}_{\pi_k(1)}^{(k)[0]}) - (\widehat{\Sigma}^{(k)[0]})^{-1}(\widehat{\mu}_{\pi_{k'}(r)}^{(k)[0]} - \widehat{\mu}_{\pi_{k'}(1)}^{(k)[0]})\|_2 - 2h - \xi \right) \\ &\geq \sum_{\substack{k \neq k' \in S \\ \pi_k(1) \neq \pi_{k'}(1)}} R \left(\|(\widehat{\Sigma}^{(k)[0]})^{-1}(\widehat{\mu}_{\pi_k(1)}^{(k)[0]} - \widehat{\mu}_{\pi_{k'}(1)}^{(k)[0]})\|_2 - 2h - \xi \right) \\ &\geq \sum_{\substack{k \neq k' \in S \\ \pi_k(1) \neq \pi_{k'}(1)}} R(c_{\Sigma}^{-1/2} \Delta - 2h - 2\xi). \end{aligned}$$

Hence

$$[2] - [2]' \geq \sum_{\substack{k \neq k' \in S \\ \pi_k(1) \neq \pi_{k'}(1)}} [R(c_{\Sigma}^{-1/2} \Delta - 2h - 2\xi) - R(2\xi + 2h)] = \sum_{\substack{k \neq k' \in S \\ \pi_k(1) \neq \pi_{k'}(1)}} R(c_{\Sigma}^{-1/2} \Delta - 4h - 4\xi).$$

Therefore,

$$\begin{aligned} [1] &= \underbrace{\sum_{\substack{k \neq k' \in S \\ \pi_k(1) = \pi_{k'}(1)}} \sum_{r=2}^R \sum_{\pi_k(r) = \pi_{k'}(r)} \|(\widehat{\Sigma}^{(k)[0]})^{-1}(\widehat{\mu}_{\pi_k(r)}^{(k)[0]} - \widehat{\mu}_{\pi_k(1)}^{(k)[0]}) - (\widehat{\Sigma}^{(k')[0]})^{-1}(\widehat{\mu}_{\pi_{k'}(r)}^{(k')[0]} - \widehat{\mu}_{\pi_{k'}(1)}^{(k')[0]})\|_2}_{[1]_1} \\ &\quad + \underbrace{\sum_{\substack{k \neq k' \in S \\ \pi_k(1) = \pi_{k'}(1)}} \sum_{r=2}^R \sum_{\pi_k(r) \neq \pi_{k'}(r)} \|(\widehat{\Sigma}^{(k)[0]})^{-1}(\widehat{\mu}_{\pi_k(r)}^{(k)[0]} - \widehat{\mu}_{\pi_k(1)}^{(k)[0]}) - (\widehat{\Sigma}^{(k')[0]})^{-1}(\widehat{\mu}_{\pi_{k'}(r)}^{(k')[0]} - \widehat{\mu}_{\pi_{k'}(1)}^{(k')[0]})\|_2}_{[1]_2}. \end{aligned}$$

Correspondingly, we can decompose $[1]'$ in the same way as $[1]' = [1]'_1 + [1]'_2$ with

$$[1]'_1 = \sum_{\substack{k \neq k' \in S \\ \pi_k(1) = \pi_{k'}(1)}} \sum_{r=2}^R \sum_{\pi_k(r) = \pi_{k'}(r)} \|(\widehat{\Sigma}^{(k)[0]})^{-1}(\widehat{\mu}_r^{(k)[0]} - \widehat{\mu}_1^{(k)[0]}) - (\widehat{\Sigma}^{(k')[0]})^{-1}(\widehat{\mu}_r^{(k')[0]} - \widehat{\mu}_1^{(k')[0]})\|_2,$$

$$[1]_2' = \sum_{\substack{k \neq k' \in S \\ \pi_k(1) = \pi_{k'}(1)}} \sum_{r=2}^R \sum_{\pi_k(r) = \pi_{k'}(r)} \|(\widehat{\Sigma}^{(k)[0]})^{-1}(\widehat{\mu}_r^{(k)[0]} - \widehat{\mu}_1^{(k)[0]}) - (\widehat{\Sigma}^{(k')[0]})^{-1}(\widehat{\mu}_r^{(k')[0]} - \widehat{\mu}_1^{(k')[0]})\|_2.$$

Note that

$$\begin{aligned} [1]_2 &= \sum_{\substack{k \neq k' \in S \\ \pi_k(1) = \pi_{k'}(1)}} \sum_{r=2}^R \sum_{\pi_k(r) \neq \pi_{k'}(r)} \left(\|(\widehat{\Sigma}^{(k)[0]})^{-1}(\widehat{\mu}_{\pi_k(r)}^{(k)[0]} - \widehat{\mu}_{\pi_k(1)}^{(k)[0]}) - (\widehat{\Sigma}^{(k)[0]})^{-1}(\widehat{\mu}_{\pi_{k'}(r)}^{(k)[0]} - \widehat{\mu}_{\pi_{k'}(1)}^{(k)[0]})\|_2 - 2h - \xi \right) \\ &= \sum_{\substack{k \neq k' \in S \\ \pi_k(1) = \pi_{k'}(1)}} \sum_{r=2}^R \sum_{\pi_k(r) \neq \pi_{k'}(r)} \left(\|(\widehat{\Sigma}^{(k)[0]})^{-1}(\widehat{\mu}_{\pi_k(r)}^{(k)[0]} - \widehat{\mu}_{\pi_{k'}(r)}^{(k)[0]})\|_2 - 2h - \xi \right) \\ &\geq \sum_{\substack{k \neq k' \in S \\ \pi_k(1) = \pi_{k'}(1)}} \sum_{r=2}^R \sum_{\pi_k(r) \neq \pi_{k'}(r)} \left(c_{\Sigma}^{-1/2} \Delta - 2h - 2\xi \right), \end{aligned}$$

hence

$$[1]_2 - [1]_2' \geq \sum_{\substack{k \neq k' \in S \\ \pi_k(1) = \pi_{k'}(1)}} \sum_{r=2}^R \sum_{\pi_k(r) \neq \pi_{k'}(r)} \left(c_{\Sigma}^{-1/2} \Delta - 4h - 4\xi \right).$$

And

$$\begin{aligned} [1]_1 &= \sum_{\substack{k \neq k' \in S \\ \pi_k(1) = \pi_{k'}(1)}} \sum_{r=2}^R \sum_{\pi_k(r) = \pi_{k'}(r)} \|(\widehat{\Sigma}^{(k)[0]})^{-1}(\widehat{\mu}_{\pi_k(r)}^{(k)[0]} - \widehat{\mu}_{\pi_k(1)}^{(k)[0]}) - (\widehat{\Sigma}^{(k')[0]})^{-1}(\widehat{\mu}_{\pi_{k'}(r)}^{(k')[0]} - \widehat{\mu}_{\pi_{k'}(1)}^{(k')[0]})\|_2 \\ &= \underbrace{\sum_{\substack{k \neq k' \in S \\ \pi_k(1) = \pi_{k'}(1) = 1}} \sum_{r=2}^R \sum_{\pi_k(r) = \pi_{k'}(r)} \|(\widehat{\Sigma}^{(k)[0]})^{-1}(\widehat{\mu}_{\pi_k(r)}^{(k)[0]} - \widehat{\mu}_{\pi_k(1)}^{(k)[0]}) - (\widehat{\Sigma}^{(k')[0]})^{-1}(\widehat{\mu}_{\pi_{k'}(r)}^{(k')[0]} - \widehat{\mu}_{\pi_{k'}(1)}^{(k')[0]})\|_2}_{[1]_{11}} \\ &\quad + \underbrace{\sum_{\substack{k \neq k' \in S \\ \pi_k(1) = \pi_{k'}(1) \neq 1}} \sum_{r=2}^R \sum_{\pi_k(r) = \pi_{k'}(r)} \|(\widehat{\Sigma}^{(k)[0]})^{-1}(\widehat{\mu}_{\pi_k(r)}^{(k)[0]} - \widehat{\mu}_{\pi_k(1)}^{(k)[0]}) - (\widehat{\Sigma}^{(k')[0]})^{-1}(\widehat{\mu}_{\pi_{k'}(r)}^{(k')[0]} - \widehat{\mu}_{\pi_{k'}(1)}^{(k')[0]})\|_2}_{[1]_{12}}. \end{aligned}$$

$$\begin{aligned}
[1]_{12} &= \sum_{\substack{k \neq k' \in S \\ \pi_k(1) = \pi_{k'}(1) \neq 1}} \sum_{r=2}^R \sum_{\pi_k(r) = \pi_{k'}(r)} (\|(\widehat{\Sigma}^{(k)[0]})^{-1}(\widehat{\mu}_{\pi_k(r)}^{(k)[0]} - \widehat{\mu}_{\pi_k(1)}^{(k)[0]}) - (\widehat{\Sigma}^{(k')[0]})^{-1}(\widehat{\mu}_{\pi_{k'}(r)}^{(k')[0]} - \widehat{\mu}_{\pi_{k'}(1)}^{(k')[0]})\|_2 - 2h - \xi) \\
&\geq - \sum_{\substack{k \neq k' \in S \\ \pi_k(1) = \pi_{k'}(1) \neq 1}} \sum_{r=2}^R \sum_{\pi_k(r) = \pi_{k'}(r)} (2h + \xi).
\end{aligned}$$

An important observation is that $[1]_{11} = [1]_{11}'$. Therefore,

$$[1]_1 - [1]_1' = [1]_{12} - [1]_{12}' \geq - \sum_{\substack{k \neq k' \in S \\ \pi_k(1) = \pi_{k'}(1) \neq 1}} \sum_{r=2}^R \sum_{\pi_k(r) = \pi_{k'}(r)} (4h + 3\xi).$$

And

$$\begin{aligned}
[1] - [1]' &= [1]_2 - [1]_2' + [1]_1 - [1]_1' \\
&\geq \sum_{\substack{k \neq k' \in S \\ \pi_k(1) = \pi_{k'}(1)}} \sum_{r=2}^R \sum_{\pi_k(r) \neq \pi_{k'}(r)} (c_{\Sigma}^{-1/2} \Delta - 4h - 4\xi) - \sum_{\substack{k \neq k' \in S \\ \pi_k(1) = \pi_{k'}(1) \neq 1}} \sum_{r=2}^R \sum_{\pi_k(r) = \pi_{k'}(r)} (4h + 3\xi).
\end{aligned}$$

Furthermore, by triangle inequality,

$$\begin{aligned}
[3] - [3]' &= 2 \sum_{\substack{k \in S \\ k' \in S^c}} \sum_{r=2}^R \|(\widehat{\Sigma}^{(k)[0]})^{-1}(\widehat{\mu}_{\pi_k(r)}^{(k)[0]} - \widehat{\mu}_{\pi_k(1)}^{(k)[0]}) - (\widehat{\Sigma}^{(k')[0]})^{-1}(\widehat{\mu}_{\pi_{k'}(r)}^{(k')[0]} - \widehat{\mu}_{\pi_{k'}(1)}^{(k')[0]})\|_2 \\
&\quad - 2 \sum_{\substack{k \in S \\ k' \in S^c}} \sum_{r=2}^R \|(\widehat{\Sigma}^{(k)[0]})^{-1}(\widehat{\mu}_r^{(k)[0]} - \widehat{\mu}_1^{(k)[0]}) - (\widehat{\Sigma}^{(k')[0]})^{-1}(\widehat{\mu}_{\pi_{k'}(r)}^{(k')[0]} - \widehat{\mu}_{\pi_{k'}(1)}^{(k')[0]})\|_2 \\
&\geq -2 \sum_{\substack{k \in S \\ k' \in S^c}} \sum_{r=2}^R \|(\widehat{\Sigma}^{(k)[0]})^{-1}(\widehat{\mu}_{\pi_k(r)}^{(k)[0]} - \widehat{\mu}_{\pi_k(1)}^{(k)[0]} - \widehat{\mu}_r^{(k')[0]} + \widehat{\mu}_1^{(k')[0]})\|_2.
\end{aligned}$$

Putting all pieces together,

$$\begin{aligned}
\text{score}(\pi) - \text{score}(\pi^*) &\geq \sum_{\substack{k \neq k' \in S \\ \pi_k(1) \neq \pi_{k'}(1)}} R(c_{\Sigma}^{-1/2} \Delta - 4h - 4\xi) + \sum_{\substack{k \neq k' \in S \\ \pi_k(1) = \pi_{k'}(1)}} \sum_{r=2}^R \sum_{\pi_k(r) \neq \pi_{k'}(r)} (c_{\Sigma}^{-1/2} \Delta - 4h - 4\xi) \\
&\quad - \sum_{\substack{k \neq k' \in S \\ \pi_k(1) = \pi_{k'}(1) \neq 1}} \sum_{r=2}^R \sum_{\pi_k(r) = \pi_{k'}(r)} (4h + 3\xi)
\end{aligned}$$

$$-2 \sum_{k \in S} \sum_{r=2}^R \|(\widehat{\Sigma}^{(k)[0]})^{-1}(\widehat{\mu}_{\pi_k(r)}^{(k)[0]} - \widehat{\mu}_{\pi_k(1)}^{(k)[0]} - \widehat{\mu}_r^{(k')[0]} + \widehat{\mu}_1^{(k')[0]})\|_2 \cdot |S^c|.$$

Denote m_r = the majority class among $\{\pi_k(r)\}_{k \in S}$ and $S_{\tilde{r}}^{(r)} = \{k \in S : \pi_k(r) = \tilde{r}\}$.

(i) **Case 1:** $|S_1^{(1)}| \leq \frac{2}{3}|S|$.

Since $\pi_k^*(1) = 1$, we have $|S_1^{(r)}| \leq \frac{2}{3}|S|$ for all $r \in [R]$, otherwise by our assumption, $m_1 = r_0$ since r_0 satisfies $|S_1^{(r_0)}| > \frac{2}{3}|S| \geq |S_1^{(1)}|$, which is a contradiction. Therefore,

$$\begin{aligned} \sum_{\substack{k \neq k' \in S \\ \pi_k(1) \neq \pi_{k'}(1)}} 1 &= 2 \sum_{r \neq r'} |S_1^{(r)}| \cdot |S_1^{(r')}| \\ &= \left(\sum_{r=1}^R |S_1^{(r)}| \right)^2 - \sum_{r=1}^R |S_1^{(r)}|^2 \\ &= |S|^2 - \sum_{r=1}^R |S_1^{(r)}|^2 \\ &\geq |S|^2 - \left(\frac{4}{9}|S|^2 + \frac{1}{9}|S|^2 \right) \\ &= \frac{4}{9}|S|^2, \end{aligned}$$

and

$$\begin{aligned} \sum_{\substack{k \neq k' \in S \\ \pi_k(1) = \pi_{k'}(1) \neq 1}} \sum_{r=2}^R \sum_{\pi_k(r) = \pi_{k'}(r)} 1 &\leq R \cdot 2 \binom{|S| - |S_1^{(1)}|}{2} \\ &\leq R(|S| - |S_1^{(1)}|)^2 \\ &\leq R|S|^2. \end{aligned}$$

Also,

$$\sum_{k \in S} \sum_{r=2}^R 1 \leq 2R|S|.$$

Hence, since $\frac{4}{9}|S| - 4D|S^c| = \frac{4}{9}(1 - \epsilon)K - 4D\epsilon K > 0$,

$$\begin{aligned} \text{score}(\pi) - \text{score}(\pi^*) &\geq \frac{4}{9}|S|^2 R(c_{\Sigma}^{-1/2} \Delta - 4h - 4\xi) - |S|^2 R(4h + 4\xi) - 2R|S||S^c|(2Dc_{\Sigma}^{1/2} \Delta + 2\xi) \\ &= |S|R \left[\left(\frac{4}{9}c_{\Sigma}^{-1/2}|S| - 4Dc_{\Sigma}^{1/2}|S^c| \right) \Delta - \frac{52}{9}|S|h - \left(\frac{52}{9}|S| + 4|S^c| \right) \xi \right] \\ &> 0. \end{aligned}$$

(ii) **Case 2:** $|S_1^{(1)}| > \frac{2}{3}|S|$.

In this case, by our assumption, we must have $m_1 = 1$. And

$$\begin{aligned}
\sum_{\substack{k \neq k' \in S \\ \pi_k(1) \neq \pi_{k'}(1)}} 1 &\geq 2|S_1^{(1)}|(|S| - |S_1^{(1)}|), \\
\sum_{\substack{k \neq k' \in S \\ \pi_k(1) = \pi_{k'}(1) \neq 1}} \sum_{r=2}^R \sum_{\pi_k(r) = \pi_{k'}(r)} 1 &\leq R \sum_{k, k'} \mathbb{1}(k \neq k' \in S, \pi_k(1) = \pi_{k'}(1) \neq 1) \\
&\leq R \cdot 2 \binom{|S| - |S_1^{(1)}|}{2} \\
&\leq R(|S| - |S_1^{(1)}|)^2 \\
&\leq R \frac{1}{2} |S_1^{(1)}|(|S| - |S_1^{(1)}|).
\end{aligned}$$

Moreover,

$$\begin{aligned}
&\sum_{k \in S} \sum_{r=2}^R \|(\widehat{\Sigma}^{(k)[0]})^{-1}(\widehat{\mu}_{\pi_k(r)}^{(k)[0]} - \widehat{\mu}_{\pi_k(1)}^{(k)[0]} - \widehat{\mu}_r^{(k)[0]} + \widehat{\mu}_1^{(k)[0]})\|_2 \cdot |S^c| \\
&\leq \sum_{\substack{k \in S \\ \pi_k(1) \neq 1}} \sum_{r=2}^R \|(\widehat{\Sigma}^{(k)[0]})^{-1}(\widehat{\mu}_{\pi_k(r)}^{(k)[0]} - \widehat{\mu}_{\pi_k(1)}^{(k)[0]} - \widehat{\mu}_r^{(k)[0]} + \widehat{\mu}_1^{(k)[0]})\|_2 \cdot |S^c| \\
&\quad + \sum_{k \in S} \sum_{\substack{\pi_k(1)=1 \\ \pi_k(r) \neq r}} \|(\widehat{\Sigma}^{(k)[0]})^{-1}(\widehat{\mu}_{\pi_k(r)}^{(k)[0]} - \widehat{\mu}_r^{(k)[0]})\|_2 \cdot |S^c| \\
&\leq (|S| - |S_1^{(1)}|)R \cdot (2c_{\Sigma}^{1/2} \Delta + 2\xi) |S^c| + \sum_{k \in S} \sum_{\substack{\pi_k(1)=1 \\ \pi_k(r) \neq r}} (Dc_{\Sigma}^{1/2} \Delta + \xi) |S^c|.
\end{aligned}$$

For r satisfying $|S_r^{(r)}| \leq \frac{1}{2}|S|$, we have

$$\begin{aligned}
\sum_{k \in S} \sum_{\substack{\pi_k(1)=1 \\ \pi_k(r) \neq r}} &\leq 2|S_1^{(1)}|, \\
\sum_{\substack{k \neq k' \in S \\ \pi_k(1) = \pi_{k'}(1)}} \sum_{\pi_k(r) \neq \pi_{k'}(r)} 1 &\geq 2 \cdot \frac{1}{2} |S| \left(\frac{2}{3} |S| - \frac{1}{2} |S| \right) = \frac{1}{6} |S|^2.
\end{aligned}$$

For r satisfying $|S_r^{(r)}| > \frac{1}{2}|S|$, we have $m_r = r$. Define $\widetilde{S}_r^{(r)} = \{k \in S : \pi_k(1) = 1, \pi_k(r) = r\}$. Note that $|\widetilde{S}_r^{(r)}| \geq \frac{2}{3}|S| - \frac{1}{2}|S| = \frac{1}{6}|S|$. Furthermore,

$$\sum_{k \in S} \sum_{\substack{\pi_k(1)=1 \\ \pi_k(r) \neq r}} \leq |S_1^{(1)}| - |\widetilde{S}_r^{(r)}|,$$

$$\sum_{\substack{k \neq k' \in S \\ \pi_k(1) = \pi_{k'}(1)}} \sum_{\pi_k(r) \neq \pi_{k'}(r)} 1 \geq 2|\tilde{S}_r^{(r)}|(|S_1^{(1)}| - |\tilde{S}_r^{(r)}|) \geq \frac{1}{3}|S|(|S_1^{(1)}| - |\tilde{S}_r^{(r)}|).$$

This implies that

$$\begin{aligned} & \text{score}(\pi) - \text{score}(\pi^*) \\ & \geq 2|S_1^{(1)}|(|S| - |S_1^{(1)}|)R(c_{\Sigma}^{-1/2}\Delta - 4h - 4\xi) - R\frac{1}{2}|S_1^{(1)}|(|S| - |S_1^{(1)}|) \cdot (4h + 3\xi) \\ & \quad + \sum_{r: |S_r^{(r)}| \leq |S|/2} \left[\frac{1}{6}|S|^2(c_{\Sigma}^{-1/2}\Delta - 4h - 4\xi) - 2|S_1^{(1)}| \cdot 2|S^c|(c_{\Sigma}^{1/2}D\Delta + \xi) \right] \\ & \quad + \sum_{r: |S_r^{(r)}| > |S|/2} \left[\frac{1}{3}|S|(|S_1^{(1)}| - |\tilde{S}_r^{(r)}|)(c_{\Sigma}^{-1/2}\Delta - 4h - 4\xi) - (|S_1^{(1)}| - |\tilde{S}_r^{(r)}|) \cdot |S^c| \cdot 2(c_{\Sigma}^{1/2}D\Delta + \xi) \right] \\ & \geq |S_1^{(1)}|(|S| - |S_1^{(1)}|)R\left(2c_{\Sigma}^{-1/2}\Delta - 10h - \frac{19}{2}\xi\right) \\ & \quad + \sum_{r: |S_r^{(r)}| \leq |S|/2} |S| \left[\left(\frac{1}{6}c_{\Sigma}^{-1/2}|S| - 4Dc_{\Sigma}^{1/2}|S^c|\right)\Delta - \frac{2}{3}|S|h - \left(\frac{2}{3}|S| + 4|S^c|\right)\xi \right] \\ & \quad + \sum_{r: |S_r^{(r)}| > |S|/2} \left[\left(\frac{1}{3}c_{\Sigma}^{-1/2}|S| - 2Dc_{\Sigma}^{1/2}|S^c|\right)\Delta - \frac{4}{3}|S|h - \left(\frac{4}{3}|S| + 2|S^c|\right)\xi \right] (|S_1^{(1)}| - |\tilde{S}_r^{(r)}|) \\ & > 0. \end{aligned}$$

E.18 Proof of Theorem 17

WLOG, consider the step $\tilde{K} \in [K]$ in the for loop and the case that $\iota = \text{identify}$ mapping from $[K]$ to $[K]$, and $\tilde{K} \in S$. Denote $\tilde{S} = S \cap [\tilde{K}]$ and $\tilde{S}^c = S^c \cap [\tilde{K}]$, hence $[\tilde{K}] = \tilde{S} \cup \tilde{S}^c$. WLOG, consider $\pi_1 = \pi_2 = \dots = \pi_{\tilde{K}-1} = \text{identify}$ from $[R]$ to $[R]$. Denote $\boldsymbol{\pi} = \{\pi_k\}_{k=1}^{\tilde{K}}$ and $\tilde{\boldsymbol{\pi}} = \{\pi_k\}_{k=1}^{\tilde{K}-1} \cup \{\tilde{\pi}_{\tilde{K}}\}$ with $\tilde{\pi}_{\tilde{K}} = \text{identify}$ from $[R]$ to $[R]$. It suffices to show that

$$\text{score}(\boldsymbol{\pi}) > \text{score}(\tilde{\boldsymbol{\pi}}), \quad (\text{E.79})$$

for any $\boldsymbol{\pi}$ with $\pi_{\tilde{K}} \neq \tilde{\pi}_{\tilde{K}} = \text{identify}$ from $[R]$ to $[R]$. If this is the case, then $\hat{\boldsymbol{\pi}} = \{\hat{\pi}_k\}_{k=1}^K$ satisfies $\hat{\pi}_k = \text{identity}$ for all $k \in S$, which completes the proof.

We focus on the derivation of (E.79) in the remaining part of the proof.

(i) Case 1: $\pi_{\tilde{K}}(1) = 1$.

$$\text{score}(\pi) - \text{score}(\pi^*) = \underbrace{\sum_{r=2}^R \sum_{k \in \tilde{S}} \|(\hat{\Sigma}^{(k)[0]})^{-1}(\hat{\boldsymbol{\mu}}_r^{(k)[0]} - \hat{\boldsymbol{\mu}}_1^{(k)[0]}) - (\hat{\Sigma}^{(\tilde{K})[0]})^{-1}(\hat{\boldsymbol{\mu}}_{\pi_{\tilde{K}}(r)}^{(\tilde{K})[0]} - \hat{\boldsymbol{\mu}}_1^{(\tilde{K})[0]})\|_2}_{[1]}$$

$$\begin{aligned}
& + \underbrace{\sum_{r=2}^R \sum_{k \in \tilde{S}^c} \|(\widehat{\Sigma}^{(k)[0]})^{-1}(\widehat{\mu}_{\pi_k(r)}^{(k)[0]} - \widehat{\mu}_{\pi_k(1)}^{(k)[0]}) - (\widehat{\Sigma}^{(\tilde{K})[0]})^{-1}(\widehat{\mu}_{\pi_{\tilde{K}}(r)}^{(\tilde{K})[0]} - \widehat{\mu}_1^{(\tilde{K})[0]})\|_2}_{[2]} \\
& - \underbrace{\sum_{r=2}^R \sum_{k \in \tilde{S}} \|(\widehat{\Sigma}^{(k)[0]})^{-1}(\widehat{\mu}_r^{(k)[0]} - \widehat{\mu}_1^{(k)[0]}) - (\widehat{\Sigma}^{(\tilde{K})[0]})^{-1}(\widehat{\mu}_r^{(\tilde{K})[0]} - \widehat{\mu}_1^{(\tilde{K})[0]})\|_2}_{[1]'} \\
& - \underbrace{\sum_{r=2}^R \sum_{k \in \tilde{S}^c} \|(\widehat{\Sigma}^{(k)[0]})^{-1}(\widehat{\mu}_{\pi_k(r)}^{(k)[0]} - \widehat{\mu}_{\pi_k(1)}^{(k)[0]}) - (\widehat{\Sigma}^{(\tilde{K})[0]})^{-1}(\widehat{\mu}_r^{(\tilde{K})[0]} - \widehat{\mu}_1^{(\tilde{K})[0]})\|_2}_{[2]'} .
\end{aligned}$$

Note that

$$\begin{aligned}
[1] - [1]' &= \#\{r \in 2 : R : \pi_{\tilde{K}}(r) \neq r\} \cdot \\
& \sum_{k \in \tilde{S}} \left[\|(\widehat{\Sigma}^{(k)[0]})^{-1}(\widehat{\mu}_r^{(k)[0]} - \widehat{\mu}_1^{(k)[0]}) - (\widehat{\Sigma}^{(\tilde{K})[0]})^{-1}(\widehat{\mu}_{\pi_{\tilde{K}}(r)}^{(\tilde{K})[0]} - \widehat{\mu}_1^{(\tilde{K})[0]})\|_2 \right. \\
& \quad \left. - \|(\widehat{\Sigma}^{(k)[0]})^{-1}(\widehat{\mu}_r^{(k)[0]} - \widehat{\mu}_1^{(k)[0]}) - (\widehat{\Sigma}^{(\tilde{K})[0]})^{-1}(\widehat{\mu}_r^{(\tilde{K})[0]} - \widehat{\mu}_1^{(\tilde{K})[0]})\|_2 \right] \\
& \geq \#\{r \in 2 : R : \pi_{\tilde{K}}(r) \neq r\} \cdot \left[\|(\Sigma^{(k)*})^{-1}(\mu_r^{(k)*} - \mu_{\pi_{\tilde{K}}(r)}^{(k)*})\|_2 - 2\xi - 2h - 2\xi - 2h \right] \\
& \geq \#\{r \in 2 : R : \pi_{\tilde{K}}(r) \neq r\} \cdot |\tilde{S}| \cdot (\Delta c_{\Sigma}^{-1/2} - 4\xi - 4h).
\end{aligned}$$

$$\begin{aligned}
[2] - [2]' &= \#\{r \in 2 : R : \pi_{\tilde{K}}(r) \neq r\} \cdot \\
& \sum_{k \in \tilde{S}^c} \left[\|(\widehat{\Sigma}^{(k)[0]})^{-1}(\widehat{\mu}_{\pi_k(r)}^{(k)[0]} - \widehat{\mu}_{\pi_k(1)}^{(k)[0]}) - (\widehat{\Sigma}^{(\tilde{K})[0]})^{-1}(\widehat{\mu}_{\pi_{\tilde{K}}(r)}^{(\tilde{K})[0]} - \widehat{\mu}_1^{(\tilde{K})[0]})\|_2 \right. \\
& \quad \left. - \|(\widehat{\Sigma}^{(k)[0]})^{-1}(\widehat{\mu}_{\pi_k(r)}^{(k)[0]} - \widehat{\mu}_{\pi_k(1)}^{(k)[0]}) - (\widehat{\Sigma}^{(\tilde{K})[0]})^{-1}(\widehat{\mu}_r^{(\tilde{K})[0]} - \widehat{\mu}_1^{(\tilde{K})[0]})\|_2 \right] \\
& \geq -\#\{r \in 2 : R : \pi_{\tilde{K}}(r) \neq r\} \cdot \sum_{k \in \tilde{S}^c} \|(\widehat{\Sigma}^{(\tilde{K})[0]})^{-1}(\widehat{\mu}_{\pi_{\tilde{K}}(r)}^{(\tilde{K})[0]} - \widehat{\mu}_r^{(\tilde{K})[0]})\|_2 \\
& \geq -\#\{r \in 2 : R : \pi_{\tilde{K}}(r) \neq r\} \cdot |\tilde{S}^c| \cdot [\|(\Sigma^{(\tilde{K})*})^{-1}(\mu_{\pi_{\tilde{K}}(r)}^{(\tilde{K})*} - \mu_r^{(\tilde{K})*})\|_2 + \xi] \\
& \geq -\#\{r \in 2 : R : \pi_{\tilde{K}}(r) \neq r\} \cdot |\tilde{S}^c| \cdot (Dc_{\Sigma}^{1/2}\Delta + \xi).
\end{aligned}$$

These imply that

$$\text{score}(\boldsymbol{\pi}) - \text{score}(\tilde{\boldsymbol{\pi}}) = [1] - [1]' + [2] - [2]'$$

$$\begin{aligned}
&\geq \#\{r \in 2 : R : \pi_{\tilde{K}}(r) \neq r\} \cdot [(|\tilde{S}|c_{\Sigma}^{-1/2} - |\tilde{S}^c|Dc_{\Sigma}^{1/2})\Delta - (4|\tilde{S}| + |\tilde{S}^c|)\xi - 4|\tilde{S}|h] \\
&> 0,
\end{aligned}$$

where we used the fact that $|\tilde{S}|/|\tilde{S}^c| \leq \frac{K_0}{K\epsilon}$.

(ii) Case 2: $\pi_{\tilde{K}}(1) \neq 1$.

$$\begin{aligned}
\text{score}(\pi) - \text{score}(\pi^*) &= \underbrace{\sum_{r=2}^R \sum_{k \in \tilde{S}} \|(\hat{\Sigma}^{(k)[0]})^{-1}(\hat{\mu}_r^{(k)[0]} - \hat{\mu}_1^{(k)[0]}) - (\hat{\Sigma}^{(\tilde{K})[0]})^{-1}(\hat{\mu}_{\pi_{\tilde{K}}(r)}^{(\tilde{K})[0]} - \hat{\mu}_{\pi_{\tilde{K}}(1)}^{(\tilde{K})[0]})\|_2}_{[1]} \\
&\quad + \underbrace{\sum_{r=2}^R \sum_{k \in \tilde{S}^c} \|(\hat{\Sigma}^{(k)[0]})^{-1}(\hat{\mu}_{\pi_k(r)}^{(k)[0]} - \hat{\mu}_{\pi_k(1)}^{(k)[0]}) - (\hat{\Sigma}^{(\tilde{K})[0]})^{-1}(\hat{\mu}_{\pi_{\tilde{K}}(r)}^{(\tilde{K})[0]} - \hat{\mu}_{\pi_{\tilde{K}}(1)}^{(\tilde{K})[0]})\|_2}_{[2]} \\
&\quad - \underbrace{\sum_{r=2}^R \sum_{k \in \tilde{S}} \|(\hat{\Sigma}^{(k)[0]})^{-1}(\hat{\mu}_r^{(k)[0]} - \hat{\mu}_1^{(k)[0]}) - (\hat{\Sigma}^{(\tilde{K})[0]})^{-1}(\hat{\mu}_r^{(\tilde{K})[0]} - \hat{\mu}_1^{(\tilde{K})[0]})\|_2}_{[1]'} \\
&\quad - \underbrace{\sum_{r=2}^R \sum_{k \in \tilde{S}^c} \|(\hat{\Sigma}^{(k)[0]})^{-1}(\hat{\mu}_{\pi_k(r)}^{(k)[0]} - \hat{\mu}_{\pi_k(1)}^{(k)[0]}) - (\hat{\Sigma}^{(\tilde{K})[0]})^{-1}(\hat{\mu}_r^{(\tilde{K})[0]} - \hat{\mu}_1^{(\tilde{K})[0]})\|_2}_{[2]'} .
\end{aligned}$$

By previous results,

$$\begin{aligned}
[1] &= \sum_{k \in \tilde{S}} \sum_{r=2}^R \|(\hat{\Sigma}^{(k)[0]})^{-1}(\hat{\mu}_r^{(k)[0]} - \hat{\mu}_1^{(k)[0]}) - (\hat{\Sigma}^{(\tilde{K})[0]})^{-1}(\hat{\mu}_{\pi_{\tilde{K}}(r)}^{(\tilde{K})[0]} - \hat{\mu}_{\pi_{\tilde{K}}(1)}^{(\tilde{K})[0]})\|_2 \\
&\geq \sum_{k \in \tilde{S}} \sum_{r=2}^R \left(\|(\hat{\Sigma}^{(k)[0]})^{-1}(\hat{\mu}_r^{(k)[0]} - \hat{\mu}_1^{(k)[0]}) - (\hat{\Sigma}^{(\tilde{K})[0]})^{-1}(\hat{\mu}_{\pi_{\tilde{K}}(r)}^{(\tilde{K})[0]} - \hat{\mu}_{\pi_{\tilde{K}}(1)}^{(\tilde{K})[0]})\|_2 - 2h - \xi \right) \\
&\geq |\tilde{S}|R \cdot \left(\|(\hat{\Sigma}^{(k)[0]})^{-1}(\hat{\mu}_{\pi_{\tilde{K}}(1)}^{(k)[0]} - \hat{\mu}_1^{(k)[0]})\|_2 - 2h - \xi \right) \\
&\geq |\tilde{S}|R \cdot \left(\|(\Sigma^{(k)*})^{-1}(\mu_{\pi_{\tilde{K}}(1)}^{(k)*} - \mu_1^{(k)*})\|_2 - \xi - 2h - \xi \right) \\
&\geq |\tilde{S}|R \cdot (c_{\Sigma}^{-1/2}\Delta - 2\xi - 2h),
\end{aligned}$$

and

$$-[1]' \geq -|\tilde{S}|R \cdot (2\xi + 2h).$$

Similar to case 1,

$$\begin{aligned}
[2] - [2]' &\geq - \sum_{r=2}^R \sum_{k \in \tilde{S}} \|(\hat{\Sigma}^{(\tilde{K})})^{-1}(\hat{\mu}_{\pi_{\tilde{K}(r)}}^{(\tilde{K})[0]} - \hat{\mu}_{\pi_{\tilde{K}(1)}}^{(\tilde{K})[0]}) - (\hat{\Sigma}^{(\tilde{K})})^{-1}(\hat{\mu}_r^{(\tilde{K})[0]} - \hat{\mu}_1^{(\tilde{K})[0]})\|_2 \\
&\geq -R|\tilde{S}^c| \cdot (2Dc_{\Sigma}^{1/2}\Delta + 2\xi).
\end{aligned}$$

Therefore,

$$\begin{aligned}
\text{score}(\boldsymbol{\pi}) - \text{score}(\tilde{\boldsymbol{\pi}}) &= [1] - [1]' + [2] - [2]' \\
&\geq R[(|\tilde{S}|c_{\Sigma}^{-1/2} - |\tilde{S}^c| \cdot 2Dc_{\Sigma}^{1/2})\Delta - (2|\tilde{S}| + 2|\tilde{S}^c|)\xi - 2|\tilde{S}|h] \\
&> 0,
\end{aligned}$$

where we used the fact that $|\tilde{S}|/|\tilde{S}^c| \leq \frac{K_0}{K\epsilon}$.