Dynamical consistency conditions for rapid turn inflation

Lilia Anguelova^{a1}, Calin Iuliu Lazaroiu^{b2}

^a Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Sofia, Bulgaria

^b Horia Hulubei National Institute for Physics and Nuclear Engineering (IFIN-HH), Bucharest-Magurele, Romania

Abstract

We derive consistency conditions for sustained slow roll and rapid turn inflation in two-field cosmological models with oriented scalar field space, which imply that inflationary models with field-space trajectories of this type are non-generic. In particular, we show that third order adiabatic slow roll, together with large and slowly varying turn rate, requires the scalar potential of the model to satisfy a certain nonlinear second order PDE, whose coefficients depend on the scalar field metric. We also derive consistency conditions for slow roll inflationary solutions in the so called "rapid turn attractor" approximation, as well as study the consistency conditions for circular rapid turn trajectories with slow roll in two-field models with rotationally invariant field space metric. Finally, we argue that the rapid turn regime tends to have a natural exit after a limited number of e-folds.

¹anguelova@inrne.bas.bg

²lcalin@theory.nipne.ro

Contents

1	Introduction	1
2	Two-field cosmological models	4
	2.1 Characteristics of an inflationary solution	6
	2.2 Slow roll and rapid turn regimes	7
	2.3 Some other useful parameters	8
3	The consistency condition for sustained slow roll with rapid turn	9
	3.1 The adapted frame	9
	3.2 Expressing η_{\parallel} and η_{\perp} in terms of θ_{φ} and c	10
	3.3 The condition for sustained rapid turn with third order slow roll	11
4	The case of rotationally invariant metrics	14
	4.1 Slow roll consistency condition for circular trajectories	16
5	A unifying approximation for rapid turn models	19
	5.1 Equations of motion in adapted frame	19
	5.2 Adapted frame parameters	20
	5.3 Consistency conditions	22
6	The characteristic angle	23
7	Conclusions	26
\mathbf{A}	Derivatives of the adapted frame	28

1 Introduction

Modern observations have established – to a very good degree of accuracy – that the present day universe is homogeneous and isotropic on large scales. This is naturally explained if one assumes that the early universe underwent a period of accelerated expansion called inflation. This idea can be realized in models where the inflationary expansion is driven by the potential energy of a number of real scalar fields called inflatons. The most studied models of this type contain a single scalar field. However, recent arguments related to quantum gravity suggest that it is more natural, or may even be necessary [1-4] to have

more than one inflaton. This has generated renewed interest in multi-field cosmological models, which had previously attracted only limited attention.

Multifield cosmological models have richer phenomenology than single field models since they allow for solutions of the equations of motion whose field-space trajectories are not (reparameterized) geodesics. Such trajectories are characterized by a non-zero *turn rate*. In the past it was thought that phenomenological viability requires small turn rate, by analogy with the slow roll approximation used in the single-field case. This assumption leads to the celebrated slow-roll slow-turn (SRST) approximation of [5,6]. However, in recent years it was understood that rapid turn trajectories can also be (linearly) perturbatively stable [7,8] and of phenomenological interest. For instance, a brief rapid turn during slow-roll inflation can induce primordial black hole generation [9–12]; moreover, trajectories with large and constant turn rate can correspond to solutions behaving as dark energy [13, 14]. There is also a variety of proposals for full-fledged rapid-turn inflation models, relying on large turn rates during the entire inflationary period [15–22].

Finding inflationary solutions in multifield models is much harder than in the singlefield case, because the background field equations form a complicated coupled system of nonlinear ODEs. Thus usually such models are either studied numerically or solved only approximately.³ Mathematically, this complicated coupled system is encoded by the so-called *cosmological equation*, a nonlinear second order geometric ODE defined on the scalar field space of the model. The latter is a connected paracompact manifold, usually called the *scalar manifold*. In turn, the cosmological equation is equivalent with a dissipative geometric dynamical system defined on the tangent bundle of that manifold. Little is known in general about this dynamical system, in particular because the scalar manifold need not be simply-connected and – more importantly – because this manifold is non-compact in most applications of physical interest and hence cosmological trajectories can "escape to infinity". The resulting dynamics can be surprisingly involved⁴ and hard to analyze even by numerical methods (see [28–30] for some nontrivial examples in two-field

 $^{^{3}}$ A notable exception is provided by models with hidden symmetry, which greatly facilitates the search for exact solutions [23–27].

⁴It is sometimes claimed that the complexity of this dynamics could be ignored, because in "phenomenologically relevant models" one should "expect that" all directions orthogonal to the physically relevant scalar field trajectory are heavy and hence can be integrated out, thus reducing the analysis to that of a single-field model. This argument is incorrect for a number of reasons. First, current phenomenological data does *not* rule out multifield dynamics. Second, such a reduction to a one field model (even when possible) relies on knowledge of an appropriate cosmological trajectory, which itself must first be found by analyzing the dynamics of the multifield model.

models), though a conceptual approach to some aspects of that dynamics was recently proposed in [31, 32].

A common approach to looking for cosmological trajectories with desirable properties is to first simplify the equations of motion by imposing various approximations (such as slow-roll to a certain order, rapid-turn and/or other conditions). This leads to an approximate system of equations, obtained by neglecting certain terms in the original ODEs. Then one attempts to solve the approximate system numerically or analytically. However, there is apriori no guarantee that a solution of the approximate system is a good approximant of a solution of the exact system for a sufficiently long period of time. In general, this will be the case only if the data which parameterizes the exact system (namely the scalar field metric and scalar potential of the model) satisfies appropriate *consistency conditions*. Despite being of fundamental conceptual importance, such consistency conditions have so far not been studied systematically in the literature.

In this paper, we investigate the problem of consistency conditions in two-field models with orientable scalar manifold for several commonly used approximations. First, we consider third order slow-roll trajectories with large but slowly varying turn rate. In this case, we show that compatibility with the equations of motion requires that the scalar potential satisfies a certain nonlinear second order PDE whose coefficients depend on the scalar field metric. This gives a nontrivial and previously unknown consistency condition that must be satisfied in the field-space regions where one can expect to find sustained rapid-turn trajectories allowing for slow roll inflation. Therefore, inflationary solutions of this type are not easy to find, implying that two-field models with such families of cosmological trajectories are non-generic. In particular, this shows that the difficulty in finding such models which was noticed in [33] is not related to supergravity, but arises on a more basic level.

We also discuss the case of rotationally invariant scalar field-space metrics. In that case, it is common to consider field space trajectories which are nearly circular as candidates for sustained rapid turn inflation. Imposing the first and second order slow roll conditions in this context leads to a certain consistency condition for compatibility with the equations of motion. This is again a PDE for the scalar potential with coefficients depending on the field space metric, which does not seem to have been widely noticed in the literature. We consider its implications for important examples in previous work.

Finally, we study in detail the consistency conditions for the approximation of [20], which subsumes many prominent rapid turn models of inflation. We show that this approximation is a special case of rapid turn with slow roll, instead of being equivalent to it. We then derive conditions for compatibility of this approximation with the equations of motion. Once again, these constrain the scalar potential and field space metric.

Throughout the paper, we assume that the field space (a.k.a. scalar manifold) of the two-field model is an *oriented* and connected paracompact surface. In our considerations, a crucial role is played by a fixed oriented frame (called the *adapted frame*) of vector fields defined on this surface which is determined by the scalar potential and field-space metric, instead of the moving oriented Frenet frame determined by the field space trajectory. The two frames are related to each other through a time-dependent rotation whose time-dependent angle we call the *characteristic angle*. We conclude our investigations by studying the time evolution of this angle. We show that, generically, this angle tends rather fast to the value $\pi \mod 2\pi$, which means that the tangent vector of the inflationary trajectory aligns with minus the gradient of the potential. This implies that the rapid turn regime has a natural exit in the generic case. Our results also suggest that it is difficult to sustain this regime for a prolonged period.

The paper is organized as follows. Section 2 recalls some basic facts about twofield cosmological models and introduces various parameters which will be used later on. Section 3 discusses the consistency condition for sustained rapid turn trajectories with third order slow roll, which results from careful analysis of the compatibility of the corresponding approximations with the equations of motion. Section 4 discusses the consistency condition for circular trajectories in rotationally-invariant models, as well as some implications for previous work on such inflationary trajectories. Section 5 discusses the approximation of [20], showing how it differs from rapid turn with second order slow roll and extracts the relevant consistency conditions. Section 6 studies the time evolution of the characteristic angle, while Section 7 presents our conclusions.

2 Two-field cosmological models

The action for *n* real scalar fields $\varphi^{I}(x^{\mu})$ minimally coupled to four-dimensional gravity is:

$$S = \int d^4x \sqrt{-\det g} \left[\frac{R(g)}{2} - \frac{1}{2} G_{IJ}(\{\varphi^I\}) \partial_\mu \varphi^I \partial^\mu \varphi^J - V(\{\varphi^I\}) \right] \quad , \tag{2.1}$$

where we took $M_{\rm Pl} = 1$. Here $g_{\mu\nu}$ is the spacetime metric (which we take to have "mostly plus" signature) and R(g) is its scalar curvature. The indices μ, ν run from from 0 to 3 and G_{IJ} is the metric on the scalar manifold (a.k.a. "scalar field space") \mathcal{M} , which is parameterized by the scalars $\{\varphi^I\}$ with I, J = 1, ..., n. The scalar potential V is a smooth real-valued function defined on \mathcal{M} , which for simplicity we take to be strictly positive. The standard cosmological Ansatze for the background metric and scalar fields are:

$$ds_g^2 = -dt^2 + a^2(t)d\vec{x}^2$$
 , $\varphi^I = \varphi^I(t)$, (2.2)

where a(t) > 0 is the scale factor. As usual, the definition of the Hubble parameter is:

$$H(t) = \frac{\dot{a}}{a} \quad , \tag{2.3}$$

where the dot denotes derivation with respect to t.

With the Ansatze above, the equations of motion for the scalars reduce to:

$$D_t \dot{\varphi}^I + 3H \dot{\varphi}^I + G^{IJ} V_J = 0 \quad , \tag{2.4}$$

where $V_J \stackrel{\text{def.}}{=} \partial_J V$ with $\partial_J \stackrel{\text{def.}}{=} \partial_{\varphi^J}$ and we defined:

$$D_t A^I \stackrel{\text{def.}}{=} \dot{\varphi}^J \nabla_J A^I = \dot{A}^I + \Gamma^I_{JK}(\varphi) \dot{\varphi}^J A^K$$

for any vector field A^{I} defined on the scalar manifold, where Γ_{JK}^{I} are the Christoffel symbols of G_{IJ} . The Einstein equations can be written as:

$$G_{IJ}\dot{\varphi}^{I}\dot{\varphi}^{J} = -2\dot{H}$$
 , $3H^{2} + \dot{H} = V$. (2.5)

In this paper we focus on two-field models, i.e. we take n = 2. In this case, the scalar manifold $(\mathcal{M}, \mathcal{G})$ is a (generally non-compact) Riemann surface, which we assume to be oriented. To define various characteristics of an inflationary solution, one introduces a frame of tangent vectors to the field space. A widely used choice is the positive Frenet frame, which consists of the tangent vector T(t) and normal vector N(t) to the trajectory $\varphi(t) = (\varphi^1(t), \varphi^2(t)) \in \mathcal{M}$ of a solution of (2.4), where the normal vector is chosen such that (T(t), N(t)) forms a positively-oriented basis of the tangent space of the scalar manifold \mathcal{M} at the point $\varphi(t)$. Let σ be an increasing proper length parameter along the solution curve φ . This parameter is determined up to a constant translation and satisfies:

$$\dot{\sigma} = \sqrt{G_{IJ}\dot{\varphi}^I\dot{\varphi}^J}$$

Then the vectors T and N are given by:

$$T^{I} = \frac{\mathrm{d}\varphi^{I}}{\mathrm{d}\sigma} = \frac{\dot{\varphi}^{I}}{\dot{\sigma}}$$

and

$$N_I = (\det G)^{1/2} \epsilon_{IJ} T^J \quad . \tag{2.6}$$

One has:

$$N_I T^I = 0$$
 , $N_I N^I = 1$, $T_I T^I = 1$, (2.7)

where $T_I = G_{IJ}T^J$, $N^I = G^{IJ}N_J$.

2.1 Characteristics of an inflationary solution

Consider the opposite relative acceleration vector:

$$\eta^{I} \stackrel{\text{def.}}{=} -\frac{1}{H\dot{\sigma}} D_{t} \dot{\varphi}^{I} \quad . \tag{2.8}$$

Expanding η in the orthonormal basis (T, N) gives:

$$\eta^I = \eta_{\parallel} T^I + \eta_{\perp} N^I \;\;,$$

with:

$$\eta_{\parallel} = -\frac{\ddot{\sigma}}{H\dot{\sigma}} \quad \text{and} \quad \eta_{\perp} = \frac{\Omega}{H} ,$$
 (2.9)

where we defined the *signed turn rate* of the trajectory φ by:

$$\Omega \stackrel{\text{def.}}{=} -N_I D_t T^I \quad . \tag{2.10}$$

The quantity η_{\parallel} is called the *second slow roll parameter* of φ , while η_{\perp} (which is sometimes denoted by ω) is called the *first turn parameter* (or signed reduced turn rate). We will see in a moment that η_{\parallel} is the second slow roll parameter of a certain one-field model determined by the trajectory φ . The *second turn parameter* is defined through:

$$\nu \stackrel{\text{def.}}{=} \frac{\dot{\eta}_{\perp}}{H\eta_{\perp}} \quad . \tag{2.11}$$

It is a measure for the rate of change per e-fold of the dimensionless turn parameter η_{\perp} .

Projecting the scalar field equations (2.4) along T^{I} gives the *adiabatic equation*:

$$\ddot{\sigma} + 3H\dot{\sigma} + V_T = 0 \quad , \tag{2.12}$$

where $V_T \stackrel{\text{def.}}{=} T^I V_I$. Projecting (2.4) along N_I gives the *entropic equation*:

$$N_I D_t T^I = -\frac{V_N}{\dot{\sigma}} \quad , \tag{2.13}$$

where $V_N \stackrel{\text{def.}}{=} N^I V_I$. Using the definition (2.10), the entropic equation (2.13) reads:

$$\Omega = \frac{V_N}{\dot{\sigma}} \quad . \tag{2.14}$$

Let $V_{\varphi}(\sigma) \stackrel{\text{def.}}{=} V(\varphi(\sigma))$. Then:

$$V_T = rac{\mathrm{d}V_{arphi}}{\mathrm{d}\sigma} = V_{arphi}'(\sigma) \ ,$$

where the prime denotes derivation with respect to σ . Hence the adiabatic equation (2.12) reads:

$$\ddot{\sigma} + 3H\dot{\sigma} + V'_{\varphi}(\sigma) = 0 \ ,$$

which shows that σ obeys the equation of motion of a one-field model with scalar potential V_{φ} . Since for an expanding universe we have H > 0, the Hubble parameter can be expressed from (2.5) as:

$$H(\sigma, \dot{\sigma}) = \frac{1}{\sqrt{6}} \sqrt{\dot{\sigma}^2 + 2V(\sigma)}$$

which is the usual formula for a single-field model.

The *adiabatic Hubble slow roll parameters* of the trajectory φ are defined as the usual Hubble slow roll-parameters [34] of the adiabatic one-field model defined above. The first three of these are:

$$\varepsilon \stackrel{\text{def.}}{=} -\frac{\dot{H}}{H^2} \quad , \quad \eta_{\parallel} = -\frac{\ddot{\sigma}}{H\dot{\sigma}} \quad , \quad \xi \stackrel{\text{def.}}{=} \frac{\ddot{\sigma}}{H^2\dot{\sigma}} \quad , \quad (2.15)$$

where for ξ we use a different sign convention than [34]. Since all results of [34] apply to the adiabatic one field models, it follows that the following relations hold on-shell:

$$\begin{aligned} \epsilon_T &= \varepsilon \left(\frac{3 - \eta_{\parallel}}{3 - \varepsilon} \right)^2 \\ \eta_{\parallel} &= \varepsilon - \frac{\varepsilon'}{\sqrt{2\varepsilon}} \\ \eta_T &= \sqrt{2\varepsilon} \frac{\eta'_{\parallel}}{3 - \varepsilon} + \frac{3 - \eta_{\parallel}}{3 - \varepsilon} (\varepsilon + \eta_{\parallel}) \end{aligned}$$

where the adiabatic first and second *potential slow roll parameters* are defined through:

$$\epsilon_T \stackrel{\text{def.}}{=} \frac{1}{2} \left(\frac{V_T}{V} \right)^2 \quad , \quad \eta_T \stackrel{\text{def.}}{=} \frac{V_{\varphi}''}{V} \quad .$$

2.2 Slow roll and rapid turn regimes

The first, second and third slow roll regimes are defined respectively by the conditions $\varepsilon \ll 1$, $|\eta_{\parallel}| \ll 1$ and $|\xi| \ll 1$. The second *order* slow roll regime is defined by the two conditions $\varepsilon \ll 1$ and $|\eta_{\parallel}| \ll 1$ taken together, while the third *order* slow roll regime is defined by the three conditions $\varepsilon \ll 1$, $|\eta_{\parallel}| \ll 1$ and $|\xi| \ll 1$ considered together.

The first order rapid turn regime is defined by the condition:

$$\eta_{\perp}^2 \gg 1 \quad , \tag{2.16}$$

while the condition $|\nu| \ll 1$ defines the second order rapid turn regime. A trajectory has sustained rapid turn if both of the conditions $\eta_{\perp}^2 \ll 1$ and $|\nu| \ll 1$ are satisfied.

In the first slow roll regime $\varepsilon \ll 1$, the second equation in (2.5) becomes:

$$3H^2 \approx V$$
 . (2.17)

Together with (2.14), this gives:

$$\eta_{\perp}^2 = \frac{\Omega^2}{H^2} \approx \frac{3V_N^2}{\dot{\sigma}^2 V} \quad . \tag{2.18}$$

2.3 Some other useful parameters

For our purpose, it will be useful to consider two other parameters, which are related to the familiar ones reviewed above. The *first IR parameter* is the ratio of the kinetic and potential energies of the scalars:

$$\kappa \stackrel{\text{def.}}{=} \frac{G_{IJ}\dot{\varphi}^I \dot{\varphi}^J}{2V} = \frac{\dot{\sigma}^2}{2V} \quad . \tag{2.19}$$

This parameter plays an important role in the IR approximation discussed in [31] and [32]. On solutions of the equations of motion, the first slow roll parameter can be written as:

$$\varepsilon = \frac{3 \dot{\sigma}^2}{\dot{\sigma}^2 + 2V} \ .$$

Hence the on-shell value of the first slow roll parameter is related to the first IR parameter through:

$$\varepsilon = \frac{3\kappa}{1+\kappa} \quad . \tag{2.20}$$

This relation shows that the first slow roll condition $\varepsilon \ll 1$ is equivalent with $\kappa \ll 1$, whereas $\varepsilon = 1$ corresponds to $\kappa = \frac{1}{2}$. In particular, the condition for inflation $\varepsilon < 1$ is equivalent with $\kappa < \frac{1}{2}$.

We also define the *conservative parameter*:

$$c \stackrel{\text{def.}}{=} \frac{H\dot{\sigma}}{\sqrt{G^{IJ}V_I V_J}} \quad , \tag{2.21}$$

which is small iff the friction term can be neglected relative to the gradient term in the equation of motion for the scalar fields. When $c \ll 1$, the motion of φ is approximately conservative in the sense that the total energy $\frac{1}{2}G_{IJ}\dot{\varphi}^{I}\dot{\varphi}^{J}$ of the scalar fields is approximately conserved; in this limit, the equations of motion for the scalars reduce to:

$$\nabla_t \varphi^I + G^{IJ} \partial_J V \approx 0 \quad ,$$

which describe the motion of a particle in the Riemannian manifold $(\mathcal{M}, \mathcal{G})$ under the influence of the potential V. We will see below that, in the second slow roll regime $|\eta_{\parallel}| \ll 1$, the rapid turn condition (2.16) is equivalent with the *conservative condition*:

$$c^2 \ll 1$$
 .

3 The consistency condition for sustained slow roll with rapid turn

In this section, we derive a constraint relating the scalar potential V and the field-space metric G_{IJ} which is necessary for existence of rapid turn solutions $(\eta_{\perp}^2 \gg 1)$ that satisfy the *third order* slow roll conditions $\varepsilon \ll 1$, $|\eta_{\parallel}| \ll 1$ and $|\xi| \ll 1$ as well as the second order condition $|\nu| \ll 1$ for slowly-varying turn rate. The last condition is usually imposed to ensure the longevity of the rapid-turn slow-roll regime (so that it could produce about 50-60 e-folds of inflation). The constraint we derive does not require any extraneous assumptions about the potential, the field space metric or the shape of the field-space trajectory.

3.1 The adapted frame

We will use a globally-defined frame (n, τ) (which we call the adapted frame) of the scalar manifold determined by V and G_{IJ} . Namely, we take n to be the unit vector field along the gradient of the potential:

$$n \stackrel{\text{def.}}{=} \frac{\nabla V}{\sqrt{G^{IJ}V_IV_J}}$$
, with components $n^K = \frac{G^{KL}V_L}{\sqrt{G^{IJ}V_IV_J}}$. (3.1)

The unit vector τ is orthogonal to n and chosen such that the basis (n, τ) is positively oriented:

$$\tau_I = (\det G)^{1/2} \epsilon_{IJ} n^J \quad . \tag{3.2}$$

We have $\tau^{I} \stackrel{\text{def.}}{=} G^{IJ} \tau_{J}$ as usual. The relation between the two bases is given by:

$$T = \cos \theta_{\varphi} n + \sin \theta_{\varphi} \tau ,$$

$$N = -\sin \theta_{\varphi} n + \cos \theta_{\varphi} \tau ,$$
(3.3)

where $\theta_{\varphi} \in (-\pi, \pi]$ is the *characteristic angle* of φ , which is defined as the angle of the rotation that takes the oriented basis (n, τ) to the oriented basis (T, N) at the point of interest on the cosmological trajectory.

3.2 Expressing η_{\parallel} and η_{\perp} in terms of θ_{φ} and c

Relation (2.9) and the adiabatic equation (2.12) give:

$$\eta_{\parallel} = 3 + \frac{V_{\sigma}}{H\dot{\sigma}} = 3 + \frac{T^I G_{IJ} n^J}{H\dot{\sigma}} \sqrt{G^{KL} V_K V_L} = 3 + \frac{\cos \theta_{\varphi}}{c} \quad , \tag{3.4}$$

where the conservative parameter c was defined in (2.21). Notice that $\eta_{\parallel} \approx 3$ for either $\cos \theta_{\varphi} \approx 0$ or $c \gg 1$. Also, the second slow roll condition $|\eta_{\parallel}| \ll 1$ requires that c < 1 and $\cos \theta_{\varphi} < 0$. In fact, $\eta_{\parallel} \approx 0$ is achieved for $c \approx -\cos \theta_{\varphi}/3$.

We next consider equation (2.9) for η_{\perp} . Using the entropic equation (2.14) and substituting N from (3.3) gives:

$$\eta_{\perp} = \frac{\Omega}{H} = \frac{N_I V^I}{\dot{\sigma} H} = -\frac{(G^{KL} V_K V_L)^{\frac{1}{2}} \sin \theta_{\varphi}}{\dot{\sigma} H} = -\frac{\sin \theta_{\varphi}}{c} \quad . \tag{3.5}$$

In the second slow roll approximation we have $c \approx -\cos \theta_{\varphi}/3$ (as explained above) and equation (3.5) gives:

$$\eta_{\perp}^2 \approx 9 \, \tan^2 \theta_{\varphi} \ . \tag{3.6}$$

Hence the rapid turn condition $\eta_{\perp}^2 \gg 1$ is equivalent with $\tan^2 \theta_{\varphi} \gg 1$ or, equivalently, with $\cos^2 \theta_{\varphi} \ll 1$. In view of (3.4), it follows that the second slow roll condition $|\eta_{\parallel}| \ll 1$ requires $c^2 \ll 1$. Note that *c* itself does not need to be very small. It is enough to have $c \sim \mathcal{O}(10^{-1})$, in order to ensure rapid turn and slow roll.

The discussion above shows that the slow roll and rapid turn regime, which is usually defined by:

$$\varepsilon \ll 1$$
 , $|\eta_{\parallel}| \ll 1$, $\eta_{\perp}^2 \gg 1$, (3.7)

is characterized equivalently by:

$$\varepsilon \ll 1$$
 , $|\eta_{\parallel}| \ll 1$, $c^2 \ll 1$. (3.8)

We next derive the consistency condition on V and G_{IJ} in this regime, supplemented by the additional requirements that:

$$|\xi| \ll 1 \quad \text{and} \quad |\nu| \ll 1 \quad , \tag{3.9}$$

where ξ is the third order slow roll parameter in (2.15), while ν is the relative rate of change of η_{\perp} as defined in (2.11). The purpose of the second condition in (3.9) is to ensure that the inequality (2.16) is satisfied for a prolonged period. (Note that $|\nu| \ll 1$ does not follow from the slow roll conditions in (3.7).)

3.3 The condition for sustained rapid turn with third order slow roll

The equations of motion (2.12) and (2.14) respectively imply (see [35-37]):

$$\frac{V_{TT}}{3H^2} = \frac{\Omega^2}{3H^2} + \varepsilon + \eta_{\parallel} - \frac{\xi}{3} ,$$

$$\frac{V_{TN}}{H^2} = \frac{\Omega}{H} \left(3 - \varepsilon - 2\eta_{\parallel} + \nu\right) .$$
(3.10)

We will study the consequences of these relations for inflation with third order slow roll, defined by:

$$\varepsilon, |\eta_{\parallel}|, |\xi| \ll 1$$
, (3.11)

while also imposing the condition:

$$|\nu| \ll 1 \quad . \tag{3.12}$$

The latter ensures a small rate of change of the large turn rate and hence longevity of the rapid turn regime.

In the regime defined by (3.11)-(3.12), equations (3.10) imply:

$$V_{TT} \approx \frac{1}{9} \frac{V_{TN}^2}{H^2}$$
 (3.13)

Recalling that $3H^2 \approx V$ (see (2.17)), relation (3.13) can be written as:

$$V_{TT} \approx \frac{1}{3} \frac{V_{TN}^2}{V} \quad . \tag{3.14}$$

To extract conditions on V and G_{IJ} , let us rewrite (3.14) in terms of V_{nn} , $V_{n\tau}$ and $V_{\tau\tau}$. Using (3.3), we have:

$$V_{TT} = T^{I}T^{J}\nabla_{I}V_{J} = V_{nn}\cos^{2}\theta_{\varphi} + 2V_{n\tau}\sin\theta_{\varphi}\cos\theta_{\varphi} + V_{\tau\tau}\sin^{2}\theta_{\varphi} ,$$

$$V_{TN} = T^{I}N^{J}\nabla_{I}V_{J} = (V_{\tau\tau} - V_{nn})\cos\theta_{\varphi}\sin\theta_{\varphi} + V_{n\tau}\left(\cos^{2}\theta_{\varphi} - \sin^{2}\theta_{\varphi}\right) . \quad (3.15)$$

With the approximations (3.8), one has $\cos \theta_{\varphi} \approx -3c$ due to (3.4). In that case $\sin \theta_{\varphi} \approx s\sqrt{1-9c^2}$, where $s = \pm 1$. Hence (3.15) becomes:

$$V_{TT} \approx 9c^2 V_{nn} - 6sc \sqrt{1 - 9c^2} V_{n\tau} + (1 - 9c^2) V_{\tau\tau} ,$$

$$V_{TN} \approx -3sc \sqrt{1 - 9c^2} (V_{\tau\tau} - V_{nn}) - (1 - 18c^2) V_{n\tau} .$$
(3.16)

Since the different components of the Hessian of V may be of different orders in the small parameter c, we cannot conclude from (3.16) that $V_{TT} \approx V_{\tau\tau}$ and $V_{TN} \approx -V_{n\tau}$. Instead,

we need two relations between V_{nn} , $V_{n\tau}$, $V_{\tau\tau}$ and c, in order to be able to solve for c and then extract a relation, that involves only the components of the Hessian of V.

To achieve this, let us compute (3.10) in terms of c in the regime (3.8):

$$\frac{V_{TT}}{3H^2} \approx \frac{\Omega^2}{3H^2} = \frac{\sin^2 \theta_{\varphi}}{3c^2} \approx \frac{1}{3c^2} - 3 ,$$

$$\frac{V_{TN}}{H^2} \approx 3\frac{\Omega}{H} = -\frac{3\sin\theta_{\varphi}}{c} \approx -\frac{3s}{c}\sqrt{1-9c^2} , \qquad (3.17)$$

where we used (3.5) together with $\sin \theta_{\varphi} \approx s\sqrt{1-9c^2}$. Since $3H^2 \approx V$ during slow roll, we have:

$$V_{TT} \approx \frac{V}{3c^2} - 3V$$
 and $V_{TN} \approx -\frac{s}{c}\sqrt{1 - 9c^2} V$. (3.18)

These expressions satisfy (3.14) automatically, as should be the case. Now substitute (3.16) in (3.18). This gives the following two relations:

$$\frac{V}{3c^2} - 3V \approx 9c^2 V_{nn} - 6sc\sqrt{1 - 9c^2} V_{n\tau} + (1 - 9c^2)V_{\tau\tau} ,$$

$$\frac{s}{c}\sqrt{1 - 9c^2} V \approx 3sc\sqrt{1 - 9c^2} (V_{\tau\tau} - V_{nn}) + (1 - 18c^2)V_{n\tau} .$$
(3.19)

Our next goal is to eliminate the small parameter c without making any assumptions about the order of any of the quantities V, V_{nn} , $V_{n\tau}$, $V_{\tau\tau}$. We begin by solving for $V_{n\tau}$ from the second equation in (3.19):

$$V_{n\tau} \approx \frac{s\sqrt{1-9c^2}}{(1-18c^2)} \left[\frac{V}{c} - 3c \left(V_{\tau\tau} - V_{nn} \right) \right] .$$
 (3.20)

Substituting this in the first equation of (3.19) gives:

$$(1 - 18c^2) \left(\frac{V}{3c^2} - 3V\right) \approx -6V + 54c^2V - 9c^2V_{nn} + (1 - 9c^2)V_{\tau\tau} \quad (3.21)$$

where we did *not* drop any term containing higher powers of c; at intermediate steps there are terms with c^4 , as well as other c^2 terms, but they all cancel exactly. Since $(1-18c^2)\left(\frac{V}{3c^2}-3V\right)=\frac{V}{3c^2}-3V-6V+54c^2V$ relation (3.21) becomes:

$$\frac{V}{3c^2} - 3V \approx -9c^2 V_{nn} + (1 - 9c^2) V_{\tau\tau} \; .$$

Thus:

$$V_{\tau\tau} \approx \frac{\frac{V}{3c^2} - 3V + 9c^2 V_{nn}}{1 - 9c^2} .$$
 (3.22)

Note that this expression for $V_{\tau\tau}$ was obtained without neglecting any terms compared to (3.19). From (3.22), we have:

$$V_{\tau\tau} \approx \frac{V}{3c^2} + 9c^2 V_{nn} + \mathcal{O}(c^4)$$
 (3.23)

Substituting (3.22) in (3.20), we find:

$$V_{n\tau} \approx \frac{3scV_{nn}}{\sqrt{1-9c^2}} \quad , \tag{3.24}$$

where the terms with V canceled exactly. We stress that the result (3.24) was obtained without neglecting any further terms, just like (3.22). Now recall that in the rapid turn regime we have $c^2 \ll 1$. Hence in this regime equations (3.22) and (3.24) imply:

$$V_{\tau\tau} \approx \frac{V}{3c^2}$$
 and $V_{n\tau} \approx 3scV_{nn}$. (3.25)

The second relation gives:

$$c \approx \frac{s}{3} \frac{V_{n\tau}}{V_{nn}} \quad . \tag{3.26}$$

Substituting this into the first relation of (3.25) gives:

$$3VV_{nn}^2 \approx V_{n\tau}^2 V_{\tau\tau} \quad . \tag{3.27}$$

Since V > 0, the last relation implies, in particular, that we have:

$$V_{\tau\tau} > 0$$
 . (3.28)

In view of (3.23), the Hessian of V takes the following form to order c^2 :

$$\begin{bmatrix} V_{\tau\tau} & V_{n\tau} \\ V_{n\tau} & V_{nn} \end{bmatrix} \approx \begin{bmatrix} 9c^2V_{nn} + \frac{V}{3c^2} & 3scV_{nn} \\ 3scV_{nn} & V_{nn} \end{bmatrix}$$

The eigenvalues of the characteristic polynomial of this matrix are:

$$\lambda_1 \approx V_{nn}$$
 and $\lambda_2 \approx \frac{V}{3c^2}$,

where we used $c^2 \ll 1$. Hence the condition to have non-negative eigenvalues is:

$$V_{nn} \ge 0$$
 .

In view of (3.25), the second eigenvalue is given by $\lambda_2 \approx V_{\tau\tau}$. This is consistent with the fact that (3.26) implies (to leading order in the slow roll and rapid turn regime) that the Hessian of V is diagonal in the basis (n, τ) .

From (3.27) we have:

$$V_{\tau\tau} \approx 3V \frac{V_{nn}^2}{V_{n\tau}^2} \quad . \tag{3.29}$$

Let us compare this with the expression for $V_{\tau\tau}$ in (3.23). Substituting (3.26) in (3.23) gives:

$$V_{\tau\tau} \approx \frac{V}{3c^2} + 9c^2 V_{nn} \approx 3V \frac{V_{nn}^2}{V_{n\tau}^2} + \frac{V_{n\tau}^2}{V_{nn}} \quad . \tag{3.30}$$

Using (3.29) and (3.30) we conclude that:

$$V_{\tau\tau}V_{nn} \gg V_{n\tau}^2 \quad . \tag{3.31}$$

Notice that the consistency condition (3.27) depends not only on V but also on the fieldspace metric G_{IJ} , as is clear from the definitions of the basis vectors n and τ .

Finally, let us discuss the special case when $V_{n\tau} = 0$. In this case, we must reconsider (3.19), since (3.26) gives $c \approx 0$. So let us take $V_{n\tau} = 0$ in (3.19):

$$\frac{V}{3c^2} - 3V \approx 9c^2 V_{nn} + (1 - 9c^2) V_{\tau\tau} ,$$

$$\frac{s}{c} \sqrt{1 - 9c^2} V \approx 3sc \sqrt{1 - 9c^2} (V_{\tau\tau} - V_{nn}) . \qquad (3.32)$$

The second relation gives:

$$V_{\tau\tau} pprox rac{V}{3c^2} + V_{nn}$$
 .

Substituting this into the first equation in (3.32) we obtain:

$$V_{nn} \approx 0 \quad , \tag{3.33}$$

where we did not neglect any terms; everything except V_{nn} canceled exactly. Thus, the consistency condition (3.27) is satisfied again, although relation (3.31) is not valid anymore.

To recapitulate the considerations of this Section: We have shown that (3.27) is a necessary condition for the existence of a prolonged inflationary period in the slow-roll and rapid-turn regime defined by (3.7).

4 The case of rotationally invariant metrics

In the literature on cosmological inflation the field-space metric G_{IJ} is often assumed to be rotationally invariant. Partly, this is motivated by the desire for simplification. It is also the case that many inflationary models which arise from string theory compactifications have a scalar field metric of this type. In view of this, we will now specialize the consistency conditions derived in the previous Section to the case of rotationally invariant metrics. For such metrics, the consistency condition extracted in the previous section can be made explicit upon using local semigeodesic coordinates on the scalar manifold $(\mathcal{M}, \mathcal{G})$, in which the metric has the form:

$$ds_G^2 = dr^2 + h^2(r)d\theta^2$$
(4.1)

with h(r) > 0. In such coordinates, the non-vanishing Christoffel symbols are:

$$\Gamma^{r}_{\theta\theta} = -hh' \quad , \quad \Gamma^{\theta}_{r\theta} = \frac{h'}{h} \quad , \qquad (4.2)$$

where the prime denotes derivation with respect to r. Let us compute the components of the Hessian, which enter the condition (3.27). Using (3.1), (3.2) and (4.1), we obtain:

$$V_{nn} = n^{I} n^{J} \nabla_{I} \nabla_{J} V$$

$$= \frac{1}{(h^{2} V_{r}^{2} + V_{\theta}^{2})} \left[h^{2} V_{r}^{2} V_{rr} + 2 V_{r} V_{\theta} V_{r\theta} - \frac{h'}{h} V_{r} V_{\theta}^{2} + \frac{1}{h^{2}} V_{\theta}^{2} V_{\theta\theta} \right], \qquad (4.3)$$
$$V_{n\tau} = n^{I} \tau^{J} \nabla_{I} \nabla_{I} \nabla_{I} V$$

$$V_{n\tau} = h T \nabla T \nabla J V$$

$$= \frac{h}{(h^2 V_r^2 + V_\theta^2)} \left[V_r V_\theta \left(V_{rr} - \frac{1}{h^2} V_{\theta\theta} \right) + \left(\frac{1}{h^2} V_\theta^2 - V_r^2 \right) V_{r\theta} - \frac{h'}{h^3} V_\theta^3 \right], \quad (4.4)$$

$$V_{\tau\tau} = \tau^I \tau^J \nabla_I \nabla_I V$$

$$= \frac{h^4}{(h^2 V_r^2 + V_{\theta}^2)} \left[V_{\theta}^2 V_{rr} - 2V_r V_{\theta} \left(V_{r\theta} - \frac{h'}{h} V_{\theta} \right) + V_r^2 (V_{\theta\theta} + hh' V_r) \right].$$
(4.5)

The result from substituting (4.3)-(4.5) in (3.27) is too messy to be illuminating without further specialization. However, the very existence of this highly nontrivial relation implies that it is not easy to achieve a long lasting period of rapid turn with slow roll. In other words, one can expect that obtaining at least 50-60 e-folds of slow-roll rapid-turn inflation is possible only for special choices of scalar potentials and field-space metrics.

Expressions (4.3)-(4.5) simplify enormously for θ -independent scalar potentials V. In that case, one finds:

$$V_{nn} = V_{rr}$$
 , $V_{n\tau} = 0$, $V_{\tau\tau} = h^3 h' V_r$

If we assume that $V_{rr} \neq 0$ then the consistency condition (3.27) cannot be satisfied anywhere in field space, regardless of the specifics of the model. This agrees with the conclusion, reached in (3.33) for the case with $V_{n\tau} = 0$, that V_{nn} has to vanish in that case for consistency.

The discussion in this section shows in particular that sustained rapid turn with third order slow roll cannot be realized in models with a rotationally invariant scalar potential, regardless of the form of the field-space trajectory.⁵

⁵Note that in the solutions of [14] (which arise from a potential of the form V = V(r)) one has

4.1 Slow roll consistency condition for circular trajectories

An Ansatz that is often used in the literature to look for sustained slow-roll and rapidturn solutions which might lead to the phenomenologically desirable 50-60 or so e-folds of inflation consists of taking $V_{\theta} \neq 0$ and considering near-circular field-space trajectories. Thus one assumes $r \approx \text{const}$, which more precisely means that one approximates $\dot{r} \approx 0$ and $\ddot{r} \approx 0$. This approximation was used for example in [33] as well as in many other references on rapid turn inflation.

However, there is a consistency condition for compatibility of this approximation with the equations of motion. To see this, let us first specialize the form of the field equations to the case of a rotationally invariant field space metric. For G_{IJ} given by (4.1), the equations for the scalar fields (2.4) become:

$$\ddot{r} - hh'\dot{\theta}^2 + 3H\dot{r} + V_r = 0 , \qquad (4.6)$$

$$\ddot{\theta} + 2\frac{h'}{h}\dot{r}\dot{\theta} + 3H\dot{\theta} + \frac{1}{h^2}V_{\theta} = 0 . \qquad (4.7)$$

We next approximate $\dot{r} \approx 0$ and $\ddot{r} \approx 0$. In this case, the second slow roll parameter takes the form:

$$\eta_{\parallel} = -\frac{\dot{r}\ddot{r} + hh'\dot{r}\dot{\theta}^2 + h^2\dot{\theta}\ddot{\theta}}{H(\dot{r}^2 + h^2\dot{\theta}^2)} \approx -\frac{\ddot{\theta}}{H\dot{\theta}} .$$

$$\tag{4.8}$$

Imposing the slow roll conditions $\varepsilon \ll 1$ and $|\eta_{\parallel}| \ll 1$ within the approximations $\dot{r} \approx 0$ and $\ddot{r} \approx 0$ for a sustained near-circular trajectory, we find that (4.6)-(4.7) reduce to:⁶

$$-hh'\dot{\theta}^2 + V_r \approx 0 \tag{4.9}$$

$$3H\dot{\theta} + \frac{V_{\theta}}{h^2} \approx 0 . aga{4.10}$$

From the second equation in (2.5) we also have:

$$3H^2 \approx V$$
 , (4.11)

because $\varepsilon \ll 1$ in the slow roll regime. Notice that (4.10) does not admit solutions with $\dot{\theta} \neq 0$ for rotationally invariant potentials. Equation (4.9) gives:

$$\dot{\theta}^2 \approx \frac{V_r}{hh'}$$
, (4.12)

 $\overline{\eta_{\parallel}^{DE}} \sim \mathcal{O}(1)$. Of course, there are no observational constraints on the parameter η_{\parallel} for dark energy. Also, as discussed in [14], the mass of the corresponding entropic perturbation is always positive in those solutions, so there is no tachyonic instability at the perturbative level. However, they do *not* provide counterexamples to the conclusion of this section that long-term slow-roll rapid-turn inflation cannot occur with a rotationally invariant potential.

⁶Note that, without the assumption $r \approx \text{const}$, the second slow roll condition $|\eta_{\parallel}| \ll 1$ does not amount to neglecting $\ddot{\theta}$ in the equations of motion, as is clear from (4.8). whereas (4.10) implies:

$$H \approx -\frac{V_{\theta}}{3h^2 \dot{\theta}} \quad . \tag{4.13}$$

Substituting (4.13) in (4.11) gives:

$$\frac{V_{\theta}^2}{3h^4\dot{\theta}^2} \approx V \quad . \tag{4.14}$$

Finally, substituting (4.12) in (4.14), we find (in agreement with [38]):

$$\frac{h'}{3h^3} \frac{V_{\theta}^2}{V_r} \approx V \quad . \tag{4.15}$$

This relation need not be satisfied everywhere in field space. But it has to be (approximately) satisfied along a slowly rolling inflationary solution of the background equations of motion with a near circular field space trajectory.

To illustrate the usefulness of the consistency condition (4.15), let us apply it to an example that was considered in [33] when looking for rapid turn solutions in the slow roll regime. To do that, we first note that the field-space metric of [33] has the form:

$$ds^{2} = h^{2}(\rho) \left(d\rho^{2} + d\theta^{2} \right) \quad . \tag{4.16}$$

This is related to (4.1) via the field redefinition:

$$\frac{dr}{d\rho} = h \; .$$

Relation (4.15) has the same form in terms of the variable ρ , namely:

$$\frac{h_{\rho}}{3h^3} \frac{V_{\theta}^2}{V_{\rho}} = V \quad , \tag{4.17}$$

where $f_{\rho} \stackrel{\text{def.}}{=} \partial_{\rho} f$ and $V_{\rho} \stackrel{\text{def.}}{=} \partial_{\rho} V$.

Now we turn to investigating whether (4.17) can be satisfied in the no-scale inspired rapid-turn model considered in [33]. In that case, the functions h and V are:

$$h_{ns} = \frac{3\alpha}{2\rho^2}$$
 , $V_{ns} = \frac{p_1^2 \theta^2 + (p_0 + p_1 \rho)^2}{8^{\alpha} \rho^{3\alpha}}$, (4.18)

where $\alpha, p_0, p_1 = \text{const.}$ Rapid turn is achieved by taking the parameter α to be small. In that limit, the leading term in the potential is:

$$V_{ns} = p_1^2 \theta^2 + (p_0 + p_1 \rho)^2 \quad . \tag{4.19}$$

On the other hand, the leading term in the expression on the left-hand side of (4.17) is:

$$\frac{h_{\rho}}{3h^3} \frac{V_{\theta}^2}{V_{\rho}}\Big|_{ns} = -\frac{16}{27} \frac{p_1^3 \rho^3}{(p_0 + p_1 \rho)} \frac{\theta^2}{\alpha^2} \quad . \tag{4.20}$$

Now recall that $\rho \approx \text{const}$ in the present context. Therefore, for finite θ and small α , the expression (4.20) diverges while (4.19) does not. Hence the two expressions could be (almost) equal numerically only for $\theta \ll 1$ and such that $\theta \sim \mathcal{O}(\alpha)$. However, these conditions are not compatible with the slow roll conditions, which for $\dot{\rho} \approx 0$ are equivalent with demanding that $\epsilon_T \ll 1$ and $\eta_T \ll 1$ (see [33]).

To show this, let us consider in more detail the slow roll parameters ϵ_T and η_T , which are given by:

$$\epsilon_T \approx \frac{1}{2h^2} \left(\frac{V_\theta}{V}\right)^2 \quad \text{and} \quad \eta_T \approx \frac{1}{h^2} \frac{V_{\theta\theta}}{V}$$
(4.21)

in the approximation $\rho \approx \text{const}$ (see [33]). For the functions in (4.18), we find:

$$\epsilon_T|_{ns} \approx \frac{8}{9} \frac{p_1^4 \rho^4}{\left[p_1^2 \theta^2 + (p_0 + p_1 \rho)^2\right]^2} \frac{\theta^2}{\alpha^2}$$
 (4.22)

and:

$$\eta_T|_{ns} \approx \frac{8}{9} \frac{p_1^2 \rho^4}{\left[p_1^2 \theta^2 + (p_0 + p_1 \rho)^2\right]} \frac{1}{\alpha^2} \quad , \tag{4.23}$$

where we did not neglect any subleading terms in small α . These equalities are approximate only due to the approximations in (4.21). Since p_0, p_1 and ρ are finite constants, the slow roll condition $\epsilon_T \ll 1$ can only be satisfied if $\theta \ll \alpha$. However, this contradicts the condition $\theta \sim \mathcal{O}(\alpha)$ found above. Conversely, requiring $\epsilon_T \ll 1$ necessarily violates the consistency condition (4.17), which encodes compatibility with the equations of motion. Therefore, the above method of achieving slow roll and rapid turn is incompatible with the equations of motion. We suspect that the putative inflationary trajectories found in loc. cit solve only the scalar field equations, but not the Einstein equations. The situation is even worse for the slow roll condition $\eta_T \ll 1$, since for small α (i.e. for rapid turn) the expression in (4.23) diverges regardless of any comparison between the magnitudes of α and θ . In other words, slow roll and rapid turn are incompatible with each other in this context.

In the example of the EGNO model considered in [33], the situation is conceptually the same as in the no-scale inspired model discussed above, although the relevant expressions are more cumbersome.

The lesson from this section is that it is not reliable to look for rapid turn solutions by tuning parameters in known slow-roll slow-turn solutions. The conceptual reason is that the latter are approximate solutions of the equations of motion *only* in some parts of their parameters spaces, and not for arbitrary parameter values (because they satisfy the equations of motion up to error terms whose magnitude depends on the parameters). As a consequence, arbitrary variations of the parameters in such solutions can violate the approximations within which they solve (approximately) the equations of motion.

5 A unifying approximation for rapid turn models

A number of prominent rapid turn models of inflation (including [15–19]) can be viewed as special cases of the unifying framework proposed in [20]. The latter uses the adapted frame and does not rely on specific choices of potential V and metric G_{IJ} . It thus appears to provide a wide class of realizations of rapid turn inflation. It turns out that the considerations of [20] rely on an approximation that is less general than slow roll, as will become evident below. We will also show that imposing this approximation leads to rather nontrivial consistency conditions for compatibility with the equations of motion.

5.1 Equations of motion in adapted frame

Let us begin by rewriting the equations of motion (2.4) in the basis (n, τ) . For this, we first expand $\dot{\varphi}^I = \dot{\sigma}T^I$ as:

$$\dot{\varphi}^I = v_n n^I + v_\tau \tau^I \quad , \tag{5.1}$$

where (see (3.3)):

$$v_n \stackrel{\text{def.}}{=} n_I \dot{\varphi}^I = \dot{\sigma} \cos \theta_{\varphi} \quad \text{and} \quad v_\tau \stackrel{\text{def.}}{=} \tau_I \dot{\varphi}^I = \dot{\sigma} \sin \theta_{\varphi} \quad .$$
 (5.2)

Therefore:

$$D_t \dot{\varphi}^I \stackrel{\text{def.}}{=} \dot{\varphi}^J \nabla_J \dot{\varphi}^I = v_n \nabla_n \dot{\varphi}^I + v_\tau \nabla_\tau \dot{\varphi}^I \quad . \tag{5.3}$$

To compute the right hand side, note that (see Appendix A):

$$\nabla_n n = \mu \tau \quad , \quad \nabla_n \tau = -\mu n \quad ,
\nabla_\tau \tau = \lambda n \quad , \quad \nabla_\tau n = -\lambda \tau \quad ,$$
(5.4)

where:

$$\lambda = -\frac{V_{\tau\tau}}{\sqrt{G^{IJ}V_IV_J}} \quad , \quad \mu = \frac{V_{n\tau}}{\sqrt{G^{IJ}V_IV_J}} \quad . \tag{5.5}$$

Substituting (5.1) and (5.4) in (5.3) gives:

$$D_t \dot{\varphi}^I = \left(\dot{v}_n - \lambda v_\tau^2 - \mu v_n v_\tau\right) n^I + \left(\dot{v}_\tau + \mu v_n^2 + \lambda v_n v_\tau\right) \tau^I \quad , \tag{5.6}$$

where we used the relation $\dot{\varphi}^I \nabla_I v_n = \dot{\varphi}^I \partial_I v_n = \dot{v}_n$. Also notice that (3.1) implies:

$$G^{IJ}V_J = \sqrt{G^{KL}V_K V_L} \ n^I \ . \tag{5.7}$$

Using (5.1), (5.6) and (5.7) we find that the projections of (2.4) along n and τ give the following equations respectively:

$$\dot{v}_n - \lambda v_\tau^2 - \mu v_n v_\tau + 3Hv_n + \sqrt{V_I V^I} = 0 , \dot{v}_\tau + \mu v_n^2 + \lambda v_n v_\tau + 3Hv_\tau = 0 .$$
(5.8)

Now let us consider the approximation within which these equations of motion were studied in [20].

5.2 Adapted frame parameters

To discuss the approximation of [20, 21], we introduce the following parameters:

$$f_n \stackrel{\text{def.}}{=} -\frac{\dot{v}_n}{Hv_n} \quad , \quad f_\tau \stackrel{\text{def.}}{=} -\frac{\dot{v}_\tau}{Hv_\tau} \quad .$$
 (5.9)

These quantities are adapted frame analogues of the slow roll parameter η_{\parallel} . However, they do *not* play the same role. To see this, let us write η_{\parallel} in terms of f_n and f_{τ} . For this, notice first that:

$$\dot{\sigma}^2 = G_{IJ} \dot{\varphi}^I \dot{\varphi}^J = v_n^2 + v_\tau^2 \quad , \tag{5.10}$$

where we used (5.1). Using the first relation in (2.9) together with (5.2) and (5.10), we find:

$$\eta_{\parallel} = \cos^2 \theta_{\varphi} f_n + \sin^2 \theta_{\varphi} f_{\tau} \quad . \tag{5.11}$$

The approximation regime in [20] is obtained by imposing the conditions:

$$|f_n| \ll 1$$
 , $|f_\tau| \ll 1$, (5.12)

or more precisely $\dot{v}_{n,\tau} \sim \mathcal{O}(\varepsilon) H v_{n,\tau}$. However, relation (5.11) shows that, although the inequalities (5.12) imply the second slow roll condition $|\eta_{\parallel}| \ll 1$, the latter does *not* imply (5.12). Hence the considerations of [20] encompass only a particular subset of trajectories which satisfy the second slow roll condition. Indeed, a more general possibility is the

following. Since rapid turn requires $\cos^2 \theta_{\varphi} \ll 1$ (as discussed below relation (3.6)), one could satisfy the slow roll condition $|\eta_{\parallel}| \ll 1$ by having $|f_n| \sim \mathcal{O}(1)$ and $|f_{\tau}| \ll 1$. It would be very interesting to investigate in more detail this new regime. In particular, its existence seems to suggest that, counter-intuitively, one can have slow roll while moving relatively fast along the gradient of the potential.

Let us now compute the quantities f_n and f_{τ} on solutions of the equations of motion. Substituting \dot{v}_n and \dot{v}_{τ} from (5.8) in (5.9) gives:

$$f_n = 3 + \frac{\sqrt{V_I V^I} - \lambda v_\tau^2 - \mu v_n v_\tau}{H v_n}$$
$$f_\tau = 3 + \frac{\mu v_n^2 + \lambda v_n v_\tau}{H v_\tau} .$$

Using (2.21), (5.2) and (5.5), the last expressions become:

$$f_n = 3 + \frac{1}{c\cos\theta_{\varphi}} - \frac{c}{H^2} \tan\theta_{\varphi} (-V_{\tau\tau}\sin\theta_{\varphi} + V_{n\tau}\cos\theta_{\varphi}) ,$$

$$f_{\tau} = 3 + \frac{c}{H^2\tan\theta_{\varphi}} (-V_{\tau\tau}\sin\theta_{\varphi} + V_{n\tau}\cos\theta_{\varphi}) .$$
(5.13)

Notice that (3.4) and (3.5) can be written respectively as:

$$\cos \theta_{\varphi} = c(\eta_{\parallel} - 3) \quad , \quad \sin \theta_{\theta} = -c\eta_{\perp} \quad ,$$
 (5.14)

which imply:

$$c^{2} = \frac{1}{\eta_{\perp}^{2} + (\eta_{\parallel} - 3)^{2}} \quad .$$
(5.15)

,

Using (5.14)-(5.15) in (5.13), we find:

$$f_{n} = \eta_{\parallel} + \frac{\eta_{\perp}^{2}}{\eta_{\parallel} - 3} + \frac{\eta_{\perp} \left[V_{\tau\tau} \eta_{\perp} + V_{n\tau} (\eta_{\parallel} - 3) \right]}{H^{2} (\eta_{\parallel} - 3) \left[\eta_{\perp}^{2} + (\eta_{\parallel} - 3)^{2} \right]} ,$$

$$f_{\tau} = 3 - \frac{(\eta_{\parallel} - 3) \left[V_{\tau\tau} \eta_{\perp} + V_{n\tau} (\eta_{\parallel} - 3) \right]}{H^{2} \eta_{\perp} \left[\eta_{\perp}^{2} + (\eta_{\parallel} - 3)^{2} \right]} .$$

_

Imposing the first and second slow roll conditions, these expressions simplify to:

$$f_{n} \approx -\frac{\eta_{\perp}^{2}}{3} - \frac{\eta_{\perp} (V_{\tau\tau}\eta_{\perp} - 3V_{n\tau})}{V(\eta_{\perp}^{2} + 9)} ,$$

$$f_{\tau} \approx 3 + \frac{9(V_{\tau\tau}\eta_{\perp} - 3V_{n\tau})}{V\eta_{\perp}(\eta_{\perp}^{2} + 9)} .$$
(5.16)

Notice that relations (5.16) imply:

$$f_{\tau} \approx -\frac{9}{\eta_{\perp}^2} f_n \quad . \tag{5.17}$$

This is in line with our observation above that one can have slow roll and rapid turn with $|f_n| \sim \mathcal{O}(1)$ and $|f_\tau| \ll 1$.

5.3 Consistency conditions

We next derive consistency conditions for the approximation given by (5.12). In view of (5.17), that approximation holds when $|f_n| \ll 1$. In this case, (5.16) implies:

$$V\eta_{\perp}(\eta_{\perp}^2 + 9) \approx 3(3V_{n\tau} - V_{\tau\tau}\eta_{\perp})$$
 (5.18)

We will show below that combining this approximate equality with the slow roll condition $\varepsilon \ll 1$ leads to a constraint on the potential.

Recall from (2.20) that $\varepsilon \ll 1$ is equivalent with $\kappa \ll 1$. Using (2.21) and (5.15), we compute:

$$\dot{\sigma}^2 = \frac{V_I V^I}{H^2 \left[\eta_\perp^2 + (\eta_\parallel - 3)^2\right]}$$

Substituting this in (2.19), we find that the first slow roll condition $\kappa \ll 1$ takes the form:

$$\frac{V_I V^I}{H^2 \left[\eta_\perp^2 + (\eta_\parallel - 3)^2\right] 2V} \ll 1$$

In the slow roll regime this reduces to:

$$\eta_{\perp}^2 + 9 \gg \frac{3}{2} \frac{V_I V^I}{V^2} \quad . \tag{5.19}$$

Requiring compatibility between (5.19) and (5.18) imposes a constraint on the potential. Namely, there must be at least one solution of the depressed cubic equation (5.18) for η_{\perp} which satisfies the inequality (5.19).

Let us discuss the resulting constraint in more detail. For convenience, we rewrite (5.18) as:

$$\eta_{\perp}^3 + p\eta_{\perp} + q \approx 0 \quad , \tag{5.20}$$

where:

$$p \stackrel{\text{def.}}{=} 9 + 3 \frac{V_{\tau\tau}}{V} \quad , \quad q \stackrel{\text{def.}}{=} -9 \frac{V_{n\tau}}{V} \; .$$

The discriminant of this equation is:

$$\Delta = -4p^3 - 27q^2 \; .$$

If $\Delta \ge 0$, then (5.20) has three real roots. To ensure that at least one of them satisfies (5.19), it is enough to require the largest root to satisfy it. According to Viete's formula, the three roots of the depressed cubic can be written as:

$$\eta_{\perp} = A \cos\left(\frac{1}{3}\arccos B + \frac{2\pi}{3}k\right) \quad , \quad k = 0, 1, 2 \quad ,$$

where $A \stackrel{\text{def.}}{=} \sqrt{\frac{4|p|}{3}}$ and $B \stackrel{\text{def.}}{=} \sqrt{\frac{27q^2}{4|p|^3}}$. Clearly, the largest root belongs to the interval $[\frac{A}{2}, A]$. Hence requiring that it satisfy (5.19) amounts to the condition $9 + \frac{A^2}{4} \gg \frac{3}{2} \frac{V_I V^I}{V^2}$, namely:

$$12 + \frac{V_{\tau\tau}}{V} \gg \frac{3}{2} \frac{V_I V^I}{V^2} \quad . \tag{5.21}$$

If $\Delta < 0$, then (5.20) has a single real root, which is given by Cardano's formula:

$$\eta_{\perp *} = \sqrt[3]{-\frac{q}{2} + \sqrt{\frac{q^2}{4} + \frac{p^3}{27}}} + \sqrt[3]{-\frac{q}{2} - \sqrt{\frac{q^2}{4} + \frac{p^3}{27}}} .$$

Hence (5.19) becomes:

$$\eta_{\perp *}^2 + 9 \gg \frac{3}{2} \frac{V_I V^I}{V^2} \quad . \tag{5.22}$$

Formulas (5.21) and (5.22) are complicated relations between V and G_{IJ} . For any given field-space metric G_{IJ} , this means that there is a highly nontrivial constraint on the potential which limits significantly the choices of V that are compatible with solutions of the equations of motion in the slow roll and rapid turn regime.

6 The characteristic angle

In this section we investigate the behavior of the characteristic angle θ_{φ} , which relates the oriented frames (T, N) and (n, τ) according to equation (3.3). For that purpose, we will rewrite the adapted-frame equations of motion (5.8) in a manner that leads to an equation determining $\theta_{\varphi}(t)$. Although that equation cannot be solved exactly without specifying the form of V and G_{IJ} , it nevertheless leads to a valuable universal insight about the duration of the rapid turn regime.

To obtain an equation for $\theta_{\varphi}(t)$, let us substitute (5.2) in the second equation of (5.8). The result is:

$$\dot{\theta}_{\varphi}\cos\theta_{\varphi} + (3 - \eta_{\parallel})H\sin\theta_{\varphi} + \dot{\sigma}(\mu\cos^2\theta_{\varphi} + \lambda\sin\theta_{\varphi}\cos\theta_{\varphi}) = 0 \quad , \tag{6.1}$$

where we used that $\ddot{\sigma} = -H\dot{\sigma}\eta_{\parallel}$ in accordance with (2.9). It is convenient to rewrite (6.1) as:

$$\dot{\theta}_{\varphi} + (3 - \eta_{\parallel})H \tan \theta_{\varphi} + \dot{\sigma}(\mu \cos \theta_{\varphi} + \lambda \sin \theta_{\varphi}) = 0 \quad . \tag{6.2}$$

Similarly, the first equation in (5.8) takes the form:

$$\dot{\theta}_{\varphi} - (3 - \eta_{\parallel})H\cot\theta_{\varphi} + \dot{\sigma}(\mu\cos\theta_{\varphi} + \lambda\sin\theta_{\varphi}) - \frac{H}{c\sin\theta_{\varphi}} = 0 \quad , \tag{6.3}$$

where we used the relation $\sqrt{V_I V^I} = \frac{\dot{\sigma}H}{c}$, in accordance with (2.21). In the slow roll regime one has $|\eta_{\parallel}| \ll 1$ and $c \approx -\cos \theta_{\varphi}/3$. Hence in the slow roll approximation equation (6.3) coincides with (6.2). In general, the difference between equations (6.3) and (6.2) is precisely the on-shell relation (3.4) between η_{\parallel} , θ_{φ} and c.

Let us now study in more detail equation (6.2) in the slow roll approximation. Using (2.21) and $c \approx -\cos \theta_{\varphi}/3$, we have:

$$\dot{\sigma} \approx -\frac{\cos\theta_{\varphi}}{3} \frac{\sqrt{V_I V^I}}{H}$$

Substituting this together with (2.17) and (5.5) in (6.2), we find that during slow roll one has:

$$\dot{\theta}_{\varphi} + \sqrt{3V} \tan \theta_{\varphi} - \frac{1}{\sqrt{3V}} \left(V_{n\tau} \cos^2 \theta_{\varphi} - V_{\tau\tau} \cos \theta_{\varphi} \sin \theta_{\varphi} \right) \approx 0 \quad . \tag{6.4}$$

Recall that rapid turn requires $\cos^2 \theta_{\varphi} \ll 1$, although $\cos \theta_{\varphi}$ itself need not be very small. So assuming that $V_{n\tau}$ is of order $V_{\tau\tau}$ or smaller, which is reasonable in view of (3.31), we can simplify (6.4) to:

$$\dot{\theta}_{\varphi} + \sqrt{3V} \tan \theta_{\varphi} + \frac{1}{\sqrt{3V}} V_{\tau\tau} \cos \theta_{\varphi} \sin \theta_{\varphi} \approx 0$$
 (6.5)

When the $V_{\tau\tau}$ term is negligible, the last equation can be written in the nice form:

$$\dot{\theta}_{\varphi} \approx \pm \frac{H}{c} , \qquad (6.6)$$

where we used the fact that $\cos \theta_{\varphi} \approx -3c$ and $\sin \theta_{\theta} \approx \pm 1$ during slow roll; of course, one should keep in mind that the parameter c defined in (2.21) is not a constant along the field-space trajectory of an inflationary solution.

Now let us consider one by one the cases when the $V_{\tau\tau}$ term in (6.5) is negligible and when it is not. In the first case, we have:

$$\dot{\theta}_{\varphi} + \sqrt{3V} \tan \theta_{\varphi} \approx 0$$
 (6.7)

To gain insight in the behavior of $\theta_{\varphi}(t)$, let us assume that, at least initially, the inflationary trajectory is close to a level set of the potential⁷, in other words that $V \approx \text{const}$ along (an initial part of) the trajectory. Then the solution of (6.7) is given by:

$$\sin \theta_{\varphi} \approx \pm \tilde{C} e^{-\sqrt{3V}t} , \qquad (6.8)$$

where $\tilde{C} = \text{const} > 0$. This means that $|\sin \theta_{\varphi}|$ tends to zero with t, with a characteristic timescale determined by $T = \frac{1}{\sqrt{3V}}$. This is of course a rather crude estimate, which relies

⁷In view of (3.3), this is equivalent with $|\theta_{\varphi}(0)| \approx \frac{\pi}{2}$.



Figure 1: Approximate time evolution of θ_{φ} and c in the second order slow roll regime, where we assume that $\theta_{\varphi}(0) = \frac{\pi}{2}$, i.e. c(0) = 0. The slow roll curve exits the rapid turn regime after the characteristic time $T \simeq \frac{1}{\sqrt{3V}}$ and enters the infrared regime of [31,32], which is characterized by $\theta_{\varphi} \approx \pi$ and $c \approx 3$.

on neglecting the $V_{\tau\tau}$ term and on the assumption $V \approx \text{const}$. Nevertheless, relation (6.8) gives useful intuition. Note that, if the initial part of the trajectory were not along a level set of the potential, then $|\sin \theta_{\varphi}|$ would decay even faster.

Now let us make an improved estimate by taking into account the $V_{\tau\tau}$ term as well. In that case and assuming $V, V_{\tau\tau} \approx \text{const}$ for simplicity, (6.5) has the following solution:

$$\cos\theta_{\varphi} \approx \pm \sqrt{\frac{\hat{C}e^{2(A+B)t} + A}{\hat{C}e^{2(A+B)t} - B}} \quad , \tag{6.9}$$

where $\hat{C} = \text{const} > 0$ and we defined:

$$A = \sqrt{3V} \qquad , \qquad B = \frac{V_{\tau\tau}}{\sqrt{3V}} \qquad . \tag{6.10}$$

Again this is not an exact result, since we obtained it by assuming that A and B are constant. But it gives important insights. Namely, the characteristic time is $T = \frac{1}{A+B} = \frac{\sqrt{3V}}{3V+V_{\tau\tau}}$. Clearly, we can vary T by varying V and $V_{\tau\tau}$. In any case, though, we have $\cos \theta_{\varphi} \to \pm 1$ as t increases, i.e. we again have $|\sin \theta_{\varphi}| \to 0$ for large enough t. If $\theta_{\varphi}(0) = \frac{\pi}{2}$, then $\dot{\theta}_{\varphi} > 0$ in accordance with (6.6). And if $\theta_{\varphi}(0) = -\frac{\pi}{2}$, then $\dot{\theta}_{\varphi} < 0$. Thus in both cases $\theta_{\varphi}(t)$ tends to $\pi \mod 2\pi$ with time, which means that the field-space trajectory tends to align with minus the gradient of the potential (see Figure 1). This suggests that the rapid turn phase is only transient, although its duration depends on the particulars of the potential and the field-space metric.

Since during slow roll we have $H \approx \sqrt{V/3}$, it follows that the number of e-folds produced during the characteristic time (in the above crude approximation with $V, V_{\tau\tau} \approx \text{const}$) is given by:

$$\mathcal{N} = \int_0^T H \,\mathrm{d}t \,\approx\, \frac{V}{3V + V_{\tau\tau}} \,\lesssim\, \frac{1}{3} \,\,, \tag{6.11}$$

where the last inequality follows from (3.28). This suggests that generically the rapid turn phase is rather short-lived, likely contained within one or only a few e-folds. Obtaining a sustained rapid turn period (capable of lasting 50 - 60 or so e-folds) would require a rather special choice of potential.

Finally, let us comment on the quantities f_n and f_{τ} , defined in (5.9). To estimate them in the crude approximation of (6.7), let us first compute the following ratios:

$$\frac{\dot{v}_n}{v_n} = -\eta_{\parallel} H - \dot{\theta}_{\varphi} \tan \theta_{\varphi} \approx H \left(3 \tan^2 \theta_{\varphi} - \eta_{\parallel} \right) \approx \frac{H}{3} \eta_{\perp}^2 \quad ,$$

$$\frac{\dot{v}_{\tau}}{v_{\tau}} = -\eta_{\parallel} H - \dot{\theta}_{\varphi} \cot \theta_{\varphi} \approx -H \left(3 + \eta_{\parallel} \right) \approx -3H \quad , \qquad (6.12)$$

where we used successively (5.2), (2.9) and (6.7), as well as the slow roll approximation. Substituting (6.12) in (5.9) gives:

$$f_n \approx -\frac{\eta_\perp^2}{3} \qquad , \qquad f_\tau \approx 3 \quad .$$
 (6.13)

This is a rather crude estimate, relying on the leading behavior of the characteristic angle $\theta_{\varphi}(t)$ for $V, V_{\tau\tau} \approx \text{const.}$ However, it suggests that satisfying the approximations (5.12) would require a rather special choice of scalar potential.

7 Conclusions

We studied consistency conditions for certain approximations commonly used to search for rapid-turn and slow-roll inflationary trajectories in two-field cosmological models with orientable scalar field space. Such consistency conditions arise from requiring compatibility between the equations of motion and the various relevant approximations. We showed that long-lasting rapid turn trajectories with third order slow roll can arise only in regions of the field space, where the scalar potential satisfies a certain nonlinear second order ODE with coefficients depending on the field space metric. We also studied the relation between the scalar potential and scalar field metric, which is necessary for the existence of slow-roll circular trajectories in two-field models with rotationally invariant scalar field metric. Moreover, we showed that the approximation proposed in [20] (which is a rather special case of the slow-roll and rapid-turn approximation) is compatible with the equations of motion only when the scalar metric and potential satisfy rather nontrivial consistency conditions, which we determined explicitly.

In principle, the various consistency relations we have obtained could be (approximately) satisfied numerically for a brief period by appropriate choices of integration constants. But they can only be maintained for a prolonged duration, if they are satisfied functionally. Hence, they constrain dramatically the classes of two-field models, for which one can hope to find long-lasting rapid-turn and slow-roll inflationary trajectories. Thus, our results show that two-field cosmological models, which allow for such trajectories, are non-generic in the class of all two-field models. In particular, this shows that the difficulty in finding such models – which was previously noticed in [33] – is already present in the absence of supersymmetry and hence is unrelated to supergravity.

We also studied the time evolution of the *characteristic angle* of cosmological trajectories in general two-field models. We argued that (in a crude approximation) a slow roll cosmological trajectory with rapid turn tends to align within a short time with the opposite of a gradient flow line of the model. Thus, generically, the solution tends to enter the gradient flow regime of the model before producing a sufficient number of e-folds. This confirms our conclusion that phenomenologically viable inflationary slow roll trajectories with long-lasting rapid turn are highly non-generic in two-field cosmological models.

The above conclusion may be a 'blessing in disguise', since finding such rare trajectories could be more predictive (than if they were generic) and might lead to some deep insights about the embedding of these effective multifield models into an underlying fundamental framework. We hope to investigate in the future what properties of the scalar potential are implied by the consistency conditions found here, as well as to look for ways of solving those conditions even if only for very special choices of scalar metric. In view of [39], it is also worth exploring the consequences of transient violations of slow roll in our context.

Acknowledgements

We thank D. Andriot, E.M. Babalic, J. Dumancic, R. Gass, S. Paban, L.C.R. Wijewardhana and I. Zavala for interesting discussions on inflation and cosmology. We are grateful to P. Christodoulidis for useful correspondence. We also thank the Stony Brook Workshop in Mathematics and Physics for hospitality. L.A. has received partial support from the Bulgarian NSF grant KP-06-N38/11. The work of C.L. was supported by grant PN 19060101/2019-2022.

A Derivatives of the adapted frame

Since n is normalized, we have $\nabla_n(n_K n^K) = 0$. Hence $\nabla_n n = n^I \nabla_I n$ does not have a component along the vector n. To compute its component along the vector τ , let us contract it with τ :

$$\tau_J n^I \nabla_I n^J = \tau_J n^I \frac{G^{JK} \nabla_I V_K}{\sqrt{V_L V^L}} + \tau_J n^I G^{JK} V_K \nabla_I \left(\frac{1}{\sqrt{V_L V^L}}\right) \quad , \tag{A.1}$$

where we used (3.1) inside $\nabla_I n^J$. Substituting (3.2) in the second term of (A.1) gives:

$$\tau_J \nabla_n n^J = \frac{\tau^K n^I \nabla_I V_K}{\sqrt{V_L V^L}} + \sqrt{\det G} \,\epsilon_{JM} \, n^M n^I V^J \nabla_I \left(\frac{1}{\sqrt{V_L V^L}}\right) = \frac{V_{n\tau}}{\sqrt{V_L V^L}} \quad , \tag{A.2}$$

where in the second equality we used the relation $\epsilon_{JM} n^M V^J = \epsilon_{JM} V^M V^J / \sqrt{V_L V^L} = 0$, which follows from (3.1). In conclusion, we have

$$\nabla_n n = \frac{V_{n\tau}}{\sqrt{V_I V^I}} \tau \quad . \tag{A.3}$$

One can prove the remaining three relations in (5.4) in a similar manner.

References

- S. Garg, C. Krishnan, Bounds on Slow Roll and the de Sitter Swampland, JHEP 11 (2019) 075, arXiv:1807.05193 [hep-th].
- [2] H. Ooguri, E. Palti, G. Shiu, C. Vafa, Distance and de Sitter Conjectures on the Swampland, Phys. Lett. B 788 (2019) 180, arXiv:1810.05506 [hep-th].
- [3] A. Achucarro, G. Palma, *The string swampland constraints require multi-field inflation*, JCAP 02 (2019) 041, arXiv:1807.04390 [hep-th].
- [4] R. Bravo, G. A. Palma, S. Riquelme, A Tip for Landscape Riders: Multi-Field Inflation Can Fulfill the Swampland Distance Conjecture, JCAP 02 (2020) 004, arXiv:1906.05772 [hep-th].
- [5] C. M. Peterson, M. Tegmark, *Testing Two-Field Inflation*, Phys. Rev. D 83 (2011) 023522, arXiv:1005.4056 [astro-ph.CO].
- C. M. Peterson, M. Tegmark, Non-Gaussianity in Two-Field Inflation, Phys. Rev. D 84 (2011) 023520, arXiv:1011.6675 [astro-ph.CO].

- [7] S. Cespedes, V. Atal, G. A. Palma, On the importance of heavy fields during inflation, JCAP 05 (2012) 008, arXiv:1201.4848 [hep-th].
- [8] T. Bjorkmo, R. Z. Ferreira, M.C.D. Marsh, Mild Non-Gaussianities under Perturbative Control from Rapid-Turn Inflation Models, JCAP 12 (2019) 036, arXiv:1908.11316 [hep-th].
- [9] G. A. Palma, S. Sypsas, C. Zenteno, Seeding primordial black holes in multi-field inflation, Phys. Rev. Lett. 125 (2020) 121301, arXiv:2004.06106 [astro-ph.CO].
- [10] J. Fumagalli, S. Renaux-Petel, J. W. Ronayne, L. T. Witkowski, *Turning in the land-scape: a new mechanism for generating Primordial Black Holes*, arXiv:2004.08369 [hep-th].
- [11] L. Anguelova, On Primordial Black Holes from Rapid Turns in Two-field Models, JCAP 06 (2021) 004, arXiv:2012.03705 [hep-th].
- [12] L. Anguelova, Primordial Black Hole Generation in a Two-field Inflationary Model, arXiv:2112.07614 [hep-th].
- [13] Y. Akrami, M. Sasaki, A. Solomon and V. Vardanyan, Multi-field dark energy: cosmic acceleration on a steep potential, Phys. Lett. B 819 (2021) 136427, arXiv:2008.13660 [astro-ph.CO].
- [14] L. Anguelova, J. Dumancic, R. Gass, L.C.R. Wijewardhana, Dark Energy from Inspiraling in Field Space, JCAP 03 (2022) 018, arXiv:2111.12136 [hep-th].
- [15] A Brown, Hyperinflation, Phys. Rev. Lett. 121 (2018) 251601, arXiv:1705.03023
 [hep-th].
- [16] S. Mizuno, S. Mukohyama, Primordial perturbations from inflation with a hyperbolic field-space, Phys. Rev. D 96 (2017) 103533, arXiv:1707.05125 [hep-th].
- [17] P. Christodoulidis, D. Roest, E. Sfakianakis, Angular inflation in multi-field αattractors, JCAP 11 (2019) 002, arXiv:1803.09841 [hep-th].
- [18] S. Garcia-Saenz, S. Renaux-Petel, J. Ronayne, Primordial fluctuations and non-Gaussianities in sidetracked inflation, JCAP 1807 (2018) 057, arXiv:1804.11279 [astro-ph.CO].

- [19] A. Achucarro, E. Copeland, O. Iarygina, G. Palma, D.G. Wang, Y. Welling, Shift-Symmetric Orbital Inflation: single field or multi-field?, Phys. Rev. D 102 (2020) 021302, arXiv:1901.03657 [astro-ph.CO].
- [20] T. Bjorkmo, The rapid-turn inflationary attractor, Phys. Rev. Lett. 122 (2019) 251301, arXiv:1902.10529 [hep-th].
- [21] T. Bjorkmo, M. C. D. Marsh, Hyperinflation generalised: from its attractor mechanism to its tension with the 'swampland conditions', JHEP 04 (2019) 172, arXiv:1901.08603 [hep-th].
- [22] V. Aragam, S. Paban, R. Rosati, *The Multi-Field, Rapid-Turn Inflationary Solution*, JHEP 03 (2021) 009, arXiv:2010.15933 [hep-th].
- [23] Y. Zhang, Y.-g. Gong, Z.-H. Zhu, Noether Symmetry Approach in multiple scalar fields Scenario, Phys. Lett. B688 (2010) 13, arXiv:0912.0067 [hep-ph].
- [24] A. Paliathanasis, M. Tsamparlis, Two scalar field cosmology: Conservation laws and exact solutions, Phys. Rev. D 90 (2014) 043529, arXiv:1408.1798 [gr-qc].
- [25] L. Anguelova, E.M. Babalic, C.I. Lazaroiu, Two-field Cosmological α-attractors with Noether Symmetry, JHEP 1904 (2019) 148, arXiv:1809.10563 [hep-th].
- [26] L. Anguelova, E.M. Babalic, C.I. Lazaroiu, Hidden symmetries of two-field cosmological models, JHEP 09 (2019) 007, arXiv:1905.01611 [hep-th].
- [27] C. I. Lazaroiu, Hesse manifolds and Hessian symmetries of multifield cosmological models, Rev. Roumaine Math. Pures Appl. 66 (2021) 2, 329-345, arXiv:2009.05117 [hep-th].
- [28] C. I. Lazaroiu, C. S. Shahbazi, Generalized two-field α-attractor models from geometrically finite hyperbolic surfaces, Nucl. Phys. B 936 (2018) 542-596, arXiv:1702.06484 [hep-th].
- [29] E. M. Babalic, C. I. Lazaroiu, Generalized α-attractor models from elementary hyperbolic surfaces, Adv. Math. Phys. 2018 (2018) 7323090, arXiv:1703.01650 [hep-th].
- [30] E. M. Babalic, C. I. Lazaroiu, Generalized α-attractors from the hyperbolic triplypunctured sphere, Nucl. Phys. B 937 (2018) 434-477, arXiv:1703.06033 [hep-th].

- [31] C. I. Lazaroiu, Dynamical renormalization and universality in classical multifield cosmological models, Nucl. Phys. B 983 (2022) 115940, arXiv:2202.13466 [hep-th].
- [32] E. M. Babalic, C. I. Lazaroiu, The infrared behavior of tame two-field cosmological models, Nucl. Phys. B 983 (2022) 115929, arXiv:2203.02297 [gr-qc].
- [33] V. Aragam, R. Chiovoloni, S. Paban, R. Rosati, I. Zavala, Rapid-turn inflation in supergravity is rare and tachyonic, JCAP 03 (2022) 002, arXiv:2110.05516 [hep-th].
- [34] A, R. Liddle, P. Parsons, J. D. Barrow, Formalizing the slow-roll approximation in inflation, Phys. Rev. D 50 (1994) 7222, arXiv:astro-ph/9408015.
- [35] A. Hetz, G. Palma, Sound Speed of Primordial Fluctuations in Supergravity Inflation, Phys. Rev. Lett. 117 (2016) 101301, arXiv:1601.05457 [hep-th].
- [36] A. Achucarro, J.-O. Gong, S. Hardeman, G. Palma, S. Patil, *Features of heavy physics in the CMB power spectrum*, JCAP 01 (2011) 030, arXiv:1010.3693 [hep-ph].
- [37] D. Chakraborty, R. Chiovoloni, O. Loaiza-Brito, G. Niz, I. Zavala, Fat inflatons, large turns and the η-problem, JCAP 01 (2020) 020, arXiv:1908.09797 [hep-th].
- [38] P. Christodoulidis, D. Roest, E. Sfakianakis, Scaling attractors in multi-field inflation, JCAP 12 (2019) 059, arXiv:1903.06116 [hep-th].
- [39] S. Bhattacharya, I. Zavala, *Sharp turns in axion monodromy: primordial black holes* and gravitational waves, arXiv:2205.06065 [astro-ph.CO].