

Ensure Differential Privacy and Convergence Accuracy in Consensus Tracking and Aggregative Games with Coupling Constraints

Yongqiang Wang

Abstract—We address differential privacy for fully distributed aggregative games with shared coupling constraints. By co-designing the generalized Nash equilibrium (GNE) seeking mechanism and the differential-privacy noise injection mechanism, we propose the first GNE seeking algorithm that can ensure both provable convergence to the GNE and rigorous ϵ -differential privacy, even with the number of iterations tending to infinity. As a basis of the co-design, we also propose a new consensus-tracking algorithm that can achieve rigorous ϵ -differential privacy while maintaining accurate tracking performance, which, to our knowledge, has not been achieved before. To facilitate the convergence analysis, we also establish a general convergence result for stochastically-perturbed nonstationary fixed-point iteration processes, which lie at the core of numerous optimization and variational problems. Numerical simulation results confirm the effectiveness of the proposed approach.

I. INTRODUCTION

In recent years, Nash equilibrium seeking in aggregative games under shared coupling constraints is gaining increased traction [1], [2], [3]. Compared with the standard Nash equilibrium seeking problem where the feasible decision sets of individual players (also called agents) are independent of each other, the incorporation of shared coupling constraints in Nash equilibrium seeking, usually called the generalized Nash equilibrium (GNE) model, captures important characteristics of noncooperative games under limited network resources. To date, the GNE seeking problem has found vast applications in economics and various engineering domains, with typical examples including power grids [4], [5], [6], optical networks [7], communication networks [8], [9], plug-in electric cars [10], [11], and mobile sensor networks [12]. In many of these applications, due to practical constraints, no central coordinator/mediator exists to collect and disperse information, and individual players can only access or observe the decisions of their immediate neighbors. Compared with the traditional case where every player can access all its competitors' decisions to precisely evaluate its cost function, which is called the *full-decision information* setting, the case where each player can only access its neighbors' decisions is called the *partial-decision information* setting, and is more challenging in that individual players lack information to compute their cost functions or gradients.

To account for the lack of information in the partial-decision information setting, players have to exchange information among local neighbors to estimate global information necessary for GNE seeking. Since the seminal work in [13], [14],

significant inroads have been made in games with partial-decision information, both for Nash equilibrium seeking without coupling constraints (see, e.g., [15], [16], [17], [18]) and GNE seeking (see, e.g., [1], [3], [19]). A key technique underlying these fully distributed algorithms is consensus tracking [20] (also called dynamic average consensus), which allows a group of agents to locally track the average of their reference inputs while each agent only has access to its own individual reference input. However, the employment of consensus tracking in GNE seeking also entails explicit sharing of (estimated) decisions in every iteration, which is problematic when involved information is sensitive. Given the competitive nature of noncooperative games, individual players' privacy should be protected to avoid exploitation by opponents. For example, in Nash-Cournot games, individual players' cost functions could be market sensitive, and should not be disclosed to competitors [21]. In fact, in many scenarios, privacy preservation is mandated by legislation. For example, in road routing games [22], California Privacy Rights Act forbids disclosing drivers' spatiotemporal information, which, otherwise, can reveal a person's activities [23]. Moreover, privacy preservation is also a crucial step to encourage participation in cooperative policies [24].

To address the privacy issue in games, several approaches have been proposed for the full-decision information setting where a coordinator exists (see, e.g., [25], [26], [27]). Recently, results have also emerged to protect privacy in the partial-decision information setting (see, e.g., [24], [28]). However, these approaches are restricted in that they either require the communication graph to satisfy certain properties, or can only protect the cost function value from being uniquely identifiable. As differential privacy (DP) has become the de facto standard for privacy protection due to its strong resilience against arbitrary post-processing and auxiliary information [29], recently [30] and [31] achieve DP in aggregative games at the cost of losing accurate convergence to the exact Nash equilibrium. By co-designing the Nash-equilibrium seeking mechanism and the DP-noise injection mechanism, our prior results [32], [33] successfully achieve both ϵ -DP and accurate convergence in coupling-constraint free Nash-equilibrium seeking. However, these results are inapplicable to GNE seeking problems, where the shared coupling constraints increase attack surfaces, and hence, pose additional challenges to privacy protection. To the best of our knowledge, privacy protection for GNE seeking is still an open problem in the partial-decision information setting.

In this paper, we propose a fully distributed GNE seeking algorithm that can ensure both accurate convergence and rigorous ϵ -DP, with the cumulative privacy budget guaranteed to be finite even when the number of iterations tends to

The work was supported in part by the National Science Foundation under Grants ECCS-1912702, CCF-2106293, CCF-2215088, and CNS-2219487.

Yongqiang Wang is with the Department of Electrical and Computer Engineering, Clemson University, Clemson, SC 29634, USA yongqiw@clemson.edu

infinity. Our approach is motivated by the observation that DP-noises enter the algorithm through inter-player interaction, and hence, its influence on convergence accuracy can be attenuated by gradually weakening inter-player interaction, which would become unnecessary anyway after convergence. To ensure that every player can still estimate global information necessary for GNE seeking, we judiciously design the weakening factor sequence for inter-player interaction and the stepsize sequence, which enables the achievement of accurate convergence even in the presence of DP-noises. We rigorously prove that the algorithm is ϵ -differentially private with a finite cumulative privacy budget, even when the number of iterations tends to infinity. Given that a key component of our GNE seeking algorithm is consensus tracking but the conventional consensus tracking algorithm will lead to cumulative and exploding variance in the presence of persistent noise [34], we first propose a new robust consensus tracking algorithm that can ensure both provable convergence and ϵ -DP under persistent DP-noise.

The main contributions are summarized as follows:

- 1) The proposed new consensus tracking approach can ensure both provable convergence accuracy and ϵ -DP, a goal that has not been achieved before to the best of our knowledge.
- 2) The proposed fully distributed GNE seeking algorithm, to our knowledge, is the first to achieve rigorous ϵ -DP in GNE seeking under partial-decision information. Note that compared with Nash equilibrium seeking without coupling constraints, the shared coupling constraints in GNE problems create additional attack surfaces and thus challenges in privacy protection.
- 3) Besides achieving rigorous ϵ -DP, the approach can simultaneously ensure accurate convergence to the GNE, which is in sharp contrast to existing DP solutions for coupling-constraint free aggregative games (e.g., [30], [31]) that have to trade accurate convergence for DP.
- 4) To facilitate convergence analysis under DP-noises, we establish convergence results for stochastically-perturbed non-stationary fixed-point iteration processes. Given that fixed-point iteration processes lie at the core of many optimization and variational problems, the results are expected to have broad ramifications.
- 5) Even without considering privacy protection, our proof techniques are fundamentally different from existing counterparts and are of independent interest. More specifically, existing proof techniques (in, e.g., [13], [31], [3], [35], [36], [37], [38], [39]) for partial-decision information games rely on the geometric decreasing of consensus errors among the players. In the proposed approach, the diminishing inter-player interaction makes it impossible to have such geometric decreasing of consensus errors, which entails new proof techniques.

The organization of the paper is as follows. Sec. II gives the problem formulation and some results for a later use. Sec. III presents a new consensus-tracking algorithm. This section also proves that the algorithm can ensure both provable accurate convergence and rigorous ϵ -DP with a finite cumulative privacy budget, even when the number of iterations

goes to infinity. Sec. IV proposes a fully distributed GNE seeking algorithm, while its convergence and privacy analysis are presented in Sec. V and Sec. VI, respectively. Sec. VII presents numerical simulation results to confirm the obtained results. Finally, Sec. VIII concludes the paper.

Notations: We define $\mathbb{N} \triangleq \{0, 1, 2, \dots\}$. We use \mathbb{R}^d to denote the Euclidean space of dimension d and \mathbb{R}_+^d to denote the set of all nonnegative vectors in \mathbb{R}^d . We write I_d for the identity matrix of dimension d , and $\mathbf{1}_d$ for the d -dimensional column vector with all entries equal to 1; in both cases we suppress the dimension when it is clear from the context. A vector is viewed as a column vector, and for a vector x , $[x]_i$ denotes its i th element. We use $\langle \cdot, \cdot \rangle$ to denote the inner product and $\|x\|$ for the standard Euclidean norm of a vector x . We use $\|x\|_1$ to represent the L_1 norm of a vector x . We write $\|A\|$ for the matrix norm induced by the vector norm $\|\cdot\|$. A^T denotes the transpose of a matrix A . Given vectors x_1, \dots, x_m , we define $\text{col}(x_1, \dots, x_m) = [x_1^T, \dots, x_m^T]^T$, $x_{-i} = \text{col}(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_m)$, and $\bar{x} = \frac{\sum_{i=1}^m x_i}{m}$. Often, we abbreviate *almost surely* by *a.s.* Given sets $\Omega_1, \Omega_2, \dots, \Omega_m$, we use $\Omega_1 \times \Omega_2 \times \dots \times \Omega_m$ or $\prod_{i=1}^m \Omega_i$ to represent the Cartesian product of these sets.

II. PROBLEM FORMULATION AND PRELIMINARIES

A. Monotone Operators

All of the following are adopted from [40]. Let \mathcal{X} and \mathcal{Y} be non-empty sets, and $2^{\mathcal{Y}}$ be the family of all subsets of \mathcal{Y} . An operator $T : \mathcal{X} \rightarrow \mathcal{Y}$ is a mapping that maps every point $x \in \mathcal{X}$ to a point $T(x)$ in \mathcal{Y} . Thus, the notion $T : \mathcal{X} \rightarrow 2^{\mathcal{Y}}$ means that T is a set-valued operator from \mathcal{X} to \mathcal{Y} , i.e., T maps every point $x \in \mathcal{X}$ to a set $T(x) \subset \mathcal{Y}$. Id denotes the identity operator. The domain of an operator T is represented by $\text{dom}T = \{x | T(x) \neq \emptyset\}$ where \emptyset denotes the empty set. The range of an operator T is defined by $\text{ran}T = \{u | \exists x, u \in T(x)\}$. The graph of T is defined as $\text{gra}T = \{(x, u) | u \in T(x)\}$, and the inverse of an operator is defined using graph as $\text{gra}T^{-1} = (u, x) | u \in T(x)$. The zero set of T is defined as $\text{zer}T = \{x | 0 \in T(x)\}$.

We consider operators that are monotone, i.e., operators satisfying $\langle x - y, u - v \rangle \geq 0$ for all (x, u) and (y, v) in $\text{gra}T$. An operator T is maximally monotone if $\text{gra}T$ is not contained in the graph of any other monotone operator. T is η -strongly monotone, with $\eta > 0$, if $\langle x - y, u - v \rangle \geq \eta \|u - v\|^2$ for all (x, u) and (y, v) in $\text{gra}T$. A set-valued operator $T : \mathbb{R}^d \rightarrow 2^{\mathbb{R}^d}$ is called restricted-strictly monotone with respect to $\mathcal{U} \subset \mathbb{R}^d$ if $\langle x^* - x, u^* - u \rangle > 0$ for all $u^* \in \mathcal{U}$, $u \in \mathbb{R}^d \setminus \mathcal{U}$, $x^* \in T(u^*)$, and $x \in T(u)$. For an operator $T : \mathbb{R}^d \rightarrow 2^{\mathbb{R}^d}$, we use $(\text{Id} + T)^{-1}$ to represent its resolvent, which is single valued and has domain equal to \mathbb{R}^d if T is maximally monotone. For two operators T_1 and T_2 , their composition is denoted as $T_1 \circ T_2$, and their sum $T_1 + T_2$ is defined as $\text{gra}(T_1 + T_2) = \{(x, y + z) | (x, y) \in \text{gra}T_1, (x, z) \in \text{gra}T_2\}$. If T_2 is single valued, then we always have $\text{zer}(T_1 + T_2) = \text{Fix}((\text{Id} + T_1)^{-1} \circ (\text{Id} - T_2))$, where $\text{Fix}T$ denotes the fixed point set $\{x \in \text{dom}T | T(x) = x\}$.

The subdifferential ∂f of a proper lower semicontinuous convex function $f : x \rightarrow 2^{\mathbb{R}^d}$ is defined as $x \rightarrow \{g \in$

$\mathbb{R}^d | f(y) \geq f(x) + \langle g, y - x \rangle, \forall y \in \text{dom} f\}$. We define the indicator function of Ω as $\iota_\Omega(x)$ ($\iota_\Omega(x) = 0$ for $x \in \Omega$ and $\iota_\Omega(x) = \infty$ for $x \notin \Omega$). ι_Ω is always a proper lower semicontinuous convex function when Ω is a closed convex set. $\partial\iota_\Omega$ is usually called the normal cone operator of Ω , and is denoted as N_Ω .

Operator T is nonexpansive if it satisfies $\|T(x) - T(y)\| \leq \|x - y\|$ for all x and y in $\text{dom}T$. It is α -averaged for $0 < \alpha < 1$ if there exists a nonexpansive operator R such that $T = (1 - \alpha)\text{Id} + \alpha R$ holds. T being α -averaged is equivalent to $\|T(x) - T(y)\|^2 \leq \|x - y\|^2 - \frac{1 - \alpha}{\alpha} \|(x - y) - (Tx - Ty)\|^2$ (or $\|T(x) - T(y)\|^2 + (1 - \alpha)\|x - y\|^2 \leq 2(1 - \alpha)\langle x - y, Tx - Ty \rangle$) for all $x, y \in \text{dom}T$ [40].

When T is $\frac{1}{2}$ -averaged, we call it firmly nonexpansive. For a maximally monotone operator, we have $(\text{Id} + T)^{-1}$ being firmly nonexpansive. The projection mapping of x onto a closed and convex set Ω , i.e., $\Pi_\Omega(x) = \arg \min_{y \in \Omega} \|x - y\|$, is firmly nonexpansive since $\Pi_\Omega(x)$ is equivalent to $(\text{Id} + N_\Omega)^{-1}(x)$, which is maximally monotone. T is called β -cocoercive for $\beta > 0$ if βT is $\frac{1}{2}$ -averaged, i.e., $\|\beta T(x) - \beta T(y)\|^2 \leq \langle x - y, Tx - Ty \rangle$. If f is convex differentiable, with its gradient ∇f being θ -Lipschitz, then ∇f is $\frac{1}{\theta}$ -cocoercive.

B. On GNE Seeking

We consider a GNE problem among a set of m players (agents) $[m] = \{1, \dots, m\}$. We index the players by $1, 2, \dots, m$. Player i is characterized by a strategy set $\Omega_i \subseteq \mathbb{R}^d$ and a cost function $J_i(x_i, \bar{x})$, which is dependent on both its own decision x_i and the aggregate of all players' decisions $\bar{x} \triangleq \frac{\sum_{i=1}^m x_i}{m}$. Since each x_i is restricted in Ω_i , the average \bar{x} is restricted to the set that is the $1/m$ -scaling of the Minkowski sum¹ of the sets Ω_i , denoted by $\bar{\Omega} = \frac{1}{m}(\Omega_1 + \Omega_2 + \dots + \Omega_m)$.

Moreover, the decisions of all players must satisfy a shared affine coupling constraint

$$\sum_{i=1}^m C_i x_i \leq \sum_{i=1}^m c_i, \quad (1)$$

where $C_i \in \mathbb{R}^{n \times m}$ and $c_i \in \mathbb{R}^n$ are local parameters only known to player i . It is worth noting that such affine coupling constraints arise in various game scenarios involving upper or lower limits of shared resources [4], [10], [41], [42], and have been a common assumption in the literature of noncooperative games [1], [2], [3], [35], [43]. For notational simplicity, we define $C \triangleq [C_1, \dots, C_m] \in \mathbb{R}^{n \times md}$ and $c \triangleq \sum_{i=1}^m c_i \in \mathbb{R}^n$.

With these notations, we can formalize the game that player i faces as the following parameterized optimization problem:

$$\begin{aligned} & \min J_i(x_i, \bar{x}) \\ \text{s.t. } & x_i \in \Omega_i \text{ and } C_i x_i - c_i \leq \sum_{j \neq i, j \in [m]} c_j - C_j x_j, \end{aligned} \quad (2)$$

where the last constraint is equivalent to (1). The constraint set Ω_i , the function $J_i(\cdot, \cdot)$, and the coupling matrices C_i as well as c_i are known to player i only.

¹A scaling tX of a set X with a scalar t is the set given by $tX = \{tx \mid x \in X\}$. A Minkowski sum of two sets X and Y is the set $X + Y = \{x + y \mid x \in X, y \in Y\}$.

Assumption 1. For every player $i \in [m]$, $J_i(x_i, \bar{x})$ is a differentiable convex function with respect to x_i given any fixed x_{-i} . $\Omega_i \subset \mathbb{R}^d$ is nonempty, compact, and convex. Moreover, the global feasible set

$$K \triangleq \Pi_{i=1}^m \Omega_i \cap \left\{ x \in \mathbb{R}^{md} \mid \sum_{i=1}^m C_i x_i \leq \sum_{i=1}^m c_i \right\} \quad (3)$$

is nonempty and satisfies Slater's constraint qualification.

A GNE of game (2) is a collective strategy $x^* = \text{col}(x_1^*, \dots, x_m^*)$ such that the following relationship holds for any $i \in [m]$ and $x \in K$:

$$J_i(x_i^*, \bar{x}^*) \leq J_i\left(z, \frac{z}{m} + \frac{\sum_{j \neq i, j \in [m]} x_j^*}{m}\right), \text{ s.t. } (z, x_{-i}^*) \in K.$$

Inspired by the recent success of the operator-theoretic approach to GNE seeking [1], [2], [3], [44], we will employ a monotone-operator based approach [40] to treat the convergence to GNE under DP-noises. More specifically, according to [1], [2], [3], a set of strategies x^* is a GNE of the game in (2) if and only if the following conditions are satisfied for any $i \in [m]$:

$$\begin{cases} 0 \in \nabla_{x_i} J(x_i^*, \bar{x}^*) + N_{\Omega_i}(x_i^*) + C_i^T \lambda_i^* \\ 0 \in -(Cx^* - c) + N_{\mathbb{R}_+^n}(\lambda_i^*). \end{cases} \quad (4)$$

Similar to [1], [2], [3], we are interested in a type of GNE called variational GNE, which has the economic interpretation of no price discrimination [45], and corresponds to a solution set of (4) with equal dual variables $\lambda_1^* = \dots = \lambda_m^*$ [1], [46]. Recently, [3] shows that the variational GNE corresponds to the zero set of the following set-valued mapping:

$$T : \begin{bmatrix} x \\ \lambda \end{bmatrix} \mapsto \begin{bmatrix} N_\Omega(x) + F(x) + \frac{C_f^T \lambda}{m} \\ N_{\mathbb{R}_+^{mn}}(\lambda) + \Pi_f \lambda - \frac{C_f x - c_f}{m} \end{bmatrix} \quad (5)$$

where $x = \text{col}(x_1, \dots, x_m)$, $\lambda = \text{col}(\lambda_1, \dots, \lambda_m)$, $C_f = \mathbf{1}_m \otimes C$, $c_f = \mathbf{1}_m \otimes c$, $\Pi_f \triangleq (I_m - \frac{\mathbf{1}\mathbf{1}^T}{m}) \otimes I_n$, and $F(x)$ is the pseudogradient of the game defined as

$$F(x) = \text{col}(\nabla_{x_1} J(x_1, \bar{x}), \dots, \nabla_{x_m} J(x_m, \bar{x})). \quad (6)$$

Similar to [3], we make the following assumption:

Assumption 2. $F(\cdot)$ in (6) is μ -cocoercive over K .

It is worth noting that when $F(\cdot)$ is Lipschitz, then $F(\cdot)$ being strongly monotone implies that it is cocoercive.

Using the definition of $F(\cdot)$, we can further define $F_i(v, u) \triangleq \nabla_v J_i(v, u)$ and $F(v, u) \triangleq \text{col}(F_1(v, u), \dots, F_m(v, u))$. It can be verified that $F(x, \mathbf{1} \otimes \bar{x})$ is equal to $F(x)$ defined in (6).

Moreover, following [1], [3], [13], we make the following assumption on the mapping $F(\cdot, \cdot)$:

Assumption 3. There exists a constant \tilde{L} such that $F(\cdot, \cdot)$ satisfies the following relationship for all $u_1, u_2 \in K$ and $v_1, v_2 \in \mathbb{R}^{md}$:

$$\|F(u_1, v_1) - F(u_2, v_2)\| \leq \tilde{L} \left\| \begin{bmatrix} u_1 \\ v_1 \end{bmatrix} - \begin{bmatrix} u_2 \\ v_2 \end{bmatrix} \right\|.$$

Using operator splitting [40], the mapping T in (5) can be split into the sum of two mappings:

$$\begin{aligned} T_1 &: \begin{bmatrix} x \\ \lambda \end{bmatrix} \mapsto \begin{bmatrix} F(x) \\ \Pi_f \lambda + \frac{c_f}{m} \end{bmatrix}, \\ T_2 &: \begin{bmatrix} x \\ \lambda \end{bmatrix} \mapsto \begin{bmatrix} N_\Omega(x) + \frac{C_f^T \lambda}{N} \\ N_{\mathbb{R}_+^{mn}}(\lambda) - \frac{C_f x}{m} \end{bmatrix}. \end{aligned} \quad (7)$$

It has been proven in [3] that the operator T_2 is maximally monotone, and that the operator T_1 is δ -cocoercive ($0 < \delta \leq \min\{1, \mu\}$ with μ from Assumption 2) and strictly monotone when $\lambda_1 = \dots = \lambda_m$ holds. Furthermore, [3] also showed that when T is restricted-strictly monotone, the variational GNE of the game in (2) is fully characterized by the zeros of the operator T :

Lemma 1. [3] *Under Assumption 1, if $\text{zer}(T) \neq \emptyset$ and $\text{col}(x^*, \lambda^*) \in \text{zer}(T)$, then x^* is a variational GNE and $\lambda_1^* = \lambda_2^* = \dots = \lambda_m^* \in \mathbb{R}_+^n$.*

We consider distributed algorithms for GNE seeking of the game in (2). Namely, no player has direct access to the average decision \bar{x} . Instead, each player has to construct a local estimate of the average through local interactions with its neighbors. We describe the local interaction using a weight matrix $L = \{L_{ij}\}$, where $L_{ij} > 0$ if player j and player i can directly communicate with each other, and $L_{ij} = 0$ otherwise. For a player $i \in [m]$, its neighbor set \mathbb{N}_i is defined as the collection of players j such that $L_{ij} > 0$. We define $L_{ii} \triangleq -\sum_{j \in \mathbb{N}_i} L_{ij}$ for all $i \in [m]$, where \mathbb{N}_i is the neighbor set of player i . Furthermore, We make the following assumption on L :

Assumption 4. *The matrix $L = \{w_{ij}\} \in \mathbb{R}^{m \times m}$ is symmetric and satisfies $\mathbf{1}^T L = \mathbf{0}^T$, $L \mathbf{1} = \mathbf{0}$, and $\|I + L - \frac{\mathbf{1}\mathbf{1}^T}{m}\| < 1$.*

Assumption 4 ensures that the interaction graph induced by L is connected, i.e., there is a path from each player to every other player. It can be verified that $\|I + L - \frac{\mathbf{1}\mathbf{1}^T}{m}\| = \max\{|1 + \rho_2|, |1 + \rho_m|\}$, where $\{\rho_i, i \in [m]\}$ are the eigenvalues of L , with $\rho_m \leq \dots \leq \rho_2 \leq \rho_1 = 0$.

We also need the following lemma for convergence analysis:

Lemma 2. [47] *Let $\{v^k\}, \{\alpha^k\}$, and $\{p^k\}$ be random non-negative scalar sequences, and $\{q^k\}$ be a deterministic non-negative scalar sequence satisfying $\sum_{k=0}^{\infty} \alpha^k < \infty$ a.s., $\sum_{k=0}^{\infty} q^k = \infty$, $\sum_{k=0}^{\infty} p^k < \infty$ a.s., and*

$$\mathbb{E}[v^{k+1} | \mathcal{F}^k] \leq (1 + \alpha^k - q^k)v^k + p^k, \quad \forall k \geq 0 \quad \text{a.s.}$$

where $\mathcal{F}^k = \{v^\ell, \alpha^\ell, p^\ell; 0 \leq \ell \leq k\}$. Then, $\sum_{k=0}^{\infty} q^k v^k < \infty$ and $\lim_{k \rightarrow \infty} v^k = 0$ hold almost surely.

C. On Differential Privacy

We use the notion of ϵ -DP for continuous bit streams [48], which has recently been applied to distributed optimization (see [49] as well as our own work [47]). To enable ϵ -DP, we use the Laplace noise mechanism, i.e., we add Laplace noise to all shared messages. For a constant $\nu > 0$, we use $\text{Lap}(\nu)$ to denote a Laplace distribution of a scalar random variable with the probability density function $x \mapsto \frac{1}{2\nu} e^{-\frac{|x|}{\nu}}$.

One can verify that $\text{Lap}(\nu)$'s mean is zero and its variance is $2\nu^2$. Inspired by the formulation of distributed optimization in [49], to facilitate DP analysis, we represent the game \mathcal{P} in (2) by three parameters (K, \mathcal{J}, L) , where K defined in (3) is the domain of decision variables, $\mathcal{J} \triangleq \{J_1, \dots, J_m\}$, and L is the inter-player interaction weight matrix. Then we define adjacency between two games as follows:

Definition 1. *Two GNE seeking problems $\mathcal{P} = (K, \mathcal{J}, L)$ and $\mathcal{P}' = (K', \mathcal{J}', L')$ are adjacent if the following conditions hold:*

- $K = K'$ and $L = L'$, i.e., the domains of decision variables and the interaction weight matrices are identical;
- there exists an $i \in [m]$ such that $J_i \neq J'_i$ but $J_j = J'_j$ for all $j \in [m], j \neq i$.

Definition 1 implies that two GNE problems are adjacent if and only if one player changes its cost function (can be in an arbitrary way) while all other game characteristics are identical.

Given a distributed GNE seeking algorithm, we represent an execution of such an algorithm as \mathcal{A} , which is an infinite sequence of the iteration variable ϑ , i.e., $\mathcal{A} = \{\vartheta^0, \vartheta^1, \dots\}$. We consider adversaries that can observe all communicated messages among the players. Therefore, the observation part of an execution is the infinite sequence of shared messages, which is denoted as \mathcal{O} . Given a distributed GNE problem \mathcal{P} and an initial state ϑ^0 , we define the observation mapping as $\mathcal{R}_{\mathcal{P}, \vartheta^0}(\mathcal{A}) \triangleq \mathcal{O}$. Given a GNE problem \mathcal{P} , observation sequence \mathcal{O} , and an initial state ϑ^0 , $\mathcal{R}_{\mathcal{P}, \vartheta^0}^{-1}(\mathcal{O})$ is the set of executions \mathcal{A} that can generate the observation \mathcal{O} .

Definition 2. (ϵ -differential privacy, adapted from [49]). *For a given $\epsilon > 0$, an iterative distributed algorithm is ϵ -differentially private if for any two adjacent \mathcal{P} and \mathcal{P}' , any set of observation sequences $\mathcal{O}_s \subseteq \mathbb{O}$ (with \mathbb{O} denoting the set of all possible observation sequences), and any initial state ϑ^0 , the following relationship always holds*

$$\mathbb{P}[\mathcal{R}_{\mathcal{P}, \vartheta^0}^{-1}(\mathcal{O}_s)] \leq e^\epsilon \mathbb{P}[\mathcal{R}_{\mathcal{P}', \vartheta^0}^{-1}(\mathcal{O}_s)], \quad (8)$$

with the probability \mathbb{P} taken over the randomness over iteration processes.

ϵ -DP ensures that an adversary having access to all shared messages cannot gain information with a significant probability of any participating player's cost function. It can be seen that a smaller ϵ means a higher level of privacy protection. It is worth noting that the considered notion of ϵ -DP is more stringent than other relaxed (approximate) DP notions such as (ϵ, δ) -DP [50], zero-concentrated DP [51], or Rényi DP [52].

III. DIFFERENTIALLY-PRIVATE CONSENSUS TRACKING

Lying at the core of our distributed GNE seeking algorithm is the consensus tracking technique. However, the conventional consensus tracking algorithm (see, e.g., [20]) is sensitive to noise in that it leads to the accumulation and explosion of noise variance in the presence of persistent information-sharing noise [34]. Hence, in this section, we first propose a new robust consensus tracking algorithm. Our basic idea is

to tailor consensus tracking for ϵ -DP by adding one stepsize parameter α^k and a decaying factor χ^k (see Algorithm 1), which enable us to suppress the influence of persistent DP-noise, and hence, ensure both goals of rigorous ϵ -DP and guaranteed convergence accuracy. To our knowledge, this is the first time that both goals are achieved in consensus tracking.

Algorithm 1: Differentially-private consensus tracking with guaranteed convergence accuracy

Parameters: Weakening factor $\chi^k > 0$ and stepsize $\alpha^k > 0$.

Every player i 's input is r_i^k . Every player i maintains one state variable x_i^k , which is initialized as $x_i^0 = r_i^0$.

for $k = 1, 2, \dots$ **do**

- a) Every player j adds persistent DP-noise ζ_j^k to its state x_j^k , and then sends the obscured state $x_j^k + \zeta_j^k$ to player $i \in \mathbb{N}_j$.
- b) After receiving $x_j^k + \zeta_j^k$ from all $j \in \mathbb{N}_i$, player i updates its state as follows:

$$x_i^{k+1} = (1 - \alpha^k)x_i^k + \chi^k \sum_{j \in \mathbb{N}_i} L_{ij}(x_j^k + \zeta_j^k - x_i^k) + r_i^{k+1} - (1 - \alpha^k)r_i^k. \quad (9)$$

We make the following assumption on the DP-noise:

Assumption 5. For every $i \in [m]$ and every k , conditional on the state x_i^k , the DP-noise ζ_i^k satisfies $\mathbb{E}[\zeta_i^k | x_i^k] = 0$ and $\mathbb{E}[\|\zeta_i^k\|^2 | x_i^k] = (\sigma_i^k)^2$ for all $k \geq 0$, and

$$\sum_{k=0}^{\infty} (\chi^k)^2 \max_{i \in [m]} (\sigma_i^k)^2 < \infty, \quad (10)$$

where $\{\chi^k\}$ is the attenuation sequence from Algorithm 1. The initial random vectors satisfy $\mathbb{E}[\|x_i^0\|^2] < \infty, \forall i \in [m]$.

A. Convergence Analysis

To prove that Algorithm 1 can ensure every x_i^k to track the average target signal $\bar{r}^k \triangleq \frac{1}{m} \sum_{i=1}^m r_i^k$, we first prove that \bar{x}^k converges almost surely to \bar{r}^k under the following assumption:

Assumption 6. For every $i \in [m]$, there exist some nonnegative sequence $\{\gamma^k\}$ and a constant C such that

$$\|r_i^{k+1} - (1 - \alpha^k)r_i^k\| \leq \gamma^k C \quad (11)$$

holds, where α^k is from Algorithm 1 and $\{\gamma^k\}$ satisfies $\lim_{k \rightarrow \infty} \frac{\alpha^k}{\gamma^k} < \infty$.

Lemma 3. Under Assumptions 4, 5, \bar{x}^k in Algorithm 1 converges a.s. to \bar{r}^k if the following conditions hold:

$$\sum_{k=0}^{\infty} \alpha^k = \infty, \quad \sum_{k=0}^{\infty} (\alpha^k)^2 < \infty.$$

Proof. According to the definitions of \bar{x}^k and \bar{r}^k , we have the following relationship based on (9):

$$\bar{x}^{k+1} = (1 - \alpha^k)\bar{x}^k + \chi^k \bar{\zeta}^k + \bar{r}^{k+1} - (1 - \alpha^k)\bar{r}^k, \quad (12)$$

where $\bar{\zeta}^k \triangleq \frac{\sum_{i=1}^m |L_{ii}| \zeta_i^k}{m}$ and we have used the symmetric property of L and the fact $L_{ii} \triangleq -\sum_{j \in \mathbb{N}_i} L_{ij}$.

The preceding relationship implies

$$\begin{aligned} \|\bar{x}^{k+1} - \bar{r}^{k+1}\|^2 &= \|(1 - \alpha^k)(\bar{x}^k - \bar{r}^k) + \chi^k \bar{\zeta}^k\|^2 \\ &= (1 - \alpha^k)^2 \|\bar{x}^k - \bar{r}^k\|^2 + (\chi^k)^2 \|\bar{\zeta}^k\|^2 \\ &\quad + 2 \langle (1 - \alpha^k)(\bar{x}^k - \bar{r}^k), \chi^k \bar{\zeta}^k \rangle. \end{aligned}$$

Using the assumption that the DP-noise ζ_i^k has zero mean and variance $(\sigma_i^k)^2$ conditional on x_i^k (see Assumption 5), taking the conditional expectation, given $\mathcal{F}^k = \{x^0, \dots, x^k\}$, we obtain the following inequality for all $k \geq 0$:

$$\begin{aligned} \mathbb{E}[\|\bar{x}^{k+1} - \bar{r}^{k+1}\|^2 | \mathcal{F}^k] &\leq (1 + (\alpha^k)^2 - 2\alpha^k) \|\bar{x}^k - \bar{r}^k\|^2 \\ &\quad + \frac{\sum_{i=1}^m L_{ii}^2 (\sigma_i^k)^2 (\chi^k)^2}{m} \end{aligned}$$

where we have used the relationship $\|\bar{\zeta}^k\|^2 \leq \frac{\sum_{i=1}^m L_{ii}^2 \|\zeta_i^k\|^2}{m}$.

Under the conditions in the lemma statement, it can be seen that $\|\bar{x}^{k+1} - \bar{r}^{k+1}\|^2$ satisfies the conditions for v^k in Lemma 2, with $o^k = (\alpha^k)^2$, $q^k = 2\alpha^k$, and $p^k = \frac{\sum_{i=1}^m L_{ii}^2 (\sigma_i^k)^2 (\chi^k)^2}{m}$. Therefore, we have $\|\bar{x}^{k+1} - \bar{r}^{k+1}\|^2$ converging a.s. to zero, and hence, \bar{x}^k converging a.s. to \bar{r}^k . ■

Lemma 4. Under Assumption 4 and two positive sequences $\{\chi^k\}$ and $\{\alpha^k\}$ satisfying $\sum_{k=0}^{\infty} (\alpha^k)^2 < \infty$ and $\sum_{k=0}^{\infty} (\chi^k)^2 < \infty$, there always exists a $T > 0$ such that $\|(1 - \alpha^k)(I - \frac{\mathbf{1}\mathbf{1}^T}{m}) + \chi^k L\| < 1 - \chi^k |\rho_2|$ holds for all $k \geq T$, where ρ_2 is the second largest eigenvalue of L .

Proof. Because L is symmetric according to Assumption 4, all of its eigenvalues are real numbers. Since L 's off-diagonal entries are non-negative and its diagonal entries L_{ii} are given by $L_{ii} = -\sum_{j \in \mathbb{N}_i} L_{ij}$, the Gershgorin circle theorem implies that all eigenvalues of L are non-positive, with one of them being equal to 0. In fact, the 0 eigenvalue is simple (i.e., all the other eigenvalues are strictly negative) from Assumption 4. Arrange the eigenvalues of L as $\rho_m \leq \rho_{m-1} \leq \dots \leq \rho_2 \leq \rho_1 = 0$. It can be verified that the eigenvalues of $(1 - \alpha^k)I + \chi^k L$ are equal to $1 - \alpha^k + \chi^k \rho_m \leq 1 - \alpha^k + \chi^k \rho_{m-1} \leq \dots \leq 1 - \alpha^k + \chi^k \rho_2 \leq 1 - \alpha^k + \chi^k \rho_1 = 1 - \alpha^k$, and the eigenvalues of $(1 - \alpha^k)(I - \frac{\mathbf{1}\mathbf{1}^T}{m}) + \chi^k L = (1 - \alpha^k)I + \chi^k L + (1 - \alpha^k)\frac{\mathbf{1}\mathbf{1}^T}{m}$ are given by $\{1 - \alpha^k + \chi^k \rho_m, 1 - \alpha^k + \chi^k \rho_{m-1}, \dots, 1 - \alpha^k + \chi^k \rho_2, 0\}$, with $|\rho_m| \geq |\rho_{m-1}| \geq \dots \geq |\rho_2| > 0$. Hence, the norm of $\|(1 - \alpha^k)(I - \frac{\mathbf{1}\mathbf{1}^T}{m}) + \chi^k L\|$ is no larger than $\max\{|1 - \alpha^k + \chi^k \rho_m|, |1 - \alpha^k + \chi^k \rho_2|\}$. Further taking into account the fact that α^k and χ^k decay to zero because they are square summable, we conclude that there always exists a $T > 0$ such that $|1 - \alpha^k + \chi^k \rho_m| = 1 - \alpha^k - \chi^k |\rho_m| < 1 - \chi^k |\rho_m|$ and $|1 - \alpha^k + \chi^k \rho_2| = 1 - \alpha^k - \chi^k |\rho_2| < 1 - \chi^k |\rho_2|$ hold for all $k \geq T$. Given $|\rho_m| \geq |\rho_2|$, we have the stated result. ■

Theorem 1. Under Assumptions 4, 5, 6, if the nonnegative sequences $\{\alpha^k\}$ and $\{\chi^k\}$ in Algorithm 1 and the nonnegative sequence $\{\gamma^k\}$ in Assumption 6 satisfy

$$\sum_{k=0}^{\infty} \alpha^k = \infty, \quad \sum_{k=0}^{\infty} \chi^k = \infty, \quad \sum_{k=0}^{\infty} (\chi^k)^2 < \infty, \quad \sum_{k=0}^{\infty} \frac{(\gamma^k)^2}{\chi^k} < \infty, \quad (13)$$

then, the following results hold almost surely:

- 1) every x_i^k in Algorithm 1 converges to $\bar{r}^k = \frac{\sum_{i=1}^m r_i^k}{m}$;
- 2) $\sum_{k=0}^{\infty} \chi^k \sum_{i=1}^m \|x_i^k - \bar{x}^k\|^2 < \infty$;

$$3) \sum_{k=0}^{\infty} \gamma^k \sum_{i=1}^m \|x_i^k - \bar{x}^k\| < \infty.$$

Proof. For the convenience of analysis, we write the iterates of x_i^k on per-coordinate expressions. More specifically, for all $\ell = 1, \dots, d$, and $k \geq 0$, we define $x^k(\ell) = [[x_1^k]_\ell, \dots, [x_m^k]_\ell]^T$ where $[x_i^k]_\ell$ represents the ℓ th element of the vector x_i^k . Similarly, we define $r^k(\ell) = [[r_1^k]_\ell, \dots, [r_m^k]_\ell]^T$ and $\zeta^k(\ell) = [[\zeta_1^k]_\ell, \dots, [\zeta_m^k]_\ell]^T$. In this per-coordinate view, (9) has the following form for all $\ell = 1, \dots, d$, and $k \geq 0$:

$$x^{k+1}(\ell) = (1 - \alpha^k)x^k(\ell) + \chi^k L x^k(\ell) + \chi^k L^0 \zeta^k(\ell) + r^{k+1}(\ell) - (1 - \alpha^k)r^k(\ell). \quad (14)$$

where L^0 is obtained by replacing all diagonal entries of L with zero.

The dynamics of \bar{x}^k is governed by

$$\begin{aligned} \bar{x}^{k+1} t_\ell &= \frac{\mathbf{1}^T}{m} x^{k+1}(\ell) = \frac{\mathbf{1}^T}{m} ((1 - \alpha^k)x^k(\ell) + \chi^k L x^k(\ell)) \\ &\quad + \frac{\mathbf{1}^T}{m} (\chi^k L^0 \zeta^k(\ell) + r^{k+1}(\ell) - (1 - \alpha^k)r^k(\ell)), \end{aligned} \quad (15)$$

where $[\bar{x}^{k+1}]_\ell$ represents the ℓ -th element of \bar{x}^{k+1} .

Assumption 4 implies $\mathbf{1}^T L = 0$, which can further simplify (15) as:

$$\begin{aligned} \bar{x}^{k+1} t_\ell &= \frac{\mathbf{1}^T}{m} ((1 - \alpha^k)x^k(\ell) + \chi^k L^0 \zeta^k(\ell) \\ &\quad + r^{k+1}(\ell) - (1 - \alpha^k)r^k(\ell)). \end{aligned} \quad (16)$$

Combining the relations in (14) and (16) yields

$$\begin{aligned} x^{k+1}(\ell) - \mathbf{1}[\bar{x}^{k+1}]_\ell &= ((1 - \alpha^k)\Pi + \chi^k L) x^k(\ell) + \chi^k \Pi L^0 \zeta^k(\ell) \\ &\quad + \Pi (r^{k+1}(\ell) - (1 - \alpha^k)r^k(\ell)). \end{aligned} \quad (17)$$

where $\Pi \triangleq I - \frac{\mathbf{1}\mathbf{1}^T}{m}$.

To simplify the notation, we define

$$W^k \triangleq (1 - \alpha^k)\Pi + \chi^k L = (1 - \alpha^k)(I - \frac{\mathbf{1}\mathbf{1}^T}{m}) + \chi^k L. \quad (18)$$

Assumption 4 ensures $W^k \mathbf{1} = 0$, and hence $W^k \mathbf{1}[\bar{x}^k]_\ell = 0$ for any $1 \leq \ell \leq d$. Subtracting $W^k \mathbf{1}[\bar{x}^k]_\ell = 0$ from the right hand side of (17) yields

$$\begin{aligned} x^{k+1}(\ell) - \mathbf{1}[\bar{x}^{k+1}]_\ell &= W^k (x^k(\ell) - \mathbf{1}[\bar{x}^k]_\ell) + \chi^k \Pi L^0 \zeta^k(\ell) \\ &\quad + \Pi (r^{k+1}(\ell) - (1 - \alpha^k)r^k(\ell)), \end{aligned} \quad (19)$$

which, further implies

$$\begin{aligned} \|x^{k+1}(\ell) - \mathbf{1}[\bar{x}^{k+1}]_\ell\|^2 &= \|W^k (x^k(\ell) - \mathbf{1}[\bar{x}^k]_\ell) + \Pi (r^{k+1}(\ell) - (1 - \alpha^k)r^k(\ell))\|^2 \\ &\quad + 2 \langle W^k (x^k(\ell) - \mathbf{1}[\bar{x}^k]_\ell) + \Pi (r^{k+1}(\ell) - (1 - \alpha^k)r^k(\ell)), \\ &\quad \chi^k \Pi L^0 \zeta^k(\ell) \rangle + \|\chi^k \Pi L^0 \zeta^k(\ell)\|^2 \\ &\leq \|W^k (x^k(\ell) - \mathbf{1}[\bar{x}^k]_\ell) + \Pi (r^{k+1}(\ell) - (1 - \alpha^k)r^k(\ell))\|^2 \\ &\quad + 2 \langle W^k (x^k(\ell) - \mathbf{1}[\bar{x}^k]_\ell) + \Pi (r^{k+1}(\ell) - (1 - \alpha^k)r^k(\ell)), \\ &\quad \chi^k \Pi L^0 \zeta^k(\ell) \rangle + (\chi^k)^2 \|L^0\|^2 \|\zeta^k(\ell)\|^2. \end{aligned} \quad (20)$$

where we used $\|\Pi\| = 1$. Using the assumption that the DP-noise ζ_i^k has zero mean and variance $(\sigma_i^k)^2$ conditional on x_i^k

(see Assumption 5), taking the conditional expectation, given $\mathcal{F}^k = \{x^0, \dots, x^k\}$, we obtain the following inequality for all $k \geq 0$:

$$\begin{aligned} &\mathbb{E} [\|x^{k+1}(\ell) - \mathbf{1}[\bar{x}^{k+1}]_\ell\|^2 | \mathcal{F}^k] \\ &\leq \|W^k (x^k(\ell) - \mathbf{1}[\bar{x}^k]_\ell) + \Pi (r^{k+1}(\ell) - (1 - \alpha^k)r^k(\ell))\|^2 \\ &\quad + (\chi^k)^2 \|L^0\|^2 \mathbb{E} [\|\zeta^k(\ell)\|^2] \\ &\leq (\|W^k\| \|x^k(\ell) - \mathbf{1}[\bar{x}^k]_\ell\| + \|r^{k+1}(\ell) - (1 - \alpha^k)r^k(\ell)\|)^2 \\ &\quad + (\chi^k)^2 \|L^0\|^2 \mathbb{E} [\|\zeta^k(\ell)\|^2]. \end{aligned} \quad (21)$$

For the first term on the right hand side of the preceding inequality, we bound it using the fact that there exists a $T \geq 0$ such that $0 < \|W^k\| \leq 1 - \chi^k |\rho_2|$ holds for all $k \geq T$ (see Lemma 4). Hence, equation (21) implies the following relationship for all $k \geq T$:

$$\begin{aligned} &\mathbb{E} [\|x^{k+1}(\ell) - \mathbf{1}[\bar{x}^{k+1}]_\ell\|^2 | \mathcal{F}^k] \leq \\ &((1 - \chi^k |\rho_2|) \|x^k(\ell) - \mathbf{1}[\bar{x}^k]_\ell\| + \|r^{k+1}(\ell) - (1 - \alpha^k)r^k(\ell)\|)^2 \\ &\quad + (\chi^k)^2 \|L^0\|^2 \mathbb{E} [\|\zeta^k(\ell)\|^2]. \end{aligned} \quad (22)$$

Next, we apply the inequality $(a + b)^2 \leq (1 + \epsilon)a^2 + (1 + \epsilon^{-1})b^2$, which is valid for any scalars a, b , and $\epsilon > 0$, to (22):

$$\begin{aligned} &\mathbb{E} [\|x^{k+1}(\ell) - \mathbf{1}[\bar{x}^{k+1}]_\ell\|^2 | \mathcal{F}^k] \leq (\chi^k)^2 \|L^0\|^2 \mathbb{E} [\|\zeta^k(\ell)\|^2] \\ &\quad + (1 + \epsilon)(1 - \chi^k |\rho_2|)^2 \|x^k(\ell) - \mathbf{1}[\bar{x}^k]_\ell\|^2 \\ &\quad + (1 + \epsilon^{-1}) \|r^{k+1}(\ell) - (1 - \alpha^k)r^k(\ell)\|^2. \end{aligned} \quad (23)$$

Setting $\epsilon = \frac{\chi^k |\rho_2|}{1 - \chi^k |\rho_2|}$ (which leads to $(1 + \epsilon) = \frac{1}{1 - \chi^k |\rho_2|}$ and $1 + \epsilon^{-1} = \frac{1}{\chi^k |\rho_2|}$) yields

$$\begin{aligned} &\mathbb{E} [\|x^{k+1}(\ell) - \mathbf{1}[\bar{x}^{k+1}]_\ell\|^2 | \mathcal{F}^k] \leq (\chi^k)^2 \|L^0\|^2 \mathbb{E} [\|\zeta^k(\ell)\|^2] \\ &\quad + (1 - \chi^k |\rho_2|) \|x^k(\ell) - \mathbf{1}[\bar{x}^k]_\ell\|^2 \\ &\quad + \frac{1}{\chi^k |\rho_2|} \|r^{k+1}(\ell) - (1 - \alpha^k)r^k(\ell)\|^2. \end{aligned} \quad (24)$$

Note that the following relations always hold: $\sum_{\ell=1}^d \|x^k(\ell) - [\bar{x}^k]_\ell \mathbf{1}\|^2 = \sum_{i=1}^m \|x_i^k - \bar{x}^k\|^2$, $\sum_{\ell=1}^d \|x^{k+1}(\ell) - x^k(\ell)\|^2 = \sum_{i=1}^m \|x_i^{k+1} - x_i^k\|^2$, $\sum_{\ell=1}^d \|\zeta^k(\ell)\|^2 = \sum_{i=1}^m \|\zeta_i^k\|^2$.

Hence, summing (24) over $\ell = 1, \dots, d$ leads to

$$\begin{aligned} &\mathbb{E} \left[\sum_{i=1}^m \|x_i^{k+1} - \bar{x}^{k+1}\|^2 | \mathcal{F}^k \right] \\ &\leq (\chi^k)^2 \|L^0\|^2 \mathbb{E} \left[\sum_{i=1}^m \|\zeta_i^k\|^2 \right] + (1 - \chi^k |\rho_2|) \sum_{i=1}^m \|x_i^k - \bar{x}^k\|^2 \\ &\quad + \frac{1}{\chi^k |\rho_2|} \sum_{i=1}^m \|r_i^{k+1} - (1 - \alpha^k)r_i^k\|^2 \\ &\leq (\chi^k)^2 \|L^0\|^2 \sum_{i=1}^m (\sigma_i^k)^2 + (1 - \chi^k |\rho_2|) \sum_{i=1}^m \|x_i^k - \bar{x}^k\|^2 \\ &\quad + \frac{1}{\chi^k |\rho_2|} \sum_{i=1}^m \|r_i^{k+1} - (1 - \alpha^k)r_i^k\|^2. \end{aligned} \quad (25)$$

Using the assumption $\|r_i^{k+1} - (1 - \alpha^k)r_i^k\| \leq \gamma^k C$ for all $i \in [m]$ from Assumption 6, we obtain

$$\frac{1}{\chi^k |\rho_2|} \sum_{i=1}^m \|r_i^{k+1} - (1 - \alpha^k)r_i^k\|^2 \leq \frac{(\gamma^k)^2}{\chi^k |\rho_2|} m C^2.$$

Submitting the preceding relationship into (25) yields

$$\begin{aligned} \mathbb{E} \left[\sum_{i=1}^m \|x_i^{k+1} - \bar{x}^{k+1}\|^2 \mathcal{F}^k \right] &\leq (\chi^k)^2 \|L^0\|^2 \sum_{i=1}^m (\sigma_i^k)^2 \\ &+ (1 - \chi^k |\rho_2|) \sum_{i=1}^m \|x_i^k - \bar{x}^k\|^2 + \frac{(\gamma^k)^2}{\chi^k |\rho_2|} m C^2. \end{aligned} \quad (26)$$

Therefore, under Assumption 5 and the conditions for χ^k and γ^k in (13), we have that the sequence $\{\sum_{i=1}^m \|x_i^k - \bar{x}^k\|^2\}$ satisfies the conditions for $\{v^k\}$ in Lemma 2, and hence, converges to zero almost surely. So x_i^k converges to \bar{x}^k almost surely. Further recalling \bar{x}^k converging a.s. to \bar{r}^k in Lemma 3 yields that x_i^k converges a.s. to \bar{r}^k .

Moreover, Lemma 2 also implies the following relation a.s.:

$$\sum_{k=0}^{\infty} \chi^k \sum_{i=1}^m \|x_i^k - \bar{x}^k\|^2 < \infty. \quad (27)$$

To prove the last statement, we invoke the Cauchy–Schwarz inequality, which ensures

$$\begin{aligned} \sum_{k=0}^{\infty} \sqrt{\chi^k \sum_{i=1}^m \|x_i^k - \bar{x}^k\|^2} \sqrt{\frac{(\gamma^k)^2}{\chi^k}} \\ \leq \sqrt{\sum_{k=0}^{\infty} \chi^k \sum_{i=1}^m \|x_i^k - \bar{x}^k\|^2} \sqrt{\sum_{k=0}^{\infty} \frac{(\gamma^k)^2}{\chi^k}}. \end{aligned}$$

Noting that the summand in the left hand side of the preceding inequality is actually $\gamma^k \sqrt{\sum_{i=1}^m \|x_i^k - \bar{x}^k\|^2}$, and the right hand side of the preceding inequality is less than infinity almost surely under the proven result in (27) and the assumption in (13), we have that $\sum_{k=0}^{\infty} \gamma^k \sqrt{\sum_{i=1}^m \|x_i^k - \bar{x}^k\|^2} < \infty$ holds almost surely. Further utilizing the relationship $\sum_{i=1}^m \|x_i^k - \bar{x}^k\| \leq \sqrt{m \sum_{i=1}^m \|x_i^k - \bar{x}^k\|^2}$ yields the stated result. ■

Next, we prove that Algorithm 1 can ensure the ϵ -DP of individual agents' tracking variables r_i^k , even when the number of iterations tends to infinity.

B. Differential-privacy Analysis

To analyze the level of DP protection on individual r_i^k , we first note that the adjacency defined in Definition 1 can be easily adapted to the consensus tracking problem by replacing J_i with r_i . Specifically, we can define two adjacent consensus-tracking problems as \mathcal{P} and \mathcal{P}' , with the only difference being one tracking target (represent it as r_i in \mathcal{P} and r'_i in \mathcal{P}' without loss of generality). From the theory of ϵ -DP, we have to characterize the sensitivity of an algorithm in order to quantify the amount of DP-noise that has to be injected to shared messages to enable a certain level of ϵ -DP. Similar to the sensitivity definition of iterative optimization algorithms

in [49], we define the sensitivity of a consensus-tracking algorithm as follows:

Definition 3. At each iteration k , for any initial state ϑ^0 and any adjacent consensus-tracking problems \mathcal{P} and \mathcal{P}' , the sensitivity of Algorithm 1 is

$$\Delta^k \triangleq \sup_{\mathcal{O} \in \mathbb{O}} \left\{ \sup_{\vartheta \in \mathcal{R}_{\mathcal{P}, \vartheta^0}^{-1}(\mathcal{O}), \vartheta' \in \mathcal{R}_{\mathcal{P}', \vartheta^0}^{-1}(\mathcal{O})} \|\vartheta^{k+1} - \vartheta'^{k+1}\|_1 \right\}. \quad (28)$$

Then, we have the following lemma:

Lemma 5. In Algorithm 1, at each iteration k , if each agent's DP-noise vector $\zeta_i^k \in \mathbb{R}^d$ consists of d independent Laplace noises with parameter ν^k such that $\sum_{k=1}^{T_0} \frac{\Delta^k}{\nu^k} \leq \bar{\epsilon}$, then Algorithm 1 is ϵ -differentially private with the cumulative privacy budget from iteration $k = 0$ to $k = T_0$ less than $\bar{\epsilon}$.

Proof. The lemma follows the same line of reasoning of Lemma 2 in [49] (also see Theorem 3 in [30]). ■

Theorem 2. Under the conditions of Theorem 1, if all elements of ζ_i^k are drawn independently from Laplace distribution $\text{Lap}(\nu^k)$ with $(\sigma_i^k)^2 = 2(\nu^k)^2$ satisfying Assumption 5, then all agents will converge a.s. to the average tracking target \bar{r}^k . Moreover,

- 1) Algorithm 1 is ϵ -differentially private with the cumulative privacy budget bounded by $\epsilon \leq \sum_{k=1}^{T_0} \frac{2\bar{C}\gamma^k}{\nu^k}$ for iterations from $k = 0$ to $k = T_0$, where $\bar{C} = \sqrt{d}C$ with C a constant from (11) and d the dimension of x_i^k . And the cumulative privacy budget is always finite for $T_0 \rightarrow \infty$ when the sequence $\{\frac{\gamma^k}{\nu^k}\}$ is summable, i.e., $\sum_{k=0}^{\infty} \frac{\gamma^k}{\nu^k} < \infty$;
- 2) Suppose that two non-negative sequences $\{\nu^k\}$ and $\{\gamma^k\}$ have a finite sequence-ratio sum $\Phi_{\gamma, \nu} \triangleq \sum_{k=1}^{\infty} \frac{\gamma^k}{\nu^k}$. Then setting the Laplace noise parameter ν^k as $\nu^k = \frac{2\bar{C}\Phi_{\gamma, \nu'}}{\epsilon} \nu^k$ ensures that Algorithm 1 is ϵ -differentially private with any cumulative privacy budget $\epsilon > 0$ even when the number of iterations goes to infinity;
- 3) In the special case where $\gamma^k = \frac{1}{k}$, setting $\nu^k = \frac{2\bar{C}\Phi}{\epsilon} k^{0.3}$ with $\Phi \triangleq \sum_{k=1}^{\infty} \frac{1}{k^{1.3}} \approx 3.93$ (which can be verified to satisfy Assumption 5) ensures that Algorithm 1 is always ϵ -differentially private for any cumulative privacy budget $\epsilon > 0$ even when the number of iterations goes to infinity.

Proof. Since the Laplace noise satisfies Assumption 5, the convergence result follows directly from Theorem 1.

To prove the three statements on the strength of ϵ -DP, we first prove that the sensitivity of Algorithm 1 satisfies $\Delta^k \leq 2\bar{C}\gamma^k$. Given two adjacent consensus-tracking problems \mathcal{P} and \mathcal{P}' , for any given fixed observation \mathcal{O} and initial state $\vartheta^0 = x^0$, the sensitivity is determined by $\|\mathcal{R}_{\mathcal{P}, x^0}^{-1}(\mathcal{O}) - \mathcal{R}_{\mathcal{P}', x^0}^{-1}(\mathcal{O})\|_1$ according to Definition 3. Since in \mathcal{P} and \mathcal{P}' , there is only one tracking target that is different, we represent this different tracking target as the i th one, i.e., r_i and r'_i in \mathcal{P} and \mathcal{P}' , respectively, without loss of generality. We define $o^k \triangleq (1 - \alpha^k)x_i^k + \sum_{j \in \mathbb{N}_i} \chi^k w_{ij}(x_j^k - x_i^k)$ and $o'^k \triangleq (1 - \alpha^k)x_i'^k + \sum_{j \in \mathbb{N}_i} \chi^k w_{ij}(x_j'^k - x_i'^k)$, which are accessible to adversaries

under r_i and r'_i , respectively (note that the sensitivity analysis is independent of DP-noise [29]). Because the observations under \mathcal{P} and \mathcal{P}' are identical, o^k and o'^k have to be the same according to the sensitivity definition. Therefore, we have the following relationship:

$$\begin{aligned} & \|\mathcal{R}_{\mathcal{P},x^0}^{-1}(\mathcal{O}) - \mathcal{R}_{\mathcal{P}',x^0}^{-1}(\mathcal{O})\|_1 \\ &= \left\| o^k + r_i^{k+1} - (1 - \alpha^k)r_i^k - \left(o'^k + r_i'^{k+1} - (1 - \alpha^k)r_i'^k \right) \right\|_1 \\ &= \left\| r_i^{k+1} - (1 - \alpha^k)r_i^k - \left(r_i'^{k+1} - (1 - \alpha^k)r_i'^k \right) \right\|_1 \\ &\leq \|r_i^{k+1} - (1 - \alpha^k)r_i^k\|_1 + \|r_i'^{k+1} - (1 - \alpha^k)r_i'^k\|_1. \end{aligned} \quad (29)$$

According to Assumption 6, in \mathcal{P} , $\|r_i^{k+1} - (1 - \alpha^k)r_i^k\| \leq \gamma^k C$ holds for all i , which implies

$$\|r_i^{k+1} - (1 - \alpha^k)r_i^k\|_1 \leq \sqrt{d} \|r_i^{k+1} - (1 - \alpha^k)r_i^k\| \leq \gamma^k \bar{C}.$$

where $\bar{C} = \sqrt{d}C$. Similar results can be obtained for \mathcal{P}' :

$$\|r_i'^{k+1} - (1 - \alpha^k)r_i'^k\|_1 \leq \sqrt{d} \|r_i'^{k+1} - (1 - \alpha^k)r_i'^k\| \leq \gamma^k \bar{C}.$$

Plugging the preceding two inequalities into (29) leads to

$$\|\mathcal{R}_{\mathcal{P},x^0}^{-1}(\mathcal{O}) - \mathcal{R}_{\mathcal{P}',x^0}^{-1}(\mathcal{O})\|_1 \leq 2\gamma^k \bar{C}.$$

From Lemma 5, the cumulative privacy budget is always less than $\sum_{k=1}^{T_0} \frac{2\bar{C}\gamma^k}{\nu^k}$. Hence, the cumulative privacy budget ϵ will always be finite even when T_0 tends to infinity if the sequence $\{\frac{\gamma^k}{\nu^k}\}$ is summable, i.e., $\sum_{k=0}^{\infty} \frac{\gamma^k}{\nu^k} < \infty$.

Lemma 5 implies that the cumulative privacy budget is inversely proportional to ν^k . Hence, the second statement can be obtained from the first statement by scaling ν^k proportionally. The result in the third statement can be obtained by specializing the selection of γ^k and ν^k sequences. ■

Note that in consensus-tracking applications such as constraint-free Nash equilibrium seeking, [30] achieves ϵ -DP by enforcing the tracking input to be summable (geometrically-decreasing, more specifically), which, however, also makes it impossible to ensure accurate convergence to the desired equilibrium. In our approach, by allowing the tracking input $r_i^{k+1} - (1 - \alpha^k)r_i^k$ to be non-summable (since we allow $\sum_{k=0}^{\infty} \gamma^k = \infty$), we achieve both accurate convergence and finite cumulative privacy budget, even when the number of iterations goes to infinity. To our knowledge, this is the first consensus tracking algorithm that can achieve both almost-sure convergence to a tracking target and rigorous ϵ -DP, even with the number of iterations going to infinity.

Remark 1. To ensure that the cumulative privacy budget $\epsilon = \sum_{k=1}^{\infty} \frac{2\bar{C}\gamma^k}{\nu^k}$ is bounded, we employ Laplace noise with parameter ν^k increasing with time (since we require the sequence $\{\frac{\gamma^k}{\nu^k}\}$ to be summable while the sequence $\{\gamma^k\}$ is non-summable). Because the strength of the shared signal is time-invariant (i.e., x_i^k), an increasing ν^k makes the relative level between noise ζ_i^k and signal x_i^k increase with time. However, since it is $\chi^k \text{Lap}(\nu^k)$ that is actually fed into the algorithm, and the increase in the noise level ν^k is outweighed by the decrease of χ^k (see Assumption 5), the actual noise fed into the algorithm still decays with time, which explains why Algorithm 1 can ensure every agent's accurate convergence.

Moreover, according to Theorem 1, the convergence will not be affected if we scale ν^k by any constant $\frac{1}{\epsilon} > 0$ to achieve any desired level of ϵ -DP, as long as ν^k (with associated variance $(\sigma_i^k)^2 = 2(\nu^k)^2$) satisfies Assumption 5.

IV. DIFFERENTIALLY-PRIVATE GNE SEEKING

Based on our differentially-private consensus tracking algorithm, we propose a differentially-private GNE seeking algorithm that can ensure both rigorous ϵ -DP and provable convergence to the GNE, which is summarized in Algorithm 2. Inspired by the distributed GNE seeking algorithm in [3], we let each player maintain three estimates of the global information that are not locally available. More specifically, σ_i^k is used to track the average \bar{x}^k of the decision variables x_i^k of all players, z_i^k is used to track the average $\bar{\lambda}^k$ of the dual variables λ_i^k of all players, and y_i^k is used to track the average \bar{d}^k of the violation of the coupling constraint, which is defined as

$$d_i^k \triangleq 2C_i \tilde{x}_i^k - C_i x_i^k - c_i \quad (30)$$

for player i (please see (31) for \tilde{x}_i^k).

Algorithm 2: Differentially-private GNE seeking algorithm with guaranteed convergence accuracy

Parameters: Weakening factor $\chi^k > 0$ and stepsizes $\alpha^k, \beta^k, \gamma^k > 0$.

Every player i maintains a decision variable x_i^k and an auxiliary variable \tilde{x}_i^k , both of which are initialized randomly in $\Omega_i \subseteq \mathbb{R}^d$. Player i also maintains a dual variable λ_i^k , which is initialized randomly in \mathbb{R}_+^n . Every player i also maintains local estimates (represented by σ_i^k , z_i^k , and y_i^k) of global information \bar{x}^k , $\bar{\lambda}^k$, and \bar{d}^k . These estimates are initialized as $\sigma_i^0 = x_i^0$, $z_i^0 = \lambda_i^0$, and $y_i^0 = 2C_i \tilde{x}_i^{-1} - C_i x_i^{-1} - c_i$, respectively.

for $k = 0, 1, \dots$ **do**

- Every player j adds persistent DP-noises ζ_j^k , ξ_j^k , and ϑ_j^k to σ_j^k , y_j^k , and z_j^k , respectively, and then sends the obscured estimates $\sigma_j^k + \zeta_j^k$, $y_j^k + \xi_j^k$, and $z_j^k + \vartheta_j^k$ to player $i \in \mathbb{N}_j$.
- After receiving $\sigma_j^k + \zeta_j^k$, $y_j^k + \xi_j^k$, and $z_j^k + \vartheta_j^k$ from all $j \in \mathbb{N}_i$, player i updates its decision variable and estimates as follows:

$$\begin{aligned} \tilde{x}_i^k &= \Pi_{\Omega_i} [x_i^k - \alpha^k (F_i(x_i^k, \sigma_i^k) + C_i^T z_i^k)], \\ y_i^{k+1} &= (1 - \gamma^k) y_i^k + \chi^k \sum_{j \in \mathbb{N}_i} L_{ij} (y_j^k + \xi_j^k - y_i^k) \\ &\quad + C_i (2\tilde{x}_i^k - x_i^k - c_i) - (1 - \gamma^k) C_i (2\tilde{x}_i^{k-1} - x_i^{k-1} - c_i), \\ \tilde{\lambda}_i^k &= \Pi_{\mathbb{R}_+^n} [\lambda_i^k + \beta^k (y_i^{k+1} - \lambda_i^k + z_i^k)], \\ x_i^{k+1} &= x_i^k + \gamma^k (\tilde{x}_i^k - x_i^k), \\ \lambda_i^{k+1} &= \lambda_i^k + \gamma^k (\tilde{\lambda}_i^k - \lambda_i^k), \\ \sigma_i^{k+1} &= (1 - \gamma^k) \sigma_i^k + \chi^k \sum_{j \in \mathbb{N}_i} L_{ij} (\sigma_j^k + \zeta_j^k - \sigma_i^k) \\ &\quad + x_i^{k+1} - (1 - \gamma^k) x_i^k, \\ z_i^{k+1} &= (1 - \gamma^k) z_i^k + \chi^k \sum_{j \in \mathbb{N}_i} L_{ij} (z_j^k + \vartheta_j^k - z_i^k) \\ &\quad + z_i^{k+1} - (1 - \gamma^k) z_i^k. \end{aligned} \quad (31)$$

where $\Pi_{\diamond}[\cdot]$ denotes the Euclidean projection of a vector onto a set \diamond .

c) **end**

Remark 2. Note that different from [28] where players use correlated noise which restricts the strength of privacy protection, here the noises $\zeta_i^k, \xi_i^k, v_i^k$ ($i = 1, 2, \dots, m$) of all players are completely independent of each other, and hence can enable strong privacy protection.

The sequence $\{\chi^k\}$, which diminishes with time, is used to suppress the influence of persistent DP-noises ζ_i^k, ξ_i^k , and v_i^k on the convergence point of the iterates. The stepsize sequences $\{\alpha^k\}, \{\beta^k\}, \{\gamma^k\}$ and attenuation sequence $\{\chi^k\}$ have to be designed appropriately to guarantee the accurate convergence of the iterate vector $x^k \triangleq \text{col}(x_1^k, \dots, x_m^k)$ to the GNE $x^* \triangleq \text{col}(x_1^*, \dots, x_m^*)$. The persistent DP noise sequences $\{\zeta_i^k\}, \{\xi_i^k\}, \{v_i^k\}, i \in [m]$ have zero-means and χ^k -bounded (conditional) variances, which are specified by the assumption below:

Assumption 7. In Algorithm 2, for every $i \in [m]$, the noise sequences $\{\zeta_i^k\}, \{\xi_i^k\}$, and $\{v_i^k\}$ are zero-mean independent random variables, and independent of $\{\sigma_i^0, z_i^0, y_i^0; i \in [m]\}$. Also, for every k , the noise collection $\{\zeta_j^k, \xi_j^k, v_j^k; j \in [m]\}$ is independent. The noise variances $(\sigma_{\zeta,i}^k)^2 = \mathbb{E}[\|\zeta_i^k\|^2]$, $(\sigma_{\xi,i}^k)^2 = \mathbb{E}[\|\xi_i^k\|^2]$, and $(\sigma_{v,i}^k)^2 = \mathbb{E}[\|v_i^k\|^2]$ satisfy

$$\sum_{k=0}^{\infty} (\chi^k)^2 \max_{i \in [m]} \{ \max \{ (\sigma_{\zeta,i}^k)^2, (\sigma_{\xi,i}^k)^2, (\sigma_{v,i}^k)^2 \} \} < \infty \quad (32)$$

The initial random vectors satisfy $\max\{\mathbb{E}[\|x_i^0\|^2], \mathbb{E}[\|z_i^0\|^2], \mathbb{E}[\|y_i^0\|^2]\} < \infty, \forall i \in [m]$.

V. CONVERGENCE ANALYSIS OF ALGORITHM 2

Our basic idea of convergence analysis is to leverage the framework of stochastically-perturbed nonstationary fixed-point iteration processes. Recently, the authors in [53] show that stochastically-perturbed nonstationary fixed-point iteration processes with nonexpansive operators can still converge to a fixed point if the iteration stepsize sequence $\{\gamma^k\}$ is lower bounded away from 0, i.e., $\liminf_{k \rightarrow \infty} \gamma^k > 0$. In our case, since the DP-noises are persistent to ensure a strong privacy protection, the stepsize has to decay to zero to ensure a finite privacy budget. Hence, we extend the result in [53] to the case where the iteration stepsize is allowed to decay to zero. Interestingly, we prove that almost sure convergence to a fixed point can still be achieved in this case. It is worth noting that our results generalize the Krasnosel'skiĭ-Mann iteration process in [40] which addresses a time-invariant operator T , and the inexact Krasnosel'skiĭ-Mann iteration process in [3], [54] which only addresses deterministic errors.

We first prove that a stochastically-perturbed nonstationary fixed-point iteration process can still ensure almost sure convergence to a fixed point, even when the iteration stepsize decays to zero:

Lemma 6. For every $k \geq 0$, let $R_k : \mathbb{R}^p \rightarrow \mathbb{R}^p$ be nonexpansive operators such that $\mathbf{F} \triangleq \bigcap_{k \in \mathbb{N}} \text{Fix} R_k \neq \emptyset$. Then, under iteration

$$\omega^{k+1} = \omega^k + \gamma^k (R_k(\omega^k) + e^k - \omega^k), \quad (33)$$

ω^k converges almost surely to some $\omega^* \in \mathbf{F}$ if the following conditions are satisfied:

- 1) The nonnegative sequence $\{\gamma^k\}$ satisfies $\sum_{k=0}^{\infty} \gamma^k (1 - \gamma^k) = +\infty$;
- 2) The error sequence $\{e^k\}$ satisfies $\sum_{k=0}^{\infty} \gamma^k \sqrt{\mathbb{E}[\|e^k\|^2 | \mathcal{F}^k]} < \infty$ almost surely, where $\mathcal{F}^k = \{\omega^\ell; 0 \leq \ell \leq k\}$;
- 3) For every $\{\omega^k\}$, $\lim_{k \rightarrow \infty} \|R_k(\omega^k) - \omega^k\|$ exists almost surely;
- 4) The cluster points of $\{\omega^k\}$ belong to \mathbf{F} .

Proof. This lemma is an adaptation of Corollary 2.7 and Theorem 3.2 of [53]. More specifically, we use the almost sure convergence of $\lim_{k \rightarrow \infty} \|R_k(x^k) - x^k\|$ in condition 3) to replace the requirement of $\liminf_{k \rightarrow \infty} \gamma^k > 0$ in Theorem 3.2 of [53], which serves the same purpose in the derivation. ■

Remark 3. Different from [53] which requires the stepsize γ^k in nonstationary fixed-point iteration processes to be bounded away from 0, Lemma 6 allows the stepsize to decay to zero. This difference is significant in that allowing the stepsize to decay to zero is key to ensuring the privacy budget to be bounded, and hence, enable rigorous ϵ -DP, as detailed later in Theorem 4.

In the remainder of the convergence analysis, we show that Algorithm 2 is indeed a stochastically-perturbed version of a nonstationary fixed-point iteration process. Hence, we can leverage Lemma 6 to establish the almost sure convergence of Algorithm 2 under persistent DP-noises. The main challenges lie in 1) finding a sequence of deterministic nonexpansive operators $R_k(\cdot)$ that have convergent properties as required in the third and fourth conditions in Lemma 6, and 2) proving that Algorithm 2 is a stochastically-perturbed version of $R_k(\cdot)$ with the amplitude of perturbation e^k satisfying the almost sure summability condition in condition 2) of Lemma 6. We address the two challenges in Sec. V-A and Sec. V-B, respectively.

A. Finding a non-perturbed fixed-point iteration process

In this subsection, we show that when the estimates σ_i^k, y_i^k , and z_i^k in Algorithm 2 are replaced with their target signals that they are designed to track, i.e., \bar{x}^k, \bar{d}^k , and $\bar{\lambda}^k$, then the iterations in Algorithm 2 reduce to a nonstationary fixed-point iteration process with involved operators being nonexpansive and satisfying the conditions 3) and 4) in Lemma 6, if $\{\alpha^k\}$ and $\{\beta^k\}$ are set as nonnegative sequences satisfying $\sum_{k=0}^{\infty} \alpha^k = \infty, \sum_{k=0}^{\infty} (\alpha^k)^2 < \infty, \sum_{k=0}^{\infty} \beta^k = \infty$, and $\sum_{k=0}^{\infty} (\beta^k)^2 < \infty$. Note that replacing σ_i^k, y_i^k , and z_i^k with \bar{x}^k, \bar{d}^k , and $\bar{\lambda}^k$ corresponds to the full-decision information setting where a central coordinator has direct access to all x_i^k, d_i^k , and λ_i^k for $i \in [m]$, and hence, can compute and disperse the respective averages to all players. The reduced algorithm is summarized in Algorithm 3:

Algorithm 3: GNE seeking with full-decision information

Parameters: Stepsizes $\alpha^k > 0$, $\beta^k > 0$, and $\gamma^k > 0$.

Every player i maintains one decision variable x_i^k , which is initialized randomly in $\Omega_i \subseteq \mathbb{R}^d$. Every player i also maintains a dual variable λ_i^k , which is initialized randomly in \mathbb{R}_+^n .

for $k = 0, 1, \dots$ **do**

a) Every player i calculates

$$\begin{aligned} \tilde{x}_i^k &= \Pi_{\Omega_i} [x_i^k - \alpha^k (F_i(x_i^k, \bar{x}^k) + C_i^T \bar{\lambda}^k)], \\ d_i^k &= 2C_i \tilde{x}_i^k - C_i x_i^k - c_i, \\ \bar{\lambda}_i^k &= \Pi_{\mathbb{R}_+^n} [\lambda_i^k + \beta^k (\bar{d}_i^k - \lambda_i^k + \bar{\lambda}_i^k)]. \end{aligned} \quad (34)$$

b) Player i updates its decision variable and dual variable as follows:

$$\begin{aligned} x_i^{k+1} &= x_i^k + \gamma^k (\tilde{x}_i^k - x_i^k), \\ \lambda_i^{k+1} &= \lambda_i^k + \gamma^k (\bar{\lambda}_i^k - \lambda_i^k). \end{aligned} \quad (35)$$

c) **end**

Lemma 7. *Algorithm 3 corresponds to a nonstationary fixed-point iteration process*

$$\omega^{k+1} = \omega^k + \gamma^k (R_k(\omega^k) - \omega^k), \quad (36)$$

where $\omega = \text{col}(x^k, \lambda^k)$, and the operators R_k are given by

$$R_k = (\text{Id} + (\Phi^k)^{-1} T_2)^{-1} \circ (\text{Id} - (\Phi^k)^{-1} T_1) \quad (37)$$

with T_1 and T_2 given in (7) and

$$\Phi^k = \begin{bmatrix} (\alpha^k)^{-1} I_{md} & -\frac{C_f^T}{m} \\ -\frac{C_f}{m} & (\beta^k)^{-1} I_{mn} \end{bmatrix}. \quad (38)$$

Moreover, when $\sum_{k=0}^{\infty} \alpha^k = \infty$, $\sum_{k=0}^{\infty} \beta^k = \infty$, $\sum_{k=0}^{\infty} (\alpha^k)^2 < \infty$, $\sum_{k=0}^{\infty} (\beta^k)^2 < \infty$, and

$$\alpha^k \leq \frac{m}{2 \max_{i \in [m]} \|C_i\|}, \quad \beta^k \leq \frac{m}{2 \max_{i \in [m]} \|C_i\|} \quad (39)$$

hold, then $\{R_k\}$ are nonexpansive operators and the conditions 3) and 4) in Lemma 6 are satisfied.

Proof. The proof that Algorithm 3 corresponds to a nonstationary fixed-point iteration process (36) follows the forward-backward operator splitting approach in [2], [3]. More specifically, Algorithm 3 can be rewritten in the following compact form:

$$\begin{aligned} \tilde{x}^k &= \Pi_{\Omega} [x^k - \alpha^k (F(x^k, \bar{x}_1^k) + C_d^T \bar{\lambda}_1^k)], \\ \bar{\lambda}^k &= \Pi_{\mathbb{R}_+^{mn}} [\lambda^k - \beta^k (\bar{d}_1^k - \lambda^k + \bar{\lambda}_1^k)], \\ x^{k+1} &= x^k + \gamma^k (\tilde{x}^k - x^k), \\ \lambda^{k+1} &= \lambda^k + \gamma^k (\bar{\lambda}^k - \lambda^k), \end{aligned} \quad (40)$$

where $\bar{x}_1^k = \mathbf{1} \otimes \bar{x}^k$, $\bar{\lambda}_1^k = \mathbf{1} \otimes \bar{\lambda}^k$, $\bar{d}_1^k = \mathbf{1} \otimes \bar{d}^k$, and $C_d = \text{diag}(C_1, \dots, C_m)$.

Using the relations $\Pi_{\Omega} = (\text{Id} + N_{\Omega})^{-1}$, $F(x^k, \bar{x}_1^k) = F(x)$, and $C_d^T \bar{\lambda}_1^k = C_d^T (\mathbf{1} \otimes \bar{\lambda}^k) = \frac{C_f^T \lambda^k}{m}$, we can obtain the following relationship from the first equality in (40):

$$(\text{Id} + N_{\Omega})(\tilde{x}^k) \ni x^k - \alpha^k \left(F(x^k) + \frac{C_f^T \lambda^k}{m} \right).$$

Given $\frac{1}{\alpha^k} N_{\Omega}(\tilde{x}^k) = N_{\Omega}(\tilde{x}^k)$, the preceding relationship implies

$$-F(x^k) \ni N_{\Omega}(\tilde{x}^k) + \frac{C_f^T \bar{\lambda}^k}{m} + \frac{1}{\alpha^k} (\tilde{x}^k - x^k) - \frac{C_f^T (\bar{\lambda}^k - \lambda^k)}{m}. \quad (41)$$

Following a similar line of derivation, using relations

$$\mathbf{1} \otimes \bar{d}^k = \frac{2C_f \tilde{x}^k - C_f x^k - c_f}{m}$$

and

$$\lambda^k - \bar{\lambda}_1^k = \left(\left(I - \frac{\mathbf{1}\mathbf{1}^T}{m} \right) \times I_n \right) \lambda^k = \Pi_f \lambda^k$$

with Π_f defined in (5), the second equality in (40) implies

$$(\text{Id} + N_{\mathbb{R}_+^{mn}})(\bar{\lambda}^k) \ni \lambda^k + \beta^k \left(\frac{2C_f \tilde{x}^k - C_f x^k - c_f}{m} - \Pi_f \lambda^k \right)$$

and further

$$-\Pi_f \lambda^k - \frac{c_f}{m} \ni N_{\mathbb{R}_+^{mn}} - \frac{C_f \tilde{x}^k}{m} - \frac{C_f (\tilde{x}^k - x^k)}{m} + \frac{\bar{\lambda}^k - \lambda^k}{\beta^k}. \quad (42)$$

Defining $\omega^k \triangleq \text{col}(x^k, \lambda^k)$, (41) and (42) can be cast in a compact form as

$$-T_1(\omega^k) \in T_2(\tilde{\omega}^k) + \Phi^k(\tilde{\omega}^k - \omega^k). \quad (43)$$

with T_1 , T_2 , and Φ^k defined in (7) and (38), respectively.

Making $\tilde{\omega}^k$ explicit in (43) leads to

$$\tilde{\omega}^k = R_k(\omega^k),$$

with the operator R_k given in (37).

Finally, combining the preceding equality with the last two equalities in (40) yields (36).

Next, we prove that R_k s are nonexpansive under the conditions in (39). For the convenience of derivation, we assume that $\alpha^k \geq \beta^k$. But the same result can be obtained if the opposite is true.

According to the Gershgorin circle theorem, the eigenvalues of Φ^k are no less than $\min_{i \in [m]} \{ \min \{ \frac{1}{\alpha^k} - \frac{\|C_i\|}{m}, \frac{1}{\beta^k} - \frac{\|C_i\|}{m} \} \}$. Under the assumption $\alpha^k \geq \beta^k$, we have that the eigenvalues of Φ^k are no less than $\min_{i \in [m]} \{ \frac{1}{\alpha^k} - \frac{\|C_i\|}{m} \}$, and further no less than $\frac{1}{2\alpha^k}$ according to (39). Hence, we have $\|\Phi^k\| \geq \frac{1}{2\alpha^k}$ and $\|(\Phi^k)^{-1}\| \leq 2\alpha^k$. Then, following a derivation similar to that of Lemma 10 in [3], we can obtain that $(\Phi^k)^{-1} T_1$ is $\frac{\delta}{2\alpha^k}$ -cocoercive (with $0 < \delta \leq \min\{1, \mu\}$ where μ is from Assumption 2) and $\text{Id} - (\Phi^k)^{-1} T_1$ is $\frac{\alpha^k}{\delta}$ -averaged. Since T_2 is maximally monotone by Lemma 3 in [3], we have that $(\Phi^k)^{-1} T_2$ is maximally monotone, and hence, the mapping $(\text{Id} - (\Phi^k)^{-1} T_2)^{-1}$ is $\frac{1}{2}$ -averaged. Therefore, using the properties that for an a_1 -averaged operator T_a and an a_2 -averaged operator T_b , the composition $T_a \circ T_b$ is $\frac{a_1 + a_2 - 2a_1 a_2}{1 - a_1 a_2}$ -averaged (Proposition 4.44 of [40]), we can obtain that R_k defined in (37) is $\frac{1}{2 - \frac{2\alpha^k}{\delta}}$ -averaged, and hence, is nonexpansive (Remark 3.43 of [40]).

To prove that for every $\{\omega^k\}$, $\lim_{k \rightarrow \infty} \|R_k(\omega^k) - \omega^k\|$ exists almost surely, we write $(\Phi^k)^{-1}$ explicitly as follows:

$$(\Phi^k)^{-1} = I_{\alpha^k, \beta^k} \left(I - U (I_s + I_{\alpha^k, \beta^k} U)^{-1} I_{\alpha^k, \beta^k} \right) \quad (44)$$

where $I_{\alpha^k, \beta^k} \triangleq \begin{bmatrix} (\alpha^k)I_{md} & 0 \\ 0 & (\beta^k)I_{mn} \end{bmatrix}$, $U \triangleq \begin{bmatrix} -\frac{C_f^T}{m} & 0 \\ 0 & -\frac{C_f}{m} \end{bmatrix}$, and $I_s \triangleq \begin{bmatrix} 0 & I_{md} \\ I_{mn} & 0 \end{bmatrix}$.

Given $\sum_{k=0}^{\infty} (\alpha^k)^2 < \infty$, $\sum_{k=0}^{\infty} (\beta^k)^2 < \infty$, it can be easily seen that both α^k and β^k have to converge to 0, and hence $(\Phi^k)^{-1}$ will converge to the zero matrix, which, in turn, implies that R^k will converge to the identity operator. Therefore, we always have that $\lim_{k \rightarrow \infty} \|R_k(\omega^k) - \omega^k\|$ exists almost surely for every ω^k . The fact that Condition 4) in Lemma 6 can be guaranteed by $\sum_{k=0}^{\infty} \alpha^k = \infty$ and $\sum_{k=0}^{\infty} \beta^k = \infty$ follows [5], [9]. ■

Remark 4. Different from the existing result in [53] which requires the stepsize to be bounded away from zero, Lemma 7 allows the stepsize to decay to zero, which is key to enable ϵ -DP with guaranteed finite cumulative privacy budget, as detailed later in Sec. VI.

B. Algorithm 2 is a stochastically-perturbed version of (36)

In this subsection, we prove that Algorithm 2 is a stochastically perturbed version of (36) with the perturbation $e(k)$ satisfying the conditions in Lemma 6. To this end, we first prove that the averages of σ_i^k , y_i^k , and z_i^k in Algorithm 2 always converge to \bar{x}^k , \bar{d}^k , and $\bar{\lambda}^k$, respectively:

Lemma 8. Under Assumptions 4, 7, $\bar{\sigma}^k = \frac{\sum_{i=1}^m \sigma_i^k}{m}$, $\bar{y}^k = \frac{\sum_{i=1}^m y_i^k}{m}$, and $\bar{z}^k = \frac{\sum_{i=1}^m z_i^k}{m}$ in Algorithm 2 converge a.s. to \bar{x}^k , $\bar{d}^k = \frac{\sum_{i=1}^m d_i^k}{m}$ with d_i^k defined in (30), and $\bar{\lambda}^k$, respectively, if $\sum_{k=0}^{\infty} \gamma^k = \infty$ and $\sum_{k=0}^{\infty} (\gamma^k)^2 < \infty$ hold. Moreover, $\sum_{k=0}^{\infty} \alpha^k \|\bar{\sigma}^k - \bar{x}^k\| < \infty$, $\sum_{k=0}^{\infty} \alpha^k \|\bar{z}^k - \bar{\lambda}^k\| < \infty$, and $\sum_{k=0}^{\infty} \alpha^k \|\bar{y}^k - \bar{d}^k\| < \infty$ hold a.s. if $\sum_{k=0}^{\infty} \frac{(\alpha^k)^2}{\gamma^k} < \infty$ is true.

Proof. The derivation of the first part follows the line of reasoning in Lemma 3, and the derivation of the second part follows the last paragraph of the proof of Theorem 1. Hence, we do not include the proof here. ■

To show that Algorithm 2 corresponds to a stochastically-perturbed version of (36), we rewrite the updates in Algorithm 2 in the following more compact form:

$$\begin{aligned} \tilde{x}^k &= \Pi_{\Omega} [x^k - \alpha^k (F(x^k, \sigma^k) + C_d^T z^k)], \\ \tilde{\lambda}^k &= \Pi_{\mathbb{R}^{mn}} [\lambda^k - \beta^k (y^k - \lambda^k + z^k)], \\ x^{k+1} &= x^k + \gamma^k (\tilde{x}^k - x^k), \\ \lambda^{k+1} &= \lambda^k + \gamma^k (\tilde{\lambda}^k - \lambda^k), \end{aligned} \quad (45)$$

with

$$\begin{aligned} y^{k+1} &= (1 - \gamma^k)y^k + \chi^k (L \otimes I_n)y^k + \chi^k (L^0 \otimes I_n)\xi^k \\ &\quad + C_d(2\tilde{x}^k - x^k + c_i) \\ &\quad - (1 - \gamma^k)C_d(2\tilde{x}^{k-1} - x^{k-1} - c_i). \end{aligned} \quad (46)$$

Recalling that (40) corresponds to the nonstationary iteration process in (36), comparing (45) and (40) yields that Algorithm 2 corresponds to the following stochastically-perturbed nonstationary fixed-point iteration process

$$\omega^{k+1} = \omega^k + \gamma^k (R_k(\omega^k) + e^k - \omega^k), \quad (47)$$

with $e^k \triangleq \text{col}(e_x^k, e_{\lambda}^k)$ given by

$$\begin{aligned} e_x^k &= \Pi_{\Omega} [x^k - \alpha^k (F(x^k, \sigma^k) + C_d^T z^k) \\ &\quad - \Pi_{\Omega} [x^k - \alpha^k (F(x^k, \bar{x}_1^k) + C_d^T \bar{\lambda}_1^k)]], \\ e_{\lambda}^k &= \Pi_{\mathbb{R}^{mn}} [\lambda^k - \beta^k (y^k - \lambda^k + z^k) \\ &\quad - \Pi_{\mathbb{R}^{mn}} [\lambda^k - \beta^k (\bar{d}_1^k - \lambda^k + \bar{\lambda}_1^k)]]. \end{aligned} \quad (48)$$

C. Combine preceding two subsections to prove the final result

In this subsection, we combine the preceding two subsections to prove that Algorithm 2 can ensure the almost sure convergence of all agents to the GNE. Following [3], we have the following results:

Proposition 1. In Algorithm 2, x_i^k , \tilde{x}_i^k , $\tilde{\lambda}_i^k$, λ_i^k can always be made bounded.

Proof. One can obtain that \tilde{x}_i^k and x_i^k are always within Ω_i since \tilde{x}_i^k is always obtained by projection onto Ω_i , and the update $x_i^k = x_i^{k-1} + \gamma^k (\tilde{x}_i^{k-1} - x_i^{k-1})$ amounts to a convex combination of elements in the convex set Ω_i . Since Ω_i is compact, we have that \tilde{x}_i^k and x_i^k are bounded. The boundedness of $\tilde{\lambda}_i^k$ can be easily enforced in a distributed manner by a superset based technique [3], which is independent of the DP-design. Under the update rule $\lambda_i^k = \lambda_i^{k-1} + \gamma^k (\tilde{\lambda}_i^{k-1} - \lambda_i^{k-1})$, a bounded $\tilde{\lambda}_i^k$ implies a bounded λ_i^k . ■

Theorem 3. Under Assumptions 1,2,3,4,7, when the nonnegative sequences $\{\alpha^k\}$ and $\{\beta^k\}$ satisfy

$$\sum_{k=0}^{\infty} \alpha^k = \infty, \sum_{k=0}^{\infty} \beta^k = \infty, \sum_{k=0}^{\infty} (\alpha^k)^2 < \infty, \sum_{k=0}^{\infty} (\beta^k)^2 < \infty, \quad (49)$$

and the nonnegative sequences $\{\gamma^k\}$ and $\{\chi^k\}$ satisfy

$$\sum_{k=0}^{\infty} \chi^k = \infty, \sum_{k=0}^{\infty} \frac{(\gamma^k)^2}{\chi^k} < \infty, \sum_{k=0}^{\infty} \frac{(\alpha^k)^2}{\gamma^k} < \infty, \sum_{k=0}^{\infty} \frac{(\beta^k)^2}{\gamma^k} < \infty, \quad (50)$$

then the iterates x_i^k for $i \in [m]$ in Algorithm 2 converge almost surely to a GNE x^* of (2).

Proof. The basic idea is to leverage Lemma 6. To this end, we characterize $\|e^k\|$, which can be verified to satisfy

$$\|e^k\| \leq \|e_x^k\| + \|e_{\lambda}^k\|. \quad (51)$$

Using the nonexpansive property of the projection operator, we have the following relationship for $\|e_x^k\|$

$$\begin{aligned} \|e_x^k\| &\leq \|x^k - \alpha^k (F(x^k, \sigma^k) + C_d^T z^k) \\ &\quad - (x^k - \alpha^k (F(x^k, \bar{x}_1^k) + C_d^T \bar{\lambda}_1^k))\| \\ &\leq \alpha^k \|F(x^k, \sigma^k) - F(x^k, \bar{x}_1^k)\| + \alpha^k \|C_d^T\| \|z^k - \bar{\lambda}_1^k\| \\ &= \alpha^k \|F(x^k, \sigma^k) - F(x^k, \bar{\sigma}_1^k) + F(x^k, \bar{\sigma}_1^k) - F(x^k, \bar{x}_1^k)\| \\ &\quad + \alpha^k \|C_d^T\| \|z^k - \bar{z}_1^k + \bar{z}_1^k - \bar{\lambda}_1^k\| \\ &\leq \alpha^k \tilde{L} \|\sigma^k - \bar{\sigma}_1^k\| + \alpha^k \tilde{L} \|\bar{\sigma}_1^k - \bar{x}_1^k\| \\ &\quad + \alpha^k \|C_d^T\| \|z^k - \bar{z}_1^k\| + \alpha^k \|C_d^T\| \|\bar{z}_1^k - \bar{\lambda}_1^k\| \\ &= \alpha^k \tilde{L} \|\sigma^k - \bar{\sigma}_1^k\| + \alpha^k \sqrt{m} \tilde{L} \|\bar{\sigma}^k - \bar{x}^k\| \\ &\quad + \alpha^k \|C_d^T\| \|z^k - \bar{z}_1^k\| + \alpha^k \sqrt{m} \|C_d^T\| \|\bar{z}^k - \bar{\lambda}^k\|, \end{aligned} \quad (52)$$

where we have used the Lipschitz condition in Assumption 3 in the last inequality.

Lemma 8 implies $\sum_{k=0}^{\infty} \alpha^k \|\bar{\sigma}^k - \bar{x}^k\| < \infty$ and $\sum_{k=0}^{\infty} \alpha^k \|\bar{z}^k - \bar{\lambda}^k\| < \infty$ a.s. Hence, we only need to consider the first and third terms on the right hand side of (52).

According to (31), the evolutions of σ_i^k and z_i^k follow

$$\sigma_i^{k+1} = (1 - \gamma^k) \sigma_i^k + \chi^k \sum_{j \in \mathbb{N}_i} L_{ij} (\sigma_j^k + \zeta_j^k - \sigma_i^k) + \gamma^k (\tilde{x}_i^k),$$

and

$$z_i^{k+1} = (1 - \gamma^k) z_i^k + \chi^k \sum_{j \in \mathbb{N}_i} L_{ij} (z_j^k + v_j^k - z_i^k) + \gamma^k (\tilde{\lambda}_i^k),$$

respectively. Therefore, given that $x_i^k, \tilde{x}_i^k, \tilde{\lambda}_i^k, \lambda_i^k$ are bounded from Proposition 1, Theorem 1 implies that $\sigma_i^k \rightarrow \bar{x}^k$ and $z_i^k \rightarrow \bar{\lambda}^k$ almost surely, and the following relations hold a.s.:

$$\sum_{k=0}^{\infty} \gamma^k \sum_{i=1}^m \|\sigma_i^k - \bar{\sigma}^k\| < \infty, \quad \sum_{k=0}^{\infty} \gamma^k \sum_{i=1}^m \|z_i^k - \bar{z}^k\| < \infty. \quad (53)$$

Combining (52) and (53) implies the following relation a.s.:

$$\sum_{k=0}^{\infty} \gamma^k \|e_x^k\| < \infty. \quad (54)$$

Following the same line of derivation, we have

$$\begin{aligned} \|e_\lambda^k\| &\leq \|\lambda^k - \beta^k (y^k - \lambda^k + z^k) \\ &\quad - (\lambda^k - \beta^k (\bar{d}_1^k - \lambda^k + \bar{\lambda}_1^k))\| \\ &\leq \beta^k \|y^k - \bar{y}^k\| + \beta^k \|\sqrt{m}\| \bar{y}^k - \bar{d}^k\| + \beta^k \|z^k - \bar{z}_1^k\| \\ &\quad + \beta^k \|\sqrt{m}\| \bar{z}^k - \bar{\lambda}^k\| \end{aligned} \quad (55)$$

Lemma 8 implies $\sum_{k=0}^{\infty} \beta^k \|\bar{z}^k - \bar{\lambda}^k\| < \infty$ and $\sum_{k=0}^{\infty} \beta^k \|\bar{y}^k - \bar{d}^k\| < \infty$ a.s. under $\sum_{k=0}^{\infty} \frac{(\beta^k)^2}{\gamma^k} < \infty$. Hence, we only need to consider the first and third terms on the right hand side of (55).

According to Theorem 1, for the dynamics y_i^k in (46), if we can prove that $\|C_d(2\tilde{x}^k - x^k - c_i) - (1 - \gamma^k)C_d(2\tilde{x}^{k-1} - x^{k-1} - c_i)\| \leq \gamma^k C$ holds for some C , then we can have $\sum_{k=0}^{\infty} \gamma^k \sum_{i=1}^m \|y_i^k - \bar{y}^k\| < \infty$ and hence, $\sum_{k=0}^{\infty} \beta^k \sum_{i=1}^m \|y_i^k - \bar{y}^k\| < \infty$.

Noticing $C_d(2\tilde{x}^k - x^k - c_i) - (1 - \gamma^k)C_d(2\tilde{x}^{k-1} - x^{k-1} - c_i) = 2C_d(\tilde{x}^k - \tilde{x}^{k-1}) - C_d(x^k - x^{k-1}) + \gamma^k C_d(2\tilde{x}^{k-1} - x^{k-1} - c_i)$, we have

$$\begin{aligned} &\|C_d(2\tilde{x}^k - x^k - c_i) - (1 - \gamma^k)C_d(2\tilde{x}^{k-1} - x^{k-1} - c_i)\| \\ &\leq \|2C_d\| \|\tilde{x}^k - \tilde{x}^{k-1}\| + \|C_d\| \|x^k - x^{k-1}\| \\ &\quad + \gamma^k \|C_d(2\tilde{x}^{k-1} - x^{k-1} - c_i)\| \\ &\leq \|2C_d\| \|\tilde{x}^k - \tilde{x}^{k-1}\| + \gamma^k \|C_d\| \|\tilde{x}^{k-1} - x^{k-1}\| \\ &\quad + \gamma^k \|C_d(2\tilde{x}^{k-1} - x^{k-1} - c_i)\|, \end{aligned} \quad (56)$$

where we have used the update rule of x_i^k in (31) in the last inequality.

To characterize $\|\tilde{x}^k - \tilde{x}^{k-1}\|$, we employ the update rule of \tilde{x}_i^k in (31). Using the nonexpansive property of projection operators, we have

$$\begin{aligned} \|\tilde{x}^k - \tilde{x}^{k-1}\| &= \|\Pi_\Omega [x^k - \alpha^k (F(x^k, \sigma^k) + C_d^T z^k)] \\ &\quad - \Pi_\Omega [x^{k-1} - \alpha^{k-1} (F(x^{k-1}, \sigma^{k-1}) + C_d^T z^{k-1})]\| \\ &\leq \|x^k - \alpha^k (F(x^k, \sigma^k) + C_d^T z^k) \\ &\quad - (x^{k-1} - \alpha^{k-1} (F(x^{k-1}, \sigma^{k-1}) + C_d^T z^{k-1}))\| \\ &\leq \|x^k - x^{k-1}\| + \alpha^k \|F(x^k, \sigma^k) + C_d^T z^k\| \\ &\quad + \alpha^{k-1} \|F(x^{k-1}, \sigma^{k-1}) + C_d^T z^{k-1}\| \end{aligned} \quad (57)$$

Using the proven results that $\sigma_i^k \rightarrow \bar{x}^k$ and $z_i^k \rightarrow \bar{\lambda}^k$ almost surely, and \bar{x}^k and $\bar{\lambda}^k$ are bounded (see Proposition 1), we have that z_i^k and σ_i^k are bounded almost surely. Therefore, the term $F(x^k, \sigma^k) + C_d^T z^k$ is bounded almost surely. Without loss of generality, we assume

$$\|F(x^k, \sigma^k) + C_d^T z^k\| \leq \bar{C}$$

for all k almost surely.

Plugging the preceding relationship into (57) leads to

$$\begin{aligned} \|\tilde{x}^k - \tilde{x}^{k-1}\| &\leq \|x^k - x^{k-1}\| + (\alpha^k + \alpha^{k-1})\bar{C} \\ &\leq \gamma^k \|\tilde{x}^k - x^k\| + (\alpha^k + \alpha^{k-1})\bar{C}, \end{aligned} \quad (58)$$

where we have used the update rule of x_i^k in (31) in the last inequality. Given $\sum_{k=0}^{\infty} \frac{(\alpha^k)^2}{\gamma^k} < \infty$ in the theory statement, we have that $(\alpha^k + \alpha^{k-1}) \leq \tilde{C}\gamma^k$ holds for some constant \tilde{C} , and hence $\|\tilde{x}^k - \tilde{x}^{k-1}\| \leq \gamma^k (\|\tilde{x}^k - x^k\| + \tilde{C})$ and further

$$\begin{aligned} &\|C_d(2\tilde{x}^k - x^k - c_i) - (1 - \gamma^k)C_d(2\tilde{x}^{k-1} - x^{k-1} - c_i)\| \\ &\leq \gamma^k \|C_d\| (\|\tilde{x}^{k-1} - x^{k-1}\| + 2\|\tilde{x}^k - x^k\| + 2\tilde{C}) \\ &\quad + \gamma^k \|C_d(2\tilde{x}^{k-1} - x^{k-1} - c_i)\| \end{aligned} \quad (59)$$

by using (56).

Combining (46) and (59), we have that the evolution of y_i^k satisfies the conditions of Theorem 1, and hence, $\sum_{k=0}^{\infty} \gamma^k \sum_{i=1}^m \|y_i^k - \bar{y}^k\| < \infty$ holds almost surely. Further invoking (53) and (55) yields the following result a.s.:

$$\sum_{k=0}^{\infty} \gamma^k \|e_\lambda^k\| < \infty. \quad (60)$$

In summary, (51), (54), and (60) mean that $\sum_{k=0}^{\infty} \gamma^k \|e^k\| < \infty$ holds deterministically under the theorem statement almost surely, and hence, $\sum_{k=0}^{\infty} \gamma^k \mathbb{E}[\|e^k\|] < \infty$ holds almost surely. Given $\sqrt{\mathbb{E}[\|e^k\|^2]} \leq \mathbb{E}[\|e^k\|]$ due to Jensen's inequality, invoking Lemma 6 yields that Algorithm 2 guarantees the convergence of all players to the GNE almost surely. ■

VI. PRIVACY ANALYSIS OF ALGORITHM 2

Similar to Definition 3, we define the sensitivity of a distributed GNE seeking algorithm to problem (2) as follows:

Definition 4. At each iteration k , for any initial state ϑ^0 and any adjacent distributed GNE problems \mathcal{P} and \mathcal{P}' , the sensitivity of a GNE seeking algorithm is

$$\Delta^k \triangleq \sup_{\mathcal{O} \in \mathcal{O}} \left\{ \sup_{\vartheta \in \mathcal{R}_{\mathcal{P}, \vartheta^0}^{-1}(\mathcal{O}), \vartheta' \in \mathcal{R}_{\mathcal{P}', \vartheta^0}^{-1}(\mathcal{O})} \|\vartheta^{k+1} - \vartheta'^{k+1}\|_1 \right\}. \quad (61)$$

Then, similar to Lemma 5, we have the following lemma:

Lemma 9. In Algorithm 2, at each iteration k , if each player adds noise vectors ζ_i^k , ξ_i^k , and ϑ_i^k to its shared messages σ_i^k , z_i^k , and y_i^k , respectively, with every noise vector consisting of independent Laplace scalar noises with parameter ν^k , such that $\sum_{k=1}^{T_0} \frac{\Delta^k}{\nu^k} \leq \bar{\epsilon}$, then the iterative distributed Algorithm 2 is ϵ -differentially private with the cumulative privacy budget for iterations from $k = 0$ to $k = T_0$ less than $\bar{\epsilon}$.

Proof. The lemma can be obtained following the same line of reasoning of Lemma 2 in [49] (also see Theorem 3 in [30]).

Theorem 4. Under the conditions of Theorem 3, and all elements of ζ_i^k , ξ_i^k , and ϑ_i^k are drawn independently from Laplace distribution $\text{Lap}(\nu^k)$ with $(\sigma_i^k)^2 = 2(\nu^k)^2$ satisfying Assumption 7, then all players in Algorithm 2 will converge almost surely to the GNE. Moreover,

- 1) Algorithm 2 is ϵ -differentially private with the cumulative privacy budget bounded by $\epsilon \leq \sum_{k=1}^{T_0} \frac{2C\gamma^k}{\nu^k}$ for iterations from $k = 0$ to $k = T_0$ where C is determined by the maximal value of the primal solution \tilde{x}_i^k and dual solution $\tilde{\lambda}_i^k$, which are always bounded as proven in the convergence proof. And the cumulative privacy budget is always finite for $T_0 \rightarrow \infty$ when the sequence $\{\frac{\gamma^k}{\nu^k}\}$ is summable;
- 2) Suppose that two non-negative sequences $\{\nu^k\}$ and $\{\gamma^k\}$ have a finite sequence-ratio sum $\Phi_{\gamma, \nu} \triangleq \sum_{k=1}^{\infty} \frac{\gamma^k}{\nu^k}$. Then setting the Laplace noise parameter ν^k as $\nu^k = \frac{2C\Phi_{\gamma, \nu}}{\epsilon} \nu^k$ ensures that Algorithm 2 is ϵ -differentially private with any cumulative privacy budget $\epsilon > 0$ even when the number of iterations goes to infinity;
- 3) In the special case where $\gamma^k = \frac{1}{k^{0.9}}$, setting $\nu^k = \frac{2C\Phi_{\gamma, \nu}}{\epsilon} k^{0.3}$ with $\Phi \triangleq \sum_{k=1}^{\infty} \frac{1}{k^{1.3}} \approx 3.93$ (which can be verified to satisfy Assumption 7) ensures that Algorithm 2 is always ϵ -differentially private for any cumulative privacy budget $\epsilon > 0$ even when the number of iterations goes to infinity.

Proof. Since the Laplace noises satisfy Assumption 7, the convergence follows directly from Theorem 3.

To prove the results on ϵ -DP, we first prove that the sensitivity of Algorithm 2 satisfies $\Delta^k \leq 2C\gamma^k$ for some constant C . Since the sensitivity is independent of DP-noises, we consider (61) in the absence of DP-noises. It can be seen that for Algorithm 2, we have $\vartheta_i = \text{col}(x_i, y_i, \lambda_i, \sigma_i, z_i)$ and $\vartheta = \text{col}(\vartheta_1, \dots, \vartheta_m)$ (note that \tilde{x}_i and $\tilde{\lambda}_i$ are intermediate variables that are fully represented by the iterations of x_i and λ_i). The observation includes σ_i^k , y_i^k , and z_i^k . Note that Theorem 3 ensures the convergence of all players to the GNE

even in the presence of DP-noises. Hence, in the absence of DP-noises, we have that every x_i^k converges to the GNE, and \tilde{x}_i^k and $\tilde{\lambda}_i^k$ are bounded in norm. Without loss of generality, we assume that \tilde{x}_i^k and $\tilde{\lambda}_i^k$ are bounded by a constant C .

Given two adjacent distributed GNE problems \mathcal{P} and \mathcal{P}' , for any given fixed observation \mathcal{O} and initial state ϑ^0 , the sensitivity is determined by $\|\mathcal{R}_{\mathcal{P}, \vartheta^0}^{-1}(\mathcal{O}) - \mathcal{R}_{\mathcal{P}', \vartheta^0}^{-1}(\mathcal{O})\|_1$ according to Definition 4. Since in \mathcal{P} and \mathcal{P}' , there is only one cost function that is different, we represent this different cost function as the i th one, i.e., J_i in \mathcal{P} and J'_i in \mathcal{P}' , without loss of generality. Because the observations under \mathcal{P} and \mathcal{P}' are identical, we have

$$\sigma_j^k = \sigma_j'^k, y_j^k = y_j'^k, z_j^k = z_j'^k \quad \forall k \geq 0, \quad \forall j \in [m]. \quad (62)$$

At a given iteration k , (62) implies the following relationship

$$x_j^\ell = x_j'^\ell, \lambda_j^\ell = \lambda_j'^\ell \quad \forall 0 \leq \ell \leq k, \quad \forall j \in [m] \quad (63)$$

based on the last two relations in (31) and the fact that the initial conditions for \mathcal{P} and \mathcal{P}' are the same.

Therefore, we have the following relationship for the sensitivity of Algorithm 2:

$$\begin{aligned} & \|\mathcal{R}_{\mathcal{P}, \vartheta^0}^{-1}(\mathcal{O}) - \mathcal{R}_{\mathcal{P}', \vartheta^0}^{-1}(\mathcal{O})\|_1 \\ &= \left\| \begin{bmatrix} x^{k+1} \\ \lambda^{k+1} \\ y^{k+1} \\ \sigma^{k+1} \\ z^{k+1} \end{bmatrix} - \begin{bmatrix} x'^{k+1} \\ \lambda'^{k+1} \\ y'^{k+1} \\ \sigma'^{k+1} \\ z'^{k+1} \end{bmatrix} \right\|_1 = \left\| \begin{bmatrix} x^{k+1} - x'^{k+1} \\ \lambda^{k+1} - \lambda'^{k+1} \\ 0 \\ 0 \\ 0 \end{bmatrix} \right\|_1 \\ &= \left\| \begin{bmatrix} x^{k+1} - x'^{k+1} \\ \lambda^{k+1} - \lambda'^{k+1} \end{bmatrix} \right\|_1, \end{aligned} \quad (64)$$

where we used the fact that the observations in (62) are the same in the second last equality.

From the update rule of x_i^k and σ_i^k in (31), and the fact that $J_j = J'_j$ for all $j \neq i$, we have $x_j^{k+1} = x_j'^{k+1}$ and $\lambda_j^{k+1} = \lambda_j'^{k+1}$ for all $j \neq i$, and hence,

$$\begin{aligned} & \|\mathcal{R}_{\mathcal{P}, \vartheta^0}^{-1}(\mathcal{O}) - \mathcal{R}_{\mathcal{P}', \vartheta^0}^{-1}(\mathcal{O})\|_1 \\ &= \left\| \begin{bmatrix} x_i^{k+1} - x_i'^{k+1} \\ \lambda_i^{k+1} - \lambda_i'^{k+1} \end{bmatrix} \right\|_1 \\ &= \left\| \begin{bmatrix} x_i^k + \gamma^k(\tilde{x}_i^k - x_i^k) - \left(x_i'^k + \gamma^k(\tilde{x}_i'^k - x_i'^k) \right) \\ \lambda_i^k + \gamma^k(\tilde{\lambda}_i^k - \lambda_i^k) - \left(\lambda_i'^k + \gamma^k(\tilde{\lambda}_i'^k - \lambda_i'^k) \right) \end{bmatrix} \right\|_1 \\ &= \left\| \begin{bmatrix} \gamma^k(\tilde{x}_i^k - \tilde{x}_i'^k) \\ \gamma^k(\tilde{\lambda}_i^k - \tilde{\lambda}_i'^k) \end{bmatrix} \right\|_1 \end{aligned} \quad (65)$$

where we used the relationship in (63) in the last equality.

Since \tilde{x}_i^k , $\tilde{\lambda}_i^k$, $\tilde{x}_i'^k$, and $\tilde{\lambda}_i'^k$ are bounded by some constant C , as discussed in the beginning of the derivation (note Algorithm 2 guarantees convergence under both \mathcal{P} and \mathcal{P}'), we have the sensitivity of Algorithm 2 being no greater than $2\gamma^k C$.

Using Lemma 9, we can obtain that the cumulative privacy budget is always less than $\sum_{k=1}^{T_0} \frac{2C\gamma^k}{\nu^k}$. Hence, the cumulative privacy budget ϵ will always be finite even when the number

of iterations T_0 tends to infinity if the sequence $\{\frac{\gamma^k}{\nu^k}\}$ is summable, i.e., $\sum_{k=0}^{\infty} \frac{\gamma^k}{\nu^k} < \infty$.

From Lemma 9, it can also be seen that the cumulative privacy budget is inversely proportional to ν^k . Hence, we can obtain the second statement from the first statement by scaling ν^k proportionally. The third statement can be obtained by specializing the selection of γ^k and ν^k sequences. ■

Note that in the DP design for coupling-constraint free Nash equilibrium seeking, to avoid the cumulative privacy budget from growing to infinity, [30] and [49] employ a summable stepsize (geometrically-decreasing stepsize, more specifically), which, however, also makes it impossible to ensure convergence to the exact desired equilibrium. In contrast, by allowing the stepsize sequence to be non-summable, our Algorithm 2 achieves both accurate convergence and ϵ -DP with a finite cumulative privacy budget, even when the number of iterations goes to infinity. To our knowledge, this is the first time that both almost-sure convergence and rigorous ϵ -DP are achieved in distributed GNE seeking, even with the number of iterations going to infinity.

Remark 5. *To ensure that the cumulative privacy budget $\epsilon = \sum_{k=1}^{\infty} \frac{2C\gamma^k}{\nu^k}$ is bounded when $k \rightarrow \infty$, our algorithm uses Laplace noise with parameter ν^k that increases with time (since we require $\{\frac{\gamma^k}{\nu^k}\}$ to be summable while $\{\gamma^k\}$ is non-summable). Such an increasing ν^k makes the relative level between DP-noises and shared signals (σ_i^k , y_i^k , and z_i^k) increase with time. However, given that it is $\chi^k \text{Lap}(\nu^k)$ that is actually fed into the algorithm, and the increasing speed of the noise level ν^k is outweighed by the decreasing speed of χ^k (see Assumption 7), the actual noise fed into the algorithm still decays with time, which makes it possible for Algorithm 2 to ensure provable convergence to the GNE. In addition, according to Theorem 3, the convergence is not affected by scaling ν^k by any constant coefficient $\frac{1}{\epsilon} > 0$ to achieve any desired level of ϵ -DP, as long as the DP-noise parameter ν^k satisfies Assumption 7.*

VII. NUMERICAL SIMULATIONS

We evaluate the performance of the proposed GNE seeking algorithm using a Nash-Cournot game recently considered in [1], [13], [55]. In the game, we consider m firms producing a homogeneous commodity competing over N markets, with a schematic presented in Fig. 1. In the schematic, we plot $N = 7$ markets (represented by M_1, \dots, M_7) and $m = 20$ firms (represented by circles). We use an edge from circle i to M_j to denote that firm i participates in market M_j .

We use $x_i \in \mathbb{R}^N$ to represent the amount of firm i 's products. If firm i does not participate in market j , then the j th entry of x_i will be forced to be 0 all the time. Hence, if firm i participates in $1 \leq n_i \leq N$ markets, then its production vector x_i will have n_i non-zero entries. For notational simplicity, we use an adjacency matrix $B_i \in \mathbb{R}^{N \times N}$ to describe the association relationship between firm i and all markets. More specifically, B_i 's off-diagonal elements are all zero, and its j th diagonal entry is one if firm i participates in market j , otherwise, its j th diagonal entry is zero. Every firm i has a

maximal capacity for each market j it participates in, which is represented by C_{ij} . Defining $\bar{C}_i \triangleq [C_{i1}, \dots, C_{iN}]^T$, the capacity constraint can be formulated as $x_i \leq \bar{C}_i$. Defining $B \triangleq [B_1, \dots, B_N]$, $Bx \in \mathbb{R}^N = \sum_{i=1}^N B_i x_i$ represents the total product supply to all markets, given firm i 's production amount x_i . We assume that every market has a maximum capacity \bar{c}_i . Defining $\bar{c} \triangleq [\bar{c}_1, \dots, \bar{c}_N]^T \in \mathbb{R}^N$, the shared coupling constraints can be formulated as $Bx \leq \bar{c}$.

As in [1], we let the commodity's price in every market M_i follow a linear inverse demand function:

$$p_i(x) = \bar{P}_i - \varsigma_i [Bx]_i,$$

where \bar{P}_i and $\varsigma_i > 0$ are constants and $[Bx]_i$ denotes the i th element of the vector Bx . It can be verified that the price value decreases when the amount of supplied commodity increases.

Representing the price vector of all markets as $p \triangleq [p_1, \dots, p_N]^T$, we have

$$p = \bar{P} - \Xi Bx,$$

where $\bar{P} \triangleq [\bar{P}_1, \dots, \bar{P}_N]^T$ and $\Xi \triangleq \text{diag}(\varsigma_1, \dots, \varsigma_N)$. The total payoff of firm i can then be expressed as $p^T B_i x_i$. Firm i 's production cost is assumed to be a quadratic function

$$\phi_i(x_i) = x_i^T Q_i x_i + q_i^T x_i,$$

where $Q_i \in \mathbb{R}^{N \times N}$ is positive definite and $q_i \in \mathbb{R}^N$.

Firm i 's local cost function, which is determined by its production cost ϕ_i and payoff, is given by

$$J_i(x_i, x) = \phi_i(x_i) - (\bar{P} - \Xi Bx)^T B_i^T x_i.$$

One can verify that the gradient of the cost function is

$$F_i(x_i, x) = 2Q_i x_i + q_i + B_i^T \Xi B_i x_i - B_i (\bar{P} - \Xi Bx).$$

It is clear that both firm i 's local cost function and gradient are dependent on other firms' actions.

In our simulation, we consider $m = 20$ firms competing over $N = 7$ markets. Since in the partial-decision information setting, each firm can only communicate with its immediate neighboring firms, we use a randomly generated local communication pattern given in Fig. 2. The maximal capacities for firm i (elements in \bar{C}_i) are randomly selected from the interval $[8, 10]$. The maximal capacities for all markets are set as $\kappa \sum_{i=1}^{20} \bar{C}_i$, with κ randomly selected from a uniform distribution on $(0, 1)$. Q_i in the production cost function is set as νI with ν randomly selected from $[1, 10]$. q_i in $\phi_i(x_i)$ is randomly selected from a uniform distribution in $[1, 2]$. In the price function, \bar{P}_i and ς_i are randomly chosen from uniform distributions in $[10, 20]$ and $[1, 3]$, respectively.

To evaluate the proposed Algorithm 2, for every firm i , we inject DP-noises ζ_i^k , ξ_i^k , and v_i^k in every shared σ_i^k , y_i^k , and z_i^k in all iterations. Each element of the noise vectors follows Laplace distribution with parameter $\nu^k = 1 + 0.1k^{0.2}$. We set the stepsizes and diminishing sequences as $\alpha^k = \beta^k = \frac{0.1}{1+0.1k}$, $\gamma^k = \frac{0.01}{1+0.1k^{0.98}}$ and $\chi^k = \frac{1}{1+0.1k^{0.9}}$, respectively, which satisfy the conditions in Theorem 3. In the evaluation, we run our algorithm for 100 times, and calculate the average and the variance of $\|x^k - x^*\|$ against the iteration index k . The result is given by the red curve and error bars in Fig. 3. For




Fig. 1. Nash-Cournot game of 20 players (firms) competing over 7 locations (markets). Each firm is represented by a circular and each market is represented by a square. An edge between firm i ($1 \leq i \leq 20$) and market j ($1 \leq j \leq 7$) means that firm i participates in market j .




Fig. 2. The randomly generated interaction pattern of the 20 firms.

comparison, we also run the existing GNE seeking algorithm proposed by Belgioioso et al. in [3] under the same noise, and the existing DP approach for Nash equilibrium seeking proposed by Ye et al. in [30] under the same cumulative privacy budget ϵ . Note that the DP approach in [30] addresses Nash equilibrium seeking problems without shared coupling constraints. We adapt its DP mechanism (geometrically decreasing stepsizes and DP-noises) to the GNE seeking problem. The evolution of the average error/variance of the approaches in [3] and [30] are given by the blue and black curves/error bars in Fig. 3, respectively. It can be seen that the proposed algorithm has a much better accuracy.

VIII. CONCLUSIONS

This paper proposes a differentially-private fully distributed algorithm for generalized Nash equilibrium seeking. Different from existing privacy solutions for coupling-constraint free aggregative games, the proposed approach allows the existence of shared coupling constraints, which increase attack surfaces, and hence, pose additional challenges to privacy protection. More interestingly, the proposed approach can ensure both provable convergence to the generalized Nash equilibrium point and rigorous ϵ -differential privacy. To our knowledge, the simultaneous achievement of both ϵ -differential privacy and provable convergence accuracy has not been achieved in generalized Nash equilibrium seeking before. As a basis of the differentially private generalized Nash equilibrium seeking algorithm, we also propose a new consensus tracking algorithm that can ensure both provable convergence accuracy and




Fig. 3. Comparison of Algorithm 2 with the existing GNE seeking algorithm by Belgioioso et al. in [3] (under the same noise) and the differential-privacy approach from Ye et al. in [30] after adaptation to GNE seeking (under the same privacy budget ϵ).

rigorous ϵ -differential privacy, which, to our knowledge, has not been reported before. The convergence analysis generalizes existing results of stochastically-perturbed nonstationary fixed-point iteration processes to the diminishing-stepsize case, which is crucial to ensure a finite privacy budget, and hence, rigorous differential privacy. Numerical simulation results confirm the effectiveness of the proposed algorithm.

REFERENCES

- [1] L. Pavel, "Distributed GNE seeking under partial-decision information over networks via a doubly-augmented operator splitting approach," *IEEE Transactions on Automatic Control*, vol. 65, no. 4, pp. 1584–1597, 2020.
- [2] P. Yi and L. Pavel, "An operator splitting approach for distributed generalized Nash equilibria computation," *Automatica*, vol. 102, pp. 111–121, 2019.
- [3] G. Belgioioso, A. Nedić, and S. Grammatico, "Distributed generalized Nash equilibrium seeking in aggregative games on time-varying networks," *IEEE Transactions on Automatic Control*, vol. 66, no. 5, pp. 2061–2075, 2021.
- [4] W. Saad, Z. Han, H. V. Poor, and T. Başar, "Game-theoretic methods for the smart grid: An overview of microgrid systems, demand-side management, and smart grid communications," *IEEE Signal Processing Magazine*, vol. 29, no. 5, pp. 86–105, 2012.
- [5] M. Zhu and E. Frattoli, "Distributed robust adaptive equilibrium computation for generalized convex games," *Automatica*, vol. 63, pp. 82–91, 2016.
- [6] M. Ye and G. Hu, "Game design and analysis for price-based demand response: An aggregate game approach," *IEEE Transactions on Cybernetics*, vol. 47, no. 3, pp. 720–730, 2016.
- [7] Y. Pan and L. Pavel, "Games with coupled propagated constraints in optical networks with multi-link topologies," *Automatica*, vol. 45, no. 4, pp. 871–880, 2009.
- [8] T. Alpcan and T. Başar, "Distributed algorithms for Nash equilibria of flow control games," in *Advances in dynamic games*. Springer, 2005, pp. 473–498.
- [9] H. Yin, U. V. Shanbhag, and P. G. Mehta, "Nash equilibrium problems with scaled congestion costs and shared constraints," *IEEE Transactions on Automatic Control*, vol. 56, no. 7, pp. 1702–1708, 2011.
- [10] Z. Ma, D. S. Callaway, and I. A. Hiskens, "Decentralized charging control of large populations of plug-in electric vehicles," *IEEE Transactions on Control Systems Technology*, vol. 21, no. 1, pp. 67–78, 2011.
- [11] Z. Ma, S. Zou, L. Ran, X. Shi, and I. A. Hiskens, "Efficient decentralized coordination of large-scale plug-in electric vehicle charging," *Automatica*, vol. 69, pp. 35–47, 2016.

- [12] M. S. Stankovic, K. H. Johansson, and D. M. Stipanovic, "Distributed seeking of Nash equilibria with applications to mobile sensor networks," *IEEE Transactions on Automatic Control*, vol. 57, no. 4, pp. 904–919, 2011.
- [13] J. Koshal, A. Nedić, and U. V. Shanbhag, "Distributed algorithms for aggregative games on graphs," *Operations Research*, vol. 64, no. 3, pp. 680–704, 2016.
- [14] F. Salehisadaghiani and L. Pavel, "Distributed Nash equilibrium seeking: A gossip-based algorithm," *Automatica*, vol. 72, pp. 209–216, 2016.
- [15] T. Tatarenko, W. Shi, and A. Nedić, "Geometric convergence of gradient play algorithms for distributed Nash equilibrium seeking," *IEEE Transactions on Automatic Control*, vol. 66, no. 11, pp. 5342–5353, 2020.
- [16] D. Gadjov and L. Pavel, "A passivity-based approach to Nash equilibrium seeking over networks," *IEEE Transactions on Automatic Control*, vol. 64, no. 3, pp. 1077–1092, 2018.
- [17] M. Ye and G. Hu, "Distributed Nash equilibrium seeking by a consensus based approach," *IEEE Transactions on Automatic Control*, vol. 62, no. 9, pp. 4811–4818, 2017.
- [18] F. Salehisadaghiani, W. Shi, and L. Pavel, "Distributed Nash equilibrium seeking under partial-decision information via the alternating direction method of multipliers," *Automatica*, vol. 103, pp. 27–35, 2019.
- [19] M. Bianchi, G. Belgioioso, and S. Grammatico, "Fast generalized Nash equilibrium seeking under partial-decision information," *Automatica*, vol. 136, p. 110080, 2022.
- [20] M. Zhu and S. Martínez, "Discrete-time dynamic average consensus," *Automatica*, vol. 46, no. 2, pp. 322–329, 2010.
- [21] S. Rassenti, S. S. Reynolds, V. L. Smith, and F. Szidarovszky, "Adaptation and convergence of behavior in repeated experimental cournot games," *Journal of Economic Behavior & Organization*, vol. 41, no. 2, pp. 117–146, 2000.
- [22] R. Dong, W. Krichene, A. M. Bayen, and S. S. Sastry, "Differential privacy of populations in routing games," in *2015 54th IEEE Conference on Decision and Control (CDC)*. IEEE, 2015, pp. 2798–2803.
- [23] T. A. Gerhart and A. Steinberg, "Proposition 24: Protecting California consumers by expanding protections, ensuring governmental oversight, and safeguarding the law from special interests," *California Initiative Review (CIR)*, vol. 2020, no. 1, p. 12, 2020.
- [24] M. Shakarami, C. De Persis, and N. Monshizadeh, "Distributed dynamics for aggregative games: Robustness and privacy guarantees," *International Journal of Robust and Nonlinear Control*, vol. 32, no. 9, pp. 5048–5069, 2022.
- [25] Y. Lu and M. Zhu, "Game-theoretic distributed control with information-theoretic security guarantees," *IFAC-PapersOnLine*, vol. 48, no. 22, pp. 264–269, 2015.
- [26] R. Cummings, M. Kearns, A. Roth, and Z. S. Wu, "Privacy and truthful equilibrium selection for aggregative games," in *International Conference on Web and Internet Economics*. Springer, 2015, pp. 286–299.
- [27] I. Shilov, H. Le Cadre, and A. Busic, "Privacy impact on generalized Nash equilibrium in peer-to-peer electricity market," *Operations Research Letters*, vol. 49, no. 5, pp. 759–766, 2021.
- [28] S. Gade, A. Winnicki, and S. Bose, "On privatizing equilibrium computation in aggregate games over networks," *IFAC-PapersOnLine*, vol. 53, no. 2, pp. 3272–3277, 2020.
- [29] C. Dwork, A. Roth *et al.*, "The algorithmic foundations of differential privacy," *Foundations and Trends in Theoretical Computer Science*, vol. 9, no. 3–4, pp. 211–407, 2014.
- [30] M. Ye, G. Hu, L. Xie, and S. Xu, "Differentially private distributed Nash equilibrium seeking for aggregative games," *IEEE Transactions on Automatic Control*, vol. 67, no. 5, pp. 2451–2458, 2021.
- [31] J. Wang, J.-F. Zhang, and X. He, "Differentially private distributed algorithms for stochastic aggregative games," *Automatica*, vol. 142, p. 110440, 2022.
- [32] Y. Wang and A. Nedić, "Differentially-private distributed algorithms for aggregative games with guaranteed convergence," *arXiv preprint arXiv:2209.01486*, 2022.
- [33] Y. Wang and T. Başar, "Ensuring both accurate convergence and differential privacy in Nash equilibrium seeking on directed graphs," *arXiv preprint arXiv:2209.04938*, 2022.
- [34] —, "Gradient-tracking based distributed optimization with guaranteed optimality under noisy information sharing," *IEEE Transactions on Automatic Control*, 2022.
- [35] F. Parise, B. Gentile, and J. Lygeros, "A distributed algorithm for almost-Nash equilibria of average aggregative games with coupling constraints," *IEEE Transactions on Control of Network Systems*, vol. 7, no. 2, pp. 770–782, 2019.
- [36] D. Gadjov and L. Pavel, "Single-timescale distributed GNE seeking for aggregative games over networks via forward-backward operator splitting," *IEEE Transactions on Automatic Control*, vol. 66, no. 7, pp. 3259–3266, 2020.
- [37] R. Zhu, J. Zhang, K. You, and T. Başar, "Asynchronous networked aggregative games," *Automatica*, vol. 136, p. 110054, 2022.
- [38] S. Liang, P. Yi, and Y. Hong, "Distributed Nash equilibrium seeking for aggregative games with coupled constraints," *Automatica*, vol. 85, pp. 179–185, 2017.
- [39] J. Lei and U. V. Shanbhag, "Distributed variable sample-size gradient-response and best-response schemes for stochastic Nash equilibrium problems," *SIAM Journal on Optimization*, vol. 32, no. 2, pp. 573–603, 2022.
- [40] H. H. Bauschke and P. L. Combettes, *Convex analysis and monotone operator theory in Hilbert spaces*. Springer, 2011, vol. 408.
- [41] N. Li, L. Chen, and M. A. Dahleh, "Demand response using linear supply function bidding," *IEEE Transactions on Smart Grid*, vol. 6, no. 4, pp. 1827–1838, 2015.
- [42] J. Barrera and A. Garcia, "Dynamic incentives for congestion control," *IEEE Transactions on Automatic Control*, vol. 60, no. 2, pp. 299–310, 2014.
- [43] D. Paccagnan, B. Gentile, F. Parise, M. Kamgarpour, and J. Lygeros, "Nash and wardrop equilibria in aggregative games with coupling constraints," *IEEE Transactions on Automatic Control*, vol. 64, no. 4, pp. 1373–1388, 2018.
- [44] G. Belgioioso, P. Yi, S. Grammatico, and L. Pavel, "Distributed generalized Nash equilibrium seeking: An operator-theoretic perspective," *IEEE Control Systems Magazine*, vol. 42, no. 4, pp. 87–102, 2022.
- [45] A. A. Kulkarni and U. V. Shanbhag, "On the variational equilibrium as a refinement of the generalized Nash equilibrium," *Automatica*, vol. 48, no. 1, pp. 45–55, 2012.
- [46] F. Facchinei and J.-S. Pang, *Finite-dimensional variational inequalities and complementarity problems*. Springer, 2003.
- [47] Y. Wang and A. Nedić, "Tailoring gradient methods for differentially-private distributed optimization," *arXiv preprint arXiv:2202.01113*, 2022.
- [48] C. Dwork, M. Naor, T. Pitassi, and G. N. Rothblum, "Differential privacy under continual observation," in *Proceedings of the forty-second ACM Symposium on Theory of Computing*, 2010, pp. 715–724.
- [49] Z. Huang, S. Mitra, and N. Vaidya, "Differentially private distributed optimization," in *Proceedings of the 2015 International Conference on Distributed Computing and Networking*, New York, NY, USA, 2015.
- [50] P. Kairouz, S. Oh, and P. Viswanath, "The composition theorem for differential privacy," in *International Conference on Machine Learning*. PMLR, 2015, pp. 1376–1385.
- [51] M. Bun and T. Steinke, "Concentrated differential privacy: Simplifications, extensions, and lower bounds," in *Theory of Cryptography Conference*. Springer, 2016, pp. 635–658.
- [52] I. Mironov, "Rényi differential privacy," in *The 30th Computer Security Foundations Symposium*. IEEE, 2017, pp. 263–275.
- [53] P. L. Combettes and J.-C. Pesquet, "Stochastic quasi-Fejér block-coordinate fixed point iterations with random sweeping," *SIAM Journal on Optimization*, vol. 25, no. 2, pp. 1221–1248, 2015.
- [54] P. L. Combettes, "Quasi-Fejérian analysis of some optimization algorithms," in *Studies in Computational Mathematics*. Elsevier, 2001, vol. 8, pp. 115–152.
- [55] D. T. A. Nguyen, D. T. Nguyen, and A. Nedić, "Distributed Nash equilibrium seeking over time-varying directed communication networks," *arXiv preprint arXiv:2201.02323*, 2022.