Identities for the k-generalized Fibonacci sequence with negative indices and its zero-multiplicity

Jonathan García * Carlos A. Gómez †
 Florian Luca ‡

November 2, 2022

Abstract

In this paper, we prove identities for members of the *k*-generalized Fibonacci sequence with negative indices and we apply these identities to deduce an exact formula for its zero-multiplicity.

Key words and phrases: k-generalized Fibonacci sequence, zero–multiplicity of linear recurrences.

Mathematics Subject Classification 2020: 11B39, 11J86

1 Introduction

One of the most famous and curious numerical sequence, due to the large number of properties and relationships with other areas [8], is the Fibonacci sequence, denoted by $\mathbf{F} := (F_n)_{n\geq 0}$. Its initial values are $F_0 = 0$, $F_1 = 1$ and obeys the recurrence $F_n = F_{n-1} + F_{n-2}$ for all $n \geq 2$. This sequence has been generalized in many ways, some by preserving the initial conditions, and others by preserving the recurrence relation. In general, a sequence $(u_n)_{n\in\mathbb{Z}} \subseteq \mathbb{C}$ is a linear recurrence sequence of order $k \in \mathbb{Z}^+$ if it satisfies the recurrence relation $u_{n+k} = a_1u_{n+k-1} + a_2u_{n+k-2} + \cdots + a_ku_n$ for all $n \geq 0$ with coefficients $a_1, \ldots, a_k \in \mathbb{C}$ and $a_k \neq 0$. We assume that k is minimal with the above property. Such a sequence $(u_n)_{n\in\mathbb{Z}}$ has an associated characteristic

^{*}Departamento de Matemáticas, Universidad del Valle, 25360 Cali, Calle 13 No 100-00, Colombia. e-mail: garcia.jonathan@correounivalle.edu.co

[†]Departamento de Matemáticas, Universidad del Valle, 25360 Cali, Calle 13 No 100-00, Colombia; Research Group ALTENUA–Universidad del Valle: Álgebra, Teoría de Números y Aplicaciones, Colciencias code: COL0017217. e-mail: carlos.a.gomez@correounivalle.edu.co

[‡]School of Mathematics, University of the Witwatersrand, Johannesburg, South Africa; Research Group in Algebraic Structures and Applications, King Abdulaziz University, Jeddah, Saudi Arabia; Max Planck Institute for Software Systems Saarbrücken, Germany and Centro de Ciencias Matemáticas UNAM, Morelia, Mexico. e-mail: florian.luca@wits.ac.za

polynomial given by $\Psi_k(z) = z^k - a_1 z^{k-1} - \cdots - a_k$. Let $\alpha_1, \ldots, \alpha_s \in \mathbb{C}$ be the distinct roots of $\Psi_k(z)$. From the theory of linear recurrence sequences (see [10, Theorem C.1]) there are polynomials $p_1(z), p_2(z), \ldots, p_s(z)$ in $\mathbb{C}[z]$, uniquely determined by the initial values u_0, \ldots, u_{k-1} , such that the formula $u_n = p_1(n)\alpha_1^n + p_2(n)\alpha_2^n + \cdots + p_s(n)\alpha_s^n$ holds for all $n \in \mathbb{Z}$. Studying certain arithmetic properties of members of linear recurrence sequences (u_n) can be performed with methods from the theory of exponential Diophantine equations by using the above expression for u_n .

In this paper, we present some identities for the members of the k-generalized Fibonacci sequence with negative indices. The k-generalized Fibonacci sequence $F^{(k)}$ is the linear recurrence sequence of order k with initial values $F_{-i}^{(k)} = 0$ for $i = 0, 1, \ldots, k - 2$ and $F_1^{(k)} = 1$, where each subsequent term is the sum of the k previous ones $F_n^{(k)} = F_{n-1}^{(k)} + F_{n-2}^{(k)} + \cdots + F_{n-k}^{(k)}$, for all $n, k \geq 2$. This sequence can be extended to all integer indices n. Since $F_n^{(k)} > 0$ for n > 0, all zeros of $\mathbf{F}^{(k)}$ correspond to non-positive indices n. We denote $H^{(k)} := (H_n^{(k)})_{n\geq 0}$, where $H_n^{(k)} = F_{-n}^{(k)}$. This sequence obeys the recurrence relation

$$H_n^{(k)} = H_{n-k}^{(k)} - H_{n-(k-1)}^{(k)} - \dots - H_{n-1}^{(k)} \quad \text{for all} \quad n \ge k, \tag{1}$$

with initial values $H_i^{(k)} = 0$ for $0 \le i \le k - 2$ and $H_{k-1}^{(k)} = 1$.

In this paper, we find identities for members of $H^{(k)}$ and apply them to deduce an exact formula for the zero–multiplicity of $H^{(k)}$.

2 The main results

2.1 Identities for $H^{(k)}$

Our identities are similars to the given by Ferguson [3], Gabai [4] or Cooper and Howard [1] for the k-generalized Fibonacci sequence with positive indices.

Theorem 1. For all $k \geq 2$, the sequence $H^{(k)}$ satisfies

- (i) For all $0 \le m \le r \le k 2$, we have $H_{mk+r}^{(k)} = 0$.
- (*ii*) For all $m \in [1, k-1]$, we have $H_{mk-1}^{(k)} = 2^{m-1}$.

(*iii*) For all $0 \le r < m \le k - 1$,

$$H_{mk+r}^{(k)} = (-1)^{r+1} \left[\binom{m-1}{r} + \binom{m}{r+1} \right] 2^{m-r-2}.$$

(iv) For all $r \in [-1, k-2]$ and $m \ge k-1$,

$$H_{mk+r}^{(k)} = \sum_{i=0}^{l} (-1)^{ik+r+1} \left[\binom{m-i-1}{ik+r} + \binom{m-i}{ik+r+1} \right] 2^{m-i(k+1)-r-2},$$

where l = m - 1 if k = 2, and $l = \lfloor m/(k - 1) \rfloor$ if k > 2.

2.2 Zero–multiplicity for $H^{(k)}$

One of the classical problems in the theory of linear recurrence sequences is the *Skolem problem*: Given the linear recurrence sequence $\mathbf{u} := (u_n)_{n \in \mathbb{Z}}$, one want to find $\mathcal{Z}(\mathbf{u}) = \{n \in \mathbb{Z} : u_n = 0\}$. The cardinality of $\mathcal{Z}(\mathbf{u})$ (when this is finite) is called the *zero-multiplicity* of \mathbf{u} . There is no known algorithm to find $\mathcal{Z}(\mathbf{u})$ in general. One of the more important results here is due to Skolem [11, 10]: If the coefficients a_1, \ldots, a_k of the linear recurrence sequence \mathbf{u} are rational, then the set $\mathcal{Z}(\mathbf{u})$ is a union of finitely many arithmetical progressions together with a finite set. Hagedorn [7] showed that if the roots of $\Psi_k(z)$ are real then $\#\mathcal{Z}(\mathbf{u}) \leq 2k - 3$.

Our current research is motived by our previous work [5]. There we obtained that

$$\mathcal{A} := \bigcup_{m=0}^{k-2} [m(k+1), (m+1)k - 2] \subseteq \mathcal{Z}(H^{(k)}).$$
(2)

Thus, $\#\mathcal{Z}(H^{(k)}) \ge k(k-1)/2$. Furthermore, we checked that

$$#\mathcal{Z}(H^{(2)}) = 1, \ #\mathcal{Z}(H^{(3)}) = 4 \text{ and } \#\mathcal{Z}(H^{(k)}) = k(k-1)/2,$$

for $k \in [4, 500]$.

Based on the above results, we proposed in [5] the following conjecture.

Conjecture 1. The zero-multiplicity $\#\mathcal{Z}(H^{(k)})$ of the k-generalized Fibonacci sequence with non-positive indices $H^{(k)}$ for $k \ge 4$ is the (k-1)st triangular number; i.e.

$$#\mathcal{Z}(H^{(k)}) = k(k-1)/2.$$

Recently, relating the 2-adic valuation of $F^{(k)}$ with the Diophantine equation $H_n^{(k)} = 0$, Young [12] showed that for all k > 500,

$$#\mathcal{Z}(H^{(k)}) \le k(k+1)/2 + \lfloor k/2 \rfloor.$$

In this paper, we confirm the above conjecture. As a consequence of this, we get that $\mathcal{Z}(\mathbf{F}^{(\mathbf{k})})$ is exactly \mathcal{A} .

3 Preliminary results

To simplify the notation, from now on we denote $H_n^{(k)} := H_n$, where k is fixed. From the recurrence relation (1), it is easy to see that

$$H_n = 2H_{n-k} - (H_{n-k} + H_{n-(k-1)} + \dots + H_{n-2} + H_{n-1})$$

= $2H_{n-k} - H_{n-k-1}$ for all $n \ge k+1$. (3)

The containment (2) is a direct consequence of the initial values of H_n and the identity (3). Indeed, inductively we see that $H_n = 0$ for all $0 \le n \le k-2$. Further

 $H_n = 2H_{n-k} - H_{n-k-1} = 0$ for all $m(k+1) \le n \le mk + (k-2)$,

with $1 \le m \le k-2$. Moreover, the length of these intervals is given by the decreasing quantity

$$mk + (k-2) - m(k+1) = k - (m+2),$$

which becomes 0 at m = k - 2.

So far, we have only summarized what we obtained in [5]. This is fundamental for what follows, since we note that the sequence H_n can be represented as lists matrices, starting with a rectangular matrix of size $(k - 1) \times (k + 1)$ and continuing with square matrices of size $k \times k$. Then, we observed interesting patterns in H_n that we were able to formulate and prove.

We start by ordering the first $k^2 - 1$ elements of the sequence H_n in matrix form $(H_{(i-1)(k+1)+(j-1)})_{ij}$ with $1 \le i \le k-1$ and $1 \le j \le k+1$. This is

$$\begin{pmatrix} \mathbf{H_0} & \cdots & \mathbf{H_{k-2}} & H_{k-1} & H_k \\ \mathbf{H_{(k+1)+0}} & \cdots & H_{(k+1)+k-2} & H_{(k+1)+k-1} & H_{(k+1)+k} \\ \mathbf{H_{2(k+1)+0}} & \cdots & H_{2(k+1)+k-2} & H_{2(k+1)+k-1} & H_{2(k+1)+k} \\ \vdots & \ddots & \vdots & \vdots & \vdots \\ \mathbf{H_{(k-2)(k+1)+0}} & \cdots & H_{(k-2)(k+1)+k-2} & H_{(k-2)(k+1)+k-1} & H_{(k-2)(k+1)+k} \end{pmatrix}$$

and we notice that the "upper triangular" part, which is in **bold**, is composed of zeros. We can also write this matrix as

$$\begin{pmatrix} 0 & 0 & \cdots & 0 & 0 & H_{k-1} & H_k \\ 0 & 0 & \cdots & 0 & H_{2k-1} & H_{2k} & H_{2k+1} \\ 0 & 0 & \cdots & H_{3k-1} & H_{3k} & H_{3k+1} & H_{3k+2} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots \\ 0 & H_{(k-1)k-1} & \cdots & H_{(k-1)k+k-5} & H_{(k-1)k+k-4} & H_{(k-1)k+k-3} & H_{(k-1)k+k-2} \end{pmatrix},$$
(4)

and we observe the following behavior in the non-zero diagonals.

Lemma 1.

- (a) For all $m \in [1, k-1]$, we have $H_{mk-1} = 2^{m-1}$.
- (b) For all $m \in [1, k-1]$, we have $H_{mk} = -(m+1)2^{m-2}$.
- (c) For all $1 \le r < m \le k 1$,

$$H_{mk+r} = -\sum_{j=r}^{m-1} 2^{m-1-j} H_{jk+r-1}$$

Proof. To show that the items of this lemma are fulfilled, we proceed by induction on m, taking into account that the upper bound for m is $k-1 \ge 1$.

(a) If m = 1, given the initial values, we have $H_{k-1} = 1 = 2^0$. Suppose that $H_{(m-1)k-1} = 2^{m-2}$ for $m \in [2, k-1]$. Then, using identity (3), it follows that

$$H_{mk-1} = 2H_{(m-1)k-1} - H_{(m-1)k-2} = 2^{m-1},$$

since $0 \le m - 2 \le k - 3$ and therefore $H_{((m-2)+1)k-2} = 0$ (see (2)).

(b) When m = 1, we have

$$H_k = H_0 - H_1 - \dots - H_{k-1} = -1 = -2 \times 2^{-1}.$$

Assume $H_{(m-1)k} = -m2^{m-3}$ and we obtain from identity (3) and item (a), that

$$H_{mk} = 2H_{(m-1)k} - H_{(m-1)k-1} = -m2^{m-2} - 2^{m-2} = -(m+1)2^{m-2}.$$

(c) Assume m = r + 1. We have by identity (3) that

$$H_{(r+1)k+r} = 2H_{rk+r} - H_{rk+r-1} = -H_{rk+r-1} = -\sum_{j=r}^{r} 2^{r-j} H_{jk+r-1},$$

because $H_{rk+r} = H_{r(k+1)} = 0$ with $1 \le r \le k-2$ (see (2)). We take

$$H_{(m-1)k+r} = -\sum_{j=r}^{m-2} 2^{m-2-j} H_{jk+r-1}$$

as the inductive hypothesis. Then

$$H_{mk+r} = 2H_{(m-1)k+r} - H_{(m-1)k+r-1}$$

= $-\left(\sum_{j=r}^{m-2} 2^{m-1-j} H_{jk+r-1}\right) - H_{(m-1)k+r-1}$
= $-\sum_{j=r}^{m-1} 2^{m-1-j} H_{jk+r-1}.$

Matrix (4) is the first element of the list of matrices that we use to organize all the elements of H_n , as we mentioned before. It is rectangular of size $(k-1) \times (k+1)$. The matrix that follows is of size $k \times k$ and includes all the non-zero elements of the last row of matrix (4), this is

$$\begin{pmatrix} H_{(k^{2}-k-1)+0} & H_{(k^{2}-k-1)+1} & \cdots & H_{(k^{2}-k-1)+k-1} \\ H_{(k^{2}-k-1)+k} & H_{(k^{2}-k-1)+k+1} & \cdots & H_{(k^{2}-k-1)+k+(k-1)} \\ \vdots & \vdots & \ddots & \vdots \\ H_{(k^{2}-k-1)+(k-1)k} & H_{(k^{2}-k-1)+(k-1)k+1} & \cdots & H_{(k^{2}-k-1)+(k-1)k+(k-1)} \end{pmatrix}^{.}$$

$$(5)$$

In general, all square matrices after matrix (4) form the sequence $\{M_b\}_{b\in\mathbb{Z}^+}$, where

$$M_b := \begin{pmatrix} H_{(bk^2-bk-1)+0} & H_{(bk^2-bk-1)+1} & \cdots & H_{(bk^2-bk-1)+k-1} \\ H_{(bk^2-bk-1)+k} & H_{(bk^2-bk-1)+k+1} & \cdots & H_{(bk^2-bk-1)+k+(k-1)} \\ \vdots & \vdots & \ddots & \vdots \\ H_{(bk^2-bk-1)+(k-1)k} & H_{(bk^2-bk-1)+(k-1)k+1} & \cdots & H_{(bk^2-bk-1)+(k-1)k+(k-1)} \end{pmatrix}.$$

Furthermore, by calling M_0 the matrix (4), we have arrived at

$$(H_n^{(k)})_{n\geq 0} = \bigcup_{b\geq 0} \{h : h \text{ is an entry of } M_b\}$$

From now on, we simplify the notation by setting

$$H_{b,jk+r} := H_{(bk^2 - bk - 1) + jk + r}$$
 for all $b \ge 0$.

Note that matrix (5) is exactly M_1 . In fact, these matrices satisfy that their first row is exactly the last row of the immediately preceding matrix. Indeed,

$$H_{b,r} = H_{b-1,(k-1)k+r}$$
 for all $0 \le r \le k-1$ and $b \ge 1$. (6)

Also,

$$H_{b,-1} = H_{b-1,(k-2)k+(k-1)} \quad \text{for all } b \ge 1 \tag{7}$$

and in particular, by containment (2),

$$H_{1,-1} = H_{0,(k-2)k+(k-1)} = H_{(k-2)(k+1)} = 0.$$

Moreover, we observe that identity (3) is still satisfied with this notation

$$H_{b,n} = 2H_{b,n-k} - H_{b,n-k-1} \tag{8}$$

for all $b \ge 1$ and $n \ge 2$, or for b = 0 and $n \ge k + 2$.

Next, we find the following patterns in all entries below the main diagonal of M_b with $b \ge 1$.

Lemma 2. Let $b \ge 1$, then

(I)
$$H_{b,jk} = \begin{cases} 2H_{b,0} - H_{b-1,(k-2)k+(k-1)}, & \text{if } j = 1, \\ 2^{j-1}H_{b,k} - \sum_{i=0}^{j-2} 2^{j-i-2}H_{b,(i+1)k-1}, & \text{if } 2 \le j \le k-1. \end{cases}$$

(II) For all $r, j \in [1, k - 1]$, it holds that

$$H_{b,jk+r} = 2^{j-1}H_{b,k+r} - \sum_{i=1}^{j-1} 2^{j-1-i}H_{b,ik+r-1}.$$

(III) Let $r, j \in [0, k-1]$ be fixed. Then

$$H_{b,jk+r} = \sum_{i=0}^{t} (-1)^i \binom{t}{i} 2^{t-i} H_{b,(j-t)k+r-i},$$

for all $t \in \left[0, \left\lfloor \frac{bk^2 - bk - 1 + jk + r}{k+1} \right\rfloor\right].$

Proof. We consider each item separately.

(I) The case j = 1, follows from identities (7) and (8). Let $2 \le j \le k-1$. We argue by induction on j. If j = 2, item (I) is immediate by identity (8). Suppose it is satisfied for j - 1; i.e.,

$$H_{b,(j-1)k} = 2^{j-2} H_{b,k} - \sum_{i=0}^{j-3} 2^{j-i-3} H_{b,(i+1)k-1}.$$
 (9)

Then, by (8) and (9),

$$H_{b,jk} = 2H_{b,(j-1)k} - H_{b,(j-1)k-1}$$

= $\left(2^{j-1}H_{b,k} - \sum_{i=0}^{j-3} 2^{j-i-2}H_{b,(i+1)k-1}\right) - H_{b,(j-1)k-1}$
= $2^{j-1}H_{b,k} - \sum_{i=0}^{j-2} 2^{j-i-2}H_{b,(i+1)k-1},$

and thus we conclude item (I) for j.

(II) If j = 1, the identity is trivial. For $j \ge 2$, we apply recursively, j - 1 times, identity (8) on $H_{b,jk+r}$. That is,

$$\begin{split} H_{b,jk+r} &= 2H_{b,(j-1)k+r} - H_{b,(j-1)k+r-1} \\ &= 2^2 H_{b,(j-2)k+r} - 2H_{b,(j-2)k+r-1} - H_{b,(j-1)k+r-1} \\ &= 2^3 H_{b,(j-3)k+r} - 2^2 H_{b,(j-3)k+r-1} - 2H_{b,(j-2)k+r-1} - H_{b,(j-1)k+r-1} \\ &\vdots \\ &= 2^{j-1} H_{b,(j-(j-1))k+r} - 2^{j-2} H_{b,(j-(j-1))k+r-1} - \dots - H_{b,(j-1)k+r-1} \end{split}$$

(III) We proceed by induction on t. If t = 0, the identity is trivial. Suppose it is satisfied for $t - 1 \ge 0$; i.e.,

$$H_{b,jk+r} = \sum_{i=0}^{t-1} (-1)^i \binom{t-1}{i} 2^{t-i-1} H_{b,(j-t+1)k+r-i}.$$

Then, by the inductive hypothesis and identity (8), it follows that

$$H_{b,jk+r} = \sum_{i=0}^{t-1} (-1)^{i} {\binom{t-1}{i}} 2^{t-i-1} \left(2H_{b,(j-t)k+r-i} - H_{b,(j-t)k+r-i-1} \right)$$
$$= (-1)^{0} {\binom{t-1}{0}} 2^{t} H_{b,(j-t)k+r} + \sum_{i=1}^{t-1} (-1)^{i} {\binom{t-1}{i}} 2^{t-i} H_{b,(j-t)k+r-i}$$
$$+ \sum_{i=1}^{t} (-1)^{i} {\binom{t-1}{i-1}} 2^{t-i} H_{b,(j-t)k+r-i}$$

$$= 2^{t} H_{b,(j-t)k+r} + \sum_{i=1}^{t-1} (-1)^{i} \left(\binom{t-1}{i} + \binom{t-1}{i-1} \right) 2^{t-i} H_{b,(j-t)k+r-i} + (-1)^{t} \binom{t-1}{t-1} 2^{t-t} H_{b,(j-t)k+r-t} = 2^{t} H_{b,(j-t)k+r} + \sum_{i=1}^{t-1} (-1)^{i} \binom{t}{i} 2^{t-i} H_{b,(j-t)k+r-i} + (-1)^{t} H_{b,(j-t)k+r-t} = \sum_{i=0}^{t} (-1)^{i} \binom{t}{i} 2^{t-i} H_{b,(j-t)k+r-i},$$

and the identity is satisfied for t. We must assume that $t \leq \left\lfloor \frac{bk^2 - bk - 1 + jk + r}{k+1} \right\rfloor$, because in this case

$$bk^{2} - bk - 1 + (j - t)k + r - i \ge bk^{2} - bk - 1 + (j - t)k + r - t \ge 0,$$

for all $0 \le i \le t$. Therefore, $H_{b,(j-t)k+r-i}$ is well defined for all $0 \le i \le t$. This completes the proof of this lemma.

The above lemmas will allow us, in Section 4, to characterize all entries of the matrices M_b and therefore all elements of H_n . For this purpose we introduce the following notation:

$$\psi(v,w) := \binom{v}{w} + \binom{v+1}{w+1}.$$
(10)

Note that

$$\psi(v, -1) = 1$$
 for all $v \neq -1$ and $\psi(v, w) = 0$ for all $w > v \ge 0$. (11)

We prove the following properties of $\psi(v, w)$.

Lemma 3. The function ψ satisfies:

- (1) $\psi(v, w) + \psi(v, w + 1) = \psi(v + 1, w + 1).$
- (2) $\sum_{i=0}^{n} \psi(v+i,v) = \psi(v+n+1,v+1).$

- (3) $\sum_{i=-1}^{n} \psi(v+i,i) = \psi(v+n+1,n).$
- (4) $\sum_{i=w}^{v} \psi(i-1, w-1) = \psi(v, w).$

(5)
$$\sum_{i=0}^{w} {\binom{u}{i}} \psi(v, w - i - 1) = \psi(u + v, w - 1).$$

Proof. To show that all items are satisfied, we use basic properties of binomial coefficients. Item (1) is immediate.

(2) We know that

$$\sum_{i=0}^{n} \psi(v+i,v) = \sum_{i=v}^{v+n} \binom{i}{v} + \sum_{i=v+1}^{v+n+1} \binom{i}{v+1} = \psi(v+n+1,v+1).$$

(3) It is clear that

$$\begin{split} \sum_{i=-1}^{n} \psi(v+i,i) &= \sum_{i=0}^{n} \binom{v+i}{i} + \sum_{i=-1}^{n} \binom{v+i+1}{i+1} \\ &= \sum_{i=v}^{v+n} \binom{i}{v} + \sum_{i=v}^{v+n+1} \binom{i}{v} \\ &= \psi(v+n+1,n). \end{split}$$

(4) We have

$$\sum_{i=w}^{v} \psi(i-1, w-1) = \sum_{i=w-1}^{v-1} \binom{i}{w-1} + \sum_{i=w}^{v} \binom{i}{w} = \psi(v, w).$$

(5) We obtain

$$\sum_{i=0}^{w} \binom{u}{i} \psi(v, w - i - 1) = \sum_{i=0}^{w-1} \binom{u}{i} \binom{v}{w - i - 1} + \sum_{i=0}^{w} \binom{u}{i} \binom{v + 1}{w - i} = \psi(u + v, w - 1).$$

4 Proof of Theorem 1

For items (i) and (ii), see containment (2) and Lemma 1, item (a), respectively. The remaining items are shown below.

4.1 Item (*iii*)

With notation (10), we must prove that

$$H_{mk+r} = (-1)^{r+1} \psi(m-1,r) 2^{m-r-2} \quad \text{for all} \quad 0 \le r < m \le k-1.$$
(12)

For this, we apply induction on r. If r = 0, by Lemma 1, item (b), we get that identity (12) is satisfied. Indeed,

$$H_{mk} = -(m+1)2^{m-2} = (-1)^1 \psi(m-1,0)2^{m-2} \quad \text{for all} \quad 1 \le m \le k-1.$$

We assume by the inductive hypothesis that identity (12) is satisfied for r-1and therefore $0 \le r-1 < m \le k-1$; i.e.,

$$H_{mk+r-1} = (-1)^r \psi(m-1, r-1) 2^{m-r-1}$$
 for all $r \le m \le k-1$.

In particular,

$$H_{jk+r-1} = (-1)^r \psi(j-1,r-1) 2^{j-r-1} \quad \text{for all} \quad r \le j \le m-1.$$
(13)

Then, by Lemma 1, item (c), Lemma 3, item (4) and identity (13), we obtain

$$H_{mk+r} = -\sum_{j=r}^{m-1} 2^{m-1-j} \left((-1)^r \psi(j-1,r-1) 2^{j-r-1} \right)$$

= $(-1)^{r+1} \left(\sum_{j=r}^{m-1} \psi(j-1,r-1) \right) 2^{m-r-2}$
= $(-1)^{r+1} \psi(m-1,r) 2^{m-r-2}$ for all $1 \le r < m \le k-1$.

This completes the proof of item (iii).

4.2 Item (*iv*)

4.2.1 Case k = 2:

Here we must prove that for all $m \ge 1$,

$$H_{2m+r} = \begin{cases} \sum_{i=0}^{m-1} \psi(m-i-1,2i-1)2^{m-3i-1}, & \text{if } r = -1, \\ -\sum_{i=0}^{m-1} \psi(m-i-1,2i)2^{m-3i-2}, & \text{if } r = 0. \end{cases}$$
(14)

For this we proceed by induction on m. Suppose that identity (14) is satisfied up to $m-1 \ge 2$ (the cases m = 1 and m = 2 are easily verified). Then using (3), (11), Lemma 3 and the fact that $\psi(0, 2m-2) = \psi(0, 2m-3) = 0$ (since $m \ge 2$), we obtain

$$\begin{split} H_{2m-1} &= 2H_{2(m-1)-1} - H_{2(m-2)} \\ &= \sum_{i=0}^{m-2} \psi(m-i-2,2i-1)2^{m-3i-1} + \sum_{i=0}^{m-3} \psi(m-i-3,2i)2^{m-3i-4} \\ &= \psi(m-2,-1)2^{m-1} + \sum_{i=1}^{m-2} (\psi(m-i-2,2i-1) + \psi(m-i-2,2i-2))2^{m-3i-1} \\ &= \psi(m-1,-1)2^{m-1} + \sum_{i=1}^{m-2} \psi(m-i-1,2i-1)2^{m-3i-1} + \psi(0,2m-3)2^{-2(m-1)} \\ &= \sum_{i=0}^{m-1} \psi(m-i-1,2i-1)2^{m-3i-1}, \end{split}$$

and

$$\begin{aligned} H_{2m} &= 2H_{2(m-1)} - H_{2(m-1)-1} \\ &= -\sum_{i=0}^{m-2} \psi(m-i-2,2i)2^{m-3i-2} - \sum_{i=0}^{m-2} \psi(m-i-2,2i-1)2^{m-3i-2} \\ &= -\psi(0,2m-2)2^{-2m+1} - \sum_{i=0}^{m-2} \left(\psi(m-i-2,2i) + \psi(m-i-2,2i-1)\right)2^{m-3i-2} \\ &= -\sum_{i=0}^{m-1} \psi(m-i-1,2i)2^{m-3i-2}. \end{aligned}$$

4.2.2 Case k > 2:

To treat this case, we present the following two lemmas.

Lemma 4. For all k > 2 and $j, r \in [0, k - 1]$, it holds that

$$H_{1,jk+r} = (-1)^r 2^{k+j-r-2} \psi(k+j-2,r-1) + (-1)^{k+r} 2^{j-r-3} \psi(k+j-3,k+r-1).$$
(15)

Proof. We apply double induction on $j, r \in [0, k-1]$.

(A) Let j = r = 0. Since k > 2, by observation (11), it follows that

$$\psi(k-2,-1) = 1$$
 and $\psi(k-3,k-1) = 0.$

Then, using item (ii), we obtain

$$H_{1,0} = H_{(k-1)k-1} = 2^{k-2} = 2^{k-2}\psi(k-2,-1) + (-1)^k 2^{-3}\psi(k-3,k-1)$$

fulfilling identity (15).

(B) Let j = 0 and $r \in [1, k - 1]$. Then, by item (*iii*), we get that

$$H_{1,r} = H_{k(k-1)+r-1} = (-1)^r \psi(k-2, r-1) 2^{k-r-2}$$

and by observation (11) it follows that

$$H_{1,r} = (-1)^r 2^{k-r-2} \psi(k-2, r-1)$$
(16)
= $(-1)^r 2^{k-r-2} \psi(k-2, r-1) + (-1)^{k+r} 2^{-r-3} \psi(k-3, k+r-1).$

Thus, identity (15) is satisfied.

(C) Let r = 0 and $j \in [1, k - 1]$. If j = 1, by Lemma 2, item (I) and observation (11), we have that identity (15) is satisfied. In fact,

$$H_{1,k} = 2H_{1,0} - H_{0,(k-2)k+(k-1)} = 2H_{1,0} = 2^{k-1}$$
(17)
= $2^{k-1}\psi(k-1,-1) + (-1)^k 2^{-2}\psi(k-2,k-1).$

If $j \in [2, k - 1]$, we note that by Lemma 2, item (I), combined with item (III) * of this same Lemma and items (A), (B) above, we get

$$H_{1,jk} = 2^{j-1}H_{1,k} - \sum_{i=0}^{j-2} 2^{j-i-2}H_{1,ik+k-1}$$

$$= 2^{j-1}H_{1,k} + \sum_{i=0}^{j-2} \left(\sum_{\ell=0}^{i} (-1)^{\ell+1} \binom{i}{\ell} 2^{j-2-\ell}H_{1,k-1-\ell}\right)$$

$$= 2^{j-1}H_{1,k} + \left(\sum_{\ell=0}^{j-2} (-1)^{\ell+1} \left(\sum_{i=\ell}^{j-2} \binom{i}{\ell}\right) 2^{j-2-\ell}H_{1,k-1-\ell}\right)$$

$$= 2^{j-1}H_{1,k} + \left(\sum_{\ell=0}^{j-2} (-1)^{\ell+1} \binom{j-1}{\ell+1} 2^{j-2-\ell}H_{1,k-1-\ell}\right)$$

$$= 2^{j-1}H_{1,k} + (-1)^{k}2^{j-3} \left(\sum_{\ell=1}^{j-1} \binom{j-1}{\ell} \psi(k-2,k-1-\ell)\right). \quad (18)$$

In addition,

$$\psi(k-2,k-1) = 0$$
 and $\binom{j-1}{\ell} = 0$ for $j \le \ell \le k$. (19)

Then, by identities (17), (18), (19), observation (11) and Lemma 3 (item 5), we obtain

$$H_{1,jk} = 2^{k+j-2} + (-1)^k 2^{j-3} \left(\sum_{\ell=0}^k \binom{j-1}{\ell} \psi(k-2,k-\ell-1) \right)$$

= $2^{k+j-2} \psi(k+j-2,-1) + (-1)^k 2^{j-3} \psi(k+j-3,k-1).$

Thus, identity (15) is satisfied.

(D) Let $r \in [1, k - 1]$ be fixed and assume as inductive hypothesis that identity (15) is satisfied up to $j - 1 \ge 0$. Then replacing r by r - 1 in the inductive hypothesis, we have

$$H_{1,ik+r-1} = (-1)^{r-1} 2^{k+i-r-1} \psi(k+i-2,r-2) + (-1)^{k+r-1} 2^{i-r-2} \psi(k+i-3,k+r-2),$$

for all $1 \le i \le j-1$ and $r-1 \in [0, k-2]$ fixed[†]. In particular, this also implies (see (11)) that

$$H_{1,k+r} = (-1)^r 2^{k-r-1} \psi(k-1,r-1) + (-1)^{k+r} 2^{-r-2} \psi(k-2,k+r-1)$$

= $(-1)^r 2^{k-r-1} \psi(k-1,r-1)$

^{*}This item is used with j = i, r = k - 1 and t = i for the elements $H_{1,ik+k-1}$. †The case r - 1 = 0 is obtained from Item (C) above.

for $r \in [1, k - 1]$ fixed. Thus, by the above two identities and Lemma 2, item (II), we get

$$H_{b,jk+r} = 2^{j-1}H_{b,k+r} - \sum_{i=1}^{j-1} 2^{j-1-i}H_{b,ik+r-1}$$

= $(-1)^r 2^{k+j-r-2}\psi(k-1,r-1)$
+ $(-1)^r 2^{k+j-r-2}\sum_{i=1}^{j-1}\psi(k+i-2,r-2)$
+ $(-1)^{k+r} 2^{j-r-3}\sum_{i=1}^{j-1}\psi(k+i-3,k+r-2)$ (20)

for all $r, j \in [1, k - 1]$, with r fixed. We note that using the identity (1) of Lemma 3, it follows that

$$\psi(k-1,r-1) + \sum_{i=1}^{j-1} \psi(k+i-2,r-2)$$

= $\psi(k,r-1) + \sum_{i=2}^{j-1} \psi(k+i-2,r-2)$
= $\psi(k+1,r-1) + \sum_{i=3}^{j-1} \psi(k+i-2,r-2)$
:
 $\psi(k+i-2,r-2)$

$$= \psi(k+j-3,r-1) + \sum_{i=j-1}^{j-1} \psi(k+i-2,r-2)$$
$$= \psi(k+j-2,r-1).$$
(21)

Furthermore, by observation (11),

$$\psi(k+i-3, k+r-2) = 0$$
 for $i < r+1$. (22)

Then, by the identities (20), (21), (22) and Lemma 3, item (2), we obtain

$$\begin{split} H_{b,jk+r} &= (-1)^r 2^{k+j-r-2} \left(\psi(k-1,r-1) + \sum_{i=1}^{j-1} \psi(k+i-2,r-2) \right) \\ &+ (-1)^{k+r} 2^{j-r-3} \sum_{i=1}^{j-1} \psi(k+i-3,k+r-2) \\ &= (-1)^r 2^{k+j-r-2} \psi(k+j-2,r-1) \\ &+ (-1)^{k+r} 2^{j-r-3} \psi(k+j-3,k+r-1). \end{split}$$

Therefore, identity (15) is satisfied.

(E) We fix $j \in [1, k-1]$ and take as inductive hypothesis the identity (15) for $r-1 \ge 0$. In items (A), (C) and (D) above, we proved that identity (15) is satisfied on H_{ik+r-1} , for fixed $r-1 \ge 0$ and all $i \in [0, k-1]$. Then

$$H_{1,ik+r-1} = (-1)^{r-1} 2^{k+i-r-1} \psi(k+i-2,r-2) + (-1)^{k+r-1} 2^{i-r-2} \psi(k+i-3,k+r-2)$$

holds for all $0 \le i \le j - 1$. Thus, by the above identity, identity (16) and Lemma 2, item (II), it follows that

$$H_{1,jk+r} = 2^{j}H_{1,r} - \sum_{i=0}^{j-1} 2^{j-1-i}H_{1,ik+r-1}$$

= $(-1)^{r}2^{k+j-r-2}\psi(k-2,r-1) + (-1)^{r}2^{k+j-r-2}\sum_{i=0}^{j-1}\psi(k+i-2,r-2)$
+ $(-1)^{k+r}2^{j-r-3}\sum_{i=0}^{j-1}\psi(k+i-3,k+r-2)$ (23)

holds for all $r, j \in [1, k-1]$, with j fixed. Now, by identity (1) of Lemma 3 and identity (21), we deduce that

$$\psi(k-2,r-1) + \sum_{i=0}^{j-1} \psi(k+i-2,r-2)$$

= $\psi(k-1,r-1) + \sum_{i=1}^{j-1} \psi(k+i-2,r-2) = \psi(k+j-2,r-1).$ (24)

Therefore, identities (22), (23), (24) and Lemma 3, item (2), lead us to

$$\begin{split} H_{1,jk+r} &= (-1)^r 2^{k+j-r-2} \psi(k+j-2,r-1) \\ &+ (-1)^{k+r} 2^{j-r-3} \sum_{i=r+1}^{j-1} \psi(k+i-3,k+r-2) \\ &= (-1)^r 2^{k+j-r-2} \psi(k+j-2,r-1) \\ &+ (-1)^{k+r} 2^{j-r-3} \sum_{i=0}^{j-r-2} \psi(k+r-2+i,k+r-2) \\ &= (-1)^r 2^{k+j-r-2} \psi(k+j-2,r-1) \\ &+ (-1)^{k+r} 2^{j-r-3} \psi(k+j-3,k+r-1), \end{split}$$

thus satisfying identity (15).

Finally, items (A), (B), (C), (D) and (E) above prove the lemma.

The following result generalizes the previous one.

Lemma 5. For all k > 2; $j, r \in [0, k - 1]$ and $b \in \mathbb{Z}^+$, it holds that

$$H_{b,jk+r} = \sum_{i=0}^{b} (-1)^{ik+r} 2^{(b-i)k+j-r-b-i-1} \psi(bk+j-b-i-1,ik+r-1).$$
(25)

Proof. Here, we proceed by induction in b. The case b = 1 follows from Lemma 4. We state as inductive hypothesis that

$$H_{b-1,jk+r} = \sum_{i=0}^{b-1} (-1)^{ik+r} 2^{(b-i-1)k+j-r-b-i} \psi((b-1)k+j-b-i,ik+r-1), \quad (26)$$

for $b-1 \ge 1$, k > 2 and $j, r \in [0, k-1]$. Next, we consider the following cases.

(A') Let $j \leq r$. Then, by Lemma 2 (item *III*) with t = j and the identities (26), (6), we have that

$$H_{b,jk+r} = \sum_{\ell=0}^{j} (-1)^{\ell} {\binom{j}{\ell}} 2^{j-\ell} H_{b-1,(k-1)k+r-\ell}$$

= $\sum_{i=0}^{b-1} (-1)^{ik+r} 2^{(b-i)k+j-r-b-i-1} \sum_{\ell=0}^{j} {\binom{j}{\ell}} \psi(bk-b-i-1,ik+r-\ell-1).$ (27)

We note that $j \leq r < ik + r$ for all $i \geq 0$, and

$$\binom{j}{\ell} = 0 \quad \text{for } j+1 \le \ell \le ik+r.$$
(28)

We use the identities (27), (28) and Lemma 3, item (5), to obtain

$$H_{b,jk+r} = \sum_{i=0}^{b-1} (-1)^{ik+r} 2^{(b-i)k+j-r-b-i-1} \sum_{\ell=0}^{ik+r} {j \choose \ell} \psi(bk-b-i-1,ik+r-\ell-1)$$
$$= \sum_{i=0}^{b-1} (-1)^{ik+r} 2^{(b-i)k+j-r-b-i-1} \psi(bk+j-b-i-1,ik+r-1).$$
(29)

In fact, b(k-2) + j - 1 < bk + r - 1 and therefore (see (11))

$$\psi(b(k-2) + j - 1, bk + r - 1) = 0.$$

So, the sum in identity (29) can be extended to b, fulfilling identity (25) in this case.

(B') Let r < j. Here, we proceed by induction on r.

• For r = 0, we apply induction on j > 0. If j = 1, by the identities (6), (7) and (26) we obtain that

$$H_{b,k} = 2H_{b,0} - H_{b,-1} = 2H_{b-1,(k-1)k} - H_{b-1,(k-2)k+(k-1)}$$
$$= \sum_{i=0}^{b-1} (-1)^{ik} 2^{(b-i)k-b-i} \psi(bk-b-i-1,ik-1)$$
$$- \sum_{i=0}^{b-1} (-1)^{ik+k-1} 2^{(b-i-1)k-b-i-1} \psi(bk-b-i-2,ik+k-2).$$

In addition,

$$\psi(bk - b - b - 1, bk - 1) = 0,$$

$$\psi(bk - b - 1, -1) = \psi(bk - b, -1) = 1$$

(see (11)) and by Lemma 3, item (1) we obtain

$$H_{b,k} = \sum_{i=0}^{b-1} (-1)^{ik} 2^{(b-i)k-b-i} \psi(bk-b-i-1,ik-1) + \sum_{i=1}^{b} (-1)^{ik} 2^{(b-i)k-b-i} \psi(bk-b-i-1,ik-2) = 2^{bk-b} \psi(bk-b-1,-1) + \sum_{i=1}^{b} (-1)^{ik} 2^{(b-i)k-b-i} (\psi(bk-b-i-1,ik-1) + \psi(bk-b-i-1,ik-2)) = \sum_{i=0}^{b} (-1)^{ik} 2^{(b-i)k-b-i} \psi(bk-b-i,ik-1)$$
(30)

satisfying identity (25).

• Now, we assume that identity (25) is satisfied for j - 1, when r = 0. We obtain that

$$H_{b,jk} = 2H_{b,(j-1)k} - H_{b,(j-1)k-1} = 2H_{b,(j-1)k} - H_{b,(j-2)k+k-1}$$
$$= -H_{b,(j-2)k+k-1} + \sum_{i=0}^{b} (-1)^{ik} 2^{(b-i)k+j-b-i-1} \psi(bk+j-b-i-2,ik-1).$$
(31)

We also note that j - 2 < k - 1, so item (A') above implies that

$$H_{b,(j-2)k+k-1} = \sum_{i=0}^{b} (-1)^{ik+k-1} 2^{(b-i-1)k+j-b-i-2} \psi(bk+j-b-i-3,ik+k-2).$$
(32)

Then, since

$$\psi(bk+j-2b-3,bk+k-2) = 0$$

$$\psi(bk+j-b-2,-1) = \psi(bk+j-b-1,-1) = 1$$

(see (11)), we use identities (31), (32) and Lemma 3, item (1) to obtain

$$\begin{split} H_{b,jk} &= 2^{bk+j-b-1}\psi(bk+j-b-2,-1) \\ &+ \sum_{i=1}^{b} (-1)^{ik} 2^{(b-i)k+j-b-i-1}\psi(bk+j-b-i-2,ik-2) \\ &+ \sum_{i=1}^{b} (-1)^{ik} 2^{(b-i)k+j-b-i-1}\psi(bk+j-b-i-2,ik-1) \\ &= \sum_{i=0}^{b} (-1)^{ik} 2^{(b-i)k+j-b-i-1}\psi(bk+j-b-i-1,ik-1). \end{split}$$

Therefore identity (25) holds in this case. This concludes the induction on j when r = 0. Thus, the base case of induction on r is done.

• Let j = r + 1. Item (A') above implies that

$$H_{b,k+\ell} = \sum_{i=0}^{b} (-1)^{ik+\ell} 2^{(b-i)k-\ell-b-i} \psi(bk-b-i,ik+\ell-1)$$
(33)

for $1 \leq \ell \leq r$. By Lemma 2, item (*III*) with t = r, and the identities (30), (33), we obtain

$$H_{b,(r+1)k+r} = \sum_{\ell=0}^{r} (-1)^{r-\ell} {r \choose r-\ell} 2^{\ell} H_{b,k+\ell}$$

$$= \sum_{\ell=0}^{r} {r \choose \ell} \left(\sum_{i=0}^{b} (-1)^{ik+r} 2^{(b-i)k-b-i} \psi(bk-b-i,ik+\ell-1) \right)$$

$$= \sum_{i=0}^{b} (-1)^{ik+r} 2^{(b-i)k-b-i} \sum_{\ell=0}^{r} {r \choose r-\ell} \psi(bk-b-i,ik+r-\ell-1).$$
(34)

Furthermore, $r \leq ik + r$ for $i \geq 0$ and

$$\binom{r}{r-\ell} = \binom{r}{\ell} = 0, \quad \text{for} \quad r+1 \le \ell \le ik+r \quad \text{when} \quad i \ge 1.$$
(35)

Then, by identities (34), (35) and Lemma 3, item (5), we arrive at

$$H_{b,(r+1)k+r} = \sum_{i=0}^{b} (-1)^{ik+r} 2^{(b-i)k-b-i} \sum_{\ell=0}^{ik+r} \binom{r}{\ell} \psi(bk-b-i,ik+r-\ell-1)$$
$$= \sum_{i=0}^{b} (-1)^{ik+r} 2^{(b-i)k-b-i} \psi(bk-b-i+r,ik+r-1).$$

The above identity verifies identity (25) in this case, and we finish the base step of induction on j and on r.

• We assume by the inductive hypothesis that identity (25) is fulfilled for r-1 and up to j-1; i.e.,

$$H_{b,\ell k+r-1} = \sum_{i=0}^{b} (-1)^{ik+r-1} 2^{(b-i)k+\ell-r-b-i} \psi(bk+\ell-b-i-1,ik+r-2)$$
(36)

for $r \leq \ell \leq j-1$. Also, by item (A') above, we obtain that identity (36) holds for $0 \leq \ell \leq r-1$. In fact, it implies that

$$H_{b,r} = \sum_{i=0}^{b} (-1)^{ik+r} 2^{(b-i)k-r-b-i-1} \psi(bk-b-i-1,ik+r-1).$$

Then, by Lemma 2, item (II) and identity (36), we arrive at

$$H_{b,jk+r} = 2^{j}H_{b,r} - \sum_{\ell=0}^{j-1} 2^{j-1-\ell}H_{b,\ell k+r-1}$$

$$= \sum_{i=0}^{b} (-1)^{ik+r}2^{(b-i)k+j-r-b-i-1}\psi(bk-b-i-1,ik+r-1)$$

$$+ \sum_{i=0}^{b} (-1)^{ik+r}2^{(b-i)k+j-r-b-i-1}\sum_{\ell=0}^{j-1}\psi(bk+\ell-b-i-1,ik+r-2)$$
(37)

for all $r, j \in [1, k - 1]^{\ddagger}$. We observe, by Lemma (3), item (1), that

$$\psi(bk - b - i - 1, ik + r - 1) + \sum_{\ell=0}^{j-1} \psi(bk + \ell - b - i - 1, ik + r - 2)$$
$$= \psi(bk - b - i, ik + r - 1) + \sum_{\ell=1}^{j-1} \psi(bk + \ell - b - i - 1, ik + r - 2)$$

$$= \psi(bk - b - i + 1, ik + r - 1) + \sum_{\ell=2}^{j-1} \psi(bk + \ell - b - i - 1, ik + r - 2)$$

$$\vdots$$

$$= \psi(bk - b - i + j - 2, ik + r - 1) + \psi(bk + j - b - i - 2, ik + r - 2)$$

$$= \psi(bk - b - i + j - 1, ik + r - 1).$$
(38)

[‡]The case r = 0 is made in the first item of this item (B').

Then, by the identities (37) and (38), we obtain

$$H_{b,jk+r} = \sum_{i=0}^{b} (-1)^{ik+r} 2^{(b-i)k+j-r-b-i-1} \left(\psi(bk-b-i-1,ik+r-1) + \sum_{\ell=0}^{j-1} \psi(bk+\ell-b-i-1,ik+r-2) \right)$$
$$= \sum_{i=0}^{b} (-1)^{ik+r} 2^{(b-i)k+j-r-b-i-1} \psi(bk-b-i+j-1,ik+r-1)$$

and identity (25) is satisfied. This finishes the induction on both j and r thus completing the proof of this lemma.

Now we return to the proof of item (iv). Replacing b(k-1) + j by m and r-1 by r in Lemma 5, we obtain that

$$H_{mk+r} = H_{(b(k-1)+j)k+r} = H_{b,jk+r+1}$$
(39)

$$=\sum_{i=0}^{b} (-1)^{ik+r+1} \psi(m-i-1,ik+r) 2^{m-i(k+1)-r-2}, \qquad (40)$$

for all $k > 2, j \in [0, k - 1], r \in [-1, k - 2]$ and $b \in \mathbb{Z}^+$. By identity (6), we have

$$H_{b,r+1} = H_{b-1,(k-1)k+r+1}$$
 for all $-1 \le r \le k-2$ and $b \in \mathbb{Z}^+$,

so we can assume that $j \in [0, k - 1)$ (in order not to repeat elements of the sequence in (39)). Then

$$\frac{m}{k-1} - 1 < \frac{m-j}{k-1} \le \frac{m}{k-1} \quad \text{for} \quad k-1 > 1,$$

therefore $\lfloor \frac{m}{k-1} \rfloor - 1 < b \leq \lfloor \frac{m}{k-1} \rfloor$ and $b = \lfloor \frac{m}{k-1} \rfloor$. In conclusion, item (iv) follows from replacing b by $\lfloor \frac{m}{k-1} \rfloor$ in identity $(40)^{\S}$.

This completes the proof of Theorem 1.

5 Proof of Conjecture 1

We distinguish two cases according to the parity of k. Furthermore, since item (i) of Theorem 1 determines the set of zeros of $H^{(k)}$ in (2) and items (ii)-(iii) don't provide zeros for $H^{(k)}$, we can assume that $m \ge k - 1$.

[§]Note that $m = b(k-1) + j \ge k - 1$.

5.1 Case k even

In this case, by item (iv) in Theorem 1, we get that for all $r \in [-1, k-2]$ and $m \ge k-1$,

$$H_{mk+r} = (-1)^{r+1} \sum_{i=0}^{l} \left[\binom{m-i-1}{ik+r} + \binom{m-i}{ik+r+1} \right] 2^{m-i(k+1)-r-2},$$

where l = m - 1 if k = 2, and $l = \lfloor m/(k-1) \rfloor$ if k > 2. So, for $n = mk + r \ge k^2 - k - 1$, we have $H_n < 0$ if n is even, while $H_n > 0$ if n is odd, given that n and r have same parity because k is even.

Thus, item (i) of Theorem 1 implies that in this case the zero-multiplicity of $H^{(k)}$ is exactly k(k-1)/2, confirming Conjecture 1 in this case.

5.2 Case k odd

Since we verified in our previous work [5] that $\#\mathcal{Z}(H^{(k)}) = k(k-1)/2$ for $k \in [4, 500]$, we will assume that $H_n = 0$, for n = mk+r, with $m \ge k-1, r \in [-1, k-2]$ and k > 500 odd.

5.2.1 A lower bound for n in terms of k

By item (iv) of Theorem 1, we have after simplifying a factor of 2^{m-r-2} , that

$$\sum_{i=0}^{l} (-1)^{i} \psi(m-i-1, ik+r) 2^{-i(k+1)} = 0, \qquad for \ l = \lfloor m/(k-1) \rfloor.$$
(41)

We note that in the sum of equation (41), the term for i = 0 is non-zero since $r \leq k-2 < k-1 \leq m$, which implies that $r \leq m-1$ and therefore $\psi(m-1,r) \neq 0$. If the case i = 0 were the only for which $\psi(m-i-1,ik+r)$ is non-zero, we would have that equation (41) has no solution and our problem of zero multiplicity for the odd k case would be solved. So, we must assume that at least two terms of the above sum (the first with i = 0 and the second with i odd) are non-zero.

Next, we take l' to be the largest index i > 0 for which the *i*-th term of the sum in equation (41) is non-zero and see that equation (41) takes the form

$$\sum_{i=0}^{l'} (-1)^i \psi(m-i-1, ik+r) 2^{-i(k+1)} = 0, \quad \text{for } 0 < l' \le \lfloor m/(k-1) \rfloor,$$

where $\psi(m - l' - 1, l'k + r)$ is non-zero.

Separating the case i = l', we get

$$\psi(m-l'-1,l'k+r) = 2^{k+1} \sum_{i=0}^{l'-1} (-1)^{l'+i} \psi(m-i-1,ik+r) 2^{(l'-i-1)(k+1)},$$

with $l' - i - 1 \ge 0$ for all i = 0, ..., l' - 1. Hence,

$$2^{k+1} \mid \psi(m-l'-1, l'k+r), \quad \text{where} \quad l' \leq \lfloor m/(k-1) \rfloor.$$

Now,

$$\psi(v,w) = \binom{v}{w} + \binom{v+1}{w+1} = \binom{v}{w} \left(1 + \frac{v+1}{w+1}\right)$$

Kummer [9] proved that $\nu_2\left(\binom{v}{w}\right)$ equals the number of carries when adding w with v - w in base 2. Here, for a nonzero integer m, $\nu_2(m)$ is the exponent of 2 in the factorization of m. In particular,

$$\nu_2\left(\binom{v}{w}\right) \le \frac{\log v}{\log 2} + 1$$

Hence,

$$\nu_2(\psi(v,w)) \le \nu_2\left(\binom{v}{w}\right) + \nu_2(v+w+2) \le \frac{\log v}{\log 2} + 1 + \frac{\log(v+w+2)}{\log 2} \le \frac{2\log v}{\log 2} + 2,$$

where for the last inequality we used the fact that we may assume that $w \leq v - 2$ (indeed, if w = v, then $\psi(v, w) = 2$, so the above bound holds while if w = v - 1, then $\psi(v, w) = 2v + 1$ is odd so the above bound again holds). Since

$$2^{k+1} \mid \psi(v, w), \text{ for } (v, w) = (m - l' - 1, l'k + r),$$

we get $\nu_2(\psi(v, w)) \ge k + 1$. Hence,

$$k+1 \le \frac{2\log v}{\log 2} + 2$$
, therefore $v \ge 2^{(k-1)/2}$

Since $m - l' - 1 \le n$, we get that

$$2^{(k-1)/2} \le n. \tag{42}$$

5.2.2 An upper bound for n in terms of k

We now review the preliminary work [5] on zero–multiplicity of $H^{(k)}$, to obtain a better upper bound for n on k, which we then combine with the above lower bound (42). In [5], we used a Binet–type formula of H_n , namely

$$H_n = f_k(\alpha_1)\alpha_1^{-(n+1)} + \dots + f_k(\alpha_{k-1})\alpha_{k-1}^{-(n+1)} + f_k(\alpha_k)\alpha_k^{-(n+1)}$$

where $\alpha_1, \ldots, \alpha_{k-1}, \alpha_k$ are all the roots of the characteristic polynomial of $F^{(k)}$, with $\alpha_1 > 1$ being the only real root and the remaining k - 1 roots lie inside the unit disk. Furthermore,

$$|\alpha_1 > |\alpha_2| \ge \dots \ge |\alpha_{k-1}| \ge |\alpha_k|$$
 and $f_k(z) := (z-1)/(2 + (k+1)(z-2))$.

Thus, if $H_n = 0$, then

$$\begin{aligned} |\Lambda| &:= \left| 1 + \left(\frac{f_k(\alpha_{k-1})}{f_k(\alpha_k)} \right) \left(\frac{\alpha_k}{\alpha_{k-1}} \right)^{n+1} \right| &= \sum_{i=1}^{k-2} \left| \frac{f_k(\alpha_i)}{f_k(\alpha_k)} \right| \left| \frac{1}{\alpha_i/\alpha_k} \right|^{n+1} \\ &< \frac{3(k-2)/(k-1)}{|\alpha_{k-2}/\alpha_k|^{n+1}} \left| \frac{1}{f_k(\alpha_k)} \right| (43) \\ &< \frac{13k(3k+1)}{|\alpha_{k-2}/\alpha_k|^{n+1}}. \end{aligned}$$

Using an argument involving lower bounds for nonzero linear forms in logarithms of complex algebraic numbers, we found

$$|\Lambda| > \exp(-5 \cdot 10^{13} \times k^7 \log(n+1)(\log k)^2).$$
(44)

The following result will be fundamental to combine inequalities (43) and (44), in order to find a better upper bound for n on k that one given in [5]. The next lemma is Theorem 1 in [6] and represents an improvement over the analogous inequality in [2].

Lemma 6. The inequality

$$\frac{|\alpha_i|}{|\alpha_j|} > 1 + \frac{1}{10k^{9.6}(\pi/e)^k} \quad \text{holds for all} \quad 1 \le i < j \le (k-1)/2$$

and all $k \geq 4$.

Hence, by inequalities (43), (44) and Lemma 6

$$n < 3 \cdot 10^{14} k^{17.6} (\log k)^3 (\pi/e)^k.$$
(45)

5.2.3 Absolute bounds for n, k and final conclusion

Combining (42) and (45), we get

$$\left(\frac{\sqrt{2}}{(\pi/e)}\right)^k < 3\sqrt{2} \cdot 10^{14} k^{17.6} (\log k)^3,$$

showing that $k \leq 790$.

By (43) and (44), we have

$$5 \cdot 10^{13} \times k^7 \log(n+1) (\log k)^2 > \log \left| \frac{f_k(\alpha_k)}{3} \right| + (n+1) \log \left| \frac{\alpha_{k-2}}{\alpha_k} \right|.$$
(46)

Thus, using the fact that $k \in [501, 789]$, we obtain computationally an upper bound on n in each case:

If
$$H_n = 0$$
 and $k \in [501, 789]$, then $n \in 2.5 \cdot 10^{45}$.

Returning to inequality (42), we have $2^{(k-1)/2} < n < 2.5 \cdot 10^{45}$ which leads to $k \leq 517$. Finally, we return once again to (46) where now we get $n < 3.5 \cdot 10^{43}$ for all odd $k \in [501, 517]$. Then, by (42), we conclude that k < 500, contradicting our initial assumption about k.

This completes the proof of Conjecture 1 for the case of k odd.

Acknowledgement. J. G. thanks the Universidad del Valle for support during his master's studies. C.A.G. was supported in part by Project 71327 (Universidad del Valle).

References

- C. Cooper and F. T. Howard, Some identities for r-Fibonacci numbers, The Fibonacci Quarterly 49 (2011) 231–243.
- [2] A. Dubickas, On then distance between two algebraic numbers, *Bulletin* of the Malaysian Mathematical Science Society **43** (2020), 3049–3064.
- [3] D. E. Ferguson, An expression for generalized Fibonacci numbers, *The Fibonacci Quarterly* 4 (1966), 270–273.
- [4] H. Gabai, Generalized Fibonacci k-sequences, The Fibonacci Quarterly 8 (1970), 31–38.
- [5] J. García, C. A. Gómez, and F. Luca, On the zero-multiplicity of the k-generalized Fibonacci sequence, Journal of Difference Equations and Applications 26 (2020), 1564–1578.
- [6] J. García, C. A. Gómez and F. Luca, On the separation of the roots of the generalized Fibonacci polynomial, arXiv:2210.16717v1 (2022), 1–15.
- [7] T. R. Hagedorn, Zeros of a real linear recurrence of degree $n \ge 4$, C. R. Math. Rep. Acad. Sci. Canada **27** (2005), 41–47.
- [8] T. Koshy, Fibonacci and Lucas Numbers with Applications, John Wiley & Sons, Inc., New York, 2001.
- [9] E. Kummer, Uber die Erganzungssatze zu den allgemeinen Reciprocitatsgesetzen, J. Reine Angew. Math. 44 (1852), 93–146.
- [10] T.N. Shorey and R. Tijdeman, Exponential Diophantine equations, Cambridge University Press, 1986; reprinted 2008.
- [11] T. Skolem, Ein Verfahren zur Behandlung gewisser exponentialer Gleichungen und diophantischer Gleichungen, Proceedings of the 8th Congress of Scandinavian Mathematicians, Stockholm (1934), 163–188.
- [12] P. T. Young, On the 2-adic valuation of generalized Fibonacci sequences, Integers 22 (2022), Article 68.