Electroencephalography and mild cognitive impairment research: A scoping review and bibliometric analysis (ScoRBA)

Short running title: EEG and mild cognitive impairment

Adi Wijaya^{*}, Noor Akhmad Setiawan⁺, Asma Hayati Ahmad[‡], Rahimah Zakaria[‡], Zahiruddin Othman[¶]

^{*}Department of Health Information Management, Universitas Indonesia Maju, Jakarta, Indonesia, , +Department of Electrical and Information Engineering, Universitas Gadjah Mada, Yogyakarta, Indonesia, ‡Department of Physiology, School of Medical Sciences, Universiti Sains Malaysia, Kubang Kerian 16150, Kelantan Malaysia, , ¶Department of Psychiatry, School of Medical Sciences, Universiti Sains Malaysia, Kubang Kerian 16150, Kelantan, Malaysia

Correspondence: Zahiruddin Othman, Department of Psychiatry, School of Medical Sciences, Universiti Sains Malaysia, Kubang Kerian 16150, Kelantan, Malaysia. E-mail: zahirkb@usm.my

Conflict of interest

The authors declare no conflict of interest.

Source of funding

This research did not receive any specific funding.

ABSTRACT

Background: Mild cognitive impairment (MCI) is often considered a precursor to Alzheimer's disease (AD) due to the high rate of progression from MCI to AD. Sensitive neural biomarkers may provide a tool for an accurate MCI diagnosis, enabling earlier and perhaps more effective treatment. Despite the availability of numerous neuroscience techniques, electroencephalography (EEG) is the most popular and frequently used tool among researchers due to its low cost and superior temporal resolution.

Objective: We conducted a scoping review of EEG and MCI between 2012 and 2022 to track the progression of research in this field.

Methods: In contrast to previous scoping reviews, the data charting was aided by co-occurrence analysis using VOSviewer, while data reporting adopted a Patterns, Advances, Gaps, Evidence of Practice, and Research Recommendations (PAGER) framework to increase the quality of the results.

Results: Event-related potentials (ERPs) and EEG, epilepsy, quantitative EEG (QEEG), and EEG-based machine learning were the research themes addressed by 2310 peer-reviewed articles on EEG and MCI.

Conclusion: Our review identified the main research themes in EEG and MCI with highaccuracy detection of seizure and MCI performed using ERP/EEG, QEEG and EEG-based machine learning frameworks.

Keywords: electroencephalography; mild cognitive impairment; VOSviewer; scoping review; bibliometric analysis

Key messages:

- Health librarians and researchers may conduct a data charting for scoping review using co-occurrence analysis.
- Health librarians may engage various review methods to promote more research and publications related to health information services.
- Researchers are encouraged to conduct further research based on the research themes identified in this study.

Introduction

The most common type of neurodegenerative dementia is Alzheimer's disease (AD), with ten million new dementia cases identified each year (World Health Organization [WHO], 2020). AD is the most common cause of dementia accounting for 60–70% of these new cases, followed by vascular dementia (VaD), dementia with Lewy bodies (DLB), and other forms of neurodegenerative illnesses (WHO, 2020).

Prior to development of dementia, two stages of decreased cognition are often defined: subjective cognitive decline (SCD), which is not corroborated by an informant or neuropsychological testing, and mild cognitive impairment (MCI), which is corroborated by an informant or neuropsychological testing (Albert et al., 2011; American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders [DSM], 2013; Jessen et al., 2014; Winblad et al., 2004; WHO, 1992). In both stages, individuals are able to undertake personal and instrumental daily activities (Gao et al., 2018; Mauri, Sinforiani, Zucchella, Cuzzoni, & Bono, 2012; Pandya, Lacritz, Weiner, Deschner, & Woon, 2017; Visser, Kester, Jolles, & Verhey, 2006). According to the Alzheimer Association (2020), 15% of MCI in individuals over 65 progress to dementia, but in a different study (Chen et al., 2020), 32% of individuals with MCI were found to have an AD at the 5-year follow-up. Older age, poor cognition, APOE 4 allele carrier status, and hypertension could enhance dementia risk (Roberts & Knopman, 2013). Thus, it is important to establish biomarkers that can identify those at high risk for dementia and, ideally, its cause.

Studies combining cognitive assessments with cerebrospinal fluid (CSF) proteins, Magnetic Resonance Imaging (MRI), and Fluorodeoxyglucose-Positron Emission Tomography (FDG-PET) have shown outstanding results in detecting MCI patients who will later develop dementia (Ottoy et al., 2019; Smailagic, Lafortune, Kelly, Hyde, & Brayne, 2018; Zhou et al., 2019). However, these biomarkers are time-consuming, expensive, and some are invasive, therefore not suitable for daily clinical practice, except in subspecialized settings. Electroencephalography (EEG) is a low-cost, noninvasive, and straightforward method that can be performed in most clinics and even in primary care settings, especially with automatic reading. In the recent decade, many studies have published the EEG's ability to diagnose MCI patients who will develop dementia (Bonanni et al., 2015; Cassani, Estarellas, San-Martin, Fraga, & Falk, 2018; Moretti, 2015a&b; Poil et al., 2013). Therefore, this study aims to perform a scoping review (Arksey & O'Malley, 2005) aided by bibliometric analysis (ScoRBA) on EEG and MCI research, and utilise the Patterns, Advances, Gaps, Evidence of Practice, and Research

Recommendations (PAGER) framework (Bradbury-Jones, Isham, Morris, & Taylor, 2019) to present our data.

Methods

Literature search and study selection

The scoping review's five steps were adapted from Arksey and O'Malley's (2005) methodology (Table 1). Step 1, which was to identify the primary research question, was deliberated in the introductory section. On September 20, 2022, a literature search using the Scopus database was carried out (step 2). We decided to use the Scopus database because it covers a large number of publications (Pranckute, 2021; Zhu & Liu, 2020). The following search phrases: "TITLE-ABS-KEY (EEG OR electroencephalogra*)" AND TITLE-ABS-KEY ("neurocognitive disorder" or "cognitive impairment" or "cognitive disorder" or "cognitive disability" OR MCI) were used. We only included journal articles that were published in the English language between 2012 and 2022 (step 3). The PRISMA flow diagram in Figure 1 exhibits the search procedure.

Step	Description		
1. Identification of Research Questions and	The research question in this paper is "What		
Related Studies	are the main themes in the EEG and mild		
	cognitive impairment research?"		
2. Identification of Related Studies	A preliminary reading of relevant studies		
	was performed to determine the keywords		
	and terms utilised in the article selection		
3. Study Selection	Peer-reviewed original articles published in		
	the Scopus-indexed journal that were written		
	in English were selected. Articles that		
	assessed the usage of EEG in patients with		
	MCI met the selection criteria of this study.		
4. Charting the data	The co-occurrence keyword was used to		
	group the data into a graphical		
	representation.		

Table 1 Five-step scoping review framework

5. Collating, summarizing, and reporting the results
5. Collating, summarizing, and reporting the results
Practice, and Research Recommendations (PAGER) framework was used to summarize, analyze, and present the results in order to enhance the quality and applicability of the literature review, as

suggested by Bradbury et al. (2019)




Figure 1 PRISMA flow diagram of the search strategy (Page et al., 2021)

Results

The initial search results from the Scopus database generated 3305 documents published from 2012 until September 2022 (Figure 1). After removing papers in languages other than English,

document types other than original articles (primary research documents), and source types other than journals, 2310 documents remained.

Publication output

The number of articles showed an increasing trend from 2012 to 2021. There was a drop in the number of articles in 2022 due to incomplete data (Figure 2).




Figure 2 The number of annual publications on EEG and MCI research published from 2012 to September 2022 in the Scopus database

Research output on EEG and MCI research covered 24 different subject areas. Medicine, neuroscience, biochemistry, genetics and molecular biology, and psychology were the four subject areas that made up the most articles (3521/4172, or 84.40% of the total). These were followed by computer science, engineering, pharmacology, toxicology and pharmaceutics, multidisciplinary, nursing, health professions, etc. Figure 3 lists the top 20 EEG and MCI research areas currently under investigation.





Figure 3 Top 20 subjects in the area of EEG and MCI research

Data Charting

Previous method of data charting using the 'narrative review' approach (Pawson, 2002) was replaced with utilizing the results from the co-occurrence keyword analysis. All the 2310 documents were downloaded from the Scopus database in Comma-Separated Values (.csv) format and analysed using VOSviewer version 1.6.17 (van Eck & Waltman, 2021) for co-occurrence keyword analysis. A total of 5002 author keywords were collected from our collection of 2310 publications. Only keywords that appeared at least five times were selected for keyword analysis. This criterion was met by 229 keywords out of 5002. Our analysis indicates that the keyword network has four distinct clusters as shown in Figure 4: cluster one (red, 80 keywords), cluster two (green, 56 keywords), cluster three (blue, 47 keywords) and cluster four (yellow, 46 keywords).



A VOSviewer

Figure 4 Co-occurrence author keywords. 229 out of 5002 keywords met the threshold of a minimum five occurrence

Based on the dominant keywords in each cluster, research on the usage of EEG in mild cognitive impairment patients can be divided into four broad clusters and themes: (1) red cluster – event-related potentials (ERPs) and EEG, (2) green cluster - epilepsy, (3) blue cluster – quantitative EEG, and (4) yellow cluster – EEG and machine learning.

DISCUSSION

The PAGER framework was utilised as the final step of our scoping assessment of the usage of EEG in mild cognitive impairment research (Bradbury-Jones et al., 2019). Table 2 summarizes the key findings of the PAGER framework literature analysis, including model themes, research advances, research gaps, evidence of practice, and research recommendations.

Patterns	Advances	Gaps	Evidence for practice	Research recommendation
ERP/EEG	Discussion on the combined	There is a need to understand	Evidence from future	Future studies to establish
(Red cluster)	EEG/ERP measures of	the causal effects between	research on EEG/ERP as	ERPs' sensitivity and
	memory as a diagnostic tool	$A\beta$ depositions and neural	biomarkers for assessing	specificity in discriminating
	for detecting MCI or	excitability.	early memory decline and	AD, MCI and cognitively
	prodromal AD.		treatment response.	intact AD-risk groups as well
				as measuring late cognitive
				ERPs during complicated
				tasks, especially in healthy
				older adults at risk for
				cognitive decline.
Epilepsy	Knowledge discovery from	There is a need to select	EEG is currently used to	Future studies to explore the
(Green cluster)	machine learning classifiers	machine learning classifiers	detect seizures and	associations between
	such as seizure localization,	and features for high-	categorise seizure types and	cognitive deficit and seizure
	which exactly points to	accuracy seizure detection	epilepsy disorders.	timing.
	affected brain lobe(s), channel	and predict cognitive		
	importance, and based on	impairment.		
	participating channels in a			
	seizure may help to predict			
	cognitive impairment			

Table 2 Results of the PAGER analysis of the EEG and mild cognitive impairment research

secondary to seizures, EEG interictal spikes (IIS), and antiepileptic medicines.

OEEG Development of simple There is a need to find clear- Evidence future Future large-scale and from neuropsychiatry diagnostic algorithms using cut EEG signs and research on QEEG as a longitudinal clinical (Blue cluster) spectral analysis of EEG to correlations with diagnostic and prognostic investigations are required to detect MCI cases. neuropathology and biomarker as well as for determine the diagnostic and neuropsychology for monitoring treatment prognostic potential of precise response in MCI cases. OEEG measurements consistent and as early functional markers of assessments. AD on a subject-by-subject

basis.

EEG Machine learning There is a need to select Evidence for from and extracting relevant machine learning classifiers research to apply EEG-based EEG-based machine the learning EEG/ERP data for cognitive and features for a high- machine (Yellow cluster) assessment. cognitive frameworks for accurate extracting the most relevant accuracy diagnosis of MCI cases. assessment.

future Future studies to utilize an machine learning learning framework for features from EEG data empirically thereby resulting in a high accuracy of cognitive assessment.

EEG/ERPs and mild cognitive impairment

The first report on using scalp EEG was in 1929 (Berger, 1929). EEG is a cost-effective and non-invasive technology that directly assesses the mean electrical activity of the brain at scalp sites with excellent temporal precision (milliseconds) (Cohen, 2017). ERPs are one of the most extensively used EEG techniques to examine brain activity in response to sensory, motor, or cognitive events. Several ERP components (e.g., N1, P2, and P3) can provide full information about low or high level cognitive activities of the brain (Luck, 2014) and have been used in research for several decades.

Three EEG measures that have been found to strongly correlate with verbal learning and memory abilities in healthy elderly people and patients with MCI or prodromal Alzheimer's disease (AD) in previous research include ERP P600 (Olichney et al., 2002, 2006 & 2013), suppression of oscillatory activity in the alpha frequency range (Fell, Ludowig, Rosburg, Axmacher, & Elger, 2008; Fellner, Bauml, & Hanslmayr, 2013; Hanslmayr, Spitzer, & Bauml, 2009; Sederberg et al., 2006; Vassileiou, Meyer, Beese, & Friederici, 2018), and crossfrequency coupling between low theta/high delta and alpha/beta activity. A study by Xia and colleagues (2020) evaluated the relationships among three previously identified electrophysiological measures: the P600 ERP and two oscillatory effects (i.e. alpha suppression and $\phi/\delta - \alpha/\beta$ coupling), and their potential as biomarkers of verbal memory functioning in healthy ageing, MCI and prodromal Alzheimer's disease. The study highlighted the importance of combining ERP and EEG oscillatory measures as predictors of verbal memory in MCI individuals. Additionally, a recent study by Meghdadi et al. (2021) demonstrated that subtle resting state EEG abnormalities could differentiate between MCI and AD groups. The study designed classifiers for differentiating individual MCI and AD participants from agematched controls and concluded that the MCI group demonstrated a moderate increase in δ , ϕ , and Theta-to-Alpha (TAR) ratio, mostly in temporal channels (Meghdadi et al., 2021).

A systematic literature review investigating ERPs as potential biomarkers of AD-related neuropathology was published recently (Paitel, Samii, & Nielson, 2021). The patterns in cognitive ERPs (\geq 150 ms post-stimulus) differentiate MCI, AD, and cognitively intact elders who carry AD risk through the Apolipoprotein-E ϵ 4 allele (ϵ 4+), from healthy older adult controls. The review also implies that the integration of ERPs into cognitive assessment may significantly improve early identification and characterization of brain dysfunction, allowing for faster differential diagnosis and intervention targeting (Paitel et al., 2021). Future studies are required to establish ERPs' sensitivity and specificity in discriminating AD, MCI and cognitively intact AD-risk groups as well as measuring late cognitive ERPs during complicated tasks, especially in healthy older adults at risk for cognitive decline.

In a more recent study, Devos et al. (2022) found that neuronal hyperexcitability is related to higher amyloid levels in cognitively healthy older persons. The smaller P3 ERP task difference suggests a less efficient stimulus processing in cognitively healthy older persons with elevated amyloid levels compared to those with non-elevated amyloid levels. These results, together with the lack of statistically significant differences in behavioural outcomes, imply that the enhanced A β deposition-induced hyperexcitability is a non-functional neuronal compensation that may later impair working memory. There is a need to understand the causal effects of A β depositions and neural excitability to confirm this observation.

Epilepsy and mild cognitive impairment

Epileptic seizures are brief bursts of excessively synchronised brain activity (Chang & Lowenstein, 2003; Fisher, Scharfman, & DeCurtis, 2014; Noachtar et al., 1999; Shorvon, 2011; Trevelyan, 2016; Wang et al., 2017). They cover a wide range of occurrences, from mild clinical manifestations, such as brief and hardly detectable loss of consciousness, to vigorous episodes of muscular shaking that may cause physical harm (Landi, Petrucco, Sicca, & Ratto, 2019). Epilepsy is diagnosed based on EEG abnormalities, including seizures and interictal epileptiform discharges (IEDs). EEG findings help to categorise seizure types and epilepsy disorders (Chen & Koubeissi, 2019).

EEG analysis of patients with recurrent seizures shows "subclinical" aberrant electrical activity between seizures. IEDs are classified based on their EEG signature: sharp waves, spikes, sharp waves/spikes-and-slow waves, and numerous spikes-and-slow waves (de Curtis & Avanzini, 2001; Fisher et al., 2014; Gotman, 1980; Kane et al., 2017; Kooi, 1966; Noachtar et al., 1999; Pillai & Sperling, 2006). However, background noise and aberrations such as eye blinks and muscle movements can contaminate EEG readings, causing electrical interference that is difficult to detect visually in longer recordings. Thus, there is a need to select machine learning classifiers and features for high-accuracy seizure detection and predict cognitive impairment (Natu, Bachute, Gite, Kotecha, & Vidyarthi, 2022).

Although seizures are the most obvious clinical manifestation of epilepsies, people with epilepsy are at risk of other health problems at a higher incidence rate than would be expected by chance (Institute of Medicine [IOM], 2012). Common co-morbidities in epilepsy include cognitive impairment such as memory, attention, and processing problems, mental health

problems such as depression and anxiety, and somatic co-morbidities such as sleep disorders and migraines (Holmes, 2015).

Most of the cognitive impairment in epilepsy is due to its aetiology. Trauma, hypoxia, ischemia, and mesial temporal sclerosis secondary to prolonged seizures are acquired illnesses that can cause epilepsy and cognitive impairment. Genetic disorders including tuberous sclerosis, Fragile X, Rett, and Dravet syndromes can also lead to epilepsy and cognitive impairment. In addition to static abnormalities induced by the underlying aetiology, cognitive impairments might be temporary due to seizures, EEG interictal spikes (IIS), and antiepileptic medicines. In many people, cognitive impairment is due to a combination of causes (Holmes, 2015).

The age of seizure onset is another factor that is associated with cognitive impairment. When compared to healthy peers, children with epilepsy are more likely to have severe cognitive impairments, which primarily affect language, semantic, motor, and visual functions (Al-Malt, Abo Hammar, Rashed, & Ragab, 2020). According to the same study, children with idiopathic epilepsy who have more nocturnal seizures than diurnal seizures have both overt and subtle cognitive abnormalities (Malt et al., 2020). Future studies are needed to explore the associations between cognitive deficit and seizure timing.

Quantitative EEG and mild cognitive impairment

The diagnosis of cognitive impairment and neurodegenerative illnesses using EEG relies heavily on the EEG expert's visual assessment. The limitations that come with this have been supplemented by methods such as quantitative EEG (QEEG) that extracts particular parameters from continuous variable data analysis. At current, studies on brain function assessment and early diagnosis of degenerative brain illnesses such as AD and AD-MCI are actively advancing (Al-Qazzaz et al., 2014; Cassani et al., 2018; Smailovic & Jelic, 2019). For instance, QEEG measurements appear to differentiate between different forms of dementia and correlate favourably with surrogate markers of AD neuropathology, making them promising low-cost and noninvasive diagnostics of AD (Smailovic & Jelic, 2018). The diagnostic and prognostic value of QEEG measurements as early functional markers of AD should be determined in future extensive longitudinal clinical studies on a subject-by-subject basis.

One technique for analysing QEEG signals is using the power spectrum, and this approach is a well-established core component of the general method utilised in clinical research. The most frequent spectral shift so far is thought to be connected to the degree of cognitive deterioration brought on by the rise in the slow wave of the EEG (Czigler et al., 2008; Engedal et al., 2020; Han & Youn, 2022; Jeong, Youn, Sung, & Kim, 2021; Ya et al., 2015). Spectral analysis of the EEG is used to develop simple diagnostic algorithms that can identify MCI. However, finding distinct EEG signals with strong neuropathology and neuropsychology correlations are necessary for reliable and accurate assessments.

In a broad frequency range, spectral- and EEG-complexity alterations can be detected even in the early stages of AD. However, applying traditional EEG analysis techniques along with QEEG may further increase the likelihood of making an early diagnosis of AD (Czigler et al., 2008; Ya et al., 2015). A recent study by Engedal et al. (2020) concluded that applying QEEG using the statistical pattern recognition (SPR) method could predict conversion to dementia in patients with SCD and MCI. The authors also suggested adding this method to other routine diagnostic methods like cognitive tests and other biomarkers to benefit from the discriminant power of the proposed method. A study by Jeong et al. (2021) demonstrated that SCD participants had more frontal delta waves and less occipital alpha1 compared to control participants. SCD is one of AD's early clinical signs and is linked to nerve degeneration. Hamilton et al. (2021) demonstrated that QEEG slowing may precede the onset of dementia in prospectively identified MCI. In addition, another study by Han and Youn (2022) used QEEG for monitoring treatment response in MCI patients. Future studies should utilise QEEG characteristics, neuroimaging, and/or machine learning to better collect and show neural network activation.

EEG-based machine learning and mild cognitive impairment

Machine learning is a set of algorithms that enables us to automatically detect patterns in the data and predict the results of measurements (Bishop, 2006; Hastie, Tibshirani, & Friedman, 2009; Murphy, 2013). It is frequently used in neural engineering and neuroscience experiments to: (1) compare different circumstances (Lemm, Schafer, & Curio, 2004; Müller et al., 2008), (2) make a diagnosis of a disease (Dauwan et al., 2018; Kumar, Dewal, & Anand, 2014), or (3) find electrophysiological alterations connected to behaviour (Buriro, Shoorangiz, Weddell, & Jones, 2018; Davidson, Jones, & Peiris, 2007). Machine learning algorithms have been developed to extract information from the EEG in order to identify different brain states and aid in the diagnosis of several conditions (such as epilepsy, AD, and schizophrenia) (Shoorangiz, Weddell, & Jones, 2021).

Several studies have described EEG-based machine learning frameworks for the diagnosis of AD, MCI, and other types of dementia. Mcbride et al. (2014) established an EEG-based

classification system using spectrum and complexity analysis. The complexity analysis includes computational activity, mobility, complexity, sample entropy, and lempel-Ziv characteristics, while the spectral analysis involves extracting features from delta, theta, alpha1, alpha2, gamma, beta1, and beta2 subbands. The support vector machine (SVM) classifier was found to have an average accuracy of 79.2% for MCI classification. Kashefpoor, Rabbani, and Barekatain (2016) extracted EEG spectral parameters using delta, theta, alpha1, alpha2, gamma, beta1, and beta2 subbands, as well as 19 spectral characteristics from a 19-channel EEG data. A correlation-based algorithm identified the best discriminative characteristics. The best classification accuracy was 88.89% utilising neuro-fuzzy (NF) and k-nearest neighbour (KNN) classifiers, (Kashefpoor et al., 2016). Kashefpoor, Rabbani, and Barekatain (2019), in a different study, employed a 19-channel EEG data and supervised dictionary learning techniques like label-consistent K-SVD (LC-KSVD) and correlation-based label-consistent K-SVD (CLC-KSVD) to diagnose MCI and achieved the best accuracy at 88.9%.

Khatun, Morshed, and Bidelman (2019) developed a single-channel EEG method for diagnosing MCI using speech-evoked brain responses. The top 25 characteristics were picked using the random forest (RF) method after temporal and spectral domain analysis extracted 590 from recorded sounds. The best classification result they found was 87.9% accuracy using logistic regression (LR) and SVM models. Yin, Cao, Siuly, and Wang (2019) also suggested a spectral-temporal analysis-based MCI diagnosis technique. This method employs spectral-temporal analysis to extract features and the 3-D evaluation algorithm to produce an optimum feature subset. In the study, the SVM classifier had 96.94% accuracy (Yin et al. 2019).

Sharma, Kolekar, Jha, & Kumar (2019) used power spectral density (PSD), skewness, kurtosis, spectral skewness, spectral kurtosis, spectral crest factor, spectral entropy (SE), and fractal dimension (FD) to develop an EEG-based MCI detection method. Open eyes, closed eyes, finger tapping test (FTT), and continuous performance test were the conditions used to collect the dataset. SVM's classification accuracy was 96.94% (Sharma et al., 2019). Later, Siuly et al. (2020) established an MCI patient-identification system using extreme learning machine (ELM), SVM, and KNN classification models. The ELM classifier achieved the best classification result, with 98.78% accuracy (Siuly et al., 2020).

The spectral-power-based task-induced intra-subject EEG variability extracted by the sequential forward selection (SFS) algorithm-based feature selection and SVM classification has been shown to have the potential to serve as a neurophysiological feature for the early detection of MCI in individuals (Trinh et al., 2021). Rossini, Miraglia, and Vecchio (2022), on the other hand, used a combination of graph analysis tools and SVM classification to identify the distinctive features of physiological/pathological brain ageing focusing on functional connectivity networks evaluated on EEG data and other biomarkers.

Hsiao et al. (2021) proposed a novel EEG feature, the kernel eigen-relative-power (KERP) feature, for achieving high classification accuracy of MCI (90.2%) when used in combination with an SVM classifier. Meanwhile, Oltu, Akşahin, and Kibaroğlu (2021) proposed an EEG-based categorization for MCI, AD, and healthy persons using discrete wavelet transform (DWT), PSD, and interhemispheric coherence for feature extraction, and Bagged Trees for classification. The study's best result was 96% (Oltu et al. 2021). Movahed and Rezaeian (2022) proposed an automatic diagnosis of MCI using 0-spectral, functional connectivity, and nonlinear EEG-based features. Ten-fold cross-validation was used to assess metrics including accuracy (AC), sensitivity (SE), specificity (SP), F1-score (F1), and false discovery rate (FDR). The best performance of the proposed framework employing the linear support vector machine (LSVM) classifier and the combination of all feature sets have been reported, with an average AC of 99.4%, SE of 98.8%, SP of 100%, F1 of 99.4%, and FDR of 0%. Based on the achieved results, it is possible to develop and use a computer-aided diagnostic (CAD) tool for clinical purposes.

Conclusion

This scoping review examined 2310 articles on EEG and MCI. The articles were divided into four broad themes, including ERP/EEG, epilepsy, QEEG, and EEG-based machine learning. According to numerous studies the ERP/EEG, QEEG and EEG-based machine learning framework can be widely used for high-accuracy seizure and MCI detector.

References

Albert, M. S., DeKosky, S. T., Dickson, D., Dubois, B., Feldman, H. H., Fox, N. C., ... Phelps, C. H. (2011). The diagnosis of mild cognitive impairment due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimer's & Dementia*, 7(3), 270–9. https://doi.org/10.1016/j.jalz.2011.03.008

Al-Malt, A. M., Abo Hammar, S. A., Rashed, K. H., & Ragab, O. A. (2020). The effect of nocturnal epileptic seizures on cognitive functions in children with idiopathic epilepsy. *The Egyptian Journal of Neurology, Psychiatry and Neurosurgery, 56*, 1-6.

Al-Qazzaz, N. K., Ali, S. H., Ahmad, S. A., Chellappan, K., Islam, M. S., & Escudero, J. (2014). Role of EEG as biomarker in the early detection and classification of dementia. *The Scientific World Journal*, *2014*, 906038. https://doi.org/10.1155/2014/906038

Alzheimer Association (2020). 2020 Alzheimer's disease facts and figures. *Alzheimer's & Dementia*, 16, 3. https://doi.org/10.1002/alz.12068

American Psychiatric Association. (2013). *Diagnostic and Statistical Manual of Mental Disorders*. 5th ed. Washington (DC): American Psychiatric Publishing.

Arksey, H., & O'Malley, L. (2005) Scoping studies: towards a methodological framework. *International Journal of Social Research Methodology*, 8(1), 19-32, https://doi.org/10.1080/1364557032000119616.

Berger, H. (1929). Über das elektrenkephalogramm des menschen. Archiv für Psychiatrie und Nervenkrankheiten, 87(1), 527–570. https://doi.org/10.1007/BF01797193

Bishop, C. M. (2006). Pattern recognition and machine learning. Springer: Berlin/Heidelberg.

Bonanni, L., Perfetti, B., Bifolchetti, S., Taylor, J. P., Franciotti, R., Parnetti, L., ... Onofrj, M. (2015). Quantitative electroencephalogram utility in predicting conversion of mild cognitive

impairment to dementia with Lewy bodies. *Neurobiology of Aging*, *36*(1), 434–445. https://doi.org/10.1016/j.neurobiolaging.2014.07.009

Bradbury-Jones, C., Isham, L., Morris, A. J., & Taylor, J. (2021). The "neglected" relationship between child maltreatment and oral health? An international scoping review of research. *Trauma, Violence & Abuse*, 22(2), 265–276. https://doi.org/10.1177/1524838019841598

Buriro, A. B., Shoorangiz, R., Weddell, S. J., & Jones, R. D. (2018). Predicting microsleepstates using EEG inter-channel relationships. *IEEE transactions on Neural Systems andRehabilitationEngineering*, 26(12),https://doi.org/10.1109/TNSRE.2018.2878587

Cassani, R., Estarellas, M., San-Martin, R., Fraga, F. J., & Falk, T. H. (2018). systematic review on resting-state EEG for Alzheimer's disease diagnosis and progression assessment. *Disease Markers*, 2018, 5174815. https://doi.org/10.1155/2018/5174815

Chang, B. S., & Lowenstein, D. H. (2003). Epilepsy. *The New England Journal of Medicine*, 349(13), 1257–1266. https://doi.org/10.1056/NEJMra022308

Chen, Y., Zhang, J., Zhang, T., Cao, L., You, Y., Zhang, C., ... Zhang, Q. (2020). Meditation treatment of Alzheimer disease and mild cognitive impairment: A protocol for systematic review. *Medicine*, *99*(10), e19313. https://doi.org/10.1097/MD.000000000019313

Chen, H., & Koubeissi, M. Z. (2019).Electroencephalography in EpilepsyEvaluation. Continuum(Minneapolis, Minn.), 25(2), 431–453.https://doi.org/10.1212/CON.0000000000000705

Cohen, M. X. (2017). Where does EEG come from and what does it mean?. *Trends in Neurosciences*, 40(4), 208–218. https://doi.org/10.1016/j.tins.2017.02.004

Institute of Medicine (IOM). (2012). *Epilepsy across the spectrum: promoting health and understanding*. Washington, DC: The National Academies Press.

Czigler, B., Csikós, D., Hidasi, Z., Anna Gaál, Z., Csibri, E., Kiss, E., ... Molnár, M. (2008). Quantitative EEG in early Alzheimer's disease patients - power spectrum and complexity features. *International Journal of Psychophysiology*, 68(1), 75–80. https://doi.org/10.1016/j.ijpsycho.2007.11.002

Dauwan, M., Linszen, M., Lemstra, A. W., Scheltens, P., Stam, C. J., & Sommer, I. E. (2018). EEG-based neurophysiological indicators of hallucinations in Alzheimer's disease: Comparison with dementia with Lewy bodies. *Neurobiology of Aging*, 67, 75–83. https://doi.org/10.1016/j.neurobiolaging.2018.03.013

Davidson, P. R., Jones, R. D., & Peiris, M. T. (2007). EEG-based lapse detection with high temporal resolution. *IEEE Transactions on Bio-Medical Engineering*, *54*(5), 832–839. https://doi.org/10.1109/TBME.2007.893452

de Curtis, M., & Avanzini, G. (2001). Interictal spikes in focal epileptogenesis. *Progress in Neurobiology*, 63(5), 541–567. https://doi.org/10.1016/s0301-0082(00)00026-5

Devos, H., Gustafson, K., Liao, K., Ahmadnezhad, P., Estes, B., Martin, L. E., ...Burns, J. M. (2022). EEG/ERP evidence of possible hyperexcitability in older adults with elevated betaamyloid. *Translational Neurodegeneration*, *11*(1), 8. https://doi.org/10.1186/s40035-022-00282-5

Engedal, K., Barca, M. L., Høgh, P., Bo Andersen, B., Winther Dombernowsky, N., Naik, M., ... Snaedal, J. (2020). The Power of EEG to Predict Conversion from Mild Cognitive Impairment and Subjective Cognitive Decline to Dementia. *Dementia and Geriatric Cognitive Disorders*, 49(1), 38–47. https://doi.org/10.1159/000508392

Fell, J., Ludowig, E., Rosburg, T., Axmacher, N., & Elger, C. E. (2008). Phase-locking within human mediotemporal lobe predicts memory formation. *NeuroImage*, *43*(2), 410–419. https://doi.org/10.1016/j.neuroimage.2008.07.021

Fellner, M. C., Bäuml, K. H., & Hanslmayr, S. (2013). Brain oscillatory subsequent memory effects differ in power and long-range synchronization between semantic and survival processing. *NeuroImage*, *79*, 361–370. https://doi.org/10.1016/j.neuroimage.2013.04.121

Fisher, R. S., Scharfman, H. E., & deCurtis, M. (2014). How can we identify ictal and interictal abnormal activity? *Advances in Experimental Medicine and Biology*, *813*, 3–23. https://doi.org/10.1007/978-94-017-8914-1_1

Gao, Q., Gwee, X., Feng, L., Nyunt, M., Feng, L., Collinson, S. L., Chong, M. S., Lim, W. S., Lee, T. S., Yap, P., Yap, K. B., & Ng, T. P. (2018). Mild cognitive impairment reversion and progression: rates and predictors in community-living older persons in the Singapore longitudinal ageing studies cohort. *Dementia and Geriatric Cognitive Disorders Extra*, 8(2), 226–237. https://doi.org/10.1159/000488936

Gotman J. (1980). Quantitative measurements of epileptic spike morphology in the human EEG. *Electroencephalography and Clinical Neurophysiology*, *48*(5), 551–557. https://doi.org/10.1016/0013-4694(80)90290-4

Hamilton, C. A., Schumacher, J., Matthews, F., Taylor, J. P., Allan, L., Barnett, N., ... Thomas,
A. J. (2021). Slowing on quantitative EEG is associated with transition to dementia in mild
cognitive impairment. *International Psychogeriatrics*, 33(12), 1321–1325.
https://doi.org/10.1017/S1041610221001083

Han, S. H., & Chul Youn, Y. (2022). Quantitative electroencephalography changes in patients with mild cognitive impairment after choline alphoscerate administration. *Journal of Clinical Neuroscience*, *102*, 42–48. https://doi.org/10.1016/j.jocn.2022.06.006

Hanslmayr, S., Spitzer, B., & Bäuml, K. H. (2009). Brain oscillations dissociate between semantic and nonsemantic encoding of episodic memories. *Cerebral cortex (New York, N.Y.: 1991)*, *19*(7), 1631–1640. https://doi.org/10.1093/cercor/bhn197

Hastie, T., Tibshirani, R., & Friedman, J. (2009). *The elements of statistical learning: data mining, inference, and prediction.* Springer Series in Statistics, 2nd ed. Springer: New York.

Holmes G. L. (2015). Cognitive impairment in epilepsy: the role of network abnormalities. *Epileptic Disorders*, *17*(2), 101–116. https://doi.org/10.1684/epd.2015.0739

Hsiao, Y. -T., Tsai, C. -F., Wu, C. -T., Trinh, T. -T., Lee, C. -Y., & Liu, Y. -H. (2021). MCI detection using kernel Eigen-relative-power features of EEG Signals. *Actuators*, *10*(7), 152. https://doi.org/10.3390/act10070152

Jeong, H. T., Youn, Y. C., Sung, H. H., & Kim, S. Y. (2021). Power spectral changes of quantitative EEG in the subjective cognitive decline: comparison of community normal control groups. *Neuropsychiatric Disease and Treatment*, *17*, 2783–2790. https://doi.org/10.2147/NDT.S320130

Jessen, F., Amariglio, R. E., van Boxtel, M., Breteler, M., Ceccaldi, M., Chételat, G., ... Subjective Cognitive Decline Initiative (SCD-I) Working Group (2014). A conceptual framework for research on subjective cognitive decline in preclinical Alzheimer's disease. *Alzheimer's & Dementia*, *10*(6), 844–852. https://doi.org/10.1016/j.jalz.2014.01.001

Kane, N., Acharya, J., Benickzy, S., Caboclo, L., Finnigan, S., Kaplan, P. W., ...van Putten,
M. (2017). A revised glossary of terms most commonly used by clinical electroencephalographers and updated proposal for the report format of the EEG findings.
Revision 2017. *Clinical Neurophysiology Practice*, 2, 170–185. https://doi.org/10.1016/j.cnp.2017.07.002

Kashefpoor, M., Rabbani, H., & Barekatain, M. (2016). Automatic Diagnosis of Mild Cognitive Impairment Using Electroencephalogram Spectral Features. *Journal of Medical Signals and Sensors*, 6(1), 25–32.

Kashefpoor, M., Rabbani, H., & Barekatain, M. (2019). Supervised dictionary learning of EEG signals for mild cognitive impairment diagnosis. *Biomedical Signal Processing and Control*, *53*, 2019, 101559. https://doi.org/10.1016/j.bspc.2019.101559

Khatun, S., Morshed, B. I., & Bidelman, G. M. (2019). A Single-Channel EEG-Based Approach to Detect Mild Cognitive Impairment via Speech-Evoked Brain Responses. *IEEE Transactions on Neural Systems and Rehabilitation Engineering*, 27(5), 1063–1070. https://doi.org/10.1109/TNSRE.2019.2911970 Kooi K. A. (1966). Voltage-time characteristics of spikes and other rapid electroencephalographic transients: semantic and morphological considerations. *Neurology*, *16*(1), 59–66. https://doi.org/10.1212/wnl.16.1.59

Kumar, Y., Dewal, M. L., & Anand, R. S. (2014). Epileptic seizure detection using DWT based fuzzy approximate entropy and support vector machine. *Neurocomputing*, *133*, 271–279 https://Doi.Org/10.1016/J.Neucom.2013.11.009

Landi, S., Petrucco, L., Sicca, F., & Ratto, G. M. (2019). Transient cognitive impairment in epilepsy. *Frontiers* in *Molecular Neuroscience*, *11*, 458. https://doi.org/10.3389/fnmol.2018.00458

Lemm, S., Schäfer, C., & Curio, G. (2004). BCI Competition 2003--Data set III: probabilistic modeling of sensorimotor mu rhythms for classification of imaginary hand movements. *IEEE Transactions on Bio-Medical Engineering*, *51*(6), 1077–1080. https://doi.org/10.1109/TBME.2004.827076

Luck, S. J. (2014). *An Introduction to the Event-Related Potential Technique*. Cambridge, MA: MIT Press.

Mauri, M., Sinforiani, E., Zucchella, C., Cuzzoni, M. G., & Bono, G. (2012). Progression to dementia in a population with amnestic mild cognitive impairment: clinical variables associated with conversion. *Functional Neurology*, *27*(1), 49–54.

McBride, J. C., Zhao, X., Munro, N. B., Smith, C. D., Jicha, G. A., Hively, L., ...Jiang, Y. (2014). Spectral and complexity analysis of scalp EEG characteristics for mild cognitive impairment and early Alzheimer's disease. *Computer Methods and Programs in Biomedicine*, *114*(2), 153–163. https://doi.org/10.1016/j.cmpb.2014.01.019

Meghdadi, A. H., Stevanović Karić, M., McConnell, M., Rupp, G., Richard, C., Hamilton, J., ... Berka, C. (2021). Resting state EEG biomarkers of cognitive decline associated with Alzheimer's disease and mild cognitive impairment. *PloS One*, *16*(2), e0244180. https://doi.org/10.1371/journal.pone.0244180 Moretti, D. V. (2015b). Association of EEG, MRI, and regional blood flow biomarkers is predictive of prodromal Alzheimer's disease. *Neuropsychiatric Disease and Treatment*, *11*, 2779–2791. https://doi.org/10.2147/NDT.S93253

Moretti, D. V. (2015a). Conversion of mild cognitive impairment patients in Alzheimer's disease: prognostic value of Alpha3/Alpha2 electroencephalographic rhythms power ratio. *Alzheimer's Research & Therapy*, 7, 80. https://doi.org/10.1186/s13195-015-0162-x

Movahed, R. A., & Rezaeian, M. (2022). Automatic diagnosis of mild cognitive impairment based on spectral, functional connectivity, and nonlinear EEG-based features. *Computational and Mathematical Methods in Medicine*, 2022, 2014001. https://doi.org/10.1155/2022/2014001

Müller, K. R., Tangermann, M., Dornhege, G., Krauledat, M., Curio, G., & Blankertz, B. (2008). Machine learning for real-time single-trial EEG-analysis: from brain-computer interfacing to mental state monitoring. *Journal of Neuroscience Methods*, *167*(1), 82–90. https://doi.org/10.1016/j.jneumeth.2007.09.022

Murphy, K. P. (2013). Machine learning: A probabilistic perspective. Cambridge; MIT Press.

Sharma, N., Kolekar, M. H., Jha, K., & Kumar, Y. (2019). EEG and cognitive biomarkers based mild cognitive impairment diagnosis. *IRBM*, 40(2), 113-121. https://doi.org/10.1016/j.irbm.2018.11.007

Natu, M., Bachute, M., Gite, S., Kotecha, K., & Vidyarthi, A. (2022). Review on epileptic seizure prediction: Machine learning and deep learning approaches. *Computational and Mathematical Methods in Medicine*, 2022, 7751263. https://doi.org/10.1155/2022/7751263

Noachtar, S., Binnie, C., Ebersole, J., Mauguière, F., Sakamoto, A., & Westmoreland, B. (1999). A glossary of terms most commonly used by clinical electroencephalographers and proposal for the report form for the EEG findings. The International Federation of Clinical Neurophysiology. *Electroencephalography and Clinical Neurophysiology. Supplement*, *52*, 21–41.

Olichney, J. M., Iragui, V. J., Salmon, D. P., Riggins, B. R., Morris, S. K., & Kutas, M. (2006).Absent event-related potential (ERP) word repetition effects in mild Alzheimer'sdisease. ClinicalNeurophysiology, 117(6),https://doi.org/10.1016/j.clinph.2006.02.022

Olichney, J. M., Morris, S. K., Ochoa, C., Salmon, D. P., Thal, L. J., Kutas, M., & Iragui, V. J. (2002). Abnormal verbal event related potentials in mild cognitive impairment and incipient Alzheimer's disease. *Journal of Neurology, Neurosurgery, and Psychiatry*, *73*(4), 377–384. https://doi.org/10.1136/jnnp.73.4.377

Olichney, J. M., Taylor, J. R., Gatherwright, J., Salmon, D. P., Bressler, A. J., Kutas, M., & Iragui-Madoz, V. J. (2008). Patients with MCI and N400 or P600 abnormalities are at very high risk for conversion to dementia. *Neurology*, *70*(19 Pt 2), 1763–1770. https://doi.org/10.1212/01.wnl.0000281689.28759.ab

Oltu, B., Akşahin, M. F., & Kibaroğlu, S. (2021). A novel electroencephalography based approach for Alzheimer's disease and mild cognitive impairment detection. *Biomedical Signal Processing and Control, 63*, 102223. https://doi.org/10.1016/j.bspc.2020.102223.

Ottoy, J., Niemantsverdriet, E., Verhaeghe, J., De Roeck, E., Struyfs, H., Somers, C., ...Staelens, S. (2019). Association of short-term cognitive decline and MCI-to-AD dementia conversion with CSF, MRI, amyloid- and ¹⁸F-FDG-PET imaging. *NeuroImage. Clinical*, 22, 101771. https://doi.org/10.1016/j.nicl.2019.101771

Page, M. J., McKenzie, J. E., Bossuyt, P. M., Boutron, I., Hoffmann, T. C., Mulrow, C. D., ... Moher, D. (2021). The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ (Clinical Research Ed.)*, *372*, n71. https://doi.org/10.1136/bmj.n71

Paitel, E. R., Samii, M. R., & Nielson, K. A. (2021). A systematic review of cognitive eventrelated potentials in mild cognitive impairment and Alzheimer's disease. *Behavioural Brain Research*, *396*, 112904. https://doi.org/10.1016/j.bbr.2020.112904 Pandya, S. Y., Lacritz, L. H., Weiner, M. F., Deschner, M., & Woon, F. L. (2017). Predictors of Reversion from Mild Cognitive Impairment to Normal Cognition. *Dementia and Geriatric Cognitive Disorders*, *43*(3-4), 204–214. https://doi.org/10.1159/000456070

Pawson, R. (2002). Evidence-based policy: In search of a method. *Evaluation*, 8(2), 157–181. https://doi.org/10.1177/1358902002008002512

Petersen R. C. (2011). Clinical practice. Mild cognitive impairment. *The New England Journal of Medicine*, *364*(23), 2227–2234. https://doi.org/10.1056/NEJMcp0910237

Pillai, J., & Sperling, M. R. (2006). Interictal EEG and the diagnosis of epilepsy. *Epilepsia*, 47 *Suppl 1*, 14–22. https://doi.org/10.1111/j.1528-1167.2006.00654.x

Poil, S. S., de Haan, W., van der Flier, W. M., Mansvelder, H. D., Scheltens, P., & Linkenkaer-Hansen, K. (2013). Integrative EEG biomarkers predict progression to Alzheimer's disease at the MCI stage. *Frontiers in Aging Neuroscience*, 5, 58. https://doi.org/10.3389/fnagi.2013.00058

Pranckute, R. (2021). Web of Science (WoS) and Scopus: The titans of bibliographic information in today's academic world. *Publications*, 9(1):12. https://doi.org/10.3390/publications9010012

Roberts, R., & Knopman, D. S. (2013). Classification and epidemiology of MCI. *Clinics in Geriatric Medicine*, 29(4), 753–772. https://doi.org/10.1016/j.cger.2013.07.003

Rossini, P. M., Miraglia, F., & Vecchio, F. (2022). Early dementia diagnosis, MCI-to-dementia risk prediction, and the role of machine learning methods for feature extraction from integrated biomarkers, in particular for EEG signal analysis. *Alzheimer's & Dementia*, 10.1002/alz.12645. Advance online publication. https://doi.org/10.1002/alz.12645

Sederberg, P. B., Schulze-Bonhage, A., Madsen, J. R., Bromfield, E. B., McCarthy, D. C., Brandt, A., ... Kahana, M. J. (2007). Hippocampal and neocortical gamma oscillations predict memory formation in humans. *Cerebral Cortex (New York, N.Y. : 1991)*, *17*(5), 1190–1196. https://doi.org/10.1093/cercor/bhl030 Sharma, N., Kolekar, M. H., Jha, K., & Kumar, Y. (2019). EEG and cognitive biomarkers based mild cognitive impairment diagnosis. *IRBM*, 40(2), 113-121. https://doi.org/10.1016/j.irbm.2018.11.007.

Shoorangiz, R., Weddell, S. J., & Jones, R. D. (2021). EEG-based machine learning: Theory and applications. In N. V. Thakor (Ed.), *Hand book of neuroengineering* (pp 1-39). Springer Nature: Singapore Pte Ltd. https://doi.org/10.1007/978-981-15-2848-4_70-1

Shorvon S. D. (2011). The etiologic classification of epilepsy. *Epilepsia*, *52*(6), 1052–1057. https://doi.org/10.1111/j.1528-1167.2011.03041.x

Siuly, S., Alcin, O. F., Kabir, E., Sengur, A., Wang, H., Zhang, Y., & Whittaker, F. (2020). A New Framework for Automatic Detection of Patients With Mild Cognitive Impairment Using Resting-State EEG Signals. *IEEE Transactions on Neural Systems and Rehabilitation Engineering*, 28(9), 1966–1976. https://doi.org/10.1109/TNSRE.2020.3013429

Smailagic, N., Lafortune, L., Kelly, S., Hyde, C., & Brayne, C. (2018). 18F-FDG PET for prediction of conversion to Alzheimer's disease dementia in people with mild cognitive impairment: An updated systematic review of test accuracy. *Journal of Alzheimer's Disease*, 64(4), 1175–1194. https://doi.org/10.3233/JAD-171125

Smailovic, U., Koenig, T., Kåreholt, I., Andersson, T., Kramberger, M. G., Winblad, B., & Jelic, V. (2018). Quantitative EEG power and synchronization correlate with Alzheimer's disease CSF biomarkers. *Neurobiology of Aging*, *63*, 88–95. https://doi.org/10.1016/j.neurobiolaging.2017.11.005

Smailovic, U., & Jelic, V. (2019). Neurophysiological Markers of Alzheimer's Disease: Quantitative EEG Approach. *Neurology and Therapy*, 8(Suppl 2), 37–55. https://doi.org/10.1007/s40120-019-00169-0

Trevelyan A. J. (2016). Do cortical circuits need protecting from themselves?. *Trends in Neurosciences*, *39*(8), 502–511. https://doi.org/10.1016/j.tins.2016.06.002

Trinh, T. T., Tsai, C. F., Hsiao, Y. T., Lee, C. Y., Wu, C. T., & Liu, Y. H. (2021). Identifying individuals with mild cognitive impairment using working memory-induced intra-subject variability of resting-state EEGs. *Frontiers in Computational Neuroscience*, *15*, 700467. https://doi.org/10.3389/fncom.2021.700467

van Eck, N. J., Waltman, L. (2021). *VOSviewer Manual: Manual for VOSviewer version 1.6.17*. Leiden: Centre for Science and Technology Studies (CWTS) of Leiden University. Retrieved from https://www.vosviewer.com/documentation/Manual_VOSviewer_1.6.17.pdf

Vassileiou, B., Meyer, L., Beese, C., & Friederici, A. D. (2018). Alignment of alpha-band desynchronization with syntactic structure predicts successful sentence comprehension. *NeuroImage*, *175*, 286–296. https://doi.org/10.1016/j.neuroimage.2018.04.008

Visser, P. J., Kester, A., Jolles, J., & Verhey, F. (2006). Ten-year risk of dementia in subjects with mild cognitive impairment. *Neurology*, *67*(7), 1201–1207. https://doi.org/10.1212/01.wnl.0000238517.59286.c5

Wang, B., Huang, L., Zhu, Y., Kundaje, A., Batzoglou, S., & Goldenberg, A. (2017). Vicus: Exploiting local structures to improve network-based analysis of biological data. *PLoS Computational Biology*, *13*(10), e1005621. https://doi.org/10.1371/journal.pcbi.1005621

Winblad, B., Palmer, K., Kivipelto, M., Jelic, V., Fratiglioni, L., Wahlund, L. O., ... Petersen,
R. C. (2004). Mild cognitive impairment--beyond controversies, towards a consensus: Report of the International Working Group on Mild Cognitive Impairment. *Journal of Internal Medicine*, 256(3), 240–246. https://doi.org/10.1111/j.1365-2796.2004.01380.x

World Health Organization (2020). *Dementia Key Facts*. Retrieved from http://www.who.int/news-room/fact-sheets/detail/dementia

World Health Organization. (1992). *The ICD-10 Classification of Mental and Behavioural Disorders: Clinical descriptions and diagnostic guidelines*. Geneva: World Health Organization.

Xia, J., Mazaheri, A., Segaert, K., Salmon, D. P., Harvey, D., Shapiro, K., ... Olichney, J. M. (2020). Event-related potential and EEG oscillatory predictors of verbal memory in mild cognitive impairment. *Brain Communications*, 2(2), fcaa213. https://doi.org/10.1093/braincomms/fcaa213

Ya, M., Xun, W., Wei, L., Ting, H., Hong, Y., & Yuan, Z. (2015). Is the electroencephalogram power spectrum valuable for diagnosis of the elderly with cognitive impairment? *International Journal of Gerontology*, *9*(4), 196–200. doi:10.1016/j.ijge.2014.07.001

Yin, J., Cao, J., Siuly, S, & Wang, H. (2019). An integrated MCI detection framework based on spectral-temporal analysis. *International Journal of Automation and Computing*, *16*, 786– 799 https://doi.org/10.1007/s11633-019-1197-4

Zhou, H., Jiang, J., Lu, J., Wang, M., Zhang, H., Zuo, C., & Alzheimer's Disease Neuroimaging Initiative (2019). Dual-model radiomic biomarkers predict development of mild cognitive impairment progression to Alzheimer's disease. *Frontiers in Neuroscience*, *12*, 1045. https://doi.org/10.3389/fnins.2018.01045

Zhu, J., & Liu, W. (2020). A tale of two databases: The use of Web of Science and Scopus in academic papers. *Scientometrics*, *123*(1):321-335. https://doi.org/10.1007/s11192-020-03387-8