

Recurrence relations and general solution of the exceptional Hermite equation

Alfred Michel Grundland¹, Danilo Latini², Ian Marquette²

¹ Centre de Recherches Mathématiques, Université de Montréal

Succ. Centre-Ville, CP6128, Montréal (Qc) H3C 3J7 Canada

Department of Mathematics and Computer Science,

Université du Québec, CP500, Trois-Rivières (Qc) G9A 5H7 Canada

² School of Mathematics and Physics, The University of Queensland,

Brisbane, QLD 4072, Australia

Abstract

Exceptional orthogonal Hermite and Laguerre polynomials have been linked to the k -step extension of harmonic and singular oscillators. The exceptional polynomials allow the existence of different supercharges from the Darboux-Crum and Krein-Adler constructions of supersymmetric quantum mechanics. They also allow the existence of different types of ladder relations and their associated recurrence relations. The existence of such relations is a unique property of these polynomials. Those relations have been used to construct 2D models which are superintegrable, and display an interesting spectrum, degeneracies and finite-dimensional unitary representations. In previous works, only the physical or polynomial part of the spectrum is discussed. It is known that the general solutions are associated with other types of recurrence/ladder relations. We plan to discuss in detail the case of the exceptional Hermite polynomials $X_2^{(1)}$ and to explicitly present new chains obtained by acting with different types of ladder operators. We will exploit a recent result by one of the authors [32], where the general analytic solution was constructed and connected with the non-degenerate confluent Heun equation. The analogue Rodrigues formulas for the general solution are constructed. The set of finite states from which the other states can be obtained algebraically is not unique but the vanishing arrow and diagonal arrow from the diagram of the 2-chain representations can be used to obtain minimal sets. These Rodrigues formulas are then exploited, not only to construct the states, polynomial and non-polynomial, in a purely algebraic way, but also to obtain coefficients from the action of ladder operators also in an algebraic manner based on further commutation relations between monomials of the generators.

PACS numbers: 03.65.Fd, 03.65.Ge

email: grundlan@crm.umontreal.ca, d.latini@uq.edu.au, i.marquette@uq.edu.au

1 Introduction

The combination of the Darboux-Crum (DC) and Krein-Adler (KA) transformations [1–4], also referred to as supersymmetric quantum mechanics [5–8], has been a useful tool in obtaining broader classes of exactly solvable quantum systems. These transformations allow us to obtain the spectra and their almost isospectral Hamiltonians. Their wavefunctions, however, differ greatly and are not necessarily expressed in terms of the hypergeometric class of special functions.

In most of the literature, Darboux-Crum and Krein-Adler transformations (the creating or deleting approach) are concerned with the polynomial (physical) part of the spectrum. It was recognized how those models are associated with various deformations of Heisenberg algebras [7,8] and how physical states are part of the representations which combine finite-and infinite-dimensional representations. Among those models, systems for which the physical part of the spectrum is related to exceptional orthogonal polynomials (EOP) [9–12] have been the object of particular attention and, in fact, allow the states to be associated with different Rodrigues recurrence relations and patterns of representations of the deformed Heisenberg algebras [13–19]. Other special functions are also related such as confluent and Heun equations which themselves have various properties and were studied in a systematic way from point of view of their isomonodromic deformations [20]. In regard of EOPs, it was demonstrated how the ladder operators are not unique and can even be used to build higher-dimensional models [21–26] related to the k-step extension of the harmonic and singular oscillators. Those systems are related to multi-indexed Hermite and Laguerre exceptional orthogonal polynomials of type III. A broad review of recent developments of this subject can be found in [9–19, 21–25]. Those models also play an essential role in the context of superintegrability and the case related to $X_2^{(1)}$ was present in earlier work [27] and solved via supersymmetric quantum mechanics (SUSYQM) [13, 28]. The models were also studied before using the Frobenius method [29, 30]. Those works have demonstrated how, in fact, one can obtain the polynomial solutions without relying on Darboux transformations.

It was demonstrated how, for isotonic oscillators, the standard ladders are associated with a broader type of representation of the polynomial Heisenberg algebra [21–24]. It

was discovered how the polynomial and non-polynomial parts play a role in the study of the quantum systems and their properties and how, in fact, there are generalized states from which different states can be generated [31], those states then form wider chains, which we will refer to as 2-chains. Recently [32], the general solutions were presented by a more careful analysis of the gap and series solutions which have been developed. Those solutions connected with confluent Heun equations were applied in the context of the theory of surfaces.

The purpose of this paper is to develop explicit expressions for the representations and the 2-chains for models related to the Hermite exceptional orthogonal polynomials $X_2^{(1)}$ for the standard ladder, and also for new types of ladders combining both Krein-Adler (KA) and Darboux-Crum (DC) transformations [21–24]. The second objective of this paper is to use the recent formula for the general analytic solutions [32] and the connection with the confluent Heun equations related to the exceptional Hermite polynomial $X_2^{(1)}$ to present the action for those states and to demonstrate how they are, in fact, more suitable for the description of those representations of the polynomial Heisenberg algebra. This allows us to present the representations in detail from different perspectives.

The paper is organized as follows. In Section 2, we recall the method for constructing polynomial Heisenberg algebras and how to define states of the 2-chains using the approach from [31]. We present an explicit action for the standard ladder operators and then, in Section 3, we extend the approach to the non-standard ladder operator involving different types of supercharges. We also point out how a further polynomial Heisenberg algebra can be generated by considering both types of ladder operator. The question of how the algebraic structure can mix different types of ladders has not been addressed in the literature. In Section 4, we use the relation between the EOP Hermite differential equation and the confluent Heun equation to obtain a recurrence relation involving the general solution obtained in [32] given by the μ_n and ν_n functions expressed in terms of an expression involving generalized hypergeometric functions. These expressions are new. We also rewrite higher-order ladders in terms of first-order ladder operators involving an explicit dependence on the eigenvalues of the Hamiltonian. We also consider the action of $\alpha_n^{(o)}$ and $\alpha_n^{(e)}$ which,

combined with μ_n and ν_n , provide the general solution of the confluent Heun equation where parameters are related to the EOP Hermite ordinary differential equation (ODE). In Section 5, we exploit the connection between the confluent Heun equations and their derivatives. We then obtain the ladders for the Heun equation with specific parameters. Section 6 is devoted to the obtention and modification of a Rodrigues analogue formula in order to present an algebraic construction of the states, polynomial and non-polynomial for $n \in \mathbb{Z}$. This allows us to demonstrate how the states are cyclic i.e. generated by a finite set of states, and how the 2-chain representation and the coefficients from the action of the ladder can also be obtained in a purely algebraic manner. This allows us to specify the form of those coefficients in a more direct way.

2 Construction of the polynomial Heisenberg algebra and the associated 2-chain representations for the standard ladder

In this section we recall several aspects of the harmonic oscillator and the deformed oscillator related to the Hermite exceptional orthogonal polynomial $X_2^{(1)}$ [13, 21–24, 28, 31] in order to explicitly obtain the action related to 2-chain representations for two different types of ladder operators. As a starting point, consider the simplest Hamiltonian

$$H_o = -\frac{d^2}{dx^2} + x^2.$$

This Hamiltonian has been the most studied, possessing the ladder operators a and a^\dagger

$$a = \frac{d}{dx} + x, \quad a^\dagger = -\frac{d}{dx} + x$$

which satisfy the Heisenberg algebra

$$[a, a^\dagger] = 2, \quad [H_o, a] = -2a, \quad [H_o, a^\dagger] = 2a^\dagger.$$

Next, we construct the solution of a time-independent Schrödinger problem $H\phi_n = E_n\phi_n$ and obtain the following well-known solution involving the Hermite polynomials $H_n(x)$ of

degree n

$$\phi_n = e^{-\frac{x^2}{2}} H_n(x).$$

The associated spectrum is discrete and given by $E_n = 2n + 1$. We also construct non-normalizable solutions below the ground state in terms of the pseudo-Hermite polynomial

$$\phi_n^- = e^{\frac{x^2}{2}} H_n(ix).$$

We label the indices alternatively as $\phi_{-n-1} = \phi_n^-$ and then include those with a negative index with the normalizable solutions and define ψ_p for $p = 0, \pm 1, \pm 2, \dots$. The linearly independent solution is then obtained via

$$\tilde{\phi}_p = \phi_p \int^x \frac{1}{\phi_p^2(z)} dz.$$

The importance of those states was, in particular, pointed out in [31]. Those states also admit recurrence relations connected with the Rodrigues Formula

$$a\psi_0 = 0$$

$$(a^\dagger)^n \psi_0 \propto \psi_n.$$

We consider a more general situation, namely when the Hamiltonian is


$$H = -\frac{d^2}{dx^2} + x^2 - \frac{16}{(1+2x^2)^2} + \frac{8}{1+2x^2}. \quad (2.1)$$

This leads to a description of a 1-step extension of the harmonic oscillator with $m = 2$, which was studied in [21–24]. In general, for a 1-step extension, there are two types of supercharges, both connecting the Hamiltonian with the harmonic oscillator (with different numbers of intermediate Hamiltonians). This ensures the existence of different ladder operators. Those ladder operators lead to a polynomial Heisenberg algebra which has been analyzed from the point of view of differential operator realizations and the action on the physical states ψ_n with $\psi_{-3}, \psi_0, \psi_1, \psi_2, \psi_3, \dots$ and the corresponding energies $-3, 3, 5, 7, \dots$. The spectrum is then given by $E_n = 2n + 3$ and the corresponding normalizable solutions are

$$\psi_n \propto \frac{e^{-\frac{1}{2}x^2}}{\mathcal{H}_2} y_n(x), \quad n = -3, 0, 1, 2, \dots \quad (2.2)$$

$$y_n(x) = \begin{cases} 1 & \text{if } n = -3, \\ -\mathcal{H}_2 H_{n+1} - 4\mathcal{H}_1 H_n & \text{if } n = 0, 1, 2, \dots, \end{cases} \quad (2.3)$$

where ψ_{-3} is the fundamental state.



Those physical states are related to exceptional orthogonal Hermite polynomials of type III [9–19, 21–25]. The ladder operators can be constructed from the supercharges

$$A = \frac{d}{dx} - x - \frac{8x}{2 + 4x^2}, \quad A^\dagger = -\frac{d}{dx} - x - \frac{8x}{2 + 4x^2}$$

using (2.1) and the intertwining relations

$$A^\dagger(H - 2) = H_o A^\dagger, \quad (H - 2)A = A H_o, \quad A A^\dagger = H + 3, \quad A^\dagger A = H_o + 5.$$

The supercharges A and A^\dagger connect the Hamiltonian H with the harmonic oscillator and provide the ladder operators for the Hamiltonian H , which is a deformation of the harmonic oscillator $b^\dagger = A a^\dagger A^\dagger$ and $b = A a A^\dagger$. Here, b and b^\dagger satisfy

$$[H, b] = -2b, \quad [H, b^\dagger] = 2b^\dagger \quad (2.5)$$

$$b^\dagger b = (H - 3)(H + 1)(H + 3), \quad b b^\dagger = (H - 1)(H + 3)(H + 5). \quad (2.6)$$

The action on the physical states $\{\psi_{-3}, \psi_0, \psi_1, \psi_2, \dots\}$, (where ψ_{-3} is a singlet as it is annihilated by b and b^\dagger , and ψ_0 is annihilated by b) and the action of b^\dagger on ψ_0 generate an infinite sequence of physical states. Various further identities can be proven and will be useful in using Rodrigues-type formulas and calculating the action of different ladders among those

$$[b, (b^\dagger)^n] = (b^\dagger)^{n-1}(6nH^2 + 4n(1 + 3n)H + 2n(-9 + 2n + 4n^2)) \quad (2.7)$$


$$[b^\dagger, b^n] = b^{n-1}(-6nH^2 + 4n(-7 + 3n)H - 2n(7 - 14n + 4n^2)). \quad (2.8)$$

We will now extend the study of those ladder operators to the action on physical and non-physical states. The general solution contains other linearly independent solutions that we denote $\bar{\psi}_n$ corresponding to all values of the energy (even the one with no physical states). Here, we present the two linearly independent solutions obtained via the Darboux transformation from the linearly independent solution of the Harmonic oscillator i.e. we take $\tilde{\psi}_p$ as defined previously with $p = 0, \pm 1, \pm 2, \pm 3, \dots$ and we define the state of the deformed oscillator as

$$\psi_p = A\phi_p, \quad \tilde{\psi}_p = A\tilde{\phi}_p. \quad (2.9)$$

However, we exclude $p \neq -3$ (which we consider because $\psi_{-3} = 0$), because this state is exactly the zero mode of the supercharge. We need to define the state via $A^\dagger \psi_{-3} = 0$ and then we add the linearly independent solution via

$$\tilde{\psi}_{-3} = \psi_{-3} \int^x \frac{1}{\psi_{-3}^2(z)} dz \quad (2.10)$$



$$b^\dagger \psi_n = \begin{cases} 2(n+3)\psi_{n+1} & n \geq 0 \\ 2i(n+1)(n+3)(n+4)\psi_{n+1} & -1 \geq n \geq -4 \\ 4i(n+1)(n+3)\psi_{n+1} & n \leq -5 \end{cases} \quad (2.12)$$

$$b^\dagger \tilde{\psi}_n = \begin{cases} 4(n+1)(n+3)\tilde{\psi}_{n+1} & n \geq 0 \\ -4\psi_0 & n = -1 \\ -2\imath\tilde{\psi}_{-1} & n = -2 \\ -2\imath\psi_{-2} & n = -3 \\ +2\imath\psi_{-3} & n = -4 \\ -2\imath(n+3)\tilde{\psi}_{n+1} & n \leq -5 \end{cases} \quad (2.13)$$

$$b\psi_n = \begin{cases} 4n(n+3)\psi_{n-1} & n \geq 0 \\ -4\imath\psi_{-2} & n = -1 \\ 0 & n = -2 \\ -2\imath(n+3)\psi_{n-1} & n \leq -3 \end{cases} \quad (2.14)$$

$$b\tilde{\psi}_n = \begin{cases} 2(n+3)\tilde{\psi}_{n-1} & n \geq 1 \\ 6\psi_{-1} & n = 0 \\ -8\imath\tilde{\psi}_{-2} & n = -1 \\ -8\imath\psi_{-3} & n = -2 \\ -\imath/2\psi_{-4} & n = -3 \\ 4\imath n(n+3)\tilde{\psi}_{n-1} & n \leq -4 \end{cases} \quad (2.15)$$

3 Construction of ladder operators with a Darboux-Crum/Krein-Adler transformation and 2-chain representations

The other supercharges are:

$$A_1 = \frac{d}{dx} + x - \frac{1}{x}, \quad A_1^\dagger = -\frac{d}{dx} + x - \frac{1}{x}$$

$$A_2 = \frac{d}{dx} + x + \frac{1}{x} - \frac{8x}{2+4x^2}, \quad A_2^\dagger = -\frac{d}{dx} + x + \frac{1}{x} - \frac{8x}{2+4x^2}.$$

The supercharges also satisfy the intertwining relation with H given by (2.1)

$$(A_2 A_1) H_o = (H + 4)(A_2 A_1), \quad H_o(A_1^\dagger A_2^\dagger) = (A_1^\dagger A_2^\dagger)(H + 4)$$

and the factorization

$$(A_2 A_1)(A_1^\dagger A_2^\dagger) = (H - 1)(H + 1) \quad (A_1^\dagger A_2^\dagger)(A_2 A_1) = (H_o - 5)(H_o - 3).$$


We can then form other ladder operators c and c^\dagger as follows:

$$c^\dagger = AA_1^\dagger A_2^\dagger, \quad c = A_2 A_1 A^\dagger \quad (3.1)$$

and

$$[H, c] = -6c, \quad [H, c^\dagger] = 6c^\dagger \quad (3.2)$$


$$c^\dagger c = (H + 3)(H - 5)(H - 7), \quad cc^\dagger = (H - 1)(H + 1)(H + 9). \quad (3.3)$$



Here we only include the non vanishing arrows. The action on the polynomial part of the physical states is c , which annihilates $\{\psi_{-3}, \psi_1, \psi_2\}$. Therefore, by acting on those zero modes c^\dagger gives three infinite chains of states. We demonstrate the further higher-order commutation relations

$$[c, (c^\dagger)^n] = (c^\dagger)^{n-1}(18nH^2 + 108n(n-1)H + 6n(-1 - 54n + 36n^2)) \quad (3.5)$$

$$[c^\dagger, c^n] = c^{n-1}(-18nH^2 + 108n(n-1)H - 6n(-1 - 54n + 36n^2)) \quad (3.6)$$



$$c^\dagger \psi_n = \begin{cases} -\psi_{n+3} & n \geq 0 \cup \{-3\} \\ 0 & n = -6 \\ 8i(n+1)(n+2)(n+3)\psi_{n+3} & n < 0 \setminus \{-3, -6\} \end{cases} \quad (3.8)$$

$$c \psi_n = \begin{cases} -8(n-1)(n-2)(n+3)\psi_{n-3} & n \geq 0 \\ -i\frac{n+3}{n}\psi_{n-3} & n < 0 \end{cases} \quad (3.9)$$

$$c^\dagger \tilde{\psi}_n = \begin{cases} -8(n+1)(n+2)(n+3)\tilde{\psi}_{n+3} & n \geq 0 \\ \psi_2 & n = -1 \\ -i\psi_1 & n = -2 \\ 4\tilde{\psi}_0 & n = -3 \\ -i\psi_{-3} & n = -6 \\ -i\tilde{\psi}_{n+3} & n \leq -4 \setminus \{-6\} \end{cases} \quad (3.10)$$

$$c \tilde{\psi}_n = \begin{cases} -\frac{n+3}{n}\tilde{\psi}_{n-3} & n \geq 3 \\ -5/2 \psi_{-1} & n = 2 \\ 4i\psi_{-2} & n = 1 \\ 12\tilde{\psi}_{-3} & n = 0 \\ i/12 \psi_{-6} & n = -3 \\ 8i(n(n^2-7)+6)\tilde{\psi}_{n-3} & n < 0 \setminus \{-3\} \end{cases} \quad (3.11)$$

We can show using a direct approach that only two types of ladders of order 3 exist (the lowest order). A third type of fifth-order differential operator can also be constructed. In general, different types of ladder operators can exist for a k-step extension of the harmonic oscillator but no classification has been performed so far. When the potential is unknown, the existence of ladder operators is, in general, a nonlinear problem, but when the potential is known, the compatibility equations are all linear and can be solved explicitly.

The two ladders of minimal order can be combined in a common algebraic structure. In view of the differential operators, one can demonstrate the following relations

$$(b^3)c^\dagger = -H^6 - 24H^5 - 205H^4 - 720H^3 - 739H^2 + 744H + 945$$

$$c(b^\dagger)^3 = -H^6 + 12H^5 - 25H^4 - 120H^3 + 34H^2 + 108H - 315.$$

Then, defining $B = b^3$ and $B^\dagger = (b^\dagger)^3$, we are able to close the relations as an operator algebra.

4 Connection with the confluent Heun equations and the general solution

In this section we apply the construction of the ladder in the context of the EOP Hermite ODE in a different form. Specifically, we transform the second-order ODE used in [32] into the confluent Heun equation with particular values of the parameters. In order to connect the Schrödinger equation with the confluent Heun equation, we consider the gauge/similarity $(\psi_{-3}^{-1}(H+3)\psi_{-3})$ and then obtain the second order ODE

$$\left(\frac{d^2}{dx^2} - \left(2x + \frac{8x}{1+2x^2}\right)\frac{d}{dx} + 2n\right)f_n(x) = 0.$$

The gauge/similarity transformation preserves the spectrum, the existence of the ladder, the commutation relations and the polynomial Heisenberg algebra generated by the ladder operators. The wavefunctions are only related via $\psi_n = f_n\psi_{-3}$. We denote

$$\tilde{b} = -\frac{d^3}{dx^3} + \left(2x + \frac{12x}{1+2x^2}\right)\frac{d^2}{dx^2} + \left(-4 + \frac{24}{(1+2x^2)^2} - \frac{16}{1+2x^2}\right)\frac{d}{dx}.$$

Similarly we get

$$\tilde{c} = -\frac{d^3}{dx^3} + \frac{12x}{1+2x^2}\frac{d^2}{dx^2} - \frac{48x^2}{(1+2x^2)^2}\frac{d}{dx}.$$

The raising operators can be obtained directly from the lowering in both cases.

4.1 First-order form of the ladder operators

In most of the literature on ladder operators of exceptional orthogonal polynomials, the ladders are expressed only in terms of higher-order differential operators. As pointed out in the context of SUSYQM and of different shape-invariant potentials, other alternative forms can be convenient and written as first-order ladder operators which depend on the Hamiltonian eigenvalues. Here we will present an alternative form by using the ODE

$$f_n'' = -2nf_n + \left(2x + \frac{8x}{1+2x^2}\right)f_n'$$

and the equation obtained by taking the derivative. This allows us to rewrite \tilde{c} , \tilde{c}^\dagger , \tilde{b} , \tilde{b}^\dagger as first-order operators depending explicitly on n

$$\tilde{b}_n = -\frac{8nx}{(1+2x^2)} + \frac{2(-3+n+4nx^2+4(n-1)x^4)}{(1+2x^2)^2} \frac{d}{dx}$$

$$\tilde{c}_n = \frac{4nx(-1+4x^4)}{(1+2x^2)^2} + \frac{2(-5+n+2(-5+2n)x^2+4(-5+n)x^4-8x^6)}{(1+2x^2)^2} \frac{d}{dx}.$$

4.2 The general solutions $\nu_n(x)$ and $\mu_n(x)$

In this section, our discussion will facilitate the computation of the action on μ_n and ν_n . We have

$$\nu_n(x) = x - \frac{1}{3}(n-5)x^3 + \sum_{k=3}^{\infty} 2^{k-1}((-1)^{k+1} \frac{(n - ((2(k-1))^2 + 1))}{(2k-1)!}) \prod_{j=1}^{k-2} (n-2(1+j)+1)x^{2k-1}$$

$$\mu_n(x) = 1 - nx^2 + \frac{n(n-10)}{6}x^4 + \sum_{k=3}^{\infty} 2^k((-1)^k \frac{n(n - ((2k-1)^2 + 1))}{(2k)!}) \prod_{j=1}^{k-2} (n-2(1+j))x^{2k}.$$

These expressions admit representations as generalized hypergeometric functions which can facilitate the action and consequently can be expressed as follows :

$$\begin{aligned} \mu_n(x) &= 1 - nx^2 + \frac{1}{6}n(n-10)x^4 - \frac{1}{6(n-2)} \left(-12 + 6n + 12nx^2 - 6n^2x^2 + 20nx^4 - 12n^2x^4 + n^3x^4 \right. \\ &\quad \left. + 24nx^2 {}_1F_1 \left(1 - \frac{n}{2}, \frac{3}{2}, x^2 \right) - 6(n-2) {}_1F_1 \left(-\frac{n}{2}, \frac{1}{2}, x^2 \right) - 24nx^2 {}_2F_2 \left(\left\{ 2, 1 - \frac{n}{2} \right\}, \left\{ 1, \frac{3}{2} \right\}, x^2 \right) \right) \\ &= {}_1F_1 \left(-\frac{n}{2}, \frac{1}{2}, x^2 \right) - \frac{4}{3}nx^4 {}_1F_1 \left(2 - \frac{n}{2}, \frac{5}{2}, x^2 \right) \end{aligned}$$

$$\begin{aligned} \nu_n(x) &= x - \frac{1}{3}(n-5)x^3 + \frac{1}{3(n-1)x} \left(3x^2 - 3nx^2 + 5x^4 - 6nx^4 + n^2x^4 - 15x^2 {}_1F_1 \left(\frac{1}{2} - \frac{n}{2}, \frac{3}{2}, x^2 \right) \right. \\ &\quad \left. + 3nx^2 {}_1F_1 \left(\frac{1}{2} - \frac{n}{2}, \frac{3}{2}, x^2 \right) + 24x^2 {}_2F_2 \left(\left\{ 2, \frac{1}{2} - \frac{n}{2} \right\}, \left\{ 1, \frac{3}{2} \right\}, x^2 \right) - 12x^2 {}_3F_3 \left(\left\{ 2, 2, \frac{1}{2} - \frac{n}{2} \right\}, \left\{ 1, 1, \frac{3}{2} \right\}, x^2 \right) \right) \end{aligned}$$

$$= (x + 2x^3)_1 F_1 \left(\frac{1}{2} - \frac{n}{2}, \frac{3}{2}, x^2 \right) + \frac{2}{3} x^3 (-1 + 2x^2)_1 F_1 \left(\frac{3}{2} - \frac{n}{2}, \frac{5}{2}, x^2 \right).$$

Generalized hypergeometric functions admit various recurrence relations and formulas. They can be used to calculate the action of b_n , b_n^\dagger , c_n , c_n^\dagger on $\nu_n(x)$ and $\mu_n(x)$

$$\tilde{b}_n \mu_n = (4n - 4n^2) \nu_{n-1}, \quad \tilde{b}_n \nu_n = (-6 + 2n) \mu_{n-1}$$

and consequently the differential operators independent of n and denoted \tilde{b} can be evaluated. The zero modes are, respectively, μ_0 , μ_1 and ν_3 . For the creation (raising) operators, we have the following recurrence relations

$$\tilde{b}_n^\dagger \mu_n = (4n + 4n^2) \nu_{n+1}, \quad \tilde{b}_n^\dagger \nu_n = (4 - 2n) \mu_{n+1}.$$

Also, from the differential operator \tilde{b}^\dagger independent of n , we obtain the same recurrence relations. The zero modes are μ_0 , μ_{-1} and ν_2 . Also, we obtain the following recurrence relations from the other set of ladder operators

$$\tilde{c}_n \mu_n = (16n - 4n^2) \nu_{n-3}, \quad \tilde{c}_n \nu_n = (-10 + 2n) \mu_{n-3},$$

$$\tilde{c}_n^\dagger \mu_n = 4(n - 1)(n + 3) \nu_{n+3}, \quad \tilde{c}_n^\dagger \nu_n = (4 - 2n) \mu_{n+3}.$$

Those results demonstrate how most of the general solutions of the EOP Hermite ODE have recurrence relations in a very simple form. Those functions μ_n and ν_n are, however, only a part of the general solution. In the next section we will add the other component of the general solution and this will explain how even when the general solution is written in this way, the action of the ladder still has an involved form.

4.3 The coefficients $\alpha_n^{(o)}$ and $\alpha_n^{(e)}$

In order to define the other linearly independent solutions of the EOP Hermite ODE we need to use the expression

$$\alpha_n(z) = M_3(n) H_n(z)$$

$$H_n(z) = M_1(n)\nu_n(z), \quad \forall \quad n = 2l - 1, l \geq 2$$

$$H_n(z) = M_2(n)\mu_n(z), \quad \forall \quad n = 2l, l \in \{0, 2, 3, 4, \dots\}$$

with

$$M_3(n) = \begin{cases} -2 & n = 1 \\ 2 & n = 2 \\ -M_1^{-1}(n) & n \in 2\mathbb{N} - 1 \setminus \{1\} \\ M_2^{-1}(n) & n \in 2\mathbb{N} \setminus \{2\} \end{cases}$$

and

$$M_1(n) = \frac{(-1)^{\frac{n+1}{2}} n! 2^{\frac{(n+1)}{2}}}{P_{(1,1)}(n) \prod_{j=1}^{\frac{(n-3)}{2}} (n - 2(j+1) + 1)}$$

$$M_2(n) = (-1)^{\frac{(n+2)}{2}} 2^{n-1} \pi^{-\frac{1}{2}} \Gamma\left(\frac{n-1}{2}\right), \quad \forall n \in 2\mathbb{N} \setminus \{2\}$$

where $P_{(1,1)}(n) = (n-1)(n-2)$. We also have

$$\alpha_1^{(o)} = -8x - 2e^{x^2} \sqrt{\pi} (1 - 2x^2) (1 - \text{Erf}(x)) = (-2\sqrt{\pi}\mu_1 - 4\nu_1),$$

$$\alpha_2^{(e)} = 4 + 8x^2 + 8\sqrt{\pi}xe^{x^2} (1 - \text{Erf}(x)) = \alpha_2^{(e)} = 4(\mu_2 + 2\sqrt{\pi}\nu_2).$$

There are further relations such as

$$\alpha_n^{(o)} = -\nu_n, \quad n = 3, 5, 7, \dots$$

$$\alpha_n^{(e)} = \mu_n, \quad n = 4, 6, 8, \dots$$

We evaluate the action of b_n , b_n^\dagger , c_n , c_n^\dagger and \tilde{b} , \tilde{b}^\dagger , \tilde{c} , \tilde{c}^\dagger , and consider the $\alpha_n^{(o)}$ for odd indices and $\alpha_n^{(e)}$ for even indices

$$\tilde{b}\alpha_1^{(o)} = 16\alpha_0^{(e)}, \quad \tilde{b}^\dagger\alpha_1^{(o)} = -2\alpha_2^{(e)}$$

with $\alpha_0 := \mu_0 \equiv \alpha_0^{(e)} \equiv \alpha_0^{(o)} = 1$. We will only list the relations for other ladder operators

$$\begin{aligned}
\tilde{b}\alpha_2^{(e)} &= 8\alpha_1^{(o)} \quad \tilde{b}^\dagger\alpha_2^{(e)} = -96\alpha_3^{(o)} \\
\tilde{b}\alpha_n^{(o)} &= (6-2n)\alpha_{n-1}^{(e)}, \quad n = 3, 5, 7, \dots \\
\tilde{b}^\dagger\alpha_n^{(o)} &= (2n-4)\alpha_{n+1}^{(e)}, \quad n = 3, 5, 7, \dots \\
\tilde{b}\alpha_n^{(e)} &= 4n(n-1)\alpha_{n-1}^{(o)}, \quad n = 4, 6, 8, \dots \\
\tilde{b}^\dagger\alpha_n^{(e)} &= -4n(n+1)\alpha_{n+1}^{(o)}, \quad n = 4, 6, 8, \dots \\
\tilde{c}\alpha_1^{(o)} &= 32\mu_{-2} - 24\sqrt{\pi}\nu_{-2}, \quad \tilde{c}^\dagger\alpha_1^{(o)} = -8\alpha_4^{(e)}, \quad \tilde{c}\alpha_2^{(e)} = -48\sqrt{\pi}\mu_{-1} + 64\nu_{-1} \\
\tilde{c}^\dagger\alpha_2^{(e)} &= -80\alpha_5^{(o)}, \quad \tilde{c}\alpha_n^{(o)} = \begin{cases} 4\alpha_0 & n = 3 \\ (10-2n)\alpha_{n-3}^{(e)} & n = 5, 7, 9, \dots \end{cases}, \quad \tilde{c}^\dagger\alpha_n^{(o)} = (2n-4)\alpha_{n+3}^{(e)} \quad n = 3, 5, \dots \\
\tilde{c}\alpha_n^{(e)} &= 4n(n-4)\alpha_{n-3}^{(o)}, \quad n = 4, 6, \dots \\
\tilde{c}^\dagger\alpha_n^{(e)} &= 4(1-n)(n+3)\alpha_{n+3}^{(o)}, \quad n = 4, 6, \dots
\end{aligned}$$

In this context, the coefficient of the action of the ladder differs greatly from the action in Sections 2 and 3 which were obtained via Darboux transformation of the eigenfunctions of the harmonic oscillator for the two linearly independent solutions. Here we used series solution and connection with generalized and hypergeometric functions. The coefficients are now connected with the functions obtained by [32], which constitute the general solution.

5 Ladders and representations in the Heun form

The purpose of this section is to connect the obtained results with the confluent Heun and Heun equations. The EOP $X_2^{(1)}$ Hermite ODE

$$\omega'' - 2\left(x + \frac{4x}{1+2x^2}\right)\omega' + 2n\omega = 0$$

can be written in the following form in terms of the variable s ($y = -2x^2$) [32]

$$\frac{d^2\omega}{ds^2} + \left(\frac{\gamma}{s} + \frac{\delta}{s-1} + \epsilon\right) \frac{d\omega}{ds} + \left(\frac{\alpha s - q}{s(s-1)}\right) \omega = 0$$

where the parameters of the confluent Heun equation take the following values : $\gamma = \frac{1}{2}$, $\alpha = q = -\frac{n}{4}$, $\epsilon = \frac{1}{2}$, $\delta = -2$. Similarly, the ladder operators can be written in terms of the variable s :

$$\begin{aligned}\bar{b} &= -16s\sqrt{-2s}\partial_s^3 - \frac{8\sqrt{-2s}(s^2 - 4s - 3)}{s-1}\partial_s^2 + \frac{4\sqrt{-2s}(s^2 - 4s - 9)}{(s-1)^2}\partial_s \\ \bar{c} &= -16s\sqrt{-2s}\partial_s^3 - \frac{24\sqrt{2s}(s+1)}{\sqrt{-s}(s-1)}\partial_s^2 - \frac{24\sqrt{-2s}(s+1)}{(s-1)^2}\partial_s.\end{aligned}$$

The functions μ and ν can also be written in term of s and expressed in term of the confluent hypergeometric function ${}_1F_1$ only

$$\begin{aligned}\bar{\nu}_n(s) &= \frac{\sqrt{-s}}{3\sqrt{2}} \left(-3(s-1) {}_1F_1 \left(\frac{1}{2} - \frac{n}{2}, \frac{3}{2}, -\frac{s}{2} \right) + s(s+1) {}_1F_1 \left(\frac{3}{2} - \frac{n}{2}, \frac{5}{2}, -\frac{s}{2} \right) \right) \\ \bar{\mu}_n(s) &= \frac{1}{n-2} \left(-(s^2 + (n+1)s - n + 2) {}_1F_1 \left(-\frac{n}{2}, \frac{1}{2}, -\frac{s}{2} \right) + (1+n)s(s+1) {}_1F_1 \left(-\frac{n}{2}, \frac{3}{2}, -\frac{s}{2} \right) \right)\end{aligned}$$

with recurrence relations

$$\begin{aligned}\bar{b}\bar{\mu}_n &= 4n(1-n)\bar{\nu}_{n-1}, & \bar{b}\bar{\nu}_n &= (2n-6)\bar{\mu}_{n-1} \\ \bar{b}^\dagger\bar{\mu}_n &= 4n(n+1)\bar{\nu}_{n+1}, & \bar{b}^\dagger\bar{\nu}_n &= (4-2n)\bar{\mu}_{n+1} \\ \bar{c}\bar{\mu}_n &= 4n(4-n)\bar{\nu}_{n-3}, & \bar{c}\bar{\nu}_n &= (2n-10)\bar{\mu}_{n-3} \\ \bar{c}^\dagger\bar{\mu}_n &= 4(n-1)(n+3)\bar{\nu}_{n+3}, & \bar{c}^\dagger\bar{\nu}_n &= (4-2n)\bar{\mu}_{n+3}.\end{aligned}$$

We now want to look at the ladder operators for the derivative of ω . The connection between the derivative of a Heun function was discussed in a recent work [32]. However recurrence relations for derivatives of the Heun polynomial appear to be an unexplored problem in the literature. Setting $v(s) = d\omega/ds$, we then perform a change of variable $s = \frac{s_1}{s_1-1}$ and $v(s_1) = (1-s_1)^{(1-\frac{n}{2})}W(s_1) = g_n W(s_1)$. This can also be done in operator form as the gauge transformation $\hat{H}_1 = g_n^{-1}\bar{H}_1 g_n$ with \hat{H}_1 acting on $W(s_1)$ as $\hat{H}_1 W(s_1) = 0$, which gives

$$(-1 + s_1)^2 s_1 \frac{d^2}{ds_1^2} W + \frac{1}{4} (6 + 4s_1(-8 + n + 6s_1 - ns_1)) \frac{d}{ds_1} W + \frac{1}{4} (n - 4)(4 + (n - 6)s_1) W = 0.$$

Here, $W(s_1)$ has the linearly independent solutions

$$\begin{aligned} \hat{\mu}_n(s_1) &= \frac{n(1 - s_1)^{\frac{n-6}{2}}}{6(n-2)} \left(3(s_1(1 - n) + n - 2) {}_1F_1 \left(1 - \frac{n}{2}; \frac{3}{2}; \frac{s_1}{2 - 2s_1} \right) \right. \\ &\quad \left. + (n + 1)s_1 {}_1F_1 \left(1 - \frac{n}{2}; \frac{5}{2}; \frac{s_1}{2 - 2s_1} \right) \right) \\ \hat{\nu}_n(s_1) &= \frac{\sqrt{\frac{1}{2-2s_1}}(1 - s_1)^{\frac{n-6}{2}}}{6\sqrt{s_1}(n-1)} \left(3((s_1^2 - 1)n + 1) {}_1F_1 \left(\frac{1}{2} - \frac{n}{2}; \frac{3}{2}; \frac{s_1}{2 - 2s_1} \right) \right. \\ &\quad \left. - s_1(n + 2)((s_1 - 1)n + 1) {}_1F_1 \left(\frac{1}{2} - \frac{n}{2}; \frac{5}{2}; \frac{s_1}{2 - 2s_1} \right) \right). \end{aligned}$$

We want to obtain the corresponding ladder operator from the same transformation i.e. taking the derivative and the above gauge transformation. In particular we consider $\frac{d}{ds} \bar{b}^\dagger$ and $\frac{d}{ds} \bar{b}$ and $v = \frac{d\omega}{ds}$. Here we only present the ladder operators b and b^\dagger acting on $\hat{\mu}$ and $\hat{\nu}$ in the s_1 variable. Explicitly, the lowering operator becomes

$$\begin{aligned} \hat{b}_n &= -16\sqrt{2}(-1 + s_1)^4 s_1^{\frac{3}{2}} \frac{d^3}{ds_1^3} + (8\sqrt{2}(-1 + s_1)^2 \sqrt{s_1}(-6 + (31 - 3n)s_1 + 3(-8 + n)s_1^2) \frac{d^2}{ds_1^2} \\ &\quad - \frac{1}{\sqrt{s_1}}(4\sqrt{2}(-1 + s_1)(-3 + s_1(81 - 12n - (-6 + n)(-38 + 3n)s_1 + 3(-8 + n)(-6 + n)s_1^2)) \frac{d}{ds_1} \\ &\quad - \frac{1}{\sqrt{s_1}}(2\sqrt{2}(3(-5 + n) + 3(-4 + n)(-15 + 2n)s_1 + (-15 + n)(-8 + n)(-4 + n)s_1^2 \\ &\quad - (-8 + n)(-6 + n)(-4 + n)s_1^3)) \end{aligned}$$

and we have

$$\begin{aligned} \hat{b}_n \hat{\mu}_n &= 4n(1 - n) \hat{\nu}_{n-1}, \quad \hat{b}_n \hat{\nu}_n = (2n - 6) \hat{\mu}_{n-1} \\ \hat{b}_n^\dagger \hat{\mu}_n &= 4n(n + 1) \hat{\nu}_{n+1}, \quad \hat{b}_n^\dagger \hat{\nu}_n = (2n - 4) \hat{\mu}_{n+1}. \end{aligned}$$

6 Rodrigues-type formula for both third ladder operators

We consider the problem of constructing algebraically all the solutions of the differential equation for the exceptional Hermite polynomial $X_2^{(1)}$. It was demonstrated how, from the ladder operators $\{b^\dagger, b\}$, $\{c^\dagger, c\}$, one can generate in a purely algebraic way the physical states for $\{\psi_n, n = -3, 0, 1, 2, \dots\}$. These can be viewed as analogous to the Rodrigues formula

$$\begin{aligned} b\psi_{-3} &= 0, & b^\dagger\psi_{-3} &= 0 \\ b\psi_0 &= 0, & \psi_n &= (b^\dagger)^n\psi_0. \end{aligned}$$

We then need two initial zero modes to induce all the physical states. Also we can then reduce the problem of finding the action of different operators to an algebraic setting. We obtain directly

$$b^\dagger\psi_n = \psi_{n+1}, \quad b\psi_n = (8n(n+2)(n+3))\psi_{n-1}.$$

Interestingly, there exists another construction and it can also be put completely into an algebraic setting

$$c\psi_{-3} = 0, \quad c\psi_1 = 0, \quad c\psi_2 = 0$$

and

$$\psi_{3n-3} = (c^\dagger)^n\psi_{-3}, \quad \psi_{3n+1} = (c^\dagger)^n\psi_1, \quad \psi_{3n+2} = (c^\dagger)^n\psi_2$$

and the action is then given by

$$c^\dagger\psi_{3n-3} = \psi_{3(n+1)-3}, \quad c^\dagger\psi_{3n+1} = \psi_{3(n+1)+1}, \quad c^\dagger\psi_{3n+2} = \psi_{3(n+1)+2}$$

$$c\psi_{3n-3} = 24n(3n-5)(3n-4)\psi_{3(n-1)-3}, \quad c\psi_{3n+1} = 24n(3n-1)(3n+4)\psi_{3(n-1)+1},$$

$$c\psi_{3n+2} = 24n(3n+1)(3n+5)\psi_{3(n-1)+2}.$$

The action is not described for either of the cases of the gap states i.e. $n = -2, -1$, the unbounded states with $n \leq -4$ or the other solutions of the Hamiltonian denoted $\bar{\psi}_n$ for $n \in \mathbb{Z}$. A recent work by one of the authors [33] discussed the idea of a recurrence relation connecting ψ_{-2} , ψ_{-1} with the other physical states. Here, we will discuss this connection using another algebraic approach. We consider the case $n = \{-3, -2, -1, 0, 1, \dots\}$, ψ_n . Using ladder operators of type b , we obtain ψ_{-3} , ψ_{-2} and ψ_0 as initial states which induce all other states

$$\psi_{-1} = (b^\dagger)\psi_{-2}, \quad \psi_n = (b^\dagger)^n\psi_0.$$

In this case, we want to induce other ψ_n with $n \leq -4$ using

$$\psi_{-4-n} = (b)^n\psi_{-4}.$$

We can then obtain the action as

$$b^\dagger\psi_{-1} = 0, \quad b\psi_{-1} = -4i\psi_{-2}, \quad b\psi_{-2} = 0.$$

Then

$$b^\dagger\psi_n = \psi_{n+1}, \quad b\psi_n = 8n(n+2)(n+3)\psi_{n-1}$$

$$b\psi_{-4-n} = \psi_{-4-(n+1)}, \quad b^\dagger\psi_{-4-n} = -8n(n+1)(n+3)\psi_{-4-(n-1)}.$$

For ψ_n with $n \in \mathbb{Z}$, we can use the c -type ladder operators as well as the initial state to generate all the states. The initial states are $\{\psi_{-6}, \psi_{-1}, \psi_{-2}\}$

$$\psi_{-3n-6} = (c)^n\psi_{-6}, \quad \psi_{3n-6} = (c^\dagger)^n\psi_{-6}, \quad \psi_{-1-3n} = (c)^n\psi_{-1}$$

$$\psi_{-1+3n} = (c^\dagger)^n\psi_{-1}, \quad \psi_{-2+3n} = (c^\dagger)^n\psi_{-2}, \quad \psi_{-2-3n} = (c)^n\psi_{-2}$$

and then the action will be given by the formula

$$\begin{aligned}
c\psi_{3n-6} &= 24(n-1)(3n-8)(3n-7)\psi_{3(n-1)-6}, & c^\dagger\psi_{3n-6} &= \psi_{3(n+1)-6} \\
c\psi_{3n-1} &= 24(n-1)(3n-2)(3n+2)\psi_{3(n-1)-1}, & c^\dagger\psi_{3n-1} &= \psi_{3(n+1)-1} \\
c\psi_{3n-2} &= 24(n-1)(3n-4)(3n+1)\psi_{3(n-1)-2}, & c^\dagger\psi_{3n-2} &= \psi_{3(n+1)-2}
\end{aligned}$$

and

$$\begin{aligned}
c^\dagger\psi_{-3n-6} &= -24n(4+3n)(5+3n)\psi_{-3(n-1)-6}, & c\psi_{-3n-6} &= \psi_{-3(n+1)-6} \\
c^\dagger\psi_{-3n-1} &= -24n(3n-5)(3n-1)\psi_{-3(n-1)-1}, & c\psi_{-3n-1} &= \psi_{-3(n+1)-1} \\
c^\dagger\psi_{-3n-2} &= -24n(3n-4)(3n+1)\psi_{-3(n-1)-2}, & c\psi_{-3n-2} &= \psi_{-3(n+1)-2}.
\end{aligned}$$

Now we turn to the problem of inducing all solutions of the Hermite equation of $X_2^{(1)}$, ψ_n and $\bar{\psi}_n$. Those have been explicitly calculated in terms of a series for $n \geq -3$. Here we will construct ψ_n and $\bar{\psi}_n \forall n \in \mathbb{Z}$ using an algebraic approach. For the b -type ladder operators we will need four states $\bar{\psi}_{-2}$, ψ_{-2} , $\bar{\psi}_{-4}$ and ψ_0

$$\begin{aligned}
\psi_{-3} &= b\bar{\psi}_{-2}, & \bar{\psi}_{-1} &= b^\dagger\bar{\psi}_{-2}, & \psi_{n-2} &= (b^\dagger)^{n+1}\bar{\psi}_{-2}, & n &\geq 1 \\
\psi_{-1} &= b^\dagger\psi_{-2}, & \bar{\psi}_{-4-n} &= (b)^n\bar{\psi}_{-4}, & \bar{\psi}_{-3} &= b^\dagger\bar{\psi}_{-4} \\
\psi_{-4-n} &= (b)^nb^\dagger\bar{\psi}_{-4}, & \bar{\psi}_n &= (b^\dagger)^n\bar{\psi}_0, & b^\dagger\bar{\psi}_n &= \bar{\psi}_{n+1} \\
b\bar{\psi}_n &= 8n(2+n)(3+n)\bar{\psi}_{n-1}.
\end{aligned}$$

From the above relations together with the others given previously we get

$$\begin{aligned}
b\psi_{n-1} &= 8(n-1)(n+1)(n+2)\psi_{n-2}, & b^\dagger\psi_{n-2} &= \psi_{n-1} \\
b\bar{\psi}_{-4-n} &= \bar{\psi}_{-4-(n+1)}, & b^\dagger\bar{\psi}_{-4-n} &= -4n(6+n)(1+2n)\bar{\psi}_{-4-(n-1)} + \psi_{-4-n} \\
b^\dagger\psi_{-4-n} &= -8n(n+1)(n+3)\psi_{-4-(n-1)}, & b\psi_{-4-n} &= \psi_{-4-(n+1)}.
\end{aligned}$$

We will now use the c -type ladder operators and the functions $\bar{\psi}_1$, $\bar{\psi}_2$, $\bar{\psi}_{-3}$, ψ_{-3} , ψ_{-6} . A pattern is obtained from the diagonal arrow and the one vanishing in the 2-chain representation obtained by the action on the states constructed using ladders and supercharges

$$\begin{aligned}
\bar{\psi}_{1-3n} &= (c^\dagger)^n \bar{\psi}_1, \quad n \geq 0, & \psi_{3n-2} &= (c^\dagger)^n b \bar{\psi}_1, \quad n \geq 0 \\
\psi_{1-3n} &= (c)^n \bar{\psi}_1, \quad n \geq 0, & \bar{\psi}_{1+3n} &= (c^\dagger)^n \bar{\psi}_1, \quad n \geq 0 \\
\bar{\psi}_{2+3n} &= (c^\dagger)^n \bar{\psi}_2, & \psi_{-1+3n} &= (c^\dagger)^n c \bar{\psi}_2, & \psi_{2-3n} &= (c)^n \bar{\psi}_2, & \bar{\psi}_{-3n-3} &= (c)^n \bar{\psi}_{-3} \\
\bar{\psi}_{3n-3} &= (c^\dagger)^n \bar{\psi}_{-3}, & \psi_{3n-3} &= (c^\dagger)^n \psi_{-3}, & \psi_{-3n-6} &= (c)^n \psi_{-6}
\end{aligned}$$

and we get the following

$$\begin{aligned}
c \bar{\psi}_{1-3n} &= 24n(3n-1)(4+3n) \bar{\psi}_{1-3(n-1)} + \psi_{3n-2}, & c \psi_{3n-2} &= 24n(3n-1)(4+3n) \bar{\psi}_{3(n-1)-2} + \psi_{3n-2} \\
c \psi_{1-3n} &= \psi_{1-3(n+1)}, & c \bar{\psi}_{2+3n} &= 24n(1+3n)(5+3n) \bar{\psi}_{2+3(n-1)} + \psi_{-1+3n} \\
c \psi_{-1+3n} &= 24(n-1)(3n-2)(3n+2) \psi_{-1+3(n-1)} \\
c \psi_{2-3n} &= \psi_{2-3(n+1)}, & c \bar{\psi}_{-3n-3} &= \bar{\psi}_{-3(n+1)} \\
c \bar{\psi}_{3n-3} &= 24n(-5+3n)(-4+3n) \bar{\psi}_{3(n-1)-3}, & c \psi_{3n-3} &= 24n(-5+3n)(-4+3n) \psi_{3(n-1)-3} \\
c \psi_{3n-6} &= \psi_{3(n+1)-6}
\end{aligned}$$

and

$$\begin{aligned}
c^\dagger \bar{\psi}_{1-3n} &= \bar{\psi}_{1-3(n+1)}, & c^\dagger \psi_{3n-2} &= \psi_{3(n+1)-2} \\
c^\dagger \psi_{1-3n} &= -24(n-1)(3n-7)(3n-2) \psi_{1-3(n-1)}, & c^\dagger \psi_{1+3n} &= \psi_{1+3(n+1)} \\
c^\dagger \bar{\psi}_{2+3n} &= \bar{\psi}_{2+3(n+1)}, & c^\dagger \psi_{-1+3n} &= \psi_{-1+3(n+1)} \\
c^\dagger \psi_{2-3n} &= (768 - 24n(68 - 45n + 9n^2)) c^{n-1} \bar{\psi}_2, & c^\dagger \bar{\psi}_{-3n-3} &= (48 - 24n(-7 + 9n^2)) \bar{\psi}_{-3(n-1)-3} \\
c^\dagger \psi_{2-3n} &= -24(n-1)(3n-8)(3n-4) \bar{\psi}_{2-3(n-1)}, & c^\dagger \bar{\psi}_{-3n-3} &= (48 - 24n(-7 + 9n^2)) \bar{\psi}_{-3(n-1)-3} \\
c^\dagger \bar{\psi}_{3n-3} &= \bar{\psi}_{3(n+1)-3}, & c^\dagger \psi_{3n-3} &= \psi_{3(n+1)-3}.
\end{aligned}$$

We have now presented a complete description of the 2 chain representations in an algebraic manner. Those results demonstrate how all the states are, in fact, obtained via a

minimal set of states, both polynomial and non-polynomial. More importantly, they transform the problem of obtaining the coefficients from a problem of higher-order differential operators acting on a non-trivial solution of the EOP Hermite ODE into a simpler problem where the commutation relations can be determined and used to build all the coefficients explicitly as third-degree polynomials in terms of n .

7 Conclusion

In this paper we obtain explicit expressions for 2-chain representations for the polynomial Heisenberg algebra of the deformed oscillator related to X_2 . This demonstrates how the polynomial, but also the non-polynomial part, play a role from a mathematical perspective. We build those 2-chain representations directly by constructing the states on \mathbb{Z} and by then constructing the linearly independent solutions using the connection with the harmonic oscillator (Hermite) ODE. Non-trivial coefficients are then obtained for the action of the standard and non-standard ladder operators, both third-order differential operators.

We present an alternative point of view by using recently introduced general solutions [32] and point out how those special functions beyond the hypergeometric are suitable for a description of those representations. We also discuss the construction in the context of the confluent form of the equation and also point out how the results can be used to build ladders for the Heun equation.

One of the main results of the paper is the construction of a Rodrigues analogue which allows us to construct all the states in an algebraic way. We then use commutator identities to calculate the action of lowering and raising operators in a purely algebraic way. All those results describe a ladder for the EOP $X_2^{(1)}$ Hermite ODE for the general solution from a different perspective related to the theory of ODE and of algebraic structures beyond Lie algebras. Those 2-chain representations are also interesting as they extend finite and infinite dimensional 1-chain representations made by the polynomial part of the solution. The connection of EOP Hermite equations [34, 35] and Heun equation [20] with Painlevé transcendents were the objectives of several works. Those connection may allow us to understand equation of the Painlevé transcendents and their properties.

Acknowledgments

AMG was supported by a research grant from NSERC of Canada. DL was supported by Australian Research Council Discovery Project DP190101529 (A/Prof. Y.-Z. Zhang). IM was supported by the Australian Research Council Fellowship FT180100099. I.M. thanks the Centre de Recherches Mathématiques, Université de Montréal for its hospitality.

References

- [1] M.G. Darboux, Sur une proposition relative aux équations linéaires, *Comptes Rendus Acad. Sci.* **94** (1882) 1456.
- [2] M. M. Crum, Associated Sturm-Liouville systems, *Q. J. Math. Oxford Ser. 2* **6**, 121 (1955).
- [3] M.G. Krein, On a continual analogue of a Christoffel formula from the theory of orthogonal polynomials, *Dokl. Akad. Nauk SSSR* **113**, 970 (1957).
- [4] V.E. Adler, On a modification of Crum's method, *Theor. Math. Phys.* **101**, 1381 (1994).
- [5] G. Junker, *Supersymmetric Methods in Quantum and Statistical Physics* (Springer, Berlin, 1996).
- [6] B. K. Bagchi, *Supersymmetry in Quantum and Classical Physics* (Chapman, Boca Raton, 2001).
- [7] A. Andrianov, F. Cannata, M. Ioffe, and D. Nishnianidze, Systems with higher-order shape invariance: spectral and algebraic properties, *Phys. Lett. A* **266** (2000) 341.
- [8] D.J. Fernández, and V. Hussin, Higher-order SUSY, linearized nonlinear Heisenberg algebras and coherent states, *J. Phys. A: Math. Gen.* **32** (1999) 3603.
- [9] D. Gomez-Ullate, N. Kamran, and R. Milson, An extension of Bochner's problem: Exceptional invariant subspaces, *J. Approx. Theory* **162**, 987 (2010).
- [10] D. Gomez-Ullate, N. Kamran, and R. Milson, An extended class of orthogonal polynomials defined by a Sturm-Liouville problem, *J. Math. Anal. Appl.* **359**, 352 (2009).
- [11] S. Odake and R. Sasaki, Infinitely many shape invariant potentials and new orthogonal polynomials, *Phys. Lett. B* **679**, 414 (2009).

- [12] C. Quesne, Solvable rational potentials and exceptional orthogonal polynomials in supersymmetric quantum mechanics, *SIGMA* **5**, 084 (2009).
- [13] J. M. Fellows and R. A. Smith, Factorization solution of a family of quantum nonlinear oscillators, *J. Phys. A* **42**, 335303 (2009).
- [14] C. Quesne, Higher-order SUSY, exactly solvable potentials, and exceptional orthogonal polynomials, *Mod. Phys. Lett. A* **26**, 1843 (2011).
- [15] S. Odake and R. Sasaki, Krein-Adler transformations for shape-invariant potentials and pseudo virtual states, *J. Phys. A: Math. Theor.* **46**, 245201 (2013).
- [16] D. Gomez-Ullate, Y. Grandati, and R. Milson, Extended Krein-Adler theorem for the translationally shape invariant potentials, *J. Math. Phys.* **55**, 043510 (2014).
- [17] D. Gomez-Ullate, Y. Grandati, and R. Milson, Rational extensions of the quantum harmonic oscillator and exceptional Hermite polynomials, *J. Phys. A: Math. Theor.* **47**, 015203 (2014).
- [18] D. Gomez-Ullate, A. Kasman, A.B.J. Kuijlaars, and R. Milson, Recurrence relations for exceptional Hermite polynomials, *J. Approx. Theor.* **204** (2016) 1.
- [19] D. Gomez-Ullate, Y. Grandati, and R. Milson, Spectral Theory of Exceptional Hermite Polynomials, arXiv:2012.02354 [math.CA].
- [20] J. Dereziński, A. Ishkhanyan and A. Latosinski, From Heun class equations to Painlevé equations, *SIGMA* **17** (2021) 056, 59 pages.
- [21] I. Marquette and C. Quesne, New families of superintegrable systems from Hermite and Laguerre exceptional orthogonal polynomials, *J. Math. Phys.* **54**, 042102 (2013).
- [22] I. Marquette and C. Quesne, Two-step rational extensions of the harmonic oscillator: exceptional orthogonal polynomials and ladder operators, *J. Phys. A: Math. Theor.* **46**, 155201 (2013).

- [23] I. Marquette and C. Quesne, New ladder operators for a rational extension of the harmonic oscillator and superintegrability of some two-dimensional systems, *J. Math. Phys.* **54**, 102102 (2013).
- [24] I. Marquette and C. Quesne, Combined state-adding and state-deleting approaches to type III multi-step rationally extended potentials: Applications to ladder operators and superintegrability, *J. Math. Phys.* **55**, 112103 (2014).
- [25] J. F. Carinena and M. S. Plyushchay, ABC of ladder operators for rationally extended quantum harmonic oscillator systems, *J. Phys A* **50** 27 275202 (2017).
- [26] D. Latini, I. Marquette, Y.-Z. Zhang, Polynomial algebras of superintegrable systems separating in Cartesian coordinates from higher order ladder operators, *Physica D: Nonlinear Phenomena* **440** 133464 (2022).
- [27] S. Gravel, Hamiltonians separable in Cartesian coordinates and third-order integrals of motion, *J. Math. Phys.* **45**, 1003 (2004).
- [28] I. Marquette, Superintegrability with third order integrals of motion, cubic algebras, and supersymmetric quantum mechanics. I. Rational function potentials, *J. Math. Phys.* **50**, 012101 (2009).
- [29] J. Sesma, The generalized quantum isotonic oscillator *J. Phys. A: Math. Theor.* **43** 185303 (2010).
- [30] J.F. Carinena, A.M. Perelomov, M.F. Ranada and M. Santander, A quantum exactly solvable non-linear oscillator related with the isotonic oscillator *J. Phys. A: Math. Theor.* **41**, 085301 (10 p.) (2008).
- [31] J.F. Carinena and M.S. Plyushchay, Ground-state isolation and discrete flows in a rationally extended quantum harmonic oscillator *Physical Review D* **94** (10) 105022 (2016).

- [32] V. Chalifour and A.M. Grundland, General solution of the exceptional Hermite differential equation and its minimal surface representation, *Ann. Henri Poincaré* **21** (2020) 3341.
- [33] G. Filipuk, A Ishkhangan and J Derezinski, On the derivatives of the Heun functions, *J. Contemp. Math. Anal.* **55** 200 (2020).
- [34] I. Marquette and C. Quesne, Connection between quantum systems involving the fourth Painlevé transcendent and k-step rational extensions of the harmonic oscillator related to Hermite exceptional orthogonal polynomial, *J. Math. Phys.* **57** (2016) 052101.
- [35] K. Zelaya, I. Marquette, and V. Hussin, Third-order ladder operators, generalized Okamoto and exceptional orthogonal polynomials, *J. Phys. A: Math. Theor.* **55** 045205 (2022).