

Two classes of subfield codes of linear codes

Xiaoqiong Ran^a, Rong Luo^a

^a*School of Mathematics, Southwest Jiaotong University, Chengdu, 610031, China*

Abstract

Recently, subfield codes of linear code over $\text{GF}(q)$ with good parameters were studied, and many optimal subfield codes were obtained. In this paper, Our mainly motivation is to generalize the results of the subfield codes of hyperoval in Ding and Heng (Finite Fields Their Appl. 56, 308-331 (2019)), and generalize the results of two families of subfield codes in Xiang and Yin (Cryptogr. Commun. 13(1), 117-127 (2021)) to p -ary where p is odd. We get the parameters and weight distribution of these subfield codes. At the same time, the parameters of their dual codes are also studied. When $m = 1$, The dual codes of these subfield codes are almost MDS code, when $m > 1$ and p odd, these dual codes are dimension-optimal with respect to the sphere-backing bound.

Keywords:

Linear code, weight distribution, subfield code, character sums.

1. Introduction

Let q be a prime power. Let $\text{GF}(q)$ be a finite fields with q elements. An $[n, k, d]$ linear code over $\text{GF}(q)$ is a k -dimensional subspace of $\text{GF}(q)^n$ with length n and minimum Hamming distance d , where n, k, d are positive integer. The dual code of C is defined by

$$C^\perp = \{y \in \text{GF}(q)^n : yx^T = 0 \text{ for all } x \in C\},$$

where yx^T is the standard inner product of two vectors x and y . Let A_i denote the number of codewords with Hamming weight i of linear code C over $\text{GF}(q)$. The weight enumerator of C is defined by $1 + A_1z + A_2z^2 + \cdots + A_nz^n$, and sequence $(1, A_1, \dots, A_n)$ is weight distribution of C . Studying weight distribution of linear code is very important. A linear code C is called t -weight code if the number of nonzero A_i in the sequence (A_1, A_2, \dots, A_n) is equal to t . A linear code C over $\text{GF}(q)$ is said to be dimension-optimal (or distance-optimal) if there does not exist an

*Corresponding author

Email addresses: Ran2021@my.swjtu.edu.cn (Xiaoqiong Ran), luorong@swjtu.edu.cn (Rong Luo)

$[n, \geq k+1, d]$ (or $[n, k, \geq d+1]$) linear code over $\text{GF}(q)$ and almost distance-optimal (or almost dimension-optimal) if $[n, k, d+1]$ optimal (or $[n, k+1, d]$) optimal.

Let $f_i(x_1, x_2, \dots, x_t) (1 \leq i \leq a)$ be an t -variable polynomial over $\text{GF}(q)$. Denote

$$G_{x_1, x_2, \dots, x_t} = \begin{pmatrix} f_1(x_1, x_2, \dots, x_t) \\ f_2(x_1, x_2, \dots, x_t) \\ \dots \\ f_a(x_1, x_2, \dots, x_t) \end{pmatrix}_{(x_1, x_2, \dots, x_t) \in \text{GF}(q)^t}$$

which is a $a \times q^t$ matrix over $\text{GF}(q)$.

Let $B = [G_{(x_1, x_2, \dots, x_t)} | A]$, we define a class of linear codes C over $\text{GF}(q)$ with generator matrix B , where A is a $a \times b$ matrix over $\text{GF}(q)$ consists of b distinct column vectors of the identity matrix I_a for $0 \leq b \leq a$.

Recently, Subfield codes of linear code over finite fields with good parameters and optimal were widely studied, which were first considered in [3, 4], but the authors do not use the name "subfield codes". The notion of subfield codes was first introduced in [[5], p.5117] and a Magma function for subfield codes is actually executed in the Magma package. With the help of Magma, some of optimal or almost optimal subfield codes were obtained.

Heng and Ding obtained some basic properties and results about subfield codes [10], they have studied the subfield codes of the two families of ovoid codes, and the parameters and the weight distributions of the subfield codes of the elliptic quadric codes and Tits ovoid codes were obtained. Meanwhile, Heng and Ding [2, 6, 7, 9] studied the subfield codes of hyperoval codes, conic codes, maximal arcs and some $[q+1, 2, q]$ MDS codes, [1] discussed two families of subfield codes of linear code when $p = 2$ and obtained some distance-optimal and almost distance-optimal binary codes with a few weights. Wang and Zheng [11] extended the work of Heng and Ding in [2], and studied more subfield codes of linear code. [8] studied a class of subfield codes, which are distance-optimal. It can be known from other's research methods, we can construct some subfield codes with good parameters and optimal (or almost optimal) from some know good linear codes. Inspired and motivated by [1, 2], we study the subfield codes of linear codes over $\text{GF}(q)$ with q is a prime power.

In the rest of this paper. Section 2 introduces some basic properties of characters, character sums, Norms function, Weil sums, subfield codes and Pless power moments over finite fields. Section 3 studies the parameters of weight distribution of two families of subfield codes from linear codes and their dual codes. The proofs are presented in Section 4. In section 5, we summarized.

2. Preliminaries

In this section, we recall the properties of characters, character sums, Weil sums, subfield codes and the Pless power moments over finite fields which will be needed in later sections.

2.1. The properties of characters and Weil sums

Let $q = p^m$ be a prime power, $\text{GF}(q)$ be the finite field with q elements and α be a primitive element of $\text{GF}(q)$. Let $\text{Tr}_{q/p}$ denote the trace function from $\text{GF}(q)$ to $\text{GF}(p)$ given by $\text{Tr}_{q/p}(x) = \sum_{i=0}^{m-1} x^{p^i}$ for all $x \in \text{GF}(q)$. Let $N_{q/p}$ denote the norm function from $\text{GF}(q)^*$ to $\text{GF}(p)^*$, which is represented as

$$N_{q/p}(x) = x \cdot x^p \cdot x^{p^2} \cdots x^{p^{m-1}} = x^{\frac{p^m-1}{p-1}} \text{ for all } x \in \text{GF}(q)^*$$

An additive character of $\text{GF}(q)$ is a nonzero function χ from $\text{GF}(q)$ to \mathbb{C}^* , such that

$$\chi(x+y) = \chi(x) + \chi(y), \text{ for all } x, y \in \text{GF}(q),$$

where \mathbb{C}^* denotes the set of all nonzero complex numbers. For any $a \in \text{GF}(q)$, the function

$$\chi_a(x) = \zeta_p^{\text{Tr}_{q/p}(ax)}, x \in \text{GF}(q),$$

define an additive character of $\text{GF}(q)$, where ζ_p is the primitive p -th root of complex unity. If $a = 0$, $\chi_0(x) = 1$ for all $x \in \text{GF}(q)$, we call it the trivial additive character of $\text{GF}(q)$. If $a = 1$, we refer χ_1 as the canonical additive character of $\text{GF}(q)$. Obviously, $\chi_a(x) = \chi_1(ax)$ for all $x \in \text{GF}(q)$. The orthogonality relation of additive characters is given as follows

$$\sum_{x \in \text{GF}(q)} \chi_a(x) = \begin{cases} q & \text{for } a = 0, \\ 0 & \text{for } a \in \text{GF}(q)^*. \end{cases}$$

Let $\text{GF}(q)^* = \text{GF}(q) \setminus \{0\}$. A multiplicative character of $\text{GF}(q)^*$ is a nonzero function ψ from $\text{GF}(q)^*$ to \mathbb{C}^* such that $\psi(xy) = \psi(x)\psi(y)$ for all $x, y \in \text{GF}(q)^*$. For any $a \in \text{GF}(q)^*$, there is a integer k , such that $a = \alpha^k$. All the multiplicative characters are given by

$$\psi_j(\alpha^k) = \zeta_{q-1}^{jk} \text{ for } k = 0, \dots, q-1,$$

where $0 \leq j \leq q-2$ and $\zeta_{q-1} = e^{\frac{2\pi i}{q-1}}$. Furthermore, $\{\psi_j : j = 0, 1, \dots, q-2\}$ is multiplicative cyclic group, ψ_0 is called the trivial multiplicative character. Let $\eta := \psi_{(q-1)/2}$, and is called the quadratic multiplicative character of $\text{GF}(q)$. The orthogonality relation of multiplicative characters is given by

$$\sum_{x \in \text{GF}(q)^*} \psi_j(x) = \begin{cases} q-1 & \text{for } j = 0, \\ 0 & \text{for } j \neq 0. \end{cases}$$

Let χ be a additive character of $\text{GF}(q)$, and ψ be a multiplicative character of $\text{GF}(q)^*$. The Gaussian sum $G(\chi, \psi)$ is given by

$$G(\chi, \psi) = \sum_{x \in \text{GF}(q)^*} \psi(x)\chi(x),$$

where $G(\eta, \chi)$ is called the quadratic sum over $\text{GF}(q)$ and χ is nontrivial [14].

Lemma 2.1 ([14], Th. 5.15). *Let $q = p^m$ with p odd. χ be the canonical additive character of $GF(q)$. Then*

$$\begin{aligned} G(\chi, \psi) &= (-1)^{m-1} (\sqrt{-1})^{(\frac{p-1}{2})^2 m} \sqrt{q} \\ &= \begin{cases} (-1)^{m-1} \sqrt{q} & \text{for } p \equiv 1 \pmod{4}, \\ (-1)^{m-1} (\sqrt{-1})^m \sqrt{q} & \text{for } p \equiv 3 \pmod{4}. \end{cases} \end{aligned}$$

Let χ be a canonical additive character of $GF(q)$, and $f(x)$ is a polynomial over $GF(q)$ with positive degree. The character sums is given by

$$\sum_{x \in GF(q)} \chi(f(x))$$

is called Weil sums. If q is odd, the Weil sums is described in the following lemma.

Lemma 2.2 ([14], Th. 5.33). *Let χ be a nontrivial additive character of $GF(q)$ with q odd, and Let $f(x) = a_2 x^2 + a_1 x + a_0 \in GF(q)[x]$ with $a_2 \neq 0$. Then*

$$\sum_{x \in F_q} \chi(f(x)) = \chi(a_0 - a_1^2 (4a_2)^{-1}) \eta(a_2) G(\eta, \chi).$$

2.2. Pless power moments

Let C is an $[n, k, d]$ linear code over $GF(p)$ with the weight distribution $(1, A_1, \dots, A_n)$, and A_i be the number of codeword of weight i for all $i = 0, 1, \dots, n$, the weight distribution of dual code C^\perp is denoted by $(1, A_1^\perp, \dots, A_n^\perp)$. Then the first four Pless power moments are given as follows [13]:

$$\begin{aligned} \sum_{i=0}^n A_i &= p^k; \\ \sum_{i=0}^n i A_i &= p^{k-1} (pn - n - A_1^\perp); \\ \sum_{i=0}^n i^2 A_i &= p^{k-2} [(p-1)n(pn - n + 1) - (2pn - p - 2n + 2)A_1^\perp + 2A_2^\perp]; \\ \sum_{i=0}^n i^3 A_i &= p^{k-3} [(p-1)n(p^2 n^2 - 2pn^2 + 3pn - p + n^2 - 3n + 2) - (3p^2 n^2 - 3p^2 n - 6pn^2 + 12pn \\ &\quad + p^2 - 6p + 3n^2 - 9n + 6)A_1^\perp + 6(pn - p - n + 2)A_2^\perp - 6A_3^\perp]. \end{aligned}$$

The following lemma give sphere packing bound.

Lemma 2.3 ([13], Th. 1.12.1). *(Sphere Packing bound) Let C be a q -ary $[n, k, d]$ code. Then*

$$q^n \geq q^k \sum_{i=0}^{\lfloor \frac{d-1}{2} \rfloor} (q-1)^i \binom{n}{i}$$

2.3. The subfield code from linear code over $GF(q)$

Let $GF(q)^m$ be a finite fields with q^m elements. Let C be an $[n, k, d]$ linear code over $GF(q)^m$ with generator matrix G , we can construct a new $[n, k', d']$ linear code over $GF(q)$ by the following method.

Take a basis of $GF(q)^m$ over $GF(q)$. Represent each entry of G as an $m \times 1$ column vector of $GF(q)^m$ with respect to this basis, and replace each entry of G with the corresponding $m \times 1$ column vector of $GF(q)^m$. In this way, G is modified into a $km \times n$ matrix over $GF(q)$, which generates the new subfield code $C^{(q)}$ over $GF(q)$ with length n . The subfield codes $C^{(q)}$ is independent of both the choice of the basis of $GF(q)^m$ over $GF(q)$ and the choice of the generator matrix G of C (see Theorems 2.1 and 2.6 in [10]).

The following two theorems give the trace representation of linear code C , which is called subfield code and a relation between the minimal distance of C^\perp and the minimal distance of $C^{(p)\perp}$, respectively.

Theorem 2.4 ([10], Th. 2.5). *Let C be an $[n, k]$ linear code over $GF(q)$. Let $G = [g_{ij}]_{1 \leq i \leq k, 1 \leq j \leq n}$ be a generator matrix of C . Then the trace representation of the subfield code $C^{(p)}$ is given by*

$$C^{(p)} = \left\{ \left(\text{Tr}_{q/p} \left(\sum_{i=1}^k a_i g_{i1} \right), \dots, \text{Tr}_{q/p} \left(\sum_{i=1}^k a_i g_{in} \right) \right) : a_1, \dots, a_k \in GF(q) \right\},$$

Theorem 2.5. *Let C be an $[n, k]$ linear code over $GF(q)$. Then the minimal distance d^\perp of C^\perp and the minimal distance $d^{(p)\perp}$ of $C^{(p)\perp}$ satisfy $d^{(p)\perp} \geq d^\perp$.*

3. The subfield codes from two families of linear codes

Let $q = p^m$ with p be a prime, and $U = \{x^2 : x \in GF(q)\}$. Let $f_{11}(x, y) = x$, $f_{12}(x, y) = y$, $f_{13}(x, y) = 1$, $f_{21}(x, y) = \text{Norm}(x)$, $f_{22}(x, y) = y$, $f_{23}(x, y) = 1$ be 2-variable polynomials over $GF(q)$. Denote

$$G_{x,y} = \begin{pmatrix} x \\ y \\ 1 \end{pmatrix}_{x \in U, y \in GF(q)} \quad \text{and} \quad \tilde{G}_{x,y} = \begin{pmatrix} \text{Norm}(x) \\ y \\ 1 \end{pmatrix}_{x \in GF(q)^*, y \in GF(q)},$$

which are $3 \times \frac{p^{2m} + p^m}{2}$ matrix and $3 \times q(q-1)$ matrix over $GF(q)$, respectively. Let

$$G_1 = \begin{pmatrix} z & 1 & 0 \\ y & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}_{x \in U, y \in GF(q)} \quad \text{and} \quad G_2 = \begin{pmatrix} \text{Norm}(x) & 1 & 0 \\ y & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}_{x \in GF(q)^*, y \in GF(q)}.$$

Two types of linear codes C_1 and C_2 over $GF(q)$ are defined, where their generator matrix are G_1 and G_2 , respectively.

The subfield codes of C_1 and C_2 are described as $C_1^{(p)}$ and $C_2^{(p)}$, respectively. By Theorem 2.4, the trace representations are given by

$$C_1^{(p)} = \{((Tr_{q/p}(ax + by) + c)_{x \in U, y \in GF(q)}, Tr_{q/p}(a), Tr_{q/p}(b)) : (a, b) \in GF(q)^2, c \in GF(p)\}$$

and

$$C_2^{(p)} = \{((Tr_{q/p}(a\text{Norm}(x) + by) + c)_{x \in GF(q)^*, y \in GF(q)}, Tr_{q/p}(a), Tr_{q/p}(b)) : a, b \in GF(q), c \in GF(p)\}$$

In the following content, our goal is to investigate the parameters and weight distribution of subfield codes $C_1^{(p)}$ with p odd, and the parameters and weight distribution of $C_2^{(p)}$ with p is prime and $m = 2$. The following lemma will be used later.

Lemma 3.1. [2] *Let $q = p^m$ be an odd prime power. Then the following hold.*

$$\begin{aligned} \#\{a \in GF(q)^* : \eta(a) = 1 \text{ and } Tr_{q/p}(a) = 0\} &= \begin{cases} \frac{p^{m-1} - 1 - (p-1)p^{\frac{m-2}{2}}(-1)^{\frac{(p-1)m}{4}}}{2} & \text{for even } m, \\ \frac{p^{m-1} - 1}{2} & \text{for odd } m. \end{cases} \\ \#\{a \in GF(q)^* : \eta(a) = 1 \text{ and } Tr_{q/p}(a) \neq 0\} &= \begin{cases} \frac{(p-1)(p^{m-1} + p^{\frac{m-2}{2}}(-1)^{\frac{(p-1)m}{4}})}{2} & \text{for even } m, \\ \frac{p^{m-1}(p-1)}{2} & \text{for odd } m. \end{cases} \\ \#\{a \in GF(q)^* : \eta(a) = -1 \text{ and } Tr_{q/p}(a) = 0\} &= \begin{cases} \frac{p^{m-1} - 1 + (p-1)p^{\frac{m-2}{2}}(-1)^{\frac{(p-1)m}{4}}}{2} & \text{for even } m, \\ \frac{p^{m-1} - 1}{2} & \text{for odd } m. \end{cases} \\ \#\{a \in GF(q)^* : \eta(a) = -1 \text{ and } Tr_{q/p}(a) \neq 0\} &= \begin{cases} \frac{(p-1)(p^{m-1} - p^{\frac{m-2}{2}}(-1)^{\frac{(p-1)m}{4}})}{2} & \text{for even } m, \\ \frac{p^{m-1}(p-1)}{2} & \text{for odd } m. \end{cases} \end{aligned}$$

The following two theorems is the main results of this paper, which give the parameters and weight distributions of subfield codes $C_1^{(p)}$ and $C_2^{(p)}$, respectively, the proofs will be presented in Section 4.

Theorem 3.2. *Let $q = p^m$ with p odd and m be a positive integer. We have the following results.*

Table 1 *The weight distribution of $C_1^{(p)}$ with m odd.*

Weight	Multiplicity
0	1
$\frac{p^{2m}+p^m}{2}$	$p-1$
$\frac{p^{2m}+p^m-p^{2m-1}-p^{m-1}}{2}$	$(p^{m-1}-1)p^m$
$\frac{p^{2m}+p^m-p^{2m-1}-p^{m-1}}{2} + 1$	$(p^m-p^{m-1})(2p^m-p)$
$\frac{p^{2m}+p^m-p^{2m-1}-p^{m-1}}{2} + 2$	$p(p^m-p^{m-1})^2$
$\frac{p^{2m}-p^{2m-1}}{2}$	$p^{m-1}-1$
$\frac{p^{2m}-p^{2m-1}}{2} + 1$	p^m-p^{m-1}
$\frac{p^{2m}+p^m-p^{2m-1}-p^{\frac{3m-1}{2}}}{2} (-1)^{\frac{(m+1)(p-1)}{4}}$	$\frac{(p^{m-1}-1)(p-1)}{2}$
$\frac{p^{2m}+p^m-p^{2m-1}-p^{\frac{3m-1}{2}}}{2} (-1)^{\frac{(m+1)(p-1)}{4}} + 1$	$\frac{p^{m-1}(p-1)^2}{2}$
$\frac{p^{2m}+p^m-p^{2m-1}+p^{\frac{3m-1}{2}}}{2} (-1)^{\frac{(m+1)(p-1)}{4}}$	$\frac{(p^{m-1}-1)(p-1)}{2}$
$\frac{p^{2m}+p^m-p^{2m-1}+p^{\frac{3m-1}{2}}}{2} (-1)^{\frac{(m+1)(p-1)}{4}} + 1$	$\frac{p^{m-1}(p-1)^2}{2}$

Table 2 The weight distribution of $C_1^{(p)}$ with m even.

Weight	Multiplicity
0	1
$\frac{p^{2m}+p^m}{2}$	$p-1$
$\frac{p^{2m}+p^m-p^{2m-1}-p^{m-1}}{2}$	$(p^{m-1}-1)p^m$
$\frac{p^{2m}+p^m-p^{2m-1}-p^{m-1}}{2} + 1$	$(p^m-p^{m-1})(2p^m-p)$
$\frac{p^{2m}+p^m-p^{2m-1}-p^{m-1}}{2} + 2$	$p(p^m-p^{m-1})^2$
$\frac{p^{2m}-p^{2m-1}+(p-1)p^{\frac{3m-2}{2}}}{2} (-1)^{\frac{m(p-1)}{4}}$	$\frac{p^{m-1}-1-(p-1)p^{\frac{m-2}{2}}(\sqrt{-1})^{\frac{(p-1)m}{2}}}{2}$
$\frac{p^{2m}-p^{2m-1}+(p-1)p^{\frac{3m-2}{2}}}{2} (-1)^{\frac{m(p-1)}{4}} + 1$	$\frac{(p-1)(p^{m-1}+p^{\frac{m-2}{2}}(\sqrt{-1})^{\frac{(p-1)m}{2}})}{2}$
$\frac{p^{2m}-p^{2m-1}-(p-1)p^{\frac{3m-2}{2}}}{2} (-1)^{\frac{m(p-1)}{4}}$	$\frac{p^{m-1}-1+(p-1)p^{\frac{m-2}{2}}(\sqrt{-1})^{\frac{(p-1)m}{2}}}{2}$
$\frac{p^{2m}-p^{2m-1}-(p-1)p^{\frac{3m-2}{2}}}{2} (-1)^{\frac{m(p-1)}{4}} + 1$	$\frac{(p-1)(p^{m-1}-p^{\frac{m-2}{2}}(\sqrt{-1})^{\frac{(p-1)m}{2}})}{2}$
$\frac{p^{2m}+p^m-p^{2m-1}-p^{\frac{3m-2}{2}}}{2} (-1)^{\frac{m(p-1)}{4}}$	$\frac{(p-1)(p^{m-1}-1-(p-1)p^{\frac{m-2}{2}}(\sqrt{-1})^{\frac{(p-1)m}{2}})}{2}$
$\frac{p^{2m}+p^m-p^{2m-1}-p^{\frac{3m-2}{2}}}{2} (-1)^{\frac{m(p-1)}{4}} + 1$	$\frac{(p-1)^2(p^{m-1}+p^{\frac{m-2}{2}}(\sqrt{-1})^{\frac{(p-1)m}{2}})}{2}$
$\frac{p^{2m}+p^m-p^{2m-1}+p^{\frac{3m-2}{2}}}{2} (-1)^{\frac{m(p-1)}{4}}$	$\frac{(p-1)(p^{m-1}-1+(p-1)p^{\frac{m-2}{2}}(\sqrt{-1})^{\frac{(p-1)m}{2}})}{2}$
$\frac{p^{2m}+p^m-p^{2m-1}+p^{\frac{3m-2}{2}}}{2} (-1)^{\frac{m(p-1)}{4}} + 1$	$\frac{(p-1)^2(p^{m-1}-p^{\frac{m-2}{2}}(\sqrt{-1})^{\frac{(p-1)m}{2}})}{2}$

(1) If $m = 1$ and p odd, then $C_1^{(p)}$ is $[\frac{p^2+p}{2} + 2, 3, \frac{p^2-p+2}{2}]$ code, and $C_1^{(p)\perp}$ is an almost MDS $[\frac{p^2+p}{2} + 2, \frac{p^2+p}{2} - 1, 3]$ code. If p is odd and $m > 1$, then $C_1^{(p)}$ is a $[\frac{p^{2m}+p^m}{2} + 2, 2m +$

- 1, $\frac{p^{2m}+p^m-p^{2m-1}-p^{\frac{3m-1}{2}}}{2}$] code. For odd m , the weight distribution of $C_1^{(p)}$ is given in Table 1.
- (2) If $m \geq 2$ is even, then $C_1^{(p)}$ is a $[\frac{p^{2m}+p^m}{2} + 2, 2m + 1, \frac{p^{2m}-p^{2m-1}-(p-1)p^{\frac{3m-2}{2}}}{2}]$ code, and its weight distribution is given in Table 2.
- (3) For any positive integer m , the dual $C_1^{(p)\perp}$ of $C_1^{(p)}$ is a dimension-optimal $[\frac{p^{2m}+p^m}{2} + 2, \frac{p^{2m}+p^m}{2} - 2m + 1, 3]$ code with respect to the sphere-packing bound and a best known code according to <http://www.codetables.de>.

Theorem 3.3. Let $q = p^m$ with p be a prime. Let $m = 2$. We have the following results.

Table 3 The weight distribution of $C_2^{(p)}$.

Weight	Multiplicity
0	1
$p^2(p^2 - 1)$	$p - 1$
$p^2(p^2 - 1) + 1$	$p - 1$
$(p^2 - p)(p^2 - 1)$	$p(p - 1)$
$(p^2 - p)(p^2 - 1) + 1$	$p(p - 1)(2p - 1)$
$(p^2 - p)(p^2 - 1) + 2$	$(p^2 - p)^2$
$p^2(p + 1)(p - 2) + 1$	$(p - 1)^2$

- (1) For any prime p and $m = 2$, $C_2^{(p)}$ is a $[p^2(p^2 - 1) + 2, 4, p^2(p + 1)(p - 2) + 1]$ code.
- (2) If $p = 2$, then the dual $C_2^{(2)\perp}$ is a $[14, 10, 2]$ almost optimal code according to the tables of best code known maintained at <http://www.codetables.de>. If $p \geq 3$ is an odd, then the dual $C_2^{(p)}$ is a $[p^2(p^2 - 1) + 2, p^2(p^2 - 1) - 2, 2]$ distance-optimal code with respect to the sphere-packing bound.

4. Proofs of the results

In this section, our main work is to prove two theorems we get in Section 3.

Proof. Theorem 3.2. The subfield code of $C_1^{(p)}$ is given by

$$C_1^{(p)} = \{((Tr_{q/p}(ax + by) + c)_{x \in U, y \in GF(q)}, Tr_{q/p}(a), Tr_{q/p}(b)) : (a, b) \in GF(q)^2, c \in GF(p)\}$$

Let χ and χ' are the canonical additive characters of $GF(q)$ and $GF(p)$, respectively. Let η and η' be the quadratic multiplicative characters of $GF(q)^*$ and $GF(p)^*$, respectively. For each

$a, b \in \text{GF}(q), c \in \text{GF}(p)$ and p odd, denote

$$\begin{aligned} N_0(a, b, c) &= \#\{(x, y) : x \in U, y \in \text{GF}(q) : \text{Tr}_{q/p}(ax + by) + c = 0\}, U = \{x^2 : x \in \text{GF}(q)\}, \\ N'_0(a, b, c) &= \#\{(x, y) \in \text{GF}(q)^2 : \text{Tr}_{q/p}(ax^2 + by) + c = 0\}, \\ N''_0(a, b, c) &= \#\{y : x = 0, y \in \text{GF}(q) : \text{Tr}_{q/p}(by) + c = 0\}. \end{aligned}$$

Through analysis, we conclude that

$$N_0(a, b, c) = \frac{N'_0(a, b, c) + N''_0(a, b, c)}{2}. \quad (1)$$

In order to calculate $N_0(a, b, c)$, we just need to calculate $N'_0(a, b, c)$ and $N''_0(a, b, c)$.

By the orthogonality relation of additive characters,

$$\begin{aligned} N'_0(a, b, c) &= \frac{1}{p} \sum_{\alpha \in \text{GF}(p)} \sum_{(x, y) \in \text{GF}(p)^2} \zeta_p^{\alpha(\text{Tr}_{q/p}(ax^2 + by) + c)} \\ &= p^{2m-1} + \frac{1}{p} \sum_{\alpha \in \text{GF}(p)^*} \sum_{(x, y) \in \text{GF}(q)^2} \zeta_p^{\alpha(\text{Tr}_{q/p}(ax^2 + by) + c)} \\ &= p^{2m-1} + \frac{1}{p} \Gamma(a, b, c), \end{aligned} \quad (2)$$

where $\Gamma(a, b, c) = \sum_{\alpha \in \text{GF}(p)^*} \sum_{(x, y) \in \text{GF}(q)^2} \zeta_p^{\alpha(\text{Tr}_{q/p}(ax^2 + by) + c)}$. We discuss the value of $\Gamma(a, b, c)$ in the following cases.

- (1) If $(a, b, c) = (0, 0, 0)$. Then $\Gamma(a, b, c) = p^{2m}(p - 1)$.
- (2) If $(a, b) = (0, 0)$ and $c \neq 0$. Then $\Gamma(a, b, c) = p^{2m} \sum_{\alpha \in \text{GF}(p)^*} \zeta_p^{\alpha c} = -p^{2m}$.
- (3) If $a = 0$ and $b \neq 0$ or $a \neq 0$ and $b \neq 0$. Then $\Gamma(a, b, c) = 0$.
- (4) If $a \neq 0$ and $b = 0$. We have

$$\begin{aligned} \Gamma(a, b, c) &= q \cdot \sum_{\alpha \in \text{GF}(p)^*} \zeta_p^{\alpha c} \sum_{x \in \text{GF}(q)} \chi(a\alpha x^2) \\ &= q \cdot \eta(a) G(\eta, \chi) \sum_{\alpha \in \text{GF}(p)^*} \zeta_p^{\alpha c} \eta(\alpha) \\ &= q \cdot \eta(a) G(\eta, \chi) \sum_{\alpha \in \text{GF}(p)^*} \chi'(\alpha c) \eta(\alpha). \end{aligned}$$

(a) If m is odd, we have $\eta(\alpha) = \eta'(\alpha)$ for any $\alpha \in \text{GF}(p)^*$. Then by Lemma 3.1 we have

$$\begin{aligned}\Gamma(a, b, c) &= q \cdot \eta(a) G(\eta, \chi) \sum_{\alpha \in \text{GF}(p)^*} \chi'(\alpha c) \eta'(\alpha) \\ &= \begin{cases} 0 & \text{if } c = 0 \\ q \cdot G(\eta, \chi) G(\eta', \chi') \eta(a) \eta'(c) & \text{if } c \neq 0 \end{cases} \\ &= \begin{cases} 0 & \text{if } c = 0, \\ q \cdot p^{\frac{m+1}{2}} (-1)^{\frac{(p-1)(m+1)}{4}} & \text{if } c \neq 0, \eta(a) \eta'(c) = 1, \\ q \cdot p^{\frac{m+1}{2}} (-1)^{\frac{(p-1)(m+1)+4}{4}} & \text{if } c \neq 0, \eta(a) \eta'(c) = -1. \end{cases}\end{aligned}$$

(b) If m is even, then $\eta(\alpha) = 1$ for any $\alpha \in \text{GF}(p)^*$. Then by Lemma 3.1 we have

$$\begin{aligned}\Gamma(a, b, c) &= q \cdot G(\eta, \chi) \eta(a) \sum_{\alpha \in \text{GF}(p)^*} \chi'(\alpha c) \\ &= \begin{cases} q \cdot (p-1) p^{\frac{m}{2}} (-1)^{\frac{m(p-1)+4}{4}} & \text{if } c = 0, \eta(a) = 1, \\ q \cdot (p-1) p^{\frac{m}{2}} (-1)^{\frac{m(p-1)}{4}} & \text{if } c = 0, \eta(a) = -1, \\ q \cdot p^{\frac{m}{2}} (-1)^{\frac{m(p-1)}{4}} & \text{if } c \neq 0, \eta(a) = 1, \\ q \cdot p^{\frac{m}{2}} (-1)^{\frac{m(p-1)+4}{4}} & \text{if } c \neq 0, \eta(a) = -1. \end{cases}\end{aligned}$$

By equation (2) and the discussions above, if m is odd, we conclude that

$$N'_0(a, b, c) = \begin{cases} p^{2m} & \text{for } (a, b, c) = (0, 0, 0), \\ 0 & \text{for } (a, b) = (0, 0), c \neq 0, \\ p^{2m-1} & \text{for } b \neq 0, \text{ or } a \neq 0, b = 0, c = 0, \\ p^{2m-1} + p^{\frac{3m-1}{2}} (-1)^{\frac{(p-1)(m+1)}{4}} & \text{for } a \neq 0, b = 0, c \neq 0, \eta(a) \eta'(c) = 1, \\ p^{2m-1} + p^{\frac{3m-1}{2}} (-1)^{\frac{(p-1)(m+1)+4}{4}} & \text{for } a \neq 0, b = 0, c \neq 0, \eta(a) \eta'(c) = -1, \end{cases} \quad (3)$$

if m is even, then

$$N'_0(a, b, c) = \begin{cases} p^{2m} & \text{for } (a, b, c) = (0, 0, 0), \\ 0 & \text{for } (a, b) = (0, 0), c \neq 0, \\ p^{2m-1} & \text{for } a = 0, b \neq 0, \text{ or } a \neq 0, b \neq 0, \\ p^{2m-1} + (p-1) p^{\frac{3m-2}{2}} (-1)^{\frac{m(p-1)+4}{4}} & \text{for } a \neq 0, b = 0, c = 0, \eta(a) = 1, \\ p^{2m-1} + (p-1) p^{\frac{3m-2}{2}} (-1)^{\frac{m(p-1)}{4}} & \text{for } a \neq 0, b = 0, c = 0, \eta(a) = -1, \\ p^{2m-1} + p^{\frac{3m-2}{2}} (-1)^{\frac{m(p-1)}{4}} & \text{for } a \neq 0, b = 0, c \neq 0, \eta(a) = 1, \\ p^{2m-1} + p^{\frac{3m-2}{2}} (-1)^{\frac{m(p-1)+4}{4}} & \text{for } a \neq 0, b = 0, c \neq 0, \eta(a) = -1, \end{cases} \quad (4)$$

By the orthogonality relation of additive characters,

$$N_0''(a, b, c) = \begin{cases} p^m & \text{for } (b, c) = (0, 0), \\ 0 & \text{for } b = 0, c \neq 0, \\ p^{m-1} & \text{for } b \neq 0, \end{cases} \quad (5)$$

by equation (1), (3)-(5) and discussions above, if m is odd, we have

$$N_0(a, b, c) = \begin{cases} \frac{p^{2m} + p^m}{2} & \text{for } (a, b, c) = (0, 0, 0), \\ 0 & \text{for } (a, b) = (0, 0), c \neq 0, \\ \frac{p^{2m-1} + p^{m-1}}{2} & \text{for } a = 0, b \neq 0, \text{ or } a \neq 0, b \neq 0, \\ \frac{p^{2m-1} + p^m}{2} & \text{for } a \neq 0, b = 0, c = 0, \\ \frac{p^{2m-1} + p^{\frac{3m-1}{2}} (-1)^{\frac{(p-1)(m+1)}{4}}}{2} & \text{for } a \neq 0, b = 0, c \neq 0, \eta(a)\eta'(c) = 1, \\ \frac{p^{2m-1} - p^{\frac{3m-1}{2}} (-1)^{\frac{(p-1)(m+1)}{4}}}{2} & \text{for } a \neq 0, b = 0, c \neq 0, \eta(a)\eta'(c) = -1, \end{cases} \quad (6)$$

if m is even, then

$$N_0(a, b, c) = \begin{cases} \frac{p^{2m} + p^m}{2} & \text{for } (a, b, c) = (0, 0, 0), \\ 0 & \text{for } (a, b) = (0, 0), c \neq 0, \\ \frac{p^{2m-1} + p^{m-1}}{2} & \text{for } a = 0, b \neq 0, \text{ or } a \neq 0, b \neq 0, \\ \frac{p^{2m-1} + p^m - (p-1)p^{\frac{3m-2}{2}} (-1)^{\frac{m(p-1)}{4}}}{2} & \text{for } a \neq 0, b = 0, c = 0, \eta(a) = 1, \\ \frac{p^{2m-1} + p^m + (p-1)p^{\frac{3m-2}{2}} (-1)^{\frac{m(p-1)}{4}}}{2} & \text{for } a \neq 0, b = 0, c = 0, \eta(a) = -1, \\ \frac{p^{2m-1} + p^{\frac{3m-2}{2}} (-1)^{\frac{m(p-1)}{4}}}{2} & \text{for } a \neq 0, b = 0, c \neq 0, \eta(a) = 1, \\ \frac{p^{2m-1} - p^{\frac{3m-2}{2}} (-1)^{\frac{m(p-1)}{4}}}{2} & \text{for } a \neq 0, b = 0, c \neq 0, \eta(a) = -1. \end{cases} \quad (7)$$

For any codeword $c_1(a, b, c)$ of $C_1^{(p)}$ with m is odd, we have

$$c_1(a, b, c) = ((Tr_{q/p}(ax + by) + c)_{x \in U, y \in GF(q)}, Tr_{q/p}(a), Tr_{q/p}(b))$$

by equation (6) the Hamming distance of codeword $c_1(a, b, c)$ is given by

$$wt(c_1(a, b, c))$$

$$= \begin{cases} 0 & \text{for } (a, b, c) = (0, 0, 0), \\ \frac{p^{2m} + p^m}{2} & \text{for } (a, b) = (0, 0), c \neq 0, \\ \frac{p^{2m} + p^m - p^{2m-1} - p^{m-1}}{2} & \text{for } a = 0, b \neq 0, Tr_{q/p}(b) = 0, \\ \frac{p^{2m} + p^m - p^{2m-1} - p^{m-1}}{2} + 1 & \text{for } a = 0, b \neq 0, Tr_{q/p}(b) \neq 0, \\ \frac{p^{2m} + p^m - p^{2m-1} - p^{m-1}}{2} & \text{for } a \neq 0, b \neq 0, Tr_{q/p}(a) = Tr_{q/p}(b) = 0, \\ \frac{p^{2m} + p^m - p^{2m-1} - p^{m-1}}{2} + 1 & \text{for } a \neq 0, b \neq 0, Tr_{q/p}(a) = 0, Tr_{q/p}(b) \neq 0, \\ \frac{p^{2m} + p^m - p^{2m-1} - p^{m-1}}{2} + 1 & \text{for } a \neq 0, b \neq 0, Tr_{q/p}(a) \neq 0, Tr_{q/p}(b) = 0, \\ \frac{p^{2m} + p^m - p^{2m-1} - p^{m-1}}{2} + 2 & \text{for } a \neq 0, b \neq 0, Tr_{q/p}(a) \neq 0, Tr_{q/p}(b) \neq 0, \\ \frac{p^{2m} - p^{2m-1}}{2} & \text{for } a \neq 0, b = 0, c = 0, Tr_{q/p}(a) = 0, \\ \frac{p^{2m} - p^{2m-1}}{2} + 1 & \text{for } a \neq 0, b = 0, c \neq 0, Tr_{q/p}(a) \neq 0, \\ \frac{p^{2m} + p^m - p^{2m-1} - p^{\frac{3m-1}{2}}(-1)^{\frac{(p-1)(m+1)}{4}}}{2} & \text{for } a \neq 0, b = 0, c \neq 0, \eta(a)\eta'(c) = 1, Tr_{q/p}(a) = 0, \\ \frac{p^{2m} + p^m - p^{2m-1} - p^{\frac{3m-1}{2}}(-1)^{\frac{(p-1)(m+1)}{4}}}{2} + 1 & \text{for } a \neq 0, b = 0, c \neq 0, \eta(a)\eta'(c) = 1, Tr_{q/p}(a) \neq 0, \\ \frac{p^{2m} + p^m - p^{2m-1} + p^{\frac{3m-1}{2}}(-1)^{\frac{(p-1)(m+1)}{4}}}{2} & \text{for } a \neq 0, b = 0, c \neq 0, \eta(a)\eta'(c) = -1, Tr_{q/p}(a) = 0, \\ \frac{p^{2m} + p^m - p^{2m-1} + p^{\frac{3m-1}{2}}(-1)^{\frac{(p-1)(m+1)}{4}}}{2} + 1 & \text{for } a \neq 0, b = 0, c \neq 0, \eta(a)\eta'(c) = -1, Tr_{q/p}(a) \neq 0, \end{cases}$$

with 1 times,
with $p - 1$ times,
with $(p^{m-1} - 1)p^m$ times,
with $(p^m - p^{m-1})(2p^m - p)$ times,
with $p(p^m - p^{m-1})^2$ times,
with $p^{m-1} - 1$ times,
with $p^m - p^{m-1}$ times
with $\frac{(p^{m-1}-1)(p-1)}{2}$ times,
with $\frac{p^{m-1}(p-1)^2}{2}$ times,
with $\frac{(p^{m-1}-1)(p-1)}{2}$ times,
with $\frac{p^{m-1}(p-1)^2}{2}$ times,

when (a, b, c) runs through $GF(q) \times GF(q) \times GF(p)$. Let a linear $[n, k, d]$ code with weight distribution $(1, A_1, \dots, A_n)$, the weight distribution of its dual is $(1, A_1^\perp, \dots, A_n^\perp)$, where A_i and A_i^\perp are the number of codewords with weight i , respectively, and $p^k = \sum_{i=0}^n A_i$, k is denoted as the

dimension of $C_1^{(p)}$, then dimension of $C_1^{(p)}$ is $2m+1$ as $A_0 = 1$.

When m is even, we can get similar results and their weight distribution by equation (7) and Lemma 3.1. We omit the details here.

By discussion above, we get the parameters and weight distribution of $C_1^{(p)}$, the dual of $C_1^{(p)\perp}$ has length $\frac{p^{2m}+p^m}{2} + 2$ and dimension $\frac{p^{2m}+p^m}{2} - 2m + 1$. From the weight distribution and the first four Pless power moments, we can obtain that $A_1^{(p)\perp} = A_2^{(p)\perp} = 0$ and $A_3^{(p)\perp} > 0$. Thus, the minimal distance of $C_1^{(p)\perp}$ is $d_1^{(p)\perp} = 3$. By the sphere-packing bound, we have

$$p^{k_1^{(p)\perp}} \left(\sum_{i=0}^{\lfloor \frac{3-1}{2} \rfloor} (p-1)^i \binom{\frac{p^{2m}+p^m}{2} + 2}{i} \right) \leq p^{\frac{p^{2m}+p^m}{2} + 2},$$

which implies that $k_1^{(p)\perp} \leq \frac{p^{2m}+p^m}{2} - 2m + 1$. Thus, the dual codes $C_1^{(p)\perp}$ are dimension-optimal according to the sphere-packing bound. We have completed the proof of Theorem 3.2. \square

Example 4.1. Let $p = 3$, m be a positive integer, $C_1^{(3)}$ is a linear code over $GF(3)$, and is presented in Theorem 3.2.

(1) Let $m = 1$, $p = 3$. Then $C_1^{(3)}$ has parameters $[8, 3, 4]$ and weight enumerator

$$1 + 4x^4 + 6x^5 + 14x^6 + 2x^7,$$

it's almost optimal code according to the tables of best code known maintained at <http://www.codetables.de>.

The dual code $C_1^{(3)\perp}$ has parameters $[8, 5, 3]$, which be an almost MDS code.

(2) Let $m = 2$, $p = 3$. Then $C_1^{(3)}$ has parameters $[47, 5, 18]$ and weight enumerator

$$1 + 2x^{18} + 2x^{19} + 8x^{28} + 18x^{30} + 90x^{31} + 108x^{32} + 4x^{36} + 8x^{37} + 2x^{45},$$

where its dual code has parameters $[47, 42, 3]$, which is dimension-optimal with respect to the sphere packing bound.

Proof. Theorem 3.3. The subfield code of $C_2^{(p)}$ is given by

$$C_2^{(p)} = \{ ((Tr_{q/p}(a\text{Norm}(x) + by) + c)_{x \in GF(q)^*, y \in GF(q)}, Tr_{q/p}(a), Tr_{q/p}(b)) : a, b \in GF(q), c \in GF(p) \}.$$

For each $a, b \in GF(q)$ and $c \in GF(p)$ where p is a prime, denote

$$N_0(a, b, c) = \#\{(x, y) : x \in GF(q)^*, y \in GF(q) : Tr_{q/p}(a\text{Norm}(x) + by) + c = 0\}.$$

By the orthogonality relation of additive characters,

$$\begin{aligned} N_0(a, b, c) &= \frac{1}{p} \sum_{\alpha \in GF(p)} \sum_{x \in GF(q)^*, y \in GF(q)} \zeta_p^{\alpha(Tr_{q/p}(a\text{Norm}(x) + by) + c)} \\ &= \frac{q(q-1)}{p} + \frac{q}{p} \sum_{\alpha \in GF(p)^*} \sum_{x \in GF(q)^*, y \in GF(q)} \zeta_p^{\alpha(Tr_{q/p}(a\text{Norm}(x) + by) + c)} \\ &= \frac{q(q-1)}{p} + \frac{1}{p} \Gamma(a, b, c), \end{aligned} \tag{8}$$

where $\Gamma(a, b, c) = \sum_{\alpha \in \text{GF}(p)^*} \sum_{x \in \text{GF}(q)^*, y \in \text{GF}(q)} \zeta_p^{\alpha(\text{Tr}_{q/p}(a\text{Norm}(x)+by)+c)}$. It seems difficult to calculate the value of exponential sums for any positive integer m , but we can get the value when $m = 2$.

Let $m = 2$, we have $\text{Norm}(x) = x^{p+1}$. Next we will discuss the value of $\Gamma(a, b, c)$.

- (1) If $(a, b, c) = (0, 0, 0)$. Then $\Gamma(a, b, c) = q(q-1)(p-1)$.
- (2) If $(a, b) = (0, 0)$ and $c \neq 0$. Then $\Gamma(a, b, c) = -q(q-1)$.
- (3) If $a = 0$ and $b \neq 0$, or $a \neq 0$ and $b \neq 0$. Then $\Gamma(a, b, c) = 0$.
- (4) If $a \neq 0, \text{Tr}_{q/p}(a) = 0$ and $b = 0$. Then

$$\begin{aligned} \Gamma(a, b, c) &= q \sum_{\alpha \in \text{GF}(p)^*} \zeta_p^{\alpha c} \sum_{x \in \text{GF}(q)^*} \zeta_p^{\text{Tr}_{q/p}(a\alpha x^{p+1})} \\ &= \begin{cases} q(q-1)(p-1) & \text{if } c = 0, \\ -q(q-1) & \text{if } c \neq 0. \end{cases} \end{aligned}$$

- (5) If $a \neq 0, \text{Tr}_{q/p}(a) \neq 0$ and $b = 0$. Then

$$\Gamma(a, b, c) = \begin{cases} -q(p+1)(p-1) & \text{if } c = 0, \\ q(p+1) & \text{if } c \neq 0. \end{cases}$$

by equation (8) and discussions above, we conclude that

$$N_0(a, b, c) = \begin{cases} p^2(p^2-1) & \text{for } (a, b, c) = (0, 0, 0), \text{ or } a \neq 0, b = c = 0, \text{Tr}_{q/p}(a) = 0 \\ 0 & \text{for } (a, b) = (0, 0), c \neq 0, \text{ or } a \neq 0, b = \text{Tr}_{q/p}(a) = 0, c \neq 0, \\ & \text{or } a \neq 0, b = c = 0, \text{Tr}_{q/p}(a) \neq 0, \\ p(p^2-1) & \text{for } b \neq 0, \\ p^2(p+1) & \text{for } a \neq 0, b = 0, \text{Tr}_{q/p}(a) = 0, c \neq 0. \end{cases} \quad (9)$$

For any codeword $c_2(a, b, c)$ of $C_2^{(p)}$. By equation (9), the Hamming weight is given by

$$wt(c_2(a, b, c)) = \begin{cases} 0 & \text{for } (a, b, c) = (0, 0, 0), \\ p^2(p^2 - 1) & \text{for } (a, b) = (0, 0), c \neq 0, \\ (p^2 - p)(p^2 - 1) & \text{for } a \neq 0, \neq 0, Tr_{q/p}(a) = Tr_{q/p}(b) = 0, \\ (p^2 - p)(p^2 - 1) + 1 & \text{for } a \neq 0, \neq 0, Tr_{q/p}(a) = 0, Tr_{q/p}(b) \neq 0, \\ (p^2 - p)(p^2 - 1) + 1 & \text{for } a \neq 0, \neq 0, Tr_{q/p}(b) = 0, Tr_{q/p}(a) \neq 0, \\ (p^2 - p)(p^2 - 1) + 2 & \text{for } a \neq 0, \neq 0, Tr_{q/p}(a) \neq 0, Tr_{q/p}(b) \neq 0, \\ (p^2 - p)(p^2 - 1) & \text{for } a = 0, b \neq 0, Tr_{q/p}(b) = 0, \\ (p^2 - p)(p^2 - 1) + 1 & \text{for } a = 0, b \neq 0, Tr_{q/p}(b) \neq 0, \\ 0 & \text{for } a \neq 0, b = Tr_{q/p}(a) = c = 0, \\ p^2(p^2 - 1) & \text{for } a \neq 0, b = Tr_{q/p}(a) = 0, c \neq 0, \\ p^2(p^2 - 1) + 1 & \text{for } a \neq 0, b = c = 0, Tr_{q/p}(a) \neq 0, \\ p^2(p + 1)(p - 2) + 1 & \text{for } a \neq 0, b = 0, Tr_{q/p}(a) \neq 0, c \neq 0 \end{cases}$$

$$= \begin{cases} 0 & \text{with } p \text{ times,} \\ p^2(p^2 - 1) & \text{with } p(p - 1) \text{ times,} \\ p^2(p^2 - 1) + 1 & \text{with } p(p - 1) \text{ times,} \\ (p^2 - p)(p^2 - 1) & \text{with } p^2(p - 1) \text{ times,} \\ (p^2 - p)(p^2 - 1) + 1 & \text{with } p^2(p - 1)(2p - 1) \text{ times,} \\ (p^2 - p)(p^2 - 1) + 2 & \text{with } p(p^2 - p)^2 \text{ times,} \\ p^2(p + 1)(p - 2) + 1 & \text{with } p(p - 1)^2 \text{ times.} \end{cases}$$

when (a, b, c) runs through $GF(q) \times GF(q) \times GF(p)$. By observing the calculation above, the Hamming weight 0 occurs p times. Thus, any codeword of $C_2^{(p)}$ repeats p times. Then the dimension of $C_2^{(p)}$ is 4.

By discussion above, we get the parameters and weight distribution of $C_2^{(p)}$, the dual is a $[p^2(p^2 - 1) + 2, p^2(p^2 - 1) - 2]$ code. From the first four Pless power moments, we get $A_1^{(p)\perp} = 0$ and $A_2^{(p)\perp} > 0$. Thus, the minimal distance $d_2^{(p)\perp} = 2$. By the sphere-packing bound, we have

$$p^{p^2(p^2-1)-2} \left(\sum_{i=0}^{\lfloor \frac{d_2^{(p)\perp}-1}{2} \rfloor} (p-1)^i \binom{p^2(p^2-1)+2}{i} \right) \leq p^{p^2(p^2-1)+2},$$

which implies that $d_2^{(p)\perp} \leq 2$. Therefore, the dual code $C_2^{(p)\perp}$ is distance-optimal according to the sphere-packing bound. The proof is completed. \square

Example 4.2. Let $C_2^{(p)}$ be the linear code over $GF(p)$, and is presented in Theorem 3.3.

- (1) Let $p = 2$. Then $C_2^{(2)}$ is a $[14, 4, 1]$ code, the dual is a $[14, 10, 2]$ best known code according to <http://www.codetables.de>.
- (2) Let $p = 3$. Then $C_2^{(3)}$ is a $[74, 4, 37]$ code, the dual is a $[74, 70, 2]$ distance-optimal code with respect to the sphere-packing bound.

5. Conclusion

In this paper, we used the results of Heng and Ding in [2] and continued the work of Xiang and Yin in [1], and mainly investigated the p -ary subfield codes of a class of linear codes. At the same time, the complete weight distribution of $C^{(p)}$ is presented in Table 1, Table 2 and Table 3. Some linear subfield codes we get in this paper are almost optimal in some cases. The parameters of the dual codes presented in this paper are dimension-optimal (or distance-optimal) according to the sphere-packing bound. From the construction of subfield codes, by selecting generator matrix G of C , we can get many optimal or almost optimal subfield codes of linear codes C over $\text{GF}(p)$. It would be interesting to study the weight distributions of subfield codes with good parameters over extended fields.

References

- [1] C. Xiang, W. Yin, Two families of subfield codes with a few weights. *Cryptogr. Commun.* 13(1), 117-127 (2021).
- [2] Z. Heng, C. Ding, The subfield codes of hyperoval and conic codes. *Finite Fields Their Appl.* 56, 308-331 (2019).
- [3] Canteaut, A., Charpin, P., Dobbertin, H., Weight divisibility of cyclic codes, highly nonlinear functions on F_{2^n} , and crosscorrelation of maximum-length sequences. *SIAM Disc. Math.* 13(1), 105-138 (2000).
- [4] Carlet, C., Charpin, P., Zinoviev, V., Codes, bent functions and permutations suitable for DES-like cryptosystems. *Des. Codes Cryptogr.* 15(2), 125-156 (1998).
- [5] Cannon, J., Bosma, W., Fieker, C., Stell, E., Handbook of magma functions, version 2. 19. Sydney (2013).
- [6] Z. Heng, C. Ding, The Subfield Codes of Some $[q+1, 2, q]$ MDS Codes. *IEEE Trans. Inf. Theory.* 68(6), 3643-3656 (2022).
- [7] Z. Heng, Q. Wang and C. Ding, Two Families of Optimal Linear Codes and Their Subfield Codes. *IEEE Trans. Inf. Theory.* 66(11), 6872-6883 (2020).
- [8] X. Wang, D. Zheng and Y. Zhang, A class of subfield codes of linear codes and their duals. *Cryptogr. Commun.* 13(1), 173-196 (2021).
- [9] Z. Heng, C. Ding, Optimal binary linear codes from maximal arcs. *IEEE Trans. Inf. Theory.* 66(9), 5387-5394 (2020).
- [10] C. Ding, Z. Heng, The Subfield Codes of Ovoid Codes. *IEEE Trans. Inf. Theory.* 65(8), 4715-4729 (2019).
- [11] X. Wang, D. Zheng, The subfield codes of several classes of linear codes. *Cryptogr. Commun.* 12(6), 1111-1131 (2020).
- [12] Griesmer, J.H., A bound for error-correcting codes. *IBM J. Res. Develop.* 4(5), 532-542 (1960).
- [13] Huffman, W.C., Pless, V., Fundamental of error-correcting codes. Cambridge Univ Press, Cambridge (2003).
- [14] Lidl, R., Niederreiter, H., Finite Fields. Cambridge Univ Press, Cambridge (1997).