

Counting and Computing Join-Endomorphisms in Lattices (Revisited)^{*}

Carlos Pinzón^c, Santiago Quintero^d, Sergio Ramírez^e, Camilo Rueda^a, Frank Valencia^{a,b,*}

^a*Department of Electronics and Computer Science
Pontificia Universidad Javeriana, Cali, Colombia*

^b*CNRS, LIX École Polytechnique de Paris, France*

^c*INRIA, École Polytechnique de Paris, France*

^d*LIX École Polytechnique de Paris, France*

^e*Universidad EAFIT, Medellín, Colombia*

Abstract

Structures involving a lattice and join-endomorphisms on it are ubiquitous in computer science. We study the cardinality of the set $\mathcal{E}(L)$ of all join-endomorphisms of a given finite lattice L . In particular, we show for \mathbf{M}_n , the discrete order of n elements extended with top and bottom, $|\mathcal{E}(\mathbf{M}_n)| = n! \mathcal{L}_n(-1) + (n+1)^2$ where $\mathcal{L}_n(x)$ is the Laguerre polynomial of degree n . We also study the following problem: Given a lattice L of size n and a set $S \subseteq \mathcal{E}(L)$ of size m , find the greatest lower bound $\prod_{\mathcal{E}(L)} S$. The join-endomorphism $\prod_{\mathcal{E}(L)} S$ has meaningful interpretations in epistemic logic, distributed systems, and Aumann structures. We show that this problem can be solved with worst-case time complexity in $O(mn)$ for distributive lattices and $O(mn + n^3)$ for arbitrary lattices. In the particular case of modular lattices, we present an adaptation of the latter algorithm that reduces its average time complexity. We provide theoretical and experimental results to support this enhancement. The complexity is expressed in terms of the basic binary lattice operations performed by the algorithm.

Keywords: join-endomorphisms, lattice cardinality, lattice algorithms.

1. Introduction

There is a long established tradition of using lattices to model structural entities in many fields of mathematics and computer science. For example, lattices

^{*}This work has been partially supported by the ECOS-NORD project FACTS (C19M03)

^{*}Corresponding author

Email addresses: carlos.pinzon@inria.fr (Carlos Pinzón), squinter@polytechnique.fr (Santiago Quintero), ssramirezr@eafit.edu.co (Sergio Ramírez), crueda@javerianacali.edu.co (Camilo Rueda), frank.valencia@lix.polytechnique.fr (Frank Valencia)

are used in concurrency theory to represent the hierarchical organization of the information resulting from agent's interactions [1]. *Mathematical morphology* (MM), a well-established theory for the analysis and processing of geometrical structures, is founded upon lattice theory [2, 3]. Lattices are also used as algebraic structures for modal and epistemic logics as well as Aumann structures (e.g., modal algebras and constraint systems [4]).

In all these and many other applications, lattice join-endomorphisms appear as fundamental. A *join-endomorphism* is a function from a lattice to itself that preserves finite joins. In MM, join-endomorphisms correspond to one of its fundamental operations; *dilations*. In modal algebra, they correspond via duality to the box modal operator. In epistemic settings, they represent belief or knowledge of agents. In fact, our own interest in lattice theory derives from using join-endomorphisms to model the perception that agents may have of a statement in a lattice of partial information [4].

For finite lattices, devising suitable algorithms to compute lattice maps with some given properties would thus be of great utility. We are interested in constructing algorithms for computing lattice join-endomorphisms. This requires, first, a careful study of the space of such maps to have a clear idea of how particular lattice structures impact on the size of the space. We are, moreover, particularly interested in computing the *maximum* join-endomorphism below a given collection of join-endomorphisms. This turns out to be important, among others, in spatial computation (and in epistemic logic) to model the distributed information (resp. distributed knowledge) available to a set of agents as conforming a group [5]. It could also be regarded as the maximum perception consistent with (or derivable from) a collection of perceptions of a group of agents.

Problem. Consider the set $\mathcal{E}(L)$ of all join-endomorphisms of a finite lattice L . The set $\mathcal{E}(L)$ can be made into a lattice by ordering join-endomorphisms point-wise wrt the order of L . We investigate the following maximization problem: *Given a lattice L of size n and a set $S \subseteq \mathcal{E}(L)$ of size m , find in $\mathcal{E}(L)$ the greatest lower bound of S , i.e. $\prod_{\mathcal{E}(L)} S$.* Simply taking $\sigma : L \rightarrow L$ with $\sigma(e) \stackrel{\text{def}}{=} \prod_L \{f(e) \mid f \in S\}$ does not solve the problem as σ may not be a join-endomorphism. Furthermore, since $\mathcal{E}(L)$ can be seen as the search space, we also consider the problem of determining its cardinality. Our main results are the following.

This paper. We present characterizations of the exact cardinality of $\mathcal{E}(L)$ for some fundamental lattices. Our contribution is to establish the cardinality of $\mathcal{E}(L)$ for the stereotypical non-distributive lattice $L = \mathbf{M}_n$. We show that $|\mathcal{E}(\mathbf{M}_n)|$ equals $r_0^n + \dots + r_n^n + r_1^{n+1} = n! \mathcal{L}_n(-1) + (n+1)^2$ where r_k^m is the number of ways to place k non-attacking rooks on an $m \times m$ board and $\mathcal{L}_n(x)$ is the Laguerre polynomial of degree n . We also present cardinality results for powerset and linear lattices that are part of the lattice theory folklore: The number of join-endomorphisms is $n^{\log_2 n}$ for powerset lattices of size n and $\binom{2n}{n}$ for linear lattices of size $n+1$. Furthermore, we provide algorithms that, given a lattice L of size n and a set $S \subseteq \mathcal{E}(L)$ of size m , compute $\prod_{\mathcal{E}(L)} S$. Our

contribution is to show that $\prod_{\varepsilon(L)} S$ can be computed with worst-case time complexity in $O(n + m \log n)$ for powerset lattices, $O(mn^2)$ for lattices of sets, and $O(nm + n^3)$ for arbitrary lattices.

2. Background: Join-Endomorphisms and Their Space

We presuppose basic knowledge of order theory [6] and use the following notions.

Partially Ordered Sets. Let (L, \sqsubseteq) be a partially ordered set (poset), and let $S \subseteq L$. We use $\bigsqcup_L S$ to denote the least upper bound (or *supremum* or *join*) of S in L , if it exists. Dually, $\prod_L S$ is the greatest lower bound (glb) (*infimum* or *meet*) of S in L , if it exists. We shall often omit the index L from \bigsqcup_L and \prod_L when no confusion arises. As usual, if $S = \{c, d\}$, $c \sqcup d$ and $c \sqcap d$ represent $\bigsqcup S$ and $\prod S$, respectively. If L has a greatest element (top) \top , and a least element (bottom) \perp , we have $\bigsqcup \emptyset = \perp$ and $\prod \emptyset = \top$. The poset L is *distributive* iff for every $a, b, c \in L$, $a \sqcup (b \sqcap c) = (a \sqcup b) \sqcap (a \sqcup c)$.

Given n , we use \mathbf{n} to denote the poset $\{1, \dots, n\}$ with the linear order $x \sqsubseteq y$ iff $x \leq y$. The poset $\bar{\mathbf{n}}$ is the set $\{1, \dots, n\}$ with the discrete order $x \sqsubseteq y$ iff $x = y$. Given a poset L , we use L_\perp for the poset that results from adding a bottom element to L . The poset L^\top is similarly defined.

Lattices. The poset L is a *lattice* iff each finite nonempty subset of L has a supremum and infimum in L , and it is a *complete lattice* iff each subset of L has a supremum and infimum in L . The lattice $\mathbf{2}^n$ is the n -fold Cartesian product of $\mathbf{2}$ ordered coordinate-wise. We define \mathbf{M}_n as the lattice $(\bar{\mathbf{n}}_\perp)^\top$. A *lattice of sets* is a set of sets ordered by inclusion and closed under finite unions and intersections. A *powerset lattice* is a lattice of sets that includes all the subsets of its top element. A lattice L is said to be *modular* if for every $a, b, c \in L$, with $a \sqsubseteq b$, the equality $a \sqcup (c \sqcap b) = (a \sqcup c) \sqcap b$ holds.

Self-maps. A *self-map* on L is a function $f : L \rightarrow L$. A self-map f is *monotonic* if $a \sqsubseteq b$ implies $f(a) \sqsubseteq f(b)$. We say that f *preserves* the join of $S \subseteq L$ iff $f(\bigsqcup S) = \bigsqcup \{f(c) \mid c \in S\}$.

Definition 1 (Downsets, Covers, Join-irreducibility [6]). Let L be a lattice and $a, b \in L$. We say that a *covers* b , written $b \prec a$ (or $a \succ b$), if $a \sqsupset b$ and there is no $c \in L$ such that $a \sqsupset c \sqsupset b$. The *down-set* of a is $\downarrow a \stackrel{\text{def}}{=} \{b \in L \mid b \sqsubseteq a\}$ and the set of elements *covered* by a is $\downarrow^1 a \stackrel{\text{def}}{=} \{b \mid b \prec a\}$. We write $b \preceq a$ for $b \prec a$ or $b = a$ and define the *cover set* of a as $\downarrow^{01} a \stackrel{\text{def}}{=} \{b \in L \mid b \preceq a\}$. An element $c \in L$ is said to be *join-irreducible* if $c = a \sqcup b$ implies $c = a$ or $c = b$. The set of all join-irreducible elements of L is $\mathcal{J}(L)$ and $\downarrow^{\mathcal{J}} c \stackrel{\text{def}}{=} \downarrow c \cap \mathcal{J}(L)$.

We shall investigate the set of all join-endomorphisms of a given lattice ordered point-wise. Notice that every finite lattice is a complete lattice.

Definition 2 (Join-endomorphisms and their space). Let L be a complete lattice. We say that a self-map is a *(lattice) join-endomorphism* iff it preserves the join of every finite subset of L . Define $\mathcal{E}(L)$ as the set of all join-endomorphisms of L . Furthermore, given $f, g \in \mathcal{E}(L)$, define $f \sqsubseteq_{\mathcal{E}} g$ iff $f(a) \sqsubseteq g(a)$ for every $a \in L$.

The following are immediate consequences of the above definition.

Proposition 1. *Let L be a complete lattice. $f \in \mathcal{E}(L)$ iff $f(\perp) = \perp$ and $f(a \sqcup b) = f(a) \sqcup f(b)$ for all $a, b \in L$. If f is a join-endomorphism of L then f is monotonic.*

Given a set $S \subseteq \mathcal{E}(L)$, where L is a finite lattice, we are interested in finding the greatest join-endomorphism in $\mathcal{E}(L)$ below the elements of S , i.e. $\prod_{\mathcal{E}(L)} S$. Since every finite lattice is also a complete lattice, the existence of $\prod_{\mathcal{E}(L)} S$ is guaranteed by the following proposition.

Proposition 2 ([7]). *If (L, \sqsubseteq) is a complete lattice, $(\mathcal{E}(L), \sqsubseteq_{\mathcal{E}})$ is a complete lattice.*

In the following sections we study the cardinality of $\mathcal{E}(L)$ for some fundamental lattices and provide efficient algorithms to compute $\prod_{\mathcal{E}(L)} S$.

3. The Size of the Function Space

The main result of this section is Theorem 1. It states the size of $\mathcal{E}(\mathbf{M}_n)$. Propositions 3 and 4 state, respectively, the size of $\mathcal{E}(L)$ for the cases when L is a powerset lattice and when L is a total order. These propositions follow from simple observations and they are part of the lattice theory folklore [8, 9, 10]. We include our original proofs of these propositions in the technical report of this paper [11].

3.1. Distributive Lattices

We begin with lattices isomorphic to $\mathbf{2}^n$. They include *finite boolean algebras* and *powerset* lattices [6]. The size of these lattices are easy to infer from the observation that the join-preserving functions on them are determined by their action on the lattices' atoms.

Proposition 3. *Suppose that $m \geq 0$. Let L be any lattice isomorphic to the product lattice $\mathbf{2}^m$. Then $|\mathcal{E}(L)| = n^{\log_2 n}$ where $n = 2^m$ is the size of L .*

Thus powerset lattices and boolean algebras have a super-polynomial, sub-exponential number of join-endomorphisms. Nevertheless, linear order lattices allow for an exponential number of join-endomorphisms given by the *central binomial coefficient*. The following proposition is also easy to prove from the observation that the join-endomorphisms over a linear order are also monotonic functions. In fact, this result appears in [8] and it is well-known among the algebraic theory community [9, 10].

Proposition 4. *Suppose that $n \geq 0$. Let L be any lattice isomorphic to the linear order lattice \mathbf{n}_\perp . Then $|\mathcal{E}(L)| = \binom{2n}{n}$.*

It is easy to prove that $\frac{4^n}{2\sqrt{n}} \leq \binom{2n}{n} \leq 4^n$ for $n \geq 1$. Together with Proposition 4, this gives us explicit exponential lower and upper bounds for $|\mathcal{E}(L)|$ when L is a linear lattice.

3.2. Non-distributive Case

The number of join-endomorphisms for some non-distributive lattices of a given size can be much bigger than that for those distributive lattices of the same size in the previous section. We will characterize this number for an archetypal non-distributive lattice in terms of Laguerre (and rook) polynomials.

Laguerre polynomials are solutions to Laguerre's second-order linear differential equation $xy'' + (1-x)y' + ny = 0$ where y' and y'' are the first and second derivatives of an unknown function y of the variable x , and n is a non-negative integer. The Laguerre polynomial of degree n in x , $\mathcal{L}_n(x)$ is given by the summation $\sum_{k=0}^n \binom{n}{k} \frac{(-1)^k}{k!} x^k$.

The lattice \mathbf{M}_n is non-distributive for any $n \geq 3$. The size of $\mathcal{E}(\mathbf{M}_n)$ can be succinctly expressed as follows.

Theorem 1. $|\mathcal{E}(\mathbf{M}_n)| = (n+1)^2 + n!\mathcal{L}_n(-1)$.

In combinatorics rook polynomials are generating functions of the number of ways to place non-attacking rooks on a board. A *rook polynomial* (for square boards) $\mathcal{R}_n(x)$ has the form $\sum_{k=0}^n x^k r(k, n)$ where the (rook) coefficient $r(k, n)$ represents the number of ways to place k non-attacking rooks on an $n \times n$ chessboard. For instance, $r(0, n) = 1$, $r(1, n) = n^2$ and $r(n, n) = n!$. In general $r(k, n) = \binom{n}{k}^2 k!$.

Rook polynomials are related to Laguerre polynomials by the equation $\mathcal{R}_n(x) = n!x^n \mathcal{L}_n(-x^{-1})$. Therefore, as a direct consequence of the above theorem, we can also characterize $|\mathcal{E}(\mathbf{M}_n)|$ in combinatorial terms as the following sum of rook coefficients.

Corollary 1. *Let $r'(n+1, n) = r(1, n+1)$ and $r'(k, n) = r(k, n)$ if $k \leq n$. Then $|\mathcal{E}(\mathbf{M}_n)| = \sum_{k=0}^{n+1} r'(k, n)$.*

We conclude this section with another pleasant correspondence between the endomorphisms in $\mathcal{E}(\mathbf{M}_n)$ and $\mathcal{R}_n(x)$. Let $f : L \rightarrow L$ be a function over a lattice (L, \sqsubseteq) . We say that f is *non-reducing* in L iff it does not map any value to a smaller one; i.e. there is no $e \in L$ such that $f(e) \sqsubset e$. The number of join-endomorphisms that are non-reducing in \mathbf{M}_n is exactly the value of the rook polynomial $\mathcal{R}_n(x)$ for $x = 1$.

Corollary 2. $\mathcal{R}_n(1) = |\{ f \in \mathcal{E}(\mathbf{M}_n) \mid f \text{ is non-reducing in } \mathbf{M}_n \}|$.

Table 1 illustrates the join-endomorphisms over the lattice \mathbf{M}_n as a union $\bigcup_{i=1}^4 \mathcal{F}_i$. Corollary 2 follows from the observation that the set of non-reducing functions in \mathbf{M}_n is equal to \mathcal{F}_4 whose size is $\mathcal{R}_n(1)$ as shown in the following proof of Theorem 1.

3.2.1. Proof of Theorem 1.

We show that $|\mathcal{E}(\mathbf{M}_n)|$ can be expressed in terms of Laguerre polynomials: $|\mathcal{E}(\mathbf{M}_n)| = (n+1)^2 + n!\mathcal{L}_n(-1)$.

Let $\mathcal{F} = \bigcup_{i=1}^4 \mathcal{F}_i$ where the mutually exclusive \mathcal{F}_i 's are defined in Table 1, and $I = \{1, \dots, n\}$. The proof is divided in two parts: (I) $\mathcal{F} = \mathcal{E}(\mathbf{M}_n)$ and (II) $|\mathcal{F}| = (n+1)^2 + n!\mathcal{L}_n(-1)$.





 <p>Let \mathcal{F}_1 be the family of functions f that for all $e \in \mathbf{M}_n$, $f(e) = \perp$.</p>	 <p>Let \mathcal{F}_2 be the family of bottom preserving functions f such that for some $e, e' \in I$: (a) $f(\top) = e$, (b) $f(e') = \perp$ or $f(e') = e$, and (c) $f(e'') = e$ for all $e'' \in I \setminus \{e'\}$.</p>
 <p>Let \mathcal{F}_3 be the family of top and bottom preserving functions f such that for some $e \in I$: (a) $f(e) = \perp$, and (b) $f(e') = \top$ for all $e' \in I \setminus \{e\}$.</p>	 <p>Let \mathcal{F}_4 be the family of top and bottom preserving functions f that for some $J \subseteq I$: (a) $f(e) = \top$ for every $e \in J$, (b) $f _{I \setminus J}$ is injective, and (c) $\text{Img}(f _{I \setminus J}) \subseteq I$.</p>

Table 1: Families $\mathcal{F}_1, \dots, \mathcal{F}_4$ of join-endomorphisms of \mathbf{M}_n . $I = \{1, \dots, n\}$. $f|_A$ is the restriction of f to a subset A of its domain. $\text{Img}(f)$ is the image of f . A function from each \mathcal{F}_i for \mathbf{M}_5 is depicted with blue arrows.

Part (I)

For $\mathcal{F} \subseteq \mathcal{E}(\mathbf{M}_n)$, it is easy to verify that each $f \in \mathcal{F}$ is a join-endomorphism.

For $\mathcal{E}(\mathbf{M}_n) \subseteq \mathcal{F}$ we show that for any function f from \mathbf{M}_n to \mathbf{M}_n if $f \notin \mathcal{F}$, then $f \notin \mathcal{E}(\mathbf{M}_n)$. Immediately, if $f(\perp) \neq \perp$ then $f \notin \mathcal{E}(\mathbf{M}_n)$.

Suppose $f(\perp) = \perp$. Let J, K, H be disjoint possibly empty sets such that $I = J \cup K \cup H$ and let $j = |J|$, $k = |K|$ and $h = |H|$. The sets J, K, H represent the

elements of I mapped by f to \top , to elements of I , and to \perp , respectively. More precisely, $\text{Img}(f|_J) = \{\top\}$, $\text{Img}(f|_K) \subseteq I$ and $\text{Img}(f|_H) = \{\perp\}$. Furthermore, for every f either (1) $f(\top) = \perp$, (2) $f(\top) \in I$ or (3) $f(\top) = \top$. For each case we show that $f \notin \mathcal{E}(\mathbf{M}_n)$.

1. $f(\top) = \perp$. Since $f \notin \mathcal{F}_1$ there is an $e \in I$ such that $f(e) \neq \perp$. We have $e \sqsubseteq \top$ but $f(e) \not\sqsubseteq f(\top)$. Then f is not monotonic. From Proposition 1 we conclude $f \notin \mathcal{E}(\mathbf{M}_n)$.
2. $f(\top) \in I$. Let K_1, K_2 be disjoint possibly empty sets such that $K_1 \cup K_2 = K$, $\text{Img}(f|_{K_1}) = \{f(\top)\}$ and $\text{Img}(f|_{K_2}) \neq \{f(\top)\}$. Notice that if $j > 0$ or $|K_2| > 0$, f is non-monotonic and then $f \notin \mathcal{E}(\mathbf{M}_n)$.

We then must have $j = 0$ and $K_2 = \emptyset$. Since $\text{Img}(f|_K) = \{f(\top)\}$ and $f \notin \mathcal{F}_2$ then $h > 1$. Therefore there must be $e_1, e_2 \in H$ such that $f(e_1) = f(e_2) = \perp$. This implies $f(e_1 \sqcup e_2) = f(\top) \neq \perp = f(e_1) \sqcup f(e_2)$, therefore $f \notin \mathcal{E}(\mathbf{M}_n)$.

3. $f(\top) = \top$.
 - 3.1. Suppose $k = 0$. Notice that $f \notin \mathcal{F}_3$ and $f \notin \mathcal{F}_4$ hence $h \neq 1$ and $h \neq 0$. Thus $h > 1$ implies that there are at least two $e_1, e_2 \in H$ such that $f(e_1) = f(e_2) = \perp$. But then $f(e_1 \sqcup e_2) = f(\top) = \top \neq \perp = f(e_1) \sqcup f(e_2)$, hence $f \notin \mathcal{E}(\mathbf{M}_n)$.
 - 3.2. Suppose $k > 0$. Assume $h = 0$. Notice that $K = I \setminus J$ and $\text{Img}(f|_K) \subseteq I$. Since f is a bottom and top preserving function and it satisfies conditions (a) and (c) of \mathcal{F}_4 but $f \notin \mathcal{F}_4$, then f must violate condition (b). Thus $f|_K$ is not injective. Then there are $a, b \in K$ such that $a \neq b$ but $f(a) = f(b)$. Then $f(a) \sqcup f(b) \neq \top = f(a \sqcup b)$. Consequently, $f \notin \mathcal{E}(\mathbf{M}_n)$.
Assume $h > 0$. There must be $e_1, e_2, e_3 \in I$ such that $f(e_1) = \perp$ and $f(e_2) = e_3$. Notice that $f(e_1) \sqcup f(e_2) = e_3 \neq \top = f(\top) = f(e_1 \sqcup e_2)$. Therefore, $f \notin \mathcal{E}(\mathbf{M}_n)$.

Part (II)

We prove that $|\mathcal{F}| = \sum_{i=1}^4 |\mathcal{F}_i| = (n+1)^2 + n! \mathcal{L}_n(-1)$. Recall that $n = |I|$.

1. $|\mathcal{F}_1| = 1$. There is only one function mapping every element in \mathbf{M}_n to \perp .
2. $|\mathcal{F}_2| = n^2 + n$. There are n possibilities to choose an element of I to map \top to. If there is an element of I mapped to \perp , for each one of the previous n options there are also n possibilities to choose such an element. Then, in this case there are n^2 functions. If no element of I is mapped to \perp , then there are n additional functions.
3. $|\mathcal{F}_3| = n$. One of the elements of I is mapped to \perp . All the other elements of I are mapped to \top . Then, there are n functions in \mathcal{F}_3 .

4. $|\mathcal{F}_4| = n!\mathcal{L}_n(-1) = \mathcal{R}_n(1)$. Let $f \in \mathcal{F}_4$ and let $J \subseteq I$ be a possibly empty set such that $\text{Img}(f|_J) = \{\top\}$ and $\text{Img}(f|_{I \setminus J}) \subseteq I$, where $f|_{I \setminus J}$ is an injective function. We shall call $j = |J|$.

For each of the $\binom{n}{j}$ possibilities for J , the elements of $I \setminus J$ are to be mapped to I by the injective function $f|_{I \setminus J}$. The number of functions $f|_{I \setminus J}$ is $\frac{n!}{j!}$. Therefore, $|\mathcal{F}_4| = \sum_{j=0}^n \binom{n}{j} \frac{n!}{j!}$. This sum equals $n!\mathcal{L}_n(-1)$ which in turn is equal to $\mathcal{R}_n(1)$.

It follows that $|\mathcal{F}| = \sum_{i=1}^4 |\mathcal{F}_i| = (n+1)^2 + n!\mathcal{L}_n(-1)$ as wanted. \square

4. Computing the Meet of Join-Endomorphisms

We shall provide efficient algorithms for the maximization problem mentioned in the introduction.

Problem 1. Given a finite lattice L of size n and $S \subseteq \mathcal{E}(L)$ of size m , find the *greatest* join-endomorphism $h : L \rightarrow L$ below all elements of S , i.e. $h = \prod_{\mathcal{E}(L)} S$.

Notice that the lattice $\mathcal{E}(L)$, which could be exponentially bigger than L (see Section 3), is not an input to the problem above.

Finding $\prod_{\mathcal{E}(L)} S$ may not be immediate. For instance, see $\prod_{\mathcal{E}(L)} S$ in Figure 1a for a small lattice of four elements and two join-endomorphisms. As already mentioned, a *naive approach* is to compute $\prod_{\mathcal{E}(L)} S$ by taking $\sigma_S(c) \stackrel{\text{def}}{=} \prod_L \{f(c) \mid f \in S\}$ for each $c \in L$. This does not work since σ_S is not necessarily a join-endomorphism as shown in Figure 1b.

A *brute force* solution to compute $\prod_{\mathcal{E}(L)} S$ can be obtained by generating the set $S' = \{g \mid g \in \mathcal{E}(L) \text{ and } g \sqsubseteq f \text{ for all } f \in S\}$ and taking its join. This approach works since $\bigsqcup S' = \prod_{\mathcal{E}(L)} S$ but as shown in Section 3, the size of $\mathcal{E}(L)$ can be super-polynomial for distributive lattices and exponential in general.

Nevertheless, one can use lattice properties to compute $\prod_{\mathcal{E}(L)} S$ efficiently. For distributive lattices, we use the inherent compositional nature of $\prod_{\mathcal{E}(L)} S$. For arbitrary lattices, we present an algorithm that uses the function σ_S in the naive approach to compute $\prod_{\mathcal{E}(L)} S$ by approximating it from above.

We will give the time complexities in terms of the number of basic binary lattice operations (i.e. meets, joins and subtractions) performed during execution.

4.1. Meet of Join-Endomorphisms in Distributive Lattices

Here we shall illustrate some pleasant compositionality properties of the infima of join-endomorphisms that can be used for computing the join-endomorphism $\prod_{\mathcal{E}(L)} S$ in a finite distributive lattice L . In what follows we assume $n = |L|$ and $m = |S|$. Then we will further improve these results by focusing on the join-irreducible elements of L when computing $\prod_{\mathcal{E}(L)} S$.

We use X^J to denote the set of tuples $(x_j)_{j \in J}$ of elements $x_j \in X$ for each $j \in J$.




Figure 1: $S = \{f, g\} \subseteq \mathcal{E}(L)$. (a) $h = \prod_{\mathcal{E}(L)} S$. (b) $\sigma_S(c) \stackrel{\text{def}}{=} f(c) \sqcap g(c)$ is not a join-endomorphism of \mathbf{M}_2 : $\sigma_S(1 \sqcup 2) \neq \sigma_S(1) \sqcup \sigma_S(2)$. (c) δ_S in Lemma 1 is not a join-endomorphism of the non-distributive lattice \mathbf{M}_3 : $\delta_S(1) \sqcup \delta_S(2) = 1 \neq \perp = \delta_S(1 \sqcup 2)$.

Lemma 1 ([12]). *Let L be a finite distributive lattice and $S = \{f_i\}_{i \in I} \subseteq \mathcal{E}(L)$. Then $\prod_{\mathcal{E}(L)} S = \delta_S$ where $\delta_S(c) \stackrel{\text{def}}{=} \prod_L \{ \bigsqcup_{i \in I} f_i(a_i) \mid (a_i)_{i \in I} \in L^I \text{ and } \bigsqcup_{i \in I} a_i \sqsupseteq c \}$.*

The above lemma basically says that $(\prod_{\mathcal{E}(L)} S)(c)$ is the greatest element in L below all possible applications of the functions in S to elements whose join is greater or equal to c . The proof that $\delta_S \sqsupseteq_{\mathcal{E}} \prod_{\mathcal{E}(L)} S$ uses the fact that join-endomorphisms preserve joins. The proof that $\delta_S \sqsubseteq_{\mathcal{E}} \prod_{\mathcal{E}(L)} S$ proceeds by showing that δ_S is a lower bound in $\mathcal{E}(L)$ of S . Distributivity of the lattice L is crucial for this direction. In fact without it $\prod_{\mathcal{E}(L)} S = \delta_S$ does not necessarily hold as shown by the following counter-example.

Example 1. Consider the non-distributive lattice \mathbf{M}_3 and $S = \{f, g\}$ defined as in Figure 1c. We obtain $\delta_S(1 \sqcup 2) = \delta_S(\top) = \perp$ and $\delta_S(1) \sqcup \delta_S(2) = 1 \sqcup \perp = 1$. Then, $\delta_S(1 \sqcup 2) \neq \delta_S(1) \sqcup \delta_S(2)$, i.e. δ_S is not a join-endomorphism.

We rewrite Problem 1 to work with only two join-endomorphisms.

Problem 2. Given a lattice L of size n and two join-endomorphisms $f, g : L \rightarrow L$, find the *greatest* join-endomorphism $h : L \rightarrow L$ below both f and g : i.e. $h = f \sqcap_{\mathcal{E}(L)} g$.

The following remark gives us compelling reason for using this simplified version of Problem 1.

Remark 1. From the associativity of the meet operator (\sqcap), given a finite lattice L and a set of join-endomorphisms $S = \{f_i \mid i \in I\} \subseteq \mathcal{E}(L)$, where $I = \{1, 2, \dots, m\}$. We have $\prod_{\mathcal{E}(L)} S = (\dots (f_1 \sqcap_{\mathcal{E}(L)} f_2) \sqcap_{\mathcal{E}(L)} \dots) \sqcap_{\mathcal{E}(L)} f_m$.

Thanks to associativity of the meet operator (\sqcap), solving Problem 1 is equivalent to solving Problem 2 for m pairs of join-endomorphisms in S .

Nevertheless, we can use Lemma 1 to provide a recursive characterization of $\sqcap_{\mathcal{E}(L)} S$ that can be used in a divide-and-conquer algorithm with lower time complexity.

Corollary 3. *Let L be a finite distributive lattice and $S = \{f, g\} \subseteq \mathcal{E}(L)$. Then $\sqcap_{\mathcal{E}(L)} S = \delta_S$ where $\delta_S(c) \stackrel{\text{def}}{=} \sqcap_L \{f(a) \sqcup_L g(b) \mid a, b \in L \text{ and } a \sqcup b \sqsupseteq c\}$.*

The above corollary bears witness to the compositional nature of $\sqcap_{\mathcal{E}(L)} S$.

Naive Algorithm A_1 . One could use Corollary 3 directly in the obvious way, along with Remark 1 to provide an algorithm for $\sqcap_{\mathcal{E}(L)} S$ by computing δ_S : i.e. computing, for a pair $f, g \in S$, the meet of elements of the form $f(a) \sqcup_L g(b)$ for every pair $a, b \in L$ such that $a \sqcup b \sqsupseteq c$. And repeating the process for the remaining functions in S . For each $c \in L$, $\delta_{\{f, g\}}(c)$ performs $O(n^2)$ meets and $O(n^2)$ joins; and this must be done m times. Thus, A_1 can compute $\sqcap_{\mathcal{E}(L)} S$ by performing $O(n \times n^2 \times m) = O(mn^3)$ binary lattice operations.

4.2. Using Subtraction and Downsets to characterize $\sqcap_{\mathcal{E}(L)} S$

In what follows we show that $\sqcap_{\mathcal{E}(L)} S$ can be computed in $O(mn^2)$ for distributive lattices. To achieve this we use the subtraction operator from co-Heyting algebras and the notion of down set¹.

Subtraction Operator. Notice that in Corollary 3 we are considering *all* pairs $a, b \in L$ such that $a \sqcup b \sqsupseteq c$. However, because of the monotonicity of join-endomorphisms, it suffices to take, for each $a \in L$, just *the least* b such that $a \sqcup b \sqsupseteq c$. In finite distributive lattices, and more generally in co-Heyting algebras [13], the *subtraction* operator $c \ominus a$ gives us exactly such a least element. The subtraction operator is uniquely determined by the property (*Galois connection*) $b \sqsupseteq c \ominus a$ iff $a \sqcup b \sqsupseteq c$ for all $a, b, c \in L$.

Down-sets. Besides using just $c \ominus a$ instead of all b 's such that $a \sqcup b \sqsupseteq c$, we can use a further simplification: Rather than including every $a \in L$, we only need to consider every a in the *down-set* of c . Recall that the down-set of c is defined as $\downarrow c = \{e \in L \mid e \sqsubseteq c\}$ (see Definition 1). This additional simplification is justified using properties of distributive lattices to show that for any $a' \in L$, such that $a' \not\sqsubseteq c$, there exists $a \sqsubseteq c$ such that $f(a) \sqcup g(c \ominus a) \sqsubseteq f(a') \sqcup g(c \ominus a')$.

The above observations lead us to the following theorem.

Theorem 2. *Let L be a finite distributive lattice, $S = \{f, g\} \subseteq \mathcal{E}(L)$ and let $h = \sqcap_{\mathcal{E}(L)} S$. Then $h(c) = \sqcap_L \{f(a) \sqcup g(c \ominus a) \mid a \in \downarrow c\}$.*

¹Recall that we give time complexities in terms of the number of basic binary lattice operations (i.e. meets, joins and subtractions) performed during execution.

The above result can be used to derive a simple recursive algorithm that, given a finite distributive lattice L and $S \subseteq \mathcal{E}(L)$, computes $\prod_{\mathcal{E}(L)} S$ in worst-case time complexity $O(mn^2)$ where $m = |S|$ and $n = |L|$. We show this algorithm next.

Algorithm 1 $\text{DMEET}(L, S)$ finds $\prod_{\mathcal{E}(L)} S = h$, where L is a finite distributive lattice, $S \subseteq \mathcal{E}(L)$, and $|S| \geq 1$

```

1: if  $S = \{f\}$  then return  $f$   $\triangleright f \in \mathcal{E}(L)$ 
2: if  $S = \{f, g\}$  then  $\triangleright f, g \in \mathcal{E}(L)$ 
3:   for each  $c \in L$  do
4:      $h(c) \leftarrow \prod_L \{f(a) \sqcup g(c \ominus a) \mid a \in \downarrow c\}$ 
5:   return  $h$ 
6: else
7:   partition  $S$  into  $S_1$  and  $S_2$  such that  $|S_1| = 2$ 
8:   return  $\text{DMEET}(L, \{\text{DMEET}(L, S_1)\} \cup S_2)$ 

```

Complexity for Distributive Lattices. Assume that L is a finite distributive lattice of size n , and that S is a subset of $\mathcal{E}(L)$ of size m . With Algorithm 1, DMEET is called m times, and for each $c \in L$ the size of $\downarrow c$ is at most of size n . Then, we must do $O(n)$ meets and $O(n)$ joins for each $c \in L$. The worst-case time complexity of DMEET is then in $O(mn^2)$.

Nevertheless, we will show how to achieve a better upper bound when considering the join-irreducible elements of the finite distributive lattice.

4.3. Using Join-Irreducible Elements

As seen from the beginning of Section 4, if we rewrite $\sigma_S(c) \stackrel{\text{def}}{=} \prod_L \{f(c) \mid f \in S\}$ where $S = \{f, g\}$, we obtain following somewhat appealing equation

$$(f \prod_{\mathcal{E}(L)} g)(a) = f(a) \prod_L g(a) \tag{1}$$

Sadly, it still *does not* hold in general, as illustrated in the lattices \mathbf{M}_2 and \mathbf{M}_3 in Figure 1b and Figure 2.

However, it turns out that we can partly use Equation 1 to obtain a better upper bound. The following lemma states that Equation 1 holds if L is distributive and $a \in \mathcal{J}(L)$.

Lemma 2. *Let L be a finite distributive lattice and $f, g \in \mathcal{E}(L)$. Then the following equation holds: $(f \prod_{\mathcal{E}(L)} g)(a) = f(a) \prod_L g(a)$ for every $a \in \mathcal{J}(L)$.*

Proof. From Theorem 2, $(f \prod_{\mathcal{E}(L)} g)(a) = \prod \{f(a') \sqcup g(a \ominus a') \mid a' \in \downarrow a\}$. Note that since $a \in \mathcal{J}(L)$ if $a' \in \downarrow a$ then $a \ominus a' = a$ when $a \neq a'$, and $a \ominus a' = \perp$ when $a = a'$. Then,

$$\begin{aligned} \{f(a') \sqcup g(a \ominus a') \mid a' \in \downarrow a\} &= \{f(a') \sqcup g(a \ominus a') \mid a' \sqsubset a\} \cup \{f(a) \sqcup g(\perp)\} \\ &= \{f(a') \sqcup g(a) \mid a' \sqsubset a\} \cup \{f(a)\} \\ &= \{f(a') \sqcup g(a) \mid \perp \sqsubset a' \sqsubset a\} \cup \{f(a), g(a)\}. \end{aligned}$$




Figure 2: $f: \dots \rightarrow$, $g: \rightarrow$, $h: \dashrightarrow$. Any $h: \mathbf{M}_3 \rightarrow \mathbf{M}_3$ s.t. $h(a) = f(a) \sqcap g(a)$ for $a \in \mathcal{J}(\mathbf{M}_3)$ is not in $\mathcal{E}(\mathbf{M}_3)$: $h(\top) = h(1 \sqcup 2) = h(1) \sqcup h(2) = 1 \neq \perp = h(2) \sqcup h(3) = h(2 \sqcup 3) = h(\top)$.

By absorption, we know that $(f(a') \sqcup g(a)) \sqcap g(a) = g(a)$. Finally, using properties of \sqcap ,

$$\begin{aligned} (f \sqcap_{\mathcal{E}(L)} g)(a) &= \bigsqcap (\{f(a') \sqcup g(a) \mid \perp \sqsubset a' \sqsubset a\} \cup \{f(a), g(a)\}) \\ &= \bigsqcap \{f(a') \sqcup g(a) \mid \perp \sqsubset a' \sqsubset a\} \sqcap f(a) \sqcap g(a) \\ &= f(a) \sqcap g(a). \end{aligned}$$

□

It is worth noting the Lemma 2 may not hold for non-distributive lattices. This is illustrated in Figure 2 with the archetypal non-distributive lattice \mathbf{M}_3 . Suppose that f and g are given as in Figure 2. Let $h = f \sqcap_{\mathcal{E}(L)} g$ with $h(a) = f(a) \sqcap g(a)$ for all $a \in \{1, 2, 3\} = \mathcal{J}(\mathbf{M}_3)$. Since h is a join-endomorphism, we would have $h(\top) = h(1 \sqcup 2) = h(1) \sqcup h(2) = 1 \neq \perp = h(2) \sqcup h(3) = h(2 \sqcup 3) = h(\top)$, a contradiction.

The following proposition will come in handy for defining join-endomorphisms in terms of join-irreducible elements. Recall that $\downarrow^{\mathcal{J}} e = \downarrow e \cap \mathcal{J}(L)$ (see Definition 1).

Proposition 5 ([7, 6]). *Let L be a lattice. If L is finite and distributive, $f \in \mathcal{E}(L)$ iff $(\forall e \in L) f(e) = \bigsqcup \{f(e') \mid e' \in \downarrow^{\mathcal{J}} e\}$.*

Lemma 2 and 5 lead us to the following characterization of meets over $\mathcal{E}(L)$.

Theorem 3. *Let L be a finite distributive lattice and $f, g \in \mathcal{E}(L)$. Then $h = f \sqcap_{\mathcal{E}(L)} g$ iff h satisfies*

$$h(a) = \begin{cases} f(a) \sqcap_L g(a) & \text{if } a \in \mathcal{J}(L) \text{ or } a = \perp \\ h(b) \sqcup_L h(c) & \text{if } b, c \in \downarrow^1 a \text{ with } b \neq c \end{cases} \quad (2)$$

Proof. The only-if direction follows from Lemma 2 and Proposition 5. For the if-direction, suppose that h satisfies Equation 2. If $h \in \mathcal{E}(L)$ the result follows from Lemma 2 and 5. To prove $h \in \mathcal{E}(L)$ from 5 it suffices to show

$$h(e) = \bigsqcup \{h(e') \mid e' \in \downarrow^{\mathcal{J}} e\} \quad (3)$$

for every $e \in L$. From Equation 2 and since f and g are monotonic, h is monotonic. If $e \in \mathcal{J}(L)$ then $h(e') \sqsubseteq h(e)$ for every $e' \in \downarrow^{\mathcal{J}} e$. Therefore, $\bigsqcup \{h(e') \mid e' \in \downarrow^{\mathcal{J}} e\} = h(e)$. If $e \notin \mathcal{J}(L)$, we proceed by induction. Assume Equation 3 holds for all $a \in \downarrow^1 e$. By definition, $h(e) = h(b) \sqcup h(c)$ for any $b, c \in \downarrow^1 e$ with $b \neq c$. Then, we have $h(b) = \bigsqcup \{h(e') \mid e' \in \downarrow^{\mathcal{J}} b\}$ and $h(c) = \bigsqcup \{h(e') \mid e' \in \downarrow^{\mathcal{J}} c\}$. Notice that $e' \in \downarrow^{\mathcal{J}} b$ or $e' \in \downarrow^{\mathcal{J}} c$ iff $e' \in \downarrow^{\mathcal{J}}(b \sqcup c)$, since L is distributive. Thus, $h(e) = h(b) \sqcup h(c) = \bigsqcup \{h(e') \mid e' \in \downarrow^{\mathcal{J}}(b \sqcup c)\} = \bigsqcup \{h(e') \mid e' \in \downarrow^{\mathcal{J}} e\}$ as wanted. \square

We conclude this section by stating the time complexity $O(n)$ to compute h in the above theorem. The time complexity is determined by the number of basic binary lattice operations (i.e. meets and joins) performed during execution.

Corollary 4. *Given a distributive lattice L of size n , and functions $f, g \in \mathcal{E}(L)$, the function $h = f \sqcap_{\mathcal{E}(L)} g$ can be computed in $O(n)$ binary lattice operations.*

Proof. If $a \in \mathcal{J}(L)$ then from Theorem 3, $h(a)$ can be computed as $f(a) \sqcap g(a)$. If $a = \perp$ then $h(a)$ is \perp . If $a \notin \mathcal{J}(L)$ and $a \neq \perp$, we pick any $b, c \in \downarrow^1 a$ such that $b \neq c$ and compute $h(a)$ recursively as $h(b) \sqcup h(c)$ by Theorem 3. We can use a lookup table to keep track of the values of $a \in L$ for which $h(a)$ has been computed, starting with all $a \in \mathcal{J}(L)$. Since $h(a)$ is only computed once for each $a \in L$, either as a meet for elements in $\mathcal{J}(L)$ or as a join otherwise, we only perform n binary lattice operations. \square

Following Remark 1 and Corollary 4, for a finite distributive lattice L of size n and a set of join-endomorphisms $S \subseteq \mathcal{E}(L)$ of size m , we can compute $\prod_{\mathcal{E}(L)} S$ with a worst-case complexity in $O(mn)$ binary lattice operations.

Profiling and Runtime

For distributive lattices, we now present some experimental results comparing the average runtime between Algorithm 1 based on Theorem 2, referred to as DMEET, and the proposed algorithm in Theorem 3, called DMEET+.

We use the algorithms presented above to compute the greatest dilation below a given set of dilations and illustrate its result for a simple image in Section 4.6.

The algorithms were implemented using Python 3, the code is available in [14]. All the results in Figure 3, were executed in a MacBook Pro (Retina, 15-inch, Mid 2014) with a Quad-Core Intel Core i7 processor, and 16 GB of 1600 MHz DDR3 RAM with a constant supply of electricity.

Figure 3 shows the average runtime of each algorithm, from 100 runs with a random pair of join-endomorphisms. For Figure 3a, we compared each algorithm




Figure 3: Comparison between an implementation of Theorem 2 (DMEET) and Theorem 3 (DMEET+).

Size	DMEET Time [s]	DMEET+ Time [s]	DMEET # \sqcup	DMEET+ # \sqcup	DMEET # \sqcap	DMEET+ # \sqcap
16	0.000246	0.000024	81	11	81	4
32	0.000971	0.000059	243	26	243	5
64	0.002659	0.000094	729	57	729	6
128	0.008735	0.000163	2187	120	2187	7
256	0.038086	0.000302	6561	247	6561	8
512	0.244304	0.000645	19683	502	19683	9
1024	1.518173	0.001468	59049	1013	59049	10

Table 2: Average runtime in seconds over powerset lattices. Number of \sqcup and \sqcap operations performed for each algorithm.

against powerset lattices of sizes between 2^2 and 2^{10} . For Figure 3b, 10 random distributive lattices of size 10 were selected. In both cases, all binary lattice operation are guaranteed a complexity in $O(1)$ to showcase the quadratic nature of DMEET compared to the linear growth of DMEET+. The time reduction from DMEET to DMEET+ is also reflected in a reduction on the number of \sqcup and \sqcap operations performed as illustrated in Table 2. For DMEET+, given a distributive lattice L of size n , $\#\sqcap = |\mathcal{J}(L)|$ and $\#\sqcup = |L| - |\mathcal{J}(L)| - 1$ (\perp is directly mapped to \perp).

4.4. Algorithms for Arbitrary Lattices

The previous algorithms may fail to produce $\prod_{\varepsilon(L)} S$ for non-distributive finite lattices. Nonetheless, for any arbitrary finite lattice L , $\prod_{\varepsilon(L)} S$ can be computed by successive approximations, starting with some self-map known to be smaller than each $f \in S$ and greater than $\prod_{\varepsilon(L)} S$. Assume a self-map $\sigma : L \rightarrow L$ such that $\sigma \sqsupseteq_{\varepsilon} \prod_{\varepsilon(L)} S$, and for all $f \in S$, $\sigma \sqsubseteq_{\varepsilon} f$. A good starting

point is $\sigma(u) = \prod\{f(u) \mid f \in S\}$, for all $u \in L$. By definition of meet operator, σ is the biggest function under all functions in S , hence $\sigma \sqsupseteq_{\mathcal{E}} \prod_{\mathcal{E}(L)} S$. The program GMEET in Algorithm 2 computes decreasing upper bounds of $\prod_{\mathcal{E}(L)} S$ by correcting σ values not conforming to the *join-endomorphism property*: $\sigma(u) \sqcup \sigma(v) = \sigma(u \sqcup v)$. The correction decreases σ and maintains the invariant $\sigma \sqsupseteq_{\mathcal{E}} \prod_{\mathcal{E}(L)} S$, as stated in Theorem 4. The proof of this statement is given in Section 4.7.

Theorem 4. *Let L be a finite lattice, $u, v \in L$, $\sigma : L \rightarrow L$ and $S \subseteq \mathcal{E}(L)$. Assume $\sigma \sqsupseteq_{\mathcal{E}} \prod_{\mathcal{E}(L)} S$ holds, and consider the following updates:*

1. *when $\sigma(u) \sqcup \sigma(v) \sqsubset \sigma(u \sqcup v)$, assign $\sigma(u \sqcup v) \leftarrow \sigma(u) \sqcup \sigma(v)$*
2. *when $\sigma(u) \sqcup \sigma(v) \not\sqsubseteq \sigma(u \sqcup v)$, assign $\sigma(u) \leftarrow \sigma(u) \sqcap \sigma(u \sqcup v)$ and also $\sigma(v) \leftarrow \sigma(v) \sqcap \sigma(u \sqcup v)$*

Let σ' be the function resulting after the update. Then, (1) $\sigma' \sqsubset \sigma$ and (2) $\sigma' \sqsupseteq_{\mathcal{E}} \prod_{\mathcal{E}(L)} S$.

Algorithm 2 GMEET finds $\sigma = \prod_{\mathcal{E}(L)} S$

```

1:  $\sigma(u) \leftarrow \prod\{f(u) \mid f \in S\}$  ▷ for all  $u \in L$ 
2: while  $u, v \in L \wedge \sigma(u) \sqcup \sigma(v) \neq \sigma(u \sqcup v)$  do
3:   if  $\sigma(u) \sqcup \sigma(v) \sqsubset \sigma(u \sqcup v)$  then ▷ case (1)
4:      $\sigma(u \sqcup v) \leftarrow \sigma(u) \sqcup \sigma(v)$ 
5:   else ▷ case (2)
6:      $\sigma(u) \leftarrow \sigma(u) \sqcap \sigma(u \sqcup v)$ 
7:      $\sigma(v) \leftarrow \sigma(v) \sqcap \sigma(u \sqcup v)$ 

```

The procedure (see Algorithm 2) loops through pairs $u, v \in L$ while there is some pair satisfying cases (1) or (2) above for the current σ . When there is, it updates σ as mentioned in Theorem 4. At the end of the loop all pairs $u, v \in L$ satisfy the join preservation property. By the invariant mentioned in the theorem, this means $\sigma = \prod_{\mathcal{E}(L)} S$.

As for the previous algorithms in this paper, the worst-time time complexity will be expressed in terms of the binary lattice operations performed during execution. Assume a fixed set S of size m . The complexity of the initialization (Line 1) of GMEET is $O(nm)$ with $n = |L|$. The value of σ for a given $w \in L$ can be updated (decreased) at most n times. Thus, there are at most n^2 updates of σ for all values of L . Finding a $w = u \sqcup v$ where $\sigma(w)$ needs an update because $\sigma(u) \sqcup \sigma(v) \neq \sigma(u \sqcup v)$ (test of the loop, Line 2) takes $O(n^2)$. Hence, the worst time complexity of the loop is in $O(n^4)$.

The program GMEET+, in Algorithm 3, uses appropriate data structures to reduce significantly the time complexity of the algorithm. Essentially, different sets are used to keep track of properties of (u, v) lattice pairs with respect to the current σ . We have a support (correct) pairs set $\text{Sup}_w = \{(u, v) \mid w = u \sqcup v \wedge \sigma(u) \sqcup \sigma(v) = \sigma(w)\}$. We also have a conflicts set $\text{Con}_w = \{(u, v) \mid w =$

Algorithm 3 GMEET+ finds $\sigma = \prod_{\varepsilon(L)} S$

```

1:  $\sigma(u) \leftarrow \prod\{f(u) \mid f \in S\}$  ▷ for all  $u \in L$ 
2: Initialize  $\text{Sup}_w, \text{Con}_w, \text{Fail}_w$ , for all  $w$ 
3: while  $w \in L$  such that  $(u, v) \in \text{Con}_w$  do ▷ some conflict set not empty
4:    $\text{Con}_w \leftarrow \text{Con}_w \setminus \{(u, v)\}$ 
5:    $\sigma(w) \leftarrow \sigma(u) \sqcup \sigma(v)$ 
6:    $\text{Fail}_w \leftarrow \text{Fail}_w \cup \text{Sup}_w$  ▷ all pairs previously in  $\text{Sup}_w$  are now failures
7:    $\text{Sup}_w \leftarrow \{(u, v)\}$ 
8:   CHECKSUPPORTS( $w$ ) ▷ for  $u \in L$ , verify property  $\text{Sup}_{w \sqcup u}$ 
9:   while  $z \in L$  such that  $(x, y) \in \text{Fail}_z$  do ▷ some failures set not empty
10:     $\text{Fail}_z \leftarrow \text{Fail}_z \setminus \{(x, y)\}$ 
11:    if  $\sigma(x) \neq \sigma(x) \sqcap \sigma(z)$  then
12:       $\sigma(x) \leftarrow \sigma(x) \sqcap \sigma(z)$  ▷  $\sigma(x)$  decreases
13:       $\text{Fail}_x \leftarrow \text{Fail}_x \cup \text{Sup}_x$  ▷ all pairs in  $\text{Sup}_x$  are now failures
14:       $\text{Sup}_x \leftarrow \emptyset$ 
15:      CHECKSUPPORTS( $x$ ) ▷ for  $u \in L$ , verify property  $\text{Sup}_{x \sqcup u}$ 
16:      if  $\sigma(y) \neq \sigma(y) \sqcap \sigma(z)$  then
17:         $\sigma(y) \leftarrow \sigma(y) \sqcap \sigma(z)$  ▷  $\sigma(y)$  decreases
18:         $\text{Fail}_y \leftarrow \text{Fail}_y \cup \text{Sup}_y$  ▷ all pairs in  $\text{Sup}_y$  are now failures
19:         $\text{Sup}_y \leftarrow \emptyset$ 
20:        CHECKSUPPORTS( $y$ ) ▷ for  $u \in L$ , verify property  $\text{Sup}_{y \sqcup u}$ 
21:      if  $\sigma(x) \sqcup \sigma(y) = \sigma(z)$  then
22:         $\text{Sup}_z \leftarrow \text{Sup}_z \cup \{(x, y)\}$  ▷  $(x, y)$  is now correct
23:      else
24:         $\text{Con}_z \leftarrow \text{Con}_z \cup \{(x, y)\}$  ▷  $(x, y)$  is now a conflict

```

$u \sqcup v \wedge \sigma(u) \sqcup \sigma(v) \sqsubset \sigma(w)$ and failures set $\text{Fail}_w = \{(u, v) \mid w = u \sqcup v \wedge \sigma(u) \sqcup \sigma(v) \not\sqsubseteq \sigma(w)\}$. Algorithm 3 updates σ as mentioned in Theorem 4 and so maintains the invariant $\sigma \sqsupseteq \prod_{\varepsilon(L)} S$. An additional invariant is that, for all w , sets $\text{Sup}_w, \text{Con}_w, \text{Fail}_w$ are pairwise disjoint. When the outer loop finishes sets Con_w and Fail_w are empty (for all w) and thus every (u, v) belongs to $\text{Sup}_{u \sqcup v}$, i.e. the resulting $\sigma = \prod_{\varepsilon(L)} S$.

Auxiliary procedure CHECKSUPPORTS(u) identifies all the pairs $(u, x) \in \text{Sup}_{u \sqcup x}$ that may no longer satisfy the join-endomorphism property $\sigma(u) \sqcup \sigma(x) = \sigma(u \sqcup x)$ because of an update to $\sigma(u)$. When this happens, it adds (u, x) to the appropriate Con, or Fail set. The time complexity of the algorithm depends on the set operations computed for each $w \in L$ chosen, either in the *conflicts* Con_w set or in the *failures* Fail_w set. When a w is selected (for some (u, v) such that $u \sqcup v = w$) the following holds: (1) at least one of $\sigma(w), \sigma(u), \sigma(v)$ is decreased, (2) some fix k number of elements are removed from or added to a set, (3) a union of two *disjoint* sets is computed, and (4) new support sets of w, u or v are calculated.

Complexity for Arbitrary Lattices. With an appropriate implementation, operations (1)-(2) take $O(1)$, and also operation (3), since sets are disjoint. Operation (4) clearly takes $O(n)$. In each loop of the (outer or inner) cycles of the algo-




Figure 4: Average performance time of GMEET+, DMEET and BRUTE-FORCE. Plots A and D use 2^n lattices, B and E distributive lattices, and C and F arbitrary lattices. Plots A-C have a fixed number of join-endomorphisms and plots D-F have a fixed lattice size.

rithm, at least one reduction of σ is computed. Furthermore, for each reduction of σ , $O(n)$ operations are performed. The maximum possible number of $\sigma(w)$ reductions, for a given w , is equal to the length d of the longest strictly decreasing chain in the lattice. The total number of possible σ reductions is thus equal to nd . The total number of operations of the algorithm is then $O(n^2d)$. In general, d could be (at most) equal to n , therefore, after initialization, worst case complexity is $O(n^3)$. The initialization (Lines 1-2) takes $O(nm) + O(n^2)$, where $m = |S|$. Worst time complexity is thus $O(mn + n^3)$. For powerset lattices, $d = \log_2 n$, thus worst time complexity in this case is $O(mn + n^2 \log_2 n)$.

Experimental Results

Here we present some experimental results showing the execution time of the proposed algorithms.

Consider Figure 4. In plots 4.A-C, the horizontal axis is the size of the lattice. In plots 4.D-F, the horizontal axis is the size of $S \subseteq \mathcal{E}(L)$. Curves in images 4.A-C plot, for each algorithm, the average execution time of 100 runs (10 for 4.A) with random sets S of size 4. Images 4.D-F, show the mean execution time of each algorithm for 100 runs (10 for 4.D) varying the number of join-endomorphisms ($|S| = 4i$, $1 \leq i \leq 8$). The lattice size is fixed: $|L| = 10$ for 4.E and 4.F, and $|L| = 2^5$ for 4.D. In all cases the lattices were randomly generated, and the parameters selected to showcase the difference between each algorithm with a sensible overall execution time. For a given lattice L and $S \subseteq \mathcal{E}(L)$, the brute-force algorithm explores the whole space $\mathcal{E}(L)$ to find all the join-endomorphism below each element of S and then computes the greatest of them. In particular, the measured spike in plot 4.C corresponds to the random lattice of seven elements with the size of $\mathcal{E}(L)$ being bigger than in the other experiments in the same figure. In our experiments we observed that for a fixed S , as the size of the lattice increases, DMEET outperforms GMEET+. This is noticeable in lattices 2^n (see 4.A). Similarly, for a fixed lattice, as the size of S

Size	A_1	GMEET	GMEET+	DMEET
16	2.01	0.00360	0.000603	0.000632
32	64.6	0.0633	0.00343	0.00181
64	1901	0.948	0.0154	0.00542
128	>600	15.4	0.0860	0.0160
256	>600	252	0.361	0.0483
512	>600	>600	2.01	0.166
1024	>600	>600	10.7	0.547

Table 3: Average time in seconds over powerset lattices with $|S| = 4$

increases GMEET+ outperforms DMEET. GMEET+ performance can actually improve with a higher number of join-endomorphisms (see 4.D) since the initial σ is usually smaller in this case.

To illustrate some performance gains, Table 3 shows the mean execution time of the algorithms discussed in this paper. We include naive algorithm A_1 outlined just after Corollary 3.

4.5. Meet of Join-Endomorphisms in Modular lattices

In the latter section we have shown that given a finite lattice L and $S \subseteq \mathcal{E}(L)$, Algorithm 3 finds $\prod_{\mathcal{E}(L)} S$ by starting at $\sigma(c) = \prod\{f(c) \mid f \in S\}$ (for all $c \in L$) and correcting the values that violates the equality $\sigma(a \sqcup b) = \sigma(a) \sqcup \sigma(b)$, for every pair $(a, b) \in L^2$. However, one may wonder whether the algorithm tries to correct a value of the function that has been already corrected. Indeed, this may happen for pairs of elements whose join is the same. The following example illustrates such a situation.

Example 2. Consider the modular lattice L and $f, g \in \mathcal{E}(L)$ as depicted in Figure 5. We want to compute $h = f \prod_{\mathcal{E}(L)} g$ represented with dashed teal arrows. Recall that Algorithm 3 initiates with the function $\sigma(c) = f(c) \prod g(c)$ for all $c \in L$ and finds h by correcting σ when $\sigma(c) \neq h(c)$. In Figure 5, x-head arrows show the maps such that $\sigma(c) \neq h(c)$. Notice that to correct the mapping of \top , it suffices to assign $\sigma(4) \sqcup \sigma(5)$ to $\sigma(\top)$, instead of considering all the pairs (c, d) such that $c \sqcup d = \top$, as the algorithm does.

The above example suggests that, in modular lattices, we can decide whether or not a function is a join-endomorphism by checking the distribution property in the covers of every element in the lattice. We shall formalize this intuition and propose an adaptation of Algorithm 3 for this particular case. We specifically shall show that for modular lattices, the algorithm should only consider pairs in the cover set, $\downarrow^{01}c$, for every $c \in L$ (see Definition 1).

4.5.1. Join-Endomorphisms in Modular Lattices

For the sake of the presentation, we present the following definition.

Definition 3. Let L be a lattice and $f : L \rightarrow L$ be a function. We say that f preserves **01-relations** if for every $a, b, c \in L$ such that $a, b \in \downarrow^{01}c$, $f(a \sqcup b) = f(a) \sqcup f(b)$ holds.




Figure 5: Join-endomorphisms $f : \cdots \rightarrow$, $g : \rightarrow$ and $h = f \circ g$ ($h : \dashrightarrow$) on modular lattice L . x-head arrows depict maps of the function $\sigma(c) = f(c) \sqcap g(c)$ s.t. $\sigma(c) \neq h(c)$.

“fgfg” The following is a well-known property of modular lattices; the cornerstone of the characterization of join-endomorphisms in this section.

Proposition 6. [15] *Let L be a lattice. L is modular iff for every $a, b \in L$, the maps $\psi_a : [a \sqcap b, b] \rightarrow [a, a \sqcup b]$ and $\varphi_b : [a, a \sqcup b] \rightarrow [a \sqcap b, b]$ defined as $\psi_a(c) = c \sqcup a$ and $\varphi_b(c) = c \sqcap b$, are mutually inverse isomorphisms.*

The following theorem states that a function is a lattice join-endomorphism if it is **01-relations** preserving.

Theorem 5. *Let L be a modular lattice and let $f : L \rightarrow L$ be a function. f is a join-endomorphism iff f is bottom and **01-relations** preserving.*

Proof. The only-if direction is straightforward. For the if direction, let f be a function such that $f(\perp) = \perp$ and, for every $a, b, c \in L$ with $a, b \in \downarrow^{01}c$, $f(a \sqcup b) = f(a) \sqcup f(b)$ holds.

By Proposition 1, to prove that f is a join-endomorphism, it suffices to show that (\star) $f(a \sqcup b) = f(a) \sqcup f(b)$ for every $a, b \in L$.

We define $n = |[a, a \sqcup b]| = |[a \sqcap b, b]|$ and $m = |[b, a \sqcup b]| = |[a \sqcap b, a]|$ where $a, b \in L$ are any pair of elements such that $a \parallel b$. To prove (\star) we proceed by induction on the size of those intervals, i.e. on both n and m .

For the base cases, if $n = m = 2$ (Figure 6a) the proof follows by the assumption on f . Without loss of generality, assume $n = 2$ and $m = 3$ as shown in Figure 6b. Since f preserves **01-relations**, we have $f(a \sqcup b) = f(a) \sqcup f(e)$ and $f(e) = f(\hat{a}) \sqcup f(b)$ where $b \sqsubset e \prec a \sqcup b$ and $a \sqcap b \sqsubset \hat{a} \prec a$. Then, we obtain $f(a \sqcup b) = f(a) \sqcup f(\hat{a}) \sqcup f(b) = f(a \sqcup \hat{a}) \sqcup f(b) = f(a) \sqcup f(b)$ as wanted.

For the inductive case, assume that (\star) holds for every $2 < n' < n$ and every $2 < m' < m$, i.e. for any intervals $[a, d]$ and $[b, e]$ of size n' and m' , respectively, where $a \sqsubset d \sqsubset a \sqcup b$ and $b \sqsubset e \sqsubset a \sqcup b$. By assumption on f , $f(a \sqcup b) = f(d) \sqcup f(e)$




Figure 6: Isomorphism on modular lattices with different size of intervals

with $a \sqsubset d \prec a \sqcup b$ and $b \sqsubset e \prec a \sqcup b$ as shown in Figure 6c. From the induction hypothesis and Proposition 6, we know that $f(d) = f(a) \sqcup f(\hat{b})$ and $f(e) = f(\hat{a}) \sqcup f(b)$ where $\hat{a} \prec a$ and $\hat{b} \prec b$. Thus, the proof follows from the fact that f preserves **01-relations**: $f(a \sqcup b) = f(a \sqcup \hat{a}) \sqcup f(b \sqcup \hat{b}) = f(a) \sqcup f(b)$.

If a and b are comparable, let us say $a \sqsubseteq b$ (the other case is analogous), the proof goes as before with $\hat{b} = d$ and $e = b$. \square

Now we have the means to present Algorithm 3 adapted to modular lattices.

4.5.2. Algorithm for Modular Lattices

Recall the sets \mathbf{Con}_w , \mathbf{Fail}_w and \mathbf{Sup}_w used by the algorithm to separate the pairs $(u, v) \in L^2$: Namely, with $w = u \sqcup v$, $(u, v) \in \mathbf{Sup}_w$ if $\sigma(u) \sqcup \sigma(v) = \sigma(w)$, $(u, v) \in \mathbf{Con}_w$ if $\sigma(u) \sqcup \sigma(v) \sqsubset \sigma(w)$ and, $(u, v) \in \mathbf{Fail}_w$ otherwise. We shall show that for any modular lattice L , we can construct these sets by using elements in the sets $\downarrow^{01}w = \downarrow^1w \cup \{w\}$ for every $w \in L$. Roughly speaking, we consider only pairs of elements (u, v) that are covered by an element w .

Observation 1. *Let L be a modular lattice, $S \subseteq \mathcal{E}(L)$ and $W = \bigcup_{w \in L} \downarrow^{01}w \subseteq L$. If we form the sets \mathbf{Con} , \mathbf{Fail} and \mathbf{Sup} with pairs of elements in W , it is an immediate consequence of Theorem 4 that Algorithm 3 terminates. Therefore, we have found a function σ' (see Theorem 4) such that $\sigma' = \prod_{\mathcal{E}(L)} S$. Moreover, Theorem 5 guarantees that such a function is a join-endomorphism.*

Experimental Results

Figure shows...

Complexity for Modular Lattices. As we have shown, the time complexity of Algorithm 3 reduces drastically with the adaptation for modular lattices. Specifically, the reduction is due to the initialization step (line 2 in Algorithm 3) of the sets \mathbf{Con} , \mathbf{Fail} and, \mathbf{Sup} . Since we are taking elements in the set $W \subseteq L$, we reduce the number of pairs that the algorithm loops through. However, if the lattice is \mathbf{M}_n , the set $\downarrow^{01}\top = \{\top, 1, \dots, n\}$; in other words, the algorithm will consider all the elements of \mathbf{M}_n . This implies that the algorithm takes no




Figure 7: Binary image I (on the left). Dilations $\delta_{s_1}, \delta_{s_2}$ for structuring elements s_1, s_2 . On the right $(\delta_{s_1} \square_{\mathcal{E}(L)} \delta_{s_2})(I)$. New pixels of the image after each operation in grey and black.

advantage of the modularity of the lattice and the worst-time time complexity is still $O(mn + n^3)$.

We conclude this section with a small example where join-endomorphisms represent dilation operators from Mathematical Morphology [2]. We use the algorithms presented above to compute the greatest dilation below a given set of dilations and illustrate its result for a simple image.

4.6. A Mathematical Morphology Example

Mathematical morphology (MM) is a theory, based on topological, lattice-theoretical and geometric concepts, for the analysis of geometric structures. Its algebraic framework comprises [2, 3, 16], among others, complete lattices together with certain kinds of morphisms, such as *dilations*, defined as *join-endomorphisms* [3]. Our results give bounds about the number of all dilations over certain specific finite lattices and also efficient algorithms to compute their infima.

A typical application of MM is image processing. Consider the space $G = \mathbb{Z}^2$. A dilation [2] by $s_i \subseteq \mathcal{P}(G)$ is a function $\delta_{s_i} : \mathcal{P}(G) \rightarrow \mathcal{P}(G)$ such that $\delta_{s_i}(X) = \{x + e \mid x \in X \text{ and } e \in s_i\}$. The dilation $\delta_{s_i}(X)$ describes the interaction of an image X with the *structuring element* s_i . Intuitively, the dilation of X by s_i is the result of superimpose s_i on every activated pixel of X , with the center of s_i aligned with the corresponding pixel of X . Then, each pixel of every superimposed s_i is included in $\delta_{s_i}(X)$.

Let L be the powerset lattice for some finite set $D \subseteq G$. It turns out that the dilation $\square_{\mathcal{E}(L)} S$ corresponds to the intersection of the structuring elements of the corresponding dilations in S . Figure 7 illustrates $\square_{\mathcal{E}(L)} S$ for the two given dilations $\delta_{s_1}(I)$ and $\delta_{s_2}(I)$ with structuring elements s_1 and s_2 over the given image I .

4.7. Proof of Theorem 4

Let L be a finite lattice, $u, v \in L$, $\sigma : L \rightarrow L$ and $S \subseteq \mathcal{E}(L)$. Assume $\sigma \sqsupseteq \square_{\mathcal{E}(L)} S$ holds, and consider the following updates:

1. when $\sigma(u) \sqcup \sigma(v) \sqsubset \sigma(u \sqcup v)$, assign $\sigma(u \sqcup v) \leftarrow \sigma(u) \sqcup \sigma(v)$

2. when $\sigma(u) \sqcup \sigma(v) \not\sqsubseteq \sigma(u \sqcup v)$, assign $\sigma(u) \leftarrow \sigma(u) \sqcap \sigma(u \sqcup v)$ and also $\sigma(v) \leftarrow \sigma(v) \sqcap \sigma(u \sqcup v)$

Let σ' be the function resulting after the update. Then, (1) $\sigma' \sqsubset \sigma$ and (2) $\sigma' \sqsupseteq \prod_{\mathcal{E}(L)} S$

Proof. Let L be a finite lattice, $S \subseteq \mathcal{E}(L)$, and $h = \prod_{\mathcal{E}(L)} S$.

For update (1), given the condition, the assignment obviously decreases $\sigma(u \sqcup v)$, so $\sigma' \sqsubset \sigma$. For the invariant, since $\sigma \sqsupseteq h$, then, $\sigma(u) \sqsupseteq h(u)$ and $\sigma(v) \sqsupseteq h(v)$, and therefore, $\sigma'(u \sqcup v) = \sigma(u) \sqcup \sigma(v) \sqsupseteq h(u) \sqcup h(v) = h(u \sqcup v)$.

For update (2), the assignments either decrease $\sigma(u)$ or $\sigma(v)$ (or both). To see why, assume the opposite, $\sigma(u) = \sigma(u) \sqcap \sigma(u \sqcup v) \rightarrow \sigma(u) \sqsubset \sigma(u \sqcup v)$, and also $\sigma(v) = \sigma(v) \sqcap \sigma(u \sqcup v) \rightarrow \sigma(v) \sqsubset \sigma(u \sqcup v)$. Therefore, $\sigma(u) \sqcup \sigma(v) \sqsubset \sigma(u \sqcup v)$, contradicting the condition for update 2. Assignments in update 2 also preserve the invariant $\sigma \sqsupseteq h$.

Assume $\sigma(u) \sqcap \sigma(u \sqcup v) \sqsubset \sigma(u)$ (otherwise the invariant holds trivially). By the invariant hypothesis for σ before the assignment, we have that $\sigma(u) \sqsupseteq h(u)$ and $\sigma(u \sqcup v) \sqsupseteq h(u \sqcup v)$. Therefore,

$$\begin{aligned} \sigma(u) \sqcap \sigma(u \sqcup v) &\sqsupseteq h(u) \sqcap h(u \sqcup v) \\ &= h(u) \sqcap (h(u) \sqcup h(v)) \\ &= h(u) \sqcup (h(u) \sqcap h(v)) \\ &= h(u) \end{aligned}$$

The proof for $\sigma'(v)$ is analogous. □

5. Conclusions and Related Work

We have shown that given a lattice L of size n and a set $S \subseteq \mathcal{E}(L)$ of size m , $\prod_{\mathcal{E}(L)} S$ can be computed in the worst-case in $O(n + m \log n)$ binary lattice operations for powerset lattices, $O(mn^2)$ for lattices of sets, and $O(nm + n^3)$ for arbitrary lattices. We illustrated the experimental performance of our algorithms and a small example from mathematical morphology.

In [17] a bit-vector representation of a lattice is discussed. This work gives algorithms of logarithmic (in the size of the lattice) complexity for join and meet operations. These results count bit-vector operations. From [8] we know that $\mathcal{E}(L)$ is isomorphic to the downset of $(P \times P^{op})$, where P is the set of join-prime elements of L , and that this, in turn, is isomorphic to the set of order-preserving functions from $(P \times P^{op})$ to $\mathbf{2}$. Therefore, for the problem of computing $\prod_{\mathcal{E}(L)} S$, we get bounds $O(m \log_2(2^{n^2})) = O(mn^2)$ for set lattices and $O(m(\log_2 n)^2)$ for powerset lattices where $n = |L|$ and $m = |S|$. This, however, assumes a bit-vector representation of a lattice isomorphic to $\mathcal{E}(L)$. Computing this representation takes time and space proportional to the size of $\mathcal{E}(L)$ [17] which could be exponential as stated in the present paper. Notice that in our algorithms the input lattice is L instead of $\mathcal{E}(L)$.

We have stated the cardinality of the set of join-endomorphisms $\mathcal{E}(L)$ for significant families of lattices. To the best of our knowledge we are the first to establish the cardinality $(n+1)^2 + n!\mathcal{L}_n(-1)$ for the lattice \mathbf{M}_n . The cardinalities $n^{\log_2 n}$ for power sets (boolean algebras) and $\binom{2n}{n}$ for linear orders can also be found in the lattice literature [8, 9, 10]. Our original proofs for these statements can be found in the technical report of this paper [11].

The lattice $\mathcal{E}(L)$ have been studied in [7]. The authors showed that a finite lattice L is distributive iff $\mathcal{E}(L)$ is distributive. A lower bound of $2^{2n/3}$ for the number of monotonic self-maps of any finite poset L is given in [18]. Nevertheless to the best of our knowledge, no other authors have studied the problem of determining the size $\mathcal{E}(L)$ nor algorithms for computing $\prod_{\mathcal{E}(L)} S$. We believe that these problems are important, as argued in the Introduction; algebraic structures consisting of a lattice and join-endomorphisms are very common in mathematics and computer science. In fact, our interest in this subject arose in the algebraic setting of spatial and epistemic constraint systems [5] where continuous join-endomorphisms, called space functions, represent knowledge and the infima of endomorphisms correspond to distributed knowledge. We showed in [5] that distributed knowledge can be computed in $O(mn^{1+\log_2(m)})$ for distributive lattices and $O(n^4)$ in general. In this paper we have provided much lower complexity orders for computing infima of join-endomorphisms. Furthermore [5] does not provide the exact cardinality of the set of space functions of a given lattice.

As future work we plan to explore in detail the applications of our work in mathematical morphology and computer music [19]. Furthermore, in the same spirit of [20] we have developed algorithms to generate distributive and arbitrary lattices. In our experiments, we observed that for every lattice L of size n we generated, $n^{\log_2 n} \leq |\mathcal{E}(L)| \leq (n+1)^2 + n!\mathcal{L}_n(-1)$ and if the generated lattice was distributive, $n^{\log_2 n} \leq |\mathcal{E}(L)| \leq \binom{2n}{n}$. We plan to establish if these inequalities hold for every finite lattice.

Acknowledgments.

We are indebted to the anonymous referees and editors of RAMiCS 2020 for helping us to improve one of the complexity bounds, some proofs, and the overall quality of the paper.

References

- [1] S. Knight, C. Palamidessi, P. Panangaden, F. D. Valencia, Spatial and Epistemic Modalities in Constraint-Based Process Calculi, in: 23rd International Conference on Concurrency Theory, Vol. 7454 of Lecture Notes in Computer Science, Springer, 2012, pp. 317–332.
- [2] I. Bloch, H. Heijmans, C. Ronse, Mathematical morphology, in: M. Aiello, I. Pratt-Hartmann, J. Van Benthem (Eds.), Handbook of Spatial Logics, Springer Netherlands, 2007, pp. 857–944.

- [3] C. Ronse, Why mathematical morphology needs complete lattices, *Signal Processing* 21 (2) (1990) 129 – 154.
- [4] M. Guzmán, S. Haar, S. Perchy, C. Rueda, F. D. Valencia, Belief, knowledge, lies and other utterances in an algebra for space and extrusion, *J. Log. Algebr. Meth. Program.* 86 (1) (2017) 107–133.
- [5] M. Guzmán, S. Knight, S. Quintero, S. Ramírez, C. Rueda, F. D. Valencia, Reasoning about Distributed Knowledge of Groups with Infinitely Many Agents, in: *CONCUR 2019 - 30th International Conference on Concurrency Theory*, Vol. 29, 2019, pp. 1–29.
- [6] B. A. Davey, H. A. Priestley, *Introduction to lattices and order*, 2nd Edition, Cambridge university press, 2002.
- [7] G. Grätzer, E. Schmidt, On the lattice of all join-endomorphisms of a lattice, *Proceedings of The American Mathematical Society - PROC AMER MATH SOC* 9 (1958) 722–722.
- [8] G. Birkhoff, *Lattice Theory*, no. v. 25, pt. 2 in *American Mathematical Society colloquium publications*, American Mathematical Society, 1967.
- [9] P. Jipsen, Relation algebras, idempotent semirings and generalized bunched implication algebras, in: *Relational and Algebraic Methods in Computer Science*, Springer International Publishing, 2017, pp. 144–158.
- [10] L. Santocanale, On Discrete Idempotent Paths, in: *Combinatorics on Words*, Vol. 11682, Springer, 2019, pp. 312–325.
- [11] S. Quintero, S. Ramírez, C. Rueda, F. D. Valencia, Counting and Computing Join-Endomorphisms in Lattices, Research report, LIX, Ecole polytechnique ; INRIA Saclay - Ile-de-France (2019).
URL <https://hal.archives-ouvertes.fr/hal-02422624>
- [12] M. Guzmán, S. Knight, S. Quintero, S. Ramírez, C. Rueda, F. Valencia, Reasoning about distributed information with infinitely many agents, *Journal of Logical and Algebraic Methods in Programming* 121 (2021) 100674. doi:<https://doi.org/10.1016/j.jlamp.2021.100674>.
- [13] G. Gierz, K. H. Hofmann, K. Keimel, J. D. Lawson, M. Mislove, D. S. Scott, *Continuous lattices and domains*, Cambridge University Press, 2003.
- [14] S. Quintero, C. Pinzón, S. Ramírez, F. Valencia (2018). [link].
URL <https://github.com/Sirquini/delta>
- [15] G. Grätzer, *Lattice Theory: Foundation*, Springer Basel, 2011.
- [16] J. Stell, Why mathematical morphology needs quantales, in: M. Wilkinson, J. Roerdink (Eds.), *International Symposium on Mathematical Morphology*, ISMM09, Institute for Mathematics and Computing Science, University of Groningen, 2009, pp. 13–16.

- [17] M. Habib, L. Nourine, Tree structure for distributive lattices and its applications, *Theoretical Computer Science* 165 (2) (1996) 391–405.
- [18] D. Duffus, V. Rodl, B. Sands, R. Woodrow, Enumeration of order preserving maps, *Order* 9 (1) (1992) 15–29.
- [19] C. Rueda, F. Valencia, On validity in modelization of musical problems by ccp, *Soft Computing* 8 (9) (2004) 641–648.
- [20] P. Jipsen, N. Lawless, Generating all finite modular lattices of a given size, *Algebra universalis* 74 (3) (2015) 253–264.