CONDITIONAL ESTIMATES FOR L-FUNCTIONS IN THE SELBERG CLASS

NEEA PALOJÄRVI AND ALEKSANDER SIMONIČ

ABSTRACT. Assuming the Generalized Riemann Hypothesis, we provide upper bounds with explicit main terms for moduli of $(\mathcal{L}'/\mathcal{L})(s)$ and $\log \mathcal{L}(s)$ for $|3/4 - \sigma| \le 1/4 - \delta$, fixed $\delta \in (0, 1/4)$ and for functions in the Selberg class. Moreover, by further assuming a polynomial Euler product representation for $\mathcal{L}(s)$, we establish uniform bounds in \mathcal{L} for $|3/4 - \sigma| \le 1/4 - 1/\log \log (q_{\mathcal{L}}|t|^{d_{\mathcal{L}}}), |1 - \sigma| \le 1/\log \log (q_{\mathcal{L}}|t|^{d_{\mathcal{L}}})$ and $\sigma = 1$, and completely explicit estimates by assuming also the strong λ -conjecture.

1. INTRODUCTION

The Selberg class of functions \mathcal{S} was introduced in [Sel92] and consists of Dirichlet series

$$\mathcal{L}(s) = \sum_{n=1}^{\infty} \frac{a(n)}{n^s},$$

where $s = \sigma + it$ for real numbers σ and t, satisfying the following axioms:

- (1) Ramanujan hypothesis. We have $a(n) \ll_{\varepsilon} n^{\varepsilon}$ for any $\varepsilon > 0$.
- (2) Analytic continuation. There exists $k \in \mathbb{N}_0$ such that $(s-1)^k \mathcal{L}(s)$ is an entire function of finite order.
- (3) Functional equation. The function $\mathcal{L}(s)$ satisfies $\mathfrak{L}(s) = \omega \overline{\mathfrak{L}(1-\bar{s})}$, where

$$\mathfrak{L}(s) = \mathcal{L}(s)Q^s \prod_{j=1}^{f} \Gamma\left(\lambda_j s + \mu_j\right)$$

with $(Q, \lambda_j) \in \mathbb{R}^2_+$, and $(\mu_j, \omega) \in \mathbb{C}^2$ with $\Re\{\mu_j\} \ge 0$ and $|\omega| = 1$.

(4) Euler product. The function $\mathcal{L}(s)$ has a product representation

$$\mathcal{L}(s) = \prod_{p} \exp\left(\sum_{k=1}^{\infty} \frac{b\left(p^{k}\right)}{p^{ks}}\right),$$

where the coefficients $b(p^k)$ satisfy $b(p^k) \ll p^{k\theta}$ for some $0 \le \theta < 1/2$.

It is well-known that the data from the functional equation is not uniquely defined by $\mathcal{L} \in \mathcal{S}$. However, one can show that the degree $d_{\mathcal{L}}$ and the conductor $q_{\mathcal{L}}$,

$$d_{\mathcal{L}} = 2 \sum_{j=1}^{f} \lambda_j, \quad q_{\mathcal{L}} = (2\pi)^{d_{\mathcal{L}}} Q^2 \prod_{j=1}^{f} \lambda_j^{2\lambda_j},$$

are indeed invariants for the class S. In the present paper we work only with these two invariants and for the sake of simplicity we introduce $\tau := q_{\mathcal{L}} |t|^{d_{\mathcal{L}}}$. There exist other invariants and the reader is referred to [Per05] and the references thereof. Notable examples of functions in S include:

- The Riemann zeta-function $\zeta(s)$. We have $d_{\zeta} = 1$ and $q_{\zeta} = 1$.
- Dirichlet L-functions $L(s, \chi)$ for a primitive characters χ modulo $q \ge 2$. We have $d_{L(s,\chi)} = 1$ and $q_{L(s,\chi)} = q$.
- Dedekind zeta-functions $\zeta_{\mathbb{K}}(s)$ for a number fields \mathbb{K} with degree $n_{\mathbb{K}} = [\mathbb{K} : \mathbb{Q}]$ and discriminant $\Delta_{\mathbb{K}}$. We have $d_{\zeta_{\mathbb{K}}} = n_{\mathbb{K}}$ and $q_{\zeta_{\mathbb{K}}} = |\Delta_{\mathbb{K}}|$.

Date: October 30, 2023.

²⁰¹⁰ Mathematics Subject Classification. 11M06, 11M26, 41A30.

Key words and phrases. Selberg class, Generalized Riemann Hypothesis, Bandlimited functions.

We know that the only element in S with the degree zero is the constant function $\mathcal{L} \equiv 1$, and that $d_{\mathcal{L}} \geq 1$ otherwise, see [Ste07, Theorem 6.1]. To simplify our considerations, we will assume that $\mathcal{L} \neq 1$ throughout this paper.

The celebrated Generalized Riemann Hypothesis (GRH) claims that $\mathcal{L}(s) \neq 0$ for $\sigma > 1/2$. The main purpose of the present paper is to provide various estimates for moduli of the logarithmic derivative $(\mathcal{L}'/\mathcal{L})(s)$ and the logarithm $\log \mathcal{L}(s)$ in different regions right of the critical line and under the assumption of GRH. These functions have connections, for example, to primes and primes in arithmetic progressions, see [Dav00, MV07]. For the Riemann zeta-function, Littlewood [Lit25] proved that the Riemann Hypothesis (RH) implies

$$\frac{\zeta'}{\zeta}(s) \ll \left((\log t)^{2-2\sigma} + 1 \right) \min\left\{ \frac{1}{|\sigma - 1|}, \log \log t \right\},\tag{1}$$

$$\log \zeta(s) \ll \frac{(\log t)^{2-2\sigma}}{\log \log t} \min\left\{\frac{1}{|\sigma-1|}, \log \log t\right\} + \log \log \log t \tag{2}$$

for $1/2 + 1/\log \log t \le \sigma \le 3/2$ and t large, see [MV07, Corollaries 13.14 and 13.16]. In particular,

$$\frac{\zeta'}{\zeta}(s) \ll \left(\log t\right)^{2-2\sigma}, \quad \log \zeta(s) \ll \frac{(\log t)^{2-2\sigma}}{\log \log t} \tag{3}$$

for $1/2 + \delta \leq \sigma \leq 1 - \delta$ and fixed $\delta \in (0, 1/4)$. In fact, Littlewood stated only (3), but it is possible to extend his techniques (see also [Tit86, Section 14.5]) to the region right of the 1-line, see [Tit86, Section 14.33]. However, the argument which is based on a slightly modified version of the Selberg moment formula [MV07, Section 13.2] is more transparent, allows generalizations¹ to other functions, and easily yields (1) and (2). Such an approach with an additional feature of estimating the sum over the non-trivial zeros with a help of bandlimited majorants was explored in [CHS23], and the present paper extends this to the Selberg class.

The shape of the aforementioned bounds has never been improved, and efforts have been placed into obtaining explicit constants for the main terms. In 2018, Lumley [Lum18, Theorems 2.1 and 2.2, Corollary 2.3] provided bounds for L(1 + it, f) for a large set of *L*-functions assuming the GRH and Ramanujan-Petersson hold for L(s, f). Recently, such result for (1) was given by Chirre and Gonçalves [CG22, Theorem 1], and for (2) by Carneiro and Chandee [CC11, Theorems 1 and 2]. Our first result is a generalization of (3) for functions in the Selberg class where we also provide explicit constants for the main terms, see Theorem 1. Note that $|a(n)| \leq C(\varepsilon)n^{\varepsilon}$ for any sufficiently small $\varepsilon > 0$ is a quantitative version of the Ramanujan hypothesis.

Theorem 1. Let $\mathcal{L} \in \mathcal{S}$ and assume the Generalized Riemann Hypothesis. Fix $\alpha \geq \frac{1}{2}\log 2$, $\delta \in (0, 1/4)$ and $\varepsilon \in (0, \min{\{\delta, 1/6\}})$. If $|a(n)| \leq C(\varepsilon)n^{\varepsilon}$, then

$$\left|\frac{\mathcal{L}'}{\mathcal{L}}(s)\right| \le \left(A(\mathcal{C}(\varepsilon), \alpha, \sigma - \varepsilon, \sigma) + \frac{\mathcal{C}(\varepsilon)2^{1+\varepsilon}\log 2}{2^{1+\varepsilon} - 2^{\sigma}} + o(1)\right) (\log \tau)^{2(1-\sigma+\varepsilon)}$$
(4)

and

$$\log \mathcal{L}(s)| \le \left(\eta(\alpha, \sigma, \tau) A(\mathcal{C}(\varepsilon), \alpha, \sigma - \varepsilon, \sigma) + \frac{\mathcal{C}(\varepsilon) 2^{1+\varepsilon} \log 2}{2^{1+\varepsilon} - 2^{\sigma}} + o(1)\right) \frac{\left(\log \tau\right)^{2(1-\sigma+\varepsilon)}}{\log \log \tau} \tag{5}$$

for $1/2 + \delta \leq \sigma \leq 1 - \delta$ as $\tau \to \infty$, where

$$A(a, \alpha, u, \sigma) := \frac{a(2\sigma - 1)\left(1 - \exp\left(-\frac{2\alpha(1 - u)}{2\sigma - 1}\right)\right)}{2\alpha(1 - u)^2}$$
(6)

and

$$\eta(\alpha,\sigma,\tau) := \frac{1}{2} \left(1 - \frac{\alpha}{(2\sigma - 1)\log\log\tau} \right)^{-1}.$$
(7)

The implied o-constants may depend on \mathcal{L} .

¹See also [IK04, Theorems 5.17 and 5.19].

The main reason that (4) and (5) are worse by an additional ε in the exponent is in the estimation of the generalized von Mangoldt function $\Lambda_{\mathcal{L}}(n)$. This arithmetic function is defined by

$$\frac{\mathcal{L}'}{\mathcal{L}}(s) = -\sum_{n=1}^{\infty} \frac{\Lambda_{\mathcal{L}}(n)}{n^s}$$

for $\sigma > 1$, and obviously $\Lambda_{\zeta}(n) = \Lambda(n)$ for $\Lambda(n)$ being the von Mangoldt function. We can see that the Euler product representation implies $\Lambda_{\mathcal{L}}(n) = b(n) \log n$ with $\Lambda_{\mathcal{L}}(n) = 0$ for n not being a prime power. Moreover, it is not hard to prove that if $|a(n)| \leq \mathcal{C}(\varepsilon)n^{\varepsilon}$ for $\varepsilon > 0$, then

$$\left|\Lambda_{\mathcal{L}}\left(p^{k}\right)\right| \leq \mathcal{C}(\varepsilon)\left(2^{k}-1\right)p^{k\varepsilon}\log p \tag{8}$$

for $\varepsilon > 0$, see [Mur08, pp. 405–406] for instance. In analogy with $\log \zeta(s)$, we define

$$\log \mathcal{L}(s) = \sum_{n=2}^{\infty} \frac{\Lambda_{\mathcal{L}}(n)}{n^s \log n}$$

for $\sigma > 1$, where the value of $\log \mathcal{L}(s)$ is that which tends to 0 as $\sigma \to \infty$ for any fixed t. For $\sigma < 2$ and $t \neq 0$, we define $\log \mathcal{L}(s)$ as the analytic continuation of the above function along the straight line from 2 + it to $\sigma + it$, provided that $\mathcal{L}(s) \neq 0$ on this segment of line. Note that GRH implies holomorphicity of $\log \mathcal{L}(z)$ on the domain $\{z \in \mathbb{C} : \Re\{z\} > 1/2\} \setminus (1/2, 1]$.

We are using (8) in our proof of Theorem 1, and the factor $p^{k\varepsilon}$ contributes ε in (4) and (5). However, we can substantially improve upon (8) by assuming that $\mathcal{L}(s)$ has a polynomial Euler product representation, i.e., axiom (4) is replaced by

$$\mathcal{L}(s) = \prod_{p} \prod_{j=1}^{m} \left(1 - \frac{\alpha_j(p)}{p^s} \right)^{-1},$$

where $m \in \mathbb{N}$ is the order of a polynomial Euler product and $\alpha_j(p) \in \mathbb{C}$, $1 \leq j \leq m$, are some functions which are defined for every prime number p. Denote by SP the class of such functions. If $\mathcal{L} \in SP$, then

$$b\left(p^{k}\right) = \frac{1}{k} \sum_{j=1}^{m} \left(\alpha_{j}(p)\right)^{k},$$

and because $|\alpha_i(p)| \leq 1$ for $1 \leq j \leq m$, see [Ste07, Lemma 2.2], we have that

$$|\Lambda_{\mathcal{L}}(n)| \le m\Lambda(n). \tag{9}$$

Of course $SP \subseteq S$ and it is conjectured that SP = S. We will show that inequality (9) indeed guarantees the expected conditional estimate for the logarithmic derivative of $\mathcal{L}(s)$, see Theorem 2. Moreover, we can state the result uniformly in \mathcal{L} , i.e., the implied *O*-constants are independent of the parameters from the axioms of the Selberg class unless otherwise indicated by $O_{(\cdot)}$. We would like to provide an opportunity to apply results in various settings and possibility to use sharp estimates. Hence, some of the results are quite long.

Theorem 2. Assume the Generalized Riemann Hypothesis. Let $\mathcal{L} \in S\mathcal{P}$ with a polynomial Euler product of order m and the order of pole of $\mathcal{L}(s)$ at s = 1 is $m_{\mathcal{L}}$. Fix $\alpha \in [0.35, 2)$ and let

$$\lambda^{-} \le \min_{1 \le j \le f} \{\lambda_j\} \le \max_{1 \le j \le f} \{\lambda_j\} \le \lambda^{+}, \quad \max_{1 \le j \le f} \{|\mu_j|\} \le \mu^{+},$$
(10)

where f, λ_j and μ_j are from the functional equation for \mathcal{L} . Then

$$\left|\frac{\mathcal{L}'}{\mathcal{L}}(s)\right| \leq \begin{cases} \widehat{A}\left(m,\alpha,\sigma\right)\left(\log\tau\right)^{2-2\sigma} - \frac{m\sigma2^{1-\sigma}}{1-\sigma} \\ + O\left(m_{\mathcal{L}}\left(\log\tau\right)^{3-4\sigma} + \frac{m}{(2\sigma-1)^{3}}\right) \\ + O\left(\frac{d_{\mathcal{L}}+m_{\mathcal{L}}}{(\log\tau)^{2\sigma-1}} + m_{\mathcal{L}}\left(\frac{q_{\mathcal{L}}}{\tau}\right)^{\frac{2}{d_{\mathcal{L}}}}\left(\log\tau\right)^{2-2\sigma}\right) & \left|\frac{3}{4} - \sigma\right| \leq \frac{1}{4} - \frac{1}{\log\log\tau}, \\ + O_{\lambda^{+},\lambda^{-}}\left(\frac{f}{(\log\tau)^{(2-\alpha)\sigma}} + \frac{f}{(\log\tau)^{2\sigma-1}}\right), \\ 2m\log\log\tau + O\left(m+m|1-\sigma|\left(\log\log\tau\right)^{2}\right) \\ + O\left(\frac{d_{\mathcal{L}}+m_{\mathcal{L}}}{\log\tau} + m_{\mathcal{L}}\left(\frac{q_{\mathcal{L}}}{\tau}\right)^{\frac{2}{d_{\mathcal{L}}}}\right) + O_{\lambda^{+},\lambda^{-}}\left(\frac{f}{\log\tau}\right) & |1-\sigma|\log\log\tau \leq 1 \end{cases}$$
(11)

and

$$\left|\log \mathcal{L}(s)\right| \leq \begin{cases} \widetilde{A}\left(m,\alpha,\sigma,\tau\right) \frac{\left(\log \tau\right)^{2-2\sigma}}{\log\log \tau} - \frac{m\eta(\alpha,\sigma,\tau)}{(1-\sigma)\log\log \tau} + m\log\left(2\log\log \tau\right) \\ + O\left(\frac{m(\log \tau)^{2-2\sigma}}{(1-\sigma)^{2}(\log\log \tau)^{2}} + \frac{m_{\mathcal{L}}(\log \tau)^{3-4\sigma}}{\log\log \tau} + \frac{m}{(2\sigma-1)^{2}}\right) \\ + O\left(\frac{d_{\mathcal{L}}+m_{\mathcal{L}}}{(\log \tau)^{2\sigma-1}\log\log \tau} + m_{\mathcal{L}}\left(\frac{q_{\mathcal{L}}}{\tau}\right)^{\frac{2}{d_{\mathcal{L}}}} \frac{(\log \tau)^{2-2\sigma}}{(\log \sigma)^{2-2\sigma}}\right) \\ + O\left(\frac{d_{\mathcal{L}}}{(\log \tau)^{1-\alpha/2}\log\log \tau}\right) + O_{\lambda^{-}}\left(\frac{f}{(\log \tau)^{2\sigma-1}\log\log \tau}\right) \\ + O_{\lambda^{+}}\left(\frac{f}{(2-\alpha)(\log \tau)^{(2-\alpha)\sigma}\log\log \tau}\right), \\ m\log\left(2\log\log \tau\right) + O\left(m + \frac{d_{\mathcal{L}}+m_{\mathcal{L}}}{\log \tau\log\log \tau} + \left(\frac{q_{\mathcal{L}}}{\tau}\right)^{\frac{2}{d_{\mathcal{L}}}} \frac{m_{\mathcal{L}}}{\log\log \tau}\right) \\ + O_{\lambda^{+},\lambda^{-}}\left(\frac{f}{\log \tau\log\log \tau}\right), \end{cases}$$

$$(12)$$

for sufficiently large τ and $|t| \ge \mu^+ / \lambda^-$. Here,

$$\widehat{A}(m,\alpha,\sigma) := A(m,\alpha,\sigma,\sigma) + \frac{e^{\alpha} + 1}{2\alpha}, \qquad (13)$$

$$\widetilde{A}(m,\alpha,\sigma,\tau) := \eta(\alpha,\sigma,\tau)A(m,\alpha,\sigma,\sigma) + \frac{e^{\alpha}+1}{4\alpha},$$
(14)

and $A(a, \alpha, u, \sigma)$ and $\eta(\alpha, \sigma, \tau)$ are defined by (6) and (7), respectively.

An immediate consequence of Theorem 2 is that GRH implies the following corollary.

Corollary 1. Assume the same assumptions as in Theorem 2. Then we have

$$\left|\frac{\mathcal{L}'}{\mathcal{L}}(s)\right| \le \left(\widehat{A}(m,\alpha,\sigma) + o(1)\right) \left(\log \tau\right)^{2-2\sigma},\tag{15}$$

$$\left|\log \mathcal{L}(s)\right| \le \left(\widetilde{A}\left(m, \alpha, \sigma, \tau\right) + o(1)\right) \frac{\left(\log \tau\right)^{2-2\sigma}}{\log\log \tau}$$
(16)

for $\mathcal{L} \in S\mathcal{P}$, $1/2 + \delta \leq \sigma \leq 1 - \delta$, $\tau \to \infty$ and fixed $\delta \in (0, 1/4)$, where the implied o-constants may depend on \mathcal{L} .

Comparing the estimates from Corollary 1 with known results reveals the following:

- Estimate (15) for $\mathcal{L}(s) = \zeta(s)$ and $\alpha = 3/4$ improves [CG22, Theorem 1]. This was already announced in [CHS23, Estimate (1.5)], although without a detailed proof.
- Estimate (16) for $\mathcal{L}(s) = \zeta(s)$ and $\alpha = 3/2$ improves [CC11, Theorems 1 and 2] in the range $0.565 \le \sigma \le 1 \delta$ for fixed $\delta \in (0, 0.435)$.
- Chirre [Chi19] obtained an estimate for $\log |L(s,\pi)|$, where $L(s,\pi)$ is an entire L-function satisfying special conditions, see [Chi19, Section 1.1] for a definition. Estimate (16) for $\mathcal{L}(s) = L(s,\pi)$, $m = d_{\mathcal{L}} \geq 3$ and $\alpha = 1.6$ improves [Chi19, Corollary 2] in the range $0.56 \leq \sigma \leq 0.999$.
- Carneiro, Chirre and Milinovich [CCM19] obtained the RH estimate for $S_{0,\sigma}(t) = \frac{1}{\pi} \arg \zeta(\sigma + it)$. Note that $|S_{0,\sigma}(t)| \leq \frac{1}{\pi} |\log \zeta(s)|$ for $\sigma \in (1/2, 1)$ and under RH. Estimate (16) for $\mathcal{L}(s) = \zeta(s)$ and $\alpha = 1.38$ then improves [CCM19, Corollary 5] in the range $0.55 \leq \sigma \leq 1 - \delta$ for fixed $\delta \in (0, 0.45)$.

It is clear by the above that our bounds are weaker for σ close to the critical line when compared with known results. It might be interesting to see how much of the results and methods from [CC11, CG22] we can recover for functions in the class SP. In addition to this, generalizing the results for $S_{n,\sigma}(t)$ from [CCM19] could also be worthwhile to perform. Recently, explicit and conditional upper bounds for $|S_{0,1/2}(t)|$ and $|S_{1,1/2}(t)|$ appeared in [Sim22b].

For s = 1 + it we are able to improve upon the constant terms from Theorem 2, see the next theorem.

Theorem 3. Let $\mathcal{L} \in S\mathcal{P}$ with a polynomial Euler product of order m and assume the Generalized Riemann Hypothesis. Take $\alpha \geq \frac{1}{2}\log 2$. Then

$$\left|\frac{\mathcal{L}'}{\mathcal{L}}(1+\mathrm{i}t)\right| \leq 2m\log\log\tau - m\left(\gamma+\alpha\right) + \frac{e^{\alpha}+1}{2\alpha} + O\left(\frac{m\left(\log\log\tau\right)^2}{\log\tau} + \frac{\mathrm{d}_{\mathcal{L}}+m_{\mathcal{L}}}{\log\tau} + m_{\mathcal{L}}\left(\frac{\mathrm{q}_{\mathcal{L}}}{\tau}\right)^{\frac{2}{\mathrm{d}_{\mathcal{L}}}}\right) + O_{\lambda^+,\lambda^-}\left(\frac{f}{\log\tau}\right) \quad (17)$$

and

$$\log \mathcal{L}(1+\mathrm{i}t)| \leq m \log \left(2 \log \log \tau\right) + m\gamma + \frac{e^{\alpha} + 1}{4\alpha \log \log \tau} + O\left(\frac{m \log \log \tau}{\log \tau} + \frac{\mathrm{d}_{\mathcal{L}} + m_{\mathcal{L}}}{\log \tau \log \log \tau} + \frac{m_{\mathcal{L}}}{\log \log \tau} \left(\frac{\mathrm{q}_{\mathcal{L}}}{\tau}\right)^{\frac{2}{\mathrm{d}_{\mathcal{L}}}}\right) + O_{\lambda^{+},\lambda^{-}}\left(\frac{f}{\log \tau \log \log \tau}\right)$$
(18)

for sufficiently large τ and $|t| \ge \mu^+/\lambda^-$, where λ^+ , λ^- and μ^+ are as in (10), and γ is the Euler-Mascheroni constant.

Estimate (17) is a generalisation of [CHS23, Theorem 5] and (18) is a generalisation of a well-known result by Littlewood [Lit25, Theorem 7] that

$$\zeta(1+\mathrm{i}t)| \le (2e^{\gamma} + o(1))\log\log t \tag{19}$$

for large t. In addition to (19), Littlewood [Lit28] also proved that

$$\frac{1}{|\zeta(1+\mathrm{i}t)|} \le \left(\frac{12e^{\gamma}}{\pi^2} + o(1)\right) \log\log t.$$
(20)

We are not able to extend (20) to the class SP because just adapting Littlewood's strategy to deal with a sum over prime numbers, see [CC11, p. 377], would require special properties for $\Lambda_{\mathcal{L}}(n)$; note that $\Lambda_{L(s,\chi)}(n) = \chi(n)\Lambda(n)$ and that $\chi(n)$ is a multiplicative function with modulus not greater than 1. This allows to deduce (20) also for Dirichlet *L*-functions. Similar although more precise methods were developed in [LLS15] to obtain explicit estimates of the same shape as (19) and (20) for $\zeta(1 + it)$ and $L(1,\chi)$, see also [Lum18]. Unfortunately, our estimate (18) for $\mathcal{L}(s) = \zeta(s)$ is slightly weaker than theirs. It might be interesting to investigate a possibility to improve (18) by adapting techniques from [LLS15].

Recently, explicit and conditional results on (ζ'/ζ) (s) or log $\zeta(s)$, in different regions right of the critical line and for other *L*-functions as well, appeared in [IMS09, LLS15, LT22, Sim22a, Sim23, CHS23]. Such results are useful, e.g., in the theory of distribution of prime numbers [BMOR18, EHP22] and for studying the Mertens function [Sim23]. One advantage of our approach is a possibility to obtain effective versions of Theorems 2 and 3, see Theorems 4 and 5. Although there is no serious obstacle in the method, we are additionally assuming also the *strong* λ -conjecture, i.e., all λ_j from the functional equation can be chosen to be equal to 1/2, in order to simplify the proof. The main results in this direction are too complicated to be written here and we refer to Section 2 for this. However, as an application of Theorem 4, we are stating "ready to use" results in the case of the Riemann zeta-function, see Corollary 2.

Corollary 2. Let $t \ge 10^6$, $1/2 + 1/\log \log t \le \sigma < 1$ and assume the Riemann Hypothesis. Then

$$\left|\frac{\zeta'}{\zeta}(s)\right| \le \widehat{A}\left(1,\frac{3}{4},\sigma\right) \left(\log t\right)^{2-2\sigma} - \frac{\sigma 2^{1-\sigma}}{1-\sigma} + 4.81\left(\log t\right)^{3-4\sigma} + \frac{0.64}{(2\sigma-1)^3} + \frac{3.76\left(\log\log t\right)^2}{(2\sigma-1)\left(\log t\right)^{2\sigma-1}}$$
(21)

and

$$\begin{aligned} |\log \zeta(s)| &\leq \widetilde{A}\left(1, \frac{3}{2}, \sigma, t\right) \frac{(\log t)^{2-2\sigma}}{\log \log t} - \frac{\eta(3/2, \sigma, t)}{(1-\sigma)\log \log t} + \log\left(2\log\log t\right) + \frac{5.831\left(\log t\right)^{2-2\sigma}}{\left(1-\sigma\right)^2\left(\log\log t\right)^2} \\ &+ \frac{1.06\left(\log t\right)^{3-4\sigma}}{\log\log t} + \frac{5.6}{(2\sigma-1)^2} + \frac{1.58\log\log t}{\left(\log t\right)^{2\sigma-1}} + \frac{10.6}{\left(\log t\right)^{\frac{1}{4}}\log\log t} + \frac{0.24\log\log t}{\sqrt{\log t}}, \end{aligned}$$
(22)

where functions $\widehat{A}(1,3/4,\sigma)$, $\widetilde{A}(1,3/2,\sigma,t)$ and $\eta(3/2,\sigma,t)$ are defined by (13), (14) and (7), respectively.

The idea of the proofs is to write the function $\mathcal{L}'(s)/\mathcal{L}(s)$ as sums over primes and a sum over zeros of the function $\mathcal{L}(s)$ using Selberg's moment formula. Then we show that this leads to a similar kind of an estimate for the function $\log \mathcal{L}(s)$, and we estimate these sums using partial summation and the Guinand–Weil exact formula for the Selberg class.

The outline of this paper is as follows. In Section 2 we formulate effective versions of Theorems 2 and 3, and also provide the proof of Corollary 2. In Section 3 we revise Selberg's moment formula for functions in the Selberg class, and in Section 4 we derive estimates for the corresponding sums over prime numbers. Section 5 is devoted to the extension of the Guinand–Weil exact formula to the Selberg class and estimation of the sum over the non-trivial zeros. The proofs of Theorems 1, 2 and 3 are presented in Section 6, while the proofs of Theorems 4 and 5 are provided in Section 7.

Acknowledgements. A.S. would like to thank N.P. and The University of Helsinki for their hospitality during his visit in June 2022. He also gratefully acknowledges the Lift-off Fellowship support from The Australian Mathematical Society. N. P. would like to thank the Emil Aaltonen Foundation for supporting her work. The authors would like to thank Tim Trudgian, Jörn Steuding and Andrés Chirre for taking time to read the manuscript.

2. Effective results

In this section we are stating explicit versions of Theorems 2 and 3 under additional assumption of the strong λ -conjecture. Moreover, we also prove Corollary 2.

Theorem 4. Let $\mathcal{L} \in S\mathcal{P}$ and assume the Generalized Riemann Hypothesis and the strong λ -conjecture. Consider the following three cases:

(1) Let $2\alpha_1 > \alpha \geq \frac{1}{2}\log 2$ and

$$\frac{1}{2} + \frac{\alpha_1}{\log\log\tau} \le \sigma < 1, \tag{23}$$

$$\sqrt{60} \exp\left(2^{1/(2-\alpha/\alpha_1)}\right) \exp\left(2^{2\alpha_1}\right) = |t| \ge t \ge \max\left\{2 \exp\left(|t_1|\right) + 1\right\}$$

$$\tau \ge \tau_0 > \max\left\{e^{\sqrt{60}}, \exp\left(2^{1/(2-\alpha/\alpha_1)}\right), \exp\left(e^{2\alpha_1}\right)\right\}, \quad |t| \ge t_0 \ge \max\left\{2\max_{1\le j\le f}\left\{|\mu_j|\right\}, 1\right\}.$$

(2) Let $\alpha \ge \log 2$, $\alpha_2 > 0$, $\alpha_3 > 0$ and

$$1 - \frac{\alpha_2}{\log\log\tau} \le \sigma \le 1 + \frac{\alpha_3}{\log\log\tau},$$

$$\tau \ge \tau_0 > \max\left\{e^{\sqrt{60}}, \exp\left(\sqrt{2}\exp\left(\frac{\alpha}{2\sigma-1}\right)\right), \exp\left(e^{\alpha+2\alpha_2}\right), \exp\left(e^{4\alpha_2}\right), \exp\left(e^{2\alpha_3}\right)\right\},$$

$$|t| \ge t_0 \ge \max\left\{2\max_{1\le j\le f}\left\{|\mu_j|\right\}, 1\right\}.$$

$$(24)$$

(3) Let $\alpha_3 > 0$, $\tau > e$ and $\sigma \ge 1 + \alpha_3 / \log \log \tau$. Under the conditions from (1) we have

$$\left|\frac{\mathcal{L}'}{\mathcal{L}}(s)\right| \leq \widehat{A}\left(m,\alpha,\sigma\right)\left(\log\tau\right)^{2-2\sigma} - \frac{m\sigma2^{1-\sigma}}{1-\sigma} + \frac{\mathfrak{a}\left(e^{\alpha}+1\right)}{\alpha}\left(\log\tau\right)^{3-4\sigma} + \frac{m\sigma2^{\frac{3}{2}-\sigma}\left(4+\left(2+\left(2\sigma-1\right)\log2\right)^{2}\right)}{8\pi(2\sigma-1)^{3}} + \left(\frac{m\left(2\left(\alpha+1\right)\sigma-1\right)\left(e^{\alpha}-1\right)}{2\pi\alpha}\right)\frac{\left(\log\log\tau\right)^{2}}{(2\sigma-1)\left(\log\tau\right)^{2\sigma-1}} + \left(\frac{2m\left(e^{\alpha}+1\right)}{\alpha}\right)\frac{\log\log\tau}{\left(\log\tau\right)^{2\sigma-1}} + \frac{\mathfrak{b}\left(e^{\alpha}+1\right)}{\alpha\left(\log\tau\right)^{2\sigma-1}} + \frac{4.3\mathrm{d}_{\mathcal{L}}\left(\sigma-\frac{1}{2}\right)\left(1+\left(\log\tau_{0}\right)^{-\frac{\alpha}{\alpha_{1}}\sigma}\right)}{\alpha\left(\log\tau\right)^{\left(2-\frac{\alpha}{\alpha_{1}}\right)\sigma}} + \frac{m_{\mathcal{L}}}{\alpha}\left(\frac{\mathrm{q}_{\mathcal{L}}}{\tau}\right)^{\frac{2}{\mathrm{d}_{\mathcal{L}}}}\left(\log\tau\right)^{2\left(1-\sigma\right)}$$
(25)

and

$$\begin{split} |\log \mathcal{L}(s)| &\leq \widetilde{A}\left(m, \alpha, \sigma, \tau\right) \frac{\left(\log \tau\right)^{2-2\sigma}}{\log \log \tau} - \frac{m\eta(\alpha, \sigma, \tau)}{(1-\sigma)\log \log \tau} + m\log\left(2\log\log \tau\right) \\ &+ m\left(\left(\nu_2\eta(\alpha, \sigma, \tau)\right)^2 \exp\left(-\frac{2\alpha(1-\sigma)}{2\sigma-1}\right)\right) \frac{\left(\log \tau\right)^{2-2\sigma}}{(1-\sigma)^2 \left(\log\log \tau\right)^2} + \frac{\mathfrak{a}\left(e^{\alpha}+1\right)\left(\log \tau\right)^{3-4\sigma}}{4\alpha\log\log \tau} \\ &+ \frac{m2^{\frac{3}{2}-\sigma}\left(-1+\sigma\left(4-\log 2\right)+\sigma^2\log 4\right)}{8\pi\left(2\sigma-1\right)^2} + \frac{m}{\nu_1^2}\exp\left(-\frac{2\alpha(1-\sigma)}{(2\sigma-1)\nu_2}\right)\left(\log \tau\right)^{\frac{2}{\nu_2}(1-\sigma)} \\ &+ m\left(\frac{1-\sigma2^{1-\sigma}}{(1-\sigma)\log 2} - \log\log 2 + \int_0^{\nu_1}\theta_1(u)du\right) + \frac{m\left(e^{\alpha}-1\right)\log\log \tau}{4\pi\alpha\left(\log \tau\right)^{2\sigma-1}} + \frac{m\left(e^{\alpha}+1\right)}{\alpha\left(\log \tau\right)^{2\sigma-1}} \\ &+ \frac{\max\{0,\mathfrak{b}\}\left(e^{\alpha}+1\right)}{2\alpha\left(\log \tau\right)^{2\sigma-1}\log\log \tau} + \frac{2m\eta\left(\alpha,\sigma,\tau\right)\exp\left(\frac{\alpha}{2\sigma-1}\right)}{\log \tau\log\log \tau} + \frac{5m\exp\left(\frac{2\alpha}{2\sigma-1}\right)\left(1+\frac{\log\log \tau}{\eta(\alpha,\sigma,\tau)}\right)}{16\pi\log^2 \tau} \end{split}$$

$$+\frac{4.3d_{\mathcal{L}}\alpha_{1}\left(1+\left(\log\tau_{0}\right)^{-\frac{\alpha}{2\alpha_{1}}}\right)}{\alpha\left(2\alpha_{1}-\alpha\right)\left(\log\tau\right)^{\left(2-\frac{\alpha}{\alpha_{1}}\right)\sigma}\log\log\tau}+\frac{m_{\mathcal{L}}\left(1+e^{\frac{\alpha}{2}}\right)}{2\alpha}\left(\frac{q_{\mathcal{L}}}{\tau}\right)^{\frac{2}{d_{\mathcal{L}}}}\frac{\left(\log\tau\right)^{2-2\sigma}}{\log\log\tau}.$$
(26)

Here, $\widehat{A}(m, \alpha, \sigma)$ and $\widetilde{A}(m, \alpha, \sigma, \tau)$ are defined by (13) and (14), respectively, and the functions $\mathfrak{a} = \mathfrak{a}(m_{\mathcal{L}}, \alpha_1, t_0)$ and $\mathfrak{b} = \mathfrak{b}(d_{\mathcal{L}}, m, m_{\mathcal{L}}, \alpha_1, t_0, \tau_0)$ are defined by (62) and (63), respectively. Additionally, $\nu_1 > 0$ and $\nu_2 > 1$, and $\eta(\alpha, \sigma, \tau)$ and $\theta_1(u)$ are defined by (7) and (29), respectively.

Under the conditions from (2) we have

$$\begin{aligned} \left| \frac{\mathcal{L}'}{\mathcal{L}}(s) \right| &\leq 2m \log \log \tau + m \left(1 - \sigma \log 2\right) + \frac{e^{\alpha} + 1}{2\alpha} \\ &+ \left(4m\theta_1(M) + \frac{(e^{\alpha} + 1)\theta_2(M)}{\alpha \log \log \tau_0} + \frac{m\sigma(\log 2)^2\theta_1\left(\frac{M\log 2}{2\log \log \tau_0}\right)}{(\log \log \tau_0)^2} \right) |1 - \sigma| \left(\log \log \tau\right)^2 \\ &+ \frac{m\sigma 2^{\frac{3}{2} - \sigma} \left(4 + \left(2 + \left(2\sigma - 1\right)\log 2\right)^2\right)}{8\pi(2\sigma - 1)^3} + \left(\frac{m\left(2\left(\alpha + 1\right)\sigma - 1\right)\left(e^{\alpha} - 1\right)}{2\pi\alpha}\right) \frac{(\log \log \tau)^2}{(2\sigma - 1)(\log \tau)^{2\sigma - 1}} \\ &+ \left(\frac{2m\left(e^{\alpha} + 1\right)}{\alpha}\right) \frac{\log \log \tau}{(\log \tau)^{2\sigma - 1}} + \frac{e^{\alpha} + 1}{\alpha} \left(\frac{\mathfrak{a}_1}{(\log \tau)^{4\sigma - 3}} + \frac{\mathfrak{b}_1}{(\log \tau)^{2\sigma - 1}}\right) \\ &+ \frac{4.3d_{\mathcal{L}}e^{2\alpha_2}\left(\frac{1}{2} + \frac{\alpha_3}{\log\log \tau_0}\right)\left(1 + \exp\left(\frac{2\alpha(\log\log \tau_0 - \alpha_2)}{\log\log \tau_0 - 2\alpha_2}\right)\right)}{\alpha\left(\log \tau\right)^2} \\ &+ \frac{m_{\mathcal{L}}e^{2\alpha_2}}{\alpha}\left(\frac{1}{2} + \frac{\alpha_3}{\log\log \tau_0}\right)\left(1 + \exp\left(\frac{2\alpha\alpha_3}{\log\log \tau_0 + 2\alpha_3}\right)\right)\left(\frac{q_{\mathcal{L}}}{\tau}\right)^{\frac{2}{d_{\mathcal{L}}}} \end{aligned}$$
(27)

and

$$\begin{aligned} |\log \mathcal{L}(s)| &\leq m \log \left(2 \log \log \tau\right) + m \left(\frac{1}{\log 2} - \log \log 2 - \sigma + \frac{1}{1 - \frac{\alpha}{\log \log \tau_0 - 2\alpha_2}}\right) \\ &+ \left(\frac{2m\theta_1(M)}{1 - \frac{\alpha}{\log \log \tau_0 - 2\alpha_2}} + 2m\theta_1(2\alpha_2) + \frac{m\sigma \left(\log 2\right)\theta_1\left(\frac{M\log 2}{2\log \log \tau_0}\right)}{\log \log \tau_0}\right)|1 - \sigma|\log \log \tau \\ &+ \frac{m2^{\frac{3}{2} - \sigma}\left(-1 + \sigma\left(4 - \log 2\right) + \sigma^2\log 4\right)}{8\pi \left(2\sigma - 1\right)^2} + \frac{m\left(e^{\alpha} - 1\right)\log \log \tau}{4\pi\alpha \left(\log \tau\right)^{2\sigma - 1}} + \frac{\left(e^{\alpha} + 1\right)\left(\log \tau\right)^{2-2\sigma}}{4\alpha \log \log \tau} \\ &+ \frac{e^{\alpha} + 1}{4\alpha \log \log \tau}\left(\frac{\mathfrak{a}_1}{\left(\log \tau\right)^{4\sigma - 3}} + \frac{2\max\{0, \mathfrak{b}_1\}}{\left(\log \tau\right)^{2\sigma - 1}}\right) + \frac{m\left(e^{\alpha} + 1\right)}{\alpha \left(\log \tau\right)^{2\sigma - 1}} \\ &+ \frac{m\exp\left(\frac{\alpha \log \log \tau_0}{\log \log \tau_0 - 2\alpha_2}\right)}{\left(1 - \frac{\alpha}{\log \log \tau_0 - 2\alpha_2}\right)\log \tau \log \log \tau} + \frac{5m\exp\left(\frac{2\alpha \log \log \tau_0}{\log \log \tau_0 - 2\alpha_2}\right)\left(1 + 2\log \log \tau\right)}{16\pi \left(\log \tau\right)^2} \\ &+ \frac{4.3d_{\mathcal{L}}\left(1 + \exp\left(\frac{2\alpha (\log \log \tau_0 - \alpha_2)}{\log \log \tau}\right)\right)}{2\alpha \left(\log \tau\right)^{2\sigma} \log \log \tau} + \frac{m\underline{\mathcal{L}}e^{2\alpha_2}\left(1 + e^{\frac{\alpha}{2}}\right)}{2\alpha \log \log \tau}\left(\frac{q_{\mathcal{L}}}{\tau}\right)^{\frac{2}{\alpha_{\mathcal{L}}}}. \end{aligned}$$

$$\tag{28}$$

Here,

$$M := 2 \max\{\alpha_2, \alpha_3\}, \quad and \quad \theta_1(u) := \frac{e^u - u - 1}{u^2}, \quad \theta_2(u) := \frac{e^u - 1}{u}$$
(29)

for u > 0. Additionally, $\mathfrak{a}_1 := \mathfrak{a}\left(m_{\mathcal{L}}, \frac{1}{2}\log\log\tau_0 - \alpha_2, t_0\right)$ and $\mathfrak{b}_1 := \mathfrak{b}\left(\mathrm{d}_{\mathcal{L}}, m, m_{\mathcal{L}}, \frac{1}{2}\log\log\tau_0 - \alpha_2, t_0, \tau_0\right)$ with \mathfrak{a} and \mathfrak{b} defined by (62) and (63), respectively.

Under the conditions from (3) we have

$$\left|\frac{\mathcal{L}'}{\mathcal{L}}(s)\right| \le \frac{m}{\alpha_3} \log\log\tau, \quad \left|\log\mathcal{L}(s)\right| \le m \log\left(\frac{1}{\alpha_3}\log\log\tau\right) + \frac{m\gamma\alpha_3}{\log\log\tau},\tag{30}$$

where γ is the Euler–Mascheroni constant.

Proof of Corollary 2. We are using the first part of Theorem 4 for $d_{\mathcal{L}} = q_{\mathcal{L}} = m = \alpha_1 = 1$ and $\tau_0 = t_0 = 10^6$, and thus $\tau = |t|$, in order to prove both (21) and (22). However, we additionally set $\alpha = 3/4$ to prove (21), and $\alpha = 3/2$ to prove (22). Note that the conditions from the case (1) of Theorem 4

are satisfied in either case. Then (21) easily follows by observing that the last three terms on the right-hand side of (25) can be omitted because their sum is negative. To prove (22), note first that $\eta = \eta(3/2, \sigma, \tau) \in [1/2, 2]$ for σ from the range (23). Then the sum of the fourth term and the seventh term on the right-hand side of (26) is not greater than

$$\left(4\nu_2^2 + \frac{(1-\sigma)^2 \left(\log\log t\right)^2}{\nu_1^2 \left(\log t\right)^{2(1-1/\nu_2)(1-\sigma)}}\right) \frac{\left(\log t\right)^{2-2\sigma}}{(1-\sigma)^2 \left(\log\log t\right)^2} \le \left(4\nu_2^2 + \frac{1}{e^2\nu_1^2 \left(1-\frac{1}{\nu_2}\right)^2}\right) \frac{\left(\log t\right)^{2-2\sigma}}{\left(1-\sigma\right)^2 \left(\log\log t\right)^2}.$$

The parameters ν_1 and ν_2 allow some optimisation on various terms. Observe that for $\sigma = 1 - 1/\log \log t$ the right-hand side of the above inequality is constant. Therefore, we choose $\nu_1 = 3.76$ and $\nu_2 = 1.13$ in order to minimize

$$4\nu_2^2 e^2 + \frac{1}{\nu_1^2 \left(1 - \frac{1}{\nu_2}\right)^2} + \int_0^{\nu_1} \theta_1(u) \mathrm{d}u.$$

The fourth term and the sixth term on the right-hand side of (22) now easily follow. The seventh term and the eight term follow after merging the ninth, tenth and fifteenth term, and the twelfth and fourteenth term on the right-hand side of (26), respectively. The remaining terms in (22) are easy to deduce.

Theorem 5. Let $\mathcal{L} \in S\mathcal{P}$ and assume the Generalized Riemann Hypothesis and the strong λ -conjecture. If $\alpha \geq \frac{1}{2} \log 2$ and

$$\tau \ge \tau_0 \ge \exp\left(e^{\alpha}\sqrt{60}\right), \quad |t| \ge t_0 \ge \max\left\{2\max_{1\le j\le f}\left\{|\mu_j|\right\}, 1\right\},$$

then

$$\left|\frac{\mathcal{L}'}{\mathcal{L}}(1+\mathrm{i}t)\right| \leq 2m\log\log\tau - m\left(\gamma+\alpha\right) + \frac{e^{\alpha}+1}{2\alpha} + \left(\frac{m\left(e^{\alpha}-1\right)\left(2\alpha+1\right)}{2\pi\alpha}\right)\frac{\left(\log\log\tau\right)^{2}}{\log\tau} + \left(\frac{2m\left(e^{\alpha}+1\right)}{\alpha}\right)\frac{\log\log\tau}{\log\tau} + \left(0.24me^{\alpha}+\frac{1}{\alpha}\left(e^{\alpha}+1\right)\left(\mathfrak{a}_{2}+\mathfrak{b}_{2}\right)\right)\frac{1}{\log\tau} + \frac{2.15d_{\mathcal{L}}\left(e^{2\alpha}+1\right)}{\alpha\log^{2}\tau} + \frac{m_{\mathcal{L}}}{\alpha}\left(\frac{q_{\mathcal{L}}}{\tau}\right)^{\frac{2}{\alpha}}$$
(31)

and

$$\begin{split} \log \mathcal{L}(1+\mathrm{i}t) &| \leq m \log \left(2 \log \log \tau\right) + m \left(\gamma + \left(1 - \frac{\log \log \tau}{\alpha}\right) \log \left(1 - \frac{\alpha}{\log \log \tau}\right) - 1\right) \\ &+ \frac{e^{\alpha} + 1}{4\alpha \log \log \tau} + \left(\frac{m \left(1 + e^{\alpha} \left(4\alpha - 1\right)\right)}{4\pi \alpha}\right) \frac{\log \log \tau}{\log \tau} \\ &+ \frac{m}{\alpha} \left(e^{\alpha} + 1 + \frac{e^{\alpha} \left(7\alpha - 4\left(\alpha^{2} + 1\right)\right) + 4}{4\pi}\right) \frac{1}{\log \tau} \\ &+ \left(\frac{\left(e^{\alpha} + 1\right)\left(\mathfrak{a}_{2} + 2\max\{0, \mathfrak{b}_{2}\}\right)}{4\alpha} + \frac{me^{\alpha} \log \log \tau_{0}}{\log \log \tau_{0} - \alpha}\right) \frac{1}{\log \tau \log \log \tau} \\ &+ \frac{5me^{2\alpha} \log \log \tau}{8\pi \log^{2} \tau} + \frac{5me^{2\alpha} \left(1 - 2\alpha\right)}{16\pi \log^{2} \tau} + \frac{2.15 \mathrm{d}_{\mathcal{L}} \left(e^{2\alpha} + 1\right)}{\alpha \left(\log \tau\right)^{2} \log \log \tau} + \frac{m_{\mathcal{L}} \left(1 + e^{\frac{\alpha}{2}}\right)}{2\alpha \log \log \tau} \left(\frac{q_{\mathcal{L}}}{\tau}\right)^{\frac{2}{\alpha}}, (32) \end{split}$$

where $\mathfrak{a}_2 := \mathfrak{a}\left(m_{\mathcal{L}}, \frac{1}{2}\log\log\tau_0, t_0\right)$ and $\mathfrak{b}_2 := \mathfrak{b}\left(\mathrm{d}_{\mathcal{L}}, m, m_{\mathcal{L}}, \frac{1}{2}\log\log\tau_0, t_0, \tau_0\right)$ with \mathfrak{a} and \mathfrak{b} defined by (62) and (63), respectively, and γ is the Euler–Mascheroni constant.

3. The Selberg moment formula for functions in ${\mathcal S}$

Selberg [Sel43, Lemma 2] discovered an interesting connection (also known as the Selberg moment formula) between the logarithmic derivative of the Riemann zeta-function and special truncated Dirichlet series. In this paper we follow the same idea. Hence, the following lemma generalizes Selberg's result to functions in the Selberg class and follows from using similar contour integral as in [MV07, the beginning of Section 13.2].

Lemma 1. Assume that $\mathcal{L} \in \mathcal{S}$ and $\mathcal{L}(1) \neq 0$. Let $q_j(k) := (k + \mu_j) / \lambda_j$ for $j \in \{1, \ldots, f\}$ and $k \in \mathbb{N}_0$. Take $x \geq 2$ and $y \geq 2$. Then

$$\frac{\mathcal{L}'}{\mathcal{L}}(s) = -\sum_{n \le xy} \frac{\Lambda_{\mathcal{L},x,y}(n)}{n^s} + m_{\mathcal{L}} \frac{(xy)^{1-s} - x^{1-s}}{(1-s)^2 \log y} + \frac{1}{\log y} \sum_{j=1}^f \sum_{k=0}^\infty \frac{x^{-q_j(k)-s} - (xy)^{-q_j(k)-s}}{(q_j(k)+s)^2} + \frac{1}{\log y} \sum_{\rho} \frac{x^{\rho-s} - (xy)^{\rho-s}}{(\rho-s)^2}$$
(33)

for $s \notin \{-q_j(k) \colon k \in \mathbb{N}_0, 1 \leq j \leq f\} \cup \{1\}$ and $s \neq \rho$, where

$$\Lambda_{\mathcal{L},x,y}(n) := \begin{cases} \Lambda_{\mathcal{L}}(n), & 1 \le n \le x, \\ \Lambda_{\mathcal{L}}(n) \frac{\log\left(xy/n\right)}{\log y}, & x < n \le xy. \end{cases}$$

Let $\sigma > 1/2$ and take

$$y = \exp\left(\frac{\alpha}{\sigma - \frac{1}{2}}\right), \quad x = y^{-1}\log^2 \tau.$$
 (34)

It is not hard to verify that $x \ge 2$ and $y \ge 2$ under the conditions (1) and (2) from Theorem 4, and also under conditions from Theorem 5. Therefore, we can use Lemma 1 for x and y which are defined by (34) in order to prove these theorems. Moreover, because these conditions are independent on the assumptions of the strong λ -conjecture or a polynomial Euler product, we can use the same approach also to prove Theorems 1, 2 and 3.

Let us bound the terms on the right-hand side of (33). The first term is estimated easily by

$$\left| \sum_{n \le xy} \frac{\Lambda_{\mathcal{L},x,y}(n)}{n^s} \right| \le \sum_{n \le x} \frac{|\Lambda_{\mathcal{L}}(n)|}{n^\sigma} + \frac{1}{\log y} \sum_{x < n \le xy} \frac{|\Lambda_{\mathcal{L}}(n)| \log \frac{xy}{n}}{n^\sigma} =: S_{\mathcal{L},x,y}(\sigma).$$
(35)

Under GRH we can estimate the sum over zeros as

$$\frac{1}{\log y} \left| \sum_{\rho} \frac{x^{\rho-s} - (xy)^{\rho-s}}{(\rho-s)^2} \right| \leq \frac{(xy)^{\frac{1}{2}-\sigma} \left(1+y^{\sigma-\frac{1}{2}}\right)}{\log y} \sum_{\rho} \frac{1}{|\rho-s|^2} \leq \frac{e^{\alpha} + 1}{\alpha} \left(\log \tau\right)^{1-2\sigma} \sum_{\gamma} \frac{\sigma - \frac{1}{2}}{\left(\sigma - \frac{1}{2}\right)^2 + (t-\gamma)^2} =: \mathcal{Z}(\sigma).$$
(36)

We bound the remaining two terms as

$$\frac{1}{\log y} \left| \sum_{j=1}^{f} \sum_{k=0}^{\infty} \frac{x^{-q_j(k)-s} - (xy)^{-q_j(k)-s}}{(q_j(k)+s)^2} \right| \leq \frac{(xy)^{-\sigma} (1+y^{\sigma})}{\log y} \sum_{j=1}^{f} \sum_{k=0}^{\infty} \left(\frac{k}{\lambda_j} + \frac{1}{2}\right)^{-2} \leq \frac{\left(\sigma - \frac{1}{2}\right) \left(1 + \exp\left(\frac{2\alpha\sigma}{2\sigma-1}\right)\right)}{\alpha \left(\log \tau\right)^{2\sigma}} \sum_{j=1}^{f} \lambda_j^2 \left(\frac{\Gamma'}{\Gamma}\right)' \left(\frac{\lambda_j}{2}\right) =: \mathcal{R}_1(\sigma) \quad (37)$$

and

$$m_{\mathcal{L}} \left| \frac{(xy)^{1-s} - x^{1-s}}{(1-s)^2 \log y} \right| \le \frac{m_{\mathcal{L}} \left(\sigma - \frac{1}{2} \right) \left(1 + \exp\left(\frac{2\alpha(\sigma-1)}{2\sigma-1} \right) \right)}{\alpha} \left(\frac{q_{\mathcal{L}}}{\tau} \right)^{\frac{2}{d_{\mathcal{L}}}} \left(\log \tau \right)^{2-2\sigma} =: \mathcal{R}_2(\sigma).$$
(38)

An obvious strategy to produce the desired bounds for \mathcal{L}'/\mathcal{L} is to employ estimates (35)–(38) into the Selberg moment formula (33), that is

$$\left|\frac{\mathcal{L}'}{\mathcal{L}}(s)\right| \le S_{\mathcal{L},x,y}(\sigma) + \mathcal{Z}(\sigma) + \mathcal{R}_1(\sigma) + \mathcal{R}_2(\sigma).$$
(39)

However, in order to do so we need to obtain suitable bounds for $S_{\mathcal{L},x,y}(\sigma)$ and $\mathcal{Z}(\sigma)$. We are going to do this in the following two sections. At this stage it is not hard to see how to estimate $\log \mathcal{L}$ with the help of \mathcal{L}'/\mathcal{L} . Under GRH we can write

$$\log \mathcal{L}(s) = \log \mathcal{L}\left(\frac{3}{2} + \mathrm{i}t\right) - \int_{\sigma}^{\frac{3}{2}} \frac{\mathcal{L}'}{\mathcal{L}} \left(\sigma' + \mathrm{i}t\right) \mathrm{d}\sigma'.$$

Because then

$$\int_{\sigma}^{\frac{3}{2}} \left| \frac{\mathcal{L}'}{\mathcal{L}} \left(\sigma' + \mathrm{i}t \right) \right| \mathrm{d}\sigma' \leq \sum_{n \leq xy} \frac{|\Lambda_{\mathcal{L},x,y}(n)|}{n^{\sigma} \log n} - \sum_{n \leq x} \frac{|\Lambda_{\mathcal{L}}(n)|}{n^{\frac{3}{2}} \log n} + \int_{\sigma}^{\frac{3}{2}} \left(\mathcal{Z} + \mathcal{R}_1 + \mathcal{R}_2 \right) \left(\sigma' \right) \mathrm{d}\sigma'$$

by Lemma 1, we obtain

$$\left|\log \mathcal{L}(s)\right| \le \widehat{S}_{\mathcal{L},x,y}(\sigma) + \mathcal{E}_x + \int_{\sigma}^{\frac{3}{2}} \left(\mathcal{Z} + \mathcal{R}_1 + \mathcal{R}_2\right)(\sigma') \,\mathrm{d}\sigma',\tag{40}$$

where

$$\widehat{S}_{\mathcal{L},x,y}(\sigma) := \sum_{n \le x} \frac{|\Lambda_{\mathcal{L}}(n)|}{n^{\sigma} \log n} + \frac{1}{\log y} \sum_{x < n \le xy} \frac{|\Lambda_{\mathcal{L}}(n)| \log \frac{xy}{n}}{n^{\sigma} \log n}, \quad \mathcal{E}_x := \sum_{n > x} \frac{|\Lambda_{\mathcal{L}}(n)|}{n^{\frac{3}{2}} \log n}.$$
(41)

We are using (40) for the estimation of $\log \mathcal{L}$. Because our treatment of $\widehat{S}_{\mathcal{L},x,y}(\sigma)$ and \mathcal{E}_x is similar to that of $S_{\mathcal{L},x,y}(\sigma)$, we will do this in the next section.

4. VARIOUS SUMS OVER PRIME NUMBERS

In this section we obtain several upper bounds for $S_{\mathcal{L},x,y}(\sigma)$, $\widehat{S}_{\mathcal{L},x,y}(\sigma)$ and \mathcal{E}_x coming from the Selberg moment formula, see (35) and (41) for definitions, according to whether $\mathcal{L} \in \mathcal{S}$ or $\mathcal{L} \in \mathcal{SP}$. Throughout this section the truth of RH is assumed.

First we estimate $S_{\mathcal{L},x,y}(\sigma)$ when $\mathcal{L} \in \mathcal{S}$.

Lemma 2. Assume the notation and conditions from Theorem 1. Then

$$S_{\mathcal{L},x,y}(\sigma) \le \left(A\left(\mathcal{C}(\varepsilon), \alpha, \sigma - \varepsilon, \sigma\right) + \frac{\mathcal{C}(\varepsilon)2^{1+\varepsilon}\log 2}{2^{1+\varepsilon} - 2^{\sigma}} + o(1)\right) \left(\log \tau\right)^{2(1-\sigma+\varepsilon)},$$

where $A(a, \alpha, u, \sigma)$ is defined by (6).

Proof. Observe that when x and y are from (34), they are not less than 2 for sufficiently large τ and $xy = \log^2 \tau$. It is clear that

$$S_{\mathcal{L},x,y}(\sigma) \leq \sum_{p \leq x} \frac{|\Lambda_{\mathcal{L}}(p)|}{p^{\sigma}} + \frac{1}{\log y} \sum_{x$$

Denote by S_1, \ldots, S_4 the terms on the right-hand side of the latter expression, written in the same order. It is not hard to estimate the last two terms since obviously

$$S_3 \ll \sum_{p \le \log \tau} \frac{\log p}{p^{2(\sigma-\varepsilon)}} \ll 1$$

and

$$S_4 \leq \sum_{k=3}^{\lfloor \frac{2\log\log\tau}{\log 2} \rfloor} \frac{\mathcal{C}(\varepsilon)2^k \log 2}{2^{k(\sigma-\varepsilon)}} + \sum_{3 \leq p \leq 7} \sum_{k=3}^{\lfloor \frac{2\log\log\tau}{\log p} \rfloor} \frac{\mathcal{C}(\varepsilon)2^k \log p}{p^{k(\sigma-\varepsilon)}} + \sum_{p \geq 11} \frac{80\mathcal{C}(\varepsilon)\log p}{p^{3(1/2-\varepsilon)}} \\ \leq \left(\frac{\mathcal{C}(\varepsilon)2^{1+\varepsilon}\log 2}{2^{1+\varepsilon} - 2^{\sigma}} + o(1)\right) (\log\tau)^{2(1-\sigma+\varepsilon)} .$$

Partial summation [RS62, Equation 4.14] and the RH estimate $\vartheta(v) - v < \frac{1}{8\pi}\sqrt{v}\log^2 v$ for $v \ge 2$ where $\vartheta(v) = \sum_{p \le v} \log p$, see [Sch76, Equation 6.5], assures that we have

$$\sum_{p \le X} \frac{\log p}{p^u} \le \frac{X^{1-u} - u2^{1-u}}{1-u} + \frac{\vartheta(X) - X}{X^u} + O\left(\log^2 X\right)$$

and

$$\sum_{X$$

for $1/2 + \delta - \varepsilon \le u \le 1 - \delta - \varepsilon$ and $2 \le X < Y$. Therefore, taking X = x, Y = xy and $u = \sigma - \varepsilon$ in the latter inequalities, using (34) and [Sch76, Equation 6.3], we have

$$\begin{split} S_{\mathcal{L},x,y}(\sigma) &\leq S_1 + S_2 + S_3 + S_4 \\ &\leq \mathcal{C}(\varepsilon) \left(\sum_{p \leq x} \frac{\log p}{p^{\sigma-\varepsilon}} + \frac{1}{\log y} \sum_{x$$

for $1/2 + \delta \leq \sigma \leq 1 - \delta$ with $\tau \to \infty$, where $A(a, \alpha, u, \sigma)$ is defined by (6), and $\alpha \geq \frac{1}{2}\log 2$, $\delta \in (0, 1/4)$ and $\varepsilon \in (0, \min\{\delta, 1/6\})$ are fixed.

Next we estimate $\widehat{S}_{\mathcal{L},x,y}(\sigma)$ when $\mathcal{L} \in \mathcal{S}$.

Lemma 3. Assume the notation and conditions from Theorem 1. Then

$$\widehat{S}_{\mathcal{L},x,y}(\sigma) \leq \left(\eta(\alpha,\sigma,\tau)A\left(\mathcal{C}(\varepsilon),\alpha,\sigma-\varepsilon,\sigma\right) + \frac{\mathcal{C}(\varepsilon)2^{1+\varepsilon}\log 2}{2^{1+\varepsilon}-2^{\sigma}} + o(1)\right) \frac{\left(\log \tau\right)^{2(1-\sigma+\varepsilon)}}{\log\log \tau},$$

where $A(a, \alpha, u, \sigma)$ and $\eta(\alpha, \sigma, \tau)$ are defined by (6) and (7), respectively.

Proof. Observe that then x and y from (34) are not less than 2 for sufficiently large τ . It is clear that

$$\widehat{S}_{\mathcal{L},x,y}(\sigma) \le \sum_{p \le x} \frac{|\Lambda_{\mathcal{L}}(p)|}{p^{\sigma} \log p} + \frac{1}{\log y} \sum_{x$$

Denote by $\hat{S}_1, \ldots, \hat{S}_4$ the terms on the right-hand side of the latter expression, written in the same order. Similarly as in the proof of Lemma 2 we have $\hat{S}_3 \ll 1$ and

$$\begin{split} \widehat{S}_4 &\leq \sum_{k=3}^{\lfloor \frac{2\log\log\tau}{\log^2} \rfloor} \frac{\mathcal{C}(\varepsilon)2^k}{k2^{k(\sigma-\varepsilon)}} + \sum_{3 \leq p \leq 7} \sum_{k=3}^{\lfloor \frac{2\log\log\tau}{\log p} \rfloor} \frac{\mathcal{C}(\varepsilon)2^k}{kp^{k(\sigma-\varepsilon)}} + \sum_{p \geq 11} \sum_{k=3}^{\infty} \frac{\mathcal{C}(\varepsilon)2^k}{kp^{(1/2-\varepsilon)k}} \\ &\leq \left(\sum_{k=3}^{\lfloor \frac{\log\log\tau}{\log^2} \rfloor} + \sum_{k=\lfloor \frac{\log\log\tau}{\log^2} \rfloor + 1} \right) \frac{\mathcal{C}(\varepsilon)2^k}{k2^{k(\sigma-\varepsilon)}} + \sum_{3 \leq p \leq 7} \left(\sum_{k=3}^{\lfloor \frac{\log\log\tau}{\log p} \rfloor} + \sum_{k=\lfloor \frac{\log\log\tau}{\log p} \rfloor + 1} \right) \frac{\mathcal{C}(\varepsilon)2^k}{kp^{k(\sigma-\varepsilon)}} + O(1) \\ &\leq \frac{\mathcal{C}(\varepsilon)2^{1+\varepsilon}}{3(2^{1+\varepsilon}-2^{\sigma})} (\log\tau)^{1-\sigma+\varepsilon} + \frac{\mathcal{C}(\varepsilon)2^{1+\varepsilon} (\log 2) (\log\tau)^{2(1-\sigma+\varepsilon)}}{(2^{1+\varepsilon}-2^{\sigma}) \log\log\tau} + o\left(\frac{(\log\tau)^{2(1-\sigma+\varepsilon)}}{\log\log\tau}\right) \\ &= \left(\frac{\mathcal{C}(\varepsilon)2^{1+\varepsilon}\log 2}{2^{1+\varepsilon}-2^{\sigma}} + o(1)\right) \frac{(\log\tau)^{2(1-\sigma+\varepsilon)}}{\log\log\tau}. \end{split}$$

Partial summation also assures that

$$\sum_{p \le X} \frac{1}{p^u} \le \frac{X^{1-u}}{(1-u)\log X} + \frac{\vartheta(X) - X}{X^u \log X} + O\left(\frac{X^{1-u}}{(1-u)^2 \log^2 X}\right)$$

and

$$\sum_{X$$

for $1/2 + \delta - \varepsilon \le u \le 1 - \delta - \varepsilon$ and $2 \le X < Y$. Therefore, taking X = x, Y = xy and $u = \sigma - \varepsilon$ in the latter inequalities, using (34) and [Sch76, Equation 6.3], we have

$$\widehat{S}_{\mathcal{L},x,y}(\sigma) \le \widehat{S}_1 + \widehat{S}_2 + \widehat{S}_3 + \widehat{S}_4$$

$$\leq \mathcal{C}(\varepsilon) \left(\sum_{p \leq x} \frac{1}{p^{\sigma-\varepsilon}} + \frac{1}{\log y} \sum_{x
$$\leq \left(\eta(\alpha, \sigma, \tau) A\left(\mathcal{C}(\varepsilon), \alpha, \sigma - \varepsilon, \sigma\right) + \frac{\mathcal{C}(\varepsilon)2^{1+\varepsilon}\log 2}{2^{1+\varepsilon} - 2^{\sigma}} + o(1) \right) \frac{(\log \tau)^{2(1-\sigma+\varepsilon)}}{\log \log \tau}$$$$

for $1/2 + \delta \le \sigma \le 1 - \delta$ with $\tau \to \infty$, and $\alpha \ge \frac{1}{2} \log 2$, $\delta \in (0, 1/4)$ and $\varepsilon \in (0, \min\{\delta, 1/6\})$ are fixed.

Now we estimate \mathcal{E}_x when $\mathcal{L} \in \mathcal{S}$ or $\mathcal{L} \in \mathcal{SP}$.

Lemma 4. If $\mathcal{L} \in S\mathcal{P}$, then

$$\frac{1}{m}\mathcal{E}_x \le \frac{2\eta\left(\alpha,\sigma,\tau\right)\exp\left(\frac{\alpha}{2\sigma-1}\right)}{\log\tau\log\log\tau} + \frac{5\exp\left(\frac{2\alpha}{2\sigma-1}\right)\left(1 + \frac{\log\log\tau}{\eta(\alpha,\sigma,\tau)}\right)}{16\pi\log^2\tau},\tag{42}$$

where $\eta(\alpha, \sigma, \tau)$ is defined by (7). If $\mathcal{L} \in \mathcal{S}$ and $\varepsilon \in (0, 1/2)$ from axiom (1) is fixed, then $\mathcal{E}_x \ll 1$. *Proof.* If x > 74, then by using [Sch76. Equation 6.2], i.e., $|\psi(u) - u| < \frac{1}{2} \sqrt{u} \log^2 u$ for u > 74, we have

Troof. If
$$x \ge 74$$
, then by using [Sch76, Equation 6.2], i.e., $|\psi(u) - u| \le \frac{1}{8\pi}\sqrt{u\log^2 u}$ for $u \ge 74$, we have
$$\frac{1}{2}\mathcal{E}_x < \sum \frac{\Lambda(n)}{2} = \int_0^\infty \frac{\mathrm{d}u}{2\pi} - \frac{\psi(x) - x}{2} + \frac{1}{2} \int_0^\infty \frac{(2 + 3\log u)(\psi(u) - u)}{(\psi(u) - u)} \mathrm{d}u$$

$$\frac{1}{m}\mathcal{E}_{x} \leq \sum_{n>x} \frac{1}{n^{\frac{3}{2}}\log n} = \int_{x} \frac{1}{u^{\frac{3}{2}}\log u} - \frac{1}{x^{\frac{3}{2}}\log x} + \frac{1}{2}\int_{x} \frac{1}{u^{\frac{5}{2}}\log^{2} u} du$$
$$\leq \frac{2}{\sqrt{x}\log x} + \frac{5(1+\log x)}{16\pi x} = \frac{2\eta(\alpha,\sigma,\tau)\exp\left(\frac{\alpha}{2\sigma-1}\right)}{\log\tau\log\log\tau} + \frac{5\exp\left(\frac{2\alpha}{2\sigma-1}\right)\left(1+\frac{\log\log\tau}{\eta(\alpha,\sigma,\tau)}\right)}{16\pi\log^{2}\tau}.$$

Further, if $2 \le x < 74$, then we can estimate

$$\frac{1}{m}\mathcal{E}_x \le \sum_{n>x} \frac{\Lambda(n)}{n^{\frac{3}{2}}\log n} = \left(\sum_{x74}\right) \frac{\Lambda(n)}{n^{\frac{3}{2}}\log n} < \left(\sum_{x10^5} \frac{1}{n^{\frac{3}{2}}} \le 0.05 + \sum_{x$$

with the last inequality directly verified for $2 \le x \le 74$ by computer.

Next we estimate $S_{\mathcal{L},x,y}(\sigma)$ when $\mathcal{L} \in S\mathcal{P}$.

Lemma 5. Assume $\mathcal{L} \in SP$. Then

$$\frac{1}{m}S_{\mathcal{L},x,y}(\sigma) \leq A(1,\alpha,\sigma,\sigma)\left(\log\tau\right)^{2-2\sigma} - \frac{\sigma2^{1-\sigma}}{1-\sigma} + \frac{\sigma2^{\frac{3}{2}-\sigma}\left(4 + (2+(2\sigma-1)\log 2)^2\right)}{8\pi(2\sigma-1)^3} + \left(\frac{(2(\alpha+1)\sigma-1)(e^{\alpha}-1)}{2\pi\alpha}\right)\frac{(\log\log\tau)^2}{(2\sigma-1)(\log\tau)^{2\sigma-1}} \quad (43)$$

for the variables α , $\sigma \neq 1$ and τ , which satisfy conditions from the cases (1) or (2) of Theorem 4, and $A(a, \alpha, u, \sigma)$ is defined by (6). Further, for σ from the range (24) we have

$$\frac{1}{m}S_{\mathcal{L},x,y}(\sigma) \leq 2\log\log\tau + 1 - \sigma\log2 + 4\theta_1(M)|1 - \sigma|\left(\log\log\tau\right)^2 + \left(\log2\right)^2\theta_1\left(\frac{M\log2}{2\log\log\tau_0}\right)\sigma|1 - \sigma| + \frac{\sigma2^{\frac{3}{2}-\sigma}\left(4 + \left(2 + (2\sigma-1)\log2\right)^2\right)}{8\pi(2\sigma-1)^3} + \left(\frac{\left(2\left(\alpha+1\right)\sigma-1\right)\left(e^{\alpha}-1\right)}{2\pi\alpha}\right)\frac{\left(\log\log\tau\right)^2}{(2\sigma-1)(\log\tau)^{2\sigma-1}}, \quad (44)$$

where M and $\theta_1(u)$ are from (29). Also,

$$\frac{1}{m}S_{\mathcal{L},x,y}(1) \le 2\log\log\tau - \gamma - \alpha + \frac{(e^{\alpha} - 1)(2\alpha + 1)(\log\log\tau)^2}{2\pi\alpha\log\tau} + \frac{0.24e^{\alpha}}{\log\tau}.$$
(45)

Proof. We know that if $\mathcal{L} \in S\mathcal{P}$, then inequality (9) holds and thus by integration by parts

$$\frac{1}{m}S_{\mathcal{L},x,y}(\sigma) \le \sum_{n \le x} \frac{\Lambda(n)}{n^{\sigma}} + \frac{1}{\log y} \sum_{x < n \le xy} \frac{\Lambda(n)\log\frac{xy}{n}}{n^{\sigma}} = \frac{x^{1-\sigma} - \sigma 2^{1-\sigma}}{1-\sigma} - \int_{2}^{x} \left(\psi(u) - u\right) \left(\frac{1}{u^{\sigma}}\right)' \mathrm{d}u$$

CONDITIONAL ESTIMATES FOR L-FUNCTIONS IN THE SELBERG CLASS

$$+ \frac{(xy)^{1-\sigma}}{\log y} \int_{1}^{y} \frac{\log u}{u^{2-\sigma}} du - \frac{1}{\log y} \int_{x}^{xy} (\psi(u) - u) \left(\frac{1}{u^{\sigma}} \log \frac{xy}{u}\right)' du \\ \leq \frac{(xy)^{1-\sigma} - x^{1-\sigma}}{(1-\sigma)^{2} \log y} - \frac{\sigma 2^{1-\sigma}}{1-\sigma} + \frac{\sigma}{8\pi} \int_{2}^{x} \frac{\log^{2} u}{u^{\frac{1}{2}+\sigma}} du + \frac{\left(\sigma + \frac{1}{\log y}\right) \left(x^{\frac{1}{2}-\sigma} - (xy)^{\frac{1}{2}-\sigma}\right) \log^{2}(xy)}{8\pi \left(\sigma - \frac{1}{2}\right)}$$

see especially estimates (3.5) and (3.7), and [CHS23, Footnote 4 on p. 4] for similar ideas. Taking into account also (34), the last inequality assures (43).

Let us now prove (45) since we are going to use this bound to prove (44). Observe that $x = e^{-2\alpha} \log^2 \tau$ since $\sigma = 1$. Therefore, if $\tau \ge \exp\left(e^{\alpha}\sqrt{60}\right)$, then $x \ge 60$ and thus (45) holds by the first line in Equation (3.6) from [CHS23].

Let us now prove (44). Firstly, assume that $\sigma \neq 1$. By the first paragraph of the proof we know that (43) holds for σ from the range (24) while $\sigma \neq 1$. Since $e^u - 1 - u \ge 0$ for all real numbers u, we have

$$A(1,\alpha,\sigma,\sigma)(\log \tau)^{2-2\sigma} - \frac{\sigma 2^{1-\sigma}}{1-\sigma} = -(\log \tau)^{2-2\sigma} \left(\frac{2\alpha}{2\sigma-1}\right) \left(\frac{\exp\left(-\frac{2\alpha(1-\sigma)}{2\sigma-1}\right) - 1 - \left(-\frac{2\alpha(1-\sigma)}{2\sigma-1}\right)}{\left(-\frac{2\alpha(1-\sigma)}{2\sigma-1}\right)^2}\right) + \frac{(\log \tau)^{2-2\sigma}}{1-\sigma} - \frac{\sigma 2^{1-\sigma}}{1-\sigma} \le \frac{(\log \tau)^{2-2\sigma}}{1-\sigma} - \frac{\sigma 2^{1-\sigma}}{1-\sigma}.$$
(46)

Even more, note that $|e^u - 1 - u| \le \theta_1(u_0) u^2$ for $|u| \le u_0$, where $\theta_1(u_0)$ is from (29). This implies

$$\left| (\log \tau)^{2-2\sigma} - (1+2(1-\sigma)\log\log\tau) \right| \le 4\theta_1(M)(1-\sigma)^2 (\log\log\tau)^2 \\ \left| \frac{\sigma 2^{1-\sigma}}{1-\sigma} - \left(\frac{\sigma}{1-\sigma} + \sigma\log 2 \right) \right| \le (\log 2)^2 \theta_1 \left(\frac{M\log 2}{2\log\log\tau_0} \right) \sigma |1-\sigma|,$$

since $|1 - \sigma| \le M/(2 \log \log \tau)$ with M from (29). Therefore, by the previous two inequalities and (46) we have

$$A(1,\alpha,\sigma,\sigma)\left(\log\tau\right)^{2-2\sigma} - \frac{\sigma^{2^{1-\sigma}}}{1-\sigma} \le 2\log\log\tau + 1 - \sigma\log2 + 4\theta_1(M)|1-\sigma|\left(\log\log\tau\right)^2 + \left(\log2\right)^2\theta_1\left(\frac{M\log2}{2\log\log\tau_0}\right)\sigma|1-\sigma|,$$

which gives (44) for $\sigma \neq 1$ from the range (24). However, estimate (44) holds also for $\sigma = 1$ by (45). The proof of Lemma 5 is thus complete.

Finally, we estimate $\widehat{S}_{\mathcal{L},x,y}(\sigma)$ when $\mathcal{L} \in S\mathcal{P}$. The method we are using here is very similar to that from the proof of Lemma 5.

Lemma 6. Assume $\mathcal{L} \in S\mathcal{P}$, $\nu_1 > 0$ and $\nu_2 > 1$, and let $\theta_1(u)$ be defined as in (29). Then, under the assumptions from the case (1) of Theorem 4, we have

$$\frac{1}{m}\widehat{S}_{\mathcal{L},x,y}(\sigma) \leq \eta(\alpha,\sigma,\tau) \left(A(1,\alpha,\sigma,\sigma) \frac{(\log \tau)^{2-2\sigma}}{\log\log \tau} - \frac{1}{(1-\sigma)\log\log \tau} \right) \\
+ \left((\nu_2\eta(\alpha,\sigma,\tau))^2 \exp\left(-\frac{2\alpha(1-\sigma)}{2\sigma-1}\right) \right) \frac{(\log \tau)^{2-2\sigma}}{(1-\sigma)^2 (\log\log \tau)^2} \\
+ \frac{1}{\nu_1^2} \exp\left(-\frac{2\alpha(1-\sigma)}{(2\sigma-1)\nu_2}\right) (\log \tau)^{\frac{2}{\nu_2}(1-\sigma)} + \log\left(2\log\log \tau\right) + \frac{1-\sigma 2^{1-\sigma}}{(1-\sigma)\log 2} - \log\log 2 \\
+ \int_0^{\nu_1} \theta_1(u) du + \frac{2^{\frac{3}{2}-\sigma} \left(-1+\sigma\left(4-\log 2\right)+\sigma^2\log 4\right)}{8\pi \left(2\sigma-1\right)^2} + \frac{(e^\alpha-1)\log\log \tau}{4\pi\alpha \left(\log \tau\right)^{2\sigma-1}}, \quad (47)$$

where $A(a, \alpha, u, \sigma)$ and $\eta(\alpha, \sigma, \tau)$ are defined by (6) and (7), respectively.

Further, under the assumptions from the case (2) of Theorem 4 and assuming that $\sigma \neq 1$, we have

$$\frac{1}{m}\widehat{S}_{\mathcal{L},x,y}(\sigma) \le \log\left(2\log\log\tau\right) + 2\left(2\eta(\alpha,\sigma,\tau)\theta_1(M) + \theta_1(2\alpha_2)\right)\left|1 - \sigma\right|\log\log\tau$$

N. PALOJÄRVI AND A. SIMONIČ

$$+\frac{1}{\log 2} - \log \log 2 - \sigma + (\log 2) \theta_1 \left(\frac{M \log 2}{2 \log \log \tau_0}\right) \sigma |1 - \sigma| + 2\eta(\alpha, \sigma, \tau) \\ + \frac{2^{\frac{3}{2} - \sigma} \left(-1 + \sigma \left(4 - \log 2\right) + \sigma^2 \log 4\right)}{8\pi \left(2\sigma - 1\right)^2} + \frac{(e^{\alpha} - 1) \log \log \tau}{4\pi \alpha \left(\log \tau\right)^{2\sigma - 1}}, \tag{48}$$

where $M = 2 \max \{\alpha_2, \alpha_3\}.$

Finally, if $\sigma = 1$, then

$$\frac{1}{m}\widehat{S}_{\mathcal{L},x,y}(1) \leq \log\left(2\log\log\tau\right) + \gamma + \left(1 - \frac{\log\log\tau}{\alpha}\right)\log\left(1 - \frac{\alpha}{\log\log\tau}\right) - 1 \\
+ \frac{\left(1 + e^{\alpha}\left(4\alpha - 1\right)\right)\log\log\tau + e^{\alpha}\left(7\alpha - 4\left(\alpha^{2} + 1\right)\right) + 4}{4\pi\alpha\log\tau}.$$
(49)

Proof. Let $\sigma \in [0, \infty) \setminus \{1\}$, $x \ge 2$ and $y \ge 2$. By partial summation and footnote 4 p. 4 in [CHS23] we have

$$\sum_{n \le x} \frac{\Lambda(n)}{n^{\sigma} \log n} \le \int_{2}^{x} \frac{\mathrm{d}u}{u^{\sigma} \log u} + \frac{\psi(x) - x}{x^{\sigma} \log x} + \frac{2^{1-\sigma}}{\log 2} + \frac{1}{8\pi} \int_{2}^{x} \frac{1 + \sigma \log u}{u^{\frac{1}{2} + \sigma}} \mathrm{d}u \tag{50}$$

and

$$\frac{1}{\log y} \sum_{x < n \le xy} \frac{\Lambda(n) \log \frac{xy}{n}}{n^{\sigma} \log n} \le \frac{(xy)^{1-\sigma}}{\log y} \int_{1}^{y} \frac{\log u \, \mathrm{d}u}{u^{2-\sigma} \log \frac{xy}{u}} - \frac{\psi(x) - x}{x^{\sigma} \log x} + \frac{1}{8\pi \log y} \int_{x}^{xy} \frac{(1+\sigma \log u) \log \frac{xy}{u} + \log u}{u^{\frac{1}{2}+\sigma}} \mathrm{d}u \qquad (51)$$

$$\le \frac{(xy)^{1-\sigma}}{\log u \log x} \int_{x}^{y} \frac{\log u}{u^{2-\sigma}} \mathrm{d}u - \frac{\psi(x) - x}{u^{\sigma} \log x} + \frac{1}{8\pi} \int_{x}^{xy} \frac{1+\sigma \log u}{u^{\frac{1}{2}+\sigma}} \mathrm{d}u + \frac{1}{8\pi \log x} \int_{x}^{xy} \frac{\log u}{\frac{1+\sigma}{2}} \mathrm{d}u. \qquad (52)$$

$$= \log x \log y \int_1^{-} u^{2-\sigma} x^{\sigma} \log x + 8\pi \int_x^{-} u^{\frac{1}{2}+\sigma} x^{\sigma} \log y \int_x^{-} u^{\frac{1}{2}+$$

We know that if $\mathcal{L} \in S\mathcal{P}$, then inequality (9) holds. Therefore, combining (50) and (52) gives

$$\frac{1}{m}\widehat{S}_{\mathcal{L},x,y}(\sigma) \leq \frac{(xy)^{1-\sigma} - x^{1-\sigma}}{(1-\sigma)^2 \log x \log y} - \frac{x^{1-\sigma}}{(1-\sigma)\log x} + \int_2^x \frac{\mathrm{d}u}{u^{\sigma}\log u} + \frac{2^{1-\sigma}}{\log 2} + \frac{1}{8\pi}\int_2^{xy} \frac{1+\sigma\log u}{u^{\frac{1}{2}+\sigma}} \mathrm{d}u + \frac{1}{8\pi\log y}\int_x^{xy} \frac{\log u}{u^{\frac{1}{2}+\sigma}} \mathrm{d}u. \quad (53)$$

The strategy to bound (53) is now very similar to that in the proof of Lemma 5, but with an exemption of the first integral in (53). Following [CC11, Appendix A], we have

$$\int_{2}^{x} \frac{\mathrm{d}u}{u^{\sigma} \log u} = \log \log x - \log \log 2 + \int_{(1-\sigma)\log 2}^{(1-\sigma)\log x} \theta_{2}(u) \mathrm{d}u$$
$$= \frac{x^{1-\sigma} - 1}{(1-\sigma)\log x} + \log \log x - \frac{2^{1-\sigma} - 1}{(1-\sigma)\log 2} - \log \log 2 + \int_{(1-\sigma)\log 2}^{(1-\sigma)\log x} \theta_{1}(u) \mathrm{d}u, \qquad (54)$$

where $\theta_2(u)$ is defined in (29). We need to estimate the integral on the right-hand side of (54) for each of the cases (23) and (24). Next we consider different cases depending on the size of σ .

Let $\sigma \in [0, 1)$. Then

$$\int_{(1-\sigma)\log 2}^{(1-\sigma)\log x} \theta_1(u) \mathrm{d}u \le \int_0^{(1-\sigma)\log x} \theta_1(u) \mathrm{d}u = \left(\int_0^{\nu_1} + \int_{\nu_1}^{\frac{(1-\sigma)\log x}{\nu_2}} + \int_{\frac{(1-\sigma)\log x}{\nu_2}}^{(1-\sigma)\log x}\right) \theta_1(u) \mathrm{d}u$$
$$\le \frac{\nu_2^2 x^{1-\sigma}}{(1-\sigma)^2 \log^2 x} + \frac{1}{\nu_1^2} x^{\frac{1-\sigma}{\nu_2}} + \int_0^{\nu_1} \theta_1(u) \mathrm{d}u. \tag{55}$$

Consider the assumptions from the case (1) of Theorem 4. Then (53), (54) and (55) imply inequality (47).

For $\sigma \geq 1 - \alpha_2 / \log \log \tau$, we also have

$$\int_{(1-\sigma)\log 2}^{(1-\sigma)\log x} \theta_1(u) \mathrm{d}u \le \theta_1(2\alpha_2) |1-\sigma|\log x \tag{56}$$

since the integral in (56) is negative for $\sigma > 1$ and $(1 - \sigma) \log x \le 2\alpha_2$. Consider the assumptions from the case (2) of Theorem 4 and assume $\sigma \ne 1$. Then similarly as in the proof of Lemma 5, (53), (54) and (56) imply (48).

As in the proof of Lemma 5, we can estimate $\widehat{S}_{\mathcal{L},x,y}(1)$ even better. We have

$$\sum_{n \le x} \frac{\Lambda(n)}{n \log n} \le \sum_{p \le x} \frac{1}{p} + \sum_{p} \sum_{k=2}^{\infty} \frac{1}{kp^k} = \sum_{p \le x} \frac{1}{p} - \sum_{p} \left(\frac{1}{p} + \log\left(1 - \frac{1}{p}\right) \right)$$
$$\le \log \log x + \gamma + \frac{\psi(x) - x}{x \log x} + \frac{3 + \log x}{4\pi\sqrt{x}}$$
(57)

for $x \ge 60$. The last inequality follows for $x \ge 599$ by [RS62, Equations 2.7 and 4.15], [Sch76, Equation 6.3] and $\vartheta(x) - x \le \psi(x) - x$, see also the proof of [Sch76, Corollary 2]. We can verify by computer that it holds also for $60 \le x \le 599$. Next, note that (51) holds also for $\sigma = 1$. Therefore,

$$\frac{1}{\log y} \sum_{x < n \le xy} \frac{\Lambda(n) \log \frac{xy}{n}}{n \log n} \le -\frac{\log \left(xy\right)}{\log y} \log \left(1 - \frac{\log y}{\log \left(xy\right)}\right) - 1 - \frac{\psi(x) - x}{x \log x} + \frac{8 + \log \left(xy\right)}{4\pi\sqrt{xy} \log y} + \frac{3 + \log x}{4\pi\sqrt{x}} - \frac{8 + \log x}{4\pi\sqrt{x} \log y}.$$
(58)

Then (34), (57) and (58) imply (49).

5. The sum over the non-trivial zeros

In this section we obtain an upper estimate for the sum from (36), which we obtained from the Selberg moment formula, over the non-trivial zeros of $\mathcal{L}(s)$. We will do this with the help of the Guinand–Weil exact formula for functions in the Selberg class, see the next lemma.

Lemma 7. Assume that $\mathcal{L} \in \mathcal{S}$ and $\mathcal{L}(1) \neq 0$. Let h(z) be a holomorphic function on a strip

$$\left\{z\in\mathbb{C}\colon -\frac{1}{2}-\varepsilon<\Im\{z\}<\frac{1}{2}+\varepsilon\right\}$$

for some $\varepsilon > 0$, such that $h(z) (1 + |z|)^{1+\delta}$ is bounded for some $\delta > 0$. Then

$$\sum_{\rho} h\left(\frac{\rho - \frac{1}{2}}{i}\right) = m_{\mathcal{L}}\left(h\left(\frac{1}{2i}\right) + h\left(-\frac{1}{2i}\right)\right) + \frac{\log Q}{\pi}\hat{h}(0) + \frac{1}{\pi}\sum_{j=1}^{f}\int_{-\infty}^{\infty}h(u)\lambda_{j}\Re\left\{\frac{\Gamma'}{\Gamma}\left(\frac{\lambda_{j}}{2} + \mu_{j} + i\lambda_{j}u\right)\right\}du - \frac{1}{2\pi}\sum_{n=2}^{\infty}\frac{1}{\sqrt{n}}\left(\Lambda_{\mathcal{L}}(n)\hat{h}\left(\frac{\log n}{2\pi}\right) + \overline{\Lambda_{\mathcal{L}}(n)}\hat{h}\left(-\frac{\log n}{2\pi}\right)\right),$$
(59)

where $\hat{h}(\xi) = \int_{-\infty}^{\infty} h(u) e^{-2\pi i u \xi} du$ denotes the Fourier transform of h(z) and the sum runs over all non-trivial zeros ρ of $\mathcal{L}(s)$.

Proof. The proof is almost identical to the proof of [Fin16, Lemma 5], so we will highlight only the main differences. Write $\mathfrak{L}(s) = \mathcal{L}(s)\mathfrak{L}_1(s)$, where $\mathfrak{L}(s)$ is from the functional equation. Then the only zeros ρ of $\mathfrak{L}(s)$ are the non-trivial zeros ρ of $\mathcal{L}(s)$, where $\rho \notin \{0,1\}$ by the assumption $\mathcal{L}(1) \neq 0$. Let $\varepsilon > 0$. By the residue theorem

$$\sum_{\rho} h\left(\frac{\rho - \frac{1}{2}}{\mathrm{i}}\right) - m_{\mathcal{L}}\left(h\left(\frac{1}{2\mathrm{i}}\right) + h\left(-\frac{1}{2\mathrm{i}}\right)\right) = \frac{1}{2\pi\mathrm{i}}\left(\int_{\gamma_1} + \int_{\gamma_2}\right) h\left(\frac{z - \frac{1}{2}}{\mathrm{i}}\right)\frac{\mathfrak{L}'}{\mathfrak{L}}(z)\mathrm{d}z,$$

where $\gamma_1 = \{z \in \mathbb{C} : \Re\{z\} = -\varepsilon/2\}$ and $\gamma_2 = \{z \in \mathbb{C} : \Re\{z\} = 1 + \varepsilon/2\}$ with the appropriate orientation, see [Fin16, p. 15]. Hence, we have obtained the first term in the right-hand side of (59). Furthermore,

$$\sum_{\rho} h\left(\frac{\rho - \frac{1}{2}}{i}\right) - m_{\mathcal{L}}\left(h\left(\frac{1}{2i}\right) + h\left(-\frac{1}{2i}\right)\right) = \frac{1}{2\pi i} \int_{\gamma_2} \left(h\left(\frac{z - \frac{1}{2}}{i}\right) \frac{\mathcal{L}'}{\mathcal{L}}(z) + h\left(\frac{\frac{1}{2} - z}{i}\right) \frac{\overline{\mathcal{L}'}(\bar{z})}{\mathcal{L}}\right) dz$$

$$+\frac{1}{2\pi \mathrm{i}}\int_{\gamma_2} \left(h\left(\frac{z-\frac{1}{2}}{\mathrm{i}}\right)\frac{\mathfrak{L}'_1}{\mathfrak{L}_1}(z) + h\left(\frac{\frac{1}{2}-z}{\mathrm{i}}\right)\frac{\overline{\mathfrak{L}'_1}}{\mathfrak{L}_1}(\bar{z})\right)\mathrm{d}z$$

by employing the functional equation, see [Fin16, p. 16]. Note that

$$\frac{\mathfrak{L}'_1}{\mathfrak{L}_1}(z) = \log Q + \sum_{j=1}^J \lambda_j \frac{\Gamma'}{\Gamma} \left(\lambda_j z + \mu_j\right).$$

The last sum in (59) comes from the first two integrals in the latter equation, see Step 3 in [Fin16, p. 17]. After moving the line of integration in the last two integrals to $\Re\{z\} = 1/2$ and noticing that $(\mathfrak{L}'_1/\mathfrak{L}_1)(z)$ is holomorphic on domain $\{z \in \mathbb{C} : \Re\{z\} > 0\}$, see Step 4 in [Fin16, p. 18], we obtain the last two remaining terms from (59).

Remark 1. Beside the conditions from Lemma 7 assume also GRH and that h(z) is real for $z \in \mathbb{R}$. Note that the last condition assures that $\overline{h(z)} = h(\overline{z})$ and also that $\overline{\hat{h}(\xi)} = \hat{h}(-\xi)$ for $\xi \in \mathbb{R}$. We can see that Lemma 7 then implies

$$\begin{split} \sum_{\gamma} h(\gamma) &= 2m_{\mathcal{L}} \Re\left\{h\left(\frac{1}{2\mathrm{i}}\right)\right\} + \frac{\log Q}{\pi} \widehat{h}(0) - \frac{1}{\pi} \sum_{n=2}^{\infty} \frac{1}{\sqrt{n}} \Re\left\{\Lambda_{\mathcal{L}}(n) \widehat{h}\left(\frac{\log n}{2\pi}\right)\right\} \\ &+ \frac{1}{\pi} \sum_{j=1}^{f} \int_{-\infty}^{\infty} h(u) \lambda_{j} \Re\left\{\frac{\Gamma'}{\Gamma}\left(\frac{\lambda_{j}}{2} + \mu_{j} + \mathrm{i}\lambda_{j}u\right)\right\} \mathrm{d}u, \end{split}$$

where the sum runs over the imaginary parts of the non-trivial zeros of $\mathcal{L}(s)$.

Lemma 8. Let $\mathcal{L} \in S\mathcal{P}$. Assume the Generalized Riemann Hypothesis and the strong λ -conjecture. Assume also that

$$\frac{1}{2} + \frac{\alpha_1}{\log\log\tau} \le \sigma \le \frac{3}{2} \tag{60}$$

for $\alpha_1 > 0$, and

$$au \ge au_0 \ge \max\left\{e^{\sqrt{60}}, \exp\left(e^{\alpha_1}\right)\right\}, \quad |t| \ge t_0 \ge \max\left\{2\max_{1\le j\le f}\left\{|\mu_j|\right\}, 1\right\}.$$

Then we have

$$\sum_{\gamma} \frac{\sigma - \frac{1}{2}}{\left(\sigma - \frac{1}{2}\right)^2 + (t - \gamma)^2} \le \frac{\log \tau}{2} + \mathfrak{a} \left(m_{\mathcal{L}}, \alpha_1, t_0\right) \left(\log \tau\right)^{2 - 2\sigma} + 2m \log \log \tau + \mathfrak{b} \left(\mathrm{d}_{\mathcal{L}}, m, m_{\mathcal{L}}, \alpha_1, t_0, \tau_0\right), \quad (61)$$

where

$$\mathfrak{a}(m_{\mathcal{L}},\alpha_1,t_0) := \frac{1}{1 - e^{-2\alpha_1}} \left(1 + \frac{4m_{\mathcal{L}}}{\left(t_0^2 - \frac{3}{4}\right)\left(1 - e^{-2\alpha_1}\right)} \right)$$
(62)

and

$$\mathfrak{b}\left(\mathrm{d}_{\mathcal{L}}, m, m_{\mathcal{L}}, \alpha_{1}, t_{0}, \tau_{0}\right) \coloneqq \frac{\mathrm{d}_{\mathcal{L}}}{\pi} \operatorname{coth}^{2}\left(\alpha_{1}\right) \int_{0}^{\infty} \frac{\log\left(1 + \frac{y}{2\pi t_{0}}\right)}{1 + y^{2}} \mathrm{d}y \\ + \frac{2m_{\mathcal{L}}\left(1 + e^{-4\alpha_{1}}\right)}{\left(t_{0}^{2} - \frac{3}{4}\right)\left(1 - e^{-2\alpha_{1}}\right)^{2}} + m\left(-1 - \gamma + \frac{\frac{4\log\log\tau_{0}}{\log\tau_{0}} + \frac{2.24}{\log\tau_{0} - 1}}{1 - (\log\tau_{0})^{-1}}\right). \quad (63)$$

Proof. The proof is similar to the one in [CHS23, Sections 4 and 7]. Indeed, we are using Lemma 7 in order to estimate the sum over zeros.

The left-hand side of (61) can be written as $\sum_{\gamma} f_a(t-\gamma)$, where $a = \sigma - 1/2$ and $f_a \colon \mathbb{R} \to \mathbb{R}$ is defined by

$$f_a(x) = \frac{a}{a^2 + x^2}$$

However, the function $f_a(x)$ does not satisfy the conditions from Lemma 7. So, as in [CHS23], let us set

$$h(s) = h_{a,\Delta}(s) = \left(\frac{a}{a^2 + s^2}\right) \left(\frac{e^{2\pi a\Delta} + e^{-2\pi a\Delta} - 2\cos\left(2\pi\Delta s\right)}{\left(e^{\pi a\Delta} - e^{-\pi a\Delta}\right)^2}\right).$$

By [CCM19, Lemma 9], for any $\Delta > 0$ the function h(s) is an entire function of exponential type of $2\pi\Delta$, $f_a(u) \leq h(u)$ for all $u \in \mathbb{R}$ and the Fourier transform $\hat{h}(\xi)$ satisfies $\hat{h}(\xi) \geq 0$ if $|\xi| \leq \Delta$, $\hat{h}(\xi) = 0$ if $|\xi| > \Delta$ and $\hat{h}(0) = \pi \coth(\pi a \Delta)$. Therefore, the function h(s) satisfies the assumptions given in Lemma 7. Employing Remark 1 for the function $z \mapsto h(t-z)$, we obtain

$$\sum_{\gamma} f_a(t-\gamma) \leq \sum_{\gamma} h(t-\gamma) \leq 2m_{\mathcal{L}} \left| h\left(t + \frac{1}{2\mathrm{i}}\right) \right| + \frac{\log Q}{\pi} \widehat{h}(0) + \frac{1}{\pi} \sum_{j=1}^f \int_{-\infty}^\infty h(u) \lambda_j \Re \left\{ \frac{\Gamma'}{\Gamma} \left(\frac{\lambda_j}{2} + \mu_j + \lambda_j(t-u)\mathrm{i} \right) \right\} \mathrm{d}u + \frac{1}{\pi} \sum_{n=2}^\infty \frac{1}{\sqrt{n}} \left| \Lambda_{\mathcal{L}}(n) \widehat{h}\left(\frac{\log n}{2\pi} \right) \right|.$$
(64)

In what follows we are taking $\Delta = \frac{1}{\pi} \log \log \tau$. Let us consider each term on the right-hand side of (64) separately.

Using similar idea as at the end of Section 4 in [CHS23], we have that the value of the first term on the right-hand side of (64) does not exceed

$$\frac{2m_{\mathcal{L}}a\left|e^{2\pi a\Delta} + e^{-2\pi a\Delta} - \cos(2t\pi\Delta)\left(e^{\pi\Delta} + e^{-\pi\Delta}\right) - i\sin(2t\pi\Delta)\left(e^{\pi\Delta} - e^{-\pi\Delta}\right)\right|}{\left|a^{2} + (t - i/2)^{2}\right|\left(e^{\pi a\Delta} - e^{-\pi a\Delta}\right)^{2}} \leq \frac{2m_{\mathcal{L}}a}{\left|a^{2} - 1/4 + t^{2}\right|} \left(\frac{2e^{(1-2a)\pi\Delta} + 1 + e^{-4\pi a\Delta}}{\left(1 - e^{-2\pi a\Delta}\right)^{2}}\right) \leq \frac{m_{\mathcal{L}}(2\sigma - 1)}{t^{2} - 3/4} \left(\frac{2(\log \tau)^{2-2\sigma} + 1 + (\log \tau)^{-2(2\sigma - 1)}}{\left(1 - (\log \tau)^{-(2\sigma - 1)}\right)^{2}}\right).$$
(65)

Let us now consider the second and third terms. Using $\Re\{(\Gamma'/\Gamma)(z)\} \leq \log |z|$ for $\Re\{z\} \geq 1/4$, see [Cha09, Lemma 2.3], we obtain

$$\Re\left\{\frac{\Gamma'}{\Gamma}\left(\frac{\lambda_j}{2} + \mu_j + \lambda_j(t-u)\mathbf{i}\right)\right\} \le \log\left|\frac{\lambda_j}{2} + \mu_j + \lambda_j(t-u)\mathbf{i}\right|$$
$$\le \log\left(2\lambda_j\pi|t|\right) + \log\left(\frac{1}{4\pi|t|} + \frac{\max_{1\le j\le f}\left\{|\mu_j|\right\}}{2\min_{1\le j\le f}\left\{\lambda_j\right\}\pi|t|} + \frac{1}{2\pi}\left|1 - \frac{u}{t}\right|\right)$$
$$\le \log\left(2\lambda_j\pi|t|\right) + \log\left(1 + \frac{|u|}{2\pi t_0}\right) \tag{66}$$

since we are assuming the strong λ -conjecture. Therefore,

$$\sum_{j=1}^{J} \lambda_j \Re\left\{\frac{\Gamma'}{\Gamma}\left(\frac{\lambda_j}{2} + \mu_j + \lambda_j(t-u)\mathbf{i}\right)\right\} \le \frac{1}{2}\log\tau - \log Q + \frac{\mathrm{d}_{\mathcal{L}}}{2}\log\left(1 + \frac{|u|}{2\pi t_0}\right).$$

Furthermore, because $\pi a \Delta \geq \alpha_1$ and $\hat{h}(0) = \pi \coth(\pi a \Delta)$, we have

$$\frac{\log Q}{\pi}\widehat{h}(0) + \frac{1}{\pi}\sum_{j=1}^{f}\int_{-\infty}^{\infty}h(u)\lambda_{j}\Re\left\{\frac{\Gamma'}{\Gamma}\left(\frac{\lambda_{j}}{2} + \mu_{j} + \lambda_{j}(t-u)\mathbf{i}\right)\right\}du$$
$$\leq \frac{\log\tau}{2} + \frac{(\log\tau)^{2-2\sigma}}{1-e^{-2\alpha_{1}}} + \frac{d_{\mathcal{L}}}{\pi}\coth^{2}\left(\alpha_{1}\right)\int_{0}^{\infty}\frac{\log\left(1+\frac{y}{2\pi t_{0}}\right)}{1+y^{2}}dy, \quad (67)$$

where we also used inequality (4.6) from [CHS23] and the fact that $0 < a \leq 1$.

We are left with the last term. Since the function $\mathcal{L}(s)$ has a polynomial Euler product of order m, we know that (9) holds. Thus, by the proof of [CHS23, Theorem 5], the fourth term does not exceed

$$m\left(\frac{2\log\log\tau - \gamma - 1 + 0.24(\log\tau)^{-1}}{(1 - (\log\tau)^{-1})^2}\right) \le 2m\log\log\tau + m\left(-1 - \gamma + \frac{1}{1 - (\log\tau_0)^{-1}}\left(\frac{4\log\log\tau_0}{\log\tau_0} + \frac{2.24}{\log\tau_0 - 1}\right)\right).$$
 (68)

Simplifying (65) by using (60), and then adding (67) and (68), we finally arrive at (61). The proof of Lemma 8 is thus complete.

Remark 2. The only instance where we used the strong λ -conjecture in the proof of Lemma 8 was in (66). Without this assumption we know that

$$\lambda_j \Re \left\{ \frac{\Gamma'}{\Gamma} \left(\frac{\lambda_j}{2} + \mu_j + \lambda_j (t - u) \mathbf{i} \right) \right\} \le \lambda_j \log \left(2\lambda_j \pi |t| \right) + \lambda_j \log \left(1 + |u| \right) + O_{\lambda^-}(1)$$

by Stirling's formula $(\Gamma'/\Gamma)(z) = \log z + O(1/|z|)$, where $|t| \ge \mu^+/\lambda^-$ is sufficiently large, and λ^- and μ^+ are from (10). Therefore, GRH implies that

$$\sum_{\gamma} \frac{\sigma - \frac{1}{2}}{\left(\sigma - \frac{1}{2}\right)^2 + (t - \gamma)^2} \le \frac{\log \tau}{2} + O\left(m_{\mathcal{L}}(\log \tau)^{2 - 2\sigma} + d_{\mathcal{L}} + m_{\mathcal{L}}\right) + 2m \log \log \tau + O_{\lambda^-}(f)$$

uniformly for $1/2 + 1/\log \log \tau \leq \sigma \leq 3/2$ and sufficiently large τ and $|t| \geq \mu^+/\lambda^-$. Furthermore, if we are considering the full Selberg class S, then additionally to the above the estimate (68) should be replaced by

$$\ll \sum_{n \le (\log \tau)^2} \frac{|\Lambda_{\mathcal{L}}(n)|}{n} \ll \sum_{p \le (\log \tau)^2} \frac{\log p}{p^{1-\varepsilon}} + \sum_{k=2}^{\lfloor \frac{2\log \log \tau}{\log 2} \rfloor} 2^{k\varepsilon} + \sum_{p \ge 3} \sum_{k=2}^{\infty} \frac{2^k \log p}{p^{k(1-\varepsilon)}} \ll (\log \tau)^{2\varepsilon} \,,$$

where $\varepsilon \in (0, 1/6)$ is fixed. Therefore, under this condition GRH implies that

$$\sum_{\gamma} \frac{\sigma - \frac{1}{2}}{\left(\sigma - \frac{1}{2}\right)^2 + (t - \gamma)^2} \le \frac{\log \tau}{2} + O\left(m_{\mathcal{L}}(\log \tau)^{2-2\sigma} + (\log \tau)^{2\varepsilon} + d_{\mathcal{L}} + m_{\mathcal{L}}\right) + O_{\lambda^-}(f)$$

uniformly for $1/2 + 1/\log \log \tau \le \sigma \le 3/2$ and sufficiently large τ and $|t| \ge \mu^+/\lambda^-$.

6. Proof of Theorems 1, 2 and 3

Proof of Theorem 1. Assume the notation and conditions from Theorem 1. Observe that fixed $\alpha \geq \frac{1}{2} \log 2$ guarantees that x and y from (34) are not less than 2 for sufficiently large τ . By (37), (38) and Remark 2 we have

$$\mathcal{Z}(\sigma) \ll \left(\log \tau\right)^{2-2\sigma}, \quad \mathcal{R}_1(\sigma) \ll 1, \quad \mathcal{R}_2(\sigma) \ll 1$$
(69)

for $1/2 + \delta \leq \sigma \leq 3/2$, where the implied constants may depend on \mathcal{L} . Taking (69) and Lemma 2 in (39) furnishes the proof of (4). In order to prove (5), take (69) into (40), together with $\mathcal{E}_x \ll 1$, see Lemma 4, and Lemma 3.

Proof of Theorem 2. Assume the notation and conditions from Theorem 2. Observe that fixed $\alpha \in [0.35, 2)$ guarantees that x and y from (34) are not less than 2 for sufficiently large τ . In what follows we are assuming that τ and $|t| \ge \mu^+ / \lambda^-$ are sufficiently large. We can see from Remark 2 that

$$\mathcal{Z}(\sigma) \leq \frac{e^{\alpha} + 1}{2\alpha} \left(\log \tau\right)^{2-2\sigma} + O\left(m_{\mathcal{L}} \left(\log \tau\right)^{3-4\sigma} + \frac{m \log \log \tau}{\left(\log \tau\right)^{2\sigma-1}} + \frac{\mathrm{d}_{\mathcal{L}} + m_{\mathcal{L}}}{\left(\log \tau\right)^{2\sigma-1}}\right) + O_{\lambda^{-}}\left(\frac{f}{\left(\log \tau\right)^{2\sigma-1}}\right)$$
(70)

uniformly for $1/2 + 1/\log \log \tau \le \sigma \le 3/2$, where $\mathcal{Z}(\sigma)$ is defined by (36). Note that

$$\mathcal{Z}(\sigma) = O\left(1 + \frac{m\log\log\tau}{\log\tau} + \frac{\mathrm{d}_{\mathcal{L}} + m_{\mathcal{L}}}{\log\tau}\right) + O_{\lambda^{-}}\left(\frac{f}{\log\tau}\right)$$

uniformly for $|1 - \sigma| \le 1/\log \log \tau$. We also have

$$\mathcal{R}_1(\sigma) \ll_{\lambda^+} \frac{f}{\left(\log \tau\right)^{(2-\alpha)\sigma}}, \quad \mathcal{R}_2(\sigma) \ll m_{\mathcal{L}} \left(\frac{q_{\mathcal{L}}}{\tau}\right)^{\frac{2}{d_{\mathcal{L}}}} \left(\log \tau\right)^{2-2\sigma}$$
(71)

uniformly for $1/2 + 1/\log \log \tau \leq \sigma \leq 3/2$, where $\mathcal{R}_1(\sigma)$ and $\mathcal{R}_2(\sigma)$ are defined by (37) and (38), respectively. In addition,

$$\mathcal{R}_1(\sigma) \ll_{\lambda^+} \frac{f}{\left(\log \tau\right)^2}, \quad \mathcal{R}_2(\sigma) \ll m_{\mathcal{L}} \left(\frac{\mathbf{q}_{\mathcal{L}}}{\tau}\right)^{\frac{2}{\mathbf{d}_{\mathcal{L}}}}$$
(72)

uniformly for $|1 - \sigma| \leq 1/\log \log \tau$. Note that

$$\frac{\left(\log\log\tau\right)^2}{\left(2\sigma-1\right)\left(\log\tau\right)^{2\sigma-1}} \ll \frac{1}{(2\sigma-1)^3}$$

. . .

uniformly for $\sigma > 1/2$. By (43) we thus have

$$S_{\mathcal{L},x,y}(\sigma) \le A(m,\alpha,\sigma,\sigma) \left(\log \tau\right)^{2-2\sigma} - \frac{m\sigma 2^{1-\sigma}}{1-\sigma} + O\left(\frac{m}{(2\sigma-1)^3}\right)$$

uniformly for $1/2 + 1/\log \log \tau \le \sigma \le 1 - 1/\log \log \tau$, where $A(a, \alpha, u, \sigma)$ is defined by (6). In addition, by (44) we have

$$S_{\mathcal{L},x,y}(\sigma) \le 2m \log \log \tau + O\left(m + m|1 - \sigma| \left(\log \log \tau\right)^2\right)$$

uniformly for $|1 - \sigma| \leq 1/\log \log \tau$. The proof of inequality (11) is complete after employing the above estimates on all terms from (39).

We are using (40) to prove the second part of Theorem 2. It follows by (70) and (71) that

$$\int_{\sigma}^{\frac{3}{2}} \mathcal{Z}(\sigma') \, \mathrm{d}\sigma' \leq \left(\frac{e^{\alpha} + 1}{4\alpha}\right) \frac{\left(\log \tau\right)^{2-2\sigma}}{\log\log \tau} \\ + O\left(\frac{m_{\mathcal{L}} \left(\log \tau\right)^{3-4\sigma}}{\log\log \tau} + \frac{m}{\left(\log \tau\right)^{2\sigma-1}} + \frac{\mathrm{d}_{\mathcal{L}} + m_{\mathcal{L}}}{\left(\log \tau\right)^{2\sigma-1}\log\log \tau}\right) + O_{\lambda^{-}}\left(\frac{f}{\left(\log \tau\right)^{2\sigma-1}\log\log \tau}\right)$$

and

$$\int_{\sigma}^{\frac{3}{2}} \mathcal{R}_{1}(\sigma') \, \mathrm{d}\sigma' \ll_{\lambda^{+}} \frac{f}{(2-\alpha) \left(\log \tau\right)^{(2-\alpha)\sigma} \log \log \tau},$$
$$\int_{\sigma}^{\frac{3}{2}} \mathcal{R}_{2}(\sigma') \, \mathrm{d}\sigma' \ll m_{\mathcal{L}} \left(\frac{q_{\mathcal{L}}}{\tau}\right)^{\frac{2}{d_{\mathcal{L}}}} \frac{\left(\log \tau\right)^{2-2\sigma}}{\log \log \tau}$$

uniformly for $1/2 + 1/\log \log \tau \leq \sigma \leq 3/2$. Because $\mathcal{R}_1(\sigma) \ll_{\lambda^+} f \cdot (\log \tau)^{-2\sigma}$ uniformly for 1 - 1 $1/\log\log\tau < \sigma < 3/2$, we have by the above estimates that

$$\int_{\sigma}^{\frac{3}{2}} \left(\mathcal{Z} + \mathcal{R}_1 + \mathcal{R}_2\right) (\sigma') \mathrm{d}\sigma' = O\left(m + \frac{\mathrm{d}_{\mathcal{L}} + m_{\mathcal{L}}}{\log \tau \log \log \tau} + \left(\frac{\mathrm{q}_{\mathcal{L}}}{\tau}\right)^{\frac{2}{\mathrm{d}_{\mathcal{L}}}} \frac{m_{\mathcal{L}}}{\log \log \tau}\right) + O_{\lambda^+,\lambda^-} \left(\frac{f}{\log \tau \log \log \tau}\right)$$

uniformly for $|1 - \sigma| \leq 1/\log \log \tau$. By (47) and (42) we have

$$\begin{aligned} \widehat{S}_{\mathcal{L},x,y}(\sigma) &\leq \eta(\alpha,\sigma,\tau)A\left(m,\alpha,\sigma,\sigma\right) \frac{\left(\log\tau\right)^{2-2\sigma}}{\log\log\tau} - \frac{m\eta(\alpha,\sigma,\tau)}{\left(1-\sigma\right)\log\log\tau} \\ &+ m\log\left(2\log\log\tau\right) + O\left(\frac{m\left(\log\tau\right)^{2-2\sigma}}{\left(1-\sigma\right)^2\left(\log\log\tau\right)^2} + \frac{m}{(2\sigma-1)^2}\right) \end{aligned}$$

and

$$\mathcal{E}_x \ll \frac{m}{\left(\log \tau\right)^{1-\frac{\alpha}{2}} \log \log \tau}$$

uniformly for $1/2 + 1/\log \log \tau \le \sigma \le 1 - 1/\log \log \tau$, where $\eta(\alpha, \sigma, \tau)$ is defined by (7). In addition, by (48) and (42) we have

$$\widehat{S}_{\mathcal{L},x,y}(\sigma) \le m \log \left(2 \log \log \tau\right) + O(m)$$

and $\mathcal{E}_x \ll m$ uniformly for $|1 - \sigma| \leq 1/\log \log \tau$. The proof of inequality (12) is complete after employing the above estimates on all terms from (40).

Proof of Theorem 3. Assume the notation and conditions from Theorem 3. Let $\sigma = 1$, and let τ and $|t| \ge \mu^+ / \lambda^-$ be sufficiently large. By (70) we have

$$\mathcal{Z}(1) \le \frac{e^{\alpha} + 1}{2\alpha} + O\left(\frac{m \log \log \tau}{\log \tau} + \frac{\mathrm{d}_{\mathcal{L}} + m_{\mathcal{L}}}{\log \tau}\right) + O_{\lambda^{-}}\left(\frac{f}{\log \tau}\right)$$

and by (45) we also have

$$S_{\mathcal{L},x,y}(1) \le 2m \log \log \tau - m (\gamma + \alpha) + O\left(\frac{m (\log \log \tau)^2}{\log \tau}\right)$$

Then (17) follows by taking the last two bounds and (72) in (39). To prove (18), note that

$$\int_{1}^{\frac{3}{2}} \left(\mathcal{Z} + \mathcal{R}_{1} + \mathcal{R}_{2}\right) \left(\sigma'\right) \mathrm{d}\sigma' \leq \frac{e^{\alpha} + 1}{4\alpha \log \log \tau} + O\left(\frac{m}{\log \tau} + \frac{\mathrm{d}_{\mathcal{L}} + m_{\mathcal{L}}}{\log \tau \log \log \tau}\right) + O_{\lambda^{-}}\left(\frac{f}{\log \tau \log \log \tau}\right)$$

$$+ O_{\lambda^{+}} \left(\frac{f}{\left(\log \tau\right)^{2} \log \log \tau} \right) + O\left(\frac{m_{\mathcal{L}}}{\log \log \tau} \left(\frac{q_{\mathcal{L}}}{\tau} \right)^{\frac{2}{d_{\mathcal{L}}}} \right),$$

see the proof of Theorem 2 for details. Also, $\mathcal{E}_x \ll m/(\log \tau \log \log \tau)$ by (42), and

$$\widehat{S}_{\mathcal{L},x,y}(1) \le m \log\left(2\log\log\tau\right) + m\gamma + O\left(\frac{m\log\log\tau}{\log\tau}\right)$$

by (49). Then (18) follows by taking the last three inequalities in (40).

7. Proof of Theorems 4 and 5

Proof of Theorem 4. The proof of both estimates from (30) is relatively simple and holds unconditionally since $\mathcal{L} \in S\mathcal{P}$ and thus

$$\left| \frac{\mathcal{L}'}{\mathcal{L}}(s) \right| \le m \sum_{p} \sum_{k=1}^{\infty} \frac{\log p}{p^{k\sigma_0}} = -m \frac{\zeta'}{\zeta}(\sigma_0) \le \frac{m}{\sigma_0 - 1},$$
$$\left| \log \mathcal{L}(s) \right| \le m \sum_{p} \sum_{k=1}^{\infty} \frac{1}{k p^{k\sigma_0}} = m \log \zeta(\sigma_0) \le m \log \frac{1}{\sigma_0 - 1} + m\gamma(\sigma_0 - 1)$$

for $\sigma \geq \sigma_0 > 1$. The second inequality in the former expression follows from [Del87] or [MV07, Exercise 11.1.1(3)], and the second inequality in the latter expression follows from [Ram16, Lemma 5.4]. When we set $\sigma_0 = 1 + \alpha_3 / \log \log \tau$, we have proved the case (30).

Next, we are going to prove (25), i.e., upper bounds for the terms $|\mathcal{L}'(s)/\mathcal{L}(s)|$ and $|\log \mathcal{L}(s)|$ under assumption (1) of Theorem 4. Assume these conditions. By (39), it is sufficient to estimate the terms $S_{\mathcal{L},x,y}(\sigma)$, $\mathcal{Z}(\sigma)$, $\mathcal{R}_1(\sigma)$ and $\mathcal{R}_2(\sigma)$ in order to estimate the term $|\mathcal{L}'(s)/\mathcal{L}(s)|$. Now $S_{\mathcal{L},x,y}(\sigma)$ can be bounded by (43) and Lemma 8 can be used in order to estimate $\mathcal{Z}(\sigma)$. To bound $\mathcal{R}_1(\sigma)$ from (37), note that

$$\exp\left(\frac{2\alpha\sigma}{2\sigma-1}\right) \le (\log\tau)^{\frac{\alpha}{\alpha_1}}$$

for $\sigma \geq 1/2 + \alpha_1/\log\log\tau$, and

$$\sum_{j=1}^{f} \lambda_j^2 \left(\frac{\Gamma'}{\Gamma}\right)' \left(\frac{\lambda_j}{2}\right) = \frac{f}{4} \left(\frac{\Gamma'}{\Gamma}\right)' \left(\frac{1}{4}\right) \le 4.3 \mathrm{d}_{\mathcal{L}}$$
(73)

by the strong λ -conjecture. Hence,

$$\mathcal{R}_{1}(\sigma) \leq \frac{4.3 \mathrm{d}_{\mathcal{L}} \left(\sigma - \frac{1}{2}\right)}{\alpha \left(\log \tau\right)^{2\sigma}} \left(1 + \left(\log \tau\right)^{\frac{\alpha}{\alpha_{1}}\sigma}\right).$$
(74)

We estimate $\mathcal{R}_2(\sigma)$ from (38) trivially by using $\sigma < 1$ leading to the estimate

$$\mathcal{R}_2(\sigma) < \frac{m_{\mathcal{L}}}{\alpha} \left(\frac{q_{\mathcal{L}}}{\tau}\right)^{\frac{2}{d_{\mathcal{L}}}} \left(\log \tau\right)^{2(1-\sigma)}.$$
(75)

Collecting all bounds for all terms, we get estimate (25) for the term $|\mathcal{L}'(s)/\mathcal{L}(s)|$. Now we are using (40) in order to prove (26), i.e., estimate for $|\log \mathcal{L}(s)|$. Hence, it is sufficient to estimate $\widehat{S}_{\mathcal{L},x,y}(\sigma)$, \mathcal{E}_x and integrals of $\mathcal{Z}(\sigma)$, $\mathcal{R}_1(\sigma)$ and $\mathcal{R}_2(\sigma)$. We are estimating $\widehat{S}_{\mathcal{L},x,y}(\sigma)$ by (47) and \mathcal{E}_x by (42). Note that Lemma 8 implies

$$\int_{\sigma}^{\frac{3}{2}} \mathcal{Z}\left(\sigma'\right) \mathrm{d}\sigma' \leq \frac{\left(e^{\alpha}+1\right) \left(\log \tau\right)^{2-2\sigma}}{4\alpha \log \log \tau} + \frac{\mathfrak{a}\left(e^{\alpha}+1\right) \left(\log \tau\right)^{3-4\sigma}}{4\alpha \log \log \tau} + \frac{m\left(e^{\alpha}+1\right)}{\alpha \left(\log \tau\right)^{2\sigma-1}} + \frac{\max\{0,\mathfrak{b}\}\left(e^{\alpha}+1\right)}{2\alpha \left(\log \tau\right)^{2\sigma-1} \log \log \tau},\tag{76}$$

and that (74) and (75) guarantee

$$\int_{\sigma}^{\frac{3}{2}} \mathcal{R}_1\left(\sigma'\right) \mathrm{d}\sigma' \le \frac{4.3 \mathrm{d}_{\mathcal{L}} \alpha_1 \left(1 + \left(\log \tau_0\right)^{-\frac{\alpha}{2\alpha_1}}\right)}{\alpha \left(2\alpha_1 - \alpha\right) \left(\log \tau\right)^{\left(2 - \frac{\alpha}{\alpha_1}\right)\sigma} \log \log \tau},\tag{77}$$

$$\int_{\sigma}^{\frac{3}{2}} \mathcal{R}_{2}(\sigma') \, \mathrm{d}\sigma' \leq \frac{m_{\mathcal{L}}\left(1 + e^{\frac{\alpha}{2}}\right)}{2\alpha} \left(\frac{q_{\mathcal{L}}}{\tau}\right)^{\frac{2}{d_{\mathcal{L}}}} \frac{\left(\log \tau\right)^{2-2\sigma}}{\log \log \tau}.$$
(78)

In the derivation of the last inequality we also used the fact that $(\sigma - 1)/(2\sigma - 1)$ is an increasing function. Collecting all bounds for all terms in (40) finally gives (26).

We are going to prove (27). Assume the conditions from the case (2) of Theorem 4. Note that

$$\frac{3}{4} < 1 - \frac{\alpha_2}{\log \log \tau_0} \le \sigma \le 1 + \frac{\alpha_3}{\log \log \tau_0} < \frac{3}{2}.$$
 (79)

Then $S_{\mathcal{L},x,y}(\sigma)$ can be bounded by (44) and Lemma 8 can be used for $\alpha_1 = \frac{1}{2} \log \log \tau_0 - \alpha_2$ in order to estimate $\mathcal{Z}(\sigma)$. We are using also

$$\left(\log \tau\right)^{2-2\sigma} \le 1 + 2\theta_2(M)|1 - \sigma|\log\log \tau,$$

where $\theta_2(u)$ and M are as in Theorem 4. The functions $\mathcal{R}_1(\sigma)$ and $\mathcal{R}_2(\sigma)$ are bounded trivially by using (73) and (79), while observing also that $\sigma/(2\sigma-1)$ is a decreasing function. Collecting all bounds for all terms in (39) gives (27). We are using (40) in order to prove (28). We are estimating $\widehat{S}_{\mathcal{L},x,y}(\sigma)$ by (48) and \mathcal{E}_x by (42), where we are using also

$$\frac{1}{2} \le \eta(\alpha, \sigma, \tau) \le \frac{1}{2} \left(1 - \frac{\alpha}{\log \log \tau_0 - 2\alpha_2} \right)^{-1}$$

Furthermore, the estimate (76) can be used with \mathfrak{a} and \mathfrak{b} replaced by \mathfrak{a}_1 and \mathfrak{b}_1 , the latter pair of functions being as in Theorem 4. Also, we are using (78), while (77) is replaced by

$$\int_{\sigma}^{\frac{3}{2}} \mathcal{R}_{1}(\sigma') \, \mathrm{d}\sigma' \leq \frac{4.3 \mathrm{d}_{\mathcal{L}} \left(1 + \exp\left(\frac{2\alpha(\log\log\tau_{0} - \alpha_{2})}{\log\log\tau_{0} - 2\alpha_{2}}\right)\right)}{2\alpha \left(\log\tau\right)^{2\sigma} \log\log\tau}$$

Collecting all bounds for all terms in (40) finally gives (28). The proof of Theorem 4 is thus complete.

Proof of Theorem 5. Note that the conditions of Theorem 5 guarantee that $x \ge 60$ and $y \ge 2$. Firstly, we are going to prove (31). We can use (45) for the estimation of $S_{\mathcal{L},x,y}(1)$. Moreover, Lemma 8 can be used for $\alpha_1 = \frac{1}{2} \log \log \tau_0$ in order to bound $\mathcal{Z}(1)$ from (36). Estimates for $\mathcal{R}_1(1)$ and $\mathcal{R}_2(1)$ from (37) and (38), respectively, can be obtained by trivial estimation while using also (73). Taking all estimates in (39) furnishes the proof of (31). Next, we need to prove also (32). We are estimating $\widehat{S}_{\mathcal{L},x,y}(1)$ by (49) and \mathcal{E}_x by (42). Furthermore, estimate (76) can be used for $\sigma = 1$ with \mathfrak{a} and \mathfrak{b} replaced by \mathfrak{a}_2 and \mathfrak{b}_2 , the latter pair of functions being as in Theorem 5. For the remaining two terms we are using (78) for $\sigma = 1$, and

$$\int_{1}^{\frac{3}{2}} \mathcal{R}_{1}\left(\sigma'\right) \mathrm{d}\sigma' \leq \frac{4.3 \mathrm{d}_{\mathcal{L}}\left(e^{2\alpha}+1\right)}{2\alpha \left(\log \tau\right)^{2} \log \log \tau}.$$

Inequality (32) now follows by (40). The proof of Theorem 5 is thus complete.

References

- [BMOR18] M. A. Bennett, G. Martin, K. O'Bryant, and A. Rechnitzer, Explicit bounds for primes in arithmetic progressions, Illinois J. Math. 62 (2018), no. 1-4, 427–532.
- [CC11] E. Carneiro and V. Chandee, Bounding $\zeta(s)$ in the critical strip, J. Number Theory **131** (2011), no. 3, 363–384. [CCM19] E. Carneiro, A. Chirre, and M. B. Milinovich, Bandlimited approximations and estimates for the Riemann
- zeta-function, Publ. Mat. **63** (2019), no. 2, 601–661.
- [Cha09] V. Chandee, Explicit upper bounds for L-functions on the critical line, Proc. Amer. Math. Soc. 137 (2009), no. 12, 4049–4063.
- [Chi19] A. Chirre, A note on entire L-functions, Bull. Braz. Math. Soc. (N.S.) 50 (2019), no. 1, 67–93.
- [CG22] A. Chirre and F. Gonçalves, *Bounding the log-derivative of the zeta-function*, Math. Z. **300** (2022), no. 1, 1041–1053.
- [CHS23] A. Chirre, M. V. Hagen, A. Simonič, Conditional estimates for the logarithmic derivative of Dirichlet Lfunctions, Indag. Math. (N.S.) (2023), DOI: 10.1016/j.indag.2023.07.005.
- [Dav00] H. Davenport, Multiplicative number theory, 3rd ed., Graduate Texts in Mathematics, Springer New York, NY, 2000.
- [Del87] H. Delange, Une remarque sur la dérivée logarithmique de la fonction zêta de Riemann, Colloq. Math. 53 (1987), no. 2, 333–335.
- [EHP22] A.-M. Ernvall-Hytönen and N. Palojärvi, Explicit bound for the number of primes in arithmetic progressions assuming the generalized riemann hypothesis, Math. Comp. 91 (2022), no. 335, 1317–1365.
- [Fin16] R. H. Finder, Variation-diminishing maximal operators and the argument of L-functions, PhD, Instituto de Matemática Pura e Aplicada, 2016.

N. PALOJÄRVI AND A. SIMONIČ

- [IK04] H. Iwaniec and E. Kowalski, Analytic number theory, American Mathematical Society Colloquium Publications 53, AMS, Providence, RI, 2004.
- [LLS15] Y. Lamzouri, X. Li, and K. Soundararajan, Conditional bounds for the least quadratic non-residue and related problems, Math. Comp. 84 (2015), no. 295, 2391–2412.
- [LT22] A. Languasco and T. S. Trudgian, Uniform effective estimates for $|L(1,\chi)|$, J. Number Theory **236** (2022), 245–260.
- [Lit25] J. E. Littlewood, On the Riemann Zeta-Function, Proc. London Math. Soc. (2) 24 (1925), no. 3, 175–201.
- [Lit28] _____, Mathematical Notes (5): On the function $1/\zeta(1+ti)$, Proc. London Math. Soc. (2) **27** (1928), no. 5, 349–357.
- [Lum18] A. Lumley, Explicit bounds for L-functions on the edge of the critical strip, J. Number Theory 188 (2018), 186–209.
- [MV07] H. L. Montgomery and R. C. Vaughan, Multiplicative number theory. I. Classical theory, Cambridge Studies in Advanced Mathematics, vol. 97, Cambridge University Press, Cambridge, 2007.
- [Mur08] M. R. Murty, Problems in analytic number theory, 2nd ed., Graduate Texts in Mathematics, vol. 206, Springer, New York, 2008, Readings in Mathematics.
- [Per05] A. Perelli, A survey of the Selberg class of L-functions. I, Milan J. Math. 73 (2005), 19–52.
- [Ram16] O. Ramaré, An explicit density estimate for Dirichlet L-series, Math. Comp. 85 (2016), no. 297, 325–356.
 [RS62] J. B. Rosser and L. Schoenfeld, Approximate formulas for some functions of prime numbers, Illinois J. Math.
- 6 (1962), 64–94. [Sch76] L. Schoenfeld, Sharper bounds for the Chebyshev functions $\theta(x)$ and $\psi(x)$. II, Math. Comp. **30** (1976), no. 134, 337–360.
- [Sel43] A. Selberg, On the normal density of primes in small intervals, and the difference between consecutive primes, Arch. Math. Naturvid. 47 (1943), no. 6, 87–105.
- [Sel92] _____, Old and new conjectures and results about a class of Dirichlet series, Proceedings of the Amalfi Conference on Analytic Number Theory (Maiori, 1989), Univ. Salerno, Salerno, 1992, pp. 367–385.
- [Sim22a] A. Simonič, Explicit estimates for $\zeta(s)$ in the critical strip under the Riemann Hypothesis, Q. J. Math. 73 (2022), no. 3, 1055–1087.
- [Sim22b] _____, On explicit estimates for S(t), $S_1(t)$, and $\zeta(1/2+it)$ under the Riemann Hypothesis, J. Number Theory **231** (2022), 464–491.
- [Sim23] _____, Estimates for L-functions in the critical strip under GRH with effective applications, Mediterr. J. Math. 20 (2023), no. 2, paper no. 87, 24 pp.
- [Ste07] J. Steuding, Value-distribution of L-functions, Lecture Notes in Mathematics, vol. 1877, Springer, Berlin, 2007.
 [Tit86] E. C. Titchmarsh, The theory of the Riemann zeta-function, 2nd ed., The Clarendon Press, Oxford University Press, New York, 1986.

DEPARTMENT OF MATHEMATICS AND STATISTICS, THE UNIVERSITY OF HELSINKI, HELSINKI, FINLAND *Email address*: neea.palojarvi@helsinki.fi

SCHOOL OF SCIENCE, THE UNIVERSITY OF NEW SOUTH WALES (CANBERRA), ACT, AUSTRALIA *Email address*: a.simonic@student.adfa.edu.au