A Black Hole Kicked At Birth: MAXI J1305-704

CHASE KIMBALL,¹ SAM IMPERATO,^{1,2} VICKY KALOGERA,¹ KYLE A. ROCHA,^{3,4} ZOHEYR DOCTOR,⁵ JEFF J. ANDREWS,^{5,6} AARON DOTTER,⁵ EMMANOUIL ZAPARTAS,^{7,8} SIMONE S. BAVERA,⁷ KONSTANTINOS KOVLAKAS,⁷ TASSOS FRAGOS,⁷ PHILIPP M. SRIVASTAVA,^{3,9} DEVINA MISRA,⁷ MENG SUN,³ AND ZEPEI XING⁷

¹Center for Interdisciplinary Exploration and Research in Astrophysics (CIERA), Department of Physics and Astronomy, Northwestern University, 1800 Sherman Avenue, Evanston, IL 60201, USA

²Department of Astronomy, Boston University, 725 Commonwealth Avenue Boston, MA 02215, USA

³Center for Interdisciplinary Exploration and Research in Astrophysics (CIERA), Northwestern University, 2145 Sheridan Road,

Evanston, IL 60208, USA

⁴Department of Physics and Astronomy, Northwestern University, 2145 Sheridan Road, Evanston, IL 60208, USA

⁵ Center for Interdisciplinary Exploration and Research in Astrophysics (CIERA), Department of Physics and Astronomy, Northwestern University, 1800 Sherman Avenue, Evanston, IL 60201, USA

⁶Department of Physics, University of Florida, 2001 Museum Rd, Gainesville, FL 32611, USA

⁷ Département d'Astronomie, Université de Genève, Chemin Pegasi 51, CH-1290 Versoix, Switzerland

⁸IAASARS, National Observatory of Athens, Vas. Pavlou and I. Metaxa, Penteli, 15236, Greece

⁹ Electrical and Computer Engineering, Northwestern University, 2145 Sheridan Road, Evanston, IL 60208, USA

ABSTRACT

When a compact object is formed in a binary, any mass lost during core collapse will impart a kick on the binary's center of mass. Asymmetries in this mass loss or neutrino emmission would impart an additional natal kick on the remnant black hole or neutron star, whether it was formed in a binary or in isolation. While it is well established that neutron stars receive natal kicks upon formation, it is unclear whether black holes do as well. Here, we consider the low-mass X-ray binary MAXI J1305-704, which has been reported to have a space velocity $\gtrsim 200$ km/s. In addition to integrating its trajectory to infer its velocity upon formation of its black hole, we account for recent estimates of its period, black hole mass, mass ratio, and donor effective temperature from photometric and spectroscopic observations. We find that if MAXI J1305-704 formed via isolated binary evolution in the thick Galactic disk, then the supernova that formed its black hole imparted a natal kick of at least 70 km/s while ejecting less than $\simeq 1 \ M_{\odot}$ with 95% confidence assuming uninformative priors on mass loss and natal kick velocity.

Keywords: Astrophysical black holes, Binary evolution, Supernovae, Natal kicks

1. INTRODUCTION

The physics underlying supernovae (SNe) and explosive mass loss may be imprinted on the velocities of the compact objects they leave behind. While any sudden mass lost from a component of a binary system will impart a velocity on the center of mass proportional to the amount of mass lost (Blaauw 1961), the compact object remnants themselves may also receive an additional "natal" kick on the order of hundreds of kilometers per second due to asymmetries in the SN mechanism (Shklovskii 1970; Janka 2012; Wongwathanarat et al. 2013). Surveys of the proper motions and heights above the Galactic disk (Lyne & Lorimer 1994; Hobbs et al. 2005; Kapil et al. 2022) have strongly suggested that neutron star remnants must experience these kicks, as have binary evolutionary analyses (e.g. Wong et al. (2010); Tauris et al. (2017) and references therein). However, the evidence that black holes (BHs) receive natal kicks is less clear.

In recent years, aided by the growing wealth of BH observations, studies have worked toward constraining BH natal kicks using a variety of methods and data sets. A number of studies have focused on the population of massive runaway stars with large space velocities, considering the possibility that they may have originated in binaries that were disrupted via the core collapse of their companions, possibly with the assistance of natal kicks (van Rensbergen et al. 1996; De Donder et al. 1997;

CharlesKimball2022@u.northwestern.edu

Dray et al. 2005; Eldridge et al. 2011; Boubert & Evans 2018; Renzo et al. 2019; Aghakhanloo et al. 2022).

Other studies have turned to X-ray binaries (XRBs) with BH accretors – aided by the growing subset of systems with known proper motions and radial velocities to constrain natal BH kicks. Some used just these kinematic constraints to estimate birth velocities of known XRBs, suggesting that they may have received kicks in excess of what they could have received due to symmetric mass loss alone (Mirabel et al. 2001; Repetto et al. 2012; Atri et al. 2019; Sánchez et al. 2021). Others combined the kinematic constraints with observational constraints on the orbital, BH, and donor properties, modeling the binary evolution and core collapse of their XRB progenitors to individually constrain mass loss and natal kicks (Willems et al. 2005; Fragos et al. 2009; Wong et al. 2012, 2014). Of these studies that considered both kinematic and binary/stellar observational constraints, most found only upper limits on the BH natal kicks with one exception: XTE J1118+40, which Fragos et al. (2009) found must have received a natal kick in excess of $\simeq 80$ km/s. Here, we focus on the case of the low-mass X-ray binary (LMXB) MAXI J1305-704 because of its high peculiar velocity ($V_{\rm pec} \simeq 80^{+30}_{-30}$ km/s), distance above the Galactic plane ($|Z| \simeq 1$ kpc), and short orbital period (P $\simeq 0.4$ days), reported in Sánchez et al. (2021). We constrain both its evolutionary and kinematic history to find a lower limit on its BH natal kick, making it only the second fully-modeled BH system that must have received a natal kick at birth.

While BH LMXBs may form dynamically via binary exchanges or single-single captures (Clark 1975; Hills 1976), here we work under the assumption that MAXI J1305-704 (hereafter J1305) formed in the Galactic field and evolved as an isolated binary. The standard isolated formation channel for BH LMXBs usually starts with an unequal-mass zero-age main sequence binary with a primary $\gtrsim 20 M_{\odot}$ that fills its Roche lobe (RL), undergoing a period of stable or unstable mass transfer (MT) – the latter case leading to a common envelope (CE) phase. If the binary survives without merging first, the primary is left as a massive stripped naked helium star in a tight, circularized, orbit with a relatively unevolved companion. Upon core collapse, the orbit may be altered or completely disrupted due to mass ejection and a possible natal kick. If it survives, the resulting detached binary will evolve until the secondary fills its Roche lobe and begins transferring mass onto the primary BH, forming a BH LMXB. In the case of J1305, with a sub-solar donor mass with a likely convective envelope, orbiting its BH with a sub-day period, the onset of RLO may have been aided by magnetic braking, which would have counteracted the tendency of the orbit to expand via mass transfer from the lighter secondary onto the heavier BH (Postnov & Yungelson 2014).

In this study, we combine forward-modeling of this evolutionary channel with backward modeling of the kinematic history of J1305 to constrain the range of natal kicks its BH may have experienced – following a method developed by Willems et al. (2005), Fragos et al. (2009), and Wong et al. (2012, 2014). While the kinematic history gives us constraints on the systemic velocity of the system at the birth of the BH, the forward modeling allows us to extract any asymmetric natal kick - in addition to the Blaauw kick from mass-loss alone - needed to attain that velocity. We start by evolving J1305's trajectory backwards through the Galactic potential to construct a distribution of potential peculiar velocities of the system at the birth of its BH. We then run a series of detailed mass transfer sequences using MESA¹ (Paxton et al. 2010, 2013, 2015, 2019) from the RLO of the secondary (RLO2) forwards, identifying potential progenitor binaries that simultaneously satisfy the observations of J1305's BH mass, mass ratio, donor effective temperature, and orbital period. We then use POSYDON (Fragos et al. 2022) to constrain the parameter space of the possible progenitors that could produce these RLO2 binaries.

We note that when modeling core collapse, we choose not to adopt a SN prescription and instead sample massloss and natal kick velocity from uniform, uninformative priors. Modern SN prescriptions typically scale BH kicks according to ejecta mass which is calculated via fitting formulae (see, e.g. Bray & Eldridge (2018); Giacobbo & Mapelli (2020) and references therein). However, BH kicks and the mechanisms that produce them are poorly understood, and here we aim to *infer* magnitude of SN mass loss and BH kicks under uninformative priors, potentially informing SN prescriptions.

Using our kinematic constraint on the total birth velocity of J1305, we then extract the range of required natal BH kicks from our forward modeling, finding that J1305's BH likely received a natal kick greater than 70 km/s at 95% confidence. In Section 2, we discuss the observational constraints on the stellar, binary, and kinematic properties of J1305. In Section 3.1 and 3.2, we describe how we reconstruct its kinematic history and constrain its possible pre- and post-SN properties, combining the results in 3.3. We discuss our conclusions in Section 4.

 $^{^1}$ Revision 11701 and the 20190503 version of the MESA software development kit

2. MAXI J1305-704

While there are hundreds of known XRB sources, only about two dozen have confirmed BH accretors. Of these, 21 are classified as LMXBs (Jonker et al. 2021), having companions less than a few solar masses. There are proper motion measurements in the literature for 16 of these, and only 12 are complete with radial velocity measurements (Atri et al. 2019), crucial to estimating peculiar velocities and inferring BH kicks. Of these systems, J1305 is particularly interesting because of its high distance above the Galactic plane, with $|Z| \approx 1$ kpc, and low orbital period (≈ 0.4 days), which is particularly helpful in constraining its progenitor properties (see Section 3.2 for details). J1305 was first discovered by the International Space Station's Monitor of All-sky X-ray Image instrument (Matsuoka et al. 2009; Mihara et al. 2011), and was first identified as an X-ray transient in Sato et al. (2012). Follow-up observations classified J1305 as an LMXB, potentially accreting onto a stellar-mass BH (Greiner et al. 2012; Kennea et al. 2012; Suwa et al. 2012; Morihana et al. 2013). The nature of its accretor was confirmed by Sánchez et al. (2021), after photometric and spectroscopic observations of J1305 in guiescence. They found that J1305 has an orbital period of $0.394^{+0.004}_{-0.004}$ days, consisting of a black hole of mass $M_{1,obs} = 8.9^{+1.6}_{-1.0} M_{\odot}$. With an observed mass ratio $q_{obs} = 0.05^{+0.02}_{-0.02}$ and donor effective temperature $T_{\rm eff,obs} = 4610^{+130}_{-160}$ K, they find that the companion is an evolved dwarf star with $M_{2,\text{obs}} = 0.43^{+0.16}_{-0.16} M_{\odot}$ and an effective temperature of $T_{\text{eff}} = 4610^{+130}_{-160}$ K. They es-timated that J1305 is at a distance of $d = 7.5^{+2.8}_{-1.4}$ kpc, with a radial velocity of $\gamma = 9^{+5}_{-5}$ km/s. Combining this together with proper motion measurements in the direction of right ascension and declination (α and δ , respectively) from GAIA (Gaia Collaboration et al. 2021) of $\Delta \alpha \cos \delta = -7.89^{+0.62}_{-0.62} \text{ mas yr}^{-1} \text{ and } \Delta \delta = -0.16^{+0.72}_{-0.72},$ they calculated a peculiar velocity with respect to the local Galactic rotation of $V_{\rm pec} = 80^{+30}_{-30}$ km/s. While we can not exclude that J1305 originated in a globular cluster, this suggests that if it indeed originated in the thick disk, then J1305 may have received a natal kick at birth.

3. CONSTRAINING THE PROGENITOR OF J1305

We mostly follow the methodology described in Wong et al. (2014), using kinematic modeling to infer the potential peculiar velocities of J1305 at the birth of its black hole, and then constrain the possible pre- and post-SN binary properties using forward modeling to extract the asymmetric natal kick on the black hole from the total imparted peculiar velocity inferred kinematically. However, rather than use a rapid population synthesis code based on fits to single-star models when evolving potential progenitors, we compute detailed binary stellar evolution sequences using MESA, and generate populations with POSYDON (Fragos et al. 2022), which is trained on MESA binary evolution grids and fully consistent with its treatment of binary stellar evolution and mass transfer. Except for where otherwise noted in relation to our treatment of core collapse, all of our parameter choices and prescriptions are as described in POSYDON v1.

3.1. Kinematics

We begin by using the observed proper motion of J1305 to constrain the total velocity imparted by the SN upon the birth of its BH. Following the methodology of Wong et al. (2014), we assume that J1305 formed in the Galactic disk and integrate its trajectory backwards to infer the systemic velocity of J1305 upon the birth of its black hole ($V_{\rm pec,birth,obs}$). In Section 3.2, we infer the possible range of pre- and post-SN properties of the binary to extract the range of natal kicks required to create this birth velocity.

We model the Milky Way with the static potential of Carlberg & Innanen (1987) and updated parameters from Kuijken & Gilmore (1989). We start with a coordinate system with axes XYZ with origin at the Galactic center, where Z=0 coincides with the Galactic mid-plane, the positive Y axis is in the direction of Galactic rotation at the location of the Sun, and the negative X-axis connects the projection of the Sun onto Galactic midplane to the Galactic center. Assuming a distance from the Sun to the Galactic center of $R_0 =$ 8.05 kpc and local rotational velocity of $\Omega_0 = 238$ km/s Honma et al. (2012), this places the Sun at $(X_{\odot}, Y_{\odot}, Z_{\odot})$ = (-8.05, 0, .03) kpc with a velocity of $(U_{\odot}, V_{\odot}, W_{\odot}) =$ (11.1, 12.24, 7.25) km/s with respect to the local standard of rest. Meanwhile, adopting the distance to J1305 of $d = 7.5^{+2.8}_{-1.4}$ kpc from the analysis in Sánchez et al. (2021), this puts J1305 at (X,Y,Z) = (-3.87, -6.15, -6.15)-0.98) kpc. Combining this with the radial velocity $\gamma = 9^{+5}_{-5}$ km/s from that analysis with the angular velocity measurements $(\Delta \alpha, \Delta \delta) = (-7.89^{+0.62}_{-0.62}, -0.16^{+0.72}_{-0.72})$ mas yr^{-1} from the GAIA Early Data Release 3 (Gaia Collaboration et al. 2021), we compute the proper motion of J1305 with respect to our local standard of rest, finding (U,V,W) = (-227, -140, 20) km/s.

In order to estimate J1305's peculiar velocity upon the birth of its black hole, we integrate its trajectory backwards in time, and sample potential peculiar velocities at moments that coincide with crossings of the Galactic mid-plane within 10 Myr of the age of a donor star from the winning RLO2 sequences. We take the donor age



Figure 1. Here we plot the possible kinematic histories of J1305. In blue we plot the trajectory corresponding to taking the median observed position and proper motion as initial conditions. In grey we draw random initial conditions from the corresponding posteriors. In the top two panels, we plot its trajectory in the X-Y and X-Z planes in Galactocentric coordinates. In the bottom panel, we plot a segment of the peculiar velocity as a function of time before the present. The rotated histogram on the right is the distribution of potential peculiar velocities at birth, inferred by sampling the trajectory at crossings of the Galactic plane that coincide (to within 10 Myrs) with donor ages from our successful RLO2 binaries.

as a proxy for black hole age under the assumption that the lifetime of the black hole progenitor is negligible with respect to the age of our winning donors. We plot the result in Figure 1, finding that J1305 likely had a peculiar velocity of $V_{\text{pec,birth,obs}} = 74^{+19}_{-11} \text{ km/s}$ just after the birth of its black hole, and treat this as an observable. Assuming that prior to the progenitor to J1305 was moving in the Galactic mid-plane with the local Galactic rotation, this gives a measure of the total velocity imparted on J1305's center of mass upon core-collapse. This assumption that J1305 was born directly in the disk at Z = 0 kpc with exactly the circular velocity at that location equates to assuming a "kinematically cold" disk. To check that our results are not affected by this assumption, we compared J1305's current peculiar velocity to that obtained when using the average velocities of systems nearby J1305's projection onto the Galactic midplane using GAIA (Gaia Collaboration et al. 2021), and find no significant quantitative difference. We also reran

the analysis using gala's MilkyWayPotential (Price-Whelan 2017; Price-Whelan et al. 2020; Bovy 2015) and found that our results are not significantly affected by our particular choice of Galactic potential.

3.2. Binary Evolution

We now use the possible pre- and post-SN parameter space of J1305's progenitor in order to extract the BH natal kick from the total imparted systemic velocity inferred from its kinematics. We do this in two steps: First we constrain the BH mass, donor mass, and period $(M_{1,\text{RLO2}}, M_{2,\text{RLO2}}, P_{\text{RLO2}})$ at the onset of mass transfer from the secondary onto the black hole. We consider a successful RLO2 system to be any binary that evolves to simultaneously satisfy observational constraints on the BH and donor masses, donor effective temperature, and orbital period of J1305 to within 2- σ of their observed values. We then use POSYDON to evolve a population of ZAMS binaries up until RLO2 while modeling and sampling over natal kick velocities, matching the



Figure 2. The viable RLO2 parameter space explored with MESA. The colored shapes mark successful binaries (at different BH masses) that evolved to simultaneously satisfy all observational constraints on the BH and donor masses, donor effective temperature, and orbital period of J1305 to within $2-\sigma$. In black are unsuccessful binaries where magnetic braking was efficient, shrinking their periods throughout mass transfer. In grey are unsuccessful binaries above the bifurcation period, expanding during mass transfer.



Figure 3. Example of a successful RLO2 system. The 1.8 M_{\odot} donor fills its Roche lobe at a period of 0.75 days with an 8 M_{\odot} BH. The orange line denotes the age of the donor at the observed period. The green shaded regions mark the 2- σ confidence intervals on the observed black hole mass, mass ratio, and donor effective temperature.



Figure 4. Posteriors over parameters and kicks of the potential progenitor binaries just before and after SN. In blue we plot the unweighted posteriors. In green, orange, purple, and red we weight these posteriors individually according to the observed $V_{\text{pec,birth,obs}}$, $M_{1,\text{obs}}$, q_{obs} , $T_{\text{eff,obs}}$. In black we plot our final result, weighted by the product of the individual weights.

results to our successful RLO2 systems and identifying potential J1305 progenitors.

The mass of the black hole at RLO2 is constrained from above by the upper bound on the observed black hole mass $M_{bh,obs} \leq 12.1 M_{\odot}$. Meanwhile, the RLO2 period is constrained from below by the requirement that the donor is not already filling its Roche lobe at ZAMS, and from above by the requirement that it will fill its Roche lobe in a Hubble time. With the exception of very low donor masses $M_{2,\text{RLO2}} \leq 0.75 M_{\odot}$, this lower bound means that J1305's orbit must have *shrunk* since RLO2 ². Since subsequent mass transfer from the donor onto the BH would expand J1305's orbit (q < 1), this is only possible with efficient magnetic braking. Assuming the companion stays tidally locked, spin angular momentum lost through magnetic braking would be compensated by the orbital angular momentum of the binary, shrinking its orbit. Since magnetic braking is inefficient in massive stars and at large periods, this adds the constraint that $M_{2,\text{RLO2}}$ is sufficiently small so that by the time it loses enough mass to enter the regime where magnetic braking is efficient, the period is still below the bifurcation period that delineates whether mass transfer will grow or shrink the orbit in the presence of magnetic braking. We find that this effectively limits $M_{2,\text{RLO2}} < 3 \ M_{\odot}$ and $P_{\text{RLO2}} < 1$ day.

From these constraints, we construct a grid of binaries – spaced uniformly in BH mass, donor mass, and orbital period at intervals of 1 M_{\odot} , 0.2 M_{\odot} , and 0.05 days, respectively – for which we calculate detailed mass transfer sequences using MESA. Figure 2 shows results for selected slices of that grid. Of these sequences, we find 12 – marked in colored shapes in Figure 2 – that simultaneously satisfy observational constraints on the masses, period, and effective donor temperature of J1305 within $2-\sigma$. We refer to these as successful RLO2 binaries. We plot unsuccessful binaries that expanded throughout mass transfer with grey circles, and those below the bifurcation period that shrank due to efficient magnetic braking with black circles. In Figure 3, we plot an example of a successful RLO2 mass transfer sequence.

Having identified the space of potential J1305 progenitors at RLO2, we can continue to narrow down the potential space of J1305 progenitors at ZAMS. We begin by evolving a nominal population of 10,000,000 ZAMS binaries via POSYDON's interpolation scheme over the MESA grids described in Fragos et al. (2022). We choose a population flat in mass ratio, with orbital separation sampled log-uniform between 5 and 10⁴ R_☉, and primary mass $M_{1,ZAMS}$ drawn from a Kroupa IMF (Kroupa 2001) between 7 and 120 M_{\odot} . It is reasonable to assume that J1305 did not undergo mass inversion and that M_2 will lose a negligible amount of mass between ZAMS and RLO2 – consistent with predictions of negligible winds in low-mass stars – so we limit $M_{2,ZAMS} < 3 M_{\odot}$ by the above argument that requires progenitors at RLO2 to



Figure 5. Here we plot the final marginalized posterior over pre-SN primary mass and kick velocity, with the final, combined weights. The dotted, dashed, and solid contours enclose central 68%, 90%, and 99% probability.

be in a regime where magnetic braking is efficient. We evolve each binary until the core-collapse of the primary (CC1), discarding systems that merged before compact object formation. Again assuming that M_2 will lose negligible mass before RLO2, we also discard any systems where the pre-SN donor mass $M_{2,\text{preSN}} > 3 M_{\odot}$, leaving a population of ~30,000 pre-SN binaries that could viably be progenitors to J1305.

Finally we evolve the viable pre-SN binaries through the core collapse of the primary and the subsequent detached binary evolution up until RLO2. Rather than assume any particular supernova prescription, we take a broad, agnostic approach to modeling the core collapse, sampling mass loss uniformly such that the remnant black hole mass is between the minimum and maximum of our winning RLO black hole masses, and natal kicks $V_{\rm kick}$ uniformly between 0 and 500 km/s. We repeat this sampling 6,000 times per binary, calculating the post-supernova orbital properties and systemic velocities according to Kalogera (1996), and evolve the resulting BH + H-rich star binaries until the donor fills its Roche lobe at RLO2. We then identify potential J1305 progenitors from this population by matching them to the cells of our successful RLO2 binaries, forming ~ 850 potential evolutionary histories for J1305. We refer to these binaries as "potential progenitors". We note that by sampling uniformly over mass loss and kick velocity at SN, our span of potential progenitors is conservative. We discuss our results and how we weigh them by the observed properties of J1305 in Section 3.3.

3.3. Constraints on the natal BH kick and progenitor properties

 $^{^2}$ While it is possible that J1305 entered RLO2 immediately after a CE, or immediately post-SN due to a fortuitous kick, this becomes less likely when accounting for observations that find that the donor is evolved. Due to the high mass of the BH progenitor relative to the donor, this would make it unlikely that the donor began mass transfer at such an early stage given we are still observing it with such a short period today.

Since not all potential progenitors to J1305 are equally successful, we weigh the distributions over our potential progenitor parameters by the observed properties of J1305. That is, given a set of observable quantities $\vec{\theta}_f$ at present-day corresponding to a progenitor binary with parameters $\vec{\theta}_i$, $p(\vec{\theta}_i)$ is weighted by the probability of the observable quantities given the electromagnetic data EM:

$$p(\vec{\theta_i}|\text{EM}) \propto p(\vec{\theta_i})p(\vec{\theta_f}|\text{EM}).$$
 (1)

For $\vec{\theta}_f \in \{M_{1,\text{obs}}, q_{\text{obs}}, T_{\text{eff,obs}}\}$ we use (asymmetric as appropriate) Gaussians to fit the corresponding posteriors reported in Sánchez et al. (2021). For $V_{\text{pec,birth,obs}}$, we use a kernel density estimation to fit the posterior shown in the bottom-right panel of Figure 1. We plot the results in Figure 4. In blue we plot the unweighted posteriors, which is the set of binaries that at some point had a BH mass, mass ratio, and effective temperature within 2σ of the observed value. In green, orange, purple, and red we weight these posteriors individually according to the observed $V_{\text{pec,birth,obs}}$, $M_{1,\text{obs}}$, $q_{\rm obs}, T_{\rm eff,obs}$. Our final results, in black, are obtained by multiplying all of the observational weights together. In Figure 5, we plot the final marginalized posterior over the pre-SN primary mass and kick velocity, weighted by all observations.

Although we allowed for arbitrarily high mass loss upon core collapse, we find that $\Delta M_{1,CC} < 0.9 \ M_{\odot}$ with 95% confidence. With low $\Delta M_{1,CC}$ the velocity imparted on the center of mass via mass loss alone would have been small, with $V_{\Delta M} = 0.7^{+1.0}_{-0.5} \ \text{km/s}$. In the absence of an additional SN kick on the remnant BH, the systemic velocities would be incompatible with the inferred $V_{\text{pec,birth,obs}}$. Indeed, once weighted by observations – and crucially the kinematic constraints on the total birth velocity – we find that $V_{\text{kick}} > 70 \ \text{km/s}$ with 95% confidence, with $V_{\text{kick}} = 91^{+23}_{-12} \ \text{km/s}$.

The presence of an SN kick may also have helped J1305 reach the short periods at RLO2 that we find to be necessary to produce it. Since the progenitor would have circularized during a common envelope phase, the post-SN orbit would have mild eccentricity – and therefore would have difficulty shedding angular momentum – in the absence of a kick and with little to no mass loss. Indeed, we find that all but the shortest-period pre-SN potential progenitors received a kick. Before weighting by the inferred $V_{\rm pec,birth}$, we find that of the progenitors that received a kick of < 10 km/s, all had pre-SN periods of <5 days, and none had post-SN eccentricities of > 0.1.

We stress that these results should be understood as constraints inferred using flat priors, and are therefore conservative. Our finding of a scenario with low massloss and non-negligible natal kick are at odds with SN prescriptions that scale BH kicks by mass-loss (Bray & Eldridge 2018; Giacobbo & Mapelli 2020) resulting in negligible kicks with low mass-loss. On the other hand, simulations in Coleman & Burrows (2022) – listed in Table 1 and discussed in Section 3.1.3 therein – find that BHs may receive natal kicks of up to $\simeq 75$ km/s due to relativistic asymmetric neutrino emission alone, with negligible contribution from matter ejecta.

4. CONCLUSIONS

We find that MAXI J1305-704 is only the second BH system that – when analyzed in the context of both its kinematic and binary/stellar history under the assumption that it was formed in the Galactic disk via isolated binary evolution, and assuming uninformative priors on the mass-loss and natal kick velocity imparted at SNrequires a natal kick to exist. Our inferred natal kick of 91^{+23}_{-12} km/s is consistent with Sánchez et al. (2021), who estimated from kinematics alone that the total center-ofmass velocity imparted to this system was 75^{+25}_{-12} km/s. It is also consistent with Atri et al. (2019), who found that the population of BH LMXB birth velocities is well fit by a normal distribution peaking around 107 km/s with a standard deviation of 16 km/s. Our result suggests that the birth velocity of J1305 could not have been provided by symmetric mass-loss alone, but requires an additional natal kick. Our 95% lower limit on this natal kick of 70 km/s is also similar to the findings of Fragos et al. (2009) for XTE J1118+40, who found that its BH received a kick of at least $\simeq 80$ km/s. With this analysis of J1305, we strengthen the case that at least some BHs receive natal kicks at birth.

We find that our results are robust against kinematic uncertainties. The inferred boost to the center of mass velocity from symmetric mass loss alone, $V_{\Delta M} = 0.7^{+1.0}_{-0.5}$ km/s, would be incompatible with even moderate birth peculiar velocities $\gtrsim 10$ km/s. We assess that our inference is not qualitatively affected by our choice of Milky Way potential nor the assumption that J1305 was born exactly in the Galactic mid-plane with a circular velocity. Further, as shown in Fragos et al. (2009) for a system with a very similar donor, the kick constraints are not sensitive to the assumed magnetic braking law. We do note that, as in all binary evolution calculations to date we assume that binaries instantly circularize upon the onset of mass transfer. This is not always a welljustified assumption and it leads to a mismatch between the donor ages produced in our population synthesis and in our individual MESA runs that matched the observed properties; it is possible that correcting for this effect

(Rocha et al. 2022, in prep.) may have a small quantitative effect on our kick constraints. Lastly, we note that at ~ 1 kpc above the Galactic mid-plane, we cannot exclude that J1305 formed dynamically before being ejected from a globular cluster, in which case our analysis is not applicable.

Through this and other past studies we have demonstrated that accounting for all observational characteristics and coupling binary evolution and kinematic modeling we can provide robust constraints on natal BH kicks and associated mass loss at formation. With an increasing sample of LMXBs with proper motion and radial velocity estimates, our goal in future work is to investigate potential statistical correlations between BH kicks and mass loss characteristics at birth, which may shed light on the natal kick mechanism.

ACKNOWLEDGEMENTS

We thank Christopher Berry, Lieke van Son, and Michael Zevin for helpful discussions and comments on the manuscript. CK is supported by the Riedel Family Fellowship. VK is grateful for support from a Guggenheim Fellowship, from CIFAR as a Senior Fellow, and from Northwestern University, including the Daniel I. Linzer Distinguished University Professorship fund. KR and DM thank the LSSTC Data Science Fellowship Program, which is funded by the LSST Corporation, NSF Cybertraining Grant No. 1829740, the Brinson Foundation, and the Gordon and Betty Moore Foundation; their participation in the program has benefited this work. ZD is grateful for support from the CIERA Board of Visitors Research Professorship. KK and EZ were partially supported by the Federal Commission for Scholarships for Foreign Students for the Swiss Government Excellence Scholarship (ESKAS No. 2021.0277 and ESKAS No. 2019.0091, respectively). ZX was supported by the Chinese Scholarship Council (CSC). This work was performed with the help of the computing resources at CIERA provided by the Quest high performance computing facility at Northwestern University – funded through NSF PHY-1726951 – which is jointly supported by the Office of the Provost, the Office for Research, and Northwestern University Information Technology.

Software: NumPy (van der Walt et al. 2011), SciPy (Virtanen et al. 2020), matplotlib (Hunter 2007), pandas (Wes McKinney 2010), POSYDON (Fragos et al. 2022)

REFERENCES

- Aghakhanloo, M., Smith, N., Andrews, J., et al. 2022, MNRAS, 516, 2142, doi: 10.1093/mnras/stac2265
- Atri, P., Miller-Jones, J. C. A., Bahramian, A., et al. 2019, MNRAS, 489, 3116, doi: 10.1093/mnras/stz2335
- Blaauw, A. 1961, BAN, 15, 265
- Boubert, D., & Evans, N. W. 2018, MNRAS, 477, 5261, doi: 10.1093/mnras/sty980
- Bovy, J. 2015, ApJS, 216, 29, doi: 10.1088/0067-0049/216/2/29
- Bray, J. C., & Eldridge, J. J. 2018, Monthly Notices of the Royal Astronomical Society, 480, 5657, doi: 10.1093/mnras/sty2230
- Carlberg, R. G., & Innanen, K. A. 1987, AJ, 94, 666, doi: 10.1086/114503
- Clark, G. W. 1975, ApJL, 199, L143, doi: 10.1086/181869
- Coleman, M. S. B., & Burrows, A. 2022, MNRAS, 517, 3938, doi: 10.1093/mnras/stac2573
- De Donder, E., Vanbeveren, D., & van Bever, J. 1997, A&A, 318, 812
- Dray, L. M., Dale, J. E., Beer, M. E., Napiwotzki, R., & King, A. R. 2005, MNRAS, 364, 59, doi: 10.1111/j.1365-2966.2005.09536.x

- Eldridge, J. J., Langer, N., & Tout, C. A. 2011, MNRAS, 414, 3501, doi: 10.1111/j.1365-2966.2011.18650.x
- Fragos, T., Willems, B., Kalogera, V., et al. 2009, Astrophysical Journal, 697, 1057, doi: 10.1088/0004-637X/697/2/1057
- Fragos, T., Andrews, J. J., Bavera, S. S., et al. 2022, arXiv e-prints, arXiv:2202.05892.

https://arxiv.org/abs/2202.05892

- Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2021, A&A, 649, A1, doi: 10.1051/0004-6361/202039657
- Giacobbo, N., & Mapelli, M. 2020, ApJ, 891, 141, doi: 10.3847/1538-4357/ab7335
- Greiner, J., Rau, A., & Schady, P. 2012, The Astronomer's Telegram, 4030, 1
- Hills, J. G. 1976, Monthly Notices of the Royal Astronomical Society, 175, 1P, doi: 10.1093/mnras/175.1.1P
- Hobbs, G., Lorimer, D. R., Lyne, A. G., & Kramer, M. 2005, Monthly Notices of the Royal Astronomical Society, 360, 974, doi: 10.1111/J.1365-2966.2005.09087.X
- Honma, M., Nagayama, T., Ando, K., et al. 2012, Publications of the Astronomical Society of Japan, 64, doi: 10.1093/pasj/64.6.136

- Hunter, J. D. 2007, Computing in Science & Engineering, 9, 90, doi: 10.1109/MCSE.2007.55
- Janka, H. T. 2012, Annual Review of Nuclear and Particle Science, 62, 407,
 - doi: 10.1146/ANNUREV-NUCL-102711-094901
- Jonker, P. G., Kaur, K., Stone, N., & Torres, M. A. P. 2021, The Astrophysical Journal, 921, 131, doi: 10.3847/1538-4357/ac2839
- Kalogera, V. 1996, ApJ, 471, 352, doi: 10.1086/177974
- Kapil, V., Mandel, I., Berti, E., & Müller, B. 2022. https://arxiv.org/abs/2209.09252
- Kennea, J. A., Yang, Y. J., Altamirano, D., et al. 2012, The Astronomer's Telegram, 4044, 1
- Kroupa, P. 2001, MNRAS, 322, 231, doi: 10.1046/j.1365-8711.2001.04022.x
- Kuijken, K., & Gilmore, G. 1989, MNRAS, 239, 571, doi: 10.1093/mnras/239.2.571
- Lyne, A. G., & Lorimer, D. R. 1994, Nature, 369, 127, doi: 10.1038/369127a0
- Matsuoka, M., Kawasaki, K., Ueno, S., et al. 2009, Publications of the Astronomical Society of Japan, 61, 999, doi: 10.1093/pasj/61.5.999
- Mihara, T., Nakajima, M., Sugizaki, M., et al. 2011, PASJ, 63, S623, doi: 10.1093/pasj/63.sp3.S623
- Mirabel, I. F., Dhawan, V., Mignani, R. P., Rodrigues, I., & Guglielmetti, F. 2001, Nature, 413, 139, doi: 10.1038/35093060
- Morihana, K., Sugizaki, M., Nakahira, S., et al. 2013, PASJ, 65, L10, doi: 10.1093/pasj/65.5.L10
- Paxton, B., Bildsten, L., Dotter, A., et al. 2010, The Astrophysical Journal Supplement Series, 192, 3, doi: 10.1088/0067-0049/192/1/3
- Paxton, B., Cantiello, M., Arras, P., et al. 2013, The Astrophysical Journal Supplement Series, 208, 4, doi: 10.1088/0067-0049/208/1/4
- Paxton, B., Marchant, P., Schwab, J., et al. 2015, The Astrophysical Journal Supplement Series, 220, 15, doi: 10.1088/0067-0049/220/1/15
- Paxton, B., Smolec, R., Schwab, J., et al. 2019, The Astrophysical Journal Supplement Series, 243, 10, doi: 10.3847/1538-4365/ab2241
- Postnov, K. A., & Yungelson, L. R. 2014, Living Reviews in Relativity, 17, 3, doi: 10.12942/lrr-2014-3
- Price-Whelan, A., Sipőcz, B., Lenz, D., et al. 2020, adrn/gala: v1.3, v1.3, Zenodo, doi: 10.5281/zenodo.4159870

- Price-Whelan, A. M. 2017, The Journal of Open Source Software, 2, doi: 10.21105/joss.00388
- Renzo, M., Zapartas, E., De Mink, S. E., et al. 2019, Astronomy and Astrophysics, 624, A66, doi: 10.1051/0004-6361/201833297
- Repetto, S., Davies, M. B., & Sigurdsson, S. 2012, Monthly Notices of the Royal Astronomical Society, 425, 2799, doi: 10.1111/j.1365-2966.2012.21549.x
- Sánchez, D. M., Rau, A., Hernández, A. A., et al. 2021, Mon. Not. Roy. Astron. Soc., 506, 581, doi: 10.1093/mnras/stab1714
- Sato, R., Serino, M., Nakahira, S., et al. 2012, The Astronomer's Telegram, 4024, 1
- Shklovskii, I. S. 1970, Soviet Ast., 13, 562
- Suwa, F., Negoro, H., Nakahira, S., et al. 2012, The Astronomer's Telegram, 4035, 1
- Tauris, T. M., et al. 2017, Astrophys. J., 846, 170, doi: 10.3847/1538-4357/aa7e89
- van der Walt, S., Colbert, S. C., & Varoquaux, G. 2011, Computing in Science & Engineering, 13, 22, doi: 10.1109/MCSE.2011.37
- van Rensbergen, W., Vanbeveren, D., & De Loore, C. 1996, A&A, 305, 825
- Virtanen, P., Gommers, R., Oliphant, T. E., et al. 2020, Nature Methods, 17, 261, doi: 10.1038/s41592-019-0686-2
- Wes McKinney. 2010, in Proceedings of the 9th Python in Science Conference, ed. Stéfan van der Walt & Jarrod Millman, 56 – 61, doi: 10.25080/Majora-92bf1922-00a
- Willems, B., Henninger, M., Levin, T., et al. 2005, The Astrophysical Journal, 625, 324, doi: 10.1086/429557
- Wong, T. W., Valsecchi, F., Ansari, A., et al. 2014,
 Astrophysical Journal, 790, 119,
 doi: 10.1088/0004-637X/790/2/119
- Wong, T. W., Valsecchi, F., Fragos, T., & Kalogera, V. 2012, Astrophysical Journal, 747, 111, doi: 10.1088/0004-637X/747/2/111
- Wong, T. W., Willems, B., & Kalogera, V. 2010, Astrophysical Journal, 721, 1689, doi: 10.1088/0004-637X/721/2/1689
- Wongwathanarat, A., Janka, H. T., & Müller, E. 2013, A&A, 552, A126, doi: 10.1051/0004-6361/201220636