

Hyperheavenly spaces and their application in Walker and para-Kähler geometries: part II

Adam Chudecki*

November 8, 2022

* Center of Mathematics and Physics, Lodz University of Technology,
Al. Politechniki 11, 90-924 Łódź, Poland, adam.chudecki@p.lodz.pl

Abstract.

4-dimensional spaces equipped with congruences of null strings are considered. It is assumed that a space admits a congruence of expanding self-dual null strings and its self-dual part of the Weyl tensor is algebraically degenerate. Different Petrov-Penrose types of such spaces are analyzed. A special attention is paid to para-Kähler Einstein spaces. All para-Kähler Einstein metrics of spaces with algebraically degenerate self-dual Weyl spinor are found in all the generality.

1 Introduction

This is the second part of the work devoted to para-Kähler and Walker geometries. In the first part [5] we considered complex and real neutral, 4-dimensional spaces equipped with a nonexpanding congruence¹ of self-dual (SD) null strings. These are spaces of the types $[\text{deg}]^n \otimes [\text{any}]$. This part is devoted to spaces with expanding SD \mathcal{C} , i.e., spaces of the types $[\text{deg}]^e \otimes [\text{any}]$.

[At this point an important remark is needed. In the present paper we use the spinorial formalism, hyperheavenly spaces (\mathcal{HH} -spaces) formalism and abbreviations introduced in Ref. [5]. We encourage the Reader of this paper to read Ref. [5] first. More detailed approach to the spinorial formalism can be found in [14, 15, 17]. A spinorial approach to Walker and sesquiWalker geometries was presented in [9, 11–13]. [4, 10] are fundamental references devoted to \mathcal{HH} -spaces. Congruences of null strings are analyzed in [7, 16, 18]. Applications of complex methods in real geometries, especially in para-Kähler geometry can be found in [1–3, 6, 8].]

If a space is equipped with a \mathcal{C}^n which is SD then the SD Weyl spinor is algebraically special. However, if SD \mathcal{C} is expanding then the SD Weyl spinor is not algebraically special in general. Thus, in what follows we assume two basic facts


- (i) a space admits an expanding congruence of null strings and the orientation is chosen in such a manner that this congruence is self-dual,

¹A congruence (a foliation) of null strings is abbreviated by \mathcal{C} . Also, an intersection of self-dual and anti-self-dual congruences of null strings which constitutes a congruence of null geodesics is abbreviated by \mathcal{I} . The list of all the abbreviations which we use in the present paper can be found in Ref. [5].

(ii) the self-dual part of the Weyl tensor is algebraically special.

In Einstein spaces (i) and (ii) are equivalent (it is a statement of the complex Goldberg-Sachs theorem, see [16]). However, in Sections 2.2 - 2.4, 3.1 and 4.1 we consider spaces with an arbitrary traceless Ricci tensor. For such spaces (i) and (ii) are not equivalent in general.

The paper is organized, as follows. In Section 2 the structure of weak (expanding) \mathcal{HH} -spaces is presented. Weak \mathcal{HH} -spaces are equipped with a single \mathcal{C} *ex definitione*. This \mathcal{C} is assumed to be expanding and it is a common property of all the spaces which metrics are presented in this paper. Then we follow the pattern of Ref. [5] (Scheme 1 shows a structure of the present paper and compares it to Ref. [5].).



Scheme 1: Diagram of weak \mathcal{HH} -spaces equipped with ASD congruences of null strings.

First we assume that a weak \mathcal{HH} -space admits also an anti-self-dual (ASD) \mathcal{C}^e (Section 2.2). The existence of two \mathcal{C} s of a different duality allows to introduce a sub-classification of spaces based on the properties of the intersection of \mathcal{C} s. Hence, there are 4 subtypes of spaces of the type $[\text{deg}]^e \otimes [\text{any}]^e$.

Then we assume that the ASD \mathcal{C} is nonexpanding (Section 2.3). There are 2 subtypes of such spaces. The next step is to equipped a space with the second ASD \mathcal{C} which is

expanding (Section 2.4). According to the results of Ref. [5] there are 7 possible subtypes of such spaces. Our considerations prove that one of these subtypes, in fact, cannot exist². The metrics presented in Sections 2.2 - 2.4 are not analyzed in details because their geometrical interpretation is not clear yet. It is only worth to mention that all of the spaces which are equipped with ASD \mathcal{C}^n are Walker spaces.

In Sections 3 and 4 the second ASD \mathcal{C} becomes nonexpanding. Thus, spaces are equipped with ASD \mathcal{C}^{nn} and they are para-Kähler. There are two different subtypes of such spaces, namely $\{[\text{deg}]^e \otimes [D]^{nn}, [++, --]\}$ and $\{[\text{deg}]^e \otimes [D]^{nn}, [++, ++]\}$.

Section 3 is devoted to the types $\{[\text{deg}]^e \otimes [D]^{nn}, [++, --]\}$. These types have been completely solved (Theorem 3.1). The metric (3.10) is a general metric of spaces of the types $\{[\text{deg}]^e \otimes [D]^{nn}, [++, --]\}$. Thorough analysis of such spaces via Petrov-Penrose classification is presented. We consider also para-Kähler Einstein spaces of the types $\{[\text{deg}]^e \otimes [D]^{nn}, [++, --]\}$ (Theorem 3.4, the metric (3.31)).

The types $\{[\text{deg}]^e \otimes [D]^{nn}, [++, ++]\}$ (Section 4) are much more complicated. We found two forms of the metric of such spaces (Theorem 4.1, metrics (4.27) and (4.28)). The geometrical difference between metrics (4.27) and (4.28) remains unknown. We are not sure if they are essentially different or the metric (4.28) is just a special case of (4.27). Thus, we do not analyze these metrics in details. They will be analyzed elsewhere.

In Section 4.2 we focus on para-Kähler Einstein spaces of the types $\{[\text{deg}]^e \otimes [D]^{nn}, [++, ++]\}$ and we arrive at the main result³ of the paper. It is Theorem 4.2 and the metric (4.43) which is a general metric of the para-Kähler Einstein spaces for which SD Weyl tensor is algebraically special. The metric (4.43) depends on 4 functions of two variables like it was proved in [3]. The types $\{[\text{III}]^e \otimes [D]^{nn}, [++, ++]\}$ and $\{[N]^e \otimes [D]^{nn}, [++, ++]\}$ have been also found and they depend on 3 and 2 functions of two variables, respectively⁴.

Concluding remarks end the paper.

Considerations are purely local. We consider complex manifolds of dimension four equipped with a holomorphic metric. Thus, coordinates are complex and functions are holomorphic. If we replace complex coordinates by real ones and holomorphic functions by real smooth ones, we arrive at 4-dimensional real neutral geometries. Hence, all metrics presented in this paper have neutral slices.

²This unexpected fact gives an answer to one of the questions formulated in *Concluding Remarks* of Ref. [5]. Namely, we asked if there are any geometrical obstacles which prevent the existence of certain subtypes listed in Tables 9-11 of Ref. [5]. The answer to this question is positive. In fact, the type $\{[\text{deg}]^e \otimes [\text{deg}]^{ne}, [++, --]\}$ does not exist.

³The question which of the results presented in this paper is the "main" result is a moot point. According to our best knowledge almost all the metrics which we found are new results (see the summary Tables 1 and 2). However, it seems that pKE metrics have the most transparent geometrical meaning and via their relation with twistor distributions [3] are physically relevant. Thus, we think that the most important results of the paper are Theorems 3.4 and 4.2. Theorem 4.2 as a more general than Theorem 3.4 deserves to be called "the main result" of the paper.

⁴We skip an analysis of the types $[D]^{ee} \otimes [D]^{nn}$. All such types were found in [3].

2 Weak expanding hyperheavenly spaces

2.1 Spaces of the types $[\text{deg}]^e \otimes [\text{any}]$

2.1.1 Structure of weak expanding hyperheavenly spaces

In this Section we gather the basic definitions and notations concerning weak hyperheavenly spaces. This Section resembles Section 2.6 of Ref. [5] but in [5] all the formulas were adapted to *weak nonexpanding hyperheavenly spaces* (types $[\text{deg}]^n \otimes [\text{any}]$). In what follows we focus on more general *weak expanding hyperheavenly spaces*, i.e., spaces of the types $[\text{deg}]^e \otimes [\text{any}]$.

Definition 2.1. *Weak hyperheavenly space (weak \mathcal{HH} -space) is a pair (\mathcal{M}, ds^2) where \mathcal{M} is a 4-dimensional complex analytic differential manifold and ds^2 is a holomorphic metric, satisfying the following conditions*

- (i) *there exists a 2-dimensional holomorphic totally null self-dual integrable distribution given by the Pfaff system*

$$m_A g^{A\dot{B}} = 0, \quad m_A \neq 0 \quad (2.1)$$

- (ii) *the SD Weyl spinor C_{ABCD} is algebraically degenerate and m_A is a multiple Penrose spinor i.e.*

$$C_{ABCD} m^A m^B m^C = 0 \quad (2.2)$$

Definition 2.1 appeared in [9] for the first time. It was also proved in [9] that the metric of a weak \mathcal{HH} -space can be brought to the form

$$\frac{1}{2} ds^2 = \phi^{-2} (-dp^{\dot{A}} dq_{\dot{A}} + Q^{\dot{A}\dot{B}} dq_{\dot{A}} dq_{\dot{B}}) = e^1 e^2 + e^3 e^4 \quad (2.3)$$

where $(q_{\dot{A}}, p_{\dot{B}})$ are local complex coordinates. Coordinates $q_{\dot{A}}$ label the null string while $p_{\dot{A}}$ are coordinates on leafs of \mathcal{C}_{m^A} . ϕ and $Q^{\dot{A}\dot{B}} = Q^{(\dot{A}\dot{B})}$ are holomorphic functions of variables $(q^{\dot{A}}, p^{\dot{B}})$. The null tetrad (e^1, e^2, e^3, e^4) defined as follows

$$\begin{aligned} [e^3, e^1] &= \phi^{-2} dq_{\dot{A}} \\ [e^4, e^2] &= -dp^{\dot{A}} + Q^{\dot{A}\dot{B}} dq_{\dot{B}} \end{aligned} \quad (2.4)$$

is called *Plebański tetrad*. The dual basis is given by

$$-\partial_{\dot{A}} = [\partial_4, \partial_2], \quad \bar{\partial}^{\dot{A}} = [\partial_3, \partial_1], \quad \implies \quad \partial_{A\dot{B}} = \sqrt{2} [\partial_{\dot{B}}, \bar{\partial}_{\dot{B}}] \quad (2.5)$$

where

$$\begin{aligned} \partial_{\dot{A}} &:= \frac{\partial}{\partial p^{\dot{A}}}, \quad \bar{\partial}_{\dot{A}} := \phi^2 \left(\frac{\partial}{\partial q^{\dot{A}}} - Q_{\dot{A}}^{\quad \dot{B}} \partial_{\dot{B}} \right) \\ \partial^{\dot{A}} &:= \frac{\partial}{\partial p_{\dot{A}}}, \quad \bar{\partial}^{\dot{A}} := \phi^2 \left(\frac{\partial}{\partial q_{\dot{A}}} + Q^{\dot{A}\dot{B}} \partial_{\dot{B}} \right) \end{aligned} \quad (2.6)$$

From the first structure equations we find the spinorial connection coefficients

$$\begin{aligned}
\Gamma_{111\dot{D}} &= 0 & \Gamma_{112\dot{D}} &= -\sqrt{2}\phi^{-1}J_{\dot{D}} \\
\Gamma_{121\dot{D}} &= \frac{3}{\sqrt{2}}\phi^{-1}J_{\dot{D}} & \Gamma_{122\dot{D}} &= -\frac{\phi}{\sqrt{2}}\left(\partial^{\dot{A}}(\phi Q_{\dot{A}\dot{D}}) - \frac{\partial\phi}{\partial q^{\dot{D}}}\right) \\
\Gamma_{221\dot{D}} &= \sqrt{2}\phi\left(Q_{\dot{D}\dot{A}}J^{\dot{A}} + \frac{\partial\phi}{\partial q^{\dot{D}}}\right) & \Gamma_{222\dot{D}} &= -\sqrt{2}\phi^2\eth^{\dot{A}}Q_{\dot{A}\dot{D}} \\
\Gamma_{\dot{A}\dot{B}1\dot{D}} &= -\sqrt{2}\phi^{-1}J_{(\dot{A}}\epsilon_{\dot{B})\dot{D}} \\
\Gamma_{\dot{A}\dot{B}2\dot{D}} &= \sqrt{2}\phi\left(\phi\partial_{(\dot{A}}Q_{\dot{B})\dot{D}} + \epsilon_{\dot{D}(\dot{B}}\left(Q_{\dot{A})\dot{C}}J^{\dot{C}} + \frac{\partial\phi}{\partial q^{\dot{A}}}\right)\right)
\end{aligned} \tag{2.7}$$

where

$$J_{\dot{A}} := \partial_{\dot{A}}\phi \equiv \frac{\partial\phi}{\partial p^{\dot{A}}} \implies J^{\dot{A}} = -\partial^{\dot{A}}\phi \equiv -\frac{\partial\phi}{\partial p_{\dot{A}}} \tag{2.8}$$

Nonzero SD curvature coefficients $C^{(i)}$ and the ASD Weyl spinor read

$$\begin{aligned}
C^{(5)} &= 0 \\
C^{(4)} &= 0 \\
C^{(3)} &= -\frac{1}{3}\phi^2\partial_{\dot{A}}\partial_{\dot{B}}Q^{\dot{A}\dot{B}} \\
\frac{1}{2}C^{(2)} &= -\frac{1}{2}\phi^5\partial^{\dot{A}}(\phi^{-3}\eth^{\dot{B}}Q_{\dot{A}\dot{B}}) - \frac{1}{2}\phi^2\eth_{\dot{A}}(\phi^{-3}\eth^{\dot{A}}\phi) \\
\frac{1}{2}C^{(1)} &= -\phi^4\eth^{\dot{A}}(\phi^{-2}\eth^{\dot{B}}Q_{\dot{A}\dot{B}}) + \phi^4(\eth^{\dot{A}}Q_{\dot{A}\dot{B}})(\partial_{\dot{C}}Q^{\dot{B}\dot{C}}) \\
C_{\dot{A}\dot{B}\dot{C}\dot{D}} &= -\phi^2\partial_{(\dot{A}}\partial_{\dot{B}}Q_{\dot{C}\dot{D}})
\end{aligned} \tag{2.9}$$

Curvature scalar \mathcal{R} and the spinorial image of the traceless Ricci tensor $C_{\dot{A}\dot{B}\dot{C}\dot{D}}$ have the form

$$\begin{aligned}
-\frac{1}{2}\mathcal{R} &= \phi^2\partial_{\dot{A}}\partial_{\dot{B}}Q^{\dot{A}\dot{B}} - 6\phi^3\partial_{\dot{A}}(\phi^{-4}\eth^{\dot{A}}\phi) \\
C_{11\dot{A}\dot{B}} &= -\phi^{-1}\partial_{\dot{A}}\partial_{\dot{B}}\phi \\
C_{12\dot{A}\dot{B}} &= -\frac{1}{2}\phi^2\partial_{(\dot{A}}\partial^{\dot{C}}Q_{\dot{B})\dot{C}} - \phi\partial_{(\dot{A}}(\phi^{-2}\eth_{\dot{B}})\phi) \\
C_{22\dot{A}\dot{B}} &= -\phi^5\partial_{(\dot{A}}\phi^{-3}\eth^{\dot{C}}Q_{\dot{B})\dot{C}} + \phi(\eth^{\dot{C}}\phi)(\partial_{\dot{C}}Q_{\dot{A}\dot{B}}) - \phi\eth_{(\dot{A}}(\phi^{-2}\eth_{\dot{B}})\phi)
\end{aligned} \tag{2.10}$$

The SD congruence of null strings $\mathcal{C}_{m^A}^e$ is generated by the spinor $m_A \sim [0, m]$, $m \neq 0$, what implies that it is spanned by the vectors (∂_2, ∂_4) . In this sense Plebański tetrad is *adapted* to $\mathcal{C}_{m^A}^e$. The expansion of $\mathcal{C}_{m^A}^e$ reads

$$M_{\dot{B}} = -\sqrt{2}m\phi^{-1}J_{\dot{B}} \neq 0 \tag{2.11}$$

In the rest of the paper we use the abbreviations

$$q^{\dot{1}} =: q, \quad q^{\dot{2}} =: p, \quad p^{\dot{1}} =: x, \quad p^{\dot{2}} =: y, \quad Q^{\dot{A}\dot{B}} =: \begin{bmatrix} \mathcal{A} & \mathcal{Q} \\ \mathcal{Q} & \mathcal{B} \end{bmatrix} \tag{2.12}$$

with these abbreviations the metric (2.3) takes the form

$$\frac{1}{2}ds^2 = \phi^{-2}(dqdy - dpdx + \mathcal{A}dp^2 - 2\mathcal{Q}dqdp + \mathcal{B}dq^2) \tag{2.13}$$

In what follows the coordinate system (q, p, x, y) is called *the hyperheavenly coordinate system*.

Note, that the metric (2.13) depends on four functions of four variables.

2.1.2 Gauge transformations

The metric (2.3) remains invariant under the following transformations of the coordinates

$$q'^{\dot{A}} = q'^{\dot{A}}(q^{\dot{M}}), \quad p'^{\dot{A}} = \lambda^{-1} D^{-1}_{\dot{B}}{}^{\dot{A}} p^{\dot{B}} + \sigma^{\dot{A}} \quad (2.14)$$

where $\lambda = \lambda(q_{\dot{B}})$ and $\sigma^{\dot{A}} = \sigma^{\dot{A}}(q_{\dot{B}})$ are arbitrary gauge functions and

$$\begin{aligned} D_{\dot{A}}{}^{\dot{B}} &:= \frac{\partial q'^{\dot{A}}}{\partial q_{\dot{B}}} = \Delta \frac{\partial q^{\dot{B}}}{\partial q'^{\dot{A}}} = \lambda \Delta \frac{\partial p'^{\dot{A}}}{\partial p_{\dot{B}}} = \lambda^{-1} \frac{\partial p^{\dot{B}}}{\partial p'^{\dot{A}}}, \quad \Delta := \det \left(\frac{\partial q'^{\dot{A}}}{\partial q_{\dot{B}}} \right) = \frac{1}{2} D_{\dot{A}\dot{B}} D^{\dot{A}\dot{B}} \\ D^{-1}_{\dot{A}}{}^{\dot{B}} &= \frac{\partial q_{\dot{A}}}{\partial q'^{\dot{B}}} = \Delta^{-1} \frac{\partial q'^{\dot{B}}}{\partial q_{\dot{A}}} = \lambda \frac{\partial p'^{\dot{B}}}{\partial p_{\dot{A}}} = \lambda^{-1} \Delta^{-1} \frac{\partial p_{\dot{A}}}{\partial p'^{\dot{B}}} \end{aligned} \quad (2.15)$$

Hence

$$\begin{aligned} D^{\dot{A}}{}_{\dot{B}} &:= D_{\dot{M}}{}^{\dot{N}} \in^{\dot{M}\dot{A}} \in_{\dot{B}\dot{N}} = -\Delta D^{-1}_{\dot{B}}{}^{\dot{A}} \\ D^{-1}_{\dot{B}}{}^{\dot{A}} &:= D^{-1}_{\dot{M}}{}^{\dot{N}} \in^{\dot{M}\dot{A}} \in_{\dot{B}\dot{N}} = -\frac{1}{\Delta} D_{\dot{B}}{}^{\dot{A}} \end{aligned} \quad (2.16)$$

Functions ϕ and $Q^{\dot{A}\dot{B}}$ transform under (2.14) as follows

$$\begin{aligned} \phi' &= \lambda^{-\frac{1}{2}} \phi \\ Q'^{\dot{A}\dot{B}} &= \lambda^{-1} D^{-1}_{\dot{R}}{}^{\dot{A}} D^{-1}_{\dot{S}}{}^{\dot{B}} Q^{\dot{R}\dot{S}} + D^{-1}_{\dot{R}}{}^{(\dot{A}} \frac{\partial p'^{\dot{B})}}{\partial q_{\dot{R}}} \end{aligned} \quad (2.17)$$

Transformations (2.14) are equivalent to the spinorial transformations

$$\begin{aligned} L^{\dot{A}}{}_{\dot{B}} &= \begin{bmatrix} \lambda^{-1} \Delta^{-\frac{1}{2}} & h \phi^2 \Delta^{\frac{1}{2}} \\ 0 & \lambda \Delta^{\frac{1}{2}} \end{bmatrix}, \quad 2h := \frac{\partial \sigma^{\dot{R}}}{\partial q'^{\dot{R}}} - \frac{1}{\lambda^2 \Delta} \frac{\partial \lambda}{\partial q^{\dot{R}}} p^{\dot{R}} \\ M^{\dot{A}}{}_{\dot{B}} &= \Delta^{\frac{1}{2}} D^{-1}_{\dot{B}}{}^{\dot{A}} \end{aligned} \quad (2.18)$$

where $L^{\dot{A}}{}_{\dot{B}}, M^{\dot{A}}{}_{\dot{B}} \in SL(2, \mathbb{C})$. Dotted and undotted spinors transform as follows

$$m'^{A_1 \dots A_n} = L^{A_1}_{R_1} \dots L^{A_n}_{R_n} m^{R_1 \dots R_n}, \quad m'^{\dot{A}_1 \dots \dot{A}_n} = M^{\dot{A}_1}_{\dot{R}_1} \dots M^{\dot{A}_n}_{\dot{R}_n} m^{\dot{R}_1 \dots \dot{R}_n} \quad (2.19)$$

2.1.3 ASD null string equations

Assume that a weak \mathcal{HH} -space admits an additional structure which is ASD \mathcal{C}^e . Let this congruence be expanding in general and let it be generated by a spinor $m_{\dot{A}}$ with an expansion given by M_A . ASD null string equations read

$$m_{\dot{A}} M_A = m^{\dot{B}} \nabla_{A\dot{A}} m_{\dot{B}} \equiv m^{\dot{B}} \partial_{A\dot{A}} m_{\dot{B}} + m^{\dot{S}} m^{\dot{B}} \Gamma_{\dot{S}\dot{B}\dot{A}\dot{A}} \quad (2.20)$$

Explicitly, Eqs. (2.20) take the form

$$\frac{\phi}{\sqrt{2}}m_{\dot{A}}M_1 = \phi m^{\dot{B}}\frac{\partial m_{\dot{B}}}{\partial p^{\dot{A}}} + m_{\dot{A}}m^{\dot{S}}J_{\dot{S}} \quad (2.21a)$$

$$\begin{aligned} \frac{1}{\sqrt{2}\phi}m_{\dot{A}}M_2 &= \phi m^{\dot{B}}\frac{\partial m_{\dot{B}}}{\partial q^{\dot{A}}} + \phi m^{\dot{S}}\frac{\partial}{\partial p^{\dot{S}}}(m^{\dot{B}}Q_{\dot{A}\dot{B}}) - \phi m^{\dot{B}}Q_{\dot{A}\dot{B}}\frac{\partial m^{\dot{C}}}{\partial p^{\dot{C}}} \\ &\quad + m_{\dot{A}}m^{\dot{B}}Q_{\dot{C}\dot{B}}J^{\dot{C}} + m_{\dot{A}}m^{\dot{S}}\frac{\partial \phi}{\partial q^{\dot{S}}} \end{aligned} \quad (2.21b)$$

Eqs. (2.21) contracted with $m^{\dot{A}}$ give ASD null string equations *sensu stricto*

$$m^{\dot{A}}m^{\dot{B}}\frac{\partial m_{\dot{A}}}{\partial p^{\dot{B}}} = 0 \quad (2.22a)$$

$$m^{\dot{A}}m^{\dot{B}}\frac{\partial m_{\dot{A}}}{\partial q^{\dot{B}}} + m^{\dot{S}}\frac{\partial \mathcal{Y}}{\partial p^{\dot{S}}} - \mathcal{Y}\left(2\frac{\partial m^{\dot{S}}}{\partial p^{\dot{S}}} + \frac{1}{\sqrt{2}}M_1 - \frac{1}{\phi}J_{\dot{S}}m^{\dot{S}}\right) = 0 \quad (2.22b)$$

$$\mathcal{Y} := m^{\dot{A}}m^{\dot{B}}Q_{\dot{A}\dot{B}}$$

ASD null string equations (2.20) remain invariant if a spinor $m_{\dot{A}}$ which generates \mathcal{C} is re-scaled. Hence, $\mathcal{C}_{m^{\dot{A}}}$ and $\mathcal{C}_{m'^{\dot{A}}}$ are the same congruences if $m^{\dot{A}} \sim m'^{\dot{A}}$. Consequently, the spinor $m_{\dot{A}}$ can be always re-scaled to the form with $m_1 = 1$ or $m_2 = 1$ so it has only one component to be determined via Eq. (2.22a). Solution of Eq. (2.22b) gives \mathcal{Y} . Having $m_{\dot{A}}$ and \mathcal{Y} one returns to Eqs. (2.21) and calculates the expansion M_A .

An expansion θ and a twist ϱ of $\mathcal{I}(\mathcal{C}_{m^{\dot{A}}}^e, \mathcal{C}_{m^{\dot{A}}})$ read

$$\theta \sim mM_1 + \frac{\sqrt{2}m}{\phi}m_{\dot{A}}\frac{\partial \phi}{\partial p_{\dot{A}}}, \quad \varrho \sim mM_1 - \frac{\sqrt{2}m}{\phi}m_{\dot{A}}\frac{\partial \phi}{\partial p_{\dot{A}}} \quad (2.23)$$

In the next Sections we follow the pattern of Ref. [5]. We equip a weak \mathcal{HH} -space with "the first" ASD \mathcal{C} generated by a spinor $m_{\dot{A}} \sim [z, 1]$, $z = z(q, p, x, y)$, with an expansion $M_A \neq 0$ (this congruence is denoted by $\mathcal{C}_{m^{\dot{A}}}$, Section 2.2). Then we consider a case with $M_A = 0$ (Section 2.3). In Section 2.4 a space is equipped with "the second" \mathcal{C} generated by a spinor $n_{\dot{A}} \sim [1, w]$, $w = w(q, p, x, y)$, with an expansion $N_A \neq 0$ (we denote this congruence by $\mathcal{C}_{n^{\dot{A}}}$). Of course, $\mathcal{C}_{m^{\dot{A}}}$ and $\mathcal{C}_{n^{\dot{A}}}$ are distinct so $m^{\dot{A}}n_{\dot{A}} \sim zw - 1 \neq 0$. Finally we arrive at a case with ASD \mathcal{C}^{nn} (Sections 3 and 4).

We believe that Scheme 1 clearly presents the structure of our article.

2.1.4 ASD congruence $\mathcal{C}_{m^{\dot{A}}}^e$

Inserting $m_{\dot{A}} \sim [z, 1]$ into Eqs. (2.22), (2.21) and (2.23) one finds

$$z_x - zz_y = 0 \quad (2.24a)$$

$$z_q - zz_p - z_y\mathcal{Z} + z\frac{\partial \mathcal{Z}}{\partial y} - \frac{\partial \mathcal{Z}}{\partial x} = 0, \quad \mathcal{Z} := \mathcal{B} + 2z\mathcal{Q} + z^2\mathcal{A} \quad (2.24b)$$

Expansion of $\mathcal{C}_{m^{\dot{A}}}^e$ reads

$$\frac{\phi}{\sqrt{2}}M_1 = -\phi z_y + z\phi_y - \phi_x \quad (2.25a)$$

$$\begin{aligned} \frac{1}{\sqrt{2}\phi}M_2 &= z\phi_p - \phi z_p - \phi_q + \phi z\frac{\partial}{\partial y}(\mathcal{Q} + z\mathcal{A}) - \phi\frac{\partial}{\partial x}(\mathcal{Q} + z\mathcal{A}) \\ &\quad + (\mathcal{Q} + z\mathcal{A})(\phi_x - \phi z_y) + (\mathcal{B} + z\mathcal{Q})\phi_y \end{aligned} \quad (2.25b)$$

Optical scalars are given by the formulas

$$\mathcal{I}(\mathcal{C}_{m^A}^e, \mathcal{C}_{m^{\dot{A}}}^e) : \quad \theta_1 \sim \phi z_y - 2z\phi_y + 2\phi_x, \quad \varrho_1 \sim z_y \quad (2.26)$$

Transformation rule for z follows from (2.19) and it yields

$$z' = \frac{p'_q - p'_p z}{q'_p z - q'_q} \quad (2.27)$$

Lemma 2.1. *Let $\mathcal{I}(\mathcal{C}_{m^A}^e, \mathcal{C}_{m^{\dot{A}}}^e)$ be nontwisting, $\varrho_1 = 0$. Then $z = 0$ and the following equations hold true*

$$\begin{aligned} \mathcal{B}_x &= 0, \quad \theta_1 \sim \phi_x, \\ M_1 &\sim \phi_x, \quad M_2 \sim -\phi \mathcal{Q}_x + \phi_x \mathcal{Q} + \phi_y \mathcal{B} - \phi_q \end{aligned} \quad (2.28)$$

Proof. If $\varrho_1 = 0$ then $z_y = 0$ and from (2.24a) it follows that $z_x = 0$. Hence, $z = z(p, q)$. Finally z can be gauged away without any loss of generality (compare (2.27)). With $z = 0$, Eqs. (2.24), (2.25) and (2.26) reduce to (2.28). ■

Remark 2.1. If $z = 0$ then the gauge (2.14) is restricted to the condition $p'_q = 0$.

Lemma 2.2. *Let $\mathcal{I}(\mathcal{C}_{m^A}^e, \mathcal{C}_{m^{\dot{A}}}^e)$ be twisting, $\varrho_1 \neq 0$. Then*

$$y = -xz + \Sigma(q, p, z) \quad (2.29)$$

where $\Sigma = \Sigma(q, p, z)$ is an arbitrary function. Moreover

$$\mathcal{B} + 2z\mathcal{Q} + z^2\mathcal{A} = \Omega(x - \Sigma_z) + z\Sigma_p - \Sigma_q, \quad \Omega = \Omega(q, p, z) \quad (2.30)$$

and

$$\theta_1 \sim \phi - 2\phi_x(x - \Sigma_z), \quad (2.31a)$$

$$M_1 \sim \phi - \phi_x(x - \Sigma_z), \quad (2.31b)$$

$$M_2 \sim z\phi_p + \frac{\phi\Sigma_p}{\Sigma_z - x} - \phi_q - \phi \frac{\partial}{\partial x}(\mathcal{Q} + z\mathcal{A}) + (\mathcal{Q} + z\mathcal{A}) \left(\phi_x - \frac{\phi}{\Sigma_z - x} \right) - \Omega\phi_z \quad (2.31c)$$

where $\Omega = \Omega(q, p, z)$ is an arbitrary function.

Proof. Multiplying (2.24a) by $dx \wedge dy \wedge dq \wedge dp$ and treating y as a function of (q, p, x, z) one arrives at (2.29). Hence

$$z_x = \frac{z}{\Sigma_z - x}, \quad z_y = \frac{1}{\Sigma_z - x}, \quad z_q = -\frac{\Sigma_q}{\Sigma_z - x}, \quad z_p = -\frac{\Sigma_p}{\Sigma_z - x} \quad (2.32)$$

Eq. (2.29) suggests that z should be treated as a new variable. Thus we pass to a new coordinate system (q, p, x, z) . Denote

$$\tilde{\mathcal{Z}} = \tilde{\mathcal{Z}}(q, p, x, z) = \tilde{\mathcal{Z}}(q, p, x, y(q, p, x, z)) = \mathcal{Z}(q, p, x, y) \quad (2.33)$$

then the derivatives of \mathcal{Z} transform as follows

$$\mathcal{Z}_y = \tilde{\mathcal{Z}}_z z_y, \quad \mathcal{Z}_x = \tilde{\mathcal{Z}}_x + \tilde{\mathcal{Z}}_z z_x, \quad \mathcal{Z}_q = \tilde{\mathcal{Z}}_q + \tilde{\mathcal{Z}}_z z_q, \quad \mathcal{Z}_p = \tilde{\mathcal{Z}}_p + \tilde{\mathcal{Z}}_z z_p \quad (2.34)$$

Eq. (2.24b) takes the form

$$\Sigma_q - z\Sigma_p + \tilde{\mathcal{Z}} + (\Sigma_z - x)\tilde{\mathcal{Z}}_x = 0 \quad (2.35)$$

which is solved by (2.30). Analogously one finds (2.31). ■

Remark 2.2. In (2.30) and (2.31) functions ϕ , \mathcal{A} , \mathcal{Q} and \mathcal{B} are understood as functions of (q, p, x, z) .

Considerations of this section should be supplemented by the transformation formula for Σ

$$\Sigma' = -\frac{\lambda^{-1}}{q'_p z - q'_q} \Sigma + \frac{p'_q - p'_p z}{q'_p z - q'_q} \sigma^{\dot{1}} + \sigma^{\dot{2}} \quad (2.36)$$

Transformation for Ω is much more complicated and its explicit form is not necessary at this stage of considerations.

2.1.5 ASD congruence $\mathcal{C}_{n\dot{A}}^e$

"The second" ASD congruence denoted by $\mathcal{C}_{n\dot{A}}^e$ is generated by a spinor $n_{\dot{A}}$ such that $n_{\dot{A}} \sim [1, w]$. ASD null string equations for $\mathcal{C}_{n\dot{A}}^e$ have the form (2.22) and (2.21) with $m_{\dot{A}} \rightarrow n_{\dot{A}}$ and $M_A \rightarrow N_A$. Hence

$$w_y - w w_x = 0 \quad (2.37a)$$

$$w_p - w w_q + w_x \mathcal{W} - w \frac{\partial \mathcal{W}}{\partial x} + \frac{\partial \mathcal{W}}{\partial y} = 0, \quad \mathcal{W} := \mathcal{A} + 2w \mathcal{Q} + w^2 \mathcal{B} \quad (2.37b)$$

Expansion of $\mathcal{C}_{n\dot{A}}^e$ reads

$$\frac{\phi}{\sqrt{2}} N_1 = \phi w_x - w \phi_x + \phi_y \quad (2.38a)$$

$$\begin{aligned} \frac{1}{\sqrt{2}\phi} N_2 &= \phi w_q - w \phi_q + \phi_p + \phi w \frac{\partial}{\partial x} (\mathcal{Q} + w \mathcal{B}) - \phi \frac{\partial}{\partial y} (\mathcal{Q} + w \mathcal{B}) \\ &\quad (\phi_y - \phi w_x)(\mathcal{Q} + w \mathcal{B}) + (\mathcal{A} + w \mathcal{Q}) \phi_x \end{aligned} \quad (2.38b)$$

Optical scalars of $\mathcal{I}(\mathcal{C}_{m\dot{A}}^e, \mathcal{C}_{n\dot{A}}^e)$ take the form

$$\mathcal{I}(\mathcal{C}_{m\dot{A}}^e, \mathcal{C}_{n\dot{A}}^e) : \quad \theta_2 \sim \phi w_x - 2w \phi_x + 2\phi_y, \quad \varrho_2 \sim w_x \quad (2.39)$$

and, finally, the transformation rule for the function w reads

$$w' = \frac{w q'_q - q'_p}{p'_p - p'_q w} \quad (2.40)$$

Lemma 2.3. *Let $\mathcal{I}(\mathcal{C}_{m\dot{A}}^e, \mathcal{C}_{n\dot{A}}^e)$ be nontwisting, $\varrho_2 = 0$. Then $w = 0$ and the following equations hold true*

$$\begin{aligned} \mathcal{A}_y &= 0, \quad \theta_2 \sim \phi_y, \\ N_1 &\sim \phi_y, \quad N_2 \sim -\phi \mathcal{Q}_y + \phi_y \mathcal{Q} + \phi_x \mathcal{A} + \phi_p \end{aligned} \quad (2.41)$$

Proof. We skip the proof due to its similarity to that of Lemma 2.1. ■

Remark 2.3. If $w = 0$ then the gauge (2.14) is restricted to the condition $q'_p = 0$.

Lemma 2.4. *Let $\mathcal{I}(\mathcal{C}_{m\dot{A}}^e, \mathcal{C}_{n\dot{A}}^e)$ be twisting, $\varrho_2 \neq 0$. Then*

$$x = -wy + H(q, p, w) \quad (2.42)$$

where $H = H(q, p, w)$ is an arbitrary function. Moreover

$$\mathcal{A} + 2w\mathcal{Q} + w^2\mathcal{B} = R(y - H_w) + H_p - wH_q, \quad R = R(q, p, w) \quad (2.43)$$

and

$$\theta_2 \sim \phi - 2\phi_y(y - H_w), \quad (2.44a)$$

$$N_1 \sim \phi - \phi_y(y - H_w), \quad (2.44b)$$

$$N_2 \sim \phi_p - w\phi_q - \frac{\phi H_q}{H_w - y} - \phi \frac{\partial}{\partial y}(\mathcal{Q} + w\mathcal{B}) + (\mathcal{Q} + w\mathcal{B}) \left(\phi_y - \frac{\phi}{H_w - y} \right) - R\phi_w \quad (2.44c)$$

where $R = R(q, p, w)$ is an arbitrary function.

Proof. We skip the proof due to its similarity to that of Lemma 2.2. ■

Remark 2.4. In (2.43) and (2.44) functions ϕ , \mathcal{A} , \mathcal{Q} and \mathcal{B} are understood as functions of (q, p, w, y) .

Note that

$$w_x = \frac{1}{H_w - y}, \quad w_y = \frac{w}{H_w - y}, \quad w_q = -\frac{H_q}{H_w - y}, \quad w_p = -\frac{H_p}{H_w - y} \quad (2.45)$$

and the transformation formula for H reads

$$H' = \frac{\lambda^{-1}}{p'_p - p'_q w} H + \frac{q'_q w - q'_p}{p'_p - p'_q w} \sigma^2 + \sigma^1 \quad (2.46)$$

The transformation formula for R is more advanced and we do not need it at this stage.

2.2 Spaces of the types $[\text{deg}]^e \otimes [\text{any}]^e$

In this Section we assume that a weak \mathcal{HH} -space admits $\mathcal{C}_{m\dot{A}}^e$.

2.2.1 Spaces of the types $\{[\text{deg}]^e \otimes [\text{any}]^e, [+ -]\}$ and $\{[\text{deg}]^e \otimes [\text{any}]^e, [- -]\}$

Theorem 2.1. *Let (\mathcal{M}, ds^2) be a complex (neutral) space of the type $\{[\text{deg}]^e \otimes [\text{any}]^e, [- -]\}$ or $\{[\text{deg}]^e \otimes [\text{any}]^e, [+ -]\}$. Then there exists a local coordinate system (q, p, x, y) such that the metric takes the form*

$$\frac{1}{2}ds^2 = \phi^{-2} (dqdy - dpdx + \mathcal{A}dp^2 - 2\mathcal{Q}dqdp + \mathcal{B}dq^2) \quad (2.47)$$

where $\phi = \phi(q, p, x, y)$, $\mathcal{A} = \mathcal{A}(q, p, x, y)$, $\mathcal{Q} = \mathcal{Q}(q, p, x, y)$, $\mathcal{B} = \mathcal{B}(q, p, y)$ are arbitrary holomorphic (real smooth) functions such that

$$\text{for the type } \{[\text{deg}]^e \otimes [\text{any}]^e, [- -]\}: \quad \phi_x = 0, \quad \phi\mathcal{Q}_x \neq \phi_y\mathcal{B} - \phi_q \quad (2.48a)$$

$$\text{for the type } \{[\text{deg}]^e \otimes [\text{any}]^e, [+ -]\}: \quad \phi_x \neq 0 \quad (2.48b)$$

Proof. If $\varrho_1 = 0$ (twist of $\mathcal{I}(\mathcal{C}_{m\dot{A}}^e, \mathcal{C}_{m\dot{A}}^e)$ vanishes) then Lemma 2.1 holds true. From (2.28) it follows that $\mathcal{B} = \mathcal{B}(q, p, y)$. Inserting this into (2.13) one arrives at (2.47).

For a subcase $[+ -]$ the expansion θ_1 is nonzero what implies that $\phi_x \neq 0$ and $M_1 \neq 0$. This proves (2.48b). For a subcase $[- -]$ we have $\phi_x = 0$, but $\phi\mathcal{Q}_x \neq \phi_y\mathcal{B} - \phi_q$, otherwise $M_2 = 0$ and a space is no longer equipped with an expanding ASD \mathcal{C} . Thus, (2.48a) is proved. ■

2.2.2 Spaces of the types $\{[\deg]^e \otimes [\text{any}]^e, [++]\}$ and $\{[\deg]^e \otimes [\text{any}]^e, [-+]\}$

Theorem 2.2. *Let (\mathcal{M}, ds^2) be a complex (neutral) space of the type $\{[\deg]^e \otimes [\text{any}]^e, [-+]\}$ or $\{[\deg]^e \otimes [\text{any}]^e, [++]\}$. Then there exists a local coordinate system (q, p, x, z) such that the metric takes the form*

$$\frac{1}{2}ds^2 = \phi^{-2} \left(-dpdx - x dqdz - z dqdx + (\Sigma_z dz + \Sigma_p dp) dq + \mathcal{A} dp^2 \right. \\ \left. - 2\mathcal{Q} dqdp - (2z\mathcal{Q} + z^2\mathcal{A} - \Omega(x - \Sigma_z) - z\Sigma_p) dq^2 \right) \quad (2.49)$$

where $\phi = \phi(q, p, x, z)$, $\mathcal{A} = \mathcal{A}(q, p, x, z)$, $\mathcal{Q} = \mathcal{Q}(q, p, x, z)$, $\Sigma = \Sigma(q, p, z)$ and $\Omega = \Omega(q, p, z)$ are arbitrary holomorphic (real smooth) functions such that

$$\text{for the type } \{[\deg]^e \otimes [\text{any}]^e, [-+]\} : \phi = F(x - \Sigma_z)^{\frac{1}{2}}, \quad F = F(q, p, z) \neq 0 \quad (2.50a)$$

$$\text{for the type } \{[\deg]^e \otimes [\text{any}]^e, [++]\} : \phi - 2\phi_x(x - \Sigma_z) \neq 0 \quad (2.50b)$$

and

$$\left[\phi - \phi_x(x - \Sigma_z) \neq 0 \text{ or } \right. \\ \left. z\phi_p + \frac{\phi\Sigma_p}{\Sigma_z - x} - \phi_q - \phi \frac{\partial}{\partial x}(\mathcal{Q} + z\mathcal{A}) + (\mathcal{Q} + z\mathcal{A}) \left(\phi_x - \frac{\phi}{\Sigma_z - x} \right) - \Omega\phi_z \neq 0 \right]$$

Proof. If twist of $\mathcal{I}(\mathcal{C}_{m^A}^e, \mathcal{C}_{m^A}^e)$ is nonzero ($\varrho_1 \neq 0$) then Lemma 2.2 holds true. Eq. (2.30) gives a solution for \mathcal{B} in terms of \mathcal{A} , \mathcal{Q} , Σ and Ω . Also, (2.29) gives y in terms of (q, p, z) . Putting this into (2.13) one arrives at (2.49).

For a subtype $[-+]$ one has $\theta_1 = 0 \implies \phi = F(q, p, z)(x - \Sigma_z)^{\frac{1}{2}}$ where $F = F(q, p, z)$ is an arbitrary nonzero function. Thus, (2.50a) is proved. For a subtype $[++]$ we have $\theta_1 \neq 0$ and ($M_1 \neq 0$ or $M_2 \neq 0$). It proves (2.50b). ■

2.3 Spaces of the types $[\deg]^e \otimes [\deg]^n$

In this Section we assume that a weak \mathcal{HH} -space admits $\mathcal{C}_{m^A}^n$. It means that $M_A = 0$. If $\varrho_1 = 0$ then from Lemma 2.1 it follows that $M_1 = 0 \implies \theta_1 = 0$. In this case we deal with the type $\{[\deg]^e \otimes [\deg]^n, [--]\}$. On the other hand, if $\varrho_1 \neq 0$ then from Lemma 2.2 it follows that $M_1 = 0 \implies \theta_1 \neq 0$. Thus, the second possible type is $\{[\deg]^e \otimes [\deg]^n, [++]\}$. Note, that the fact that $M_A = 0$ automatically implies that ASD Weyl spinor is algebraically special.

Theorem 2.3. *Let (\mathcal{M}, ds^2) be a complex (neutral) space of the type $\{[\deg]^e \otimes [\deg]^n, [--]\}$. Then there exists a local coordinate system (q, p, x, y) such that the metric takes the form*

$$\frac{1}{2}ds^2 = \phi^{-2} \left(dqdy - dpdx + \mathcal{A} dp^2 + \mathcal{B} dq^2 - 2(\phi^{-1}(\phi_y \mathcal{B} - \phi_q)x + S) dqdp \right) \quad (2.51)$$

where $\phi = \phi(q, p, y)$, $\mathcal{A} = \mathcal{A}(q, p, x, y)$, $\mathcal{B} = \mathcal{B}(q, p, y)$ and $S = S(q, p, y)$ are arbitrary holomorphic (real smooth) functions.

Proof. If twist of $\mathcal{I}(\mathcal{C}_{m^A}^e, \mathcal{C}_{m^A}^n)$ vanishes ($\varrho_1 = 0$) then Lemma 2.1 holds true. Condition $M_A = 0$ implies $\phi = \phi(q, p, y)$, $\mathcal{B} = \mathcal{B}(q, p, y)$ and $\mathcal{Q} = \phi^{-1}(\phi_y \mathcal{B} - \phi_q)x + S$ where $S = S(q, p, y)$ is an arbitrary function. Inserting this into (2.13) one arrives at (2.51). ■

Theorem 2.4. *Let (\mathcal{M}, ds^2) be a complex (neutral) space of the type $\{[deg]^e \otimes [deg]^n, [++]\}$. Then there exists a local coordinate system (q, p, x, z) such that the metric takes the form*

$$\begin{aligned} \frac{1}{2}ds^2 = F^{-2}(x - \Sigma_z)^{-2} \Big\{ & - (dp + zdq)dx - (x - \Sigma_z)dqdz + \mathcal{A}(dp + zdq)^2 \\ & - \left(2T(x - \Sigma_z)^2 + \frac{2}{F}(x - \Sigma_z)(\Omega F_z - zF_p + F_q) \right. \\ & \left. - \Omega\Sigma_{zz} + z\Sigma_{zp} - \Sigma_{zq} \right) (dp + zdq)dq + (x - \Sigma_z)\Omega dq^2 \Big\} \end{aligned} \quad (2.52)$$

where $\mathcal{A} = \mathcal{A}(q, p, x, z)$, $F = F(q, p, z)$, $T = T(q, p, z)$, $\Omega = \Omega(q, p, z)$ and $\Sigma = \Sigma(q, p, z)$ are arbitrary holomorphic (real smooth) functions.

Proof. If twist of $\mathcal{I}(\mathcal{C}_{m^A}^e, \mathcal{C}_{m^A}^n)$ is nonzero ($\varrho_1 \neq 0$) then Lemma 2.2 holds true. From (2.31a) one finds that condition $M_1 = 0$ implies

$$\phi = F(q, p, z)(x - \Sigma_z) \quad (2.53)$$

where F is an arbitrary nonzero function. Thus, θ_1 is automatically nonzero. Eq. $M_2 = 0$ is quite easy to solve and it yields

$$\mathcal{Q} + z\mathcal{A} = T(x - \Sigma_z)^2 + \frac{1}{F}(\Omega F_z - zF_p + F_q)(x - \Sigma_z) + \frac{1}{2}(\Sigma_p + z\Sigma_{zp} - \Sigma_{zq} - \Omega\Sigma_{zz}) \quad (2.54)$$

where $T = T(q, p, z)$ is an arbitrary function. Formula (2.54) is a solution for \mathcal{Q} in terms of \mathcal{A} , Σ , Ω , F and T . Inserting (2.54) into (2.30) one gets a solution for \mathcal{B} . Collecting all the results and inserting them into (2.13) one arrives at (2.52). \blacksquare

Function $F = F(q, p, z)$ (compare (2.53)) transforms under (2.14) as follows

$$F' = \frac{\Delta\lambda^{\frac{1}{2}}F}{q'_q - q'_p z} \quad (2.55)$$

2.4 Spaces of the types $[deg]^e \otimes [deg]^{ne}$

In this Section we consider weak \mathcal{HH} -spaces equipped with ASD \mathcal{C}^{ne} . To be more precise, we consider spaces of the types $\{[deg]^e \otimes [deg]^n, [--]\}$ (the metric (2.51)) and $\{[deg]^e \otimes [deg]^n, [++]\}$ (the metric (2.52)) and we equip these spaces with "the second" ASD congruence, namely $\mathcal{C}_{n^A}^e$. In what follows the metrics (2.51) and (2.52) are background for further considerations.

The classification of spaces equipped with SD \mathcal{C}^e and ASD \mathcal{C}^{ne} presented in [5] lists 7 subtypes of such spaces. These subtypes characterize by the following properties of $\mathcal{I}s$: $[--, ++]$, $[--, +-]$, $[--, -+]$, $[+, -]$, $[+, -+]$, $[+, +-]$ and $[+, ++]$. In Section 2.4.4 we prove that the subtype $\{[deg]^e \otimes [deg]^{ne}, [++, -+]\}$ cannot, in fact, exist⁵.

Remark 2.5. The eighth possibility (types $\{[deg]^e \otimes [deg]^{ne}, [--, --]\}$) also cannot exist. Indeed, for such types the function ϕ is a function of (q, p, y) only (compare Theorem 2.3). Moreover, $\theta_2 = 0$ what implies $\phi_y = 0$ (compare Lemma 2.3). Consequently, $\phi = \phi(q, p) \implies J_A = 0$ and SD $\mathcal{C}_{m^A}^e$ is nonexpanding what is a contradiction.

⁵It does not mean that SD \mathcal{C}^e and ASD \mathcal{C}^{ne} cannot intersect in such a manner that properties of $\mathcal{I}s$ are $[+, -+]$. It is possible within the types $\{[I]^e \otimes [deg]^{ne}, [++, -+]\}$ but we do not consider such types in this article.

2.4.1 Spaces of the types $\{[\mathbf{deg}]^e \otimes [\mathbf{deg}]^{ne}, [--, ++]\}$, $\{[\mathbf{deg}]^e \otimes [\mathbf{deg}]^{ne}, [--, +-]\}$ and $\{[\mathbf{deg}]^e \otimes [\mathbf{deg}]^{ne}, [--, -+]\}$

In this Section the metric (2.51) is a background.

Theorem 2.5. *Let (\mathcal{M}, ds^2) be a complex (neutral) space of the type $\{[deg]^e \otimes [deg]^{ne}, [--, +-]\}$. Then there exists a local coordinate system (q, p, x, y) such that the metric takes the form*

$$\frac{1}{2}ds^2 = \phi^{-2} \left(dqdy - dpdx + \mathcal{A}dp^2 + \mathcal{B}dq^2 - 2(\phi^{-1}(\mathcal{B}\phi_y - \phi_q)x + S) dqdp \right) \quad (2.56)$$

where $\phi = \phi(q, p, y)$, $\mathcal{A} = \mathcal{A}(q, p, x)$, $\mathcal{B} = \mathcal{B}(q, p, y)$ and $S = S(q, p, y)$ are arbitrary holomorphic (real smooth) functions such that $\phi_y \neq 0$.

Proof. Consider the metric (2.51) equipped with $\mathcal{C}_{n\dot{A}}^e$ such that $\mathcal{I}(\mathcal{C}_{m\dot{A}}^e, \mathcal{C}_{n\dot{A}}^e) = \mathcal{I}^{+-}$. From Lemma 2.3 one finds that $\mathcal{A}_y = 0$ and $\theta_2 \sim \phi_y \neq 0$. The condition $\phi_y \neq 0$ automatically implies that $N_1 \neq 0$ so $\mathcal{C}_{n\dot{A}}^e$ is necessarily expanding. ■

Spaces for which twist ϱ_2 of $\mathcal{I}(\mathcal{C}_{m\dot{A}}^e, \mathcal{C}_{n\dot{A}}^e)$ is nonzero are a little more complicated. For such spaces Theorem 2.3 and Lemma 2.4 hold true simultaneously. Let us consider the case $\mathcal{I}(\mathcal{C}_{m\dot{A}}^e, \mathcal{C}_{n\dot{A}}^e) = \mathcal{I}^{++}$ first.

Theorem 2.6. *Let (\mathcal{M}, ds^2) be a complex (neutral) space of the type $\{[deg]^e \otimes [deg]^{ne}, [--, ++]\}$. Then there exists a local coordinate system (q, p, w, y) such that the metric takes the form*

$$\begin{aligned} \frac{1}{2}ds^2 = \phi^{-2} \{ & (dq + wdp)dy + (y - H_w)dpdw - H_q dqdp + \mathcal{B}dq^2 \\ & - 2(\phi^{-1}(\mathcal{B}\phi_y - \phi_q)(H - wy) + S) (dq + wdp)dp \\ & + (-w^2\mathcal{B} + R(y - H_w) - wH_q) dp^2 \} \end{aligned} \quad (2.57)$$

where $\phi = \phi(q, p, y)$, $\mathcal{B} = \mathcal{B}(q, p, y)$, $S = S(q, p, y)$, $R = R(q, p, w)$ and $H = H(q, p, w)$ are arbitrary holomorphic (real smooth) functions such that

$$\phi - 2\phi_y(y - H_w) \neq 0 \quad (2.58a)$$

and

$$\left[\phi - \phi_y(y - H_w) \neq 0 \quad \text{or} \right] \quad (2.58b)$$

$$\begin{aligned} \phi_p - w\phi_q - \frac{\phi H_q}{H_w - y} - \phi \frac{\partial}{\partial y} (\phi^{-1}(\mathcal{B}\phi_y - \phi_q)(H - wy) + S + w\mathcal{B}) \\ + (\phi^{-1}(\mathcal{B}\phi_y - \phi_q)(H - wy) + S + w\mathcal{B}) \left(\phi_y - \frac{\phi}{H_w - y} \right) \neq 0 \end{aligned} \quad (2.58c)$$

Proof. Consider the metric (2.51) equipped with $\mathcal{C}_{n\dot{A}}^e$ such that $\mathcal{I}(\mathcal{C}_{m\dot{A}}^e, \mathcal{C}_{n\dot{A}}^e) = \mathcal{I}^{++}$. Co-ordinate x is replaced by (2.42). Formula (2.43) gives a solution for \mathcal{A} in terms of ϕ , \mathcal{B} , S , R and H , namely

$$\mathcal{A} = -2w(\phi^{-1}(\phi_y\mathcal{B} - \phi_q)(H - wy) + S) - w^2\mathcal{B} + R(y - H_w) + H_p - wH_q \quad (2.59)$$

Putting (2.59) and (2.42) into (2.51) one arrives at (2.57).

From Theorem 2.3 it follows that $\phi_x = 0$. Thus, $0 = \phi_x = \phi_w w_x \implies \phi_w = 0$. With $\phi_w = 0$, Eqs. (2.44a)-(2.44c) reduce to (2.58a)-(2.58c). ■

Theorem 2.7. *Let (\mathcal{M}, ds^2) be a complex (neutral) space of the type $\{[deg]^e \otimes [deg]^{ne}, [-, -, -+]\}$. Then there exists a local coordinate system (q, p, w, y) such that the metric takes the form*

$$\frac{1}{2}ds^2 = y^{-1}(dqdy + w dpdy + y dpdw + \mathcal{B} dq^2 + (w\mathcal{B} - 2S) dpdq + (-2wS + yR) dp^2) \quad (2.60)$$

where $\mathcal{B} = \mathcal{B}(q, p, y)$, $S = S(q, p, y)$ and $R = R(q, p, w)$ are arbitrary holomorphic (real smooth) functions.

Proof. Consider the metric (2.57). Condition $\theta_2 = 0$ yields $\phi - 2\phi_y(y - H_w) = 0$. Because $\phi = \phi(q, p, y)$, $\phi_y \neq 0$ and $H = H(q, p, w)$ one finds that $H_{ww} = 0 \implies H = H_1(q, p)w + H_2(q, p)$. Fast analysis of the transformation formula (2.46) proves that both H_1 and H_2 can be gauged away. With $H = 0$ one gets $\phi = \tilde{\phi}y^{\frac{1}{2}}$ where $\tilde{\phi}$ is an arbitrary nonzero function of (q, p) . From the transformation formula (2.17) it follows that $\tilde{\phi}$ can be brought to 1 without any loss of generality. Thus, $\phi = y^{\frac{1}{2}}$ and the metric (2.57) reduces to (2.60). ■

2.4.2 Spaces of the types $\{[deg]^e \otimes [deg]^{ne}, [++, --]\}$ and $\{[deg]^e \otimes [deg]^{ne}, [++, +-]\}$

In this Section spaces of the types $\{[deg]^e \otimes [deg]^n, [++]\}$ are equipped with \mathcal{C}_{nA}^e such that $\mathcal{I}(\mathcal{C}_{mA}^e, \mathcal{C}_{nA}^e) = \mathcal{I}^{+-}$ or $\mathcal{I}(\mathcal{C}_{mA}^e, \mathcal{C}_{nA}^e) = \mathcal{I}^{--}$. For such spaces Theorem 2.4 and Lemma 2.3 hold true simultaneously. Hence, the metric (2.52) is a background.

Theorem 2.8. *Let (\mathcal{M}, ds^2) be a complex (neutral) space of the type $\{[deg]^e \otimes [deg]^{ne}, [++, +-]\}$. Then there exists a local coordinate system (q, p, x, z) such that the metric takes the form*

$$\begin{aligned} \frac{1}{2}ds^2 = F^{-2}(x - \Sigma_z)^{-2} \Big\{ & - (dp + zdq)dx - (x - \Sigma_z)dqdz + \mathcal{A}(dp + zdq)^2 \\ & - \left(2T(x - \Sigma_z)^2 + \frac{2}{F}(x - \Sigma_z)(\Omega F_z - zF_p + F_q) \right. \\ & \left. - \Omega\Sigma_{zz} + z\Sigma_{zp} - \Sigma_{zq} \right) (dp + zdq)dq + (x - \Sigma_z)\Omega dq^2 \Big\} \end{aligned} \quad (2.61)$$

where $\mathcal{A} = \mathcal{A}(q, p, x)$, $F = F(q, p, z)$, $T = T(q, p, z)$, $\Omega = \Omega(q, p, z)$ and $\Sigma = \Sigma(q, p, z)$ are arbitrary holomorphic (real smooth) functions such that $F_z \neq 0$ or $\Sigma_{zz} \neq 0$.

Proof. Consider the metric (2.52) with constraint Eqs. (2.41). Formulas (2.41) need to be transformed from the coordinate system (q, p, x, y) to the coordinate system (q, p, x, z) . Hence, $0 = \mathcal{A}_y = \mathcal{A}_z z_y \implies \mathcal{A}_z = 0$. Moreover, $\theta_2 \neq 0$ implies $0 \neq \phi_y = [F_z(x - \Sigma_z) - F\Sigma_{zz}]z_y$. Thus, $F_z \neq 0$ or $\Sigma_{zz} \neq 0$. Also, $\theta_2 \neq 0$ automatically implies that $N_1 \neq 0$ so \mathcal{C}_{nA} is necessarily expanding. ■

Theorem 2.9. *Let (\mathcal{M}, ds^2) be a complex (neutral) space of the type $\{[deg]^e \otimes [deg]^{ne}, [++, --]\}$. Then there exists a local coordinate system (q, p, x, z) such that the metric takes the form*

$$\begin{aligned} \frac{1}{2}ds^2 = x^{-2} \Big\{ & - (dp + zdq)dx - x dqdz + \mathcal{A}(dp + zdq)^2 \\ & - 2Tx^2(dp + zdq)dq + x\Omega dq^2 \Big\} \end{aligned} \quad (2.62)$$

where $\mathcal{A} = \mathcal{A}(q, p, x)$, $T = T(q, p, z)$ and $\Omega = \Omega(q, p, z)$ are arbitrary holomorphic (real smooth) functions such that $T_z \neq 0$.

Proof. Consider the metric (2.61) such that $\theta_2 = 0$. Thus, $F_z = 0$ and $\Sigma_{zz} = 0$. From the transformation formulas (2.55) and (2.36) it follows that F can be brought to 1 and Σ can be gauged away without any loss of generality. Hence, one arrives at (2.62).

From (2.54) one finds $\mathcal{Q} + z\mathcal{A} = Tx^2$. Also, $y = -xz$ (compare (2.29) with $\Sigma = 0$). Because $\theta_2 = 0$ then $N_1 = 0$ (compare (2.41)). Thus, N_2 must be nonzero otherwise \mathcal{C}_{nA} is nonexpanding. From (2.41) we have

$$N_2 \sim \mathcal{A} - x\mathcal{Q}_y = \mathcal{A} - x\frac{\partial}{\partial y}(Tx^2 + \frac{y}{x}\mathcal{A}) = x^2T_z \quad (2.63)$$

Hence, $T_z \neq 0$. ■

2.4.3 Spaces of the types $\{[\deg]^e \otimes [\deg]^{ne}, [++, ++]\}$

For the most complicated types $\{[\deg]^e \otimes [\deg]^{ne}, [++, ++]\}$ twists of $\mathcal{I}(\mathcal{C}_{mA}^e, \mathcal{C}_{nA}^e)$ and $\mathcal{I}(\mathcal{C}_{mA}^e, \mathcal{C}_{m\dot{A}}^e)$ are nonzero, $\varrho_1 \neq 0$, $\varrho_2 \neq 0$. Thus, both Lemmas 2.2 and 2.4 hold true. In particular, Eqs. (2.29) and (2.42) hold true simultaneously. Hence

$$x = \frac{H - w\Sigma}{1 - zw}, \quad y = \frac{\Sigma - zH}{1 - zw}, \quad H = H(q, p, w), \quad \Sigma = \Sigma(q, p, z) \quad (2.64)$$

Additional constraints are Eqs. (2.30), (2.43), $M_A = 0$ (compare (2.31b)-(2.31c)), $N_A \neq 0$ (compare (2.44b)-(2.44c)), $\theta_1 \neq 0$ and $\theta_2 \neq 0$ (compare (2.31a) and (2.44a)).

Theorem 2.10. *Let (\mathcal{M}, ds^2) be a complex (neutral) space of the type $\{[\deg]^e \otimes [\deg]^{ne}, [++, ++]\}$. Then there exists a local coordinate system (q, p, w, z) such that the metric takes the form*

$$\begin{aligned} \frac{1}{2}ds^2 = \phi^{-2} \Big\{ & dqdy - dpdx - 2\mathbf{a}(dp + zdq)dq + \mathbf{b}dq^2 \\ & + \frac{\mathbf{c} - w^2\mathbf{b} - 2w\mathbf{a}(1 - wz)}{(1 - wz)^2}(dp + zdq)^2 \Big\} \end{aligned} \quad (2.65)$$

where $x, y, \phi, \mathbf{a}, \mathbf{b}$ and \mathbf{c} stand for abbreviations

$$\begin{aligned} x &:= \frac{H - w\Sigma}{1 - zw}, \quad y := \frac{\Sigma - zH}{1 - zw}, \quad \phi := F(x - \Sigma_z) \\ \mathbf{a} &:= T(x - \Sigma_z)^2 + \frac{1}{F}(x - \Sigma_z)(\Omega F_z - zF_p + F_q) + \frac{1}{2}(\Sigma_p - \Omega\Sigma_{zz} + z\Sigma_{zp} - \Sigma_{zq}) \\ \mathbf{b} &:= \Omega(x - \Sigma_z) + z\Sigma_p - \Sigma_q \\ \mathbf{c} &:= R(y - H_w) + H_p - wH_q \end{aligned}$$

and $\Omega = \Omega(q, p, z)$, $\Sigma = \Sigma(q, p, z)$, $R = R(q, p, w)$, $T = T(q, p, z)$, $H = H(q, p, w)$ and $F = F(q, p, z)$ are holomorphic (real smooth) functions such that

$$\frac{\phi}{H_w - y} + \frac{2\phi_z(1 - wz)}{\Sigma_z - x} \neq 0 \quad (2.66a)$$

and

$$\left[\frac{\phi}{H_w - y} + \frac{\phi_z(1 - wz)}{\Sigma_z - x} \neq 0 \quad \text{or} \right] \quad (2.66b)$$

$$\begin{aligned} & -\frac{\phi H_q}{H_w - y} - w\phi_q + \phi_p - \phi \frac{1 - wz}{\Sigma_z - x} \frac{\partial}{\partial z} \left(\mathbf{a} + \frac{w\mathbf{b} - z\mathbf{c}}{1 - wz} \right) - \phi_w R \\ & - \frac{\phi}{H_w - y} \left(\mathbf{a} + \frac{w\mathbf{b} - z\mathbf{c}}{1 - wz} \right) + \frac{\phi_z}{\Sigma_z - x} (w\Omega(x - \Sigma_z) + (1 - wz)(\mathbf{a} - \Sigma_p)) \neq 0 \end{aligned} \quad (2.66c)$$

Proof. Equation $M_1 = 0$ implies $\phi = F(q, p, z)(x - \Sigma_z)$, $F \neq 0$ and equation $M_2 = 0$ implies (2.54). From Eqs. (2.30), (2.43) and (2.54) one finds

$$\mathcal{A} = \frac{\mathbf{c} - 2w\mathbf{a}(1 - zw) - w^2\mathbf{b}}{(1 - zw)^2}, \quad \mathcal{Q} = \mathbf{a} - z\mathcal{A}, \quad \mathcal{B} = \mathbf{b} - 2z\mathbf{a} + z^2\mathcal{A} \quad (2.67)$$

Inserting ϕ and (2.67) into (2.13) with x and y replaced according to (2.64) one arrives at (2.65).

The expansion θ_1 is nonzero (it follows from Eq. $M_1 = 0$). Moreover, $\theta_2 \neq 0$ and at least one of the N_1 or N_2 must be nonzero. Formulas (2.44a)-(2.44c) transformed into coordinate system (q, p, w, z) lead to the Eqs. (2.66a)-(2.66c). ■

2.4.4 Spaces of the types $\{[\deg]^e \otimes [\deg]^{ne}, [++, -+]\}$

The last types which remained to be considered are types $\{[\deg]^e \otimes [\deg]^{ne}, [++, -+]\}$. The metric of spaces of such types can be obtained from the metric (2.65) by demanding that $\theta_2 = 0$. It gives the condition

$$\frac{\phi}{H_w - y} + \frac{2\phi_z(1 - wz)}{\Sigma_z - x} = 0 \quad (2.68)$$

where $\phi = F(q, p, z)(x - \Sigma_z)$; x and y are given by (2.64). Explicitly, (2.68) yields

$$F(x - \Sigma_z)^2 + 2(1 - wz)(y - H_w) \left(F_z(x - \Sigma_z) + F \left(\frac{w(x - \Sigma_z)}{1 - zw} - \Sigma_{zz} \right) \right) = 0 \quad (2.69)$$

Elimination of x and y in (2.69) via (2.64) gives an equation for $F = F(q, p, z)$, $\Sigma = \Sigma(q, p, z)$ and $H = H(q, p, w)$.

Before we deal with Eq. (2.69) we point out an additional relation, very helpful in further analysis. For the types $\{[\deg]^e \otimes [\deg]^{ne}, [++, -+]\}$ conditions $M_1 = 0$ (2.31b) and $\theta_2 = 0$ (2.44a) hold true simultaneously. A solution of Eq. $M_1 = 0$ reads $\phi = F(q, p, z)(x - \Sigma_z)$ (in the coordinate system (q, p, x, z)) and a solution of Eq. $\theta_2 = 0$ yields $\phi = G(q, p, w)(H_w - y)^{\frac{1}{2}}$, $G \neq 0$ (in the coordinate system (q, p, w, y)). Both these solutions give an alternative forms of ϕ . Thus, we arrive at the consistency condition

$$F(q, p, z) \left(\frac{H - w\Sigma}{1 - zw} - \Sigma_z \right) = G(q, p, w) \left(H_w - \frac{\Sigma - zH}{1 - zw} \right)^{\frac{1}{2}} \quad (2.70)$$

Of course, (2.69) and (2.70) are equivalent but it is algebraically problematic to find F , Σ , H and G using only (2.69) or only (2.70). In what follows we cleverly juggle with both (2.69) and (2.70).

Lemma 2.5. *Equation (2.68) holds true only if $\phi = 0$.*

Proof. We use condition (2.70) to eliminate factor $(H_w - y)$ in (2.69) and we put an explicit form of x into (2.69). After some algebraic work we find that Eq. (2.69) reduces to the following condition

$$\tilde{a}w^2 + \tilde{b}w + \tilde{c} = (1 - zw)G^2 - H \left(\tilde{f}_z + w(2\tilde{f} - z\tilde{f}_z) \right) \quad (2.71)$$

where

$$\begin{aligned} \tilde{c} &:= -2\tilde{f}\Sigma_{zz} - \tilde{f}_z\Sigma_z, \quad \tilde{b} := -2z\tilde{c} - 2\tilde{f}\Sigma_z - \tilde{f}_z\Sigma \\ \tilde{a} &:= -z^2\tilde{c} - z\tilde{b} - 2\tilde{f}\Sigma, \quad \tilde{f} := F^2 \end{aligned} \quad (2.72)$$

Note, that \tilde{a} , \tilde{b} , \tilde{c} and \tilde{f} are functions of (q, p, z) only. Thus, the left hand side of (2.71) is a second order polynomial in w . Acting on (2.71) by $\partial_w^3 \partial_z^2$ one finds that (i) $H_{ww} = 0$ or (ii) $\tilde{f}_{zzz} = 0$.

(i), case $H_{ww} = 0$. From (2.46) it follows that if $H_{ww} = 0$ then H can be gauged away without any loss of generality. With $H = 0$, formula (2.71) implies that \tilde{a} , \tilde{b} and \tilde{c} are all linear in z . After simple calculations one arrives at the solutions $\tilde{c} = \tilde{c}(q, p)$, $\tilde{a} = \tilde{\alpha}z$ and $\tilde{b} = -\tilde{c}z - \tilde{\alpha}$ where $\tilde{\alpha} = \tilde{\alpha}(q, p)$. The last step is to put these solutions into definition of \tilde{a} . Finally, one finds that $\tilde{f}\Sigma = 0$. Because H and Σ cannot simultaneously vanish, $\tilde{f} = 0$. It yields $F = 0$ and, eventually, $\phi = 0$.

(ii), case $\tilde{f}_{zzz} = 0$. If $\tilde{f}_{zzz} = 0$ then $\tilde{f} = \tilde{m}(q, p)z^2 + \tilde{n}(q, p)z + \tilde{s}(q, p)$ and $\tilde{a}_{zz} = \tilde{b}_{zz} = \tilde{c}_{zz} = 0$. Hence, $\tilde{c} = -\tilde{\alpha}z - \tilde{\beta}$, $\tilde{b} = \tilde{\gamma}z + \tilde{\delta}$, $\tilde{a} = \tilde{\mu}z + \tilde{\nu}$ where $\tilde{\alpha}$, $\tilde{\beta}$, $\tilde{\gamma}$, $\tilde{\delta}$, $\tilde{\mu}$ and $\tilde{\nu}$ are functions of (q, p) only. Inserting these formulas into (2.72) one arrives at the conditions

$$2\tilde{f}\Sigma_{zz} + \tilde{f}_z\Sigma_z = \tilde{\alpha}z + \tilde{\beta} \quad (2.73a)$$

$$2\tilde{f}\Sigma_z + \tilde{f}_z\Sigma = 2\tilde{\alpha}z^2 + (2\tilde{\beta} - \tilde{\gamma})z - \tilde{\delta} \quad (2.73b)$$

$$2\tilde{f}\Sigma = \tilde{\alpha}z^3 + (\tilde{\beta} - \tilde{\gamma})z^2 - (\tilde{\delta} + \tilde{\mu})z - \tilde{\nu} \quad (2.73c)$$

(2.73b) - $\partial_z(2.73c)$ gives $\tilde{f}_z\Sigma = \tilde{\alpha}z^2 - \tilde{\gamma}z - \tilde{\mu}$. Using this result to eliminate factor $\tilde{f}_z\Sigma$ in (2.73b) one finds the condition $2\tilde{f}\Sigma_z = \tilde{\alpha}z^2 + 2\tilde{\beta}z + \tilde{\mu} - \tilde{\delta}$. Inserting all these formulas into (2.73a) we get $\tilde{f}\Sigma_{zz} = 0$. Thus, $\tilde{f} = 0 \implies \phi = 0$, or $\Sigma_{zz} = 0$.

If $\Sigma_{zz} = 0$ then Σ can be gauged away without any loss of generality (compare transformation formula (2.36)). $\Sigma = 0$ implies $\tilde{a} = \tilde{b} = \tilde{c} = 0$ and (2.71) simplifies to the formula $H(\tilde{m} + \tilde{n}w + \tilde{s}w^2)$. Because Σ and H cannot be simultaneously equal 0, then $\tilde{m} = \tilde{n} = \tilde{s} = 0$. Finally, $\tilde{f} = 0 \implies \phi = 0$. Hence, the proof is completed. \blacksquare

Corollary 2.1. *Spaces of the types $\{[\deg]^e \otimes [\deg]^{ne}, [++, -+]\}$ do not exist.*

Proof. Eq. (2.68) is a necessary condition for spaces of the types $\{[\deg]^e \otimes [\deg]^{ne}, [++, -+]\}$ to exist. According to Lemma 2.5, Eq. (2.68) implies $\phi = 0$ what is a contradiction. \blacksquare

3 Spaces of the types $\{[\deg]^e \otimes [\mathbf{D}]^{nn}, [++, --]\}$

3.1 Non-Einstein case

3.1.1 General results

In this Section we finally arrive at the first class of para-Kähler spaces. To be more precise we find a general metric of spaces of the types $\{[\deg]^e \otimes [\mathbf{D}]^{nn}, [++, --]\}$. Such a metric can be extracted from Theorem 2.9 by demanding that the expansion of $\mathcal{C}_{n\bar{A}}$ vanishes. It is equivalent to the fact that $T_z = 0 \iff T = T(q, p)$. Hence, the metric takes the form (2.13) with

$$\phi = x, \quad \mathcal{A} = \mathcal{A}(q, p, x), \quad \mathcal{Q} = \frac{y}{x} \mathcal{A} + x^2 T, \quad T = T(q, p) \quad (3.1)$$

$$\mathcal{B} = \frac{y^2}{x^2} \mathcal{A} + 2yxT + x\Omega, \quad \Omega = \Omega(q, p, z), \quad z := -\frac{y}{x}$$

The metric (2.13) with (3.1) remains invariant under transformations (2.14) such that

$$\begin{aligned}\sigma^{\dot{A}} &= 0, \quad q' = q'(q), \quad p' = p'(q, p), \quad \lambda^{-1} = p_p'^2 \\ x' &= p_p' x, \quad y' = \frac{p_p'}{q_q'}(p_q' x + p_p' y), \quad z' = \frac{1}{q_q'}(p_p' z - p_q')\end{aligned}\tag{3.2}$$

Metrical functions \mathcal{A} , T and Ω transform under (3.2) as follows

$$\begin{aligned}\mathcal{A}' &= \mathcal{A} + \frac{p_{pp}'}{p_p'} x, \quad T' = \frac{1}{p_p' q_q'} T, \\ p_p' \Omega' &= \frac{p_p'^2}{q_q'^2} \Omega - \frac{p_p' p_{pp}'}{q_q'^2} z^2 + \frac{1}{q_q'} \frac{\partial}{\partial q} \left(\frac{p_p'^2}{q_q'} \right) z - \frac{p_p'}{q_q'} \frac{\partial}{\partial q} \left(\frac{p_q'}{q_q'} \right)\end{aligned}\tag{3.3}$$

For further purposes it is necessary to calculate conformal curvature coefficients, curvature scalar and traceless Ricci tensor. Curvature scalar reads

$$\mathcal{R} = -2x^2 \left(\mathcal{A}_{xx} + \frac{1}{x} \Omega_{zz} - \frac{4\mathcal{A}_x}{x} + \frac{6\mathcal{A}}{x^2} \right)\tag{3.4}$$

Nonzero ASD curvature coefficients are⁶

$$C_{ii\dot{z}\dot{z}} = \frac{\mathcal{R}}{12}, \quad C_{iii\dot{z}} = z \frac{\mathcal{R}}{4}, \quad C_{iiii} = z^2 \frac{\mathcal{R}}{2},\tag{3.5}$$

SD conformal curvature coefficients takes the form

$$\begin{aligned}C^{(3)} &= \frac{\mathcal{R}}{6} + 2\mathcal{A} - 2x\mathcal{A}_x = -\frac{x}{3}(\Omega_{zz} + x\mathcal{A}_{xx} + 2\mathcal{A}_x) \\ -x^{-4}C^{(2)} &= 4T_p x - \Omega_{zp} - z\mathcal{A}_{xp} - \frac{z}{x}\mathcal{A}_p + \frac{1}{x}\mathcal{A}_q + \mathcal{A}_{xq} + T(x\Omega_{zz} - x^2\mathcal{A}_{xx} + 6\mathcal{A}) \\ \frac{1}{2}x^{-6}C^{(1)} &= -x\Omega_{pp} + 2Tx^2\Omega_{zp} + T^2x^2(-x\Omega_{zz} - x^2\mathcal{A}_{xx} + 6x\mathcal{A}_x - 12\mathcal{A}) \\ &\quad + \Omega_z(\mathcal{A}_q - z\mathcal{A}_p - 2Tx^2\mathcal{A}_x + 4Tx\mathcal{A} + 2x^2T_p) + T_q(2x^2\mathcal{A}_x - 4x\mathcal{A}) \\ &\quad + T_p(-4Tx^3 - 2zx^2\mathcal{A}_x + 4zx\mathcal{A}) - 2T_{pq}x^2 + 2zx^2T_{pp} - 2Tzx^2\mathcal{A}_{xp} \\ &\quad + \mathcal{A}_p(\Omega + 8Txz) + x\mathcal{A}_x\Omega_p - \mathcal{A}_{qq} + 2z\mathcal{A}_{pq} - 8Tx\mathcal{A}_q + 2Tx^2\mathcal{A}_{xq} - z^2\mathcal{A}_{pp}\end{aligned}\tag{3.6}$$

Nonzero coefficients of the traceless Ricci tensor are

$$-\frac{4}{x}C_{12i\dot{z}} = -\frac{2}{zx}C_{12i\dot{i}} = \alpha, \quad \frac{1}{zx^3}C_{22i\dot{i}} = \frac{2}{x^3}C_{22i\dot{z}} = \beta\tag{3.7}$$

where

$$\begin{aligned}\alpha &:= \Omega_{zz} - x\mathcal{A}_{xx} + 2\mathcal{A}_x \\ \beta &:= \partial_q(x\mathcal{A}_x - 3\mathcal{A}) - z\partial_p(x\mathcal{A}_x - 3\mathcal{A}) + x\Omega_{pz} + \frac{1}{2}Tx\mathcal{R}\end{aligned}\tag{3.8}$$

Equivalently

$$C_{AB\dot{C}\dot{D}} = (x^3\beta\delta_A^2\delta_B^2 - x\alpha\delta_A^1\delta_B^2)m_{(\dot{C}n_{\dot{D})}), \quad m_{\dot{A}} = [z, 1], n_{\dot{A}} = [1, 0]\tag{3.9}$$

⁶Obviously, the condition $2C_{iii\dot{z}}^2 - 3C_{iiii}C_{ii\dot{z}\dot{z}} = 0$ for the ASD Weyl spinor to be of the type [D] is identically satisfied.

If $\alpha \neq 0$ then the traceless Ricci tensor has four different eigenvectors and two double eigenvalues (complex type is $[2N_1 - 2N]_2$, see [19]). If $\alpha = 0$ and $\beta \neq 0$ then the traceless Ricci tensor has two different eigenvectors and one quadruple eigenvalue (complex type $^{(2)}[4N]_2^a$). Algebraic types of the traceless Ricci tensor in neutral spaces are slightly more complicated and we do not discuss this issue here (for details see [6, 7]).

Theorem 3.1. *Let (\mathcal{M}, ds^2) be a complex (neutral) space of the type $\{[deg]^e \otimes [D]^{nn}, [++, --]\}$ ($\{[deg]^e \otimes [D_r]^{nn}, [++, --]\}$). Then there exist a local coordinate system (q, p, x, z) such that the metric takes the form*

$$\frac{1}{2}ds^2 = x^{-2} \left\{ - (dp + zdq)dx - x dqdz + \mathcal{A}(dp + zdq)^2 - 2T_0x^2(dp + zdq)dq + x\Omega dq^2 \right\} \quad (3.10)$$

where $T_0 = \{-1, 0, 1\}$ is a constant, $\mathcal{A} = \mathcal{A}(q, p, x)$ and $\Omega = \Omega(q, p, z)$ are arbitrary holomorphic (real smooth) functions such that

$$\Omega_{zz} + x\mathcal{A}_{xx} - 4\mathcal{A}_x + \frac{6\mathcal{A}}{x} \neq 0 \quad (3.11)$$

Proof. Inserting (3.1) into (2.13) one arrives at (3.10). From (3.3) it follows that the function T can be brought to a constant value, namely $T = \{-1, 0, 1\}$. The curvature scalar \mathcal{R} given by (3.4) must be nonzero, otherwise the ASD conformal curvature vanishes. Hence, (3.11) is valid. \blacksquare

As long as $C^{(3)} \neq 0$ and $2C^{(2)}C^{(2)} - 3C^{(3)}C^{(1)} \neq 0$ the metric (3.10) is of the type $\{[II]^e \otimes [D]^{nn}, [++, --]\}$.

3.1.2 Spaces of the type $\{[D]^e \otimes [D]^{nn}, [++, --]\}$

If $C^{(3)} \neq 0$ and $2C^{(2)}C^{(2)} - 3C^{(3)}C^{(1)} = 0$ the metric (3.10) is of the type $\{[D]^e \otimes [D]^{nn}, [++, --]\}$. However, the condition $2C^{(2)}C^{(2)} - 3C^{(3)}C^{(1)} = 0$ written explicitly is even more complicated than its counterpart for the space of the type $[D]^n \otimes [D]^{nn}$ (Eq. (3.20) of Ref. [5]). It will be analyzed elsewhere.

Remark 3.1. An example of a (non-Einstein) space of the type $\{[D]^e \otimes [D]^{nn}, [++, --]\}$ cannot be obtained even if one additionally assumes that the traceless Ricci tensor has two eigenvectors. Indeed, conditions $2C^{(2)}C^{(2)} - 3C^{(3)}C^{(1)} = 0$ and $\alpha = 0$ imply $\beta = 0$. Hence, a space becomes an Einstein space. Such a solution reads

$$\mathcal{A} = \frac{\Lambda}{3} + \frac{x^3}{(a_0p + b_0q + c_0)^3}, \quad \Omega = T = 0 \quad (3.12)$$

where a_0 is an arbitrary constant, $b_0 = \{-1, 0, 1\}$ and c_0 is a constant which is nonzero only if both a_0 and b_0 vanish. The metric given by (3.12) reads

$$\frac{1}{2}ds^2 = x^{-2} \left(dqdy - dpdx + \left(\frac{\Lambda}{3x} + \frac{x^2}{(a_0p + b_0q + c_0)^3} \right) (xdp - ydq)^2 \right) \quad (3.13)$$

To be more precise, solution (3.13) is a metric of the type $\{[D]^{ee} \otimes [D]^{nn}, [++, --, --, ++]\}$. It belongs to one of the classes described in [3].

3.1.3 Spaces of the type $\{[\text{III}]^e \otimes [\text{D}]^{nn}, [++, --]\}$

In this Section we do not assume that the function T is gauged to a constant value, like it was done in Theorem 3.1. Thus, $T = T(q, p)$ but the gauge function $p' = p'(q, p)$ is still arbitrary at our disposal.

The type $\{[\text{III}]^e \otimes [\text{D}]^{nn}, [++, --]\}$ is given by $C^{(3)} = 0$, $C^{(2)} \neq 0$ and $\mathcal{R} \neq 0$. The condition $C^{(3)} = 0$ implies

$$\mathcal{A} = Ax + B + \frac{C}{x}, \quad \Omega = -Az^2 + Mz + N \quad (3.14)$$

where A , B , C , M and N are functions of (q, p) . The transformation formula for A reads

$$p'_p A = A + \frac{p'_{pp}}{p'_p} \quad (3.15)$$

Thus, A can be gauged away without any loss of generality. However, from now on the gauge (3.2) is restricted to the transformations such that

$$p' = g(q)p + h(q) \quad (3.16)$$

The fact that Ω becomes a first order polynomial in z (and $z = -y/x$) allows us to return to the hyperheavenly coordinate system (q, p, x, y) . In terms of (q, p, x, y) functions \mathcal{A} , \mathcal{Q} and \mathcal{B} take the form

$$\begin{aligned} \mathcal{A} &= B + \frac{C}{x}, \quad \mathcal{Q} = B \frac{y}{x} + C \frac{y}{x^2} + Tx^2 \\ \mathcal{B} &= B \frac{y^2}{x^2} + C \frac{y^2}{x^3} + 2Txy - My + Nx \end{aligned} \quad (3.17)$$

Hence

$$\begin{aligned} -\frac{\mathcal{R}}{12} &= B + \frac{2C}{x} \\ -x^{-4}C^{(2)} &= B_p \frac{y}{x^2} + (B_q + 4TC) \frac{1}{x} + 4T_p x + 6TB - M_p \\ \alpha &= -4x^{-2}C \\ \beta &= -\frac{4}{x^2}(xC_q + yC_p) + x(M_p - 6TB) - 3B_p \frac{y}{x} - 3(4TC + B_q) \end{aligned} \quad (3.18)$$

(Coefficient $C^{(1)}$ is more complicated and we do not need its explicit form at this stage). Thus one formulates

Theorem 3.2. *Let (\mathcal{M}, ds^2) be a complex (neutral) space of the type $\{[\text{III}]^e \otimes [\text{D}]^{nn}, [++, --]\}$ ($\{[\text{III}_r]^e \otimes [\text{D}_r]^{nn}, [++, --]\}$). Then there exist a local coordinate system (q, p, x, y) such that the metric takes the form*

$$\begin{aligned} \frac{1}{2}ds^2 &= x^{-2} \left\{ dqdy - dpdx + \left(B + \frac{C}{x} \right) dp^2 - 2 \left(B \frac{y}{x} + C \frac{y}{x^2} + Tx^2 \right) dqdp \right. \\ &\quad \left. + \left(B \frac{y^2}{x^2} + C \frac{y^2}{x^3} + 2Txy - My + Nx \right) dq^2 \right\} \end{aligned} \quad (3.19)$$

where $B = B(q, p)$, $C = C(q, p)$, $M = M(q, p)$, $N = N(q, p)$ and $T = T(q, p)$ are arbitrary holomorphic (real smooth) functions such that

$$(B \neq 0 \text{ or } C \neq 0) \quad (3.20a)$$

and

$$(B_p \neq 0 \text{ or } T_p \neq 0 \text{ or } B_q + 4TC \neq 0 \text{ or } 6TB - M_p \neq 0) \quad (3.20b)$$

If $C \neq 0$ the traceless Ricci tensor has four eigenvectors and if $C = 0$ the traceless Ricci tensor has two eigenvectors.

Proof. The proof follows immediately from (3.17), (3.18) and (2.13). ■

3.1.4 Spaces of the type $\{[N]^e \otimes [D]^{nn}, [++, --]\}$

Theorem 3.3. Let (\mathcal{M}, ds^2) be a complex (neutral) space of the type $\{[N]^e \otimes [D]^{nn}, [++, --]\}$ ($\{[N_r]^e \otimes [D_r]^{nn}, [++, --]\}$). Then there exist a local coordinate system (q, p, x, y) such that the metric takes the form

$$\begin{aligned} \frac{1}{2}ds^2 = x^{-2} \Big\{ & dqdy - dpdx + \left(B + \frac{C}{x}\right) dp^2 - 2 \left(B \frac{y}{x} + C \frac{y}{x^2} + T_0 x^2\right) dqdp \\ & + \left(B \frac{y^2}{x^2} + C \frac{y^2}{x^3} + 2T_0 xy - 6T_0 Bpy + Nx\right) dq^2 \Big\} \end{aligned} \quad (3.21)$$

where $T = T_0 = \{-1, 0, 1\}$ as a constant, $B = B(q)$, $C = C(q, p)$ and $N = N(q, p)$ are arbitrary holomorphic (real smooth) functions such that

$$(4CT_0 = -B_q) \quad (3.22a)$$

and

$$(B \neq 0 \text{ or } C \neq 0) \quad (3.22b)$$

and

$$\begin{aligned} (C_{pp} \neq 0 \text{ or } 4C_{pq} + 3BB_q \neq 0 \text{ or } 24T_0^2 B^2 p - N_{pp} - 3T_0 B_q \neq 0 \text{ or} \\ 2NC_p - 2CN_p - 2C_{qq} - 3pBB_{qq} \neq 0 \text{ or } B_{qq} - 2T_0 pBB_q \neq 0) \end{aligned} \quad (3.22c)$$

If $C \neq 0$ the traceless Ricci tensor has four eigenvectors and if $C = 0$ the traceless Ricci tensor vanishes.

Proof. For the type $\{[N]^e \otimes [D]^{nn}, [++, --]\}$ the coefficient $C^{(2)}$ vanishes. Hence, we obtain the conditions (compare (3.18))

$$B = B(q), \quad T = T(q), \quad B_q + 4TC = 0, \quad M_p = 6TB \quad (3.23)$$

From (3.3) and (3.16) one finds the transformation for T

$$T' = \frac{1}{gq'_q} T \quad (3.24)$$

Because $T = T(q)$ then it can be gauged to a constant value, $T = T_0 = \{-1, 0, 1\}$. From now on the gauge functions g and q'_q are restricted to such that $gq'_q = 1$.

With $T = T_0$ we find $M = 6T_0 Bp + m(q)$ where m is an arbitrary function which transform as follows

$$m' = gm - 6T_0 Bh + 3g_q \quad (3.25)$$

Obviously, m can be gauged away without any loss of generality. Hence, $M = 6T_0Bp$ and (3.21) is proved. The condition (3.22b) assures that the curvature scalar \mathcal{R} is nonzero. The coefficient $C^{(1)}$ takes the form

$$C^{(1)} = -C_{pp} \frac{z^2}{x} + \left(2C_{pq} + \frac{3}{2}BB_q \right) \frac{z}{x} + (24T_0^2B^2p - N_{pp} - 3T_0B_q) x \quad (3.26)$$

$$+ \left(NC_p - CN_p - C_{qq} - \frac{3}{2}pBB_{qq} \right) \frac{1}{x} + \frac{3}{2}B_{qq} - 3T_0pBB_q$$

Thus, $C^{(1)} \neq 0$ implies (3.22c).

Finally, from (3.18) one finds that $C \neq 0 \iff \alpha \neq 0$. However, if $C = 0$ then $\alpha = \beta = 0$ and the traceless Ricci tensor vanishes. ■

3.2 Einstein case

Results from Section 3.1.1 can be easily specialized to the Einstein case. $\alpha = \beta = 0$ and $\mathcal{R} = -4\Lambda \neq 0$ imply the solutions for \mathcal{A} , Ω and T

$$\mathcal{A} = Ax^3 + Bx + \frac{\Lambda}{3}, \quad \Omega = -Bz^2 + Mz + N, \quad T = \frac{M_p - 2B_q}{2\Lambda} \quad (3.27)$$

Transformation formula for B reads

$$p'_p B' = B + \frac{p'_{pp}}{p'_p} \quad (3.28)$$

Thus, B can be gauged away without any loss of generality but from now on $p' = g(q)p + h(q)$. With $B = 0$ the metrical functions ϕ , \mathcal{A} , \mathcal{Q} and \mathcal{B} take the form

$$\phi = x, \quad \mathcal{A} = Ax^3 + \frac{\Lambda}{3}, \quad \mathcal{Q} = Ayx^2 + \frac{\Lambda}{3} \frac{y}{x} + \frac{M_p}{2\Lambda} x^2, \quad (3.29)$$

$$\mathcal{B} = Ay^2x + \frac{\Lambda}{3} \frac{y^2}{x^2} + \frac{M_p}{\Lambda} xy + Nx - My$$

and

$$C_{i i \dot{2} \dot{2}} = -\frac{\Lambda}{3}, \quad C_{i i i \dot{2}} = \Lambda \frac{y}{x}, \quad C_{i i i i} = -2\Lambda \frac{y^2}{x^2} \quad (3.30)$$

$$C^{(3)} = -4Ax^3$$

$$-x^{-5}C^{(2)} = 4A_p y + 4A_q x + \frac{2}{\Lambda} M_{pp}$$

(Coefficient $C^{(1)}$ is much more complicated and its explicit form is not necessary at this stage.)

Theorem 3.4. *Let (\mathcal{M}, ds^2) be an Einstein complex (neutral) space of the type $\{[deg]^e \otimes [D]^{nn}, [++, --]\}$ ($\{[deg]^e \otimes [D_r]^{nn}, [++, --]\}$). Then there exist a local coordinate system (q, p, x, y) such that the metric takes the form*

$$\frac{1}{2}ds^2 = x^{-2} \left\{ dqdy - dpdx + \left(Ax^3 + \frac{\Lambda}{3} \right) dp^2 - 2 \left(Ayx^2 + \frac{\Lambda}{3} \frac{y}{x} + \frac{M_p}{2\Lambda} x^2 \right) dqdp \right. \quad (3.31)$$

$$\left. + \left(Ay^2x + \frac{\Lambda}{3} \frac{y^2}{x^2} + \frac{M_p}{\Lambda} xy + Nx - My \right) dq^2 \right\}$$

where $\Lambda \neq 0$ is a cosmological constant, $A = A(q, p)$, $M = M(q, p)$ and $N = N(q, p)$ are arbitrary holomorphic (real smooth) functions.

Proof. Inserting (3.29) into (2.13) one arrives at (3.31). ■

As long as $A \neq 0$ and $2C^{(2)}C^{(2)} - 3C^{(3)}C^{(1)} \neq 0$ the metric (3.31) is of the type $\{[II]^e \otimes [D]^{nn}, [++, --]\}$. We skip an analysis of the type $[D]^{ee} \otimes [D]^{nn}$ (all such types were found in [3]).

Theorem 3.5. *Let (\mathcal{M}, ds^2) be an Einstein complex (neutral) space of the type $\{[III]^e \otimes [D]^{nn}, [++, --]\}$ ($\{[III_r]^e \otimes [D_r]^{nn}, [++, --]\}$). Then there exist a local coordinate system (q, p, x, y) such that the metric takes the form*

$$\begin{aligned} \frac{1}{2}ds^2 = x^{-2} \Big\{ & dqdy - dpdx + \frac{\Lambda}{3}dp^2 - 2 \left(\frac{\Lambda}{3} \frac{y}{x} + \frac{M_p}{2\Lambda} x^2 \right) dqdp \\ & + \left(\frac{\Lambda}{3} \frac{y^2}{x^2} + \frac{M_p}{\Lambda} xy + Nx - My \right) dq^2 \Big\} \end{aligned} \quad (3.32)$$

where $\Lambda \neq 0$ is a cosmological constant, $M = M(q, p)$ and $N = N(q, p)$ are arbitrary holomorphic (real smooth) functions such that $M_{pp} \neq 0$.

Proof. $C^{(3)} = 0$ implies $A = 0$ and $C^{(2)} \neq 0$ implies $M_{pp} \neq 0$ (compare (3.30)). ■

Theorem 3.6. *Let (\mathcal{M}, ds^2) be an Einstein complex (neutral) space of the type $\{[I]^e \otimes [D]^{nn}, [++, --]\}$ ($\{[N_r]^e \otimes [D_r]^{nn}, [++, --]\}$). Then there exist a local coordinate system (q, p, x, y) such that the metric takes the form*

$$\begin{aligned} \frac{1}{2}ds^2 = x^{-2} \Big\{ & dqdy - dpdx + \frac{\Lambda}{3}dp^2 - 2 \left(\frac{\Lambda}{3} \frac{y}{x} + \frac{M_0}{2\Lambda} x^2 \right) dqdp \\ & + \left(\frac{\Lambda}{3} \frac{y^2}{x^2} + \frac{M_0}{\Lambda} xy + Nx - M_0py \right) dq^2 \Big\} \end{aligned} \quad (3.33)$$

where $\Lambda \neq 0$ is a cosmological constant, $M_0 = \{-1, 0, 1\}$ is a constant and $N = N(q, p)$ is an arbitrary holomorphic (real smooth) function such that $3N_{pp} - 2M_0^2p \neq 0$.

Proof. $C^{(2)} = 0$ implies $M_{pp} = 0$. Hence, $M = m(q)p + n(q)$. Transformation laws for m and n read

$$\begin{aligned} q'_q g m' &= m \\ q'_q n' &= n - \frac{h}{g} m + \frac{q'_q}{g^2} \frac{\partial}{\partial q} \left(\frac{g^2}{q'_q} \right) \end{aligned} \quad (3.34)$$

Arbitrariness of the gauge functions q'_q and g allows to gauge away n and brought m to a constant value, $m = M_0 = \{-1, 0, 1\}$. Hence, $M = M_0p$. Coefficient $C^{(1)}$ reads

$$-\frac{1}{2}x^{-7}C^{(1)} = N_{pp} - \frac{2}{3}M_0^2p \quad (3.35)$$

Thus, $3N_{pp} - 2M_0^2p \neq 0$. ■

4 Spaces of the types $\{[\mathbf{deg}]^e \otimes [\mathbf{D}]^{nn}, [++, ++]\}$

4.1 Non-Einstein case

In this Section we consider the second class of para-Kähler spaces, i.e., spaces of the types $\{[\mathbf{deg}]^e \otimes [\mathbf{D}]^{nn}, [++, ++]\}$. Twists ϱ_1 and ϱ_2 are nonzero within this class. Thus,

Lemmas 2.2 and 2.4 hold true simultaneously. Hence, the coordinate system (q, p, w, z) seems to be the optimal one. For the sake of transparency of the considerations we proceed according to the following strategy⁷

- (i) We deal with Eqs. $M_1 = 0$ (2.31b) and $N_1 = 0$ (2.44b). Each of these equations gives a solution for ϕ . Consequently, we find ϕ and an equation which we call "the first consistency condition".
- (ii) We solve the first consistency condition.
- (iii) We consider Eqs. (2.30) and (2.43) with $M_2 = 0$ (2.31c) and $N_2 = 0$ (2.44c). These are four conditions for three functions \mathcal{A} , \mathcal{Q} and \mathcal{B} . We obtain solutions for \mathcal{A} , \mathcal{Q} and \mathcal{B} and a single equation which we call "the second consistency condition".
- (iv) We solve the second consistency condition.

If (i)-(iv) are completed we are ready to write down the metric. Note, that equations $M_1 = 0$ and $N_1 = 0$ automatically imply that $\theta_1 \neq 0$ and $\theta_2 \neq 0$.

4.1.1 The first consistency condition

From equation $M_1 = 0$ it follows that $\phi = F(q, p, z)(x - \Sigma_z)$ and this solution is written in the coordinate system (q, p, x, z) . Analogously, from equation $N_1 = 0$ one finds that $\phi = G(q, p, w)(H_w - y)$ and this solution is expressed in the coordinate system (q, p, w, y) . Both these solutions can be easily transformed into the coordinate system (q, p, w, z) via (2.64). Thus we find the first consistency condition

$$F(q, p, z) \left(\frac{H - w\Sigma}{1 - zw} - \Sigma_z \right) = G(q, p, w) \left(H_w - \frac{\Sigma - zH}{1 - zw} \right) \quad (4.1)$$

Transformation formulas for $\Sigma(q, p, z)$, $H(q, p, w)$ and $F(q, p, z)$ are given by (2.36), (2.46) and (2.55), respectively. Transformation for $G(q, p, w)$ reads

$$G' = \frac{\lambda^{\frac{1}{2}} \Delta G}{p'_p - p'_q w} \quad (4.2)$$

Playing algebraically with (4.1) we find two conditions helpful in further analysis

$$F_{zz} H_{ww} = -\Sigma_{zz} G_{ww} \quad (4.3a)$$

$$G^3 H_{ww} = -F^3 \Sigma_{zz} \quad (4.3b)$$

(we skip the proofs of (4.3a) and (4.3b) as they are quite straightforward).

Lemma 4.1. *If $\Sigma_{zz} = 0 \iff H_{ww} = 0$ then the functions F , Σ , G and H can be brought to the form*

$$\Sigma = 0, \quad H = w, \quad F = 1, \quad G = w \quad (4.4)$$

without any loss of generality.

⁷Alternatively, the metric of spaces of the types $\{[\deg]^e \otimes [\mathbf{D}]^{nn}, [++, ++]\}$ can be obtained from Theorem 2.10 by demanding that \mathcal{C}_{nA} is nonexpanding. However, such an approach is more complicated.

Proof. If $\Sigma_{zz} = 0$ then from (4.3b) it follows that $H_{ww} = 0$ and vice versa. Thus, the equivalence $\Sigma_{zz} = 0 \iff H_{ww} = 0$ is proved.

If $\Sigma_{zz} = 0$ then Σ is linear in z with coefficients which depend on (q, p) only. Hence, the gauge functions $\sigma^{\dot{A}}$ can be used to gauge away Σ (compare (2.36)). After this step the gauge (2.14) is restricted to such that $\sigma^{\dot{A}} = 0$. Function H becomes linear in w . Consequently, it can be simplified to the form $H = w$ (compare (2.46)). The choice $H = w$ imposes additional conditions on gauge (2.14), namely, $q'_p = 0$ and $\lambda^{-1} = q'_q$. With $\Sigma = 0$ and $H = w$ the first consistency condition (4.1) reduces to the equation $wF(q, p, z) = G(q, p, w)$. Thus, $F = F(q, p)$. Finally, F can be brought to 1 (compare (2.55)) what yields $G = w$. ■

Remark 4.1. The choice (4.4) leaves us with the gauge (2.14) restricted to the conditions

$$\begin{aligned} q' &= q'(q), \quad \frac{dq'}{dq} := f(q), \quad \lambda = \frac{1}{f}, \quad p' = f^{\frac{1}{2}}p + h(q), \\ x' &= f^{\frac{1}{2}}x, \quad y' = y + f^{-\frac{1}{2}}p'_q x, \quad \sigma^{\dot{A}} = 0 \end{aligned} \quad (4.5)$$

where $f = f(q) \neq 0$ and $h = h(q)$ are arbitrary gauge functions.

Remark 4.2. Note, that the choice (4.4) implies

$$\phi = x; \quad x = \frac{w}{1 - zw}, \quad y = -\frac{zw}{1 - zw} \iff z = -\frac{y}{x}, \quad w = \frac{x}{1 - y} \quad (4.6)$$

Remark 4.3. The choice (4.4) is not the only one possible for the case with $\Sigma_{zz} = H_{ww} = 0$. However, the other choices (including "the most symmetric one", namely $\Sigma = H = 1$, $F = z - 1$, $G = 1 - w$) lead to metrics of more complicated form.

Lemma 4.2. *The following statements are equivalent*

- (i) $\Sigma_{zz} = 0$
- (ii) $C_{11\dot{A}\dot{B}} = 0$

Proof. If $\Sigma_{zz} = 0$ then (4.4) holds true and $\phi = x$. Thus, $-\phi C_{11\dot{A}\dot{B}} = \partial_{\dot{A}}\partial_{\dot{B}}x = \partial_{\dot{A}}(\delta_{\dot{B}}^{\dot{1}}) = 0$. It proves implication (i) \implies (ii).

If $C_{11\dot{A}\dot{B}} = 0$ then ϕ takes the form $\phi = J_{\dot{A}}p^{\dot{A}} + \kappa$ where $J_{\dot{A}} = J_{\dot{A}}(q, p) \neq 0$ and $\kappa = \kappa(q, p)$ are arbitrary functions. $J_{\dot{A}}$ must be nonzero, otherwise SD \mathcal{C} is nonexpanding what is a contradiction. From (2.14) we find the transformation formulas for $J_{\dot{A}}$ and κ

$$J'_{\dot{A}} = \lambda^{\frac{1}{2}} J_{\dot{B}} \frac{\partial q'_{\dot{A}}}{\partial q_{\dot{B}}}, \quad \kappa' = \lambda^{-\frac{1}{2}} \kappa - J'_{\dot{A}} \sigma^{\dot{A}} \quad (4.7)$$

Hence, one puts $J_1 = 1$, $J_2 = 0 = \kappa$ without any loss of generality. It implies $\phi = x$. It corresponds to the solution $\phi = F(x - \Sigma_z)$ with $F = 1$ and $\Sigma_z = 0$. Thus, the proof of implication (ii) \implies (i) is completed. ■

From Lemma (4.2) it follows that the case $\Sigma_{zz} = H_{ww} = 0$ (\implies (4.4)) is not a general case because it implies $C_{11\dot{A}\dot{B}} = 0$. Hence, the traceless Ricci tensor is not general anymore. Thus, (4.4) does not lead to the general metric of para-Kähler, type $\{[\text{deg}]^e \otimes [\text{D}]^{nn}, [++, ++]\}$ spaces. However, the choice (4.4) will be treated as a starting point for para-Kähler, Einstein spaces of the types $\{[\text{deg}]^e \otimes [\text{D}]^{nn}, [++, ++]\}$ (see Section 4.2).

In the rest of this Section we assume $\Sigma_{zz} \neq 0$, $H_{ww} \neq 0$.

Lemma 4.3. Assume $\Sigma_{zz} \neq 0$, $F_{zz} \neq 0$, $G_{ww} \neq 0$ and $H_{ww} \neq 0$. Then functions Σ , F , G and H can be brought to the form

$$\Sigma = F = A(q, p) z^{\frac{1}{2}}, \quad -G = H = A(q, p) w^{\frac{1}{2}} \quad (4.8)$$

without any loss of generality. $A = A(p, q)$ is an arbitrary nonzero function.

Proof. If $\Sigma_{zz} \neq 0$ and $H_{ww} \neq 0$ then from (4.3b) it follows that $\Sigma_{zz} = \tilde{\alpha}/F^3$, $H_{ww} = -\tilde{\alpha}/G^3$, where $\tilde{\alpha} = \tilde{\alpha}(q, p)$ is an arbitrary nonzero function. Eq. (4.3a) reduces to $G^3 G_{ww} = F^3 F_{zz} =: \tilde{\beta}(q, p)$ where $\tilde{\beta}$ is an arbitrary nonzero function. Hence, $\Sigma_{zz} = a F_{zz}$ and $H_{ww} = -a G_{ww}$ where $a := \tilde{\alpha}/\tilde{\beta}$. Thus

$$\Sigma = aF + bz + c, \quad H = -aG + mw + n \quad (4.9)$$

where $a \neq 0$, b , c , m and n are functions of (q, p) . Inserting (4.9) into consistency condition (4.1) one arrives at the equation

$$\begin{aligned} w(-aF^2 + mF - cF + zaFF_z) + nF - aFF_z - bF = \\ = z(-bG - aG^2 + nG + awGG_w) - cG - aGG_w + mG \end{aligned} \quad (4.10)$$

$\partial_w^2(4.10)$ implies $m = c$ and $\partial_z^2(4.10)$ gives $n = b$. Inserting these conditions into (4.10) and remembering that $a \neq 0$ we get

$$w(zFF_z - F^2) - FF_z = z(wGG_w - G^2) - GG_w \quad (4.11)$$

From (4.11) it follows that $\partial_z^3(F^2) = 0$ and $\partial_w^3(G^2) = 0$. Thus, $F^2 = Mz^2 + 2Az + N$ and $G^2 = Pw^2 + 2Bw + S$, where M , A , N , P , B and S are arbitrary functions of (q, p) . Feeding (4.11) with these forms of F^2 and G^2 one finds $B = A$, $P = N$ and $S = M$. Finally, we arrive at the solutions

$$\begin{aligned} F^2 &= Mz^2 + 2Az + N, \quad \Sigma = aF + bz + c, \\ G^2 &= Nw^2 + 2Aw + M, \quad H = -aG + cw + b \end{aligned} \quad (4.12)$$

Transformation formulas for a , b , c , M , A and N reads

$$\begin{aligned} a' &= \Delta^{-1} \lambda^{-\frac{3}{2}} a \\ b' &= \lambda^{-1} \Delta^{-1} (q'_q b + q'_p c) + \sigma^1 \\ c' &= \lambda^{-1} \Delta^{-1} (p'_q b + p'_p c) + \sigma^2 \\ M' &= \lambda (q_q'^2 M + 2q'_q q'_p A + q_p'^2 N) \\ N' &= \lambda (p_q'^2 M + 2p'_q p'_p A + p_p'^2 N) \\ A' &= \lambda (q'_q p'_q M + p'_p q'_p N + (q'_q p'_p + q'_p p'_q) A) \end{aligned} \quad (4.13)$$

Thorough analysis of (4.13) proves that one puts $a = 1$ and $b = c = M = N = 0$ without any loss of generality. Re-defining A , namely $\sqrt{2A} \rightarrow A$, we arrive at (4.8). \blacksquare

Remark 4.4. Note, that (4.8) implies

$$x = \frac{Aw^{\frac{1}{2}}}{1 + z^{\frac{1}{2}}w^{\frac{1}{2}}}, \quad y = \frac{Az^{\frac{1}{2}}}{1 + z^{\frac{1}{2}}w^{\frac{1}{2}}}, \quad \phi = -\frac{A^2}{2} \frac{1 - z^{\frac{1}{2}}w^{\frac{1}{2}}}{1 + z^{\frac{1}{2}}w^{\frac{1}{2}}} \quad (4.14)$$

Lemma 4.3 holds true if $F_{zz} \neq 0$ and $G_{ww} \neq 0$. The case $F_{zz} = 0$ and $G_{ww} = 0$ must be analyzed separately.

Lemma 4.4. *Assume $\Sigma_{zz} \neq 0$, $H_{ww} \neq 0$, $F_{zz} = 0$ and $G_{ww} = 0$. Then the functions F , Σ , H and G can be brought to the form*

$$F = 1, \quad G = -w, \quad \Sigma = Az^2, \quad H = \frac{A}{w} \quad (4.15)$$

without any loss of generality. $A = A(p, q)$ is an arbitrary nonzero function.

Proof. $F_{zz} = 0$ implies $F = a(q, p)z + b(q, p)$. From (2.55) it follows that without any loss of generality F can be brought to a constant value, namely $F = 1$. Eq. (4.3b) reduces to $G^3 H_{ww} = -\Sigma_{zz}$. Thus, Σ_{zz} is a function of (q, p) only so $\Sigma = A(q, p)z^2 + B(q, p)z + C(q, p)$, $A \neq 0$. From the formula (2.36) it follows that B and C can be gauged away, leaving us with $\Sigma = Az^2$. With these forms of F and Σ , solutions for G and H can be easily obtained from (4.1). ■

Remark 4.5. Note, that (4.15) implies

$$x = \frac{1 + zw}{w}A, \quad y = -\frac{z}{w}A, \quad \phi = \frac{1 - zw}{w}A \quad (4.16)$$

4.1.2 The second consistency condition

The next step is to solve the null string equations (2.24b) and (2.37b). Solutions of these equations are given by (2.30) (expressed in the coordinate system (q, p, x, z)) and (2.43) (expressed in the coordinate system (q, p, w, y)).

Vanishing of the expansions of \mathcal{C}_{m^A} and \mathcal{C}_{n^A} implies $M_2 = 0$ (2.31c) and $N_2 = 0$ (2.44c). Solution of Eq. (2.31c) is given by the formula (2.54) (compare the proof of Theorem 2.4). Solution of Eq. (2.44c) is obtained analogously. Thus

$$\mathcal{Q} + z\mathcal{A} = \mathfrak{t}, \quad \mathcal{Q} + w\mathcal{B} = \mathfrak{e} \quad (4.17)$$

\mathfrak{t} and \mathfrak{e} stand for abbreviations

$$\begin{aligned} \mathfrak{t} &:= T(x - \Sigma_z)^2 + \frac{1}{F}(\Omega F_z - zF_p + F_q)(x - \Sigma_z) + \frac{1}{2}(\Sigma_p - \Omega\Sigma_{zz} + z\Sigma_{zp} - \Sigma_{zq}) \\ \mathfrak{e} &:= E(y - H_w)^2 + \frac{1}{G}(RG_w + wG_q - G_p)(y - H_w) - \frac{1}{2}(RH_{ww} + H_q + wH_{wq} - H_{wp}) \end{aligned} \quad (4.18)$$

where $T = T(q, p, z)$, $E = E(q, p, w)$, $\Omega = \Omega(q, p, z)$ and $R = R(q, p, w)$. Combining (4.17) with (2.30) and (2.43) one arrives at the solutions for \mathcal{A} , \mathcal{B} and \mathcal{Q}

$$\begin{aligned} \mathcal{A} &= -w(\mathcal{Q} + \mathfrak{e}) + R(y - H_w) + H_p - wH_q \equiv \frac{1}{z}(\mathfrak{t} - \mathcal{Q}) \\ \mathcal{B} &= -z(\mathcal{Q} + \mathfrak{t}) + \Omega(x - \Sigma_z) + z\Sigma_p - \Sigma_q \equiv \frac{1}{w}(\mathfrak{e} - \mathcal{Q}) \\ (1 - wz)\mathcal{Q} &= \mathfrak{t} + zw\mathfrak{e} - z(R(y - H_w) + H_p - wH_q) \\ &= \mathfrak{e} + wz\mathfrak{t} - w(\Omega(x - \Sigma_z) + z\Sigma_p - \Sigma_q) \end{aligned} \quad (4.19)$$

Clearly, the second consistency condition reads

$$(1 - wz)(\mathfrak{t} - \mathfrak{e}) - z(R(y - H_w) + H_p - wH_q) + w(\Omega(x - \Sigma_z) + z\Sigma_p - \Sigma_q) = 0 \quad (4.20)$$

Lemma 4.5. Assume that (4.4) holds true. Then (4.19) and (4.20) imply

$$\begin{aligned}(1 - wz)^3 \mathcal{Q} &= zR - w^2 \Omega \\ (1 - wz)^3 \mathcal{A} &= -R + w^3 \Omega \\ (1 - wz)^3 \mathcal{B} &= -z^2 R + w\Omega + w(1 - wz)^2 V\end{aligned}\tag{4.21}$$

where $R = R(q, p, w)$, $\Omega = \Omega(q, p, z)$ and $V = V(q, p)$ are arbitrary functions.

Proof. With (4.4) and (4.6), formulas (4.18) simplify considerably and they yield

$$\mathfrak{t} = \frac{w^2 T}{(1 - zw)^2}, \quad \mathfrak{e} = \frac{E}{(1 - zw)^2} - \frac{R}{w(1 - zw)}\tag{4.22}$$

Consistency condition (4.20) gives

$$T + \Omega = \frac{1}{w^2} \left(E - \frac{1}{w} R \right) =: V(q, p) \implies T = V - \Omega, \quad E = w^2 V + \frac{1}{w} R\tag{4.23}$$

Thus, Ω , R and V remain as independent functions. Inserting (4.23) into (4.22) and then into (4.19) one gets

$$\begin{aligned}(1 - wz)^3 \mathcal{Q} &= z(R + w^3 V) - w^2(\Omega - V) \\ (1 - wz)^3 \mathcal{A} &= -(R + w^3 V) + w^3(\Omega - V) \\ (1 - wz)^3 \mathcal{B} &= -z^2(R + w^3 V) + w(\Omega - V) + w(1 - wz)^2 V\end{aligned}\tag{4.24}$$

Re-defining functions Ω and R according to $R + w^3 V \rightarrow R$, $\Omega - V \rightarrow \Omega$ one arrives at (4.21). ■

Lemma 4.6. Assume that (4.8) holds true. Then (4.19) and (4.20) imply

$$\begin{aligned}(1 + z^{\frac{1}{2}} w^{\frac{1}{2}})^2 (1 - wz) \mathcal{Q} &= \frac{1}{2} \Omega A w z^{-\frac{1}{2}} + \frac{1}{2} R A z w^{-\frac{1}{2}} + \frac{1}{4} V (1 + zw) (1 - z^{\frac{1}{2}} w^{\frac{1}{2}})^2 \\ &\quad + (2 + z^{\frac{1}{2}} w^{\frac{1}{2}} + 2wz) (1 - z^{\frac{1}{2}} w^{\frac{1}{2}}) (A_p z^{\frac{1}{2}} - A_q w^{\frac{1}{2}}) \\ z(1 + z^{\frac{1}{2}} w^{\frac{1}{2}})^2 (1 - wz) \mathcal{A} &= -\frac{1}{2} R A z w^{-\frac{1}{2}} - \frac{1}{2} \Omega A w^2 z^{\frac{1}{2}} - \frac{1}{2} V z w (1 - w^{\frac{1}{2}} z^{\frac{1}{2}})^2 \\ &\quad + 3A_q z w^{\frac{3}{2}} (1 - z^{\frac{1}{2}} w^{\frac{1}{2}}) + A_p z w^{\frac{1}{2}} (1 - z^{\frac{1}{2}} w^{\frac{1}{2}}) (1 - z^{\frac{1}{2}} w^{\frac{1}{2}} + zw) \\ w(1 + z^{\frac{1}{2}} w^{\frac{1}{2}})^2 (1 - wz) \mathcal{B} &= -\frac{1}{2} \Omega A w z^{-\frac{1}{2}} - \frac{1}{2} R A z^2 w^{\frac{1}{2}} - \frac{1}{2} V z w (1 - z^{\frac{1}{2}} w^{\frac{1}{2}})^2 \\ &\quad - 3A_p w z^{\frac{3}{2}} (1 - z^{\frac{1}{2}} w^{\frac{1}{2}}) - A_q w z^{\frac{1}{2}} (1 - z^{\frac{1}{2}} w^{\frac{1}{2}}) (1 - z^{\frac{1}{2}} w^{\frac{1}{2}} + zw)\end{aligned}\tag{4.25}$$

where $R = R(q, p, w)$, $\Omega = \Omega(q, p, z)$, $A = A(q, p) \neq 0$ and $V = V(q, p)$ are arbitrary functions.

Proof. We skip the proof due to its similarity to that of Lemma 4.5 (although it is much more tedious). ■

Lemma 4.7. Assume that (4.15) holds true. Then (4.19) and (4.20) imply

$$\begin{aligned}(1 - wz) \mathcal{Q} &= \frac{(1 + zw)(1 - zw)^2}{w^2} A^2 V - A\Omega - \frac{z}{w^2} A R - \frac{z(1 - zw)(2 + zw)}{2w} A_p \\ (1 - wz) \mathcal{A} &= -\frac{2(1 - zw)^2}{w} A^2 V + w A \Omega + \frac{1}{w^2} A R - (1 - wz) A_q + \frac{(1 - wz)(1 + 2zw)}{w} A_p \\ (1 - wz) \mathcal{B} &= -\frac{2z(1 - zw)^2}{w^2} A^2 V + \frac{1}{w} A \Omega + \frac{z^2}{w^2} A R + \frac{z^2(1 - zw)}{w} A_p\end{aligned}\tag{4.26}$$

where $R = R(q, p, w)$, $\Omega = \Omega(q, p, z)$, $A = A(q, p) \neq 0$ and $V = V(q, p)$ are arbitrary functions.

Proof. We skip the proof due to its similarity to that of Lemma 4.5 (although it is much more tedious). ■

4.1.3 The metrics

Theorem 4.1. *Let (\mathcal{M}, ds^2) be a complex (neutral) space of the type $\{[deg]^e \otimes [D]^{nn}, [++, ++]\}$ ($\{[deg]^e \otimes [D_r]^{nn}, [++, ++]\}$). Then there exist a local coordinate system (q, p, w, z) such that the metric takes the form*

$$\begin{aligned} \frac{1}{2}ds^2 = & \frac{1}{(1 - w^{\frac{1}{2}}z^{\frac{1}{2}})^2} \left\{ z^{-\frac{1}{2}}A(dq + wdp)dz - w^{-\frac{1}{2}}A(dp + zdq)dw \right. \\ & - \frac{1}{1 - wz} \left[R(dp + zdq)^2 + \Omega(dq + wdp)^2 \right. \\ & \quad \left. + (1 - w^{\frac{1}{2}}z^{\frac{1}{2}})^2 V(dp + zdq)(dq + wdp) \right. \\ & \quad \left. - (1 - w^{\frac{1}{2}}z^{\frac{1}{2}})(2z^{\frac{1}{2}}A_p - 2w^{\frac{1}{2}}A_q)(dp + zdq)(dq + wdp) \right] \left. \right\} \end{aligned} \quad (4.27)$$

or

$$\begin{aligned} \frac{1}{2}ds^2 = & \frac{1}{(1 - wz)^2} \left\{ wA(dq + wdp)dz - A(dp + zdq)dw \right. \\ & - \frac{1}{1 - wz} \left[R(dp + zdq)^2 + w\Omega(dq + wdp)^2 \right. \\ & \quad \left. + (1 - wz)^2 V(dp + zdq)(dq + wdp) \right. \\ & \quad \left. - (1 - wz)w(zA_p - A_q)(dp + zdq)(dq + wdp) \right] \left. \right\} \end{aligned} \quad (4.28)$$

where $R = R(q, p, w)$, $\Omega = \Omega(q, p, z)$, $V = V(q, p)$ and $A = A(q, p) \neq 0$ are arbitrary holomorphic (real smooth) functions.

Proof. Inserting (4.14) and (4.25) into (2.13) and re-defining the functions according to

$$\frac{2}{A^3} \rightarrow A, \quad \frac{4V}{2A^4} \rightarrow V, \quad \frac{2w^{-\frac{1}{2}}R}{A^3} \rightarrow R, \quad \frac{2z^{-\frac{1}{2}}\Omega}{A^3} \rightarrow \Omega \quad (4.29)$$

one arrives at (4.27).

Similarly, inserting (4.16) and (4.26) into (2.13) and re-defining the functions according to

$$-\frac{1}{A} \rightarrow A, \quad -\frac{R}{A} \rightarrow R, \quad -\frac{\Omega}{A} \rightarrow \Omega, \quad 2V \rightarrow V \quad (4.30)$$

one arrives at (4.28). ■

We end this section at this point. Detailed analysis of the metrics (4.27) and (4.28) will be presented elsewhere. The main question is whether the metrics (4.27) and (4.28) are essentially different or (4.28) is just a special case of (4.27). The answer to this question remains unknown and as long as it is unknown it does not make any sense to discuss the structure of the traceless Ricci tensor or Petrov-Penrose types of the corresponding spaces.

4.2 Einstein case

In this Section we focus on pKE spaces of the types $\{[\deg]^e \otimes [D]^{nn}, [++, ++]\}$. We have to impose the conditions $C_{AB\dot{A}\dot{B}} = 0$, $\mathcal{R} = -4\Lambda \neq 0$. The first step has been already done in Section 4.1 (Lemmas 4.1 and 4.2). Namely, we proved that the condition $C_{11\dot{A}\dot{B}} = 0$ implies (4.6) and (4.21).

The second step is to consider the conditions $C_{12\dot{A}\dot{B}} = 0$. After some straightforward but tedious work one finds

$$\begin{aligned} \frac{2}{y(1-y)}C_{12\dot{1}\dot{1}} &= \frac{4}{x(2y-1)}C_{12\dot{1}\dot{2}} = -\frac{2}{x^2}C_{12\dot{2}\dot{2}} = \alpha \\ \alpha &:= \Omega_{zz} + \frac{2w}{1-zw}\Omega_z + wR_{ww} - \frac{2(1-2zw)R_w}{1-zw} - \frac{6zR}{1-zw} \end{aligned} \quad (4.31)$$

The curvature scalar reads

$$-\frac{1}{2}(1-zw)\mathcal{R} = w\Omega_{zz} - w^2R_{ww} + 4wR_w - 6R \quad (4.32)$$

Equations $C_{12\dot{A}\dot{B}} = 0$ and $\mathcal{R} = -4\Lambda$ yield

$$(1-zw)(\Omega_{zz} + wR_{ww}) + 2w\Omega_z - 6zR - 2(1-2zw)R_w = 0 \quad (4.33a)$$

$$(1-zw)2\Lambda = w\Omega_{zz} - w^2R_{ww} + 4wR_w - 6R \quad (4.33b)$$

$\partial_z^2(4.33b)$ gives $\Omega_{zzzz} = 0$ and $\partial_w^2(4.33b)$ yields $R_{wwww} = 0$. Thus, Ω and R are third order polynomials in z and w , respectively. Further calculations are straightforward and they lead to the formulas

$$\Omega = -\frac{\Lambda}{3}z^3 + Bz^2 + Cz + M, \quad R = (M-A)w^3 + Cw^2 + Bw - \frac{\Lambda}{3} \quad (4.34)$$

where A , B , C and M are arbitrary functions of (q, p) . We leave the coordinate system (q, p, w, z) and we return to the hyperheavenly coordinate system (q, p, x, y) (compare (4.6)). Feeding (4.21) with (4.6) and (4.34) one arrives at the formulas

$$\begin{aligned} \mathcal{A} &= Ax^3 - Cx^2 - Bx(1-2y) + \frac{\Lambda}{3}(1-3y+3y^2) \\ \mathcal{Q} &= Ax^2y - By(1-y) - Mx^2 + \frac{\Lambda}{3}\frac{y(1-y)(1-2y)}{x} \\ \mathcal{B} &= Axy^2 - Cy(1-y) + Mx(1-2y) + \frac{\Lambda}{3}\frac{y^2(1-y)^2}{x^2} + Vx \end{aligned} \quad (4.35)$$

Equations $C_{22\dot{A}\dot{B}} = 0$ are much more tedious. To calculate $C_{22\dot{A}\dot{B}}$ we need quantities $\mathfrak{D}^{\dot{A}}Q_{\dot{A}\dot{B}}$. They read

$$\begin{aligned} \Xi_{\dot{1}} &:= x^{-2}\mathfrak{D}^{\dot{A}}Q_{\dot{A}\dot{1}} = \mathcal{B}_p + \mathcal{Q}_q - V(Cx^2 - \Lambda y(1-y)) \\ \Xi_{\dot{2}} &:= x^{-2}\mathfrak{D}^{\dot{A}}Q_{\dot{A}\dot{2}} = -\mathcal{Q}_p - \mathcal{A}_q + xV(2Bx + \Lambda(2y-1)) \end{aligned} \quad (4.36)$$

Finally we get

$$\begin{aligned} \frac{1}{x^3y(1-y)}C_{22\dot{1}\dot{1}} &= \frac{2}{x^4(2y-1)}C_{22\dot{1}\dot{2}} = -\frac{1}{x^5}C_{22\dot{2}\dot{2}} = \beta, \\ \beta &:= 2\Lambda V - 2B_q - C_p \end{aligned} \quad (4.37)$$

Conditions $C_{22\dot{A}\dot{B}} = 0$ imply the solution for V in terms of B and C

$$V = \frac{2B_q + C_p}{2\Lambda} \quad (4.38)$$

Inserting \mathcal{A} , \mathcal{Q} and \mathcal{B} ((4.35) with V given by (4.38)) into (2.9) one finds

$$\begin{aligned} C^{(3)} &= -4Ax^3 \\ x^{-5}C^{(2)} &= -4A_p y - 4A_q x + 4M_p + 2C_q + \frac{2}{\Lambda}B(2B_q + C_p) \\ \frac{1}{2}x^{-6}C^{(1)} &= \frac{\partial \Xi_{\dot{2}}}{\partial q} - \frac{\partial \Xi_{\dot{1}}}{\partial p} - \mathcal{A} \frac{\partial \Xi_{\dot{1}}}{\partial x} - \mathcal{B} \frac{\partial \Xi_{\dot{2}}}{\partial y} - \mathcal{Q} \left(\frac{\partial \Xi_{\dot{2}}}{\partial x} + \frac{\partial \Xi_{\dot{1}}}{\partial y} \right) \\ &\quad + (\mathcal{A}_x + \mathcal{Q}_y)\Xi_{\dot{1}} + (\mathcal{B}_y + \mathcal{Q}_x)\Xi_{\dot{2}} \end{aligned} \quad (4.39)$$

The ASD part of the conformal curvature read

$$\begin{aligned} -x^{-2}C_{iiii} &= \frac{2y^2(1-y)^2}{x^4}\Lambda \\ -x^{-2}C_{iii\dot{2}} &= \frac{(1-y)(2y^2-y)}{x^3}\Lambda \\ -x^{-2}C_{ii\dot{2}\dot{2}} &= \frac{6y^2-6y+1}{3x^2}\Lambda \\ -x^{-2}C_{i\dot{2}\dot{2}\dot{2}} &= \frac{1-2y}{x}\Lambda \\ -x^{-2}C_{\dot{2}\dot{2}\dot{2}\dot{2}} &= 2\Lambda \end{aligned} \quad (4.40)$$

Equivalently, (4.40) can be rewritten in the form

$$-x^{-2}C_{\dot{A}\dot{B}\dot{C}\dot{D}}\xi^{\dot{A}}\xi^{\dot{B}}\xi^{\dot{C}}\xi^{\dot{D}} = 2\Lambda \left(\xi - \frac{y}{x} \right)^2 \left(\xi - \frac{y-1}{x} \right)^2 = 2\Lambda(\xi + z)^2 \left(\xi + \frac{1}{w} \right)^2 \quad (4.41)$$

where $\xi^{\dot{A}} = [1, \xi]$ is a spinor. Formula (4.41) confirms that ASD conformal curvature is of the type [D] and necessarily $\Lambda \neq 0$. Otherwise ASD conformal curvature vanishes.

Transformation formulas for the functions A , B , C and M are essential for further considerations. The gauge (2.14) is restricted to the transformations such that (4.5) holds true. Finally one arrives at the formulas

$$\begin{aligned} f^{\frac{3}{2}}A' &= A \\ f^{\frac{1}{2}}B' &= B - \Lambda f^{-\frac{1}{2}}p'_q \\ fC' &= C + 2f^{-\frac{1}{2}}p'_q B - \Lambda f^{-1}p_q'^2 \\ fM' &= f^{-\frac{1}{2}}M + f^{-1}p_q'^2 B + f^{-1}p'_q C - \frac{\Lambda}{3}f^{-2}p_q'^3 \end{aligned} \quad (4.42)$$

Now we are ready to formulate the main theorem of this article.

Theorem 4.2. *Let (\mathcal{M}, ds^2) be an Einstein complex (neutral) space of the type $\{[deg]^e \otimes [D]^{nn}, [++, ++]\}$ ($\{[deg]^e \otimes [D_r]^{nn}, [++, ++]\}$). Then there exist a local coordinate system (q, p, x, y) such that the metric takes the form*

$$\begin{aligned} \frac{1}{2}ds^2 &= x^{-2} \left\{ dqdy - dpdx + \left(Ax^3 - Cx^2 - Bx(1-2y) + \frac{\Lambda}{3}(1-3y+3y^2) \right) dp^2 \right. \\ &\quad - 2 \left(Ax^2y - By(1-y) - Mx^2 + \frac{\Lambda}{3} \frac{y(1-y)(1-2y)}{x} \right) dqdp \\ &\quad \left. + \left(Axy^2 - Cy(1-y) + Mx(1-2y) + \frac{\Lambda}{3} \frac{y^2(1-y)^2}{x^2} + \frac{2B_q + C_p}{2\Lambda} x \right) dq^2 \right\} \end{aligned} \quad (4.43)$$

where $\Lambda \neq 0$ is a cosmological constant, $A = A(q, p)$, $B = B(q, p)$, $C = C(q, p)$ and $M = M(q, p)$ are arbitrary holomorphic (real smooth) functions.

Proof. Inserting (4.35) and (4.38) into (2.13) one arrives at (4.43). ■

As long as $A \neq 0$ and $2C^{(2)}C^{(2)} - 3C^{(3)}C^{(1)} \neq 0$ the metric (4.43) is of the type $\{[II]^e \otimes [D]^{nn}, [++, ++]\}$. As before, we skip an analysis of the type $[D]^{ee} \otimes [D]^{nn}$.

Theorem 4.3. *Let (\mathcal{M}, ds^2) be an Einstein complex (neutral) space of the type $\{[III]^e \otimes [D]^{nn}, [++, ++]\}$ ($\{[III_r]^e \otimes [D_r]^{nn}, [++, ++]\}$). Then there exist a local coordinate system (q, p, x, y) such that the metric takes the form*

$$\begin{aligned} \frac{1}{2}ds^2 = x^{-2} \Big\{ & dqdy - dpdx + \left(-Cx^2 - Bx(1-2y) + \frac{\Lambda}{3}(1-3y+3y^2) \right) dp^2 \\ & - 2 \left(-By(1-y) - Mx^2 + \frac{\Lambda}{3} \frac{y(1-y)(1-2y)}{x} \right) dqdp \\ & + \left(-Cy(1-y) + Mx(1-2y) + \frac{\Lambda}{3} \frac{y^2(1-y)^2}{x^2} + \frac{2B_q + C_p}{2\Lambda} x \right) dq^2 \Big\} \end{aligned} \quad (4.44)$$

where $\Lambda \neq 0$ is a cosmological constant, $B = B(q, p)$, $C = C(q, p)$ and $M = M(q, p)$ are arbitrary holomorphic (real smooth) functions such that

$$2M_p + C_q + \frac{1}{\Lambda} B(2B_q + C_p) \neq 0 \quad (4.45)$$

Proof. $C^{(3)} = 0$ implies $A = 0$ and $C^{(2)} \neq 0$ yields (4.45) (compare (4.39)). ■

Type $\{[N]^e \otimes [D]^{nn}, [++, ++]\}$ is a little more complicated. Eq. $C^{(2)} = 0$ implies

$$2M_p + C_q + \frac{1}{\Lambda} B(2B_q + C_p) = 0 \quad (4.46)$$

Coefficient $C^{(1)}$ takes relatively simple form

$$\frac{1}{2}x^{-7}C^{(1)} = -\frac{1}{2\Lambda}\partial_p^2(2B_q + C_p) - \partial_p(MB) + \frac{1}{3}C(B_q + 2C_p) + \frac{\Lambda}{3}M_q \quad (4.47)$$

Those, who prefer the hyperheavenly coordinate system must be satisfied with the metric (4.44) restricted to the conditions (4.46) and $C^{(1)} \neq 0$. Otherwise, one formulates

Theorem 4.4. *Let (\mathcal{M}, ds^2) be an Einstein complex (neutral) space of the type $\{[N]^e \otimes [D]^{nn}, [++, ++]\}$ ($\{[N_r]^e \otimes [D_r]^{nn}, [++, ++]\}$). Then there exist a local coordinate system (q, t, x, y) such that the metric takes the form*

$$\begin{aligned} \frac{1}{2}ds^2 = x^{-2} \Big\{ & dqdy - (P_t dt + P_q dq)dx \\ & + \left(- \left(t - \frac{B^2}{\Lambda} \right) x^2 - Bx(1-2y) + \frac{\Lambda}{3}(1-3y+3y^2) \right) (P_t dt + P_q dq)^2 \\ & - 2 \left(-By(1-y) - \left(\frac{N}{2} + \frac{B^3}{3\Lambda^2} \right) x^2 + \frac{\Lambda}{3} \frac{y(1-y)(1-2y)}{x} \right) dq(P_t dt + P_q dq) \\ & + \left[- \left(t - \frac{B^2}{\Lambda} \right) y(1-y) + \left(\frac{N}{2} + \frac{B^3}{3\Lambda^2} \right) x(1-2y) + \frac{\Lambda}{3} \frac{y^2(1-y)^2}{x^2} \right. \\ & \left. + \frac{1}{2\Lambda} \left(2B_q + \frac{1}{P_t} - \frac{2B_t}{P_t}(2P_q - N_t) \right) x \right] dq^2 \Big\} \end{aligned} \quad (4.48)$$

where $\Lambda \neq 0$ is a cosmological constant, $B := \Lambda(P_q - N_t)$; $N = N(q, t)$ and $P = P(q, t)$ are arbitrary holomorphic (real smooth) functions such that $P_t \neq 0$ and

$$\begin{aligned} & -\frac{1}{2\Lambda} \frac{1}{P_t} \frac{\partial}{\partial t} \left(\frac{1}{P_t} \frac{\partial}{\partial t} \left(2B_q - 2\frac{P_q}{P_t} B_t + \frac{1}{P_t} - \frac{2}{\Lambda} \frac{1}{P_t} B B_t \right) \right) - \frac{1}{2} \frac{1}{P_t} \partial_t (NB) \\ & + \frac{\Lambda}{6} \left(N_q - \frac{P_q}{P_t} N_t \right) - \frac{4t}{3\Lambda} \frac{1}{P_t} B B_t + \frac{2t}{3} \frac{1}{P_t} + \frac{t}{3} \left(B_q - \frac{P_q}{P_t} B_t \right) - \frac{2}{3\Lambda} \frac{1}{P_t} B^2 \neq 0 \end{aligned} \quad (4.49)$$

Proof. Substitute $2M =: N + \frac{2B^3}{3\Lambda^2}$ and $C =: t - \frac{B^2}{\Lambda}$ into (4.46) (N and t are new functions of (q, p)). Eq. (4.46) simplifies to the form

$$N_p + t_q + \frac{B}{\Lambda} t_p = 0 \quad (4.50)$$

Multiplying (4.50) by $dp \wedge dq$ and treating (q, t) as independent variables we find

$$\tilde{B}(q, t) = \Lambda(\tilde{P}_q - \tilde{N}_t), \quad N(q, p) =: \tilde{N}(q, t), \quad p =: \tilde{P}(q, t) \quad (4.51)$$

Note that

$$\frac{\partial B(q, p)}{\partial q} = \tilde{B}_q + \tilde{B}_t t_q, \quad \frac{\partial B(q, p)}{\partial p} = \tilde{B}_t t_p, \quad t_p = \frac{1}{\tilde{P}_t}, \quad t_q = -\frac{\tilde{P}_q}{\tilde{P}_t} \quad (4.52)$$

Similar formulas hold true for derivatives of $C(q, p)$ and $N(q, p)$. Inserting (4.51) and (4.52) into (4.44), passing to the coordinate system (q, t, x, y) and dropping tildes one arrives at (4.48).

$C^{(1)}$ transformed into the coordinate system (q, t, x, y) takes the form

$$\begin{aligned} \frac{1}{2} x^{-7} C^{(1)} = & -\frac{1}{2\Lambda} \frac{1}{P_t} \frac{\partial}{\partial t} \left(\frac{1}{P_t} \frac{\partial}{\partial t} \left(2B_q - 2\frac{P_q}{P_t} B_t + \frac{1}{P_t} - \frac{2}{\Lambda} \frac{1}{P_t} B B_t \right) \right) - \frac{1}{2} \frac{1}{P_t} \partial_t (NB) \\ & + \frac{\Lambda}{6} \left(N_q - \frac{P_q}{P_t} N_t \right) - \frac{4t}{3\Lambda} \frac{1}{P_t} B B_t + \frac{2t}{3} \frac{1}{P_t} + \frac{t}{3} \left(B_q - \frac{P_q}{P_t} B_t \right) - \frac{2}{3\Lambda} \frac{1}{P_t} B^2 \end{aligned} \quad (4.53)$$

(where we already dropped tildes over B , P and N). Because $C^{(1)} \neq 0$, (4.49) is proved. \blacksquare

5 Concluding remarks.

This article is the second part of the research programme devoted to para-Kähler, para-Kähler Einstein and Walker geometries. In this part we analyzed complex and real neutral, 4-dimensional spaces equipped with a single, expanding congruence of SD null strings. Thus, a weak and expanding \mathcal{HH} -space was a "starting point" for further considerations. Then we equipped a weak \mathcal{HH} -space with ASD congruences of null strings according to the following scheme: $\mathcal{C}^e \rightarrow \mathcal{C}^n \rightarrow \mathcal{C}^{ne} \rightarrow \mathcal{C}^{nn}$ (compare Scheme 1). Finally we arrived at pK-spaces. The path from weak \mathcal{HH} -spaces to pK-spaces is full of geometrically interesting spaces equipped with \mathcal{C} s and \mathcal{I} s of different properties. The metrics of all these spaces were found in all the generality. The results are gathered in the Table 1 (the metric marked by * has been already known, the rest of the metrics are new results).

Type	Metric	Functions in the metric
Spaces with $C_{ab} \neq 0$		
$[\text{deg}]^e \otimes [\text{any}]$	(2.13)*	4 functions of 4 variables
$\{[\text{deg}]^e \otimes [\text{any}]^e, [++]\}$	(2.49)	3 functions of 4 variables, 2 functions of 3 variables
$\{[\text{deg}]^e \otimes [\text{any}]^e, [+ -]\}$	(2.47)	3 functions of 4 variables, 1 function of 3 variables
$\{[\text{deg}]^e \otimes [\text{any}]^e, [- +]\}$	(2.49)	2 functions of 4 variables, 3 functions of 3 variables
$\{[\text{deg}]^e \otimes [\text{any}]^e, [--]\}$	(2.47)	2 functions of 4 variables, 2 functions of 3 variables
$\{[\text{deg}]^e \otimes [\text{any}]^n, [++]\}$	(2.52)	1 function of 4 variables, 4 functions of 3 variables
$\{[\text{deg}]^e \otimes [\text{any}]^n, [--]\}$	(2.51)	1 function of 4 variables, 3 functions of 3 variables
$\{[\text{deg}]^e \otimes [\text{deg}]^{ne}, [- -, ++]\}$	(2.57)	5 functions of 3 variables
$\{[\text{deg}]^e \otimes [\text{deg}]^{ne}, [- -, +-]\}$	(2.56)	4 functions of 3 variables
$\{[\text{deg}]^e \otimes [\text{deg}]^{ne}, [- -, - +]\}$	(2.60)	3 functions of 3 variables
$\{[\text{deg}]^e \otimes [\text{deg}]^{ne}, [++, ++]\}$	(2.65)	6 functions of 3 variables
$\{[\text{deg}]^e \otimes [\text{deg}]^{ne}, [++, - +]\}$	-	such spaces do not exist
$\{[\text{deg}]^e \otimes [\text{deg}]^{ne}, [++, +-]\}$	(2.61)	5 functions of 3 variables
$\{[\text{deg}]^e \otimes [\text{deg}]^{ne}, [++, --]\}$	(2.62)	3 functions of 3 variables

Table 1: Summary of main results.

The strongest result of this paper concerns pKE-spaces. We found all pKE metrics which are algebraically special (see Table 2). The most general space which belong to such class is a space of the type $\{[\text{II}]^e \otimes [\text{D}]^{nn}, [++, ++]\}$. The metric of such a space depends on 4 functions of 2 variables. The metrics of the types $\{[\text{III}, \text{N}]^e \otimes [\text{D}]^{nn}, [++, ++]\}$ also were found in all the generality.

Slightly less general sub-types of pKE-spaces are types $\{[\text{II}, \text{III}, \text{N}]^e \otimes [\text{D}]^{nn}, [++, --]\}$. Some of these types (namely, $\{[\text{III}, \text{N}]^e \otimes [\text{D}]^{nn}, [++, --]\}$) were found earlier in [8] (the metric (5.16) of [8]). However, we did not know it yet in [8] that the metric (5.16) is equipped with \mathcal{I} s with properties $[++, --]$. On the other hand, the metric of the type $\{[\text{II}]^e \otimes [\text{D}]^{nn}, [++, --]\}$ is a new result.

The only type of pKE-space which metric is still unknown⁸ is the type $[\text{I}] \otimes [\text{D}]^{nn}$. The third part of our work will be devoted to space of such a type.

⁸According to our best knowledge there is no even a single explicit example of such space.

Type	Metric	Functions in the metric
Spaces with $C_{ab} \neq 0$		
$\{[\text{II}]^e \otimes [\text{D}]^{nn}, [++, --]\}$	(3.10)	2 functions of 3 variables, one constant
$\{[\text{III}]^e \otimes [\text{D}]^{nn}, [++, --]\}$	(3.19)	5 functions of 2 variables
$\{[\text{N}]^e \otimes [\text{D}]^{nn}, [++, --]\}$	(3.21)	2 functions of 2 variables, 1 function of 1 variable, one constant
$\{[\text{II}]^e \otimes [\text{D}]^{nn}, [++, ++]\}$	(4.27) or (4.28)	2 functions of 3 variables, 2 functions of 2 variables
Spaces with $C_{ab} = 0$		
$\{[\text{II}]^e \otimes [\text{D}]^{nn}, [++, --]\}$	(3.31)	3 functions of 2 variables
$\{[\text{III}]^e \otimes [\text{D}]^{nn}, [++, --]\}$	(3.32)*	2 functions of 2 variables
$\{[\text{N}]^e \otimes [\text{D}]^{nn}, [++, --]\}$	(3.33)*	1 functions of 2 variables, one constant
$\{[\text{II}]^e \otimes [\text{D}]^{nn}, [++, ++]\}$	(4.43)	4 functions of 2 variables
$\{[\text{III}]^e \otimes [\text{D}]^{nn}, [++, ++]\}$	(4.44)	3 functions of 2 variables
$\{[\text{N}]^e \otimes [\text{D}]^{nn}, [++, ++]\}$	(4.48)	2 functions of 2 variables

Table 2: Summary of main results. Para-Kähler spaces.

References

- [1] An D. and Nurowski P., *Twistor space for rolling bodies*, Comm. Math. Phys., 326(2), 393-414 (2014)
- [2] Bor G., Hernández Lamonedá L. and Nurowski P., *The dancing metric, G2-symmetry and projective rolling*, Trans. Amer. Math. Soc., 370(6), 4433-4481 (2018)
- [3] Bor G., Makhmali O. and Nurowski P., *Para-Kähler-Einstein 4-manifolds and non-integrable twistor distributions*, Geometriae Dedicata 216, 9 (2022)
- [4] Boyer C.P., Finley III J.D. and Plebański J.F., *Complex general relativity, \mathcal{H} and \mathcal{HH} spaces - a survey to one approach*, General Relativity and Gravitation. Einstein Memorial Volume, ed. A. Held (Plenum, New York) vol.2. pp. 241-281 (1980)
- [5] Chudecki A., *Hyperheavenly spaces and their application in Walker and para-Kähler geometries: Part I*, Journal of Geometry and Physics 179, 104591
- [6] Chudecki A., *Classification of the traceless Ricci tensor in 4-dimensional pseudo-Riemannian spaces of neutral signature*, Acta Physica Polonica B, Vol. 48, No. 1, 53-74
- [7] Chudecki A., *On geometry of congruences of null strings in 4-dimensional complex and real pseudo-Riemannian spaces*, Journal of Mathematical Physics **58**, 112502, (2017)
- [8] Chudecki A., *On some examples of para-Hermite and para-Kähler Einstein spaces with $\Lambda \neq 0$* , Journal of Geometry and Physics 112, 175-196 (2017)
- [9] Chudecki A. and Przanowski M., *From hyperheavenly spaces to Walker and Osserman spaces I*, Classical and Quantum Gravity, **25**, 145010, (2008)

- [10] Finley III J.D. and Plebański J.F., *The intrinsic spinorial structure of hyperheavens*, Journal of Mathematical Physics, **17**, 2207 (1976)
- [11] Law P.R. and Matsushita Y., *A spinor approach to Walker geometry*, Communications of Mathematical Physics 282, 577–623 (2008)
- [12] Law P.R. and Matsushita Y., *Algebraically special, real alpha-geometries*, Journal of Geometry and Physics, 61, 2064–2080 (2011)
- [13] Law P.R. and Matsushita Y., *Real AlphaBeta-geometries and Walker geometry*, Journal of Geometry and Physics, 65, 35–44 (2013)
- [14] Penrose R., *A spinor approach to general relativity*, Ann. Phys. **10**, 171 (1960)
- [15] Plebański J.F., *Spinors, tetrads and forms*, unpublished monograph of Centro de Investigacion y Estudios Avanzados del IPN, Mexico 14 (1974)
- [16] Plebański J.F. and Hacyan S., *Null geodesic surfaces and Goldberg - Sachs theorem in complex Riemannian spaces*, J. Math. Phys., **16**, 2403 (1975)
- [17] Plebański J.F. and Przanowski M., *Null tetrad, spinor and helicity formalisms for all 4-dimensional Riemannian manifolds. I. Null tetrads and spinors*, Acta Phys. Polon., **B19** 805 (1988)
- [18] Plebański J.F. and Rózga K., *The optics of null strings*, J. Math. Phys., **25**, 1930 (1984)
- [19] Przanowski M. and Plebański J. F., *Generalized Goldberg-Sachs theorems in complex and real space-times. I.*, Acta Physica Polonica, Vol. B10 (1979)