

Vertex operator for generalised Kac–Moody algebras associated to the two-sphere and the two-torus

February 17, 2023

Rutwig Campoamor-Stursberg^{1*}, Michel Rausch de Traubenberg^{2†}

¹ Instituto de Matemática Interdisciplinar and Dpto. Geometría y Topología, UCM, E-28040 Madrid, Spain

³ Université de Strasbourg, CNRS, IPHC UMR7178, F-67037 Strasbourg Cedex, France

* Email: rutwig@ucm.es

† Email: Michel.Rausch@iphc.cnrs.fr

Abstract

We pursue our study of generalised Kac-Moody and Virasoro algebras defined on compact homogeneous manifolds. Extending the well-known Vertex operator in the case of the two-torus or the two-sphere, we obtain explicit bosonic realisations of the semi-direct product of the extension of Kac-Moody and Virasoro algebras on $\mathbb{S}^1 \times \mathbb{S}^1$ and \mathbb{S}^2 , respectively. As for the fermionic realisation previously constructed, in order to have well defined algebras, we introduce, beyond the usual normal ordering prescription, a regulator and regularise infinite sums by means of the Riemann ζ -function.

keywords: Kac-Moody; Virasoro algebras; bosons realization; regularisation.

PACS numbers: 02.20 Tw; 03.65 Fd; 11.25Hf

It is well known that Kac-Moody [1, 2, 3, 4, 5, 6] and Virasoro [7] algebras are intimately related to the one-dimensional sphere \mathbb{S}^1 . Recently, a class of infinite dimensional Lie algebras associated to compact real manifolds \mathcal{M} , with \mathcal{M} either a compact Lie group G_c or a compact homogeneous space G_c/H (H being a closed subgroup of G_c) has been studied [8]. Extensions of Kac-Moody algebras of this type have also been considered in [9, 10, 11] for $\mathcal{M} = \mathbb{S}^2$ and $\mathcal{M} = \mathbb{S}^1 \times \cdots \times \mathbb{S}^1$, as well as in [12, 13] (in the first of these articles, the algebra associated to $\mathbb{S}^1 \times \cdots \times \mathbb{S}^1$ is called a multi-loop algebra). The problem of the existence of central extensions, as well as their representation theory in the context of structural properties and representations of either G_c or \mathcal{M} were inspected in [8]. Following this analysis, it was realised in [14] that the Kac-Moody algebra and the corresponding Virasoro algebras associated to $\mathcal{M} = \mathbb{S}^2$ or $\mathcal{M} = \mathbb{S}^1 \times \mathbb{S}^1$ can indeed be constructed in a natural manner from the usual Kac-Moody and Virasoro algebras, a fact that enables us to derive an explicit fermion realisation.

In this note we pursue our study of Kac-Moody and Virasoro algebras associated to \mathbb{S}^2 and $\mathbb{S}^1 \times \mathbb{S}^1$ and derive an explicit boson realisation, which can be interpreted as a natural extension of the vertex construction [15, 16], along the lines of [14].

We first recall some properties of simply laced Lie algebras [4]. Let \mathfrak{g} be a compact simply laced Lie algebra of rank r and let Σ be the set of roots of \mathfrak{g} . Since \mathfrak{g} is simply laced, all roots have the same length and are normalised such that their square equals two. Furthermore, for any two roots α, β such that $\alpha + \beta$ is a root, we have $\alpha \cdot \beta = -1$. In the Cartan-Weyl basis $\mathfrak{g} = \{H^i, E^\alpha, i = 1, \dots, r, \alpha \in \Sigma\}$, the Lie brackets take the form

$$[H^i, H^j] = 0, \quad [H^i, E_\alpha] = \alpha^i E_\alpha, \quad [E_\alpha, E_\beta] = \begin{cases} \epsilon(\alpha, \beta) E_{\alpha+\beta} & \text{if } \alpha + \beta \in \Sigma \\ \alpha \cdot H & \text{if } \alpha + \beta = 0 \\ 0 & \text{otherwise} \end{cases} \quad (1)$$

where $\epsilon(\alpha, \beta) = \pm 1$. Let $\Lambda_R(\mathfrak{g})$ be the root lattice of \mathfrak{g} . As for all $\alpha, \beta \in \Lambda_R(\mathfrak{g})$, we have $\alpha \cdot \beta \in \mathbb{Z}$, and the root lattice is said to be an integer lattice. Since \mathfrak{g} is compact, the scalar product is Euclidean and the lattice is called Euclidean (with $\alpha \cdot \beta$ being the corresponding scalar product for $\alpha, \beta \in \Lambda_R(\mathfrak{g})$). The key point in the construction below, as shown by Frenkel and Kac [15] (see also [17]), is to extend this sign function $\epsilon(\alpha, \beta)$ for any $\alpha, \beta \in \Lambda_R(\mathfrak{g})$. This extension satisfies

$$\epsilon(\alpha, \beta) = (-1)^{\alpha \cdot \beta + \alpha \cdot \alpha \beta \cdot \beta} \epsilon(\beta, \alpha), \quad \epsilon(\alpha, \beta) \epsilon(\alpha + \beta, \gamma) = \epsilon(\alpha, \beta + \gamma) \epsilon(\beta, \gamma), \quad (2)$$

for any $\alpha, \beta \in \Lambda_R(\mathfrak{g})$. A lattice satisfying these requirements is called an ϵ -lattice. In fact, for any simply laced Lie algebra \mathfrak{g} , the root lattice $\Lambda_R(\mathfrak{g})$ is an ϵ -lattice (see *e.g.* [18] for details). The ϵ -function is not uniquely defined, and it admits a kind of \mathbb{Z}_2 -gauge invariance, as the new function

$$\epsilon'(\alpha, \beta) = \eta_\alpha \eta_\beta \eta_{\alpha+\beta} \epsilon(\alpha, \beta),$$

satisfies condition (2), where $\eta_\alpha, \eta_\beta, \eta_{\alpha+\beta} = \pm 1$. Fixing the gauge enables to chose an ϵ -function such that

$$\epsilon(0, \alpha) = \epsilon(\alpha, 0) = \epsilon(\alpha, -\alpha) = 1 \quad \epsilon(\alpha, \beta) = \epsilon(-\beta, -\alpha) = \epsilon(-\alpha, \alpha + \beta), \quad (3)$$

Details can be found for instance in [6].

Now we turn to the boson realisation of the Kac-Moody and Virasoro algebras associated to the two-sphere and the two-torus. We first introduce $r = \text{rk}(\mathfrak{g})$ scalar fields Q^i and their associated momenta P^i . Instead of parameterising the two-torus (resp. the two-sphere) by the angular variables $(\varphi_1, \varphi_2) \in [0, 2\pi[\times [0, 2\pi[$ (resp. $(\varphi, u) \in [0, 2\pi[\times [-1, 1]$ —with $u = \cos \theta$), we consider in our two-dimensional manifolds the local coordinates (z, φ) (resp. (z, u)) with $z \in \mathbb{C}$, as done in [14], in order to use techniques of Conformal Field Theory. With these notations, we introduce for the two-torus $\mathbb{S}^1 \times \mathbb{S}^1$

$$\begin{aligned} Q^i(z, \varphi) &= \sum_{n \in \mathbb{Z}} \left[q_n^i - i p_n^i \ln z + i \sum_{m > 0} \left(\frac{1}{m} \alpha_{m,n}^i z^{-m} - \frac{1}{m} \alpha_{-m,n}^i z^m \right) \right] e^{-in\varphi}, \\ &= q^i(\varphi) - i p^i(\varphi) \ln z + i \sum_{m > 0} \frac{1}{m} \left(\alpha_m^i(\varphi) z^{-m} - \alpha_{-m}^i(\varphi) z^m \right) \\ P^i(z, \varphi) &= iz \frac{dQ^i(z, \varphi)}{dz} = \sum_{n \in \mathbb{Z}} \left[p_n^i + \sum_{m > 0} \left(\alpha_{m,n}^i z^{-m} + \alpha_{-m,n}^i z^m \right) \right] e^{-in\varphi}, \\ &= p^i(\varphi) + \sum_{m > 0} \left(\alpha_m^i(\varphi) z^{-m} + \alpha_{-m}^i(\varphi) z^m \right) \end{aligned} \quad (4)$$

and for the two-sphere \mathbb{S}^2

$$\begin{aligned} Q^i(z, u) &= \sum_{\ell \geq 0} \left(q_\ell^i - i p_\ell^i \ln z \right) Q_{\ell 0}(u) + i \sum_{m > 0} \sum_{\ell \geq m} \frac{1}{m} \left(\alpha_{\ell m}^i z^{-m} - \alpha_{\ell-m}^i z^m \right) Q_{\ell, m}(u) \\ &= q^i(u) - i p^i(u) \ln z + i \sum_{m > 0} \frac{1}{m} \left(\alpha_m^i(u) z^{-m} - \alpha_{-m}^i(u) z^m \right) \\ P^i(z, u) &= iz \frac{dQ^i(z, u)}{dz} = \sum_{\ell \geq 0} p_\ell^i Q_{\ell 0}(u) + \sum_{m > 0} \sum_{\ell \geq m} \left(\alpha_{\ell m}^i z^{-m} + \alpha_{\ell-m}^i z^m \right) Q_{\ell, m}(u), \\ &= p^i(u) + \sum_{m > 0} \left(\alpha_m^i(u) z^{-m} + \alpha_{-m}^i(u) z^m \right) \end{aligned} \quad (5)$$

where $Q_{\ell m}$ are the normalised associated Legendre functions (see [14] for notations), and

$$\mathcal{B}_m = \left\{ Q_{\ell m}, \ell \geq |m| \right\}, \quad \forall m \in \mathbb{Z}, \quad (6)$$

constitutes a Hilbert basis for the square integrable functions on $[-1, 1]$. We observe that the second and fourth lines in (4) and (5) correspond to the usual mode expansions considered in the context of strings. In other words, we assume in our construction that the string modes $(\alpha_{m,p}^i$ and $q^i)$ have mode expansions in the φ -direction for the two-torus, and in the u -direction for the two-sphere. This is the key observation to obtain, from the usual Virasoro and Kac-Moody algebras, a corresponding analogue on $\mathbb{S}^1 \times \mathbb{S}^1$ and \mathbb{S}^2 . This mode expansion has also been considered in [14] to construct fermion realisations of generalised Kac-Moody and Virasoro algebras. Note finally that, in the case of the two-torus, the construction is dissymmetric in both directions. This is in fact a consequence of our ordering prescription (see (9) below), but is in accordance with unitarity of representations of generalised Kac-Moody algebras, as shown in [8].

We assume the quantisation relations

$$\begin{aligned} [\alpha_{m,p}^i, \alpha_{n,q}^j] &= m\delta^{ij}\delta_{m+n}\delta_{p+q}, & [p_m^i, q_n^j] &= -i\delta^{ij}\delta_{m+n} \quad \text{for } \mathbb{S}^1 \times \mathbb{S}^1 \\ [\alpha_{\ell,m}^i, \alpha_{\ell',m'}^j] &= m\delta^{ij}\delta_{\ell\ell'}\delta_{m+m'}, & [p_\ell^i, q_{\ell'}^j] &= -i\delta^{ij}\delta_{\ell\ell'} \quad \text{for } \mathbb{S}^2 \end{aligned} \quad (7)$$

as well as the reality conditions

$$\begin{aligned} (\alpha_{m,p}^i)^\dagger &= \alpha_{-m,-p}^i, & (p_m^i)^\dagger &= p_{-m}^i, & (q_m^i)^\dagger &= q_{-m}^i \quad \text{for } \mathbb{S}^1 \times \mathbb{S}^1 \\ (\alpha_{\ell,m}^i)^\dagger &= \alpha_{\ell,-m}^i, & (p_\ell^i)^\dagger &= p_\ell^i, & (q_\ell^i)^\dagger &= q_\ell^i \quad \text{for } \mathbb{S}^2. \end{aligned} \quad (8)$$

Note that in equations (7) and (8) we don't have the $(-1)^m$ coefficients for the two-sphere, as it is the case for fermions in [14]. This is simply due to the fact that in the expansion (5), we have substituted $Q_{\ell,m}(u) = (-1)^m Q_{\ell,-m}(u)$ by $(-1)^m Q_{\ell,-m}(u)$ for negative values of m , thus reabsorbing the sign factor. Defining the order relation

$$\begin{aligned} (m, p) > 0 &\iff m > 0 \quad \text{for } \mathbb{S}^1 \times \mathbb{S}^1 \\ (\ell, m) > 0 &\iff m > 0 \quad \text{for } \mathbb{S}^2 \end{aligned} \quad (9)$$

the annihilation operators are $\alpha_{m,p}^i, m > 0, p_p^i, \forall p \in \mathbb{Z}$ for the two-torus and $\alpha_{\ell,m}^i, m > 0, \ell \geq m, p_\ell^i, \ell \geq 0$ for the two-sphere. As usual in Quantum Field Theory, to avoid divergences we define the normal ordering prescription

$$\begin{aligned} \circ \alpha_{m,p}^i \alpha_{n,q}^j \circ &= \begin{cases} \alpha_{n,q}^j \alpha_{m,p}^i & m > 0 \quad \forall q \in \mathbb{Z} \\ \alpha_{m,p}^i \alpha_{n,q}^j & m < 0 \quad \forall q \in \mathbb{Z} \end{cases} & \circ p_m^i q_n^j \circ &= q_n^j p_m^i \quad \text{for } \mathbb{S}^1 \times \mathbb{S}^1 \\ \circ \alpha_{\ell,m}^i \alpha_{\ell',m'}^j \circ &= \begin{cases} \alpha_{\ell',m'}^j \alpha_{\ell,m}^i & m > 0 \quad \ell \geq |m| \\ \alpha_{\ell,m}^i \alpha_{\ell',m'}^j & m < 0 \quad \ell \geq |m| \end{cases} & \circ p_\ell^i q_{\ell'}^j \circ &= q_{\ell'}^j p_\ell^i \quad \text{for } \mathbb{S}^2. \end{aligned} \quad (10)$$

However, as already observed in [14], the normal ordering prescription (10) is still not enough to avoid divergences. To circumvent this difficulty and obtain an appropriate regularisation procedure, we define the ϵ -regularised bosons (with $\epsilon > 0$)

$$\begin{aligned} Q_\epsilon^i(z, \varphi) &= \sum_{n \in \mathbb{Z}} \left[q_n^i - ip_n^i \ln z + i \sum_{m > 0} \left(\frac{1}{m} \alpha_{m,n}^i z^{-m} - \frac{1}{m} \alpha_{-m,n}^i z^m \right) \right] e^{-in\varphi} e^{-\epsilon(|n| - \frac{1}{2})} \\ &\quad \text{for } \mathbb{S}^1 \times \mathbb{S}^1 \\ Q_\epsilon^i(z, u) &= \sum_{\ell \geq 0} \left(q_\ell^i - ip_\ell^i \ln z \right) Q_{\epsilon,\ell,0}(u) + i \sum_{m > 0} \sum_{\ell \geq m} \frac{1}{m} \left(\alpha_{\ell m}^i z^{-m} - \alpha_{\ell-m}^i z^m \right) Q_{\epsilon,\ell,m}(u) \\ &\quad \text{for } \mathbb{S}^2 \end{aligned} \quad (11)$$

with $Q_{\epsilon,\ell,m}$ the ϵ -regularised associated Legendre functions [14]. A completely analogous expression is obtained for the P_ϵ^i s. Associated to these ϵ -regularised bosons we define the functions

$$\begin{aligned} \delta_\epsilon(\theta - \varphi) &= \sum_{m \in \mathbb{Z}} e^{-im(\theta - \varphi)} e^{-2\epsilon(|m| - \frac{1}{2})} \quad \text{for } \mathbb{S}^1 \times \mathbb{S}^1 \\ \delta_\epsilon^m(u - v) &= \sum_{\ell \geq |m|} Q_{\epsilon,\ell,m}(u) Q_{\epsilon,\ell,m}(v) \quad \text{for } \mathbb{S}^2. \end{aligned} \quad (12)$$

When $\epsilon \rightarrow 0$, on the one hand the regularised bosons (11) reduce to either (4) or (5), while on the other hand, the functions (12) reduce to the usual δ -distributions on either $\mathbb{S}^1 \times \mathbb{S}^1$ or \mathbb{S}^2 (in the second case, the δ -distribution is associated to the basis (6)). However, as the generators of the Kac-Moody and the Virasoro algebras on the two-torus and the two-sphere are bilinear in the bosons (see below), the Wick theorem involves a double contraction, and in particular a term of the type $\delta_\epsilon(\varphi - \theta)^2$ in the case of the two-torus. This term is perfectly defined whilst the limit for $\epsilon \rightarrow 0$ is not defined. This situation has already been encountered in [14], when constructing a fermion realisation. We recall here the salient points of our regularisation procedure. In particular, to obtain a well-defined limit, we propose regularisation prescription. This prescription is done in two steps. In the first step, we substitute (in the case of the two-torus) $\delta_\epsilon(\theta - \varphi)^2$ by $\delta_{\epsilon_1}(\theta - \varphi)\delta_{\epsilon_2}(\theta - \varphi)$ and take the limit $\epsilon_1 \rightarrow 0, \epsilon_2 \rightarrow 0$ separately. This leads to

$$\lim_{\epsilon_1 \rightarrow 0} \delta_{\epsilon_1}(\theta - \varphi)\delta_{\epsilon_2}(\theta - \varphi) = \delta(\theta - \varphi)\delta_{\epsilon_2}(\theta - \varphi) = \delta(\theta - \varphi)\delta_{\epsilon_2}(0)$$

with $\delta_{\epsilon_2}(0)$ finite. In a second step, the divergent limit $\epsilon_2 \rightarrow 0$ for $\delta_{\epsilon_2}(0)$ is regularised using the ζ -regularisation (where ζ denotes the Riemann ζ -function) [14]. A similar regularisation process is valid for the two-sphere. In both cases, we introduce the regulated expressions [14]

$$\begin{aligned} \delta_{\text{reg}}(0) &= \lim_{\epsilon \rightarrow 0} \delta_\epsilon(0)|_{\text{reg}} = 1 \quad \text{for } \mathbb{S}^1 \times \mathbb{S}^1 \\ \delta_{\text{reg}}^m(0) &= \lim_{\epsilon \rightarrow 0} \delta_\epsilon^m(0)|_{\text{reg}} = 1 \quad \text{for } \mathbb{S}^2. \end{aligned} \quad (13)$$

Using the normal ordering prescription (10), the quantisation relation (7) and applying the Wick theorem (omitting the regular term — analogously, in all expressions below we shall omit the regular term), we obtain

$$\begin{aligned} Q_\epsilon^i(z, \theta)Q_\epsilon^i(w, \varphi) &= -\delta^{ij} \ln(z - w)\delta_\epsilon(\theta - \varphi), \quad |z| > |w| \\ P_\epsilon^i(z, \theta)Q_\epsilon^i(w, \varphi) &= -\frac{i\delta^{ij}z}{z - w}\delta_\epsilon(\theta - \varphi), \quad |z| > |w| \\ P_\epsilon^i(z, \theta)P_\epsilon^i(w, \varphi) &= \frac{\delta^{ij}zw}{(z - w)^2}\delta_\epsilon(\theta - \varphi), \quad |z| > |w| \end{aligned} \quad (14)$$

for the two-torus, while for the two-sphere we have

$$\begin{aligned} Q_\epsilon^i(z, u)Q_\epsilon^i(w, v) &= \delta^{ij}(-\ln z\delta_\epsilon^0(u - v) + \sum_{m>0} \sum_{\ell \geq m} \frac{1}{m} \left(\frac{w}{z}\right)^m \delta_\epsilon^m(u - v)) \\ &\equiv -\delta^{ij}L_\epsilon(z, w, u, v), \quad |z| > |w| \\ P_\epsilon^i(z, u)Q_\epsilon^i(w, v) &= -i\delta^{ij} \sum_{m \geq 0} \left(\frac{w}{z}\right)^m \delta_\epsilon^m(u - v) \\ &\equiv -i\delta^{ij}\Delta_\epsilon(z, w, u, v), \quad |z| > |w| \\ P_\epsilon^i(z, u)P_\epsilon^i(w, v) &= \delta^{ij} \sum_{m \geq 0} m \left(\frac{w}{z}\right)^m \delta_\epsilon^m(u - v). \quad |z| > |w| \end{aligned} \quad (15)$$

In the limit $\epsilon \rightarrow 0$, the equations (14) and (15) are not singular, and in both cases we obtain

$$\begin{aligned} Q^i(z, x)Q^i(w, y) &= -\delta^{ij} \ln(z - w)\delta(x - y), \quad |z| > |w| \\ P^i(z, x)Q^i(w, y) &= -\frac{i\delta^{ij}z}{z - w}\delta(x - y), \quad |z| > |w| \\ P^i(z, x)P^i(w, y) &= \frac{\delta^{ij}zw}{(z - w)^2}\delta(x - y), \quad |z| > |w| \end{aligned} \quad (16)$$

where $(x, y) = (\varphi, \theta)$ for $\mathbb{S}^1 \times \mathbb{S}^1$ and $(x, y) = (u, v)$ for \mathbb{S}^2 . The Hilbert space associated to the bosons (4) and (5) is built from the vacuum state $|0\rangle$ defined by

$$\begin{aligned} \alpha_{m,n}^i|0\rangle &= 0, \quad m > 0, \quad p_n^i|0\rangle = 0, \quad \forall n \in \mathbb{Z} \quad \text{for } \mathbb{S}^1 \times \mathbb{S}^1 \\ \alpha_{\ell,m}^i|0\rangle &= 0, \quad m > 0, \quad p_\ell^i|0\rangle = 0, \quad \forall \ell \geq 0 \quad \text{for } \mathbb{S}^2 \end{aligned} \quad (17)$$

Introduce now $\lambda \in \Lambda_W(\mathfrak{g})$ the weight lattice of \mathfrak{g} ($\Lambda_\Sigma(\mathfrak{g}) \subset \Lambda_W(\mathfrak{g})$) and define

$$\begin{aligned} |\lambda, \varphi\rangle &= e^{i\lambda \cdot q(\varphi)} |0\rangle, & q^i(\varphi) &= \sum_{n \in \mathbb{Z}} q_n^i e^{-in\varphi} & \text{for } \mathbb{S}^1 \times \mathbb{S}^1 \\ |\lambda, u\rangle &= e^{i\lambda \cdot q(u)} |0\rangle, & q^i(u) &= \sum_{\ell \in \mathbb{N}} q_\ell^i Q_{\ell,0}(u) & \text{for } \mathbb{S}^2 \end{aligned} \quad (18)$$

which are normalised as follows

$$\begin{aligned} \langle \lambda, \varphi | \mu, \theta \rangle &= \delta_{\mu\lambda} \delta(\varphi - \theta) & \text{for } \mathbb{S}^1 \times \mathbb{S}^1 \\ \langle \lambda, u | \mu, v \rangle &= \delta_{\mu\lambda} \delta(u - v) & \text{for } \mathbb{S}^2. \end{aligned}$$

We introduce the vertex operator

$$U_\epsilon^\alpha(z, x) = z^{\alpha \cdot \alpha / 2} \circ e^{i\alpha \cdot Q_\epsilon(z, x)} \circ \quad (19)$$

with $x = \varphi$ for $\mathbb{S}^1 \times \mathbb{S}^1$ and $x = u$ for \mathbb{S}^2 . The limit $\epsilon \rightarrow 0$ of (19) is well defined. This procedure constitutes a direct generalisation of the Vertex operator considered in string theory [19, 20, 21, 6]. Acting on the states (18), the vertex operator is single valued, as λ belongs to the weight lattice (see *e.g.* [6]). Again, using the Wick theorem, we obtain

$$\begin{aligned} P_\epsilon^i(z, \varphi) U_\epsilon^\alpha(w, \theta) &= \alpha^i \frac{z}{z-w} \delta_\epsilon(\varphi - \theta) U_\epsilon^\alpha(w, \theta) & \text{for } \mathbb{S}^1 \times \mathbb{S}^1 \\ P_\epsilon^i(z, u) U_\epsilon^\alpha(w, v) &= \alpha^i \Delta_\epsilon(z, w, u, v) U_\epsilon^\alpha(w, v) & \text{for } \mathbb{S}^2 \end{aligned} \quad |z| > |w|. \quad (20)$$

In the limit $\epsilon \rightarrow 0$, both relations above are regular and lead to

$$P^i(z, x) U^\alpha(w, y) = \alpha^i \frac{z}{z-w} \delta(x - y) U^\alpha(w, y), \quad |z| > |w| \quad (21)$$

where $(x, y) = (\theta, \varphi)$ for $\mathbb{S}^1 \times \mathbb{S}^1$ and $(x, y) = (u, v)$ for \mathbb{S}^2 . Finally, pushing the creation operators of U_ϵ^β to the left of the annihilation operators of U_ϵ^α , using (10), we obtain (for $|z| > |w|$)

$$\begin{aligned} U_\epsilon^\alpha(z, \theta) U_\epsilon^\beta(w, \varphi) &= \circ (z-w)^{\alpha \cdot \beta \delta_\epsilon(\theta - \varphi)} e^{i\alpha \cdot Q_\epsilon(z, \theta) + i\beta \cdot Q_\epsilon(w, \varphi)} \circ & \text{for } \mathbb{S}^1 \times \mathbb{S}^1 \\ U_\epsilon^\alpha(z, u) U_\epsilon^\beta(w, v) &= \circ e^{\alpha \cdot \beta L_\epsilon(z, w, u, v)} e^{i\alpha \cdot Q_\epsilon(z, \theta) + i\beta \cdot Q_\epsilon(w, \varphi)} \circ & \text{for } \mathbb{S}^2 \end{aligned}$$

It is known that, in the construction of the usual Kac-Moody algebras, the Vertex operator has to be corrected by a two-cocycle (see *e.g.* [6] for the case of usual Kac-Moody algebras). Analogously, we define (associated to the vacuum specified by the momenta $\lambda \in \Lambda_W(\mathfrak{g})$, see (18))

$$c_\alpha(x) = \sum_{\beta \in \Lambda_R(\mathfrak{g})} \epsilon(\alpha, \beta) |\beta + \lambda, x\rangle \langle \beta + \lambda, x|$$

with $x = \varphi$ for $\mathbb{S}^1 \times \mathbb{S}^1$ and $x = u$ for \mathbb{S}^2 . Next we introduce

$$\hat{c}_\alpha(x) = e^{iq(x) \cdot \alpha} c_\alpha(x) = \sum_{\beta \in \Lambda_R(\mathfrak{g})} \epsilon(\alpha, \beta) |\alpha + \beta + \lambda, x\rangle \langle \beta + \lambda, x| \quad (22)$$

where $q^i(\varphi)$ and $q^i(u)$ are defined in (18). A direct computation using (2) shows that

$$\hat{c}_\alpha(x) \hat{c}_\beta(y) = (-1)^{\alpha \cdot \beta + \alpha \cdot \alpha \cdot \beta \cdot \beta} \hat{c}_\beta(y) \hat{c}_\alpha(x) = \epsilon(\alpha, \beta) \hat{c}_{\alpha+\beta}(y) \delta(x - y) \quad (23)$$

and

$$\hat{c}_\alpha^\dagger(x) = \hat{c}_{-\alpha}(x).$$

If we now define $\hat{U}_\epsilon^\alpha(z, x) = U_\epsilon^\alpha(z, x) c_\alpha(x)$ and regularise $\hat{U}_\epsilon^\alpha(z, x) \hat{U}_\epsilon^\beta(w, y)$, we obtain for the two-torus (for $|z| > |w|$)

$$\begin{aligned} \hat{U}_\epsilon^\alpha(z, \theta) \hat{U}_\epsilon^\beta(w, \varphi) &= \epsilon(\alpha, \beta) \circ (z-w)^{\alpha \cdot \beta \delta_\epsilon(\theta - \varphi)} e^{i\alpha \cdot Q_\epsilon(z, \theta) + i\beta \cdot Q_\epsilon(w, \varphi)} \circ \\ &\quad \hat{c}_{\alpha+\beta}(\varphi) \delta(\theta - \varphi) \\ &= \epsilon(\alpha, \beta) \circ (z-w)^{\alpha \cdot \beta \delta_\epsilon(0)} e^{i\alpha \cdot Q_\epsilon(z, \theta) + i\beta \cdot Q_\epsilon(w, \varphi)} \circ \\ &\quad \hat{c}_{\alpha+\beta}(\varphi) \delta(\theta - \varphi) \\ &\quad \downarrow \text{reg} \\ \hat{U}^\alpha(z, \theta) \hat{U}^\beta(w, \varphi) &= \epsilon(\alpha, \beta) (z-w)^{\alpha \cdot \beta} \circ e^{i\alpha \cdot Q(z, \theta) + i\beta \cdot Q(w, \varphi)} \circ \hat{c}_{\alpha+\beta}(\varphi) \delta(\theta - \varphi), \end{aligned} \quad (24)$$

For the two-sphere, a similar procedure leads to (for $z| > |w|$)

$$\hat{U}^\alpha(z, u) \hat{U}^\beta(z, v) = \epsilon(\alpha, \beta) (z - w)^{\alpha \cdot \beta} \circ e^{i\alpha \cdot Q(z, x) + i\beta \cdot Q(w, y)} \circ \hat{c}_{\alpha+\beta}(v) \delta(u - v) . \quad (25)$$

Assuming the mode expansion (here we change slightly the conventions with respect to (5) since the expansion for \mathbb{S}^2 involves $Q_{\ell, m}$ and not $Q_{\ell, -m}$ for negative values of m)

$$\left. \begin{aligned} P^i(z, \varphi) &= \sum_{m \in \mathbb{Z}} \sum_{p \in \mathbb{Z}} H_{m, p}^i z^{-m} e^{-ip\varphi} \\ \hat{U}^\alpha(z, \varphi) &= \sum_{m \in \mathbb{Z}} \sum_{p \in \mathbb{Z}} E_{\alpha, m, p} z^{-m} e^{-ip\varphi} \end{aligned} \right\} \text{ for } \mathbb{S}^1 \times \mathbb{S}^1$$

$$\left. \begin{aligned} P^i(z, u) &= \sum_{m \in \mathbb{Z}} \sum_{\ell \geq |m|} H_{m, p}^i z^{-m} Q_{\ell m}(u) \\ \hat{U}^\alpha(z, u) &= \sum_{m \in \mathbb{Z}} \sum_{\ell \geq |m|} E_{\alpha, \ell, m} z^{-m} Q_{\ell m}(u) \end{aligned} \right\} \text{ for } \mathbb{S}^2 \quad (26)$$

and extracting the modes

$$\left. \begin{aligned} H_{m, p}^i &= \oint \frac{dz}{2i\pi} z^m \int_0^{2\pi} \frac{d\varphi}{2\pi} e^{ip\varphi} P^i(z, \varphi) \\ E_{\alpha, m, p} &= \oint \frac{dz}{2i\pi} z^m \int_0^{2\pi} \frac{d\varphi}{2\pi} e^{ip\varphi} \hat{U}^\alpha(z, \varphi) \end{aligned} \right\} \text{ for } \mathbb{S}^1 \times \mathbb{S}^1$$

$$\left. \begin{aligned} H_{\ell, m}^i &= \oint \frac{dz}{2i\pi} z^m \int_{-1}^1 \frac{du}{2} P^i(z, u) Q_{\ell m}(u) \\ E_{\alpha, \ell, m} &= \oint \frac{dz}{2i\pi} z^m \int_{-1}^1 \frac{du}{2} \hat{U}^\alpha(z, u) Q_{\ell m}(u) \end{aligned} \right\} \text{ for } \mathbb{S}^2 \quad (27)$$

Applying standard techniques of Conformal Field Theory, *i.e.*, integration in the complex z - and w -planes with adapted contours of integration (see *e.g.* [6, 22]) with the φ and θ or the u and v integrations being trivial, the equations (16), (21), (24) and (25) lead to

$$\begin{aligned} [H_{m_1, m_2}^i, H_{n_1, n_2}^j] &= m_1 \delta^{ij} \delta_{m_1+n_1} \delta_{m_2+n_2} , \\ [H_{m_1, m_2}^i, E_{\alpha, n_1, n_2}] &= \alpha^i E_{\alpha, m_1+n_1, m_2+n_2} , \\ [E_{\alpha, m_1, m_2}, E_{\beta, n_1, n_2}] &= \begin{cases} \epsilon(\alpha, \beta) E_{\alpha+\beta, m_1+n_1, m_2+n_2} & \text{if } \alpha + \beta \in \Sigma \\ \alpha \cdot H_{m_1+n_1, m_2+n_2} + m_1 \delta_{m_1+n_1} \delta_{m_2+n_2} & \text{if } \alpha + \beta = 0 \\ 0 & \text{otherwise} \end{cases} \end{aligned} \quad (28)$$

for the two-torus, and to

$$\begin{aligned} [H_{\ell_1, m_1}^i, H_{\ell_2, m_2}^j] &= (-1)^{m_1} m_1 \delta^{ij} \delta_{\ell_1, \ell_2} \delta_{m_1+m_2} , \\ [H_{\ell_1, m_1}^i, E_{\alpha, \ell_2, m_2}] &= c_{\ell_1, m_1, \ell_2, m_2}^{\ell_3, m_3} \alpha^i E_{\alpha, \ell_3, m_3} , \\ [E_{\alpha, \ell_1, m_1}, E_{\beta, \ell_2, m_2}] &= \begin{cases} \epsilon(\alpha, \beta) c_{\ell_1, m_1, \ell_2, m_2}^{\ell_3, m_3} E_{\alpha+\beta, \ell_3, m_3} & \text{if } \alpha + \beta \in \Sigma \\ c_{\ell_1, m_1, \ell_2, m_2}^{\ell_3, m_3} \alpha \cdot H_{\ell_3, m_3} + (-1)^{m_1} m_1 \delta_{\ell_1, \ell_2} \delta_{m_1+m_2} & \text{if } \alpha + \beta = 0 \\ 0 & \text{otherwise} \end{cases} \end{aligned} \quad (29)$$

for the two-sphere, where the coefficients $c_{\ell_1, m_1, \ell_2, m_2}^{\ell_3, m_3}$ (related to the Clebsch-Gordan coefficients) are given in [8]. We have thus obtained an explicit bosonic realisation of the Kac-Moody algebras associated to the two-torus or the two-sphere with a central charge $k = 1$ defined in [8]. We emphasise that, as stated

previously, the Kac-Moody algebra associated to the two-torus is dissymmetric in both directions. This is a direct consequence of our construction procedure (see the ordering relation (10)).

We now introduce the generators of the Virasoro algebra associated to the free bosons on the two-sphere and the two-torus

$$\begin{aligned} T_\epsilon(z, \varphi) &= \frac{1}{2} \sum_{i=1}^r \circ P_\epsilon^i(z, \varphi) P_\epsilon^i(z, \varphi) \circ & \text{for } \mathbb{S}^1 \times \mathbb{S}^1 \\ T_\epsilon(z, u) &= \frac{1}{2} \sum_{i=1}^r \circ P_\epsilon^i(z, u) P_\epsilon^i(z, u) \circ & \text{for } \mathbb{S}^2 \end{aligned} \quad (30)$$

and define the mode expansion, when $\epsilon \rightarrow 0$, by

$$\begin{aligned} T(z, \varphi) &= \sum_{n \in \mathbb{Z}} \sum_{p \in \mathbb{Z}} L_{m,p} z^{-n} e^{-ip\varphi} & \text{for } \mathbb{S}^1 \times \mathbb{S}^1 \\ T(z, u) &= \sum_{m \in \mathbb{Z}} \sum_{\ell \geq |m|} L_{\ell,m} z^{-m} Q_{\ell m}(u) & \text{for } \mathbb{S}^2 \end{aligned} \quad (31)$$

Computing the products $T_\epsilon T_\epsilon$ and using the Wick theorem, we proceed to consider all possible contractions, where only the double contractions lead to divergences. The latter are regularised by the procedure described above. After regularisation, performing all possible contractions with omission of the regular part, we obtain

$$\begin{aligned} T(z, \varphi) T(w, \theta) &= \frac{r}{2} \frac{(zw)^2}{(z-w)^4} \delta(\varphi - \theta) + \frac{zw}{(z-w)^2} 2T(w, \varphi) \delta(\varphi - \theta) \\ &\quad + \frac{z}{z-w} w \partial_w T(w, \varphi) \delta(\varphi - \theta) , \quad |z| > |w| \end{aligned}$$

for the two-torus, and

$$\begin{aligned} T(z, u) T(w, v) &= \frac{r}{2} \frac{(zw)^2}{(z-w)^4} \delta(u - v) + \frac{zw}{(z-w)^2} 2T(w, v) \delta(u - v) \\ &\quad + \frac{z}{z-w} w \partial_w T(w, v) \delta(u - v) , \quad |z| > |w| \end{aligned}$$

for the two-sphere. Again using adapted contours of integration in the z - and w - plane (see *e.g.* [6, 22]), the mode expansion (30) leads to

$$\begin{aligned} [L_{m,p}, L_{n,q}] &= (m-n) L_{n+m,p+q} + \frac{r}{12} m(m^2-1) \delta_{m+n} \delta_{p+q} \\ &\quad \text{for } \mathbb{S}^1 \times \mathbb{S}^1 \\ [L_{\ell_1, m_1}, L_{\ell_2, m_2}] &= (m_1 - m_2) c_{\ell_1, m_1, \ell_2, m_2}^{\ell_3, m_1+m_2} L_{\ell_3, m_1+m_2} \\ &\quad + (-1)^{m_1} \frac{r}{12} m_1(m_1^2-1) \delta_{\ell_1 \ell_2} \delta_{m_1+m_2} \quad \text{for } \mathbb{S}^2 . \end{aligned}$$

Thus we have obtained an explicit bosonic realisation of the Virasoro algebras associated to the two-torus or the two-sphere with a central charge $c = r$ (the number of bosons, *i.e.*, the rank of \mathfrak{g}). This construction constitutes a the direct extension of that performed on the circle \mathbb{S}^1 , but adapted to the two-dimensional manifolds $\mathbb{S}^1 \times \mathbb{S}^1$ and \mathbb{S}^2 .

In order to derive the action of the Virasoro algebra on the Kac-Moody algebra (computing $T_\epsilon P_\epsilon$ and $T_\epsilon U_\epsilon$) for the two-sphere and the two-torus, we apply again the Wick theorem and the regularisation procedure (only $T_\epsilon U_\epsilon$ needs to be regularised). After the mentioned regularisation, a routine computation leads to

$$\begin{aligned} T(z, x) P^i(w, y) &= \left(\frac{zw}{(z-w)^2} P^i(w, y) + \frac{z}{z-w} w \partial_w P^i(w, y) \right) \delta(x - y) , \quad |z| > |w| \\ T(z, x) U^\alpha(w, y) &= \left(\frac{z^2}{(z-w)^2} U^\alpha(w, y) + \frac{z}{z-w} w \partial_w U^\alpha(w, y) \right) \delta(x - y) , \quad |z| > |w| \end{aligned}$$

where $x = \theta, y = \varphi$ for $\mathbb{S}^1 \times \mathbb{S}^1$ and $x = u, y = v$ for \mathbb{S}^2 . After extraction of the modes we arrive at the expressions

$$\begin{aligned} \begin{aligned} [L_{m,p}, H_{n,q}^i] &= -nH_{m+n,p+q}^i \\ [L_{m,p}, E_{\alpha,n,q}] &= -nE_{\alpha,m+n,p+q} \end{aligned} & \quad \text{for } \mathbb{S}^1 \times \mathbb{S}^1 \\ \\ \begin{aligned} [L_{\ell_1,m_1}, H_{\ell_2,m_2}^i] &= -m_2 c_{\ell_1,m_1,\ell_2,m_2}^{\ell_3,m_1+m_2} H_{\ell_3,m_1+m_2}^i \\ [L_{\ell_1,m_1}, E_{\alpha,\ell_2,m_2}] &= -m_2 c_{\ell_1,m_1,\ell_2,m_2}^{\ell_3,m_1+m_2} E_{\alpha,\ell_3,m_1+m_2} \end{aligned} & \quad \text{for } \mathbb{S}^2 \end{aligned} \quad (32)$$

It follows that (28) (resp. (29)) and (32), (32) correspond to a bosonic realisation of the semi-direct product of the Kac-Moody and Virasoro algebras of the two-torus and the two-sphere, that we denote $\text{Vir}(\mathbb{S}^1 \times \mathbb{S}^1) \ltimes \hat{\mathfrak{g}}(\mathbb{S}^1 \times \mathbb{S}^1)$ and $\text{Vir}(\mathbb{S}^2) \ltimes \hat{\mathfrak{g}}(\mathbb{S}^2)$, respectively. These two algebras can be seen as a direct generalisation of the usual Kac-Moody and Virasoro algebras $\text{Vir} \ltimes \hat{\mathfrak{g}}$. The case of the two-torus and the two-sphere are structurally different, as only for the case of the two-torus the embedding $\text{Vir} \ltimes \hat{\mathfrak{g}} \subset \text{Vir}(\mathbb{S}^1 \times \mathbb{S}^1) \ltimes \hat{\mathfrak{g}}(\mathbb{S}^1 \times \mathbb{S}^1)$ holds, *i.e.*, we have the semi-direct product of the usual Virasoro and Kac-Moody algebras as a subalgebra. This observation is a simple consequence of the geometrical properties of the two-torus and the two-sphere, where only the former can be written as a Cartesian product of one-spheres. So, taking the two-torus with one of its radii tending to zero (hence corresponding of some ‘kind’ of compactification), reproduces $\text{Vir} \ltimes \hat{\mathfrak{g}}$ in this limit. Furthermore, in the same limit, the mode expansions (4) lead to the usual modes in String Theory.

Finally, we would like to comment briefly on our regularisation procedure. This observation follows from what we already mentioned in the fermionic realisation obtained in [14]. Indeed, the fact that in the case of the two-torus we obtain the usual Kac-Moody and Virasoro algebras as a subalgebra of their analogue on the two-torus, *but with the correct central charges* ($k = 1, c = r$), legitimates *a posteriori* the regularisation related to ζ -regularisation prescription.

The Fock space obtained acting on the vacuum (17) with the creation operators leads to a unitary representation of $\text{Vir}(\mathbb{S}^1 \times \mathbb{S}^1) \ltimes \hat{\mathfrak{g}}(\mathbb{S}^1 \times \mathbb{S}^1)$ and $\text{Vir}(\mathbb{S}^2) \ltimes \hat{\mathfrak{g}}(\mathbb{S}^2)$. Finally, observe that the generators $-L_{0,0}$ of $\text{Vir}(\mathbb{S}^1 \times \mathbb{S}^1)$ and $\text{Vir}(\mathbb{S}^2)$ correspond to the differential operators noted by d in [8]. In addition, in the case of the two-torus, only one differential operator is obtained and only one central charge is non-vanishing. This is in accordance with the property of unitarity of the representations of $\hat{\mathfrak{g}}(\mathbb{S}^1 \times \cdots \times \mathbb{S}^1)$, which implies only one non-vanishing central charge.

We would like to further comment on our regularisation procedure. The main point to obtain a bosonic realisation of the Kac-Moody and Virasoro algebras associated to the two-torus and the two-sphere is related to the consideration of an infinite number of bosons corresponding to a suitable expansion of the r usual bosons of the circle \mathbb{S}^1 on an extra-dimension associated to the two-sphere or the two-torus. (See (4) and (5)). Of course, as we have an infinite number of bosons, new divergences appear. These new divergences are suitably treated with our Riemann ζ -regularisation. As explained in [14], it may seem that this regularisation leads to contradictions. Consider the case of the two-sphere. In this case, the infinite number of bosons is simply related to the mode expansion of the usual bosons of \mathbb{S}^1 $q^i(\varphi), p^i(\varphi), \alpha_m^i(\varphi)$ (see (4)). Introducing regulated bosons, the ζ -function regularisation treats with care infinite sums related to the modes in the φ -direction. If now we take the large r limit (*i.e.*, consider an infinite number of bosons $Q^i(\theta, \varphi), i = 1, \dots, r$), the central charge of the Virasoro algebra diverges and *is not* regularised by our procedure. This means that we *cannot* consider the large r limit, and the only infinities which are allowed are related to the mode expansion in the φ -direction.

We now focus on the two-torus. The Hilbert space is then constructed in an easy and standard manner, acting on the vacuum $|0\rangle$ with creation operators $\alpha_{-mp}^i, m > 0, p \in \mathbb{Z}$. Differently, if we consider the boson operators $P_m^i(\varphi)$ and $P^i(z, \varphi)$, caution needs to be taken. In particular, if one considers the state $\alpha_m^i(\varphi)|0\rangle = |i, m, \varphi\rangle, m < 0$ or $P^i(z, \varphi)|0\rangle = |i, z, \varphi\rangle$, a direct computation shows that

$$\langle i, m, \varphi | j, p, \theta \rangle = \delta_{ij} \delta_{mp} \delta(\varphi - \theta), \quad \langle i, z, \varphi | j, w, \theta \rangle = \delta_{ij} \frac{zw}{(z-w)^2} \delta(\varphi - \theta).$$

This construction extends naturally to the n -particle state, but as it is usual in QFT, the normal ordering prescription and the regularisation procedure have to be applied (in particular for the latter when we have operators at the same points). Thus, the normal ordering prescription (possibly with the additional ζ -regularisation) must be considered when computing N-points functions.

Some comments are in order for the case of the two-torus. The requirement to introduce a ζ -regularisation comes from our normal ordering prescription (10) and order relation (9). Indeed, the generators of the Kac-Moody algebra of the two-torus (a similar analysis holds for the generators of the Virasoro algebra) are parameterised by the roots of \mathfrak{g} and two relative numbers. The two latter numbers correspond to the Taylor expansion of the generators of the simple Lie algebra \mathfrak{g} along the two loops of the two-torus. Stated differently, the necessity of our regularisation procedure is simply the consequence of the fact that we have to order two couples of numbers (m, p) and (n, q) . One may wonder whether it is possible to avoid this regularisation procedure. Actually, an alternative construction of the representation theory of the toroidal algebras (*i.e.*, the Kac-Moody algebra of the two-torus) and its quantum extension were studied in [23, 24] (the second paper dealing specifically with the quantum version of the toroidal algebra). In these two papers there is no need to define a regularisation prescription. This results from a different presentation of the (quantum) toroidal algebra, as in both cases the presentation of the algebra is defined through the mode expansion of the affine (quantum affine) Lie algebra along the circle (in the second article the authors add central extensions to the loop affine algebra). In particular, in [23] a vertex operator was constructed along these lines. Thus, to the roots α of $\hat{\mathfrak{g}}$ a Heisenberg algebra generated by $\alpha(m)$ with α a root of $\hat{\mathfrak{g}}$ and $m \in \mathbb{Z} \setminus \{0\}$ is introduced:

$$[\alpha(m), \beta(n)] = m\delta_{m+n}\alpha \cdot \beta \quad (33)$$

(In our case we can easily introduce such a Heisenberg algebra by setting $\alpha(m, p) = \alpha_i \alpha_{m, p}^i$ with α a root of \mathfrak{g} with component α_i) and where $\alpha(0)$ is the analogue of the p^i in our construction. Then within the Heisenberg algebra $\alpha(m), m \neq 0$ a Vertex representation of the toroidal algebra was considered (the operator e^α of their construction is the analogue of \hat{c}_α in our construction). This construction is in fact very close to ours, with two fundamental differences. In first place, as the expansion involves only one number (the presentation of the algebra is made from the affine Lie algebra), standard techniques of Conformal Field Theory can be used without ζ -regularisation. Secondly, there is an important difference between simple Lie algebras and affine Lie algebras: for the former the determinant of the Cartan matrix is non-zero, whilst for the latter the corank of the Cartan matrix is one. In particular, for affine Lie algebras we have imaginary (or null) roots (generally denoted $n\delta$, with $n \in \mathbb{Z} \setminus \{0\}$) such that $\delta \cdot \delta = 0$. This means in particular that the Heisenberg algebra (33) is degenerate, and consequently we have zero-norm states in the representation space. Thus the representation obtained in [23] is non unitary, even if $\hat{\mathfrak{g}}$ is an affine Lie algebra associated to a compact Lie algebra. Let us notice that the operators associated to $n\delta$ are basically related to new central charges (beyond $k = 1$) since $\alpha(n\delta), n \in \mathbb{Z}$ are central (see eq.[33]) and are not considered in our analysis. An interesting question that emerges from this context and deserves further investigation is the precise relationship between the two constructions. It is worthy to be recalled that the construction of [23] has been extended to n -tori, together with an extension of the Virasoro algebra close to our construction (again having the roots of $\hat{\mathfrak{g}}$ and not those of \mathfrak{g} as starting point)[25].

Let us finally mention that the infinitely many bosons corresponding to the mode expansion of the bosons on the two-torus $P^i(z, \varphi), i = 1, \dots, d$ enable us to construct in a straightforward way the Hilbert space (some care has to be taken only when we consider bosons operators with an infinite number of modes). In this construction, there is no need to proceed to any normal ordering or regularisation prescription since the action of the bosons modes α_{mp}^i on the Fock space is well defined. In other words, a normal ordering and regularisation prescription are needed when one tries to define a Kac-Moody and a Virasoro algebra associated to the two-torus, in particular for the computation of the central charge. Having regulated the algebra, it turns out the Fock space associated to the bosons is naturally a unitary representation of the Kac-Moody and Virasoro algebras of the two-torus. This means that we have, prior to a bosonic realisation of our algebra, a unitary representation. Thus to make contact with the representation space of [23], one may wonder whether one can extract the zero-norm states and, after extraction of these states, the two resulting representations spaces coincide.

In this note we have extended in a natural manner the usual vertex realisation of Kac-Moody and Virasoro

algebras to their analogues associated to the two-torus and the two-sphere. The crucial point to obtain non-trivial central extensions is the normal ordering prescription, where creation operators are moved to the left of annihilation operators. This normal ordering prescription has to be supplemented by a regularisation procedure to regularise infinite divergent sums. We finally observe that the proposed construction extends naturally to the n -torus, and possibly to more complicate manifolds. In the case of higher dimensional tori, we have the canonical embedding chain

$$\mathrm{Vir}(\mathbb{T}^n) \ltimes \mathfrak{g}(\mathbb{T}^n) \subset \mathrm{Vir}(\mathbb{T}^{n-1}) \ltimes \mathfrak{g}(\mathbb{T}^{n-1}) \subset \cdots \subset \mathrm{Vir}(\mathbb{S}^1 \times \mathbb{S}^1) \ltimes \hat{\mathfrak{g}}(\mathbb{S}^1 \times \mathbb{S}^1) \subset \mathrm{Vir} \ltimes \hat{\mathfrak{g}}.$$

This series of embeddings again follows easily from the geometrical properties of n -tori, as $\mathbb{T}^n = \mathbb{T}^{n-1} \times \mathbb{S}^1$, and can be properly understood if we consider successively the limits for the various radii tending to zero. Besides the apparently cumbersome alternative construction with the ζ -regularisation, the procedure presented in this work suggests a quite generic approach that is potentially of use for constructing bosonic realisations of the semi-direct product of the extension of Kac-Moody and Virasoro algebras on higher-dimensional compact manifolds, and where the usual approaches are of difficult implementation. Work in this direction is currently in progress.

Acknowledgements. The authors thank P. Sorba, A. Marrani, Ch. Schubert and G. Bossard for valuable suggestions and comments. We are grateful to the anonymous reviewer for many helpful comments and for pointing out reference [24]. RCS acknowledges financial support by the research grant PID2019-106802GB-I00/AEI/10.13039/501100011033 (AEI/ FEDER, UE).

References

- [1] V. G. Kac, “Simple graded Lie algebras of finite growth,” [Func. Anal. Appl.](#) **1** (1967) 82–83.
- [2] V. G. Kac, [Infinite Dimensional Lie Algebras](#). 3rd ed. Cambridge University Press: Cambridge, MA, 1990.
- [3] R. V. Moody, “Lie algebras associated with generalized Cartan matrices,” [Bull. Amer. Math. Soc.](#) **73** (1967) 217–221.
- [4] I. G. Macdonald, “Kac-Moody-algebras..” Lie Algebras and Related Topics, proc. semin., windsor/ont. 1984, cms conf. proc. 5, 69-109 (1986)., 1986.
- [5] A. Pressley and G. Segal, [Loop Groups](#). Oxford University Press: Oxford, 1986.
- [6] P. Goddard and D. I. Olive, “Kac-Moody and Virasoro Algebras in relation to Quantum Physics,” [Int. J. Mod. Phys. A](#) **1** (1986) 303–404.
- [7] A. Belavin, A. Polyakov, and A. Zamolodchikov, “Infinite conformal symmetry in two-dimensional quantum field theory,” [Nuclear Physics B](#) **241** (1984) no. 2, 333–380.
- [8] R. Campoamor-Stursberg, M. de Montigny, and M. Rausch de Traubenberg, “An overview of generalised Kac-Moody algebras on compact real manifolds,” [J. Geom. Phys.](#) **180** (2022) 104624, [arXiv:2105.05461 \[math-ph\]](#).
- [9] I. Bars, “Local charge algebras in quantum chiral models and gauge theories.” In *Vertex Operators in Mathematics and Physics*, Ed. J. Lepowsky, S. Mandelstam and I. M. Singer (Springer, Berlin 1984), pp 373–391.
- [10] R. Høegh-Krohn and B. Torresani, “Classification and construction of quasisimple Lie algebras,” [J. Funct. Anal.](#) **89** (1990) no. 1, 106–136.
- [11] L. Frappat, E. Ragoucy, P. Sorba, F. Thuillier, and H. Hogaasen, “Generalized Kac-Moody algebras and the diffeomorphism group of a closed surface,” [Nucl. Phys. B](#) **334** (1990) 250–264.

- [12] P.-M. Ho, Y. Matsuo, and S. Shiba, “Lorentzian Lie (3-)algebra and toroidal compactification of M/string theory,” [JHEP **03** \(2009\) 045](#), [arXiv:0901.2003 \[hep-th\]](#).
- [13] K. Harada, P.-M. Ho, Y. Matsuo, and A. Watanabe, “Dimensional oxidization on coset space,” [JHEP **10** \(2020\) 198](#), [arXiv:2005.13936 \[hep-th\]](#).
- [14] R. Campoamor-Stursberg and M. Rausch de Traubenberg, “Fermion realisations of generalised Kac–Moody algebras associated to the two-sphere and the two-torus,” [arXiv:2211.04199 \[math-ph\]](#).
- [15] I. B. Frenkel and V. G. Kac, “Basic Representations of Affine Lie Algebras and Dual Resonance Models,” [Invent. Math. **62** \(1980\) 23–66](#).
- [16] G. Segal, “Unitarity Representations of Some Infinite Dimensional Groups,” [Commun. Math. Phys. **80** \(1981\) 301–342](#).
- [17] W. A. de Graaf, [Lie Algebras: Theory and Algorithms](#), vol. 56 of [North-Holland Mathematical Library](#). North-Holland Publishing Co., Amsterdam, 2000.
- [18] P. Goddard and D. Olive, “Algebras, lattices and strings,” [Proceedings, Vertex Operators in Mathematics and Physics **3** \(1985\) 51–96](#).
- [19] S. Fubini and G. Veneziano, “Duality in operator formalism,” [Nuovo Cim. A **67** \(1970\) 29–47](#).
- [20] S. Fubini, D. Gordon, and G. Veneziano, “A general treatment of factorization in dual resonance models,” [Phys. Lett. B **29** \(1969\) 679–682](#).
- [21] J.-L. Gervais, “Operator expression for the Koba and Nielsen multi-Veneziano formula and gauge identities,” [Nucl. Phys. B **21** \(1970\) 192–204](#).
- [22] P. Di Francesco, P. Mathieu, and D. Senechal, [Conformal Field Theory](#). Graduate Texts in Contemporary Physics, Springer: New York, 1997.
- [23] R. V. Moody, S. E. Rao, and T. Yokonuma, “Toroidal Lie algebras and vertex representations,” [Geom. Dedicata **35** \(1990\) no. 1-3, 283–307](#). <https://doi.org/10.1007/BF00147350>.
- [24] B. Feigin, M. Jimbo, T. Miwa, and E. Mukhin, “Representations of quantum toroidal \mathfrak{gl}_n ,” [J. Algebra **380** \(2013\) 78–108](#). <https://doi-org.scd-rproxy.u-strasbg.fr/10.1016/j.jalgebra.2012.12.029>.
- [25] S. Eswara Rao and R. V. Moody, “Vertex representations for n -toroidal Lie algebras and a generalization of the Virasoro algebra,” [Commun. Math. Phys. **159** \(1994\) no. 2, 239–264](#).