Particle identification with the cluster counting technique for the IDEA drift chamber

C. Caputo^a, G. Chiarello^b, A. Corvaglia^b, F. Cuna^{b,c}, B. D'Anzi^{d,e,*}, N. De Filippis^{e,f}, W. Elmetenawee^e, E. Gorini^{b,c}, F. Grancagnolo^b, M. Greco^{b,c}, S. Gribanov^g, K. Johnson^h, M. Maggi^e, A. Miccoli^b, M. Panareo^{b,c}, A. Popov^g, M. Primavera^b, A. Taliercio^a, G. F. Tassielli^e, A. Ventura^b, S. Xin^{i,j}

^aInstitut de recherche en mathématique et physique, Université Catholique de Louvain, Chemin du Cyclotron 2/L7.01.01, Louvain-la-Neuve, 1348, Belgium

^dUniversità degli Studi di Bari Aldo Moro, Piazza Umberto I, 1, Bari BA, 70121, Italy

^eIstituto Nazionale di Fisica Nucleare, Via Giovanni Amendola, 173, Bari BA, 70126, Italy

^fPolitecnico di Bari, Via Edoardo Orabona, 4, Bari BA, 70126, Italy

⁸No affiliation. Russian Federation

^hFlorida State University, 600 W College Ave, Tallahassee FL, 32306, United States

ⁱChinese Academy of Sciences (CAS), 19A Yuquan Road, Shijing District, Beijing, 100049, China

¹*Philote States* (CAS), 174 Nuquan Road, Shijing District, Beijing, 100049, China ¹*Philote Academy of Sciences* (CAS), 174 Nuquan Road, Shijing District, Beijing, 100049, China ¹*Institute of High Energy Physics, 19B. Yuquan Road, Shijing District, Beijing, 100049, China* ¹*Philote Academy of Sciences* (CAS), 174 Nuquan Road, Shijing District, Beijing, 100049, China ¹*Institute of High Energy Physics, 19B. Yuquan Road, Shijing District, Beijing, 100049, China* ¹*Philote Academy of Sciences* (CAS), 174 Nuquan Road, Shijing District, Beijing, 100049, China ¹*Philote Academy of Sciences* (CAS), 174 Nuquan Road, Shijing District, Beijing, 100049, China ¹*Philote Academy of Sciences* (CAS), 174 Nuquan Road, Shijing District, Beijing, 100049, China ¹*Philote Academy of Sciences* (CAS), 174 Nuquan Road, Shijing District, Beijing, 100049, China ¹*Philote Academy of Sciences* (CAS), 174 Nuquan Road, Shijing District, Beijing, 100049, China ¹*Philote Academy of Sciences* (CAS), 174 Nuquan Road, Shijing District, Beijing, 100049, China ¹*Philote Academy of Sciences* (CAS), 174 Nuquan Road, Shijing District, Beijing, 100049, China ¹*Philote Academy of Sciences* (CAS), 174 Nuquan Road, Shijing District, Beijing, 100049, China ¹*Philote Academy of Sciences* (CAS), 174 Nuquan Road, Shijing District, Beijing, 100049, China ¹*Philote Academy of Sciences* (CAS), 174 Nuque, Nuque

and offers a more statistically robust procedure to infer mass information. This is particularly true for He-based drift chambers with low density of the primary ionization clusters. The technique consists in singling out, in ever recorded detector signal, the isolated structures related to the arrival on the anode wire of the electrons belonging to a single primary ionization act. The goodness of this method relies on the number of primary ionizations being independent from cluster size and gas gain fluctuations and being insensitive to highly ionizing δ -rays. It is going

the future plan for optimizing and validating the CC algorithms and the CC technique for PID.

2. Analytical calculations and Simulations

From analytical calculations, we infer an excellent K/π separation for He/iC₄H₁₀ 90/10 gas mixture over the entire momentum range except for 0.85 GeV (blue lines inFigure 1), which can be easily recovered with a modest (100 ps resolution over approximately 2 m path length) time of flight measurement. On the other hand, starting from detailed studies

^bIstituto Nazionale di Fisica Nucleare, Via Arnesano, 0, Lecce LE, 73100, Italy

^cUniversità del Salento, Piazza Tancredi, 7, Lecce LE, 73100, Italy

^{*}Corresponding author. Email address: brunella.danzi@ba.infn.it (B. D'Anzi)

Preprint submitted to Nuclear Inst. and Methods in Physics Research, A



Figure 1: Analytic evaluation of particle separation capabilities achievable with dE/dx (dashed curves) and dN/dx (solid curves). The region between 0.85 GeV/c and 1.05 GeV/c, where a different technique is needed, is highlighted in yellow.



Figure 2: Particle separation capability results from a simulation of the ionization process in 200 drift cells, 1 cm wide, in He/iC_4H_{10} 90/10 gas mixture that has been performed both in Garfield++ (top) and in Garfield-modelled Geant4 (bottom) by using the energy loss method (left) and CC technique (right).

of Garfield++ simulation results about the ionization process in He-based gas mixtures, an algorithm which reproduces the cluster number and cluster size distributions using the energy deposit information by Geant4 has been developed under the assumption of a 100% CC efficiency [6]. From Figure 2, we observe that the dN_{cl}/dx improves particle separation capabilities and Garfield++ is in good agreement with analytical calculations up to 20 GeV/c, then falls much more rapidly at higher momenta.

3. Cluster counting technique applied to beam test data

To validate the simulations results, a first beam test, using a 165 GeV/c muon beam on a setup made of 1 cm and 2 cm size drift tubes, equipped with Mo, Al, and W sense wires in the 10 - 40 μm diameter range, has been performed at CERN/H8 beam line by collecting data with the gas mixtures He/iC₄H₁₀ 90/10 and 80/20, by making a scan in high voltage and angle between the wire direction and ionizing tracks. Fulfilling the main requirements for an efficient CC on the readout electronics (1 GHz bandwidth and at least a 1 GS/s sampling rate at 12 bits), entirely based on a DRS system [7], has been crucial. The most efficient CC algorithms are the Running Template Algorithm (RTA), scanning the waveform with an electron peak template



Figure 3: Example of real waveforms collected from 2 cm size drift tubes (90/10 He/iC₄H₁₀). The blue and red arrows represent the cluster and electron peaks found with the DERIV and RTA algorithms.



Figure 4: Distribution of the number of clusters for 1 cm size drift tubes in He:IsoB 90/10 fitted with a gaussian.

and its iterative subtraction, and the Derivative Algorithm (DE-RIV), imposing cuts on the waveform amplitude, the first and second derivatives (see Figure 3). They define the limiting effects for a fully efficient CC (space charge effect + attachment + recombination) which is approximately 80% in He/iC₄H₁₀ 90/10 gas mixture. Figure 4 shows the distribution of the number of clusters for a particular configuration. The gaussian fit indicates a mean value μ very close to the expected one and a sigma almost equal to the square root of μ , confirming the Poisson nature of the distribution.

4. Conclusion and Future Perspectives

Using preliminary CC algorithms gave promising results in reproducing the expected physical distributions. As a next step, the experimental setup will undergo new tests in muon beams with momenta over the full range of interest for the future lepton machines, to assess the particle identification capabilities of the cluster counting approach with optimized algorithms.

References

- M. Benedikt, et al., Fcc-ee the lepton collider : Future circular collider conceptual design-report, Eur. Phys. J. Spec. Top. 228 (2) (2019) 261–623.
- [2] F. Bedeschi, A detector concept proposal for a circular e+e- collider, in: PoS. ICHEP2020 Conf., 2021, p. 819.
- [3] G. F. Tassielli, A proposal of a drift chamber for the idea experiment for a future e+e- collider, in: PoS. ICHEP2020 Conf., 2021, p. 877.
- [4] H. Fischle, J. Heintze, B. Schmidt, Experimental determination of ionization cluster size distributions in counting gases, Nuclear Instruments and Methods in Physics Research A 301 (1991) 202–214.
- [5] R. G. Kepler, C. A. D'Andlau, F. W. B., L. F. Hansen, Relativistic increase of energy loss by ionization in gases, Il Nuovo Cimento VII (1) (1958).
- [6] F. Cuna, N. De Filippis, F. Grancagnolo, G. F. Tassielli, Simulation of particle identification with the cluster counting technique, in: Proc. LCWS2021 Work., 2021.
- [7] S. Ritt, R. Dinapoli, U. Hartmann, Application of the drs chip for fast waveform digitizing, Nuclear Instruments and Methods in Physics Research Section A 623 (1) (2010) 486–488.