

COMBINATORIAL BASES OF STANDARD MODULES OF TWISTED AFFINE LIE ALGEBRAS IN TYPES $A_{2l-1}^{(2)}$ AND $D_{l+1}^{(2)}$: RECTANGULAR HIGHEST WEIGHTS

MARIJANA BUTORAC AND SLAVEN KOŽIĆ

ABSTRACT. We consider the standard modules of rectangular highest weights of affine Lie algebras in types $A_{2l-1}^{(2)}$ and $D_{l+1}^{(2)}$. By using vertex algebraic techniques we construct the combinatorial bases for standard modules and their principal subspaces and parafermionic spaces. Finally, we compute the corresponding character formulae and, as an application, we obtain two new families of combinatorial identities.

1. INTRODUCTION

The connection between Rogers–Ramanujan-type sums and characters of parafermionic spaces was originally established in the work of J. Lepowsky and M. Primc [LP85] by constructing the vertex operator bases of parafermionic spaces for the affine Lie algebra $A_1^{(1)}$. Later on, such combinatorial bases for the parafermionic spaces of standard modules of rectangular highest weights for all untwisted affine Lie algebras were found by G. Georgiev [G95] (type $A_l^{(1)}$) and by M. Primc and the authors [BuKoP21] (types $B_l^{(1)}$ – $G_2^{(1)}$). These bases were used to prove the generalized Rogers–Ramanujan-type sums obtained by A. Kuniba, T. Nakanishi and J. Suzuki [KuNS93] and D. Gepner [Gep87]. Generalizing this approach, M. Okado and R. Takenaka [OTa22, Ta22] found bases for the parafermionic spaces of standard modules of highest weights $k\Lambda_0$ for twisted affine Lie algebras. Moreover, using these bases they proved character formulae conjectured by G. Hatayama, A. Kuniba, M. Okado, T. Takagi and Z. Tsuboi [HKOTT02].

In this paper, we construct the combinatorial bases for the parafermionic spaces of standard modules of *rectangular* highest weights, i.e. of weights of the form

$$\Lambda = k_0\Lambda_0 + k_j\Lambda_j, \quad \text{where} \quad k_0, k_j \in \mathbb{Z}_{\geq 0} \quad \text{and} \quad k = k_0 + k_j > 0, \quad (1.1)$$

for twisted affine Lie algebras of types $A_{2l-1}^{(2)}$ and $D_{l+1}^{(2)}$, where $j = 1$ (resp. $j = l$) in type $A_{2l-1}^{(2)}$ (resp. $D_{l+1}^{(2)}$). As with [BuKoP21, OTa22, G95], the construction consists of several steps, as follows.

First, we obtain the quasi-particle bases of the principal subspaces $W_{L^{\hat{\nu}}(\Lambda)}$ of standard modules $L^{\hat{\nu}}(\Lambda)$ for the highest weight Λ as in (1.1). As with the corresponding bases in the case $\Lambda = k\Lambda_0$, found by the first author and C. Sadowski [BuSa19], they are expressed in terms of monomials of twisted quasi-particles applied on the highest weight vector of $L^{\hat{\nu}}(\Lambda)$, such that the quasi-particle energies satisfy certain difference and initial conditions. However, in this case, the quasi-particle monomials satisfy different initial

(M. Butorac) FACULTY OF MATHEMATICS, UNIVERSITY OF RIJEKA, RADMILE MATEJČIĆ 2, 51000 RIJEKA, CROATIA

(S. Kožić) DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, UNIVERSITY OF ZAGREB, BIJENIČKA CESTA 30, 10000 ZAGREB, CROATIA

E-mail addresses: mbutorac@math.uniri.hr, kslaven@math.hr.

2010 *Mathematics Subject Classification.* 17B67 (Primary) 17B69, 05A19 (Secondary).

Key words and phrases. Principal subspace, Combinatorial basis, Standard module, Parafermionic space.

conditions, which we determine by employing results of J. Lepowsky [L85] and B. Bakalov and V. Kac [BK04] on the lattice construction of level one twisted modules. Despite an apparent analogy between the cases $\Lambda = k\Lambda_0$ and $\Lambda = k_0\Lambda_0 + k_j\Lambda_j$, the latter requires a certain new ingredient in the proof of linear independence. More specifically, we use the so-called simple current operator from Li's papers [Li96] and [Li12], which is a certain linear bijection between level one twisted modules from [L85]. In addition, we also construct quasi-particle bases for the principal subspaces of generalized Verma modules.

Next, we use the quasi-particle bases for principal subspaces to construct the combinatorial bases for standard modules of rectangular highest weights in types $A_{2l-1}^{(2)}$ and $D_{l+1}^{(2)}$, thus generalizing the aforementioned results of M. Okado and R. Takanaka [OTa22]. As before, the proof of linear independence relies on Li's simple current operators. Finally, we employ the parafermionic projection to obtain the combinatorial bases of parafermionic spaces for all rectangular highest weights in types $A_{2l-1}^{(2)}$ and $D_{l+1}^{(2)}$.

At the end, we use all these bases to compute the character formulae of the corresponding standard modules and their principal subspaces and parafermionic spaces. In particular, we obtain the parafermionic character formulae which coincide with those of D. Gepner [Gep22]. Furthermore, by comparing the quasi-particle bases for the principal subspaces of the generalized Verma modules with their Poincaré–Birkhoff–Witt bases, we discover two new families of combinatorial identities which correspond to types $A_{2l-1}^{(2)}$ and $D_{l+1}^{(2)}$. The identities of such form often appear in different areas of mathematics; see, e.g., the papers by J. Fulman [Fu00] and J. Hua [Hu00].

2. PRELIMINARIES

2.1. Level one standard modules. Let \mathfrak{g} be a complex simple Lie algebra of type A_{2l-1} or D_{l+1} with the Cartan subalgebra \mathfrak{h} and the root system R , equipped with the standard nondegenerate symmetric invariant bilinear form $\langle \cdot, \cdot \rangle$. Using this form one can identify \mathfrak{h} and its dual \mathfrak{h}^* via $\langle \alpha, h \rangle = \alpha(h)$ for $\alpha \in \mathfrak{h}^*$ and $h \in \mathfrak{h}$. We assume that the form is rescaled so that $\langle \alpha, \alpha \rangle = 2$ when α is a simple root. The simple roots $\alpha_1, \dots, \alpha_{\text{rk}\mathfrak{g}}$, where $\text{rk}\mathfrak{g}$ stands for the rank of \mathfrak{g} , generate the root lattice Q and the fundamental weights $\lambda_1, \dots, \lambda_{\text{rk}\mathfrak{g}}$ generate the weight lattice P of \mathfrak{g} . Let ν be the Dynkin diagram automorphism of Q of order 2 as in Figure 1.




FIGURE 1. Finite Dynkin diagrams and the action of the automorphism ν

Let us recall the construction of standard modules of twisted affine Lie algebra $\widehat{\mathfrak{g}}[\widehat{\nu}]$ associated to \mathfrak{g} and ν from [L85] (see also [BK04, CLM14, DL96, FLM87, FLM88, Li96]). The central extension \widehat{Q} of the root lattice Q by the cyclic group $\langle -1 \rangle = \{\pm 1\}$,

$$1 \longrightarrow \langle -1 \rangle \longrightarrow \widehat{Q} \xrightarrow{2} Q \longrightarrow 1 \quad (2.1)$$

is determined by the commutator map $C_0(\alpha, \beta) = (-1)^{\langle \alpha, \beta \rangle}$, i.e. by $aba^{-1}b^{-1} = C_0(\bar{a}, \bar{b})$ for $a, b \in \widehat{Q}$. We fix a section $e : Q \rightarrow \widehat{Q}$ normalized so that $e_0 = 1$ and $\bar{e}_\alpha = \alpha$ for all $\alpha \in Q$ and

$$e_\alpha e_\beta = \epsilon_{C_0}(\alpha, \beta) e_{\alpha+\beta} \quad \text{for all } \alpha, \beta \in Q.$$

Here $\epsilon_{C_0}(\cdot, \cdot)$ denotes the normalized 2-cocycle $\epsilon_{C_0} : Q \times Q \rightarrow \langle -1 \rangle$ defined as in [PeSa18].

Let $\widehat{\nu}$ be the lifting of ν to the involutive automorphism of \widehat{Q} such that

$$\overline{\widehat{\nu}a} = \nu\bar{a} \quad \text{for all } a \in \widehat{Q} \quad \text{and} \quad \widehat{\nu}a = a \quad \text{if } \nu\bar{a} = \bar{a}.$$

By the same symbol we denote the extension of the automorphism $\widehat{\nu}$ to the lattice vertex operator algebra V_Q . Moreover, for any k we write $\widehat{\nu} = \widehat{\nu}^{\otimes k}$ for the extension of $\widehat{\nu}$ to the vertex operator algebra $V_Q^{\otimes k}$. Again, we have $\widehat{\nu}^2 = 1$ (cf. [BK04, L85]).

For any integer m set

$$\mathfrak{h}_{(m)} = \{x \in \mathfrak{h} \mid \nu(x) = (-1)^m x\}.$$

This gives us the decomposition of the Cartan subalgebra

$$\mathfrak{h} = \mathfrak{h}_{(0)} \oplus \mathfrak{h}_{(1)}, \quad \text{where} \quad \mathfrak{h}_{(0)} = \bigoplus_{i=1}^l \mathbb{C}(\alpha_i + \nu\alpha_i) \quad \text{and} \quad \mathfrak{h}_{(1)} = \bigoplus_{i=1}^l \mathbb{C}(\alpha_i - \nu\alpha_i).$$

For any $\alpha \in \mathfrak{h}$ and $m \in \mathbb{Z}$ we write $\alpha_{(m)}$ for the projection of α onto $\mathfrak{h}_{(m)}$.

Let

$$\widehat{\mathfrak{h}}[\nu]^\pm = \mathfrak{h}_{(0)} \otimes t^{\pm 1} \mathbb{C}[t^{\pm 1}] \oplus \mathfrak{h}_{(1)} \otimes t^{\pm 1/2} \mathbb{C}[t^{\pm 1}].$$

Consider the Heisenberg subalgebra

$$\widehat{\mathfrak{h}}[\nu]_{\frac{1}{2}\mathbb{Z}} = \widehat{\mathfrak{h}}[\nu]^+ \oplus \widehat{\mathfrak{h}}[\nu]^- \oplus \mathbb{C}c$$

of the twisted affine Lie algebra

$$\widetilde{\mathfrak{h}}[\nu] = \coprod_{m \in \mathbb{Z}} \mathfrak{h}_{(m)} \otimes t^{m/2} \oplus \mathbb{C}c \oplus \mathbb{C}d = \mathfrak{h}_{(0)} \otimes \mathbb{C}[t, t^{-1}] \oplus \mathfrak{h}_{(1)} \otimes t^{1/2} \mathbb{C}[t, t^{-1}] \oplus \mathbb{C}c \oplus \mathbb{C}d.$$

Furthermore, let $S[\nu]$ be the irreducible $\widehat{\mathfrak{h}}[\nu]_{\frac{1}{2}\mathbb{Z}}$ -module

$$S[\nu] = U(\widetilde{\mathfrak{h}}[\nu]) \otimes_{U_+} \mathbb{C}, \quad \text{where} \quad U_+ = U(\prod_{m \in \frac{1}{2}\mathbb{Z}_{\geq 0}} \mathfrak{h}_{(2m)} \otimes t^m \oplus \mathbb{C}c \oplus \mathbb{C}d).$$

Note that $S[\nu]$ is linearly isomorphic to the space of symmetric algebra $S(\widehat{\mathfrak{h}}[\nu]^-)$.

Form the induced \widehat{Q} -module

$$U_T = \mathbb{C}[\widehat{Q}] \otimes_{\mathbb{C}[\widehat{N}]} T \cong \mathbb{C}[Q/N],$$

where \widehat{N} is the subgroup of \widehat{Q} obtained by pulling back $N = \mathfrak{h}_{(1)} \cap Q$, and T denotes the one dimensional \widehat{N} -module \mathbb{C} with character $\tau : \widehat{N} \rightarrow \mathbb{C}^\times$. Note that U_T can be regarded as an $\widetilde{\mathfrak{h}}[\nu]$ -module with

$$d \cdot a = -\frac{1}{2} \langle \bar{a} + \nu\bar{a}, \bar{a} \rangle a.$$

Moreover, the $\widetilde{\mathfrak{h}}[\nu]$ -module U_T is isomorphic to $\mathbb{C}[\pi_0 Q] \otimes T$, where π_0 denotes the projection from \mathfrak{h} onto $\mathfrak{h}_{(0)}$ (cf. [CLM14, L85]).

Let V_Q^T be the irreducible $\widehat{\nu}$ -twisted module for V_Q given by

$$V_Q^T = S[\nu] \otimes U_T.$$

For each $e_\alpha \in \widehat{Q}$, we consider $\widehat{\nu}$ -twisted vertex operator

$$Y^{\widehat{\nu}}(\iota(e_\alpha), z) = 2^{-\frac{\langle \alpha, \alpha \rangle}{2}} E^-(-\alpha, z) E^+(-\alpha, z) e_\alpha z^{\alpha_{(0)} + \frac{\langle \alpha_{(0)}, \alpha_{(0)} \rangle}{2} - \frac{\langle \alpha, \alpha \rangle}{2}},$$

where

$$E^\pm(-\alpha, z) = \exp \left(\sum_{m \in \pm \frac{1}{2}\mathbb{Z}_+} \frac{-\alpha_{(2m)}(m)}{m} z^{-m} \right).$$

The component operators $x_\alpha^{\widehat{\nu}}(m)$ are defined by

$$Y^{\widehat{\nu}}(\iota(e_\alpha), z) = \sum_{m \in \frac{1}{2}\mathbb{Z}} x_\alpha^{\widehat{\nu}}(m) z^{-m - \frac{\langle \alpha, \alpha \rangle}{2}}.$$

By [L85], they satisfy

$$x_{\nu\alpha}^{\widehat{\nu}}(m) = (-1)^{2m} x_\alpha^{\widehat{\nu}}(m) \quad \text{for all } m \in \frac{1}{2}\mathbb{Z}. \quad (2.2)$$

We shall use the symbol $\widehat{\nu}$ to denote the lifting of the corresponding automorphism on \mathfrak{h} to an automorphism of \mathfrak{g} , which is given by

$$\widehat{\nu}x_\alpha = \begin{cases} \epsilon_{C_0}(\alpha, \alpha)x_{\nu\alpha}, & \text{if } \mathfrak{g} \text{ is of type } A_{2l-1}, \\ x_{\nu\alpha}, & \text{if } \mathfrak{g} \text{ is of type } D_{l+1}. \end{cases}$$

For any $m \in \mathbb{Z}$ set

$$\mathfrak{g}_{(m)} = \{x \in \mathfrak{g} \mid \widehat{\nu}(x) = (-1)^m x\}.$$

We shall denote by $x_{(m)}$ the projection of the element $x \in \mathfrak{g}$ onto $\mathfrak{g}_{(m)}$. Let $\widetilde{\mathfrak{g}}[\widehat{\nu}]$ be the $\widehat{\nu}$ -twisted affine Lie algebra associated with \mathfrak{g} and $\widehat{\nu}$,

$$\widetilde{\mathfrak{g}}[\widehat{\nu}] = \widehat{\mathfrak{g}}[\widehat{\nu}] \oplus \mathbb{C}d, \quad \text{where} \quad \widehat{\mathfrak{g}}[\widehat{\nu}] = \coprod_{m \in \frac{1}{2}\mathbb{Z}} \mathfrak{g}_{(2m)} \otimes t^m \oplus \mathbb{C}c.$$

Its Lie brackets are given by

$$[x \otimes t^m, y \otimes t^n] = [x, y] \otimes t^{m+n} + \langle x, y \rangle m \delta_{m+n, 0} c, \quad [c, \widetilde{\mathfrak{g}}[\widehat{\nu}]] = 0, \quad [d, x \otimes t^n] = nx \otimes t^n,$$

for $x \in \mathfrak{g}_{(2m)}$, $y \in \mathfrak{g}_{(2n)}$ and $m, n \in \frac{1}{2}\mathbb{Z}$. The Lie algebra $\widetilde{\mathfrak{g}}[\widehat{\nu}]$ is isomorphic to the twisted affine Lie algebra of type $A_{2l-1}^{(2)}$ or $D_{l+1}^{(2)}$ with respect to the choice of Q and ν (cf. [K90]).

The representation of $\widetilde{\mathfrak{h}}[\nu]$ on V_Q^T uniquely extends to the representation of $\widetilde{\mathfrak{g}}[\widehat{\nu}]$ by

$$(x_\alpha)_{(2m)} \otimes t^m \mapsto x_\alpha^{\widehat{\nu}}(m) \quad \text{for } m \in \frac{1}{2}\mathbb{Z}, \alpha \in Q.$$

Moreover, we have $V_Q^T \cong L^{\widehat{\nu}}(\Lambda_0)$, where $L^{\widehat{\nu}}(\Lambda_0)$ is the level one standard $\widetilde{\mathfrak{g}}[\widehat{\nu}]$ -module of the highest weight Λ_0 with the highest weight vector $v_{\Lambda_0} = v_{L^{\widehat{\nu}}(\Lambda_0)} = 1 \otimes 1$ (cf. [L85]). Let

$$\gamma = \pi_0 \lambda_j, \quad (2.3)$$

where, as in (1.1), we set $j = 1$ (resp. $j = l$) for \mathfrak{g} of type A_{2l-1} (resp. D_{l+1}). The element $\gamma \in \mathfrak{h}_{(0)}$ satisfies $\langle \gamma, \alpha \rangle \in \frac{1}{2}Q$ for all $\alpha \in Q$. Following [L85] (see also [CLM14]) we obtain the level one standard $\widetilde{\mathfrak{g}}[\widehat{\nu}]$ -module $L^{\widehat{\nu}}(\Lambda_j)$ of the highest weight Λ_j by defining a new representation of $\widetilde{\mathfrak{h}}[\nu]$ on V_Q^T as follows. The elements of $\mathfrak{h}_{(0)}$ act on U_T by

$$\alpha^\gamma \cdot b \otimes t = \langle \alpha, \gamma + \bar{b} \rangle b \otimes t$$

for $b \in \widehat{Q}$, $t \in T$, $\alpha \in \mathfrak{h}_{(0)}$, the operators z^{α^γ} act by

$$z^{\alpha^\gamma} \cdot b \otimes t = z^{\langle \alpha, \gamma + \bar{b} \rangle} b \otimes t$$

and the Heisenberg algebra $\widehat{\mathfrak{h}}[\nu]_{\frac{1}{2}\mathbb{Z}}$ acts trivially. A γ -shifted $\widehat{\nu}$ -twisted vertex operator $Y^{\widehat{\nu}, \gamma}(\iota(e_\alpha), z)$ is defined by

$$Y^{\widehat{\nu}, \gamma}(\iota(e_\alpha), z) = Y^{\widehat{\nu}}(\iota(e_\alpha), z) z^{\langle \alpha_{(0)}, \gamma \rangle} = 2^{-\frac{\langle \alpha, \alpha \rangle}{2}} E^-(-\alpha, z) E^+(-\alpha, z) e_\alpha z^{\alpha_{(0)}^\gamma + \frac{\langle \alpha_{(0)}, \alpha_{(0)} \rangle}{2} - \frac{\langle \alpha, \alpha \rangle}{2}}$$

(cf. [CLM14]). The component operators $x_\alpha^{\hat{\nu},\gamma}(m)$, $m \in \frac{1}{2}\mathbb{Z}$, are given by

$$Y^{\hat{\nu},\gamma}(\iota(e_\alpha), z) = \sum_{m \in \frac{1}{2}\mathbb{Z}} x_\alpha^{\hat{\nu},\gamma}(m) z^{-m-1} = x_\alpha^{\hat{\nu},\gamma}(z),$$

and they satisfy

$$x_\alpha^{\hat{\nu},\gamma}(m) = x_\alpha^{\hat{\nu}}(m + \langle \alpha_{(0)}, \gamma \rangle). \quad (2.4)$$

Let $\hat{\nu}_\gamma$ be the automorphism of Lie algebra \mathfrak{g} such that

$$\hat{\nu}_\gamma h = \nu h \quad \text{for all } h \in \mathfrak{h} \quad \text{and} \quad \hat{\nu}_\gamma x_\alpha = (-1)^{2\langle \alpha, \gamma \rangle} \hat{\nu} x_\alpha \quad \text{for all } \alpha \in Q.$$

Then $\hat{\nu}_\gamma^2 = 1$ and the $\hat{\nu}_\gamma$ -twisted affine Lie algebra $\tilde{\mathfrak{g}}[\hat{\nu}_\gamma]$ associated to \mathfrak{g} coincides with $\tilde{\mathfrak{g}}[\hat{\nu}]$. Furthermore, from [L85, Thm. 10.1] (see also [CLM14, Thm. 4.2]) follows that the new representation of $\tilde{\mathfrak{h}}[\nu]$ on $V_Q^{T,\gamma}$ uniquely extends to the representation of $\tilde{\mathfrak{g}}[\hat{\nu}]$ by

$$(x_\alpha)_{(2m)} \otimes t^m \mapsto x_\alpha^{\hat{\nu},\gamma}(m) \quad \text{for all } m \in \frac{1}{2}\mathbb{Z}, \alpha \in Q.$$

It can be easily checked that we have the isomorphism of $\tilde{\mathfrak{g}}[\hat{\nu}]$ -modules $V_Q^{T,\gamma} \cong L^{\hat{\nu}}(\Lambda_j)$. We denote the highest weight vector of $L^{\hat{\nu}}(\Lambda_j)$ by v_{Λ_j} . By [Li96, Prop. 5.4], $V_Q^{T,\gamma}$ is a $\nu_\gamma \hat{\nu}$ -twisted V_Q -module, where $\nu_\gamma = e^{-2\pi\sqrt{-1}\gamma(0)}$ is an automorphism of V_Q .

2.2. Higher level standard modules. Let $k \in \mathbb{N}$ be fixed. The tensor product $(V_Q^{T,\gamma})^{\otimes k}$ of k copies of $\hat{\nu}_\gamma$ -twisted V_Q -modules $V_Q^{T,\gamma}$ possesses a structure of $\hat{\nu}_\gamma$ -twisted module for the vertex operator algebra $V_Q^{\otimes k}$ with vertex operators

$$Y^{\hat{\nu},\gamma}(v_1 \otimes \cdots \otimes v_k, z) = Y^{\hat{\nu},\gamma}(v_1, z) \otimes \cdots \otimes Y^{\hat{\nu},\gamma}(v_k, z).$$

The operators $x_\alpha^{\hat{\nu},\gamma}(m)$ on $(V_Q^{T,\gamma})^{\otimes k}$ are given by

$$Y^{\hat{\nu},\gamma}(x_\alpha(-1) \cdot (\mathbf{1} \otimes \cdots \otimes \mathbf{1}), z) = \sum_{m \in \frac{1}{2}\mathbb{Z}} x_\alpha^{\hat{\nu},\gamma}(m) z^{-m-1}.$$

This construction gives the diagonal action of the sublattice $kQ \subset Q$ on $(V_Q^{T,\gamma})^{\otimes k}$:

$$k\alpha \mapsto \rho(k\alpha) = e^\alpha \otimes \cdots \otimes e^\alpha, \quad \alpha \in Q. \quad (2.5)$$

To simplify the notation, in the rest of the paper we will omit the symbol γ and write $x^{\hat{\nu}}(z)$ for $x^{\hat{\nu},\gamma}(z)$ as well, as it will be clear from the context whether γ is 0 or $\pi_0 \lambda_j$. We shall need the following relations on the standard $\tilde{\mathfrak{g}}[\hat{\nu}]$ -module $L^{\hat{\nu}}(\Lambda)$, which are consequence of the adjoint action of e_α on $\tilde{\mathfrak{g}}$ (cf. [K90], see also [OTa22, P94]):

$$e_\alpha d e_\alpha^{-1} = d + \alpha - \frac{\langle \alpha_{(0)}, \alpha_{(0)} \rangle}{2} c, \quad (2.6)$$

$$e_\alpha c e_\alpha^{-1} = c, \quad (2.7)$$

$$e_\alpha h_{(0)} e_\alpha^{-1} = h_{(0)} - \langle \alpha_{(0)}, h_{(0)} \rangle c, \quad h \in \mathfrak{h}, \quad (2.8)$$

$$e_\alpha h_{(2s)}(s) e_\alpha^{-1} = h_{(2s)}(s), \quad s \neq 0, \quad (2.9)$$

$$e_\alpha x_\beta^{\hat{\nu}}(s) e_\alpha^{-1} = x_\beta^{\hat{\nu}}(s - \langle \beta_{(0)}, \alpha_{(0)} \rangle). \quad (2.10)$$

For any positive integer r consider the twisted vertex operators

$$x_{r\alpha_i}^{\hat{\nu}}(z) = Y^{\hat{\nu}}(x_{\alpha_i}(-1)^r \mathbf{1}, z) = \sum_{m \in \frac{1}{2}\mathbb{Z}} x_{r\alpha_i}^{\hat{\nu}}(m) z^{-m-r} = (Y^{\hat{\nu}}(x_{\alpha_i}(-1) \mathbf{1}, z))^r, \quad (2.11)$$

associated with the vector $x_{\alpha_i}(-1)^r \mathbf{1} \in L(k\Lambda_0)$ (cf. [Li96]). They satisfy the commutator formula for twisted vertex operators,

$$[x_{\alpha}^{\widehat{\nu}}(z_1), x_{r\beta}^{\widehat{\nu}}(z_2)] = \sum_{s \geq 0} \frac{(-1)^s}{2s!} \left(\frac{d}{dz_1} \right)^s z_2^{-1} \sum_{q \in \mathbb{Z}/2\mathbb{Z}} \delta \left((-1)^q \frac{z_1^{\frac{1}{2}}}{z_2^{\frac{1}{2}}} \right) Y^{\widehat{\nu}}(\nu^q x_{\alpha}(s) x_{\beta}(-1)^r \mathbf{1}, z_2).$$

For any $h \in \mathfrak{h}$ and $m \in \frac{1}{2}\mathbb{Z}$ we have

$$[h_{(2m)}(m), x_{r\alpha}^{\widehat{\nu}}(z)] = r \langle h_{(2m)}, \alpha_{(-2m)} \rangle z^m x_{r\alpha}^{\widehat{\nu}}(z). \quad (2.12)$$

For any dominant integral weight $\lambda \in (\mathfrak{h}_{(0)})^*$ let U_{λ} be the finite-dimensional irreducible $\mathfrak{g}_{(0)}$ -module of highest weight λ . The generalized Verma module $N^{\widehat{\nu}}(\Lambda)$ associated with U_{λ} is defined by $N^{\widehat{\nu}}(\Lambda) = U(\widehat{\mathfrak{g}}[\widehat{\nu}]) \otimes_{U(\widehat{\mathfrak{g}}[\widehat{\nu}] \geq 0)} U_{\lambda}$, where the action of the Lie algebra

$$\widehat{\mathfrak{g}}[\widehat{\nu}]^{\geq 0} = \bigoplus_{n \in \frac{1}{2}\mathbb{Z}_{\geq 0}} (\mathfrak{g}_{(2n)} \otimes t^n) \oplus \mathbb{C}c$$

on U_{λ} is given by

$$\mathfrak{g}_{(2n)} \otimes t^n \cdot u = 0 \quad \text{for all } n > 0 \quad \text{and} \quad c \cdot u = ku \quad \text{for all } u \in U_{\lambda}.$$

Recall that the Poincaré–Birkhoff–Witt theorem for the universal enveloping algebra gives the vector space isomorphism

$$N^{\widehat{\nu}}(\Lambda) \cong U(\widehat{\mathfrak{g}}[\widehat{\nu}]^{<0}) \otimes_{\mathbb{C}} U_{\lambda}, \quad \text{where} \quad \widehat{\mathfrak{g}}[\widehat{\nu}]^{<0} = \bigoplus_{n \in \frac{1}{2}\mathbb{Z}_{<0}} (\mathfrak{g}_{(2n)} \otimes t^n).$$

Let V denote the generalized Verma module $N^{\widehat{\nu}}(\Lambda)$ or its unique simple quotient $L^{\widehat{\nu}}(\Lambda)$. Then there exists a linear map

$$\begin{aligned} Y^{\widehat{\nu}} : \mathfrak{g}(-1) \otimes \mathbf{1} &\rightarrow (\text{End } V)[[z^{\frac{1}{2}}, z^{-\frac{1}{2}}]], \\ Y^{\widehat{\nu}}(x(-1) \otimes \mathbf{1}, z) &= x^{\widehat{\nu}}(z) = \sum_{n \in \frac{1}{2}\mathbb{Z}} x^{\widehat{\nu}}(n) z^{-n-1}, \end{aligned}$$

which defines a structure of $\widehat{\nu}$ -twisted $N(k\Lambda_0)$ -module over V ; see, e.g., [Li96]. Its coefficients $x^{\widehat{\nu}}(n)$ satisfy the identity in (2.2) as well.

2.3. Principal subspaces. Consider the nilpotent Lie subalgebra

$$\mathfrak{n} = \bigoplus_{\alpha \in R_+} \mathbb{C}x_{\alpha}$$

of \mathfrak{g} generated by all root vectors x_{α} which correspond to positive roots $\alpha \in R_+$. Let

$$\widetilde{\mathfrak{n}}[\widehat{\nu}] = \overline{\mathfrak{n}}[\widehat{\nu}] \oplus \mathbb{C}c, \quad \text{where} \quad \overline{\mathfrak{n}}[\widehat{\nu}] = \prod_{m \in \frac{1}{2}\mathbb{Z}, \alpha \in R_+} \mathbb{C}x_{\alpha(2m)} \otimes t^m,$$

be the corresponding subalgebra of $\widehat{\mathfrak{g}}[\widehat{\nu}]$. Denote by v_{Λ} the highest weight vector of $V = N^{\widehat{\nu}}(\Lambda), L^{\widehat{\nu}}(\Lambda)$. Following [FeSt94], define the *principal subspace* of $V = N^{\widehat{\nu}}(\Lambda), L^{\widehat{\nu}}(\Lambda)$ by

$$W_V^T = U(\overline{\mathfrak{n}}[\widehat{\nu}])v_{\Lambda}.$$

The principal subspace W_V^T coincides with

$$U(\overline{\mathfrak{n}}_{\alpha_l}[\widehat{\nu}])U(\overline{\mathfrak{n}}_{\alpha_{l-1}}[\widehat{\nu}]) \cdots U(\overline{\mathfrak{n}}_{\alpha_1}[\widehat{\nu}])v_{\Lambda}, \quad \text{where} \quad \overline{\mathfrak{n}}_{\alpha_i}[\widehat{\nu}] = \prod_{m \in \frac{1}{2}\mathbb{Z}} \mathbb{C}x_{\alpha_i(2m)} \otimes t^m,$$

(see [BuSa19, Lemma 3.1] and [G96, Lemma 3.1]). In addition, as with the untwisted case in [BuKo22, Sect. 5], for $V = N^{\widehat{\nu}}(\Lambda)$ we have the isomorphism of $\bar{\mathfrak{n}}[\widehat{\nu}]$ -modules

$$W_{N^{\widehat{\nu}}(\Lambda)}^T \cong U(\bar{\mathfrak{n}}[\widehat{\nu}]_{<0}), \quad \text{where} \quad \bar{\mathfrak{n}}[\widehat{\nu}]_{<0} = \coprod_{m \in \frac{1}{2}\mathbb{Z}_{<0}, \alpha \in R_+} \mathbb{C}x_{\alpha(2m)} \otimes t^m. \quad (2.13)$$

3. QUASI-PARTICLE BASES OF PRINCIPAL SUBSPACES

From now on, we consider the rectangular highest weights Λ , as defined by (1.1). We use the realization of the standard $\widehat{\mathfrak{g}}[\widehat{\nu}]$ -module $L^{\widehat{\nu}}(\Lambda) \cong U(\widehat{\mathfrak{g}}[\widehat{\nu}]) \cdot v_{\Lambda}$ as a submodule of $(V_Q^T)^{\otimes k}$ with the highest weight vector

$$v_{\Lambda} = v_{\Lambda_{j_k}} \otimes \cdots \otimes v_{\Lambda_{j_1}}, \quad \text{where} \quad j_s = \begin{cases} 0 & \text{for } 1 \leq s \leq k_0, \\ j & \text{for } k_0 + 1 \leq s \leq k. \end{cases} \quad (3.1)$$

3.1. Twisted quasi-particles. For positive integers r and $i = 1, \dots, l$ and $m \in \frac{1}{2}\mathbb{Z}$ we define a *quasi-particle* of *charge* r , *color* i and *energy* $-m$ as the coefficient $x_{r\alpha_i}^{\widehat{\nu}}(m)$ of the twisted vertex operator $x_{r\alpha_i}^{\widehat{\nu}}(z)$, as given by (2.11) (cf. [BuSa19, G96]). Our goal is to construct the so-called quasi-particle bases of the principal subspaces, i.e. the bases consisting of some monomials of quasi-particles applied on the highest weight vector. Charges of quasi-particles which appear in the basis of the principal subspaces $W_{L^{\widehat{\nu}}(\Lambda)}^T$ will be less than or equal to k since we have the integrability relations

$$x_{(k+1)\alpha_i}^{\widehat{\nu}}(z) = 0 \quad (3.2)$$

on the standard $\widehat{\mathfrak{g}}[\widehat{\nu}]$ -modules $L^{\widehat{\nu}}(\Lambda)$ of level k (cf. [Li96]). On the other hand, the elements of basis for $W_{N^{\widehat{\nu}}(\Lambda)}^T$ will contain quasi-particle of all charges.

By (2.11), it follows that in the case of $W_{N^{\widehat{\nu}}(\Lambda)}^T$ all quasi-particles satisfy

$$x_{r\alpha_i}^{\widehat{\nu}}(z)v_{\Lambda} \in \begin{cases} z^{-\frac{r}{2}} W_{N^{\widehat{\nu}}(\Lambda)}^T [[z^{1/2}]], & \text{when } \nu\alpha_i \neq \alpha_i, \\ W_{N^{\widehat{\nu}}(\Lambda)}^T [[z]], & \text{when } \nu\alpha_i = \alpha_i, \end{cases} \quad (3.3)$$

while in the case of $W_{L^{\widehat{\nu}}(\Lambda)}^T$ we have

$$x_{r\alpha_i}^{\widehat{\nu}}(z)v_{\Lambda} \in \begin{cases} z^{-\frac{r}{2} + \frac{1}{2} \sum_{s=1}^r \delta_{i,j_s}} W_{L^{\widehat{\nu}}(\Lambda)}^T [[z^{1/2}]], & \text{when } \nu\alpha_i \neq \alpha_i, \\ W_{L^{\widehat{\nu}}(\Lambda)}^T [[z]], & \text{when } \nu\alpha_i = \alpha_i. \end{cases} \quad (3.4)$$

The quasi-particle bases of principal subspaces will be expressed in the form

$$\begin{aligned} bv_{\Lambda} &= b^{\widehat{\nu}}(\alpha_l) \cdots b^{\widehat{\nu}}(\alpha_1) v_{\Lambda} \\ &= \underbrace{x_{n_{r_l^{(1)},l}\alpha_l}^{\widehat{\nu}}(m_{r_l^{(1)},l}) \cdots x_{n_{1,l}\alpha_l}^{\widehat{\nu}}(m_{1,l})}_{b^{\widehat{\nu}}(\alpha_l)} \cdots \underbrace{x_{n_{r_1^{(1)},1}\alpha_1}^{\widehat{\nu}}(m_{r_1^{(1)},1}) \cdots x_{n_{1,1}\alpha_1}^{\widehat{\nu}}(m_{1,1})}_{b^{\widehat{\nu}}(\alpha_1)} v_{\Lambda}, \end{aligned} \quad (3.5)$$

where for $i = 1, \dots, l$ and $t \in \mathbb{N}$ we have

$$0 \leq n_{r_i^{(1)},i} \leq \cdots \leq n_{1,i}, \quad r_i^{(1)} \geq r_i^{(2)} \geq \cdots \geq r_i^{(t)} \geq 0, \quad m_{r_i^{(1)},i} \leq \cdots \leq m_{1,i}.$$

Here $r_i^{(p)}$ for $1 \leq p \leq t$ represent the parts of the partition $\mathcal{R}_i = (r_i^{(1)}, r_i^{(2)}, \dots, r_i^{(t)})$ of some fixed positive number r_i . The partition \mathcal{R}_i is conjugate to the partition $\mathcal{R}'_i = (n_{r_i^{(1)},i}, \dots, n_{1,i})$. We now recall terminology from [BuSa19, G96]. For every color $i = 1, \dots, l$, the number r_i is said to be a *color-type*, a partition \mathcal{R}'_i a *charge-type*, a partition \mathcal{R}_i a *dual-charge-type* and a finite sequence of energies $\mathcal{E}_i = (m_{r_i^{(1)},i}, \dots, m_{1,i})$ an *energy-type* of the monomial $b_i = b^{\widehat{\nu}}(\alpha_i)$. Generalizing this notions to the quasi-particle monomial $b =$

$b^{\hat{\nu}}(\alpha_l) \cdots b^{\hat{\nu}}(\alpha_1)$ we define its *color-type* $\mathcal{C} = (r_l, \dots, r_1)$, *charge-type* $\mathcal{R}' = (\mathcal{R}'_l, \dots, \mathcal{R}'_1)$, *dual-charge-type* $\mathcal{R} = (\mathcal{R}_l, \dots, \mathcal{R}_1)$ and *energy-type* $\mathcal{E} = (\mathcal{E}_l, \dots, \mathcal{E}_1)$. Moreover, its *total charge* $\text{chg } b$ and *total energy* $\text{en } b$ are defined by

$$\text{chg } b = \sum_{i=1}^l r_i \quad \text{and} \quad \text{en } b = -m_{r_l^{(1)}, l} - \cdots - m_{1, l} - \cdots - m_{r_1^{(1)}, 1} - \cdots - m_{1, 1}.$$

Let M_{QP} denote the set of all quasi-particle monomials as in (3.5). We introduce the linear order " $<$ " on the set of all monomials from M_{QP} of fixed degree and $\mathfrak{h}_{(0)}$ -weight as follows. Let b and \bar{b} be quasi-particle monomials of color-types \mathcal{C} and $\bar{\mathcal{C}}$, charge-types \mathcal{R}' and $\bar{\mathcal{R}}'$ and energy-types \mathcal{E} and $\bar{\mathcal{E}}$, respectively. Then we write $bv_{\Lambda} < \bar{b}v_{\Lambda}$ if one of the following conditions holds:

- (1) $\text{chg } b > \text{chg } \bar{b}$
- (2) $\text{chg } b = \text{chg } \bar{b}$ and $\mathcal{C} < \bar{\mathcal{C}}$
- (3) $\mathcal{C} = \bar{\mathcal{C}}$ and $\text{en } b < \text{en } \bar{b}$
- (4) $\mathcal{C} = \bar{\mathcal{C}}$, $\text{en } b = \text{en } \bar{b}$ and $\mathcal{R}' < \bar{\mathcal{R}}'$
- (5) $\mathcal{R}' = \bar{\mathcal{R}}'$ and $\mathcal{E} < \bar{\mathcal{E}}$,

where for integer sequences we write $(x_p, \dots, x_1) < (y_r, \dots, y_1)$ if there exists s such that

$$x_1 = y_1, \dots, x_{s-1} = y_{s-1} \quad \text{and} \quad s = p + 1 < r \quad \text{or} \quad x_s < y_s. \quad (3.6)$$

3.2. Quasi-particle bases of principal subspaces. By (3.3) and (3.4), the vectors of the form (3.5), such that for $i = 1, \dots, l$ their quasi-particle energies satisfy

$$m_{p,i} \leq -\mu_i n_{p,i} - \sum_{s=1}^{n_{p,i}} \mu_i \delta_{i,j_s} \quad \text{with} \quad 1 \leq p \leq r_i^{(1)} \quad \text{and} \quad \mu_i := \frac{1}{2} \langle \alpha_{i(0)}, \alpha_{i(0)} \rangle,$$

span entire principal subspace. We shall now strengthen the constraints above by using relations among quasi-particles. First, by employing the commutator formula for twisted vertex operators on $N^{\hat{\nu}}(\Lambda)$ we generalize [BuSa19, Lemma 4.2] as follows.

Lemma 3.1. *Let n_1, n_2 be positive integers and $\alpha, \beta \in R_+$ positive simple roots.*

a) *If $\langle \alpha, \beta \rangle = -1 = \langle \nu\alpha, \beta \rangle$, we have*

$$(z_1 - z_2)^{\min\{n_1, n_2\}} x_{n_1\alpha}^{\hat{\nu}}(z_1) x_{n_2\beta}^{\hat{\nu}}(z_2) = (z_1 - z_2)^{\min\{n_1, n_2\}} x_{n_2\beta}^{\hat{\nu}}(z_2) x_{n_1\alpha}^{\hat{\nu}}(z_1).$$

b) *If $\langle \alpha, \beta \rangle = -1$ and $\langle \nu\alpha, \beta \rangle = 0$, we have*

$$(z_1^{1/2} - z_2^{1/2})^{\min\{n_1, n_2\}} x_{n_1\alpha}^{\hat{\nu}}(z_1) x_{n_2\beta}^{\hat{\nu}}(z_2) = (z_1^{1/2} - z_2^{1/2})^{\min\{n_1, n_2\}} x_{n_2\beta}^{\hat{\nu}}(z_2) x_{n_1\alpha}^{\hat{\nu}}(z_1).$$

The interaction between quasi-particles of the same color [BuSa19, Lemma 4.1] can be summarized as follows.

Lemma 3.2. *Let $i = 1, \dots, l$ and $v \in W_V^T$ for $V = N^{\hat{\nu}}(\Lambda), L^{\hat{\nu}}(\Lambda)$. For any charges n_1, n_2 such that $n_1 \geq n_2$ and integer $M = m_1 + m_2$ the $2\mu_i n_2$ vectors*

$$x_{n_1\alpha_i}^{\hat{\nu}}(m_1) x_{n_2\alpha_i}^{\hat{\nu}}(m_2) v, \quad x_{n_1\alpha_i}^{\hat{\nu}}(m_1 - 1) x_{n_2\alpha_i}^{\hat{\nu}}(m_2 + 1) v, \dots \\ \dots, x_{n_1\alpha_i}^{\hat{\nu}}(m_1 - 2\mu_i n_2 + 1) x_{n_2\alpha_i}^{\hat{\nu}}(m_2 + 2\mu_i n_2 - 1) v$$

can be expressed as a finite linear combination of the monomial vectors

$$x_{n_1\alpha_i}^{\hat{\nu}}(s_1) x_{n_2\alpha_i}^{\hat{\nu}}(s_2) v \quad \text{such that} \quad s_1 + s_2 = M \quad \text{and} \quad s_1 \leq m_1 - 2\mu_i n_2 \quad \text{or} \quad s_2 < m_2$$

and monomial vectors which contain a quasi-particle of color i and charge $n_1 + 1$. Moreover, if $n_1 = n_2$, the monomial vectors

$$x_{n_1\alpha_i}^{\hat{\nu}}(m_1) x_{n_1\alpha_i}^{\hat{\nu}}(m_2) v \quad \text{such that} \quad m_1 + m_2 = M \quad \text{and} \quad m_2 - 2\mu_i n_1 < m_1 \leq m_2$$

can be expressed as a finite linear combination of monomial vectors

$$x_{n_1\alpha_i}^{\widehat{\nu}}(s_1)x_{n_1\alpha_i}^{\widehat{\nu}}(s_2)v \quad \text{such that} \quad s_1 + s_2 = M \quad \text{and} \quad s_1 \leq s_2 - 2\mu_i n_1$$

and monomial vectors which contain a quasi-particle of color i and charge $n_1 + 1$.

Denote by $B_{W_{N^{\widehat{\nu}}(\Lambda)}}$ the set of all quasi-particle monomials of the form (3.5) such that their energies $m_{p,i}$ satisfy the following difference conditions for all $i = 1, \dots, l$:

$$m_{p,i} \leq (1 - 2p)\mu_i n_{p,i} - \langle \alpha_{i(0)}, \alpha_{i-1(0)} \rangle \sum_{q=1}^{r_{i-1}^{(1)}} \min \{n_{q,i-1}, n_{p,i}\} - \sum_{s=1}^{n_{p,i}} \mu_i \delta_{i,j_s}, \quad (3.7)$$

$$m_{p+1,i} \leq m_{p,i} - 2\mu_i n_{p,i} \quad \text{if} \quad n_{p+1,i} = n_{p,i}, \quad (3.8)$$

where we set $r_0^{(1)} := 0$ and j_s are given by (3.1). As for the standard module, we denote by $B_{W_{L^{\widehat{\nu}}(\Lambda)}}$ the subset of $B_{W_{N^{\widehat{\nu}}(\Lambda)}}$ such that its monomials contain only quasi-particles of charge less than or equal to k ; recall (3.2). For $V = N^{\widehat{\nu}}(\Lambda)$, $L^{\widehat{\nu}}(\Lambda)$ we have

Theorem 3.3. *For any highest weight Λ as in (1.1) the set $\mathcal{B}_{W_V} = \{bv_\Lambda \mid b \in B_{W_V}\}$ forms a basis of the principal space W_V^T .*

The proof that \mathcal{B}_{W_V} spans the principal subspace goes in parallel with [BuKo22, Sect. 5.5] and [G96, Sect. 5]. Furthermore, the linear independence of $\mathcal{B}_{W_{N^{\widehat{\nu}}(\Lambda)}}$ can be verified by arguing as in [Bu14, Sect. 3]. As for the set $\mathcal{B}_{W_{L^{\widehat{\nu}}(\Lambda)}}$, in comparison with [BuSa19] (where the analogous basis for the highest weights of the form $k\Lambda_0$ is established), the proof of linear independence in the rectangular case requires a certain new ingredient, the map $[\pi_0\lambda_j]$ which connects modules $L^{\widehat{\nu}}(\Lambda_0)$ and $L^{\widehat{\nu}}(\Lambda_j)$. In Subsection 3.3 below, we recall the Georgiev projection $\pi_{\mathcal{R}}$ and the aforementioned map $[\pi_0\lambda_j]$. Finally, in Subsection 3.4 we employ them to finalize the proof.

3.3. Georgiev's projection and the simple current map. Let \mathcal{R} be the dual-charge-type of the quasi-particle monomial from (3.5). Consider Georgiev's projection $\pi_{\mathcal{R}}$ of the principal subspace $W_{L^{\widehat{\nu}}(\Lambda)}^T$ on the tensor product space

$$W_{L^{\widehat{\nu}}(\Lambda_{j_k})}^T(r_l^{(k)}, \dots, r_1^{(k)}) \otimes \cdots \otimes W_{L^{\widehat{\nu}}(\Lambda_{j_1})}^T(r_l^{(1)}, \dots, r_1^{(1)}) \subset (W_{L^{\widehat{\nu}}(\Lambda_j)}^T)^{\otimes k_j} \otimes (W_{L^{\widehat{\nu}}(\Lambda_0)}^T)^{\otimes k_0},$$

where $W_{L^{\widehat{\nu}}(\Lambda_{j_s})}^T(r_l^{(s)}, \dots, r_1^{(s)})$, $s = 1, \dots, k$, denote the $\mathfrak{h}_{(0)}$ -weight subspaces of the level one principal subspace $W_{L^{\widehat{\nu}}(\Lambda_{j_s})}^T$ consisting of vectors of charges $r_l^{(s)}, \dots, r_1^{(s)}$; see [BuSa19, G96] for more details. We will also use the symbol $\pi_{\mathcal{R}}$ to denote the generalization of the projection to the space of formal power series with coefficients in $(W_{L^{\widehat{\nu}}(\Lambda_j)}^T)^{\otimes k_j} \otimes (W_{L^{\widehat{\nu}}(\Lambda_0)}^T)^{\otimes k_0}$. As in [BuSa19, Sect. 5.1], one can use the integrability relations (3.2) at the level $k = 1$, $x_{2\alpha_i}^{\widehat{\nu}}(z)^2 = 0$ for $i = 1, \dots, l$, to express the projection of the monomial (3.5) as the coefficient of the product of the corresponding vertex operators applied on the highest weight vector (3.1).

Denote by $[\pi_0\lambda_j]$ the identity map on the vector space V_Q^T endowed with two different structures of level one standard $\widehat{\mathfrak{g}}[\widehat{\nu}]$ -modules, $L^{\widehat{\nu}}(\Lambda_0)$ and $L^{\widehat{\nu}}(\Lambda_j)$ (cf. [DLiM96, Li96, Li12]). By the construction of these modules follows that, for the suitably normalized highest weight vector v_{Λ_j} , we have for any root α , $m \in \frac{1}{2}\mathbb{Z}$ and $i = 1, \dots, l$ the identities

$$[\pi_0\lambda_j] v_{\Lambda_0} = v_{\Lambda_j}, \quad (3.9)$$

$$x_{\alpha}^{\widehat{\nu}}(m) [\pi_0\lambda_j] = [\pi_0\lambda_j] x_{\alpha}^{\widehat{\nu}}(m + \langle \lambda_{j(0)}, \alpha_{(0)} \rangle), \quad (3.10)$$

$$[\pi_0\lambda_j] \circ e_{\alpha_i} = e_{\alpha_i} \circ [\pi_0\lambda_j]. \quad (3.11)$$

Suppose b is the quasi-particle monomial from (3.5). The equalities (3.9) and (3.10) imply that $\pi_{\mathcal{R}}b([\pi_0\lambda_j]v_{\Lambda_0})^{\otimes k_j} \otimes v_{\Lambda_0}^{\otimes k_0}$ coincides with the coefficient $([\pi_0\lambda_j]^{\otimes k_j} \otimes 1^{\otimes k_0}) (\pi_{\mathcal{R}}b^+v_{\Lambda_0}^{\otimes k})$ of the variables

$$\begin{aligned} & z_{r_i^{(1)},i}^{-m_{r_i^{(1)},i}} z_{r_i^{(1)},i}^{-n_{r_i^{(1)},i}} \cdots z_{r_i^{(k_0+1)+1,i}}^{-m_{r_i^{(k_0+1)+1,i}}} z_{r_i^{(k_0+1)+1,i}}^{-n_{r_i^{(k_0+1)+1,i}}} \cdots \\ & \cdots z_{r_i^{(k_0+1)},i}^{-m_{r_i^{(k_0+1)},i}} z_{r_i^{(k_0+1)},i}^{-n_{r_i^{(k_0+1)},i}} + \langle \lambda_{j(0)}, \alpha_{i(0)} \rangle (n_{r_i^{(k_0+1)},i} - k_0) \cdots z_{1,i}^{-m_{1,i}} z_{1,i}^{-n_{1,i}} + \langle \lambda_{j(0)}, \alpha_{i(0)} \rangle (n_{1,i} - k_0) \end{aligned}$$

in the image of the (generalized) Georgiev projection

$$\pi_{\mathcal{R}} \left(x_{n_{r_l^{(1)},l} \alpha_l}^{\widehat{\nu}} (z_{r_l^{(1)},l}) \cdots x_{n_{1,1} \alpha_1}^{\widehat{\nu}} (z_{1,1}) ([\pi_0\lambda_j]v_{\Lambda_0})^{\otimes k_j} \otimes v_{\Lambda_0}^{\otimes k_0} \right).$$

Here we denote by b^+ the quasi-particle monomial

$$b^+ = b_{\alpha_l}^{\widehat{\nu}^+} \cdots b_{\alpha_1}^{\widehat{\nu}^+} \quad (3.12)$$

such that its factors $b_{\alpha_i}^{\widehat{\nu}^+}$, $i = 1, \dots, l$, are given by

$$\begin{aligned} b_{\alpha_i}^{\widehat{\nu}^+} &= x_{n_{r_i^{(1)},i} \alpha_i} (m_{r_i^{(1)},i}) \cdots x_{n_{r_i^{(k_0+1)+1,i} \alpha_i} (m_{r_i^{(k_0+1)+1,i}}) \cdots \\ & \cdots x_{n_{r_i^{(k_0+1)},i} \alpha_i} (m_{r_i^{(k_0+1)},i} + \langle \lambda_{j(0)}, \alpha_{i(0)} \rangle (n_{r_i^{(k_0+1)},i} - k_0)) \cdots \\ & \cdots x_{n_{1,i} \alpha_i} (m_{1,i} + \langle \lambda_{j(0)}, \alpha_{i(0)} \rangle (n_{1,i} - k_0)). \end{aligned} \quad (3.13)$$

Finally, it remains to observe that by (3.7) and (3.8) the energies of $b^+v_{k\Lambda_0}$ satisfy the initial and difference conditions for the quasi-particle basis of principal subspace $W_{L^{\widehat{\nu}}(k\Lambda_0)}^T$, as given by [BuSa19, Thm. 5.1].

3.4. Proof of linear independence. We are now ready to complete the proof of Theorem 3.3. Suppose that the quasi-particle monomial b of charge-type \mathcal{R}' and dual-charge-type \mathcal{R} is the smallest monomial, with respect to the linear ordering from Subsection 3.1, in the linear combination (indexed by some finite set A),

$$\sum_{a \in A} c_a b_a v_{\Lambda} = 0. \quad (3.14)$$

We can assume that all quasi-particle monomials $b_a \in B_{W_{L^{\widehat{\nu}}(\Lambda)}}$ are of the same color-type and that the scalars $c_a \in \mathbb{C}$ are nonzero. Let us apply the projection $\pi_{\mathcal{R}}$ to (3.14). By its definition, all nonzero summands which appear in

$$\sum_{a \in A} c_a \pi_{\mathcal{R}} b_a v_{\Lambda} = 0$$

contain monomials b_a of the same charge-type \mathcal{R}' . Using (3.9) we find

$$0 = \sum_{a \in A} c_a \pi_{\mathcal{R}} b_a ([\pi_0\lambda_j]v_{\Lambda_0})^{\otimes k_j} \otimes v_{\Lambda_0}^{\otimes k_0} = ([\pi_0\lambda_j]^{\otimes k_j} \otimes 1^{\otimes k_0}) \sum_{a \in A} c_a \pi_{\mathcal{R}} b_a^+ v_{\Lambda_0}^{\otimes k}$$

with b_a^+ as in (3.12) and (3.13). Dropping the invertible operator $[\pi_0\lambda_j]^{\otimes k_j} \otimes 1^{\otimes k_0}$, we get

$$\sum_{a \in A} c_a \pi_{\mathcal{R}} b_a^+ v_{\Lambda_0}^{\otimes k} = 0.$$

Thus, using the quasi-particle basis for the weights of the form $k\Lambda_0$, as given by [BuSa19, Thm. 5.1], we obtain $c_a = 0$, so that the assertion of Theorem 3.3 follows.

4. COMBINATORIAL BASES OF STANDARD MODULES AND PARA-FERMIONIC SPACES

In this section, we use the quasi-particle bases for principal subspaces to construct combinatorial bases of standard modules and their parafermionic spaces. Throughout the section, Λ denotes the rectangular highest weight; recall (1.1).

4.1. Standard modules. Let $B_{U(\widehat{\mathfrak{h}}[\nu]^-)}$ be the Poincaré–Birkhoff–Witt basis of the irreducible $\widehat{\mathfrak{h}}[\nu]_{\frac{1}{2}\mathbb{Z}}$ -module $U(\widehat{\mathfrak{h}}[\nu]^-)$ of level k which consists of all elements $h = h_{\alpha_l} \cdots h_{\alpha_1}$ such that for $i = 1, \dots, l$ we have

$$h_{\alpha_i} = (\alpha_{i(2s_{t_i,i})}(-s_{t_i,i}))^{r_{t_i,i}} \cdots (\alpha_{i(2s_{1,i})}(-s_{1,i}))^{r_{1,i}}, \quad \text{where} \quad (4.1)$$

$$t_i \in \mathbb{Z}_{\geq 0} \quad \text{and} \quad r_{p,i}, s_{p,i} \in \mathbb{N} \quad \text{are such that} \quad s_{1,i} \leq \dots \leq s_{t_i,i} \quad \text{for all} \quad p = 1, \dots, t_i.$$

We shall now generalize the definition of the linear order from Subsection 3.1 to the set

$$\left\{ e_\mu h b v_\Lambda \mid \mu \in Q, h \in B_{U(\widehat{\mathfrak{h}}[\nu]^-)}, b \in M'_{QP} \right\}, \quad (4.2)$$

where M'_{QP} denotes the subset of M_{QP} which consists of all vectors of the form (3.5) such that their quasi-particle charges are less than or equal to $k - 1$. Let $e_\mu h b v_\Lambda$ and $e_{\mu'} \bar{h} \bar{b} v_\Lambda$ be any two elements of the set (4.2) of the same degree and $\mathfrak{h}_{(0)}$ -weight. We shall write $e_\mu h b v_\Lambda < e_{\mu'} \bar{h} \bar{b} v_\Lambda$ if we have $b v_\Lambda < \bar{b} v_\Lambda$ (i.e., in other words, if one of the conditions (1)–(5) holds) or if the energy types of b and \bar{b} coincide and we have $h < \bar{h}$. The order on $B_{U(\widehat{\mathfrak{h}}[\nu]^-)}$ is defined as follows. For any two elements $h = h_{\alpha_l} \cdots h_{\alpha_1}$ and $\bar{h} = \bar{h}_{\alpha_l} \cdots \bar{h}_{\alpha_1}$ of the basis $B_{U(\widehat{\mathfrak{h}}[\nu]^-)}$ we write $h < \bar{h}$ if one of the conditions holds:

- (a) $(r_{t_l,l}, \dots, r_{1,1}) < (\bar{r}_{t_l,l}, \dots, \bar{r}_{1,1})$
- (b) $(r_{t_l,l}, \dots, r_{1,1}) = (\bar{r}_{t_l,l}, \dots, \bar{r}_{1,1})$ and $(s_{t_l}, \dots, s_1) < (\bar{s}_{t_l}, \dots, \bar{s}_1)$.

The integer sequences in (a) and (b) have the same meaning as in (4.1) and we compare them as in (3.6).

In order to construct the quasi-particle basis of standard module $L^{\widehat{\nu}}(\Lambda)$ we need the canonical isomorphism of d -graded vector spaces given by Lepowsky–Wilson in [LW84],

$$\begin{aligned} U(\widehat{\mathfrak{h}}[\nu]^-) \otimes L^{\widehat{\nu}}(\Lambda)^{\widehat{\mathfrak{h}}[\nu]^+} &\xrightarrow{\cong} L^{\widehat{\nu}}(\Lambda) \\ h \otimes u &\longmapsto h \cdot u, \end{aligned} \quad (4.3)$$

where $L^{\widehat{\nu}}(\Lambda)^{\widehat{\mathfrak{h}}[\nu]^+}$ denotes the *vacuum subspace* of standard module $L^{\widehat{\nu}}(\Lambda)$, i.e.

$$L^{\widehat{\nu}}(\Lambda)^{\widehat{\mathfrak{h}}[\nu]^+} = \left\{ v \in L^{\widehat{\nu}}(\Lambda) \mid \widehat{\mathfrak{h}}[\nu]^+ \cdot v = 0 \right\}. \quad (4.4)$$

Moreover, the construction relies the vertex operator formula

$$\frac{1}{p!} (2zx_{\alpha}^{\widehat{\nu}}(z))^p = \frac{1}{q!} \epsilon_{C_0}(\alpha, -\alpha)^{-q} E^{-}(-\alpha, z) (2zx_{-\alpha}^{\widehat{\nu}}(z))^q E^{+}(-\alpha, z) e_{\alpha} z^{\mu_{\alpha} + \alpha_{(0)}}, \quad (4.5)$$

where $p, q \geq 0$ are such that $k = p + q$ and $\mu_{\alpha} = \langle \alpha_{(0)}, \alpha_{(0)} \rangle / 2$ (cf. [OTa22]). Formula (4.5) implies that the quasi-particle of color i and charge k acts on $L^{\widehat{\nu}}(\Lambda)$ as an operator

$$x_{k\alpha_i}^{\widehat{\nu}}(m) = e_{\alpha_i}^k h \quad \text{for some} \quad h \in U(\widehat{\mathfrak{h}}[\nu]).$$

Consequently, we do not need quasi-particles of the highest charge k to form the basis of the standard module. Let $Q_{(0)} = \{\alpha_{(0)} \mid \alpha \in Q\}$. We have

Theorem 4.1. *For any highest weight Λ as in (1.1) the set*

$$\mathcal{B}_{L^{\widehat{\nu}}(\Lambda)} = \left\{ e_\mu h b v_\Lambda \mid \mu \in Q_{(0)}, h \in B_{U(\widehat{\mathfrak{h}}[\nu]^-)}, b \in B_{W_{L^{\widehat{\nu}}(\Lambda)}} \cap M'_{QP} \right\}$$

forms a basis of the standard module $L^{\widehat{\nu}}(\Lambda)$.

Proof. The proof that $\mathcal{B}_{L^{\hat{\nu}}(\Lambda)}$ spans the standard module $L^{\hat{\nu}}(\Lambda)$ relies on the quasi-particle relations (4.5) and Lemmas 3.1 and 3.2. It can be carried out by arguing as in [BuKoP21, Lemma 2.3], so we omit its details.

As for the linear independence, its proof employs the Georgiev-type projection

$$\pi_{\mathcal{R}_{\alpha_1}} : (L(\Lambda))_{r_1} \rightarrow L(\Lambda_{j_k})_{r_1^{(1)}} \otimes \dots \otimes L(\Lambda_{j_1})_{r_1^{(k)}},$$

with respect to the decomposition

$$\begin{aligned} L^{\hat{\nu}}(\Lambda) &= \coprod_{r \in \mathbb{Z}} L^{\hat{\nu}}(\Lambda)_r \subset L^{\hat{\nu}}(\Lambda_{j_k}) \otimes \dots \otimes L^{\hat{\nu}}(\Lambda_{j_1}), \quad \text{where} \\ L^{\hat{\nu}}(\Lambda)_r &= \coprod_{r_2, \dots, r_l \in \mathbb{Z}} L^{\hat{\nu}}(\Lambda)_{k\Lambda|_{\mathfrak{h}(0)} + r_l\alpha_l + \dots + r_2\alpha_2 + r\alpha_1}. \end{aligned} \quad (4.6)$$

The projection $\pi_{\mathcal{R}_{\alpha_1}}$, as well as its generalization to the space of formal power series with coefficients in (4.6) (which we again denote by the same symbol), are uniquely determined by the dual-charge type $\mathcal{R}_{\alpha_1} = (r_1^{(1)}, r_1^{(2)}, \dots, r_1^{(k)})$ and the color-type $r_1 = \sum_{t=1}^{k_{\alpha_1}} r_1^{(t)}$ with respect to the color $i = 1$. Consider the projection of the vector

$$\begin{aligned} &e_{\mu} (\alpha_{l(2s_{t_l,l})}(-s_{t_l,l}))^{r_{t_l,l}} \dots (\alpha_{1(2s_{1,1})}(-s_{1,1}))^{r_{1,1}} \\ &\times x_{n_{r_l^{(1)},l}\alpha_l}^{\hat{\nu}}(m_{r_l^{(1)},l}) \dots x_{n_{1,l}\alpha_l}^{\hat{\nu}}(m_{1,l}) \dots x_{n_{r_1^{(1)},1}\alpha_1}^{\hat{\nu}}(m_{r_1^{(1)},1}) \dots x_{n_{1,1}\alpha_1}^{\hat{\nu}}(m_{1,1}) v_{\Lambda}, \end{aligned}$$

such that its submonomial with respect to color $i = 1$,

$$b^{\hat{\nu}}(\alpha_1) = x_{n_{r_1^{(1)},1}\alpha_1}^{\hat{\nu}}(m_{r_1^{(1)},1}) \dots x_{n_{1,1}\alpha_1}^{\hat{\nu}}(m_{1,1}),$$

is of dual-charge-type \mathcal{R}_{α_1} . It coincides with the corresponding coefficient in

$$\begin{aligned} &\pi_{\mathcal{R}_{\alpha_1}} e_{\mu} (\alpha_{l,-}(w_{t_l,l}))^{r_{t_l,l}} \dots (\alpha_{1,-}(w_{1,1}))^{r_{1,1}} \\ &\times x_{n_{r_l^{(1)},l}\alpha_l}^{\hat{\nu}}(z_{r_l^{(1)},l}) \dots x_{n_{1,l}\alpha_l}^{\hat{\nu}}(z_{1,l}) \dots x_{n_{r_1^{(1)},1}\alpha_1}^{\hat{\nu}}(z_{r_1^{(1)},1}) \dots x_{n_{1,1}\alpha_1}^{\hat{\nu}}(z_{1,1}) v_{\Lambda}, \end{aligned}$$

where $\alpha_{i,-}(z) = \sum_{m < 0} \alpha_{i(2m)}(m) z^{-m-1}$ for $i = 1, \dots, l$.

Suppose that there exists a finite linear combination

$$\sum c_{\mu,h,b} e_{\mu} h b v_{\Lambda} = 0 \quad (4.7)$$

of vectors $e_{\mu} h b v_{\Lambda} \in \mathcal{B}_{L^{\hat{\nu}}(\Lambda)}$ of the same degree and $\mathfrak{h}(0)$ -weight such that all coefficients $c_{\mu,h,b} \in \mathbb{C}$ are nonzero. Choose any monomial vector in (4.7) of the form

$$e_{\mu} h b v_{\Lambda} \quad \text{with} \quad \text{chg } b(\alpha_1) = r_1 \quad (4.8)$$

and the maximal charge-type \mathcal{R}'_{α_1} . As before, in (4.8) we write $b(\alpha_1)$ for the submonomial of b in color $i = 1$. Its dual-charge-type $\mathcal{R}_{\alpha_1} = (r_1^{(1)}, \dots, r_1^{(p)})$ in color $i = 1$, where $p < k$ and $r_1 = r_1^{(1)} + \dots + r_1^{(p)}$, uniquely determines the Georgiev-type projection $\pi_{\mathcal{R}_{\alpha_1}}$ (with $r_1^{(t)} = 0$ for $t > p$). The projection $\pi_{\mathcal{R}_{\alpha_1}}$ ensures that only nonzero summands in

$$\sum c_{\mu,h,b} \pi_{\mathcal{R}_{\alpha_1}} e_{\mu} h b v_{\Lambda} = 0 \quad (4.9)$$

are those of the form as in (4.8). Indeed, from

$$e_{\alpha_1} (v_{\Lambda_{j_k}} \otimes \dots \otimes v_{\Lambda_{j_1}}) = e_{\alpha_1} v_{\Lambda_{j_k}} \otimes \dots \otimes e_{\alpha_1} v_{\Lambda_{j_1}},$$

and

$$e_{\mu} h b v_{\Lambda} \in \coprod_{\substack{r_1, \dots, r_{k-1} \in \mathbb{Z} \\ r_k > 0}} L^{\hat{\nu}}(\Lambda_{j_k})_{r_1} \otimes \dots \otimes L^{\hat{\nu}}(\Lambda_{j_1})_{r_k},$$

follows that $\pi_{\mathcal{R}_{\alpha_1}}$ annihilates all vectors $e_{\mu} h b v_{\Lambda}$ in (4.7) such that $\text{chg } b(\alpha_1) < r_1$.

By using the identities (3.9)–(3.11) and arguing as in Subsection 3.4 we write (4.9) as

$$\sum c_{\mu,h,b} \pi_{\mathcal{R}_{\alpha_1}} e_{\mu} h b^+ v_{k\Lambda_0} = 0, \quad (4.10)$$

where the monomials b^+ are found as in (3.12) and (3.13). Hence we can now proceed with the iterated use of the constant terms of twisted Δ -maps from [Li96] and the bijective map e_{α_1} as in [BuSa19, Sect. 5.4], until we reduce (4.10) to a linear combination of the vectors of the form $\pi_{\mathcal{R}_{\alpha_1}} e_{\mu} h b v_{k\Lambda_0}$ which satisfy the initial and difference conditions and $\text{ch}g b(\alpha_1) = 0$; see [BuSa19, Thm. 5.1] for more details. Next, we apply the same procedure for the remaining colors $i = 2, \dots, l$. Finally, we find that all coefficients $c_{\mu,h,b}$ in (4.7) are zero, thus verifying the linear independence of the set $\mathcal{B}_{L^{\hat{\nu}}(\Lambda)}$. \square

4.2. Vacuum spaces. Following J. Lepowsky and R. Wilson [LW81], for a quasi-particle monomial of charge-type $\mathcal{R}' = (n_{r_l^{(1)},l}, \dots, n_{1,1})$ we define \mathcal{Z} -operators

$$\begin{aligned} \mathcal{Z}_{\mathcal{R}'}(z_{r_l^{(1)},l}, \dots, z_{1,1}) &= E^-(\alpha_l, z_{r_l^{(1)},l})^{n_{r_l^{(1)},l}/k} \cdots E^-(\alpha_1, z_{1,1})^{n_{1,1}/k} x_{\mathcal{R}'}^{\hat{\nu}}(z_{r_l^{(1)},l}, \dots, z_{1,1}) \\ &\quad \times E^+(\alpha_l, z_{r_l^{(1)},l})^{n_{r_l^{(1)},l}/k} \cdots E^+(\alpha_1, z_{1,1})^{n_{1,1}/k}, \end{aligned}$$

where

$$x_{\mathcal{R}'}^{\hat{\nu}}(z_{r_l^{(1)},l}, \dots, z_{1,1}) = x_{n_{r_l^{(1)},l}\alpha_l}^{\hat{\nu}}(z_{r_l^{(1)},l}) \cdots x_{n_{1,1}\alpha_1}^{\hat{\nu}}(z_{1,1}).$$

We shall write the above formal Laurent series as

$$\mathcal{Z}_{\mathcal{R}'}(z_{r_l^{(1)},l}, \dots, z_{1,1}) = \sum_{m_{r_l^{(1)},l}, \dots, m_{1,1} \in \mathbb{Z}} \mathcal{Z}_{\mathcal{R}'}(m_{r_l^{(1)},l}, \dots, m_{1,1}) z_{r_l^{(1)},l}^{-m_{r_l^{(1)},l}} \cdots z_{1,1}^{-m_{1,1}}.$$

Its coefficients act on the vacuum space, i.e. we have

$$\mathcal{Z}_{\mathcal{R}'}(m_{r_l^{(1)},l}, \dots, m_{1,1}): L^{\hat{\nu}}(\Lambda)^{\hat{\mathfrak{h}}[\nu]^+} \rightarrow L^{\hat{\nu}}(\Lambda)^{\hat{\mathfrak{h}}[\nu]^+}. \quad (4.11)$$

By (2.9) we also have the action of the Weyl group translations e_{α} on the vacuum space,

$$e_{\alpha}: L^{\hat{\nu}}(\Lambda)^{\hat{\mathfrak{h}}[\nu]^+} \rightarrow L^{\hat{\nu}}(\Lambda)^{\hat{\mathfrak{h}}[\nu]^+}.$$

Consider the projection

$$\pi^{\hat{\mathfrak{h}}[\nu]^+}: L^{\hat{\nu}}(\Lambda) \rightarrow L^{\hat{\nu}}(\Lambda)^{\hat{\mathfrak{h}}[\nu]^+} \quad (4.12)$$

defined by the direct sum decomposition

$$L^{\hat{\nu}}(\Lambda) = L^{\hat{\nu}}(\Lambda)^{\hat{\mathfrak{h}}[\nu]^+} \oplus \hat{\mathfrak{h}}[\nu]^- U(\hat{\mathfrak{h}}[\nu]^-) \cdot L^{\hat{\nu}}(\Lambda)^{\hat{\mathfrak{h}}[\nu]^+}.$$

By (4.11) it follows that the action of projection (4.12) satisfies

$$\pi^{\hat{\mathfrak{h}}[\nu]^+}: x_{\mathcal{R}'}^{\hat{\nu}}(z_{r_l^{(1)},l}, \dots, z_{1,1}) v_{\Lambda} \mapsto \mathcal{Z}_{\mathcal{R}'}(z_{r_l^{(1)},l}, \dots, z_{1,1}) v_{\Lambda}.$$

By employing the isomorphism (4.3) along with the projection (4.12), one easily obtains the following consequence of Theorem 4.1:

Theorem 4.2. *For any highest weight Λ as in (1.1) the set of vectors*

$$e_{\mu} \mathcal{Z}_{\mathcal{R}'}(m_{r_l^{(1)},l}, \dots, m_{1,1}) v_{\Lambda},$$

where $\mu \in Q_{(0)}$ and the charge-type \mathcal{R}' and the energy-type $(m_{r_l^{(1)},l}, \dots, m_{1,1})$ satisfy the initial and difference conditions for $\mathcal{B}_{W_{L^{\hat{\nu}}(\Lambda)}}$, forms a basis of the vacuum space $L^{\hat{\nu}}(\Lambda)^{\hat{\mathfrak{h}}[\nu]^+}$.

4.3. Parafermionic spaces. We now extend the definition of the parafermionic space of highest weight $k\Lambda_0$ from [OTa22] (see also [BuKoP21, G95] for the untwisted case) to any rectangular highest weight (1.1). The *parafermionic space* is defined as the quotient

$$L^{\hat{\nu}}(\Lambda)_{kQ}^{\hat{\mathfrak{h}}[\nu]^+} = L^{\hat{\nu}}(\Lambda)^{\hat{\mathfrak{h}}[\nu]^+} / \text{span} \left\{ (\rho(k\alpha) - 1) \cdot v \mid \alpha \in Q, v \in L^{\hat{\nu}}(\Lambda)^{\hat{\mathfrak{h}}[\nu]^+} \right\}, \quad (4.13)$$

where the map ρ is given by (2.5). Denote by $\pi_{kQ}^{\hat{\mathfrak{h}}[\nu]^+}$ the canonical projection

$$\pi_{kQ}^{\hat{\mathfrak{h}}[\nu]^+} : L^{\hat{\nu}}(\Lambda)^{\hat{\mathfrak{h}}[\nu]^+} \rightarrow L^{\hat{\nu}}(\Lambda)_{kQ}^{\hat{\mathfrak{h}}[\nu]^+}.$$

We have

$$L^{\hat{\nu}}(\Lambda)_{kQ}^{\hat{\mathfrak{h}}[\nu]^+} \cong \coprod_{\mu \in \Lambda + Q/kQ} L^{\hat{\nu}}(\Lambda)_{\mu}^{\hat{\mathfrak{h}}[\nu]^+}.$$

As in [OTa22], we assign to quasi-particle monomials of charge-type $\mathcal{R}' = (n_{r_l^{(1)},l}, \dots, n_{1,1})$ the Ψ -operators

$$\Psi_{\mathcal{R}'}(z_{r_l^{(1)},l}, \dots, z_{1,1}) = \mathcal{Z}_{\mathcal{R}'}(z_{r_l^{(1)},l}, \dots, z_{1,1}) z_{r_l^{(1)},l}^{-n_{r_l^{(1)},l} \alpha_{l(0)}/k} \dots z_{1,1}^{-n_{1,1} \alpha_{1(0)}/k} \epsilon_{\alpha_l}^{-n_{1,1} \alpha_l/k} \dots \epsilon_{\alpha_1}^{-n_{1,1} \alpha_1/k},$$

where $\epsilon_{\alpha} := \epsilon_{C_0}(\alpha, \cdot)$ is the 2-cocycle on the weight lattice P . In parallel with [OTa22, Lemma 12], we have

$$\begin{aligned} \Psi_{n_r \beta_r, \dots, n_1 \beta_1}(z_r, \dots, z_1) &= \prod_{1 \leq p < s \leq r} \left((z_s^{1/2} - z_p^{1/2})^{\langle n_s \beta_s, n_p \beta_p \rangle/k} (z_s^{1/2} + z_p^{1/2})^{\langle n_s \nu \beta_s, n_p \beta_p \rangle/k} \right. \\ &\quad \left. \times \epsilon_{C_0}(\beta_s, \beta_p)^{n_s n_p/k} \right) \Psi_{n_r \beta_r}(z_r) \dots \Psi_{n_1 \beta_1}(z_1). \end{aligned} \quad (4.14)$$

Since the Ψ -operators commute both with the action of the Heisenberg subalgebra $\hat{\mathfrak{h}}[\nu]_{\frac{1}{2}\mathbb{Z}}$ and with the action of $\rho(kQ)$, their coefficients $\psi_{\mathcal{R}'}(m_{r_l^{(1)},l}, \dots, m_{1,1})$, given by

$$\Psi_{\mathcal{R}'}(z_{r_l^{(1)},l}, \dots, z_{1,1}) = \sum_{m_{r_l^{(1)},l}, \dots, m_{1,1}} \psi_{\mathcal{R}'}(m_{r_l^{(1)},l}, \dots, m_{1,1}) z_{r_l^{(1)},l}^{-m_{r_l^{(1)},l} \alpha_{l(0)}/k} \dots z_{1,1}^{-m_{1,1} \alpha_{1(0)}/k},$$

are induced operators on the quotient (4.13). The summation on the right-hand side goes over all sequences $(m_{r_l^{(1)},l}, \dots, m_{1,1})$ such that $m_{p,i} \in \frac{n_{p,i}}{k} \langle \alpha_{i(0)}, \mu \rangle + \mu_i \mathbb{Z}$ for all $1 \leq p \leq r_i^{(1)}$ when $\psi_{\mathcal{R}'}(m_{r_l^{(1)},l}, \dots, m_{1,1})$ acts on the μ -weight subspace of $L^{\hat{\nu}}(\Lambda)^{\hat{\mathfrak{h}}[\nu]^+}$. The following relations, which connect the coefficients of Ψ -operators and \mathcal{Z} -operators, follow from [OTa22, Lemma 10] (see also [BuKoP21, Lemma 3.2]):

$$\begin{aligned} \mathcal{Z}_{\mathcal{R}'}(m_{r_l^{(1)},l}, \dots, m_{1,1}) \Big|_{L^{\hat{\nu}}(\Lambda)_{\mu}^{\hat{\mathfrak{h}}[\nu]^+}} &= \prod_{i=1}^l \prod_{p=1}^{r_i^{(1)}} \left(\epsilon_{C_0}(-\tfrac{1}{k} n_{p,i} \alpha_i, \mu)^{-1} \right. \\ &\quad \left. \times \psi_{\mathcal{R}'}(m_{r_l^{(1)},l} + \tfrac{1}{k} n_{r_l^{(1)},l} \langle \alpha_{l(0)}, \mu \rangle, \dots, m_{1,1} + \tfrac{1}{k} n_{1,1} \langle \alpha_{1(0)}, \mu \rangle) \Big|_{L^{\hat{\nu}}(\Lambda)_{\mu}^{\hat{\mathfrak{h}}[\nu]^+}} \right). \end{aligned}$$

Finally, Theorem 4.2 implies

Theorem 4.3. *For any highest weight Λ as in (1.1) the set of vectors*

$$\begin{aligned} \pi_{kQ}^{\hat{\mathfrak{h}}[\nu]^+} \mathcal{Z}_{\mathcal{R}'}(m_{r_l^{(1)},l}, \dots, m_{1,1}) v_{\Lambda} \\ = \psi_{\mathcal{R}'}(m_{r_l^{(1)},l} + \tfrac{1}{k} n_{r_l^{(1)},l} \langle \alpha_{l(0)}, \Lambda \rangle, \dots, m_{1,1} + \tfrac{1}{k} n_{1,1} \langle \alpha_{1(0)}, \Lambda \rangle) v_{\Lambda}, \end{aligned}$$

where the charge-type \mathcal{R}' and the energy-type $(m_{r_l^{(1)}}, \dots, m_{1,1})$ satisfy the initial and difference conditions for $\mathcal{B}_{W_{L^{\hat{\nu}}(\Lambda)}}$, forms a basis of the parafermionic space $L^{\hat{\nu}}(\Lambda)_{kQ}^{\hat{\mathfrak{h}}[\nu]^+}$.

5. CHARACTER FORMULAE AND COMBINATORIAL IDENTITIES

In this section, we use our main results, Theorems 3.3, 4.1, 4.2 and 4.3 to compute the character formulae of the corresponding bases. Furthermore, as an application, in the case of the principal subspace of the generalized Verma module we obtain two new families of combinatorial identities. Throughout the entire section, Λ denotes an arbitrary rectangular highest weight, i.e. the weight of the form (1.1).

5.1. Principal subspaces. Let $V = N^{\hat{\nu}}(\Lambda), L^{\hat{\nu}}(\Lambda)$. We define the character of the principal subspace W_V^T by

$$\text{ch } W_V^T = \sum_{m, r_1, \dots, r_l \geq 0} \dim W_{V(m, r_1, \dots, r_l)}^T q^m y_1^{r_1} \cdots y_l^{r_l},$$

where $W_{V(m, r_1, \dots, r_l)}^T$ is the weight subspace of the Cartan subalgebra $\mathfrak{h}_{(0)} \oplus \mathbb{C}c \oplus \mathbb{C}d$ of weight $-m\delta + r_1\alpha_1 + \cdots + r_l\alpha_l$.

For the dual-charge-type of quasi-particle monomial as in (3.5) let $p_i^{(s)}$ be the number of quasi-particles of color i and charge $s = 1, \dots, t$ which appear in the monomial. Furthermore, we organize the numbers $p_i^{(s)}$ into the sequence $\mathcal{P} = (\mathcal{P}_1, \dots, \mathcal{P}_l)$ such that $\mathcal{P}_i = (p_i^{(1)}, \dots, p_i^{(t)})$. Also, we write

$$(a; q)_r = \prod_{i=1}^r (1 - aq^{i-1}) \quad \text{for } r \geq 0 \quad \text{and} \quad (a; q)_\infty = \prod_{i \geq 1} (1 - aq^{i-1}).$$

In addition, to express the character formulae we shall need the following notation

$$\frac{1}{(q^{\mu_i}; q^{\mu_i})_r} = \sum_{s \geq 0} p_r(s) q^{\mu_i s},$$

where $p_r(s)$ is the number of partitions of s with most r parts. By arguing as in [G96, Sect.5] and using the difference conditions (3.7) and (3.8), which determine the quasi-particle basis of the principal subspace of standard module via Theorem 3.3, we find

Theorem 5.1. *For the affine Lie algebras $A_{2l-1}^{(2)}$ and $D_{l+1}^{(2)}$ we have*

$$\text{ch } W_{L^{\hat{\nu}}(\Lambda)}^T = \sum_{\mathcal{P}} \frac{q^{\frac{1}{2} \sum_{i,r=1}^l \sum_{m,n=1}^k \langle \alpha_i(0), \alpha_r(0) \rangle \min\{m,n\} p_i^{(m)} p_r^{(n)} + \tilde{p}_j}}{\prod_{i=1}^l \prod_{s=1}^k (q^{\mu_i}; q^{\mu_i})_{p_i^{(s)}}} \prod_{i=1}^l y_i^{n_i},$$

where $\tilde{p}_j = \sum_{s=k_0+1}^k (s - k_0) \mu_j p_j^{(s)}$ and $n_i = \sum_{s=1}^k s p_i^{(s)}$. The sum goes over all finite sequences $\mathcal{P} = (\mathcal{P}_1, \dots, \mathcal{P}_l)$ of lk nonnegative integers.

The character formula for the principal subspace of the generalized Verma module is obtained analogously:

Theorem 5.2. *For the affine Lie algebras $A_{2l-1}^{(2)}$ and $D_{l+1}^{(2)}$ we have*

$$\text{ch } W_{N^{\hat{\nu}}(\Lambda)}^T = \sum_{\mathcal{P}} \frac{q^{\frac{1}{2} \sum_{i,r=1}^l \sum_{m,n=1}^t \langle \alpha_i(0), \alpha_r(0) \rangle \min\{m,n\} p_i^{(m)} p_r^{(n)}}}{\prod_{i=1}^l \prod_{s=1}^t (q^{\mu_i}; q^{\mu_i})_{p_i^{(s)}}} \prod_{i=1}^l y_i^{n_i},$$

where $n_i = \sum_{s=1}^t s p_i^{(s)}$, and the sum goes over all finite sequences $\mathcal{P} = (\mathcal{P}_1, \dots, \mathcal{P}_l)$ of nonnegative integers with finite support.

Due to the vector space isomorphism (2.13), we can compute the character of $W_{N^{\hat{\nu}}(\Lambda)}^T$ from the Poincaré–Birkhoff–Witt basis of $U(\tilde{\mathfrak{n}}_+^{\leq 0})$ as well, thus getting

$$\text{ch } W_{N^{\hat{\nu}}(\Lambda)}^T = \frac{1}{\prod_{\alpha \in R_+} (\alpha; q^{\langle \alpha_{(0)}, \alpha_{(0)} \rangle / 2})_{\infty}}. \quad (5.1)$$

On the right-hand side we used the notation

$$(\alpha; q)_{\infty} = (qy_1^{a_1} y_2^{a_2} \cdots y_l^{a_l}; q)_{\infty},$$

for any positive root $\alpha \in R_+$ such that $\alpha_{(0)} = a_1 \alpha_{1(0)} + \cdots + a_l \alpha_{l(0)}$. Theorem 5.2 and the character formula (5.1) imply the following generalization of the identities from [Bu16]:

Corollary 5.3. *For the affine Lie algebras $A_{2l-1}^{(2)}$ and $D_{l+1}^{(2)}$ we have*

$$\frac{1}{\prod_{\alpha \in R_+} (\alpha; q^{\langle \alpha_{(0)}, \alpha_{(0)} \rangle / 2})_{\infty}} = \sum_{\mathcal{P}} \frac{q^{\frac{1}{2} \sum_{i,r=1}^l \sum_{m,n=1}^t \langle \alpha_{i(0)}, \alpha_{r(0)} \rangle \min\{m,n\} p_i^{(m)} p_r^{(n)}}}{\prod_{i=1}^l \prod_{s=1}^t (q^{\mu_i}; q^{\mu_i})_{p_i^{(s)}}} \prod_{i=1}^l y_i^{n_i},$$

where $n_i = \sum_{s=1}^t s p_i^{(s)}$, and the sum goes over all finite sequences $\mathcal{P} = (\mathcal{P}_1, \dots, \mathcal{P}_l)$ of nonnegative integers with finite support.

5.2. Standard modules. Let $\mathcal{R} = (r_l^{(1)}, \dots, r_1^{(k-1)})$ be the dual-charge-type of a quasi-particle monomial $b \in \mathcal{B}_{L^{\hat{\nu}}(\Lambda)}$. Consider the sequences $\mathcal{P}_i = (p_i^{(1)}, \dots, p_i^{(k-1)})$ given by

$$\mathcal{P}_i = (r_i^{(1)} - r_i^{(2)}, r_i^{(2)} - r_i^{(3)}, \dots, r_i^{(k-2)} - r_i^{(k-1)}, r_i^{(k-1)}) \quad \text{for } i = 1, \dots, l. \quad (5.2)$$

Note that the number $p_i^{(m)}$ equals the number of quasi-particles of color i and charge m in the quasi-particle monomial b . We define the character of $L^{\hat{\nu}}(\Lambda)$ by

$$\text{ch } L^{\hat{\nu}}(\Lambda) = \sum_{m, r_1, \dots, r_l \geq 0} \dim L^{\hat{\nu}}(\Lambda)_{(m, r_1, \dots, r_l)} q^m y_1^{r_1} \cdots y_l^{r_l},$$

where $L^{\hat{\nu}}(\Lambda)_{(m, r_1, \dots, r_l)}$ is the weight subspace of weight $-m\delta + r_1\alpha_1 + \cdots + r_l\alpha_l$, with respect to the Cartan subalgebra $\mathfrak{h}_{(0)} \oplus \mathbb{C}c \oplus \mathbb{C}d$. In a similar way, we can define the character of the vacuum space $L^{\hat{\nu}}(\Lambda)^{\hat{\mathfrak{h}}[\nu]^+}$. By the isomorphism in (4.3) we have

$$\text{ch } L^{\hat{\nu}}(\Lambda) = \frac{1}{\prod_{i=1}^l (q^{\mu_i}; q^{\mu_i})_{\infty}} \text{ch } L^{\hat{\nu}}(\Lambda)^{\hat{\mathfrak{h}}[\nu]^+}.$$

Hence Theorems 4.1 and 4.2 imply

Theorem 5.4. *For the affine Lie algebras $A_{2l-1}^{(2)}$ and $D_{l+1}^{(2)}$ we have*

$$\begin{aligned} \text{ch } L^{\hat{\nu}}(\Lambda) &= \frac{1}{\prod_{i=1}^l (q^{\mu_i}; q^{\mu_i})_{\infty}} \sum_{\mathcal{P}} \frac{q^{\frac{1}{2} \sum_{i,r=1}^l \sum_{m,n=1}^{k-1} \langle \alpha_{i(0)}, \alpha_{r(0)} \rangle \min\{m,n\} p_i^{(m)} p_r^{(n)} + \tilde{p}_j}}{\prod_{i=1}^l \prod_{s=1}^{k-1} (q^{\mu_i}; q^{\mu_i})_{p_i^{(s)}}} \\ &\quad \times \sum_{\alpha \in Q} q^{\langle \alpha_{(0)}, \frac{k}{2} \alpha_{(0)} + \Lambda + \sum_{s=1}^{k-1} s p_i^{(s)} \alpha_{i(0)} \rangle} \prod_{i=1}^l y_i^{h_{\lambda_i} (k\alpha_{(0)} + \Lambda + \sum_{s=1}^{k-1} s p_i^{(s)} \alpha_{i(0)})}, \end{aligned}$$

where the first sum goes over all finite sequences $\mathcal{P} = (\mathcal{P}_1, \dots, \mathcal{P}_l)$ of $l(k-1)$ nonnegative integers, $\tilde{p}_j = \sum_{s=k_0+1}^{k-1} (s - k_0) \mu_j p_j^{(s)}$ and h_{λ_i} are the fundamental coweights of $\mathfrak{g}_{(0)}$.

As an illustration of the theorem, we now give some examples of character formulae.

Example 5.5. For the affine Lie algebra $A_5^{(2)}$ we have

$$\begin{aligned}
ch L^{\hat{\nu}}(\Lambda_1) &= \frac{1}{\prod_{i=1}^3 (q^{\mu_i}; q^{\mu_i})_{\infty}} \sum_{\alpha \in Q} q^{\langle \alpha_{(0)}, \frac{1}{2}\alpha_{(0)} + \Lambda_1 \rangle} \prod_{i=1}^3 y_i^{h_{\lambda_i}(\alpha_{(0)} + \Lambda_1)}, \\
ch L^{\hat{\nu}}(2\Lambda_1) &= \frac{1}{\prod_{i=1}^3 (q^{\mu_i}; q^{\mu_i})_{\infty}} \sum_{(\mathcal{P}_1, \mathcal{P}_2, \mathcal{P}_3)} \frac{q^{\frac{1}{2}(p_1^{(1)^2} - p_1^{(1)}p_2^{(1)} + p_2^{(1)^2} - 2p_2^{(1)}p_3^{(1)} + 2p_3^{(1)^2}) + \frac{1}{2}p_1^{(1)}}}{\prod_{i=1}^3 (q^{\mu_i}; q^{\mu_i})_{p_i^{(1)}}} \\
&\quad \times \sum_{\alpha \in Q} q^{\langle \alpha_{(0)}, \alpha_{(0)} + 2\Lambda_1 + \sum_{i=1}^3 p_i^{(1)}\alpha_{i(0)} \rangle} \prod_{i=1}^3 y_i^{h_{\lambda_i}(2\alpha_{(0)} + 2\Lambda_1 + \sum_{i=1}^3 p_i^{(1)}\alpha_{i(0)})}, \\
ch L^{\hat{\nu}}(\Lambda_0 + \Lambda_1) &= \frac{1}{\prod_{i=1}^3 (q^{\mu_i}; q^{\mu_i})_{\infty}} \sum_{(\mathcal{P}_1, \mathcal{P}_2, \mathcal{P}_3)} \frac{q^{\frac{1}{2}(p_1^{(1)^2} - p_1^{(1)}p_2^{(1)} + p_2^{(1)^2} - 2p_2^{(1)}p_3^{(1)} + 2p_3^{(1)^2})}}{\prod_{i=1}^3 (q^{\mu_i}; q^{\mu_i})_{p_i^{(1)}}} \\
&\quad \times \sum_{\alpha \in Q} q^{\langle \alpha_{(0)}, \alpha_{(0)} + \Lambda_0 + \Lambda_1 + \sum_{i=1}^3 p_i^{(1)}\alpha_{i(0)} \rangle} \prod_{i=1}^3 y_i^{h_{\lambda_i}(2\alpha_{(0)} + \Lambda_0 + \Lambda_1 + \sum_{i=1}^3 p_i^{(1)}\alpha_{i(0)})}.
\end{aligned}$$

Example 5.6. For the affine Lie algebra $D_3^{(2)}$ we have

$$\begin{aligned}
ch L^{\hat{\nu}}(\Lambda_2) &= \frac{1}{\prod_{i=1}^2 (q^{\mu_i}; q^{\mu_i})_{\infty}} \sum_{\alpha \in Q} q^{\langle \alpha_{(0)}, \frac{1}{2}\alpha_{(0)} + \Lambda_2 \rangle} \prod_{i=1}^2 y_i^{h_{\lambda_i}(\alpha_{(0)} + \Lambda_2)}, \\
ch L^{\hat{\nu}}(2\Lambda_2) &= \frac{1}{\prod_{i=1}^2 (q^{\mu_i}; q^{\mu_i})_{\infty}} \sum_{(\mathcal{P}_1, \mathcal{P}_2)} \frac{q^{\frac{1}{2}(2p_1^{(1)^2} - 2p_1^{(1)}p_2^{(1)} + p_2^{(1)^2}) + \frac{1}{2}p_2^{(1)}}}{\prod_{i=1}^2 (q^{\mu_i}; q^{\mu_i})_{p_i^{(1)}}} \\
&\quad \times \sum_{\alpha \in Q} q^{\langle \alpha_{(0)}, \alpha_{(0)} + 2\Lambda_2 + \sum_{i=1}^2 p_i^{(1)}\alpha_{i(0)} \rangle} \prod_{i=1}^2 y_i^{h_{\lambda_i}(2\alpha_{(0)} + 2\Lambda_2 + \sum_{i=1}^2 p_i^{(1)}\alpha_{i(0)})}, \\
ch L^{\hat{\nu}}(\Lambda_0 + \Lambda_2) &= \frac{1}{\prod_{i=1}^2 (q^{\mu_i}; q^{\mu_i})_{\infty}} \sum_{(\mathcal{P}_1, \mathcal{P}_2)} \frac{q^{\frac{1}{2}(2p_1^{(1)^2} - 2p_1^{(1)}p_2^{(1)} + p_2^{(1)^2})}}{\prod_{i=1}^2 (q^{\mu_i}; q^{\mu_i})_{p_i^{(1)}}} \\
&\quad \times \sum_{\alpha \in Q} q^{\langle \alpha_{(0)}, \alpha_{(0)} + \Lambda_0 + \Lambda_2 + \sum_{i=1}^2 p_i^{(1)}\alpha_{i(0)} \rangle} \prod_{i=1}^2 y_i^{h_{\lambda_i}(2\alpha_{(0)} + \Lambda_0 + \Lambda_2 + \sum_{i=1}^2 p_i^{(1)}\alpha_{i(0)})}.
\end{aligned}$$

5.3. Parafermionic spaces. As in [OTa22], to define the character of the parafermionic space we use the parafermionic grading operator D on $L^{\hat{\nu}}(\Lambda)_{kQ}^{\hat{h}[\nu]^+}$ given by

$$D = -d - D^{\hat{h}[\nu]^+} \quad \text{and} \quad D^{\hat{h}[\nu]^+} \Big|_{L^{\hat{\nu}}(\Lambda)_{\mu}^{\hat{h}[\nu]^+}} = \frac{\langle \mu_{(0)}, \mu_{(0)} \rangle}{2k} - \frac{\langle \Lambda, \Lambda \rangle}{2k}. \quad (5.3)$$

Note that by (5.3) the conformal energy of the basis vector

$$\psi_{\mathcal{R}'}(\mathcal{E})v_{\Lambda} = \psi_{(n_{r_l^{(1)}, l}, \dots, n_{1,1})}(m_{r_l^{(1)}, l}, \dots, m_{1,1})v_{\Lambda}, \quad (5.4)$$

which corresponds to the quasi-particle monomial

$$x_{n_{r_l^{(1)}, l}, l}^{\hat{\nu}}(m_{r_l^{(1)}, l}) \dots x_{n_{1,1}, l}^{\hat{\nu}}(m_{1, l}) \dots x_{n_{r_1^{(1)}, 1}, 1}^{\hat{\nu}}(m_{r_1^{(1)}, 1}) \dots x_{n_{1,1}, 1}^{\hat{\nu}}(m_{1,1}),$$

is equal to

$$- \sum_{i=1}^l \sum_{u=1}^{r_i^{(1)}} m_{u,i} - \frac{k_j}{k} \sum_{t=1}^{k-1} t p_j^{(t)}$$

$$- \sum_{i=1}^l \sum_{u=1}^{r_i^{(1)}} \left(\frac{n_{u,i}^2 \mu_i}{k} + \frac{1}{k} \left\langle n_{u,i} \alpha_{i(0)}, \sum_{s=1}^{u-1} n_{s,i} \alpha_{i(0)} + \sum_{p=1}^{i-1} \sum_{s=1}^{r_p^{(1)}} n_{s,p} \alpha_{p(0)} \right\rangle \right),$$

where the second summand is due to the identity

$$-\frac{1}{k} \left\langle \sum_{t=1}^{k-1} t p_j^{(t)} \alpha_{j(0)}, \Lambda \right\rangle = -\frac{k_j}{k} \mu_j \sum_{t=1}^{k-1} t p_j^{(t)}.$$

Define the character of the parafermionic space $L^{\hat{\nu}}(\Lambda)_{kQ}^{\hat{h}[\nu]^+}$ by

$$\text{ch } L^{\hat{\nu}}(\Lambda)_{kQ}^{\hat{h}[\nu]^+} = \sum_{m, r_1, \dots, r_l} \dim L^{\hat{\nu}}(\Lambda)_{kQ}^{\hat{h}[\nu]^+}_{(m, r_1, \dots, r_l)} q^m,$$

where $L^{\hat{\nu}}(\Lambda)_{kQ}^{\hat{h}[\nu]^+}_{(m, r_1, \dots, r_l)}$ is the weight space spanned by monomial vectors (5.4) of conformal energy $-m$ and color-type (r_1, \dots, r_l) .

Recall the notation from (5.2) and introduce the following expressions:

$$D_{\mathcal{P}}(q) = \frac{1}{\prod_{i=1}^l \prod_{s=1}^{k-1} (q^{\mu_i}; q^{\mu_i})_{p_i^{(s)}}}, \quad B_{\mathcal{P}}(q) = q^{\mu_j \sum_{t=k_0+1}^{k-1} (t-k_0) p_j^{(t)}} q^{-\mu_j \frac{k_j}{k} \sum_{t=1}^{k-1} t p_j^{(t)}},$$

$$G_{\mathcal{P}}(q) = q^{\frac{1}{2} \sum_{i,r=1}^l \sum_{m,n=1}^{k-1} \langle \alpha_{i(0)}, \alpha_{r(0)} \rangle (\min\{m,n\} - \frac{mn}{k}) p_i^{(m)} p_r^{(n)}}.$$

From the above considerations and Theorem 4.3 we get

Theorem 5.7. *For the affine Lie algebras $A_{2l-1}^{(2)}$ and $D_{l+1}^{(2)}$ we have*

$$\text{ch } L^{\hat{\nu}}(\Lambda)_{kQ}^{\hat{h}[\nu]^+} = \sum_{\mathcal{P}} D_{\mathcal{P}}(q) G_{\mathcal{P}}(q) B_{\mathcal{P}}(q), \quad (5.5)$$

where the sum goes over all finite sequences $\mathcal{P} = (\mathcal{P}_1, \dots, \mathcal{P}_l)$ of $l(k-1)$ nonnegative integers such that $\mathcal{P}_i = (p_i^{(1)}, \dots, p_i^{(k-1)})$.

It is worth noting that the character formula (5.5) coincides with [Gep22, Eq. (14)]. At the end, let us give some examples of the parafermionic character formulae.

Example 5.8. *For the affine Lie algebra $A_5^{(2)}$ we have*

$$\text{ch } L^{\hat{\nu}}(2\Lambda_1)_{2Q}^{\hat{h}[\nu]^+} = \sum_{\mathcal{P}=(\mathcal{P}_1, \mathcal{P}_2, \mathcal{P}_3)} \frac{q^{\frac{1}{4}(p_1^{(1)2} - p_1^{(1)} p_2^{(1)} + p_2^{(1)2} - 2p_2^{(1)} p_3^{(1)} + 2p_3^{(1)2})}}{\prod_{i=1}^3 (q^{\mu_i}; q^{\mu_i})_{p_i^{(1)}}},$$

$$\text{ch } L^{\hat{\nu}}(\Lambda_0 + \Lambda_1)_{2Q}^{\hat{h}[\nu]^+} = \sum_{\mathcal{P}=(\mathcal{P}_1, \mathcal{P}_2, \mathcal{P}_3)} \frac{q^{\frac{1}{4}(p_1^{(1)2} - p_1^{(1)} p_2^{(1)} + p_2^{(1)2} - 2p_2^{(1)} p_3^{(1)} + 2p_3^{(1)2}) - \frac{1}{4} p_1^{(1)}}}{\prod_{i=1}^3 (q^{\mu_i}; q^{\mu_i})_{p_i^{(1)}}}.$$

Example 5.9. *For the affine Lie algebra $D_3^{(2)}$ we have*

$$\text{ch } L^{\hat{\nu}}(2\Lambda_2)_{2Q}^{\hat{h}[\nu]^+} = \sum_{\mathcal{P}=(\mathcal{P}_1, \mathcal{P}_2)} \frac{q^{\frac{1}{4}(2p_1^{(1)2} - 2p_1^{(1)} p_2^{(1)} + p_2^{(1)2})}}{\prod_{i=1}^2 (q^{\mu_i}; q^{\mu_i})_{p_i^{(1)}}},$$

$$\text{ch } L^{\hat{\nu}}(\Lambda_0 + \Lambda_2)_{2Q}^{\hat{h}[\nu]^+} = \sum_{\mathcal{P}=(\mathcal{P}_1, \mathcal{P}_2)} \frac{q^{\frac{1}{4}(2p_1^{(1)2} - 2p_1^{(1)} p_2^{(1)} + p_2^{(1)2}) - \frac{1}{4} p_2^{(1)}}}{\prod_{i=1}^2 (q^{\mu_i}; q^{\mu_i})_{p_i^{(1)}}}.$$

ACKNOWLEDGEMENT

The authors are grateful to Mirko Primc for numerous helpful discussions. The authors would also like to thank Ole Warnaar for bringing to their attention some useful references on combinatorial identities. This work has been supported in part by Croatian Science Foundation under the project UIP-2019-04-8488. The first author is partially supported by the QuantiXLie Centre of Excellence, a project cofinanced by the Croatian Government and European Union through the European Regional Development Fund - the Competitiveness and Cohesion Operational Programme (Grant KK.01.1.1.01.0004).

REFERENCES

- [BK04] B. N. Bakalov, V. G. Kac, *Twisted modules over lattice vertex algebras*, Heinz-Dietrich Doebner and Vladimir K. Dobrev, editors, *Lie theory and its applications in physics V*, pages 3–26. World Scientific (2004); [arXiv:math/0402315v3 \[math.QA\]](#).
- [Bu14] M. Butorac, *Combinatorial bases of principal subspaces for the affine Lie algebra of type $B_2^{(1)}$* , *J. Pure Appl. Algebra* **218** (2014), 424–447; [arXiv:1212.5920 \[math.QA\]](#).
- [Bu16] M. Butorac, *Quasi-particle bases of principal subspaces for the affine Lie algebras of types $B_l^{(1)}$ and $C_l^{(1)}$* , *Glas. Mat. Ser. III* **51** (2016), 59–108; [arXiv:1505.00450 \[math.QA\]](#).
- [BuKo22] M. Butorac, S. Kožić, *Principal subspaces for the affine Lie algebras in types D , E and F* , *J. Algebraic Combin.* **56** (2022), 1063–1096; [arXiv:1902.10794 \[math.QA\]](#).
- [BuKoP21] M. Butorac, S. Kožić, M. Primc, *Parafermionic bases of standard modules for affine Lie algebras*, *Math. Z.* **298** (2021), 1003–1032; [arXiv:2002.00435v2 \[math.QA\]](#).
- [BuSa19] M. Butorac, C. Sadowski, *Combinatorial bases of principal subspaces of modules for twisted affine Lie algebras of type $A_{2l-1}^{(2)}$, $D_l^{(2)}$, $E_6^{(2)}$ and $D_4^{(3)}$* , *New York J. Math.* **25** (2019), 71–106; [arXiv:1803.06733v2 \[math.QA\]](#).
- [CLM14] C. Calinescu, J. Lepowsky, A. Milas, *Vertex-algebraic structure of principal subspaces of standard $A_2^{(2)}$ -modules, I*, *Internat. J. Math.* **25** (2014) 1450063; [arXiv:1402.3026 \[math.QA\]](#).
- [DL96] C. Dong, J. Lepowsky, *The algebraic structure of relative twisted vertex operators*, *J. Pure Appl. Algebra* **110** (1996), 259–295; [arXiv:q-alg/9604022](#).
- [DLiM96] C. Dong, H.-S. Li, G. Mason, *Simple currents and extensions of vertex operator algebras*, *Comm. Math. Phys.* **180** (1996), 671–707; [arXiv:q-alg/9504008](#).
- [FeSt94] A. V. Stoyanovsky, B. L. Feigin, *Functional models of the representations of current algebras and semi-infinite Schubert cells*, *Funktsional Anal. i Prilozhen.* **28** (1) (1994), 68–90, 96 (in Russian); translation in *Funct. Anal. Appl.* **28** (1) (1994), 55–72; preprint B. L. Feigin, A. V. Stoyanovsky, *Quasi-particles models for the representations of Lie algebras and geometry of flag manifold*, [arXiv:hep-th/9308079](#).
- [FHL93] I. Frenkel, Y.-Z. Huang, J. Lepowsky, *On Axiomatic Approaches to Vertex Operator Algebras and Modules*, *Mem. Amer. Math. Soc.* **104**, No. 494 (1993), 64 pages.
- [FLM87] I. B. Frenkel, J. Lepowsky, A. Meurman, *Vertex operator calculus*, *Mathematical aspects of string theory* (San Diego, Calif., 1986), 150–188, *Adv. Ser. Math. Phys.*, 1, *World Sci. Publishing, Singapore*, 1987.
- [FLM88] I. Frenkel, J. Lepowsky, A. Meurman, *Vertex operator algebras and the monster*, *Pure and Appl. Math.*, Academic Press, Boston, 1988.
- [Fu00] J. Fulman, *The Rogers-Ramanujan identities, the finite general linear groups, and the Hall-Littlewood polynomials*, *Proc. Amer. Math. Soc.* **128** (2000), 17–25.
- [G96] G. Georgiev, *Combinatorial constructions of modules for infinite-dimensional Lie algebras, I. Principal subspace*, *J. Pure Appl. Algebra* **112** (1996), 247–286; [arXiv:hep-th/9412054](#).
- [G95] G. Georgiev, *Combinatorial constructions of modules for infinite dimensional Lie algebras, II. Parafermionic space*, preprint [arXiv:q-alg/9504024](#).
- [Gep87] D. Gepner, *New conformal field theories associated with Lie algebras and their partition functions*, *Nuclear Phys. B* **290** (1987), 10–24.
- [Gep22] D. Gepner, *Generalized Rogers-Ramanujan expressions for some non-singlet twisted affine algebras*, *Modern Phys. Lett. A* **37** (2022), 2250105; [arXiv:2204.10601 \[hep-th\]](#).
- [Hu00] J. Hua, *Counting Representations of Quivers over Finite Fields*, *J. Algebra* **226** (2000), 1011–1033.

- [HKOTT02] G. Hatayama, A. Kuniba, M. Okado, T. Takagi and T. Tsuboi, *Path, Crystals and Fermionic Formulae*, MathPhys Odyssey 2001, 205–272, Prog. Math. Phys. **23**, Birkhäuser Boston, MA, 2002.
- [K90] V. G. Kac, *Infinite Dimensional Lie Algebras*, 3rd ed., Cambridge University Press, Cambridge, 1990.
- [KuNS93] A. Kuniba, T. Nakanishi, J. Suzuki, *Characters in conformal field theories from thermodynamic Bethe Ansatz*, Modern Phys. Lett. A **8** (1993), 1649–1659; [arXiv:hep-th/9301018](#).
- [L85] J. Lepowsky, *Calculus of twisted vertex operators*, Proc. Nat. Acad. Sci. U.S.A. **82** (1985), 8295–8299.
- [LP85] J. Lepowsky, M. Primc, *Structure of the standard modules for the affine Lie algebra $A_1^{(1)}$* , Contemp. Math. **46**, Amer. Math. Soc., Providence, 1985.
- [LW81] J. Lepowsky, R. L. Wilson, *A new family of algebras underlying the Rogers-Ramanujan identities and generalizations*, Proc. Nat. Acad. Sci. U.S.A. **78** (1981), 7254–7258.
- [LW84] J. Lepowsky, R. L. Wilson, *The structure of standard modules, I: Universal algebras and the Rogers-Ramanujan identities*, Invent. Math. **77** (1984), 199–290; **II**, *The case A_1 , principal gradation*, Invent. Math. **79** (1985), 417–442.
- [Li96] H. Li, *Local systems of twisted vertex operators, vertex operator superalgebras and twisted modules*, Contemp. Math. **193** (1996), 203–236; [arXiv:q-alg/9504022](#).
- [Li12] H. Li, *Twisted modules and pseudo-endomorphisms*, Algebra Colloq. **19** (2012), 219–336; [arXiv:1004.0864 \[math.QA\]](#).
- [OTa22] M. Okado, R. Takenaka, *Parafermionic bases of standard modules for twisted affine Lie algebras of type $A_{2l-1}^{(2)}$, $D_l^{(2)}$, $E_6^{(2)}$ and $D_4^{(3)}$* , Algebr. Represent. Theory (2022), <https://doi.org/10.1007/s10468-022-10145-2>; [arXiv:2109.08892 \[math.RT\]](#).
- [PeSa18] M. Penn, C. Sadowski, *Vertex-algebraic structure of principal subspaces of the basic modules for twisted Affine Kac-Moody Lie algebras of type $A_{2n-1}^{(2)}$, $D_n^{(2)}$, $E_6^{(2)}$* , J. Algebra **496** (2018), 242–291.
- [P94] M. Primc, *Vertex operator construction of standard modules for $A_n^{(1)}$* , Pacific J. Math. **162** (1994), 143–187.
- [Ta22] R. Takenaka, *Vertex algebraic construction of modules for twisted affine Lie algebras of type $A_{2l}^{(2)}$* , [arXiv:2205.05271 \[math.RT\]](#).