

**LOWER BOUNDS FOR RANKIN-SELBERG L -FUNCTIONS ON
THE EDGE OF THE CRITICAL STRIP**

QIAO ZHANG

ABSTRACT. Let F be a number field, and let π_1 and π_2 be distinct unitary cuspidal automorphic representations of $\mathrm{GL}_{n_1}(\mathbb{A}_F)$ and $\mathrm{GL}_{n_2}(\mathbb{A}_F)$ respectively. In this paper, we derive new lower bounds for the Rankin-Selberg L -function $L(s, \pi_1 \times \tilde{\pi}_2)$ along the edge $\Re s = 1$ of the critical strip in the t -aspect. The corresponding zero-free region for $L(s, \pi_1 \times \tilde{\pi}_2)$ is also determined.

1. INTRODUCTION

The value distributions of L -functions on the edge of the critical strip are important in number theory. In particular, the lower bounds for the L -values along the edge $\Re s = 1$ are closely related to the determination of zero-free regions of the corresponding L -function.

In this paper, we are interested in the lower bounds of Rankin-Selberg L -functions along the edge of the critical strip. More precisely, let F/\mathbb{Q} be a number field of degree n_F , and let π_1 and π_2 be unitary cuspidal automorphic representations of $\mathrm{GL}_{n_1}(\mathbb{A}_F)$ and $\mathrm{GL}_{n_2}(\mathbb{A}_F)$ respectively. Assume that the central characters of π_1 and π_2 are unitary and normalized so that they are trivial on the diagonally embedded copy of the positive reals. Note that, with this normalization, the Rankin-Selberg L -functions $L(s, \pi_1 \times \tilde{\pi}_1)$ and $L(s, \pi_2 \times \tilde{\pi}_2)$ both have simple poles at $s = 1$. The main subject of this paper is the lower bound for the Rankin-Selberg L -function $L(1 + it, \pi_1 \times \tilde{\pi}_2)$, which in turn yields a new zero-free region for $L(s, \pi_1 \times \tilde{\pi}_2)$.

In 1980, Shahidi [10] showed that $L(1 + it, \pi_1 \times \tilde{\pi}_2) \neq 0$ for every $t \in \mathbb{R}$. Since then, there has been substantial progress on the determination of lower bounds and zero-free regions for these Rankin Selberg L -functions, especially in the following two scenarios. On the one hand, if either π_1 or π_2 is self-dual, then it is known, by the work of Moreno [8] in 1985 and Sarnak [9] in 2004, that $L(s, \pi_1 \times \tilde{\pi}_2)$ has the classical zero-free region of the de la Vallée-Poussin type, namely

$$\sigma > 1 - \frac{c}{\log\left(\mathfrak{c}(\pi_1) \mathfrak{c}(\pi_2)(|t| + 2)\right)}$$

where $\mathfrak{c}(\pi_1)$ and $\mathfrak{c}(\pi_2)$ are analytic conductors of π_1 and π_2 respectively. On the other hand, the special case with $\pi_1 = \pi_2$ has also attracted much attention. In 2018, Goldfeld and Li [4] obtained the lower bound

$$L(1 + it, \pi \times \tilde{\pi}) \gg \frac{1}{(\log(|t| + 2))^3}$$

2020 *Mathematics Subject Classification.* Primary 11M26; Secondary 11F66.

Key words and phrases. Rankin-Selberg L -function, Lower bound, Zero-free region.

where π is an irreducible cuspidal unramified representation of $\mathrm{GL}_n(\mathbb{A}_{\mathbb{Q}})$ with $n \geq 2$, tempered almost everywhere. This lower bound immediately leads to the zero-free region for $L(s, \pi \times \tilde{\pi})$

$$\sigma > 1 - \frac{c_{\pi}}{(\log(|t| + 2))^5} \quad |t| \geq 1$$

These results were improved by Humphries and Brumley [5] in 2019 and generalized to an arbitrary number field F , with the lower bound

$$L(1 + it, \pi \times \tilde{\pi}) \gg_{\pi} \frac{1}{\log(|t| + 2)}$$

and the corresponding zero-free region for $L(s, \pi \times \tilde{\pi})$

$$(1) \quad \sigma > 1 - \frac{c_{\pi}}{\log(|t| + 2)} \quad |t| \geq 1$$

Recently in 2022, Humphries and Thorner [6] made explicit the dependence of the constant c_{π} upon π , and gave a more precise form for (1) given by

$$\sigma > 1 - \frac{c}{\log\left(\mathfrak{c}(\pi)^n (|t| + e)^{n^2 n_F}\right)}$$

for some effective absolute constant $c > 0$.

The study of general Rankin-Selberg L -functions without the self-duality assumption is much more difficult, and our knowledge is rather limited. A lower bound

$$L(1 + it, \pi_1 \times \tilde{\pi}_2) \gg \frac{1}{\left(\mathfrak{c}(\pi_1) \mathfrak{c}(\pi_2) (|t| + 2)\right)^{N_{\pi_1, \pi_2}}}$$

and the corresponding zero-free region

$$\sigma > 1 - \frac{c}{\left(\mathfrak{c}(\pi_1) \mathfrak{c}(\pi_2) (|t| + 2)\right)^{N_{\pi_1, \pi_2}}}$$

were obtained in 2006 by Brumley [1] in connection with the strong multiplicity one theorem, and by Gelbert and Lapid [3] for a broad family of L -functions. In 2013, Brumley [2] gave an explicit value for N_{π_1, π_2} . More precisely, if $\pi_1 \neq \pi_2$, then he showed in [2, Theorem A.1] that

$$(2) \quad L(1 + it, \pi_1 \times \tilde{\pi}_2) \gg_{F, n_1, n_2, \varepsilon} \mathfrak{c}(\Pi_t \times \tilde{\Pi}_t)^{-\frac{1}{2}(1 - \frac{1}{n_1 + n_2}) - \varepsilon}$$

where (see (6))

$$\mathfrak{c}(\Pi_t \times \tilde{\Pi}_t) = \mathfrak{c}(\pi_1 \times \tilde{\pi}_1) \mathfrak{c}(\pi_2 \times \tilde{\pi}_2) \mathfrak{c}(\pi_1 \times \tilde{\pi}_2, 1 + it) \mathfrak{c}(\tilde{\pi}_1 \times \pi_2, 1 - it)$$

In particular, by the estimate (7)

$$\mathfrak{c}(\Pi_t \times \tilde{\Pi}_t) \leq \left(\mathfrak{c}(\pi_1) \mathfrak{c}(\pi_2)\right)^{2n_1 + 2n_2} (|t| + 2)^{2n_F n_1 n_2}$$

the lower bound (2) can be re-written as

$$(3) \quad L(1 + it, \pi_1 \times \tilde{\pi}_2) \gg \left(\mathfrak{c}(\pi_1) \mathfrak{c}(\pi_2)\right)^{-(n_1 + n_2 - 1) - \varepsilon} (|t| + 2)^{-n_F n_1 n_2 (1 - \frac{1}{n_1 + n_2}) - \varepsilon}$$

Further, Brumley [2] also derived the corresponding zero-free regions for $L(s, \pi_1 \times \tilde{\pi}_2)$; in fact, it is shown that these lower bounds still hold in these zero-free regions.

In this paper, we seek to improve upon the above results of Brumley in the t -aspect, in particular the lower bound (3), and obtain the following theorem.

Theorem 1.1. *Let F/\mathbb{Q} be a number field of degree n_F , and let π_1 and π_2 be unitary cuspidal automorphic representations of $\mathrm{GL}_{n_1}(\mathbb{A}_F)$ and $\mathrm{GL}_{n_2}(\mathbb{A}_F)$ respectively. Assume that $\pi_1 \neq \pi_2$, and that the central characters of π_1 and π_2 are unitary and normalized so that they are trivial on the diagonally embedded copy of the positive reals. Then we have*

$$L(\sigma + it, \pi_1 \times \tilde{\pi}_2) \gg \left(\mathfrak{c}(\pi_1) \mathfrak{c}(\pi_2) \right)^{-(n_1+n_2-1)-\varepsilon} (|t| + 2)^{-\frac{n_F n_1 n_2}{2} \left(1 - \frac{1}{n_1+n_2}\right) - \varepsilon}$$

where the \gg -constant depends upon F , n_1 , n_2 and ε only.

Remark 1.2. Brumley [2] used the notation $L(s, \pi_1 \times \tilde{\pi}_2)$ to denote the complete Rankin-Selberg L -function. Hence our (finite) L -function $L(s, \pi_1 \times \tilde{\pi}_2)$ is what he denoted by $L^\infty(s, \pi_1 \times \tilde{\pi}_2)$.

Remark 1.3. The work of Brumley [2] also considered the case $\pi_1 = \pi_2$, and similar results were obtained for the lower bounds and zero-free regions. As it is clear from Section 3, our arguments can also be applied to derive the corresponding improvements of these results in the t -aspect. However, these results have now been superseded by the above mentioned works of Goldfeld and Li [4], of Humphries and Brumley [5], and of Humphries and Thorne [6]. Hence we skip the relevant discussions in this paper.

Applying the convexity bound of Li [7], we can also derive from the above lower bound the corresponding zero-free region. Since the arguments are essentially the same as the first part of the proof to [2, Theorem A.1], we omit the details here and only present our result in the following corollary.

Corollary 1.4. *Under the assumptions of Theorem 1, for every $\varepsilon > 0$ there exists a constant $c = c(F, n_1, n_2, \varepsilon) > 0$ such that*

$$L(s, \pi_1 \times \tilde{\pi}_2) \gg_{F, n_1, n_2, \varepsilon} \left(\mathfrak{c}(\pi_1) \mathfrak{c}(\pi_2) \right)^{-(n_1+n_2-1)-\varepsilon} (|t| + 2)^{-\frac{dn_1 n_2}{2} \left(1 - \frac{1}{n_1+n_2}\right) - \varepsilon}$$

in the region

$$\left\{ s = \sigma + it \mid \sigma \geq 1 - \frac{c}{\left(\mathfrak{c}(\pi_1) \mathfrak{c}(\pi_2) \right)^{(n_1+n_2-1)+\varepsilon} (|t| + 2)^{\frac{dn_1 n_2}{2} \left(1 - \frac{1}{n_1+n_2}\right) + \varepsilon}} \right\}$$

Acknowledgements. The author would like to thank Dorian Goldfeld for stimulating this research, and for many insightful discussions.

2. PRELIMINARY

Let F be a number field of degree n_F , and let π_1 and π_2 be unitary cuspidal automorphic representations of $\mathrm{GL}_{n_1}(\mathbb{A}_F)$ and $\mathrm{GL}_{n_2}(\mathbb{A}_F)$ respectively. Assume that the central characters of π_1 and π_2 are unitary and normalized so that they are trivial on the diagonally embedded copy of the positive reals.

Since the Rankin-Selberg L -functions $L(s, \pi_1 \times \tilde{\pi}_1)$ and $L(s, \pi_2 \times \tilde{\pi}_2)$ both have a simple pole at $s = 1$, we may write their Laurent expansions at $s = 1$ as

$$L(s, \pi_1 \times \tilde{\pi}_1) = \sum_{k=-1}^{\infty} A_k (s-1)^k$$

$$L(s, \pi_2 \times \tilde{\pi}_2) = \sum_{k=-1}^{\infty} B_k (s-1)^k$$

Lemma 2.1. *For every $\varepsilon > 0$ there exists a constant $C'_1 = C'_1(\varepsilon) > 0$ such that*

$$\begin{aligned} |A_{-1}| &\leq C'_1 \mathfrak{c}(\pi_1)^\varepsilon & |A_0| &\leq C'_1 \mathfrak{c}(\pi_1)^\varepsilon \\ |B_{-1}| &\leq C'_1 \mathfrak{c}(\pi_2)^\varepsilon & |B_0| &\leq C'_1 \mathfrak{c}(\pi_2)^\varepsilon \end{aligned}$$

For simplicity, write

$$(4) \quad C_1 = C'_1 \mathfrak{c}(\pi_1)^\varepsilon \mathfrak{c}(\pi_2)^\varepsilon$$

then we have

$$|A_{-1}| \leq C_1 \quad |A_0| \leq C_1 \quad |B_{-1}| \leq C_1 \quad |B_0| \leq C_1$$

Proof. The lemma follows from Li's convexity bounds [7]; see also [2]. \square

Further, for every $t \in \mathbb{R}$ consider the isobaric representation

$$\Pi_t = (\pi_1 \otimes |\det|^{it/2}) \boxplus (\pi_2 \otimes |\det|^{-it/2})$$

then we have the factorization of L -functions

$$(5) \quad L(s, \Pi_t \times \tilde{\Pi}_t) = L(s, \pi_1 \times \tilde{\pi}_1) L(s, \pi_2 \times \tilde{\pi}_2) L(s+it, \pi_1 \times \tilde{\pi}_2) L(s-it, \tilde{\pi}_1 \times \pi_2)$$

and so the corresponding factorization of analytic conductors

$$(6) \quad \mathfrak{c}(\Pi_t \times \tilde{\Pi}_t) = \mathfrak{c}(\pi_1 \times \tilde{\pi}_1) \mathfrak{c}(\pi_2 \times \tilde{\pi}_2) \mathfrak{c}(\pi_1 \times \tilde{\pi}_2, 1+it) \mathfrak{c}(\tilde{\pi}_1 \times \pi_2, 1-it)$$

In particular, by the bound [1, (8)] we have

$$(7) \quad \mathfrak{c}(\Pi_t \times \tilde{\Pi}_t) \leq \left(\mathfrak{c}(\pi_1) \mathfrak{c}(\pi_2) \right)^{2n_1+2n_2} (|t|+2)^{2n_F n_1 n_2}$$

By the factorization (5), since π_1 and π_2 are assumed distinct, the Rankin-Selberg L -function $L(s, \Pi_t \times \tilde{\Pi}_t)$ has a double pole at $s=1$ and is holomorphic everywhere else. Accordingly, we write the Laurent series representation of $L(s, \Pi_t \times \tilde{\Pi}_t)$ at $s=1$ as

$$L(s, \Pi_t \times \tilde{\Pi}_t) = \sum_{k=-2}^{\infty} r_k (s-1)^k$$

Lemma 2.2. *For every $\varepsilon > 0$, there exists a constant $C'_2 = C'_2(\varepsilon) > 0$ such that*

$$\begin{aligned} |r_{-1}| + |r_{-2}| &\geq C'_2 \mathfrak{c}(\Pi_t \times \tilde{\Pi}_t)^{-\frac{1}{2}(1-\frac{1}{n_1+n_2})-\varepsilon} \\ &\geq C'_2 \left(\mathfrak{c}(\pi_1) \mathfrak{c}(\pi_2) \right)^{-(n_1+n_2-1)-\varepsilon} (|t|+2)^{-n_F n_1 n_2 (1-\frac{1}{n_1+n_2})-\varepsilon} \end{aligned}$$

For simplicity, write

$$(8) \quad C_2 = C'_2 \left(\mathfrak{c}(\pi_1) \mathfrak{c}(\pi_2) \right)^{-(n_1+n_2-1)-\varepsilon}$$

then we have

$$|r_{-1}| + |r_{-2}| \geq C_2 (|t|+2)^{-n_F n_1 n_2 (1-\frac{1}{n_1+n_2})-\varepsilon}$$

Proof. This follows immediately from [1, Theorem 3]. \square

Lemma 2.3. *Let $|t| \geq 1$. Then we have*

$$|r_{-1}| + |r_{-2}| \leq 3C_1^2 \left| L(1 + it, \pi_1 \times \tilde{\pi}_2) \right|^2 + C_1^2 \left| \frac{d}{dt} \left(\left| L(1 + it, \pi_1 \times \tilde{\pi}_2) \right|^2 \right) \right|$$

where C_1 is as given in (4).

Proof. By the factorization (5), we have

$$r_{-2} = A_{-1} B_{-1} \left| L(1 + it, \pi_1 \times \tilde{\pi}_2) \right|^2$$

so by Lemma 2.1 we have

$$\begin{aligned} |r_{-2}| &= |A_{-1}| |B_{-1}| \left| L(1 + it, \pi_1 \times \tilde{\pi}_2) \right|^2 \\ &\leq C_1^2 \left| L(1 + it, \pi_1 \times \tilde{\pi}_2) \right|^2 \end{aligned}$$

Also, by the factorization (5) we have

$$\begin{aligned} r_{-1} &= (A_{-1}B_0 + A_0B_{-1}) \left| L(1 + it, \pi_1 \times \tilde{\pi}_2) \right|^2 \\ &\quad + 2A_{-1} B_{-1} \Re \left(L'(1 + it, \pi_1 \times \tilde{\pi}_2) \cdot \overline{L(1 + it, \pi_1 \times \tilde{\pi}_2)} \right) \\ &= (A_{-1}B_0 + A_0B_{-1}) \left| L(1 + it, \pi_1 \times \tilde{\pi}_2) \right|^2 \\ &\quad + A_{-1} B_{-1} \frac{d}{dt} \left(\left| L(1 + it, \pi_1 \times \tilde{\pi}_2) \right|^2 \right) \end{aligned}$$

so again by Lemma 2.1 we have

$$\begin{aligned} |r_{-1}| &\leq (|A_{-1}| |B_0| + |A_0| |B_{-1}|) \left| L(1 + it, \pi_1 \times \tilde{\pi}_2) \right|^2 \\ &\quad + |A_{-1}| |B_{-1}| \left| \frac{d}{dt} \left(\left| L(1 + it, \pi_1 \times \tilde{\pi}_2) \right|^2 \right) \right| \\ &\leq 2C_1^2 \left| L(1 + it, \pi_1 \times \tilde{\pi}_2) \right|^2 + C_1^2 \left| \frac{d}{dt} \left(\left| L(1 + it, \pi_1 \times \tilde{\pi}_2) \right|^2 \right) \right| \end{aligned}$$

Hence our lemma follows as we combine all the above estimates. \square

3. PROOF OF THEOREM 1

In this section, we present the proof of Theorem 1, and will freely use the notations introduced in Section 2.

The theorem is well known if $n_1 = n_2 = 1$, so henceforth we assume that either $n_1 \geq 2$ or $n_2 \geq 2$. For simplicity, henceforth we write

$$\theta = \frac{n_F n_1 n_2}{2} \left(1 - \frac{1}{n_1 + n_2} \right) \geq \frac{1}{2}$$

Let $\varepsilon > 0$. By [2, Theorem A.1], there exists a constant $C'_\varepsilon > 0$ such that

$$\left| L(1 + it, \pi_1 \times \tilde{\pi}_2) \right| \geq C'_\varepsilon \mathfrak{c}(\Pi_t \times \tilde{\Pi}_t)^{-\frac{1}{2} \left(1 - \frac{1}{n_1 + n_2} \right) - \frac{\varepsilon}{2(n_1 + n_2 + n_F n_1 n_2)}}$$

Recall the bound (7)

$$\mathbf{c}(\Pi_t \times \tilde{\Pi}_t) \leq \left(\mathbf{c}(\pi_1) \mathbf{c}(\pi_2) \right)^{2n_1+2n_2} (|t|+2)^{2n_F n_1 n_2}$$

then over the interval $[-1, 1]$ the above lower bound gives

$$(9) \quad \begin{aligned} \left| L(1+it, \pi_1 \times \tilde{\pi}_2) \right| &\geq C'_\varepsilon \left(\mathbf{c}(\pi_1) \mathbf{c}(\pi_2) \right)^{-(n_1+n_2-1)-\varepsilon} (|t|+2)^{-2\theta-\varepsilon} \\ &\geq \frac{C'_\varepsilon}{3^\theta} \left(\mathbf{c}(\pi_1) \mathbf{c}(\pi_2) \right)^{-(n_1+n_2-1)-\varepsilon} (|t|+2)^{-\theta-\varepsilon} \end{aligned}$$

Now we claim that

$$(10) \quad \left| L(1+it, \pi_1 \times \tilde{\pi}_2) \right| \geq C_\varepsilon (|t|+2)^{-\theta-\varepsilon} \quad (t \in \mathbb{R})$$

where

$$C_\varepsilon = \min \left\{ \frac{C'_\varepsilon}{3^\theta} \left(\mathbf{c}(\pi_1) \mathbf{c}(\pi_2) \right)^{-(n_1+n_2-1)-\varepsilon}, \sqrt{\frac{C_2}{6C_1^2 (2\theta+2\varepsilon)}} \right\}$$

and C_1 and C_2 are the constants defined in (4) and (8) respectively. Note in particular that we have

$$C_\varepsilon \gg \left(\mathbf{c}(\pi_1) \mathbf{c}(\pi_2) \right)^{-(n_1+n_2-1)-\varepsilon}$$

where the \gg -constant depends upon F , n_1 , n_2 and ε only.

By the construction of C_ε and the estimate (9), the claim (10) is obviously valid over the interval $[-1, 1]$. Now assume that there indeed exists some point $|t_0| > 1$ such that

$$(11) \quad \left| L(1+it_0, \pi_1 \times \tilde{\pi}_2) \right| < C_\varepsilon (|t_0|+2)^{-\theta-\varepsilon}$$

Without loss of generality, we may assume that $t_0 > 1$. For simplicity, henceforth we write

$$g(t) = \left| L(1+it, \pi_1 \times \tilde{\pi}_2) \right|^2$$

then our claim (10) becomes

$$g(t) \geq C_\varepsilon^2 (|t|+2)^{-2\theta-2\varepsilon} \quad (t \in \mathbb{R})$$

and the assumption (11) becomes

$$(12) \quad g(t_0) < C_\varepsilon^2 (t_0+2)^{-2\theta-2\varepsilon}$$

Define

$$\mathcal{S}_1 = \left\{ 0 \leq t < t_0 \mid g(t) \geq C_\varepsilon^2 (t+2)^{-2\theta-2\varepsilon} \right\}$$

Since $[0, 1] \subseteq \mathcal{S}_1$, we have $\mathcal{S}_1 \neq \emptyset$, so the supremum $t_1 = \sup(\mathcal{S}_1)$ exists and is finite, and we have

$$t_1 \geq 1 \quad g(t_1) = C_\varepsilon^2 (t_1+2)^{-2\theta-2\varepsilon}$$

Also, we define

$$\mathcal{S}_2 = \left\{ t > t_0 \mid g(t) \geq C_\varepsilon^2 (t+2)^{-2\theta-2\varepsilon} \right\} \cup \{\infty\} \quad t_2 = \inf(\mathcal{S}_2)$$

so $t_0 \in (t_1, t_2)$ and we have

$$(13) \quad g(t) < C_\varepsilon^2 (t+2)^{-2\theta-2\varepsilon} \quad (t_1 < t < t_2)$$

Further, if $t_2 < \infty$, then we also have

$$g(t_2) = C_\varepsilon^2 (t_2 + 2)^{-2\theta-2\varepsilon}$$

Combining Lemmas 2.2 and 2.3, for $t \geq 1$ we have

$$C_2(t+2)^{-2\theta-\varepsilon} \leq |r_{-1}| + |r_{-2}| \leq 3C_1^2 g(t) + C_1^2 |g'(t)|$$

Hence we have

$$(14) \quad |g'(t)| \geq \frac{C_2}{C_1^2} (t+2)^{-2\theta-\varepsilon} - 3g(t)$$

In particular, over the interval (t_1, t_2) , by (13) we have

$$g(t) < C_\varepsilon^2 (|t|+2)^{-2\theta-2\varepsilon} \leq \frac{C_2}{6C_1^2} (|t|+2)^{-2\theta-2\varepsilon}$$

so the estimate (14) gives

$$\begin{aligned} |g'(t)| &\geq \frac{C_2}{C_1^2} (t+2)^{-2\theta-\varepsilon} - 3g(t) \\ &\geq \frac{C_2}{C_1^2} (t+2)^{-2\theta-\varepsilon} - 3 \cdot \frac{C_2}{6C_1^2} (t+2)^{-2\theta-2\varepsilon} \\ &\geq \frac{C_2}{2C_1^2} (t+2)^{-2\theta-\varepsilon} \quad (t_1 < t < t_2) \end{aligned}$$

In particular, we have $g'(t) \neq 0$ over (t_1, t_2) , so the derivative $g'(t)$ must keep the same sign in this interval. If $g'(t) > 0$ over (t_1, t_2) , then $g(t)$ is a strictly increasing function, and in particular

$$g(t_0) > g(t_1) = C_\varepsilon^2 (t_1 + 2)^{-2\theta-2\varepsilon} > C_\varepsilon^2 (t_0 + 2)^{-2\theta-2\varepsilon}$$

in contradiction to the assumption (12). Hence we have $g'(t) < 0$ over (t_1, t_2) , and so

$$(15) \quad -g'(t) = |g'(t)| \geq \frac{C_2}{2C_1^2} (t+2)^{-2\theta-\varepsilon} \quad (t_1 < t < t_2)$$

If $t_2 < \infty$, then by the choices of t_1 and t_2 we have

$$\begin{aligned} \int_{t_1}^{t_2} \frac{C_2}{2C_1^2} (t+2)^{-2\theta-\varepsilon} dt &\leq \int_{t_1}^{t_2} (-g'(t)) dt = g(t_1) - g(t_2) \\ &= C_\varepsilon^2 (t_1 + 2)^{-2\theta-2\varepsilon} - C_\varepsilon^2 (t_2 + 2)^{-2\theta-2\varepsilon} \\ &= \int_{t_1}^{t_2} C_\varepsilon^2 (2\theta + 2\varepsilon)(t+2)^{-2\theta-2\varepsilon-1} dt \end{aligned}$$

which is simply

$$(16) \quad \int_{t_1}^{t_2} C_\varepsilon^2 \frac{2\theta + 2\varepsilon}{(t+2)^{1+\varepsilon}} (t+2)^{-2\theta-\varepsilon} dt \geq \int_{t_1}^{t_2} \frac{C_2}{2C_1^2} (t+2)^{-2\theta-\varepsilon} dt$$

On the other hand, over the interval (t_1, t_2) , by the construction of the constant C_ε we have

$$C_\varepsilon^2 \frac{2\theta + 2\varepsilon}{(t+2)^{1+\varepsilon}} \leq \frac{C_2}{6C_1^2(2\theta + 2\varepsilon)} \cdot \frac{2\theta + 2\varepsilon}{1+2} < \frac{C_2}{2C_1^2}$$

so we should have

$$\int_{t_1}^{t_2} C_\varepsilon^2 \frac{2\theta + 2\varepsilon}{(t+2)^{1+\varepsilon}} (t+2)^{-2\theta-\varepsilon} dt < \int_{t_1}^{t_2} \frac{C_2}{2C_1^2} (t+2)^{-2\theta-\varepsilon} dt$$

a contradiction to (16). Hence we must have $t_2 = \infty$. In particular, this implies that we can take t_0 to be any value in the infinite interval (t_1, ∞) .

In summary, combining (13) and (15) gives

$$g(t) < C_\varepsilon^2 (t+2)^{-2\theta-2\varepsilon} \quad -g'(t) \geq \frac{C_2}{2C_1^2} (t+2)^{-2\theta-\varepsilon} \quad (t > t_1)$$

so for every $t_0 > t_1$ we have

$$\begin{aligned} C_\varepsilon^2 (t_1+2)^{-2\theta-2\varepsilon} = g(t_1) &\geq g(t_1) - g(t_0) = \int_{t_1}^{t_0} (-g'(t)) dt \\ &\geq \int_{t_1}^{t_0} \frac{C_2}{2C_1^2} (t+2)^{-2\theta-\varepsilon} dt = \frac{C_2}{2C_1^2} \frac{(t+2)^{1-2\theta-\varepsilon}}{1-2\theta-\varepsilon} \Big|_{t_1}^{t_0} \\ &= \frac{C_2}{2C_1^2} \frac{(t_1+2)^{1-2\theta-\varepsilon} - (t_0+2)^{1-2\theta-\varepsilon}}{2\theta+\varepsilon-1} \end{aligned}$$

Since $\theta \geq 1/2$, we have $1-2\theta-\varepsilon > 0$, so letting $t_0 \rightarrow \infty$ gives

$$C_\varepsilon^2 (t_1+2)^{-2\theta-2\varepsilon} \geq \frac{C_2}{2C_1^2} \frac{(t_1+2)^{1-2\theta-\varepsilon}}{2\theta+\varepsilon-1}$$

This implies that

$$C_\varepsilon^2 \geq \frac{C_2}{2C_1^2} \frac{(t_1+2)^{1+\varepsilon}}{2\theta+\varepsilon-1}$$

and so

$$\begin{aligned} (t_1+2)^{1+\varepsilon} &\leq C_\varepsilon^2 \cdot \frac{2C_1^2}{C_2} \cdot (2\theta+\varepsilon-1) \\ &\leq \frac{C_2}{6C_1^2(2\theta+2\varepsilon)} \cdot \frac{2C_1^2}{C_2} \cdot (2\theta+\varepsilon-1) \\ &= \frac{2\theta+\varepsilon-1}{3(\theta+\varepsilon)} < 1 \end{aligned}$$

which is impossible. Hence there cannot exist a point $|t_0| > 1$ satisfying (12), and so the claim (10) is valid for every $t \in \mathbb{R}$, i.e.,

$$\left| L(1+it, \pi_1 \times \tilde{\pi}_2) \right| \geq C_\varepsilon (|t|+2)^{-\theta-\varepsilon} \quad (t \in \mathbb{R})$$

This completes our proof of Theorem 1.

REFERENCES

- [1] F. Brumley, *Effective Multiplicity One on GL_n and Narrow Zero-Free Regions for Rankin-Selberg L -Functions*, Amer. J. Math. **128** (2006), no. 6, 1455–1474. MR2275908.
- [2] E. Lapid, *On the Harish-Chandra Schwartz space of $G(F)\backslash G(A)$* , With an appendix by F. Brumley, in “Automorphic representations and L -functions”, Tata Inst. Fundam. Res. Stud. Math. **22**, 335–377, Tata Inst. Fund. Res., Mumbai, 2013. MR3156857.
- [3] S. Gelbart and E. Lapid, *Lower bounds for L -functions at the edge of the critical strip*, Amer. J. Math. **128** (2006), no. 3, 619–638. MR2230919.
- [4] D. Goldfeld and X. Li, *A Standard Zero Free Region for Rankin-Selberg L -Functions*, Int. Math. Res. Not. IMRN **2018** (2018) (22), 7067–7136. MR3878594.
- [5] P. Humphries and F. Brumley, *Standard zero-free regions for Rankin-Selberg L -functions via sieve theory*, Math. Z. **292** (2019), no. 3–4, 1105–1122. MR3980284.
- [6] P. Humphries and J. Thorner, *Towards a GL_n variant of the Hoheisel phenomenon*, Trans. Amer. Math. Soc. **375** (2022), no. 3, 1801–1824. MR4378080.

- [7] X. Li, *Upper bounds on L -functions at the edge of the critical strip*, Int. Math. Res. Not. IMRN **2010** (2010), no. 4, 727–755. MR 2595006.
- [8] C. J. Moreno, *Analytic proof of the strong multiplicity one theorem*, Amer. J. Math. **107** (1985), no. 1, 163–206. MR0778093.
- [9] P. C. Sarnak, *Nonvanishing of L -functions on $\Re(s) = 1$* , in “Contributions to automorphic forms, geometry, and number theory”, 719–732, Johns Hopkins Univ. Press, Baltimore, MD, 2004. MR2058625.
- [10] F. Shahidi, *On nonvanishing of L -functions*, Bull. Amer. Math. Soc. (N.S.) **2** (1980), no. 3, 462–464. MR0561534.

DEPARTMENT OF MATHEMATICS, TEXAS CHRISTIAN UNIVERSITY, FORT WORTH, TX 76129

Email address: q.zhang@tcu.edu