

Decoherence of a 2-Path System by Infrared Photons

Colby DeLisle^{1,2,*} and P.C.E. Stamp^{1,2,3,†}

¹*Department of Physics and Astronomy, University of British Columbia,
6224 Agricultural Rd., Vancouver, B.C., Canada V6T 1Z1*

²*Pacific Institute of Theoretical Physics, University of British Columbia,
6224 Agricultural Rd., Vancouver, B.C., Canada V6T 1Z1*

³*Theoretical Astrophysics, Cahill, California Institute of Technology,
1200 E. California Boulevard, MC 350-17, Pasadena CA 91125, USA*

(Dated: November 14, 2022)

We calculate the decoherence caused by photon emission for a charged particle travelling through an interferometer; the decoherence rate gives a quantitative measure of how much “which-path” quantum information is gained by the electromagnetic field. We isolate the quantum information content of both leading and sub-leading soft photons, and show that it can be extracted entirely from information about the endpoints of the particle’s paths. When infrared dressing is properly incorporated, the leading order soft photons give no contribution to decoherence, and carry no quantum information. The sub-leading soft photons carry finite which-path information, and the sub-leading contribution to decoherence takes an extremely simple, time-independent form depending only on the size of the interferometer.

I. INTRODUCTION

Our subject in this paper is the well-known “2-path” problem in quantum mechanics [1], in the present case for a charged particle coupled to the electromagnetic (EM) field. We thus assume a pointlike particle with electric charge e , which is travelling through an interferometer, and is put into a superposition of trajectories falling into two distinct classes [2], which we label L and R . We will sometimes refer to this particle as an electron, although it could of course be some other charged particle of larger mass.

An external observer then has the choice of either (i) measuring which path the particle has followed, or (ii) letting it simply pass through the interferometer unmolested by any measuring system. As is well known, in the former case no interference is seen between the 2 paths, whereas in the latter case, a standard interference pattern is seen on any screen which intercepts the 2 paths.

However, it is also well known that on its way through the interferometer, the particle couples to the electromagnetic (EM) field, which may be either at temperature $T = 0$ (in its ground state) or at finite T in the rest frame of the interferometer. In either case, the EM field is able to, in effect, passively “measure” which path the particle has followed; in quantum-information-theoretic language, it can act as a “witness” as to which path is followed. This mechanism is of course quite general, and is usually called “environmental decoherence” [3–6]; because of it one expects partial or total destruction of the interference between the 2 paths.

The particular case of decoherence for 2-path systems in which electrons coupled to photons has been repeatedly discussed in the literature [7–9]. In this paper we

wish to revisit this topic, and clarify the role of low frequency infrared (IR) photons in bringing about the decoherence. Our main result will be explained in more detail below – we show that decoherence actually comes not from the leading IR divergent terms in the effective electron-photon coupling (which actually give zero contribution), but rather the sub-leading terms.

We stress that in any real experiment there will be other processes also causing decoherence – these include coupling of the electron to any gas particles in the experiment, as well as the long-range interaction of the electron with “charge fluctuators” in the apparatus. The former can be understood using scattering theory [10], and the latter using a “spin bath” theory [11]. We will ignore these contributions in what follows, since we are primarily interested in the EM decoherence mechanism.

The topic of IR divergences, and their influence on the quantum electronic dynamics, of course has a long history, dating all the way back to the work of Bloch and Nordsieck [12]. Early work clarified the ways in which the leading divergences coming from radiative corrections to scattering processes [12–14] were cancelled by divergences from virtual soft particles, leading to the derivation of multiplicative “soft factors” (see, eg., refs. [15–17], and also refs. [18–21]) in scattering processes. These factors, when correctly handled, give finite multiplicative corrections to scattering cross-sections.

One can write soft factors for any electron-photon scattering process. Here we are interested in the 3-point electron-photon scattering process (see Fig. 1), in which the incoming state is an electron with 4-momentum p^a , and the outgoing state is an electron and a photon, with momenta $p^a - q^a$ and q^a respectively. The “soft photon theorem” then says that one can write the fully renormalized vertex for this process in the form

$$\Lambda(p, q) = K_2(p) [\mathcal{S}_{(0)}(q) + \mathcal{S}_{(1)}(q)] + \mathcal{O}(|\mathbf{q}|) \quad (1)$$

where $K_2(p)$ is the fully renormalized 2-point propagator

* cdelisle@phas.ubc.ca

† stamp@phas.ubc.ca, stamp@tapir.caltech.edu

for the electron alone, with no photon emission. Here the divergent leading soft factor $\mathcal{S}_{(0)}(q) \sim \mathcal{O}(1/|\mathbf{q}|)$, where \mathbf{q} is the 3-momentum, and the sub-leading factor $\mathcal{S}_{(1)}(q) \sim \mathcal{O}(1)$.




FIG. 1. The 3-point electron-photon vertex $\Lambda(p, q)$ written, for small $|\mathbf{q}|$, as a product of the single electron propagator $K_2(p)$ and the “soft factor” $\mathcal{S}(q)$; compare eq. (1) in the text.

Later work has shown that this result can also be understood by using either an eikonal analysis [22–24], or a coherent state representation for the coupled electron-photon system [25–30]. The latter representation, in which the system is represented by “dressed” electron states, is quite illuminating, for two reasons.

First, it highlights that the naïve photon vacuum is non-unique; there are infinitely many field configurations with zero energy. Thus, one needs to be careful in trying to calculate decoherence: the decoherence rate one finds will diverge if one evaluates scattering processes between the usual photon Fock states, due to transitions between these superselected vacua, but will be convergent if one instead uses dressed states [31, 32]. This is physically rather obvious – the electrons are at no point ever undressed, so one should always work with dressed states.

Second, the coherent state representation affords a nice way of separating the leading contributions to the scattering processes from all sub-leading contributions, first discussed in perturbative calculations [33–36] (this separation is also done in a very clear way in the eikonal expansion [22, 23]). The separation of terms will be crucial in what follows, since what we wish to establish here is that the leading IR contributions to the soft factor in QED make no contribution at all to decoherence. It is instead the sub-leading contributions that are responsible for the lowest-energy decoherence in QED.

To show this we will use the recent discovery that the leading and sub-leading “soft photon factors” are encoded at the endpoints of charged particle worldlines [37]. We shall see that we can use this result to isolate the quantum information content of both leading and sub-leading soft photons in our model; and we shall see that the former is zero, and the latter is finite. What is more, it is not necessary to go to any asymptotic limit to show this – we will see that even in some experiment in which all processes are confined to a finite spacetime region, these results still hold – the leading term still

gives identically zero, and the sub-leading contribution is time-independent, depending only on the size of the interferometer.

The results we obtain for the total amount of decoherence is in fact consistent with a result derived two decades ago by Breuer and Petruccione [9]. It is satisfying that their result – which was found without worrying about the subtleties of the different contributions to soft factors – be confirmed, since its derivation was clearly based on physical arguments. However it is also satisfying to be able to unravel the different soft contributions – from leading and sub-leading terms – to the decoherence functional, entirely in terms of the boundary (endpoint) currents. This method makes for a simple derivation, since we don’t have to worry too much about the details of the paths between the endpoints. It also allows for a rather novel and transparent interpretation of decoherence in the infrared.

A. Summary & Structure of the Paper

The rest of the paper proceeds as follows. First, in Section II we introduce what is by now a standard semiclassical interferometry model. We describe the assumed geometry of the interferometer, and discuss the way in which we evaluate decoherence in the model using the decoherence functional Γ .

In Section III we discuss relevant features of the momentum space electromagnetic current $j^a(q)$ which appears in the decoherence functional. This will allow us to review results from our recent work [37], which showed that boundary terms in the electromagnetic current encode the “soft factors” appearing in both the leading and sub-leading soft photon theorems.

We then show in Section IV that superpositions of the leading soft factor generically cause the decoherence functional Γ to diverge in the infrared – even for finite-time experiments. We must deal with these infrared divergences properly in order to obtain sensible results for the quantum information content of the photon field, so Section IV also discusses how we ought to “dress” the time evolution operator in our semiclassical model in order to render the decoherence functional infrared-finite. After adding the appropriate dressing, the leading soft photons cease to obtain any quantum information about the matter particle.

Section V contains the evaluation of the corrected expression for Γ in our model. We show that sub-leading soft photons still contribute to Γ , and isolate the contribution coming from the difference in the sub-leading soft current on each branch of the superposition. The sub-leading contribution to decoherence turns out to be time-independent, and to depend only upon the spatial extent of the model interferometer.

The paper concludes in Section VII with a discussion of our results, and of some open questions.

Here we use the mostly-negative metric signature,

$\eta_{ab} \equiv \text{diag}(1, -1, -1, -1)$, units in which $\hbar = c = \epsilon_0 = 1$, and a bar over a quantity will indicate its complex conjugate.

II. INTERFEROMETER MODEL

Let us begin by introducing the simple model we will use to describe a charged particle traversing an interferometer - this model is fairly standard in quantum optics [38]. The spatial trajectories taken by the superposed particle are illustrated in Figure 2. The particle enters the apparatus along the \hat{x} direction with four-velocity \dot{X}_2^a ; upon reaching the point X_1^a at time $t = 0$, the particle is put into a superposition, in which it is either kicked into the \hat{y} direction to follow the trajectory $X_L^a(s)$, or allowed to proceed in the \hat{x} direction and follow the trajectory $X_R^a(s)$. Here s is the proper time experienced by the particle as it evolves along the trajectories.




FIG. 2. Top-down view of the two-path geometry traversed by the charged particle.

As the system then evolves, that branch of the superposition following the trajectory $X_L^a(s)$ (the solid line in Figure 2) proceeds with the four-velocity \dot{X}_1^a to the point X_L^a , and then with four-velocity \dot{X}_2^a until it reaches the detector \mathcal{D} . The branch following the trajectory $X_R^a(s)$ (dashed line) proceeds with the four-velocity \dot{X}_2^a to the point X_R^a , and then with four-velocity \dot{X}_1^a to the detector. On both paths, the speed of the particle is taken to be constant at $v \equiv l/\tau$, where l is the length of one side of the interferometer, so that the particle reaches either X_L^a or X_R^a at time $t = \tau$ and reaches the detector at $t = 2\tau$.

As noted in the introduction, Fig. 2 is schematic in the sense that in reality, a *single* path shown going along the left or right trajectory actually represents the set of *all* paths passing through the left arm of an interferometer (this shorthand way of representing entire classes of path appears to have originated in Feynman's celebrated description of two-path experiments [1]).

In the computations that follow however, we will use a semiclassical model of interferometry, consisting of a simple discrete superposition of two classical electromagnetic

currents, j_L^a and j_R^a in the interferometer. The electromagnetic current of a pointlike particle with charge e , which follows either of the trajectories $X_{L/R}^a(s)$, is

$$j_{L/R}^a(x) = e \int_{s_i}^{s_f} ds \dot{X}_{L/R}^a(s) \delta^{(4)}(x - X_{L/R}(s)), \quad (2)$$

where again s is the particle's proper time, and $\dot{X}_{L/R}^a(s) \equiv \frac{d}{ds} X_{L/R}(s)$ its four-velocity. The bounds on this integral are chosen such that the initial and final proper times coincide with the beginning and end of the experiment, $X_{L/R}^0(s_i) = 0$ and $X_{L/R}^0(s_f) = 2\tau$.

Assuming the photon field is in vacuum at the time the experiment starts, under time evolution the electromagnetic environment will be placed into a superposition of states $|L\rangle$ and $|R\rangle$ which will be coherent photon states, sourced by either the L or R current:

$$|L/R\rangle \equiv \mathcal{T} e^{-i \int d^4x j_{L/R}^a(x) \hat{A}_a(x)} |0\rangle \quad (3)$$

These coherent states are obtained by the action of the interaction-picture time evolution operator acting on the photon vacuum $|0\rangle$, and here \mathcal{T} denotes time-ordering. In writing this expression we have assumed as in (2) that the currents $j_{L/R}^a(x)$ have support only for the duration of the experiment (between s_i and s_f), so we can let the integral $\int d^4x$ run over all values of $t \in (-\infty, +\infty)$.

The state of the matter+radiation system as the particle enters the detector \mathcal{D} is taken to be

$$\frac{1}{\sqrt{2}} [|\psi_L\rangle |L\rangle + |\psi_R\rangle |R\rangle], \quad (4)$$

with $|\psi_{L/R}\rangle$ the states of the matter particle after traversing the L/R arm of the interferometer. As the particle enters the detector, its position space probability distribution is then

$$P(x) = \frac{1}{2} |\psi_L(x)|^2 + \frac{1}{2} |\psi_R(x)|^2 + \frac{1}{2} \langle R|L \rangle \psi_L(x) \bar{\psi}_R(x) + \frac{1}{2} \langle L|R \rangle \bar{\psi}_L(x) \psi_R(x), \quad (5)$$

where we have defined the position space wavefunctions $\psi_{L/R}(x) \equiv \langle x | \psi_{L/R} \rangle$.

The quantity governing the quantum coherence of the particle in the two-path experiment is then the inner product between the photon coherent states,

$$\langle R|L \rangle \equiv e^{i\Theta - \Gamma}; \quad \Theta, \Gamma \in \mathbb{R}, \quad \Gamma \geq 0. \quad (6)$$

Here, $\Gamma \equiv -\text{Re}(\ln \langle R|L \rangle)$, and $\Theta \equiv \text{Im}(\ln \langle R|L \rangle)$. Because the coherent states $|L/R\rangle$ are fully determined by the classical currents $j_{L/R}^a$, we can think of the inner product (6) as a functional of these currents,

$$\langle R|L \rangle \equiv \langle R|L \rangle [j_L, j_R]. \quad (7)$$

Consequently $\Gamma \equiv \Gamma[j_L, j_R]$ and $\Theta \equiv \Theta[j_L, j_R]$ are functionals of the classical currents as well.

The quantity Γ will be of particular importance to us, and is called the *decoherence functional* [5, 7, 39, 40]. It will be a functional of the entire currents along the L and R paths, and is entirely responsible for the suppression of interference terms when measuring the matter state, as it alone governs the modulus of the electromagnetic inner product

$$|\langle R|L\rangle| = e^{-\Gamma}. \quad (8)$$

The electromagnetic decoherence functional was first derived long ago [40, 41], and has been discussed more recently e.g. by Ford [8], and by Breuer and Petruccione [9]. At zero temperature it takes the form

$$\Gamma[j_L, j_R] = \frac{1}{2} \int \frac{d^3q}{(2\pi)^3 2\omega} P_{ab} \delta j^a(q) \delta j^b(-q), \quad (9)$$

where $\delta j^a \equiv j_L^a - j_R^a$ is the difference in electromagnetic currents along the two superposed paths, $q^a \equiv \omega(1, \hat{n})$, with \hat{n} an angular unit vector, is a null four-momentum, and $\int \frac{d^3q}{(2\pi)^3} \equiv \oint \frac{dS^2(\hat{n})}{(2\pi)^2} \int_0^\infty \frac{d\omega}{2\pi} \omega^2$. Finally, the transverse projector P_{ab} can be written in terms of the momentum q^a as $P_{jk} \equiv \delta_{jk} - \frac{q_j q_k}{\omega^2}$, with $P_{0a} = 0$.

Note that Γ as written is a functional of currents which may have support for times $t \in (-\infty, +\infty)$. However we have restricted to finite-time dynamics here by restricting the range of the proper time integrals in (2).

III. FORM OF ELECTROMAGNETIC CURRENTS & DECOHERENCE FUNCTIONAL

Let us now further discuss the matter currents we will insert into the decoherence functional (9); we will wish to understand in particular the various contributions to the decoherence functional – which is central to any analysis of decoherence, for any system – coming from the endpoints of the particle’s paths through the interferometer.

What appears in the decoherence functional is actually the Fourier transform of the current, $j^a(q) \equiv \int d^4x e^{iq \cdot x} j^a(x)$, which using (2) is

$$j^a(q) = e \int ds \dot{X}^a(s) e^{iq \cdot X(s)}. \quad (10)$$

This current can be rewritten as

$$j^a(q) = e \int ds \dot{X}^a \left(\frac{1}{iq \cdot \dot{X}} \right) \frac{d}{ds} e^{iq \cdot X}, \quad (11)$$

so that integrating by parts in s gives

$$j^a(q) = -ie \int ds \frac{d}{ds} \left(e^{iq \cdot X} \frac{\dot{X}^a}{q \cdot \dot{X}} \right) + ie \int ds e^{iq \cdot X} \frac{d}{ds} \left(\frac{\dot{X}^a}{q \cdot \dot{X}} \right). \quad (12)$$

Discarding the boundary term appearing here [37, 42, 43] ensures that the current is conserved, $q_a j^a(q) = 0$, leaving us with the following corrected expression for the particle current in momentum space:

$$j^a(q) = ie \int ds e^{iq \cdot X} \frac{d}{ds} \left(\frac{\dot{X}^a}{q \cdot \dot{X}} \right) \quad (13)$$

Notice that the form (13) of the matter current has the property that the current vanishes when the particle is not accelerating ($\dot{X}^a = 0$). Plugging the current into the decoherence functional (9), we note immediately the important fact that *there is no decoherence without acceleration*.

A. Soft Electromagnetic Currents

Recently, we showed [37] that the electromagnetic current of a point particle (13) possesses two pieces which are entirely localized to the boundary of the worldline the particle traces out in space-time. These boundary terms turn out to be the parts of the current which dominate at low frequencies. In particular we showed that, expanding (13) in powers of q , we get

$$ij^a(-q) = e \left[\Delta \mathcal{S}_{(0)}^a(q, m\dot{X}) + \Delta \mathcal{S}_{(1)}^a(q, X, m\dot{X}) \right] + \mathcal{O}(q). \quad (14)$$

If the proper time integral in (13) runs from s_i to s_f , then the Δ in this expression has the meaning

$$\Delta f(s) \equiv f(s_f) - f(s_i) = \int_{s_i}^{s_f} \partial_s f(s), \quad (15)$$

showing that these contributions live on the endpoints of the particle worldline, at s_i and s_f , and we have defined the leading and sub-leading electromagnetic “soft factors”

$$\mathcal{S}_{(0)}^a(q, p) \equiv \frac{p^a}{q \cdot p}, \quad \mathcal{S}_{(1)}^a(q, x, p) \equiv i \frac{q_b J^{ba}}{q \cdot p}. \quad (16)$$

Here, $J^{ab} \equiv 2p^{[a} x^{b]}$ is the angular momentum tensor. Ref. [37] further showed that these boundary terms explain the factorization of the leading and sub-leading soft photon theorems at tree level.

We emphasize that these boundary terms on the worldline are *not* necessarily located at asymptotic coordinate times $t \rightarrow \pm\infty$. They exist and take the form (14) for any choice of bounds on the proper time integral in (13).

From eq. (14) it is then clear that we can always split the point particle current into three pieces,

$$j^a(q) \equiv j_{div}^a(q) + j_{sub}^a(q) + j_{hard}^a(q), \quad (17)$$

with

$$\begin{aligned} j_{div}^a(q) &\equiv ie\Delta \frac{\dot{X}^a}{q \cdot \dot{X}} \\ j_{sub}^a(q) &\equiv e\Delta \frac{q_b [X^a \dot{X}^b - \dot{X}^a X^b]}{q \cdot \dot{X}} \\ j_{hard}^a(q) &\equiv j^a(q) - j_{div}^a(q) - j_{sub}^a(q). \end{aligned} \quad (18)$$

We interpret the currents j_{div}^a , j_{sub}^a , and j_{hard}^a as the sources of leading soft radiation, sub-leading soft radiation, and hard radiation, respectively. In this work, we then *define* “leading soft radiation” as radiation which is sourced by j_{div}^a , “sub-leading soft radiation” as radiation which is sourced by j_{sub}^a , and “hard radiation” as radiation which is sourced by j_{hard}^a . This interpretation is illustrated in Figure 3.

We note immediately that the leading soft current j_{div}^a is *infrared divergent*, as it goes like $1/\omega$ and blows up as $\omega \rightarrow 0$. The sub-leading soft current j_{sub}^a is however $\mathcal{O}(1)$ and perfectly infrared-finite, and the hard current j_{hard}^a is $\mathcal{O}(\omega)$ and vanishes completely in the soft $\omega \rightarrow 0$ limit. Additionally we note that each of these currents is individually conserved, i.e.,

$$q_a j_{div}^a(q) = q_a j_{sub}^a(q) = q_a j_{hard}^a(q) = 0. \quad (19)$$

The identification of these soft currents will allow us to rather directly assess the contribution to the decoherence functional (9) from leading and sub-leading soft photon modes. In particular we will expand the decoherence functional (9) in terms of the splitting (17) to get

$$\begin{aligned} \Gamma &= \frac{e^2}{2} \int \frac{d^3q}{(2\pi)^3 2\omega} P_{ab} \times \\ &[\delta j_{div}^a(q) + \delta j_{sub}^a(q) + \delta j_{hard}^a(q)] \times \\ &[\delta j_{div}^b(-q) + \delta j_{sub}^b(-q) + \delta j_{hard}^b(-q)], \end{aligned} \quad (20)$$

displaying clearly the contributions from the leading and sub-leading soft currents to Γ .

We will see later on that this expression for Γ is unphysical, since it contains divergent terms that in the end make no contribution. The true physical form for Γ will be given in eq. (32) below.

B. Decoherence Functional for Interferometer

The general form of a decoherence functional Γ is independent of the geometry of the system – but we wish to have a specific form for our 2-path system. Later we will see what is specific to our interferometric setup, and what is more generally true.

To do this, we first write the electromagnetic currents along the left and right trajectories as

$$\begin{aligned} j_L^a(q) &= ie \int ds e^{iq \cdot X_L(s)} \partial_s \left[\frac{\dot{X}_L^a(s)}{q \cdot \dot{X}_L(s)} \right], \\ j_R^a(q) &= ie \int ds e^{iq \cdot X_R(s)} \partial_s \left[\frac{\dot{X}_R^a(s)}{q \cdot \dot{X}_R(s)} \right]. \end{aligned} \quad (21)$$




FIG. 3. Left – two charged particle trajectories which source different hard radiation, but the same soft radiation. Right – two trajectories which source different hard and soft radiation, due to differences in the soft currents j_{div}^a and j_{sub}^a at the worldline boundary. The hard radiation is shown schematically in purple, the leading soft radiation in red, and the sub-leading soft radiation in orange.

The difference between the two currents on the superposed paths,

$$\delta j^a(q) \equiv j_L^a(q) - j_R^a(q) \quad (22)$$

can then be approximated in the geometry illustrated in Figure 2 as

$$\delta j^a(q) = ie [e^{iq \cdot X_i} - e^{iq \cdot X_L} - e^{iq \cdot X_R}] \left[\frac{\dot{X}_1^a}{q \cdot \dot{X}_1} - \frac{\dot{X}_2^a}{q \cdot \dot{X}_2} \right]. \quad (23)$$

Notice that the current difference consists of terms associated with the points X_i^a , X_L^a , and X_R^a . These are the space-time events at which the particle accelerates and would radiate photons.

If we assume that the particle travels non-relativistically – by which we mean that

$$v \equiv \frac{l}{\tau} \ll 1 \quad (24)$$

in units with $c = 1$ – then we can assume that the spatial parts of the phases appearing in (23) can be neglected to leading order. This is just the well-known dipole approximation [46]; we are assuming that the radiation emitted during the experiment will have a wavelength that is long compared to the spatial extent of the interferometer, and so cannot resolve the spatial separation of the points at which the particle radiates.

In this approximation, the current difference can be further simplified to

$$\delta j^a(q) \approx ie (1 - 2e^{i\omega\tau}) \left[\frac{\dot{X}_1^a}{q \cdot \dot{X}_1} - \frac{\dot{X}_2^a}{q \cdot \dot{X}_2} \right]. \quad (25)$$

Plugging this current difference into the decoherence

functional gives

$$\Gamma \approx \frac{e^2}{4(2\pi)^3} \int_{\lambda}^{\Omega} \frac{d\omega}{\omega} (5 - 4 \cos \omega\tau) \times \oint dS^2(\hat{n}) \omega^2 \left[2 \frac{\dot{X}_1 \cdot \dot{X}_2}{(q \cdot \dot{X}_1)(q \cdot \dot{X}_2)} - \frac{1}{(q \cdot \dot{X}_1)^2} - \frac{1}{(q \cdot \dot{X}_2)^2} \right]. \quad (26)$$

Note that the sphere integral here is independent of ω , since the term in square brackets goes like $\frac{1}{\omega^2}$ and this gets multiplied by the factor ω^2 from the integration measure.

Note here that although the decoherence functional formally includes a frequency integration from $\omega = 0$ to $\omega \rightarrow \infty$, we have included a hard ultraviolet cutoff Ω in order to characterize the size of the “scattering” regions $X_{\{i,L,R\}}^a$ inside of which the particle experiences acceleration. An ultraviolet cutoff of this sort also emerges naturally if one smears out the superposed trajectories we are considering, allowing the currents to have support along worldtubes of finite width $\sim 1/\Omega$, rather than on perfectly-localized one-dimensional worldlines [9]. The length scale $l_0 \equiv 1/\Omega$ is then the shortest length scale relevant to our analysis.

The result (26) for Γ incorporates all the soft contributions – what we will now show is that the contribution from the leading divergent contributions is actually zero.

IV. CONTRIBUTION FROM LEADING DIVERGENT TERM

In order for the integral (26) to be well-defined, we have also been forced to introduce an infrared cutoff λ . The integrand of (26) diverges like $1/\omega$ as $\omega \rightarrow 0$, so the frequency integral goes to infinity like $-\ln \lambda$ as we take the infrared cutoff $\lambda \rightarrow 0$. Taken naively, this leads one to predict an infinite amount of decoherence, since $\Gamma \rightarrow \infty$ in the IR limit; this is clearly unphysical.

In earlier discussion of IR divergences, noted in the introduction, it was necessary to introduce complicated devices (eg., measuring systems designed to observe IR photons) to deal with these divergences, and show how they were either cancelled or otherwise eliminated. However in what follows we will show how our understanding of soft currents allows these divergences to be handled in a fairly straightforward way, which simply reduces their contribution to Γ to zero.

A. Infrared Divergences

Begin by splitting the current (25) into its relevant soft ($\mathcal{O}(\omega^{-1})$ and $\mathcal{O}(1)$) and hard ($\mathcal{O}(\omega)$) pieces. In this

model we have $\delta j^a \equiv \delta j_{div}^a + \delta j_{sub}^a + \delta j_{hard}^a$, with

$$\begin{aligned} \delta j_{div}^a(q) &\approx -ie \left[\frac{\dot{X}_1^a}{q \cdot \dot{X}_1} - \frac{\dot{X}_2^a}{q \cdot \dot{X}_2} \right] \\ \delta j_{sub}^a(q) &\approx 2e\omega\tau \left[\frac{\dot{X}_1^a}{q \cdot \dot{X}_1} - \frac{\dot{X}_2^a}{q \cdot \dot{X}_2} \right] \\ \delta j_{hard}^a(q) &\approx 2ie (1 - e^{i\omega\tau} + i\omega\tau) \left[\frac{\dot{X}_1^a}{q \cdot \dot{X}_1} - \frac{\dot{X}_2^a}{q \cdot \dot{X}_2} \right] \end{aligned} \quad (27)$$

in the dipole approximation.

From the general form of the decoherence functional, eq. (20), we see that Γ contains a term involving only the leading soft current j_{div}^a :

$$\begin{aligned} \Gamma &\supset \frac{1}{2} \int \frac{d^3q}{(2\pi)^3} 2\omega P_{ab} \delta j_{div}^a(q) \delta j_{div}^b(-q) \\ &= \frac{e^2}{4(2\pi)^3} \int_{\lambda}^{\Omega} \frac{d\omega}{\omega} \times \oint dS^2(\hat{n}) \omega^2 \left[2 \frac{\dot{X}_1 \cdot \dot{X}_2}{(q \cdot \dot{X}_1)(q \cdot \dot{X}_2)} - \frac{1}{(q \cdot \dot{X}_1)^2} - \frac{1}{(q \cdot \dot{X}_2)^2} \right]. \end{aligned} \quad (28)$$

It is this term that causes the infrared divergence when taking $\lambda \rightarrow 0$, seen here as the logarithmically divergent integral $\int_{\lambda}^{\Omega} \frac{d\omega}{\omega}$. The infrared divergence is caused by a superposition of the leading soft current, i.e. $\Gamma \rightarrow \infty$ because $\delta j_{div}^a \neq 0$. The resulting infinite amount of decoherence should thus be understood as coming from information loss into leading soft photon modes.

B. Infrared Dressing

If this result turned out to be the end of the story, then the leading-order soft photons would always make a perfect which-path measurement of the charged particle in any experiment of the sort shown in Figure 2. This is obviously wrong: if it were the case, we would have $\Gamma \rightarrow \infty$ for *any* choice of experimental parameters (e, l, τ) , and from eq. (5) we see that there would *never* be any observable interference pattern once the particle reaches the detector! Because quantum interference is observed in Nature, we must reject this result.

This conclusion is not new, and indeed has informed all investigations of IR divergences in QED since Bloch and Nordsieck [12]. That one needs to properly handle these divergences is also obvious in the eikonal formulation of QED [22, 23] and was a motivating factor in the coherent state formulation of QED in terms of dressed states [25–31]. In the coherent state formulation, the IR divergences are eliminated by dressing the electron, so that soft photons appear already in the time evolution operator.

It turns out to be incredibly simple to effect IR dressing in our treatment. We show in Appendix A that the net effect of applying a “minimal” implementation of the

dressed formalism, which we define in that appendix, is that we can simply set the divergent soft current j_{div}^a to zero everywhere in our calculations – the physical result of IR dressing can then be interpreted as the total decoupling of j_{div}^a from the Maxwell field. After incorporating the dressing, when studying the electromagnetic radiation caused by the charged particle moving through our model interferometer, what we do is define *dressed coherent states* in the same way as we defined the coherent radiation states in eq. (3), but now explicitly subtracting out the divergent current j_{div}^a , i.e., we write:

$$||L/R\rangle\rangle \equiv \mathcal{T}e^{-i \int d^4x [j_{L/R}^a - j_{L/R,div}^a](x)\hat{A}_a(x)}|0\rangle \quad (29)$$

We can then define a *dressed decoherence functional* in the same way we defined Γ in the “undressed” case, but using the dressed coherent states instead,

$$|\langle\langle R|L\rangle\rangle| \equiv e^{-\Gamma_{dressed}}. \quad (30)$$

The dressed decoherence functional $\Gamma_{dressed}$ can be obtained from the undressed decoherence functional (20) by simply setting the divergent current j_{div}^a to zero, giving

$$\Gamma_{dressed} = \frac{e^2}{2} \int \frac{d^3q}{(2\pi)^3 2\omega} P_{ab} \times \quad (31)$$

$$[\delta j_{sub}^a(q) + \delta j_{hard}^a(q)] \times$$

$$[\delta j_{sub}^b(-q) + \delta j_{hard}^b(-q)].$$

The expression (31) makes it clear that in the dressed formalism, the leading soft current no longer plays a role in decoherence of the matter state. Because infrared divergences necessitate this dressing, we conclude that leading soft photons do not cause decoherence, and therefore do not carry any quantum information about the matter.

V. SUB-LEADING SOFT PHOTONS

After having “dressed away” the leading soft photon contributions to decoherence, we can now begin our study of the remaining sub-leading contributions. We will do this by once more evaluating the decoherence functional, but this time using the dressed expression (31).

A. True Dressed Decoherence Functional

We begin by inserting the sub-leading and hard parts of the current difference from (27) into (31) gives

$$\Gamma_{dressed} \approx \frac{2e^2}{(2\pi)^3} \int_0^\Omega \frac{d\omega}{\omega} (1 - \cos \omega\tau) \times \quad (32)$$

$$\oint dS^2(\hat{n}) \omega^2 \left[2 \frac{\dot{X}_1 \cdot \dot{X}_2}{(q \cdot \dot{X}_1)(q \cdot \dot{X}_2)} - \frac{1}{(q \cdot \dot{X}_1)^2} - \frac{1}{(q \cdot \dot{X}_2)^2} \right].$$

This expression incorporates both the sub-leading current difference $\delta j_{sub}^a(q)$, and the hard current difference $\delta j_{hard}^a(q)$; and cross-terms between them. We stress that since the contribution from the divergent leading photon terms to Γ is zero, eq. (32) should be regarded as the correct expression for the decoherence functional.

We see that after incorporating the infrared dressing, the integrand of the frequency integral now goes to zero as $\omega \rightarrow 0$, killing off the divergence we found before, and rendering the integral finite so that we can evaluate it properly, even with the infrared cutoff λ now taken to zero. The dressing is working as intended, and we note that since there are no further infrared divergences, there seems to be no immediate need to add any sort of “sub-leading dressing.”

To arrive at (32), we have again assumed that the particle travels through the interferometer non-relativistically, $v \ll 1$, which, as discussed earlier, allowed us to invoke the dipole approximation, in which we ignored spatial parts of the phases appearing in the decoherence functional. In almost any realistic interferometer, $v \ll 1$; thus, when evaluating $\Gamma_{dressed}$ we can now also assume that $\Omega\tau \gg 1$, i.e. that the time of completion of the experiment, τ , is much greater than the time it takes light to cross the “scattering” regions at the corners of the interferometer, at which the particle accelerates, so that $\tau \gg 1/\Omega$.

We evaluate the expression (32) using these approximations in Appendix B1, with the result

$$\Gamma_{dressed} \approx \frac{e^2 v^2}{3\pi^2} \ln \Omega\tau. \quad (33)$$

which agrees with previous calculations of electromagnetic decoherence [9, 45].

Note also that the decoherence is dipolar as expected from electromagnetic radiation – we can write $\Gamma_{dressed} \propto (\partial_t e\delta x)^2$, where $e\delta x$ is the difference in electric dipole moments between the L and R branches.

B. Purely Sub-Leading Contribution

In addition to computing the full amount of photon decoherence in our model, we would now like to focus on the decoherence coming only from the sub-leading terms in our dressed expression (32) for $\Gamma_{dressed}$.

In the general form (31) of the dressed decoherence functional, we see that the purely sub-leading term $P_{ab}\delta j_{sub}^a\delta j_{sub}^b$, the cross terms $P_{ab}\delta j_{sub}^a\delta j_{hard}^b$, and the purely hard part $P_{ab}\delta j_{hard}^a\delta j_{hard}^b$ are all of different orders in $\omega\tau$, and so can be classified according to this order. The first purely sub-leading term gives decoherence caused purely by superpositions of the sub-leading soft current. Writing this contribution as Γ_{sub} , we then

have

$$\begin{aligned}\Gamma_{sub} &\equiv \frac{1}{2} \int \frac{d^3q}{(2\pi)^3 2\omega} P_{ab} \delta j_{sub}^a(q) \delta j_{sub}^b(-q) \\ &= \frac{e^2}{(2\pi)^3} \int_0^\Omega d\omega \omega \tau^2 \times \\ &\oint dS^2(\hat{n}) \omega^2 \left[2 \frac{\dot{X}_1 \cdot \dot{X}_2}{(q \cdot \dot{X}_1)(q \cdot \dot{X}_2)} - \frac{1}{(q \cdot \dot{X}_1)^2} - \frac{1}{(q \cdot \dot{X}_2)^2} \right],\end{aligned}\quad (34)$$

where we have used the expression for δj_{sub}^a found in eq. (27), and where we see that this result is nothing but the lowest order term in the expansion of the integrand of (32) in powers of $\omega\tau$.

The quantity Γ_{sub} has an intuitive interpretation. Imagine allowing the electromagnetic field to couple *only to the sub-leading soft current* j_{sub}^a during our hypothetical interferometry experiment. The resulting states of the electromagnetic radiation sourced by the sub-leading current on the L/R branches would then be

$$|l/r\rangle \equiv \mathcal{T} e^{-i \int d^4x j_{L/R,sub}^a(x) \hat{A}_a(x)} |0\rangle, \quad (35)$$

and Γ_{sub} would encode the modulus of the inner product between these two states as

$$|\langle r|l\rangle| \equiv e^{-\Gamma_{sub}}. \quad (36)$$

Γ_{sub} is then a decoherence functional in its own right, and is the natural measure of the decoherence caused by the electromagnetic field's direct response to a superposition of sub-leading soft currents.

We evaluate the sub-leading contribution (34) in Appendix B 2, with the result

$$\Gamma_{sub} \approx \frac{e^2}{3\pi^2} \Omega^2 v^2 \tau^2. \quad (37)$$

Because $v \equiv l/\tau$, this expression is equivalent to

$$\Gamma_{sub} \approx \frac{e^2}{3\pi^2} \Omega^2 l^2 \equiv \frac{e^2}{3\pi^2} \left(\frac{l}{l_0}\right)^2, \quad (38)$$

where we have used the fundamental length scale $l_0 \equiv 1/\Omega$ defined by our ultraviolet cutoff.

The result (38) is then that the purely subleading contributions to $\Gamma_{dressed}$ have a remarkable form: they only depend upon the strength of the coupling e of the particle to the electromagnetic field, and on the size of the interferometer l , measured in units of l_0 , the size of a ‘‘pixel’’ in the implicit coarse-graining of space-time implied by our use of an ultraviolet cutoff. Unlike the expression (33) for the total amount of decoherence, this purely sub-leading soft contribution does not depend directly upon the particle's velocity v or on the time τ taken to carry out the experiment.

VI. DISCUSSION

Interpreting the result (38) is mostly straightforward. The full electromagnetic field causes an amount of decoherence given by (33). The amount of this decoherence attributable solely to the difference between the sub-leading soft radiation emitted by each branch of the matter superposition is given by (38), which shows that the sub-leading soft contribution grows quadratically in l , the size of the interferometer. This makes sense, because sub-leading soft photons are by definition excitations of the electromagnetic field with a very long wavelength, and are capable of resolving only large-scale details of the matter system. The interpretation of (38) is then: the larger the matter system, the more capable sub-leading soft photons are of making quantum measurements of it. For larger experiment geometries, more of the quantum information broadcast by the charged particle then ends up being stored in sub-leading soft modes, relative to the hard modes. Keep in mind of course that we have obtained our results using the dipole approximation, in which we assume that the wavelength of any emitted radiation is still much larger than the size of the experiment. It would be interesting to go beyond this approximation in future work.

It is also interesting to reflect upon the different roles played in our model by the leading and sub-leading soft photon modes. Leading soft radiation was completely removed from our discussion by the incorporation of infrared dressing, meaning that leading soft photons do not carry quantum information about the matter. Sub-leading soft photons in contrast do carry quantum information away from the charged particle, causing an amount of decoherence given by (38).

We could, if we wanted, also consider ‘‘dressing away’’ the sub-leading modes [44]. In our semiclassical model we could try adding sub-leading dressing in the same way as we dressed away the leading modes, by setting j_{sub}^a to zero everywhere. However, although we *can* do this, it isn't clear why we would *need* to; infrared divergences appear to mandate that we dress away j_{div}^a , but the contribution of j_{sub}^a is perfectly well-behaved. Furthermore the soft currents j_{div}^a and j_{sub}^a have been shown [37] to encode the leading and sub-leading soft photon theorems *at tree level*. While the leading soft photon theorem is exact at tree level, the sub-leading soft photon theorem in general receives loop corrections [47–50]. It then seems prudent to explore quantum corrections to our simple semiclassical model, if we hope to fully understand whether any sub-leading dressing is required, and what form it might take.

VII. ACKNOWLEDGEMENTS

We thank J. Wilson-Gerow for discussions of this work, which was supported by NSERC in Canada.

Appendix A: Dressing, in Detail

In this appendix, we will illustrate the usual “dressed” formalism [25–30] using a scattering amplitude involving only a single charged particle in QED. This simple example suffices for our needs, since our interferometry model too involves only a single charged particle. We then show how the dressing works in our finite-time model, proving that a minimal form of infrared dressing can be accomplished simply by setting the divergent soft current j_{div}^a to zero everywhere.

1. Dressed Scattering

In textbook QED, the scattering amplitude for a charged particle to transition from the momentum eigenstate $|p_i\rangle$ to the momentum eigenstate $|p_f\rangle$, while the photon field transitions from $|\alpha\rangle$ to $|\beta\rangle$ is

$$\langle p_f, \beta | \hat{S} | p_i, \alpha \rangle, \quad (\text{A1})$$

where \hat{S} is the QED S-matrix. Applying infrared dressing to this formalism amounts to making the replacement

$$\langle p_f, \beta | \hat{S} | p_i, \alpha \rangle \rightarrow \langle p_f, \beta | e^{-\hat{R}_f} \hat{S} e^{\hat{R}_i} | p_i, \alpha \rangle, \quad (\text{A2})$$

where the “dressing operator,” which acts on the photon Hilbert space, is

$$\hat{R}_{i/f} \equiv \sum_{h=+,-} \int \frac{d^3q}{(2\pi)^3 \sqrt{2\omega}} \left[F_{i/f}^h(q) \hat{a}_h^\dagger(q) - \bar{F}_{i/f}^h(q) \hat{a}_h(q) \right], \quad (\text{A3})$$

in which $\hat{a}_h^\dagger(q)$ and $\hat{a}_h(q)$ respectively create and annihilate a photon with momentum $q^a \equiv \omega(1, \hat{n})$ and helicity h , and obey

$$\left[\hat{a}_h(q), \hat{a}_{h'}^\dagger(q') \right] = \delta_{hh'} (2\pi)^3 \delta^{(3)}(\vec{q} - \vec{q}'). \quad (\text{A4})$$

We have also defined

$$F_{i/f}^h(q) \equiv e^{\frac{\epsilon_h(q) \cdot p_{i/f}}{q \cdot p_{i/f}}} \phi(q, p_{i/f}), \quad (\text{A5})$$

with ϵ_h^a the polarization vector corresponding to the helicity h , and $\phi(q, p)$ is a mostly arbitrary function, which must go smoothly to 1 as $|\vec{q}| \rightarrow 0$ in order for the dressing to properly remove the infrared divergence we saw in the main text.

To remove the divergence, it suffices to choose $\phi(q, p)$ nonzero only in a neighborhood of $|\vec{q}| = 0$, where the divergence occurs. Such a choice would however necessitate the introduction of an arbitrary new energy scale “ Λ ” to our model, in order to characterize the size of the momentum neighborhood involved in the dressing. There is no physical energy scale in our interferometry model with which we could identify Λ , so instead we make a

more conservative choice involving no new parameters at all, simply letting

$$\phi(q, p) = 1, \quad (\text{A6})$$

a choice which has been referred to elsewhere as “minimal” dressing [51]. With this choice, the frequency integral in (A3) should be understood as extending from the infrared cutoff λ up to the ultraviolet cutoff Ω , with λ eventually being taken to zero. We will use this choice of $\phi(q, p)$ in what follows, and we will see that the minimal dressing has a very clean interpretation in terms of the infrared divergent soft matter current j_{div}^a , making it a natural choice in light of the deep interplay between the soft currents and infrared radiation [37].

Evolving states using the dressed S-matrix $e^{-\hat{R}_f} \hat{S} e^{\hat{R}_i}$ instead of the bare S-matrix \hat{S} allows one to avoid unphysical infrared divergent decoherence rates. This has been shown already in scattering calculations [31], and we will now show how it works in our finite-time interferometry model. In particular we want to dress the interaction picture time evolution operator, which we used in eq. (3), as

$$\begin{aligned} \mathcal{T} e^{-i \int d^4x j^a(x) \hat{A}_a(x)} \\ \rightarrow e^{-\hat{R}_f} \left[\mathcal{T} e^{-i \int d^4x j^a(x) \hat{A}_a(x)} \right] e^{\hat{R}_i}, \end{aligned} \quad (\text{A7})$$

with $\hat{R}_{i/f}$ given by (A3), but now with the factors $F_{i/f}^h(q)$ of (A5) evaluated at the beginning and end of the interferometry experiment (at proper times $s_{i/f}$), rather than at the asymptotic times $t \rightarrow \pm\infty$ relevant in scattering scenarios. We will now show that in order to effect minimal dressing, using (A6), we need only set the divergent part j_{div}^a of the electromagnetic current j^a appearing here to zero. That is, we will show that

$$\begin{aligned} e^{-\hat{R}_f} \left[\mathcal{T} e^{-i \int d^4x j^a(x) \hat{A}_a(x)} \right] e^{\hat{R}_i} \\ = \mathcal{T} e^{-i \int d^4x [j^a - j_{div}^a](x) \hat{A}_a(x)}. \end{aligned} \quad (\text{A8})$$

The mechanism by which minimal infrared dressing removes infrared divergences is thus the wholesale decoupling of the Maxwell field from the infrared divergent soft current j_{div}^a .

2. Minimal Dressing Removes the Divergent Soft Current

To prove eq. (A8), let us begin with the second line and work our way backward. The second term in the exponent there is

$$i \int d^4x j_{div}^a(x) \hat{A}_a(x), \quad (\text{A9})$$

with j_{div}^a defined by eq. (18). Using the usual mode expansion of $\hat{A}_a(x)$,

$$\hat{A}_a(x) = \sum_{h=+,-} \int \frac{d^3q}{(2\pi)^3 \sqrt{2\omega}} [\epsilon_a^h \hat{a}_h^\dagger(q) e^{iq \cdot x} + \bar{\epsilon}_a^h \hat{a}_h(q) e^{-iq \cdot x}], \quad (\text{A10})$$

we can rewrite (A9) as

$$\begin{aligned} & i \int d^4x j_{div}^a(x) \sum_{h=+,-} \int \frac{d^3q}{(2\pi)^3 \sqrt{2\omega}} [\epsilon_a^h \hat{a}_h^\dagger e^{iq \cdot x} + \bar{\epsilon}_a^h \hat{a}_h e^{-iq \cdot x}] \\ &= i \sum_{h=+,-} \int \frac{d^3q}{(2\pi)^3 \sqrt{2\omega}} [\epsilon_a^h \hat{a}_h^\dagger j_{div}^a(q) + \bar{\epsilon}_a^h \hat{a}_h j_{div}^a(-q)] \\ &= \sum_{h=+,-} \int \frac{d^3q}{(2\pi)^3 \sqrt{2\omega}} [\bar{\epsilon}_a^h \hat{a}_h - \epsilon_a^h \hat{a}_h^\dagger] \frac{\dot{X}^a(s_f)}{q \cdot \dot{X}(s_f)} - (s_f \rightarrow s_i). \end{aligned} \quad (\text{A11})$$

Comparing to eqs. (A3) and (A5), this is nothing but

$$-\hat{R}_f + \hat{R}_i, \quad (\text{A12})$$

with $p_{i/f}^a = m \dot{X}^a(s_{i/f})$ and the choice of minimal dressing $\phi(q, p) = 1$.

This result allows us to conclude that

$$\begin{aligned} & \mathcal{T} e^{-i \int d^4x j^a - j_{div}^a(x) \hat{A}_a(x)} \\ &= \mathcal{T} e^{-i \int d^4x j^a(x) \hat{A}_a(x) - \hat{R}_f + \hat{R}_i}. \end{aligned} \quad (\text{A13})$$

We can rewrite this using the form (2) of the position space current $j^a(x)$ as

$$\mathcal{T} e^{-ie \int_{s_i}^{s_f} ds \dot{X}^a(s) \hat{A}_a(X(s)) - \hat{R}_f + \hat{R}_i}, \quad (\text{A14})$$

in order to see explicitly that the time ordering of operators using the coordinate time t gives the same result as ordering operators using the proper time s . Because the dressing operators here act at the beginning and end of the experiment – i.e. $\hat{R}_{i/f}$ contains a contribution only when the proper time is $s_{i/f}$ – we can thus pull \hat{R}_f out from under the time ordering operator \mathcal{T} to the left, and we can pull \hat{R}_i out to the right, leaving

$$e^{-\hat{R}_f} \left[\mathcal{T} e^{-ie \int_{s_i}^{s_f} ds \dot{X}^a(s) \hat{A}_a(X(s))} \right] e^{\hat{R}_i}, \quad (\text{A15})$$

or equivalently

$$e^{-\hat{R}_f} \left[\mathcal{T} e^{-i \int d^4x j^a(x) \hat{A}_a(x)} \right] e^{\hat{R}_i}, \quad (\text{A16})$$

proving eq. (A8).

Appendix B: Integrals

This appendix includes further detail on the evaluation of the dressed decoherence functional $\Gamma_{dressed}$, and the sub-leading soft contribution Γ_{sub} to this, which are defined in the main text. Our approach to these calculations is similar to that of Breuer & Petruccione [9].

1. The Dressed Decoherence Functional

First write $\Gamma_{dressed}$ (eq. (32)) as

$$\Gamma_{dressed} \equiv \frac{2e^2}{(2\pi)^3} \mathcal{I}_\omega \mathcal{I}_{\hat{n}} \quad (\text{B1})$$

with

$$\mathcal{I}_\omega \equiv \int_0^\Omega d\omega \frac{(1 - \cos \omega\tau)}{\omega} \quad (\text{B2})$$

and

$$\begin{aligned} \mathcal{I}_{\hat{n}} &\equiv \int dS^2(\hat{n}) \omega^2 \left[2 \frac{\dot{X}_1 \cdot \dot{X}_2}{(q \cdot \dot{X}_1)(q \cdot \dot{X}_2)} - \frac{1}{(q \cdot \dot{X}_1)^2} - \frac{1}{(q \cdot \dot{X}_2)^2} \right]. \end{aligned} \quad (\text{B3})$$

We will evaluate each of these integrals one at a time.

The frequency integral \mathcal{I}_ω can be written as (using $\varpi \equiv \omega\tau$)

$$\int_0^{\Omega\tau} d\varpi \frac{(1 - \cos \varpi)}{\varpi}. \quad (\text{B4})$$

We can then use the following identity [52] involving the cosine integral,

$$\begin{aligned} \text{Ci}(\Omega\tau) &\equiv - \int_{\Omega\tau}^\infty d\varpi \frac{\cos \varpi}{\varpi} \\ &= \gamma_{EM} + \ln \Omega\tau - \int_0^{\Omega\tau} d\varpi \frac{(1 - \cos \varpi)}{\varpi}, \end{aligned} \quad (\text{B5})$$

where $\gamma_{EM} \approx 0.577$ is the Euler-Mascheroni constant, from which it follows that \mathcal{I}_ω is

$$\int_0^{\Omega\tau} d\varpi \frac{(1 - \cos \varpi)}{\varpi} = \gamma + \ln \Omega\tau - \text{Ci}(\Omega\tau). \quad (\text{B6})$$

In the limit $\Omega\tau \gg 1$ discussed in the main text, the logarithm dominates this expression, giving the approximate result

$$\mathcal{I}_\omega \approx \ln \Omega\tau. \quad (\text{B7})$$

The angular integral $\mathcal{I}_{\hat{n}}$ is a sum of three terms with integrals of the form

$$\begin{aligned} & \int dS^2(\hat{n}) \omega^2 \frac{\dot{X}_A \cdot \dot{X}_B}{(q \cdot \dot{X}_A)(q \cdot \dot{X}_B)} \\ &= \int dS^2(\hat{n}) \frac{(1 - \vec{v}_A \cdot \vec{v}_B)}{(1 - \hat{n} \cdot \vec{v}_A)(1 - \hat{n} \cdot \vec{v}_B)}, \end{aligned} \quad (\text{B8})$$

with $A, B \in \{1, 2\}$. The Lorentz-invariance of this integral allows us to evaluate it in any reference frame. For convenience we can choose the frame in which $\vec{v}_B = 0$, bringing $\mathcal{I}_{\hat{n}}$ into the simple form

$$\int dS^2(\hat{n}) \frac{1}{(1 - \hat{n} \cdot \vec{v}_{AB})} = \frac{4\pi}{v_{AB}} \tanh^{-1} v_{AB}, \quad (\text{B9})$$

where v_{AB} is the relative velocity,

$$v_{AB} \equiv \sqrt{1 - \frac{1}{(\dot{X}_A \cdot \dot{X}_B)^2}}. \quad (\text{B10})$$

We can use the result (B9) to see that the angular integral (B3) is

$$\mathcal{I}_{\hat{n}} = 8\pi \left[\frac{1}{v_{12}} \tanh^{-1} v_{12} - 1 \right]. \quad (\text{B11})$$

In the geometry illustrated in Figure 2, the relative velocity v_{12} is approximately

$$v_{12} \approx \sqrt{2}v, \quad (\text{B12})$$

in the case of non-relativistic speeds, $v \ll 1$, where again $v \equiv l/\tau$. We can further approximate

$$\frac{1}{v_{12}} \tanh^{-1} v_{12} = 1 + \frac{1}{3}v_{12}^2 + \dots \approx 1 + \frac{2}{3}v^2 \quad (\text{B13})$$

In this limit, the angular integral is

$$\mathcal{I}_{\hat{n}} \approx \frac{16\pi}{3}v^2. \quad (\text{B14})$$

Plugging eq. (B7) and eq. (B14) into eq. (B1) yields eq. (33),

$$\Gamma_{dressed} \approx \frac{4e^2v^2}{3\pi^2} \ln \Omega\tau. \quad (\text{B15})$$

2. The Sub-Leading Soft Contribution to the Decoherence Functional

The sub-leading soft part of the dressed decoherence functional, given by equation (34), can also be written in the form of a frequency integral multiplied by an angular integral:

$$\Gamma_{sub} \equiv \frac{e^2}{(2\pi)^3} \mathcal{I}'_{\omega} \mathcal{I}_{\hat{n}} \quad (\text{B16})$$

Here, the angular integral $\mathcal{I}_{\hat{n}}$ is the same one we have already evaluated, so we can simply use the result (B14). The frequency integral is different, however; we have

$$\mathcal{I}'_{\omega} \equiv \int_0^{\Omega} d\omega \omega\tau^2 = \frac{1}{2}\Omega^2\tau^2. \quad (\text{B17})$$

Plugging these results into (B16) gives the result (37)

$$\Gamma_{sub} \approx \frac{e^2}{3\pi^2} \Omega^2 v^2 \tau^2. \quad (\text{B18})$$

-
- [1] R.P. Feynman, R.B. Leighton, and M. Sands, *The Feynman Lectures on Physics*, vol. 3, Addison-Wesley, (1965).
- [2] To rigorously divide the matter paths into discrete classes requires a little work – see A. Auerbach, S. Kivelson, and D. Nicole, *Path Decomposition for Multidimensional Tunneling*, PRL **53**, 411 (1984); A. Auerbach, S. Kivelson, *The path decomposition expansion and multidimensional tunneling*, Nucl. Phys **B 257**, 799 (1985), for non-relativistic discussions appropriate to, e.g., a 2-path system.
- [3] E. Joos and H.D. Zeh, *The emergence of classical properties through interactions with the environment*, Zeitschrift für Physik, **B59**, 223 (1985).
- [4] E. Joos et al., *Decoherence and the appearance of a classical world in quantum theory* (2nd ed.), Springer (2003)
- [5] H.-P. Breuer and F. Petruccione, *The Theory of Open Quantum Systems*, Oxford University Press (2002).
- [6] P.C.E. Stamp, *The decoherence puzzle*, Studies in History and Philosophy of Science Part B: Studies in History and Philosophy of Modern Physics **37**, 467 (2006).
- [7] A. Stern, Y. Aharonov, and Y. Imry, *Phase uncertainty and loss of interference: A general picture*, Phys. Rev. **A41**, 3436 (1990).
- [8] L.H. Ford, *Electromagnetic vacuum fluctuations and electron coherence*, Phys. Rev. **D 47**, 5571 (1993).
- [9] H.-P. Breuer and F. Petruccione, *Destruction of quantum coherence through emission of bremsstrahlung*, Phys. Rev. **A 63**, 032102 (2001).
- [10] see, eg., K. Hornberger, J. E. Sipe, and M. Arndt, *Theory of decoherence in a matter wave Talbot-Lau interferometer*, Phys. Rev. A **70**, 053608 (2004).
- [11] N.V. Prokof'ev and P.C.E. Stamp, *Theory of the Spin Bath*, Rep. Prog. Phys. **63**, 669 (2000).
- [12] F. Bloch and A. Nordsieck, *Note on the radiation field of the electron*, Phys. Rev. **52** (1937) 54
- [13] V. V. Sudakov, *Vertex Parts at Very High Energies in Quantum Electrodynamics*, Sov. Phys. J.E.T.P. **3**, 65 (1956).
- [14] A.A. Abrikosov, *The Infrared Catastrophe in Quantum Electrodynamics*, Sov. Phys. J.E.T.P. **3**, 71 (1956).
- [15] T. Murota, *On radiative corrections due to soft photons*, Prog. Th. Phys. **24**, 1109 (1960).
- [16] T. Kinoshita, *Note on the infrared catastrophe*, Prog. Theor. Phys. **5**, 1045 (1950)
- [17] N. Nakanishi, *General theory of infrared divergence*, Prog. Theor. Phys. **19**, 159 (1958)
- [18] D.R. Yennie, S.C. Frautschi, and H. Suura, *The infrared divergence phenomenon and high-energy processes*, Ann. Phys. **13**, 379 (1961).
- [19] T. Kinoshita, *Mass singularities of Feynman amplitudes*, J. Math. Phys. **3**, 650 (1962)

- [20] G. Grammer and D.R. Yennie, *improved treatment for the infrared divergence problem in Quantum Electrodynamics*, Phys. Rev. **D 8**, 4332 (1973).
- [21] S. Weinberg, *Infrared photons and gravitons*, Phys. Rev. **140**, B516 (1965).
- [22] E.S. Fradkin, *Application of functional methods in quantum field theory and quantum statistics. (I). Divergence-free field theory with local non-linear interaction*, Nucl. Phys. **49**, 624 (1963)
- [23] E.S. Fradkin, *Application of functional methods in quantum field theory and quantum statistics. (II)*, Nucl. Phys. **76**, 588 (1966)
- [24] J. Wilson-Gerow, C. DeLisle and P.C.E. Stamp, *A functional approach to soft graviton scattering and BMS charges*, Class. Quant. Grav. **35**, 164001 (2018)
- [25] V. Chung, *Infrared Divergence in Quantum Electrodynamics*, Phys. Rev. **140**, B1110 (1965).
- [26] T.W.B. Kibble, *Coherent soft-photon states and infrared divergences. I. classical currents*, Lect. Theor. Phys. **D11**, 387 (1969).
- [27] T.W.B. Kibble, *Coherent soft-photon states and infrared divergences. II. mass shell singularities of Green's functions*, Phys. Rev. **173**, 1527 (1968).
- [28] T.W.B. Kibble, *Coherent soft-photon states and infrared divergences. III. asymptotic states and reduction formulas*, Phys. Rev. **174**, 1882 (1968).
- [29] T.W.B. Kibble, *Coherent soft-photon states and infrared divergences. IV. the scattering operator*, Phys. Rev. **175**, 1624 (1968).
- [30] P.P. Kulish and L.D. Faddeev, *Asymptotic conditions and infrared divergences in quantum electrodynamics*, Theor. Math. Phys. **4**, 745 (1970)
- [31] D. Carney, L. Chaurette, D. Neuenfeld, and G. Semenoff, *On the need for soft dressing*, JHEP 2018, 121, (2018).
- [32] G.W. Semenoff, *Entanglement and the Infrared*, Dobrev, V. (eds) Lie Theory and Its Applications in Physics, LT 2019, Springer Proceedings in Mathematics & Statistics, vol **335**, Springer, Singapore (2020).
- [33] F.E. Low, *Scattering of light of very low frequency by systems of spin 1/2*, Phys. Rev. **96**, 1428 (1954).
- [34] F.E. Low, *Bremsstrahlung of very low-energy quanta in elementary particle collisions*, Phys. Rev. **110**, 974 (1958).
- [35] T.H. Burnett and N.M. Kroll, *Extension of the Low soft photon theorem*, Phys. Rev. Lett. **20**, 86 (1968).
- [36] M. Gell-Mann and M.L. Goldberger, *Scattering of low-energy photons by particles of spin 1/2*, Phys. Rev. **96**, 1433 (1954).
- [37] C. DeLisle, J. Wilson-Gerow, and P.C.E. Stamp, *Soft theorems from boundary terms in the classical point particle currents*, JHEP 2021, 290 (2021).
- [38] G. Grynberg, A. Aspect, and C. Fabre, *Introduction to Quantum Optics*, Cambridge University Press (2010).
- [39] Decoherence functionals are described in, eg., A. Schmid, J. Low Temp. Phys. **49**, 609 (1982).
- [40] R.P. Feynman and F.L. Vernon, *Theory of a general quantum system interacting with a linear dissipative system*, Ann. Phys. (NY) **24**, 118 (1963).
- [41] R.P. Feynman and A.R. Hibbs, *Quantum Mechanics and Path Integrals*, Dover (1965).
- [42] V. Dinu, T. Heinzl, and A. Ilderton, *Infrared divergences in plane wave backgrounds*, Phys. Rev. D **86**, 085037 (2012).
- [43] A. Laddha and A. Sen, *Logarithmic Terms in the Soft Expansion in Four Dimensions*, JHEP 2018, 56 (2018).
- [44] S. Choi and R. Akhoury, *Subleading soft dressings of asymptotic states in QED and perturbative quantum gravity*, JHEP 2019, 31 (2019).
- [45] D. Neuenfeld, *Infrared-safe scattering without photon vacuum transitions and time-dependent decoherence*, JHEP 2021, 189 (2021).
- [46] J.D. Jackson, *Classical Electrodynamics*, 3rd ed., Wiley (1999).
- [47] V. Del Duca, *High-energy Bremsstrahlung Theorems for Soft Photons*, Nucl. Phys. B **345**, 369 (1990).
- [48] Z. Bern, S. Davies, and J. Nohle, *On Loop Corrections to Subleading Soft Behavior of Gluons and Gravitons*, Phys. Rev. D **90**, 085015 (2014).
- [49] S. He, Y.-t. Huang, and C. Wen, *Loop Corrections to Soft Theorems in Gauge Theories and Gravity*, JHEP 2014, 115 (2014).
- [50] B. Sahoo and A. Sen, *Classical and Quantum Results on Logarithmic Terms in the Soft Theorem in Four Dimensions*, JHEP 2019, 86 (2019).
- [51] D. Kapec, M. Perry, A.-M. Raclariu, and A. Strominger, *Infrared Divergences in QED, Revisited*, Phys. Rev. D **96**, 085002, (2017).
- [52] I.S. Gradshteyn and I.M. Ryzhik, *Table of Integrals, Series, and Products*, Academic Press, New York (1980).