

A sub-Neptune planet around TOI-1695 discovered and characterized with SPIRou and TESS

Flavien Kiefer^{1,2*}, G. Hébrard^{2,3}, E. Martioli^{4,2}, E. Artigau⁵, R. Doyon⁵, J.-F. Donati⁶, C. Cadieux⁵, A. Carmona⁷, D. R. Ciardi⁸, P. I. Cristofari⁶, L. de Almeida⁴, P. Figueira^{11,12}, E. Gaidos²⁸, E. Gonzales⁹, A. Lecavelier², K. G. Stassun¹⁰, L. Arnold¹³, B. Benneke¹⁴, I. Boisse¹⁵, X. Bonfils⁷, N. J. Cook⁵, P. Cortés-Zuleta¹⁵, X. Delfosse⁷, J. Dias do Nascimento²², M. Fausnaugh¹⁶, W. Fong¹⁶, P. Fouqué⁵, T. Forveille⁷, J. Gomes da Silva²⁵, K. Hesse¹⁶, Á. Kóspál^{29,30,31,32}, H. Lewis¹⁷, C.-F. Liu²⁷, J. H. C. Martins²⁵, M. Paegert¹⁸, S. Seager^{19,20,21}, H. Shang²⁷, J. D. Twicken^{22,23}, T. Vandal⁵, S. Vinatier¹, T. Widemann^{1,26}, J. N. Winn²⁴

(Affiliations can be found after the references)

Received ; accepted

ABSTRACT

TOI-1695 is a V-mag=13 M-dwarf star from the northern hemisphere at 45 pc from the Sun, around which a 3.134-day periodic transit signal from a super-Earth candidate was identified in TESS photometry. With a transit depth of 1.3 mmag, the radius of candidate TOI-1695.01 was estimated by the TESS pipeline to be $1.82 R_{\oplus}$ with an equilibrium temperature of ~ 620 K. We successfully detect a reflex motion of the star and establish it is due to a planetary companion at an orbital period consistent with the photometric transit period thanks to a year-long radial-velocity monitoring of TOI-1695 by the SPIRou infrared spectropolarimeter. We use and compare different methods to reduce and analyse those data. We report a $5.5\text{-}\sigma$ detection of the planetary signal, giving a mass of $5.5 \pm 1.0 M_{\oplus}$ and a radius of $2.03 \pm 0.18 R_{\oplus}$. We derive a mean equilibrium planet temperature of 590 ± 90 K. The mean density of this small planet of $3.6 \pm 1.1 \text{ g cm}^{-3}$ is similar ($1.7\text{-}\sigma$ lower) than that of the Earth. It leads to a non-negligible fraction of volatiles in its atmosphere with $f_{\text{H,He}} = 0.28^{+0.46}_{-0.23}\%$ or $f_{\text{water}} = 23 \pm 12\%$. TOI-1695 b is a new sub-Neptune planet at the border of the M-dwarf radius valley that can help test formation scenarios for super-Earth/sub-Neptune-like planets.

Key words. Planets and satellite: detection – Planets and satellite: fundamental parameters – Planets and satellite: individual: TOI-1695 b – techniques: photometric – techniques: radial velocities

1. Introduction

Over the next few years, new space- and ground-based observatories (e.g. the James Webb Space Telescope and ESO's extremely large telescope) will be paramount to extend the current boundaries of planetary science. These will boost the exploration of the atmospheres of exoplanets to an unprecedented level of precision, and probe the gas content of super-Earth and terrestrial exoplanets. It is thus of key importance to identify new low-mass exoplanets amenable to future atmospheric characterisation.

A central feature of small exoplanets ($R < 4 R_{\oplus}$) at small orbital separation ($P < 100$ days) is the radius valley (Owen & Wu 2013; Fulton et al. 2017; Cloutier & Menou 2020) separating solid super-Earths from gaseous sub-Neptunes. One possible explanation for this dichotomy is runaway evaporation or grinding of the atmospheres of low-mass exoplanets by a strong XUV stellar radiation (Owen & Wu 2013). Alternative explanations are core-powered mass loss (Ginzburg et al. 2018), impact erosion by planetesimals (Shuvalov 2009), or formation of distinct rocky and non-rocky planet populations with delayed gas accretion (Lee et al. 2014; Lee & Connors 2021). Around solar-like stars, photoevaporation and core-powered mass loss scenarios predict that the valley location shifts towards lower radius with lower stellar mass and larger orbital period (Van Eylen et al. 2018; Fulton & Petigura 2018; Martinez et al. 2019; Wu 2019; Gupta et al. 2022).

Around low-mass stars, Cloutier & Menou (2020) find that the slope of the valley around M dwarfs in the radius-period diagram is inverse to the valley slope around FGK stars, leading to increasing radius of the transition with orbital period. This relationship is more compatible with a gas-depleted planet formation scenario in which solid cores increase in size with increasing distance to the star (Lee et al. 2014; Lopez & Rice 2018). Similarly than for solar-like stars, for planets around low-mass stars, photoevaporation and core-powered mass loss are also expected to shape a radius gap with a negative slope in $R_p - P$ diagram. However this slope is expected to be closer and closer to zero with decreasing stellar mass (Gupta et al. 2022).

Here we characterize a candidate planet with a super-Earth/sub-Neptune size, that was detected and identified around the M2-type star TOI-1695, with the NASA Transiting Exoplanet Survey Satellite (TESS; Ricker et al. 2015). With an initially estimated radius of $1.8 R_{\oplus}$ and an orbital period of 3.134 days, TOI-1695.01 lies in a radius/separation region where the question of formation and evolution of the atmosphere of super-Earth or sub-Neptune around low-mass star can be tackled. Investigating whether or not such a key planet harbors a significant atmosphere may help explain which scenario is dominantly shaping the radius valley depending on the host star mass.

In this paper, we seek to first establish the planetary nature of TOI-1695.01, and to characterize its mass and radius by combining TESS photometric measurements with radial velocities (RV) variations observed with the infrared (0.98–2.45 μm) SpectroPo-

* flavien.kiefer@obspm.fr

larimetre InfraRouge (SPIRou; Donati et al. 2020) installed on the 3.6-m Canada France Hawaii Telescope (CFHT).

In Section 2 we review the observations available for TOI-1695, including those from TESS and SPIRou. In Section 3 we characterize the host star using spectroscopy, spectrophotometry and spectropolarimetry. In Section 4 we characterize the planet properties through a joint fit of TESS and SPIRou data. In Section 5 we discuss the results. We summarize and conclude in Section 6.

2. Observations

2.1. TESS lightcurves

TOI-1695, also known as TIC-422756130.01 in the TESS Input Catalog (TIC), was covered by TESS sectors 18, 19, 24, 25 and 52 between November 2019 and June 2022 with 120-s cadence. In the data from these observing campaigns, a signal at a 3.13-day period is found using the Box-Least-Square (BLS) algorithm (Kovács et al. 2002) with 34 transits found, among which 32 were already identified in the data validation timeseries (DVT hereafter; Twicken et al. 2018; Li et al. 2019) of sectors 18 to 25, allowing the event TOI-1695.01 to be identified. We retrieved and used in the rest of this study the Science Processing Operations Center Pipeline (SPOC; Jenkins et al. 2016) simple aperture photometry with presearch data conditioning (PDCSAP; Smith et al. 2012; Stumpe et al. 2014). The full SPOC PDCSAP lightcurve with all identified transits is shown in Fig. 1.

We first performed a preliminary fit of the TESS PDCSAP data of TOI-1695 by a photometric batman model (Kreidberg 2015) with a third-degree polynomial modeling of the continuum of all individual transits. It has a transit depth of ~ 1000 ppm for a total transit duration of 1.2 hour and a period of 3.134319 ± 0.000026 days. It was pointed out¹ that the SPOC photometry pipeline was prone to over-estimate the background level and over-correct for background in crowded fields in the TESS primary mission (sectors 1-26). The pipeline was updated to prevent this in the extended mission (sectors 27+). We estimate that the combined transit search of the TOI 1695 data for sectors 18, 19, 24, 25 and 52 would have over-estimated the transit depth (planet radius) by approximately 2.8% (1.4%). This is a minor correction compared to uncertainties ($\sim 8.9\%$) resulting from the full analysis performed in Section 4.

Using TRICERATOPS (Giacalone & Dressing 2020), we examined possible sources of false-positive (FP) transit detection in the TOI-1695 lightcurve, from the system itself and from nearby background systems within the TESS aperture. The full list of possible causes of an FP is given in Table 1 of Giacalone et al. (2021). We show the aperture used in Sector 18 and possible contaminating sources around TOI-1695 in Fig. A.1. This source is located in a crowded region with 411 neighbors with $T\text{-mag} < 18$ at less than $200''$. At most 8 stars identified in the different TESS apertures could be contributing to the transit signal with TIC IDs 422756137, 422756132, 422756120, 629325853, 422756126, 422756145, 422756114, and 422756147. They have a flux ratio with TOI-1695 in TESS passband between 5.3 and 7.4 mag. We find an FP probability of transit from TOI-1695 itself of 0.18% and an FP probability of a background star pollution of 0.10%. These probabilities accounts for the contrast curve around TOI-1695 obtained in Section 2.3. According to the criteria defined by Giacalone et al. (2021), TOI-1695 b is thus a

"validated planet". Nevertheless, complementary follow-up with RV, as performed in the present paper, is necessary to independently validate its planetary nature and measure its mass.

Beyond the transit signal, a 15-day modulation with an average amplitude of ~ 175 ppm seems to be present in the PDCSAP light curves. A periodogram of the TESS PDCSAP data produced with *lightcurve* (Lightcurve Collaboration et al. 2018) is shown in Fig. 2. This 15-day periodic signal is in fact mainly seen in the Sector 18 with an amplitude of ~ 700 ppm (Fig. 1). We discard effects from variable background stars, since given the sources identified above within TOI-1695 apertures, such amplitude of variations in the lightcurve would require those sources to have proper oscillations $> 10\%$ which is rare (McQuillan et al. 2012). If not of instrumental origin, which we cannot fully exclude, this fluctuating signal could well be due to spots, in which case the 15-day period would be linked with the rotation period of the star. We discuss this possibility in more details and confront it to other observations in Section 3.4.




Fig. 1. The full TESS lightcurve separated in sectors (gray dots) and binned with a 0.1-day timestep (black). The identified transit locations are shown with orange vertical lines.




Fig. 2. Periodogram of the TESS SPOC lightcurve with a peak at ~ 15 days for a 175 ppm signal.

¹ https://archive.stsci.edu/missions/teess/doc/teess_drn/teess_sector_27_drn38_v02.pdf

2.2. SPIRou spectra

2.2.1. Observations

SPIRou is a near-infrared high-resolution (980-2450 nm; $R=75,000$) high-stability ($\sim 1 \text{ m s}^{-1}$) velocimeter and spectropolarimeter installed at the Cassegrain focus of the 3.6-m CFHT at Maunakea (Donati et al. 2020). TOI-1695 was observed between Dec. 2020 and Jan. 2022 with SPIRou as part of the large program SPIRou Legacy Survey (SLS; id P42, PI: Jean-François Donati). It was observed at 45 observing epochs with 4 polarimetric exposures per epoch, totaling 180 SPIRou spectra. Table F.1 presents the observation log associated with each exposure. All SPIRou observations of TOI-1695 consist of polarization sequences, each split into 4 sub-exposures associated with different rhomb retarder configurations, from which we retrieve the corresponding Stokes I (unpolarised) and V (circularly polarized) spectra of the star at each epoch (Donati et al. 2020). The sequence number 1-4 of any sub-exposure is also given in Table F.1. Science data are acquired on fibers A and B fed by the 2 orthogonal states of the selected polarization, while fiber C is used for simultaneously recording the spectrum of the SPIRou Fabry-Pérot RV reference, as described in Hobson et al. (2021).

A calibration sequence, including flats, darks, and spectra of comparison lamps (Fabry-Pérot, UNE), is performed every afternoon and morning, preceding and following each night of observation with SPIRou. A master-dark is constructed from a subset of well characterized darks. The SPIRou detector software also archives ramp-files, that are detector images in ADU/s, during which creation, a correction for non-linearity is performed. Early-type stars (A-type) are observed nightly as telluric absorption standards. Bright inactive cool stars are also regularly observed as RV standards. These data are used to calibrate the measurements extracted from the SPIRou spectra. The detailed calibration sequence is described in Cook et al. (2022, submitted).

2.2.2. APERO reduction

The raw data were first reduced with the SPIRou reduction software, A PipelinE to Reduce Observations or APERO, with version 0.6.132 (Cook et al., submitted), or v6 hereafter. APERO first corrects detector effects, removes constant background thermal component, and identify bad pixels and cosmic ray impacts. It then calculates the position of the 49 of the 50 echelle spectral orders² and optimally extracts spectra from fibers A, B and C into 2D order-separated e2ds and 1D order-merged s1d spectra. A blaze function is derived from the flat-field exposures.

The wavelengths are calibrated on the spectrum collected on Fiber C. An absolute calibration of wavelengths with respect to the Solar System barycentric rest frame using the current Barycentric Earth RV (BERV) and the Barycentric Julian Date (BJD) of each exposure is performed by the code *barycorrpy* (Kanodia & Wright 2018; Wright & Eastman 2014). Finally, for each exposure (*e.fits* file) APERO calculates the corresponding spectrum of the telluric transmission out of the whole collection of standard star observations carried out with SPIRou since 2018 using PCA algorithm (Artigau et al. 2014) and divides it out, producing a telluric-corrected *t.fits* spectrum. APERO also calculates the Stokes V spectra using the method of Donati et al. (1997), as described in detail in Martioli et al. (2020).

A more advanced version of the APERO pipeline, version 0.7.194, was released during the writing of this work (Artigau et al. 2022; Cadieux et al. 2022), hereafter called v7. We used

both the v6 and v7 versions in our study, and compare them in Section 4.

2.2.3. RV derivation

The RV were derived from the telluric corrected *t.fits* spectra using the line-by-line (LBL) algorithm (Artigau et al. 2022). As in Bouchy et al. (2001), LBL requires a template spectrum with S/N as large as possible, since a derivative of the template is used to determine the RV of each spectral line. For TOI-1695, with only 40 epochs, the combined spectrum produced by APERO reaches a S/N much lower than other bright standard stars monitored with SPIRou. Instead of TOI-1695, we thus used as template another target with a similar M2V spectral type, GL15A ($T_{\text{eff}}=3600 \text{ K}$; $\log g=4.8 \text{ cgs units}$; $[M/H]=-0.16 \text{ dex}$; Passegger et al. 2019; Cristofari et al. 2022b). Table 2 shows the stellar parameters that we find for TOI-1695. We combined 1040 spectra of GL15A with for each a S/N ~ 350 at 1670 nm leading to a template with S/N close to 10 000.

The second and third derivatives of each spectral line are used as a proxy for the full width at half maximum (FWHM) and bisector span (BIS) (Artigau et al. 2022). Their variations with respect to RV are studied in Section 3.3.

Calibration drifts are derived for all RV measurement epochs using the simultaneous Fabry-Pérot on fiber C. They lead to drift-corrected (DC hereafter) RV. An average zero-point correction (ZPC, hereafter) is also derived from the monitoring of RV standards (Cadieux et al. 2022). Our reference RVs in the rest of the paper will be those corrected from drift and zero-point, but we do a comparison of RV data reduction with or without drift and zero-point correction when analysing the planet signal in Section 4.

All RV data, FWHM and BIS used for TOI-1695 and reduced with versions v6 and v7 are available in Tables G.1 and G.2, respectively. Their variations are inspected in more details in Section 3.3. In both cases, we ignored the measurement of epoch #44 (JD-2459605) that has only a single sub-exposure. With the v6, we also ignored the measurements of epoch #33 (JD-2459591) for which the drift calculation failed.

2.3. Imaging

As part of a standard process to validate transiting exoplanets and to assess the possible contamination of bound or unbound companions on the derived planetary radii (Ciardi et al. 2015), TOI-1695 was observed with infrared high-resolution adaptive optics (AO) imaging at Keck Observatory with the NIRC2 instrument on Keck-II behind the natural guide star AO system (Wizinowich et al. 2000). The observations were made on 2020 May 28 UT in the standard 3-point dither pattern that is used with NIRC2 to avoid the lower left quadrant of the detector which is typically noisier than the other three quadrants. The dither pattern step size was $3''$ and was repeated twice, with each dither offset from the previous dither by $0.5''$. The camera was in the narrow-angle mode with a full field of view of $\sim 10''$ and a pixel scale of approximately $0.01''$ per pixel. The observations were made in the narrow-band Br- γ filter ($\lambda_o=2.17\mu\text{m}$; $\Delta\lambda=0.03\mu\text{m}$) with an integration time of 4 s with one coadd per frame for a total of 36 s on target.

The AO data were processed and analyzed with a custom set of IDL tools. The science frames were flat-fielded and sky-subtracted. The flat fields were generated from a median of dark subtracted flats taken on-sky, and the flats were normalized such

² APERO does not extract the bluest order due to low S/N.

that the median value of the flats is unity. Sky frames were generated from the median average of the 9 dithered science frames; each science image was then sky-subtracted and flat-fielded. The reduced science frames were combined into a single combined image using an intra-pixel interpolation that conserves flux, shifts the individual dithered frames by the appropriate fractional pixels, and median-coadds the frames. The final resolution of the combined dithers was determined from the FWHM of the point spread function to 0.054".

The sensitivities of the final combined AO image were determined by injecting simulated sources azimuthally around the primary target every 20° at separations of integer multiples of the central source's FWHM (Furlan et al. 2017). The brightness of each injected source was scaled until standard aperture photometry detected it with 5 σ significance. The resulting brightness of the injected sources relative to the target set the contrast limits at that injection location. The final 5 σ limit at each separation was determined from the average of all of the determined limits at that separation and the uncertainty on the limit was set by the RMS dispersion of the azimuthal slices at a given radial distance.

No additional companions to within the limits of the data were detected (see Fig. 3). With contrast sensitivities of ~ 3.5 mag at 0.06" (2.7 au) and ~ 7 mag at 0.5" (22 au), the NIR AO observations indicate that there are likely no stellar companions down to $\sim M_6 - L_9$ (see E. Mamajek's compilation of Mean Dwarf Stellar Colors version 2021.03.02³).




Fig. 3. Companion sensitivity for the Keck NIR adaptive optics imaging. The black points represent the 5 σ limits and are separated in steps of 1 FWHM ($\sim 0.054''$); the purple represents the azimuthal dispersion (1 σ) of the contrast determinations (see text). The inset image is of the primary target showing no additional companions to within 3" of the target.

3. Stellar characterization

3.1. Stellar parameters

We use four different methods to measure the stellar parameters of the star, including the effective temperature T_{eff} , the surface gravity $\log g$, the metallicity $[M/H]$, the micro-turbulent v_{micro} , the macro-turbulent velocity v_{macro} , the bolometric flux F_{bol} , the bolometric luminosity L_{bol} , the stellar mass $M_{\star}$, and the stellar

Table 1. Main astrometric and photometric data of TOI-1695.

Parameter	Unit	Value	Ref.
TIC		422756130	
Gaia DR2/EDR3		534988616816537728	
2MASS		J01274094+7217472	
WISE		J012741.12+721747.6	
RA	ICRS (J2000)	01:27:40.973	1
DEC	ICRS (J2000)	+72:17:47.186	1
PM RA μ_{α}	mas yr ⁻¹	71.628 $\pm$ 0.013	1
PM DEC μ_{δ}	mas yr ⁻¹	40.450 $\pm$ 0.019	1
Epoch		2016.0	1
Gaia DR3 parallax	(mas)	22.226 $\pm$ 0.014	1
distance	(pc)	45.131 $\pm$ 0.068	1
GAIA G_{BP}	mag	13.3064 $\pm$ 0.0040	1
GAIA G	mag	12.1275 $\pm$ 0.0011	1
GAIA G_{RP}	mag	11.0464 $\pm$ 0.0022	1
TESS T	mag	11.0294 $\pm$ 0.0074	
2MASS J	mag	9.640 $\pm$ 0.024	2
2MASS H	mag	8.984 $\pm$ 0.028	2
2MASS K	mag	8.818 $\pm$ 0.021	2
WISE 3.4 μ m	mag	8.684 $\pm$ 0.024	3
WISE 4.6 μ m	mag	8.61 $\pm$ 0.02	3
WISE 12 μ m	mag	8.511 $\pm$ 0.027	3
WISE 22 μ m	mag	8.40 $\pm$ 0.29	3

Notes. (1) Gaia Collaboration et al. (2021); (2) Cutri et al. (2003); (3) Wright et al. (2010).

radius $R_{\star}$. The methods of spectral Energy Distribution (SED) and of the TESS-exoFOP are summarized in Section 3.1.1. The methods of comparison to synthetic spectra, based on SPIRou spectra, are explained in Section 3.1.2. All results are summarized and compared in Table 2.

3.1.1. Spectral Energy Distribution

We performed an analysis of the broadband spectral energy distribution (SED) of the star together with the *Gaia* EDR3 parallax (with no systematic offset applied; see in particular Stassun & Torres 2021), in order to determine an empirical measurement of the stellar radius, following the procedures described in Stassun & Torres (2016); Stassun et al. (2017, 2018). We retrieved the JHK_S magnitudes from 2MASS, the W1–W4 magnitudes from WISE, and the $G_{BP}G_{RP}$ magnitudes from *Gaia*, as well as the NUV flux from GALEX. Together, the available photometry spans the stellar SED over the wavelength range 0.2–22 μ m (see Fig. 4). All photometric data are summarized in Table 1.

We performed a fit using NextGen stellar atmosphere models (Hauschildt et al. 1999), with the free parameters being the effective temperature (T_{eff}) and metallicity ($[Fe/H]$), as well as the extinction A_V , which we limited to maximum line-of-sight value from the Galactic dust maps of Schlegel et al. (1998). The resulting fit (Fig. 4) has a best-fit $A_V = 0.02 \pm 0.02$, $T_{\text{eff}} = 3630 \pm 50$ K, and $[Fe/H] = 0.0 \pm 0.5$, with a reduced χ^2 of 1.3. It also fit well with the calibrated mean *Gaia* BP/RP spectrum available for this star in *Gaia* DR3 (Montegriffo et al. 2022) as shown in Fig. 4. Integrating the (unreddened) model SED gives the bolometric flux at Earth, $F_{\text{bol}} = 6.82 \pm 0.24 \times 10^{-10}$ erg s⁻¹ cm⁻², which with the *Gaia* parallax gives directly the bolometric luminosity, $L_{\text{bol}} = 0.0431 \pm 0.0015 L_{\odot}$. Taking the F_{bol} and T_{eff} together with the *Gaia* parallax gives the stellar radius, $R_{\star} = 0.525 \pm 0.017 R_{\odot}$. In addition, we can estimate the stellar mass from the empirical M_K relations of Mann et al. (2019a), giving $M_{\star} = 0.539 \pm 0.027 M_{\odot}$. These mass and radius lead to $\log g = 4.72 \pm 0.14$ in cgs units.

³ https://www.pas.rochester.edu/~emamajek/EEM_dwarf_UBVIJHK_colors_Teff.txt

The stellar parameters from the TESS project can be found in the TESS-exoFOP and are also reported in Table 2. TOI-1695 belongs to the specially curated list of cool dwarfs of the TIC (Muirhead et al. 2018; Stassun et al. 2019). As explained in Stassun et al. (2019), the effective temperature T_{eff} is retrieved from the cool dwarfs catalog of TIC v7 (Stassun et al. 2018). The radius is derived from T_{eff} , Gaia magnitudes and Gaia parallax using equation (4) of Stassun et al. (2019). The mass is obtained from the spline-interpolation of an empirical mass- T_{eff} relationship.

The parameters derived from both methods are consistent within uncertainties.




Fig. 4. Spectral energy distribution of TOI-1695. Red symbols represent the observed photometric measurements, where the horizontal bars represent the effective width of the passband. Blue symbols are the model fluxes from the best-fit NextGen atmosphere model (black). The green curve is the published Gaia DR3 spectrum for this star.

3.1.2. Comparison to synthetic spectra

We also derive the stellar parameters for TOI-1695 from a high-resolution template spectrum built from the tens of individual spectra acquired with SPIRou. The method used consists in the comparison of a grid of synthetic spectra to the template spectrum. We use two independent methods, both based on MARCS models (Gustafsson et al. 2008).

The first method used is based on a process briefly described below (more details in Cristofari et al. 2022b,a). This process was tested and calibrated on reference stars. The synthetic spectra used for this analysis were computed from the MARCS model atmospheres with the Turbospectrum radiative transfer code (Plez 2012), for a wide range of effective temperatures (T_{eff}), surface gravities ($\log g$) and metallicities ($[M/H]$). The models and template spectrum are compared by computing a χ^2 value in key regions, containing lines that are both well reproduced by the models and sensitive to the stellar parameters. We therefore retrieve a three-dimensional grid of χ^2 values, and perform a paraboloid fit to identify the best fitting parameters and estimate error bars on these values. For TOI-1695, the parameters estimated with this method are $T_{\text{eff}} = 3627 \pm 31$ K, $\log g = 4.60 \pm 0.05$ dex and $[M/H] = 0.10 \pm 0.10$ dex. A fit of the star's projected rotational velocity $v \sin i$, fixing the micro-turbulent velocity to $v_{\text{micro}} = 1 \text{ km s}^{-1}$, the macro-turbulence velocity to $v_{\text{macro}} = 0 \text{ km s}^{-1}$ and the resolution linewidth of SPIRou

to $\text{FWHM} \sim 4.3 \text{ km s}^{-1}$, leads to a $v \sin i = 1.9 \pm 0.2 \text{ km s}^{-1}$. Given that v_{macro} is fixed to 0, we can only interpret this measurement as an upper-limit on $v \sin i$, therefore $< 2.5 \text{ km s}^{-1}$ at 1-sigma, i.e. at most on the order of the pixel-scale of the SPIRou detector ($\sim 2.3 \text{ km s}^{-1}$; Donati et al. 2020). Modeled and observed spectral lines are compared together in Fig. B.1.

The second method used is described in greater detail in section 4.2 in Martioli et al. (2022). We calculated a grid of 650,000 synthetic spectra generated with MOOG (Snedden et al. 2012) covering T_{eff} from 2900 K to 4100 K with 50 K step, $\log g$ from 3.5 to 5.3 with 0.1 dex step, $[M/H]$ from -0.6 to 0.5 with 0.11 dex step, alpha from -0.2 to 0.4 with 0.06 dex step and v_{mic} from 0.5 km s^{-1} to 4 km s^{-1} with 0.35 km s^{-1} step. We modified the Apache Point Observatory Galactic Evolution Experiment (APOGEE) line list (Smith et al. 2021) by selecting only the lines that are most sensitive to T_{eff} and $\log g$ within the spectral range from $1.5 \mu\text{m}$ to $1.7 \mu\text{m}$. We searched the grid for the best fit parameters, which are adopted as priors in the iSpec integrator (Blanco-Cuaresma 2019) to calculate the synthetic spectrum that best fits our SPIRou data within the domain of the selected lines. We used the MOOG code with the MARCS GES model (Gustafsson et al. 2008) and Asplund et al. (2009) solar abundances. In total, 78 spectra were compared. The parameters of TOI-1695 estimated with this method are $T_{\text{eff}} = 3711 \pm 47$ K, $\log g = 4.53 \pm 0.13$ dex, $[M/H] = 0.00 \pm 0.08$ dex, and $v_{\text{micro}} = 0.3 \pm 0.7 \text{ km s}^{-1}$.

The values derived with these two methods agree well together, with moreover good agreement with the SED fitting at 2σ . In the rest of the analysis, we thus adopt the values $T_{\text{eff}} = 3650 \pm 40$ K, $\log g = 4.72 \pm 0.14$, and $[Fe/H] = 0.0 \pm 0.1$ dex. We also adopt the $M_{\star} = 0.54 \pm 0.03 M_{\odot}$ and $R_{\star} = 0.53 \pm 0.02 R_{\odot}$ derived from the SED fitting (Section 3.1.1 above) that rely directly on the spectrophotometry of the star with robust empirical relations.

3.2. Polarimetry

An independent polarimetric reduction and LSD analysis of TOI-1695 SPIRou data using the Libre-Esprit (LE) pipeline (Donati et al. 1997, 2020), leads to the polarimetric longitudinal field, B_{ℓ} , for TOI-1695 shown in Fig. C.1 and given in Table G.3. The B_{ℓ} measurements are consistent with those obtained in the classical APERO reduction (e.g. Martioli et al. 2022). At a given epoch, B_{ℓ} is directly determined from the Stokes V and Stokes I profiles (Donati et al. 1997). We find that the star has a weak polarimetric variability and magnetic activity. The Stokes V profiles have a $\chi_r^2 = 0.8$ for a constant model, while the "null polarization" N (Donati et al. 1997) has a $\chi_r^2 = 1$. Any variability in the data is thus fully consistent with the error bars.

A power spectrum of the B_{ℓ} data (Fig. 5) shows peaks beyond 20 days, all with a false-alarm probability (FAP) $> 10\%$. Thus no peculiar polarimetric signal is significantly detected for TOI-1695 given the current precision. The period of ~ 48 days seems nevertheless to dominate the power-spectrum at small frequencies. We fitted the data with a quasi-periodic Gaussian process (GP; see e.g. Haywood et al. 2014; Aigrain et al. 2015) using the same codes as in Donati et al. (2017), at an initial period of 48 days with typical decay-time $\tau_{\text{decay}} = 500$ days and smoothing factor $\gamma_{\text{smooth}} = 0.6$. We obtain an almost flat GP model with a marginal likelihood difference $\Delta \ln \mathcal{L} = 2$ compared to a constant model. The GP model is plotted over the longitudinal field on Fig. C.1.

We can thus conclude that (i) we do not see trace of a 15 days signal as reported in Section 2.1, (ii) the star was, if active, passing through a quiet phase during SPIRou observations, and

Table 2. Spectral parameters of the star derived from spectral template matching with the method from (Cristofari et al. 2022b) and de Almeida et al. (in prep.), with the SED fitting method (Stassun & Torres 2016; Stassun et al. 2017, 2018), and with the TESS project stellar parameters derivation method (Stassun et al. 2018).

Parameter	Unit	TurboSpectrum+MARCS	Moog+MARCS	SED	TESS Input Catalog v8 ^f
T_{eff}	K	3627±31	3711±47	3630±50	3575±157
$\log(g)$		4.60±0.05	4.53±0.13	4.5±0.5	4.723±0.009
[M/H]	dex	0.10±0.10	0.00±0.08	0.0±0.5	
$v \sin i$	km s ⁻¹	<2.5 ^a			
v_{micro}	km s ⁻¹	fixed to 1.0	0.3±0.7		
F_{bol}	10 ⁻¹⁰ erg s ⁻¹ cm ⁻²			6.82±0.24 ^b	
L_{bol}	L _⊙			0.0431±0.0015 ^c	0.039±0.009
$M_{\star}$	M _⊙			0.54±0.03 ^d	0.51±0.02
$R_{\star}$	R _⊙			0.53±0.02 ^e	0.52±0.02

Notes.

^(a) At 3-sigma and fixing $v_{\text{macro}}=0$ km s⁻¹ (more details in Section 3.1.2).

^(b) From direct integration of SED.

^(c) Luminosity derived from the bolometric flux and the parallax.

^(d) Mass determined from Mann et al. (2019b) empirical relations.

^(e) Radius determined from the bolometric flux and the parallax.

^(f) Taken from Stassun et al. (2019). See text for details.

(iii) more data are needed to confirm whether the hinted 48-days period could be a trace of the star’s rotation signal.

3.3. RV variations and stellar activity effects

We searched for correlations between RVs and the activity indicators, bisector span (BIS) and FWHM variations, obtained from the two APERO version v6 & v7 studied in this work. The Pearson correlation coefficient for RV–BIS are $R_{v6}=0.14$ (p -value = 0.07) and $R_{v7}=-0.22$ (p -value = 0.004). The Pearson correlation coefficient for RV–FWHM is $R_{v6}=-0.13$ (p -value = 0.10) and $R_{v7}=0.06$ (p -value = 0.46). The only significant slope that we detect ($1.7-\sigma$) is for the FWHM and RV derived from the v6, of -0.62 ± 0.37 m s⁻¹ per m s⁻¹. It disappears with the v7 suggesting that some instrumental signal was corrected from the SPIRou data with the last implementations in the pipeline.

We thus find no significant correlations of astrophysical origin between RV and the activity indicators FWHM and BIS. This is compatible with the weak magnetic activity deduced from the almost flat longitudinal field B_{ℓ} .

The periodograms of the BIS and FWHM compared to the periodogram of the RVs in Fig. 6 show indeed no significant peaks with FAP<5% at periods <20 days, especially none around 3.13 days, the orbital period of the candidate planet detected with TESS. Peaks with FAP<1% appear at periods beyond 20 days, some of which coincide with large peaks of the window-function.

On the other hand, in the RV periodogram we have a positive detection of the candidate planet signal at 3.13 days with a FAP~1%. Two other strong peaks on both sides of this signal are due to aliases with the observing window frequencies. Indeed, most of the structures in the near periods of 3 days significantly weakens after fitting out a 3.13-days Keplerian to the RVs. As seen in the RV residuals periodogram in Fig. 7, residual RV variations then mainly concentrate around ~16 days. Interestingly, it is close to the 15-day period spotted in the lightcurve continuum variations of TESS sector 18 (Section 2.1). This would suggest an origin intrinsic to TOI-1695 itself or linked to its direct environment, even if no corresponding signal is detected in polarimetry. Studies of the TOI-1695’s surrounding in Sec-

tion 2.3 exclude however that the RV variations could originate from background stars at less than 3’’ distance.

Inspecting the periodogram of the LBL RV derived using the APERO v7 version (Fig. D.1) leads to similar conclusions. Only the power at low frequencies is significantly reduced from the RV, FWHM and the BIS variation, leaving only insignificant peaks, some of which are most likely bias of the Earth orbital period with the ~20-days gap period. There is indeed some power at long period ~365 days which could be linked to yearly changes in the instrument resolution.

3.4. Concluding remark on the rotation period of TOI-1695

The TESS photometry and SPIRou RVs show traces of signals with periods within 14–19 days. A rotation period of the star with a period in this range would be in agreement with the M2V spectral type found for TOI-1695 (see e.g. Goulding et al. 2012; Goulding 2013; Basri et al. 2011; Moutou et al. 2017). There are few other observations that agree or not with such stellar rotation period.

Firstly, photometry and RV do not strictly agree with each other on the period of modulation. Photometry rather show a periodogram peak at 14.8 days. The LBL RV calculated from the APERO v6 data rather show a peak at 16 days. If they are calculated from the APERO v7 version the peak even shifts to ~18 days.

Secondly, if the rotation period of the star were equal to 15 days, following the hinted photometric variations in Section 2.1, the stellar radius of $0.53 R_{\odot}$ would imply a $v \sin i$ equal to 1.8 km s⁻¹, or smaller if the stellar spin and planet orbit are misaligned. This could agree with the $v \sin i$ of 1.9 ± 0.2 km s⁻¹ that was obtained in Section 3, implying in this case that the equatorial plane of the star is aligned with the orbital plane of TOI-1695 b.

Thirdly, the SPIRou polarimetry of the star does not detect a 15-day periodicity of the star’s magnetic field during the timespan of the SPIRou observations. If due to stellar rotation, we would have seen a stronger field and detected the rotational modulation in polarimetry.




Fig. 5. Periodogram of the longitudinal magnetic field variations, with dominant frequencies highlighted in blue. The window function is shown in orange, with a peak close to the synodic orbital period of the Moon.




Fig. 6. From top to bottom, RV, FWHM and BIS Lomb-Scargle periodograms with FAP levels 0.01 and 0.05 indicated as dotted and dashed lines for data extracted with the APERO v6 version. They are compared to the window function (red solid line). The vertical orange solid line indicates the 3.134-days period of the TESS transit signal. The moon synodic orbital period (29.53 days) is also shown as a green solid lines.

Finally, we retrieved available public photometry data to search for periodic signals indicative of stellar rotation. As of 10 September 2022, TOI-1695 was observed by the All Sky-Automated Survey for SuperNovae (ASAS-SN Shappee et al. 2014; Kochanek et al. 2017) at 2043 epochs. ASAS-SN images are obtained through a Sloan *g* filter (and previously, for 879 epochs, in *V*-band). A Lomb-Scargle periodogram analysis shows a signal at 52.4 days in *V*-band, but no significant signal in *g*-band. We also retrieved 328 and 490 epochs of photometry in Sloan *g* and *r*-band from the Zwicky Transient Factory archive




Fig. 7. Periodogram of the residuals O-C after fitting a 3.134-days Keplerian to the RVs. Same color coding as in Fig. 6.

(Bellm et al. 2019; Masci et al. 2019). We find a $FAP < 1\%$ signal at 27 days in the *r*-band data but not the *g*-band data. In principle, the 27-day signal could be an upper harmonic of the ≈ 52 -day signal obtained from ASAS-SN. But no hint for a 15-day modulation. The absence of confirmatory detections in the other bands does not allow us to draw a firm conclusion about the rotation period of TOI-1695 from these data.

We conclude that the 15-day signal seen in TESS sector 18, as well as the 16 (or 18) days signal seen in the SPIRou LBL RVs, are likely not related to the rotation of the star. Polarimetry, spectroscopy, photometry and RV agree nevertheless on a star rotation period larger than 15 days. Moreover polarimetry, *V*-band and *r*-band photometry suggest a rotation period in the range 20–52 days. Such long rotation period is found for M dwarfs with a stellar age > 1 Gyr (Engle & Guinan 2018). We thus infer that TOI-1695 is more than 1 Gyr old.

4. Joint analysis of TESS and SPIRou data

We use here the same recipe as Martioli et al. (2022) to fit the combined photometric and velocimetric data from respectively TESS and SPIRou recorded for TOI-1695. We run a Markov-Chain Monte-Carlo (MCMC) algorithm with the *emcee* (Foreman-Mackey et al. 2013) routine with prior distribution shown in Table 3. We fix the quadratic limb darkening parameters u_0 & u_1 with respect to Claret (2017) and Claret & Southworth (2022), for a $T_{\text{eff}} = 3600$ K. We initialize 32 walkers, sample parameters space on 20,000 steps, and burn the 10,000 first samples. The posterior distributions of parameters are listed in Table 4.

We apply the fit on different analysis schemes of the LBL RV, using both v6 and v7 reductions (RV_{v6} and RV_{v7} hereafter). This allows comparing the quality of the RV series extracted from both APERO versions and qualifying the improvements brought

Table 3. Prior distribution of fitted or fixed parameters entering the MCMC sampling.

Parameter	Unit	Prior type	Prior distribution
Stellar parameters			
$M_\star$	$M_\odot$	Normal	$\mathcal{N}(0.54, 0.03)$
$R_\star$	$R_\odot$	Normal	$\mathcal{N}(0.53, 0.02)$
u_0		Fixed	$0.23^\dagger$
u_1		Fixed	$0.38^\dagger$
Keplerian parameters			
K	km s^{-1}	Uniform	$\mathcal{U}(0,10)$
T_0	BTJD	Uniform	$1791 + \mathcal{U}(0.2, 0.8)$
P	day	Uniform	$\mathcal{U}(2.51, 3.76)$
a	$R_\star$	Uniform	$\mathcal{U}(10, 23)$
R_p	R_J	Uniform	$\mathcal{U}(0.02, 0.04)$
I_p	$^\circ$	Uniform	$\mathcal{U}(70, 110)$
e		Uniform or Fixed	$\mathcal{U}(0, 1)$ or 0.0
ω	$^\circ$	Uniform or Fixed	$\mathcal{U}(0, 360)$ or 90
Gaussian process parameters			
$\sigma_{\text{white,phot}}$	ppm	Uniform	$\mathcal{U}(-10^{100}, 10^{100})$
$\tau_{\text{decay,phot}}$	day	Uniform	$\mathcal{U}(0.25, 1000)$
$\gamma_{\text{smooth,phot}}$		Uniform	$\mathcal{U}(0.1, 1.5)$
P_{phot}	days	Uniform	$\mathcal{U}(10, 20)$
A_{phot}	ppm	Uniform	$\mathcal{U}(10^{-4}, 2000)$
$\sigma_{\text{white,RV}}$	m s^{-1}	Uniform	$\mathcal{U}(-10^{100}, 10^{100})$
$\tau_{\text{decay,RV}}$	day	Uniform	$\mathcal{U}(6, 1000)$
$\gamma_{\text{smooth,RV}}$		Uniform	$\mathcal{U}(0.1, 1.5)$
P_{RV}	days	Uniform	$\mathcal{U}(10, 20)$
A_{RV}	m s^{-1}	Uniform	$\mathcal{U}(0, 100)$

Notes.

^(†) Based on Claret (2017); Claret & Southworth (2022).

by the new pipeline. For both versions, fits without drift correction nor zero-point correction were poor in terms of χ^2 and the O-C residuals. So we performed fits assuming:

- DC+ZPC: accounting for the long-term calibration drift of the RV and a zero-point correction (see Section 2 for details);
- DC+ZPC+GP: adding a quasi-periodic (QP) Gaussian Process (GP) to the fit, as explained in more details below.

The result of the joint Keplerian fit of both lightcurve and RVs are all compared in Table 4, assuming circular orbit.

In the DC+ZPC scheme, the 3.134-days RV signal is found with a S/N larger than $5-\sigma$. This allows us to confirm the detection of the planet signal with a corresponding companion mass in the super-Earth/sub-Neptune regime $\sim 6 M_\oplus$. Although compatible at $0.7-\sigma$, the semi-amplitude of the Keplerian model fitted to the RV_{v7} ($3.6 \pm 0.7 \text{ m s}^{-1}$) is smaller than that of the RV_{v6} ($4.1 \pm 0.7 \text{ m s}^{-1}$). On the other hand, the RV-residuals dispersion is slightly smaller for RV_{v6} than for RV_{v7} with respective reduced χ^2 of 0.8 and 1.1, and absolute O-C dispersion of 6 m s^{-1} and 7 m s^{-1} . We also noted a difference of $\sim 460 \text{ m s}^{-1}$ in systemic RV which we identified to a different treatment of the calibration between the two versions. The difference in the Keplerian solutions and residuals, implying systematic variations unaccounted for in both, motivated us into digging further with the GP analysis of those datasets.

The quasi-periodic Gaussian process that we used combines a squared exponential (SE) and a periodic kernel with the code

george (Ambikasaran et al. 2015) to remove the periodic pattern in the RV and the TESS-LC identified in Sections 2.1 and 3.3. The GP is trained separately on the residuals RVs and on the photometric time series excluding the transits. The priors used on the GP parameters are added to Table 3. We were careful not to allow the decay-time of the SE kernel to be smaller than twice the orbital period of TOI-1695 b, thus preventing the GP to overfit and absorb some of the orbital signal present in the RV, along the recommendations drawn in Angus et al. (2018). For optimising the GP hyper-parameters more efficiently using an MCMC, the photometric lightcurves were rebinned to time steps of 0.25 days. Fig. 8 and 9 show the GP modeling of the TESS lightcurve and the SPIRou RV.

The QP GP fit of the lightcurve converged to a period of $13.0^{+3.4}_{-0.7}$ days. For the RV_{v6} it converged to a period of $16.2^{+0.6}_{-3.9}$ days and for the RV_{v7} to $18.3^{+0.5}_{-4.1}$ days. Once applying and then removing the GP from the RV and photometric time series, we fit again the orbital and transit signals from the RV and LC. The DC+ZPC+GP analysis leads to the smallest RV residual dispersion with reduced χ^2 of 0.6 and 0.5 respectively of RV_{v6} and RV_{v7} . The flux dispersion ~ 2008 ppm is constant over all schemes, which is not surprising given that the lightcurve continuum is always detrended around the transits epochs. The bayesian information criterion (BIC) among all schemes is clearly in favour of the DC+ZPC+GP scheme with the smallest BIC. The posterior distributions although different are compatible with the DC+ZPC solutions. As expected, the model semi-amplitudes of RV_{v6} ($3.95 \pm 0.66 \text{ m s}^{-1}$) and RV_{v7} ($3.63 \pm 0.68 \text{ m s}^{-1}$) are smaller (only slightly for RV_{v7}) than without modeling the systematic variations. The Keplerian fit of the RV_{v7} -GP residuals leads to the best residual dispersion among all analysed datasets, $\sigma_{\text{O-C}} = 4.8 \text{ m s}^{-1}$, but with similar planetary parameters than without fitting the GP.

As an additional check, we run a fit of the data in the DC+ZPC scheme, but considering a non-zero eccentricity. The eccentricity obtained when letting it freely vary is consistent with 0 at $1.3-\sigma$ with $e < 0.2$. More importantly, at maximum likelihood, the BIC is larger than for the zero-eccentricity case. This validates that fixing the eccentricity to zero in the other fits is a correct, most parsimonious approach, consistent with a tidally-circularized orbit of the planet, and moreover not generating significant systematic errors on the other orbital parameters.

Varying the limb darkening coefficients led to wide posterior distributions of u_0 and u_1 with unrealistic values $u_0 \sim 0.5 \pm 0.4$ and $u_1 \sim 0.9 \pm 0.7$. Although compatible within errors with the common values assumed in the rest of the analysis, they lead to less accurate planet radius estimation with highly biased posteriors on semi-major axis and radius. Notably, the median of the radius posterior distribution is $1.77 \pm 0.21 R_\oplus$, but the maximum a posteriori point estimate is $\sim 1.99 R_\oplus$. This shows that the present dataset is not able on its own to refine the limb darkening coefficients. It is more reasonable to rely on the known values of the limb darkening coefficients of an M2 star.

Comparing all the different reduction/analysis schemes, we retain the solution with the smallest BIC. Finally, it is the DC+ZPC+GP derived from the APERO v7 version that leads to the best solution. It can be trusted that the GP did not overfit the RV signal as the planetary parameters are similar than if not fitting a GP. The corner plot of the posterior distribution of all varied and derived Keplerian parameters is shown in Fig. E.1. The best fitting model of the transit curve is shown in Fig. 10, and the best fitting models of the phase-folded LBL RV is shown in Fig. 11.




Fig. 8. Gaussian process fit of the lightcurve with a quasi-periodic kernel. Grey points are raw PDCSAP flux TESS data, and black points are the binned data with $\Delta t_{\text{bin}}=0.25$ days.




Fig. 9. Gaussian process fit of the SPIRou RV in the case of the APERO v6 reduction (top) and the v7 reduction (bottom) with a quasi-periodic kernel.

It results in a companion on an orbit of 3.1343 days with a planetary radius of $2.03 \pm 0.18 R_{\oplus}$ and a planet mass of $5.5 \pm 1.0 M_{\oplus}$. This implies a sub-Earth planet density of $3.6 \pm 1.1 \text{ g cm}^{-3}$, and an equilibrium temperature of $590 \pm 90 \text{ K}$.

5. Discussion on TOI-1695 b mass and radius

We find that TOI-1695 b is a sub-Neptune planet with an average density $\sim 3.6 \text{ g cm}^{-3}$ smaller than the Earth density. The mass and radius of TOI-1695 b are compared to other super-Earth/sub-Neptunes in Fig. 12, as obtained thanks to the *pyexor* package (Zeng et al. 2021; Francesco et al. 2022). TOI-1695 b sits in-between two similar sub-Neptunes, Wolf-503 b (Peterson et al. 2018; Polanski et al. 2021) and TOI-270 d (Van Eylen et al. 2021; Günther et al. 2019) in terms of mass, radius and temperature, at the border of the radius valley but on the sub-Neptune side of it. It also has similar mass and radius as K2-3 b (Kosiarek et al. 2019) although warmer (463 K for K2-3 b).




Fig. 10. Results from fits to the TESS light curve, with all the observed transits stacked, with the model taken from the maximum a posteriori point estimation, based on maximizing the posterior probability distributions for each parameter. At this point, $T_0=1791.5206 \text{ BTJD}$, $P=3.1342799$ days, $a=21.1 R_{\star}$, $R_p=0.034 R_{\star}$ & $I_p=89.4^{\circ}$.




Fig. 11. Phase-folded modelisation of the RV variation seen with SPIRou of the star TOI-1695 with a 3.13 days circular orbit. The data used here are derived from the APERO v7 version. Grey dots gather all the RV data and black dots are binned with a time step of 0.05 days.

5.1. The possible composition of the planet

Assuming a pure silicate core leads to a planet with a significant gas envelope in the formalism developed by Zeng et al. (2021). An M-R diagram comparing TOI-1695 b to other super-Earths and sub-Neptunes is shown in Fig. 12. Using the *smint* code (Piaulet et al. 2021) based on the formalism of Lopez & Fortney (2014); Zeng et al. (2016); Aguichine et al. (2021), we considered three different scenarios:

- a planet with H/He envelope and a solid interior leads to $f_{\text{H/He}}=0.28^{+0.46}_{-0.23}\%$;
- a 100% silicate interior with pure water on top of it, we derive a water envelope with a mass fraction of $f_{\text{H}_2\text{O}}=55^{+29}_{-30}\%$;

Table 4. MCMC results of the photometry (TESS) + spectroscopy (SPIRou) joint fit in the different RV-reduction schemes using APERO v6 and v7 versions, which are presented in Section 4. For all parameters, except those of GPs, the main value is the median and the uncertainty is the average distance of the median to the 16th and 84th percentiles. For the GP hyper-parameters, since the posteriors are asymmetrical, the main value is given by the mode, and the uncertainties are the distance between this mode and the 16th and 84th percentiles.

Param.	Unit	with APERO v6		with APERO v7	
		DC + ZPC	DC + ZPC + GP	DC + ZPC	DC + ZPC + GP
Fitted parameters					
K	m/s	4.12±0.69	3.95±0.66	3.69±0.65	3.63±0.68
t_c	BTJD	1791.5205±0.0013	1791.5204±0.0013	1791.5204±0.0013	1791.5203±0.0012
P	day	3.1342788±0.0000077	3.1342793±0.0000077	3.1342788±0.0000078	3.1342790±0.0000072
a_p	$R_\star$	16.3±5.2	16.7±5.0	15.6±5.3	16.3±5.1
$R_p/R_\star$		0.0349±0.0026	0.0344±0.0025	0.0351±0.0028	0.0344±0.0027
I_c	°	89.9±3.5	89.8±3.3	90.0±3.8	89.8±3.5
u_0		0.22	0.22	0.22	0.22
u_1		0.39	0.39	0.39	0.39
γ	m s ^{−1}	−59921.62±0.47	−59921.64±0.47	−59461.93±0.47	−59461.96±0.45
e		0.0	0.0	0.0	0.0
ω	°	90	90	90	90
Planet intrinsic parameters					
M_p	$M_\oplus$	6.2±1.1	6.0±1.0	5.6±1.0	5.5±1.0
R_p	$R_\oplus$	2.06±0.17	2.03±0.17	2.07±0.18	2.03±0.18
ρ_p	$\rho_\oplus$	0.71±0.22	0.71±0.21	0.63±0.20	0.66±0.21
$T_{\rm eq}$	K	580±90	575±86	590±100	590±90
$T_{\rm irr}$	K	900±140	890±130	920±160	900±140
S_p	$S_\oplus$	27±17	26±16	30±17	27±17
Goodness of fit diagnostics					
$\chi^2_{\rm r, RV}$		0.79	0.66	1.16	0.49
O-C _{RV}	m s ^{−1}	5.9	5.39	7.09	4.75
O-C _{phot}	ppm	2007	2006	2010	2008
BIC		−70137	−70160	−70049	−70179
$^a \Delta$ BIC		28	25	21	20
GP hyper-parameters					
$\sigma_{\rm white, phot}$	ppm		98±19		100±18
$\tau_{\rm decay, phot}$	day		7.2 ^{+2.3} _{−5.1}		6.1 ^{+3.2} _{−4.7}
$\gamma_{\rm smooth, phot}$			0.34 ^{+0.11} _{−0.06}		0.36 ^{+0.15} _{−0.08}
$P_{\rm phot}$	days		13.0 ^{+3.4} _{−0.7}		13.0 ^{+4.1} _{−0.8}
$A_{\rm phot}$	ppm		407 ⁺⁴⁶ _{−33}		405 ⁺⁴⁸ _{−37}
$\sigma_{\rm white, RV}$	m s ^{−1}		0.52 ^{+0.53} _{−0.37}		0.58 ^{+0.61} _{−0.39}
$\tau_{\rm decay, RV}$	day		370 ⁺⁴²⁰ _{−316}		32 ⁺⁹⁵ _{−18}
$\gamma_{\rm smooth, RV}$			0.91 ^{+0.42} _{−0.50}		0.38 ^{+0.38} _{−0.16}
$P_{\rm RV}$	days		16.2 ^{+0.6} _{−3.9}		18.3 ^{+0.5} _{−4.1}
$A_{\rm RV}$	m s ^{−1}		3.5 ^{+2.6} _{−1.3}		5.0 ^{+1.8} _{−1.1}

Notes.

^(a) The ΔBIC is calculated between the model without any signal (planet or GP) and the model considered in the given column header.

- a water-world with unfixed iron content in the core of the planet leads to an iron-to-silicate ratio $f'_{core}=47\pm30\%$ and a water envelope with a mass fraction of $f_{H_2O}=23\pm12\%$.

All of these scenarios agree on the evidence that, as for other sub-Neptune exoplanets of similar radius and mass, as in particular K2-3 b, a fraction of the planet mass should be in the form of a gaseous envelope. With $T_{eq}=590\pm90$ K TOI-1695 b has an equilibrium temperature just below the water critical point, theoretically allowing water to be liquid if a surface pressure of about

200 atm can be sustained. However, if a significant atmospheric layer is present, the temperature will also certainly increase with pressure. Thus water cannot be in a liquid phase on the surface of this planet, as was shown for GJ 3470 b whose irradiation temperature is similar (Piette & Madhusudhan 2020).




Fig. 12. Mass-radius plot obtained based on the `pyExoRaMa` code (Zeng et al. 2021; Francesco et al. 2022). The red, green and blue solid lines represent the mass-radius relation for respectively pure Fe, silicate, and H_2O core. The yellow/brown-colored z -contours (see Zeng et al. 2021 for details) with black solid lines represent the radius inflation due to gas in an envelope surrounding the planet core corresponding to pure silicate. The comparison planets (black circles) are taken from the *TepCat* database (Southworth 2011), selecting only exoplanets around M and K host-stars. TOI-1695 b is shown as a blue star.

5.2. Atmospheric mass loss or gas-depleted formation?

The worlds with radius $\sim 2R_\oplus$ are reputed to be underabundant around Solar-type stars due to atmospheric loss forming rather $R_p < 1.7R_\oplus$ planets (Kite & Schaefer 2021; Rogers & Owen 2021). Yet, TOI-1695 b, as well as few others, exhibit such characteristics. Because it is located around an M-dwarf, TOI-1695 b has undergone different irradiation conditions than planets around Solar-type stars and followed a different evolution with respect to photoevaporation.

Using the equation 15 from Lecavelier Des Etangs (2007), we measure an order of magnitude for the evaporation rate for TOI-1695 b of $dm/dt \sim 10^{10} \text{ g s}^{-1}$. This evaporation rate indicates that if a H/He envelope is still present today in the atmosphere of this planet (less than 1% of the planet mass), it has to be, in this scenario, the remains of the evaporation of a much more massive envelope. If the gas represented about 10% of the mass of the initial planet, this primitive atmosphere would have had a lifetime shorter than 1.5 Gyr. This is about the minimum age of the system that we inferred from spectroscopic, photometric and activity analysis of TOI-1695 in Section 3.

A lifetime of 10 Gyr is obtained for an envelope with 70% of the mass of the core, almost doubling the initial planet mass. Although possible, in this case the large mass loss rate requires a lot of fine tuning on the envelope mass and the age of the planet to explain our observations. It would make this detection fortuitous and thus unlikely if photoevaporation is responsible for the actual values of the mass and radius of TOI-1695 b.

Therefore, if the age of this system is closer to ~ 1 Gyr, then photoevaporation could be at the origin of a shallow H/He at-




Fig. 13. The R_p - P_{orb} diagram for exoplanets around cool stars ($T_{\text{eff}} < 4000 \text{ K}$) taken from the NASA exoplanet archive. The solid and dash lines represent respectively the Cloutier & Menou (2020) and Martinez et al. (2019) empirical radius valley (see text). The shaded region shows exoplanets that from their gas content could be able to test gas-depleted formation and thermally-driven mass loss scenarios. TOI-1695 b is highlighted as a thick-lined black circle.

mosphere; but if the age of TOI-1695 is rather beyond 5 Gyr, then this planet was more likely formed as is: with a small atmosphere, and survived because it formed far away and migrated inward. In this case the scenario proposed by gas-depleted formation (Lee et al. 2014) is preferred.

This is illustrated in Fig. 13. It compares TOI-1695 b to other similar exoplanets around cool stars ($T_{\text{eff}} < 4000 \text{ K}$) in an R_p - P_{orb} diagram and confront it to rocky to non-rocky empirical transitions consistent with gas-depleted formation (Cloutier & Menou 2020) and thermally-driven atmospheric escape (Martinez et al. 2019). The latter includes photoevaporation (Lopez & Rice 2018), as well as core-powered mass loss (Gupta et al. 2022) mechanisms, each predicting a similar negative slope of the radius valley in R_p - P_{orb} space for low-mass stars. In contrast, the gas-depleted formation scenario can produce larger rocky super-Earths at longer orbital periods, resulting in a positive slope of the rocky/non rocky transition. Exoplanets in the shaded region of Fig. 13 are interesting candidates to test those formation/evolution scenarios.

Determining the rocky/gaseous nature of exoplanets in the region enclosed by those two empirical R_p - P_{orb} relations is therefore of key importance to determine which scenario dominates over the other around M dwarfs. Here, TOI-1695 b as non-rocky tends to favour the gas-depleted formation scenario. However it is located close to the thermally-driven mass loss limit, and given the age uncertainty on the host star, the photoevaporation or core-powered mass loss scenarios cannot be rejected.

5.3. Prospect for atmospheric observations

With a J magnitude of 9.6 for the star TOI-1695, the transits of TOI-1695 b can be observed in a search for atmospheric signatures. To characterize the potential for the detection of atmospheric absorption lines, we can evaluate the ratio of the atmospheric absorption depth to the noise in the transit light curve. Of

course, the noise level will depend on the wavelength, telescope, instrument, etc., and the depth of the atmospheric absorption depends on the species, abundances, oscillator strength and number of the searched lines, etc. Therefore, now only a relative S/N ratio of detection can be calculated for an exoplanet atmosphere to be compared with other planets to be observed for the search of the same signature with the same instrument.

Here, we have made the calculation in the J band, which is currently available from space and ground-based facilities, and which is close to the H₂O band observed in a large number of exoplanets (e.g., Fraine et al. 2014; Benneke et al. 2019; Tsiaras et al. 2019; Mikal-Evans et al. 2021). The atmospheric absorption depth, that is the fraction of the stellar flux that is absorbed by the atmosphere during the transit, is proportional to the area of the absorbing layer, which is given by the scale height of the atmosphere (H) multiplied by $2\pi R_p$, where R_p is the planet radius, and inversely proportional to the stellar disk area (πR_{star}^2). The noise is assumed to be proportional to the square root of the stellar flux given by $F_J \propto 10^{-0.4m_J}$, where m_J is the star J magnitude. The atmospheric scale height is given by $kT/\mu g$, where T is the atmosphere temperature, k the Boltzmann constant, μ the mean molar molecular mass, and g is the planet gravity. Finally, with $g \propto M_p/R_p^2$, where M_p is the mass of the planet, the signal-to-noise ratio is given by $S/N \propto HR_p/R_{star}^2 \sqrt{F_J}$, in agreement with the TSM (transmission spectroscopic metric) as defined by Kempton et al. (2018) (see also, Cointepas et al. 2021).

We calculated the S/N ratio expected for all known exoplanets transiting an M star, and normalized them to the value of 100 for AU Mic b as done in Martioli et al. (2022). We used the catalog of exoplanets published in the Exoplanets Encyclopedia on the 18th July, 2022 (Schneider et al. 2011). For the J magnitudes, we used the tabulated values when available, or calculated theoretical values from the V magnitudes and the stars effective temperatures assuming a black-body spectrum. We considered only the M-type stars of the catalogue; a star is considered to be an M-type star if it is tabulated with this stellar type or if no stellar type is given in the catalogue and its effective temperature is between 2200 K and 3900 K.

The result is shown in Fig. 14 where we plotted the S/N ratio of the atmospheric signatures and the TSM in the J band as a function of the planetary mass for known exoplanets orbiting M type stars with masses between 2 and 10 $M_\oplus$. With an S/N of 4 of TOI-1695 b, in the 2 to 10 Earth mass range, several exoplanets yield a much better S/N. Even if we calculate the same plot for the S/N in the V band, with a stellar type M2V and a V magnitude of 13.0, the situation is barely improved and the conclusion remains the same. Nevertheless, TOI-1695 b has a TSM of 48, a decent S/N ratio to be reached in only 10 h of observation with JWST.

We determined also the emission signature metric (ESM; eq. 4 in Kempton et al. 2018) of TOI-1695 b in the K-band, assuming a day-side temperature of the planet of $1.1 \times T_{eq}$. We found an ESM of 3, implying an S/N=3 for a JWST detection of the planet in the K-band during a secondary eclipse. This is again significantly below the threshold for defining the best "emission" targets, but still challenging in term of detectability.

Thus, even though TOI-1695 b is not the best target in terms of precision, its atmosphere could still be characterized with a limited observation time on large telescopes such as JWST.




Fig. 14. Signal-to-noise ratio (left axis) of the atmospheric signatures and TSM (right axis) in the J band as a function of the planetary mass for exoplanets orbiting M type stars with masses between 2 and 10 Earth mass. The S/N ratios have been normalized to a reference S/N of 100 for AU Mic b as done in Martioli et al. (2022).

6. Summary and conclusion

Thanks to RV follow-up observations with SPIRou at CFHT, we have established the planetary nature of a companion discovered by TESS around the star TOI-1695 with an orbital period of 3.134 days.

We have shown that TOI-1695 b is a sub-Neptune planet, with a mass of $5.5 \pm 1.0 M_\oplus$ and a radius of $2.03 \pm 0.18 R_\oplus$. We found hints of a supplementary weak variability in both photometry ($P \sim 12$ -15 days) and RV ($P \sim 16$ -19 days), but not present in activity indicators (polarimetry, FWHM, bisector span). These activity indicators moreover do not show signs of any clear modulation below a period of ~ 20 days. It implies that TOI-1695 is likely to be a slow rotator, older than 1 Gyr. The origin of the supplementary variability in RV and photometry is still uncertain.

The density of TOI-1695 b is $3.6 \pm 1.1 \text{ g cm}^{-3}$ suggesting the existence of an atmospheric layer on top of the solid bulk of the planet. If made of H/He, the gas envelope would have a low mass fraction $f_{H/He} = 0.28^{+0.46}_{-0.12}\%$. If containing water and varying its core composition, the planet may have a water mass fraction of $23 \pm 12\%$. This confirms that TOI-1695 b stands in the sub-Neptune domain of the radius valley.

The detection prospects of the atmosphere of TOI-1695 b with current and future instruments, such as the JWST and ELTs, with a TSM of 48 are relatively encouraging. Its atmosphere has a S/N ratio of detection lower than several other similar targets, and its characterization is thus not among the easiest. Nevertheless, its radius and orbital period, compared to the rocky/gaseous bounds of the different formation/evolution scenarios of super-Earth and sub-Neptune, make it an interesting case for testing those scenarios.

Acknowledgements. The authors gratefully thank the anonymous referee for his corrections and suggestions that led to improve the content of this article. The authors wish to recognize and acknowledge the very significant cultural role and reverence that the summit of MaunaKea has always had within the indigenous Hawaiian community. We are most fortunate to have the opportunity to conduct observations from this mountain. This paper includes data collected by the TESS mission, which are publicly available from the Mikulski Archive for Space Telescopes (MAST). Funding for the TESS mission is provided by NASA's Science Mission directorate. STScI is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-26555. We acknowledge funding from the French National Research Agency (ANR) under contract number ANR18CE310019 (SPLASH). F.K. acknowledges

support from the Université Paris Sciences et Lettres under the DIM-ACAV program Origines et conditions d'apparition de la vie. E. M. acknowledge funding from the Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG) under the project number APQ-02493-22. J.-F.D. acknowledges funding from the European Research Council (ERC) under the H2020 research & innovation programme (grant agreement #740651 NewWorlds). This work is partly supported by the French National Research Agency in the framework of the Investissements d'Avenir program (ANR-15-IDEX-02), through the funding of the "Origin of Life" project of the Grenoble-Alpes University. J.H.C.M. is supported in the form of a work contract funded by Fundação para a Ciência e Tecnologia (FCT) with the reference DL 57/2016/CP1364/CT0007; and also supported from FCT through national funds and by FEDER-Fundo Europeu de Desenvolvimento Regional through COMPETE2020-Programa Operacional Competitividade e Internacionalização for these grants UIDB/04434/2020 & UIDP/04434/2020, PTDC/FIS-AST/32113/2017 & POCI-01-0145-FEDER-032113, PTDC/FIS-AST/28953/2017 & POCI-01-0145-FEDER-028953, PTDC/FIS-AST/29942/2017. J.H.C.M. also acknowledges the support from FCT - Fundação para a Ciência e a Tecnologia through national funds and by FEDER through COMPETE2020 - Programa Operacional Competitividade e Internacionalização by these grants: UID/FIS/04434/2019; UIDB/04434/2020; UIDP/04434/2020; PTDC/FIS-AST/32113/2017 & POCI-01-0145-FEDER-032113; PTDC/FIS-AST/28953/2017 & POCI-01-0145-FEDER-028953.

References

- Aguichine, A., Mousis, O., Deleuil, M., & Marcq, E. 2021, *ApJ*, 914, 84
- Aigrain, S., Llama, J., Ceillier, T., et al. 2015, *MNRAS*, 450, 3211
- Ambikasaran, S., Foreman-Mackey, D., Greengard, L., Hogg, D. W., & O’Neil, M. 2015, *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 38, 252
- Angus, R., Morton, T., Aigrain, S., Foreman-Mackey, D., & Rajpaul, V. 2018, *MNRAS*, 474, 2094
- Artigau, É., Astudillo-Defru, N., Delfosse, X., et al. 2014, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 9149, Observatory Operations: Strategies, Processes, and Systems V, ed. A. B. Peck, C. R. Benn, & R. L. Seaman, 914905
- Artigau, É., Cadieux, C., Cook, N. J., et al. 2022, arXiv e-prints, arXiv:2207.13524
- Asplund, M., Grevesse, N., Sauval, A. J., & Scott, P. 2009, *ARA&A*, 47, 481
- Basri, G., Walkowicz, L. M., Batalha, N., et al. 2011, *AJ*, 141, 20
- Bellm, E. C., Kulkarni, S. R., Graham, M. J., et al. 2019, *PASP*, 131, 018002
- Benneke, B., Wong, I., Piaulet, C., et al. 2019, *ApJ*, 887, L14
- Blanco-Cuaresma, S. 2019, *MNRAS*, 486, 2075
- Bouchy, F., Pepe, F., & Queloz, D. 2001, *A&A*, 374, 733
- Cadieux, C., Doyon, R., Plotnikov, M., et al. 2022, *The Astronomical Journal*, 164, 96
- Ciardi, D. R., Beichman, C. A., Horch, E. P., & Howell, S. B. 2015, *ApJ*, 805, 16
- Claret, A. 2017, *A&A*, 600, A30
- Claret, A. & Southworth, J. 2022, arXiv e-prints, arXiv:2206.11098
- Cloutier, R. & Menou, K. 2020, *AJ*, 159, 211
- Cointepas, M., Almenara, J. M., Bonfils, X., et al. 2021, *A&A*, 650, A145
- Cristofari, P. I., Donati, J. F., Masseron, T., et al. 2022a, *MNRAS*, 516, 3802
- Cristofari, P. I., Donati, J. F., Masseron, T., et al. 2022b, *MNRAS*, 511, 1893
- Donati, J. F., Kouach, D., Moutou, C., et al. 2020, *MNRAS*, 498, 5684
- Donati, J. F., Semel, M., Carter, B. D., Rees, D. E., & Collier Cameron, A. 1997, *MNRAS*, 291, 658
- Donati, J. F., Yu, L., Moutou, C., et al. 2017, *MNRAS*, 465, 3343
- Engle, S. G. & Guinan, E. F. 2018, *Research Notes of the American Astronomical Society*, 2, 34
- Foreman-Mackey, D., Hogg, D. W., Lang, D., & Goodman, J. 2013, *PASP*, 125, 306
- Fraine, J., Deming, D., Benneke, B., et al. 2014, *Nature*, 513, 526
- Francesco, A., Mario, D., Li, Z., & Alessandro, S. 2022, *Research Notes of the American Astronomical Society*, 6, 28
- Fulton, B. J. & Petigura, E. A. 2018, *AJ*, 156, 264
- Fulton, B. J., Petigura, E. A., Howard, A. W., et al. 2017, *AJ*, 154, 109
- Furlan, E., Ciardi, D. R., Everett, M. E., et al. 2017, *AJ*, 153, 71
- Giacalone, S. & Dressing, C. D. 2020, triceratops: Candidate exoplanet rating tool, *Astrophysics Source Code Library*, record ascl:2002.004
- Giacalone, S., Dressing, C. D., Jensen, E. L. N., et al. 2021, *AJ*, 161, 24
- Ginzburg, S., Schlichting, H. E., & Sari, R. 2018, *MNRAS*, 476, 759
- Goulding, N. T. 2013, PhD thesis, University of Hertfordshire, UK
- Goulding, N. T., Barnes, J. R., Pinfield, D. J., et al. 2012, *MNRAS*, 427, 3358
- Günther, M. N., Pozuelos, F. J., Dittmann, J. A., et al. 2019, *Nature Astronomy*, 3, 1099
- Gupta, A., Nicholson, L., & Schlichting, H. E. 2022, *MNRAS*, 516, 4585
- Gustafsson, B., Edvardsson, B., Eriksson, K., et al. 2008, *A&A*, 486, 951
- Hauschildt, P. H., Allard, F., & Baron, E. 1999, *ApJ*, 512, 377
- Haywood, R. D., Collier Cameron, A., Queloz, D., et al. 2014, *MNRAS*, 443, 2517
- Hobson, M. J., Bouchy, F., Cook, N. J., et al. 2021, *A&A*, 648, A48
- Jenkins, J. M., Twicken, J. D., McCauliff, S., et al. 2016, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 9913, Software and Cyberinfrastructure for Astronomy IV, ed. G. Chiozzi & J. C. Guzman, 99133E
- Kanodia, S. & Wright, J. 2018, *Research Notes of the American Astronomical Society*, 2, 4
- Kempton, E. M. R., Bean, J. L., Louie, D. R., et al. 2018, *PASP*, 130, 114401
- Kite, E. S. & Schaefer, L. 2021, *ApJ*, 909, L22
- Kochanek, C. S., Shappee, B. J., Stanek, K. Z., et al. 2017, *PASP*, 129, 104502
- Kosiarek, M. R., Crossfield, I. J. M., Hardegree-Ullman, K. K., et al. 2019, *AJ*, 157, 97
- Kovács, G., Zucker, S., & Mazeh, T. 2002, *A&A*, 391, 369
- Kreidberg, L. 2015, *PASP*, 127, 1161
- Lecavelier Des Etangs, A. 2007, *A&A*, 461, 1185
- Lee, E. J., Chiang, E., & Ormel, C. W. 2014, *ApJ*, 797, 95
- Lee, E. J. & Connors, N. J. 2021, *ApJ*, 908, 32
- Li, J., Tenenbaum, P., Twicken, J. D., et al. 2019, *PASP*, 131, 024506
- Lightcurve Collaboration, Cardoso, J. V. d. M., Hedges, C., et al. 2018, *Lightkurve: Kepler and TESS time series analysis in Python*, *Astrophysics Source Code Library*
- Lopez, E. D. & Fortney, J. J. 2014, *ApJ*, 792, 1
- Lopez, E. D. & Rice, K. 2018, *MNRAS*, 479, 5303
- Mann, A. W., Dupuy, T., Kraus, A. L., et al. 2019a, *ApJ*, 871, 63
- Mann, A. W., Dupuy, T., Kraus, A. L., et al. 2019b, *ApJ*, 871, 63
- Martinez, C. F., Cunha, K., Ghezzi, L., & Smith, V. V. 2019, *ApJ*, 875, 29
- Martoli, E., Hébrard, G., Fouqué, P., et al. 2022, *A&A*, 660, A86
- Martoli, E., Hébrard, G., Moutou, C., et al. 2020, *A&A*, 641, L1
- Masci, F. J., Laher, R. R., Rusholme, B., et al. 2019, *PASP*, 131, 018003
- McQuillan, A., Aigrain, S., & Roberts, S. 2012, *A&A*, 539, A137
- Mikal-Evans, T., Crossfield, I. J. M., Benneke, B., et al. 2021, *AJ*, 161, 18
- Montegriffo, P., De Angeli, F., Andrae, R., et al. 2022, arXiv e-prints, arXiv:2206.06205
- Moutou, C., Hébrard, E. M., Morin, J., et al. 2017, *MNRAS*, 472, 4563
- Muirhead, P. S., Dressing, C. D., Mann, A. W., et al. 2018, *AJ*, 155, 180
- Owen, J. E. & Wu, Y. 2013, *ApJ*, 775, 105
- Pasegger, V. M., Schweitzer, A., Shulyak, D., et al. 2019, *A&A*, 627, A161
- Peterson, M. S., Benneke, B., David, T. J., et al. 2018, *AJ*, 156, 188
- Piaulet, C., Benneke, B., Rubenzahl, R. A., et al. 2021, *AJ*, 161, 70
- Piette, A. A. & Madhusudhan, N. 2020, *ApJ*, 904, 154
- Plez, B. 2012, *Turbospectrum: Code for spectral synthesis*, *Astrophysics Source Code Library*, record ascl:1205.004
- Polanski, A. S., Crossfield, I. J. M., Burt, J. A., et al. 2021, *AJ*, 162, 238
- Ricker, G. R., Winn, J. N., Vanderspek, R., et al. 2015, *Journal of Astronomical Telescopes, Instruments, and Systems*, 1, 014003
- Rogers, J. G. & Owen, J. E. 2021, *MNRAS*, 503, 1526

- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
- Schneider, J., Dedieu, C., Le Sidaner, P., Savalle, R., & Zolotukhin, I. 2011, *A&A*, 532, A79
- Shappee, B. J., Prieto, J. L., Grupe, D., et al. 2014, *ApJ*, 788, 48
- Shuvalov, V. 2009, *MAPS*, 44, 1095
- Smith, J. C., Stumpe, M. C., Van Cleve, J. E., et al. 2012, *PASP*, 124, 1000
- Smith, V. V., Bizyaev, D., Cunha, K., et al. 2021, *AJ*, 161, 254
- Snedden, C., Bean, J., Ivans, I., Lucatello, S., & Sobeck, J. 2012, MOOG: LTE line analysis and spectrum synthesis, Astrophysics Source Code Library, record ascl:1202.009
- Southworth, J. 2011, *MNRAS*, 417, 2166
- Stassun, K. G., Collins, K. A., & Gaudi, B. S. 2017, *AJ*, 153, 136
- Stassun, K. G., Corsaro, E., Pepper, J. A., & Gaudi, B. S. 2018, *AJ*, 155, 22
- Stassun, K. G., Oelkers, R. J., Paegert, M., et al. 2019, *AJ*, 158, 138
- Stassun, K. G. & Torres, G. 2016, *AJ*, 152, 180
- Stassun, K. G. & Torres, G. 2021, *ApJ*, 907, L33
- Stumpe, M. C., Smith, J. C., Catanzarite, J. H., et al. 2014, *PASP*, 126, 100
- Tsiaras, A., Waldmann, I. P., Tinetti, G., Tennyson, J., & Yurchenko, S. N. 2019, *Nature Astronomy*, 3, 1086
- Twicken, J. D., Catanzarite, J. H., Clarke, B. D., et al. 2018, *PASP*, 130, 064502
- Van Eylen, V., Agentoft, C., Lundkvist, M. S., et al. 2018, *MNRAS*, 479, 4786
- Van Eylen, V., Astudillo-Defru, N., Bonfils, X., et al. 2021, *MNRAS*, 507, 2154
- Wizinowich, P., Acton, D. S., Shelton, C., et al. 2000, *PASP*, 112, 315
- Wright, J. T. & Eastman, J. D. 2014, *PASP*, 126, 838
- Wu, Y. 2019, *ApJ*, 874, 91
- Zeng, L., Jacobsen, S. B., Hyung, E., et al. 2021, *ApJ*, 923, 247
- Zeng, L., Sasselov, D. D., & Jacobsen, S. B. 2016, *ApJ*, 819, 127

-
- ¹ LESIA, Observatoire de Paris, Université PSL, CNRS, Sorbonne Université, Université Paris Cité, 5 place Jules Janssen, 92195 Meudon, France
- ² Sorbonne Université, CNRS, UMR 7095, Institut d’Astrophysique de Paris, 98 bis bd Arago, 75014 Paris, France
- ³ Observatoire de Haute Provence, St Michel l’Observatoire, France
- ⁴ Laboratório Nacional de Astrofísica, Rua Estados Unidos 154, 37504-364, Itajubá - MG, Brazil
- ⁵ Université de Montréal, Département de Physique, IREX, Montréal, QC H3C 3J7, Canada
- ⁶ Institut de Recherche en Astrophysique et Planétologie, Université de Toulouse, CNRS, IRAP/UMR 5277, 14 avenue Edouard Belin, 31400, Toulouse, France
- ⁷ Université Grenoble Alpes, CNRS, IPAG, 38000 Grenoble, France
- ⁸ NASA Exoplanet Science Institute, Caltech/IPAC, Pasadena, CA 91125, USA
- ⁹ Department of Astronomy and Astrophysics, University of California, Santa Cruz, CA 95064, USA
- ¹⁰ Department of Physics and Astronomy, Vanderbilt University, Nashville, TN 37235, USA
- ¹¹ Instituto de Astrofísica e Ciências do Espaço, Universidade do Porto, CAUP, Rua das Estrelas, P-4150-762 Porto, Portugal
- ¹² European Southern Observatory, Alonso de Cordova, Vitacura, Santiago, Chile
- ¹³ Canada-France-Hawaii Telescope, CNRS, 96743 Kamuela, Hawai’i, USA
- ¹⁴ Institut de Recherche sur les Exoplanètes, Département de Physique, Université de Montréal, 1375 Avenue Thérèse-Lavoie-Roux, Montreal, QC H2V 0B3, Canada
- ¹⁵ Aix Marseille Univ, CNRS, CNES, LAM, Marseille, France
- ¹⁶ Department of Physics and Kavli Institute for Astrophysics and Space Research, Massachusetts Institute of Technology, Cambridge, MA 02139, USA
- ¹⁷ Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD, 21218, USA
- ¹⁸ Center for Astrophysics, Harvard & Smithsonian, 60 Garden St, Cambridge, MA 02138, USA
- ¹⁹ MIT Kavli Institute for Astrophysics and Space Research, Massachusetts Institute of Technology, Cambridge, MA 02139, USA
- ²⁰ Earth and Planetary Sciences, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, MA 02139, USA
- ²¹ Department of Aeronautics and Astronautics, MIT, 77 Massachusetts Avenue, Cambridge, MA 02139, USA
- ²² NASA Ames Research Center, Moffett Field, CA 94035, USA
- ²³ SETI Institute, Mountain View, CA 94043, USA
- ²⁴ Department of Astrophysical Sciences, Peyton Hall, 4 Ivy Lane, Princeton, NJ 08544, USA
- ²⁵ Instituto de Astrofísica e Ciências do Espaço, Universidade do Porto, CAUP, Rua das Estrelas, 4150-762, Porto, Portugal
- ²⁶ UVSQ, Université Paris-Saclay, 78047 Guyancourt, France
- ²⁷ Institute of Astronomy and Astrophysics, Academia Sinica, Taipei 10617, Taiwan
- ²⁸ Department of Earth Sciences, University of Hawai’i at Manoa, Honolulu, Hawaii 96822 USA
- ²⁹ Konkoly Observatory, Research Centre for Astronomy and Earth Sciences, Eötvös Loránd Research Network (ELKH), Konkoly-Thege Miklós út 15-17, H-1121 Budapest, Hungary
- ³⁰ CSFK, MTA Centre of Excellence, Budapest, Konkoly Thege Miklós út 15-17, H-1121, Hungary
- ³¹ Max-Planck-Institute for Astronomy, Königstuhl 17, D-69117 Heidelberg, Germany
- ³² ELTE Eötvös Loránd University, Institute of Physics, Pázmány Péter sétány 1/A, H-1117 Budapest, Hungary
- ³³ Universidade Federal do Rio Grande do Norte (UFRN), 59078-970, Natal, RN, Brazil
- ³⁴ Center for Astrophysics | Harvard & Smithsonian, 60 Garden Street, Cambridge, MA 02138, USA

Appendix A: TESS aperture and field-of-view




Fig. A.1. Aperture used to retrieve the photometry of TOI-1695 from the SPOC in sector 18, compared to the background sources identified by Gaia within $200''$ (left), and the background resolved sources on the full-frame image (right). This image is an output from TRICERATOPS (see text).

Appendix B: Spectral modelisation of the star




Fig. B.1. Stellar lines spectral fitting with MARCS model (see Section 3 for explanations), with CO (top), Mn (bottom-left), Na (bottom-middle) and OH (bottom-right) lines. The best fitting model is plotted in green solid line.

Appendix C: SPIRou polarimetry time series




Fig. C.1. Longitudinal magnetic field variations (red points). A Gaussian process best fit with a period of 48 d with a decay time of 500 days and a smoothing factor commonly fixed to 0.6 day is also shown in blue.

Appendix D: Periodograms of RV, FWHM and BIS obtained with the APERO v7 version




Fig. D.1. Same as Fig. 6 but obtained with the APERO v7.

Appendix E: Corner plot of the MCMC posterior distributions for the transit and RV fit




Fig. E.1. MCMC fit posteriors of the SPIRou and TESS data, in the scheme where RVs are drift corrected (DC), zero-point corrected (ZPC) and a GP with $P_{\text{GP}}=14.8$ days is fitted out from the RVs and the lightcurve. The blue solid line spots the median of the posterior probability density function (pdf), and the black dotted lines locate the 16 and 84 percentiles of the posterior pdf.

Appendix F: Observation logs.

Table F.1. Observations log of SPIRou observations of TOI-1695. The † symbol indicates data that were excluded from the drift corrected data reduced with the APERO v6 version. The ★ symbol indicates data removed from the drift corrected and zero point corrected data reduced with the APERO v7 version. This is discussed in Section 2.

Epoch	UT Date	BJD -2,459,000	BERV (km s ⁻¹)	Exp. time (sec)	Airmass	S/N at 1670 nm	H ₂ O rel. abs.	up?	Moon angle (°)	phase	Pol. seq. number
1	2020-12-25T07:09:48	208.794938	-9.5505	1204	1.69	142	4.17	T	62	0.80	1
1	2020-12-25T07:30:20	208.809194	-9.5649	1204	1.71	140	4.36	T	62	0.80	2
1	2020-12-25T07:50:52	208.823449	-9.5787	1204	1.75	145	4.54	T	62	0.80	3
1	2020-12-25T08:11:23	208.837704	-9.5920	1204	1.79	141	4.62	T	62	0.80	4
2	2020-12-30T06:17:47	213.758646	-10.7086	1204	1.66	135	2.10	T	64	-1.00	1
2	2020-12-30T06:38:19	213.772901	-10.7233	1204	1.68	135	2.20	T	64	-1.00	2
2	2020-12-30T06:58:50	213.787157	-10.7376	1204	1.70	137	2.25	T	64	-1.00	3
2	2020-12-30T07:19:22	213.801412	-10.7516	1204	1.73	133	2.40	T	64	-1.00	4
3	2021-09-15T11:39:36	472.980910	15.1838	1204	1.67	142	2.10	F	117	0.77	1
3	2021-09-15T12:00:07	472.995168	15.1704	1204	1.65	145	2.11	F	117	0.77	2
3	2021-09-15T12:20:33	473.009361	15.1570	1204	1.65	145	2.12	F	117	0.77	3
3	2021-09-15T12:40:59	473.023551	15.1436	1204	1.65	147	2.09	F	117	0.77	4
4	2021-09-19T09:19:11	476.883601	14.7815	1204	1.90	148	3.08	T	90	0.97	1
4	2021-09-19T09:39:37	476.897791	14.7706	1204	1.84	152	3.09	T	91	0.97	2
4	2021-09-19T10:00:09	476.912048	14.7589	1204	1.79	150	3.04	T	89	0.97	3
4	2021-09-19T10:20:35	476.926239	14.7468	1204	1.75	152	2.80	T	89	0.97	4
5	2021-09-22T08:51:52	479.864766	14.3903	1204	1.95	155	1.22	T	70	-0.98	1
5	2021-09-22T09:12:23	479.879022	14.3797	1204	1.89	156	1.20	T	70	-0.98	2
5	2021-09-22T09:32:49	479.893215	14.3684	1204	1.83	153	1.12	T	70	-0.98	3
5	2021-09-22T09:53:15	479.907407	14.3565	1204	1.78	153	1.06	T	70	-0.98	4
6	2021-09-23T09:02:08	480.871949	14.2419	1204	1.91	150	4.95	T	65	-0.94	1
6	2021-09-23T09:22:40	480.886206	14.2308	1204	1.84	148	4.74	T	65	-0.94	2
6	2021-09-23T09:43:06	480.900398	14.2190	1204	1.79	146	3.13	T	65	-0.94	3
6	2021-09-23T10:03:32	480.914590	14.2067	1204	1.75	143	4.74	T	64	-0.94	4
7	2021-09-24T08:40:33	481.857005	14.1059	1204	1.97	154	1.61	T	61	-0.88	1
7	2021-09-24T09:00:59	481.871199	14.0953	1204	1.90	150	1.58	T	60	-0.88	2
7	2021-09-24T09:21:25	481.885388	14.0840	1204	1.84	153	1.54	T	60	-0.88	3
7	2021-09-24T09:41:51	481.899581	14.0721	1204	1.79	153	1.46	T	60	-0.88	4
8	2021-10-14T10:23:45	501.929496	10.2267	1204	1.65	152	3.84	T	106	0.73	1
8	2021-10-14T10:44:17	501.943753	10.2116	1204	1.65	149	3.85	T	106	0.73	2
8	2021-10-14T11:04:49	501.958010	10.1966	1204	1.65	148	4.00	T	106	0.73	3
8	2021-10-14T11:25:15	501.972201	10.1818	1204	1.66	153	4.05	F	106	0.73	4
9	2021-10-15T10:03:28	502.915441	10.0123	1204	1.65	156	2.77	T	98	0.73	1
9	2021-10-15T10:24:00	502.929698	9.9972	1204	1.65	155	2.75	T	98	0.82	2
9	2021-10-15T10:44:26	502.943889	9.9821	1204	1.65	156	2.59	T	98	0.82	3
9	2021-10-15T11:04:52	502.958081	9.9672	1204	1.65	152	2.54	T	98	0.82	4
10	2021-10-16T10:56:00	503.951960	9.7415	1204	1.65	130	0.85	T	92	0.89	1
10	2021-10-16T11:16:32	503.966216	9.7265	1204	1.66	145	0.85	T	92	0.89	2
10	2021-10-16T11:37:04	503.980473	9.7118	1204	1.68	141	0.85	T	90	0.89	3
10	2021-10-16T11:57:30	503.994665	9.6976	1204	1.70	134	0.88	T	90	0.89	4
11	2021-10-18T10:20:43	505.927524	9.2950	1204	1.65	137	3.05	T	77	0.98	1
11	2021-10-18T10:41:15	505.941780	9.2798	1204	1.65	136	3.07	T	77	0.98	2
11	2021-10-18T11:01:47	505.956036	9.2647	1204	1.66	142	3.17	T	77	0.98	3
11	2021-10-18T11:22:13	505.970227	9.2499	1204	1.67	143	3.15	T	77	0.98	4
12	2021-10-20T10:23:54	507.929789	8.8095	1204	1.65	145	5.63	T	66	-0.99	1
12	2021-10-20T10:44:20	507.943982	8.7943	1204	1.65	145	5.79	T	66	-0.99	2
12	2021-10-20T11:04:46	507.958173	8.7793	1204	1.66	142	5.82	T	66	-0.99	3
12	2021-10-20T11:25:12	507.972368	8.7646	1204	1.68	139	6.01	T	66	-0.99	4
13	2021-10-21T10:33:20	508.936373	8.5569	1204	1.65	148	3.10	T	61	-0.97	1
13	2021-10-21T10:53:46	508.950565	8.5418	1204	1.66	147	2.98	T	61	-0.97	2
13	2021-10-21T11:14:18	508.964820	8.5268	1204	1.68	144	2.88	T	61	-0.97	3
13	2021-10-21T11:34:44	508.979012	8.5123	1204	1.70	147	3.01	T	61	-0.97	4
14	2021-10-22T10:44:04	509.943857	8.3007	1204	1.66	111	2.15	T	58	-0.93	1
14	2021-10-22T11:04:36	509.958114	8.2856	1204	1.67	111	2.17	T	58	-0.93	2

Table F.1: continued from previous page

Epoch	UT Date	BJD	BERV (km s ⁻¹)	Exp. time (sec)	Airmass	S/N at 1670 nm	H ₂ O rel. abs.	on-sky	Moon (°)	phase	Pol. seq. number
14	2021-10-22T11:25:02	509.972306	8.2710	1204	1.69	110	2.26	T	58	-0.93	3
14	2021-10-22T11:45:28	509.986497	8.2567	1204	1.72	110	2.28	T	58	-0.93	4
15	2021-10-23T10:12:50	510.922184	8.0731	1204	1.65	149	3.08	T	56	-0.93	1
15	2021-10-23T10:33:16	510.936376	8.0579	1204	1.65	145	3.12	T	56	-0.87	2
15	2021-10-23T10:53:47	510.950632	8.0427	1204	1.66	143	3.27	T	56	-0.87	3
15	2021-10-23T11:14:14	510.964823	8.0279	1204	1.68	146	3.27	T	56	-0.87	4
16	2021-10-24T11:20:08	511.968953	7.7703	1204	1.69	120	1.76	T	56	-0.80	1
16	2021-10-24T11:40:34	511.983145	7.7560	1204	1.72	127	1.84	T	56	-0.80	2
16	2021-10-24T12:01:00	511.997337	7.7422	1204	1.76	118	1.84	T	56	-0.80	3
17	2021-10-26T09:50:42	513.906896	7.3218	1204	1.65	147	2.36	T	61	-0.72	1
17	2021-10-26T10:11:14	513.921151	7.3063	1204	1.65	147	2.33	T	61	-0.72	2
17	2021-10-26T10:31:40	513.935343	7.2910	1204	1.66	145	2.45	T	61	-0.63	3
17	2021-10-26T10:52:06	513.949535	7.2759	1204	1.67	139	2.57	T	61	-0.63	4
18	2021-11-13T06:36:44	531.772487	2.4459	1204	1.76	131	1.99	T	88	0.69	1
18	2021-11-13T06:57:10	531.786678	2.4312	1204	1.73	138	2.15	T	88	0.69	2
18	2021-11-13T07:17:42	531.800934	2.4160	1204	1.70	143	2.20	T	88	0.69	3
18	2021-11-13T07:38:08	531.815126	2.4005	1204	1.68	144	2.26	T	88	0.69	4
19	2021-11-17T09:58:59	535.912967	1.1234	1204	1.71	115	5.06	T	62	0.96	1
19	2021-11-17T10:19:31	535.927223	1.1089	1204	1.75	79	5.31	T	62	0.96	2
19	2021-11-17T10:40:03	535.941479	1.0948	1204	1.79	83	5.98	T	62	0.99	3
19	2021-11-17T11:00:29	535.955670	1.0815	1204	1.84	91	6.18	T	62	0.99	4
20	2021-11-20T08:16:08	538.841557	0.3204	1204	1.65	147	1.30	T	56	-0.99	1
20	2021-11-20T08:36:40	538.855808	0.3044	1204	1.65	147	1.29	T	56	-0.99	2
20	2021-11-20T08:57:12	538.870065	0.2886	1204	1.66	149	1.22	T	56	-0.99	3
20	2021-11-20T09:17:38	538.884257	0.2731	1204	1.68	148	1.19	T	56	-0.99	4
21	2021-11-21T08:28:22	539.850042	0.0164	1204	1.65	157	1.07	T	57	-0.96	1
21	2021-11-21T08:48:54	539.864298	0.0006	1204	1.66	157	1.08	T	57	-0.96	2
21	2021-11-21T09:09:25	539.878554	-0.0151	1204	1.67	155	1.10	T	57	-0.96	3
21	2021-11-21T09:29:52	539.892747	-0.0303	1204	1.69	152	1.15	T	57	-0.96	4
22	2021-12-09T07:21:22	557.803364	-5.2062	1204	1.65	119	2.49	T	97	0.32	1
22	2021-12-09T07:41:54	557.817620	-5.2219	1204	1.66	116	2.48	T	97	0.32	2
22	2021-12-09T08:02:26	557.831874	-5.2373	1204	1.68	117	2.51	T	97	0.32	3
22	2021-12-09T08:22:52	557.846067	-5.2524	1204	1.70	119	2.48	T	97	0.32	4
23	2021-12-11T06:42:23	559.776247	-5.7413	1204	1.65	118	2.95	T	82	0.53	1
23	2021-12-11T07:02:54	559.790503	-5.7572	1204	1.65	120	2.92	T	82	0.53	2
23	2021-12-11T07:23:26	559.804759	-5.7729	1204	1.65	128	2.90	T	82	0.53	3
23	2021-12-11T07:43:52	559.818950	-5.7884	1204	1.67	123	2.98	T	80	0.53	4
24	2021-12-12T06:41:13	560.775428	-6.0201	1204	1.65	139	2.53	T	74	0.63	1
24	2021-12-12T07:01:45	560.789685	-6.0359	1204	1.65	139	2.57	T	75	0.63	2
24	2021-12-12T07:22:17	560.803939	-6.0516	1204	1.66	140	2.69	T	75	0.63	3
24	2021-12-12T07:42:43	560.818130	-6.0670	1204	1.67	140	2.77	T	75	0.63	4
25	2021-12-13T07:08:52	561.794605	-6.3189	1204	1.65	113	3.42	T	69	0.72	1
25	2021-12-13T07:29:24	561.808859	-6.3345	1204	1.66	117	3.72	T	69	0.72	2
25	2021-12-13T07:49:55	561.823115	-6.3497	1204	1.68	125	4.17	T	69	0.72	3
25	2021-12-13T08:10:22	561.837306	-6.3646	1204	1.70	120	4.21	T	69	0.72	4
26	2021-12-14T06:15:31	562.757530	-6.5535	1204	1.65	152	3.18	T	64	0.81	1
26	2021-12-14T06:36:02	562.771786	-6.5693	1204	1.65	154	2.98	T	64	0.81	2
26	2021-12-14T06:56:29	562.785977	-6.5849	1204	1.65	154	2.88	T	64	0.81	3
26	2021-12-14T07:16:55	562.800168	-6.6005	1204	1.66	155	2.85	T	64	0.81	4
27	2021-12-15T06:25:47	563.764642	-6.8348	1204	1.65	123	0.67	T	60	0.88	1
27	2021-12-15T06:46:19	563.778898	-6.8506	1204	1.65	119	0.70	T	60	0.88	2
27	2021-12-15T07:06:50	563.793154	-6.8662	1204	1.65	114	0.71	T	60	0.88	3
27	2021-12-15T07:27:17	563.807345	-6.8815	1204	1.67	127	0.72	T	60	0.88	4
28	2021-12-16T07:14:59	564.798782	-7.1435	1204	1.66	140	1.26	T	57	0.93	1
28	2021-12-16T07:35:30	564.813036	-7.1587	1204	1.68	144	1.28	T	57	0.93	2
28	2021-12-16T07:56:02	564.827292	-7.1736	1204	1.70	135	1.34	T	57	0.93	3
28	2021-12-16T08:16:28	564.841483	-7.1880	1204	1.73	138	1.42	T	57	0.93	4
29	2021-12-18T06:47:02	566.779331	-7.6582	1204	1.65	150	3.30	T	56	0.99	1
29	2021-12-18T07:07:34	566.793582	-7.6736	1204	1.66	160	3.32	T	56	0.99	2
29	2021-12-18T07:28:06	566.807838	-7.6887	1204	1.68	157	3.44	T	56	0.99	3
29	2021-12-18T07:48:32	566.822028	-7.7035	1204	1.70	161	3.50	T	56	0.99	4

Table F.1: continued from previous page

Epoch	UT Date	BJD	BERV (km s ⁻¹)	Exp. time (sec)	Airmass	S/N at 1670 nm	H ₂ O rel. abs.	on-sky	Moon (°)	phase	Pol. seq. number
30	2022-01-06T06:29:30	585.766520	-12.2163	1204	1.70	137	2.39	T	92	0.17	1
30	2022-01-06T06:49:57	585.780715	-12.2299	1204	1.72	138	2.69	T	92	0.17	2
30	2022-01-06T07:10:28	585.794967	-12.2431	1204	1.76	107	2.66	T	92	0.17	3
30	2022-01-06T07:30:54	585.809157	-12.2556	1204	1.80	110	2.67	T	92	0.17	4
31	2022-01-08T06:41:25	587.774707	-12.6328	1204	1.72	148	0.96	T	77	0.36	1
31	2022-01-08T07:01:57	587.788963	-12.6458	1204	1.76	151	1.01	T	77	0.36	2
31	2022-01-08T07:22:28	587.803217	-12.6583	1204	1.80	151	1.06	T	77	0.36	3
31	2022-01-08T07:43:00	587.817472	-12.6702	1204	1.85	148	1.13	T	78	0.36	4
32	2022-01-10T06:10:19	589.753025	-13.0040	1204	1.69	152	2.61	T	66	0.56	1
32	2022-01-10T06:30:51	589.767280	-13.0175	1204	1.72	154	2.68	T	66	0.56	2
32	2022-01-10T06:51:22	589.781535	-13.0305	1204	1.75	153	2.69	T	65	0.56	3
32	2022-01-10T07:11:48	589.795727	-13.0429	1204	1.80	154	2.67	T	65	0.56	4
33†	2022-01-12T06:36:43	591.771269	-13.3958	1204	1.74	133	0.99	T	58	0.74	1
33†	2022-01-12T06:57:14	591.785525	-13.4083	1204	1.78	129	1.05	T	58	0.74	2
33†	2022-01-12T07:17:46	591.799780	-13.4202	1204	1.83	133	1.12	T	58	0.74	3
33†	2022-01-12T07:38:12	591.813970	-13.4314	1204	1.89	139	1.15	T	58	0.74	4
34	2022-01-13T05:02:35	592.705859	-13.5142	1204	1.65	159	1.20	T	56	0.82	1
34	2022-01-13T05:23:06	592.720111	-13.5284	1204	1.66	154	1.26	T	56	0.82	2
34	2022-01-13T05:43:38	592.734367	-13.5423	1204	1.68	157	1.33	T	56	0.82	3
34	2022-01-13T06:04:04	592.748557	-13.5559	1204	1.70	158	1.34	T	56	0.82	4
35	2022-01-15T06:31:35	594.767572	-13.9225	1204	1.75	146	2.28	T	58	0.94	1
35	2022-01-15T06:52:07	594.781827	-13.9346	1204	1.80	136	2.24	T	58	0.94	2
35	2022-01-15T07:12:33	594.796017	-13.9461	1204	1.85	152	2.27	T	58	0.94	3
35	2022-01-15T07:33:05	594.810273	-13.9569	1204	1.91	159	2.30	T	58	0.94	4
36	2022-01-18T05:51:37	597.739680	-14.3897	1204	1.71	151	1.27	T	71	-1.00	1
36	2022-01-18T06:12:09	597.753935	-14.4024	1204	1.74	151	1.30	T	71	-1.00	2
36	2022-01-18T06:32:35	597.768126	-14.4146	1204	1.78	152	1.35	T	71	-1.00	3
36	2022-01-18T06:53:01	597.782317	-14.4261	1204	1.83	152	1.40	T	71	-1.00	4
37	2022-01-19T06:18:59	598.758635	-14.5620	1204	1.76	145	1.30	T	77	-0.98	1
37	2022-01-19T06:39:31	598.772889	-14.5738	1204	1.80	149	1.37	T	77	-0.98	2
37	2022-01-19T07:00:03	598.787145	-14.5849	1204	1.86	153	1.42	T	77	-0.98	3
37	2022-01-19T07:20:29	598.801335	-14.5953	1204	1.92	153	1.45	T	77	-0.98	4
38	2022-01-20T05:59:54	599.745337	-14.7018	1204	1.73	152	2.81	F	84	-0.94	1
38	2022-01-20T06:20:26	599.759591	-14.7140	1204	1.77	149	3.02	F	84	-0.94	2
38	2022-01-20T06:40:58	599.773850	-14.7256	1204	1.82	152	3.20	T	84	-0.94	3
38	2022-01-20T07:01:24	599.788038	-14.7364	1204	1.87	151	3.39	T	84	-0.94	4
39	2022-01-21T05:56:31	600.742939	-14.8467	1204	1.73	144	3.24	F	91	-0.88	1
39	2022-01-21T06:16:58	600.757130	-14.8588	1204	1.77	142	3.33	F	91	-0.88	2
39	2022-01-21T06:37:24	600.771321	-14.8703	1204	1.82	147	3.49	F	91	-0.88	3
39	2022-01-21T06:57:50	600.785511	-14.8811	1204	1.87	146	3.59	F	91	-0.88	4
40	2022-01-22T05:55:23	601.742089	-14.9888	1204	1.74	119	1.64	F	98	-0.81	1
40	2022-01-22T06:15:54	601.756344	-15.0008	1204	1.78	123	1.77	F	98	-0.81	2
40	2022-01-22T06:36:26	601.770599	-15.0121	1204	1.82	118	1.85	F	98	-0.81	3
40	2022-01-22T06:56:52	601.784791	-15.0228	1204	1.88	104	1.92	F	98	-0.81	4
41	2022-01-23T06:03:58	602.748012	-15.1322	1204	1.76	151	2.34	F	105	-0.72	1
41	2022-01-23T06:24:30	602.762267	-15.1438	1204	1.80	146	2.28	F	105	-0.72	2
41	2022-01-23T06:45:02	602.776521	-15.1547	1204	1.86	144	2.25	F	105	-0.72	3
41	2022-01-23T07:05:28	602.790713	-15.1648	1204	1.92	147	2.33	F	105	-0.72	4
42	2022-01-24T06:19:05	603.758450	-15.2746	1204	1.80	157	2.52	F	112	-0.62	1
42	2022-01-24T06:39:37	603.772704	-15.2855	1204	1.85	156	2.70	F	111	-0.62	2
42	2022-01-24T07:00:08	603.786959	-15.2956	1204	1.92	156	2.90	F	111	-0.62	3
42	2022-01-24T07:20:34	603.801150	-15.3049	1204	1.99	155	3.20	F	111	-0.62	4
43	2022-01-25T06:29:43	604.765791	-15.4096	1204	1.84	124	2.11	F	116	-0.51	1
43	2022-01-25T06:50:15	604.780045	-15.4199	1204	1.90	124	2.18	F	115	-0.51	2
43	2022-01-25T07:10:47	604.794300	-15.4294	1204	1.97	129	2.23	F	117	-0.51	3
43	2022-01-25T07:31:13	604.808491	-15.4380	1204	2.05	124	2.32	F	117	-0.51	4
44†★	2022-01-26T06:37:52	605.774581	-15.5406	652	1.90	75	4.28	F	121	-0.40	1
45	2022-01-27T05:39:40	606.730932	-15.6273	1204	1.74	128	2.54	F	122	-0.29	1
45	2022-01-27T06:00:12	606.745188	-15.6388	1204	1.78	124	2.42	F	122	-0.29	2
45	2022-01-27T06:20:44	606.759443	-15.6497	1204	1.83	123	2.38	F	122	-0.29	3
45	2022-01-27T06:41:10	606.773634	-15.6599	1204	1.89	122	2.42	F	122	-0.29	4

Table F.1: continued from previous page

Epoch	UT Date	BJD	BERV (km s ⁻¹)	Exp. time (sec)	Airmass	S/N at 1670 nm	H ₂ O rel. abs.	on-sky	Moon (°)	phase	Pol. seq. number
46	2022-01-28T05:44:06	607.733950	-15.7455	1204	1.76	143	0.51	F	123	-0.19	1
46	2022-01-28T06:04:37	607.748206	-15.7567	1204	1.80	116	0.50	F	123	-0.19	2
46	2022-01-28T06:25:09	607.762461	-15.7673	1204	1.86	118	0.52	F	123	-0.19	3
46	2022-01-28T06:45:35	607.776651	-15.7771	1204	1.92	107	0.55	F	123	-0.19	4

Appendix G: RV and polarimetry data tables

Table G.1. SPIRou LBL RV observations of TOI-1695 obtained with the APERO v6 version. The FWHM and BIS proxies are derived from respectively the second and third derivatives of LBL profiles (see text for explanation).

BJD -2,459,000	v_r (km s ⁻¹)	σ_{v_r} (km s ⁻¹)	FWHM (km s ⁻¹)	eFWHM (km s ⁻¹)	BIS (km s ⁻¹)	eBIS (km s ⁻¹)
208.79493816	-59.9476	0.0066	5.527	0.014	-0.053	0.027
208.80919391	-59.9474	0.0064	5.526	0.014	-0.036	0.027
208.82344915	-59.9404	0.0063	5.524	0.014	-0.029	0.026
208.83770450	-59.9471	0.0064	5.529	0.014	-0.041	0.027
213.75864561	-59.9192	0.0060	5.533	0.013	-0.018	0.025
213.77290070	-59.9255	0.0061	5.512	0.013	-0.030	0.025
213.78715749	-59.9290	0.0060	5.532	0.013	-0.024	0.025
213.80141198	-59.9199	0.0063	5.533	0.013	-0.022	0.026
472.98091030	-59.9349	0.0059	5.485	0.013	-0.014	0.024
472.99516782	-59.9289	0.0057	5.486	0.012	-0.005	0.024
473.00936133	-59.9328	0.0057	5.478	0.012	0.007	0.024
473.02355075	-59.9301	0.0057	5.475	0.012	-0.007	0.024
476.88360128	-59.9372	0.0063	5.474	0.014	0.018	0.026
476.89779088	-59.9340	0.0061	5.462	0.013	-0.011	0.025
476.91204768	-59.9288	0.0062	5.456	0.013	-0.029	0.025
476.92623878	-59.9308	0.0062	5.456	0.013	-0.003	0.025
479.86476628	-59.9310	0.0058	5.457	0.013	-0.012	0.024
479.87902237	-59.9284	0.0056	5.453	0.012	-0.011	0.024
479.89321505	-59.9282	0.0057	5.457	0.012	-0.015	0.023
479.90740723	-59.9338	0.0057	5.465	0.012	-0.020	0.023
480.87194906	-59.9313	0.0061	5.475	0.013	-0.031	0.025
480.88620644	-59.9303	0.0061	5.465	0.013	-0.049	0.026
480.90039831	-59.9250	0.0060	5.470	0.013	-0.035	0.025
480.91458968	-59.9281	0.0060	5.477	0.013	-0.012	0.025
481.85700461	-59.9358	0.0059	5.457	0.013	-0.018	0.025
481.87119917	-59.9356	0.0058	5.443	0.013	-0.032	0.024
481.88538804	-59.9370	0.0058	5.425	0.012	-0.011	0.024
481.89958051	-59.9333	0.0058	5.433	0.012	-0.024	0.024
501.92949642	-59.9339	0.0060	5.478	0.013	-0.026	0.025
501.94375341	-59.9382	0.0059	5.485	0.013	-0.022	0.025
501.95800989	-59.9363	0.0058	5.487	0.013	-0.008	0.024
501.97220128	-59.9305	0.0058	5.480	0.013	0.003	0.024
502.91544136	-59.9373	0.0059	5.476	0.013	-0.039	0.025
502.92969804	-59.9416	0.0059	5.474	0.013	-0.009	0.024
502.94388941	-59.9353	0.0059	5.477	0.013	-0.042	0.025
502.95808148	-59.9427	0.0060	5.490	0.013	-0.006	0.025
503.95195991	-59.9548	0.0061	5.492	0.013	-0.020	0.025
503.96621637	-59.9445	0.0056	5.502	0.012	-0.029	0.023
503.98047314	-59.9413	0.0057	5.487	0.012	-0.013	0.023
503.99466510	-59.9429	0.0058	5.498	0.013	-0.015	0.024
505.92752409	-59.9379	0.0062	5.508	0.013	-0.029	0.026
505.94177953	-59.9400	0.0061	5.498	0.013	-0.025	0.025
505.95603647	-59.9418	0.0061	5.473	0.013	-0.027	0.025
505.97022731	-59.9358	0.0060	5.514	0.013	-0.007	0.025
507.92978917	-59.9286	0.0064	5.469	0.014	-0.026	0.027
507.94398159	-59.9208	0.0063	5.472	0.014	-0.019	0.026

Table G.1: continued from previous page

BJD -2,459,000	v_r (km s ⁻¹)	σ_{v_r} (km s ⁻¹)	FWHM (km s ⁻¹)	eFWHM (km s ⁻¹)	BIS (km s ⁻¹)	eBIS (km s ⁻¹)
507.95817331	-59.9222	0.0063	5.471	0.014	-0.022	0.026
507.97236802	-59.9235	0.0064	5.469	0.014	-0.017	0.026
508.93637316	-59.9311	0.0060	5.495	0.013	-0.017	0.025
508.95056467	-59.9371	0.0059	5.496	0.013	-0.016	0.025
508.96482027	-59.9373	0.0060	5.499	0.013	-0.022	0.025
508.97901207	-59.9280	0.0060	5.519	0.013	-0.013	0.025
509.94385745	-59.9250	0.0073	5.490	0.016	0.000	0.030
509.95811414	-59.9231	0.0072	5.489	0.015	-0.036	0.030
509.97230573	-59.9348	0.0072	5.496	0.016	-0.034	0.030
509.98649743	-59.9347	0.0072	5.526	0.016	-0.013	0.030
510.92218401	-59.9222	0.0062	5.490	0.014	-0.023	0.026
510.93637590	-59.9131	0.0062	5.518	0.014	-0.017	0.026
510.95063248	-59.9140	0.0063	5.500	0.014	-0.029	0.027
510.96482346	-59.9139	0.0061	5.487	0.014	-0.020	0.026
511.96895286	-59.9232	0.0067	5.531	0.014	-0.021	0.027
511.98314483	-59.9198	0.0064	5.518	0.013	-0.027	0.026
511.99733669	-59.9306	0.0067	5.518	0.014	-0.016	0.027
513.90689610	-59.9232	0.0057	5.491	0.012	-0.026	0.024
513.92115105	-59.9252	0.0056	5.496	0.012	-0.015	0.023
513.93534260	-59.9270	0.0057	5.502	0.012	-0.039	0.024
513.94953504	-59.9273	0.0059	5.495	0.013	-0.040	0.025
531.77248693	-59.9328	0.0069	5.526	0.015	-0.034	0.029
531.78667804	-59.9338	0.0063	5.496	0.014	-0.031	0.026
531.80093355	-59.9376	0.0061	5.510	0.013	-0.062	0.025
531.81512557	-59.9268	0.0061	5.518	0.013	-0.048	0.025
535.91296729	-59.9410	0.0077	5.490	0.016	-0.036	0.031
535.92722294	-59.9531	0.0112	5.538	0.024	-0.070	0.045
535.94147939	-59.9292	0.0106	5.537	0.023	-0.101	0.044
535.95567034	-59.9257	0.0097	5.554	0.021	-0.073	0.040
538.84155732	-59.9325	0.0056	5.485	0.012	-0.038	0.024
538.85580844	-59.9317	0.0057	5.501	0.012	-0.050	0.024
538.87006465	-59.9355	0.0057	5.501	0.012	-0.036	0.024
538.88425656	-59.9297	0.0056	5.496	0.012	-0.027	0.024
539.85004231	-59.9332	0.0056	5.489	0.012	-0.040	0.024
539.86429831	-59.9276	0.0056	5.476	0.012	-0.042	0.023
539.87855411	-59.9345	0.0056	5.501	0.012	-0.048	0.023
539.89274681	-59.9326	0.0057	5.473	0.012	-0.045	0.023
557.80336435	-59.9221	0.0072	5.554	0.016	-0.026	0.030
557.81761950	-59.9161	0.0074	5.555	0.016	-0.035	0.030
557.83187445	-59.9272	0.0073	5.553	0.016	-0.028	0.030
557.84606670	-59.9196	0.0071	5.541	0.015	-0.040	0.029
559.77624676	-59.9350	0.0077	5.541	0.016	-0.060	0.031
559.79050318	-59.9408	0.0074	5.561	0.016	-0.036	0.030
559.80475871	-59.9346	0.0068	5.564	0.015	-0.054	0.028
559.81894984	-59.9353	0.0069	5.564	0.015	-0.039	0.029
560.77542835	-59.9428	0.0063	5.539	0.014	-0.033	0.026
560.78968516	-59.9381	0.0062	5.520	0.013	-0.013	0.025
560.80393897	-59.9466	0.0061	5.533	0.013	-0.032	0.025
560.81812999	-59.9446	0.0060	5.516	0.013	-0.034	0.025
561.79460490	-59.9219	0.0078	5.555	0.016	-0.032	0.031
561.80885940	-59.9272	0.0075	5.585	0.016	-0.040	0.031
561.82311469	-59.9326	0.0073	5.569	0.016	-0.039	0.030
561.83730619	-59.9371	0.0074	5.555	0.016	-0.028	0.031
562.75753023	-59.9328	0.0060	5.527	0.013	-0.027	0.025
562.77178552	-59.9345	0.0058	5.518	0.013	-0.030	0.025
562.78597711	-59.9320	0.0058	5.512	0.013	-0.041	0.024
562.80016759	-59.9267	0.0058	5.511	0.013	-0.027	0.024
563.76464181	-59.9453	0.0066	5.546	0.014	-0.005	0.026
563.77889829	-59.9400	0.0067	5.536	0.014	-0.050	0.027
563.79315376	-59.9405	0.0067	5.531	0.014	-0.023	0.027
563.80734474	-59.9301	0.0061	5.527	0.013	-0.049	0.025

Table G.1: continued from previous page

BJD -2,459,000	v_r (km s ⁻¹)	σ_{v_r} (km s ⁻¹)	FWHM (km s ⁻¹)	eFWHM (km s ⁻¹)	BIS (km s ⁻¹)	eBIS (km s ⁻¹)
564.79878234	-59.9410	0.0061	5.565	0.013	-0.035	0.025
564.81303560	-59.9329	0.0060	5.521	0.013	-0.018	0.024
564.82729206	-59.9388	0.0061	5.534	0.013	-0.035	0.025
564.84148262	-59.9299	0.0061	5.542	0.013	-0.049	0.025
566.77933148	-59.9446	0.0063	5.522	0.014	-0.055	0.026
566.79358211	-59.9428	0.0061	5.519	0.013	-0.032	0.025
566.80783755	-59.9503	0.0062	5.536	0.013	-0.024	0.026
566.82202828	-59.9385	0.0061	5.540	0.013	-0.046	0.025
585.76652048	-59.9380	0.0064	5.551	0.014	-0.013	0.027
585.78071470	-59.9383	0.0064	5.539	0.014	-0.010	0.026
585.79496722	-59.9291	0.0079	5.575	0.017	-0.037	0.033
585.80915704	-59.9367	0.0075	5.578	0.016	-0.035	0.032
587.77470698	-59.9397	0.0056	5.512	0.012	-0.011	0.023
587.78896268	-59.9365	0.0056	5.478	0.012	-0.007	0.023
587.80321708	-59.9375	0.0056	5.510	0.012	-0.012	0.023
587.81747208	-59.9402	0.0055	5.499	0.012	-0.000	0.023
589.75302522	-59.9403	0.0062	5.508	0.013	-0.007	0.026
589.76728010	-59.9335	0.0062	5.490	0.013	-0.039	0.025
589.78153548	-59.9306	0.0062	5.519	0.013	-0.026	0.025
589.79572667	-59.9318	0.0061	5.489	0.013	-0.007	0.025
592.70585888	-59.9231	0.0061	5.495	0.013	-0.009	0.025
592.72011124	-59.9234	0.0061	5.505	0.013	-0.022	0.025
592.73436680	-59.9242	0.0060	5.519	0.013	-0.035	0.025
592.74855726	-59.9241	0.0060	5.522	0.013	-0.021	0.024
594.76757159	-59.9434	0.0066	5.536	0.014	-0.018	0.027
594.78182683	-59.9326	0.0069	5.530	0.015	-0.035	0.028
594.79601747	-59.9382	0.0063	5.512	0.014	-0.014	0.026
594.81027260	-59.9363	0.0060	5.521	0.013	-0.031	0.025
597.73968050	-59.9398	0.0057	5.508	0.012	-0.001	0.023
597.75393471	-59.9394	0.0057	5.503	0.012	-0.001	0.023
597.76812613	-59.9385	0.0056	5.519	0.012	-0.018	0.023
597.78231695	-59.9362	0.0056	5.506	0.012	-0.019	0.023
598.75863479	-59.9294	0.0062	5.511	0.013	-0.033	0.025
598.77288920	-59.9320	0.0060	5.521	0.013	-0.019	0.025
598.78714471	-59.9302	0.0059	5.520	0.013	-0.016	0.025
598.80133531	-59.9404	0.0060	5.517	0.013	-0.026	0.025
599.74533714	-59.9339	0.0064	5.475	0.014	-0.027	0.026
599.75959124	-59.9358	0.0061	5.505	0.013	0.003	0.025
599.77384994	-59.9391	0.0060	5.506	0.013	-0.006	0.025
599.78803765	-59.9324	0.0060	5.508	0.013	-0.036	0.024
600.74293878	-59.9394	0.0063	5.538	0.014	-0.025	0.026
600.75712977	-59.9377	0.0061	5.523	0.013	-0.039	0.025
600.77132087	-59.9407	0.0061	5.536	0.013	-0.016	0.025
600.78551106	-59.9409	0.0062	5.526	0.013	-0.015	0.026
601.74208946	-59.9335	0.0076	5.549	0.016	-0.040	0.030
601.75634424	-59.9354	0.0071	5.541	0.015	-0.027	0.029
601.77059913	-59.9397	0.0073	5.561	0.015	-0.008	0.029
601.78479052	-59.9290	0.0083	5.598	0.018	-0.004	0.034
602.74801173	-59.9299	0.0060	5.518	0.013	-0.015	0.025
602.76226731	-59.9290	0.0060	5.533	0.013	-0.007	0.025
602.77652129	-59.9253	0.0060	5.495	0.013	-0.017	0.025
602.79071298	-59.9300	0.0058	5.500	0.013	0.000	0.024
603.75844953	-59.9340	0.0063	5.482	0.014	-0.018	0.026
603.77270410	-59.9382	0.0061	5.485	0.013	-0.026	0.025
603.78695928	-59.9429	0.0061	5.484	0.013	-0.002	0.025
603.80115015	-59.9489	0.0061	5.510	0.013	-0.025	0.025
604.76579074	-59.9210	0.0071	5.562	0.015	-0.022	0.029
604.78004541	-59.9281	0.0069	5.557	0.015	-0.029	0.028
604.79430017	-59.9295	0.0064	5.539	0.014	-0.025	0.027
604.80849134	-59.9340	0.0067	5.559	0.015	-0.037	0.028
606.73093229	-59.9328	0.0070	5.566	0.015	-0.007	0.029

Table G.1: continued from previous page

BJD -2,459,000	v_r (km s ⁻¹)	σ_{v_r} (km s ⁻¹)	FWHM (km s ⁻¹)	eFWHM (km s ⁻¹)	BIS (km s ⁻¹)	eBIS (km s ⁻¹)
606.74518775	-59.9433	0.0071	5.549	0.015	-0.025	0.029
606.75944270	-59.9392	0.0071	5.557	0.015	-0.032	0.029
606.77363376	-59.9428	0.0069	5.539	0.015	-0.026	0.029
607.73395032	-59.9287	0.0062	5.514	0.013	-0.039	0.025
607.74820557	-59.9363	0.0074	5.545	0.016	-0.034	0.030
607.76246093	-59.9318	0.0070	5.533	0.015	-0.021	0.028
607.77665138	-59.9256	0.0078	5.564	0.016	-0.024	0.030

Table G.2. SPIRou LBL RV observations of TOI-1695 obtained with the APERO v7 version.

BJD -2,459,000	v_r (km s ⁻¹)	σ_{v_r} (km s ⁻¹)	FWHM (km s ⁻¹)	FWHM error (km s ⁻¹)	BIS (km s ⁻¹)	BIS error (km s ⁻¹)
208.79493816	-59.4531	0.0084	5.8675	0.018	-0.002	0.033
208.80919391	-59.4712	0.0064	5.7880	0.014	-0.011	0.027
208.82344915	-59.4654	0.0063	5.7882	0.014	-0.004	0.027
208.83770450	-59.4662	0.0065	5.7769	0.014	-0.016	0.028
213.75864561	-59.4454	0.0062	5.7935	0.013	0.001	0.026
213.77290070	-59.4544	0.0061	5.7817	0.013	0.010	0.026
213.78715749	-59.4551	0.0061	5.7930	0.013	-0.005	0.026
213.80141198	-59.4483	0.0063	5.7954	0.014	0.004	0.027
472.98091030	-59.4657	0.0057	5.7911	0.012	-0.007	0.024
472.99516782	-59.4577	0.0056	5.7730	0.012	0.006	0.024
473.00936133	-59.4672	0.0056	5.7632	0.012	0.018	0.024
473.02355075	-59.4615	0.0056	5.7823	0.012	-0.004	0.023
476.88360128	-59.4672	0.0062	5.7740	0.014	0.018	0.027
476.89779088	-59.4646	0.0061	5.7706	0.013	-0.005	0.026
476.91204768	-59.4671	0.0060	5.7695	0.013	-0.006	0.026
476.92623878	-59.4618	0.0060	5.7687	0.013	-0.007	0.026
479.86476628	-59.4563	0.0057	5.7578	0.012	-0.008	0.024
479.87902237	-59.4511	0.0057	5.7536	0.012	-0.009	0.024
479.89321505	-59.4542	0.0056	5.7515	0.012	-0.005	0.024
479.90740723	-59.4604	0.0056	5.7654	0.012	-0.003	0.024
480.87194906	-59.4620	0.0058	5.7839	0.013	-0.009	0.025
480.88620644	-59.4627	0.0058	5.7716	0.013	-0.029	0.025
480.90039831	-59.4544	0.0058	5.7647	0.013	-0.028	0.025
480.91458968	-59.4555	0.0059	5.7744	0.013	0.008	0.025
481.85700461	-59.4616	0.0058	5.7705	0.013	-0.012	0.025
481.87119917	-59.4676	0.0057	5.7683	0.012	-0.016	0.024
481.88538804	-59.4636	0.0057	5.7367	0.012	-0.010	0.024
481.89958051	-59.4597	0.0056	5.7474	0.012	-0.020	0.024
501.92949642	-59.4617	0.0059	5.7822	0.013	-0.018	0.025
501.94375341	-59.4641	0.0059	5.7733	0.013	-0.015	0.025
501.95800989	-59.4665	0.0059	5.7847	0.013	0.007	0.025
501.97220128	-59.4564	0.0059	5.7808	0.013	0.003	0.025
502.91544136	-59.4598	0.0061	5.7711	0.013	-0.018	0.026
502.92969804	-59.4637	0.0060	5.7617	0.013	0.003	0.026
502.94388941	-59.4621	0.0060	5.7726	0.013	-0.012	0.026
502.95808148	-59.4689	0.0060	5.7901	0.013	0.005	0.026
503.95195991	-59.4779	0.0060	5.7833	0.013	-0.009	0.025
503.96621637	-59.4718	0.0055	5.7844	0.012	-0.015	0.023
503.98047314	-59.4717	0.0055	5.7966	0.012	-0.013	0.023
503.99466510	-59.4689	0.0057	5.7794	0.012	-0.003	0.024
505.92752409	-59.4635	0.0060	5.7910	0.013	-0.026	0.025
505.94177953	-59.4672	0.0061	5.7887	0.013	-0.022	0.026
505.95603647	-59.4689	0.0059	5.7718	0.013	-0.009	0.025
505.97022731	-59.4604	0.0058	5.7931	0.013	-0.000	0.025
507.92978917	-59.4545	0.0062	5.7577	0.014	-0.007	0.027
507.94398159	-59.4490	0.0061	5.7651	0.013	-0.006	0.026

Table G.2: continued from previous page

BJD -2,459,000	v_r (km s ⁻¹)	σ_{v_r} (km s ⁻¹)	FWHM (km s ⁻¹)	FWHM error (km s ⁻¹)	BIS (km s ⁻¹)	BIS error (km s ⁻¹)
507.95817331	-59.4504	0.0061	5.7491	0.013	-0.013	0.026
507.97236802	-59.4451	0.0062	5.7546	0.013	-0.013	0.026
508.93637316	-59.4568	0.0059	5.7890	0.013	-0.008	0.025
508.95056467	-59.4624	0.0059	5.7848	0.013	-0.015	0.025
508.96482027	-59.4631	0.0060	5.7795	0.013	-0.021	0.025
508.97901207	-59.4581	0.0059	5.7913	0.013	-0.021	0.025
509.94385745	-59.4552	0.0070	5.7879	0.015	-0.009	0.030
509.95811414	-59.4496	0.0070	5.7608	0.015	-0.027	0.030
509.97230573	-59.4670	0.0070	5.7817	0.015	-0.019	0.030
509.98649743	-59.4686	0.0070	5.8111	0.015	-0.013	0.030
510.92218401	-59.4565	0.0058	5.7855	0.013	-0.022	0.025
510.93637590	-59.4478	0.0059	5.7871	0.013	-0.015	0.025
510.95063248	-59.4534	0.0060	5.7824	0.013	-0.028	0.025
510.96482346	-59.4489	0.0059	5.7599	0.013	-0.021	0.025
511.96895286	-59.4491	0.0065	5.8208	0.014	-0.014	0.028
511.98314483	-59.4547	0.0062	5.8100	0.013	-0.028	0.026
511.99733669	-59.4650	0.0066	5.8064	0.014	-0.018	0.028
513.90689610	-59.4572	0.0056	5.7917	0.012	-0.034	0.024
513.92115105	-59.4618	0.0055	5.7801	0.012	-0.025	0.023
513.93534260	-59.4638	0.0056	5.7952	0.012	-0.033	0.024
513.94953504	-59.4601	0.0057	5.7967	0.012	-0.033	0.024
531.77248693	-59.4610	0.0069	5.8188	0.015	-0.000	0.029
531.78667804	-59.4636	0.0065	5.7800	0.014	-0.015	0.027
531.80093355	-59.4657	0.0061	5.7979	0.013	-0.027	0.025
531.81512557	-59.4576	0.0061	5.7894	0.013	-0.010	0.025
535.91296729	-59.4633	0.0073	5.7752	0.016	-0.006	0.031
535.92722294	-59.4559	0.0103	5.8492	0.023	-0.040	0.044
535.94147939	-59.4496	0.0099	5.8508	0.022	-0.044	0.043
535.95567034	-59.4333	0.0090	5.8318	0.020	-0.048	0.039
538.84155732	-59.4670	0.0056	5.7769	0.012	-0.027	0.024
538.85580844	-59.4628	0.0057	5.7716	0.012	-0.033	0.024
538.87006465	-59.4751	0.0057	5.7695	0.012	-0.034	0.024
538.88425656	-59.4688	0.0057	5.7742	0.012	-0.017	0.024
539.85004231	-59.4689	0.0056	5.7647	0.012	-0.030	0.023
539.86429831	-59.4651	0.0055	5.7558	0.012	-0.000	0.023
539.87855411	-59.4692	0.0055	5.7668	0.012	-0.021	0.023
539.89274681	-59.4692	0.0056	5.7679	0.012	-0.020	0.024
557.80336435	-59.4724	0.0070	5.8099	0.015	0.011	0.030
557.81761950	-59.4710	0.0071	5.8210	0.015	0.026	0.030
557.83187445	-59.4715	0.0070	5.8075	0.015	0.021	0.030
557.84606670	-59.4715	0.0069	5.8090	0.015	0.018	0.029
559.77624676	-59.4681	0.0073	5.7912	0.016	-0.006	0.031
559.79050318	-59.4750	0.0071	5.8125	0.015	0.005	0.030
559.80475871	-59.4731	0.0066	5.8147	0.014	-0.006	0.028
559.81894984	-59.4692	0.0069	5.8231	0.015	-0.004	0.029
560.77542835	-59.4681	0.0063	5.7951	0.014	-0.010	0.027
560.78968516	-59.4721	0.0060	5.7848	0.013	0.007	0.025
560.80393897	-59.4731	0.0059	5.8008	0.013	-0.007	0.025
560.81812999	-59.4709	0.0058	5.7846	0.013	-0.013	0.025
561.79460490	-59.4529	0.0075	5.8266	0.017	-0.008	0.032
561.80885940	-59.4580	0.0074	5.8299	0.016	-0.009	0.032
561.82311469	-59.4722	0.0072	5.8262	0.016	-0.014	0.031
561.83730619	-59.4668	0.0071	5.8188	0.016	-0.000	0.030
562.75753023	-59.4580	0.0060	5.7945	0.013	-0.023	0.025
562.77178552	-59.4651	0.0059	5.8029	0.013	-0.007	0.025
562.78597711	-59.4584	0.0058	5.7839	0.013	-0.031	0.025
562.80016759	-59.4530	0.0057	5.7840	0.013	-0.010	0.024
563.76464181	-59.4655	0.0063	5.8150	0.014	0.009	0.026
563.77889829	-59.4657	0.0066	5.7974	0.014	-0.016	0.027
563.79315376	-59.4632	0.0066	5.7884	0.014	-0.009	0.027
563.80734474	-59.4589	0.0061	5.7839	0.013	-0.027	0.025

Table G.2: continued from previous page

BJD -2,459,000	v_r (km s ⁻¹)	σ_{v_r} (km s ⁻¹)	FWHM (km s ⁻¹)	FWHM error (km s ⁻¹)	BIS (km s ⁻¹)	BIS error (km s ⁻¹)
564.79878234	-59.4646	0.0060	5.8243	0.013	-0.004	0.025
564.81303560	-59.4591	0.0059	5.8107	0.013	0.004	0.025
564.82729206	-59.4647	0.0062	5.8055	0.013	-0.011	0.026
564.84148262	-59.4614	0.0060	5.8159	0.013	-0.006	0.025
566.77933148	-59.4741	0.0067	5.7850	0.015	-0.022	0.029
566.79358211	-59.4721	0.0065	5.7605	0.014	-0.004	0.028
566.80783755	-59.4771	0.0067	5.7811	0.015	0.007	0.029
566.82202828	-59.4656	0.0065	5.7960	0.014	-0.033	0.028
585.76652048	-59.4594	0.0068	5.8175	0.015	-0.001	0.029
585.78071470	-59.4601	0.0067	5.7761	0.014	-0.006	0.028
585.79496722	-59.4660	0.0080	5.8302	0.017	0.018	0.034
585.80915704	-59.4590	0.0077	5.8197	0.017	-0.008	0.033
587.77470698	-59.4634	0.0057	5.7748	0.012	0.005	0.024
587.78896268	-59.4610	0.0056	5.7658	0.012	-0.001	0.023
587.80321708	-59.4624	0.0056	5.7784	0.012	-0.000	0.023
587.81747208	-59.4612	0.0055	5.7682	0.012	0.013	0.023
589.75302522	-59.4619	0.0061	5.7736	0.013	-0.014	0.026
589.76728010	-59.4594	0.0060	5.7631	0.013	-0.023	0.026
589.78153548	-59.4571	0.0059	5.7856	0.013	-0.014	0.025
589.79572667	-59.4578	0.0060	5.7794	0.013	0.005	0.025
591.77126900	-59.4521	0.0060	5.8048	0.013	-0.036	0.025
591.78552487	-59.4560	0.0060	5.7901	0.013	-0.032	0.025
591.79978033	-59.4588	0.0060	5.7753	0.013	-0.023	0.025
591.81397029	-59.4540	0.0057	5.7920	0.012	-0.010	0.024
592.70585888	-59.4500	0.0063	5.7629	0.014	-0.020	0.026
592.72011124	-59.4573	0.0062	5.7706	0.013	-0.017	0.026
592.73436680	-59.4582	0.0061	5.7728	0.013	-0.030	0.026
592.74855726	-59.4583	0.0063	5.7868	0.014	-0.001	0.026
594.76757159	-59.4715	0.0069	5.7864	0.015	-0.002	0.029
594.78182683	-59.4658	0.0071	5.7823	0.015	-0.019	0.030
594.79601747	-59.4743	0.0065	5.7616	0.014	0.007	0.028
594.81027260	-59.4677	0.0063	5.7637	0.014	0.003	0.027
597.73968050	-59.4661	0.0056	5.7792	0.012	-0.004	0.024
597.75393471	-59.4646	0.0056	5.7788	0.012	-0.007	0.023
597.76812613	-59.4661	0.0054	5.7850	0.012	-0.009	0.023
597.78231695	-59.4655	0.0054	5.7863	0.012	-0.011	0.023
598.75863479	-59.4584	0.0063	5.7973	0.014	-0.017	0.026
598.77288920	-59.4642	0.0062	5.8018	0.013	-0.015	0.026
598.78714471	-59.4606	0.0062	5.7775	0.013	-0.002	0.026
598.80133531	-59.4692	0.0060	5.7917	0.013	-0.017	0.025
599.74533714	-59.4562	0.0061	5.7671	0.013	-0.015	0.026
599.75959124	-59.4670	0.0059	5.7894	0.013	0.009	0.025
599.77384994	-59.4649	0.0058	5.7783	0.013	0.000	0.025
599.78803765	-59.4605	0.0058	5.7853	0.013	-0.025	0.025
600.74293878	-59.4719	0.0064	5.8115	0.014	-0.029	0.027
600.75712977	-59.4654	0.0063	5.7874	0.014	-0.045	0.027
600.77132087	-59.4672	0.0062	5.7938	0.014	-0.013	0.026
600.78551106	-59.4695	0.0062	5.7853	0.014	-0.011	0.026
601.74208946	-59.4553	0.0070	5.8273	0.015	-0.025	0.029
601.75634424	-59.4568	0.0068	5.8079	0.015	-0.027	0.029
601.77059913	-59.4661	0.0070	5.8272	0.015	-0.015	0.029
601.78479052	-59.4454	0.0078	5.8485	0.017	-0.004	0.033
602.74801173	-59.4522	0.0060	5.7973	0.013	-0.012	0.025
602.76226731	-59.4602	0.0058	5.8083	0.013	-0.001	0.024
602.77652129	-59.4490	0.0058	5.7821	0.013	-0.000	0.024
602.79071298	-59.4566	0.0057	5.7914	0.012	-0.006	0.024
603.75844953	-59.4634	0.0061	5.7663	0.013	-0.009	0.026
603.77270410	-59.4690	0.0059	5.7794	0.013	-0.016	0.025
603.78695928	-59.4704	0.0059	5.7685	0.013	0.001	0.025
603.80115015	-59.4784	0.0059	5.7862	0.013	-0.002	0.025
604.76579074	-59.4525	0.0069	5.8172	0.015	-0.018	0.029

Table G.2: continued from previous page

BJD -2,459,000	v_r (km s ⁻¹)	σ_{v_r} (km s ⁻¹)	FWHM (km s ⁻¹)	FWHM error (km s ⁻¹)	BIS (km s ⁻¹)	BIS error (km s ⁻¹)
604.78004541	-59.4554	0.0067	5.8201	0.015	-0.038	0.029
604.79430017	-59.4523	0.0065	5.8048	0.014	-0.021	0.028
604.80849134	-59.4634	0.0067	5.8235	0.015	-0.025	0.028
606.73093229	-59.4486	0.0072	5.8098	0.016	-0.018	0.031
606.74518775	-59.4544	0.0072	5.7984	0.016	-0.045	0.031
606.75944270	-59.4588	0.0073	5.8186	0.016	-0.035	0.031
606.77363376	-59.4635	0.0072	5.8029	0.016	-0.031	0.031

Table G.3. B_ℓ and null polarization N observations of TOI-1695.

BJD -2,459,000	B_ℓ (G)	σ_B (G)	N	σ_N	EqW (Å)
473.0016398	-4.12	4.99	5.00	4.87	1.651
476.9045374	-3.21	3.17	1.46	3.36	1.954
479.8851662	1.73	3.12	1.11	3.26	1.944
480.8918336	-1.24	3.43	-2.74	3.55	1.930
481.8775293	-2.05	3.12	-6.80	3.40	1.942
501.9502817	-1.85	3.24	-1.71	3.30	1.813
502.9355799	1.16	3.08	0.02	3.17	1.898
503.9727203	-1.88	3.70	3.94	3.71	1.802
505.9486172	3.73	3.60	6.75	3.73	1.792
507.9496392	0.35	3.70	-6.78	3.82	1.830
508.9567142	6.69	3.39	-0.73	3.51	1.915
509.9643374	-2.14	5.07	-4.97	5.20	1.793
510.9424300	1.40	3.29	-5.95	3.33	1.955
513.9267436	3.88	3.30	-1.46	3.37	1.918
531.7939318	3.00	3.49	2.10	3.66	1.930
535.9308136	-2.69	8.20	-2.33	7.68	1.738
538.8621726	0.39	3.11	2.17	3.19	1.961
539.8701053	-2.49	2.92	-0.01	3.14	1.946
557.8238866	0.52	5.16	-3.23	5.19	1.752
559.7975022	-0.53	4.85	-4.68	4.96	1.785
560.7960801	-3.17	3.54	-1.15	3.60	1.877
561.8158711	4.28	4.91	-5.31	4.91	1.792
562.7782783	0.94	3.15	-2.67	3.18	1.932
563.7854884	3.39	4.40	-4.50	4.45	1.897
564.8188796	0.44	3.96	-2.45	4.08	1.890
566.8006265	0.62	3.32	-1.53	3.40	1.890
585.7837539	0.41	4.69	0.73	6.50	1.776
589.7735483	-3.04	3.03	-1.98	3.38	1.975
591.7921350	1.51	3.68	1.54	3.72	1.947
592.7261387	-3.25	2.99	2.83	3.11	1.967
594.7893179	-7.56	3.65	-9.50	4.00	1.613
597.7601225	-2.01	4.08	-0.30	4.21	1.586
599.7656805	0.53	2.87	-4.19	2.98	1.895
600.7635277	3.07	4.20	-1.51	4.24	1.655
601.7610720	6.95	5.31	5.46	5.43	1.739
602.7681757	1.18	3.65	-6.27	3.77	1.712
603.7787482	1.05	3.64	-1.39	3.82	1.704
606.7510680	6.67	5.46	4.70	5.63	1.662