

Effective field theory in light of relative entropy

Qing-Hong Cao^{1,2,3}, Naoto Kan⁴, and Daiki Ueda³

¹ *Department of Physics and State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing 100871, China*

² *Collaborative Innovation Center of Quantum Matter, Beijing 100871, China*

³ *Center for High Energy Physics, Peking University, Beijing 100871, China*

⁴ *Department of Physics, Osaka University, Toyonaka, Osaka 560-0043, Japan*

E-mail: qinghongcao@pku.edu.cn, naotokan000@gmail.com,
ueda@pku.edu.cn

ABSTRACT: We study constraints on the effective field theory (EFT) from the relative entropy between two theories: we refer to these as target and reference theories. The consequence of the non-negativity of the relative entropy is investigated by choosing some reference theories for a given target theory involving field theories, quantum mechanical models, etc. It is found that the constraints on EFTs, e.g., the single massless scalar field with the dimension-eight operator, and SMEFT dimension-eight $SU(N)$ gauge bosonic operators, are consistent with the positivity bounds from the unitarity and causality when the higher-derivative operators are generated by the interaction between heavy and light fields. The constraints on Einstein-Maxwell theory with higher-derivative operators from the non-negativity of relative entropy are also investigated. The constraints on such EFTs from the relative entropy hold under an assumption that perturbative corrections from the interaction involving higher-derivative operators of light fields are not dominant in the EFTs. The consequence of this study on the weak gravity conjecture and the second law of thermodynamics is also discussed.

Contents

1	Introduction	1
2	Entropy constraint on Euclidean effective action	3
2.1	Main idea	3
2.2	Examples of reference theories	7
2.3	Inequalities satisfied by Euclidean effective action in field theory	11
2.3.1	Non-interacting reference theory	11
2.3.2	Massive free field reference theory	16
2.3.3	Infinite heavy mass reference theory	18
2.4	Summary of entropy constraints	20
3	Top-down approach: relative entropy in various theories	21
3.1	Gaussian distribution functions	21
3.2	A spin system in one dimension	23
3.3	A tree level UV completion of single massless scalar field theory	25
3.4	A tree level UV completion of single massless scalar field theory with linear term	27
3.5	Neutral bosons interacting with photon	33
3.5.1	Dilaton	33
3.5.2	Axion	35
3.6	Massless scalar field with a shift symmetry	37
3.7	Euler-Heisenberg theory	39
3.7.1	Heavy charged fermion field	40
3.7.2	Heavy charged scalar field	42
3.8	Gravitationally coupled massive scalar field at tree level	44
3.9	Dimension-six four-fermion operators from tree-level UV completions	46
3.10	Tree level UV completions with unstable field	48
4	Bottom-up approach: bounds on EFTs	51
4.1	Single massless scalar field with dimension-eight operator	52
4.2	SMEFT dimension-eight gauge bosonic operators	56
4.2.1	Operator Basis	56
4.2.2	Linear combination of single field strength quartics	58
4.2.3	Bounds from non-interacting reference theory	62
4.3	Einstein-Maxwell theory with higher-derivative operators	67
4.4	Summary of bottom-up approach	73

5	Weak gravity conjecture and entropy constraint	73
5.1	Corrections to thermodynamic entropy	73
5.2	Entropy constraints and weak-gravity-conjecture	76
6	Implication of entropy constraints	78
6.1	Unitary time-evolution	78
6.2	Causality	79
6.3	Second law of thermodynamics	79
7	Summary	81
A	Wick rotation	82
B	Relative entropy under field redefinition	86
C	Calculation of the second law of thermodynamics	87
D	Einstein-Maxwell theory with higher-derivative operators under field redefinition	88

1 Introduction

Effective field theory (EFT) is a fundamental framework for describing low-energy phenomena. Information about the high energy regime is transferred to the EFT by integrating out heavy degrees of freedom and can be extracted by determination of the parameters of the EFTs. Extracting the nature of the information about the high-energy regime would be a significant scientific goal, and the EFT approach is actively studied from both experimental and theoretical points of view.

From an experimental point of view, there is growing attention to the EFT approach to describe physics beyond the Standard Model (SM). The CERN large hadron collider (LHC) has discovered the Higgs boson [1, 2] and strengthened the foundation of the SM. Overwhelming evidence and hints require physics beyond the SM. Still, the intensive searches for new particles at the weak scale or heavier have yet to find convincing evidence of such new particles. In these circumstances, information about the new particles is transferred to the EFT involving the SM fields by integrating out the new particles. The EFTs such as the Standard Model Effective Field Theory (SMEFT) [3–10] are actively studied in this situation. Various observables provide constraints on the SMEFT Wilson coefficients [11–35], which could point us to the UV completion of the SMEFT in the future.

From a theoretical point of view, to exclude particular EFTs, the Weak Gravity Conjecture (WGC) [36] (see also [37] for a review) is actively studied. The string theory yields a vast landscape of four-dimensional EFTs [38]. In contrast to the landscape, the set of EFTs which cannot be generated

from quantum gravity is called the swampland [39]¹. Predicting quantitative probability distribution about what kind of EFTs belong to the landscape is a challenge. A simpler version of this challenge is suggested as the WGC, i.e., a rule to distinguish the landscape from the swampland. A mild version of the WGC states that the $U(1)$ charge-to-mass ratio of extremally charged black holes is larger than unity in any gravitational EFT that admits a consistent UV completion [36, 48]. Some attempted derivations for this statement have been made using black holes and entropy consideration [48–50] or positivity bounds [51] from unitarity and causality [52, 53]. In particular, Refs. [48–50] are based on a positivity of entropy difference between Einstein-Maxwell theories with and without perturbative corrections from the higher-dimensional operators. These works imply a close connection of the positive entropy difference with the positivity bounds from unitarity and causality. Although the WGC is suggested in the context of quantum gravity, the methodology to exclude particular EFTs that cannot be UV completed is useful in various EFTs with and without gravity, and such a close connection naturally leads us to consider a new approach to constraints on the EFTs.

Recently, inspired by the connection between the entropy and positivity bounds, a new approach [54] has been proposed to constrain the EFTs by a property of the relative entropy [55–57]. The relative entropy defined by two probability distribution functions is a non-negative quantity, which is often used as a distance-like concept between the two probability distribution functions. In Ref. [54], consequences of the non-negativity of the relative entropy have been studied by defining probability distribution function for various theories. They mainly considered the distance between theories with and without the interaction between heavy and light degrees of freedom. They showed that the relative entropy yields constraints on some EFTs such as the single massless scalar field with the dimension-eight operator, dimension-eight $SU(N)$ gauge bosonic operators in the SMEFT, and Einstein-Maxwell theory with higher-derivative operators when the higher-derivative operators are generated by the interaction between heavy and light fields. These arguments for the constraints on the Wilson coefficients hold under an assumption that perturbative corrections from the interaction involving higher-derivative operators of light fields are not dominant in the EFTs. The connection of the non-negativity of the relative entropy with the WGC and the second law of thermodynamics is also discussed.

The key role of the distance-like concept in the constraints on the EFTs would lead to an interest in considering the relative entropy between a given theory and various theories. For example, in Euclidean space, the distances between a given point and various points yield information about the coordinate of the given point. Similarly, we can evaluate the relative entropy between the given theory and several theories and study their distances. We refer to the given theory that one wants to extract its information as a *target theory*. Also, we refer to the other theories as *reference theories*. The relative entropy between the target and reference theories would provide various information about the target theory depending on the reference theory. The appropriate reference theory should be selected depending on the information one wants to extract. Then, the reference theory generally describes quite different physics from the target theory. In Ref. [54], the theory with the interaction between

¹Criteria of the swampland are studied as the swampland conjectures [36, 40–47]. The WGC is one of the swampland conjectures.

heavy and light degrees of freedom denotes the target theory, and the theory without the interaction is a reference theory. In this paper, we refer to the reference theory of Ref. [54] as the *non-interacting reference theory*.

In this paper, we provide the details of Ref. [54] and update the results in Ref. [54] by considering more target theories and new reference theories. We provide some new reference theories such as *massive free field reference theory*, which also yields the constraints on perturbative corrections from the heavy degrees of freedom to the Euclidean effective action. Each reference theory would have different advantages depending on the target theory. For each reference theory, we provide calculation methods of the relative entropy between the target theory and the reference theory. The relative entropy is calculated by the Euclidean path integral method, and therefore our following discussions are based on the validity of the Euclidean path integral method. We adopt the top-down approach for consistency checks and evaluate the relative entropy for various target theories containing heavy degrees of freedom. Also, we adopt the bottom-up approach and investigate the consequence of the non-negativity of relative entropy in EFTs such as the single massless scalar field with the dimension-eight operator, SMEFT dimension-eight $SU(N)$ gauge bosonic operators, and Einstein-Maxwell theory with higher-derivative operators. In addition, we will discuss connections of this study with some inequality such as causality, the second law of thermodynamics, the WGC, etc.

This paper is organized as follows. In Sec. 2, we review the details of the main idea of the entropy constraint on EFT of Ref. [54] and provide the procedures to calculate the relative entropy by introducing some new reference theories. In Sec. 3 we follow the top-down approach and consider various target theories to perform consistency checks of the entropy constraint. In Sec. 4 we follow the bottom-up approach and provide the bounds on some EFTs from the relative entropy. In Secs. 5 and 6, we discuss connections between the entropy constraint and some inequalities in physics. We finish with the summary of the paper in Sec. 7.

2 Entropy constraint on Euclidean effective action

In this section, for the sake of being self-contained, we start with a review of the entropy constraint [54] and then update the discussion of Ref. [54]. Inequalities satisfied by the Euclidean effective actions of the two different theories or systems are provided from the non-negativity of the relative entropy. In Sec. 2.1, we explain the main idea of the entropy constraint in two ways: the field theoretical approach and the quantum mechanical approach. In Sec. 2.2, some reference theories addressed in this paper are listed. In Sec. 2.3, we focus on the field theory and provide some inequalities satisfied by the Euclidean effective action. In Sec. 2.4, we summarize some properties of the entropy constraint.

2.1 Main idea

The relative entropy is defined by two probability distribution functions ρ_R and ρ_T as follows:

$$S(\rho_R||\rho_T) \equiv \text{Tr} [\rho_R \ln \rho_R - \rho_R \ln \rho_T], \quad (2.1)$$




Figure 1. A schematic picture of the main idea. Relative entropy is a distance-like quantity between two probability distribution functions. For a given probability distribution function of the target theory ρ_T , the relative entropy between ρ_T and reference probability distribution functions ρ_R yields inequalities about ρ_T . Various information about ρ_T can be extracted by changing ρ_R . The appropriate reference probability distribution function is selected, depending on the information one wants to extract. For example, for a reference theory ρ_{R1} , the relative entropy represents the constraints on the EFTs generated from the target theory; see Secs. 3 and 4, and for ρ_{R2} , the relative entropy denotes the second law of thermodynamics in the system described by the target theory; see Sec. 6.3, and so on.

where ρ_R and ρ_T satisfy $\rho_{R,T} = \rho_{R,T}^\dagger$, and $\text{Tr}[\rho_{R,T}] = 1$ because they are probability distribution functions. One of the important properties of the relative entropy is non-negativity. For convenience, we provide brief proof of the non-negativity of the relative entropy. Consider a convex function $f(x)$, which satisfies $f(x_R) - f(x_T) \leq (x_R - x_T) \cdot df(x_T)/dx$. For $f(x) \rightarrow x \ln x$, $x \rightarrow \rho_R$, and $x_T \rightarrow \rho_T$, the definition of convex function yields

$$S(\rho_R || \rho_T) = \text{Tr} [\rho_R \ln \rho_R - \rho_R \ln \rho_T] \geq 0. \quad (2.2)$$

Note here that, in Eq. (2.2), the equality holds if and only if $\rho_R = \rho_T$ by the definition of convex function. Therefore, the relative entropy characterizes differences between two probability distribution functions ρ_R and ρ_T and is often used as a distance between ρ_R and ρ_T even though it is not a symmetric function of the two sets of probabilities $S(\rho_R || \rho_T) \neq S(\rho_T || \rho_R)$.

In the entropy constraint [54], we define the probability distribution function ρ_T for the theory or system T which is the target from which one wants to extract its information. In this work, we mainly focus on perturbation corrections generated by heavy degrees of freedom of the theory T and attempt to extract the information about their properties. We refer to the theory T as the *target theory*. On the other hand, we define ρ_R for a reference system R, which is an auxiliary system to extract the information about the theory T by comparing T with R. We refer to the theory R as the *reference*

theory. The main idea of the entropy constraint is to *evaluate the relative entropy between the target theory and suitable reference theory and extract the information about the target theory*. In Fig. 1, we schematically describes the main idea of the entropy constraint. In Ref. [54], a non-interacting theory are chosen as the reference theory, and a connection between the non-negativity of the relative entropy and the positivity bounds on EFTs has been studied. It should be noted that the reference theory is on the same Hilbert space as the target theory but is generally not relevant to the target theory and describes different physics from the target theory. The point is that the extracted information from the non-negativity of the relative entropy changes depending on the reference theory, even if the target theory does not change. In other words, we need to choose the suitable reference theory depending on the information one wants to extract from the target theory.

First, consider the system described by the field theory and evaluate the relative entropy between two theories, R and T by defining the probability distribution function of the two theories. Although its definition is not unique, in this paper, we mainly consider probability distribution functions defined as follows:

$$P_R[\Phi] \equiv \frac{e^{-I_R[\Phi]}}{Z_R}, \quad P_T[\Phi] \equiv \frac{e^{-I_T[\Phi]}}{Z_T}, \quad (2.3)$$

where I_R and I_T are Euclidean actions of the theories, R and T, respectively, Φ denotes degrees of freedom of the field theoretical dynamics, and the partition functions are defined as

$$Z_R \equiv \int d[\Phi] e^{-I_R[\Phi]}, \quad Z_T \equiv \int d[\Phi] e^{-I_T[\Phi]}. \quad (2.4)$$

The relative entropy between theories R and T is calculated as follows:

$$S(P_R||P_T) \equiv \int d[\Phi] [P_R[\Phi] \ln P_R[\Phi] - P_R[\Phi] \ln P_T[\Phi]] \quad (2.5)$$

$$= \int d[\Phi] [P_R[\Phi] (-\ln Z_R - I_R[\Phi]) - P_R[\Phi] (-\ln Z_T - I_T[\Phi])] \quad (2.6)$$

$$= -\ln Z_R + \ln Z_T + \int d[\Phi] P_R[\Phi] (I_T[\Phi] - I_R[\Phi]) \quad (2.7)$$

$$= W_R - W_T + \langle I_T - I_R \rangle_R \geq 0, \quad (2.8)$$

where $W_R \equiv -\ln Z_R$, $W_T \equiv -\ln Z_T$, and $\langle I_T - I_R \rangle_R \equiv \int d[\Phi] P_R[\Phi] (I_T[\Phi] - I_R[\Phi])$. In the second line, $\ln P_{R,T} = -\ln Z_{R,T} - I_{R,T}[\Phi]$ is used, and $\int d[\Phi] P_{R,T} = 1$ yields the third line. The last line arises from the non-negativity of the relative entropy. Therefore, it follows from Eq. (2.8) that the upper bound on $W_T - W_R$ is expressed as

$$\langle I_T - I_R \rangle_R \geq W_T - W_R. \quad (2.9)$$

Similar to the above procedures, another choice of the relative entropy is calculated as follows:

$$S(P_T||P_R) \equiv \int d[\Phi] [P_T[\Phi] \ln P_T[\Phi] - P_T[\Phi] \ln P_R[\Phi]]$$

$$\begin{aligned}
&= \int d[\Phi] [P_T[\Phi] (-\ln Z_T - I_T[\Phi]) - P_T[\Phi] (-\ln Z_R - I_R[\Phi])] \\
&= -\ln Z_T + \ln Z_R - \int d[\Phi] P_T[\Phi] (I_T[\Phi] - I_R[\Phi]) \\
&= W_T - W_R - \langle I_T - I_R \rangle_T \geq 0,
\end{aligned} \tag{2.10}$$

where $\langle I_T - I_R \rangle_T \equiv \int d[\Phi] P_T[\Phi] (I_T[\Phi] - I_R[\Phi])$. Then, Eq. (2.10) yields the lower bound on $W_T - W_R$ as

$$W_T - W_R \geq \langle I_T - I_R \rangle_T. \tag{2.11}$$

The Euclidean actions I_R and I_T are determined by the Wick-rotated Lagrangian and boundary conditions, and the above explanations are valid even for finite temperature systems. Our strategy is to extract the information about I_T from the inequalities of (2.9) and (2.11) by choosing the suitable reference theory I_R .

Next, consider the system described by quantum mechanical dynamics and evaluate the relative entropy between the two systems R and T. The above discussions do not rely on the Lorentz symmetry, so similar inequalities are derived for the quantum mechanical dynamics. Equation (2.3) corresponds to the following density operators.

$$\rho_R \equiv \frac{e^{-\beta H_R}}{Z_R}, \quad \rho_T \equiv \frac{e^{-\beta H_T}}{Z_T}, \tag{2.12}$$

where H_R and H_T are Hamiltonians of R and T, respectively, β is an inverse temperature of the systems, and the partition functions are defined as

$$Z_R \equiv \text{Tr}[e^{-\beta H_R}], \quad Z_T \equiv \text{Tr}[e^{-\beta H_T}]. \tag{2.13}$$

The above definition of the probability distribution functions of Eq. (2.12) is one example, and other choices are also possible. We will show different choices in a later section. Equation (2.2), (2.12), and (2.13) yield the relative entropy between theories R and T as follows:

$$\begin{aligned}
S(\rho_R || \rho_T) &\equiv \text{Tr} [\rho_R \ln \rho_R - \rho_R \ln \rho_T] \\
&= \text{Tr} [\rho_R (-\ln Z_R - \beta H_R) - \rho_R (-\ln Z_T - \beta H_T)] \\
&= -\ln Z_R + \ln Z_T + \beta \cdot \text{Tr} [\rho_R (H_T - H_R)] \\
&= W_R - W_T + \beta \cdot \langle H_T - H_R \rangle_R \geq 0,
\end{aligned} \tag{2.14}$$

where $W_R = -\ln Z_R$, $W_T = -\ln Z_T$, and $\langle H_T - H_R \rangle_R \equiv \text{Tr} [\rho_R (H_T - H_R)]$. In the second line, $\ln \rho_{R,T} = -\ln Z_{R,T} - \beta H_{R,T}$ is used, and $\text{Tr}[\rho_{R,T}] = 1$ yields the third line. The last line arises from the non-negativity of the relative entropy. Similar to the field theoretical approach, Eq. (2.14) yields the upper bound on $W_T - W_R$ as

$$\beta \cdot \langle H_T - H_R \rangle_R \geq W_T - W_R. \tag{2.15}$$

Another choice of the relative entropy is also calculated as follows:

$$\begin{aligned}
S(\rho_T || \rho_R) &\equiv \text{Tr} [\rho_T \ln \rho_T - \rho_T \ln \rho_R] \\
&= \text{Tr} [\rho_T (-\ln Z_T - \beta H_T) - \rho_T (-\ln Z_R - \beta H_R)] \\
&= -\ln Z_T + \ln Z_R - \beta \cdot \text{Tr} [\rho_T (H_T - H_R)] \\
&= W_T - W_R - \beta \cdot \langle H_T - H_R \rangle_T \geq 0,
\end{aligned} \tag{2.16}$$

where $\langle H_T - H_R \rangle_T \equiv \text{Tr} [\rho_T (H_T - H_R)]$. Eq. (2.16) yields the lower bound on $W_T - W_R$ as

$$W_T - W_R \geq \beta \cdot \langle H_T - H_R \rangle_T. \tag{2.17}$$

Consequently, we obtain the lower and upper bounds on $W_T - W_R$ from the non-negativity of the relative entropy by both the field theoretical and quantum mechanical approaches. In the next section, we provide some examples of the reference theory to derive constraints on perturbative corrections from heavy degrees of freedom to the Euclidean effective action of the target theory.

2.2 Examples of reference theories

For the target theories consisting of heavy and light fields, we consider EFTs generated by integrating out the heavy fields. Throughout this section, Φ 's and ϕ 's denote the heavy and light fields in the field theoretical dynamics, respectively. The Euclidean action of the target theory is expressed as follows:

$$I_T[\phi, \Phi] = I_0[\phi, \Phi] + I_I[\phi, \Phi], \tag{2.18}$$

where I_0 does not include the interaction between Φ 's and ϕ 's, and I_I is the interacting term. From Eq. (2.3), for the background fields, ϕ 's, the probability distribution functions of the theories R and T are defined as follows:

$$P_R[\Phi] \equiv \frac{e^{-I_R[\phi, \Phi]}}{Z_R[\phi]}, \quad P_T[\Phi] \equiv \frac{e^{-I_T[\phi, \Phi]}}{Z_T[\phi]}, \tag{2.19}$$

with the partition functions,

$$Z_R[\phi] \equiv \int d[\Phi] e^{-I_R[\phi, \Phi]}, \quad Z_T[\phi] \equiv \int d[\Phi] e^{-I_T[\phi, \Phi]}. \tag{2.20}$$

The relative entropy between P_R and P_T is calculated as in Eq. (2.8) and (2.10). The path integral is performed only over the dynamical heavy field because ϕ is the background field.

Even for dynamical light fields ϕ 's², the probability distribution functions of the theories R and T can be defined as follows:

$$P_R[\phi, \Phi] \equiv \frac{e^{-I_R[\phi, \Phi]}}{Z_R[\tilde{\phi}_R, \tilde{\Phi}_R]}, \quad P_T[\phi, \Phi] \equiv \frac{e^{-I_T[\phi, \Phi]}}{Z_T[\tilde{\phi}_T, \tilde{\Phi}_T]}, \tag{2.21}$$

²We have to be careful with the validity of the Euclidean path integral over ϕ 's. When we treat ϕ 's as the dynamical fields, the path integral needs to be performed around a local minimum. If not, the saddle point approximation breaks down. We need not require such validity if ϕ 's are the background fields.

where $(\tilde{\phi}_R, \tilde{\Phi}_R)$ and $(\tilde{\phi}_T, \tilde{\Phi}_T)$ are sets of classical solutions of I_R and I_T , respectively. The partition functions are given by

$$Z_R[\tilde{\phi}_R, \tilde{\Phi}_R] \equiv \int d[\phi]d[\Phi] e^{-I_R[\phi, \Phi]}, \quad Z_T[\tilde{\phi}_T, \tilde{\Phi}_T] \equiv \int d[\phi]d[\Phi] e^{-I_T[\phi, \Phi]}. \quad (2.22)$$

The relative entropy is calculated as in Eq. (2.8) by replacing $d[\Phi]$ with $d[\phi]d[\Phi]$ as follows:

$$\begin{aligned} S(P_R||P_T) &\equiv \int d[\phi]d[\Phi] [P_R[\phi, \Phi] \ln P_R[\phi, \Phi] - P_R[\phi, \Phi] \ln P_T[\phi, \Phi]] \\ &= \int d[\phi]d[\Phi] [P_R[\phi, \Phi] (-\ln Z_R - I_R[\phi, \Phi]) - P_R[\phi, \Phi] (-\ln Z_T - I_T[\phi, \Phi])] \\ &= -\ln Z_R + \ln Z_T + \int d[\phi]d[\Phi] P_R[\phi, \Phi] (I_T[\phi, \Phi] - I_R[\phi, \Phi]) \\ &= W_R - W_T + \langle I_T - I_R \rangle_R \geq 0, \end{aligned} \quad (2.23)$$

where $Z_R \equiv Z_R[\tilde{\phi}_R, \tilde{\Phi}_R]$, $Z_T \equiv Z_T[\tilde{\phi}_T, \tilde{\Phi}_T]$, $W_R \equiv -\ln Z_R[\tilde{\phi}_R, \tilde{\Phi}_R]$, $W_T \equiv -\ln Z_T[\tilde{\phi}_T, \tilde{\Phi}_T]$, and

$$\langle I_T - I_R \rangle_R \equiv \int d[\phi]d[\Phi] P_R[\phi, \Phi] (I_T[\phi, \Phi] - I_R[\phi, \Phi]). \quad (2.24)$$

For the dynamical light fields, the relative entropy of (2.10) is given by

$$\begin{aligned} S(P_T||P_R) &\equiv \int d[\phi]d[\Phi] [P_T[\phi, \Phi] \ln P_T[\phi, \Phi] - P_T[\phi, \Phi] \ln P_R[\phi, \Phi]] \\ &= \int d[\phi]d[\Phi] [P_T[\phi, \Phi] (-\ln Z_T - I_T[\phi, \Phi]) - P_T[\phi, \Phi] (-\ln Z_R - I_R[\phi, \Phi])] \\ &= -\ln Z_T + \ln Z_R - \int d[\phi]d[\Phi] P_T[\phi, \Phi] (I_T[\phi, \Phi] - I_R[\phi, \Phi]) \\ &= W_T - W_R - \langle I_T - I_R \rangle_T \geq 0, \end{aligned} \quad (2.25)$$

where

$$\langle I_T - I_R \rangle_T \equiv \int d[\phi]d[\Phi] P_T[\phi, \Phi] (I_T[\phi, \Phi] - I_R[\phi, \Phi]). \quad (2.26)$$

Equations (2.23) and (2.25) are the same as the form of Eq. (2.8) and (2.10), respectively, which do not depend on whether the light fields are dynamical or not. To clarify procedures of the wave function renormalization of the light fields, we assume the dynamical light fields in Sec. 3.6, 3.7, and 4 but the light background fields in the other sections.

In the following, we list some reference theories to derive information about the target theory. The first three examples are relevant to the constraints on the corrections to W_T from Φ 's. In particular, the first reference theory plays an important role in deriving the constraints on EFTs in the bottom-up approach, which is discussed in Sec. 4. The last one is connected with the second law of thermodynamics; see Sec. 6.3.

- Non-interacting reference theory — In Ref. [54], the Euclidean action of the reference theory is defined as a non-interacting theory as follows:

$$I_{\text{NI}}[\phi, \Phi] \equiv I_0[\phi, \Phi], \quad (2.27)$$

where I_0 is the same as the first term of Eq. (2.18). We refer to this reference theory as the non-interacting reference theory (NIRT). For the background light fields ϕ 's, the probability distribution function of the NIRT is defined as

$$P_{\text{NI}}[\Phi] \equiv \frac{e^{-I_{\text{NI}}[\phi, \Phi]}}{Z_{\text{NI}}[\phi]}, \quad (2.28)$$

with the partition function,

$$Z_{\text{NI}}[\phi] \equiv \int d[\Phi] e^{-I_{\text{NI}}[\phi, \Phi]}. \quad (2.29)$$

For the dynamical light fields ϕ 's, we define the probability distribution function of the NIRT as

$$P_{\text{NI}}[\phi, \Phi] \equiv \frac{e^{-I_{\text{NI}}[\phi, \Phi]}}{Z_{\text{NI}}[\tilde{\phi}_{\text{NI}}, \tilde{\Phi}_{\text{NI}}]}, \quad (2.30)$$

where the partition function is given by

$$Z_{\text{NI}}[\tilde{\phi}_{\text{NI}}, \tilde{\Phi}_{\text{NI}}] \equiv \int d[\phi] d[\Phi] e^{-I_{\text{NI}}[\phi, \Phi]}, \quad (2.31)$$

where $\tilde{\phi}_{\text{NI}}$ and $\tilde{\Phi}_{\text{NI}}$ are classical solutions of I_{NI} .

- Massive free field reference theory — We propose a reference theory defined by an Euclidean action,

$$I_{\text{MF}}[\phi, \Phi] \equiv I_\phi[\phi] + I_\Phi[\Phi], \quad (2.32)$$

where $I_\phi[\phi] \equiv I_0[\phi, 0]$. I_Φ denotes only the kinetic and mass terms of Φ , and its mass term is the same as that of I_0 . In contrast to the NIRT, the self-interacting terms of Φ do not include in I_{MF} . We refer this reference theory as the massive free field reference theory (MFFRT). When ϕ 's are assumed to be background fields, we perform the path integral only over Φ 's. Then, the probability distribution functions of the MFFRT is defined as follows:

$$P_{\text{MF}}[\Phi] \equiv \frac{e^{-I_{\text{MF}}[\phi, \Phi]}}{Z_{\text{MF}}[\phi]}, \quad (2.33)$$

with the partition functions,

$$Z_{\text{MF}}[\phi] \equiv \int d[\Phi] e^{-I_{\text{MF}}[\phi, \Phi]}, \quad (2.34)$$

where solutions of I_{MF} for the heavy fields take zero values. For the dynamical light fields ϕ 's, the probability distribution functions of the MFFRT is defined as follows:

$$P_{\text{MF}}[\phi, \Phi] \equiv \frac{e^{-I_{\text{MF}}[\phi, \Phi]}}{Z_{\text{MF}}[\tilde{\phi}_{\text{MF}}, \tilde{\Phi}_{\text{MF}}]}, \quad (2.35)$$

where the partition function is given by

$$Z_{\text{MF}}[\tilde{\phi}_{\text{MF}}, \tilde{\Phi}_{\text{MF}}] \equiv \int d[\phi] d[\Phi] e^{-I_{\text{MF}}[\phi, \Phi]}, \quad (2.36)$$

where $\tilde{\phi}_{\text{MF}}$ and $\tilde{\Phi}_{\text{MF}}$ are classical solutions of I_{MF} . $\tilde{\Phi}_{\text{MF}}$ can take zero values by absorbing the plane wave solutions into quantum fluctuations.

- Infinite heavy mass reference theory — As the reference theory, we consider a theory with the same form of the action as the target theory with the infinite mass of Φ ,

$$I_{\text{IH}}[\phi, \Phi] \equiv \lim_{m_{\Phi} \rightarrow \infty} I_{\text{T}}[\phi, \Phi]. \quad (2.37)$$

We refer to this reference theory as the infinite heavy mass reference theory (IHMRT). Throughout this work, for the IHMRT, we focus on the background light fields and perform the path integral only over Φ 's. The probability distribution of the IHMRT is defined as,

$$P_{\text{IH}}[\Phi] \equiv \frac{e^{-I_{\text{IH}}[\phi, \Phi]}}{Z_{\text{IH}}[\phi]}, \quad (2.38)$$

with the partition function,

$$Z_{\text{IH}}[\phi] \equiv \int d[\Phi] e^{-I_{\text{IH}}[\phi, \Phi]}. \quad (2.39)$$

In Sec. 3, we study the IHMRT only in the tree level calculations.

- Thermal reference theory — Consider a system consisting of a thermodynamic system S and a heat bath system B. We assume both heavy and light degrees of freedom are included in the thermodynamic system S. The Hamiltonian of the whole system is expressed as $H_{\text{T}} = H_{\text{S}} + H_{\text{B}} + H_{\text{SB}}$, where H_{S} is the Hamiltonian of the thermodynamic system S, and H_{B} is that of the heat bath system B. The interacting term H_{SB} denotes the interaction between S and B and can generally depends on time. At the initial time, assume the quantum state of the whole system is expressed as

$$\rho_{\text{T}} \equiv \rho_{\text{ini}} = \rho_{\text{ini}, \text{S}} \otimes e^{-\beta H_{\text{B}}} / Z_{\text{B}}(\beta), \quad (2.40)$$

where $\rho_{\text{ini}, \text{S}}$ is the initial state of S, and $e^{-\beta H_{\text{B}}} / Z_{\text{B}}(\beta)$ is that of B, β is an inverse temperature of the heat bath system at the initial time, and $Z_{\text{B}}(\beta) \equiv \text{Tr}_{\text{B}}[e^{-\beta H_{\text{B}}}]$ is defined by tracing over the heat bath degrees of freedom. Note here that the specific form of $\rho_{\text{ini}, \text{S}}$ is irrelevant to this discussion. We assume the probability distribution of the target theory is defined by the initial

state of the whole system. After the time evolution described by a unitary operator U , the final state of the whole system is expressed as

$$\rho_{\text{fin}} \equiv U \rho_{\text{ini}} U^\dagger. \quad (2.41)$$

By tracing out the heat bath degrees of freedom, the final state of S is calculated as

$$\rho_{\text{fin},S} \equiv \text{Tr}_B[\rho_{\text{fin}}]. \quad (2.42)$$

Then, define the reference probability distribution function as follows:

$$\rho_R \equiv U^\dagger \rho_{\text{fin},S} \otimes e^{-\beta H_B} / Z_B(\beta) U. \quad (2.43)$$

We refer to this reference theory as the thermal reference theory in this work. The thermal reference theory is useful to see a connection between the non-negativity of relative entropy and the second law of thermodynamics [58, 59]. In the other reference theories discussed before, it is supposed that the target and reference theory do not include the heat bath degrees of freedom. However, in Sec. 6.3, we will demonstrate that the relative entropy between the target and reference theories does not change even if the heat bath degrees of freedom are added to both the theories.

One of our main interests is the constraints on EFT generated by the target theory, and the above first three reference theories are mainly considered in the following sections. Here, we would like to emphasize that the above definitions of the probability distributions of the target theory and reference theory are not unique, and are part of examples.

2.3 Inequalities satisfied by Euclidean effective action in field theory

We have discussed the general properties of the relative entropy between two probability distribution functions so far. In this section, we focus on the systems described by field theoretical dynamics and provide inequalities satisfied by heavy field corrections to the Euclidean effective action by using the non-negativity of the relative entropy. For each reference theory, we provide inequalities satisfied by the Euclidean effective action of the target theory in the following.

2.3.1 Non-interacting reference theory

By introducing an auxiliary parameter g , we define,

$$I_g[\phi, \Phi] \equiv I_0[\phi, \Phi] + g \cdot I_I[\phi, \Phi]. \quad (2.44)$$

By changing the parameter g , the target and reference theories are given as,

$$I_T[\phi, \Phi] = \lim_{g \rightarrow 1} I_g[\phi, \Phi], \quad I_{\text{NI}}[\phi, \Phi] = \lim_{g \rightarrow 0} I_g[\phi, \Phi]. \quad (2.45)$$

For the background light fields, the partition function and effective action of I_g are respectively defined as follows:

$$Z_g[\phi] \equiv \int d[\Phi] e^{-I_g[\phi, \Phi]}, \quad (2.46)$$

$$W_g[\phi] \equiv -\ln Z_g[\phi], \quad (2.47)$$

From Eqs. (2.8) and (2.10), by defining a probability distribution function,

$$P_g[\Phi] \equiv \frac{e^{-I_g[\phi, \Phi]}}{Z_g[\phi]}, \quad (2.48)$$

the relative entropy between P_0 and P_g is calculated as follows:

$$S(P_0||P_g) = W_0[\phi] - W_g[\phi] + g \cdot \langle I_I \rangle_{g=0} \geq 0 \quad \Rightarrow \quad W_g[\phi] - W_0[\phi] \leq g \cdot \langle I_I \rangle_{g=0}, \quad (2.49)$$

$$S(P_g||P_0) = W_g[\phi] - W_0[\phi] - g \cdot \langle I_I \rangle_g \geq 0 \quad \Rightarrow \quad g \cdot \langle I_I \rangle_g \leq W_g[\phi] - W_0[\phi], \quad (2.50)$$

with

$$\langle I_I \rangle_g \equiv \int d[\Phi] I_I[\phi, \Phi] \cdot \frac{e^{-I_g[\phi, \Phi]}}{Z_g[\phi]} = \frac{\partial W_g[\phi]}{\partial g}, \quad (2.51)$$

where the partial derivative means differentiating by g while keeping ϕ . Equations (2.49) and (2.50) yield

$$g \cdot \langle I_I \rangle_{g=0} \geq W_g[\phi] - W_0[\phi] \geq g \cdot \langle I_I \rangle_g \quad \Rightarrow \quad \langle I_I \rangle_{\text{NI}} \geq W_{\text{T}}[\phi] - W_{\text{NI}}[\phi] \geq \langle I_I \rangle_{\text{T}} \quad \text{for } g = 1. \quad (2.52)$$

Here, we used $W_{\text{T}} = W_{g=1}$, $W_{\text{NI}} = W_0$, and the following relations.

$$\begin{aligned} \langle I_I \rangle_{g=0} &= \left(\frac{\partial W_g[\phi]}{\partial g} \right)_{g=0} \\ &= \int d[\Phi] I_I[\phi, \Phi] \cdot \frac{e^{-I_0[\phi, \Phi]}}{Z_0[\phi]} \\ &= \int d[\Phi] I_I[\phi, \Phi] \cdot \frac{e^{-I_{\text{NI}}[\phi, \Phi]}}{Z_{\text{NI}}[\phi]} \\ &= \langle I_I \rangle_{\text{NI}}, \end{aligned} \quad (2.53)$$

$$\begin{aligned} \langle I_I \rangle_{g=1} &= \left(\frac{\partial W_g[\phi]}{\partial g} \right)_{g=1} \\ &= \int d[\Phi] I_I[\phi, \Phi] \cdot \frac{e^{-I_{g=1}[\phi, \Phi]}}{Z_{g=1}[\phi]} \\ &= \int d[\Phi] I_I[\phi, \Phi] \cdot \frac{e^{-I_{\text{T}}[\phi, \Phi]}}{Z_{\text{T}}[\phi]} \\ &= \langle I_I \rangle_{\text{T}}, \end{aligned} \quad (2.54)$$

where, in particular, Eq. (2.53) denotes the Feynman diagrams of Fig. 2. Note here that the Euclidean effective action $W_{\text{NI}}[\phi]$ generally includes the corrections from the self-interacting terms of Φ . For ease of understanding, let us schematically express the Euclidean effective actions as follows:

$$W_{\text{NI}}[\phi] = I_0[\phi, 0] + (\text{vacuum energy})', \quad (2.55)$$

$$W_T[\phi] = I_0[\phi, 0] + (\text{vacuum energy})' + (\text{renormalizable terms of } \phi) + (\text{non-renormalizable terms of } \phi), \quad (2.56)$$

where $I_T[\phi, 0] = I_0[\phi, 0]$ and $I_{NI}[\phi, 0] = I_0[\phi, 0]$ are used. Here, $(\text{vacuum energy})'$ denotes the vacuum energy coming from the dynamical fields, which may include corrections from the self-interacting terms of Φ . Note that $(\text{vacuum energy})'$ is independent of the background fields ϕ 's. Also, the third term of the right-hand side of Eq. (2.56) denotes the corrections from the interacting term I_I to the renormalizable terms of ϕ , and their fourth term is the corrections from the interacting term I_I to the non-renormalizable terms of ϕ . Therefore, $W_T[\phi] - W_{NI}[\phi]$ represents the perturbative corrections from the interaction between heavy and light degrees of freedom to the Euclidean effective action of the target theory other than the vacuum energy as follows:

$$W_T[\phi] - W_{NI}[\phi] = (\text{renormalizable terms of } \phi) + (\text{non-renormalizable terms of } \phi). \quad (2.57)$$

The point is that the right-hand side of Eq. (2.57) does not include ϕ independent terms. Equations (2.52) and (2.57) imply that *the expectation values of the interaction yield bounds on the perturbative corrections from the interacting term I_I to the Euclidean effective action of the target theory*. For example, $W_T[\phi] - W_{NI}[\phi]$ is increased in the theory with $\langle I_I \rangle_T \geq 0$ but decreased in the theory with $\langle I_I \rangle_{NI} \leq 0$. In Ref. [54], the theory satisfying $\langle I_I \rangle_{NI} \leq 0$ is referred to as the *non-positive interacting* theory.

For convenience, we explain the meaning of the upper bound of Eq. (2.52). Expand W_g with respect to g as follows:

$$\begin{aligned} W_g[\phi] &= W_0[\phi] + g \cdot \left(\frac{\partial W_g[\phi]}{\partial g} \right)_{g=0} + \frac{g^2}{2} \cdot \left(\frac{\partial^2 W_g[\phi]}{\partial g^2} \right)_{g=0} + \mathcal{O}(g^3), \\ &= W_{NI}[\phi] + g \cdot \langle I_I \rangle_{NI} + \frac{g^2}{2} \cdot \left(\frac{\partial^2 W_g[\phi]}{\partial g^2} \right)_{g=0} + \mathcal{O}(g^3), \\ &= W_{NI}[\phi] + g \cdot \langle I_I \rangle_{NI} + \Delta W_g^{(2)}, \end{aligned} \quad (2.58)$$

where $W_0[\phi] = W_{NI}[\phi]$ is used, and we defined the corrections for the second or higher order for g as

$$\Delta W_g^{(2)} \equiv \frac{g^2}{2} \cdot \left(\frac{\partial^2 W_g}{\partial g^2} \right)_{g=0} + \mathcal{O}(g^3). \quad (2.59)$$

Combining Eq. (2.58), and the upper bound of Eq. (2.52), we obtain

$$\Delta W_g^{(2)} \leq 0 \quad \Rightarrow \quad \Delta W_{g=1}^{(2)} \leq 0 \quad \text{for } g = 1. \quad (2.60)$$

Note here that $g \cdot \langle I_I \rangle_{NI}$ cancels in the upper bound of Eq. (2.52). Consequently, the upper bound of Eq. (2.52) means that the Euclidean effective action decreases by the second or higher order corrections for the interaction I_I . Also, according to Eq. (2.53), the non-positive interacting theory is a class of theories in which the Euclidean effective action is unchanged, or reduced at the first order of the interaction. For the non-positive interacting theory, the sign of the shift of the Euclidean effective action is the same as that of the second or higher order corrections for the interaction. In other words,




Figure 2. Feynman diagrams for $\langle I_I \rangle_{\text{NI}}$ at the tree and one-loop level. The vertices denote the interacting term I_I . The solid and dashed lines are the heavy and the background light fields, respectively. The left and right diagrams denote tree and one-loop level corrections, respectively. In the non-interacting reference theory, the classical solution of the heavy field may take a non-zero value because of the linear term of Φ , and the left diagram generally does not vanish but can be eliminated by redefinitions of Φ ; see Sec. 3.4.

the non-positive interaction, i.e., $\langle I_I \rangle_{\text{NI}} \leq 0$, is a sufficient condition to reduce the Euclidean effective action by the interaction I_I .

Focusing on the NIRT, we explain an important property of the relative entropy, i.e., the invariance of Eq. (2.9) under the field redefinition to eliminate the linear term of Φ . Consider a target theory with the linear term in the Euclidean space,

$$I_T[\phi, \Phi] = I_0^{\text{lin}}[\phi, \Phi] + I_I[\phi, \Phi], \quad (2.61)$$

where I_0^{lin} is assumed to involve the linear term of Φ , and I_I is the interacting term. Assume the classical solution of I_0^{lin} for Φ takes v , where indices of the classical solution, such as Lorentz indices, are omitted. Also, the classical solution of I_T for Φ takes $v + f(\phi)$, where $f(\phi)$ depends on the light field ϕ because of the interacting term I_I . Note here that f vanishes in the limit of $I_I \rightarrow 0$. Define the action of NIRT as,

$$I_{\text{NI}}[\phi, \Phi] \equiv I_0^{\text{lin}}[\phi, \Phi]. \quad (2.62)$$

At the tree level, the Euclidean effective actions of the target and reference theories are respectively calculated as follows:

$$W_T[\phi] = I_0^{\text{lin}}[\phi, v + f(\phi)] + I_I[\phi, v + f(\phi)], \quad (2.63)$$

$$W_{\text{NI}}[\phi] = I_0^{\text{lin}}[\phi, v]. \quad (2.64)$$

The shift of the Euclidean effective action is calculated as

$$W_T[\phi] - W_{\text{NI}}[\phi] = I_0^{\text{lin}}[\phi, v + f(\phi)] + I_I[\phi, v + f(\phi)] - I_0^{\text{lin}}[\phi, v]. \quad (2.65)$$

The expectation value of the interaction I_I in the Euclidean space is also calculated as

$$\langle I_I \rangle_{\text{NI}} = \int d[\Phi] P_{\text{NI}}[\Phi] I_I[\phi, \Phi] = I_I[\phi, v], \quad (2.66)$$

where $P_{\text{NI}}[\Phi] \equiv e^{-I_{\text{NI}}}/Z_{\text{NI}}[\phi]$ with $Z_0[\phi] \equiv \int d[\Phi] e^{-I_0^{\text{lin}}}$. Equations (2.65), (2.66), and (2.52) yield,

$$\begin{aligned} W_{\text{T}}[\phi] - W_{\text{NI}}[\phi] &\leq \langle I_I \rangle_{\text{NI}} \\ &\Rightarrow I_0^{\text{lin}}[\phi, v + f(\phi)] + I_I[\phi, v + f(\phi)] - I_I[\phi, v] - I_0^{\text{lin}}[\phi, v] \leq 0. \end{aligned} \quad (2.67)$$

Next, consider a field redefinition $\Phi \equiv \eta + v$. Equation (2.61) is expressed as

$$\begin{aligned} I'_{\text{T}}[\phi, \eta] &\equiv I_{\text{T}}[\phi, \eta + v] \\ &= I_0^{\text{lin}}[\phi, \eta + v] + I_I[\phi, \eta + v] \\ &= \left(I_0^{\text{lin}}[\phi, \eta + v] + I_I[\phi, v] \right) + (I_I[\phi, \eta + v] - I_I[\phi, v]). \end{aligned} \quad (2.68)$$

Here, we define

$$I'_0[\phi, \eta] \equiv I_0^{\text{lin}}[\phi, \eta + v] + I_I[\phi, v], \quad (2.69)$$

$$I'_I[\phi, \eta] \equiv I_I[\phi, \eta + v] - I_I[\phi, v], \quad (2.70)$$

where I'_0 does not include the linear term of η . Also define the action of NIRT as

$$I'_{\text{NI}}[\phi, \Phi] \equiv I'_0[\phi, \Phi]. \quad (2.71)$$

At the tree level, the Euclidean effective actions of I'_{T} and I'_{NI} are respectively calculated as follows:

$$W'_{\text{T}}[\phi] = I'_0[\phi, f(\phi)] + I'_I[\phi, f(\phi)] = I_0^{\text{lin}}[\phi, v + f(\phi)] + I_I[\phi, v] + I'_I[\phi, f(\phi)], \quad (2.72)$$

$$W'_{\text{NI}}[\phi] = I_0^{\text{lin}}[\phi, v] + I_I[\phi, v]. \quad (2.73)$$

Note here that the classical solution of I'_{T} for η takes $f(\phi)$ because that of I_{T} for $\Phi = v + \eta$ is $v + f(\phi)$. Similarly, the classical solution of I'_{NI} for η takes a zero value because that of I_{NI} for $\Phi = v + \eta$ is v . Then, the shift of the Euclidean effective action is calculated as

$$\begin{aligned} W'_{\text{T}}[\phi] - W'_{\text{NI}}[\phi] &= I_0^{\text{lin}}[\phi, v + f(\phi)] + I_I[\phi, v] + I'_I[\phi, f(\phi)] - I_0^{\text{lin}}[\phi, v] - I_I[\phi, v] \\ &= I_0^{\text{lin}}[\phi, v + f(\phi)] + I'_I[\phi, f(\phi)] - I_0^{\text{lin}}[\phi, v]. \end{aligned} \quad (2.74)$$

The expectation value of the interaction I'_I in the Euclidean space is calculated as

$$\langle I'_I \rangle_{\text{NI}} = \int d[\eta] P'_{\text{NI}}[\eta] I'_I[\phi, \eta] = 0, \quad (2.75)$$

where $P'_{\text{NI}}[\eta] \equiv e^{-I'_{\text{NI}}[\phi, \eta]}/Z'_{\text{NI}}[\phi]$ with $Z'_{\text{NI}}[\phi] \equiv \int d[\eta] e^{-I'_{\text{NI}}[\phi, \eta]}$. Equations (2.74), (2.75), and Eq. (2.52) yield,

$$W'_{\text{T}}[\phi] - W'_{\text{NI}}[\phi] \leq \langle I'_I \rangle_{\text{NI}}$$

$$\begin{aligned}
&\Rightarrow I_0^{\text{lin}}[\phi, v + f(\phi)] + I_1'[\phi, f(\phi)] - I_0^{\text{lin}}[\phi, v] \\
&= I_0^{\text{lin}}[\phi, v + f(\phi)] + I_1[\phi, v + f(\phi)] - I_1[\phi, v] - I_0^{\text{lin}}[\phi, v] \leq 0.
\end{aligned} \tag{2.76}$$

This result is the same as Eq. (2.67). Consequently, it is found that the upper bound of Eq. (2.52) is invariant under the field redefinition to remove the linear term of Φ . This result means that, at the tree level, theories can be the non-positive interacting theory by the field redefinition. In Sec. 3.4, we will show an example of this result. Since the calculations of the relative entropy becomes easier by the field redefinition to remove the linear term of Φ , in Sec. 4, we often use the procedures of Eq. (2.68), (2.69), and (2.70).

We comment on some properties of the NIRT. One of the main features of the NIRT is that the ϕ -independent terms cancel in $W_{\text{T}}[\phi] - W_{\text{NI}}[\phi]$ as in Eq. (2.57). In the context of the positivity bounds on EFTs, a class of EFTs that corrections to the renormalizable terms can be removed by the redefinition of the light fields, has been actively studied. For such a class of theories, the cancellation of the ϕ -independent terms is convenient to derive the constraints on the correction to higher-derivative terms by using Eq. (2.52)³. Indeed, in Sec. 3 and 4, we will provide constraints on the higher-dimensional operators of such theories. We will see that the results are consistent with the positivity bounds obtained by the analyticity, causality and unitarity.

2.3.2 Massive free field reference theory

We rewrite the action I_0 as follows:

$$I_0[\phi, \Phi] = I_0[\phi, 0] + I_\Phi[\Phi] + I_S[\Phi], \tag{2.77}$$

where I_Φ is the action of free field Φ , and I_S is the self-interacting term of Φ . By introducing an auxiliary parameters g , we define an action as follows:

$$J_g[\phi, \Phi] \equiv I_0[\phi, 0] + I_\Phi[\Phi] + g \cdot (I_S[\Phi] + I_I[\phi, \Phi]). \tag{2.78}$$

The action of Eq. (2.78) satisfies,

$$\lim_{g \rightarrow 0} J_g[\phi, \Phi] = I_{\text{MF}}[\phi, \Phi], \quad \lim_{g \rightarrow 1} J_g[\phi, \Phi] = I_{\text{T}}[\phi, \Phi]. \tag{2.79}$$

When the light fields are the background fields, the partition function and effective actions of J_g are defined as follows:

$$z_g[\phi] \equiv \int d[\Phi] e^{-J_g[\phi, \Phi]}, \tag{2.80}$$

$$w_g[\phi] \equiv -\ln z_g[\phi]. \tag{2.81}$$

From Eqs. (2.8) and (2.10), by defining a probability distribution function,

$$p_g[\Phi] \equiv \frac{e^{-J_g[\phi, \Phi]}}{z_g[\phi]}, \tag{2.82}$$

³Even in finite temperature systems, field independent terms cancel in $W_{\text{T}}[\phi] - W_{\text{NI}}[\phi]$.

the relative entropy between p_0 and p_g is calculated as follows:

$$S(p_0||p_g) = w_0[\phi] - w_g[\phi] + g \cdot \langle I_S + I_I \rangle_{g=0} \geq 0 \Rightarrow w_g[\phi] - w_0[\phi] \leq g \cdot \langle I_S + I_I \rangle_{g=0}, \quad (2.83)$$

$$S(p_g||p_0) = w_g[\phi] - w_0[\phi] - g \cdot \langle I_S + I_I \rangle_g \geq 0 \Rightarrow g \cdot \langle I_S + I_I \rangle_g \leq w_g[\phi] - w_0[\phi], \quad (2.84)$$

with

$$\langle I_S + I_I \rangle_g \equiv \int d[\Phi] (I_S[\Phi] + I_I[\phi, \Phi]) \cdot \frac{e^{-J_g[\phi, \Phi]}}{z_g[\phi]} = \frac{\partial w_g[\phi]}{\partial g}, \quad (2.85)$$

where the partial derivative denotes differentiating by g while keeping ϕ . Combining Eq. (2.83) and (2.84), we obtain

$$\begin{aligned} g \cdot \langle I_S + I_I \rangle_{g=0} &\geq w_g[\phi] - w_0[\phi] \geq g \cdot \langle I_S + I_I \rangle_g \\ \Rightarrow \langle I_S + I_I \rangle_{\text{MF}} &\geq W_{\text{T}}[\phi] - W_{\text{MF}}[\phi] \geq \langle I_S + I_I \rangle_{\text{T}} \text{ for } g = 1. \end{aligned} \quad (2.86)$$

Here, we used $W_{\text{T}} = w_{g=1}$, $W_{\text{MF}} = w_0$, and the following relations,

$$\begin{aligned} \langle I_S + I_I \rangle_{g=0} &= \left(\frac{\partial w_g[\phi]}{\partial g} \right)_{g=0} \\ &= \int d[\Phi] (I_S[\Phi] + I_I[\phi, \Phi]) \cdot \frac{e^{-J_0[\phi, \Phi]}}{z_0[\phi]} \\ &= \int d[\Phi] (I_S[\Phi] + I_I[\phi, \Phi]) \cdot \frac{e^{-I_{\text{MF}}[\phi, \Phi]}}{Z_{\text{MF}}[\phi]} \\ &= \langle I_S + I_I \rangle_{\text{MF}}, \end{aligned} \quad (2.87)$$

$$\begin{aligned} \langle I_S + I_I \rangle_{g=1} &= \left(\frac{\partial w_g[\phi]}{\partial g} \right)_{g=1} \\ &= \int d[\Phi] (I_S[\Phi] + I_I[\phi, \Phi]) \cdot \frac{e^{-J_{g=1}[\phi, \Phi]}}{z_{g=1}[\phi]} \\ &= \int d[\Phi] (I_S[\Phi] + I_I[\phi, \Phi]) \cdot \frac{e^{-I_{\text{T}}[\phi, \Phi]}}{Z_{\text{T}}[\phi]} \\ &= \langle I_S + I_I \rangle_{\text{T}}, \end{aligned} \quad (2.88)$$

where $z_0 = Z_{\text{MF}}$ and $z_{g=1} = Z_{\text{T}}$ are used.

In particular, the Feynman diagrams for $\langle I_S \rangle_{\text{MF}}$ and $\langle I_I \rangle_{\text{MF}}$ are shown in Fig. 3 and 4, respectively. The Euclidean effective actions are schematically expressed as follows:

$$W_{\text{MF}}[\phi] = I_0[\phi, 0] + (\text{vacuum energy}), \quad (2.89)$$

$$W_{\text{T}}[\phi] = I_0[\phi, 0] + (\text{vacuum energy}) + (\text{corrections from } \Phi). \quad (2.90)$$

where (vacuum energy) denotes the vacuum energy coming from Φ loop effects, and the third term of the right-hand side of Eq. (2.90) denotes the perturbative corrections from Φ other than the vacuum energy. The shift of the Euclidean effective action is given by

$$W_{\text{T}}[\phi] - W_{\text{MF}}[\phi] = (\text{corrections from } \Phi). \quad (2.91)$$

Note here that the (corrections from Φ) may include ϕ independent terms because the self-interacting term of Φ in I_T appears. Therefore, $W_T - W_{\text{MF}}$ also includes the correction from the self-interacting terms of Φ in contrast to $W_T - W_{\text{NI}}$, and the right-hand side of Eq. (2.86) may include the field-independent terms. The point is that the inequality of (2.86) provides different information about the target theory from the NIRT. Some applications are provided in the later section.

Here, we consider the meaning the upper bound of (2.86). Similar to the case of the NIRT, expand w_g for g as follows:

$$\begin{aligned} w_g[\phi] &= w_0[\phi] + g \cdot \left(\frac{\partial w_g[\phi]}{\partial g} \right)_{g=0} + \frac{g^2}{2} \cdot \left(\frac{\partial^2 w_g[\phi]}{\partial g^2} \right)_{g=0} + \mathcal{O}(g^3), \\ &= W_{\text{MF}}[\phi] + g \cdot \langle I_S + I_I \rangle_{\text{MF}} + \frac{g^2}{2} \cdot \left(\frac{\partial^2 w_g[\phi]}{\partial g^2} \right)_{g=0} + \mathcal{O}(g^3), \\ &= W_{\text{MF}}[\phi] + g \cdot \langle I_S + I_I \rangle_{\text{MF}} + \Delta w_g^{(2)}, \end{aligned} \quad (2.92)$$

where $w_0[\phi] = W_{\text{MF}}[\phi]$ is used, and we defined

$$\Delta w_g^{(2)} \equiv \frac{g^2}{2} \cdot \left(\frac{\partial^2 w_g[\phi]}{\partial g^2} \right)_{g=0} + \mathcal{O}(g^3). \quad (2.93)$$

From Eq. (2.92) and (2.86), we get

$$\Delta w_g^{(2)} \leq 0. \quad (2.94)$$

Therefore, the inequality of (2.86) means that the Euclidean effective action decrease by the second or higher order corrections for I_S and I_I .

2.3.3 Infinite heavy mass reference theory

The upper bound of Eq. (2.9) is calculated as,

$$\langle I_T - \lim_{m_\Phi \rightarrow \infty} I_T \rangle_{\text{IH}} \geq W_T[\phi] - W_{\text{IH}}[\phi], \quad (2.95)$$

where $W_{\text{IH}}[\phi] = -\ln Z_{\text{IH}}[\phi]$. The left-hand side of Eq. (2.95) is defined as follows:

$$\langle I_T - \lim_{m_\Phi \rightarrow \infty} I_T \rangle_{\text{IH}} = \int d[\Phi] P_{\text{IH}}[\Phi] \left(I_T[\phi, \Phi] - \lim_{m_\Phi \rightarrow \infty} I_T[\phi, \Phi] \right). \quad (2.96)$$

For simplicity, we focus on the tree level UV completions. Using the saddle point approximations, we obtain

$$Z_{\text{IH}}[\phi] = \int d[\Phi] e^{-I_{\text{IH}}[\phi, \Phi]} = e^{-I_{\text{IH}}[\phi, \tilde{\Phi}_{\text{IH}}]}, \quad (2.97)$$

$$Z_T[\phi] = \int d[\Phi] e^{-I_T[\phi, \Phi]} = e^{-I_T[\phi, \tilde{\Phi}_T]}, \quad (2.98)$$

$$\int d[\Phi] P_{\text{IH}}[\Phi] (I_T[\phi, \Phi] - I_{\text{NI}}[\phi, \Phi]) = I_T[\phi, \tilde{\Phi}_{\text{IH}}] - \lim_{m_\Phi \rightarrow \infty} I_T[\phi, \tilde{\Phi}_{\text{IH}}]. \quad (2.99)$$




Figure 3. Feynman diagrams for $\langle I_I \rangle_{\text{MF}}$ at the tree and one-loop level. The vertices denote the interacting term I_I . The solid and dashed lines are the heavy and the background light fields, respectively. In contrast to the non-interacting reference theory, the classical solution of the heavy field can take zero value since the reference theory is the massive free field theory. The left diagram is a tree-level correction, which vanishes because of zero values of the classical solution in the reference theory. The right diagram denotes a one-loop level correction, which generally does not vanish.




Figure 4. Feynman diagrams for $\langle I_S \rangle_{\text{MF}}$ at the one-loop level. The vertices denote the self-interacting term I_S . The solid line denotes the heavy field. The right diagram vanishes because of the classical solution of the reference theory, but the left diagram may remain.

Similar to the NIRT, $\langle I_T - \lim_{m_\Phi \rightarrow \infty} I_T \rangle_{\text{IH}}$ does not vanish when the linear term of Φ in I_{IH} remains. However, $\langle I_T - \lim_{m_\Phi \rightarrow \infty} I_T \rangle_{\text{IH}} = 0$ holds after the field redefinition of Φ such that ϕ independent terms are removed in Φ_{IH} . For $\langle I_T - \lim_{m_\Phi \rightarrow \infty} I_T \rangle_{\text{IH}} = 0$, one obtains as follows:

$$0 \geq W_T[\phi] - W_{\text{IH}}[\phi]. \quad (2.100)$$

The above right-hand side denotes the correction from Φ to the Euclidean effective action of the target theory.

2.4 Summary of entropy constraints

In Secs. 2.2 and 2.3, based on the Euclidean path integral method, we provide procedures to calculate the relative entropy in the field theory. Given the target theory, the relative entropy can be calculated by the procedures explained in Sec. 2.3. Before summarizing the properties of the relative entropy, we list the assumptions used to derive the results in this section in the following way.

- (I) **Hermiticity of probability distribution functions** — We assume the target and reference theory are represented by the Hermitian probability distribution functions. To derive the non-negativity of the relative entropy in Eq. (2.2), we used the Hermiticity of probability distribution functions, i.e., $\rho_{R,T} = \rho_{R,T}^\dagger$. The non-negativity of the relative entropy can be broken when this condition is not satisfied.
- (II) **Validity of Euclidean path integral method** — We assume the EFTs are generated from the solution of the local minimum. As shown in Sec. 3.10, the non-negativity of the relative entropy can be broken when the Euclidean path integral method is not valid, i.e., the saddle point approximation does not work because of the solution not being the local minimum.

Under these assumptions, for the NIRT and MFFRT, we obtained the properties that do not depend on the details of the target theories as follows:

- **Non-interacting reference theory** — The upper and lower bounds on the perturbative corrections from the interaction between heavy and light degrees of freedom to the Euclidean effective action is denoted as Eq. (2.52),

$$\langle I_I \rangle_{NI} \geq W_T[\phi] - W_{NI}[\phi] \geq \langle I_I \rangle_T,$$

where $W_T - W_{NI}$ is the perturbative correction to renormalizable and unrenormalizable terms for ϕ , and the background independent terms are canceled. The upper bound of Eq. (2.52) is rewritten as Eq. (2.60),

$$\Delta W_g^{(2)} \leq 0.$$

This inequality means that the Euclidean effective action decreases by the second or higher order corrections for the interaction between heavy and light degrees of freedom. In general, $\langle I_I \rangle_{NI}$ takes a non-zero value, but it can vanish at the tree level by the redefinition to eliminate the linear term of Φ in I_T . The relative entropy is invariant under such a field redefinition.

- **Massive free field reference theory** — The upper and lower bounds on the perturbative corrections from the self-interaction of heavy fields and the interaction between heavy and light degrees of freedom to the Euclidean effective action is denoted as Eq. (2.86),

$$\langle I_S + I_I \rangle_{MF} \geq W_T[\phi] - W_{MF}[\phi] \geq \langle I_S + I_I \rangle_T.$$

For $W_T - W_{\text{NI}}$, the ϕ independent vacuum energy from the heavy field loop effects cancels, but the ϕ independent perturbative corrections from the self-interacting term of Φ may be included. The upper bound of Eq. (2.86) is rewritten as Eq. (2.94),

$$\Delta w_g^{(2)} \leq 0. \quad (2.101)$$

This inequality means that the Euclidean effective action decreases by the second or higher order corrections for the self-interaction of heavy fields and the interaction between heavy and light degrees of freedom. On the left-hand side of the above inequality, ϕ independent perturbative corrections from the self-interacting term of Φ are generally included.

In the next section, we will calculate the relative entropies for various target theories and check the non-negativity of the relative entropy, i.e., the above properties. Also, we often face a situation where the EFT is known while the target UV theory is unknown. We refer to this situation as the bottom-up approach and consider such a situation in Sec. 4. In Sec. 4, we focus on a class of EFTs that the corrections to non-higher derivative terms are removed by field redefinitions and apply the NIRT to derive the constraints on such EFTs. The NIRT is more convenient than the MFFRT in the bottom-up approach because the background independent terms vanish in $W_T - W_{\text{NI}}$. In Sec. 4, we provide the constraints on some EFTs under an assumption that the corrections from the interactions involving higher-derivative operators of the light fields are not dominant in the EFTs.

3 Top-down approach: relative entropy in various theories

In this section, for consistency checks of Sec. 2, we evaluate Eqs. (2.52), (2.86), and (2.95) in various theories. In particular, we focus on the upper bound on $W_T - W_R$, which is relevant to the positivity bound on EFTs. We adopt the top-down approach, i.e., UV theories including heavy degrees of freedom are assumed to be known, and evaluate the inequalities in the previous sections. As a pedagogical example, we first consider probability distribution functions described by the Gaussian distribution function. In Sec. 3.2, the constraint on a quantum mechanical model consisting of spins is studied. In the other examples, we focus on weakly coupled field theoretical dynamics and consider up to four-derivative operators. In Sec. 3.10, we will also explain that the non-negativity of the relative entropy can be violated when the Euclidean path integral method does not work, i.e., the saddle point approximations are not valid.

3.1 Gaussian distribution functions

Firstly, we consider a target system described by the Gaussian distribution function defined as follows:

$$I_T[x, X] \equiv m^2 \cdot x^2 + M^2 \cdot X^2 + c \cdot x \cdot X, \quad [\text{Target}] \quad (3.1)$$

where x and X are not fields, but variables corresponding to the light and heavy degrees of freedom, respectively, and m , M , and c are real constant parameters. Although this target theory is neither a field theoretical nor a quantum mechanical model, this is a pedagogical example to understand the

procedure to calculate the relative entropy. We assume x is not an integral variable and behaves as a background field. The free theory and the interaction between x and X are respectively denoted as

$$I_0[x, X] \equiv m^2 \cdot x^2 + M^2 \cdot X^2, \quad (3.2)$$

$$I_I[x, X] \equiv c \cdot x \cdot X. \quad (3.3)$$

For this target theory, the NIRT is the same as the MFFRT. The action of the NIRT and MFFRT is defined as

$$I_R[x, X] \equiv I_0[x, X]. \quad \text{[Reference]} \quad (3.4)$$

Then, probability distribution functions are defined as follows:

$$P_R[X] \equiv \frac{1}{Z_R[x]} e^{-I_R[x, X]}, \quad P_T[X] \equiv \frac{1}{Z_T[x]} e^{-I_T[x, X]}, \quad (3.5)$$

with the partition functions

$$Z_R[x] \equiv \int_{-\infty}^{\infty} dX e^{-I_R[x, X]}, \quad Z_T[x] \equiv \int_{-\infty}^{\infty} dX e^{-I_T[x, X]}. \quad (3.6)$$

From these partition functions, the effective actions are given by

$$W_R[x] \equiv -\ln Z_R[x], \quad W_T[x] \equiv -\ln Z_T[x]. \quad (3.7)$$

By introducing an auxiliary parameter g , we define,

$$I_g[x, X] \equiv I_0[x, X] + g \cdot I_I[x, X]. \quad (3.8)$$

By changing g , the target and reference theory are reproduced as follows:

$$I_T[x, X] = \lim_{g \rightarrow 1} I_g[x, X], \quad I_R[x, X] = \lim_{g \rightarrow 0} I_g[x, X]. \quad (3.9)$$

The partition function and effective action of I_g are respectively calculated as follows:

$$Z_g[x] \equiv \int d[X] e^{-I_g[x, X]} = e^{-m^2 x^2} \sqrt{\frac{\pi}{M^2}} \cdot e^{g^2 \cdot c^2 x^2 / 4M^2}, \quad (3.10)$$

$$W_g[x] \equiv -\ln Z_g[x] = -\frac{g^2 \cdot c^2 x^2}{4M^2} - \ln \left[e^{-m^2 x^2} \sqrt{\frac{\pi}{M^2}} \right]. \quad (3.11)$$

By defining a probability distribution function,

$$P_g[X] \equiv \frac{e^{-I_g[x, X]}}{Z_g[x]}, \quad (3.12)$$

the expectation value of the interaction is also calculated as

$$\langle I_I \rangle_{g=0} = \int_{-\infty}^{\infty} dX P_0[X] \cdot I_I[x, X] = 0. \quad (3.13)$$

From Eqs. (2.8), (3.11), and (3.13), the relative entropy between P_0 and P_g is given by

$$\begin{aligned}
S(P_0||P_g) &\equiv \int_{-\infty}^{\infty} dX (P_0 \ln P_0 - P_0 \ln P_g), \\
&= -\ln Z_0[x] + \ln Z_g[x] + g \cdot \langle I_I \rangle_{g=0}, \\
&= -\ln Z_0[x] + \ln Z_g[x], \\
&= W_0[x] - W_g[x], \\
&= \frac{g^2 \cdot c^2 x^2}{4M^2} \geq 0.
\end{aligned} \tag{3.14}$$

Note here that the relative entropy is invariant under the field redefinition of X . The definition of the interaction of Eq. (3.3) is not invariant under the redefinition of X , but the relative entropy, i.e., the integral of the Gaussian distributions, do not change under the field redefinition. By taking to be $g = 1$, we obtain the relative entropy between P_R and P_T as follows:

$$S(P_R||P_T) = W_R[x] - W_T[x] = \frac{c^2 x^2}{4M^2} \geq 0, \tag{3.15}$$

where $P_R = P_0$, $P_T = P_{g=1}$, $W_R = W_{g=0}$, and $W_T = W_{g=1}$ are used. The above inequality represents the upper bound of Eq. (2.52) and (2.86). The relative entropy takes a positive value, as we expected. The above procedure of calculation of the relative entropy is the same as the field theoretical dynamics.

3.2 A spin system in one dimension

The entropy inequality is derived even in the quantum mechanical model. Let us consider a spin system in one dimension defined by a Hamiltonian,

$$H_T \equiv -J \sum_{i=1}^{N/2} \sigma_{2i-1} \sigma_{2i} - \mu M \sum_{i=1}^N \sigma_i, \quad [\textbf{Target}] \tag{3.16}$$

where $\sigma_i = \pm 1$ denotes a spin on site i , J is a coupling characterizing exchange interactions, N is the number of sites, μ is a magnetic moment, and M is an external magnetic field. We assume N is even so that the system consists of $N/2$ pairs of adjacent sites. Then, H_0 and H_I are defined as follows:

$$H_0 \equiv -J \sum_{i=1}^{N/2} \sigma_{2i-1} \sigma_{2i}, \tag{3.17}$$

$$H_I \equiv -\mu M \sum_{i=1}^N \sigma_i. \tag{3.18}$$

The Hamiltonian of the NIRT is defined as

$$H_{NI} \equiv H_0. \quad [\textbf{Reference}] \tag{3.19}$$

The density operators of the target and reference systems are respectively given by

$$\rho_{\text{NI}} \equiv \frac{e^{-\beta H_{\text{NI}}}}{Z_{\text{NI}}(\beta)}, \quad \rho_{\text{T}} \equiv \frac{e^{-\beta H_{\text{T}}}}{Z_{\text{T}}(\beta)}, \quad (3.20)$$

with the inverse temperature β , and the partition functions

$$Z_{\text{NI}}(\beta) \equiv \text{Tr}[e^{-\beta H_{\text{NI}}}], \quad Z_{\text{T}}(\beta) \equiv \text{Tr}[e^{-\beta H_{\text{T}}}]. \quad (3.21)$$

The effective actions are given by

$$W_{\text{NI}}(\beta) \equiv -\ln Z_{\text{NI}}(\beta), \quad W_{\text{T}}(\beta) \equiv -\ln Z_{\text{T}}(\beta). \quad (3.22)$$

By introducing the parameter g , we define

$$H_g \equiv H_0 + g \cdot H_{\text{I}}. \quad (3.23)$$

The target and reference theory are reproduced as follows:

$$H_{\text{T}} = \lim_{g \rightarrow 1} H_g, \quad H_{\text{R}} = \lim_{g \rightarrow 0} H_g. \quad (3.24)$$

The partition function and effective action of H_g are respectively given as follows:

$$Z_g(\beta) \equiv \text{Tr}[e^{-\beta H_g}] = \left(2\{e^{\beta J} \cosh(2\beta g \mu M) + e^{-\beta J}\} \right)^{N/2}, \quad (3.25)$$

$$W_g(\beta) \equiv -\ln Z_g(\beta). \quad (3.26)$$

By defining a density operator

$$\rho_g(\beta) \equiv \frac{e^{-\beta H_g}}{Z_g(\beta)}, \quad (3.27)$$

the expectation value of the interaction is calculated as

$$\text{Tr}[\rho_0 H_{\text{I}}] = 0. \quad (3.28)$$

From Eq. (2.14), the relative entropy between ρ_0 and ρ_g is given by

$$\begin{aligned} S(\rho_0 || \rho_g) &\equiv \text{Tr}[\rho_0 \ln \rho_0 - \rho_0 \ln \rho_g], \\ &= -\ln Z_0(\beta) + \ln Z_g(\beta) + g \cdot \text{Tr}[\rho_0 H_{\text{I}}], \\ &= -\ln Z_0(\beta) + \ln Z_g(\beta), \\ &= W_0(\beta) - W_g(\beta), \\ &= -\frac{N}{2} \ln \left[\frac{e^{\beta J} + e^{-\beta J}}{e^{\beta J} \cosh(2\beta g \mu M) + e^{-\beta J}} \right] \geq 0. \end{aligned} \quad (3.29)$$

The first line denotes the definition of the relative entropy, and in the third line, $\text{Tr}[\rho_0 H_{\text{I}}] = 0$ is used.

By taking to be $g = 1$, the relative entropy between ρ_{NI} and ρ_{T} is given by

$$S(\rho_{\text{NI}} || \rho_{\text{T}}) = W_{\text{NI}}(\beta) - W_{\text{T}}(\beta) = -\frac{N}{2} \ln \left[\frac{e^{\beta J} + e^{-\beta J}}{e^{\beta J} \cosh(2\beta \mu M) + e^{-\beta J}} \right] \geq 0, \quad (3.30)$$

where $\rho_{\text{NI}} = \rho_0$, $\rho_{\text{T}} = \rho_{g=1}$, $W_{\text{NI}} = W_{g=0}$, and $W_{\text{T}} = W_{g=1}$ are used. This result represents the upper bound of Eq. (2.52). We see that the external magnetic field decreases the Euclidean effective action of the target system because of the non-negativity of the relative entropy.

3.3 A tree level UV completion of single massless scalar field theory

Consider a theory in Minkowski space:

$$I_T[\phi, \Phi] \equiv \int d^4x \left(\frac{1}{2}(\partial_\mu \phi \partial^\mu \phi) + \frac{1}{2}(\partial_\mu \Phi \partial^\mu \Phi) - \frac{m^2}{2}\Phi^2 + \frac{\alpha}{\Lambda} \cdot \Phi(\partial_\mu \phi \partial^\mu \phi) \right), \text{ [Target]} \quad (3.31)$$

where ϕ denotes a massless scalar field, Φ is a heavy scalar field with mass m , and α/Λ is a dimensionful coupling constant. The above theory involving a linear term of Φ will be studied later. The action in the Euclidean space is expressed as

$$\begin{aligned} I_T^{(E)}[\phi, \Phi] &= \int (d^4x)_E \left(\frac{1}{2}(\partial_I \phi_E \partial_I \phi_E) + \frac{1}{2}(\partial_I \Phi_E \partial_I \Phi_E) + \frac{m^2}{2}\Phi_E^2 + \frac{\alpha}{\Lambda} \cdot \Phi_E(\partial_I \phi_E \partial_I \phi_E) \right), \\ &= \int (d^4x)_E \left(-\frac{1}{2}(\partial_\mu \phi \partial^\mu \phi) + \frac{1}{2}(\partial_I \Phi_E \partial_I \Phi_E) + \frac{m^2}{2}\Phi_E^2 - \frac{\alpha}{\Lambda} \cdot \Phi_E(\partial_\mu \phi \partial^\mu \phi) \right), \end{aligned} \quad (3.32)$$

where, in the second line, we assume the background field ϕ of the Euclidean space are defined from that of the Minkowski space; see Appendix A. We define the actions I_0 and I_I in the Euclidean space as follows:

$$I_0^{(E)}[\phi, \Phi] \equiv I_T^{(E)}[0, \Phi] + I_T^{(E)}[\phi, 0] = \int (d^4x)_E \left(-\frac{1}{2}(\partial_\mu \phi \partial^\mu \phi) + \frac{1}{2}(\partial_I \Phi_E \partial_I \Phi_E) + \frac{m^2}{2}\Phi_E^2 \right), \quad (3.33)$$

$$I_I^{(E)}[\phi, \Phi] \equiv I_T^{(E)}[\phi, \Phi] - I_0^{(E)}[\phi, \Phi] = -\frac{\alpha}{\Lambda} \cdot \int (d^4x)_E \Phi_E(\partial_\mu \phi \partial^\mu \phi). \quad (3.34)$$

For the background field ϕ , by integrating out Φ_E at the tree level, the partition function and effective action are defined as

$$Z_T[\phi] \equiv \int d[\Phi_E] e^{-I_T^{(E)}[\phi, \Phi_E]}, \quad (3.35)$$

$$W_T[\phi] \equiv -\ln Z_T[\phi], \quad (3.36)$$

In this target theory, the action of the MFFRT is the same as that of the NIRT. For each reference theory in Sec. 2.2, we consider the inequalities of (2.52), (2.86), and (2.95) in the following.

- **Non-interacting reference theory** — The action of the NIRT in the Euclidean space is expressed as follows:

$$I_{\text{NI}}^{(E)}[\phi, \Phi] \equiv \int (d^4x)_E \left(-\frac{1}{2}(\partial_\mu \phi \partial^\mu \phi) + \frac{1}{2}(\partial_I \Phi_E \partial_I \Phi_E) + \frac{m^2}{2}\Phi_E^2 \right). \text{ [Reference]} \quad (3.37)$$

The partition function and effective action of the NIRT are defined as

$$Z_{\text{NI}}[\phi] \equiv \int d[\Phi_E] e^{-I_{\text{NI}}^{(E)}[\phi, \Phi]}, \quad (3.38)$$

$$W_{\text{NI}}[\phi] \equiv -\ln Z_{\text{NI}}[\phi]. \quad (3.39)$$

By using the parameter g , we define

$$I_g^{(\text{E})}[\phi, \Phi] \equiv I_0^{(\text{E})}[\phi, \Phi] + g \cdot I_1^{(\text{E})}[\phi, \Phi]. \quad (3.40)$$

The target and reference theories are respectively expressed as follows:

$$I_{\text{T}}^{(\text{E})}[\phi, \Phi] = \lim_{g \rightarrow 1} I_g^{(\text{E})}[\phi, \Phi], \quad I_{\text{NI}}^{(\text{E})}[\phi, \Phi] = \lim_{g \rightarrow 0} I_g^{(\text{E})}[\phi, \Phi]. \quad (3.41)$$

The partition function and effective action of $I_g^{(\text{E})}$ are respectively calculated as follows:

$$Z_g[\phi] \equiv \int d[\Phi_{\text{E}}] e^{-I_g^{(\text{E})}[\phi, \Phi]} = e^{-I_g^{(\text{E})}[\phi, \tilde{\Phi}_g]} \quad (3.42)$$

$$W_g[\phi] \equiv I_g^{(\text{E})}[\phi, \tilde{\Phi}_g] = \int (d^4x)_{\text{E}} \left(-\frac{1}{2} (\partial_\mu \phi \partial^\mu \phi) - g^2 \cdot \frac{\alpha^2}{2m^2 \Lambda^2} (\partial_\mu \phi \partial^\mu \phi)^2 \right), \quad (3.43)$$

where $\tilde{\Phi}_g$ is the classical solution of $I_g^{(\text{E})}$ and is calculated as

$$\tilde{\Phi}_g = g \cdot \frac{\alpha}{m^2 \Lambda} (\partial_\mu \phi \partial^\mu \phi). \quad (3.44)$$

By defining the probability distribution

$$P_g[\Phi] \equiv \frac{e^{-I_g^{(\text{E})}[\phi, \Phi]}}{Z_g[\phi]}, \quad (3.45)$$

the expectation value of the interacting term is calculated as

$$\begin{aligned} \langle I_1 \rangle_{g=0} &\equiv \int d[\Phi_{\text{E}}] P_0[\Phi_{\text{E}}] I_1^{(\text{E})}[\phi, \Phi_{\text{E}}] \\ &= I_1^{(\text{E})}[\phi, \tilde{\Phi}_{g=0}] \\ &= I_1^{(\text{E})}[\phi, 0] = 0. \end{aligned} \quad (3.46)$$

From Eqs. (2.8) and (3.43), the relative entropy between P_0 and P_g is calculated as follows:

$$\begin{aligned} S(P_0 || P_g) &\equiv \int d[\Phi_{\text{E}}] (P_0 \ln P_0 - P_0 \ln P_g) \\ &= -\ln Z_0[\phi] + \ln Z_g[\phi] + g \cdot \int d[\Phi_{\text{E}}] P_0[\Phi_{\text{E}}] I_1^{(\text{E})}[\phi, \Phi_{\text{E}}] \\ &= -\ln Z_0[\phi] + \ln Z_g[\phi] \\ &= W_0[\phi] - W_g[\phi] \\ &= g^2 \cdot \frac{\alpha^2}{2m^2 \Lambda^2} \int (d^4x)_{\text{E}} (\partial_\mu \phi \partial^\mu \phi)^2 \geq 0. \end{aligned} \quad (3.47)$$

The first line is the definition of the relative entropy, and in the third line, $\langle I_1 \rangle_{g=0} = 0$ is used. By taking to be $g = 1$, the relative entropy between P_{NI} and P_{T} is given by

$$S(P_{\text{NI}} || P_{\text{T}}) = W_{\text{NI}}[\phi] - W_{\text{T}}[\phi] = \frac{\alpha^2}{2m^2 \Lambda^2} \int (d^4x)_{\text{E}} (\partial_\mu \phi \partial^\mu \phi)^2 \geq 0 \Rightarrow \frac{\alpha^2}{2m^2 \Lambda^2} \geq 0, \quad (3.48)$$

where $P_{\text{NI}} = P_0$, $P_{\text{T}} = P_{g=1}$, $W_{\text{NI}} = W_0$, and $W_{\text{T}} = W_{g=1}$ are used. We see that this result is consistent with the upper bound of Eqs. (2.52) and (2.86). In this target theory, $W_{\text{T}} - W_{\text{NI}}$ denotes the dimension-eight operator, and the non-negativity of the relative entropy yields a constraint on its Wilson coefficient.

- **Infinite heavy mass reference theory** — The action of the IHMRT in the Euclidean space is expressed as follows:

$$I_{\text{IH}}^{(\text{E})}[\phi, \Phi] \equiv \lim_{m \rightarrow \infty} I_{\text{T}}[\phi, \Phi]. \quad [\text{Reference}] \quad (3.49)$$

The classical solution of I_{IH} satisfies $\tilde{\Phi}_{\text{IH}} = 0$. Using the saddle point approximations, at the tree level, we obtain

$$Z_{\text{IH}}[\phi] \equiv \int d[\Phi_{\text{E}}] e^{-I_{\text{IH}}^{(\text{E})}[\phi, \Phi]} = e^{-I_{\text{IH}}^{(\text{E})}[\phi, 0]}, \quad (3.50)$$

$$W_{\text{IH}}[\phi] \equiv -\ln Z_{\text{IH}}[\phi] = I_{\text{IH}}^{(\text{E})}[\phi, 0], \quad (3.51)$$

$$\langle I_{\text{T}} - \lim_{m \rightarrow \infty} I_{\text{T}} \rangle_{\text{IH}} = 0. \quad (3.52)$$

From Eqs. (3.43) and (3.51), the difference of the Euclidean effective action is given by

$$W_{\text{T}}[\phi] - W_{\text{IH}}[\phi] = -\frac{\alpha^2}{2m^2\Lambda^2} \int (d^4x)_{\text{E}} (\partial_\mu \phi \partial^\mu \phi)^2 \leq 0 \Rightarrow \frac{\alpha^2}{2m^2\Lambda^2} \geq 0, \quad (3.53)$$

where $W_{\text{T}} = W_{g=1}$ and Eq. (3.51) are used. The above inequality represents Eq. (2.95) and is consistent with the non-negativity of the relative entropy. Therefore, even in the IHMRT, the non-negativity of the relative entropy yields the constraint on the Wilson coefficient.

3.4 A tree level UV completion of single massless scalar field theory with linear term

Consider the theory of Eq. (3.31) with a linear term of Φ in the Minkowski space:

$$I_{\text{T}}[\phi, \Phi] \equiv \int d^4x \left(\frac{1}{2} (\partial_\mu \phi \partial^\mu \phi) + \frac{1}{2} (\partial_\mu \Phi \partial^\mu \Phi) - \frac{m^2}{2} (\Phi - v)^2 + \frac{\alpha}{\Lambda} \cdot \Phi (\partial_\mu \phi \partial^\mu \phi) \right), \quad [\text{Target}] \quad (3.54)$$

where v denotes a dimensionful parameter. The action in the Euclidean space is expressed as

$$\begin{aligned} I_{\text{T}}^{(\text{E})}[\phi, \Phi] &= \int (d^4x)_{\text{E}} \left(\frac{1}{2} (\partial_I \phi_{\text{E}} \partial_I \phi_{\text{E}}) + \frac{1}{2} (\partial_I \Phi_{\text{E}} \partial_I \Phi_{\text{E}}) + \frac{m^2}{2} (\Phi_{\text{E}} - v)^2 + \frac{\alpha}{\Lambda} \cdot \Phi_{\text{E}} (\partial_I \phi_{\text{E}} \partial_I \phi_{\text{E}}) \right), \\ &= \int (d^4x)_{\text{E}} \left(-\frac{1}{2} (\partial_\mu \phi \partial^\mu \phi) + \frac{1}{2} (\partial_I \Phi_{\text{E}} \partial_I \Phi_{\text{E}}) + \frac{m^2}{2} (\Phi_{\text{E}} - v)^2 - \frac{\alpha}{\Lambda} \cdot \Phi_{\text{E}} (\partial_\mu \phi \partial^\mu \phi) \right). \end{aligned} \quad (3.55)$$

Similar to the previous example, ϕ is the background field. As discussed in Appendix A, the background field of the Euclidean space is defined from that of the Minkowski space. The actions I_0 and I_{I} in the Euclidean space are respectively defined as,

$$I_0^{(\text{E})}[\phi, \Phi] \equiv I_{\text{T}}^{(\text{E})}[0, \Phi] + \left(I_{\text{T}}^{(\text{E})}[\phi, 0] - \int (d^4x)_{\text{E}} \frac{m^2 v^2}{2} \right),$$

$$= \int (d^4x)_E \left(-\frac{1}{2}(\partial_\mu \phi \partial^\mu \phi) + \frac{1}{2}(\partial_I \Phi_E \partial_I \Phi_E) + \frac{m^2}{2}(\Phi_E - v)^2 \right), \quad (3.56)$$

$$I_I^{(E)}[\phi, \Phi] \equiv I_T^{(E)}[\phi, \Phi] - I_0^{(E)}[\phi, \Phi] = -\frac{\alpha}{\Lambda} \cdot \int (d^4x)_E \Phi_E (\partial_\mu \phi \partial^\mu \phi). \quad (3.57)$$

The partition function and effective action are defined as

$$Z_T[\phi] \equiv \int d[\Phi_E] e^{-I_T^{(E)}[\phi, \Phi_E]}, \quad (3.58)$$

$$W_T[\phi] \equiv -\ln Z_T[\phi]. \quad (3.59)$$

For each reference theory in Sec. 2.2, we consider the constraints on the Euclidean effective action in the following way.

- **Non-interacting reference theory** — The action of NIRT in the Euclidean space is defined as follows:

$$I_{NI}^{(E)}[\phi, \Phi] \equiv \int (d^4x)_E \left(-\frac{1}{2}(\partial_\mu \phi \partial^\mu \phi) + \frac{1}{2}(\partial_I \Phi_E \partial_I \Phi_E) + \frac{m^2}{2}(\Phi_E^2 - v)^2 \right) \cdot [\text{Reference}] \quad (3.60)$$

Note here that the linear term of Φ_E arises from the third term of the right-hand side. The solution of the equation of motion of $I_{NI}^{(E)}$ for Φ_E is calculated as $\tilde{\Phi}_{NI} = v$. The partition function and effective action of the NIRT are respectively defined as follows:

$$Z_{NI}[\phi] \equiv \int d[\Phi_E] e^{-I_{NI}^{(E)}[\phi, \Phi_E]}, \quad (3.61)$$

$$W_{NI}[\phi] \equiv -\ln Z_{NI}[\phi]. \quad (3.62)$$

By introducing the parameter g , we define

$$I_g^{(E)}[\phi, \Phi] \equiv I_0^{(E)}[\phi, \Phi] + g \cdot I_I^{(E)}[\phi, \Phi]. \quad (3.63)$$

The target and reference theories are given by,

$$I_T^{(E)}[\phi, \Phi] = \lim_{g \rightarrow 1} I_g^{(E)}[\phi, \Phi], \quad I_{NI}^{(E)}[\phi, \Phi] = \lim_{g \rightarrow 0} I_g^{(E)}[\phi, \Phi]. \quad (3.64)$$

The partition function and effective action of $I_g^{(E)}$ are respectively calculated as follows:

$$Z_g[\phi] \equiv \int d[\Phi_E] e^{-I_g^{(E)}[\phi, \Phi_E]} = e^{-I_g^{(E)}[\phi, \tilde{\Phi}_g]}, \quad (3.65)$$

$$W_g[\phi] \equiv -\ln Z_g[\phi] = \int (d^4x)_E \left(-\frac{1}{2}(\partial_\mu \phi \partial^\mu \phi) - g^2 \cdot \frac{\alpha^2}{2m^2 \Lambda^2} (\partial_\mu \phi \partial^\mu \phi)^2 - g \cdot \frac{\alpha}{\Lambda} \cdot v (\partial_\mu \phi \partial^\mu \phi) \right), \quad (3.66)$$

where $\tilde{\Phi}_g$ is the classical solution of $I_g^{(E)}$ and is calculated as follows:

$$\tilde{\Phi}_g = v + g \cdot \frac{\alpha}{m^2 \Lambda} (\partial_\mu \phi \partial^\mu \phi). \quad (3.67)$$

By defining the probability distribution function as

$$P_g[\Phi] \equiv \frac{e^{-I_g^{(E)}[\phi, \Phi]}}{Z_g[\phi]}, \quad (3.68)$$

the expectation value of the interacting term is calculated as

$$\begin{aligned} \langle I_I \rangle_{g=0} &= \int d[\Phi_E] P_0[\Phi_E] I_I^{(E)}[\phi, \Phi_E] \\ &= I_I^{(E)}[\phi, \tilde{\Phi}_0] \\ &= -\frac{\alpha}{\Lambda} \cdot \int (d^4x)_E v(\partial_\mu \phi \partial^\mu \phi). \end{aligned} \quad (3.69)$$

From Eqs. (2.8), (3.66), and (3.69), the relative entropy between P_0 and P_g is calculated as follows:

$$\begin{aligned} S(P_0||P_g) &\equiv \int d[\Phi_E] (P_0[\Phi_E] \ln P_0[\Phi_E] - P_0[\Phi_E] \ln P_g[\Phi_E]) \\ &= -\ln Z_0[\phi] + \ln Z_g[\phi] + g \cdot \int d[\Phi_E] P_g[\Phi_E] I_I^{(E)}[\phi, \Phi_E] \\ &= -\ln Z_0[\phi] + \ln Z_g[\phi] + g \cdot \langle I_I \rangle_{g=0} \\ &= W_0[\phi] - W_g[\phi] - g \cdot \frac{\alpha}{\Lambda} \cdot \int (d^4x)_E v(\partial_\mu \phi \partial^\mu \phi) \\ &= g^2 \cdot \frac{\alpha^2}{2m^2 \Lambda^2} \int (d^4x)_E (\partial_\mu \phi \partial^\mu \phi)^2 \geq 0, \end{aligned} \quad (3.70)$$

where the first line is the definition of the relative entropy, Eq. (3.69) was used in the third line, and $g \cdot \langle I_I \rangle_{g=0}$ cancels in the last line. By taking to be $g = 1$, the relative entropy between P_{NI} and P_{T} is given by

$$S(P_{\text{NI}}||P_{\text{T}}) = W_{\text{NI}}[\phi] - W_{\text{T}}[\phi] + \langle I_I \rangle_{\text{NI}} = \frac{\alpha^2}{2m^2 \Lambda^2} \int (d^4x)_E (\partial_\mu \phi \partial^\mu \phi)^2 \geq 0 \Rightarrow \frac{\alpha^2}{2m^2 \Lambda^2} \geq 0, \quad (3.71)$$

where $P_{\text{NI}} = P_0$, $P_{\text{T}} = P_{g=1}$, $W_{\text{NI}} = W_0$, $W_{\text{T}} = W_{g=1}$, and $\langle I_I \rangle_{\text{NI}} = \langle I_I \rangle_{g=0}$ are used. The above inequality yields a constraint on the Wilson coefficient of the dimension-eight operator, which is consistent with the non-negativity of the relative entropy.

We also show that the above result holds even if $\langle I_I \rangle_{\text{NI}}$ is eliminated by a redefinition of Φ . We have already seen this fact in Sec. 2.3.1 generically. By defining a new field η as $\Phi \equiv \eta + v$, the action of Eq. (3.55) is expressed as,

$$\begin{aligned} I_{\text{T}}'^{(E)}[\phi, \eta] &\equiv I_{\text{T}}^{(E)}[\phi, \eta + v] \\ &= \int (d^4x)_E \left(-\frac{1}{2}(\partial_\mu \phi \partial^\mu \phi) + \frac{1}{2}(\partial_I \eta_E \partial_I \eta_E) + \frac{1}{2}m^2 \eta_E^2 \right) \end{aligned}$$

$$-\frac{\alpha}{\Lambda} \cdot v(\partial_\mu \phi \partial^\mu \phi) - \frac{\alpha}{\Lambda} \cdot \eta_E(\partial_\mu \phi \partial^\mu \phi) \Big). \quad [\mathbf{Target}] \quad (3.72)$$

Note here that the liner term of η does not arise. Define the actions I'_0 and I'_1 in the Euclidean space as follows:

$$I'^{(E)}_0[\phi, \eta] \equiv \int (d^4x)_E \left(-\frac{1}{2}(\partial_\mu \phi \partial^\mu \phi) + \frac{1}{2}(\partial_I \eta_E \partial_I \eta_E) + \frac{m^2}{2} \eta_E^2 - \frac{\alpha}{\Lambda} \cdot v(\partial_\mu \phi \partial^\mu \phi) \right), \quad (3.73)$$

$$I'^{(E)}_1[\phi, \eta] \equiv -\frac{\alpha}{\Lambda} \cdot \int (d^4x)_E \eta_E (\partial_\mu \phi \partial^\mu \phi). \quad (3.74)$$

The partition function and effective action of $I'^{(E)}_T$ are defined as

$$Z'_T[\phi] \equiv \int d[\eta_E] e^{-I'^{(E)}_T[\phi, \eta_E]}, \quad (3.75)$$

$$W'_T[\phi] \equiv -\ln Z'_T[\phi]. \quad (3.76)$$

The action of NIRT for $I'^{(E)}_T$ is defined as follows:

$$\begin{aligned} I'^{(E)}_{\text{NI}}[\phi, \eta] &\equiv I'^{(E)}_0[\phi, \eta] \\ &= \int (d^4x)_E \left(-\frac{1}{2}(\partial_\mu \phi \partial^\mu \phi) + \frac{1}{2}(\partial_I \eta_E \partial_I \eta_E) + \frac{1}{2} m^2 \eta_E^2 - \frac{\alpha}{\Lambda} \cdot \eta_E (\partial_\mu \phi \partial^\mu \phi) \right), \quad [\mathbf{Reference}] \end{aligned} \quad (3.77)$$

where the liner term of η does not arise. The partition function and effective action of the NIRT are respectively defined as follows:

$$Z'_{\text{NI}}[\phi] \equiv \int d[\eta_E] e^{-I'^{(E)}_{\text{NI}}[\phi, \eta_E]}, \quad (3.78)$$

$$W'_{\text{NI}}[\phi] \equiv -\ln Z'_{\text{NI}}[\phi]. \quad (3.79)$$

By introducing the parameter g , we define

$$I'^{(E)}_g[\phi, \eta] \equiv I'^{(E)}_0[\phi, \eta] + g \cdot I'^{(E)}_1[\phi, \eta]. \quad (3.80)$$

The target and reference theories are expressed as follows:

$$I'^{(E)}_T[\phi, \eta] = \lim_{g \rightarrow 1} I'^{(E)}_g[\phi, \eta], \quad I'^{(E)}_{\text{NI}}[\phi, \eta] = \lim_{g \rightarrow 0} I'^{(E)}_g[\phi, \eta]. \quad (3.81)$$

The partition function and effective action of $I'^{(E)}_g$ are respectively calculated as follows:

$$Z'_g[\phi] \equiv \int d[\eta_E] e^{-I'^{(E)}_g[\phi, \eta_E]} = e^{-I'^{(E)}_g[\phi, \tilde{\eta}_g]}, \quad (3.82)$$

$$W'_g[\phi] \equiv I'^{(E)}_g[\phi, \tilde{\eta}_g] = \int (d^4x)_E \left(-\frac{1}{2} \left(1 + 2 \cdot \frac{\alpha}{\Lambda} \cdot v \right) (\partial_\mu \phi \partial^\mu \phi) - g^2 \cdot \frac{\alpha^2}{2\Lambda^2 m^2} (\partial_\mu \phi \partial^\mu \phi)^2 \right), \quad (3.83)$$

where $\tilde{\eta}_g$ is a classical solution of $I'_g{}^{(\text{E})}$ and is given by

$$\tilde{\eta}_g = g \cdot \frac{\alpha}{m^2 \Lambda} (\partial_\mu \phi \partial^\mu \phi). \quad (3.84)$$

By defining the probability distribution function as

$$P'_g[\eta] \equiv \frac{e^{-I'_g{}^{(\text{E})}[\phi, \eta]}}{Z'_g[\phi]}, \quad (3.85)$$

the expectation value of the interacting term is calculated as

$$\begin{aligned} \langle I_1 \rangle_{g=0} &= \int d[\eta_E] P'_0[\eta_E] I_1^{(\text{E})}[\phi, \eta_E] \\ &= I_1^{(\text{E})}[\phi, \tilde{\eta}_0] \\ &= 0. \end{aligned} \quad (3.86)$$

From Eqs. (2.8), (3.83), and (3.86), the relative entropy between P_0 and P_g is calculated as follows:

$$\begin{aligned} S(P'_0 || P'_g) &\equiv \int d[\eta_E] (P'_0[\eta_E] \ln P'_0[\eta_E] - P'_0[\eta_E] \ln P'_g[\eta_E]) \\ &= -\ln Z'_0[\phi] + \ln Z'_g[\phi] + g \cdot \int d[\eta_E] P_0[\eta_E] I_1^{(\text{E})}[\phi, \eta_E] \end{aligned} \quad (3.87)$$

$$= -\ln Z'_0[\phi] + \ln Z'_g[\phi] \quad (3.88)$$

$$\begin{aligned} &= W'_0[\phi] - W'_g[\phi] \\ &= g^2 \cdot \frac{\alpha^2}{2\Lambda^2 m^2} \int (d^4 x)_E (\partial_\mu \phi \partial^\mu \phi)^2 \geq 0, \end{aligned} \quad (3.89)$$

where the first line is the definition of the relative entropy, and $\langle I_1 \rangle_{g=0} = 0$ is used in the third line. By taking to be $g = 1$, the relative entropy between $P'_{\text{NI}} \equiv P'_0$ and $P'_T \equiv P'_{g=1}$ is given by

$$S(P'_{\text{NI}} || P'_T) = W'_{\text{NI}}[\phi] - W'_T[\phi] = \frac{\alpha^2}{2\Lambda^2 m^2} \int (d^4 x)_E (\partial_\mu \phi \partial^\mu \phi)^2 \geq 0 \Rightarrow \frac{\alpha^2}{2\Lambda^2 m^2} \geq 0, \quad (3.90)$$

where $W'_{\text{NI}} \equiv W'_0$, and $W'_T \equiv W'_{g=1}$ are used. This result is the same as Eq. (3.71), and we found that Eq. (2.9) is invariant under the field redefinition to remove the linear term of Φ . Therefore, it is found that the constraint on the EFT does not depend on the condition of vanishing the linear term. As explained in the details in Sec. 2.3.1, this is because the linear term proportional to g cancels in the relative entropy as shown in Eq. (2.60).

- **Massive free field reference theory** — The action of the MFFRT in the Euclidean space is defined as follows:

$$I_{\text{MF}}^{(\text{E})}[\phi, \Phi] \equiv \int (d^4 x)_E \left(-\frac{1}{2} (\partial_\mu \phi \partial^\mu \phi) + \frac{1}{2} (\partial_I \Phi_E \partial_I \Phi_E) + \frac{m^2}{2} \Phi_E^2 + \frac{m^2 v^2}{2} \right) \cdot [\text{Reference}] \quad (3.91)$$

Note here that the MFFRT does not include self-interacting terms of Φ_E . The solution of the equation of motion of $I_{\text{MF}}^{(\text{E})}$ for Φ_E is calculated as $\tilde{\Phi}_{\text{MF}} = 0$. The partition function and effective action of the MFFRT are respectively defined as follows:

$$Z_{\text{MF}}[\phi] \equiv \int d[\Phi_E] e^{-I_{\text{MF}}^{(\text{E})}[\phi, \Phi_E]}, \quad (3.92)$$

$$W_{\text{MF}}[\phi] \equiv -\ln Z_{\text{MF}}[\phi]. \quad (3.93)$$

By introducing the parameter g , we define

$$J_g^{(\text{E})}[\phi, \Phi] \equiv I_0^{(\text{E})}[\phi, 0] + I_\Phi^{(\text{E})}[\Phi] + g \cdot \left(I_S^{(\text{E})}[\Phi] + I_I^{(\text{E})}[\phi, \Phi] \right), \quad (3.94)$$

with

$$I_\Phi^{(\text{E})}[\Phi] \equiv \int (d^4x)_E \left(\frac{1}{2} (\partial_I \Phi_E \partial_I \Phi_E) + \frac{m^2}{2} \Phi_E^2 \right), \quad (3.95)$$

$$I_S^{(\text{E})}[\Phi] \equiv -v \cdot m^2 \int (d^4x)_E \Phi_E. \quad (3.96)$$

The target and reference theories are expressed as follows:

$$I_T^{(\text{E})}[\phi, \Phi] = \lim_{g \rightarrow 1} J_g^{(\text{E})}[\phi, \Phi], \quad I_{\text{MF}}^{(\text{E})}[\phi, \Phi] = \lim_{g \rightarrow 0} J_g^{(\text{E})}[\phi, \Phi]. \quad (3.97)$$

The partition function and effective action of $J_g^{(\text{E})}$ are respectively calculated as follows:

$$z_g[\phi] \equiv \int d[\Phi_E] e^{-J_g^{(\text{E})}[\phi, \Phi_E]} = e^{-J_g^{(\text{E})}[\phi, \tilde{\Phi}_g]}, \quad (3.98)$$

$$w_g[\phi] \equiv -\ln z_g[\phi]$$

$$= \int (d^4x)_E \left(-\frac{1}{2} (\partial_\mu \phi \partial^\mu \phi) - g^2 \cdot \frac{\alpha^2}{2m^2 \Lambda^2} (\partial_\mu \phi \partial^\mu \phi)^2 - g^2 \cdot \frac{\alpha}{\Lambda} \cdot v (\partial_\mu \phi \partial^\mu \phi) + (1 - g^2) \cdot \frac{m^2 v^2}{2} \right), \quad (3.99)$$

where $\tilde{\Phi}_g$ is the classical solution of $J_g^{(\text{E})}$ and is given as

$$\tilde{\Phi}_g = g \cdot v + g \cdot \frac{\alpha}{m^2 \Lambda} (\partial_\mu \phi \partial^\mu \phi). \quad (3.100)$$

By defining the probability distribution function as

$$p_g[\Phi] \equiv \frac{e^{-J_g^{(\text{E})}[\phi, \Phi]}}{z_g[\phi]}, \quad (3.101)$$

we obtain

$$\begin{aligned} \langle I_S + I_I \rangle_{g=0} &\equiv \int d[\Phi_E] p_0[\Phi_E] (I_S[\Phi_E] + I_I[\phi, \Phi_E]) \\ &= I_S[\tilde{\Phi}_0] + I_I[\phi, \tilde{\Phi}_0] \end{aligned}$$

$$= 0, \quad (3.102)$$

where $\tilde{\Phi}_0 = 0$ holds from Eq. (3.100). From Eqs. (2.8), (3.99), and (3.102), the relative entropy between p_0 and p_g is calculated as follows:

$$\begin{aligned} S(p_0||p_g) &\equiv \int d[\Phi_E] (p_0[\Phi_E] \ln p_0[\Phi_E] - p_0[\Phi_E] \ln p_g[\Phi_E]) \\ &= -\ln z_0[\phi] + \ln z_g[\phi] + g \cdot \int d[\Phi_E] p_0[\Phi_E] (I_S[\Phi_E] + I_I[\phi, \Phi_E]) \\ &= -\ln z_0[\phi] + \ln z_g[\phi] \\ &= w_0[\phi] - w_g[\phi] \\ &= g^2 \cdot \frac{1}{2m^2} \int (d^4x)_E \left(\frac{\alpha}{\Lambda} (\partial_\mu \phi \partial^\mu \phi) + m^2 v^2 \right)^2 \geq 0. \end{aligned} \quad (3.103)$$

By taking to be $g = 1$, the relative entropy between P_{MF} and P_{T} is given by

$$S(P_{\text{MF}}||P_{\text{T}}) = W_{\text{MF}}[\phi] - W_{\text{T}}[\phi] = \frac{1}{2m^2} \int (d^4x)_E \left(\frac{\alpha}{\Lambda} (\partial_\mu \phi \partial^\mu \phi) + m^2 v^2 \right)^2 \geq 0, \quad (3.104)$$

where $P_{\text{MF}} = p_0$, $P_{\text{T}} = p_{g=1}$, $W_{\text{MF}} = w_0$, and $W_{\text{T}} = w_1$ are used. This inequality denotes Eq. (2.86) and is consistent with the non-negativity of relative entropy. Because of the self-interacting term of Φ , the shift of the Euclidean effective action includes the term independent of ϕ .

3.5 Neutral bosons interacting with photon

We consider heavy neutral bosons such as the dilaton and the axion. For each model, we evaluate the relative entropy in the following way.

3.5.1 Dilaton

The action of the dilaton in the Minkowski space is expressed as

$$I_{\text{T}}[A, \phi] \equiv \int d^4x \left[-\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{1}{2} (\partial_\mu \phi)^2 - \frac{m_\phi^2}{2} \phi^2 + \frac{\phi}{f_\phi} F_{\mu\nu} F^{\mu\nu} \right], \quad [\text{Target}] \quad (3.105)$$

where m_ϕ and f_ϕ are the mass and the decay constant of the heavy neutral scalar boson, respectively, and $F_{\mu\nu}$ is the field strength of photon field defined by $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$. Based on the procedure in Appendix A, the action in the Euclidean space is obtained as

$$\begin{aligned} I_{\text{T}}^{(\text{E})}[A, \phi_E] &= \int (d^4x)_E \left(\frac{1}{4} F_{E,IJ} F_{E,IJ} + \frac{1}{2} \partial_I \phi_E \partial_I \phi_E + \frac{m_\phi^2}{2} (\phi_E)^2 - \frac{\phi_E}{f_\phi} F_{E,IJ} F_{E,IJ} \right), \\ &= \int (d^4x)_E \left(\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{1}{2} \partial_I \phi_E \partial_I \phi_E + \frac{m_\phi^2}{2} (\phi_E)^2 - \frac{\phi_E}{f_\phi} F_{\mu\nu} F^{\mu\nu} \right), \end{aligned} \quad (3.106)$$

where $F_{\mu\nu}$ is a background field in the Minkowski space. Then, I_0 and I_I are defined as follows:

$$I_0^{(E)}[A, \phi_E] \equiv I_T^{(E)}[A, 0] + I_T^{(E)}[0, \phi_E] = \int (d^4x)_E \left(\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{1}{2} \partial_I \phi_E \partial_I \phi_E + \frac{m_\phi^2}{2} (\phi_E)^2 \right), \quad (3.107)$$

$$I_I^{(E)}[A, \phi_E] \equiv I_T^{(E)}[A, \phi_E] - I_0[A, \phi_E] = \int (d^4x)_E \left(-\frac{\phi_E}{f_\phi} F_{\mu\nu} F^{\mu\nu} \right). \quad (3.108)$$

The partition function and effective action are defined as

$$Z_T[A] \equiv \int d[\phi_E] e^{-I_T^{(E)}[A, \phi_E]}, \quad (3.109)$$

$$W_T[A] \equiv -\ln Z_T[A], \quad (3.110)$$

In this target theory, the NIRT is the same as the MFFRT, and the action of the NIRT and MFFRT in the Euclidean space is defined as

$$I_{NI}^{(E)}[A, \phi_E] \equiv \int (d^4x)_E \left(\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{1}{2} \partial_I \phi_E \partial_I \phi_E + \frac{m_\phi^2}{2} (\phi_E)^2 \right). \quad [\textbf{Reference}] \quad (3.111)$$

The solution of the equation of motion of $I_{NI}^{(E)}$ is calculated as $\tilde{\phi}_{NI} = 0$. The partition function and effective action of the reference theory are respectively defined as follows:

$$Z_{NI}[A] \equiv \int d[\phi_E] e^{-I_{NI}^{(E)}[A, \phi_E]}, \quad (3.112)$$

$$W_{NI}[A] \equiv -\ln Z_{NI}[A]. \quad (3.113)$$

By introducing the parameter g , we define

$$I_g^{(E)}[A, \phi] \equiv I_0^{(E)}[A, \phi] + g \cdot I_I^{(E)}[A, \phi]. \quad (3.114)$$

The target theory and reference theories are expressed as follows:

$$I_T^{(E)}[A, \phi] = \lim_{g \rightarrow 1} I_g^{(E)}[A, \phi], \quad I_{NI}^{(E)}[A, \phi] = \lim_{g \rightarrow 0} I_g^{(E)}[A, \phi]. \quad (3.115)$$

The partition function and effective action of $I_g^{(E)}$ are respectively calculated as follows:

$$Z_g[A] \equiv \int d[\phi_E] e^{-I_g^{(E)}[A, \phi_E]} = e^{-I_g^{(E)}[A, \tilde{\phi}_g]}, \quad (3.116)$$

$$W_g[A] \equiv -\ln Z_g[A] = \int (d^4x)_E \left(\frac{1}{4} F_{\mu\nu} F^{\mu\nu} - g^2 \cdot \frac{1}{2f_\phi^2 m_\phi^2} (F_{\mu\nu} F^{\mu\nu})^2 \right), \quad (3.117)$$

where the solution of the equation of motion of $I_g^{(E)}$ for ϕ_E with the heavy mass is calculated as

$$\tilde{\phi}_g = g \cdot \frac{1}{f_\phi m_\phi^2} F_{\mu\nu} F^{\mu\nu}. \quad (3.118)$$

By defining the probability distribution function

$$P_g[\phi] \equiv \frac{e^{-I_g^{(\text{E})}[A, \phi]}}{Z_g[A]}, \quad (3.119)$$

we obtain

$$\begin{aligned} \langle I_1 \rangle_{g=0} &= \int d[\phi_E] P_0[\phi_E] I_1[A, \phi_E] \\ &= I_1[A, \tilde{\phi}_0] \\ &= 0, \end{aligned} \quad (3.120)$$

where $\tilde{\phi}_0 = 0$ holds from Eq. (3.118). From Eqs. (2.8), (3.117), and (3.118), the relative entropy between P_0 and P_g is calculated as follows:

$$\begin{aligned} S(P_0||P_g) &\equiv \int d[\phi_E] (P_0[\phi_E] \ln P_0[\phi_E] - P_0[\phi_E] \ln P_g[\phi_E]) \\ &= -\ln Z_0[A] + \ln Z_g[A] + g \cdot \int d[\phi_E] P_0[\phi_E] I_1[A, \phi_E] \\ &= -\ln Z_0[A] + \ln Z_g[A] \\ &= W_0[A] - W_g[A] \\ &= g^2 \cdot \frac{1}{2f_\phi^2 m_\phi^2} \int (d^4x)_E (F_{\mu\nu} F^{\mu\nu})^2 \geq 0, \end{aligned} \quad (3.121)$$

where the first line is the definition of the relative entropy, Eq. (3.120) is used in the third line, the fourth line is the definition of the effective action, and Eq. (3.117) and the non-negativity of the relative entropy are used in the last line. By taking to be $g = 1$, the relative entropy between P_{NI} and P_{T} is given by

$$S(P_{\text{NI}}||P_{\text{T}}) = W_{\text{NI}}[A] - W_{\text{T}}[A] = \frac{1}{2f_\phi^2 m_\phi^2} \int (d^4x)_E (F_{\mu\nu} F^{\mu\nu})^2 \geq 0 \Rightarrow \frac{1}{2f_\phi^2 m_\phi^2} \geq 0, \quad (3.122)$$

where $P_{\text{NI}} = P_0$, $P_{\text{T}} = P_{g=1}$, $W_{\text{NI}} = W_0$, and $W_{\text{T}} = W_{g=1}$ are used. The relative entropy denotes the dimension-eight term and yields the constraints on the Wilson coefficients of the dimension-eight operator.

3.5.2 Axion

The action of the axion in the Minkowski space is expressed as

$$I_{\text{T}}[A, a] \equiv \int d^4x \left[-\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{1}{2} (\partial_\mu a)^2 - \frac{m_a^2}{2} a^2 + \frac{a}{f_a} F_{\mu\nu} \tilde{F}^{\mu\nu} \right], \quad [\text{Target}] \quad (3.123)$$

where m_a and f_a are the mass and the decay constant of the heavy neutral pseudo-scalar boson, respectively, and the dual field strength is defined as $\tilde{F}^{\mu\nu} = \frac{1}{2} \epsilon^{\mu\nu\rho\sigma} F_{\rho\sigma}$. The action in the Euclidean space is given by

$$I_{\text{T}}^{(\text{E})}[A, a_E] = \int (d^4x)_E \left(\frac{1}{4} F_{E,IJ} F_{E,IJ} + \frac{1}{2} \partial_I a_E \partial_I a_E + \frac{m_a^2}{2} (a_E)^2 - i \frac{a_E}{f_a} F_{E,IJ} \tilde{F}_{E,IJ} \right),$$

$$= \int (d^4x)_E \left(\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{1}{2} \partial_I a_E \partial_I a_E + \frac{m_a^2}{2} (a_E)^2 - \frac{a_E}{f_a} F_{\mu\nu} \tilde{F}^{\mu\nu} \right), \quad (3.124)$$

where the background field of the Euclidean space $F_{E,IJ}$ is defined from that of the Minkowski space; see Appendix A. We define I_0 and I_I as follows:

$$I_0^{(E)}[A, a_E] \equiv I_T^{(E)}[A, 0] + I_T^{(E)}[0, a_E] = \int (d^4x)_E \left(\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{1}{2} \partial_I a_E \partial_I a_E + \frac{m_a^2}{2} (a_E)^2 \right), \quad (3.125)$$

$$I_I^{(E)}[A, a_E] \equiv I_T^{(E)}[A, a_E] - I_0[A, a_E] = \int (d^4x)_E \left(-\frac{a_E}{f_a} F_{\mu\nu} \tilde{F}^{\mu\nu} \right). \quad (3.126)$$

The partition function and effective action are defined as

$$Z_T[A] \equiv \int d[a_E] e^{-I_T^{(E)}[A, a_E]}, \quad (3.127)$$

$$W_T[A] \equiv -\ln Z_T[A]. \quad (3.128)$$

The action of the NIRT and MFFRT in the Euclidean space is defined as

$$I_{\text{NI}}^{(E)}[A, a_E] \equiv \int (d^4x)_E \left(\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{1}{2} \partial_I a_E \partial_I a_E + \frac{m_a^2}{2} (a_E)^2 \right). \quad [\text{Reference}] \quad (3.129)$$

The solution of the equation of motion of $I_{\text{NI}}^{(E)}$ is calculated as $\tilde{a}_{\text{NI}} = 0$. The partition function and effective action of the reference theory are defined as

$$Z_{\text{NI}}[A] \equiv \int d[a_E] e^{-I_{\text{NI}}^{(E)}[A, a_E]}, \quad (3.130)$$

$$W_{\text{NI}}[A] \equiv -\ln Z_{\text{NI}}[A]. \quad (3.131)$$

By introducing the parameter g , we define

$$I_g^{(E)}[A, a] \equiv I_0^{(E)}[A, a] + g \cdot I_I^{(E)}[A, a]. \quad (3.132)$$

The target and reference theories are expressed as follows:

$$I_T^{(E)}[A, a] = \lim_{g \rightarrow 1} I_g^{(E)}[A, a], \quad I_{\text{NI}}^{(E)}[A, a] = \lim_{g \rightarrow 0} I_g^{(E)}[A, a]. \quad (3.133)$$

The partition function and effective action of $I_g^{(E)}$ are respectively calculated as follows:

$$Z_g[A] \equiv \int d[a_E] e^{-I_g^{(E)}[A, a_E]} = e^{-I_g^{(E)}[A, \tilde{a}_g]}, \quad (3.134)$$

$$W_g[A] \equiv -\ln Z_g[A] = \int (d^4x)_E \left(\frac{1}{4} F_{\mu\nu} F^{\mu\nu} - g^2 \cdot \frac{1}{2f_a^2 m_a^2} (F_{\mu\nu} \tilde{F}^{\mu\nu})^2 \right), \quad (3.135)$$

where the solution of the equation of motion of $I_g^{(E)}$ for a_E is calculated as

$$\tilde{a}_g = g \cdot \frac{1}{f_a m_a^2} F_{\mu\nu} \tilde{F}^{\mu\nu}. \quad (3.136)$$

By defining the probability distribution function as

$$P_g[a] \equiv \frac{e^{-I_g^{(\text{E})}[A,a]}}{Z_g[A]}, \quad (3.137)$$

the expectation value of the interaction is calculated as follows:

$$\begin{aligned} \langle I_I \rangle_{g=0} &\equiv \int d[a_E] P_0[a_E] I_I^{(\text{E})}[A, a_E] \\ &= I_I^{(\text{E})}[A, \tilde{a}_0] \\ &= 0, \end{aligned} \quad (3.138)$$

where $\tilde{a}_0 = 0$ holds from Eq. (3.136). From Eqs. (2.8), (3.135), and (3.138), the relative entropy between P_0 and P_g is calculated as follows:

$$\begin{aligned} S(P_0||P_g) &\equiv \int d[a_E] (P_0[a_E] \ln P_0[a_E] - P_0[a_E] \ln P_g[a_E]) \\ &= -\ln Z_0[A] + \ln Z_g[A] + g \cdot \int d[a_E] P_0[a_E] I_I^{(\text{E})}[A, a_E] \\ &= -\ln Z_0[A] + \ln Z_g[A] \\ &= W_0[A] - W_g[A] \\ &= g^2 \cdot \frac{1}{2f_a^2 m_a^2} \int (d^4 x)_E (F_{\mu\nu} \tilde{F}^{\mu\nu})^2 \geq 0, \end{aligned} \quad (3.139)$$

where the first line is the definition of the relative entropy, Eq. (3.138) is used in the third line, the definition of the effective action is used in the fourth line, and Eq. (3.135) is used in the last line. By taking to be $g = 1$, the relative entropy between P_{NI} and P_{T} is given by

$$S(P_{\text{NI}}||P_{\text{T}}) = W_{\text{NI}}[A] - W_{\text{T}}[A] = \frac{1}{2f_a^2 m_a^2} \int (d^4 x)_E (F_{\mu\nu} \tilde{F}^{\mu\nu})^2 \geq 0 \Rightarrow \frac{1}{2f_a^2 m_a^2} \geq 0, \quad (3.140)$$

where $P_{\text{NI}} = P_0$, $P_{\text{T}} = P_{g=1}$, $W_{\text{NI}} = W_0$, and $W_{\text{T}} = W_{g=1}$ are used. The relative entropy denotes the dimension-eight term in Eq. (3.135), and the non-negativity of relative entropy yields the constraint on the Wilson coefficient of the dimension-eight operator.

3.6 Massless scalar field with a shift symmetry

Let us consider an action in the Minkowski space,

$$I_{\text{T}}[\phi, \psi, \bar{\psi}] \equiv \int d^4 x \left(\frac{1}{2} \partial_\mu \phi \partial^\mu \phi + \bar{\psi} (i \not{\partial} - m) \psi - \frac{c}{\Lambda} \bar{\psi} \not{\partial} \phi \gamma_5 \psi \right), \quad [\text{Target}] \quad (3.141)$$

where ψ is the heavy Dirac fermion, ϕ is the massless scalar field, c is a dimensionless coupling constant, and Λ is some energy scale. The interacting and non-interacting terms of I_{T} is given by

$$I_0[\phi, \psi, \bar{\psi}] \equiv I_{\text{T}}[\phi, 0, 0] + I_{\text{T}}[0, \psi, \bar{\psi}] = \int d^4 x \left(\frac{1}{2} \partial_\mu \phi \partial^\mu \phi + \bar{\psi} (i \not{\partial} - m) \psi \right), \quad (3.142)$$




Figure 5. Left panel: Feynman diagrams of $\langle I_1 \rangle_{g=0}$ for the massless scalar field with a shift symmetry at the one-loop level. Right panel: Feynman diagrams of $\langle I_1 \rangle_{g=0}$ for the Euler-Heisenberg theory with the charged fermion field at the one-loop level. The solid lines denote the heavy fermion field.

$$I_1[\phi, \psi, \bar{\psi}] \equiv I_T[\phi, \psi, \bar{\psi}] - I_0[\phi, \psi, \bar{\psi}] = \int d^4x \left(-\frac{c}{\Lambda} \bar{\psi} \not{\partial} \phi \gamma_5 \psi \right). \quad (3.143)$$

By introducing the parameter g , define an action $I_g \equiv I_0 + g \cdot I_1$. Then, the target theory is expressed as $I_T = \lim_{g \rightarrow 1} I_g$. To clarify the procedure of the wave function renormalization, we also perform the path integral over ϕ . By calculating one-loop diagrams and performing the Wick rotation, the partition function of I_g is obtained as

$$\begin{aligned} Z_g[\tilde{\phi}] &\equiv \int d[\phi] d[\psi] d[\bar{\psi}] e^{-I_g^{(E)}[\phi, \psi, \bar{\psi}]} \\ &= \int d[\phi] \exp \left[- \int (d^4x)_E \left(-\frac{1}{2} (1 + g^2 \cdot \alpha_0) (\partial_\mu \phi \partial^\mu \phi) - g^4 \cdot \frac{c^4}{12\pi^2 \Lambda^4} (\partial_\mu \phi \partial^\mu \phi)^2 \right) \right] \\ &= \exp \left[- \int (d^4x)_E \left(-\frac{1}{2} (1 + g^2 \cdot \alpha_0) (\partial_\mu \tilde{\phi}' \partial^\mu \tilde{\phi}') - g^4 \cdot \frac{c^4}{12\pi^2 \Lambda^4} (\partial_\mu \tilde{\phi}' \partial^\mu \tilde{\phi}')^2 \right) \right] \\ &= \exp \left[- \int (d^4x)_E \left(-\frac{1}{2} (\partial_\mu \tilde{\phi} \partial^\mu \tilde{\phi}) - g^4 \cdot \frac{c^4}{12\pi^2 \Lambda^4} (\partial_\mu \tilde{\phi} \partial^\mu \tilde{\phi})^2 \right) \right], \end{aligned} \quad (3.144)$$

where the first line is the definition of the partition function, $\tilde{\phi}'$ is a classical solution satisfying $\partial \partial \tilde{\phi}' = 0$, and the quantum correction to the kinetic term of the scalar field is eliminated in the last line by the redefinition of the field, i.e., $\tilde{\phi}' = (1 - g^2 \cdot \alpha_0/2) \tilde{\phi}$. In Eq. (3.144), two-loop effects are neglected. Also, the vacuum energy is omitted because it cancels in the relative entropy. We also note that dimension-six terms and other dimension-eight terms, i.e., $(\partial \partial \tilde{\phi})^2$ and $(\partial \partial \partial \tilde{\phi})^2$, are generated, but they are now eliminated by a background field satisfying $\partial \tilde{\phi} = \text{const.}$. From Eq. (3.144), the effective action of I_g is obtained as follows:

$$W_g[\tilde{\phi}] \equiv -\ln Z_g[\tilde{\phi}] = \int (d^4x)_E \left(-\frac{1}{2} (\partial_\mu \tilde{\phi} \partial^\mu \tilde{\phi}) - g^4 \cdot \frac{c^4}{12\pi^2 \Lambda^4} (\partial_\mu \tilde{\phi} \partial^\mu \tilde{\phi})^2 \right). \quad (3.145)$$

The Euclidean effective action of the target theory is given by

$$W_T[\tilde{\phi}] = \lim_{g \rightarrow 1} W_g[\tilde{\phi}]. \quad (3.146)$$

By taking the limit of $g = 0$, the action of the NIRT and MFFRT in the Minkowski space is defined as

$$I_{\text{NI}}[\phi, \psi, \bar{\psi}] \equiv \lim_{g \rightarrow 0} I_g[\phi, \psi, \bar{\psi}] = \int d^4x \left(\frac{1}{2} \partial_\mu \phi \partial^\mu \phi + \bar{\psi} (i \not{\partial} - m) \psi \right). \quad [\text{Reference}] \quad (3.147)$$

From Eqs. (3.144), and (3.145), the partition function and effective action of the reference theory are obtained as follows:

$$Z_{\text{NI}}[\tilde{\phi}] = \lim_{g \rightarrow 0} Z_g[\tilde{\phi}], \quad (3.148)$$

$$W_{\text{NI}}[\tilde{\phi}] = \lim_{g \rightarrow 0} W_g[\tilde{\phi}] = \int (d^4x)_{\text{E}} \frac{1}{2} (\partial_\mu \tilde{\phi} \partial^\mu \tilde{\phi}). \quad (3.149)$$

By defining a probability distribution function as $P_g[\phi, \psi, \bar{\psi}] \equiv e^{-I_g[\phi, \psi, \bar{\psi}]} / Z_g[\tilde{\phi}]$, the derivative of W_g with respect to g is calculated as follows:

$$\begin{aligned} \left(\frac{dW_g}{dg} \right)_{g=0} &= \left(\frac{\partial W_g}{\partial g} \right)_{g=0} + \int (d^4x)_{\text{E}} \left(\frac{\delta W_g}{\delta \tilde{\phi}'} \right)_{g=0} \left(\frac{d\tilde{\phi}'}{dg} \right)_{g=0} \\ &= \left(\frac{\partial W_g}{\partial g} \right)_{g=0} \\ &= \int d[\phi] d[\psi] d[\bar{\psi}] P_0[\psi, \bar{\psi}] I_{\text{I}}[\phi, \psi, \bar{\psi}] \\ &= \langle I_{\text{I}} \rangle_{g=0} = 0, \end{aligned} \quad (3.150)$$

where the partial derivative means differentiating by g while keeping $\tilde{\phi}'$, $(d\tilde{\phi}'/dg)_{g=0} = 0$ is used in the second line, and $(dW_g/dg)_{g=0}$ is used in the last line. The Feynman diagram for $\langle I_{\text{I}} \rangle_{g=0}$ is shown in the left panel of Fig. 5. From Eqs. (2.23), (3.145), and (3.150), the relative entropy between P_0 and P_g is calculated as follows:

$$\begin{aligned} S(P_0||P_g) &\equiv \int d[\phi] d[\psi] d[\bar{\psi}] (P_0[\phi, \psi, \bar{\psi}] \ln P_0[\phi, \psi, \bar{\psi}] - P_0[\phi, \psi, \bar{\psi}] \ln P_g[\phi, \psi, \bar{\psi}]) \\ &= -\ln Z_0[\tilde{\phi}] + \ln Z_g[\tilde{\phi}] + g \cdot \langle I_{\text{I}} \rangle_{g=0} \\ &= W_0[\tilde{\phi}] - W_g[\tilde{\phi}] \\ &= g^4 \cdot \frac{c^4}{12\pi^2 \Lambda^4} \int (d^4x)_{\text{E}} \left(\partial_\mu \tilde{\phi} \partial^\mu \tilde{\phi} \right)^2 \geq 0, \end{aligned} \quad (3.151)$$

where the first line is the definition of the relative entropy, Eq. (3.150) is used in the third line, and Eq. (3.145) is used in the last. By taking $g = 1$, the relative entropy between P_{NI} and P_{T} is given by

$$S(P_{\text{NI}}||P_{\text{T}}) = W_{\text{NI}}[\tilde{\phi}] - W_{\text{T}}[\tilde{\phi}] = \frac{c^4}{12\pi^2 \Lambda^4} \int (d^4x)_{\text{E}} \left(\partial_\mu \tilde{\phi} \partial^\mu \tilde{\phi} \right)^2 \geq 0 \Rightarrow \frac{c^4}{12\pi^2 \Lambda^4} \geq 0, \quad (3.152)$$

where $P_{\text{NI}} = P_0$, $P_{\text{T}} = P_{g=1}$, $W_{\text{NI}} = W_0$, and $W_{\text{T}} = W_{g=1}$ are used. Consequently, the relative entropy is non-negative in the UV action of (3.146). Although the above procedure is the top-down approach, we will consider a bound on the coefficient of the operator $(\partial\phi\partial\phi)^2$ in a bottom-up way in Sec. 4.1.

3.7 Euler-Heisenberg theory

We consider the Euler-Heisenberg theory, where the heavy particle is $U(1)$ charged field, and the light one is $U(1)$ gauge field. For the charged scalar field, we show that the constraint on the Wilson coefficients arises by implementing a suitable gauge fixing procedure.

3.7.1 Heavy charged fermion field

The action in the Mikowski space is described by

$$I_T[A, \psi, \bar{\psi}] \equiv \int d^4x \left[-\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \bar{\psi}(i\mathcal{D} - m)\psi \right], \quad [\text{Target}] \quad (3.153)$$

where ψ is the charged fermion field, $D_\mu = \partial_\mu + ieA_\mu$ is the covariant derivative, and $F_{\mu\nu}$ is the field strength of photon defined by $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$. We define I_0 and I_1 as follows:

$$I_0[A, \psi, \bar{\psi}] \equiv I_T[A, 0, 0] + I_T[0, \psi, \bar{\psi}] = \int d^4x \left[-\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \bar{\psi}(i\mathcal{D} - m)\psi \right], \quad (3.154)$$

$$I_1[A, \psi, \bar{\psi}] \equiv I_T[A, \psi, \bar{\psi}] - I_0[A, \psi, \bar{\psi}] = -e \cdot \int d^4x A_\mu \bar{\psi} \gamma^\mu \psi. \quad (3.155)$$

By introducing an auxiliary parameter g , define $I_g \equiv I_0 + g \cdot I_1$. Note here that the target theory is obtained as $I_T = \lim_{g \rightarrow 1} I_g$. To clarify the procedure of wave function renormalization, we also perform the path integral over the $U(1)$ gauge field. By calculating one-loop diagrams and performing the Wick rotation, the partition function of I_g is obtained in the following way [60].

$$\begin{aligned} Z_g[\bar{A}] &\equiv \int d[A] d[\psi] d[\bar{\psi}] e^{-I_g^{(E)}[A, \psi, \bar{\psi}]}, \\ &= \int d[A] \exp \left[- \int (d^4x)_E \left(\frac{1}{4} \left(1 + \alpha_0 \frac{g^2 e^2}{4! \pi^2} \right) F_{\mu\nu} F^{\mu\nu} - \alpha_2 \frac{g^2 e^2}{5! \pi^2 m^2} \partial^\mu F_{\mu\nu} \partial_\rho F^{\rho\nu} \right. \right. \\ &\quad \left. \left. - \alpha_4 \frac{g^2 e^2}{6! \pi^2 m^4} \partial^\mu F_{\mu\nu} \square \partial_\rho F^{\rho\nu} - \gamma_{4,1} \frac{g^4 e^4}{6! \pi^2 m^4} (F_{\mu\nu} F^{\mu\nu})^2 - \gamma_{4,2} \frac{g^4 e^4}{6! \pi^2 m^4} (F_{\mu\nu} \tilde{F}^{\mu\nu})^2 + \mathcal{O}(m^{-6}) \right) \right], \\ &= \exp \left[- \int (d^4x)_E \left(\frac{1}{4} \left(1 + \alpha_0 \frac{g^2 e^2}{4! \pi^2} \right) \bar{F}'_{\mu\nu} \bar{F}'^{\mu\nu} - \alpha_2 \frac{g^2 e^2}{5! \pi^2 m^2} \partial^\mu \bar{F}'_{\mu\nu} \partial_\rho \bar{F}'^{\rho\nu} \right. \right. \\ &\quad \left. \left. - \alpha_4 \frac{g^2 e^2}{6! \pi^2 m^4} \partial^\mu \bar{F}'_{\mu\nu} \square \partial_\rho \bar{F}'^{\rho\nu} - \gamma_{4,1} \frac{g^4 e^4}{6! \pi^2 m^4} (\bar{F}'_{\mu\nu} \bar{F}'^{\mu\nu})^2 - \gamma_{4,2} \frac{g^4 e^4}{6! \pi^2 m^4} (\bar{F}'_{\mu\nu} \widetilde{\bar{F}'^{\mu\nu}})^2 + \mathcal{O}(m^{-6}) \right) \right], \\ &= \exp \left[- \int (d^4x)_E \left(\frac{1}{4} \bar{F}_{\mu\nu} \bar{F}^{\mu\nu} - \gamma_{4,1} \frac{g^4 e^4}{6! \pi^2 m^4} (\bar{F}_{\mu\nu} \bar{F}^{\mu\nu})^2 - \gamma_{4,2} \frac{g^4 e^4}{6! \pi^2 m^4} (\bar{F}_{\mu\nu} \widetilde{\bar{F}^{\mu\nu}})^2 + \mathcal{O}(m^{-6}) \right) \right], \end{aligned} \quad (3.156)$$

where $\bar{F}'_{\mu\nu} \equiv \partial_\mu \bar{A}'_\nu - \partial_\nu \bar{A}'_\mu$ with the classical solution $\bar{A}'_\mu \equiv (1 + \alpha_0 g^2 e^2 / 4! \pi^2) \bar{A}_\mu$, and the last line is obtained by choosing a background field satisfying $\partial_\mu \bar{F}^{\mu\nu} = 0$. Here, we omit the vacuum energy in the Euclidean effective action because it cancels in the relative entropy. The effective action of I_g is calculated as follows:

$$\begin{aligned} W_g[\bar{A}] &\equiv -\ln Z_g[\bar{A}] \\ &= \int (d^4x)_E \left(\frac{1}{4} \bar{F}_{\mu\nu} \bar{F}^{\mu\nu} - \gamma_{4,1} \frac{g^4 e^4}{6! \pi^2 m^4} (\bar{F}_{\mu\nu} \bar{F}^{\mu\nu})^2 - \gamma_{4,2} \frac{g^4 e^4}{6! \pi^2 m^4} (\bar{F}_{\mu\nu} \widetilde{\bar{F}^{\mu\nu}})^2 + \mathcal{O}(m^{-6}) \right), \end{aligned} \quad (3.157)$$

with [60]

$$\alpha_2 = -1, \quad \alpha_4 = \frac{9}{14}, \quad \gamma_{4,1} = \frac{1}{2}, \quad \gamma_{4,2} = \frac{7}{8}. \quad (3.158)$$

The Euclidean effective action of the target theory is calculated as

$$W_T[\bar{A}] = \lim_{g \rightarrow 1} W_g[\bar{A}]. \quad (3.159)$$

By taking the limit of $g = 0$, the action of the NIRT and MFFRT in the Minkowski space is defined as

$$I_{\text{NI}}[A, \psi, \bar{\psi}] \equiv \lim_{g \rightarrow 0} I_g[A, \psi, \bar{\psi}] = \int d^4x \left[-\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \bar{\psi}(i\not{\partial} - m)\psi \right]. \quad \text{[Reference]} \quad (3.160)$$

Therefore, the partition function and effective action of the reference theory are respectively obtained as follows:

$$Z_{\text{NI}}[\bar{A}] = \lim_{g \rightarrow 0} Z_g[\bar{A}], \quad (3.161)$$

$$W_{\text{NI}}[\bar{A}] = \lim_{g \rightarrow 0} W_g[\bar{A}] = \int (d^4x)_{\text{E}} \frac{1}{4} \bar{F}_{\mu\nu} \bar{F}^{\mu\nu}. \quad (3.162)$$

By defining the probability distribution function $P_g[A, \psi, \bar{\psi}] \equiv e^{-I_g[A, \psi, \bar{\psi}]} / Z_g[\bar{A}]$, the derivative of W_g with respect to g is given by

$$\begin{aligned} \left(\frac{dW_g}{dg} \right)_{g=0} &= \left(\frac{\partial W_g}{\partial g} \right)_{g=0} + \int (d^4x)_{\text{E}} \left(\frac{\delta W_g}{\delta \bar{A}'} \right)_{g=0} \left(\frac{d\bar{A}'}{dg} \right)_{g=0} \\ &= \left(\frac{\partial W_g}{\partial g} \right)_{g=0} \\ &= \int d[A] d[\psi] d[\bar{\psi}] P_0[\psi, \bar{\psi}] I_{\text{I}}[A, \psi, \bar{\psi}] \\ &= \langle I_{\text{I}} \rangle_{g=0} = 0, \end{aligned} \quad (3.163)$$

where the partial derivative means differentiating by g while keeping \bar{A}' , $(d\bar{A}'/dg)_{g=0}$ is used in the second line, and $(dW_g/dg)_{g=0} = 0$ is used in the last line. The Feynman diagram for $\langle I_{\text{I}} \rangle_{g=0}$ is shown in the right panel of Fig. 5. From Eqs. (2.23), (3.157), and (3.163), the relative entropy between P_0 and P_g is calculated as follows:

$$\begin{aligned} S(P_0 || P_g) &\equiv \int d[A] d[\psi] d[\bar{\psi}] (P_0[A, \psi, \bar{\psi}] \ln P_0[A, \psi, \bar{\psi}] - P_0[A, \psi, \bar{\psi}] \ln P_g[A, \psi, \bar{\psi}]) \\ &= -\ln Z_0[\bar{A}] + \ln Z_g[\bar{A}] + g \cdot \int d[A] d[\psi] d[\bar{\psi}] P_0[\psi, \bar{\psi}] I_{\text{I}}[A, \psi, \bar{\psi}] \\ &= -\ln Z_0[\bar{A}] + \ln Z_g[\bar{A}] \\ &= W_0[\bar{A}] - W_g[\bar{A}] \end{aligned}$$

$$= \int (d^4x)_E \left[\gamma_{4,1} \frac{g^4 e^4}{6! \pi^2 m^4} (\bar{F}_{\mu\nu} \bar{F}^{\mu\nu})^2 + \gamma_{4,2} \frac{g^4 e^4}{6! \pi^2 m^4} (\bar{F}_{\mu\nu} \tilde{\bar{F}}^{\mu\nu})^2 + \mathcal{O}(m^{-6}) \right] \geq 0, \quad (3.164)$$

where the first line is the definition of the relative entropy, Eq. (3.163) is used in the third line, the fourth line is the definition of the effective action, and Eq. (3.157) is used in the last line. By taking $g = 1$, the relative entropy between P_{NI} and P_{T} is given by

$$\begin{aligned} S(P_{\text{NI}}||P_{\text{T}}) &= W_{\text{NI}}[\bar{A}] - W_{\text{T}}[\bar{A}] \\ &= \int (d^4x)_E \left[\gamma_{4,1} \frac{e^4}{6! \pi^2 m^4} (\bar{F}_{\mu\nu} \bar{F}^{\mu\nu})^2 + \gamma_{4,2} \frac{e^4}{6! \pi^2 m^4} (\bar{F}_{\mu\nu} \tilde{\bar{F}}^{\mu\nu})^2 + \mathcal{O}(m^{-6}) \right] \geq 0, \end{aligned} \quad (3.165)$$

where $P_{\text{NI}} = P_0$, $P_{\text{T}} = P_{g=1}$, $W_{\text{NI}} = W_0$, and $W_{\text{T}} = W_{g=1}$ are used. According to Eq. (3.158), it is clear that the right-hand side of Eq. (3.165) takes negative value up to dimension-eight terms. This result is consistent with the non-negativity of the relative entropy.

3.7.2 Heavy charged scalar field

The action of the massive charged scalar field in the Minkowski space is described by

$$I_{\text{T}}[A, \Phi] \equiv \int d^4x \left(-\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + D_\mu \Phi D^\mu \Phi^* - m^2 |\Phi|^2 \right), \quad [\text{Target}] \quad (3.166)$$

where Φ is the charged massive scalar field. We define I_0 and I_{I} as follows:

$$I_0[A, \Phi] \equiv I_{\text{T}}[A, 0] + I_{\text{T}}[0, \Phi] = \int d^4x \left(-\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \partial_\mu \Phi \partial^\mu \Phi^* - m^2 |\Phi|^2 \right), \quad (3.167)$$

$$I_{\text{I}}[A, \Phi] \equiv I_{\text{T}}[A, \Phi] - I_0[A, \Phi] = \int d^4x \left(-ieA^\mu (\partial_\mu \Phi) \Phi^* + ieA^\mu \Phi (\partial_\mu \Phi^*) + e^2 A_\mu A^\mu |\Phi|^2 \right). \quad (3.168)$$

By introducing a parameter g , define an action as $I_g \equiv I_0 + g \cdot I_{\text{I}}$. Note here that the interaction I_{I} includes the first and second order of e , and the order of g differs from that of e . Similar to the massive fermion, the Euclidean effective action of the target theory is obtained as,

$$\begin{aligned} W_{\text{T}}[\bar{A}] &= \lim_{g \rightarrow 1} W_g[\bar{A}] \\ &= \int (d^4x)_E \left(\frac{1}{4} \bar{F}_{\mu\nu} \bar{F}^{\mu\nu} - \gamma_{4,1} \frac{e^4}{6! \pi^2 m^4} (\bar{F}_{\mu\nu} \bar{F}^{\mu\nu})^2 - \gamma_{4,2} \frac{e^4}{6! \pi^2 m^4} (\bar{F}_{\mu\nu} \tilde{\bar{F}}^{\mu\nu})^2 + \mathcal{O}(m^{-6}) \right), \end{aligned} \quad (3.169)$$

with [60]

$$\alpha_2 = \frac{37}{8}, \quad \alpha_4 = \frac{159}{56}, \quad \gamma_{4,1} = \frac{261}{32}, \quad \gamma_{4,2} = \frac{243}{32}. \quad (3.170)$$

By taking the limit of $g = 0$, the action of the NIRT and MFFRT in the Minkowski space is defined as

$$I_{\text{NI}}[A, \Phi] = \lim_{g \rightarrow 0} I_g[A, \Phi] = \int d^4x \left(-\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \partial_\mu \Phi \partial^\mu \Phi^* - m^2 |\Phi|^2 \right). \quad [\text{Reference}] \quad (3.171)$$

Then, we obtain as follows:

$$Z_{\text{NI}}[\bar{A}] = \lim_{g \rightarrow 0} Z_g[\bar{A}] \quad (3.172)$$

$$W_{\text{NI}}[\bar{A}] = \lim_{g \rightarrow 0} W_g[\bar{A}] = \int (d^4x)_E \frac{1}{4} \bar{F}_{\mu\nu} \bar{F}^{\mu\nu}. \quad (3.173)$$

By defining a probability distribution function as $P_g[A, \Phi] \equiv e^{-I_g^{(\text{E})}[A, \Phi]} / Z_g[\bar{A}]$, the expectation value of the interaction is given by

$$\begin{aligned} g \cdot \langle I_I \rangle_{g=0} &= g \cdot \int d[A] d[\Phi] P_0[A, \Phi] I_I^{(\text{E})}[A, \Phi] \\ &= \int (d^4x)_E \left[\left(\frac{\delta W_g}{\delta \bar{A}'_\mu} \right)_{g=0} \bar{A}_\mu + \left(\frac{\delta W_g}{\delta \bar{A}'_\mu \bar{A}'^\mu} \right)_{g=0} \bar{A}_\mu \bar{A}^\mu \right] \\ &= \int (d^4x)_E \left(\frac{\delta W_g}{\delta \bar{A}'_\mu \bar{A}'^\mu} \right)_{g=0} \bar{A}_\mu \bar{A}^\mu, \end{aligned} \quad (3.174)$$

where the term proportional to \bar{A}_μ vanishes because of the Lorentz symmetry. The Feynman diagram for $\langle I_I \rangle_{g=0}$ is shown in Fig. 6. By taking the gauge-fixing condition $\bar{A}_\mu \bar{A}^\mu = 0$, which is called the non-linear gauge [61], $\langle I_I \rangle_{g=0}$ can vanish. From Eqs. (3.164) and (3.174), the relative entropy between P_{NI} and P_{T} is given by

$$\begin{aligned} S(P_{\text{NI}} || P_{\text{T}}) &= W_{\text{NI}}[\bar{A}] - W_{\text{T}}[\bar{A}] + \langle I_I \rangle_{g=0} \\ &= W_{\text{NI}}[\bar{A}] - W_{\text{T}}[\bar{A}] \\ &= \int (d^4x)_E \left[\gamma_{4,1} \frac{e^4}{6! \pi^2 m^4} (\bar{F}_{\mu\nu} \bar{F}^{\mu\nu})^2 + \gamma_{4,2} \frac{e^4}{6! \pi^2 m^4} (\bar{F}_{\mu\nu} \widetilde{\bar{F}}^{\mu\nu})^2 + \mathcal{O}(m^{-6}) \right] \geq 0, \end{aligned} \quad (3.175)$$

where the first line is the definition of the relative entropy, Eq. (3.174) and the gauge fixing condition $\bar{A}_\mu \bar{A}^\mu = 0$ are used in the second line, and Eq. (3.157) is used in the last line. According to Eq. (3.170), it is clear that the right-hand side of Eq. (3.175) takes a negative value up to dimension-eight terms. This result is consistent with the non-negativity of the relative entropy.

The non-negativity of the relative entropy holds even for the massive charged vector field. Since the interaction between the massive charged vector and $U(1)$ gauge fields arise from the covariant derivative of the kinetic term, Eq. (3.174) holds even for the massive charged vector field. By using the non-linear gauge $\bar{A}_\mu \bar{A}^\mu = 0$, Eq. (3.175) holds for the massive charged vector field with [60],

$$\alpha_2 = \frac{37}{8}, \quad \alpha_4 = \frac{159}{56}, \quad \gamma_{4,1} = \frac{261}{32}, \quad \gamma_{4,2} = \frac{243}{32}. \quad (3.176)$$

Consequently, the inequality of (3.175) holds even for the massive charged vector field.




Figure 6. Feynman diagrams of $\langle I_1 \rangle_{g=0}$ for the Euler-Heisenberg theory with the charged scalar field at the one-loop level. The dotted line denotes the heavy scalar field. The left and right diagrams represent $(\delta W_g / \delta A_\mu)_{A=0}$ and $(\delta W_g / \delta (A_\mu A^\mu))_{A_\mu A^\mu=0}$, respectively. The right one is a tadpole-like diagram for the composite field $A_\mu A^\mu$.

3.8 Gravitationally coupled massive scalar field at tree level

Let us consider a theory [49] in the Minkowski space:

$$I_T[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A_\mu, \Phi] \equiv \int d^4x \sqrt{-g} \left(\frac{M_{\text{Pl}}^2}{2} R - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} - (a_\Phi R + b_\Phi F_{\mu\nu} F^{\mu\nu}) \Phi + \frac{1}{2} g^{\mu\nu} \partial_\mu \Phi \partial_\nu \Phi - \frac{1}{2} m_\Phi^2 \Phi^2 \right), \quad [\text{Target}] \quad (3.177)$$

where $R_{\mu\nu\rho\sigma}$ is the Riemann tensor, R is the scalar curvature of the metric $g_{\mu\nu}$, $F_{\mu\nu}$ is the field strength of $U(1)$ gauge field, Φ is a massive real scalar field, and a_Φ, b_Φ are dimensionful coupling constants. The action in the Euclidean space is expressed as

$$I_T^{(\text{E})}[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A_\mu, \Phi_E] = \int (d^4x)_E \sqrt{g_E} \left(-\frac{M_{\text{Pl}}^2}{2} R + \frac{1}{4} F_{\mu\nu} F^{\mu\nu} + (a_\Phi R + b_\Phi F_{\mu\nu} F^{\mu\nu}) \Phi_E + \frac{1}{2} g_{E,IJ} \partial_I \Phi_E \partial_J \Phi_E + \frac{1}{2} m_\Phi^2 (\Phi_E)^2 \right), \quad (3.178)$$

where the R and $F_{\mu\nu}$ are defined from background fields in the Minkowski space; see Appendix A. We define the actions $I_0^{(\text{E})}$ and $I_1^{(\text{E})}$ in the Euclidean space as follows:

$$I_0^{(\text{E})}[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A_\mu, \Phi_E] = I_T^{(\text{E})}[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A_\mu, 0] + I_T^{(\text{E})}[g_{\mu\nu}; 0, 0, \Phi_E] \\ = \int (d^4x)_E \sqrt{g_E} \left(-\frac{M_{\text{Pl}}^2}{2} R + \frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{1}{2} g_{E,IJ} \partial_I \Phi_E \partial_J \Phi_E + \frac{1}{2} m_\Phi^2 (\Phi_E)^2 \right), \quad (3.179)$$

$$I_1^{(\text{E})}[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A_\mu, \Phi_E] = I_T^{(\text{E})}[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A_\mu, \Phi_E] - I_0^{(\text{E})}[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A_\mu, \Phi_E] \\ = \int (d^4x)_E \sqrt{g_E} (a_\Phi R + b_\Phi F_{\mu\nu} F^{\mu\nu}) \Phi_E. \quad (3.180)$$

The action of the NIRT is the same as that of the MFFRT and is defined as

$$I_{\text{NI}}^{(\text{E})}[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A_\mu, \Phi_E] = \int (d^4x)_E \sqrt{g_E} \left(-\frac{M_{\text{Pl}}^2}{2} R + \frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{1}{2} g_{E,IJ} \partial_I \Phi_E \partial_J \Phi_E + \frac{1}{2} m_\Phi^2 (\Phi_E)^2 \right). \quad (3.181)$$

Note here that the NIRT does not include the interaction between Φ and $A_\mu, R_{\mu\nu\rho\sigma}$, but the interaction between $g_{\mu\nu}$ and Φ . By introducing a parameter g , define an action as $I_g^{(E)} \equiv I_0^{(E)} + g \cdot I_I^{(E)}$. Then, the target theory is reproduced as $I_T^{(E)} = \lim_{g \rightarrow 1} I_g^{(E)}$. The target and reference theories are expressed as follows:

$$I_T^{(E)}[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A_\mu, \Phi_E] = \lim_{g \rightarrow 1} I_g^{(E)}[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A_\mu, \Phi_E], \quad (3.182)$$

$$I_{NI}^{(E)}[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A_\mu, \Phi_E] = \lim_{g \rightarrow 0} I_g^{(E)}[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A_\mu, \Phi_E]. \quad (3.183)$$

The partition function and effective action of $I_g^{(E)}$ are respectively calculated as follows:

$$\begin{aligned} Z_g[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A_\mu] &\equiv \int d[\Phi_E] e^{-I_g^{(E)}[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A_\mu, \Phi_E]} \\ &= \exp\left(\int (d^4x)_E \sqrt{g_E} \left(-\frac{M_{Pl}^2}{2} R + \frac{1}{4} F_{\mu\nu} F^{\mu\nu} - g^2 \cdot \frac{1}{2m_\Phi^2} (a_\Phi R + b_\Phi F_{\mu\nu} F^{\mu\nu})^2\right)\right), \end{aligned} \quad (3.184)$$

$$\begin{aligned} W_g[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A_\mu] &\equiv -\ln Z_g[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A_\mu] \\ &= \int (d^4x)_E \sqrt{g_E} \left(-\frac{M_{Pl}^2}{2} R + \frac{1}{4} F_{\mu\nu} F^{\mu\nu} - g^2 \cdot \frac{1}{2m_\Phi^2} (a_\Phi R + b_\Phi F_{\mu\nu} F^{\mu\nu})^2\right). \end{aligned} \quad (3.185)$$

By defining a probability distribution function as

$$P_g[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A_\mu, \Phi_E] \equiv \frac{e^{-I_g^{(E)}[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A_\mu, \Phi_E]}}{Z_g[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A_\mu]}, \quad (3.186)$$

we obtain

$$\begin{aligned} \langle I_I \rangle_{g=0} &= \int d[\Phi_E] P_0[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A_\mu, \Phi_E] I_I^{(E)}[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A_\mu, \Phi_E] \\ &= I_I^{(E)}[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A_\mu, 0] \\ &= 0. \end{aligned} \quad (3.187)$$

From Eqs. (2.8), (3.185), and (3.187), the relative entropy between P_0 and P_g is calculated as follows:

$$\begin{aligned} S(P_0||P_g) &\equiv \int d[\Phi_E] (P_0 \ln P_0 - P_0 \ln P_g) \\ &= -\ln Z_0[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A_\mu] + \ln Z_g[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A_\mu] + g \cdot \langle I_I \rangle_{g=0} \\ &= -\ln Z_0[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A_\mu] + \ln Z_g[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A_\mu] \\ &= W_0[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A_\mu] - W_g[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A_\mu] \\ &= g^2 \cdot \frac{1}{2m_\Phi^2} \int (d^4x)_E \sqrt{g_E} (a_\Phi R + b_\Phi F_{\mu\nu} F^{\mu\nu})^2 \geq 0, \end{aligned} \quad (3.188)$$

where the first line is the definition of the relative entropy, Eq. (3.187) is used in the third line, the fourth line is the definition of the effective actions, and Eq. (3.185) is used in the last line. By taking $g = 1$, the relative entropy between P_{NI} and P_{T} is given by

$$\begin{aligned} S(P_{\text{NI}}||P_{\text{T}}) &= W_{\text{NI}}[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A_\mu] - W_{\text{T}}[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A_\mu] \\ &= \frac{1}{2m_\Phi^2} \int (d^4x)_{\text{E}} \sqrt{g_{\text{E}}} (a_\Phi R + b_\Phi F_{\mu\nu} F^{\mu\nu})^2 \geq 0, \end{aligned} \quad (3.189)$$

where $P_{\text{NI}} = P_0$, $P_{\text{T}} = P_{g=1}$, $W_{\text{NI}} = W_0$, and $W_{\text{T}} = W_{g=1}$ are used. This result represents the inequalities of (2.52) and (2.86). Therefore, even in the gravitational theory, the relative entropy yields constraints on the EFT.

3.9 Dimension-six four-fermion operators from tree-level UV completions

Let us consider a tree-level UV completion of dimension-six four-fermion operators in the Minkowski space:

$$I_{\text{T}}[\psi, \bar{\psi}, \Phi_i] \equiv \int d^4x (\bar{\psi}(i\not{\partial} - m)\psi - (-1)^{s_i} M^2 \Phi_i^2 + a_i \Phi_i (\bar{\psi} \Gamma_i \psi)), \quad [\text{Target}] \quad (3.190)$$

where the kinetic terms of the heavy degrees of freedom are omitted for simplicity, and the sum over the index i is not performed. Here, we defined as

$$\Gamma_1 = \hat{1}, \quad \Gamma_2 = \gamma^\mu, \quad \Gamma_3 = i\gamma_5, \quad \Gamma_4 = \gamma^\mu \gamma_5, \quad \Gamma_5 = \sigma^{\mu\nu}, \quad (3.191)$$

$$s_1 = 0, \quad s_2 = 1, \quad s_3 = 0, \quad s_4 = 1, \quad s_5 = 0, \quad (3.192)$$

where $\sigma^{\mu\nu} \equiv i[\gamma^\mu, \gamma^\nu]/2$. The sign s_i is determined so that the solution of the equation of motion for Φ_i becomes a local minimum of the Euclidean action. Otherwise, the validity of the Euclidean path integral method is violated. We choose the time components of the background field $\bar{\psi} \Gamma_i \psi$ to be zero values since the non-negativity can be broken when the probability distribution functions of Eq. (2.2) are not Hermitian. Note here that the background field of the Euclidean space is defined by that of the Minkowski space. The massive fields are defined as follows:

$$\Phi_1 = \phi_{\text{S}}, \quad \Phi_2 = A_V^\mu, \quad \Phi_3 = \phi_P, \quad \Phi_4 = A_A^\mu, \quad \Phi_5 = T^{\mu\nu}. \quad (3.193)$$

Then, I_0 and I_{I} are given by,

$$I_0[\psi, \bar{\psi}, \Phi_i] \equiv \int d^4x (\bar{\psi}(i\not{\partial} - m)\psi - (-1)^{s_i} M^2 \Phi_i^2), \quad (3.194)$$

$$I_{\text{I}}[\psi, \bar{\psi}, \Phi_i] \equiv \int d^4x a_i \Phi_i (\bar{\psi} \Gamma_i \psi). \quad (3.195)$$

For this target theory, the NIRT is the same as the MFFRT, and the action of the NIRT and MFFRT is defined as

$$I_{\text{NI}}[\psi, \bar{\psi}, \Phi_i] \equiv I_0[\psi, \bar{\psi}, \Phi_i]. \quad [\text{Reference}] \quad (3.196)$$

By using the auxiliary parameter g , define $I_g \equiv I_0 + g \cdot I_I$. Then, the target and reference theories are reproduced as follows:

$$I_T[\psi, \bar{\psi}, \Phi_i] = \lim_{g \rightarrow 1} I_g[\psi, \bar{\psi}, \Phi_i], \quad I_{NI}[\psi, \bar{\psi}, \Phi_i] = \lim_{g \rightarrow 0} I_g[\psi, \bar{\psi}, \Phi_i]. \quad (3.197)$$

After the Wick rotation $I_g \rightarrow I_g^{(E)}$, the partition function and effective action of $I_g^{(E)}$ are respectively calculated as follows:

$$Z_g[\psi, \bar{\psi}] \equiv \int d[\Phi_{E,i}] e^{-I_g^{(E)}[\psi, \bar{\psi}, \Phi_{E,i}]}, \quad (3.198)$$

$$W_g[\psi, \bar{\psi}] \equiv -\ln Z_g[\psi, \bar{\psi}] = \int (d^4x)_E \left(\bar{\psi}(i\not{\partial} + m)\psi - g^2 \cdot (-1)^{s_i} \frac{a_i^2}{4M^2} (\bar{\psi}\Gamma_i\psi)^2 \right). \quad (3.199)$$

By defining a probability distribution function as,

$$P_g[\Phi_i] \equiv \frac{e^{-I_g^{(E)}[\psi, \bar{\psi}, \Phi_{E,i}]}}{Z_g[\psi, \bar{\psi}]}, \quad (3.200)$$

the expectation value of the interaction is calculated as

$$\begin{aligned} \langle I_I \rangle_{g=0} &\equiv \int d[\Phi_{E,i}] P_0[\Phi_i] I_I^{(E)}[\psi, \bar{\psi}, \Phi_{E,i}] \\ &= I_I^{(E)}[\psi, \bar{\psi}, 0] \\ &= 0. \end{aligned} \quad (3.201)$$

From Eqs. (2.8), (3.199), and (3.201), the relative entropy between P_0 and P_g is calculated as

$$\begin{aligned} S(P_0||P_g) &\equiv \int d[\Phi_{E,i}] (P_0[\Phi_{E,i}] \ln P_0[\Phi_{E,i}] - P_0[\Phi_{E,i}] \ln P_g[\Phi_{E,i}]) \\ &= -\ln Z_0[\psi, \bar{\psi}] + \ln Z_g[\psi, \bar{\psi}] + g \cdot \langle I_I \rangle_{g=0} \\ &= W_0[\psi, \bar{\psi}] - W_g[\psi, \bar{\psi}] \\ &= g^2 \cdot (-1)^{s_i} \frac{a_i^2}{4M^2} \int (d^4x)_E (\bar{\psi}\Gamma_i\psi)^2 \geq 0 \end{aligned} \quad (3.202)$$

By taking to be $g = 1$, we obtain the relative entropy between P_{NI} and P_T as follows.

$$S(P_{NI}||P_T) = W_{NI}[\psi, \bar{\psi}] - W_T[\psi, \bar{\psi}] = (-1)^{s_i} \frac{a_i^2}{4M^2} \int (d^4x)_E (\bar{\psi}\Gamma_i\psi)^2 \geq 0, \quad (3.203)$$

where $P_{NI} = P_0$, $P_T = P_{g=1}$, $W_{NI} = W_0$, and $W_T = W_{g=1}$ are used, and $(-1)^{s_i} (\bar{\psi}\Gamma_i\psi)^2$ takes positive values because the time components of $\bar{\psi}\Gamma_i\psi$ are assumed to be zero. This result is consistent with Ref. [62] and the non-negativity of the relative entropy.

3.10 Tree level UV completions with unstable field

So far, we have studied target theories where the Euclidean path integral method is valid, i.e., the saddle point approximation works well. In the following, we consider target theories with unstable auxiliary fields where the saddle point approximation is not valid in the Euclidean path integral method.

- Unstable dilaton-like particle — Consider a target theory defined by the action in the Minkowski space as follows:

$$I_T[A, \Phi] \equiv \int d^4x \left(-\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + m_\Phi^2 \Phi^2 + \frac{1}{f_\Phi} \Phi F_{\rho\sigma} F^{\rho\sigma} \right), \quad [\mathbf{Target}] \quad (3.204)$$

where m_Φ and f_Φ are the mass and the decay constant of the heavy auxiliary field, respectively, and $F_{\mu\nu}$ is the field strength of photon field. The action in the Euclidean space is given by

$$I_T^{(E)}[A, \Phi_E] = \int (d^4x)_E \left(\frac{1}{4} F_{\mu\nu} F^{\mu\nu} - m_\Phi^2 \Phi_E^2 - \frac{1}{f_\Phi} \Phi_E F_{\rho\sigma} F^{\rho\sigma} \right), \quad (3.205)$$

where $F_{\mu\nu}$ is assumed to be a background field. Note here that the auxiliary field is unstable because of the negative mass term. The non-interacting and interacting terms in the Minkowski space are respectively expressed as follows:

$$I_0[A, \Phi] \equiv \int d^4x \left(-\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + m_\Phi^2 \Phi^2 \right) \quad (3.206)$$

$$I_I[A, \Phi] \equiv \int d^4x \left(\frac{1}{f_\Phi} \Phi F_{\mu\nu} F^{\mu\nu} \right) \quad (3.207)$$

For this target theory, the NIRT is the same as the MFFRT, and the action of the NIRT and MFFRT is defined as

$$I_{NI}[A, \Phi] \equiv I_0[A, \Phi]. \quad [\mathbf{Reference}] \quad (3.208)$$

By using the auxiliary parameter g , define $I_g \equiv I_0 + g \cdot I_I$. Then, the target and reference theories are reproduced as follows:

$$I_T[A, \Phi] = \lim_{g \rightarrow 1} I_g[A, \Phi], \quad I_{NI}[A, \Phi] = \lim_{g \rightarrow 0} I_g[A, \Phi]. \quad (3.209)$$

The solution of the equation of motion of $I_g^{(E)}$ for Φ_E is calculated as

$$\tilde{\Phi}_g = -g \cdot \frac{1}{2m_\Phi^2 f_\Phi} F_{\mu\nu} F^{\mu\nu}, \quad (3.210)$$

which is not a local maximum of $I_g^{(E)}$. Therefore, the saddle point approximation around this solution is not valid, and the Euclidean effective action cannot be calculated. In other words, the calculation procedures of the relative entropy of this work are not applicable. To check that

the non-negativity of the relative entropy is violated, evaluate the effective action of the target theory in the Minkowski space. Consider a field redefinition of Φ as follows:

$$\Phi \rightarrow \Phi - g \cdot \frac{1}{2m_\Phi^2 f_\Phi} F_{\mu\nu} F^{\mu\nu}.$$

By performing the above field redefinition, I_g yields the effective action in the Minkowski space as follows:

$$W_g^{(M)}[A] = \int d^4x \left(-\frac{1}{4} F_{\mu\nu} F^{\mu\nu} - g^2 \cdot \frac{1}{4m_\Phi^2 f_\Phi^2} (F_{\mu\nu} F^{\mu\nu})^2 \right). \quad (3.211)$$

After the Wick rotation, we obtain the Euclidean effective action as follows:

$$W_g[A] = \int (d^4x)_E \left(\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + g^2 \cdot \frac{1}{4m_\Phi^2 f_\Phi^2} (F_{\mu\nu} F^{\mu\nu})^2 \right). \quad (3.212)$$

Note here that the second term of the right-hand side is positive. From Eqs. (2.8), and (3.212), the relative entropy between P_0 and P_g is calculated as follows:

$$\begin{aligned} S(P_0||P_g) &\equiv W_0[A] - W_g[A] + g \cdot \langle I_1 \rangle_{g=0} \\ &= -g^2 \cdot \frac{1}{4m_\Phi^2 f_\Phi^2} \int (d^4x)_E (F_{\mu\nu} F^{\mu\nu})^2, \end{aligned} \quad (3.213)$$

where $\langle I_1 \rangle_{g=0} = (\partial W_g / \partial g)_{g=0} = 0$ is used. By taking $g = 1$, we obtain

$$S(P_{\text{NI}}||P_{\text{T}}) = W_{\text{NI}}[A] - W_{\text{T}}[A] = -\frac{1}{4m_\Phi^2 f_\Phi^2} \int (d^4x)_E (F_{\mu\nu} F^{\mu\nu})^2, \quad (3.214)$$

where $P_{\text{NI}} = P_0$, $P_{\text{T}} = P_{g=1}$, $W_{\text{NI}} = W_0$, and $W_{\text{T}} = W_{g=1}$ are used. The right-hand side of the above equation takes a negative value, which is inconsistent with the non-negativity of the relative entropy. This is because the Euclidean path integral method does not work, and the relative entropy in the Euclidean space can not be defined in this theory.

- Doublet of real, shift-symmetric, massless scalar fields theory — Consider a target theory [63] in the Minkowski space,

$$I_{\text{T}}[\phi, X] \equiv \int d^4x \left(\frac{1}{2} (\partial_\mu \phi_i \partial^\mu \phi_i) + m_X^2 X_{\mu\nu} X^{\mu\nu} - \frac{\epsilon^{il}}{M} (\partial_\mu \phi_i \partial_\nu \phi_l) X^{\mu\nu} \right), \quad [\text{Target}] \quad (3.215)$$

where ϕ_i , $i = 1, 2$ is a doublet of real, shift-symmetric, massless scalar fields, $X_{\mu\nu}$ is an auxiliary field, and $\epsilon^{12} = -\epsilon^{21} = 1$. Similar to the previous example, the auxiliary field is unstable because of the mass term. The non-interacting and interacting terms in the Minkowski space are respectively expressed as follows:

$$I_0[\phi, X] \equiv \int d^4x \left(\frac{1}{2} (\partial_\mu \phi_i \partial^\mu \phi_i) + m_X^2 X_{\mu\nu} X^{\mu\nu} \right) \quad (3.216)$$

$$I_I[\phi, X] \equiv \int d^4x \left(-\frac{\epsilon^{il}}{M} (\partial_\mu \phi_i \partial_\nu \phi_l) X^{\mu\nu} \right). \quad (3.217)$$

For this target theory, the NIRT is the same as the MFFRT, and the action of the NIRT and MFFRT is defined as

$$I_{\text{NI}}[\phi, X] \equiv I_0[\phi, X]. \quad [\textbf{Reference}] \quad (3.218)$$

By the auxiliary parameter g , define $I_g \equiv I_0 + g \cdot I_I$. The target and reference theories are reproduced as follows:

$$I_{\text{T}}[\phi, X] = \lim_{g \rightarrow 1} I_g[\phi, X], \quad I_{\text{NI}}[\phi, X] = \lim_{g \rightarrow 0} I_g[\phi, X]. \quad (3.219)$$

The solution of the equation of motion of $I_g^{(\text{E})}$ for $X_{\mu\nu}$ is calculated as

$$\tilde{X}_{g,\mu\nu} = -g \cdot \frac{\epsilon^{il}}{2m_X^2 M} (\partial_\mu \phi_i \partial_\nu \phi_l), \quad (3.220)$$

which is a classical solution not being a local minimum because of the sign of the mass term of Eq. (3.215). Therefore, in the Euclidean path integral method, the saddle point approximation around this solution is not valid, and the relative entropy can not be calculated. To check the violation of the non-negativity of the relative entropy, evaluate the effective action of the target theory in the Minkowski space. By performing the following field redefinition,

$$X_{\mu\nu} \rightarrow X_{\mu\nu} + g \cdot \frac{\epsilon^{il}}{2m_X^2 M} (\partial_\mu \phi_i \partial_\nu \phi_l), \quad (3.221)$$

I_g yields the effective action in the Minkowski space as follows:

$$W_g^{(\text{M})}[\phi] = \int d^4x \left(\frac{1}{2} (\partial_\mu \phi_i \partial^\mu \phi_i) + g^2 \cdot \frac{1}{4m_X^2 M^2} \epsilon^{il} \epsilon^{kj} (\partial_\mu \phi_i \partial^\mu \phi_j) (\partial_\nu \phi_k \partial^\nu \phi_l) \right). \quad (3.222)$$

After the Wick rotation, we obtain the Euclidean effective action as follows:

$$W_g[\phi] = \int (d^4x)_{\text{E}} \left(-\frac{1}{2} (\partial_\mu \phi_i \partial^\mu \phi_i) - g^2 \cdot \frac{1}{4m_X^2 M^2} \epsilon^{il} \epsilon^{kj} (\partial_\mu \phi_i \partial^\mu \phi_j) (\partial_\nu \phi_k \partial^\nu \phi_l) \right). \quad (3.223)$$

From Eq. (2.8), and (3.223), the relative entropy between P_0 and P_g is expressed as follows:

$$\begin{aligned} S(P_0||P_g) &\equiv W_0[\phi] - W_g[\phi] + g \cdot \langle I_I \rangle_{g=0} \\ &= g^2 \cdot \frac{1}{4m_X^2 M^2} \int (d^4x)_{\text{E}} \epsilon^{il} \epsilon^{kj} (\partial_\mu \phi_i \partial^\mu \phi_j) (\partial_\nu \phi_k \partial^\nu \phi_l), \end{aligned} \quad (3.224)$$

where $\langle I_I \rangle_{g=0} = (\partial W_g / \partial g)_{g=0} = 0$ is used. By taking $g = 1$, we obtain

$$S(P_{\text{NI}}||P_{\text{T}}) = W_{\text{NI}}[\phi] - W_{\text{T}}[\phi] = \frac{1}{4m_X^2 M^2} \int (d^4x)_{\text{E}} \epsilon^{il} \epsilon^{kj} (\partial_\mu \phi_i \partial^\mu \phi_j) (\partial_\nu \phi_k \partial^\nu \phi_l), \quad (3.225)$$

where $P_{\text{NI}} = P_0$, $P_{\text{T}} = P_{g=1}$, $W_{\text{NI}} = W_0$, and $W_{\text{T}} = W_{g=1}$ are used. By choosing a background field, e.g., $\partial_\mu \tilde{\phi}_1 = (0, 1, 0, 0)$, and $\partial_\mu \tilde{\phi}_2 = (0, 0, 1, 0)$, we obtain

$$\epsilon^{il} \epsilon^{kj} (\partial_\mu \tilde{\phi}_i \partial^\mu \tilde{\phi}_j) (\partial_\nu \tilde{\phi}_k \partial^\nu \tilde{\phi}_l) = 2 \left((\partial_\mu \tilde{\phi}_1 \partial^\mu \tilde{\phi}_2)^2 - (\partial_\mu \tilde{\phi}_1 \partial^\mu \tilde{\phi}_1) (\partial_\nu \tilde{\phi}_2 \partial^\nu \tilde{\phi}_2) \right) = -2. \quad (3.226)$$

Then, the right-hand side of Eq. (3.225) takes a negative value, which is inconsistent with the non-negativity of the relative entropy. This is because the Euclidean path integral method is not valid because of the sign of the mass term of Eq. (3.215), and the relative entropy in the Euclidean space can not be defined in this theory.

4 Bottom-up approach: bounds on EFTs

We often face situations where the UV theory involving the heavy degrees of freedom is unknown in contrast to the top-down approach of the previous section. In this section, we consider zero-temperature systems and take a bottom-up approach, where the UV theory involving the interactions between heavy and light degrees of freedom are unknown. We focus on a class of EFTs, where the corrections to the non-higher derivative terms can be removed by a field redefinition of the background fields, and the entropy constraint on such EFTs is provided by focusing on the NIRT. As explained later, the dimension-eight term of a single massless scalar field, the SMEFT dimension-eight $SU(N)$ gauge bosonic operators, and Einstein-Maxwell theory with higher-derivative operators belong to such a class of theories due to the existence of symmetries.

Before presenting the details of the calculations, we list the assumptions used to derive the results in this section in the following.

- (I) Hermiticity of probability distribution functions — We assume the target and reference theory are represented by the Hermitian probability distribution functions. To derive the non-negativity of the relative entropy in Eq. (2.2), we used the Hermiticity of probability distribution functions, i.e., $\rho_{\text{R,T}} = \rho_{\text{R,T}}^\dagger$. The non-negativity of the relative entropy can be broken when this condition is not satisfied. This is the reason why the time components of the background field are chosen as zero in Sec. 3.9.
- (II) Validity of Euclidean path integral method — We assume the EFTs are generated from the solution of the local minimum. As shown in Sec. 3.10, the non-negativity of the relative entropy can be broken when the Euclidean path integral method is not valid, i.e., the saddle point approximation does not work because of the solution not being the local minimum.
- (III) Higher-derivative operators generated from the interaction between heavy and light fields — We assume the higher-derivative operators of the EFTs are generated from the interactions between heavy and light fields. The interaction of the UV theory is generally expressed as,

$$I_1[\phi, \Phi] = \int (d^4x)_{\text{E}} \mathcal{O}[\Phi] \otimes J[\phi], \quad (4.1)$$

where $\mathcal{O}[\Phi] \otimes J[\phi]$ generally involves summations over some indices in $\mathcal{O}[\Phi]$ and $J[\phi]$, e.g., Lorentz indices.

- (IV) Leading order in the interaction between heavy and light fields — We assume $J[\phi]$ does not include the higher-derivative operators⁴. This assumption is quantitatively reasonable because the higher-dimensional operator in $J[\phi]$ is suppressed by a heavier mass than Φ as follows:

$$J[\phi] = J_{\text{dim-4}}[\phi] + \sum_{i=5, \dots} \frac{1}{\Lambda^{i-4}} J_{\text{dim-}i}[\phi], \quad (4.2)$$

where $J_{\text{dim-4}}$ and $J_{\text{dim-}i}$ respectively denote operators up to dimension-four and dimension- i operators, and Λ is a mass scale satisfying $M \ll \Lambda$, where M is the mass of Φ . Note here that this assumption does not prohibit higher-dimensional interacting terms. For example, in Secs. 3.3, 3.5.1, and 3.5.2, we discussed $J[\phi] \propto (\partial_\mu \phi \partial^\mu \phi)$, $J[A] \propto F_{\mu\nu} F^{\mu\nu}$, and $J[A] \propto F_{\mu\nu} \tilde{F}^{\mu\nu}$, respectively, and the interacting terms I_1 were the dimension-five operator. Also, we assume the renormalizable terms in I_1 are dominant effects on the Euclidean effective action, and the non-renormalizable terms are negligible when I_1 includes both renormalizable and non-renormalizable terms. In other words, we consider the leading order of $1/\Lambda$ expansion for the interaction effects on the EFTs and assume as follows:

$$J[\phi] \simeq J_{\text{dim-}i}[\phi]/\Lambda^{i-4}, \quad (4.3)$$

where i denotes the leading order of $1/\Lambda$ expansion.

The first two assumptions are also imposed in Sec. 2; see Sec. 2.4. The main assumptions in this section are the third and fourth ones.

In the following sections, we focus on two cases: tree-level UV completion and loop-level UV completion. For the tree-level UV completion, we assume the tree level effects dominate the perturbative corrections from the heavy degrees of freedom to the Euclidean effective action. On the other hand, for the loop-level UV completion, we assume the loop level effects dominate the perturbative corrections to the Euclidean effective action. For some examples, i.e., the single massless scalar field with the dimension-eight term, SMEFT dimension-eight gauge bosonic operators, and Einstein-Maxwell theory with higher-derivative terms, we provide constraints from the relative entropy in the following way.

4.1 Single massless scalar field with dimension-eight operator

Consider an EFT described by a single massless scalar field theory with a dimension-eight operator as follows:

$$W_{\text{T}}[\phi] = \int (d^4x)_{\text{E}} \left[-\frac{1}{2} (\partial_\mu \phi \partial^\mu \phi) - \frac{c}{M^4} (\partial_\mu \phi \partial^\mu \phi)^2 \right]. \quad (4.4)$$

⁴For example, $(\partial\phi)^4$, $(F_{\mu\nu} F^{\mu\nu})^2$, etc. belong to the higher-derivative operators, which are the higher-dimensional operators. Note that the Einstein-Hilbert term is allowed to be J by Assumption (IV).

Because of the shift symmetry: $\phi \rightarrow \phi + \text{const.}$, Eq. (4.4) involves only the kinetic term, i.e., the non-higher derivative term, as the renormalizable term, and corrections to the kinetic term can be removed by redefining ϕ . Let us stand in the bottom-up approach and assume the second term of Eq. (4.4) is generated by integrating out the heavy fields of the theory of Eq. (2.44). According to Assumption (IV), $J[\phi]$ can be $\partial_\mu \phi$ or $\partial_\mu \phi \partial^\mu \phi$, which preserve the shift symmetry, but $\partial_\mu \phi$ effects on $\langle I_1 \rangle_{g=0}$ vanish because $\langle I_1 \rangle_{g=0}$ preserves the Lorentz symmetry. When we suppose that the EFT is generated by integrating out heavy degrees of freedom, the first order corrections for g to the Euclidean effective action are expressed as

$$\begin{aligned} g \cdot \langle I_1 \rangle_{g=0} &= g \cdot \left(\frac{\partial W_g}{\partial g} \right)_{g=0} \\ &= \int (d^4x)_E \left(\frac{\delta W_g}{\delta J} \right)_{J=0} J[\phi] \\ &\propto \int (d^4x)_E (\partial_\mu \phi \partial^\mu \phi), \end{aligned} \quad (4.5)$$

where $(\delta W_g / \delta J)_{J=0}$ denotes a tadpole-like diagram for the composite field $J[\phi]$ and does not depend on space-time. For both tree and loop-level UV completions, we consider the constraints on the Wilson coefficient of the dimension-eight operator in the following way.

- Tree-level UV completion — Consider the EFT generated by the tree-level UV completion. The partition function of $I_0 + g \cdot I_1$ is generally calculated as follows:

$$\begin{aligned} Z_g[\tilde{\phi}] &\equiv \int d[\phi] d[\Phi] e^{-I_g[\phi, \Phi]} \\ &= \int d[\phi] \exp \left[- \int (d^4x)_E \left(-\frac{1}{2} (1 + \alpha_2^{\text{tree}}) (\partial_\mu \phi \partial^\mu \phi) - \beta_2^{\text{tree}} (\partial_\mu \phi \partial^\mu \phi)^2 \right) \right] \\ &= \exp \left[- \int (d^4x)_E \left(-\frac{1}{2} (1 + \alpha_2^{\text{tree}}) (\partial_\mu \tilde{\phi}' \partial^\mu \tilde{\phi}') - \beta_2^{\text{tree}} (\partial_\mu \tilde{\phi}' \partial^\mu \tilde{\phi}')^2 \right) \right] \\ &= \exp \left[- \int (d^4x)_E \left(-\frac{1}{2} (\partial_\mu \tilde{\phi} \partial^\mu \tilde{\phi}) - \beta_2^{\text{tree}} \cdot (1 + \alpha_2^{\text{tree}})^{-2} (\partial_\mu \tilde{\phi} \partial^\mu \tilde{\phi})^2 \right) \right], \end{aligned} \quad (4.6)$$

where α_2^{tree} and β_2^{tree} denote the second or higher order corrections for g . Note here that β_2^{tree} does not include the first order correction for g because of Eq. (4.5). We assumed α_1^{tree} , α_2^{tree} , and β_2^{tree} are generated at the tree level. Also, in the second line, according to the procedure in Eqs. (2.68), (2.69), and (2.70), the first order correction for g is eliminated in α_2^{tree} and absorbed into the definition of ϕ . As discussed in Sec. 3.6, the dimension-six operators and other dimension-eight operators, e.g., $(\partial \partial \phi)^2$ and $(\partial \partial \partial \phi)^2$, generally arise, but they are eliminated by the background field satisfying $\partial_\mu \tilde{\phi}' = \text{const.}$, where $\tilde{\phi}'$ denotes the classical solution of the effective action. To remove the dimension-six operators, we choose the background fields as follows:

$$\tilde{\phi}' = (1 + \alpha_2^{\text{tree}})^{-1/2} \cdot \tilde{\phi}, \quad (4.7)$$

with $\partial_\mu \tilde{\phi} = \text{const.}$. From Eq. (4.6), the Euclidean effective actions are calculated as follows:

$$W_g[\tilde{\phi}] \equiv -\ln Z_g[\tilde{\phi}] = \int (d^4x)_E \left(-\frac{1}{2}(\partial_\mu \tilde{\phi} \partial^\mu \tilde{\phi}) - \beta_2^{\text{tree}} \cdot (1 + \alpha_2^{\text{tree}})^{-2} (\partial_\mu \tilde{\phi} \partial^\mu \tilde{\phi})^2 \right), \quad (4.8)$$

$$W_0[\tilde{\phi}] = \lim_{g \rightarrow 0} W_g[\tilde{\phi}] = \int (d^4x)_E \left(-\frac{1}{2}(\partial_\mu \tilde{\phi} \partial^\mu \tilde{\phi}) \right). \quad (4.9)$$

Combining Eq. (4.8) and (4.9), the shift of the Euclidean effective action by the interacting term is given by

$$W_g[\tilde{\phi}] - W_0[\tilde{\phi}] = -\beta_2^{\text{tree}} \cdot (1 + \alpha_2^{\text{tree}})^{-2} \int (d^4x)_E (\partial_\mu \tilde{\phi} \partial^\mu \tilde{\phi})^2. \quad (4.10)$$

Also, Eq. (4.8) yields following relations:

$$\begin{aligned} \left(\frac{dW_g}{dg} \right)_{g=0} &= \left(\frac{\partial W_g}{\partial g} \right)_{g=0} + \int (d^4x)_E \left(\frac{\delta W_g}{\delta \tilde{\phi}'} \right) \cdot \left(\frac{d\tilde{\phi}'}{dg} \right)_{g=0} \\ &= \left(\frac{\partial W_g}{\partial g} \right)_{g=0} \\ &= \langle I_1 \rangle_{g=0} = 0, \end{aligned} \quad (4.11)$$

where the partial derivative means differentiating by g while keeping $\tilde{\phi}'$, and $(d\tilde{\phi}'/dg)_{g=0} = 0$ is used because α_2^{tree} in Eq. (4.7) denotes the second or higher order corrections for g . Combining Eq. (2.49), (4.10), and (4.11), we obtain

$$\begin{aligned} S(P_0||P_g) &= W_0[\tilde{\phi}] - W_g[\tilde{\phi}] + g \langle I_1 \rangle_{g=0} \\ &= \beta_2^{\text{tree}} \cdot (1 + \alpha_2^{\text{tree}})^{-2} \int (d^4x)_E (\partial_\mu \tilde{\phi} \partial^\mu \tilde{\phi})^2 \geq 0. \end{aligned} \quad (4.12)$$

By taking $g = 1$, Eq. (4.12) represents the relative entropy between the reference and target theories and yields the following inequality.

$$S(P_{\text{NI}}||P_{\text{T}}) = \left(\beta_2^{\text{tree}} \cdot (1 + \alpha_2^{\text{tree}})^{-2} \right)_{g=1} \cdot \int (d^4x)_E (\partial_\mu \tilde{\phi} \partial^\mu \tilde{\phi})^2 \geq 0 \Rightarrow \left(\beta_2^{\text{tree}} \cdot (1 + \alpha_2^{\text{tree}})^{-2} \right)_{g=1} \geq 0, \quad (4.13)$$

where $P_{\text{NI}} = P_0$ and $P_{\text{T}} = P_{g=1}$ are used. From Eq. (4.8), this inequality represents the constraint on the coefficient of the dimension-eight operator of the effective action of the target theory.

- Loop-level UV completion — Consider the EFT generated by the loop-level UV completion. The partition function of the theory $I_0 + g \cdot I_1$ is calculated as follows:

$$Z_g[\tilde{\phi}] \equiv \int d[\phi] d[\Phi] e^{-I_g[\phi, \Phi]}$$

$$\begin{aligned}
&= \int d[\phi] \exp \left[- \int (d^4x)_E \left(-\frac{1}{2} (1 + \alpha_1^{\text{loop}} + \alpha_2^{\text{loop}}) (\partial_\mu \phi \partial^\mu \phi) - \beta_2^{\text{loop}} (\partial_\mu \phi \partial^\mu \phi)^2 \right) + E_{\text{vac}}^\Phi \right] \\
&= \exp \left[- \int (d^4x)_E \left(-\frac{1}{2} (1 + \alpha_1^{\text{loop}} + \alpha_2^{\text{loop}}) (\partial_\mu \tilde{\phi}' \partial^\mu \tilde{\phi}') - \beta_2^{\text{loop}} (\partial_\mu \tilde{\phi}' \partial^\mu \tilde{\phi}')^2 + E_{\text{vac}} \right) \right] \\
&= \exp \left[- \int (d^4x)_E \left(-\frac{1}{2} (1 + \alpha_1^{\text{loop}}) (\partial_\mu \tilde{\phi} \partial^\mu \tilde{\phi}) - \beta_2^{\text{loop}} (\partial_\mu \tilde{\phi} \partial^\mu \tilde{\phi})^2 + E_{\text{vac}} \right) \right],
\end{aligned} \tag{4.14}$$

where α_1^{loop} is the first order correction for g , α_2^{loop} and β_2^{loop} are the second or higher order correction for g , E_{vac}^Φ is the vacuum energy coming from the one-loop level correction of Φ , and E_{vac} is the vacuum energy of Φ and ϕ . We neglect two-loop effects and assume α_1^{loop} , α_2^{loop} , and β_2^{loop} are generated from the one-loop corrections of Φ . Note here that α_1^{loop} cannot be removed by redefining Φ in contrast to the tree-level UV completion. Similar to the tree-level UV completion, dimension-six and other dimension-eight operators are eliminated by suitable $\tilde{\phi}'$, which represents the classical solution of the effective action. We choose the background field as follows:

$$\tilde{\phi}' = \left(1 - \frac{1}{2} \alpha_2^{\text{loop}} \right) \cdot \tilde{\phi}, \tag{4.15}$$

with $\partial_\mu \tilde{\phi} = \text{const.}$ to remove $(\partial \partial \phi)^2$ and $(\partial \partial \partial \phi)^2$. From Eq. (4.128), the Euclidean effective actions are given by

$$W_g[\tilde{\phi}] \equiv -\ln Z_g[\tilde{\phi}] = \int (d^4x)_E \left(-\frac{1}{2} (1 + \alpha_1^{\text{loop}}) (\partial_\mu \tilde{\phi} \partial^\mu \tilde{\phi}) - \beta_2^{\text{loop}} (\partial_\mu \tilde{\phi} \partial^\mu \tilde{\phi})^2 + E_{\text{vac}} \right), \tag{4.16}$$

$$W_0[\tilde{\phi}] = \lim_{g \rightarrow 0} W_g[\tilde{\phi}] = \int (d^4x)_E \left(-\frac{1}{2} (\partial_\mu \tilde{\phi} \partial^\mu \tilde{\phi}) + E_{\text{vac}} \right). \tag{4.17}$$

From Eq. (4.16) and (4.17), the shift of the Euclidean effective action is calculated as

$$W_g[\tilde{\phi}] - W_0[\tilde{\phi}] = -\frac{1}{2} \alpha_1^{\text{loop}} \cdot \int (d^4x)_E (\partial_\mu \tilde{\phi} \partial^\mu \tilde{\phi}) - \beta_2^{\text{loop}} \int (d^4x)_E (\partial_\mu \tilde{\phi} \partial^\mu \tilde{\phi})^2. \tag{4.18}$$

Also, Eq. (4.16) yields the following relations.

$$\begin{aligned}
\left(\frac{dW_g}{dg} \right)_{g=0} &= \left(\frac{\partial W_g}{\partial g} \right)_{g=0} + \int (d^4x)_E \left(\frac{\delta W_g}{\delta \tilde{\phi}'} \right) \cdot \left(\frac{d\tilde{\phi}'}{dg} \right)_{g=0} \\
&= \left(\frac{\partial W_g}{\partial g} \right)_{g=0} \\
&= \langle I_1 \rangle_{g=0} \\
&= -\frac{1}{2} \frac{d\alpha_1^{\text{loop}}}{dg} \cdot \int (d^4x)_E (\partial_\mu \tilde{\phi} \partial^\mu \tilde{\phi}),
\end{aligned} \tag{4.19}$$

where $(d\tilde{\phi}'/dg)_{g=0} = 0$ is used in the second line, and the line is derived from Eq. (4.16). Note here that α_1^{loop} denotes the first order correction for g and satisfies a relation of the form $g \cdot (d\alpha_1^{\text{loop}}/dg) = \alpha_1^{\text{loop}}$. Combining Eqs. (2.49), (4.18), and (4.19), we obtain

$$\begin{aligned} S(P_0||P_g) &= W_0[\tilde{\phi}] - W_g[\tilde{\phi}] + g\langle I_I \rangle_{g=0} \\ &= W_0[\tilde{\phi}] - W_g^{\text{non-lin}}[\tilde{\phi}] \\ &= \beta_2^{\text{loop}} \int (d^4x)_E (\partial_\mu \tilde{\phi} \partial^\mu \tilde{\phi})^2 \geq 0, \end{aligned} \quad (4.20)$$

where $g \cdot (d\alpha_1^{\text{loop}}/dg) = \alpha_1^{\text{loop}}$ was used, and we defined as follows:

$$\begin{aligned} W_g^{\text{non-lin}}[\tilde{\phi}] &\equiv W_g[\tilde{\phi}] - g\langle I_I \rangle_{g=0} \\ &= \int (d^4x)_E \left(-\frac{1}{2} (\partial_\mu \tilde{\phi} \partial^\mu \tilde{\phi}) - \beta_2^{\text{loop}} (\partial_\mu \tilde{\phi} \partial^\mu \tilde{\phi})^2 + E_{\text{vac}} \right). \end{aligned} \quad (4.21)$$

By taking $g = 1$, Eq. (4.20) represents the relative entropy between the reference and target theories and yields the following inequality,

$$S(P_{\text{NI}}||P_{\text{T}}) = \beta_2^{\text{loop}}|_{g=1} \cdot \int (d^4x)_E (\partial_\mu \tilde{\phi} \partial^\mu \tilde{\phi})^2 \geq 0 \Rightarrow \beta_2^{\text{loop}}|_{g=1} \geq 0, \quad (4.22)$$

where $P_{\text{NI}} = P_0$ and $P_{\text{T}} = P_{g=1}$ are used. This inequality represents the constraint on the dimension-eight operator generated at the loop-level in the target theory.

According to Eq. (4.13) and (4.22), for both tree and loop level-UV completion, the relative entropy between the reference and target theories denotes the dimension-eight operator effects on the effective action. By demanding $\partial_\mu \partial^\mu \tilde{\phi} = 0$ with constant $\partial^\mu \tilde{\phi}$, after the Wick rotation, the non-negativity of the relative entropy gives rise to a constraint on Eq. (4.4) as follows:

$$S(P_{\text{NI}}||P_{\text{T}}) = \frac{c}{M^4} \int (d^4x)_E (\partial_\mu \tilde{\phi} \partial^\mu \tilde{\phi})^2 \geq 0 \Rightarrow \frac{c}{M^4} \geq 0. \quad (4.23)$$

Consequently, the coefficient c must be positive to respect the entropy constraint for both tree and loop-level UV completion. This result is consistent with the positivity bound from the unitarity and causality [51].

4.2 SMEFT dimension-eight gauge bosonic operators

In this section, using the technique provided in Sec. 2, we consider the entropy constraint on the SMEFT dimension-eight gauge bosonic operators. We list the operator basis and evaluate its constraints in the following way.

4.2.1 Operator Basis

In this section, we consider the EFT described by the following Lagrangian in the Minkowski space,

$$\mathcal{L}_{\text{SMEFT}} = -\frac{1}{4} B_{\mu\nu} B^{\mu\nu} - \frac{1}{4} W_{\mu\nu}^I W^{I,\mu\nu} - \frac{1}{4} G_{\mu\nu}^a G^{a,\mu\nu} + \frac{1}{M^4} \sum_i c_i \mathcal{O}_i, \quad (4.24)$$

where $F_{\mu\nu}^a \equiv \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + g f^{abc} A_\mu^b A_\nu^c$ is the field strength of the gauge field A_μ^a , and g denotes the gauge coupling of $SU(N)$. The gauge fields for $U(1)_Y$ hypercharge, $SU(2)_L$ weak isospin, and $SU(3)_C$ color are expressed as B_μ , W_μ^I , and G_μ^a , respectively. Also, their gauge field strengths are defined as

$$B_{\mu\nu} = \partial_\mu B_\nu - \partial_\nu B_\mu, \quad (4.25)$$

$$W_{\mu\nu}^I = \partial_\mu W_\nu^I - \partial_\nu W_\mu^I + g_2 \epsilon^{IJK} W_\mu^J W_\nu^K, \quad (4.26)$$

$$G_{\mu\nu}^a = \partial_\mu G_\nu^a - \partial_\nu G_\mu^a + g_3 f^{abc} G_\mu^b G_\nu^c, \quad (4.27)$$

where g_2 , and g_3 denote the gauge couplings of $SU(2)_L$, and $SU(3)_C$, respectively. The Greek letters stand for Lorentz indices, the Italic letters represent $SU(N)$ color indices, and totally antisymmetric and symmetric structure constants are defined by

$$[T^a, T^b] = i f^{abc} T^c, \quad (4.28)$$

$$\{T^a, T^b\} = \delta^{ab} \frac{\hat{1}}{N} + d^{abc} T^c, \quad (4.29)$$

with the generator T^a of $SU(N)$ Lie algebra. In general, the CP violating renormalizable term arises, but we assume such terms are removed by some mechanism, e.g., axion-like degrees of freedom in the UV theory. The last term of Eq. (4.24) denotes the SMEFT dimension-eight gauge bosonic operators. The basis of independent dimension-eight gauge bosonic operators for the gauge fields B_μ , W_μ^I , and G_μ^a are listed in the following [9, 10, 64, 65].

- $U(1)_Y$ — The single field strength quartics with the gauge fields of the $U(1)_Y$ hypercharge are listed as follows:

$$\mathcal{O}_1^{B^4} = (B_{\mu\nu} B^{\mu\nu})(B_{\rho\sigma} B^{\rho\sigma}), \quad (4.30)$$

$$\mathcal{O}_2^{B^4} = (B_{\mu\nu} \tilde{B}^{\mu\nu})(B_{\rho\sigma} \tilde{B}^{\rho\sigma}), \quad (4.31)$$

$$\tilde{\mathcal{O}}_1^{B^4} = (B_{\mu\nu} B^{\mu\nu})(B_{\rho\sigma} \tilde{B}^{\rho\sigma}). \quad (4.32)$$

- $SU(2)_L$ — The single field strength quartics with the gauge fields of the $SU(2)_L$ weak isospin are listed as follows:

$$\mathcal{O}_1^{W^4} = (W_{\mu\nu}^I W^{I,\mu\nu})(W_{\rho\sigma}^J W^{J,\rho\sigma}), \quad (4.33)$$

$$\mathcal{O}_2^{W^4} = (W_{\mu\nu}^I \tilde{W}^{I,\mu\nu})(W_{\rho\sigma}^J \tilde{W}^{J,\rho\sigma}), \quad (4.34)$$

$$\mathcal{O}_3^{W^4} = (W_{\mu\nu}^I W^{J,\mu\nu})(W_{\rho\sigma}^I W^{J,\rho\sigma}), \quad (4.35)$$

$$\mathcal{O}_4^{W^4} = (W_{\mu\nu}^I \tilde{W}^{J,\mu\nu})(W_{\rho\sigma}^I \tilde{W}^{J,\rho\sigma}), \quad (4.36)$$

$$\tilde{\mathcal{O}}_1^{W^4} = (W_{\mu\nu}^I W^{I,\mu\nu})(W_{\rho\sigma}^J \tilde{W}^{J,\rho\sigma}), \quad (4.37)$$

$$\tilde{\mathcal{O}}_2^{W^4} = (W_{\mu\nu}^I W^{J,\mu\nu})(W_{\rho\sigma}^I \tilde{W}^{J,\rho\sigma}). \quad (4.38)$$

- $SU(3)_C$ — The single field strength quartics with the gauge fields of the $SU(3)_C$ color are listed as follows:

$$\mathcal{O}_1^{G^4} = (G_{\mu\nu}^a G^{a,\mu\nu})(G_{\rho\sigma}^b G^{b,\rho\sigma}), \quad (4.39)$$

$$\mathcal{O}_2^{G^4} = (G_{\mu\nu}^a \tilde{G}^{a,\mu\nu})(G_{\rho\sigma}^b \tilde{G}^{b,\rho\sigma}), \quad (4.40)$$

$$\mathcal{O}_3^{G^4} = (G_{\mu\nu}^a G^{b,\mu\nu})(G_{\rho\sigma}^a G^{b,\rho\sigma}), \quad (4.41)$$

$$\mathcal{O}_4^{G^4} = (G_{\mu\nu}^a \tilde{G}^{b,\mu\nu})(G_{\rho\sigma}^a \tilde{G}^{b,\rho\sigma}), \quad (4.42)$$

$$\mathcal{O}_5^{G^4} = d^{abe} d^{cde} (G_{\mu\nu}^a G^{b,\mu\nu})(G_{\rho\sigma}^c G^{d,\rho\sigma}), \quad (4.43)$$

$$\mathcal{O}_6^{G^4} = d^{abe} d^{cde} (G_{\mu\nu}^a \tilde{G}^{b,\mu\nu})(G_{\rho\sigma}^c \tilde{G}^{d,\rho\sigma}), \quad (4.44)$$

$$\tilde{\mathcal{O}}_1^{G^4} = (G_{\mu\nu}^a G^{a,\mu\nu})(G_{\rho\sigma}^b \tilde{G}^{b,\rho\sigma}), \quad (4.45)$$

$$\tilde{\mathcal{O}}_2^{G^4} = (G_{\mu\nu}^a G^{b,\mu\nu})(G_{\rho\sigma}^a \tilde{G}^{b,\rho\sigma}), \quad (4.46)$$

$$\tilde{\mathcal{O}}_3^{G^4} = d^{abe} d^{cde} (G_{\mu\nu}^a G^{b,\mu\nu})(G_{\rho\sigma}^c \tilde{G}^{d,\rho\sigma}). \quad (4.47)$$

In addition to the dimension-eight operators, the dimension-six gauge bosonic operators may arise as follows:

$$\mathcal{O}^{W^3} = \epsilon^{IJK} W_\mu^{I,\nu} W_\nu^{J,\rho} W_\rho^{K,\mu}, \quad (4.48)$$

$$\tilde{\mathcal{O}}^{W^3} = \epsilon^{IJK} W_\mu^{I,\nu} W_\nu^{J,\rho} \tilde{W}_\rho^{K,\mu}, \quad (4.49)$$

$$\mathcal{O}^{G^3} = f^{abc} G_\mu^{a,\nu} G_\nu^{b,\rho} G_\rho^{c,\mu}, \quad (4.50)$$

$$\tilde{\mathcal{O}}^{G^3} = f^{abc} G_\mu^{a,\nu} G_\nu^{b,\rho} \tilde{G}_\rho^{c,\mu}. \quad (4.51)$$

As discussed later, we can remove the above dimension-six operator corrections to the Euclidean effective action by choosing suitable background fields. Throughout this section, CP -violating operators are denoted with a tilde.

4.2.2 Linear combination of single field strength quartics

For later convenience, we summarize building blocks to calculate the relative entropy. The dimension-eight gauge bosonic operators with a single type of the $SU(N)$ gauge field are expressed as [9, 10, 64, 65]

$$\mathcal{O}_1^{F^4} = (F_{\mu\nu}^a F^{a,\mu\nu})(F_{\rho\sigma}^b F^{b,\rho\sigma}), \quad (4.52)$$

$$\mathcal{O}_2^{F^4} = (F_{\mu\nu}^a \tilde{F}^{a,\mu\nu})(F_{\rho\sigma}^b \tilde{F}^{b,\rho\sigma}), \quad (4.53)$$

$$\mathcal{O}_3^{F^4} = (F_{\mu\nu}^a F^{b,\mu\nu})(F_{\rho\sigma}^a F^{b,\rho\sigma}), \quad (4.54)$$

$$\mathcal{O}_4^{F^4} = (F_{\mu\nu}^a \tilde{F}^{b,\mu\nu})(F_{\rho\sigma}^a \tilde{F}^{b,\rho\sigma}), \quad (4.55)$$

$$\mathcal{O}_5^{F^4} = d^{abe} d^{cde} (F_{\mu\nu}^a F^{b,\mu\nu})(F_{\rho\sigma}^c F^{d,\rho\sigma}), \quad (4.56)$$

$$\mathcal{O}_6^{F^4} = d^{abe} d^{cde} (F_{\mu\nu}^a \tilde{F}^{b,\mu\nu})(F_{\rho\sigma}^c \tilde{F}^{d,\rho\sigma}), \quad (4.57)$$

$$\mathcal{O}_7^{F^4} = d^{ace} d^{bde} (F_{\mu\nu}^a F^{b,\mu\nu})(F_{\rho\sigma}^c F^{d,\rho\sigma}), \quad (4.58)$$

$$\mathcal{O}_8^{F^4} = d^{ace} d^{bde} (F_{\mu\nu}^a \tilde{F}^{b,\mu\nu}) (F_{\rho\sigma}^c \tilde{F}^{d,\rho\sigma}), \quad (4.59)$$

$$\tilde{\mathcal{O}}_1^{F^4} = (F_{\mu\nu}^a F^{a,\mu\nu}) (F_{\rho\sigma}^b \tilde{F}^{b,\rho\sigma}), \quad (4.60)$$

$$\tilde{\mathcal{O}}_2^{F^4} = (F_{\mu\nu}^a F^{b,\mu\nu}) (F_{\rho\sigma}^a \tilde{F}^{b,\rho\sigma}), \quad (4.61)$$

$$\tilde{\mathcal{O}}_3^{F^4} = d^{abe} d^{cde} (F_{\mu\nu}^a F^{b,\mu\nu}) (F_{\rho\sigma}^c \tilde{F}^{d,\rho\sigma}), \quad (4.62)$$

$$\tilde{\mathcal{O}}_4^{F^4} = d^{ace} d^{bde} (F_{\mu\nu}^a F^{b,\mu\nu}) (F_{\rho\sigma}^c \tilde{F}^{d,\rho\sigma}), \quad (4.63)$$

where the totally antisymmetric and symmetric structure constants were defined in Eq. (4.28) and (4.29). We assume the above higher dimensional operators are produced by integrating out the heavy degrees of freedom through the interaction of Eq. (4.1). We apply the entropy inequalities of Sec. 2 and provide constraints on the Wilson coefficients of these operators.

To derive the bounds on the Wilson coefficients, we need the classical solution of the leading-order equation of motion of Eq. (4.24),

$$\partial^\mu F_{\mu\nu}^a + g f^{abc} A^{\mu,b} F_{\mu\nu}^c = 0. \quad (4.64)$$

Although the result of the inequalities derived from the relative entropy would depend on the choice of classical solutions, for simplicity, we focus on a class of solutions defined as follows:

$$\bar{A}_\mu^a = u_1^a \epsilon_{1,\mu} w_1 + u_2^a \epsilon_{2,\mu} w_2, \quad (4.65)$$

where $u_{1,2}$ is a constant real vector in $SU(N)$ color space, $\epsilon_{1,2}$ is a constant four-vector, and $w_{1,2}$ is an arbitrary Cartesian coordinate in spacetime satisfying $\partial_\mu w_1 = l_\mu$ and $\partial_\mu w_2 = k_\mu$ with l_μ and k_μ being constant four-vectors. In this work, we consider the classical solutions satisfying a condition $f^{abc} u_1^a u_2^b = 0$, and then Eq. (4.64) can be expressed as

$$\partial^\mu \bar{F}_{\mu\nu}^a = 0, \quad (4.66)$$

where $\bar{F}_{\mu\nu}^a = \partial_\mu \bar{A}_\nu^a - \partial_\nu \bar{A}_\mu^a$. The effects from the dimension-six operators on the effective action also vanish by this condition. Substituting Eq. (4.65) into the field and dual field strengths yields

$$\bar{F}_{\mu\nu}^a \bar{F}^{b,\mu\nu} = 2 \left[u_1^a u_1^b \cdot A_F + u_2^a u_2^b \cdot B_F + \left(u_1^a u_2^b + u_2^a u_1^b \right) \cdot \Gamma_F \right], \quad (4.67)$$

$$\bar{F}_{\mu\nu}^a \tilde{\bar{F}}^{b,\mu\nu} = 2 \left(u_1^a u_2^b + u_2^a u_1^b \right) \cdot \Delta_F, \quad (4.68)$$

with

$$A_F \equiv \epsilon_1^2 l^2 - (\epsilon_1 \cdot l)^2, \quad (4.69)$$

$$B_F \equiv \epsilon_2^2 k^2 - (\epsilon_2 \cdot k)^2, \quad (4.70)$$

$$\Gamma_F \equiv (\epsilon_1 \cdot \epsilon_2)(l \cdot k) - (\epsilon_1 \cdot k)(\epsilon_2 \cdot l), \quad (4.71)$$

$$\Delta_F \equiv \epsilon^{\mu\nu\rho\sigma} \epsilon_{1,\mu} l_\nu \epsilon_{2,\rho} k_\sigma. \quad (4.72)$$

From Eqs. (4.52)-(4.63), and (4.67)-(4.68), the SMEFT operators involving the background fields are expressed as follows:

$$\mathcal{O}_1^{F^4} = 4 \left[u_1^2 A_F + u_2^2 B_F + 2(u_1 \cdot u_2) \Gamma_F \right]^2, \quad (4.73)$$

$$\mathcal{O}_2^{F^4} = 16(u_1 \cdot u_2)^2 \Delta_F^2, \quad (4.74)$$

$$\begin{aligned} \mathcal{O}_3^{F^4} = 4 \Big[& (u_1^2)^2 \mathbf{A}_F^2 + (u_2^2)^2 \mathbf{B}_F^2 + 2(u_1 \cdot u_2)^2 \mathbf{A}_F \mathbf{B}_F \\ & + 2((u_1 \cdot u_2)^2 + u_1^2 u_2^2) \Gamma_F^2 + 4(u_1 \cdot u_2)(u_1^2 \mathbf{A}_F + u_2^2 \mathbf{B}_F) \Gamma_F \Big], \end{aligned} \quad (4.75)$$

$$\mathcal{O}_4^{F^4} = 8[(u_1 \cdot u_2)^2 + u_1^2 u_2^2] \Delta_F^2, \quad (4.76)$$

$$\begin{aligned} \mathcal{O}_5^{F^4} = 4 \Big[& \mathbf{A}_F^2 V_F^2 + \mathbf{B}_F^2 W_F^2 + 4\Gamma_F^2 U_F^2 \\ & + 2\mathbf{A}_F \mathbf{B}_F V_F \cdot W_F + 4\mathbf{A}_F \Gamma_F V_F \cdot U_F + 4\mathbf{B}_F \Gamma_F W_F \cdot U_F \Big], \end{aligned} \quad (4.77)$$

$$\mathcal{O}_6^{F^4} = 16U_F^2 \Delta_F^2, \quad (4.78)$$

$$\begin{aligned} \mathcal{O}_7^{F^4} = 4 \Big[& \mathbf{A}_F^2 V_F^2 + \mathbf{B}_F^2 W_F^2 + 2\Gamma_F^2 (U_F^2 + V_F \cdot W_F) \\ & + 2\mathbf{A}_F \mathbf{B}_F U_F^2 + 4\mathbf{A}_F \Gamma_F V_F \cdot U_F + 4\mathbf{B}_F \Gamma_F W_F \cdot U_F \Big], \end{aligned} \quad (4.79)$$

$$\mathcal{O}_8^{F^4} = 8(V_F \cdot W_F + U_F^2) \Delta_F^2, \quad (4.80)$$

$$\tilde{\mathcal{O}}_1^{F^4} = 8(u_1 \cdot u_2)(\mathbf{A}_F u_1^2 + 2\Gamma_F(u_1 \cdot u_2) + \mathbf{B}_F u_2^2) \Delta_F, \quad (4.81)$$

$$\tilde{\mathcal{O}}_2^{F^4} = 8[(u_1 \cdot u_2)(\Gamma_F(u_1 \cdot u_2) + \mathbf{B}_F u_2^2) + u_1^2(\mathbf{A}_F(u_1 \cdot u_2) + \Gamma_F u_2^2)] \Delta_F, \quad (4.82)$$

$$\tilde{\mathcal{O}}_3^{F^4} = 8(\mathbf{A}_F V_F \cdot U_F + 2\Gamma_F U_F^2 + \mathbf{B}_F U_F \cdot W_F) \Delta_F, \quad (4.83)$$

$$\tilde{\mathcal{O}}_4^{F^4} = 8[\mathbf{A}_F V_F \cdot U_F + \mathbf{B}_F U_F \cdot W_F + \Gamma_F(U_F^2 + V_F \cdot W_F)] \Delta_F, \quad (4.84)$$

where we defined as $U_F^a = d^{abc} u_1^b u_2^c$, $V_F^a = d^{abc} u_1^b u_2^c$, and $W_F^a = d^{abc} u_2^b u_1^c$. Then, the SMEFT operator effects on the effective action of the target theory in the Minkowski space are expressed as

$$\begin{aligned} \frac{1}{M^4} \sum_i \int d^4 x c_i \mathcal{O}_i = \frac{1}{M^4} \int d^4 x \Big[& a_F \cdot \mathbf{A}_F^2 + b_F \cdot \mathbf{B}_F^2 + c_F \cdot \Gamma_F^2 + d_F \cdot \Delta_F^2 + e_F \cdot \mathbf{A}_F \mathbf{B}_F \\ & + f_F \cdot \mathbf{A}_F \Gamma_F + g_F \cdot \mathbf{B}_F \Gamma_F + h_F \cdot \mathbf{A}_F \Delta_F + i_F \cdot \mathbf{B}_F \Delta_F + j_F \cdot \Gamma_F \Delta_F \Big], \end{aligned} \quad (4.85)$$

where the second and last lines are obtained by substituting Eq. (4.73)-(4.84) into the first line, and we defined linear combinations of the Wilson coefficients as follows:

$$a_F = 4(u_1^2)^2 c_1^{F^4} + 4(u_2^2)^2 c_3^{F^4} + 4V_F^2 c_5^{F^4} + 4V_F^2 c_7^{F^4}, \quad (4.86)$$

$$b_F = 4(u_2^2)^2 c_1^{F^4} + 4(u_2^2)^2 c_3^{F^4} + 4W_F^2 c_5^{F^4} + 4W_F^2 c_7^{F^4}, \quad (4.87)$$

$$c_F = 8 \left[2c_1^{F^4} (u_1 \cdot u_2)^2 + c_3^{F^4} ((u_1 \cdot u_2)^2 + u_1^2 u_2^2) + 2c_5^{F^4} U_F^2 + c_7^{F^4} (U_F^2 + V_F \cdot W_F) \right], \quad (4.88)$$

$$d_F = 8 \left[2c_2^{F^4} (u_1 \cdot u_2)^2 + c_4^{F^4} ((u_1 \cdot u_2)^2 + u_1^2 u_2^2) + 2c_6^{F^4} U_F^2 + c_8^{F^4} (U_F^2 + V_F \cdot W_F) \right], \quad (4.89)$$

$$e_F = 8u_1^2 u_2^2 c_1^{F^4} + 8(u_1 \cdot u_2)^2 c_3^{F^4} + 8V_F \cdot W_F c_5^{F^4} + 8U_F^2 c_7^{F^4}, \quad (4.90)$$

$$f_F = 16u_1^2(u_1 \cdot u_2)c_1^{F^4} + 16(u_1 \cdot u_2)u_1^2c_3^{F^4} + 16V_F \cdot U_F c_5^{F^4} + 16V_F \cdot U_F c_7^{F^4}, \quad (4.91)$$

$$g_F = 16u_2^2(u_1 \cdot u_2)c_1^{F^4} + 16u_2^2(u_1 \cdot u_2)c_3^{F^4} + 16W_F \cdot U_F c_5^{F^4} + 16U_F \cdot W_F c_7^{F^4}, \quad (4.92)$$

$$h_F = 8u_1^2(u_1 \cdot u_2)(\tilde{c}_1^{F^4} + \tilde{c}_2^{F^4}) + 8V_F \cdot U_F(\tilde{c}_3^{F^4} + \tilde{c}_4^{F^4}), \quad (4.93)$$

$$i_F = 8u_2^2(u_1 \cdot u_2)(\tilde{c}_1^{F^4} + \tilde{c}_2^{F^4}) + 8U_F \cdot W_F(\tilde{c}_3^{F^4} + \tilde{c}_4^{F^4}), \quad (4.94)$$

$$j_F = 8 \left[2\tilde{c}_1^{F^4}(u_1 \cdot u_2)^2 + \tilde{c}_2^{F^4}((u_1 \cdot u_2)^2 + u_1^2 u_2^2) + 2\tilde{c}_3^{F^4}U_F^2 + \tilde{c}_4^{F^4}(U_F^2 + V_F \cdot W_F) \right]. \quad (4.95)$$

For $U(1)_Y$, $SU(2)_L$, and $SU(3)_C$, the above quantities are respectively listed as follows:

- $U(1)_Y$ — For the $U(1)_Y$ gauge field B_μ , Eqs. (4.86)-(4.95) are calculated as

$$a_B = b_B = c_B/4 = e_B/2 = f_B/4 = g_B/4 = 4c_1^{B^4}, \quad (4.96)$$

$$d_B = 16c_2^{B^4}, \quad (4.97)$$

$$h_B = i_B = j_B/2 = 8\tilde{c}_1^{B^4}. \quad (4.98)$$

- $SU(2)_L$ — For the $SU(2)_L$ gauge field W_μ^I , Eqs. (4.86)-(4.95) are calculated as

$$a_W = b_W = c_W/4 = e_W/2 = f_W/4 = g_W/4 = 4(u_1^2)^2(c_1^{W^4} + c_3^{W^4}), \quad (4.99)$$

$$d_W = 16(u_1^2)^2(c_2^{W^4} + c_4^{W^4}), \quad (4.100)$$

$$h_W = i_W = j_W/2 = 8(u_1^2)^2(\tilde{c}_1^{W^4} + \tilde{c}_2^{W^4}), \quad (4.101)$$

- $SU(3)_C$ — Using an identity for $SU(N)$,

$$f^{abe} f^{cde} = \frac{2}{N} (\delta^{ac} \delta^{bd} - \delta^{ad} \delta^{bc}) + d^{ace} d^{bde} - d^{bce} d^{ade}, \quad (4.102)$$

we obtain

$$V_F W_F = U_F^2 + \frac{2}{N} [(u_1 \cdot u_2)^2 - u_1^2 u_2^2]. \quad (4.103)$$

For $SU(3)_C$, additional identities hold as follows:

$$3d^{abe} d^{cde} - f^{ace} f^{bde} - f^{ade} f^{bce} = \delta^{ac} \delta^{bd} + \delta^{ad} \delta^{bc} - \delta^{ab} \delta^{cd}, \quad (4.104)$$

$$3(d^{abe} d^{cde} + d^{ace} d^{bde} + d^{ade} d^{bce}) = \delta^{ab} \delta^{cd} + \delta^{ac} \delta^{bd} + \delta^{ad} \delta^{bc}. \quad (4.105)$$

From the above identities, we obtain

$$V_G^2 = \frac{1}{3}u_1^4, \quad W_G^2 = \frac{1}{3}u_2^4, \quad U_G^2 = \frac{1}{3}u_1^2 u_2^2, \quad (4.106)$$

$$V_G \cdot U_G = \frac{1}{3}u_1^2(u_1 \cdot u_2), \quad W_G \cdot U_G = \frac{1}{3}u_2^2(u_1 \cdot u_2), \quad V_G \cdot W_G = \frac{1}{3}(-u_1^2 u_2^2 + 2(u_1 \cdot u_2)^2), \quad (4.107)$$

$$2U_G^2 + V_G W_G = \frac{1}{3}(u_1^2 u_2^2 + 2(u_1 \cdot u_2)^2). \quad (4.108)$$

Also, from the above identities, $\mathcal{O}_7^{G^4}$, $\mathcal{O}_8^{G^4}$, and $\tilde{\mathcal{O}}_4^{G^4}$ can be rewritten by the other operators, so we omit these operators. From Eqs. (4.86)-(4.95), and (4.106)-(4.108), we obtain

$$a_G = 4(u_1^2)^2 \left(c_1^{G^4} + c_3^{G^4} + \frac{1}{3}c_5^{G^4} + \frac{1}{3}c_7^{G^4} \right), \quad (4.109)$$

$$b_G = 4(u_2^2)^2 \left(c_1^{G^4} + c_3^{G^4} + \frac{1}{3}c_5^{G^4} + \frac{1}{3}c_7^{G^4} \right), \quad (4.110)$$

$$c_G = 8 \left[(2c_1^{G^4} + c_3^{G^4})(u_1 \cdot u_2)^2 + c_3^{G^4} u_1^2 u_2^2 + 2c_5^{G^4} U_G^2 \right], \quad (4.111)$$

$$d_G = 8 \left[(2c_2^{G^4} + c_4^{G^4})(u_1 \cdot u_2)^2 + c_4^{G^4} u_1^2 u_2^2 + 2c_6^{G^4} U_G^2 \right], \quad (4.112)$$

$$e_G = 8u_1^2 u_2^2 \left(c_1^{G^4} - \frac{1}{3}c_5^{G^4} \right) + 8(u_1 \cdot u_2)^2 \left(c_3^{G^4} + \frac{2}{3}c_5^{G^4} \right), \quad (4.113)$$

$$f_G = 16u_1^2 (u_1 \cdot u_2) \left(c_1^{G^4} + c_3^{G^4} + \frac{1}{3}c_5^{G^4} \right), \quad (4.114)$$

$$g_G = 16u_2^2 (u_1 \cdot u_2) \left(c_1^{G^4} + c_3^{G^4} + \frac{1}{3}c_5^{G^4} \right), \quad (4.115)$$

$$h_G = 8u_1^2 (u_1 \cdot u_2) \left(\tilde{c}_1^{G^4} + \tilde{c}_2^{G^4} + \frac{1}{3}\tilde{c}_3^{G^4} \right), \quad (4.116)$$

$$i_G = 8u_2^2 (u_1 \cdot u_2) \left(\tilde{c}_1^{G^4} + \tilde{c}_2^{G^4} + \frac{1}{3}\tilde{c}_3^{G^4} \right), \quad (4.117)$$

$$j_G = 8 \left[(2\tilde{c}_1^{G^4} + \tilde{c}_2^{G^4})(u_1 \cdot u_2)^2 + \tilde{c}_2^{G^4} u_1^2 u_2^2 + 2\tilde{c}_3^{G^4} U_G^2 \right]. \quad (4.118)$$

In the next section, we derive some constraints on the SMEFT Wilson coefficients from the above building blocks.

4.2.3 Bounds from non-interacting reference theory

We consider the NIRT as the reference theory, and evaluate the relative entropy. We assume the dimension-eight operators listed in the previous sections are generated through the interaction between heavy and light fields, i.e., $I_1[A_\mu^a, \Phi] = \int d^4x \mathcal{O}[\Phi] \otimes J[A_\mu^a]$. Then, the first order corrections for g to the Euclidean effective action are expressed as

$$\begin{aligned} g \cdot \langle I_1 \rangle_{g=0} &= g \cdot \left(\frac{\partial W_g}{\partial g} \right)_{g=0} \\ &= \int (d^4x)_E \left(\frac{\delta W_g}{\delta J} \right)_{J=0} J[A_\mu^a], \end{aligned} \quad (4.119)$$

where $(\delta W_g / \delta J)_{J=0}$ is a tadpole-like diagram for the composite field $J[A_\mu^a]$, e.g., Fig. 6, and does not depend on space-time. Because $J[A_\mu^a]$ does not include the higher-dimensional operators according to Assumption (IV), there are two cases: (i) $J[A_\mu^a]$ preserves the gauge symmetry or (ii) not. For case (i), Eq. (4.119) is proportional to $\int (d^4x)_E F_{\mu\nu}^a F^{a,\mu\nu}$ because $\langle I_1 \rangle_{g=0}$ preserves the Lorentz symmetry from the definition, and $(\delta W_g / \delta J)_{J=0}$ does not depend on space-time. In general, the CP violating

term arises, but we omit such terms because they can be removed by some mechanism, such as axion-like degrees of freedom in the UV theory. For case (ii), $J[A_\mu^a]$ can be proportional to A_μ^a , and $A_\mu^a A_\nu^a$ because of the covariant derivative of the kinetic term. According to Assumption (IV), we focus on the leading order of the interacting term, which are corrections from the kinetic terms of heavy fields. However, $J[A_\mu^a] \propto A_\mu^a$ effects on Eq. (4.119) vanish because $\langle I_1 \rangle_{g=0}$ keeps the Lorentz symmetry. Although terms proportional to $\int (d^4x)_E A_\mu^a A^{a,\mu}$ in Eq. (4.119) may remain, they can be eliminated by implementing the gauge fixing condition, the so-called non-linear gauge [61]. Therefore, we focus on the case of $\langle I_1 \rangle_{g=0} \propto \int (d^4x)_E F_{\mu\nu}^a F^{a,\mu\nu}$ below. For each tree and loop-level UV completions, the constraints on the SMEFT Wilson coefficients from the relative entropy are evaluated as follows:

- Tree-level UV completion — Consider the SMEFT operators generated by the tree-level UV completions. The partition function of the theory $I_0 + g \cdot I_1$ is generally calculated as follows:

$$\begin{aligned}
Z_g[\bar{A}] &\equiv \int d[A] d[\Phi] e^{-I_g[A, \Phi]} \\
&= \int d[A] \exp \left[- \int (d^4x)_E \left(\frac{1}{2} (1 + \alpha_2^{\text{tree}}) F_{\mu\nu}^a F^{a,\mu\nu} - \sum_i \beta_{i,2}^{\text{tree}} \mathcal{O}_i[A] \right) \right] \\
&= \exp \left[- \int (d^4x)_E \left(\frac{1}{2} (1 + \alpha_2^{\text{tree}}) \bar{F}_{\mu\nu}^a \bar{F}^{a,\mu\nu} - \sum_i \beta_{i,2}^{\text{tree}} \mathcal{O}_i[\bar{A}] \right) \right] \\
&= \exp \left[- \int (d^4x)_E \left(\frac{1}{2} \bar{F}_{\mu\nu}^a \bar{F}^{a,\mu\nu} - \sum_i \beta_{i,2}^{\text{tree}} \cdot (1 + \alpha_2^{\text{tree}})^{-2} \mathcal{O}_i[\bar{A}] \right) \right], \quad (4.120)
\end{aligned}$$

where $\mathcal{O}_i[A]$ is the dimension-eight SMEFT operators, α_2^{tree} and $\beta_{i,2}^{\text{tree}}$ denote the second or higher order corrections for g , and $\beta_{i,2}^{\text{tree}}$ does not include the first order correction for g because of Assumption (IV). The corrections α_2^{tree} and $\beta_{i,2}^{\text{tree}}$ are generated at the tree level. Following the procedure in Eq. (2.68), (2.69), and (2.70), the first order correction for g is eliminated in α_2^{tree} . The background field \bar{A}_μ^a denotes the classical solution of the effective action. We choose the background fields as follows:

$$\bar{A}_\mu^a = (1 + \alpha_2^{\text{tree}})^{-1/2} \cdot \bar{A}_\mu^a, \quad (4.121)$$

with $\bar{F}_{\mu\nu}^a = \text{const.}$ In general, the dimension-six operators arise in Eq. (4.120) but can be eliminated by choosing the suitable background fields of Eq. (4.65). From Eq. (4.120), the Euclidean effective actions are calculated as follows:

$$W_g[\bar{A}] \equiv -\ln Z_g[\bar{A}] = \int (d^4x)_E \left(\frac{1}{2} \bar{F}_{\mu\nu}^a \bar{F}^{a,\mu\nu} - \sum_i \beta_{i,2}^{\text{tree}} \cdot (1 + \alpha_2^{\text{tree}})^{-2} \mathcal{O}_i[\bar{A}] \right), \quad (4.122)$$

$$W_0[\bar{A}] = \lim_{g \rightarrow 0} W_g[\bar{A}] = \int (d^4x)_E \left(\frac{1}{2} \bar{F}_{\mu\nu}^a \bar{F}^{a,\mu\nu} \right), \quad (4.123)$$

From Eqs. (4.122) and (4.123), the shift of the Euclidean effective action is calculated as follows:

$$W_g[\bar{A}] - W_0[\bar{A}] = - \sum_i \beta_{i,2}^{\text{tree}} \cdot (1 + \alpha_2^{\text{tree}})^{-2} \int (d^4x)_E \mathcal{O}_i[\bar{A}]. \quad (4.124)$$

From Eq. (4.122), the first order corrections for g is also calculated as

$$\begin{aligned} \left(\frac{dW_g}{dg} \right)_{g=0} &= \left(\frac{\partial W_g}{\partial g} \right)_{g=0} + \int (d^4x)_E \left(\frac{\delta W_g}{\delta \bar{A}'} \right) \cdot \left(\frac{d\bar{A}'}{dg} \right)_{g=0} \\ &= \left(\frac{\partial W_g}{\partial g} \right)_{g=0} \\ &= \langle I_I \rangle_{g=0} = 0, \end{aligned} \quad (4.125)$$

where $(d\bar{A}'/dg)_{g=0} = 0$ holds because of Eq. (4.121). From Eqs. (2.49), (4.124), and (4.125), we obtain

$$\begin{aligned} S(P_0||P_g) &= W_0[\bar{A}] - W_g[\bar{A}] + g\langle I_I \rangle_{g=0} \\ &= W_0[\bar{A}] - W_g[\bar{A}] \\ &= \sum_i \beta_{i,2}^{\text{tree}} \cdot (1 + a_2^{\text{tree}})^{-2} \int (d^4x)_E \mathcal{O}_i[\bar{A}] \geq 0. \end{aligned} \quad (4.126)$$

Note here that the relative entropy does not change even if we add the same term to both W_0 and W_g simultaneously⁵. By taking $g = 1$, Eq. (4.126) represents the relative entropy between the reference and target theories and yields the following inequality.

$$S(P_{\text{NI}}||P_{\text{T}}) = \sum_i (\beta_{i,2}^{\text{tree}} \cdot (1 + a_2^{\text{tree}})^{-2})_{g=1} \int (d^4x)_E \mathcal{O}_i[\bar{A}] \geq 0, \quad (4.127)$$

where $P_{\text{NI}} = P_0$ and $P_{\text{T}} = P_{g=1}$ are used. The right-hand side of this inequality denotes the linear combination of the coefficients of the dimension-eight operators of the target theory.

- Loop-level UV completion — Consider the SMEFT operators generated by the loop-level UV completion. The partition function of the theory $I_0 + g \cdot I_I$ is generally calculated as follows⁶:

$$\begin{aligned} Z_g[\bar{A}] &\equiv \int d[A] d[\Phi] e^{-I_g[A, \Phi]} \\ &= \int d[A] \exp \left[- \int (d^4x)_E \left(\frac{1}{2} \left(1 + \alpha_1^{\text{loop}} + \alpha_2^{\text{loop}} \right) F_{\mu\nu}^a F^{a, \mu\nu} - \sum_i \beta_{2,i}^{\text{loop}} \mathcal{O}_i[A] + E_{\text{vac}}^\Phi \right) \right] \\ &= \exp \left[- \int (d^4x)_E \left(\frac{1}{2} \left(1 + \alpha_1^{\text{loop}} + \alpha_2^{\text{loop}} \right) \bar{F}_{\mu\nu}^a \bar{F}^{a, \mu\nu} - \sum_i \beta_{2,i}^{\text{loop}} \mathcal{O}_i[\bar{A}] + E_{\text{vac}} \right) \right] \\ &= \exp \left[- \int (d^4x)_E \left(\frac{1}{2} \left(1 + \alpha_1^{\text{loop}} \right) \bar{F}_{\mu\nu}^a \bar{F}^{a, \mu\nu} - \sum_i \beta_{2,i}^{\text{loop}} \mathcal{O}_i[\bar{A}] + E_{\text{vac}} \right) \right]. \end{aligned} \quad (4.128)$$

where $\mathcal{O}_i[A]$ is the dimension-eight SMEFT operators, α_1^{loop} is the first order correction for g , α_2^{loop} and $\beta_{2,i}^{\text{loop}}$ are the second or higher order correction for g , E_{vac}^Φ is the vacuum energy

⁵We can also put the same boundary terms to both W_0 and W_g .

⁶Strictly speaking, Eq. (4.128) holds in the limit of $g = 1$ or $g = 0$. This is because the gauge symmetry may be broken by the auxiliary parameter g . However, this subtle point does not affect the following discussions because we focus only on the reference ($g = 0$) and target ($g = 1$) theories, where the gauge symmetry is restored.

coming from the one-loop level correction of Φ , and E_{vac} is the vacuum energy of Φ and A_μ^a . α_1^{loop} , α_2^{loop} , and $\beta_{2,i}^{\text{loop}}$ are generated from the one-loop corrections of Φ . In Eq. (4.128), we neglected two-loop corrections. Note here that α_1^{loop} cannot be removed by redefining Φ in contrast to the tree-level UV completion. The background field \bar{A}_μ^a denotes Eq. (4.65), which is the solution of the effective action. We choose the background field as follows:

$$\bar{A}_\mu^a = \left(1 - \frac{1}{2}\alpha_2^{\text{loop}}\right) \bar{A}_\mu^a, \quad (4.129)$$

where $\bar{F}_{\mu\nu}^a = \text{const.}$ to remove the dimension-six operators; see Eq. (4.65). From Eq. (4.128), the Euclidean effective actions are obtained as follows:

$$W_g[\bar{A}] \equiv -\ln Z_g[\bar{A}] = \int (d^4x)_{\text{E}} \left(\frac{1}{2} \left(1 + \alpha_1^{\text{loop}}\right) \bar{F}_{\mu\nu}^a \bar{F}^{a,\mu\nu} - \sum_i \beta_{2,i}^{\text{loop}} \mathcal{O}_i[\bar{A}] + E_{\text{vac}} \right), \quad (4.130)$$

$$W_0[\bar{A}] = \lim_{g \rightarrow 0} W_g[\bar{A}] = \int (d^4x)_{\text{E}} \left(\frac{1}{2} \bar{F}_{\mu\nu}^a \bar{F}^{a,\mu\nu} + E_{\text{vac}} \right). \quad (4.131)$$

The shift of the Euclidean effective action is calculated as follows:

$$W_g[\bar{A}] - W_0[\bar{A}] = \int (d^4x)_{\text{E}} \left(\frac{1}{2} \alpha_1^{\text{loop}} \bar{F}_{\mu\nu}^a \bar{F}^{a,\mu\nu} - \sum_i \beta_{2,i}^{\text{loop}} \mathcal{O}_i[\bar{A}] \right). \quad (4.132)$$

Also, the first order corrections for g is calculated as

$$\begin{aligned} \left(\frac{dW_g}{dg} \right)_{g=0} &= \left(\frac{\partial W_g}{\partial g} \right)_{g=0} + \int (d^4x)_{\text{E}} \left(\frac{\delta W_g}{\delta \bar{A}'} \right) \cdot \left(\frac{d\bar{A}'}{dg} \right)_{g=0} \\ &= \left(\frac{\partial W_g}{\partial g} \right)_{g=0} \\ &= \langle I_1 \rangle_{g=0} \\ &= \frac{1}{2} \frac{d\alpha_1^{\text{loop}}}{dg} \int (d^4x)_{\text{E}} \bar{F}_{\mu\nu}^a \bar{F}^{a,\mu\nu}, \end{aligned} \quad (4.133)$$

where $(d\bar{A}'/dg)_{g=0} = 0$ holds from Eq. (4.129). This relation represents Eq. (4.119). From Eqs. (2.49), (4.132), and (4.133), we obtain

$$\begin{aligned} S(P_0||P_g) &= W_0[\bar{A}] - W_g[\bar{A}] + g \langle I_1 \rangle_{g=0} \\ &= W_0[\bar{A}] - W_g^{\text{non-lin}}[\bar{A}] \\ &= \sum_i \beta_{2,i}^{\text{loop}} \int (d^4x)_{\text{E}} \mathcal{O}_i[\bar{A}] \geq 0. \end{aligned} \quad (4.134)$$

where $g \cdot (d\alpha_1^{\text{loop}}/dg) = \alpha_1^{\text{loop}}$ was used, and we defined as follows:

$$W_g^{\text{non-lin}}[\bar{A}] \equiv W_g[\bar{A}] - g \langle I_1 \rangle_{g=0}$$

$$= \int (d^4x)_E \left(\frac{1}{2} \bar{F}_{\mu\nu}^a \bar{F}^{a,\mu\nu} - \sum_i \beta_{2,i}^{\text{loop}} \mathcal{O}_i[\bar{A}] + E_{\text{vac}} \right). \quad (4.135)$$

Similar to the tree-level UV completion, we can also add the same boundary terms to both W_0 and $W_g^{\text{non-lin}}$ because it cancels in the relative entropy. By taking $g = 1$, Eq. (4.134) represents the relative entropy between the reference and target theories and yields the following inequality.

$$S(P_{\text{NI}} || P_{\text{T}}) = \sum_i \beta_{2,i}^{\text{loop}}|_{g=1} \int (d^4x)_E \mathcal{O}_i[\bar{A}] \geq 0, \quad (4.136)$$

where $P_{\text{NI}} = P_0$ and $P_{\text{T}} = P_{g=1}$ are used. This inequality yields the constraint on the dimension-eight operators generated at the one-loop level.

Consequently, for both tree and loop-level UV completions, it is found that the relative entropy denotes the linear combination of the dimension-eight operators generated from the interacting terms, i.e., Eqs. (4.127) and (4.136). Therefore, after Wick rotation, the inequalities (4.127) and (4.136) give rise to constraints on the dimension-eight operators of Eq. (4.24) as follows:

$$\frac{1}{M^4} \sum_i \int (d^4x)_E c_i \mathcal{O}_i[\bar{A}] \geq 0. \quad (4.137)$$

Equation (4.137) and (4.85) yield bounds as follows:

$$\begin{aligned} & a_F \cdot A_F^2 + b_F \cdot B_F^2 + c_F \cdot \Gamma_F^2 + d_F \cdot \Delta_F^2 \\ & + e_F \cdot A_F B_F + f_F \cdot A_F \Gamma_F + g_F \cdot B_F \Gamma_F + h_F \cdot A_F \Delta_F + i_F \cdot B_F \Delta_F + j_F \cdot \Gamma_F \Delta_F \geq 0. \end{aligned} \quad (4.138)$$

The quantities A_F , B_F , Γ_F , and Δ_F are independent each other, and the inequality of (4.138) yields the following inequalities.

$$a_F \geq 0, \quad b_F \geq 0, \quad c_F \geq 0, \quad d_F \geq 0, \quad (4.139)$$

$$4a_F \cdot d_F - h_F^2 \geq 0, \quad 4b_F \cdot d_F - i_F^2 \geq 0, \quad 4c_F \cdot d_F - j_F^2 \geq 0, \quad (4.140)$$

$$a_F - \frac{h_F^2}{4d_F} - \frac{\left(f_F - \frac{h_F \cdot j_F}{2d_F}\right)^2}{4\left(c_F - \frac{j_F^2}{4d_F}\right)} \geq 0, \quad b_F - \frac{i_F^2}{4d_F} - \frac{\left(g_F - \frac{i_F \cdot j_F}{2d_F}\right)^2}{4\left(c_F - \frac{j_F^2}{4d_F}\right)} \geq 0, \quad (4.141)$$

$$\begin{aligned} & 4 \left[a_F - \frac{h_F^2}{4d_F} - \frac{\left(f_F - \frac{h_F \cdot j_F}{2d_F}\right)^2}{4\left(c_F - \frac{j_F^2}{4d_F}\right)} \right] \cdot \left[b_F - \frac{i_F^2}{4d_F} - \frac{\left(g_F - \frac{i_F \cdot j_F}{2d_F}\right)^2}{4\left(c_F - \frac{j_F^2}{4d_F}\right)} \right] \\ & \geq \left[e_F - \frac{h_F \cdot i_F}{2d_F} - \frac{\left(f_F - \frac{h_F \cdot j_F}{2d_F}\right) \left(g_F - \frac{i_F \cdot j_F}{2d_F}\right)}{2\left(c_F - \frac{j_F^2}{4d_F}\right)} \right]^2. \end{aligned} \quad (4.142)$$

For $U(1)_Y$, $SU(2)_L$, and $SU(3)_C$ gauge fields, the above inequalities are listed as follows:

- $U(1)_Y$ — Substituting Eqs. (4.96)-(4.98) into the inequalities of (4.139)-(4.142), for the $U(1)_Y$ gauge field B_μ , we obtain the following constraints.

$$c_1^{B^4} \geq 0, \quad c_2^{B^4} \geq 0, \quad 4c_1^{B^4} c_2^{B^4} \geq (\tilde{c}_1^{B^4})^2. \quad (4.143)$$

Note here that the bounds of Eqs. (4.141) and (4.142) vanish by substituting Eqs. (4.96)-(4.98).

- $SU(2)_L$ — From Eqs. (4.99)-(4.101), and the inequalities of (4.139)-(4.142), for the $SU(2)_L$ gauge field W_μ^I , we obtain the following constraints.

$$c_1^{W^4} + c_3^{W^4} \geq 0, \quad c_2^{W^4} + c_4^{W^4} \geq 0, \quad 4(c_1^{W^4} + c_3^{W^4})(c_2^{W^4} + c_4^{W^4}) \geq (\tilde{c}_1^{W^4} + \tilde{c}_2^{W^4})^2. \quad (4.144)$$

Similar to the case of the $U(1)_Y$ gauge field, the bounds of Eqs. (4.141) and (4.142) vanish by substituting Eqs. (4.99), (4.100), and (4.101).

- $SU(3)_C$ — For simplicity, we assume $u_1^2 = u_2^2 = 1$ and $u_1 \cdot u_2 = \cos \xi$. By considering the two cases of $\cos^2 \xi = 0$ and 1, from Eqs. (4.109)-(4.118), and the inequalities of (4.139)-(4.142), we obtain

$$3c_1^{G^4} + 3c_3^{G^4} + c_5^{G^4} \geq 0, \quad (4.145)$$

$$3c_3^{G^4} + 2c_5^{G^4} \geq 0, \quad (4.146)$$

$$3c_2^{G^4} + 3c_4^{G^4} + c_6^{G^4} \geq 0, \quad (4.147)$$

$$3c_4^{G^4} + 2c_6^{G^4} \geq 0, \quad (4.148)$$

$$4 \left(3c_1^{G^4} + 3c_3^{G^4} + c_5^{G^4} \right) \left(3c_2^{G^4} + 3c_4^{G^4} + c_6^{G^4} \right) \geq \left(3\tilde{c}_1^{G^4} + 3\tilde{c}_2^{G^4} + \tilde{c}_3^{G^4} \right)^2, \quad (4.149)$$

$$4 \left(3c_3^{G^4} + 2c_5^{G^4} \right) \left(3c_4^{G^4} + 2c_6^{G^4} \right) \geq \left(3\tilde{c}_2^{G^4} + 2\tilde{c}_3^{G^4} \right)^2, \quad (4.150)$$

$$2c_1^{G^4} + c_3^{G^4} \geq 0. \quad (4.151)$$

The above first six bounds are the same as the positivity bounds from unitarity and causality considerations in Ref. [65], and the last inequality newly arises from the entropy constraint. Since, however, Eq. (4.145) is derived from Eqs. (4.146) and (4.151), the results of the entropy constraints are consistent with the positivity bounds from unitarity and causality considerations in Ref. [65].

4.3 Einstein-Maxwell theory with higher-derivative operators

Consider the Einstein-Maxwell theory with higher-derivative operators in the Minkowski space as follows:

$$W_{\text{EM}} = \int d^4x \sqrt{-g} \left(\frac{M_{\text{Pl}}^2}{2} R - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{\alpha_1}{4M_{\text{Pl}}^4} (F_{\mu\nu} F^{\mu\nu})^2 + \frac{\alpha_2}{4M_{\text{Pl}}^4} (F_{\mu\nu} \tilde{F}^{\mu\nu})^2 + \frac{\alpha_3}{2M_{\text{Pl}}^2} F_{\mu\nu} F_{\rho\sigma} R^{\mu\nu\rho\sigma} \right), \quad (4.152)$$

where other operators up to four-derivative are eliminated by the field redefinition of $g_{\mu\nu}$; see Ref. [49] and Appendix D. Also, the Gauss-Bonnet combination, i.e., $R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} - 4R_{\mu\nu}R^{\mu\nu} + R^2$, is a topological term that does not contribute to the extremal black hole entropy in four dimensions, so we omit it throughout this section. Similar to the previous subsections, we focus on the field theoretical description, which breaks down at some high energy scale Λ_{QFT} . Generically, Λ_{QFT} is smaller than the Planck scale $\Lambda_{\text{QFT}} \ll M_{\text{Pl}}$. In this work, we do not consider the dynamics of stringy particles in the high energy regime beyond Λ_{QFT} because the ordinary field theoretical descriptions break down by infinitely many local fields.⁷ According to Assumption (III), we consider the higher-derivative operators generated from the target UV theory defined by $I_{\text{T}}[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A, \Phi]$, where $g_{\mu\nu}$ is the metric of space-time, $R_{\mu\nu\rho\sigma}$ is the Riemann tensor, A_μ is the $U(1)$ gauge boson, and Φ is the heavy degrees of freedom. For this EFT, define the non-interacting and interacting terms as follows:

$$I_0[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A, \Phi] \equiv I_{\text{T}}[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A, 0] + I_{\text{T}}[g_{\mu\nu}; 0, 0, \Phi], \quad (4.153)$$

$$I_1[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A, \Phi] \equiv I_{\text{T}}[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A, \Phi] - I_0[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A, \Phi], \quad (4.154)$$

where the cosmological constant is omitted because it cancels in the relative entropy. These definitions are also adopted in Sec. 3.8. It should be noted that the action I_0 does not include the interaction between Φ and $A_\mu, R_{\mu\nu\rho\sigma}$, but the interaction between $g_{\mu\nu}$ and Φ . Although gravitational operators such as $R_{\mu\nu}^2$ are generated from I_0 , such operators up to four-derivative can be eliminated by the field redefinition of $g_{\mu\nu}$; see Appendix D.

We assume the dimension-eight operators of Eq. (4.152) are generated through the interaction defined in Eq. (4.1). Then, the first order corrections for g to the Euclidean effective action are expressed as follows:

$$\begin{aligned} g \cdot \langle I_1 \rangle_{g=0} &= g \cdot \left(\frac{\partial W_g}{\partial g} \right)_{g=0} \\ &= \int (d^4x)_{\text{E}} \sqrt{g} \left(\frac{\delta W_g}{\delta J} \right)_{J=0} J[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A_\mu], \end{aligned} \quad (4.155)$$

where $(\delta W_g / \delta J)_{J=0}$ is a tadpole-like diagram for the composite field J . Similar to the SMEFT, $J[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A_\mu]$ does not include the higher-derivative operators according to Assumption (IV), so there are two cases: (i) $J[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A_\mu]$ preserves the gauge symmetry or (ii) not. For case (i), Eq. (4.155) is proportional to $\int (d^4x)_{\text{E}} \sqrt{g} F_{\mu\nu} F^{\mu\nu}$ or $\int (d^4x)_{\text{E}} \sqrt{g} R$ because $\langle I_1 \rangle$ is invariant under general coordinate transformations. We assume the interaction I_1 does not involve the CP violating terms. For case (ii), $J[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A_\mu] \propto A_\mu$, or $A_\mu A_\nu$ because of the covariant derivative of the kinetic term. According to Assumption (IV), we focus on the leading order of the interacting term, which arises from the kinetic terms of the heavy charged fields. Then, $J \propto A_\mu$ effects on Eq. (4.155) vanish from the invariance of $\langle I_1 \rangle_{g=0}$ under the general coordinate transformations. Also, a term proportional to $\int (d^4x)_{\text{E}} A_\mu A^\mu$ is generated in $\langle I_1 \rangle_{g=0}$ by $J \propto A_\mu A_\nu$ effects on Eq. (4.155) because

⁷The graviton accompanied by Regge states can break the positivity bounds [53] on the Wilson coefficients of Eq. (4.152), but such a scenario would be beyond the applicability of procedures of this section based on the field theory.

of the invariance of the general coordinate transformations but can be eliminated by implementing the non-linear gauge fixing condition $A_\mu A^\mu = 0$. Then, the first order corrections of the interaction to the effective action are expressed as follows:

$$\begin{aligned}
g \cdot \langle I_1 \rangle_{g=0} &= g \cdot \left(\frac{\partial W_g}{\partial g} \right)_{g=0} \\
&= \int (d^4x)_E \sqrt{g} \left(\frac{\delta W_g}{\delta J} \right)_{J=0} J[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A_\mu] \\
&\propto \int (d^4x)_E \sqrt{g} F_{\mu\nu} F^{\mu\nu} \text{ or } \int (d^4x)_E \sqrt{g} R.
\end{aligned} \tag{4.156}$$

For each tree and loop-level UV completions, the constraints on the EFT from the relative entropy are evaluated as follows:

- **Tree-level UV completion** — Consider the EFT generated at the tree-level UV completion. The partition function of the theory $I_0 + g \cdot I_1$ is generally calculated as follows:

$$\begin{aligned}
Z_g[\bar{g}_{\mu\nu}, \bar{A}] &\equiv \int d[g]d[A]d[\Phi] e^{-I_g[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A, \Phi]} \\
&= \int d[g]d[A] \exp \left[- \int (d^4x)_E \sqrt{g} \left(- \frac{M_{\text{Pl}}^2}{2} (1 + \alpha_{2,R}^{\text{tree}}) R + \frac{1}{4} (1 + \alpha_{2,F}^{\text{tree}}) F_{\mu\nu} F^{\mu\nu} \right. \right. \\
&\quad \left. \left. - \beta_{2,1}^{\text{tree}} (F_{\mu\nu} F^{\mu\nu})^2 - \beta_{2,2}^{\text{tree}} (F_{\mu\nu} \widetilde{F}^{\mu\nu})^2 - \beta_{2,3}^{\text{tree}} F_{\mu\nu} F_{\rho\sigma} R^{\mu\nu\rho\sigma} \right) \right] \\
&= \exp \left[- \int (d^4x)_E \sqrt{g'} \left(- \frac{M_{\text{Pl}}^2}{2} (1 + \alpha_{2,R}^{\text{tree}}) \bar{R}' + \frac{1}{4} (1 + \alpha_{2,F}^{\text{tree}}) \bar{F}'_{\mu\nu} \bar{F}'^{\mu\nu} \right. \right. \\
&\quad \left. \left. - \beta_{2,1}^{\text{tree}} (\bar{F}'_{\mu\nu} \bar{F}'^{\mu\nu})^2 - \beta_{2,2}^{\text{tree}} (\bar{F}'_{\mu\nu} \widetilde{\bar{F}}'^{\mu\nu})^2 - \beta_{2,3}^{\text{tree}} \bar{F}'_{\mu\nu} \bar{F}'_{\rho\sigma} \bar{R}'^{\mu\nu\rho\sigma} \right) \right] \\
&= \exp \left[- \int (d^4x)_E \sqrt{\bar{g}} \left(- \frac{M_{\text{Pl}}^2}{2} \bar{R} + \frac{1}{4} \bar{F}_{\mu\nu} \bar{F}^{\mu\nu} \right. \right. \\
&\quad \left. \left. - \beta_{2,1}^{\text{tree}} \left(1 + \frac{2}{3} \alpha_{2,R}^{\text{tree}} - 2 \alpha_{2,F}^{\text{tree}} \right) (\bar{F}_{\mu\nu} \bar{F}^{\mu\nu})^2 - \beta_{2,2}^{\text{tree}} (1 + 2 \alpha_{2,R}^{\text{tree}} - 2 \alpha_{2,F}^{\text{tree}}) (\bar{F}_{\mu\nu} \widetilde{\bar{F}}^{\mu\nu})^2 \right. \right. \\
&\quad \left. \left. - \beta_{2,3}^{\text{tree}} \left(1 + \frac{1}{3} \alpha_{2,R}^{\text{tree}} - \alpha_{2,F}^{\text{tree}} \right) \bar{F}_{\mu\nu} \bar{F}_{\rho\sigma} \bar{R}^{\mu\nu\rho\sigma} \right) \right],
\end{aligned} \tag{4.157}$$

where $\alpha_{2,R}^{\text{tree}}$, $\alpha_{2,F}^{\text{tree}}$, $\beta_{2,1}^{\text{tree}}$, $\beta_{2,2}^{\text{tree}}$ and $\beta_{2,3}^{\text{tree}}$ ⁸ denote the second or higher order corrections for g , and $\beta_{2,1}^{\text{tree}}$, $\beta_{2,2}^{\text{tree}}$ and $\beta_{2,3}^{\text{tree}}$ do not include the first order correction for g because of Eq. (4.156). The corrections are assumed to be generated at the tree level. According to the procedure in Eq. (2.68), (2.69), and (2.70), the first order correction for g is eliminated in $\alpha_{2,R}^{\text{tree}}$ and $\alpha_{2,F}^{\text{tree}}$. Since the gravitational operators only involving the Riemann tensors can be removed by the redefinition of $g_{\mu\nu}$; see Appendix D, and the Riemann-squared operator effects on the extremal black hole entropy can be dropped in four dimensions, we omit such irrelevant terms. The

⁸From causality consideration, in Ref. [66, 67], it is argued that the tree level contribution to $\beta_{2,3}^{\text{tree}}$ requires stringy particles in UV theories. If we focus on the field theoretical descriptions, there is no contribution to $\beta_{2,3}^{\text{tree}}$ at the tree level.

background fields \overline{A}'_μ and $\overline{g}'_{\mu\nu}$ denote the classical solutions of the effective action. We choose the background field as follows:

$$\overline{A}'_\mu = \left(1 + \frac{1}{2} \left(\frac{4}{3} \alpha_{2,R}^{\text{tree}} - \alpha_{2,F}^{\text{tree}} \right)\right) \overline{A}_\mu, \quad (4.158)$$

$$\overline{g}'_{\mu\nu} = \left(1 - \frac{1}{3} \alpha_{2,R}^{\text{tree}}\right) \overline{g}_{\mu\nu}, \quad \overline{g}'^{\mu\nu} = \left(1 + \frac{1}{3} \alpha_{2,R}^{\text{tree}}\right) \overline{g}^{\mu\nu}. \quad (4.159)$$

From Eq. (4.157), the effective actions are obtained as follows:

$$\begin{aligned} W_g[\overline{g}_{\mu\nu}, \overline{A}] &\equiv -\ln Z_g[\overline{g}_{\mu\nu}, \overline{A}] \\ &= \int (d^4x)_E \sqrt{\overline{g}} \left(-\frac{M_{\text{Pl}}^2}{2} \overline{R} + \frac{1}{4} \overline{F}_{\mu\nu} \overline{F}^{\mu\nu} \right. \\ &\quad - \beta_{2,1}^{\text{tree}} \left(1 + \frac{2}{3} \alpha_{2,R}^{\text{tree}} - 2\alpha_{2,F}^{\text{tree}} \right) (\overline{F}_{\mu\nu} \overline{F}^{\mu\nu})^2 - \beta_{2,2}^{\text{tree}} (1 + 2\alpha_{2,R}^{\text{tree}} - 2\alpha_{2,F}^{\text{tree}}) (\overline{F}_{\mu\nu} \widetilde{\overline{F}}^{\mu\nu})^2 \\ &\quad \left. - \beta_{2,3}^{\text{tree}} \left(1 + \frac{1}{3} \alpha_{2,R}^{\text{tree}} - \alpha_{2,F}^{\text{tree}} \right) \overline{F}_{\mu\nu} \overline{F}_{\rho\sigma} \overline{R}^{\mu\nu\rho\sigma} \right), \end{aligned} \quad (4.160)$$

$$W_0[\overline{g}_{\mu\nu}, \overline{A}] = \lim_{g \rightarrow 0} W_g[\overline{g}_{\mu\nu}, \overline{A}] = \int (d^4x)_E \sqrt{\overline{g}} \left(-\frac{M_{\text{Pl}}^2}{2} \overline{R} + \frac{1}{4} \overline{F}_{\mu\nu} \overline{F}^{\mu\nu} \right). \quad (4.161)$$

Note here that solutions \overline{A}_μ and $\overline{g}_{\mu\nu}$ include the effects of the higher-derivative terms but the first order correction for the higher-derivative terms vanishes in W_0 by using the equation of motion. From Eq. (4.160) and (4.161), $W_g[\overline{g}_{\mu\nu}, \overline{A}] - W_0[\overline{g}_{\mu\nu}, \overline{A}]$ denotes the shift of the Euclidean effective action by the higher-derivative terms. Also, from Eq. (4.160), the first order correction for g is calculated as

$$\begin{aligned} \left(\frac{dW_g}{dg} \right)_{g=0} &= \left(\frac{\partial W_g}{\partial g} \right)_{g=0} + \int (d^4x)_E \sqrt{\overline{g}} \left(\left(\frac{\delta W_g}{\delta \overline{A}'^\mu} \right) \cdot \left(\frac{d\overline{A}'^\mu}{dg} \right)_{g=0} + \left(\frac{\delta W_g}{\delta \overline{g}'^{\mu\nu}} \right) \cdot \left(\frac{d\overline{g}'^{\mu\nu}}{dg} \right)_{g=0} \right) \\ &= \left(\frac{\partial W_g}{\partial g} \right)_{g=0} \\ &= \langle I_1 \rangle_{g=0} = 0, \end{aligned} \quad (4.162)$$

where $(d\overline{A}'^\mu/dg)_{g=0} = 0$ and $(d\overline{g}'^{\mu\nu}/dg)_{g=0} = 0$ are used from Eq. (4.158) and (4.159). From Eq. (2.49) and (4.162), we obtain the relative entropy,

$$\begin{aligned} S(P_0||P_g) &= W_0[\overline{g}_{\mu\nu}, \overline{A}] - W_g[\overline{g}_{\mu\nu}, \overline{A}] + g \langle I_1 \rangle_{g=0} \\ &= W_0[\overline{g}_{\mu\nu}, \overline{A}] - W_g[\overline{g}_{\mu\nu}, \overline{A}] \geq 0. \end{aligned} \quad (4.163)$$

Note here that we can put the same boundary terms to both W_0 and W_g because of its cancellation in the relative entropy. By taking $g = 1$, it is found that the relative entropy yields the negative shift of the effective action by the higher derivative terms generated at the tree level.

- Loop-level UV completion — Consider the EFT generated by the loop-level UV completion. The partition function of the theory $I_0 + g \cdot I_1$ is generally calculated as follows⁹:

$$\begin{aligned}
Z_g[\bar{g}_{\mu\nu}, \bar{A}] &= \int d[g]d[A]d[\Phi]e^{-I_g[g_{\mu\nu}; R_{\mu\nu\rho\sigma}, A, \Phi]} \\
&\equiv \int d[g]d[A]\exp\left[-\int (d^4x)_E \sqrt{g}\left(\Lambda_{0,\Phi}^{\text{loop}} - \frac{M_{\text{Pl}}^2}{2}(1 + \alpha_{1,R}^{\text{loop}} + \alpha_{2,R}^{\text{loop}})R\right.\right. \\
&\quad \left.+\frac{1}{4}(1 + \alpha_{1,F}^{\text{loop}} + \alpha_{2,F}^{\text{loop}})F_{\mu\nu}F^{\mu\nu}\right. \\
&\quad \left.-\beta_{2,1}^{\text{loop}}(F_{\mu\nu}F^{\mu\nu})^2 - \beta_{2,2}^{\text{loop}}(F_{\mu\nu}\tilde{F}^{\mu\nu})^2 - \beta_{2,3}^{\text{loop}}F_{\mu\nu}F_{\rho\sigma}R^{\mu\nu\rho\sigma}\right)\Big] \\
&= \exp\left[-\int (d^4x)_E \sqrt{\bar{g}}\left(\Lambda_{0,\Phi}^{\text{loop}} - \frac{M_{\text{Pl}}^2}{2}(1 + \alpha_{1,R}^{\text{loop}} + \alpha_{2,R}^{\text{loop}})\bar{R}' + \frac{1}{4}(1 + \alpha_{1,F}^{\text{loop}} + \alpha_{2,F}^{\text{loop}})\bar{F}'_{\mu\nu}\bar{F}'^{\mu\nu}\right.\right. \\
&\quad \left.-\beta_{2,1}^{\text{loop}}(\bar{F}'_{\mu\nu}\bar{F}'^{\mu\nu})^2 - \beta_{2,2}^{\text{loop}}(\bar{F}'_{\mu\nu}\tilde{\bar{F}}'^{\mu\nu})^2 - \beta_{2,3}^{\text{loop}}\bar{F}'_{\mu\nu}\bar{F}'_{\rho\sigma}\bar{R}'^{\mu\nu\rho\sigma}\right. \\
&\quad \left.+(\text{correction from } R \text{ and } F_{\mu\nu}F^{\mu\nu})\right)\Big] \\
&= \exp\left[-\int (d^4x)_E \sqrt{\bar{g}}\left(\Lambda_{0,\Phi}^{\text{loop}} - \frac{M_{\text{Pl}}^2}{2}(1 + \alpha_{1,R}^{\text{loop}})\bar{R} + \frac{1}{4}(1 + \alpha_{1,F}^{\text{loop}})\bar{F}_{\mu\nu}\bar{F}^{\mu\nu}\right.\right. \\
&\quad \left.-\beta_{2,1}^{\text{loop}}(\bar{F}_{\mu\nu}\bar{F}^{\mu\nu})^2 - \beta_{2,2}^{\text{loop}}(\bar{F}_{\mu\nu}\tilde{\bar{F}}^{\mu\nu})^2 - \beta_{2,3}^{\text{loop}}\bar{F}_{\mu\nu}\bar{F}_{\rho\sigma}\bar{R}^{\mu\nu\rho\sigma}\right. \\
&\quad \left.+(\text{correction from } R \text{ and } F_{\mu\nu}F^{\mu\nu})\right)\Big], \tag{4.164}
\end{aligned}$$

where $\alpha_{2,R}^{\text{loop}}, \alpha_{2,F}^{\text{loop}}, \beta_{2,1}^{\text{loop}}, \beta_{2,2}^{\text{loop}}$ and $\beta_{2,3}^{\text{loop}}$ are the second or higher order corrections for g , $\alpha_{1,R}^{\text{loop}}$ and $\alpha_{1,F}^{\text{loop}}$ are the first order corrections for g , and $\Lambda_{0,\Phi}^{\text{loop}}$ is the vacuum energy coming from Φ . The last term of Eq. (4.164) denotes corrections from light fields in $M_{\text{Pl}}^2 R/2$ and $F_{\mu\nu}F^{\mu\nu}/4$ at the one loop level. Since these corrections do not depend on g , they cancel in the relative entropy. We neglect two-loop effects in Eq. (4.164). The background fields \bar{A}'_μ and $\bar{g}'^{\mu\nu}$ denote the classical solution of the effective action. We choose the background field as follows:

$$\bar{A}'_\mu = \left(1 + \frac{1}{2}\left(\frac{4}{3}\alpha_{2,R}^{\text{loop}} - \alpha_{2,F}^{\text{loop}}\right)\right)\bar{A}_\mu, \tag{4.165}$$

$$\bar{g}'^{\mu\nu} = \left(1 - \frac{1}{3}\alpha_{2,R}^{\text{loop}}\right)\bar{g}^{\mu\nu}, \quad \bar{g}'^{\mu\nu} = \left(1 + \frac{1}{3}\alpha_{2,R}^{\text{loop}}\right)\bar{g}^{\mu\nu}. \tag{4.166}$$

The effective actions are obtained as follows:

$$\begin{aligned}
W_g[\bar{g}_{\mu\nu}, \bar{A}] &= \int (d^4x)_E \sqrt{\bar{g}}\left(\Lambda_{0,\Phi}^{\text{loop}} - \frac{M_{\text{Pl}}^2}{2}(1 + \alpha_{1,R}^{\text{loop}})\bar{R} + \frac{1}{4}(1 + \alpha_{1,F}^{\text{loop}})\bar{F}_{\mu\nu}\bar{F}^{\mu\nu}\right. \\
&\quad \left.-\beta_{2,1}^{\text{loop}}(\bar{F}_{\mu\nu}\bar{F}^{\mu\nu})^2 - \beta_{2,2}^{\text{loop}}(\bar{F}_{\mu\nu}\tilde{\bar{F}}^{\mu\nu})^2 - \beta_{2,3}^{\text{loop}}\bar{F}_{\mu\nu}\bar{F}_{\rho\sigma}\bar{R}^{\mu\nu\rho\sigma}\right)
\end{aligned}$$

⁹Similar to the SMEFT, strictly speaking, Eq. (4.164) holds in the limit of $g = 1$ or 0 . However, this subtle point does not affect the following discussions because we focus only on the reference ($g = 0$) and target ($g = 1$) theories.

$$+ (\text{correction from } R \text{ and } F_{\mu\nu}F^{\mu\nu}) \Big), \quad (4.167)$$

$$\begin{aligned} W_0[\bar{g}_{\mu\nu}, \bar{A}] &= \lim_{g \rightarrow 0} W_g[\bar{g}_{\mu\nu}, \bar{A}] \\ &= \int (d^4x)_E \sqrt{\bar{g}} \left(\Lambda_{0,\Phi}^{\text{loop}} - \frac{M_{\text{Pl}}^2}{2} \bar{R} + \frac{1}{4} \bar{F}_{\mu\nu} \bar{F}^{\mu\nu} + (\text{correction from } R \text{ and } F_{\mu\nu}F^{\mu\nu}) \right). \end{aligned} \quad (4.168)$$

Similar to the tree-level UV completion, the first order correction for the higher-derivative terms vanishes in W_0 by the equation of motion. Also, from Eq. (4.167), the first order correction for g is calculated as

$$\begin{aligned} \left(\frac{dW_g}{dg} \right)_{g=0} &= \left(\frac{\partial W_g}{\partial g} \right)_{g=0} + \int (d^4x)_E \sqrt{\bar{g}'} \left(\left(\frac{\delta W_g}{\delta \bar{A}'} \right) \cdot \left(\frac{d\bar{A}'}{dg} \right)_{g=0} + \left(\frac{\delta W_g}{\delta \bar{g}'_{\mu\nu}} \right) \cdot \left(\frac{d\bar{g}'_{\mu\nu}}{dg} \right)_{g=0} \right) \\ &= \left(\frac{\partial W_g}{\partial g} \right)_{g=0} \\ &= \langle I_1 \rangle_{g=0} \\ &= \int (d^4x)_E \sqrt{\bar{g}} \left(-\frac{M_{\text{Pl}}^2}{2} \frac{d\alpha_{1,R}^{\text{loop}}}{dg} \bar{R} + \frac{1}{4} \frac{d\alpha_{1,F}^{\text{loop}}}{dg} \bar{F}_{\mu\nu} \bar{F}^{\mu\nu} \right), \end{aligned} \quad (4.169)$$

where $(d\bar{A}'_{\mu}/dg)_{g=0} = 0$ and $(d\bar{g}'_{\mu\nu}/dg)_{g=0} = 0$ hold from Eq. (4.165) and (4.166). Note here that the last term of Eq. (4.167) does not depend on g . From Eq. (2.49), (4.167), (4.168) and (4.169), we obtain

$$\begin{aligned} S(P_0||P_g) &= W_0[\bar{g}_{\mu\nu}, \bar{A}] - W_g[\bar{g}_{\mu\nu}, \bar{A}] + g \langle I_1 \rangle_{g=0} \\ &= W_0[\bar{g}_{\mu\nu}, \bar{A}] - W_g^{\text{non-lin}}[\bar{g}_{\mu\nu}, \bar{A}] \geq 0, \end{aligned} \quad (4.170)$$

where we used $g \cdot (d\alpha_{1,R}^{\text{loop}}/dg) = \alpha_{1,R}^{\text{loop}}$ and $g \cdot (d\alpha_{1,F}^{\text{loop}}/dg) = \alpha_{1,F}^{\text{loop}}$ and defined the effective action without the first order corrections for g as follows:

$$\begin{aligned} W_g^{\text{non-lin}}[\bar{g}_{\mu\nu}, \bar{A}] &= \int (d^4x)_E \sqrt{\bar{g}} \left(\Lambda_{0,\Phi}^{\text{loop}} - \frac{M_{\text{Pl}}^2}{2} \bar{R} + \frac{1}{4} \bar{F}_{\mu\nu} \bar{F}^{\mu\nu} \right. \\ &\quad \left. - \beta_{2,1}^{\text{loop}} (\bar{F}_{\mu\nu} \bar{F}^{\mu\nu})^2 - \beta_{2,2}^{\text{loop}} (\bar{F}_{\mu\nu} \tilde{\bar{F}}^{\mu\nu})^2 - \beta_{2,3}^{\text{loop}} \bar{F}_{\mu\nu} \bar{F}_{\rho\sigma} \bar{R}^{\mu\nu\rho\sigma} \right. \\ &\quad \left. + (\text{correction from } R \text{ and } F_{\mu\nu}F^{\mu\nu}) \right). \end{aligned} \quad (4.171)$$

$W_0[\bar{g}_{\mu\nu}, \bar{A}] - W_g^{\text{non-lin}}[\bar{g}_{\mu\nu}, \bar{A}]$ denotes the corrections from the higher-derivative terms, and the inequality of (4.170) means that the Euclidean effective action decreases by the higher-derivative operators. It should be noted that the one-loop correction from R and $F_{\mu\nu}F^{\mu\nu}$ cancels in Eq. (4.170). By taking $g = 1$ in Eq. (4.170), we found that the relative entropy yields the negative shift of the effective action by the higher derivative terms generated at the one-loop level. Especially for loop-level UV completions involving massive charged particles with large charge-to-mass ratios, this result is consistent with Ref. [53].

It is found that, for both tree and loop-level UV completion, the non-negativity of the relative entropy yields the negative shift of the Euclidean effective action by the higher-derivative terms. This argument holds when the assumptions at the beginning of this section are valid. In the context of the WGC, the negative shift of the Euclidean effective action by the higher-derivative terms is demonstrated in a wide range of theories from unitarity and causality considerations [53]. Therefore, the results of the relative entropy considerations are consistent with that of the unitarity and causality, especially for loop-level UV completions involving massive charged particles with large charge-to-mass ratios and tree-level UV completions. As discussed in the next chapter, this result is closely related to the WGC.

4.4 Summary of bottom-up approach

In this section, we focused on a class of EFTs where the corrections to the leading terms can be removed by redefining light fields. For example, the single massless scalar field with the dimension-eight operator, the SMEFT dimension-eight gauge bosonic operators, and the Einstein-Maxwell theory with higher-derivative operators belong to such a class of EFTs. Our arguments in this section are based on assumptions summarized at the beginning of this section. In particular, Assumptions (III) and (IV) are the main assumptions in this section. Under the assumptions, we found that the relative entropy is the linear combination of the higher-derivative operators generated from the interactions I_I . Strictly speaking, for the tree and loop level UV completions, we derive the following relations,

$$\begin{aligned} S(P_{\text{NI}}||P_{\text{T}}) &= W_{\text{NI}}[\tilde{\phi}] - W_{\text{T}}[\tilde{\phi}] + \langle I_I \rangle_{\text{NI}} \\ &= (\text{linear combination of higher-derivative operators}) \geq 0. \end{aligned} \quad (4.172)$$

For each EFTs, this inequality yields the constraints on the Wilson coefficients of the higher-derivative operators, which are consistent with the positivity bounds from unitarity and causality.

5 Weak gravity conjecture and entropy constraint

We discuss a connection between the entropy constraints and the WGC. We focus on the NIRT and consider the perturbative corrections to the Euclidean effective action from the interactions between heavy and light degrees of freedom. In Sec. 5.1, we investigate a consequence of the non-negativity of the relative entropy to the corrections to thermodynamic entropy. In Sec. 5.2, we explain a relation between the shift of mass of the extremal black hole by the perturbative corrections of interaction and the non-negativity of the relative entropy. We will also comment on a connection between this work and Ref. [49].

5.1 Corrections to thermodynamic entropy

The perturbative corrections to the thermodynamic entropy have been actively studied in the context of the WGC [48]. We summarize the standard thermodynamic relations with the notation in Ref. [48] and investigate a connection between the non-negativity of the relative entropy and the corrections to thermodynamic entropy. The free energy of the thermodynamic system is defined as

$$\beta \cdot G \equiv \beta \cdot (M - \beta^{-1} \cdot S - Q \cdot \mu), \quad (5.1)$$

where G is the free energy, β is the inverse temperature, S is the thermodynamic entropy, Q is the charge such as $U(1)$ charge, and μ is the chemical potential. The first law of thermodynamics is expressed as follows:

$$dG = -S \cdot dT - Q \cdot d\mu. \quad (5.2)$$

From Eq. (5.1), we obtain

$$dG = dM - T \cdot dS - S \cdot dT - Q \cdot d\mu - \mu \cdot dQ. \quad (5.3)$$

Combining Eq. (5.2) and (5.3), we obtain

$$dM = T \cdot dS + \mu \cdot dQ \Rightarrow \beta = \left(\frac{\partial S}{\partial M} \right)_Q, \quad \mu = \left(\frac{\partial M}{\partial Q} \right)_S. \quad (5.4)$$

Now, let us assume the free energy of the thermodynamic system is shifted by perturbative effects, e.g., corrections from heavy degrees of freedom, as follows:

$$G(T, \mu, 0) \rightarrow G(T, \mu, \epsilon) \equiv G(T, \mu, 0) + \epsilon \cdot \Delta G(T, \mu), \quad (5.5)$$

where ϵ is an auxiliary parameter to characterize the perturbative corrections. Note here that ϵ is not the same as g in Sec. 2 because ΔG involve the first or higher order corrections of g . Then, the thermodynamic entropy shift by the perturbative effects is defined as follows:

$$(\Delta S)_{\beta, \mu} \equiv S(T, \mu, \epsilon) - S(T, \mu, 0) = \epsilon \cdot \left(\frac{\partial S}{\partial \epsilon} \right)_{\beta, \mu} + \mathcal{O}(\epsilon^2). \quad (5.6)$$

The leading correction of ϵ to the thermodynamic entropy is expressed as

$$\left(\frac{\partial S}{\partial \epsilon} \right)_{\beta, \mu} = \left(\frac{\partial S}{\partial M} \right)_{Q, \epsilon} \cdot \left(\frac{\partial M}{\partial \epsilon} \right)_{\beta, \mu} + \left(\frac{\partial S}{\partial Q} \right)_{M, \epsilon} \cdot \left(\frac{\partial Q}{\partial \epsilon} \right)_{\beta, \mu} + \left(\frac{\partial S}{\partial \epsilon} \right)_{M, Q}. \quad (5.7)$$

Also, using the triple product rule, we obtain

$$\left(\frac{\partial S}{\partial Q} \right)_{M, \epsilon} = - \left(\frac{\partial S}{\partial M} \right)_{Q, \epsilon} \cdot \left(\frac{\partial M}{\partial Q} \right)_{S, \epsilon}. \quad (5.8)$$

Combining Eqs. (5.4), (5.7), and (5.8), we obtain

$$\left(\frac{\partial S}{\partial \epsilon} \right)_{\beta, \mu} = \beta \cdot \left(\frac{\partial M}{\partial \epsilon} \right)_{\beta, \mu} - \beta \cdot \mu \cdot \left(\frac{\partial Q}{\partial \epsilon} \right)_{\beta, \mu} + \left(\frac{\partial S}{\partial \epsilon} \right)_{M, Q}. \quad (5.9)$$

On the other hand, Eq. (5.1) yields

$$\left(\frac{\partial(\beta \cdot G)}{\partial \epsilon} \right)_{\beta, \mu} = \beta \cdot \left(\frac{\partial M}{\partial \epsilon} \right)_{\beta, \mu} - \left(\frac{\partial S}{\partial \epsilon} \right)_{\beta, \mu} - \beta \cdot \mu \cdot \left(\frac{\partial Q}{\partial \epsilon} \right)_{\beta, \mu}. \quad (5.10)$$

From Eqs. (5.9) and (5.10), we obtain

$$\left(\frac{\partial S}{\partial \epsilon}\right)_{M,Q} = - \left(\frac{\partial(\beta \cdot G)}{\partial \epsilon}\right)_{\beta,\mu}. \quad (5.11)$$

Consequently, the leading perturbative correction of ϵ to thermodynamic entropy at fixed energy and charge is expressed as follows:

$$\epsilon \cdot \left(\frac{\partial S}{\partial \epsilon}\right)_{M,Q} = -\epsilon \cdot \left(\frac{\partial(\beta \cdot G)}{\partial \epsilon}\right)_{\beta,\mu} = -\beta \cdot (\epsilon \cdot \Delta G(T, \mu)), \quad (5.12)$$

where the most right-hand side of Eq. (5.12) denotes the leading perturbative correction of ϵ to the Euclidean effective action, which is relevant to the non-negativity of the relative entropy. The point is that Eq. (5.12) takes a positive value for the negative free energy shift.

In Eq. (2.52) of Sec. 2.3, we have provided the lower and upper bounds on the perturbative corrections to the Euclidean effective action from the interaction between the heavy and light degrees of freedom. Note here that the derivation of Eq. (2.52) does not depend on whether the temperature of the system is zero or not. The perturbative corrections to the free energy of the thermodynamic system are expressed as follows:

$$\beta \cdot (\epsilon \cdot \Delta G) \equiv W_T[\beta, \phi] - W_{\text{NI}}[\beta, \phi], \quad (5.13)$$

where the left-hand side denotes the most right-hand side of Eq. (5.12). Combing Eqs. (2.52) and (5.13), we obtain the following inequality.

$$\langle I_I \rangle_{\text{NI}} \geq \beta \cdot (\epsilon \cdot \Delta G) \geq \langle I_I \rangle_{\text{T}}. \quad (5.14)$$

From Eqs. (5.12) and (5.14), we obtain

$$-\langle I_I \rangle_{\text{T}} \geq \epsilon \cdot \left(\frac{\partial S}{\partial \epsilon}\right)_{M,Q} \geq -\langle I_I \rangle_{\text{NI}}. \quad (5.15)$$

Consequently, the relative entropy yields the upper and lower bounds on the thermodynamic entropy shift at fixed energy and charge by the interaction between heavy and light degrees of freedom.

For the class of EFTs in Sec. 4, under the assumptions summarized at the beginning of Sec. 4, the relative entropy denotes the shift of the Euclidean effective action, i.e., the free energy at zero temperature, consisting of the linear combination of higher-derivative operators as follows:

$$S(P_{\text{NI}}||P_{\text{T}}) = -\beta \cdot (\epsilon \cdot \Delta G) = (\text{linear combination of higher-derivative operators}) \geq 0. \quad (5.16)$$

Combining Eq. (5.12) and (5.16), we obtain

$$\epsilon \cdot \left(\frac{\partial S}{\partial \epsilon}\right)_{M,Q} = S(P_{\text{NI}}||P_{\text{T}}) \geq 0, \quad (5.17)$$

which is evaluated at zero temperature. Consequently, for a class of EFTs in Sec. 4, it is found that the non-negativity of relative entropy yields the positive shift of the thermodynamic entropy of zero temperature system at fixed energy and charge by the higher-derivative operators. It should be noted that, in contrast to the inequalities of (5.15), this argument is derived from the assumptions summarized at the beginning of Sec. 4. Therefore, Eq. (5.15) is applicable in more theories than Eq. (5.17).

5.2 Entropy constraints and weak-gravity-conjecture

In Ref. [48], by using general thermodynamic considerations, it has demonstrated that the perturbations decrease the minimal energy of thermodynamic systems at a fixed charge when the perturbative correction to the thermodynamic entropy of the left-hand side of Eq. (5.12) is positive. For the sake of self-contained, we briefly review Ref. [48] and then explain the connection between the WGC-like behavior in a shift of the extremal black hole mass and the relative entropy.

From Eq. (5.6), and the third law of thermodynamics, i.e., $\lim_{T \rightarrow 0} S(T, Q, \epsilon) = 0$, we obtain

$$\lim_{T \rightarrow 0} T \left(\frac{\partial S(T, Q, \epsilon)}{\partial \epsilon} \right)_{\beta, Q} = 0. \quad (5.18)$$

Equations (5.1) and (5.2) yield the following thermodynamic relations:

$$M(T, \mu, \epsilon) = G(T, \mu, \epsilon) + T \cdot S + \mu \cdot Q, \quad (5.19)$$

$$S(T, \mu) = - \left(\frac{\partial G}{\partial T} \right)_{\mu, \epsilon}, \quad (5.20)$$

$$Q(T, \mu, \epsilon) = - \left(\frac{\partial G}{\partial \mu} \right)_{\beta, \epsilon}. \quad (5.21)$$

By using the above relations, the energy shift at fixed temperature and charge is calculated as follows:

$$\begin{aligned} \left(\frac{\partial M}{\partial \epsilon} \right)_{\beta, Q} &= \left(\frac{\partial}{\partial \epsilon} (G + T \cdot S + \mu \cdot Q) \right)_{\beta, Q} \\ &= \left(\frac{\partial G}{\partial \mu} \right)_{\beta, \epsilon} \cdot \left(\frac{\partial \mu}{\partial \epsilon} \right)_{\beta, Q} + \left(\frac{\partial G}{\partial \epsilon} \right)_{\beta, \mu} + T \cdot \left(\frac{\partial S}{\partial \epsilon} \right)_{\beta, Q} + Q \cdot \left(\frac{\partial \mu}{\partial \epsilon} \right)_{\beta, Q} \\ &= \left(\frac{\partial G}{\partial \epsilon} \right)_{\beta, \mu} + T \cdot \left(\frac{\partial S}{\partial \epsilon} \right)_{\beta, Q}. \end{aligned} \quad (5.22)$$

Combining Eqs. (5.18) and (5.22), we obtain

$$\lim_{T \rightarrow 0} \left(\frac{\partial M}{\partial \epsilon} \right)_{\beta, Q} = \lim_{T \rightarrow 0} \left(\frac{\partial G}{\partial \epsilon} \right)_{\beta, \mu}. \quad (5.23)$$

From Eqs. (5.12) and (5.23), we obtain

$$\lim_{T \rightarrow 0} \left(\frac{\partial M}{\partial \epsilon} \right)_{\beta, Q} = \lim_{T \rightarrow 0} -T \cdot \left(\frac{\partial S}{\partial \epsilon} \right)_{M, Q}. \quad (5.24)$$

Since the minimum energy of the system is defined as

$$M_{\text{ext}}(Q, \epsilon) \equiv \lim_{T \rightarrow 0} M(T, Q, \epsilon), \quad (5.25)$$

Eq. (5.24) is expressed as follows:

$$\lim_{T \rightarrow 0} \left(\frac{\partial M}{\partial \epsilon} \right)_{\beta, Q} = \lim_{M \rightarrow M_{\text{ext}}(Q, \epsilon)} -T(M, Q, \epsilon) \cdot \left(\frac{\partial S}{\partial \epsilon} \right)_{M, Q}. \quad (5.26)$$

Therefore, we see that the perturbations decrease the minimal energy of thermodynamic systems at a fixed charge when the perturbative correction to the thermodynamic entropy is positive [48]. Particularly for the charged-BH described by the Einstein-Maxwell theory, Q is the $U(1)$ charge, and the minimum mass of the BH is $M_{\text{ext}} = \sqrt{2}M_{\text{Pl}}Q$. As studied in a large amount of literature [36, 49, 53, 63, 68, 69], a state with a charge-to-mass ratio larger than unity is motivated by the thought experiment of decay of the extremal black hole. When the minimum mass of BH at fixed charge decrease by the perturbative correction from the higher dimensional operators, asymptotically large extremal black holes described by the Einstein-Maxwell theory with higher derivative operators are allowed to decay to a state whose charge exceeds its mass, e.g., extremal black hole. In this work, we refer to the behavior that the minimum energy of a thermodynamic system at fixed charge decreases by the perturbative correction as *WGC-like behavior*. Note here that, in this work, we use this terminology for general thermodynamic systems with and without gravity.

For the EFTs in Sec. 4, the non-negativity of the relative entropy yields the negative free energy shift at zero temperature within the validity of the field theoretical descriptions. From Eq. (5.12), this result means that the thermodynamic entropy shift at fixed energy and charge by the higher-derivative operators is positive in the extremal limit of $M = M_{\text{ext}}$, i.e., the zero temperature limit. Then, the right-hand side of Eq. (5.26) is negative, and the WGC-like behavior arises. For the Einstein-Maxwell theory, as studied in Ref. [53], the unitarity and causality considerations also yield the positive BH entropy shift at fixed energy and charge by the higher-derivative operators generated in a wide range of theories in the extremal limit. Therefore, the entropy constraints yield consistent results with the unitarity and causality considerations [53], especially for loop-level UV completions involving massive charged particles with large charge-to-mass ratios $z = \sqrt{2}M_{\text{Pl}}|q|/m$ and tree-level UV completions. The entropy constraint implies that the extremal black hole would behave as a state with $z \geq 1$ even if massive charged particles with $z \geq 1$ do not exist. Our argument is applicable when the thermodynamic relations and the assumptions at the beginning of Sec. 4 are held. The amplitude considerations in Ref. [53] are also applicable beyond the field theoretical descriptions under the assumption that the higher spin states Reggeizing graviton exchange are subdominant. However, our discussions in Sec. 4 rely on the field theory, so it is not clear if our results are applicable beyond the field theoretical descriptions. Detailed studies of the applicability of this study are expected in the future.

Here, we comment on a relation between Ref. [49] and this work. In Ref. [49], it has been demonstrated that the Euclidean effective action decreases by higher-derivative operators generated at the tree level. For convenience, we briefly review it by using the notation of this work. Using the saddle point approximation, we obtain

$$I_0[\tilde{\phi}_0, 0] = I_g[\tilde{\phi}_0, 0] \geq I_g[\tilde{\phi}_g, \tilde{\Phi}_g], \quad (5.27)$$

where $\tilde{\phi}_0$ is the classical solution of I_0 , $\tilde{\phi}_g$ and $\tilde{\Phi}_g$ are that of I_g , and $I_g[\tilde{\phi}_0, 0] = I_0[\tilde{\phi}_0, 0]$ holds because the interaction I_I vanishes for $\Phi = 0$. The inequality of (5.27) arises because $\tilde{\Phi}_g$ denotes the local minimum of I_g and would take a small value by heavy field mass suppression. By taking $g = 1$ in Eq. (5.27), we obtain

$$W_{\text{NI}}[\tilde{\phi}_{\text{NI}}] = I_{\text{NI}}[\tilde{\phi}_{\text{NI}}, 0] \geq I_{\text{T}}[\tilde{\phi}_{\text{T}}, \tilde{\Phi}_{\text{T}}] = W_{\text{T}}[\tilde{\phi}_{\text{T}}], \quad (5.28)$$

where $I_{\text{NI}} = I_0$, $I_{\text{T}} = I_{g=1}$, $\tilde{\phi}_{\text{NI}} = \tilde{\phi}_0$, $\tilde{\phi}_{\text{T}} = \tilde{\phi}_{g=1}$, and $\tilde{\Phi}_{\text{T}} = \tilde{\Phi}_{g=1}$ are used, and $\tilde{\Phi}_{\text{T}}$ is expressed as a linear combination of $\tilde{\phi}_{\text{T}}$. The effective action W_{NI} does not generate the higher-dimensional operators, but the W_{T} yields them through the interacting term between ϕ and Φ . Therefore, the inequality (5.27) means that the Euclidean effective action decreases by higher-dimensional operators generated at the tree level. In other words, at fixed temperature β , the free energy decreases by higher-dimensional operators generated at the tree level. It should be noted that in Eq. (5.27), we used the following relation.

$$\lim_{\tilde{\phi}_g \rightarrow \tilde{\phi}_0, \tilde{\Phi}_g \rightarrow 0} I_g[\tilde{\phi}_g, \tilde{\Phi}_g] = I_0[\tilde{\phi}_0, 0] = \lim_{g \rightarrow 0} I_g[\tilde{\phi}_g, \tilde{\Phi}_g], \quad (5.29)$$

where procedures of the leftmost and rightmost sides are respectively performed by Ref. [49], and this work and Ref. [54]. The point is that Ref. [49] also derives the WGC-like behavior by comparing the NIRT and target theories. It should be emphasized that this work is essentially the same as Ref. [49] at the tree level.

6 Implication of entropy constraints

We found that the non-negativity of relative entropy yields constraints on the class of EFTs in Sec. 4, which are consistent with the conventional positivity bounds [51, 53, 65]. In this section, we investigate relations of the non-negativity of the relative entropy with the unitary time evolution of the system, causality, and the second law of thermodynamics.

6.1 Unitary time-evolution

The relative entropy, i.e., $S(\rho_{\text{T}}||\rho_{\text{R}}) = \text{Tr}[\rho_{\text{T}} \ln \rho_{\text{T}} - \rho_{\text{T}} \ln \rho_{\text{R}}]$, is a non-negative quantity, which is a consequence of the Hermiticity of the probability distribution functions ρ_{T} and ρ_{R} . In the quantum mechanical approach of the entropy constraints in Sec. 2.1, to derive the bounds on EFTs, we have focused on the probability distribution functions described as follows:

$$\rho_{\text{T}} \equiv \frac{e^{-\beta H_{\text{T}}}}{Z_{\text{T}}}, \quad \rho_{\text{R}} \equiv \frac{e^{-\beta H_{\text{R}}}}{Z_{\text{R}}}, \quad (6.1)$$

with the partition functions $Z_{\text{T}} \equiv \text{Tr}[e^{-\beta H_{\text{T}}}]$ and $Z_{\text{R}} \equiv \text{Tr}[e^{-\beta H_{\text{R}}}]$. In Sec. 3 and 4, we have demonstrated that the positivity bounds on some EFTs are derived by the above type of probability distribution functions. Since the non-negativity of the relative entropy is based on the Hermiticity of ρ_{T} and ρ_{R} , the Hermiticity of the Hamiltonians H_{T} and H_{R} is also assumed to derive the entropy

constraints on EFTs. Therefore, the time evolution of the target theory defined by H_T is unitary, and the unitary S matrix is ensured because of the Hermiticity of H_T . The point is that the entropy constraint is based on the unitary time evolution of the target theory, i.e., the optical theorem in the target theory. The conventional positivity bounds from amplitude considerations [51, 65] rely on the optical theorem.

6.2 Causality

We discuss the connection between the non-negativity of the relative entropy and the causality [51] by using a simple example. Consider the EFT defined by the Euclidean Lagrangian as,

$$\mathcal{L}^{(E)}[\phi] = \frac{1}{2}(\partial_I \phi \partial_I \phi) - \frac{c}{M^4}(\partial_I \phi \partial_I \phi)^2. \quad (6.2)$$

By setting $\phi = \tilde{\phi} + \varphi$, with φ the small dynamical perturbation, expand the Lagrangian by the perturbation as follows:

$$\begin{aligned} \mathcal{L}^{(E)} = & \frac{1}{2} \left[\delta_{IJ} - \frac{c}{6M^4} \left(4(\partial_I \tilde{\phi})^2 \delta_{IJ} + 8(\partial_I \tilde{\phi})(\partial_J \tilde{\phi}) \right) \right] (\partial_I \tilde{\phi} \partial_J \tilde{\phi}) \\ & + \left[1 - \frac{4c}{M^4} (\partial_J \tilde{\phi})^2 \right] \partial_I \tilde{\phi} \partial_I \varphi \\ & + \frac{1}{2} \left[\delta_{IJ} - \frac{c}{M^4} \left(4(\partial_I \tilde{\phi})^2 \delta_{IJ} + 8(\partial_I \tilde{\phi})(\partial_J \tilde{\phi}) \right) \right] (\partial_I \varphi \partial_J \varphi) + \mathcal{O}(\varphi^3). \end{aligned} \quad (6.3)$$

To ensure the validity of the Euclidean path integral around $\tilde{\phi}$, the last line of Eq. (6.3) yields the following relation.

$$k^2 - \frac{c}{M^4} (4q^2 k^2 + 8(q \cdot k)^2) \geq 0 \Rightarrow v_E^2 \leq 1 - \frac{8c(q \cdot k)^2}{|\vec{k}|^2(M^4 - 4cq^2)}, \quad (6.4)$$

where φ is expanded in plane waves as $\varphi \propto e^{ik \cdot x}$, $v_E^2 \equiv -k_4^2/|\vec{k}|^2$ denotes the speed of propagation of φ in the Euclidean space, and $\partial_I \tilde{\phi} = q_I$ is a constant vector. As discussed in Sec. 4.1, the non-negativity of relative entropy yields $c \geq 0$. Therefore, combining the above validity of the Euclidean path integral, and the non-negativity of relative entropy, we obtain the upper bound on v_E as follows,

$$v_E^2 \leq 1, \quad (6.5)$$

which ensures causality in the Minkowski space. Consequently, the non-negativity of relative entropy yields causality in the EFT. It should be noted that the entropy constraints are based on the Euclidean path integral method, and the validity of the Euclidean path integral around the classical solution is assumed. As explained in Sec. 3.10, the non-negativity of relative entropy may be broken when the Euclidean path integral is performed around a point not being a local minimum.

6.3 Second law of thermodynamics

The non-negativity of relative entropy is closely related to the second law of thermodynamics [58, 59]. We demonstrate that a simple derivation of the second law of thermodynamics is contained in the

entropy constraints of this study. Adopt the thermal reference theory of Sec. 2.2 as the reference theory, and consider the system consisting of a thermodynamic system S and a heat bath system B. We suppose that the Hamiltonian of the whole system is defined as

$$H_T = H_S + H_B + H_{SB}, \quad (6.6)$$

where H_S is the Hamiltonian of S, and H_B is that of B. The interacting term H_{SB} denotes the interaction between S and B, which is generally a time-dependent operator. Note here that the system S can involve both light and heavy degrees of freedom.

Now, let us assume that the initial quantum state of the whole system is defined as

$$\rho_{\text{ini}} \equiv \rho_{\text{ini},S} \otimes e^{-\beta H_B} / Z_B(\beta), \quad (6.7)$$

where $\rho_{\text{ini},S}$ is the initial state of S, and $e^{-\beta H_B} / Z_B(\beta)$ is that of B at an inverse temperature β . By tracing over the heat bath degrees of freedom, the partition function of B is defined as

$$Z_B(\beta) \equiv \text{Tr}_B[e^{-\beta H_B}]. \quad (6.8)$$

After a time-evolution of the whole system described by a unitary operator U , the final state of the whole system is expressed as

$$\rho_{\text{fin}} = U \rho_{\text{ini}} U^\dagger. \quad (6.9)$$

By tracing out the heat bath degrees of freedom, the final state of S is expressed as

$$\rho_{\text{fin},S} \equiv \text{Tr}_B[\rho_{\text{fin}}]. \quad (6.10)$$

Note here that the time evolution of S is not unitary because the heat bath system is traced out.

Following Sec. 2.2, define the probability distribution functions of the target theory and reference theory as follows.

$$\rho_T \equiv \rho_{\text{ini}}, \quad (6.11)$$

$$\rho_R \equiv U^\dagger \rho_{\text{fin},S} \otimes e^{-\beta H_B} / Z_B(\beta) U. \quad (6.12)$$

The relative entropy between ρ_T and ρ_R is calculated as

$$S(\rho_T || \rho_R) = S(\rho_{\text{fin},S}) - S(\rho_{\text{ini},S}) - \beta \cdot Q \geq 0, \quad (6.13)$$

where $S(\rho) \equiv -\text{Tr}_S[\rho \ln \rho]$ denotes the von Neumann entropy of S, and $Q \equiv \text{Tr}[H_B \rho_{\text{ini}}] - \text{Tr}[H_B \rho_{\text{fin}}]$ is the heat exchange between S and B. The details of the derivations of Eq. (6.13) are summarized in Appendix C. Equation (6.13) is the Clausius inequality and denotes the second law of thermodynamics. Consequently, we obtain the second law of thermodynamics in the above simple setup from the non-negativity of relative entropy.

Lastly, for convenience, we would like to discuss the connection between the target system of this section and that of Sec. 2.3. Although the heat bath system is not included in Sec. 2.3, we can

freely add it to the systems. Let us take the case of NIRT as an example and consider the probability distribution functions of the target and reference theories ρ'_T and ρ'_{NI} , where the heat bath system is not included. We assume ρ'_T and ρ'_{NI} denote the probability distribution functions of the target and reference theories in Sec. 2.3. By including the heat bath system $e^{-\beta H_B}/Z_B(\beta)$, one can define as follows.

$$\rho_T \equiv \rho'_T \otimes e^{-\beta H_B}/Z_B(\beta), \quad (6.14)$$

$$\rho_{NI} \equiv \rho'_{NI} \otimes e^{-\beta H_B}/Z_B(\beta), \quad (6.15)$$

where ρ_T is the same as the probability distribution of Eq. (6.11), and ρ'_T and ρ'_{NI} represent the probability distribution in Sec. 2.3. Then, the relative entropy between ρ_T and ρ_{NI} is expressed as follows,

$$S(\rho_T||\rho_{NI}) = \text{Tr}[\rho_T \ln \rho_T - \rho_T \ln \rho_{NI}] = S(\rho'_T||\rho'_{NI}), \quad (6.16)$$

$$S(\rho_{NI}||\rho_T) = \text{Tr}[\rho_{NI} \ln \rho_{NI} - \rho_{NI} \ln \rho_T] = S(\rho'_{NI}||\rho'_T). \quad (6.17)$$

The right-hand sides of Eq. (6.16) and (6.17) are the same quantity that was discussed in Sec. 2.3. The point is that the above derivation of the second law of thermodynamics is included as a special case of the entropy constraints of this work. Here, it should be noted that the corrections to thermodynamic entropy, i.e., the left-hand side of Eq. (5.12) is a different quantity from the entropy shift due to the time-evolution discussed in this section. It is remarkable that the non-negativity of both entropy shifts is derived from the relative entropy depending on the choice of the reference theory.

7 Summary

We studied constraints on the EFTs by evaluating the relative entropy between the target and reference theories. In addition to providing the details of Ref. [54], we updated the results in Ref. [54] by considering more theories and new reference theories. Firstly, in Sec. 2, we reviewed the details of the main idea of the entropy constraint of Ref. [54] and provided the procedures to calculate the relative entropy by introducing some new reference theories, i.e., massive free reference theory, and infinite mass reference theory. Following the procedures of Sec. 2, in Secs. 3-6, we evaluated the relative entropy and investigated the physical consequence of the non-negativity of the relative entropy in various EFTs.

In Sec. 3, we adopted the top-down approach and evaluated the relative entropy in various theories involving field theories, quantum mechanical models, and Gaussian distribution functions. The various examples satisfy the non-negativity of the relative entropy, which yields non-trivial constraints on the EFTs when the heavy degrees of freedom are integrated out. In Sec. 3.10, we also discussed some examples where the non-negativity of the relative entropy is violated. In contrast to Sec. 3, in Sec. 4, we adopted the bottom-up approach, i.e., the EFTs are provided while the UV theories are unknown, and investigated the consequence of the non-negativity of the relative entropy. We focused on a class of EFTs where the perturbative corrections to the non-higher derivative operators can be removed

by redefinitions of the light fields, e.g., the single massless scalar field with the dimension-eight operator, SMEFT dimension-eight $SU(N)$ gauge bosonic operators and Einstein-Maxwell theory with higher-derivative operators. Under the four assumptions, i.e., Hermiticity of probability distribution functions, the validity of Euclidean path integral method, higher-derivative operators generated from the interaction between heavy and light fields, and leading order of the interaction between heavy and light fields; see the beginning of Sec. 4, we found that the non-negativity of the relative entropy yields the constraints on such EFTs, which are consistent with the positivity bounds from the unitarity and causality. These constraints are derived for each of the tree-level and loop-level UV completions; see Sec. 4.

In Secs. 5 and 6, we discussed connections between the non-negativity of the relative entropy and various inequalities in physics, i.e., the WGC-like behavior, unitary time evolution, causality, and second law of thermodynamics. In particular, the entropy constraints on the Einstein-Maxwell theory with higher-derivative operators imply that the minimum mass of the black hole at fixed charge decrease by the higher-derivative operators generated from the interactions between heavy and light fields. This argument is applicable when the thermodynamic relations and the assumptions at the beginning of Sec. 4 are held. It should be noted that the assumptions include that corrections from the interaction involving higher-derivative operators of light fields are not dominant in the EFTs. These results about Einstein-Maxwell theory rely on the field theoretical descriptions in Sec. 4, so it is not clear if our results are applicable to UV theories involving stringy particles. Detailed studies of the applicability of this study are expected in the future. Also, our relative entropy consideration yields the second law of thermodynamics in a simple setup depending on the reference theory; see Sec. 6.3.

Our entropy consideration is applied to various theories, but further applications to more theories are required to understand its validity range. In particular, a derivation of the WGC-like behavior is subject to the assumptions at the beginning in Sec. 4, and further studies based on various UV theories are essential to confirm the consistency of our results under the assumptions. We also note that our entropy constraint is a different approach from the conventional unitarity and causality considerations, and therefore applicable theories would be different from each other.

In summary, we conclude that the relative entropy consideration provides a unified understanding of various inequalities in physics in addition to yielding a new approach to constraints on EFTs, and further studies are expected in various theories.

Acknowledgement

The work is supported in part by the National Science Foundation of China under Grant Nos. 11675002, 11635001, 11725520, and 12235001.

A Wick rotation

We provide how to obtain the Euclidean effective action from the Minkowski one. We denote the Euclidean indices by I, J, \dots and the Minkowski indices by μ, ν, \dots . The Euclidean effective action

$I_E[\phi_E, g_{E,IJ}]$ is obtained by the Wick rotation as follows:

$$I_E[\phi_E, g_{E,IJ}] \equiv \frac{1}{i} I[\phi, g_{\mu\nu}]_{t=-i\tau}, \quad (\text{A.1})$$

where $I[\phi, g_{\mu\nu}]$ is the effective action in the Minkowski space defined as

$$I[\phi, g_{\mu\nu}] = \int \sqrt{-g} d^4x \mathcal{L}[\phi, g_{\mu\nu}]. \quad (\text{A.2})$$

Here, \mathcal{L} is the Lagrangian, ϕ is the background light fields in the Minkowski space, $g_{\mu\nu}$ is a metric tensor in the Minkowski signature, and $g = \det g^{\mu\nu}$ is a determinant of the metric tensor.

Let us consider a transformation of coordinates: $x^\mu = (x^0 = t, \vec{x}) \rightarrow x'^\mu = (x'^0, \vec{x}') = (\lambda t, \vec{x})$, where λ is a constant. For convenience, we summarize the transformed scalar, vector, and metric tensor fields as follows:

$$\phi'(t', \vec{x}) = \phi(t, \vec{x}), \quad (\text{A.3})$$

$$A'^\mu(t', \vec{x}) = \frac{\partial x'^\mu}{\partial x^\alpha} A^\alpha(t, \vec{x}) = (\lambda \cdot A^0(x), A^i(x)), \quad (\text{A.4})$$

$$A'_\mu(t', \vec{x}) = \frac{\partial x^\alpha}{\partial x'^\mu} A_\alpha(t, \vec{x}) = (\lambda^{-1} \cdot A_0(x), A_i(x)), \quad (\text{A.5})$$

$$g'^{\mu\nu}(t', \vec{x}) = \frac{\partial x'^\mu}{\partial x^\alpha} \frac{\partial x'^\nu}{\partial x^\beta} g^{\alpha\beta}(t, \vec{x}) = \begin{pmatrix} \lambda^2 g^{00}(x) & \lambda g^{01}(x) & \lambda g^{02}(x) & \lambda g^{03}(x) \\ \lambda g^{10}(x) & g^{11}(x) & g^{12}(x) & g^{13}(x) \\ \lambda g^{20}(x) & g^{21}(x) & g^{22}(x) & g^{23}(x) \\ \lambda g^{30}(x) & g^{31}(x) & g^{32}(x) & g^{33}(x) \end{pmatrix}, \quad (\text{A.6})$$

$$g'_{\mu\nu}(t', \vec{x}) = \frac{\partial x^\alpha}{\partial x'^\mu} \frac{\partial x^\beta}{\partial x'^\nu} g_{\alpha\beta}(t, \vec{x}) = \begin{pmatrix} \lambda^{-2} g_{00}(x) & \lambda^{-1} g_{01}(x) & \lambda^{-1} g_{02}(x) & \lambda^{-1} g_{03}(x) \\ \lambda^{-1} g_{10}(x) & g_{11}(x) & g_{12}(x) & g_{13}(x) \\ \lambda^{-1} g_{20}(x) & g_{21}(x) & g_{22}(x) & g_{23}(x) \\ \lambda^{-1} g_{30}(x) & g_{31}(x) & g_{32}(x) & g_{33}(x) \end{pmatrix}. \quad (\text{A.7})$$

Also, the completely antisymmetric tensor is transformed as follows:

$$\frac{1}{\sqrt{-g'}} \epsilon'^{\mu\nu\rho\sigma} = J \cdot \sqrt{\lambda^2} \frac{\partial x'^\mu}{\partial x^\alpha} \frac{\partial x'^\nu}{\partial x^\beta} \frac{\partial x'^\rho}{\partial x^\gamma} \frac{\partial x'^\sigma}{\partial x^\delta} \frac{1}{\sqrt{-g}} \epsilon^{\alpha\beta\gamma\delta}, \quad (\text{A.8})$$

where $\epsilon_{\mu\nu\rho\sigma} = \epsilon^{\mu\nu\rho\sigma}$, $\epsilon_{0123} = +1$, a determinant of the metric tensor $g = \det g_{\mu\nu}$ is calculated as $g = \lambda^2 g'$, and $J = \lambda^{-1}$ is defined as

$$J \equiv \epsilon_{\mu\nu\rho\sigma} \frac{\partial x^\mu}{\partial x'^0} \frac{\partial x^\nu}{\partial x'^1} \frac{\partial x^\rho}{\partial x'^2} \frac{\partial x^\sigma}{\partial x'^3}. \quad (\text{A.9})$$

Also $\epsilon'^{\mu\nu\rho\sigma}$ satisfies the following relation.

$$J = \epsilon'^{\mu\nu\rho\sigma} \frac{\partial x^0}{\partial x'^\mu} \frac{\partial x^1}{\partial x'^\nu} \frac{\partial x^2}{\partial x'^\rho} \frac{\partial x^3}{\partial x'^\sigma}. \quad (\text{A.10})$$

By the above transformation, the covariant volume element is not changed as follows:

$$\sqrt{-g} d^4x = \sqrt{-\lambda^2 g'} \cdot \lambda^{-1} d^4x' = \sqrt{-g'} d^4x'. \quad (\text{A.11})$$

The action in the coordinate x' is obtained as

$$I[\phi, g_{\mu\nu}] = \int \sqrt{-g'} d^4 x' \mathcal{L}[\phi', g'_{\mu\nu}] = \lambda^{-1} \int \sqrt{-\lambda^2 g'} d^4 x' \mathcal{L}[\phi', g'_{\mu\nu}]. \quad (\text{A.12})$$

The Wick rotation is performed after taking $\lambda = i$. Then, we obtain the analytically continued Euclidean action as

$$I^{(\text{E})}[\phi_{\text{E}}, g_{\text{E},IJ}] = \frac{1}{i} I[\phi, g_{\mu\nu}]_{t=-i\tau} = - \int \sqrt{g_{\text{E}}} (d^4 x)_{\text{E}} \mathcal{L}[\phi_{\text{E}}, g_{\text{E},IJ}], \quad (\text{A.13})$$

where we defined the Euclidean quantities as

$$x_{\text{E}}^I = (\tau, \vec{x}_{\text{E}}) \equiv x'^{\mu} = (t', \vec{x}') = (it, \vec{x}), \quad (\text{A.14})$$

$$\partial_{\text{E},I} = (\partial_{\tau}, \vec{\partial}_{\text{E}}) \equiv \partial'_{\mu} = (\partial'_{t'}, \vec{\partial}') = (-i\partial_t, \vec{\partial}), \quad (\text{A.15})$$

$$(d^4 x)_{\text{E}} \equiv d^4 x' = i d^4 x, \quad (\text{A.16})$$

$$\sqrt{g_{\text{E}}} \equiv \sqrt{g'}|_{\lambda=i} = \sqrt{-g}. \quad (\text{A.17})$$

We also define the Euclidean scalar, vector and metric fields as follows:

$$\phi_{\text{E}}(x_{\text{E}}) \equiv \phi'(t', \vec{x}) = \phi(t, \vec{x}) = \phi(x), \quad (\text{A.18})$$

$$A_{\text{E}}^I = (A_{\text{E},\tau}, \vec{A}_{\text{E}}) \equiv A'^{\mu} = (iA^0, \vec{A}), \quad (\text{A.19})$$

$$g_{\text{E},IJ} = -g'_{\mu\nu} = \begin{pmatrix} g_{00} & ig_{0j} \\ ig_{i0} & -g_{ij} \end{pmatrix}, \quad g_{\text{E}}^{IJ} = -g'^{\mu\nu} = \begin{pmatrix} g^{00} & -ig^{0j} \\ -ig^{i0} & -g^{ij} \end{pmatrix}, \quad (\text{A.20})$$

$$\epsilon_{\text{E},IJKL} = \epsilon_{\text{E}}^{IJKL} = \epsilon_{\mu\nu\rho\sigma} = \epsilon^{\mu\nu\rho\sigma} \quad (\text{A.21})$$

where the metrics in the flat Euclidean space become the Kronecker delta δ_{IJ} and δ^{IJ} .

Here, we consider two examples: a single massless scalar field theory with a dimension-eight term and the Euler-Heisenberg theory in the following.

- Massless scalar field theory with a dimension-eight term — First, let us consider an effective action in the Minkowski space:

$$I_c[\phi] = \int d^4 x \left(\frac{1}{2} \partial_{\mu} \phi \partial^{\mu} \phi + \frac{c}{M^4} (\partial_{\mu} \phi \partial^{\mu} \phi)^2 \right), \quad (\text{A.22})$$

where ϕ denotes a background field, and a metric tensor $g_{\mu\nu} = g^{\mu\nu} = \text{diag.}(+1, -1, -1, -1)$.

By the analytic continuation, the Euclidean effective action is obtained as

$$\begin{aligned} I_c[\phi] &= \int d^4 x' \sqrt{-g'} \left(\frac{1}{2} g'^{\mu\nu} \partial'_{\mu} \phi' \partial'_{\nu} \phi' + \frac{c}{M^4} (g'^{\mu\nu} \partial'_{\mu} \phi' \partial'_{\nu} \phi')^2 \right) \\ &= \int \lambda^{-1} d^4 x' \sqrt{-\lambda^2 g'} \left(\frac{1}{2} g'^{\mu\nu} \partial'_{\mu} \phi' \partial'_{\nu} \phi' + \frac{c}{M^4} (g'^{\mu\nu} \partial'_{\mu} \phi' \partial'_{\nu} \phi')^2 \right) \\ &\rightarrow -i \int (d^4 x)_{\text{E}} \sqrt{g_{\text{E}}} \left(-\frac{1}{2} g_{\text{E}}^{IJ} \partial_{\text{E},I} \phi_{\text{E}} \partial_{\text{E},J} \phi_{\text{E}} + \frac{c}{M^4} (g_{\text{E}}^{IJ} \partial_{\text{E},I} \phi_{\text{E}} \partial_{\text{E},J} \phi_{\text{E}})^2 \right) \end{aligned}$$

$$\begin{aligned}
&= i \int (d^4x)_E \sqrt{g_E} \left(\frac{1}{2} \partial_{E,I} \phi_E \partial_{E,I} \phi_E - \frac{c}{M^4} (\partial_{E,I} \phi_E \partial_{E,I} \phi_E)^2 \right) \\
&= i \int (d^4x)_E \left(-\frac{1}{2} (\partial_\mu \phi \partial^\mu \phi) - \frac{c}{M^4} (\partial_\mu \phi \partial^\mu \phi)^2 \right) \\
&= i I_c^{(E)}[\phi_E],
\end{aligned} \tag{A.23}$$

where $g_E = +1$, $\partial_{E,I} \phi_E = (-i \cdot \partial_t \phi, \partial_i \phi)$, and the metric tensor with the Euclidean signature is the Kronecker delta, so we omitted them in the last line for short.

- Dimension-eight $U(1)$ gauge bosonic operator — Consider a effective action in Minkowski space:

$$I_e[A] = \int d^4x \left(-\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{c_1}{M^4} (F_{\mu\nu} F^{\mu\nu})^2 + \frac{c_2}{M^4} (F_{\mu\nu} \tilde{F}^{\mu\nu})^2 + \frac{c_3}{M^4} (F_{\mu\nu} F^{\mu\nu}) (F_{\mu\nu} \tilde{F}^{\mu\nu}) \right), \tag{A.24}$$

where $\tilde{F}^{\mu\nu} = \epsilon^{\mu\nu\rho\sigma} F_{\rho\sigma}/2$. Then, the Euclidean effective action is obtained as

$$\begin{aligned}
I_e[A] &= \int \sqrt{-g} d^4x \left(-\frac{1}{4} g^{\mu\rho} g^{\nu\sigma} F_{\mu\nu} F_{\rho\sigma} + \frac{c_1}{M^4} (g^{\mu\rho} g^{\nu\sigma} F_{\mu\nu} F_{\rho\sigma})^2 + \frac{c_2}{M^4} \left(\frac{1}{\sqrt{-g}} \epsilon^{\mu\nu\rho\sigma} F_{\mu\nu} F_{\rho\sigma} \right)^2 \right. \\
&\quad \left. + \frac{c_3}{M^4} (g^{\mu\rho} g^{\nu\sigma} F_{\mu\nu} F_{\rho\sigma}) \left(\frac{1}{\sqrt{-g}} \epsilon^{\mu\nu\rho\sigma} F_{\mu\nu} F_{\rho\sigma} \right) \right) \\
&= \int \sqrt{-g'} d^4x' \left(-\frac{1}{4} g'^{\mu\rho} g'^{\nu\sigma} F'_{\mu\nu} F'_{\rho\sigma} + \frac{c_1}{M^4} (g'^{\mu\rho} g'^{\nu\sigma} F'_{\mu\nu} F'_{\rho\sigma})^2 + \frac{c_2}{M^4} \left(\frac{1}{\sqrt{-g'}} \epsilon'^{\mu\nu\rho\sigma} F'_{\mu\nu} F'_{\rho\sigma} \right)^2 \right. \\
&\quad \left. + \frac{c_3}{M^4} (g'^{\mu\rho} g'^{\nu\sigma} F'_{\mu\nu} F'_{\rho\sigma}) \left(\frac{1}{\sqrt{-g'}} \epsilon'^{\mu\nu\rho\sigma} F'_{\mu\nu} F'_{\rho\sigma} \right) \right) \\
&\rightarrow -i \int \sqrt{g_E} (d^4x)_E \left(-\frac{1}{4} g_{E,IK} g_{E, JL} F_{E,IJ} F_{E,KL} + \frac{c_1}{M^4} (g_{E,IK} g_{E, JL} F_{E,IJ} F_{E,KL})^2 \right. \\
&\quad \left. - \frac{c_2}{M^4} \left(\frac{1}{\sqrt{g_E}} \epsilon_{E,IJKL} F_{E,IJ} F_{E,KL} \right)^2 - i \frac{c_3}{M^4} (g_{E,IK} g_{E, JL} F_{E,IJ} F_{E,KL}) \left(\frac{1}{\sqrt{g_E}} \epsilon_{E,IJKL} F_{E,IJ} F_{E,KL} \right) \right) \\
&= -i \int (d^4x)_E \left(-\frac{1}{4} g_{E,IK} g_{E, JL} F_{E,IJ} F_{E,KL} + \frac{c_1}{M^4} (g_{E,IK} g_{E, JL} F_{E,IJ} F_{E,KL})^2 \right. \\
&\quad \left. - \frac{c_2}{M^4} (\epsilon_{E,IJKL} F_{E,IJ} F_{E,KL})^2 - i \frac{c_3}{M^4} (g_{E,IK} g_{E, JL} F_{E,IJ} F_{E,KL}) (\epsilon_{E,IJKL} F_{E,IJ} F_{E,KL}) \right) \\
&= i \int (d^4x)_E \left(\frac{1}{4} F^{\mu\nu} F_{\mu\nu} - \frac{c_1}{M^4} (F^{\mu\nu} F_{\mu\nu})^2 - \frac{c_2}{M^4} (\epsilon^{\mu\nu\rho\sigma} F_{\mu\nu} F_{\rho\sigma})^2 + \frac{c_3}{M^4} (F^{\mu\nu} F_{\mu\nu}) (\epsilon^{\mu\nu\rho\sigma} F_{\mu\nu} F_{\rho\sigma}) \right) \\
&= i I_e^{(E)}[A],
\end{aligned} \tag{A.25}$$

where we used following relation:

$$F_{\mu\nu}^E = \frac{\partial x^\alpha}{\partial x'^\mu} \frac{\partial x^\beta}{\partial x'^\nu} F_{\alpha\beta} \bigg|_{\lambda=i} = \begin{pmatrix} 0 & -iF_{01} & -iF_{02} & -iF_{03} \\ -iF_{10} & 0 & F_{12} & F_{13} \\ -iF_{20} & F_{21} & 0 & F_{23} \\ -iF_{30} & F_{31} & F_{32} & 0 \end{pmatrix}. \tag{A.26}$$

B Relative entropy under field redefinition

To check the invariant formulation under the field redefinition, consider a target theory described by the following action in Euclidean space.

$$I_T \equiv \int (d^4x)_E \left(\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + m_A^2 \phi_A^2 - \frac{1}{M} \phi_A F_{\rho\sigma} F^{\rho\sigma} \right), \quad (\text{B.1})$$

where ϕ_A is an auxiliary field. We define the non-interacting and interacting terms as follows:

$$I_0 \equiv \int (d^4x)_E \left(\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + m_A^2 \phi_A^2 \right), \quad (\text{B.2})$$

$$I_I \equiv - \int (d^4x)_E \left(\frac{1}{M} \phi_A F_{\rho\sigma} F^{\rho\sigma} \right). \quad (\text{B.3})$$

By defining an action as $I_g \equiv I_0 + g \cdot I_I$ with the parameter g , we obtain as follows:

$$Z_g[A] \equiv \int d[\phi_A] e^{-I_g}, \quad (\text{B.4})$$

$$W_g[A] \equiv -\ln Z_g[A] = \int (d^4x)_E \left(\frac{1}{4} F_{\mu\nu} F^{\mu\nu} - g^2 \cdot \frac{1}{4m_A^2 M^2} (F_{\rho\sigma} F^{\rho\sigma})^2 \right). \quad (\text{B.5})$$

The expectation value of the interaction I_I is calculated as

$$\langle I_I \rangle_{g=0} = \left(\frac{\partial W_g}{\partial g} \right)_{g=0} = \int d[\phi_A] P_0 I_I = 0. \quad (\text{B.6})$$

Therefore, the relative entropy is calculated as

$$\begin{aligned} S(P_0 || P_g) &= W_0[A] - W_g[A] + g \langle I_I \rangle_{g=0} \\ &= W_0[A] - W_g[A] \\ &= g^2 \cdot \int (d^4x)_E \left(\frac{1}{4m_A^2 M^2} (F_{\rho\sigma} F^{\rho\sigma})^2 \right) \geq 0, \end{aligned} \quad (\text{B.7})$$

where $P_0 \equiv e^{-I_0}/Z_0[A]$ and $P_g \equiv e^{-I_g}/Z_g[A]$ are used. Here, consider the following field redefinition:

$$\phi_A \rightarrow \phi_A + g \cdot \frac{1}{2m_A^2 M} F_{\rho\sigma} F^{\rho\sigma}. \quad (\text{B.8})$$

Under this field redefinition, the actions are rewritten as

$$I_0 \rightarrow I'_0 = \int (d^4x)_E \left(\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + m_A^2 \phi_A^2 + g \cdot \frac{1}{M} \phi_A F_{\rho\sigma} F^{\rho\sigma} + g^2 \cdot \frac{1}{4m_A^2 M^2} (F_{\rho\sigma} F^{\rho\sigma})^2 \right), \quad (\text{B.9})$$

$$g \cdot I_I \rightarrow g \cdot I'_I = g \cdot I_I - g^2 \cdot \int (d^4x)_E \frac{1}{2m_A^2 M^2} (F_{\rho\sigma} F^{\rho\sigma})^2, \quad (\text{B.10})$$

$$I_g \rightarrow I'_g = \int (d^4x)_E \left(\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + m_A^2 \phi_A^2 - g^2 \cdot \frac{1}{4m_A^2 M^2} (F_{\rho\sigma} F^{\rho\sigma})^2 \right). \quad (\text{B.11})$$

Then, the relative entropy is also rewritten as

$$S(P_0||P_g) \rightarrow S(P'_0||P'_g) \quad (\text{B.12})$$

where $P'_0 \equiv e^{-I'_0}/Z'_0[A]$ and $P'_g \equiv e^{-I'_g}/Z'_g[A]$ with $Z'_0[A] \equiv \int d[\phi_A] e^{-I'_0}$ and $Z'_g[A] \equiv \int d[\phi_A] e^{-I'_g}$. Then, the relative entropy $S(P'_0||P'_g)$ is calculated as

$$\begin{aligned} S(P'_0||P'_g) &= \int d[\phi_A] (P'_0 \ln P'_0 - P'_0 \ln P'_g) \\ &= -\ln Z'_0[A] + \ln Z'_g[A] + \int d[\phi_A] P'_0 (I'_g - I'_0) \\ &= W'_0[A] - W'_g[A] - g^2 \cdot \frac{1}{2m_A^2 M^2} (F_{\rho\sigma} F^{\rho\sigma})^2 - g \cdot \int d[\phi_A] P'_0 \int (d^4x)_E \frac{1}{M} \phi_A (F_{\rho\sigma} F^{\rho\sigma}) \\ &= W'_0[A] - W'_g[A] \\ &= g^2 \cdot \int (d^4x)_E \left(\frac{1}{4m_A^2 M^2} (F_{\rho\sigma} F^{\rho\sigma})^2 \right) \geq 0, \end{aligned} \quad (\text{B.14})$$

where we used following relations.

$$\int d[\phi_A] P'_0 \int (d^4x)_E \frac{1}{M} \phi_A (F_{\rho\sigma} F^{\rho\sigma}) = -g \cdot \frac{1}{2m_A^2 M^2} (F_{\rho\sigma} F^{\rho\sigma})^2, \quad (\text{B.15})$$

$$W'_0[A] = W_0[A], \quad (\text{B.16})$$

$$W'_g[A] = W_g[A]. \quad (\text{B.17})$$

From Eq. (B.7) and (B.14), we found that the relative entropy is invariant under the field redefinition. Therefore, the relative entropy is invariant under the field redefinition once I_0 and I_1 are defined.

C Calculation of the second law of thermodynamics

We provide details of the calculation of the Clausius inequality in Eq. (6.13). The target and reference systems given in Eq. (6.11) and (6.12) are defined as

$$\rho_T \equiv \rho_{\text{ini}}, \quad (\text{C.1})$$

$$\rho_R \equiv U^\dagger \rho_{\text{fin},S} \otimes e^{-\beta H_B} / Z_B(\beta) U, \quad (\text{C.2})$$

respectively. The relative entropy between ρ_T and ρ_R is given by

$$S(\rho_T||\rho_R) = \text{Tr}[\rho_T \ln \rho_T - \rho_T \ln \rho_R] \quad (\text{C.3})$$

$$= \text{Tr}[\rho_T \ln \rho_T] - \text{Tr}[\rho_R \ln \rho_R] + \text{Tr}[\rho_R \ln \rho_R] - \text{Tr}[\rho_T \ln \rho_R]. \quad (\text{C.4})$$

We calculate each term on the right-hand side as follows:

$$\text{Tr}[\rho_T \ln \rho_T] = \text{Tr} \left[\rho_{\text{ini},S} \otimes e^{-\beta H_B} / Z_B(\beta) \ln \rho_{\text{ini},S} \otimes e^{-\beta H_B} / Z_B(\beta) \right]$$

$$= \text{Tr}_S [\rho_{\text{ini},S} \ln \rho_{\text{ini},S}] + \text{Tr}_B \left[e^{-\beta H_B} / Z_B(\beta) \ln e^{-\beta H_B} / Z_B(\beta) \right], \quad (\text{C.5})$$

$$\begin{aligned} \text{Tr}[\rho_R \ln \rho_R] &= \text{Tr} \left[U^\dagger \rho_{\text{fin},S} \otimes e^{-\beta H_B} / Z_B(\beta) U \ln U^\dagger \rho_{\text{fin},S} \otimes e^{-\beta H_B} / Z_B(\beta) U \right] \\ &= \text{Tr} \left[\rho_{\text{fin},S} \otimes e^{-\beta H_B} / Z_B(\beta) \ln \rho_{\text{fin},S} \otimes e^{-\beta H_B} / Z_B(\beta) \right] \\ &= \text{Tr}_S [\rho_{\text{fin},S} \ln \rho_{\text{fin},S}] + \text{Tr}_B \left[e^{-\beta H_B} / Z_B(\beta) \ln e^{-\beta H_B} / Z_B(\beta) \right] \\ &= \text{Tr}_S [\rho_{\text{fin},S} \ln \rho_{\text{fin},S}] - \ln Z_B(\beta) - \beta \cdot \text{Tr} \left[H_B \rho_{\text{ini},S} \otimes e^{-\beta H_B} / Z_B(\beta) \right], \end{aligned} \quad (\text{C.6})$$

$$\begin{aligned} \text{Tr}[\rho_T \ln \rho_R] &= \text{Tr}[\rho_{\text{ini},S} \otimes e^{-\beta H_B} / Z_B(\beta) \ln U^\dagger \rho_{\text{fin},S} \otimes e^{-\beta H_B} / Z_B(\beta) U] \\ &= \text{Tr}[U \rho_{\text{ini},S} \otimes e^{-\beta H_B} / Z_B(\beta) U^\dagger \ln \rho_{\text{fin},S} \otimes e^{-\beta H_B} / Z_B(\beta)] \\ &= \text{Tr}[U \rho_{\text{ini},S} \otimes e^{-\beta H_B} / Z_B(\beta) U^\dagger \ln \rho_{\text{fin},S}] + \text{Tr}[U \rho_S \otimes e^{-\beta H_B} / Z_B(\beta) U^\dagger \ln e^{-\beta H_B} / Z_B(\beta)] \\ &= \text{Tr}_S [\rho_{\text{fin},S} \ln \rho_{\text{fin},S}] + \text{Tr}[U \rho_{\text{ini},S} \otimes e^{-\beta H_B} / Z_B(\beta) U^\dagger \ln e^{-\beta H_B} / Z_B(\beta)] \\ &= \text{Tr}_S [\rho_{\text{fin},S} \ln \rho_{\text{fin},S}] - \ln Z_B(\beta) - \beta \cdot \text{Tr} \left[U^\dagger H_B U \rho_{\text{ini},S} \otimes e^{-\beta H_B} / Z_B(\beta) \right]. \end{aligned} \quad (\text{C.7})$$

Then we obtain the Clausius inequality from the non-negativity of the relative entropy:

$$S(\rho_T || \rho_R) = \text{Tr}_S [\rho_{\text{ini},S} \ln \rho_{\text{ini},S}] - \text{Tr}_S [\rho_{\text{fin},S} \ln \rho_{\text{fin},S}] - \beta \cdot (\text{Tr} [H_B \rho_{\text{ini}}] - \text{Tr} [H_B \rho_{\text{fin}}]) \quad (\text{C.8})$$

$$= S(\rho_{\text{fin},S}) - S(\rho_{\text{ini},S}) - \beta \cdot Q \geq 0, \quad (\text{C.9})$$

where $S(\rho) \equiv -\text{Tr}_S [\rho \ln \rho]$ denotes the von Neumann entropy of S, and $Q \equiv \text{Tr} [H_B \rho_{\text{ini}}] - \text{Tr} [H_B \rho_{\text{fin}}]$ is the heat exchange between S and B.

D Einstein-Maxwell theory with higher-derivative operators under field redefinition

We explain the field redefinitions to derive Eq. (4.152) following the procedures and notations of Ref. [49]. The higher-derivative operators up to the four derivative terms of the Einstein-Maxwell theory are expressed as

$$\begin{aligned} \mathcal{L} &= \frac{M_{\text{Pl}}^2}{2} R - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} \\ &\quad + c_1 R^2 + c_2 R_{\mu\nu} R^{\mu\nu} + c_3 R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} \\ &\quad + c_4 R F_{\mu\nu} F^{\mu\nu} + c_5 R_{\mu\nu} F^{\mu\rho} F^\nu{}_\rho + c_6 R_{\mu\nu\rho\sigma} F^{\mu\nu} F^{\rho\sigma}, \\ &\quad + c_7 F_{\mu\nu} F^{\mu\nu} F_{\rho\sigma} F^{\rho\sigma} + c_8 F_{\mu\nu} F^{\nu\rho} F_{\rho\sigma} F^{\sigma\mu}, \end{aligned} \quad (\text{D.1})$$

where terms involving $\nabla_\rho F_{\mu\nu}$ or $\nabla_\mu F^{\mu\nu}$ vanish. Also, the Gauss-Bonnet combination, i.e., $R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} - 4R_{\mu\nu} R^{\mu\nu} + R^2$, is a total derivative and vanishes for the extremal black hole in four dimensions. Thus, in four dimensions, Eq. (D.1) is expressed as

$$\begin{aligned} \mathcal{L} &= \frac{M_{\text{Pl}}^2}{2} R - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} \\ &\quad + (c_1 - c_3) R^2 + (c_2 + 4c_3) R_{\mu\nu} R^{\mu\nu} \\ &\quad + c_4 R F_{\mu\nu} F^{\mu\nu} + c_5 R_{\mu\nu} F^{\mu\rho} F^\nu{}_\rho + c_6 R_{\mu\nu\rho\sigma} F^{\mu\nu} F^{\rho\sigma} \end{aligned}$$

$$+ c_7 F_{\mu\nu} F^{\mu\nu} F_{\rho\sigma} F^{\rho\sigma} + c_8 F_{\mu\nu} F^{\nu\rho} F_{\rho\sigma} F^{\sigma\mu}, \quad (\text{D.2})$$

$$\begin{aligned} &= \frac{M_{\text{Pl}}^2}{2} R - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} \\ &+ c_{13} R^2 + c_{23} R_{\mu\nu} R^{\mu\nu} \\ &+ c_4 R F_{\mu\nu} F^{\mu\nu} + c_5 R_{\mu\nu} F^{\mu\rho} F^{\nu\rho} + c_6 R_{\mu\nu\rho\sigma} F^{\mu\nu} F^{\rho\sigma} \\ &+ \left(c_7 + \frac{c_8}{2} \right) (F_{\mu\nu} F^{\mu\nu})^2 + \frac{c_8}{4} (F_{\mu\nu} \tilde{F}^{\mu\nu})^2, \end{aligned} \quad (\text{D.3})$$

where $c_{13} \equiv c_1 - c_3$, $c_{23} \equiv c_2 + 4c_3$, and we used

$$F_{\mu\nu} F^{\nu\rho} F_{\rho\sigma} F^{\sigma\mu} = \frac{1}{2} (F_{\mu\nu} F^{\mu\nu})^2 + \frac{1}{4} (F_{\mu\nu} \tilde{F}^{\mu\nu})^2. \quad (\text{D.4})$$

Consider a field redefinition of $g_{\mu\nu}$ [49],

$$g_{\mu\nu} \rightarrow g_{\mu\nu} + \delta g_{\mu\nu}, \quad (\text{D.5})$$

where

$$\delta g_{\mu\nu} = r_1 R_{\mu\nu} + r_2 g_{\mu\nu} R + r_3 M_{\text{Pl}}^{-2} F_{\mu\rho} F_{\nu}{}^{\rho} + r_4 M_{\text{Pl}}^{-2} g_{\mu\nu} F_{\rho\sigma} F^{\rho\sigma}, \quad (\text{D.6})$$

with a set of four constants r_i . Under this field redefinition, the coefficients of the higher-derivative operator in four dimensions are shifted as follows:

$$c_{13} \rightarrow c_{13} - \frac{M_{\text{Pl}}^2}{4} r_1 - \frac{M_{\text{Pl}}^2}{2} r_2, \quad (\text{D.7})$$

$$c_{23} \rightarrow c_{23} + \frac{M_{\text{Pl}}^2}{2} r_1, \quad (\text{D.8})$$

$$c_4 \rightarrow c_4 + \frac{1}{8} r_1 - \frac{1}{4} r_3 - \frac{1}{2} r_4, \quad (\text{D.9})$$

$$c_5 \rightarrow c_5 - \frac{1}{2} r_1 + \frac{1}{2} r_3, \quad (\text{D.10})$$

$$c_6 \rightarrow c_6, \quad (\text{D.11})$$

$$c_7 \rightarrow c_7 + \frac{M_{\text{Pl}}^{-2}}{8} r_3, \quad (\text{D.12})$$

$$c_8 \rightarrow c_8 - \frac{M_{\text{Pl}}^{-2}}{2} r_3. \quad (\text{D.13})$$

Equation (4.152) is derived by choosing the set of four constants as follows:

$$r_1 = -\frac{2}{M_{\text{Pl}}^2} c_{23}, \quad (\text{D.14})$$

$$r_2 = \frac{2}{M_{\text{Pl}}^2} \left(c_{13} + \frac{1}{2} c_{23} \right), \quad (\text{D.15})$$

$$r_3 = -2 \left(c_5 + \frac{1}{M_{\text{Pl}}^2} c_{23} \right), \quad (\text{D.16})$$

$$r_4 = 2 \left(\frac{1}{4M_{\text{Pl}}^2} c_{23} + c_4 + \frac{1}{2} c_5 \right). \quad (\text{D.17})$$

After the above field redefinitions, Eq. (D.3) is rewritten as follows:

$$\begin{aligned} \mathcal{L} &= \frac{M_{\text{Pl}}^2}{2} R - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} + c_6 R_{\mu\nu\rho\sigma} F^{\mu\nu} F^{\rho\sigma} \\ &+ \left(\frac{1}{4M_{\text{Pl}}^4} c_{23} + \frac{1}{4M_{\text{Pl}}^2} c_5 + \frac{1}{2} (2c_7 + c_8) \right) (F_{\mu\nu} F^{\mu\nu})^2 + \left(\frac{1}{4M_{\text{Pl}}^4} c_{23} + \frac{1}{4M_{\text{Pl}}^2} c_5 + \frac{1}{4} c_8 \right) (F_{\mu\nu} \tilde{F}^{\mu\nu})^2, \\ &= \int d^4x \sqrt{-g} \left(\frac{M_{\text{Pl}}^2}{2} R - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{\alpha_1}{4M_{\text{Pl}}^4} (F_{\mu\nu} F^{\mu\nu})^2 + \frac{\alpha_2}{4M_{\text{Pl}}^4} (F_{\mu\nu} \tilde{F}^{\mu\nu})^2 + \frac{\alpha_3}{2M_{\text{Pl}}^2} F_{\mu\nu} F_{\rho\sigma} R^{\mu\nu\rho\sigma} \right), \end{aligned} \quad (\text{D.18})$$

with

$$\alpha_1 \equiv c_{23} + M_{\text{Pl}}^2 \cdot c_5 + 2M_{\text{Pl}}^4 \cdot (2c_7 + c_8), \quad (\text{D.19})$$

$$\alpha_2 \equiv c_{23} + M_{\text{Pl}}^2 \cdot c_5 + M_{\text{Pl}}^4 \cdot c_8, \quad (\text{D.20})$$

$$\alpha_3 \equiv 2M_{\text{Pl}}^2 \cdot c_6. \quad (\text{D.21})$$

References

- [1] **ATLAS** Collaboration, G. Aad et al., *Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC*, *Phys. Lett. B* **716** (2012) 1–29, [[arXiv:1207.7214](#)].
- [2] **CMS** Collaboration, S. Chatrchyan et al., *Observation of a New Boson at a Mass of 125 GeV with the CMS Experiment at the LHC*, *Phys. Lett. B* **716** (2012) 30–61, [[arXiv:1207.7235](#)].
- [3] B. Grzadkowski, M. Iskrzynski, M. Misiak, and J. Rosiek, *Dimension-Six Terms in the Standard Model Lagrangian*, *JHEP* **10** (2010) 085, [[arXiv:1008.4884](#)].
- [4] B. Henning, X. Lu, T. Melia, and H. Murayama, *2, 84, 30, 993, 560, 15456, 11962, 261485, ...: Higher dimension operators in the SM EFT*, *JHEP* **08** (2017) 016, [[arXiv:1512.03433](#)]. [Erratum: *JHEP* **09**, 019 (2019)].
- [5] E. E. Jenkins, A. V. Manohar, and M. Trott, *Renormalization Group Evolution of the Standard Model Dimension Six Operators I: Formalism and lambda Dependence*, *JHEP* **10** (2013) 087, [[arXiv:1308.2627](#)].
- [6] E. E. Jenkins, A. V. Manohar, and M. Trott, *Renormalization Group Evolution of the Standard Model Dimension Six Operators II: Yukawa Dependence*, *JHEP* **01** (2014) 035, [[arXiv:1310.4838](#)].
- [7] R. Alonso, E. E. Jenkins, A. V. Manohar, and M. Trott, *Renormalization Group Evolution of the Standard Model Dimension Six Operators III: Gauge Coupling Dependence and Phenomenology*, *JHEP* **04** (2014) 159, [[arXiv:1312.2014](#)].
- [8] I. Brivio and M. Trott, *The Standard Model as an Effective Field Theory*, *Phys. Rept.* **793** (2019) 1–98, [[arXiv:1706.08945](#)].
- [9] H.-L. Li, Z. Ren, J. Shu, M.-L. Xiao, J.-H. Yu, and Y.-H. Zheng, *Complete set of dimension-eight operators in the standard model effective field theory*, *Phys. Rev. D* **104** (2021), no. 1 015026, [[arXiv:2005.00008](#)].
- [10] C. W. Murphy, *Dimension-8 operators in the Standard Model Effective Field Theory*, *JHEP* **10** (2020) 174, [[arXiv:2005.00059](#)].
- [11] Z. Han and W. Skiba, *Effective theory analysis of precision electroweak data*, *Phys. Rev. D* **71** (2005) 075009, [[hep-ph/0412166](#)].
- [12] A. Pomarol and F. Riva, *Towards the Ultimate SM Fit to Close in on Higgs Physics*, *JHEP* **01** (2014) 151, [[arXiv:1308.2803](#)].
- [13] T. Corbett, O. J. P. Eboli, J. Gonzalez-Fraile, and M. C. Gonzalez-Garcia, *Robust Determination of the Higgs Couplings: Power to the Data*, *Phys. Rev. D* **87** (2013) 015022, [[arXiv:1211.4580](#)].
- [14] J. Ellis, V. Sanz, and T. You, *The Effective Standard Model after LHC Run I*, *JHEP* **03** (2015) 157, [[arXiv:1410.7703](#)].
- [15] B. Dumont, *Higgs, supersymmetry and dark matter after Run I of the LHC*. PhD thesis, Grenoble U., 2014. [[arXiv:1411.3465](#)].
- [16] T. Corbett, O. J. P. Éboli, J. Gonzalez-Fraile, and M. C. Gonzalez-Garcia, *Determining Triple Gauge Boson Couplings from Higgs Data*, *Phys. Rev. Lett.* **111** (2013) 011801, [[arXiv:1304.1151](#)].
- [17] W.-F. Chang, W.-P. Pan, and F. Xu, *Effective gauge-Higgs operators analysis of new physics associated*

- with the Higgs boson, *Phys. Rev. D* **88** (2013), no. 3 033004, [[arXiv:1303.7035](#)].
- [18] J. Elias-Miro, J. R. Espinosa, E. Masso, and A. Pomarol, *Higgs windows to new physics through $d=6$ operators: constraints and one-loop anomalous dimensions*, *JHEP* **11** (2013) 066, [[arXiv:1308.1879](#)].
 - [19] E. Boos, V. Bunichev, M. Dubinin, and Y. Kurihara, *Higgs boson signal at complete tree level in the SM extension by dimension-six operators*, *Phys. Rev. D* **89** (2014) 035001, [[arXiv:1309.5410](#)].
 - [20] J. Ellis, V. Sanz, and T. You, *Complete Higgs Sector Constraints on Dimension-6 Operators*, *JHEP* **07** (2014) 036, [[arXiv:1404.3667](#)].
 - [21] A. Falkowski and F. Riva, *Model-independent precision constraints on dimension-6 operators*, *JHEP* **02** (2015) 039, [[arXiv:1411.0669](#)].
 - [22] L. Berthier, M. Bjørn, and M. Trott, *Incorporating doubly resonant W^\pm data in a global fit of SMEFT parameters to lift flat directions*, *JHEP* **09** (2016) 157, [[arXiv:1606.06693](#)].
 - [23] S. Banerjee, R. S. Gupta, J. Y. Reiness, S. Seth, and M. Spannowsky, *Towards the ultimate differential SMEFT analysis*, *JHEP* **09** (2020) 170, [[arXiv:1912.07628](#)].
 - [24] A. Biekötter, R. Gomez-Ambrosio, P. Gregg, F. Krauss, and M. Schönherr, *Constraining SMEFT operators with associated $h\gamma$ production in weak boson fusion*, *Phys. Lett. B* **814** (2021) 136079, [[arXiv:2003.06379](#)].
 - [25] A. Efrati, A. Falkowski, and Y. Soreq, *Electroweak constraints on flavorful effective theories*, *JHEP* **07** (2015) 018, [[arXiv:1503.07872](#)].
 - [26] L. Silvestrini and M. Valli, *Model-independent Bounds on the Standard Model Effective Theory from Flavour Physics*, *Phys. Lett. B* **799** (2019) 135062, [[arXiv:1812.10913](#)].
 - [27] S. Descotes-Genon, A. Falkowski, M. Fedele, M. González-Alonso, and J. Virto, *The CKM parameters in the SMEFT*, *JHEP* **05** (2019) 172, [[arXiv:1812.08163](#)].
 - [28] J. Aebischer, J. Kumar, P. Stangl, and D. M. Straub, *A Global Likelihood for Precision Constraints and Flavour Anomalies*, *Eur. Phys. J. C* **79** (2019), no. 6 509, [[arXiv:1810.07698](#)].
 - [29] T. Hurth, S. Renner, and W. Shepherd, *Matching for FCNC effects in the flavour-symmetric SMEFT*, *JHEP* **06** (2019) 029, [[arXiv:1903.00500](#)].
 - [30] J. Aebischer, C. Bobeth, A. J. Buras, and J. Kumar, *SMEFT ATLAS of $\Delta F = 2$ transitions*, *JHEP* **12** (2020) 187, [[arXiv:2009.07276](#)].
 - [31] R. Aoude, T. Hurth, S. Renner, and W. Shepherd, *The impact of flavour data on global fits of the MFV SMEFT*, *JHEP* **12** (2020) 113, [[arXiv:2003.05432](#)].
 - [32] D. A. Faroughy, G. Isidori, F. Wilsch, and K. Yamamoto, *Flavour symmetries in the SMEFT*, *JHEP* **08** (2020) 166, [[arXiv:2005.05366](#)].
 - [33] A. Falkowski and K. Mimouni, *Model independent constraints on four-lepton operators*, *JHEP* **02** (2016) 086, [[arXiv:1511.07434](#)].
 - [34] A. Falkowski, M. González-Alonso, and K. Mimouni, *Compilation of low-energy constraints on 4-fermion operators in the SMEFT*, *JHEP* **08** (2017) 123, [[arXiv:1706.03783](#)].
 - [35] A. Falkowski, M. González-Alonso, and O. Naviliat-Cuncic, *Comprehensive analysis of beta decays within and beyond the Standard Model*, *JHEP* **04** (2021) 126, [[arXiv:2010.13797](#)].

- [36] N. Arkani-Hamed, L. Motl, A. Nicolis, and C. Vafa, *The String landscape, black holes and gravity as the weakest force*, *JHEP* **06** (2007) 060, [[hep-th/0601001](#)].
- [37] D. Harlow, B. Heidenreich, M. Reece, and T. Rudelius, *The Weak Gravity Conjecture: A Review*, [arXiv:2201.08380](#).
- [38] W. Taylor and Y.-N. Wang, *The F-theory geometry with most flux vacua*, *JHEP* **12** (2015) 164, [[arXiv:1511.03209](#)].
- [39] C. Vafa, *The String landscape and the swampland*, [hep-th/0509212](#).
- [40] T. Banks and N. Seiberg, *Symmetries and Strings in Field Theory and Gravity*, *Phys. Rev. D* **83** (2011) 084019, [[arXiv:1011.5120](#)].
- [41] H. Ooguri and C. Vafa, *On the Geometry of the String Landscape and the Swampland*, *Nucl. Phys. B* **766** (2007) 21–33, [[hep-th/0605264](#)].
- [42] T. W. Grimm, E. Palti, and I. Valenzuela, *Infinite Distances in Field Space and Massless Towers of States*, *JHEP* **08** (2018) 143, [[arXiv:1802.08264](#)].
- [43] H. Ooguri and C. Vafa, *Non-supersymmetric AdS and the Swampland*, *Adv. Theor. Math. Phys.* **21** (2017) 1787–1801, [[arXiv:1610.01533](#)].
- [44] B. Freivogel and M. Kleban, *Vacua Morphulism*, [arXiv:1610.04564](#).
- [45] G. Obied, H. Ooguri, L. Spodyneiko, and C. Vafa, *De Sitter Space and the Swampland*, [arXiv:1806.08362](#).
- [46] H. Ooguri, E. Palti, G. Shiu, and C. Vafa, *Distance and de Sitter Conjectures on the Swampland*, *Phys. Lett. B* **788** (2019) 180–184, [[arXiv:1810.05506](#)].
- [47] S. K. Garg and C. Krishnan, *Bounds on Slow Roll and the de Sitter Swampland*, *JHEP* **11** (2019) 075, [[arXiv:1807.05193](#)].
- [48] G. Goon and R. Penco, *Universal Relation between Corrections to Entropy and Extremality*, *Phys. Rev. Lett.* **124** (2020), no. 10 101103, [[arXiv:1909.05254](#)].
- [49] C. Cheung, J. Liu, and G. N. Remmen, *Proof of the Weak Gravity Conjecture from Black Hole Entropy*, *JHEP* **10** (2018) 004, [[arXiv:1801.08546](#)].
- [50] C. Cheung, J. Liu, and G. N. Remmen, *Entropy Bounds on Effective Field Theory from Rotating Dyonic Black Holes*, *Phys. Rev. D* **100** (2019), no. 4 046003, [[arXiv:1903.09156](#)].
- [51] A. Adams, N. Arkani-Hamed, S. Dubovsky, A. Nicolis, and R. Rattazzi, *Causality, analyticity and an IR obstruction to UV completion*, *JHEP* **10** (2006) 014, [[hep-th/0602178](#)].
- [52] B. Bellazzini, M. Lewandowski, and J. Serra, *Positivity of Amplitudes, Weak Gravity Conjecture, and Modified Gravity*, *Phys. Rev. Lett.* **123** (2019), no. 25 251103, [[arXiv:1902.03250](#)].
- [53] Y. Hamada, T. Noumi, and G. Shiu, *Weak Gravity Conjecture from Unitarity and Causality*, *Phys. Rev. Lett.* **123** (2019), no. 5 051601, [[arXiv:1810.03637](#)].
- [54] Q.-H. Cao and D. Ueda, *Entropy Constraint on Effective Field Theory*, [arXiv:2201.00931](#).
- [55] S. Kullback and R. A. Leibler, *On Information and Sufficiency*, *The Annals of Mathematical Statistics* **22** (1951), no. 1 79 – 86.
- [56] H. Umegaki, *Conditional expectation in an operator algebra. IV. Entropy and information*, *Kodai*

Mathematical Seminar Reports **14** (1962), no. 2 59 – 85.

- [57] A. Wehrl, *General properties of entropy*, *Rev. Mod. Phys.* **50** (Apr, 1978) 221–260.
- [58] H. Tasaki, *Jarzynski Relations for Quantum Systems and Some Applications*, *arXiv e-prints* (Sept., 2000) cond-mat/0009244, [[cond-mat/0009244](#)].
- [59] T. SAGAWA, *Second law-like inequalities with quantum relative entropy: An introduction*, *Lectures on Quantum Computing, Thermodynamics and Statistical Physics* (Oct, 2012) 125–190.
- [60] J. Quevillon, C. Smith, and S. Touati, *Effective action for gauge bosons*, *Phys. Rev. D* **99** (2019), no. 1 013003, [[arXiv:1810.06994](#)].
- [61] Y. Nambu, *Quantum electrodynamics in nonlinear gauge*, *Prog. Theor. Phys. Suppl. E* **68** (1968) 190–195.
- [62] A. Adams, A. Jenkins, and D. O’Connell, *Signs of analyticity in fermion scattering*, [arXiv:0802.4081](#).
- [63] N. Arkani-Hamed, Y.-t. Huang, J.-Y. Liu, and G. N. Remmen, *Causality, Unitarity, and the Weak Gravity Conjecture*, [arXiv:2109.13937](#).
- [64] A. Y. Morozov, *MATRIX OF MIXING OF SCALAR AND VECTOR MESONS OF DIMENSION $D \leq 8$ IN QCD. (IN RUSSIAN)*, *Sov. J. Nucl. Phys.* **40** (1984) 505.
- [65] G. N. Remmen and N. L. Rodd, *Consistency of the Standard Model Effective Field Theory*, *JHEP* **12** (2019) 032, [[arXiv:1908.09845](#)].
- [66] D. Li, D. Meltzer, and D. Poland, *Conformal Bootstrap in the Regge Limit*, *JHEP* **12** (2017) 013, [[arXiv:1705.03453](#)].
- [67] N. Afkhami-Jeddi, S. Kundu, and A. Tajdini, *A Conformal Collider for Holographic CFTs*, *JHEP* **10** (2018) 156, [[arXiv:1805.07393](#)].
- [68] T. Banks, M. Johnson, and A. Shomer, *A Note on Gauge Theories Coupled to Gravity*, *JHEP* **09** (2006) 049, [[hep-th/0606277](#)].
- [69] M. Montero, *A Holographic Derivation of the Weak Gravity Conjecture*, *JHEP* **03** (2019) 157, [[arXiv:1812.03978](#)].