The Dissipation of the Solar Nebula Constrained by Impacts and Core Cooling in Planetesimals

Alison C. Hunt¹, Karen J. Theis², Mark Rehkämper³, Gretchen K. Benedix^{4, 5}, Rasmus **Andreasen6 and Maria Schönbächler1**

¹Institute of Geochemistry and Petrology, ETH Zürich, Clausiusstrasse 25, 8092 Zürich, Switzerland.

2 School of Earth and Environmental Sciences, The University of Manchester, Manchester, M13 9PL, UK.

Now at: The Photon Science Institute, Alan Turing Building, The University of Manchester, Manchester, M13 9PL, UK.

³³ Department of Earth Science & Engineering, Imperial College London, London, SW7 2AZ, UK.

4 Space Science and Technology Centre, School of Earth and Planetary Sciences, Curtin University, GPO Box U1987, Perth, WA 6845, Australia.

5 Department of Earth and Planetary Sciences, Western Australian Museum, Locked Bag 49 Welshpool DC, Western Australia 6986, Australia.

6 Department of Geoscience, Aarhus University, Høegh-Guldbergs Gade 2, 8000 Aarhus C, Denmark.

***Correspondence and requests for materials should be addressed to:** [alison.hunt@erdw.ethz.ch;](mailto:alison.hunt@erdw.ethz.ch) +41 (0)44 632 4643; +41 (0)44 632 1376 (fax)

Accepted by Nature Astronomy, 01/04/2022

Abstract

Rapid cooling of planetesimal cores has been inferred for several iron meteorite parent bodies based on metallographic cooling rates, and linked to the loss of their insulating mantles during impacts. However, the timing of these disruptive events is poorly constrained. Here, we used the short-lived 107Pd-107Ag decay system to date rapid core cooling by determining Pd-Ag ages for iron meteorites. We show closure times for the iron meteorites equate to cooling in the timeframe ~7.8-11.7 Myr after CAI, and indicate that an energetic inner Solar System persisted at this time. This likely results from the dissipation of gas in the protoplanetary disk, after which the damping effect of gas drag ceases. An early giant planet instability between 5–14 Myr after CAI could have reinforced this effect. This correlates well with the timing of impacts recorded by the Pd-Ag system for iron meteorites.

Main Text

Iron meteorites are thought to represent the once heated interior portions of planetesimals and are some of the earliest-formed bodies in our Solar System¹. As such, they are survivors of the many dynamical processes that have shaped Solar System architecture, including dissipation of the protoplanetary disk, and runaway growth, migration and reorganisation of the giant planets^{2,3}. Importantly, high and variable metallographic cooling rates among many iron meteorite groups, including the IIAB, IIIAB and IVAs, indicate that these metals cooled in bodies that had been stripped of their insulating silicate mantles^{4,5}. Dating core crystallisation can therefore elucidate the timing of impact processes, which in turn constrains the evolution of our early Solar System. The short-lived ¹⁰⁷Pd-¹⁰⁷Ag decay system $(t_{1/2}$ ~6.5 Myr) provides a tool for this. Palladium, which is highly siderophile, partitions into the metal phase during crystallisation, and chalcophile Ag is strongly sequestered into sulfides. This leads to Pd-Ag fractionation during crystallisation of a cooling core resulting in samples from a single body with variable Pd/Ag ratios, thereby allowing us to derive parent-body initial $107Pd/108Pd$ ratios from isochron slopes. The ages for parent-body cooling through the closure temperature of the Pd-Ag system (~820-970 K) ⁶ can be derived from these initials.

Early studies of iron meteorites using the Pd-Ag chronometer did not fully account for the exposure of samples to galactic cosmic rays (GCR)⁷⁻⁹, which can disturb the system through secondary neutron capture reactions by Pd isotopes that lead to a decrease in ¹⁰⁷Ag/¹⁰⁹Ag. This occurs primarily via neutron capture on ¹⁰⁸Pd and subsequent β[—]decay to 109 Ag, and requires correction to achieve meaningful ages¹⁰. After GCR correction, initial ¹⁰⁷Pd/¹⁰⁸Pd ratios of individual meteorites from the iron meteorite groups IIAB, IIIAB and IVA imply core cooling between ~7.2 Myr and 10.7 Myr after the formation of calcium-aluminium rich inclusions (CAIs)^{6,11,12}, relative to the Solar System initial (SSI)¹⁰⁷Pd/¹⁰⁸Pd determined from carbonaceous chondrites¹³. Additionally, a detailed investigation of the partially-differentiated IAB irons yielded two isochrons that signified a later closure of the Pd-Ag system between \sim 14.9–18.7 Myr after CAI⁹. However, those data were not corrected for GCR effects. Here, we focus on parent body initial 107Pd/108Pd ratios to reassess the thermal evolution of planetesimals and constrain dynamic processes occurring in the early Solar System. We report new Pd-Ag data and use Pt isotopes to correct for the effects of GCR in the IAB, IIAB and IIIAB iron meteorites.

Results and Pd-Ag ages

We determined Pd-Ag and Pt isotope compositions for 10 IIAB and IIIAB iron meteorites and Pt isotope data for a further 8 IAB samples where Pd-Ag data were previously published⁹ (Table 1; Methods; Supplementary Materials). Data are presented in the epsilon notation (i.e., ε¹⁹ⁱPt/¹⁹⁵Pt is the deviation in parts per 10,000 from the average of two bracketing standards). Platinum isotope values for the iron meteorites range from ~0.0 to 0.4 for ϵ^{196} Pt (Table 1; Extended Data Fig. 1; Supplementary Materials) and compare well to published values¹⁴⁻¹⁶. Platinum isotope ratios fit well to the modelled effects for neutron capture, reaffirming that Pt isotopes are a powerful neutron dosimeter.

Corrections for GCR effects were applied to ε^{107} Ag (Methods; Table 2) and parent-body isochrons were calculated (Table 3; Extended Data Fig. 2). Corrected data define a single IAB isochron with a slope of 1.51 (± 0.16) x10⁻⁵, which is significantly different to the two previously determined isochrons⁹. The IIABs yield an isochron with a slope of 1.70 (±0.48) x10⁻⁵, which is within uncertainty of previous GCR-corrected sample-specific isochrons for the group¹¹. The IIIAB samples together define an isochron with a slope of 2.21 (± 0.57) x10⁻⁵, which overlaps with sample-specific $107Pd/108Pd$ ratios determined for this group^{6,11}. The GCR-corrected initial ¹⁰⁷Pd/¹⁰⁸Pd ratios for the IABs, IIAB and IIIABs are within uncertainty of each other, indicating that the Pd-Ag system closed in a similar time-frame for these bodies (Fig. 1).

To determine Pd-Ag ages, the SSI ¹⁰⁷Pd/¹⁰⁸Pd is needed. Two distinct initial ¹⁰⁷Pd/¹⁰⁸Pd ratios have been determined based on the IVA iron meteorite Muonionalusta (~3.5 x10⁻⁵; ¹²) and carbonaceous chondrites (\sim 5.9 x10⁻⁵; ¹³; see Methods for discussion of SSI). To consider the full range of SSI ¹⁰⁷Pd/¹⁰⁸Pd ratios, we calculate Pd-Ag ages for the meteorite groups studied here relative to both (Table 3).

Discussion

The presence of silicate inclusions and chemical data indicate the IAB iron meteorites had a complex thermal evolution and can be divided into sub-groups. The IABs studied here are members of the main group (MG) and 'Low-Au Low-Ni' (sLL) sub-group, which may be genetically related¹⁷ (Table 1). It has been argued that the IABs formed in multiple impact melt pools on a single parent body¹⁷. Alternatively, they are products of internal heating followed by a catastrophic impact and reassembly of the parent body¹⁸. In support of the latter, samples from IAB sub-groups yield identical ¹⁸²Hf-¹⁸²W ages indicating contemporaneous metal–silicate separation^{14,19}. After the new correction for GCR effects, Pd-Ag data for the IABs define a single isochron indicating that cooling on the parent asteroid occurred in several sub-groups simultaneously. Small pools of metal formed at different times would not be expected to cool simultaneously and hence our new isochron likely represents a single body-wide cooling event, lending further support to a model of internal heating followed by breakup of the parent body.

Importantly, heating of the IAB parent asteroid produced a body which only partially differentiated at 6.0 ± 0.8 Myr¹⁴. The question remains whether the body was impacted while at or close to its peak temperature, thereby effectively halting differentiation¹⁸, or whether late accretion of the parent asteroid limited the availability of ²⁶AI and allowed only subdued internal heating¹⁴. Thermal models explored a scenario where the body cooled slowly until \sim 10-12 Myr when the catastrophic breakup occurred¹⁴. However, the model used Pd-Ag data not corrected for GCR effects 9 to constrain the timing of the break-up, and hence a reassessment is necessary.

Our new IAB isochron corresponds to an age of 12.8 $^{+3.1}/_{4.6}$ or 7.9 $^{+1.0}/_{1.1}$ Myr after CAI (Table 3). Fast cooling rates are expected for the IAB parent body to keep silicate inclusions suspended in the metal portions¹⁷. Hence, a catastrophic impact should be closely followed by closure of the Pd-Ag system. Integrating this with cooling at 12.8 +3.1/-4.6 Myr suggests a delay between the body reaching its peak temperature at \sim 6 Myr and the catastrophic impact (Extended Data Fig. 3). Closure ages from the $1291-129Xe$ system for a silicate inclusion from Campo del Cielo indicate a scrambling event that occurred later than \sim 11 Myr after CAI²⁰. This agrees well with evidence of resetting from the Hf-W system at 14.9±2.8 and 11.4±2.2 Myr, respectively, in both the genetically related winonaite meteorites and a further silicate inclusion from Campo del Cielo²¹⁻²³. This resetting must be impact-related and its identification in both winonaites and IABs indicates a large-scale process that likely corresponds to the catastrophic break-up of the body. Together with the Pd-Ag system these chronometers suggest an impact event between ~11 and 13.6 Myr after CAI. Assuming that the IAB parent was impacted at this time yields the necessary fast cooling rates of between instantaneous and ~102 K/Myr (Table 3; Methods). In this scenario, the thermal model and precursor parameters determined previously remain applicable to the IAB parent body¹⁴.

Alternatively, a cooling age of 7.9 $^{+1.0}$ /-1.1 Myr (calculated relative to the SSI $^{107}Pd/^{108}Pd$ of \sim 3.5 x10⁻⁵) implies that the parent body was impacted when at or near its peak temperature (Extended Data Fig. 3). However, this scenario does not agree with the later resetting ages of silicate inclusions discussed above²⁰⁻²² because the closure temperature of the Hf-W system

($~1150~K^{24}$) is higher than that of the Pd-Ag system ($~820$ -970 K 6). Hence, a thermal event that affected the Hf-W chronometer should also reset the Pd-Ag system. This discrepancy is unlikely to represent disturbance to the Hf-W chronometer. Rapid cooling precludes significant diffusive exchange between silicate inclusions and host IAB metals²². Although GCR can affect Hf-W ages, this is unlikely to be the case here because Campo del Cielo irons experienced only limited GCR exposure (this study, 25). A model incorporating cooling at 12.8 $^{+3.1}/_{4.6}$ Myr is therefore most appropriate for the IAB parent body evolution.

Implications for the Solar System initial 107Pd/108Pd

Importantly, the observation that the Hf-W and Pd-Ag systems cannot be reconciled using a SSI $^{107}Pd^{108}Pd$ of \sim 3.5 x10⁻⁵ is robust regardless of the formation history of the IAB parent body. Campo del Cielo is a member of the IAB-MG, which crystallised, and hence cooled, as a single body¹⁷. An isochron using only the IAB-MG yields an initial $107Pd/108Pd$ of (1.61 ± 0.23) x10⁻⁵, identical within uncertainty to the IAB isochron with all samples. Using a SSI $107Pd/108Pd$ of (3.5±0.1) x10⁻⁵ implies closure of the Pd-Ag system for the MG at 7.3^{+1.3}/-1.5 Myr after CAI. Irrespective of whether this defines metal cooling after a local impact or a body-wide catastrophic event, the irons should then not include silicates with a Hf-W closure age of 11.4 \pm 2.2 Myr^{22,23}. Therefore, we suggest that the SSI ¹⁰⁷Pd/¹⁰⁸Pd ratio of ~3.5 x10⁻⁵ is too low and we focus mainly on the SSI ratio derived from carbonaceous chondrites in the following discussion.

Evolution of iron meteorite parent bodies

The Pd-Ag ages obtained here of 11.7 $+3.6/$ _{-5.9} Myr after CAI for the IIABs, and 9.2 $+3.5$ /. $_{5.7}$ Myr for the IIIABs, suggest these bodies cooled in the same time period (Table 3; Fig. 1). The IVAs also yield a similar cooling age $(7.8 \text{ }^{+3.0}/4.4, ^{-12})$. The timings of metal-silicate differentiation and core cooling of an undisrupted asteroid are primarily a function of size and time of accretion²⁶. The comparable accretion and core crystallisation ages¹ and cooling rates (Table 3; Methods) of these bodies may therefore imply that they had similar sizes. However,

high and variable metallographic cooling rates among the IIAB, IIIAB and IVA irons indicate that these metals cooled in bodies that had been stripped of their insulating silicate mantles in 'hit-and-run' style impacts4,5. Intriguingly, we find that the IIAB, IIIAB and IVA asteroids were impacted in broadly the same timeframe (Fig. 1), suggesting an energetic Solar System environment persisted at this time.

Early Solar System dynamics

Nucleosynthetic isotope variations indicate a dichotomy between non-carbonaceous (NC) and carbonaceous-type (CC) asteroids²⁷. The IIAB, IIIAB and IVA asteroids are NC and likely accreted in the inner Solar System¹. All three indicate impact-related cooling and this provides evidence for energetic events with the ability to affect multiple locations in the inner disk at \sim 7.8-11.7 Myr after CAI. Additional evidence for an energetic state in the NC reservoir²⁵ includes the IAB impact event identified above that occurred in a similar timeframe, albeit 5.0 $^{+1.0}$ /_{-1.1} Myr later than on the IVA body (Table 3). Other early-formed asteroids, including the Hchondrite parent body, the ungrouped Nedagolla iron meteorite, and the IID and IVB irons which originated in the outer disk (CC) reservoir, may also show evidence of impact processes in this interval^{5,11,28-30}. Below, we discuss mechanisms that can lead to an energetic inner disk environment between ~7.8-11.7 Myr after CAI.

The presence and subsequent dissipation of the gas disk exerts one of the strongest controls on early Solar System orbital dynamics. The presence of gas in a protoplanetary disk dampens the eccentricities of planetesimals and stabilises the system. As the gas disperses, the damping effect of gas drag ceases and energetic collisions capable of disrupting planetesimals are predicted³¹⁻³³ (Fig. 2). Imaging of young stellar clusters suggest the overall lifetime of circumstellar disks is \sim 6–8 Myr³⁴, and the gas in our protoplanetary disk likely survived no more than \sim 4 Myr after CAI¹. Whilst this is earlier than the impact timeframe discussed here, models of planetesimal dynamics after gas dispersion show that bodies initially have dynamically stable orbits, with eccentricity and inclination building up over several Myr^{31,33}. Therefore, impacts occurring between \sim 7.8 and 11.7 Myr may be attributable to the dissipation of the gas disk. The first ~10-20 Myr after gas dissipation are the most important for both the number and mean velocity of collisions 31 . Further, this mechanism generates a chaotic asteroid belt for several Myr and thereby also provides a viable explanation for later impacts on the IAB parent body. The disk-wide nature of this mechanism also provides an explanation for impacts in the CC reservoir.

Giant planets also act in multiple ways to influence the orbits of nearby planetesimals. Rapid runaway growth is predicted to perturb the orbits of nearby planetesimals to high eccentricities and gravitationally scatter them into the inner Solar System, causing widespread mixing with the potential for more frequent collisions^{35,36}. Additionally, Jupiter and Saturn are suggested to have undergone rapid gas-driven in- and then outward migration (the 'Grand Tack' model), leading to excitation of the asteroid belt and high velocity impacts in the inner Solar System^{37,38}. The formation times of chondrules and their distinct nucleosynthetic fingerprints suggest the NC and CC reservoirs remained isolated until ~3-5 Myr after CAI, constraining the earliest time of giant planet migration^{1,37}. However, impact events between ~7.8-11.7 Myr are more than ~3 Myr later than the runaway growth and hypothesised migration phase of the giant planets, and therefore cannot be attributed to these events. Additionally, these mechanisms are not expected to lead to bombardment events that lasted longer than \sim 0.5-1 Myr^{37,39,40}, whereas the energetic state of the inner Solar System recorded by the Pd-Ag system persisted for at least 3.9 $+2.3/3.1$ Myr after the closure of the IVA core (Table 3). This longevity problem persists if using the lower SSI 107 Pd/ 108 Pd ratio of ~3.5 x10⁻⁵, which yields impacts at ~2.9-6.8 Myr after CAI that broadly coincide with the migration of the giant planets. In this case, a second mechanism such as runaway growth must also be invoked. Further, these two events together do not provide a framework for understanding the later impact recorded by the IABs.

Additionally, a giant planet instability (the 'Nice' model) could provide the necessary energy to the asteroid belt and inner Solar System (Fig. 2, Scenario B). This model can explain many features of our Solar System² and was initially assumed to occur later in Solar System history thereby providing an explanation for the Late Heavy Bombardment (LHB)⁴¹. However,

recent studies have questioned both the evidence for a LHB and the dynamic consequences of delaying the instability⁴²⁻⁴⁴. An early giant planet instability occurring ≤100 Myr after gas dissipation better matches the architecture of our Solar System^{42,45-48}. Although the precise timing of the instability is difficult to determine, models which predict the small size of Mars advocate an instability \sim 5–14 Myr after CAI $47,48$. Impacts on the iron parent bodies between ~7.8-11.7 Myr agree well with such a scenario and may provide the first evidence from the meteorite record for the occurrence and timing of an early giant planet instability. Such longlived and Solar System wide events also provide an explanation for later impacts including on the IAB parent body, and impacts across the disk, including both the NC and CC reservoirs. Additionally, the effect of gas dissipation may act as the trigger for the early giant planet instability⁴⁹. Hence, these two mechanisms may have acted together to create an energetic Solar System between ~7.8-11.7 Myr after CAI.

Methods

Eighteen aliquots of thirteen iron meteorites from the IAB, IIAB and IIIAB groups were analysed for this study (Table 1). Samples were cut using a CBN blade and subsequently polished using SiC paper to remove remaining weathered areas and fusion crust. Any visible troilite inclusions were physically removed from the metals. Palladium-silver isotope data for the IAB irons were published previously in 9 , and Pt isotope data for Coahuila are from 15 . Material for Pd-Ag and Pt isotope analyses was taken from adjacent sampling locations, except for Negrillos, Caddo County and Canyon Diablo. Platinum isotope data for these samples are taken from the literature and indicate that they are unexposed or only weakly exposed to the effects of GCR (Table 1, 14,16).

Platinum isotope measurements

Platinum isotope ratios were determined according to the methods described by ⁵⁰. In brief, samples were placed in ethanol in an ultrasonic bath for 5 minutes and then leached for 5 minutes in cold 2 M HCl before digestion with a 2:1 mix of concentrated $HNO₃$ and HCl (~8 ml g^{-1} of sample). Samples were refluxed and dried, and then taken up in concentrated HCl. At this stage, 0.3 g aliquots for Pt isotope analyses were taken and dried. The aliquots were refluxed again with a 2:1 mix of concentrated $HNO₃$ and HCI, and then re-dissolved in preparation for the anion exchange procedure. This consisted of two stages and utilised Biorad AG® 1-X8 resin (200 – 400 mesh, chloride form). The first stage separated Pt from major matrix constituents, while the second stage separated Pt from Ir⁵⁰. The second stage was repeated until samples had ¹⁹¹Ir/¹⁹⁵Pt ratios \leq 0.13, which is below the tolerance limit for Ir isotope tailing onto Pt isotopes during mass spectrometry, as determined for the instrument housed at ETH Zürich⁵⁰. Osmium, which generates isobaric interferences on Pt isotopes, was then volatilised using perchloric acid.

Platinum isotope ratios were measured at ETH Zürich using a Thermo Scientific Neptune *Plus* operated in low-resolution mode and fitted with a Cetac Aridus II desolvating nebuliser and standard H cones, following the procedure of ⁵⁰. Samples were diluted to yield an ion beam intensity of \sim 2 x10⁻¹⁰ A for ¹⁹⁴Pt, which equates to \sim 150 ng g⁻¹ Pt and required ~170 ng per analysis. Analyses were corrected for instrumental mass bias using the exponential law and were internally normalized to $^{198}Pt/^{195}Pt$ ('8/5') = 0.2145 ¹⁶. Sample measurements were bracketed by the NIST SRM 3140 Pt standard solution and data are presented in the epsilon notation (i.e., $\epsilon^{19i}Pt^{195}Pt =$ deviation in parts per 10,000 from the average of the two bracketing standards). Reproducibility for Pt isotope analyses was assessed using the iron meteorite North Chile as an in-house standard. Repeat analyses of multiple aliquots of this material indicate a 2 S.D. uncertainty of 0.73 for ϵ^{192} Pt, 0.15 for ϵ^{194} Pt and 0.09 for ε^{196} Pt (n=19; ⁵⁰). Platinum blanks are in the range 0.3 – 1.0 ng g⁻¹ sample.

Pd-Ag Measurements

Sample masses for Pd-Ag analyses are given in Supplementary Materials Table 1. Samples for Pd-Ag analyses were leached in cold concentrated aqua regia for 5 minutes, followed by digestion in concentrated aqua regia (\sim 8 ml g -1) on a hotplate. They were then dried and refluxed in concentrated HCl (40 ml), before adding $20 - 40$ ml H₂O to verify that the sample was fully in solution. Approximately 9% of the sample solution was taken at this stage to determine Pd and Ag concentrations by isotope dilution, following an established method⁸. Samples were then refluxed and dried again in HCl to ensure no traces of $HNO₃$ remained. The ion exchange technique for Ag purification followed a published protocol⁵¹ and utilised a three-stage column procedure with anion and cation exchange resin. The chemical procedure removes elements that cause interferences on Ag isotopes, including isobaric (Pd, Cd and Ru) and molecular interferences (Cu, Zn, Ga, Ge, Sr, Y, Zr, Nb and Mo), such that all were below the tolerance limits for accurate Ag isotope analyses $9,51$. The yields of the chemical separation procedure were checked for each sample individually and they were always close to 100%.

Silver isotope analyses were performed by MC-ICP-MS on either a Nu Plasma instrument in the School of Earth, Atmospheric and Environmental Science at The University of Manchester or a VG Axiom in the Department of Terrestrial Magnetism at the Carnegie Institution of Washington (Table 1). The details are described in 9 and 51 , respectively, and a short summary is given here. Both instruments were operated with a glass cyclonic spray chamber. A Pd NIST SRM 3138 standard solution was added to all samples and standards just prior to analysis to correct for instrumental mass bias⁵¹, and was mixed to yield a signal intensity ratio of 1 for $107\text{Ag}/108\text{Pd}$. Additionally, sample measurements were bracketed by the NIST SRM 978a Ag - NIST SRM 3138 Pd standard solution mixture, with the Pd and Ag concentrations of the standard matched to the samples to within 20% or better. Instrumental mass bias was corrected using the exponential law with normalisation to a $^{108}Pd^{105}Pd$ value of 1.18899 ⁷. Data are presented in the epsilon notation and a 107 Ag/ 109 Ag ratio of 1.079760 13 was used for the NIST SRM 978a Ag standard.

For analyses conducted at The University of Manchester, the typical daily reproducibility of a 100 ppb Ag standard solution for the 107 Ag/ 109 Ag ratio was ~0.25 ε (2 S.D. 9 . Additionally, the external reproducibility of sample measurements was assessed by repeatedly passing the NIST SRM 978a Ag standard, the USGS standard Cody Shale SCo-1, and the Hawaiian basalt KOO49 through the chemical procedure. These materials yielded external reproducibilities of between 0.1 and 0.6 ε (2 S.D.⁹), and therefore a conservative external uncertainty of ± 0.6 ε has been applied to all sample data acquired at The University of Manchester. Similar tests were carried out at the Carnegie Institution⁵¹ and resulted in an external uncertainty of ±0.5 ε . Silver blanks for sample digestion and column chemistry were <0.254 ng.

Correction for Galactic Cosmic Ray Effects to Ag isotopes

New ε¹⁰⁷Ag data for the IIAB and IIIAB irons vary from ~37 to 103, while ¹⁰⁸Pd/¹⁰⁹Ag ratios range between 261 and 679 (Table 1). There is a positive correlation between ϵ^{107} Ag and ¹⁰⁸Pd/¹⁰⁹Ag for all meteorite groups, however, a regression cannot be fitted through our new data for the IIAB or IIIAB iron meteorites before corrections for GCR (Extended Data Fig. 2). The uncorrected IAB samples define an initial $^{107}Pd/^{108}Pd$ ratio of 1.06 (\pm 0.36) x10⁻⁵ with a MSWD of 45 9 .

Galactic cosmic rays (GCR) can disturb the 107 Pd- 107 Ag chronometer through reactions with thermal neutrons¹⁰. This must be corrected before meaningful age information can be obtained. Secondary neutron capture by Pd isotopes leads to a decrease in 107 Ag/ 109 Ag primarily via neutron capture on ¹⁰⁸Pd and subsequent β ^{-decay} to ¹⁰⁹Ag. Additionally, ¹⁰⁶Pd is also susceptible to neutron capture followed by β ^{-decay} to ¹⁰⁷Ag. The GCR-induced shift in 107 Ag/ 109 Ag is therefore linearly correlated with Pd/Ag and higher 108 Pd/ 109 Ag ratios lead to larger negative shifts in ¹⁰⁷Ag/¹⁰⁹Ag, i.e., the slope of data not corrected for GCR effects would be too shallow, if one could even be determined because of the scatter¹⁰. Longer exposure ages, bigger pre-atmospheric radii, and increasing shielding depths all lead to larger GCR offsets.

It has been demonstrated that Pt isotopes are a powerful neutron dosimeter for correcting the effects of GCR in short-lived decay systems, e.g., ¹⁸²Hf-¹⁸²W ¹⁴. They have also been applied as a dosimeter for neutron capture in the 107 Pd- 107 Ag system 11,12 . New Pt isotope data fit well to the modelled effects for neutron capture (Extended Data Fig. 1). Model trendlines are shown for Ir/Pt ratios appropriate for our samples (Supplementary Materials Table 1). These results further confirm that neutron capture reactions are the cause of Pt isotope excesses and that there are no detectable nucleosynthetic isotope variations between iron meteorites.

Following the method of 11 (given as equations 1-3 below), data for the IAB, IIAB and IIIAB irons were assessed for the effects of neutron capture. The correction relies on the ¹⁰⁸Pd/¹⁰⁹Ag ratio and neutron dose (ε¹⁹⁶Pt) of the sample in question. Measured ε¹⁹⁶Pt can be related to neutron capture corrections to ϵ^{107} Ag (ϵ^{107} Ag_{GCR}) by a linear regression:

$$
\epsilon^{107} \text{Ag}_{\text{GCR}} = \text{Offset} + \text{Slope} \times \epsilon^{196} \text{Pt} \tag{1}
$$

where *Offset* and *Slope* are functions of the measured ¹⁰⁸Pd/¹⁰⁹Ag ratio¹¹. The *Offset* defines neutron capture shifts on Ag isotopes and is described by:

$$
Offset = -0.023 + 0.0024 \times \left(\frac{108}{109} \text{G} \right) - 4.76 \times 10^{-7} \times \left(\frac{108}{109} \text{G} \right)^2 \tag{2}
$$

while the *Slope* is defined by:

Slope =
$$
1.539 - 0.226 \times \frac{108}{109} \text{Gy}
$$
 (3)

The *Offset and Slope* are then used in Eq. (1) to determine ε¹⁰⁷Ag_{GCR}. For the samples included in this study ϵ^{107} Ag_{GCR} ranges from 2.3 to -49.2, with the largest offsets recorded in the IIIAB irons (Table 2).

Solar System Initial ¹⁰⁷Pd/¹⁰⁸Pd

In order to accurately determine Pd-Ag ages, the Solar System initial (SSI) ¹⁰⁷Pd/¹⁰⁸Pd is needed. Recent estimates are based on either carbonaceous chondrites¹³ or the IVA iron Muonionalusta^{11,12}, which has resulted in a range of initial ratios. By equating the ⁵³Mn-⁵³Cr and ¹⁰⁷Pd-¹⁰⁷Ag decay systems, which both date volatile depletion, carbonaceous chondrites define a SSI $^{107}Pd^{108}Pd$ of (5.9±2.2) x10^{-5 13}. This ratio is among the highest initials yet determined and includes a relatively large uncertainty. Values for the initial ¹⁰⁷Pd/¹⁰⁸Pd based on Muonionalusta range from \sim (2.8–6.6) x10^{-5 11,12}, but the use of this meteorite to define a SSI ¹⁰⁷Pd/¹⁰⁸Pd is hampered by ambiguity over its absolute Pb-Pb age, determined from a troilite⁵². Recently, the Pd-Ag systematics of Muonionalusta were re-examined and two possible SSI $^{107}Pd/^{108}Pd$ ratios of (3.5±0.1) and (6.6±0.4) x10⁻⁵ were determined¹², dependent on the 238U/235U ratio of the Muonionalusta troilite. In order to consider the full range of SSI values, we calculate Pd-Ag ages for the iron meteorite groups studied here relative to both the lowest recent estimate based on Muonionalusta¹² (~3.5 x10⁻⁵) and the value indicated from carbonaceous chondrites¹³ (~5.9 x10⁻⁵), which also encompasses the highest ratio estimated from Muonionalusta. To circumvent the ambiguity over the SSI ¹⁰⁷Pd/¹⁰⁸Pd, closure times can also be calculated relative to the value determined for the IVA irons (107 Pd/ 108 Pd ~2.57 x10⁻⁵, $12:$ Table 3).

Calculation of cooling rates

The closure temperature of the Pd-Ag system is dependent on cooling rate, Ni and FeS content of the meteorite, and is estimated to be in the range ~820 - 970 K for the iron meteorites studied here 6 . Fast cooling rates are necessary for the IAB parent body in order to keep silicate inclusions suspended in the metal portions¹⁷. Assuming that the IAB parent was at its peak temperature of 1470 K¹⁴ when impacted at \sim 11 Myr (see Main Text) yields the expected fast cooling rates of between instantaneous and ~102 K/Myr (Table 3). Alternatively, a cooling age of 7.9^{+1.0}/-_{1.1} (calculated relative to an initial SSI ¹⁰⁷Pd/¹⁰⁸Pd of ~3.5 x 10^{-5 12}) implies that the parent body was impacted when at or near its peak temperature at 6.0 ± 0.8 Myr. This yields cooling rates of between instantaneous and ~135 K/Myr.

Modelling indicates the IIAB and IIIAB parent bodies accreted \leq ~0.4 Myr after CAI¹. Core formation on the IIAB parent body followed at 0.7 ± 0.3 Myr, based on the Hf-W chronometer¹⁶. Assuming a liquidus temperature of \sim 1570 K for the IIABs⁵³ and a steady rate of cooling between core formation and closure of the Pd-Ag chronometer indicates cooling rates of ~278-69 K/Myr for cooling at 6.8 $^{+2.3}$ / $_{3.1}$ Myr, or ~156-40 K/Myr for cooling at 11.7 $^{+3.6}$ $/$ -5.9 Myr (Table 3). Metallographic cooling rates for the IIABs suggest a much slower process (0.8-10 K/Myr⁵⁴). However, it is important to note that these apply to cooling at ~670-770 K, and that newer techniques have led to upward revisions of rates by factors of 10-50 5 .

The IIIABs experienced core formation at 1.2 \pm 0.3 Myr with a predicted liquidus temperature of \sim 1650 K^{16,53}. This yields a very wide range of cooling rates, from instantaneous-121 K/Myr for Pd-Ag closure at 4.3 $+2.2$ / $_{-2.8}$ Myr to 415-58 K/Myr for closure at 9.2 $+3.5$ / $_{-5.7}$ Myr (Table 3). Cooling rates determined here for the IIIAB irons compare well with those derived previously using the Pd-Ag system $6,11$. They also overlap with recent estimates of metallographic cooling rates obtained for this body (338 – 56 K/Myr ⁵), particularly for a Pd-Ag closure time of ~9.2 Myr after CAI.

Acknowledgements

This work was supported by the European Research Council under the European Union's Seventh Framework Programme (FP7/2007–2013/ERC Grant agreement n˚ [279779], MS). We gratefully acknowledge funding from STFC (ST/F002157/1 and ST/J001260/1, MR; ST/J001643/1, MS) and funding from the Swiss National Science Foundation (Project 200020 179129, MS). AH wishes to thank Mattias Ek and Manuela Fehr for laboratory assistance at ETH Zürich during this study. MS would like to thank Rick Carlson and Mary Horan (DTM, Carnegie Institution) for their support and the opportunity to analyse Ag isotopes in May 2006. We also thank Caroline Smith and Deborah Cassey (Natural History Museum, London), and Julie Hoskin (Smithsonian Institution National Museum of Natural History) for the loan of samples used in this work.

Author Contributions

M.S. designed the study. A.H., K.T. and M.S. prepared samples for isotope analyses and conducted the isotopic measurements. All authors were involved in the data interpretation and writing of the manuscript.

Competing Interests Statement

The authors declare no competing interests.

Supplementary Information is available for this paper in the form of one Table as a pdf.

Data Availability

All data are available within this manuscript and its Supplementary information, or available from the authors on request.

TABLES, MAIN ARTICLE

Table 1. Concentration data for Pd and Ag, isotope data for Ag, and ε¹⁹⁶Pt for IAB, IIAB and

IIIAB iron meteorites.

Uncertainties for ε^{107} Ag are based on repeat analyses of reference materials (see Methods). Uncertainties (2 S. D.) on ¹⁰⁸Pd/¹⁰⁹Ag ratios are estimated to be ~1 % for data collected at the University of Manchester and 5 % for data collected at the Carnegie Institution of Washington. Platinum isotope uncertainties are based on repeat analyses of an in-house standard (see Methods). a IAB sub-groups from ¹⁷. See main text for definitions. ^b Terrestrial ¹⁰⁷Ag/¹⁰⁹Ag ratio from ¹³. ^c Palladium-silver data taken from ⁹. All Pd-Ag isotope data measured at the University of Manchester, except ^d Carnegie Institution of Washington. Platinum isotope data taken from ^e Hunt, et al. ¹⁴, ^f Hunt, et al. ¹⁵ and ^g Kruijer, et al. ¹⁶.

Meteorite	$\overline{\epsilon^{107}}$ Ag _{meas}	$±2$ S.D.	ϵ^{107} Aggcr	$± 2$ S.D.	$\overline{\epsilon^{107}}$ Ag _{pre-exp}	$± 2$ S.D.
IAB						
Campo del Cielo	15.8	0.6	-3.1	3.5	18.9	3.5
Canyon Diablo	1.4	0.6	0.1	1.0	1.3	1.1
Odessa 1	2.2	0.6	-5.0	1.5	7.2	1.6
Odessa 2	11.7	0.6	-11.9	3.9	23.6	4.0
Goose Lake 1	7.4	0.6	-1.6	2.1	9.0	2.2
Goose Lake 2	8.6	0.6	-1.7	2.2	10.3	2.3
Toluca	1.4	0.6	-1.6	1.2	3.0	1.3
Caddo County	0.1	0.6	0.1	0.6	0.0	0.9
IIAB						
Coahuila 1	53.4	0.6	-0.2	4.8	53.7	4.8
Coahuila 2	50.6	0.5	-0.2	3.9	50.8	3.9
Negrillos	52.7	0.5	2.3	2.9	50.4	2.9
North Chile 1	55.0	0.6	-8.2	5.7	63.2	5.7
North Chile 2	53.1	0.5	-8.2	5.6	61.2	5.6
Sikhote Alin 1	59.9	0.6	-20.7	8.7	80.6	8.7
Sikhote Alin 2	36.9	0.6	-14.7	6.2	51.6	6.2
IIIAB						
Boxhole	103.5	0.6	-37.1	13.7	140.6	13.7
Henbury	77.7	0.6	-49.2	11.9	126.8	11.9
Thunda	83.3	0.6	-3.7	8.2	87.0	8.2

Table 2. Correction to ε¹⁰⁷Ag for GCR effects for IAB, IIAB and IIIAB iron meteorites.

Table 3. Parent body initial ¹⁰⁷Pd/¹⁰⁸Pd ratios and closure times of the Pd-Ag system relative to CAI and IVA irons.

^a Time after CAI and resultant cooling rates calculated using a SSI 107 Pd/ 108 Pd of 5.9 (\pm 2.2) x 10^{-5 13}. b Time after CAI calculated using a SSI ¹⁰⁷Pd/¹⁰⁸Pd of 3.5 (\pm 0.1) x 10^{-5 12}. See Extended Methods for calculation of cooling rates. ^c Initial ¹⁰⁷Pd/¹⁰⁸Pd and cooling rate for the IVA iron meteorites from ¹².

FIGURE LEGENDS, MAIN ARTICLE

Figure 1. The timing of core cooling and impacts constrained by the Pd-Ag system. a. Parent body initial ¹⁰⁷Pd/¹⁰⁸Pd ratios for the IAB, IIAB and IIIAB iron meteorite groups (Table 3). Literature data for the IVA irons¹² are also shown. b. Age after CAI for core cooling, calculated relative to the carbonaceous chondrite SSI^{13} . Iron meteorite group symbols as in panel (a). The grey and purple bands represent, respectively, the approximate timings for CAI formation and the impacts recorded by the Pd-Ag system. The timeframe suggested for the 'Grand Tack' is indicated by the size of the Jupiter symbol $1,37$. The intervals suggested for gas present in the disk, and an early giant planet instability that predicts a small Mars^{47,48}, are also shown.

Figure 2. Cartoon illustrating the evolution of differentiated iron meteorite bodies in the early Solar System (not to scale). Top panel: Parent bodies accrete and differentiate within the first \sim 3 Myr after CAI¹⁶. The disk consists of gas and dust (coloured blue) and is truncated in the region of Jupiter, preventing significant exchange between the non-carbonaceous (NC) and carbonaceous (CC) reservoirs. Later, the cores of the IIAB, IIIAB and IVA bodies were exposed by impact events, leading to fast cooling and closure of the Pd-Ag system between ~7.8 - 11.7 Myr. Two scenarios, acting alone or in combination, provide mechanisms for an energetic inner Solar System. Scenario A: When the gas disk dissipates, planetesimals are gradually excited until they begin to undergo high velocity impacts 31 . Gas dissipation is completed before this time. Scenario B: An early giant planet instability is triggered at ~7.8 - 11.7 Myr, leading to reorganisation of the inner Solar System and impact events⁴⁷.

EXTENDED DATA FIGURE LEGENDS

Extended Data Figure 1. Covariation of ε¹⁹²Pt and ε¹⁹⁶Pt for the IAB, IIAB and IIIAB iron meteorites. The GCR model calculations¹⁰ are shown for Ir/Pt ratios measured for these groups (dashed lines; Ir/Pt ratios for individual samples are given in Supplementary Materials Table 1). Uncertainties are 2 S.D.

Extended Data Figure 2. Isochron diagrams for a) IAB, b) IIAB, and c) IIIAB iron meteorites for the Pd-Ag system. In each case, ϵ^{107} Ag is plotted against 108 Pd/ 109 Ag for both GCRcorrected (filled symbols) and GCR-uncorrected data (unfilled symbols). Uncertainties on GCR-uncorrected data are 2 S.D. and within the size of the symbol (Table 1), while uncertainties for ε^{107} Ag for GCR-corrected data represent the propagated 2 S.D. uncertainties of the GCR correction (Table 2). Isochrons are determined using GCRcorrected data and ISOPLOT⁵⁵. *denotes GCR-corrected data from source¹¹, shown for comparison but not included in the regressions.

Extended Data Figure 3. IAB parent body evolution for both metal cooling times. In both scenarios the parent body accreted relatively late at ~1.4 Myr and then underwent limited metal-silicate differentiation at ~ 6.0 Myr after CAI¹⁴. For metal cooling at ~ 12.8 Myr, the catastrophic impact event occurred in the timeframe ~11 – 13.6 Myr. This was followed by fast cooling and closure of the Pd-Ag system at \sim 12.8 ^{+3.1}/_{-4.6} Myr. For metal cooling at \sim 7.9 Myr after CAI the body was disrupted at about 6 Myr, while at or near its peak temperature. It then cooled quickly, leading to closure of the Pd-Ag system at 7.9 ^{+1.0}/-_{1.1} Myr.

References

- 1 Kruijer, T. S., Burkhardt, C., Budde, G. & Kleine, T. Age of Jupiter inferred from the distinct genetics and formation times of meteorites. *Proc. Natl. Acad. Sci. U.S.A.* **114**, 6712-6716 (2017).
- 2 Tsiganis, K., Gomes, R., Morbidelli, A. & Levison, H. F. Origin of the orbital architecture of the giant planets of the Solar System. *Nature* **435**, 459-461 (2005).
- 3 Walsh, K. J. & Morbidelli, A. The effect of an early planetesimal-driven migration of the giant planets on terrestrial planet formation. *Astron. Astrophys.* **526**, A126 (2011).
- 4 Yang, J., Goldstein, J. I. & Scott, E. R. D. Iron meteorite evidence for early formation and catastrophic disruption of protoplanets. *Nature* **446**, 888 (2007).
- 5 Goldstein, J. I., Scott, E. R. D. & Chabot, N. L. Iron meteorites: Crystallization, thermal history, parent bodies, and origin. *Chemie der Erde - Geochemistry* **69**, 293- 325 (2009).
- 6 Matthes, M., van Orman, J. A. & Kleine, T. Closure temperature of the Pd-Ag system and the crystallization and cooling history of IIIAB iron meteorites. *Geochim. Cosmochim. Acta* **285**, 193-206 (2020).
- 7 Kelly, W. R. & Wasserburg, G. J. Evidence for the existence of ¹⁰⁷Pd in the early solar system. *Geophys. Res. Lett.* **5**, 1079-1082 (1978).
- 8 Woodland, S. J. *et al.* Accurate measurement of silver isotopic compositions in geological materials including low Pd/Ag meteorites. *Geochim. Cosmochim. Acta* **69**, 2153-2163 (2005).
- 9 Theis, K. J. *et al.* Palladium–silver chronology of IAB iron meteorites. *Earth Planet. Sci. Lett.* **361**, 402-411 (2013).
- 10 Leya, I. & Masarik, J. Thermal neutron capture effects in radioactive and stable nuclide systems. *Meteorit. Planet. Sci.* **48**, 665-685 (2013).
- 11 Matthes, M., Fischer-Gödde, M., Kruijer, T. S., Leya, I. & Kleine, T. Pd–Ag chronometry of iron meteorites: Correction of neutron capture-effects and application to the cooling history of differentiated protoplanets. *Geochim. Cosmochim. Acta* **169**, 45-62 (2015).
- 12 Matthes, M., Fischer-Gödde, M., Kruijer, T. S. & Kleine, T. Pd-Ag chronometry of IVA iron meteorites and the crystallization and cooling of a protoplanetary core. *Geochim. Cosmochim. Acta* **220**, 82-95 (2018).
- 13 Schönbächler, M., Carlson, R. W., Horan, M. F., Mock, T. D. & Hauri, E. H. Silver isotope variations in chondrites: Volatile depletion and the initial 107Pd abundance of the solar system. *Geochim. Cosmochim. Acta* **72**, 5330-5341 (2008).
- 14 Hunt, A. C. *et al.* Late metal–silicate separation on the IAB parent asteroid: Constraints from combined W and Pt isotopes and thermal modelling. *Earth Planet. Sci. Lett.* **482**, 490-500 (2018).
- 15 Hunt, A. C., Ek, M. & Schönbächler, M. Platinum isotopes in iron meteorites: Galactic cosmic ray effects and nucleosynthetic homogeneity in the *p-*process isotope 190Pt and the other platinum isotopes. *Geochim. Cosmochim. Acta* **216**, 82-95 (2017).
- 16 Kruijer, T. S. *et al.* Protracted core formation and rapid accretion of protoplanets. *Science* **344**, 1150-1154 (2014).
- 17 Wasson, J. T. & Kallemeyn, G. W. The IAB iron-meteorite complex: A group, five subgroups, numerous grouplets, closely related, mainly formed by crystal segregation in rapidly cooling melts. *Geochim. Cosmochim. Acta* **66**, 2445-2473 (2002).
- 18 Benedix, G. K., McCoy, T. J., Keil, K. & Love, S. G. A petrologic study of the IAB iron meteorites: Constraints on the formation of the IAB-Winonaite parent body. *Meteorit. Planet. Sci.* **35**, 1127-1141 (2000).
- 19 Hilton, C. D. & Walker, R. J. New implications for the origin of the IAB main group iron meteorites and the isotopic evolution of the noncarbonaceous (NC) reservoir. *Earth Planet. Sci. Lett.* **540**, 116248 (2020).
- 20 Pravdivtseva, O., Meshik, A., Hohenberg, C. M. & Kurat, G. I-Xe ages of Campo del Cielo silicates as a record of the complex early history of the IAB parent body. *Meteorit. Planet. Sci.* **48**, 2480-2490 (2013).
- 21 Schulz, T., Munker, C., Mezger, K. & Palme, H. Hf-W chronometry of primitive achondrites. *Geochim. Cosmochim. Acta* **74**, 1706-1718 (2010).
- 22 Schulz, T., Münker, C., Palme, H. & Mezger, K. Hf–W chronometry of the IAB iron meteorite parent body. *Earth Planet. Sci. Lett.* **280**, 185-193 (2009).
- 23 Kruijer, T. S., Kleine, T., Fischer-Gödde, M., Burkhardt, C. & Wieler, R. Nucleosynthetic W isotope anomalies and the Hf–W chronometry of Ca–Al-rich inclusions. *Earth Planet. Sci. Lett.* **403**, 317-327 (2014).
- 24 Kleine, T. *et al.* Hf–W thermochronometry: Closure temperature and constraints on the accretion and cooling history of the H chondrite parent body. *Earth Planet. Sci. Lett.* **270**, 106-118 (2008).
- 25 Worsham, E. A., Bermingham, K. R. & Walker, R. J. Characterizing cosmochemical materials with genetic affinities to the Earth: Genetic and chronological diversity within the IAB iron meteorite complex. *Earth Planet. Sci. Lett.* **467**, 157-166 (2017).
- 26 Lichtenberg, T., Golabek, G. J., Gerya, T. V. & Meyer, M. R. The effects of short-lived radionuclides and porosity on the early thermo-mechanical evolution of planetesimals. *Icarus* **274**, 350-365 (2016).
- 27 Trinquier, A., Birck, J.-L. & Allègre, C. Widespread 54Cr Heterogeneity in the Inner Solar System. *Astrophys. J.* **655**, 1179 (2007).
- 28 Wasson, J. T. & Huber, H. Compositional trends among IID irons; their possible formation from the P-rich lower magma in a two-layer core. *Geochim. Cosmochim. Acta* **70**, 6153-6167 (2006).
- 29 Spitzer, F., Burkhardt, C., Pape, J. & Kleine, T. Collisional mixing between inner and outer solar system planetesimals inferred from the Nedagolla iron meteorite. *Meteorit. Planet. Sci.* **57,** 261-276 (2022).
- 30 Hellmann, J. L., Kruijer, T. S., Van Orman, J. A., Metzler, K. & Kleine, T. Hf-W chronology of ordinary chondrites. *Geochim. Cosmochim. Acta* **258**, 290-309 (2019).
- 31 Davison, T. M., O'Brien, D. P., Ciesla, F. J. & Collins, G. S. The early impact histories of meteorite parent bodies. *Meteorit. Planet. Sci.* **48**, 1894-1918 (2013).
- 32 Morbidelli, A., Lunine, J. I., O'Brien, D. P., Raymond, S. N. & Walsh, K. J. Building Terrestrial Planets. *Annu. Rev. Earth Planet. Sci.* **40**, 251-275 (2012).
- 33 Walsh, K. J. & Levison, H. F. Planetesimals to terrestrial planets: Collisional evolution amidst a dissipating gas disk. *Icarus* **329**, 88-100 (2019).
- 34 Haisch, K. E., Lada, E. A. & Lada, C. J. Disk Frequencies and Lifetimes in Young Clusters. *Astrophys. J. Lett.* **553**, L153 (2001).
- 35 Ikoma, M., Nakazawa, K. & Emori, H. Formation of Giant Planets: Dependences on Core Accretion Rate and Grain Opacity. *Astrophys. J.* **537**, 1013-1025 (2000).
- 36 Raymond, S. N. & Izidoro, A. Origin of water in the inner Solar System: Planetesimals scattered inward during Jupiter and Saturn's rapid gas accretion. *Icarus* **297**, 134-148 (2017).
- 37 Johnson, B. C., Walsh, K. J., Minton, D. A., Krot, A. N. & Levison, H. F. Timing of the formation and migration of giant planets as constrained by CB chondrites. *Science Advances* **2**, e1601658 (2016).
- 38 Walsh, K. J., Morbidelli, A., Raymond, S. N., O'Brien, D. P. & Mandell, A. M. A low mass for Mars from Jupiter's early gas-driven migration. *Nature* **475**, 206 (2011).
- 39 Turrini, D., Magni, G. & Coradini, A. Probing the history of Solar system through the cratering records on Vesta and Ceres. *Mon. Not. R. Astron. Soc.* **413**, 2439-2466 (2011).
- 40 Turrini, D., Coradini, A. & Magni, G. Jovian early bombardment: Planetesimal erosion in the inner asteroid belt. *Astrophys. J.* **750**, 8 (2012).
- 41 Gomes, R., Levison, H. F., Tsiganis, K. & Morbidelli, A. Origin of the cataclysmic Late Heavy Bombardment period of the terrestrial planets. *Nature* **435**, 466-469 (2005).
- 42 Nesvorný, D., Vokrouhlický, D., Bottke, W. F. & Levison, H. F. Evidence for very early migration of the Solar System planets from the Patroclus–Menoetius binary Jupiter Trojan. *Nature Astronomy* **2**, 878-882 (2018).
- 43 Brasser, R., Morbidelli, A., Gomes, R., Tsiganis, K. & Levison, H. F. Constructing the secular architecture of the solar system II: the terrestrial planets. *Astron. Astrophys.* **507**, 1053-1065 (2009).
- 44 Boehnke, P. & Harrison, T. M. Illusory Late Heavy Bombardments. *Proc. Natl. Acad. Sci. U.S.A.*, 201611535 (2016).
- 45 Quarles, B. & Kaib, N. Instabilities in the Early Solar System Due to a Self-gravitating Disk. *Astron. J.* **157** (2019).
- 46 Ribeiro, R. d. S. *et al.* Dynamical evidence for an early giant planet instability. *Icarus* **339**, 113605 (2020).
- 47 Clement, M. S., Kaib, N. A., Raymond, S. N. & Walsh, K. J. Mars' growth stunted by an early giant planet instability. *Icarus* **311**, 340-356 (2018).
- 48 Clement, M. S., Kaib, N. A., Raymond, S. N., Chambers, J. E. & Walsh, K. J. The early instability scenario: Terrestrial planet formation during the giant planet instability, and the effect of collisional fragmentation. *Icarus* **321**, 778-790 (2019).
- 49 Liu, B., Raymond, S. N. & Jacobson, S. B. Early Solar System instability triggered by dispersal of the gaseous disk. *Nature* **In Press** (2022).
- 50 Hunt, A. C., Ek, M. & Schönbächler, M. Separation of Platinum from Palladium and Iridium in Iron Meteorites and Accurate High-Precision Determination of Platinum

Isotopes by Multi-Collector ICP-MS. *Geostand. Geoanalytical Res.* **41**, 633-647 (2017).

- 51 Schönbächler, M., Carlson, R. W., Horan, M. F., Mock, T. D. & Hauri, E. H. High precision Ag isotope measurements in geologic materials by multiple-collector ICPMS: An evaluation of dry versus wet plasma. *Int. J. Mass Spectrom.* **261**, 183-191 (2007).
- 52 Blichert-Toft, J., Moynier, F., Lee, C.-T. A., Telouk, P. & Albarède, F. The early formation of the IVA iron meteorite parent body. *Earth Planet. Sci. Lett.* **296**, 469-480 (2010).
- 53 Chabot, N. L. Sulfur contents of the parental metallic cores of magmatic iron meteorites. *Geochim. Cosmochim. Acta* **68**, 3607-3618 (2004).
- 54 Randich, E. & Goldstein, J. I. Cooling rates of seven hexahedrites. *Geochim. Cosmochim. Acta* **42**, 221-233 (1978).
- 55 Ludwig, K. R. ISOPLOT 3.00. A geochronological toolkit for Microsoft Excel. *Berkeley Geochronological Center Spec. Publ. No. 4*, 70 pp (2003).

Hunt, A. C., et al. (2017). Platinum isotopes in iron meteorites: Galactic cosmic ray effects and nucleosynthetic homogeneity in the p-process isotope 190Pt and the other platinum isotopes. Geochimica et Cosmochimica Acta

REFERENCES

Hunt, A. C., et al. (2018). Late metal–silicate separation on the IAB parent asteroid: Constraints from combined W and Pt isotopes and thermal modelling. Earth and Planetary Science Letters 482: 490-500.

Kruijer, T. S., et al. (2014). Protracted core formation and rapid accretion of protoplanets. Science 344(6188): 1150-1154.

Pernicka, E. and J. T. Wasson (1987). Ru, Re, Os, Pt and Au in iron meteorites. Geochimica et Cosmochimica Acta 51(6): 1717-1726.

Petaev, M. I. and S. B. Jacobsen (2004). Differentiation of metal-rich meteoritic parent bodies: I. Measurements of PGEs, Re, Mo, W, and Au in meteoritic Fe-Ni metal. Meteoritics & Planetary Science 39(10): 1685-1697.

REERRICES
HAMEL AL COSI Lise medi-alliest separation on the lab prient stereoid. Constant from combined will of resonge and them and only lise to the process lead planetary Seirce Letters 482.495-500.
Hunk, A.C. et al. (20 Wasson, J. T. and G. W. Kallemeyn (2002). The IAB iron-meteorite complex: A group, five sub-groups, numerous grouplets, closely related, mainly formed by crystal segregation in rapidly cooling melts. Geochimica et Cosmochi

a o o o ↓

Theis, K. J., et al. (2013). Palladium–silver chronology of IAB iron meteorites. Earth and Planetary Science Letters 361: 402-411.

Wasson, J. T. (1999). Trapped melt in IIIAB irons; solid/liquid elemental partitioning during the fractionation of the IIIAB magma. Geochimica et Cosmochimica Acta 63(18): 2875-2889.

Wasson, J. T., et al. (2007). Formation of IIAB iron meteorites. Geochimica et Cosmochimica Acta 71(3): 760-781.

Schönbächler, M., et al. (2008). Silver isotope variations in chondrites: Volatile depletion and the initial 107Pd abundance of the solar system. Geochimica et Cosmochimica Acta 72(21): 5330-5341.