

On R-matrix identities related to elliptic anisotropic spin Ruijsenaars-Macdonald operators

M. Matushko^{◇*}

A. Zotov^{◇•}

◇ – *Steklov Mathematical Institute of Russian Academy of Sciences,
Gubkina str. 8, 119991, Moscow, Russia*

* – *Center for Advanced Studies, Skoltech, 143026, Moscow, Russia*

• – *Institute for Theoretical and Mathematical Physics,
Lomonosov Moscow State University, Moscow, 119991, Russia*

e-mails: matushko@mi-ras.ru, zotov@mi-ras.ru

Abstract

We propose and prove a set of identities for GL_M elliptic R -matrix (in the fundamental representation). In the scalar case ($M = 1$) these are elliptic function identities derived by S.N.M. Ruijsenaars as necessary and sufficient conditions for his kernel identity underlying construction of integral solutions to quantum spinless Ruijsenaars-Schneider model. In this respect the result of the present paper can be considered as the first step towards constructing solutions of quantum eigenvalue problem for the anisotropic spin Ruijsenaars model.

Contents

1	Introduction	2
2	Spin operators and R-matrix identities	3
3	New R-matrix identities	6
3.1	Statement	6
3.2	Proof of the Theorem for $k = 1$	7
3.3	Auxiliary lemmas	8
3.4	Proof of the Theorem for $k > 1$	9
3.5	Examples	9
4	Appendix A: elliptic functions	10
5	Appendix B: proof of auxiliary lemmas	12
	References	18

1 Introduction

In his paper [15] S.N.M. Ruijsenaars introduced a set of commuting elliptic difference operators

$$D_k = \sum_{|I|=k} \prod_{\substack{i \in I \\ j \notin I}} \phi(z_j - z_i) \prod_{i \in I} p_i, \quad k = 1, \dots, N, \quad (1.1)$$

where $\phi(z) = \phi(\hbar, z)$ is the elliptic Kronecker function (A.2). The sum in (1.1) is over all subsets I of size k of $\{1, \dots, N\} = \mathcal{N}$. The shift operators p_i act on a function $f(z_1, \dots, z_N)$ of complex variables $\mathbf{z} = z_1, \dots, z_N$ as follows:

$$(p_i f)(z_1, z_2, \dots, z_N) = \exp\left(-\eta \frac{\partial}{\partial z_i}\right) f(z_1, \dots, z_N) = f(z_1, \dots, z_i - \eta, \dots, z_N). \quad (1.2)$$

The operators (1.1) are called the Ruijsenaars-Macdonald operators since in the trigonometric limit they reproduce the Macdonald operators.

The proof of mutual commutativity of D_k was shown in [15] to be equivalent to the following set of functional equations:

$$\sum_{|I|=k} \left(\prod_{\substack{i \in I \\ j \notin I}} \phi(z_j - z_i) \phi(z_i - z_j - \eta) - \prod_{\substack{i \in I \\ j \notin I}} \phi(z_i - z_j) \phi(z_j - z_i - \eta) \right) = 0, \quad k = 1, \dots, N, \quad (1.3)$$

and the function $\phi(z)$ (A.2) was shown to satisfy (1.3).

Next, it was shown in [16] that the kernel identity $(D_k(\mathbf{x}) - D_k(-\mathbf{y}))\Psi(\mathbf{x}, \mathbf{y}) = \mathbf{0}$ holds true for $\Psi(\mathbf{x}, \mathbf{y})$ given by a certain product (and/or ratio) of elliptic gamma functions. It allows to construct integral solutions to quantum eigenproblem for operators D_k [6, 9, 7]. This approach was later extended to a more general class of operators [1]. The proof of the kernel identity is performed by a direct substitution which provides a set of identities for the function ϕ . Namely, it was proved in [16] that the kernel identities is valid iff the following set of relations for $2N$ complex variables ($\mathbf{x} = x_1, \dots, x_N$ and $\mathbf{y} = y_1, \dots, y_N$) holds:

$$\sum_{\substack{I \subset \mathcal{N} \\ |I|=k}} \left(\prod_{\substack{i \in I \\ j \in I^c}} \phi(x_i - x_j) \prod_{\substack{i \in I \\ j \in \mathcal{N}}} \phi(y_j - x_i) - \prod_{\substack{i \in I \\ j \in I^c}} \phi(y_j - y_i) \prod_{\substack{i \in I \\ j \in \mathcal{N}}} \phi(y_i - x_j) \right) = 0, \quad k = 1, \dots, N, \quad (1.4)$$

where I^c means the complement of a set I in \mathcal{N} . It was shown in [16] that the elliptic Kronecker function (A.2) satisfies (1.4). For example, for $k = 1$ (1.4) yields

$$\sum_{i=1}^N \left(\prod_{j \neq i}^N \phi(x_i - x_j) \prod_{j=1}^N \phi(y_j - x_i) - \prod_{j \neq i}^N \phi(y_j - y_i) \prod_{j=1}^N \phi(y_i - x_j) \right) = 0. \quad (1.5)$$

Under the substitution $x_i = z_i$, $y_i = z_i - \eta$ the latter identity reproduces the first one (with $k = 1$) from (1.3). For $k > 1$ the substitution $x_i = z_i$, $y_i = z_i - \eta$ provides particular case of (1.3) corresponding to $\hbar = 2\eta$.

Let us mention that the identities (1.4) arise in the context of Bäcklund transformations in the classical (elliptic) Ruijsenaars-Schneider model [6]. Also, (1.5) provides the so-called self-dual form of the Ruijsenaars-Schneider model coming from the multi-pole ansatz for intermediate long-wave equation [17].

Purpose of the paper is to extend the identities (1.4) to R -matrix identities in the following sense. The elliptic Baxter-Belavin [2, 3] R -matrix (A.12) in the fundamental representation of GL_M Lie group can be considered as matrix generalization of the Kronecker function (A.2) because (A.2) is a particular case of (A.12) corresponding to $M = 1$. R -matrices do not commute (in the general case). Our aim is to find noncommutative analogue of (1.4) which reproduce (1.4) in the $M = 1$ case.

In our previous paper [13] (see also [14] for applications to long-range spin chains) we introduced anisotropic spin generalization for the operators (1.1). Similarly to the scalar case [15] we deduced and proved R -matrix generalizations of the identities (1.3). These R -matrix identities are necessary and sufficient conditions for commutativity of spin Ruijsenaars-Macdonald operators. In this paper we propose and prove R -matrix generalizations for the identities (1.4) from [16]:

$$\begin{aligned}
& \sum_{1 \leq i_1 < \dots < i_k \leq N} \left(\overrightarrow{\prod_{l_k=i_k+1}^N R_{i_k l_k}^h} \overrightarrow{\prod_{\substack{l_{k-1}=i_{k-1}+1 \\ l_{k-1} \neq i_k}}^N R_{i_{k-1} l_{k-1}}^h} \dots \overrightarrow{\prod_{\substack{l_1=i_1+1 \\ l_1 \neq i_2 \dots i_k}}^N R_{i_1 l_1}^h} \times \right. \\
& \quad \times \overrightarrow{\prod_{j_1=N+1}^{2N} R_{j_1 i_1}^h} \overrightarrow{\prod_{j_2=N+1}^{2N} R_{j_2 i_2}^h} \dots \overrightarrow{\prod_{j_k=N+1}^{2N} R_{j_k i_k}^h} \times \\
& \quad \times \overrightarrow{\prod_{\substack{m_k=1 \\ m_k \neq i_1 \dots i_{k-1}}}^{i_k-1} R_{i_k m_k}^h} \overrightarrow{\prod_{\substack{m_{k-1}=1 \\ m_{k-1} \neq i_1 \dots i_{k-2}}}^{i_{k-1}-1} R_{i_{k-1} m_{k-1}}^h} \dots \overrightarrow{\prod_{m_1=1}^{i_1-1} R_{i_1 m_1}^h} \Big) - \\
& - \sum_{N+1 \leq j_1 < \dots < j_k \leq 2N} \left(\overrightarrow{\prod_{l_k=j_k+1}^{2N} R_{l_k j_k}^h} \overrightarrow{\prod_{\substack{l_{k-1}=j_{k-1}+1 \\ l_{k-1} \neq j_k}}^{2N} R_{l_{k-1} j_{k-1}}^h} \dots \overrightarrow{\prod_{\substack{l_1=j_1+1 \\ l_1 \neq j_2 \dots j_k}}^{2N} R_{l_1 j_1}^h} \times \right. \\
& \quad \times \overrightarrow{\prod_{i_1=1}^N R_{j_1 i_1}^h} \overrightarrow{\prod_{i_2=1}^N R_{j_2 i_2}^h} \dots \overrightarrow{\prod_{i_k=1}^N R_{j_k i_k}^h} \times \\
& \quad \times \overrightarrow{\prod_{\substack{m_k=N+1 \\ m_k \neq j_1 \dots j_{k-1}}}^{j_k-1} R_{m_k j_k}^h} \overrightarrow{\prod_{\substack{m_{k-1}=N+1 \\ m_{k-1} \neq j_1 \dots j_{k-2}}}^{j_{k-1}-1} R_{m_{k-1} j_{k-1}}^h} \dots \overrightarrow{\prod_{m_1=N+1}^{j_1-1} R_{m_1 j_1}^h} \Big) = 0.
\end{aligned} \tag{1.6}$$

Notations are explained in the next Section. This result is the first step towards constructing the kernel identity and solutions to quantum problem of the (anisotropic) spin Ruijsenaars model. The latter is quite nontrivial problem since it requires a definition of matrix generalization of elliptic gamma function. We hope to clarify these questions in our future publications.

2 Spin operators and R-matrix identities

Here we recall some notations and statements from [13]. Some additional notations are introduced for the purposes of the present paper.

R -matrices and Yang-Baxter equations. Let \mathcal{H} be a vector space $\mathcal{H} = (\mathbb{C}^M)^{\otimes N}$. The elliptic Baxter-Belavin R -matrix [2, 3] is a linear map (A.12) $R_{ij}^h(z) \in \text{End}(\mathcal{H})$ acting non-trivially in the i -th and j -th tensor components of \mathcal{H} only and satisfying the quantum Yang-Baxter equation:

$$R_{ij}^h(u)R_{ik}^h(u+v)R_{jk}^h(v) = R_{jk}^h(v)R_{ik}^h(u+v)R_{ij}^h(u) \quad (2.1)$$

for any distinct integers $1 \leq i, j, k \leq N$. Also, for any distinct integers $1 \leq i, j, k, l \leq N$

$$[R_{ij}^h(u), R_{kl}^{h'}(v)] = 0. \quad (2.2)$$

Consider a pair I, J of disjoint subsets in $\{1, \dots, N\}$. The elements in I, J are in increasing ordering, that is for $J = \{j_1, j_2, \dots, j_k\}$ the elements j_m are in increasing order $j_1 < j_2 < \dots < j_k$. Similarly, for $I = \{i_1, i_2, \dots, i_l\}$ we have $i_1 < i_2 < \dots < i_l$. We will use notations $\overrightarrow{\prod_{j=1}^N} R_{ij}$ and $\overleftarrow{\prod_{j=1}^N} R_{ij}$, where the arrows mean the ordering. For example, $\overrightarrow{\prod_{j=1}^N} R_{ij} = R_{i1}R_{i2}\dots R_{iN}$ and $\overleftarrow{\prod_{j=1}^N} R_{ij} = R_{iN}R_{i,N-1}\dots R_{i1}$.

Introduce the following notation:

$$\mathcal{R}_{I,J} = \overleftarrow{\prod_{\substack{i_1 \in I \\ i_1 < j_1}} R_{i_1,j_1}} (z_{i_1} - z_{j_1}) \overleftarrow{\prod_{\substack{i_2 \in I \\ i_2 < j_2}} R_{i_2,j_2}} (z_{i_2} - z_{j_2}) \dots \overleftarrow{\prod_{\substack{i_k \in I \\ i_k < j_k}} R_{i_k,j_k}} (z_{i_k} - z_{j_k}). \quad (2.3)$$

Using the property (2.2) it is easy to rewrite it as

$$\mathcal{R}_{I,J} = \overrightarrow{\prod_{\substack{j_l \in J \\ j_l > i_l}} R_{i_l,j_l}} (z_{i_l} - z_{j_l}) \overrightarrow{\prod_{\substack{j_{l-1} \in J \\ j_{l-1} > i_{l-1}}} R_{i_{l-1},j_{l-1}}} (z_{i_{l-1}} - z_{j_{l-1}}) \dots \overrightarrow{\prod_{\substack{j_1 \in J \\ j_1 > i_1}} R_{i_1,j_1}} (z_{i_1} - z_{j_1}). \quad (2.4)$$

In a similar way we also introduce

$$\mathcal{R}'_{I,J} = \overrightarrow{\prod_{\substack{j_l \in J \\ j_l < i_l}} R_{i_l,j_l}} (z_{i_l} - z_{j_l}) \overrightarrow{\prod_{\substack{j_{l-1} \in J \\ j_{l-1} < i_{l-1}}} R_{i_{l-1},j_{l-1}}} (z_{i_{l-1}} - z_{j_{l-1}}) \dots \overrightarrow{\prod_{\substack{j_1 \in J \\ j_1 < i_1}} R_{i_1,j_1}} (z_{i_1} - z_{j_1}) = \quad (2.5)$$

$$= \overleftarrow{\prod_{\substack{i_1 \in I \\ i_1 > j_1}} R_{i_1,j_1}} (z_{i_1} - z_{j_1}) \overleftarrow{\prod_{\substack{i_2 \in I \\ i_2 > j_2}} R_{i_2,j_2}} (z_{i_2} - z_{j_2}) \dots \overleftarrow{\prod_{\substack{i_k \in I \\ i_k > j_k}} R_{i_k,j_k}} (z_{i_k} - z_{j_k}). \quad (2.6)$$

For any disjoint subsets A, B, C of $\{1, 2, \dots, N\}$ the following identities hold true:

$$\mathcal{R}_{C,A \cup B} \mathcal{R}_{B,A} = \mathcal{R}_{B \cup C,A} \mathcal{R}_{C,B}, \quad (2.7)$$

$$\mathcal{R}'_{A,B} \mathcal{R}'_{A \cup B,C} = \mathcal{R}'_{B,C} \mathcal{R}'_{A,B \cup C}. \quad (2.8)$$

The latter identities follow from the Yang-Baxter equation (2.1).

Elliptic R -matrix satisfies the unitarity property

$$R_{ij}^h(z)R_{ji}^h(-z) = \text{Id } \phi(\hbar, z)\phi(\hbar, -z), \quad (2.9)$$

Below we also use R -matrices in a slightly different normalization:

$$R_{ij}^h(z) = \phi(\hbar, z)\bar{R}_{ij}^h(z). \quad (2.10)$$

Then

$$\bar{R}_{ij}^h(z)\bar{R}_{ji}^h(-z) = \text{Id}. \quad (2.11)$$

For the products defined above we have:

$$\mathcal{R}_{I,J}\mathcal{R}'_{J,I} = \text{Id} \prod_{\substack{i \in I, j \in J \\ i < j}} \phi(h, z_i - z_j) \phi(h, z_j - z_i) \quad (2.12)$$

and

$$\mathcal{R}'_{I,J}\mathcal{R}_{J,I} = \text{Id} \prod_{\substack{i \in I, j \in J \\ i > j}} \phi(h, z_i - z_j) \phi(h, z_j - z_i). \quad (2.13)$$

Similarly,

$$\bar{\mathcal{R}}_{I,J}\bar{\mathcal{R}}'_{J,I} = \bar{\mathcal{R}}'_{J,I}\bar{\mathcal{R}}_{I,J} = \text{Id}. \quad (2.14)$$

Finally, the elliptic R -matrix (A.12) is skew-symmetric:

$$R_{12}^h(z) = -R_{21}^{-h}(-z). \quad (2.15)$$

Below by writing $R_{12}^{-h}(z)$ we assume $R_{21}^h(-z)$.

In this paper we also use the following notations. For disjoint sets $I = \{i_1 < i_2 < \dots < i_k\}$ and $J = \{j_1 < j_2 < \dots < j_l\}$ we denote by $\mathcal{Y}_{I,J}^h$ the following product of R -matrices

$$\mathcal{Y}_{I,J}^h = \overrightarrow{\prod}_{j \in J} R_{i_1,j}^h \dots \overrightarrow{\prod}_{j \in J} R_{i_k,j}^h. \quad (2.16)$$

Due to (2.2) it can be written as

$$\mathcal{Y}_{I,J}^h = \overrightarrow{\prod}_{i \in I} R_{i,j_1}^h \dots \overrightarrow{\prod}_{i \in I} R_{i,j_l}^h. \quad (2.17)$$

In what follows we use the auxiliary statement based on the Yang-Baxter equation (2.1)-(2.2).

Lemma 2.1 *Let $I = \{i_1 < \dots < i_{m-1} < i_m = a < i_{m+1} < \dots < i_k\}$ and $J = \{j_1 < \dots < j_{n-1} < j_n = b < j_{n+1} < \dots < j_l\}$ be disjoint sets. Then the following relations hold true:*

$$\mathcal{Y}_{I,J}^h R_{b,j_{n+1}}^h \dots R_{b,j_l}^h = R_{b,j_{n+1}}^h \dots R_{b,j_l}^h \mathcal{Y}_{I,J \setminus \{b\}}^h \mathcal{Y}_{I,\{b\}}^h \quad (2.18)$$

and

$$\mathcal{Y}_{I,J}^h R_{i_{m+1},a}^h \dots R_{i_k,a}^h = R_{i_{m+1},a}^h \dots R_{i_k,a}^h \mathcal{Y}_{I \setminus \{a\},J}^h \mathcal{Y}_{\{a\},J}^h. \quad (2.19)$$

The proof is given in the Appendix.

Spin operators are defined as follows:

$$\mathcal{D}_k = \sum_{|I|=k} \prod_{\substack{i \in I^c, \\ j \in I}} \phi(z_i - z_j) \cdot \bar{\mathcal{R}}_{I^c,I} \cdot \mathbf{p}_I \cdot \bar{\mathcal{R}}'_{I,I^c}, \quad \mathbf{p}_I = \prod_{i \in I} p_i, \quad (2.20)$$

where p_i are the shift operators (1.2) and the bars over \mathcal{R} mean that in the definitions (2.3), (2.5) we use R -matrices normalized as in (2.10)-(2.11). For $M = 1$ the operators (2.20) reproduce the Ruijsenaars-Macdonald operators (1.1). The operators of type (2.20) were introduced in [10] in trigonometric case with $M = 2$ based on the $U_q(\mathfrak{gl}_2)$ R -matrix.

Let us also remark that using the so-called freezing trick the spin operators (2.20) provide elliptic integrable long-range spin chains [14]. In the classical mechanics the operators (1.1) are the Hamiltonians of the elliptic Ruijsenaars-Schneider model, while the (anisotropic) spin operators are quantum analogues of the Hamiltonians of (relativistic) interacting tops [19, 5]. Description of relativistic tops can be found in [11, 8].

Commutativity of \mathcal{D}_k turns out to be equivalent to the following set of R -matrix identities:

$$\sum_{|I|=k} \left(\mathcal{R}_{I^c, I} \cdot \mathcal{R}'_{I_-, I^c} \cdot \mathcal{R}_{I_-, I^c} \cdot \mathcal{R}'_{I^c, I} - \mathcal{R}_{I, I^c} \cdot \mathcal{R}'_{I^c, I} \cdot \mathcal{R}_{I^c, I} \cdot \mathcal{R}'_{I, I^c} \right) = 0, \quad (2.21)$$

which hold true for any $k = 1, \dots, N$. Here we use the notation $\mathcal{R}_{I_-, J} = \mathbf{p}_I \mathcal{R}_{I, J} \mathbf{p}_I^{-1}$. Derivation of (2.21) as well as its proof and explicit examples can be found in [13]. For $M = 1$ (2.21) turn into identities (1.3).

3 New R-matrix identities

3.1 Statement

It is convenient to define a set of $2N$ complex variables z_1, \dots, z_{2N} , which unifies \mathbf{x} and \mathbf{y} :

$$z_i = \begin{cases} x_i & \text{for } i \in \mathcal{N}_1, \quad \mathcal{N}_1 = \{1, \dots, N\}, \\ y_i & \text{for } i \in \mathcal{N}_2, \quad \mathcal{N}_2 = \{N+1, \dots, 2N\}, \end{cases} \quad (3.1)$$

The R -matrices act on the space $\mathcal{H} = (\mathbb{C}^M)^{\otimes 2N}$, and the following short notations are assumed:

$$R_{ij}^h = \begin{cases} R_{ij}^h(x_i - x_j), & \text{if } i, j \in \mathcal{N}_1, \\ R_{ij}^h(x_i - y_j), & \text{if } i \in \mathcal{N}_1, j \in \mathcal{N}_2, \\ R_{ij}^h(y_i - x_j), & \text{if } i \in \mathcal{N}_2, j \in \mathcal{N}_1, \\ R_{ij}^h(y_i - y_j), & \text{if } i, j \in \mathcal{N}_2. \end{cases} \quad (3.2)$$

Theorem *The following identities hold for any $k = 1, \dots, N$:*

$$\sum_{\substack{I \subset \mathcal{N}_1 \\ |I|=k}} \mathcal{R}_{I, I^c}^h \mathcal{Y}_{\mathcal{N}_2, I}^h \mathcal{R}_{I, I^c}^h + (-1)^{k-1} \sum_{\substack{J \subset \mathcal{N}_2 \\ |J|=k}} \mathcal{R}_{J, J^c}^{-h} \mathcal{Y}_{\mathcal{N}_1, J}^{-h} \mathcal{R}_{J, J^c}^{'-h} = 0, \quad (3.3)$$

where $R_{ij} = R_{ij}^h(z_i - z_j)$ is the elliptic R -matrix (A.12) acting in the i -th and j -th tensor components, $i, j = 1, \dots, 2N$ according to the rule¹ (3.1)-(3.2). In the $M = 1$ case the R -matrix identities (3.3) turn into (1.4).

The idea of the proof is as follows. Denote the l.h.s. of (3.3) as $\mathcal{F}[k, N] = \mathcal{F}_1[k, N] - \mathcal{F}_2[k, N]$, where $\mathcal{F}_1[k, N], \mathcal{F}_2[k, N]$ are the first and the second sum in (3.3) respectively²:

$$\mathcal{F}_1[k, N] = \sum_{\substack{I \subset \mathcal{N}_1 \\ |I|=k}} \mathcal{R}_{I, I^c}^h \mathcal{Y}_{\mathcal{N}_2, I}^h \mathcal{R}_{I, I^c}^h, \quad (3.4)$$

¹In the first sum of (3.3) the set I^c is a complement of I in \mathcal{N}_1 . Similarly, in the second sum of (3.3) the set J^c is a complement of J in \mathcal{N}_2 .

²Sometimes we omit the indices k, N and write \mathcal{F} instead of $\mathcal{F}[k, N]$.

$$\mathcal{F}_2[k, N] = (-1)^k \sum_{\substack{J \subset \mathcal{N}_2 \\ |J|=k}} \mathcal{R}_{J, J^c}^{-\hbar} \mathcal{Y}_{\mathcal{N}_1, J}^{-\hbar} \mathcal{R}_{J, J^c}^{\prime -\hbar}. \quad (3.5)$$

More explicitly,

$$\begin{aligned} \mathcal{F}_1[k, N] = & \sum_{1 \leq i_1 < \dots < i_k \leq N} \left(\overrightarrow{\prod_{l_k=i_k+1}^N R_{i_k l_k}^{\hbar}} \overrightarrow{\prod_{\substack{l_{k-1}=i_{k-1}+1 \\ l_{k-1} \neq i_k}}^N R_{i_{k-1} l_{k-1}}^{\hbar}} \dots \overrightarrow{\prod_{\substack{l_1=i_1+1 \\ l_1 \neq i_2 \dots i_k}}^N R_{i_1 l_1}^{\hbar}} \times \right. \\ & \times \overrightarrow{\prod_{j_1=N+1}^{2N} R_{j_1 i_1}^{\hbar}} \overrightarrow{\prod_{j_2=N+1}^{2N} R_{j_2 i_2}^{\hbar}} \dots \overrightarrow{\prod_{j_k=N+1}^{2N} R_{j_k i_k}^{\hbar}} \times \\ & \times \left. \overrightarrow{\prod_{\substack{m_k=1 \\ m_k \neq i_1 \dots i_{k-1}}}^{i_k-1} R_{i_k m_k}^{\hbar}} \overrightarrow{\prod_{\substack{m_{k-1}=1 \\ m_{k-1} \neq i_1 \dots i_{k-2}}}^{i_{k-1}-1} R_{i_{k-1} m_{k-1}}^{\hbar}} \dots \overrightarrow{\prod_{m_1=1}^{i_1-1} R_{i_1 m_1}^{\hbar}} \right). \end{aligned} \quad (3.6)$$

Let us also rewrite expression (3.5) through R -matrices depending on \hbar (but not on $-\hbar$) using the skew-symmetry property (2.15):

$$\begin{aligned} \mathcal{F}_2[k, N] = & \sum_{N+1 \leq j_1 < \dots < j_k \leq 2N} \left(\overrightarrow{\prod_{l_k=j_k+1}^{2N} R_{l_k j_k}^{\hbar}} \overrightarrow{\prod_{\substack{l_{k-1}=j_{k-1}+1 \\ l_{k-1} \neq j_k}}^{2N} R_{l_{k-1} j_{k-1}}^{\hbar}} \dots \overrightarrow{\prod_{\substack{l_1=j_1+1 \\ l_1 \neq j_2 \dots j_k}}^{2N} R_{l_1 j_1}^{\hbar}} \times \right. \\ & \times \overrightarrow{\prod_{i_1=1}^N R_{j_1 i_1}^{\hbar}} \overrightarrow{\prod_{i_2=1}^N R_{j_2 i_2}^{\hbar}} \dots \overrightarrow{\prod_{i_k=1}^N R_{j_k i_k}^{\hbar}} \times \\ & \times \left. \overrightarrow{\prod_{\substack{m_k=N+1 \\ m_k \neq j_1 \dots j_{k-1}}}^{j_k-1} R_{m_k j_k}^{\hbar}} \overrightarrow{\prod_{\substack{m_{k-1}=N+1 \\ m_{k-1} \neq j_1 \dots j_{k-2}}}^{j_{k-1}-1} R_{m_{k-1} j_{k-1}}^{\hbar}} \dots \overrightarrow{\prod_{m_1=N+1}^{j_1-1} R_{m_1 j_1}^{\hbar}} \right). \end{aligned} \quad (3.7)$$

In this way (3.3) takes the form (1.6).

Consider \mathcal{F} as (matrix valued) function of $z_1 = x_1$. Our strategy is to use Lemma 4.1 from the Appendix similarly to the proof of (1.4) in [16]. We are going to show that \mathcal{F} is an entire function of z_1 . The behaviour of R -matrices on the lattice of periods $\mathbb{Z} \oplus \tau\mathbb{Z}$ is nontrivial. It is given by (A.15). For this reason we use the approach from [13]. Namely, we consider $\mathcal{F}(z_1)$ on the "large" torus with periods M and $M\tau$. Due to (A.10) $Q^M = \Lambda^M = 1_M$ the quasi-periodic behaviour of R -matrices (and \mathcal{F}) on the large torus is simplified and we may use Lemma 4.1. First, we are going to prove the absence of poles (in the variable z_1) in the fundamental parallelogram $0 \leq z_1 < 1$, $0 \leq z_1 < \tau$ and then extend the proof to the large torus.

The proof is performed by induction in k and N . For $k = 1$ the identity (3.1) can be deduced from the higher order R -matrix identities based on the associative Yang-Baxter equation.

3.2 Proof of the Theorem for $k = 1$

The proof of (1.5) can be performed straightforwardly using (A.7), which is a higher order generalization of the addition formula (A.5). Details can be found in the Appendix of [17]. Here we prove (3.3) for

$k = 1$ in a similar way. For this purpose we recall that the elliptic R -matrix in the fundamental representation (A.12) satisfies the so-called associative Yang-Baxter equation [4]:

$$R_{12}^u R_{23}^{u'} = R_{13}^{u'} R_{12}^{u-u'} + R_{23}^{u'-u} R_{13}^u, \quad R_{ab}^h = R_{ab}^h(z_a - z_b). \quad (3.8)$$

In [18] the higher order analogue of (3.8) was derived. Here we use it in a slightly different form suggested in [13]:

$$\prod_{i=1}^{\overrightarrow{n}} R_{a,i}^{u_i}(w_i) = \sum_{m=1}^n \prod_{j=m+1}^{\overrightarrow{n}} R_{m,j}^{u_j}(w_j - w_m) \cdot R_{a,m}^U(w_m) \cdot \prod_{j=1}^{\overrightarrow{m-1}} R_{m,j}^{u_j}(w_j - w_m), \quad (3.9)$$

where $U = \sum_{k=1}^n u_k$ and $a \notin \{1 \dots n\}$, $n \in \mathbb{Z}_+$. Let us fix $a = 1$ together with replacing the set $\{1 \dots n\}$ with $\{2 \dots n+1\}$. Then (3.9) takes the form

$$\prod_{i=2}^{\overrightarrow{n+1}} R_{1,i}^{u_i}(w_i) = \sum_{m=2}^{n+1} \prod_{j=m+1}^{\overrightarrow{n+1}} R_{m,j}^{u_j}(w_j - w_m) \cdot R_{1,m}^U(w_m) \cdot \prod_{j=2}^{\overrightarrow{m-1}} R_{m,j}^{u_j}(w_j - w_m). \quad (3.10)$$

Set $n = 2N - 1$ and identify $w_i = z_1 - z_i$ for $i = 2, \dots, 2N$ (recall (3.1)-(3.2)). Also, put $u_2 = \dots = u_N = \hbar$ and $u_{N+1} = \dots = u_{2N} = -\hbar$. Then $U = -\hbar$. After all substitutions in the l.h.s. of (3.10) one gets $R_{12}^h \dots R_{1N}^h \cdot R_{1,N+1}^{-\hbar} \dots R_{1,2N}^{-\hbar}$. Writing down the r.h.s. of (3.10) finally yields

$$\begin{aligned} & \sum_{i=1}^N R_{i,i+1}^h \dots R_{i,N}^h \cdot R_{N+1,i}^h \dots R_{2N,i}^h \cdot R_{i,1}^h \dots R_{i,i-1}^h - \\ & - \sum_{i=1}^N R_{N+i+1,N+i}^h \dots R_{2N,N+i}^h \cdot R_{N+i,1}^h \dots R_{N+i,N}^h \cdot R_{N+1,N+i}^h \dots R_{N+i-1,N+i}^h = 0, \end{aligned} \quad (3.11)$$

where we used the skew-symmetry of R -matrices (2.15). The obtained relation (3.11) is (3.3) for $k = 1$.

3.3 Auxiliary lemmas

Here we formulate two lemmas useful for the proof of the Theorem. The proof of Lemmas is given in the Appendix.

Lemma 3.1 Consider $\mathcal{F} = \mathcal{F}[k, N]$ as function of $z_a = x_a$. It has no poles at $x_1, \dots, x_{a-1}, x_{a+1}, \dots, x_N$.

Lemma 3.2 Consider $\mathcal{F} = \mathcal{F}[k, N]$ as function of $z_a = x_a$. The following relation holds for any $b \in \mathcal{N}_2$:

$$\text{Res}_{z_a=z_b} \mathcal{F}[k, N] = -G \cdot \mathcal{F}[k-1, N-1] \cdot H, \quad b = N+1, \dots, 2N, \quad (3.12)$$

where

$$G = R_{a,a+1}^h \dots R_{a,N}^h \cdot R_{b+1,b}^h \dots R_{2N,b}^h, \quad (3.13)$$

$$H = R_{N+1,a}^h \dots R_{b-1,a}^h \cdot R_{b,1}^h \dots R_{b,a-1}^h P_{ab} \quad (3.14)$$

and by $\mathcal{F}[k-1, N-1]$ we mean the expression $\mathcal{F} = \mathcal{F}_1 - \mathcal{F}_2$ (3.4)-(3.5) taken for $k-1$ and depending on $2(N-1)$ variables $\{x_1, \dots, x_N\} \setminus \{x_a\}$, $\{y_1, \dots, y_N\} \setminus \{y_b\}$.

The latter statement is applied to the proof of the Theorem by setting $a = 1$.

3.4 Proof of the Theorem for $k > 1$

Let us prove the main Theorem (3.3).

Proof. The proof is by induction in k . For $k = 1$ the statement was proved above in (3.11). Suppose (3.3) is true for $k - 1$. We need to prove it for k .

Main idea is the same as in the scalar case. We are going to use Lemma 4.1 given in the Appendix. For this purpose consider the quasi-periodic properties of \mathcal{F} as function of z_1 . Due to (A.15) any R -matrix of the form $R_{1k}^h = R_{1k}^h(z_1, z_k) = R_{1k}^h(z_1 - z_k)$, $k = 2, \dots, 2N$ is transformed as follows:

$$\begin{aligned} R_{1k}^h(z_1 + 1, z_k) &= Q_1^{-1} R_{1k}^h(z_1, z_k) Q_1, \\ R_{1k}^h(z_1 + \tau, z_k) &= \exp(-2\pi i \frac{\hbar}{M}) \Lambda_1^{-1} R_{1k}^h(z_1, z_k) \Lambda_1, \end{aligned} \quad (3.15)$$

where $Q_1 = Q \otimes 1_M \otimes \dots \otimes 1_M = Q \otimes 1_M^{\otimes(2N-1)}$ and similarly for Λ_1 . For R -matrices of the form $R_{k1}^h = R_{k1}^h(z_k, z_1) = R_{k1}^h(z_k - z_1)$ we have:

$$R_{k1}^h(z_k, z_1 + 1) = R_{k1}^h(z_k - 1, z_1) = Q_k R_{k1}^h(z_1) Q_k^{-1} = Q_1^{-1} R_{12}^h(z_k, z_1) Q_1 \quad (3.16)$$

and in the same way

$$R_{k1}^h(z_k, z_1 + \tau) = \exp(2\pi i \frac{\hbar}{M}) \Lambda_1^{-1} R_{k1}^h(z_k, z_1) \Lambda_1, \quad (3.17)$$

Thus, any R -matrix which contains index 1 is transformed as (3.15) or (3.16)-(3.17). The number of R -matrices of the forms R_{1k}^h or R_{k1}^h is different in different terms. But the difference of these numbers is the same for all the terms: the number of R -matrices of the form R_{1k}^h is less than those of the form R_{k1}^h by k . Therefore,

$$\mathcal{F}(z_1 + 1) = Q_1^{-1} \mathcal{F}(z_1) Q_1, \quad (3.18)$$

$$\mathcal{F}(z_1 + \tau) = \exp(2\pi i \frac{k\hbar}{M}) \Lambda_1^{-1} \mathcal{F}(z_1) \Lambda_1.$$

In order to use Lemma 4.1 let us extend the torus to the large one with periods $M, M\tau$. Then due to $Q^M = \Lambda^M = 1_M$ we have

$$\mathcal{F}(z_1 + M) = \mathcal{F}(z_1), \quad (3.19)$$

$$\mathcal{F}(z_1 + M\tau) = \exp(2\pi i k\hbar) \mathcal{F}(z_1).$$

Next, notice that $\mathcal{F}(z_1)$ is an entire function of z_1 on the fundamental parallelogram ($0 \leq z_1 < 1$, $0 \leq z_1 < \tau$). Indeed, due to (A.14) all possible poles inside the fundamental parallelogram are at points x_2, \dots, x_N and y_1, \dots, y_N . By Lemma 3.1 the poles at points x_2, \dots, x_N are absent. The absence of poles y_1, \dots, y_N easily follows from Lemma 3.2 and the induction assumption (which implies $\mathcal{F}[k-1, N-1] = 0$). Finally, by taking residue $\text{Res}_{z_1=z_k}$, $k = 2, \dots, 2N$ of both sides of (3.18) we conclude that the absence of poles in the fundamental parallelogram is extended to the absence of poles on the large torus. Then all requirements of Lemma 4.1 are fulfilled and the Theorem is proved. ■

3.5 Examples

Let us give a few explicit examples for the identities (1.6).

Example. $N = 2$, $k = 1$:

$$R_{12}^h R_{31}^h R_{41}^h + R_{32}^h R_{42}^h R_{21}^h - R_{43}^h R_{31}^h R_{32}^h - R_{41}^h R_{42}^h R_{34}^h = 0. \quad (3.20)$$

Example. $N = 3, k = 1$:

$$\begin{aligned} & R_{12}^h R_{13}^h R_{41}^h R_{51}^h R_{61}^h + R_{23}^h R_{42}^h R_{52}^h R_{62}^h R_{21}^h + R_{43}^h R_{53}^h R_{63}^h R_{31}^h R_{32}^h \\ & - R_{54}^h R_{64}^h R_{41}^h R_{42}^h R_{43}^h - R_{65}^h R_{51}^h R_{52}^h R_{53}^h R_{45}^h - R_{61}^h R_{62}^h R_{63}^h R_{46}^h R_{56}^h = 0. \end{aligned} \quad (3.21)$$

Example. $N = 3, k = 2$:

$$\begin{aligned} & R_{23}^h R_{13}^h R_{41}^h R_{51}^h R_{61}^h R_{42}^h R_{52}^h R_{62}^h + R_{12}^h R_{41}^h R_{51}^h R_{61}^h R_{43}^h R_{53}^h R_{63}^h R_{32}^h \\ & + R_{42}^h R_{52}^h R_{62}^h R_{43}^h R_{53}^h R_{63}^h R_{31}^h R_{21}^h - R_{65}^h R_{64}^h R_{41}^h R_{42}^h R_{43}^h R_{51}^h R_{52}^h R_{53}^h \\ & - R_{54}^h R_{41}^h R_{42}^h R_{43}^h R_{61}^h R_{62}^h R_{63}^h R_{56}^h - R_{51}^h R_{52}^h R_{53}^h R_{61}^h R_{62}^h R_{63}^h R_{46}^h R_{45}^h = 0. \end{aligned} \quad (3.22)$$

Let us make a comment on the trigonometric and rational degenerations. All trigonometric and rational R -matrices considered in [13, 14] satisfy the obtained set of identities. Details will be given in a separate publication.

Another remark is that in trigonometric and rational cases the scalar identities (1.4) can be extended to the case when the number of variables \mathbf{x} is different from the number of variables \mathbf{y} , so that we have $|\mathbf{x}| = N_1, |\mathbf{y}| = N_2$ and $N_1 \neq N_2$. This can be explained as follows. In the trigonometric or rational case one can consider a limit $x_i \rightarrow \infty$, which decrease the number of variables \mathbf{x} by one. Similar arguments work for R -matrix identities if the limit $\lim_{x_i \rightarrow \infty} R_{ij}(x_i - x_j)$ (or $\lim_{x_i \rightarrow \infty} R_{i,N+j}(x_i - y_j)$) is well defined. It is not true for all possible degenerations of the elliptic R -matrix. At the same time it is true for a number of widely known cases. For example, it is obviously true for the rational Yang's R -matrix $R_{12}^h(z) = 1_M \otimes 1_M \hbar^{-1} + P_{12} z^{-1}$. Then $\lim_{z \rightarrow \infty} R_{12}^h(z) = 1_M \otimes 1_M \hbar^{-1}$. Therefore, the limit $x_i \rightarrow \infty$ being applied to some identity provides another identity, where all R -matrices containing x_i are replaced by the scalar factor \hbar^{-1} . We will consider these cases in our future works.

4 Appendix A: elliptic functions

Elliptic functions. We use the following theta-function:

$$\vartheta(z) = \vartheta(z|\tau) = - \sum_{k \in \mathbb{Z}} \exp \left(\pi i \tau \left(k + \frac{1}{2} \right)^2 + 2\pi i \left(z + \frac{1}{2} \right) \left(k + \frac{1}{2} \right) \right), \quad \text{Im}(\tau) > 0. \quad (\text{A.1})$$

It is odd $\vartheta(-z) = -\vartheta(z)$ and has simple zero at $z = 0$. The Kronecker elliptic function is defined as

$$\phi(z, u) = \frac{\vartheta'(0)\vartheta(z+u)}{\vartheta(z)\vartheta(u)} = \phi(u, z). \quad (\text{A.2})$$

As function of z it has simple pole at $z = 0$ and $\text{Res}_{z=0} \phi(z, u) = 1$. The quasi-periodic properties on the lattice of periods $\Gamma = \mathbb{Z} \oplus \mathbb{Z}\tau$ (of elliptic curve \mathbb{C}/Γ):

$$\vartheta(z+1) = -\vartheta(z), \quad \vartheta(z+\tau) = -e^{-\pi i \tau - 2\pi i z} \vartheta(z), \quad (\text{A.3})$$

so that

$$\phi(z+1, u) = \phi(z, u), \quad \phi(z+\tau, u) = e^{-2\pi i u} \phi(z, u). \quad (\text{A.4})$$

The function (A.2) satisfies the following addition formula:

$$\phi(z_1, u_1)\phi(z_2, u_2) = \phi(z_1, u_1 + u_2)\phi(z_2 - z_1, u_2) + \phi(z_2, u_1 + u_2)\phi(z_1 - z_2, u_1) \quad (\text{A.5})$$

and the identity

$$\phi(z, u)\phi(z, -u) = \wp(z) - \wp(u). \quad (\text{A.6})$$

where $\wp(x)$ – is the Weierstrass \wp -function. The higher order analogue of (A.5) is as follows:

$$\prod_{i=1}^n \phi(w_i, u_i) = \sum_{i=1}^n \phi\left(w_i, \sum_{l=1}^n u_l\right) \prod_{j \neq i}^n \phi(w_j - w_i, u_j), \quad n \in \mathbb{Z}_+. \quad (\text{A.7})$$

R-matrix. Define the set of M^2 functions:

$$\varphi_a(z, \omega_a + \hbar) = \exp\left(2\pi i \frac{a_2 z}{M}\right) \phi(z, \omega_a + \hbar), \quad \omega_a = \frac{a_1 + a_2 \tau}{M}, \quad (\text{A.8})$$

where $M \in \mathbb{Z}_+$ is an integer number and $a = (a_1, a_2) \in \mathbb{Z}_M \times \mathbb{Z}_M$.

For construction of the elliptic Baxter-Belavin R -matrix [2, 3] (see also [12]) introduce the following matrix basis in $\text{Mat}(M, \mathbb{C})$:

$$T_\alpha = \exp\left(\alpha_1 \alpha_2 \frac{\pi i}{M}\right) Q^{\alpha_1} \Lambda^{\alpha_2}, \quad \alpha = (\alpha_1, \alpha_2) \in \mathbb{Z}_M \times \mathbb{Z}_M, \quad (\text{A.9})$$

where $Q, \Lambda \in \text{Mat}(M, \mathbb{C})$ are given by

$$Q_{kl} = \delta_{kl} \exp\left(\frac{2\pi i}{M} k\right), \quad \Lambda_{kl} = \delta_{k-l+1=0 \bmod M}, \quad Q^M = \Lambda^M = 1_M. \quad (\text{A.10})$$

For example, $T_0 = T_{(0,0)} = 1_M$ is the identity $M \times M$ matrix. The matrices Q and Λ satisfy relations

$$\Lambda^{a_2} Q^{a_1} = \exp\left(\frac{2\pi i}{M} a_1 a_2\right) Q^{a_1} \Lambda^{a_2}, \quad a_{1,2} \in \mathbb{Z}. \quad (\text{A.11})$$

Then $T_\alpha T_\beta = \kappa_{\alpha,\beta} T_{\alpha+\beta}$, $\kappa_{\alpha,\beta} = \exp\left(\frac{\pi i}{M} (\alpha_2 \beta_1 - \alpha_1 \beta_2)\right)$, where $\alpha + \beta = (\alpha_1 + \beta_1, \alpha_2 + \beta_2)$.

Using (A.8) and (A.9) introduce the Baxter-Belavin elliptic (\mathbb{Z}_M symmetric) R -matrix:

$$R_{12}^h(x) = \frac{1}{M} \sum_{\alpha} \varphi_{\alpha}(x, \frac{\hbar}{M} + \omega_{\alpha}) T_{\alpha} \otimes T_{-\alpha} \in \text{Mat}(M, \mathbb{C})^{\otimes 2}. \quad (\text{A.12})$$

The \mathbb{Z}_M symmetry means that

$$(Q \otimes Q) R_{12}^h(x) = R_{12}^h(x) (Q \otimes Q), \quad (\Lambda \otimes \Lambda) R_{12}^h(x) = R_{12}^h(x) (\Lambda \otimes \Lambda). \quad (\text{A.13})$$

The R -matrix (A.12) has simple poles in both variables (\hbar or z) with the following residues:

$$\text{Res}_{z=0} R_{12}^h(z) = P_{12}, \quad \text{Res}_{\hbar=0} R_{12}^h(z) = 1_M \otimes 1_M. \quad (\text{A.14})$$

The quasi-periodic behaviour on the lattice of periods is as follows:

$$R_{12}^h(z+1) = (Q^{-1} \otimes 1_M) R_{12}^h(z) (Q \otimes 1_M), \quad (\text{A.15})$$

$$R_{12}^h(z+\tau) = \exp\left(-2\pi i \frac{\hbar}{M}\right) (\Lambda^{-1} \otimes 1_M) R_{12}^h(z) (\Lambda \otimes 1_M).$$

More properties can be found in the Appendix of [13].

Proof of identities. Main idea in proving identities from this paper is formulated in the Lemma below. The same Lemma was used in [16]. Although it is widely known in the theory of elliptic functions we briefly recall its proof for readers convenience.

Lemma 4.1 *Let $f(z)$ be an entire function on elliptic curve $\Sigma_\tau = \mathbb{C}/(\mathbb{Z} \oplus \tau\mathbb{Z})$ with the following quasi-periodic properties on the lattice of periods $\mathbb{Z} \oplus \tau\mathbb{Z}$:*

$$\begin{aligned} f(z+1) &= f(z), \\ f(z+\tau) &= e^{2\pi i \alpha} f(z), \quad \alpha \in \mathbb{C}. \end{aligned} \tag{A.16}$$

Then this function is equal to zero if $\alpha \notin \mathbb{Z} \oplus \tau\mathbb{Z}$ (that is $\alpha \neq n_1 + n_2\tau$, $n_1, n_2 \in \mathbb{Z}$).

Proof. Consider the function

$$\tilde{f}(z) = e^{2\pi i \beta z} f(z), \quad \beta \in \mathbb{R}. \tag{A.17}$$

It has the following properties:

$$\begin{aligned} \tilde{f}(z+1) &= e^{2\pi i \beta} \tilde{f}(z), \\ \tilde{f}(z+\tau) &= e^{2\pi i (\alpha + \beta\tau)} \tilde{f}(z). \end{aligned} \tag{A.18}$$

By fixing the real valued parameter β as

$$\beta = -\frac{\text{Im}(\alpha)}{\text{Im}(\tau)} = -\frac{\alpha - \bar{\alpha}}{\tau - \bar{\tau}} \tag{A.19}$$

we come to

$$\begin{aligned} \tilde{f}(z+1) &= g_1 \tilde{f}(z), \quad g_1 = \exp\left(-2\pi i \frac{\alpha - \bar{\alpha}}{\tau - \bar{\tau}}\right), \\ \tilde{f}(z+\tau) &= g_\tau \tilde{f}(z), \quad g_\tau = \exp\left(2\pi i (\text{Re}(\alpha) + \beta \text{Re}(\tau))\right) = \exp\left(2\pi i \frac{\tau \bar{\alpha} - \alpha \bar{\tau}}{\tau - \bar{\tau}}\right). \end{aligned} \tag{A.20}$$

In the case $\alpha = n_1 + n_2\tau$ for some $n_1, n_2 \in \mathbb{Z}$ the function \tilde{f} becomes double periodic, i.e. $g_1 = g_\tau = 1$. Any entire double-periodic function on elliptic curve is a constant (since elliptic curve is compact). Therefore, due to (A.17), (A.19) we obtain $f(z) = \text{const} \cdot \exp(-2\pi i z \frac{\alpha - \bar{\alpha}}{\tau - \bar{\tau}})$.

If $\alpha \notin \mathbb{Z} \oplus \tau\mathbb{Z}$ then $g_1 \neq 1$ and/or $g_\tau \neq 1$. At the same time $|g_1| = |g_\tau| = 1$. Then due to the Liouville theorem (the one which says that every bounded entire holomorphic function on elliptic curve is constant) we conclude: $\tilde{f}(z) = \text{const}$. The latter constant equals zero since $g_1 \neq 1$ and/or $g_\tau \neq 1$. ■

5 Appendix B: proof of auxiliary lemmas

Proof of Lemma 2.1

Proof. Consider the l.h.s. of (2.18):

$$\mathcal{Y}_{I,J}^h R_{b,j_{n+1}}^h \cdots R_{b,j_l}^h = \overrightarrow{\prod}_{j \in J} R_{i_1,j}^h \cdots \overrightarrow{\prod}_{j \in J} R_{i_k,j}^h \cdot R_{b,j_{n+1}}^h \cdots R_{b,j_l}^h. \tag{B.1}$$

Move $R_{b,j_{n+1}}^h$ to the left using the Yang-Baxter equation:

$$\begin{aligned}
\overrightarrow{\prod}_{j \in J} R_{i_k, j}^h \cdot R_{b, j_{n+1}}^h &= \left(R_{i_k, j_1}^h \cdots R_{i_k, j_{n-1}}^h R_{i_k, b}^h R_{i_k, j_{n+1}}^h \cdots R_{i_k, j_l}^h \right) R_{b, j_{n+1}}^h = \\
&= R_{i_k, j_1}^h \cdots R_{i_k, j_{n-1}}^h R_{i_k, b}^h R_{i_k, j_{n+1}}^h R_{b, j_{n+1}}^h \cdots R_{i_k, j_l}^h \\
&= R_{i_k, j_1}^h \cdots R_{i_k, j_{n-1}}^h R_{b, j_{n+1}}^h R_{i_k, j_{n+1}}^h R_{i_k, b}^h \cdots R_{i_k, j_l}^h = \\
&= R_{b, j_{n+1}}^h R_{i_k, j_1}^h \cdots R_{i_k, j_{n-1}}^h R_{i_k, j_{n+1}}^h R_{i_k, b}^h \cdots R_{i_k, j_l}^h.
\end{aligned} \tag{B.2}$$

In the second and the last line of (B.2) we used (2.2), and in the third line we used (2.1). In the same way, by moving the product $R_{b, j_{n+1}}^h \cdots R_{b, j_l}^h$ to the left through $\overrightarrow{\prod}_{j \in J} R_{i_k, j}^h$ one obtains:

$$\overrightarrow{\prod}_{j \in J} R_{i_k, j}^h \cdot R_{b, j_{n+1}}^h \cdots R_{b, j_l}^h = R_{b, j_{n+1}}^h \cdots R_{b, j_l}^h \cdot \overrightarrow{\prod}_{\substack{j \in J \\ j \neq b}} R_{i_k, j}^h \cdot R_{i_k, b}^h. \tag{B.3}$$

By applying (B.3) to each product $\overrightarrow{\prod}_{j \in J} R_{i_l, j}^h$ in (B.1) and by moving each $R_{i_l, b}^h$ to the right we get (2.18).
■

Proof of Lemma 3.1

Proof. Arguments of R -matrices of the form $x_a - x_b$ appear in \mathcal{F}_1 , and there are no such arguments in \mathcal{F}_2 . Therefore,

$$\text{Res}_{x_a=x_b} \mathcal{F} = \text{Res}_{x_a=x_b} \mathcal{F}_1, \quad b = 1, \dots, a-1, a+1, \dots, N, \tag{B.4}$$

and we need to prove³

$$\text{Res}_{x_a=x_b} \mathcal{F}_1 = 0. \tag{B.5}$$

Consider the expression \mathcal{F}_1 (3.4). Due to the structure of poles (A.14) a nonzero input into the l.h.s. of (B.5) is provided by the terms (in \mathcal{F}_1) which contain R_{ab}^h or R_{ba}^h . It happens in two cases:

1. $a \in I$ and $b \in I^c$;
2. $b \in I$ and $a \in I^c$.

Let $a < b$. Fix some set J of $(k-1)$ elements which does not contain a and b . Denote by $J^\bullet = J^c \setminus \{a, b\}$. In the first case we choose $I = J \cup \{a\}$ and $I^c = J^\bullet \cup \{b\}$ and in the second case $I = J \cup \{b\}$ and $I^c = J^\bullet \cup \{a\}$. We are going to show that the residue of the first one is exactly cancelled out by the residue of the second one:

$$\begin{aligned}
&\text{Res}_{z_a=z_b} \mathcal{R}_{J \cup \{a\}, J^\bullet \cup \{b\}}^h \cdot \mathcal{Y}_{\mathcal{N}_2, J \cup \{a\}}^h \cdot \mathcal{R}_{J \cup \{a\}, J^\bullet \cup \{b\}}^h = \\
&- \text{Res}_{z_a=z_b} \mathcal{R}_{J \cup \{b\}, J^\bullet \cup \{a\}}^h \cdot \mathcal{Y}_{\mathcal{N}_2, J \cup \{b\}}^h \cdot \mathcal{R}_{J \cup \{b\}, J^\bullet \cup \{a\}}^h.
\end{aligned} \tag{B.6}$$

³We should notice that the structure of terms in the sum \mathcal{F}_1 excludes appearance of higher order poles. This is why we are looking for simple poles only.

In order to prove this we separate in the l.h.s. of (B.6) those R -matrices which contain index a or b using relations (2.7)-(2.8) and (2.18):

$$\mathcal{R}_{J \cup \{a\}, J \bullet \cup \{b\}}^h = \mathcal{R}_{\{a\}, J \cup J \bullet \cup \{b\}}^h \cdot \mathcal{R}_{J \cup J \bullet, \{b\}}^h \cdot \mathcal{R}_{J, J \bullet}^h \left(\mathcal{R}_{J \bullet, \{b\}}^h \right)^{-1} \left(\mathcal{R}_{\{a\}, J}^h \right)^{-1}, \quad (\text{B.7})$$

$$\mathcal{Y}_{\mathcal{N}_2, J \cup \{a\}}^h = R_{a, j_{n+1}}^h \cdots R_{a, j_{k-1}}^h \cdot \mathcal{Y}_{\mathcal{N}_2, J}^h \cdot \mathcal{Y}_{\mathcal{N}_2, \{a\}}^h \cdot \left(R_{a, j_{n+1}}^h \cdots R_{a, j_{k-1}}^h \right)^{-1}, \quad (\text{B.8})$$

$$\mathcal{R}'_{J \cup \{a\}, J \bullet \cup \{b\}}^h = \left(\mathcal{R}'_{\{a\}, J}^h \right)^{-1} \left(\mathcal{R}'_{J \bullet, \{b\}}^h \right)^{-1} \mathcal{R}'_{J, J \bullet}^h \cdot \mathcal{R}'_{J \cup J \bullet, \{b\}}^h \cdot \mathcal{R}'_{\{a\}, J \cup J \bullet \cup \{b\}}^h. \quad (\text{B.9})$$

Then using (B.7)-(B.9) we simplify the l.h.s. of (B.6) before taking the residue:

$$\begin{aligned} \mathcal{R}_{J \cup \{a\}, J \bullet \cup \{b\}}^h \mathcal{Y}_{\mathcal{N}_2, J \cup \{a\}}^h \mathcal{R}'_{J \cup \{a\}, J \bullet \cup \{b\}}^h &= \mathcal{R}_{\{a\}, J \cup J \bullet \cup \{b\}}^h \cdot \mathcal{R}_{J \cup J \bullet, \{b\}}^h \cdot \mathcal{R}_{J, J \bullet}^h \left(\mathcal{R}_{J \bullet, \{b\}}^h \right)^{-1} \\ \mathcal{Y}_{\mathcal{N}_2, J}^h \cdot \mathcal{Y}_{\mathcal{N}_2, \{a\}}^h \left(\mathcal{Y}_{\{a\}, J}^h \right)^{-1} \cdot \left(\mathcal{R}'_{J \bullet, \{b\}}^h \right)^{-1} &\mathcal{R}'_{J, J \bullet}^h \cdot \mathcal{R}'_{J \cup J \bullet, \{b\}}^h \cdot \mathcal{R}'_{\{a\}, J \cup J \bullet \cup \{b\}}^h. \end{aligned} \quad (\text{B.10})$$

Here we reduced $\left(\mathcal{R}_{\{a\}, J}^h \right)^{-1}$ from (B.7) and $R_{a, j_{n+1}}^h \cdots R_{a, j_{k-1}}^h$ from (B.8) and also used that

$$\left(R_{a, j_{n+1}}^h \cdots R_{a, j_{k-1}}^h \right)^{-1} \left(\mathcal{R}'_{\{a\}, J}^h \right)^{-1} = \left(\mathcal{Y}_{\{a\}, J}^h \right)^{-1}.$$

Next, we move $\left(\mathcal{R}_{J \bullet, \{b\}}^h \right)^{-1}$ to the right through $\mathcal{Y}_{\mathcal{N}_2, J}^h \mathcal{Y}_{\mathcal{N}_2, \{a\}}^h \left(\mathcal{Y}_{\{a\}, J}^h \right)^{-1}$ since it contains pairwise distinct indices:

$$\begin{aligned} \mathcal{R}_{J \cup \{a\}, J \bullet \cup \{b\}}^h \mathcal{Y}_{\mathcal{N}_2, J \cup \{a\}}^h \mathcal{R}'_{J \cup \{a\}, J \bullet \cup \{b\}}^h &= \mathcal{R}_{\{a\}, J \cup J \bullet \cup \{b\}}^h \cdot \mathcal{R}_{J \cup J \bullet, \{b\}}^h \cdot \mathcal{R}_{J, J \bullet}^h \\ \mathcal{Y}_{\mathcal{N}_2, J}^h \cdot \mathcal{Y}_{\mathcal{N}_2, \{a\}}^h \left(\mathcal{Y}_{\{a\}, J}^h \right)^{-1} \cdot \left(\mathcal{R}_{J \bullet, \{b\}}^h \right)^{-1} &\left(\mathcal{R}'_{J \bullet, \{b\}}^h \right)^{-1} \mathcal{R}'_{J, J \bullet}^h \cdot \mathcal{R}'_{J \cup J \bullet, \{b\}}^h \cdot \mathcal{R}'_{\{a\}, J \cup J \bullet \cup \{b\}}^h. \end{aligned} \quad (\text{B.11})$$

Now we take the residue at $z_a = z_b$ by replacing $R_{a, b}^h$ in $\mathcal{R}_{\{a\}, J \cup J \bullet \cup \{b\}}^h$ with the permutation operator P_{ab} and move it to the right:

$$\begin{aligned} \text{Res}_{z_a=z_b} \mathcal{R}_{J \cup \{a\}, J \bullet \cup \{b\}}^h \mathcal{Y}_{\mathcal{N}_2, J \cup \{a\}}^h \mathcal{R}'_{J \cup \{a\}, J \bullet \cup \{b\}}^h &= \\ R_{a, a+1}^h \cdots R_{a, b-1}^h P_{ab} R_{a, b+1}^h \cdots R_{a, N}^h \cdot R_{b-1, b}^h \cdots R_{a+1, b}^h R_{a-1, b}^h \cdots R_{1, b}^h \cdot \mathcal{R}_{J, J \bullet}^h \cdot \mathcal{Y}_{\mathcal{N}_2, J}^h \cdot \mathcal{Y}_{\mathcal{N}_2, \{a\}}^h \\ \left(\mathcal{Y}_{\{a\}, J}^h \right)^{-1} \cdot \left(\mathcal{R}_{J \bullet, \{b\}}^h \right)^{-1} \left(\mathcal{R}'_{J \bullet, \{b\}}^h \right)^{-1} &\mathcal{R}'_{J, J \bullet}^h \mathcal{R}'_{J \cup J \bullet, \{b\}}^h \mathcal{R}'_{\{a\}, J \cup J \bullet \cup \{b\}}^h = \\ = \underbrace{R_{a, a+1}^h \cdots R_{a, b-1}^h}_{\text{}} \cdot R_{b, b+1}^h \cdots R_{b, N}^h \cdot \underbrace{R_{b-1, a}^h \cdots R_{a+1, a}^h}_{\text{}} &R_{a-1, a}^h \cdots R_{1, a}^h \cdot \mathcal{R}_{J, J \bullet}^h \cdot \mathcal{Y}_{\mathcal{N}_2, J}^h \cdot \mathcal{Y}_{\mathcal{N}_2, \{b\}}^h \cdot \\ \left(\mathcal{Y}_{\{b\}, J}^h \right)^{-1} \cdot \left(\mathcal{R}_{J \bullet, \{a\}}^h \right)^{-1} \left(\mathcal{R}'_{J \bullet, \{a\}}^h \right)^{-1} &\mathcal{R}'_{J, J \bullet}^h \cdot P_{ab} \cdot \mathcal{R}'_{J \cup J \bullet, \{b\}}^h \mathcal{R}'_{\{a\}, J \cup J \bullet \cup \{b\}}^h = \\ = \prod_{j=a+1}^{b-1} \phi(h, z_a - z_j) \phi(h, z_j - z_a) \cdot \mathcal{R}_{\{b\}, J \cup J \bullet \cup \{a\}}^h &\mathcal{R}_{J \cup J \bullet, \{a\}}^h \cdot \mathcal{R}_{J, J \bullet}^h \cdot \mathcal{Y}_{\mathcal{N}_2, J}^h \cdot \mathcal{Y}_{\mathcal{N}_2, \{b\}}^h \cdot \\ \left(\mathcal{Y}_{\{b\}, J}^h \right)^{-1} \cdot \left(\mathcal{R}_{J \bullet, \{a\}}^h \right)^{-1} \left(\mathcal{R}'_{J \bullet, \{a\}}^h \right)^{-1} &\mathcal{R}'_{J, J \bullet}^h \cdot P_{ab} \cdot \mathcal{R}'_{J \cup J \bullet, \{b\}}^h \mathcal{R}'_{\{a\}, J \cup J \bullet \cup \{b\}}^h. \end{aligned} \quad (\text{B.12})$$

In the last equality we reduced the underbraced factors which give $\prod_{j=a+1}^{b-1} \phi(h, z_a - z_j) \phi(h, z_j - z_a)$ due to (2.9). Also, following the definitions (2.3) and (2.4) we performed transformations $R_{b,b+1}^h \dots R_{b,N}^h = \mathcal{R}_{\{b\}, J \cup J^\bullet \cup \{a\}}^h$ and $R_{a-1,a}^h \dots R_{1,a}^h = \mathcal{R}_{J \cup J^\bullet, \{a\}}^h$.

In a similar way we transform the r.h.s. of (B.6):

$$\begin{aligned} \text{Res}_{z_a=z_b} \mathcal{R}_{J \cup \{b\}, J^\bullet \cup \{a\}}^h \mathcal{Y}_{\mathcal{N}_2, J \cup \{b\}}^h \mathcal{R}'_{J \cup \{b\}, J^\bullet \cup \{a\}}^h &= \text{Res}_{z_a=z_b} \mathcal{R}_{\{b\}, J \cup J^\bullet \cup \{a\}}^h \cdot \mathcal{R}_{J \cup J^\bullet, \{a\}}^h \cdot \mathcal{R}_{J, J^\bullet}^h \\ \mathcal{Y}_{\mathcal{N}_2, J}^h \cdot \mathcal{Y}_{\mathcal{N}_2, \{b\}}^h \left(\mathcal{Y}_{\{b\}, J}^h \right)^{-1} \cdot \left(\mathcal{R}_{J^\bullet, \{a\}}^h \right)^{-1} \left(\mathcal{R}'_{J^\bullet, \{a\}}^h \right)^{-1} \mathcal{R}'_{J, J^\bullet}^h \cdot \mathcal{R}_{J \cup J^\bullet, \{a\}}^h \cdot \mathcal{R}_{\{b\}, J \cup J^\bullet \cup \{a\}}^h &= \\ = -\mathcal{R}_{\{b\}, J \cup J^\bullet \cup \{a\}}^h \cdot \mathcal{R}_{J \cup J^\bullet, \{a\}}^h \cdot \mathcal{R}_{J, J^\bullet}^h \cdot \mathcal{Y}_{\mathcal{N}_2, J}^h \cdot \mathcal{Y}_{\mathcal{N}_2, \{b\}}^h \left(\mathcal{Y}_{\{b\}, J}^h \right)^{-1} \cdot \left(\mathcal{R}_{J^\bullet, \{a\}}^h \right)^{-1} \left(\mathcal{R}'_{J^\bullet, \{a\}}^h \right)^{-1} \mathcal{R}'_{J, J^\bullet}^h & \\ R_{N,a}^h \dots R_{b+1,a}^h R_{b-1,a}^h \dots R_{a+1,a}^h \cdot R_{b,1}^h \dots R_{b,a-1}^h \cdot P_{ab} \cdot R_{b,a+1}^h \dots R_{b,b-1}^h & \end{aligned} \quad (\text{B.13})$$

and move permutation P_{ab} to the left:

$$\begin{aligned} \text{Res}_{z_a=z_b} \mathcal{R}_{J \cup \{b\}, J^\bullet \cup \{a\}}^h \mathcal{Y}_{\mathcal{N}_2, J \cup \{b\}}^h \mathcal{R}'_{J \cup \{b\}, J^\bullet \cup \{a\}}^h &= -\mathcal{R}_{\{b\}, J \cup J^\bullet \cup \{a\}}^h \cdot \mathcal{R}_{J \cup J^\bullet, \{a\}}^h \cdot \mathcal{R}_{J, J^\bullet}^h \\ \mathcal{Y}_{\mathcal{N}_2, J}^h \cdot \mathcal{Y}_{\mathcal{N}_2, \{b\}}^h \left(\mathcal{Y}_{\{b\}, J}^h \right)^{-1} \cdot \left(\mathcal{R}_{J^\bullet, \{a\}}^h \right)^{-1} \left(\mathcal{R}'_{J^\bullet, \{a\}}^h \right)^{-1} \mathcal{R}'_{J, J^\bullet}^h \cdot P_{ab} & \\ R_{N,b}^h \dots R_{b+1,b}^h \underbrace{R_{b-1,b}^h \dots R_{a+1,b}^h}_{\text{}} \cdot R_{a,1}^h \dots R_{a,a-1}^h \cdot \underbrace{R_{b,a+1}^h \dots R_{b,b-1}^h}_{\text{}} &= \\ -\mathcal{R}_{\{b\}, J \cup J^\bullet \cup \{a\}}^h \cdot \mathcal{R}_{J \cup J^\bullet, \{a\}}^h \cdot \mathcal{R}_{J, J^\bullet}^h \cdot \mathcal{Y}_{\mathcal{N}_2, J}^h \cdot \mathcal{Y}_{\mathcal{N}_2, \{b\}}^h \left(\mathcal{Y}_{\{b\}, J}^h \right)^{-1} \cdot \left(\mathcal{R}_{J^\bullet, \{a\}}^h \right)^{-1} \left(\mathcal{R}'_{J^\bullet, \{a\}}^h \right)^{-1} & \\ \mathcal{R}'_{J, J^\bullet}^h \cdot P_{ab} \cdot \mathcal{R}'_{J \cup J^\bullet, \{b\}}^h \mathcal{R}'_{\{a\}, J \cup J^\bullet \cup \{b\}}^h \prod_{j=a+1}^{b-1} \phi(h, z_a - z_j) \phi(h, z_j - z_a). & \end{aligned} \quad (\text{B.14})$$

In the last equality we reduced the underbraced factors which give $\prod_{j=a+1}^{b-1} \phi(h, z_a - z_j) \phi(h, z_j - z_a)$ due to (2.9) and used the definitions (2.5) and (2.6). The r.h.s. of (B.14) equals minus r.h.s. of (B.12). This finishes the proof. \blacksquare

Proof of Lemma 3.2

Proof. The proof is performed for \mathcal{F}_1 and \mathcal{F}_2 separately, that is the relation (3.12) holds true for \mathcal{F}_1 and \mathcal{F}_2 separately. The poles at $z_a = z_b$ are contained in the middle products $\mathcal{Y}_{\mathcal{N}_2, I}^h$ and $\mathcal{Y}_{\mathcal{N}_1, J}^{-h}$ in (3.4)-(3.5) for such I and J that $a \in I$ and $b \in J$.

The calculation for \mathcal{F}_1 is straightforward. For any set I which contains a the product $\mathcal{Y}_{\mathcal{N}_2, I}^h$ has only one item with R_{ba}^h and thus has simple pole at $z_a = z_b$. Due to the property (A.14) one should replace R_{ba} with the permutation $-P_{ab}$ and put down $z_a = z_b$. To transform the residue to the form (3.12) we move the permutation operator to the right.

Firstly, we rewrite each item in \mathcal{F}_1 separating R -matrices containing indices a and b . For this purpose relations (2.7)-(2.8) and (2.18)-(2.19) are used. Let us fix

$$I = \{i_1 < \dots < i_{m-1} < i_m = a < i_{m+1} < \dots < i_k\}$$

and transform \mathcal{R}_{I,I^c}^h using (2.7):

$$\begin{aligned}\mathcal{R}_{I,I^c}^h &= \mathcal{R}_{\{a\},\mathcal{N}_1 \setminus \{a\}}^h \cdot \mathcal{R}_{I \setminus \{a\},I^c}^h \cdot \left(\mathcal{R}_{\{a\},I \setminus \{a\}}^h \right)^{-1} = \\ &= R_{a,a+1}^h \cdots R_{a,N}^h \cdot \mathcal{R}_{I \setminus \{a\},I^c}^h \cdot \left(R_{a,i_{m+1}}^h \cdots R_{a,i_k}^h \right)^{-1}.\end{aligned}\tag{B.15}$$

Similar transformations are performed using (2.8):

$$\begin{aligned}\mathcal{R}'_{I,I^c}^h &= \left(\mathcal{R}'_{\{a\},I \setminus \{a\}}^h \right)^{-1} \cdot \mathcal{R}'_{I \setminus \{a\},I^c}^h \cdot \mathcal{R}'_{\{a\},\mathcal{N}_1 \setminus \{a\}}^h = \\ &= \left(R_{a,i_1}^h \cdots R_{a,i_{m-1}}^h \right)^{-1} \cdot \mathcal{R}'_{I \setminus \{a\},I^c}^h \cdot R_{a,1}^h \cdots R_{a,a-1}^h.\end{aligned}\tag{B.16}$$

By applying consistently (2.18)-(2.19) to $\mathcal{Y}_{\mathcal{N}_2,I}^h$ the latter is rewritten as follows:

$$\begin{aligned}\mathcal{Y}_{\mathcal{N}_2,I}^h &= R_{a,i_{m+1}}^h \cdots R_{a,i_k}^h \cdot R_{b+1,b}^h \cdots R_{2N,b}^h \cdot \mathcal{Y}_{\mathcal{N}_2 \setminus \{b\},I \setminus \{a\}}^h \cdot \mathcal{Y}_{\{b\},I \setminus \{a\}}^h \cdot \mathcal{Y}_{\mathcal{N}_2 \setminus \{b\},\{a\}}^h \cdot \\ &\cdot R_{ba}^h \cdot \left(R_{a,i_{m+1}}^h \cdots R_{a,i_k}^h \cdot R_{b+1,b}^h \cdots R_{2N,b}^h \right)^{-1}.\end{aligned}\tag{B.17}$$

Let us calculate the residue of \mathcal{F}_1 at $z_a = z_b$. The nontrivial part is $\mathcal{Y}_{\mathcal{N}_2,I}^h$, while the items \mathcal{R}_{I,I^c}^h and \mathcal{R}'_{I,I^c}^h has no pole. Plugging

$$\mathcal{Y}_{\{b\},I \setminus \{a\}}^h = R_{b,i_1}^h \cdots R_{b,i_{m-1}}^h R_{b,i_{m+1}}^h \cdots R_{b,i_k}^h$$

and

$$\mathcal{Y}_{\mathcal{N}_2 \setminus \{b\},\{a\}}^h = R_{N+1,a}^h \cdots R_{b-1,a}^h R_{b+1,a}^h \cdots R_{2N,a}^h$$

into (B.17) one obtains:

$$\begin{aligned}\text{Res}_{z_a=z_b} \mathcal{Y}_{\mathcal{N}_2,I}^h &= -R_{a,i_{m+1}}^h \cdots R_{a,i_k}^h \cdot R_{b+1,b}^h \cdots R_{2N,b}^h \cdot \mathcal{Y}_{\mathcal{N}_2 \setminus \{b\},I \setminus \{a\}}^h \cdot \\ &R_{b,i_1}^h \cdots R_{b,i_{m-1}}^h R_{b,i_{m+1}}^h \cdots R_{b,i_k}^h \cdot R_{N+1,a}^h \cdots R_{b-1,a}^h R_{b+1,a}^h \cdots R_{2N,a}^h \cdot \\ &P_{ab} \left(R_{a,i_{m+1}}^h \cdots R_{a,i_k}^h \cdot R_{b+1,b}^h \cdots R_{2N,b}^h \right)^{-1}.\end{aligned}\tag{B.18}$$

In order to simplify we move the permutation operator P_{ab} to the right and then reduce the underlined items:

$$\begin{aligned}\text{Res}_{z_a=z_b} \mathcal{Y}_{\mathcal{N}_2,I}^h &= -R_{a,i_{m+1}}^h \cdots R_{a,i_k}^h \cdot R_{b+1,b}^h \cdots R_{2N,b}^h \cdot \mathcal{Y}_{\mathcal{N}_2 \setminus \{b\},I \setminus \{a\}}^h \cdot \\ &R_{b,i_1}^h \cdots R_{b,i_{m-1}}^h \underline{R_{b,i_{m+1}}^h \cdots R_{b,i_k}^h} \cdot R_{N+1,a}^h \cdots R_{b-1,a}^h \underline{R_{b+1,a}^h \cdots R_{2N,a}^h} \cdot \\ &\left(\underline{R_{b,i_{m+1}}^h \cdots R_{b,i_k}^h \cdot R_{b+1,a}^h \cdots R_{2N,a}^h} \right)^{-1} P_{ab} = \\ &-R_{a,i_{m+1}}^h \cdots R_{a,i_k}^h \cdot R_{b+1,b}^h \cdots R_{2N,b}^h \cdot \mathcal{Y}_{\mathcal{N}_2 \setminus \{b\},I \setminus \{a\}}^h \cdot R_{b,i_1}^h \cdots R_{b,i_{m-1}}^h \cdot R_{N+1,a}^h \cdots R_{b-1,a}^h \cdot P_{ab}.\end{aligned}\tag{B.19}$$

Indeed, expression $R_{b,i_{m+1}}^h \cdots R_{b,i_k}^h$ commutes with $R_{N+1,a}^h \cdots R_{b-1,a}^h$ due to (2.2) since each multiplier has pairwise different indices. That is the underlined terms are cancelled.

Now we are going to calculate the residue at $z_a = z_b$ of each summand containing index $a \in I$ in \mathcal{F}_1 given in (3.4). We use (B.15), (B.16) and (B.19):

$$\text{Res}_{z_a=z_b} \mathcal{R}_{I,I^c}^h \cdot \mathcal{Y}_{\mathcal{N}_2,I}^h \cdot \mathcal{R}_{I,I^c}^h = -R_{a,a+1}^h \cdots R_{a,N}^h \cdot \mathcal{R}_{I \setminus \{a\},I^c}^h \cdot \underbrace{R_{b+1,b}^h \cdots R_{2N,b}^h}_{\mathcal{Y}_{\mathcal{N}_2 \setminus \{b\},I \setminus \{a\}}^h} \cdot \mathcal{Y}_{\mathcal{N}_2 \setminus \{b\},I \setminus \{a\}}^h. \quad (\text{B.20})$$

$$R_{b,i_1}^h \cdots R_{b,i_{m-1}}^h \cdot R_{N+1,a}^h \cdots R_{b-1,a}^h \cdot P_{ab} \cdot \left(R_{a,i_1}^h \cdots R_{a,i_{m-1}}^h \right)^{-1} \cdot \mathcal{R}_{I \setminus \{a\},I^c}^h \cdot R_{a,1}^h \cdots R_{a,a-1}^h.$$

Next, we move P_{ab} to the right and the underbraced item to the left through $\mathcal{R}_{I \setminus \{a\},I^c}^h$ due to (2.15) since it does not contain indices $b, b+1, \dots, 2N$:

$$\begin{aligned} \text{Res}_{z_a=z_b} \mathcal{R}_{I,I^c}^h \cdot \mathcal{Y}_{\mathcal{N}_2,I}^h \cdot \mathcal{R}_{I,I^c}^h &= -R_{a,a+1}^h \cdots R_{a,N}^h \cdot R_{b+1,b}^h \cdots R_{2N,b}^h \cdot \mathcal{R}_{I \setminus \{a\},I^c}^h \cdot \mathcal{Y}_{\mathcal{N}_2 \setminus \{b\},I \setminus \{a\}}^h \\ \underline{R_{b,i_1}^h \cdots R_{b,i_{m-1}}^h} \cdot R_{N+1,a}^h \cdots R_{b-1,a}^h \cdot \left(\underline{R_{a,i_1}^h \cdots R_{a,i_{m-1}}^h} \right)^{-1} \cdot \mathcal{R}_{I \setminus \{a\},I^c}^h \cdot R_{b,1}^h \cdots R_{b,a-1}^h \cdot P_{ab} &= \\ = -G \cdot \mathcal{R}_{I \setminus \{a\},I^c}^h \cdot \mathcal{Y}_{\mathcal{N}_2 \setminus \{b\},I \setminus \{a\}}^h \cdot R_{N+1,a}^h \cdots R_{b-1,a}^h \cdot \mathcal{R}_{I \setminus \{a\},I^c}^h \cdot R_{b,1}^h \cdots R_{b,a-1}^h \cdot P_{ab}. \end{aligned} \quad (\text{B.21})$$

In the last equality we combined the first terms into G using the definition (3.13) and reduced the underlined items. Then moving $R_{N+1,a}^h \cdots R_{b-1,a}^h$ through $\mathcal{R}_{I \setminus \{a\},I^c}^h$ to the right and using the definition (3.14) one obtains:

$$\text{Res}_{z_a=z_b} \mathcal{R}_{I,I^c}^h \cdot \mathcal{Y}_{\mathcal{N}_2,I}^h \cdot \mathcal{R}_{I,I^c}^h = -G \cdot \mathcal{R}_{I \setminus \{a\},I^c}^h \cdot \mathcal{Y}_{\mathcal{N}_2 \setminus \{b\},I \setminus \{a\}}^h \cdot \mathcal{R}_{I \setminus \{a\},I^c}^h \cdot H. \quad (\text{B.22})$$

This yields

$$\text{Res}_{z_a=z_b} \mathcal{F}_1[k, N] = -G \cdot \mathcal{F}_1[k-1, N-1] \cdot H. \quad (\text{B.23})$$

The same calculation can be performed for \mathcal{F}_2 since it differs from \mathcal{F}_1 by only changing I to J , \mathcal{N}_2 to \mathcal{N}_1 , $\hbar \rightarrow -\hbar$ and the total sign. Therefore, we use the result of (B.22):

$$\text{Res}_{z_a=z_b} \mathcal{R}_{J,J^c}^{-\hbar} \mathcal{Y}_{\mathcal{N}_1,J}^{-\hbar} \mathcal{R}_{J,J^c}^{-\hbar} = -G' \cdot \mathcal{R}_{J \setminus \{b\},J^c}^{-\hbar} \cdot \mathcal{Y}_{\mathcal{N}_1 \setminus \{a\},J \setminus \{b\}}^{-\hbar} \cdot \mathcal{R}_{J \setminus \{b\},J^c}^{-\hbar} \cdot H', \quad (\text{B.24})$$

where

$$G' = R_{b,b+1}^{-\hbar} \cdots R_{b,2N}^{-\hbar} \cdot R_{a+1,a}^{-\hbar} \cdots R_{N,a}^{-\hbar}, \quad (\text{B.25})$$

$$H' = R_{1,b}^{-\hbar} \cdots R_{a-1,b}^{-\hbar} \cdot R_{a,N+1}^{-\hbar} \cdots R_{a,b-1}^{-\hbar} P_{ab}. \quad (\text{B.26})$$

We change each $R_{i,j}^{-\hbar}$ to $-R_{j,i}^{\hbar}$ in (B.25) and (B.26) using (2.15):

$$G' = (-1)^{3N-a-b} R_{b+1,b}^{\hbar} \cdots R_{2N,b}^{\hbar} \cdot R_{a,a+1}^{\hbar} \cdots R_{a,N}^{\hbar} = (-1)^{a+b-N} G, \quad (\text{B.27})$$

$$H' = (-1)^{a-1+b-N-1} R_{b,1}^{\hbar} \cdots R_{b,a-1}^{\hbar} \cdot R_{N+1,a}^{\hbar} \cdots R_{b-1,a}^{\hbar} P_{ab} = (-1)^{a+b-N-1} H. \quad (\text{B.28})$$

Due to (B.27)-(B.28) one can change G' in (B.24) to G and H' to H thus providing

$$\text{Res}_{z_a=z_b} \mathcal{F}_2[k, N] = -G \cdot \mathcal{F}_2[k-1, N-1] \cdot H. \quad (\text{B.29})$$

This finishes the proof. ■

Acknowledgments

We are grateful to A. Levin, A. Liashyk and S. Khoroshkin for useful discussions. The work of M. Matushko was partially supported by Russian Science Foundation (project 20-41-09009).

References

- [1] F. Atai, M. Hallnäs, E. Langmann, *Source identities and kernel functions for deformed (quantum) Ruijsenaars models*, Lett. Math. Phys. 104 (2014) 811–835; arXiv:1311.4433 [math-ph].
M. Hallnäs, E. Langmann, M. Noumi, H. Rosengren, *From Kajihara’s transformation formula to deformed Macdonald-Ruijsenaars and Noumi-Sano operators*, Selecta Mathematica New Series (2022) 28:24; arXiv:2105.01936 [math.QA].
- [2] R.J. Baxter, *Partition function of the eight-vertex lattice model*, Ann. Phys. 70 (1972) 193–228.
- [3] A.A. Belavin, *Dynamical symmetry of integrable quantum systems*, Nucl. Phys. B, 180 (1981) 189–200.
- [4] S. Fomin, A.N. Kirillov, *Quadratic algebras, Dunkl elements, and Schubert calculus*, Advances in geometry; Prog. in Mathematics book series, 172 (1999) 147–182.
A. Polishchuk, *Classical Yang–Baxter equation and the A^∞ -constraint*, Advances in Mathematics, 168:1 (2002) 56—95; arXiv:math/0008156 [math.AG].
- [5] A. Grekov, I. Sechin, A. Zotov, *Generalized model of interacting integrable tops*, JHEP 10 (2019) 081; arXiv:1905.07820 [math-ph].
E.S. Trunina, A.V. Zotov, *Multi-pole extension of the elliptic models of interacting integrable tops*, Theoret. and Math. Phys., 209:1 (2021), 1331–1356, arXiv:2104.08982 [math-ph].
- [6] M. Hallnäs, S. Ruijsenaars, *Kernel functions and Bäcklund transformations for relativistic Calogero-Moser and Toda systems*, J. Math. Phys. 53, 123512 (2012); arXiv:1206.3786 [nlin.SI].
- [7] S. Kharchev, S. Khoroshkin, *Wave function for $GL(n, R)$ hyperbolic Sutherland model II. Dual Hamiltonians*, arXiv:2108.05393 [math.RT].
- [8] T. Krasnov, A. Zotov, *Trigonometric integrable tops from solutions of associative Yang-Baxter equation*, Annales Henri Poincaré, 20:8 (2019) 2671–2697; arXiv:1812.04209 [math-ph].
- [9] Y. Komori, M. Noumi, J. Shiraishi, *Kernel Functions for Difference Operators of Ruijsenaars Type and Their Applications*, SIGMA 5 (2009), 054; arXiv:0812.0279 [math.QA].
- [10] J. Lamers, *Resurrecting the partially isotropic Haldane-Shastry model*, Phys. Rev. B 97 (2018) 214416; arXiv:1801.05728 [cond-mat.str-el].
J. Lamers, V. Pasquier, D. Serban, *Spin-Ruijsenaars, q -deformed Haldane-Shastry and Macdonald polynomials*, arXiv:2004.13210 [math-ph].
- [11] A. Levin, M. Olshanetsky, A. Zotov, *Relativistic classical integrable tops and quantum R -matrices*, JHEP 07 (2014) 012; arXiv:1405.7523 [hep-th].
- [12] A.M. Levin, M.A. Olshanetsky, A.V. Zotov, *Quantum Baxter-Belavin R -matrices and multidimensional Lax pairs for Painlevé VI*, Theoret. and Math. Phys. 184:1 (2015) 924–939; arXiv:1501.07351 [math-ph].
- [13] M. Matushko, A. Zotov, *Anisotropic spin generalization of elliptic Macdonald-Ruijsenaars operators and R -matrix identities*, arXiv:2201.05944 [math.QA].
- [14] M. Matushko, A. Zotov, *Elliptic generalization of integrable q -deformed Haldane-Shastry long-range spin chain*, arXiv:2202.01177 [math-ph].
I. Sechin, A. Zotov, *R -matrix-valued Lax pairs and long-range spin chains*, Physics Letters B, 781:10 (2018) 1–7; arXiv:1801.08908 [math-ph].
- [15] S.N.M. Ruijsenaars, *Complete integrability of relativistic Calogero-Moser systems and elliptic function identities*, Commun. Math. Phys. 110:2 (1987) 191–213.

- [16] S. Ruijsenaars, *Zero-Eigenvalue Eigenfunctions for Differences of Elliptic Relativistic Calogero–Moser Hamiltonians*, Theoret. and Math. Phys., 146:1 (2006), 25—33.
- [17] A. Zabrodin, A. Zotov, *Self–dual form of Ruijsenaars–Schneider models and ILW equation with discrete Laplacian*, Nuclear Physics B, 927 (2018), 550–565; arXiv:1711.01036 [math-ph].
- [18] A.V. Zotov, *Calogero–Moser model and R-matrix identities*, Theoret. and Math. Phys. 197:3 (2018) 1755–1770.
- [19] A.V. Zotov, *Relativistic interacting integrable elliptic tops*, Theoret. and Math. Phys., 201:2 (2019) 1563–1578, arXiv: 1910.08246.
I.A. Sechin, A.V. Zotov, *Integrable system of generalized relativistic interacting tops*, Theoret. and Math. Phys., 205:1 (2020), 1292–1303; arXiv:2011.09599 [math-ph].
E. Trunina, A. Zotov, *Lax equations for relativistic $GL(NM, C)$ Gaudin models on elliptic curve*, arXiv: 2204.06137 [nlin.SI].