The combinatorial code and the graph rules of Dale networks

Nikola Milićević and Vladimir Itskov

Department of Mathematics, The Pennsylvania State University, University Park, PA 16802.

Abstract

We describe the combinatorics of equilibria and steady states of neurons in threshold-linear networks that satisfy the Dale's law. The combinatorial code of a Dale network is characterized in terms of two conditions: (i) a condition on the network connectivity graph, and (ii) a spectral condition on the synaptic matrix. We find that in the weak coupling regime the combinatorial code depends only on the connectivity graph, and not on the particulars of the synaptic strengths. Moreover, we prove that the combinatorial code of a weakly coupled network is a sublattice, and we provide a learning rule for encoding a sublattice in a weakly coupled excitatory network. In the strong coupling regime we prove that the combinatorial code of a generic Dale network is intersection-complete and is therefore a convex code, as is common in some sensory systems in the brain.

Key words. Dale's law, recurrent neural networks, combinatorial neural codes *MSC codes.* 92B20

Contents

1	Introduction	1
2	Preliminaries: The combinatorial codes of Dale networks	2
3		$\frac{4}{6}$
4	Discussion	9
5	Proofs 5.1 Necessary matrix theory results 5.2 The proofs	
6	Appendix: Stability of Linear-Threshold Rate Dynamics	20

1 Introduction

We study the equilibria and steady states of threshold-linear networks that satisfy the Dale's law. Threshold-linear networks have been extensively studied in [13, 24, 12, 7, 5, 9, 19], and a number of results regarding the stable fixed points of these networks were obtained, especially in the case of a symmetric synaptic matrix. Networks that satisfy the Dale's law have been previously investigated in the context of large random networks [21, 1, 15] (this list is very incomplete), where

the statistical properties such as the spectrum, the number of equilibria and phase transitions have been investigated. In contrast, we are interested in describing the exact combinatorics of the neural code of these networks, as opposed to investigating statistical features.

To this end we developed a method for understanding the combinatorial codes of threshold-linear networks that obey the Dale's law. A combinatorial code is the collection of patterns of neuronal activation at the equilibria that is possible on a given network. In particular, this discards the details of the firing rates, and only keeps track of what neurons are co-active. It turns out its possible to directly translate features of the connectivity graph into the combinatorial code of a Dale network. We show that the connectivity features completely determine the combinatorial neural code in the weak coupling regime. In the strong coupling regime the code is described in terms of connectivity and certain spectral conditions on the excitatory subnetworks. We also show that these combinatorial codes are *convex*, that is they are compatible with patterns of overlaps of convex receptive fields that are common in many sensory systems of the brain [4].

This paper is organized as follows. In Section 2 we introduce the necessary background for threshold-linear networks and Dale's law and define the combinatorial and stable combinatorial codes. In Section 3 we state our main results, while Section 4 has a brief discussion. The proofs are relegated to Section 5. Finally, Section 6 contains some necessary results concerning the stability of fixed points of threshold-linear networks.

2 Preliminaries: The combinatorial codes of Dale networks

We consider a standard firing rate model of a recurrent neural network of n neurons, where the dynamics of the firing rates $x_i(t) \ge 0$ is described by the differential equations

$$\dot{x}_i = -x_i + \left[\sum_{j=1}^n W_{ij}x_j + b_i\right]_+, \quad i = 1, \dots, n,$$
 (1)

and $[y]_{+} = \max(0, y)$ is the ReLU transfer function.

We assume that this excitatory-inhibitory network respects the *Dale's law* [10], whereas the neurons are either excitatory (denoted as \mathcal{E}) or inhibitory (denoted as \mathcal{I}), and the synaptic weights satisfy the following sign constrains:

excitatory synapses:
$$i \in \mathcal{E} \implies W_{ij} \ge 0, \quad \forall j = 1 \dots n$$

inhibitory synapses: $i \in \mathcal{I} \implies W_{ij} \le 0, \quad \forall j = 1 \dots n$.

Furthermore, we assume that the diagonal entries of W are zeros, $W_{ii} = 0$ and denote the collection of all $n \times n$ Dale matrices by \mathbb{D}_n .

Following a common architecture of the neocortex, we also assume that the excitatory neurons "broadcast" the output, while the activity of the inhibitory neurons is not observable *directly* outside of the network. We thus consider the setup where the inputs to the network are excitatory, while only the excitatory output can be "read" from the network (Figure 1). We shall call such a network *a Dale network*.




Figure 1: A Dale recurrent network.

We are interested in the combinatorics of the excitatory output of Dale networks. A combinatorial code is the set of possible patterns of neural activation at the fixed points (or steady states). For a firing rate vector $x \in \mathbb{R}^n_{\geq 0}$, we consider the *excitatory support*, i.e. the set of active excitatory neurons:

$$\operatorname{supp}_+ x = \{i \in \mathcal{E} \mid x_i > 0\} \subset \mathcal{E}.$$

Recall that $x^* \in \mathbb{R}^n$ is a fixed point of a network (1) if $x(t) = x^*$ is a constant solution. For a given Dale network (1), we denote the set of excitatory supports of all the possible fixed points as

$$\operatorname{FP}_{+}(W, b) = \left\{ \operatorname{supp}_{+} x^{*} \mid x^{*} \in \mathbb{R}_{\geq 0}^{n} \text{ is a fixed point of } (1) \right\}$$

Here the plus sign highlights the difference from a somewhat different definition in [18], which considers the combinatorics of all fixed points, which were previously investigated.

The *combinatorial code* of a Dale synaptic matrix W is the collection of all possible excitatory support in all fixed points in response to all possible inputs:

$$\mathcal{C}(W) \stackrel{\text{def}}{=} \bigcup_{b \in \mathbb{R}^n_{\ge 0}} FP_+(W, b).$$
(2)

The *stable combinatorial code* is the set of the excitatory supports of asymptotically stable fixed points:

 $\mathcal{SC}(W) \stackrel{\text{def}}{=} \bigcup_{b \in \mathbb{R}^n_{\geq 0}} \{ \sup_{y \in \mathbb{R}^n} x^* \mid x^* \in \mathbb{R}^n_{\geq 0} \text{ is an asymptotically stable fixed point} \}.$

It turns out that the combinatorial codes of Dale networks can be completely described in terms of connectivity and spectral radius of the synaptic matrix. We also show that the combinatorial code is always intersection complete, which implies it is also always a convex combinatorial code [4].

3 The main results

Here we state the main results, while all the proofs of the theorems are provided in Section 5. To simplify the mathematical analyses of the network, we make the following mild assumption.

Ground Assumption. The synaptic matrix W of the network (1) satisfies the condition that for every non-empty subset of neurons $\sigma \subset [n]$ the principal submatrix $(I - W)_{\sigma}$ is non-singular.

Note that for a square $n \times n$ matrix A and a subset $\sigma \subset [n]$ we denote by A_{σ} the appropriate principal submatrix. In all our results the Ground Assumption is always implicitly assumed. Note that the set of matrices that do *not* have this property has measure zero, thus this assumption is generically satisfied in any network without fine-tuning.

3.1 The role of excitatory-inhibitory connectivity in shaping the combinatorial code

We first observe that to understand the combinatorial code, one can streamline the excitatoryinhibitory connectivity to its "essential features" as follows.

Theorem 3.1. Let $W \in \mathbb{D}_n$ be a Dale matrix with a set of excitatory neurons \mathcal{E} and inhibitory neurons \mathcal{I} . Let $m = |\mathcal{E}| + 1$ and let $W' \in \mathbb{D}_m$ be any Dale matrix such that $W'_{\mathcal{E}} = W_{\mathcal{E}}$ and for all $i \in \mathcal{E}$,

$$W'_{im} = \begin{cases} -1, & \exists j \in \mathcal{I} \text{ with } W_{ij} \neq 0, \\ 0, & \forall j \in \mathcal{I} \text{ with } W_{ij} = 0. \end{cases}$$

Then $\mathcal{C}(W) = \mathcal{C}(W')$.

In other words, the combinatorial code $\mathcal{C}(W)$ remains unchanged if we replace all the inhibitory neurons with a single inhibitory neuron that mimics the connectivity of the entire inhibitory population to each excitatory neuron. Note that here the numerical values of the inhibitory-excitatory weights of the synaptic matrix W do not affect the combinatorial code, even though they may determine the stability of the appropriate fixed points. Furthermore, the following result states that the connections from excitatory to inhibitory neurons have no influence over the combinatorial code $\mathcal{C}(W)$, while they still may determine the stability of the appropriate fixed points, as well as other dynamical properties.

Theorem 3.2. Let $W \in \mathbb{D}_n$ be a Dale matrix with a set of excitatory neurons \mathcal{E} and inhibitory neurons \mathcal{I} , and let $W' \in \mathbb{D}_n$ be such that

$$W'_{ij} = \begin{cases} 0, & \forall i \in \mathcal{I}, \, \forall j \in \mathcal{E} \\ W_{ij}, & \text{otherwise} \end{cases}$$

Then $\mathcal{C}(W) = \mathcal{C}(W')$.

3.2 The characterization of the combinatorial code of a Dale network

Theorems 3.1 and 3.2 imply that only excitatory - excitatory synapses and a particular feature of the inhibitory-excitatory connectivity play a role in shaping the combinatorial code. To describe these features we make the following definitions.

Definition 3.3. Let W be a Dale matrix, and \mathcal{E} and \mathcal{I} denote the set of its excitatory and inhibitory neurons respectively.

• An excitatory connectivity graph of W is a directed graph $G_{\mathcal{E}}$ whose vertices are the excitatory neurons \mathcal{E} , and whose arcs are defined as

$$i \to j \iff W_{ii} \neq 0$$
, where $i \neq j \in \mathcal{E}$.

• The un-inhibited set $\beta \subset \mathcal{E}$ is the subset of excitatory neurons that do not receive any recurrent inhibition, i.e.

$$\beta \stackrel{\text{def}}{=} \{ j \in \mathcal{E} \mid \forall i \in \mathcal{I}, W_{ji} = 0 \}.$$

• For a directed graph G on a set of vertices \mathcal{E} and a subset $\beta \subset \mathcal{E}$, we define a code of a pair (G, β) as

$$\operatorname{code}(G,\beta) \stackrel{\text{def}}{=} \{ \sigma \subset \mathcal{E} \mid t(\sigma) \cap \beta \subset \sigma \},\$$

where

$$t(\sigma) \stackrel{\text{def}}{=} \bigcup_{i \in \sigma} \left\{ j \in \mathcal{E} \mid i \to j \right\}$$

denotes the set of all the synaptic targets of a subset of excitatory neurons σ .

The following theorem describes the combinatorial code of a Dale network.

Theorem 3.4. Let W be a Dale matrix, and $\sigma \subset \mathcal{E}$ be a non-empty subset of excitatory neurons. Then $\sigma \in \mathcal{C}(W)$ if and only if the following two conditions are both satisfied:

- (i) (the spectral condition) $\rho(W_{\beta_{\sigma}}) < 1$,
- (ii) (the graph condition) $\sigma \in \operatorname{code}(G_{\mathcal{E}}, \beta)$,

where $W_{\beta\sigma}$ denotes the synaptic weights of the excitatory sub-network on the subset $\beta_{\sigma} = \beta \cap \sigma$, and $\rho(W_{\beta\sigma})$ denotes the spectral radius of the matrix $W_{\beta\sigma}$.

We say that the network (1) is weakly coupled if the Frobenius matrix norm

$$||W||_F = \sqrt{\operatorname{trace}(W^T W)}$$

of its synaptic matrix W is smaller than 1. It is natural to consider the weak coupling regime separately, as in this regime the spectral condition in Theorem 3.4 is always satisfied.

Theorem 3.5. Let W be a nonsingular Dale matrix that is weakly coupled, i.e. $||W||_F < 1$. Then every fixed point is a steady state, and the combinatorial code is completely described by the graph condition:

$$\mathcal{C}(W) = \mathcal{SC}(W) = \operatorname{code}(G_{\mathcal{E}}, \beta).$$

Furthermore, for all $b \in \mathbb{R}^n_{\geq 0}$ there is a unique globally exponentially stable fixed point of (1).

The above theorem implies that the combinatorial code of weakly coupled networks is completely determined by the connectivity features *alone*, and does not depend on the strengths of the synapses as long as the network is in the weak coupling regime. The following cautionary example illustrates two Dale matrices that are *not* weakly coupled. These matrices have the same connectivity, but they

exhibit different eigenvalue spectra of excitatory subnetworks, resulting in different combinatorial codes. Consider Dale matrices

$$W = \begin{pmatrix} 0 & 2 & 0 & -2 \\ 1 & 0 & 2 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 \end{pmatrix}, \text{ and } U = \begin{pmatrix} 0 & 3 & 0 & -2 \\ 0.5 & 0 & 0.5 & 0 \\ 0 & 0.5 & 0 & 0 \\ 1 & 1 & 1 & 0 \end{pmatrix}$$

and note that both these matrices have the same excitatory connectivity graph $G_{\mathcal{E}}$ and the same excitatory neurons $\beta = \{2, 3\}$ that do not receive inhibition. Let $\sigma = \{1, 2, 3\}$, thus $\beta_{\sigma} = \{2, 3\}$ and observe that $\rho(W_{\beta_{\sigma}}) = \sqrt{2} > 1$, thus by Theorem 3.4 $\sigma \notin \mathcal{C}(W)$. On the other hand $\rho(U_{\beta_{\sigma}}) = 0.5 < 1$. Furthermore, because $\sigma = \mathcal{E}$, the graph condition $t(\sigma) \cap \beta \subset \sigma$ is satisfied and thus by Theorem 3.4, $\sigma \in \mathcal{C}(U)$.

3.3 Combinatorial codes of Dale recurrent networks are convex

Perhaps the most surprising implication of Theorem 3.4 is that Dale networks naturally produce convex combinatorial codes. To explain the relevant background, we first motivate and define convex combinatorial codes.

There are two complimentary viewpoints on what determines neural activity in sensory systems. One viewpoint is that the brain represents information via patterns of neural activity that arise as a result of neural dynamics. A different perspective is that the neural activity in sensory neural systems is induced by external stimuli, whereby each neuron responds to a given stimulus according to its own receptive field. These two pictures must agree with each other, that is the patterns of neural activity resulting from the dynamics should be compatible with those allotted by the patterns of the receptive fields.

We define a receptive field of an individual neuron as a subset $U \subset \mathbb{R}^d$ in a stimulus space \mathbb{R}^d , such that the firing rate x(t) of a given neuron is non-zero at the times when the stimulus is in the region U. Given a collection of receptive fields $\mathcal{U} = \{U_i\}$ of a population of neurons \mathcal{E} , consider the code of \mathcal{U} , that describes all possible intersection patterns of the receptive fields U_i as

$$\operatorname{code}(\mathcal{U}) \stackrel{\text{def}}{=} \{ \sigma \subset \mathcal{E} \mid R_{\sigma} \neq \emptyset \},\$$

where

$$R_{\sigma} \stackrel{\text{def}}{=} \left(\bigcap_{i \in \sigma} U_i \right) \setminus \bigcup_{j \notin \sigma} U_j$$

denotes the region in the stimulus space where each of the neurons in σ is activated and no other considered neuron is active, and $R_{\emptyset} \stackrel{\text{def}}{=} \mathbb{R}^d \setminus \bigcup_{j \in \mathcal{E}} U_j$. The regions R_{σ} partition the stimulus space \mathbb{R}^d (see Figure 2).




Figure 2: An example of a collection of receptive fields $\mathcal{U} = \{U_1, U_2, U_3, U_4\}$ and its code, $\mathcal{C} = \text{code}(\mathcal{U}) = \{\emptyset, 2, 3, 4, 12, 13, 23, 34, 123, 134\}$. Here we denote a codeword $\{i_1, i_2, \ldots, i_k\} \in \mathcal{C}$ by the string $i_1 i_2 \ldots i_k$; for example, $\{1, 3, 4\}$ is abbreviated to 134. Note that all U_i are convex (and thus connected), but the regions R_{σ} are typically non-convex for non-maximal subsets σ . For example R_{13} is neither convex nor connected.


Note that $\operatorname{code}(\mathcal{U})$ is the set of all possible pattens of neuronal activation that are compatible with the receptive fields \mathcal{U} . The same patterns should be allowed by the neural network dynamics. We associate a combinatorial neural code $\mathcal{C}(W)$ to a Dale network (1), and we interpret the fixed points of the excitatory neurons (both stable and unstable) as our model for (transiently) activated stimulus representations. In order to reconcile this interpretation with the receptive field perspective, the combinatorial code of the network must be the same as the intersection patterns of the receptive fields:

$$\mathcal{C}(W) = \operatorname{code}(\mathcal{U}).$$

Recall that a set $U \subset \mathbb{R}^d$ is called *convex* if for any two points $x, y \in U$ the line segment [x, y] also lies in U. A number of sensory areas in the brain possess convex receptive fields; a very incomplete list of such areas includes the hippocampus, the primary visual cortex, the primary auditory cortex, etc. Following [6, 4, 16, 8], we define convex combinatorial codes as follows.

Definition 3.6. A combinatorial code $\mathcal{C} \subset 2^{[n]}$ is *open convex* if $\mathcal{C} = \text{code}(\mathcal{U})$ for a collection $\mathcal{U} = \{U_i\}$ of open convex subsets in a Euclidean space \mathbb{R}^d , for some $d \ge 1$.

It has been previously established in [6, 11, 4] that not every combinatorial code is open convex. The smallest example of a nonconvex code is illustrated in Figure 3. The topological properties that prevent a combinatorial code from being convex were studied in [4, 8, 17]. In fact, a randomly chosen¹ combinatorial code on a large number of neurons is non-convex with a high probability.



Following [4], recall that a combinatorial code C is called Figure 3: A non-convex *intersection-complete*, if intersection of any two codewords in C also code $C = \{1, 2, 13, 23\}$. belongs to C:

$$\sigma, \nu \in \mathcal{C} \implies \sigma \cap \nu \in \mathcal{C}.$$

 $^{^{1}}$ This, of course, requires a proper definition of the probability distribution on the set of all codes that we omit here.

If a code is intersection-complete and also has the property that a union of two codewords is a codeword, then such code is a *sublattice* of $2^{\mathcal{E}}$ [22]. It has been previously established in [4] that intersection-complete codes are open convex.

Theorem 3.7. [4] Suppose $\mathcal{C} \subset 2^{[n]}$ is an intersection-complete code. Then \mathcal{C} is open convex.

We use this result to show that the combinatorial codes of a Dale network are convex. Consider a directed graph $G_{\mathcal{E}}$, whose vertices are the excitatory neurons \mathcal{E} , and whose edges are derived from the the excitatory connectivity as in Definition 3.3. First we observe the following.

Proposition 3.8. The code $code(G_{\mathcal{E}},\beta)$ is a sublattice of the Boolean lattice $2^{\mathcal{E}}$.

This observation, combined with Theorem 3.5 implies that the combinatorial code of a weakly coupled network is a sublattice. If the network is *not* weakly coupled, then the spectral condition in Theorem 3.4 may prevent the code of the network from being a sublattice, however the code $\mathcal{C}(W)$ of a Dale network remains intersection-complete.

Theorem 3.9. For a Dale synaptic matrix W, its code C(W) is intersection-complete and is thus convex.

The proof of this theorem is given in Section 5.2 (page 19). We suggest that this theorem may provide a new explanation for the prevalence of convex receptive fields of excitatory neurons in recurrent circuits of sensory systems.

3.4 Constructing a Dale network from a combinatorial code

Can one encode an arbitrary combinatorial code as the combinatorial code of a Dale network? Theorem 3.9 implies that the code needs to be intersection-complete, to be realized on a generic Dale network. In addition, Proposition 3.8 tells us that if one wants to build a weakly coupled network with a prescribed code, this code needs to be a sublattice. It turns out that any sublattice can be encoded on a weakly coupled Dale network.

Theorem 3.10. Given a sublattice $\mathcal{C} \subset 2^{\mathcal{E}}$ with $\emptyset, \mathcal{E} \in \mathcal{C}$, define a map $c : 2^{\mathcal{E}} \to \mathcal{C}$ as

$$c(\sigma) \stackrel{\text{def}}{=} \bigcap_{\substack{\nu \in \mathcal{C}, \\ \nu \supset \sigma}} \nu, \tag{3}$$

and consider a directed graph G_c , whose vertices are \mathcal{E} and whose edges are defined via the rule

$$i \to j \iff j \in c(\{i\}) \text{ and } i \neq j.$$
 (4)

Then $\mathcal{C} = \operatorname{code}(G_c, \mathcal{E}).$

This theorem, combined with Theorem 3.5 translates into the following

Corollary 3.11. Given a sublattice $C \subset 2^{\mathcal{E}}$ with $\emptyset, \mathcal{E} \in C$, one can find a weakly coupled network W of excitatory neurons with C = C(W).

Here the *learning rule* for an excitatory synaptic matrix W amounts to assigning small nonzero excitatory synaptic weights according to the directed arcs in the graph G_c defined in equations (3),(4).

The following example illustrates the procedure described above to construct a graph of connectivity from a given code. Consider a sublattice $C = \{\emptyset, \{4\}, \{2, 4\}, \{3, 4\}, \}$

 $\{2,3,4\},\{1,2,3,4\}\}$. It is straightforward to verify that G_c is the graph in Figure 4 and thus $C = \text{code}(G_c,\{1,2,3,4\}).$



Figure 4: The graph G_c on four vertices such that $code(G_c, [4]) = \{\emptyset, \{4\}, \{2, 4\}, \{3, 4\}, \{2, 3, 4\}, [4]\}.$

4 Discussion

We investigated threshold-linear recurrent networks that satisfy the Dale's law and established what features of the synaptic connectivity are responsible for determining their combinatorial codes. Theorem 3.4 describes the combinatorial code in terms of the connectivity graph and the spectral radii of the appropriate sub-matrices. In the case of weakly coupled networks, the spectral conditions are always satisfied and the combinatorial code depends only on the features in $\operatorname{code}(G,\beta)$ that are derived from the connectivity graph (Theorem 3.5). In this situation, all the fixed points are stable, moreover the combinatorial code is a sublattice. We also proved that any sub-lattice can be encoded as a combinatorial code of an excitatory network (Corollary 3.11).

If the network is *not* weakly coupled, the spectral condition in Theorem 3.4 may prevent the combinatorial code from being a sublattice. Intuitively, this is because the union of two codewords in C(W) may not be in C(W), as the spectral radius of the appropriate larger matrix may (or may not) increase. We have proven in Theorem 3.9 that every combinatorial code is intersection-complete and therefore convex. Can any intersection-complete code be encoded on a Dale network? The answer to this question is currently unknown, as it requires a better understanding of the interplay of the spectral and the graph conditions in Theorem 3.4.

Finally, we hypothesize that the result (Theorem 3.9) that the combinatorial codes of Dale recurrent networks are always convex may provide a natural explanation for the ubiquity of convex receptive fields in many sensory systems in the mammalian brain.

5 Proofs

In this section we provide proofs for our main results stated in Section 3. First, we recall the conditions for having a fixed point of the dynamics of (1).

Lemma 5.1. [7, Proposition 2.1] For a threshold linear network in eq. (1), characterized by (W, b), a point $x^* \in \mathbb{R}^n$ is a fixed point with support $\sigma \subset [n]$ if and only if the following conditions are all satisfied

1. $(I - W)_{\sigma} x_{\sigma}^* = b_{\sigma}$, 2. $x_{\sigma}^* > 0$, and 3. $b_{\overline{\sigma}} \leq -W_{\overline{\sigma}\sigma} x_{\sigma}^*$.

Here b_{σ} the restriction of a vector b to the indices in σ , $\overline{\sigma}$ denotes the complement in [n] to a subset σ , $W_{\overline{\sigma}\sigma}$ is the rectangular submatrix of W, restricted to the rows in $\overline{\sigma}$ and columns in σ , and $(I - W)_{\sigma} = (I - W)_{\sigma\sigma}$ is the principal submatrix restricted to the subset σ . Note that if $(I - W)_{\sigma}$ is invertible, then there is at most one fixed point with support σ and it is given by $x_{\sigma}^* = (I_{\sigma} - W_{\sigma})^{-1}b_{\sigma}$.

A key ingredient in our proofs is the observation that the first condition in Lemma 5.1 can be translated to the language of semipositive matrices (Definition 5.5 below). From there we make use of the theory M-matrices (Definition 5.2 below). Note that the machinery of semipositive matrices has been previously used in a different context of recurrent networks with Heaviside transfer function in [23].

5.1 Necessary matrix theory results

First, we recall some definitions and results from the theory of *M*-matrices, following [3]. An $m \times n$ matrix *A* is *positive* (*nonnegative*) if $A_{ij} > 0$ ($A_{ij} \ge 0$) for all *i* and *j*. If *A* is positive (nonnegative) we denote this by A > 0 ($A \ge 0$). For a vector $x \in \mathbb{R}^n$, we write x > 0, x < 0, $x \ge 0$, or $x \le 0$ if all the entries of corresponding vector *x* satisfy the appropriate inequality.

Definition 5.2. [3, Chapter 6] A square matrix A is a Z-matrix if $A_{ij} \leq 0$ for all $i \neq j$. A Z-matrix A is called an M-matrix if A = sI - B, where I denotes the identity matrix, $B_{ij} \geq 0$ for all $i \neq j$, and the scalar s is not smaller than the spectral radius of the matrix $B, s \geq \rho(B)$.

The following simple corollary of the Perron-Frobenius theorem will be used later.

Lemma 5.3. Assume that A = I - W, where $W \ge 0$. Then the following two statements are equivalent:

- (i) The matrix A is a non-singular M-matrix.
- (ii) The spectral radius of W is smaller than one: $\rho(W) < 1$.

Proof. Note that $(ii) \implies (i)$ is immediate. To prove the opposite direction observe that by the Perron-Frobenius theorem [14], $\lambda = \rho(W) \ge 0$ is an eigenvalue of the non-negative matrix W. If A = I - W is an M-matrix, then $\rho(W) \le 1$. If $\lambda = \rho(W) = 1$, then the matrix A = I - W is singular, which contradicts condition (i), thus $(i) \implies (ii)$.

We shall also make use of the following.

Lemma 5.4. [3, Chapter 2, Corollary 1.6] Let A be a square nonnegative matrix. Suppose that B is a principal submatrix of A. Then $\rho(B) \leq \rho(A)$.

Definition 5.5. A square matrix A is *semipositive* if $\exists x > 0$ such that Ax > 0.

Lemma 5.6. Let A be a non-singular $n \times n$ matrix. Then A is semipositive if and only if there exists a vector x > 0 such that $Ax \ge 0$.

Proof. The forward direction is immediate. For the converse suppose that $\exists x > 0$ such that $Ax \ge 0$. Let $a = ||A^{-1}||_{op}$ be the operator norm of A^{-1} , $x_m = \min_{1 \le i \le n}(x_i)$, $\varepsilon = \frac{x_m}{2a\sqrt{n}}$ and $\overline{\varepsilon} \in \mathbb{R}^n$ be the column vector whose all components are equal to ε . Then $||\overline{\varepsilon}||_2 = \frac{x_m}{2a}$ by construction. Furthermore, $||A^{-1}\overline{\varepsilon}||_2 \le a||\overline{\varepsilon}||_2 = a\frac{x_m}{2a} = \frac{x_m}{2}$. In particular, $\sum_{i=1}^n (A^{-1}\overline{\varepsilon})_i^2 \le \frac{x_m}{4}$ for all $1 \le i \le n$. Therefore, $(A^{-1}\overline{\varepsilon})_i^2 \le \frac{x_m^2}{4}$ which implies $|(A^{-1}\overline{\varepsilon})_i| \le \frac{x_m}{2}$ for all $1 \le i \le n$. Thus, $x + A^{-1}\overline{\varepsilon} > 0$ because of the way x_m was defined. Finally, $A(x + A^{-1}\overline{\varepsilon}) = Ax + \overline{\varepsilon} \ge 0 + \overline{\varepsilon} > 0$ and thus A is semipositive. \Box

The following example illustrates the necessity of the hypothesis that A is a non-singular matrix in Lemma 5.6. Consider a singular matrix $A = \begin{pmatrix} 1 & -1 & 0 \\ -1 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$. Let $x = \begin{pmatrix} 1 & 1 & 1 \end{pmatrix}^T$. Then note

that $Ax \ge 0$. Now let $y \in \mathbb{R}^3_{>0}$. Suppose that Ay > 0. This implies that $y_1 > y_2$ and $y_2 > y_1$ which cannot be. Thus A is not semipositive.

There are numerous characterizations of M-matrices; one can find several dozen conditions on a matrix that are all equivalent to being an M-matrix [3]. However, our proofs rely only on the following observation.

Theorem 5.7. [3, Chapter 6, Theorem 2.3] Let A be a Z-matrix. Then A is a non-singular M-matrix if and only if A is semipositive.

5.2 The proofs

In order to prove Theorem 3.1 a few preliminary results are necessary. Namely Theorem 5.9 and Corollary 5.10. We also introduce the following notation for convenience.

Definition 5.8. For the dynamical system in (1) we denote the set of supports of all the fixed points by FP(W, b) and we let $FP(W) \stackrel{\text{def}}{=} \bigcup_{b>0} FP(W, b)$.

Theorem 5.9. Let $W \in \mathbb{D}_n$ and $W' \in \mathbb{D}_{n+1}$ be Dale matrices such that $W'_{[n]} = W$, and the last (n+1)-st column of W' is inhibitory and satisfies the following condition:

$$\exists i \in \mathcal{E} \text{ with } W'_{i(n+1)} < 0 \implies \exists j \in \mathcal{I} \text{ with } W_{ij} < 0.$$
(5)

Then $\mathcal{C}(W) = \mathcal{C}(W')$.

Proof. The proof needs that both matrices W and W' satisfy the Ground Assumption (page 4). Since being non-singular is a fine-tuned condition for matrices, this is generically true. Furthermore, we can explicitly construct a W' in such a way that it satisfies the Ground Assumption and the hypothesis in the statement of the theorem. In particular we can choose the (n + 1)-st row of W'to be the zero vector. This will make it so that for any $\sigma \subset [n + 1]$, $(I - W')_{\sigma}$ is non-singular, assuming that W was already satisfying the Ground Assumption. We now proceed with the proof.

Suppose that $\sigma \in \mathcal{C}(W)$. Then, there exists $\tau \subset \mathcal{I}$ such that $\nu = \sigma \sqcup \tau \in FP(W)$, and (by Lemma 5.1) a vector $b \in \mathbb{R}^n_{\geq 0}$ such that $x_{\nu} = (I_{\nu} - W_{\nu})^{-1}b_{\nu} > 0$ and $0 \leq b_{\overline{\nu}} \leq -W_{\overline{\nu}\nu}x_{\nu}$, where

 $\overline{\nu} = [n] \setminus \nu$. Let $\nu' = \nu \cup \{n+1\}$ and define $y \in \mathbb{R}_{>0}^{|\nu|+1}$ as follows. The first $|\nu|$ entries of y are those of x_{ν} , i.e. $y_{\nu} = x_{\nu} \in \mathbb{R}_{>0}^{|\nu|}$, and the last entry is defined as

$$y_{|\nu|+1} = \left| \sum_{k \in \nu} (I - W')_{(n+1)k} x_k \right| + 1.$$

Then, by construction we have $(I - W')_{\nu'} y \ge 0$. Furthermore, $-W'_{\overline{\nu'}\nu'} y \ge -W_{\overline{\nu}\nu} x_{\nu} \ge 0$. Let $b' \in \mathbb{R}^{n+1}_{\ge 0}$ be defined by $b'_{\nu'} = (I - W')_{\nu'} y$ and $b'_{\overline{\nu'}} = 0$. Then, by Lemma 5.1 $\nu' = \sigma \sqcup \tau \cup \{n+1\} \in FP(W', b') \subset FP(W')$, and thus $\sigma \in \mathcal{C}(W')$.

Now suppose that $\sigma \in \mathcal{C}(W')$. Then there exists a $\tau' \subset \mathcal{I} \sqcup \{n+1\}$ such that $\nu = \sigma \sqcup \tau' \in FP(W')$. Since $\nu \in FP(W')$, there exists a $b' \in \mathbb{R}^{n+1}_{\geq 0}$ such that

$$x \stackrel{\text{def}}{=} (I_{\nu} - W'_{\nu})^{-1} b'_{\nu} > 0, \text{ and } 0 \leq b'_{[n+1]\setminus\nu} \leq -W'_{([n+1]\setminus\nu)\nu} x,$$

by Lemma 5.1. We have two cases to consider; either $\tau' \subset \mathcal{I}$ or $(n+1) \in \tau'$.

In the case that $\tau' \subset \mathcal{I}$, observe that $(I_{\nu} - W'_{\nu})^{-1} = (I_{\nu} - W_{\nu})^{-1}$ and that $W'_{([n+1]\setminus\nu)\nu}$ is almost identical to $W_{([n]\setminus\nu)\nu}$; $W'_{([n+1]\setminus\nu)\nu}$ can be obtained from $W_{([n]\setminus\nu)\nu}$ by adding an extra row of entries induced from the (n+1)-st row of W'. Define $b \in \mathbb{R}^n_{\geq 0}$ by $b_{\nu} = (I - W)_{\nu}x = (I - W')_{\nu}x = b'_{\nu}$ and $b_{[n]\setminus\nu} = 0$. Then by construction

$$x = (I_{\nu} - W_{\nu})^{-1} b_{\nu} > 0$$
, and $0 \le b_{[n] \setminus \nu} \le -W_{([n] \setminus \nu)\nu} x$,

and thus $\nu = \sigma \sqcup \tau' \in FP(W, b) \subset FP(W)$ by Lemma 5.1. Therefore $\sigma \in \mathcal{C}(W)$.

In the case that $(n + 1) \in \tau'$, let α be the set of indices of nonzero rows of $W'_{\mathcal{E}\tau'}$. For each $i \in \alpha$ let $j_i \in \mathcal{I}$ be such that $W'_{ij_i} = W_{ij_i} < 0$ (We can always choose j_i to not equal n + 1 by the hypothesis of the theorem, and thus we can assume $j_i \in \mathcal{I}$). Let $\mu = \sigma \sqcup \mathcal{I}$ and define

$$c \stackrel{\text{def}}{=} \max \left\{ \begin{array}{l} \max_{i \in \sigma \cap \alpha} -\frac{1}{W_{ij_i}} \Big| \sum_{k \in \sigma} (I - W)_{ik} x_k \Big|, \\ \max_{i \in ([n] \setminus \mu) \cap \alpha} -\frac{1}{W_{ij_i}} \Big| \sum_{k \in \sigma} -W_{ik} x_k \Big| \end{array} \right\}.$$

Let $y \in \mathbb{R}_{>0}^{|\mu|}$ be defined by $y_{\sigma} = x_{\sigma}$ and $y_j = 1 + c$ for $j \in \mathcal{I}$. Then for $i \in \sigma \cap \alpha$, we have

$$\begin{split} ((I-W)_{\mu}y)_{i} &= \sum_{k\in\mu} (I-W)_{ik}y_{k} = \sum_{k\in\sigma} (I-W)_{ik}y_{k} + \sum_{k\in\mathcal{I}} (I-W)_{ik}y_{k} = \\ &= \sum_{k\in\sigma} (I-W)_{ik}x_{k} + \sum_{k\in\mathcal{I}} (I-W)_{ik} + c(I-W)_{ij_{i}} + c\sum_{j_{i}\neq k\in\mathcal{I}} (I-W)_{ik} \geqslant \\ &\geqslant \sum_{k\in\sigma} (I-W)_{ik}x_{k} - W_{ij_{i}} + (-W_{ij_{i}})(-\frac{1}{W_{ij_{i}}}) \left| \sum_{k\in\sigma} (I-W)_{ik}x_{k} \right| + 0 \geqslant \\ &\geqslant 0 - W_{ij_{i}} > 0. \end{split}$$

For $i \in \sigma \setminus \alpha$, we have

$$((I - W)_{\mu}y)_{i} = \sum_{k \in \mu} (I - W)_{ik}y_{k} = \sum_{k \in \sigma} (I - W)_{ik}y_{k} + \sum_{k \in \mathcal{I}} (I - W)_{ik}y_{k} = \sum_{k \in \sigma} (I - W)_{ik}x_{k} + \sum_{k \in \mathcal{I}} (I - W)_{ik}y_{k} \ge \sum_{k \in \sigma} (I - W)_{ik}x_{k} + 0 > 0,$$

because the *i*-th row of $W'_{\mathcal{E}\tau'}$ is zero for $i \notin \alpha$ by assumption and x was by hypothesis such that for $i \in \sigma \setminus \alpha$, $((I - W')_{\nu} x)_i = \sum_{k \in \nu} (I - W')_{ik} x_k = \sum_{k \in \sigma} (I - W')_{ik} x_k + \sum_{k \in \tau'} (I - W')_{ik} x_k = \sum_{k \in \sigma} (I - W)_{ik} x_k + 0 > 0.$

For $i \in \mathcal{I}$, we have

$$((I-W)_{\mu}y)_{i} = \sum_{k\in\mu} (I-W)_{ik}y_{k} = \sum_{k\in\sigma} (I-W)_{ik}y_{k} + \sum_{k\in\mathcal{I}} (I-W)_{ik}y_{k} =$$
$$= \sum_{k\in\sigma} (I-W)_{ik}x_{k} + \sum_{k\in\mathcal{I}} (I-W)_{ik} + c(I-W)_{ii} + c\sum_{i\neq k\in\mathcal{I}} (I-W)_{ik} \geqslant$$
$$\geqslant \sum_{k\in\sigma} (I-W)_{ik}x_{k} + (I-W)_{ii} + \left|\sum_{k\in\sigma} -W_{ik}x_{k}\right| + 0 \geqslant$$
$$\geqslant 0+1>0.$$

Together, all these inequalities yield that $(I - W)_{\mu} y > 0$ and $(I_{\mu} - W_{\mu})^{-1} (I - W)_{\mu} y = y > 0$. On the other hand, for $i \in ([n] \setminus \mu) \cap \alpha$, we have

$$(-W_{([n]\setminus\mu)\mu}y)_i = (-W_{([n]\setminus\mu)\mu}y)_i = \sum_{k\in\mu} -W_{ik}y_k =$$

$$= \sum_{k\in\sigma} -W_{ik}y_k + \sum_{k\in\mathcal{I}} -W_{ik}y_k = \sum_{k\in\sigma} -W_{ik}x_k + \sum_{k\in\mathcal{I}} -W_{ik} + c\sum_{k\in\mathcal{I}} -W_{ik} =$$

$$= \sum_{k\in\sigma} -W_{ik}x_k - W_{ij_i} + \sum_{j_i\neq k\in\mathcal{I}} -W_{ik} + c(-W_{ij_i}) + c\sum_{j_i\neq k\in\mathcal{I}} -W_{ik} \ge$$

$$\geqslant \sum_{k\in\sigma} -W_{ik}x_k - W_{ij_i} + 0 + (-W_{ij_i})(-\frac{1}{W_{ij_i}}) \left|\sum_{k\in\sigma} -W_{ik}x_k\right| + 0 \ge$$

$$\geqslant -W_{ij_i} > 0.$$

Furthermore, for $i \in ([n] \setminus \mu) \setminus \alpha$, we have

$$(-W_{([n]\backslash\mu)\mu}y)_i = \sum_{k\in\mu} -W_{ik}y_k =$$
$$= \sum_{k\in\sigma} -W_{ik}y_k + \sum_{k\in\mathcal{I}} -W_{ik}y_k \ge \sum_{k\in\sigma} -W_{ik}x_k + 0 \ge 0,$$

because the *i*-th row of $W'_{\mathcal{E}\tau'}$ is zero for $i \notin \alpha$ by assumption and x was by hypothesis such that for $i \in ([n] \setminus \mu) \setminus \alpha \subset \mathcal{E}$,

$$(-W'_{([n+1]\setminus\nu)\nu}x)_i = \sum_{k\in\nu} -W'_{ik}x_k = \sum_{k\in\sigma} -W'_{ik}x_k + \sum_{k\in\tau'} -W'_{ik}x_k = \sum_{k\in\sigma} -W_{ik}x_k + 0 \ge 0.$$

Let $b \in \mathbb{R}^n_{\geq 0}$ be defined by $b_{\mu} = (I - W)_{\mu} y$ and $b_{[n] \setminus \mu} = 0$. Then, by construction $\mu = \sigma \sqcup \mathcal{I} \in FP(W, b) \subset FP(W)$ by Lemma 5.1. Therefore $\sigma \in \mathcal{C}(W)$.

Corollary 5.10. Let $W \in \mathbb{D}_n$, and $W' \in \mathbb{D}_{n+1}$ be Dale matrices such that $W'_{[n]} = W$, and the last (n + 1)-st column of W' is inhibitory and such that for all $i \in \mathcal{E}$, $W_{i(n+1)} < 0$ if and only if the *i*-th row vector of $W_{\mathcal{E}}$ is nonzero. Let W'' be a Dale matrix obtained by deleting all the inhibitory columns and rows of W' except the (n + 1)-st row and column. Then $\mathcal{C}(W) = \mathcal{C}(W'')$.

Proof. By Theorem 5.9 we have that $\mathcal{C}(W) = \mathcal{C}(W')$. Now we can reindex the inhibitory columns and rows so that the (n + 1)-st column of W' becomes the first inhibitory column. Then we start deleting all the other inhibitory rows and columns one at a time and at the end we obtain the matrix W''. The code will be preserved at each deletion step by Theorem 5.9 and thus $\mathcal{C}(W) = \mathcal{C}(W'')$. Note that W' can be generically chosen such that W' and at each deletion step all relevant matrices satisfy the Ground Assumption, which was necessary in the proof of Theorem 5.9.

Proof of Theorem 3.1. Note that we can obtain W' from W by following the procedure from Corollary 5.10 where we obtained W''. The result thus follows.

Before proceeding to prove Theorem 3.2, we make a few observations in Proposition 5.11 and Corollary 5.12 which will simplify our proof strategy.

Proposition 5.11. Let $W \in \mathbb{D}_n$, $\sigma \subset \mathcal{E}$ and $\tau \subset \mathcal{I}$. If $\sigma \sqcup \tau \in FP(W)$, then $\sigma \sqcup \tau' \in FP(W)$ for any $\tau \subset \tau' \subset \mathcal{I}$.

Proof. Suppose that $\nu = \sigma \sqcup \tau \in FP(W)$ and let $\tau \subset \tau' \subset \mathcal{I}, \nu' = \sigma \sqcup \tau'$. Then, there exists a $b \in \mathbb{R}^n_{\geq 0}$ such that

$$x \stackrel{\text{def}}{=} (I_{\nu} - W_{\nu})^{-1} b_{\nu} > 0$$
, and $0 \leq b_{\overline{\nu}} \leq -W_{\overline{\nu}\nu} x$,

by Lemma 5.1. Define

$$c \stackrel{\text{\tiny def}}{=} \max_{i \in \tau' \setminus \tau} \left| \sum_{k \in \sigma} (I - W)_{ik} x_k \right|.$$

Let $x' \in \mathbb{R}_{>0}^{|\nu'|}$ be defined by $x'_{\nu} = x$ and $x'_i = 1 + c$ for $i \in \nu' \setminus \nu$. Then for $i \in \nu$ we have

$$((I-W)_{\nu'}x')_i = \sum_{k \in \nu'} (I-W)_{ik} x'_k =$$
$$= \sum_{k \in \nu} (I-W)_{ik} x'_k + \sum_{k \in \nu' \setminus \nu} (I-W)_{ik} x'_k \ge$$
$$\ge \sum_{k \in \nu} (I-W)_{ik} x_k + 0 = b_i \ge 0.$$

For $i \in \nu' \setminus \nu$ we have

$$((I-W)_{\nu'}x')_i = \sum_{k\in\nu'}(I-W)_{ik}x'_k =$$

$$= \sum_{k\in\sigma}(I-W)_{ik}x'_k + \sum_{k\in\tau}(I-W)_{ik}x'_k + \sum_{k\in\tau'\setminus\tau}(I-W)_{ik}x'_k \ge$$

$$= \sum_{k\in\sigma}(I-W)_{ik}x'_k + 0 + (I-W)_{ii}x'_i + \sum_{i\neq k\in\tau'\setminus\tau}(I-W)_{ik}x'_k \ge$$

$$\ge \sum_{k\in\sigma}(I-W)_{ik}x_k + 1 + c + 0 \ge \sum_{k\in\sigma}(I-W)_{ik}x_k + 1 + \left|\sum_{k\in\sigma}(I-W)_{ik}x_k\right| \ge 1 > 0.$$

Furthermore, for $i \in \overline{\nu'} \subset \overline{\nu}$ we have

$$(-W_{\overline{\nu'}\nu'}x')_i = \sum_{k\in\nu'} -W_{ik}x'_k = \sum_{k\in\nu} -W_{ik}x'_k + \sum_{k\in\tau'\setminus\tau} -W_{ik}x'_k \ge \sum_{k\in\nu} -W_{ik}x_k + 0 \ge b_i \ge 0.$$

Define $b' \in \mathbb{R}^n_{\geq 0}$ by $b'_{\nu'} = (I - W)_{\nu'} x'$ and $b'_{\overline{\nu'}} = 0$. Then by construction and by Lemma 5.1, $\nu' = \sigma \sqcup \tau' \in FP(W, b') \subset FP(W)$.

We thus immediately have the following corollary.

Corollary 5.12. Let $W \in \mathbb{D}_n$. Then $\sigma \in \mathcal{C}(W)$ if and only if $\sigma \sqcup \mathcal{I} \in FP(W)$.

We now prove Theorem 3.2.

Proof of Theorem 3.2. The proof relies on W satisfying the Ground Assumption, however we will also need W' to satisfy the Ground Assumption. Because of the way W' was constructed from W, this is the case. Indeed, observe that for $\sigma \subset [n]$, we have a block matrix decomposition $(I-W')_{\sigma} = \begin{pmatrix} (I-W)_{\mathcal{E}\cap\sigma} & -W_{(\mathcal{E}\cap\sigma)}(\mathcal{I}\cap\sigma) \\ 0 & (I-W)_{\mathcal{I}\cap\sigma} \end{pmatrix}$. Therefore $\det((I-W')_{\sigma}) = \det((I-W)_{\mathcal{E}\cap\sigma}) \det((I-W)_{\mathcal{I}\cap\sigma}) \neq 0$, by the Ground Assumption for W. Having this in mind, we proceed with the proof.

Let $\sigma \in \mathcal{C}(W)$. Then there exists a $\tau \subset \mathcal{I}$ such that $\nu = \sigma \sqcup \tau \in FP(W)$. Thus, there exists a $b \in \mathbb{R}^n_{\geq 0}$ such that

$$x \stackrel{\text{def}}{=} (I_{\nu} - W_{\nu})^{-1} b_{\nu} > 0, \text{ and } - W_{\overline{\nu}\nu} x \ge b_{\overline{\nu}} \ge 0,$$

by Lemma 5.1. Observe that because of the way W' was defined we have that

$$(I - W')_{ij} = \begin{cases} 0, & i \in \mathcal{I}, j \in \mathcal{E} \\ (I - W)_{ij}, & \text{otherwise} \end{cases}$$

Therefore $y \stackrel{\text{def}}{=} (I - W')_{\nu} x \ge (I - W)_{\nu} x = b_{\nu} \ge 0$. Observe that $x = (I_{\nu} - W'_{\nu})^{-1} y = (I_{\nu} - W_{\nu})^{-1} b_{\nu}$. Furthermore, $-W'_{\overline{\nu}\nu} x \ge -W_{\overline{\nu}\nu} x \ge 0$. Define $b' \in \mathbb{R}^n_{\ge 0}$ by $b'_{\nu} = y$ and $b'_{\overline{\nu}} = 0$. Then by construction $\nu = \sigma \sqcup \tau \in FP(W', b') \subset FP(W')$ by Lemma 5.1. Therefore $\sigma \in \mathcal{C}(W')$.

Now suppose that $\sigma \in \mathcal{C}(W')$. By Corollary 5.12 it follows that $\nu = \sigma \sqcup \mathcal{I} \in FP(W')$. Thus, there exists a $b' \in \mathbb{R}^n_{\geq 0}$ such that

$$x' \stackrel{\text{def}}{=} (I_{\nu} - W'_{\nu})^{-1} b'_{\nu} > 0, \text{ and } - W'_{\overline{\nu}\nu} x' \ge b'_{\overline{\nu}} \ge 0,$$

by Lemma 5.1. Define

$$c \stackrel{\text{\tiny def}}{=} \max_{i \in \mathcal{I}} \left| \sum_{k \in \sigma} (I - W)_{ik} x'_k \right|.$$

Let $x \in \mathbb{R}_{>0}^{|\nu|}$ be defined by $x_{\sigma} = x'_{\sigma}$ and $x_i = x'_i + 1 + c$ for all $i \in \mathcal{I}$. Then, for $i \in \sigma$ we have

$$((I-W)_{\nu}x)_{i} = \sum_{k\in\nu} (I-W)_{ik}x_{k} =$$

$$= \sum_{k\in\nu} (I-W')_{ik}x'_{k} + (1+c)\sum_{k\in\mathcal{I}} (I-W)_{ik} \ge$$

$$\ge \sum_{k\in\nu} (I-W')_{ik}x'_{k} + 0 \ge 0.$$

For $i \in \mathcal{I}$ we have

$$((I-W)_{\nu}x)_{i} = \sum_{k\in\nu}(I-W)_{ik}x_{k} = \sum_{k\in\sigma}(I-W)_{ik}x_{k} + \sum_{k\in\mathcal{I}}(I-W)_{ik}x_{k} =$$
$$= \sum_{k\in\sigma}(I-W)_{ik}x'_{k} + (I-W)_{ii}x_{i} + \sum_{i\neq k\in\mathcal{I}}(I-W)_{ik}x_{k} \ge$$
$$\ge \sum_{k\in\sigma}(I-W)_{ik}x'_{k} + (x'_{i}+1+c) + 0 \ge$$
$$\ge \sum_{k\in\sigma}(I-W)_{ik}x'_{k} + 1 + \left|\sum_{k\in\sigma}(I-W)_{ik}x'_{k}\right| \ge 1 > 0.$$

Furthermore, for $i \in \overline{\nu} = \mathcal{E} \setminus \sigma$ we have

$$(-W_{\overline{\nu}\nu}x)_i = \sum_{k\in\nu} -W_{ik}x_k = \sum_{k\in\nu} -W'_{ik}x'_k + (1+c)\sum_{k\in\mathcal{I}} -W'_{ik} \ge \sum_{k\in\nu} -W'_{ik}x'_k + 0 \ge 0.$$

Define $b \in \mathbb{R}^n_{\geq 0}$ by $b_{\nu} = (I - W)_{\nu} x$ and $b_{\overline{\nu}} = 0$. Then by construction and by Lemma 5.1, $\nu = \sigma \sqcup \mathcal{I} \in FP(W, b) \subset FP(W)$. Therefore, $\sigma \in \mathcal{C}(W)$.

From Theorems 3.1 and 3.2 we have the following corollary.

Corollary 5.13. Let $W \in \mathbb{D}_n$, with $\mathcal{E} = \{1, 2, ..., m\}$, $|\mathcal{I}| = k \ge 1$, m + k = n. Let $W' \in \mathbb{D}_{m+1}$ be a Dale matrix obtained from W in the following way: $W'_{ij} = W_{ij}$ for $i, j \in \mathcal{E}$, and for all $i \in \mathcal{E}$, $W'_{i(m+1)} < 0$ if and only if the *i*-th row of $W_{\mathcal{EI}}$ is a nonzero row vector, and the (m + 1)-st row of W' is zero. Then $\mathcal{C}(W) = \mathcal{C}(W')$.

Before we proceed to prove Theorem 3.4, we will need the following lemma.

Lemma 5.14. Let A be an $n \times m$ matrix, $n \leq m$, such that $\exists \sigma \subset [m]$ with $|\sigma| = n$ and that $A_{[n]\sigma}$ is an invertible Z-matrix, and $A_{[n]([m]\setminus\sigma)} \leq 0$. Here $A_{[n]\tau}$ is a submatrix of A obtained by deleting columns outside of τ . Then $\exists x \in \mathbb{R}^m_{>0}$ such that $Ax \geq 0$ if and only if $A_{[n]\sigma}$ is an M-matrix.

Proof. We can assume without loss of generality that $\sigma = [n]$, otherwise we can re-index. Suppose that $\exists x \in \mathbb{R}_{>0}^m$ such that $Ax \ge 0$. For all $i \in [n]$ we then have $(Ax)_i = \sum_{k \in [m]} A_{ik}x_k = \sum_{k \in [n]} A_{ik}x_k + \sum_{k \in [m] \setminus [n]} A_{ik}x_k \ge 0$, from where it follows that $\sum_{k \in [n]} A_{ik}x_k = A_{i[n]}x_{[n]} \ge 0$ because $A_{i([m] \setminus [n])} \le 0$ by assumption. Thus, $A_{[n]}x_{[n]} \ge 0$ and thus $A_{[n]}$ is semipositive by Lemma 5.6. Since it is also a Z-matrix by assumption, we have that it is an M-matrix by Theorem 5.7.

Conversely, suppose that $A_{[n]}$ is an *M*-matrix. Then, since it is also invertible by assumption, from Theorem 5.7 we have that $A_{[n]}$ is semipositive. Therefore, $\exists y \in \mathbb{R}^n_{>0}$ such that $A_{[n]}y \ge 0$ by Lemma 5.6. Note that for all $i \in [n], \sum_{k \in [m] \setminus [n]} A_{ik} \le 0$. Define

$$M \stackrel{\text{def}}{=} \min_{i \in [n]} \sum_{k \in [n]} A_{ik} y_k, \ N \stackrel{\text{def}}{=} \min_{i \in [n]} \sum_{k \in [m] \setminus [n]} A_{ik}.$$

Observe that since $A_{[n]}$ is invertible by assumption and y > 0 it cannot be that $A_{[n]}y = 0$. Thus, M > 0. Suppose that N = 0. Since, $A_{[n]([m]\setminus[n])} \leq 0$ by assumption, this implies that $A_{[n]([m]\setminus[n])} = 0$. Therefore, we can define $x \in \mathbb{R}_{>0}^m$ by $x_{[n]} = y$ and $x_j = 1$ for all $j \in [m]\setminus[n]$. Then, by construction $Ax \ge 0$. Now suppose that $N \ne 0$, that is N < 0 and define $x \in \mathbb{R}_{>0}^m$ by $x_{[n]} = y$ and $x_j = \frac{-M}{N}$ for all $j \in [m] \setminus [n]$. Then for all $i \in [n]$ we have

$$(Ax)_i = \sum_{k \in [m]} A_{ik} x_k = \sum_{k \in [n]} A_{ik} x_k + \sum_{k \in [m] \setminus [n]} A_{ik} x_k =$$
$$= \sum_{k \in [n]} A_{ik} y_k - \frac{M}{N} \sum_{k \in [m] \setminus [n]} A_{ik} \ge M - \frac{M}{N} N = M - M = 0.$$

Therefore $Ax \ge 0$.

Proof of Theorem 3.4. By Theorems 3.1 and 3.2, we can assume that there is only one inhibitory neuron, let us say the n-th neuron, and that the n-th row of W is all 0.

Suppose that $\sigma \in \mathcal{C}(W)$. As observed in Corollary 5.12, $\sigma \in \mathcal{C}(W)$ if and only if $\tau = \sigma \sqcup \{n\} \in FP(W)$. By Lemma 5.1, $\tau \in FP(W)$ if and only if $\exists x \in \mathbb{R}_{>0}^{|\tau|}$ such that the following two conditions are satisfied:

$$(I - W)_{\tau} x \ge 0 \tag{6a}$$

$$0 \leqslant -W_{\overline{\tau}\tau}x. \tag{6b}$$

Observe that for any vector $y \in \mathbb{R}_{>0}^{|\tau|}$, for $i \in \beta_{\sigma}$ we have

$$((I-W)_{\tau}y)_i = \sum_{k \in \tau} (I-W)_{ik} y_k = \sum_{k \in \sigma} (I-W)_{ik} y_k,$$

since $(I - W)_{in} = 0$. Thus, condition (6a) forces us to have $x_{\sigma} > 0$ be such that $(I - W)_{\beta_{\sigma}\sigma}x_{\sigma} \ge 0$. Because $(I - W)_{\beta_{\sigma}}$ is a non-singular Z-matrix and $(I - W)_{\beta_{\sigma}(\sigma\setminus\beta_{\sigma})} \le 0$, by Lemma 5.14 we have that this is equivalent to $(I - W)_{\beta_{\sigma}}$ being an *M*-matrix, that is $\rho(W_{\beta_{\sigma}}) < 1$. Furthermore, note that $\overline{\tau} = \mathcal{E}\setminus\sigma$. Thus, condition (6b) is equivalent that for all $i \in \mathcal{E}\setminus\sigma$ we need to have $(-W_{(\mathcal{E}\setminus\sigma)\tau}x)_i = \sum_{k\in\tau} -W_{ik}x_k \ge 0$. For $i \in \mathcal{E}\setminus\sigma$, if $W_{in} = 0$ the only possibility is that $W_{i\sigma} = 0$ for the inequality to be true.

Thus to be able to find an $x \in \mathbb{R}^n_{>0}$ for which condition (6b) is satisfied we need that for all $i \in \mathcal{E} \setminus \sigma$, $W_{i\sigma} = 0$ or $W_{in} < 0$. This implies that $t(\sigma) \cap \beta \subset \sigma$.

Now suppose the following two conditions are both satisfied:

$$\rho(W_{\beta_{\sigma}}) < 1 \tag{7a}$$

$$t(\sigma) \cap \beta \subset \sigma. \tag{7b}$$

Condition (7a) means that $(I - W)_{\beta_{\sigma}}$ is an *M*-matrix. Because $(I - W)_{\beta_{\sigma}(\sigma \setminus \beta)} \leq 0$, by Lemma 5.14 this means that there $\exists x \in \mathbb{R}_{>0}^{|\sigma|}$ such that $(I - W)_{\beta_{\sigma}\sigma}x \geq 0$. Let $\tau = \sigma \sqcup \{n\}$, and define

$$\begin{split} c \stackrel{\text{def}}{=} \max \left\{ \begin{array}{l} \max_{i \in \sigma \setminus \beta_{\sigma}} -\frac{1}{W_{in}} \Big| \sum_{k \in \sigma} (I - W)_{ik} x_k \Big|, \\ \max_{i \in (\mathcal{E} \setminus \sigma) \setminus \beta_{\sigma}} -\frac{1}{W_{in}} \Big| \sum_{k \in \sigma} -W_{ik} x_k \Big|, \ \left| \sum_{k \in \sigma} -W_{nk} x_k \right| \right\}. \end{split} \right. \end{split}$$

Define $y \in \mathbb{R}_{>0}^{|\tau|}$ by $y_{\sigma} = x$ and $y_{|\tau|} = 1 + c$. Then, for $i \in \beta_{\sigma}$ we have

$$((I - W)_{\tau} y)_{i} = \sum_{k \in \tau} (I - W)_{ik} y_{k} = \sum_{k \in \sigma} (I - W)_{ik} y_{k} + (I - W)_{in} y_{|\tau|} =$$
$$= \sum_{k \in \sigma} (I - W)_{ik} x_{k} - W_{in} (1 + c) \ge 0 + 0(1 + c) = 0.$$

For $i \in \sigma \setminus \beta_{\sigma}$ we have

$$((I - W)_{\tau}y)_{i} = \sum_{k \in \tau} (I - W)_{ik} y_{k} = \sum_{k \in \sigma} (I - W)_{ik} y_{k} + (I - W)_{in} y_{|\tau|} =$$
$$= \sum_{k \in \sigma} (I - W)_{ik} x_{k} - W_{in} (1 + c) = \sum_{k \in \sigma} (I - W)_{ik} x_{k} - W_{in} - W_{in} c \ge$$
$$\ge \sum_{k \in \sigma} (I - W)_{ik} x_{k} - W_{in} + W_{in} \frac{1}{W_{in}} \left| \sum_{k \in \sigma} (I - W)_{ik} x_{k} \right| \ge -W_{in} > 0.$$

Similarly if i = n, we will also have that $((I - W)_{\tau}y)_i \ge 0$ because of the definition of $y_{|\tau|}$, Therefore we have that $(I - W)_{\tau}y \ge 0$. Furthermore, $t(\sigma) \cap \beta \subset \sigma$ implies that for all $i \in \mathcal{E} \setminus \sigma$, $W_{i\sigma} = 0$ or $W_{in} < 0$. Note that $\overline{\tau} = \mathcal{E} \setminus \sigma$. Thus for $i \in (\mathcal{E} \setminus \sigma) \cap \beta_{\sigma}$, and thus $W_{in} = 0$, we have that it must be that $W_{i\sigma} = 0$ and therefore

$$(-W_{\overline{\tau}\tau}y)_i = \sum_{k \in \tau} -W_{ik}y_k = \sum_{k \in \sigma} -W_{ik}y_k - W_{in}y_{|\tau|} = 0 - 0(1+c) = 0 \ge 0.$$

On the other hand for $i \in (\mathcal{E} \setminus \sigma) \setminus \beta_{\sigma}$ we have that $W_{in} < 0$ and thus

$$(-W_{\overline{\tau}\tau}y)_i = \sum_{k\in\sigma} -W_{ik}y_k = \sum_{k\in\sigma} -W_{ik}y_k - W_{in}y_{|\tau|} = \sum_{k\in\sigma} -W_{ik}x_k - W_{in}(1+c) = \sum_{k\in\sigma} -W_{ik}x_k - W_{in} - W_{in}c \ge \sum_{k\in\sigma} -W_{ik}x_k - W_{in} + W_{in}\frac{1}{W_{in}}\left|\sum_{k\in\sigma} -W_{ik}x_k\right| \ge \sum_{k\in\sigma} -W_{ik}x_k - W_{in} > 0.$$

Thus, we have that $0 \leq -W_{\overline{\tau}\tau}y$ and therefore $\tau = \sigma \sqcup \{n\} \in FP(W)$ by Lemma 5.1. Hence $\sigma \in \mathcal{C}(W)$.

We now prove Theorem 3.5, mainly relying on results stated in Section 6.

Proof of Theorem 3.5. Since $||W||_F < 1$, this implies that $\rho(W_\eta) < 1$ for any principal submatrix W_η , $\eta \subset [n]$. Therefore by Theorem 3.4, $\mathcal{C}(W) = \operatorname{code}(G_{\mathcal{E}}, \beta)$. Furthermore, $||W||_F < 1$ implies that $W \in \mathcal{L}$ (Definition 6.1) by Lemma 6.2. Since the Ground Assumption is assumed to hold, by Proposition 6.3, for all $b \in \mathbb{R}^n_{\geq 0}$ (1) has a unique globally exponentially stable fixed point. In particular, if $\sigma \in \mathcal{C}(W)$, by definition $\sigma \sqcup \tau$ is a support of a fixed point of (1) for some $\tau \subset \mathcal{I}$ and some input $b \in \mathbb{R}^n_{\geq 0}$. This fixed point is unique and asymptotically (exponentially) stable as argued above. Therefore $\sigma \in \mathcal{SC}(W)$. The inclusion $\mathcal{SC}(W) \subset \mathcal{C}(W)$ is by definition and thus $\mathcal{C}(W) = \mathcal{SC}(W)$.

The proof of Proposition 3.8 is straightforward and it gives us the important Corollary 5.15.

Proof of Proposition 3.8. It's easy to see that $\emptyset, \mathcal{E} \in \operatorname{code}(G, \beta)$. To show that $\operatorname{code}(G, \beta)$ is respected by intersections and unions assume that $t(\sigma_i) \cap \beta \subset \sigma_i$ for $i = 1, 2, \sigma_i \subset \mathcal{E}$. Since $t(\sigma_1 \cap \sigma_2) \subset t(\sigma_1) \cap t(\sigma_2)$, we obtain that

$$t(\sigma_1 \cap \sigma_2) \cap \beta \subset t(\sigma_1) \cap t(\sigma_1) \cap \beta \subset \sigma_1 \cap \sigma_2.$$

Similarly,

$$t(\sigma_1 \cup \sigma_2) \cap \beta = (t(\sigma_1) \cup t(\sigma_2)) \cap \beta = (t(\sigma_1) \cap \beta) \cup (t(\sigma_2) \cap \beta) \subset \sigma_1 \cup \sigma_2.$$

Corollary 5.15. Let $W \in \mathbb{D}_n$. The code $\mathcal{C}(W)$ is closed under intersections.

Proof. Suppose that $\sigma, \tau \in \mathcal{C}(W)$. By Theorem 3.4 we have that $\rho(W_{\beta_{\sigma}}), \rho(W_{\beta_{\tau}}) < 1$ and that $\sigma, \tau \in \operatorname{code}(G_{\mathcal{E}}, \beta)$. Note that $W_{\beta_{\sigma\cap\tau}}$ is a principal submatrix of $W_{\beta_{\sigma}}$ which is nonnegative. Thus, by Lemma 5.4, $\rho(W_{\beta_{\sigma\cap\tau}}) < 1$. Furthermore, $\operatorname{code}(G_{\mathcal{E}}, \beta)$ is closed under intersections by Proposition 3.8 and thus $\sigma \cap \tau \in \operatorname{code}(G_{\mathcal{E}}, \beta)$. Thus, by Theorem 3.4 it follows that $\sigma \cap \tau \in \mathcal{C}(W)$.

By Corollary 5.15 and Theorem 3.7 we immediately get Theorem 3.9.

We now proceed to prove Theorem 3.10. Given a code $C \subset 2^{\mathcal{E}}$ that is a sublattice with $\emptyset, \mathcal{E} \in C$, recall the definition of $c : 2^{\mathcal{E}} \to C$ and the graph (G_c, \mathcal{E}) from Theorem 3.10. Let $t_c(\sigma)$ denote the targets of $\sigma \subset \mathcal{E}$ in the graph (G_c, \mathcal{E}) . By definition, one sees that $c(\emptyset) = \emptyset$, $c(\mathcal{E}) = \mathcal{E}$, and $\sigma \subset c(\sigma)$. We immediately make the following observations that will help us prove Theorem 3.10.

Lemma 5.16. Let $C \subset 2^{\mathcal{E}}$ be a sublattice with $\emptyset, \mathcal{E} \in C$. Then the following are true.

1. Let $\sigma, \tau \subset \mathcal{E}$. Then $c(\sigma \cup \tau) = c(\sigma) \cup c(\tau)$,

2.
$$C = \{ \sigma \subset \mathcal{E} \mid \sigma = c(\sigma) \}.$$

Proof. For part 1, we first show that $c(\sigma) \cup c(\tau) \subset c(\sigma \cup \tau)$. Note that since $\sigma \subset \sigma \cup \tau$ it follows that $c(\sigma) \subset c(\sigma \cup \tau)$. Similarly $c(\tau) \subset c(\sigma \cup \tau)$. Therefore $c(\sigma) \cup c(\tau) \subset c(\sigma \cup \tau)$. Now we show the other inclusion. Observe that since C is a lattice, it follows that $c(\sigma), c(\tau)$ and $c(\sigma) \cup c(\tau) \in C$. Furthermore, $\sigma \cup \tau \subset c(\sigma) \cup c(\tau)$ and therefore by definition $c(\sigma \cup \tau) \subset c(\sigma) \cup c(\tau)$.

To prove part 2, suppose that $\sigma \in C$. By definition,

$$c(\sigma) = \bigcap_{\nu \in C, \sigma \subset \nu} \nu.$$

Since $\sigma \in C$, it follows that

$$\bigcap_{\nu\in\mathcal{C},\sigma\subset\nu}\nu=\sigma,$$

and thus $c(\sigma) = \sigma$. Now suppose that $\sigma \subset \mathcal{E}$ is such that

$$c(\sigma) = \bigcap_{\nu \in C, \sigma \subset \nu} \nu = \sigma.$$

Since C is finite and is closed under finite intersections, it follows that $\sigma \in C$.

Lemma 5.17. The arcs in G_c are a transitive relation on \mathcal{E} .

Proof. Let $i, j, k \in \mathcal{E}$ be such that $i \to j$ and $j \to k$. In other words, we have $j \in c(i)$ and $k \in c(j)$. Let $\nu \in C$ be such that $i \in \nu$. Then, by definition $j \in \nu$. Since $k \in c(j)$, by definition this implies that $k \in \nu$. Therefore, by definition $i \to k$.

Lemma 5.18. Let G be any graph whose transitive closure is G_c . Then $\operatorname{code}(G, \mathcal{E}) = \operatorname{code}(G_c, \mathcal{E})$

Proof. Let $t(\sigma), t_c(\sigma)$ be the targets of $\sigma \subset \mathcal{E}$, in G and G_c respectively. By definition $\operatorname{code}(G, \mathcal{E}) = \{\sigma \subset \mathcal{E} \mid t(\sigma) \subset \sigma\}$ and $\operatorname{code}(G_c, \mathcal{E}) = \{\sigma \subset \mathcal{E} \mid t_c(\sigma) \subset \sigma\}$. By assumption $t(\sigma) \subset t_c(\sigma)$ for all $\sigma \subset \mathcal{E}$. Thus for any $\sigma \subset \mathcal{E}$ if $t_c(\sigma) \subset \sigma$ then $t(\sigma) \subset \sigma$. Therefore $\operatorname{code}(G_c, \mathcal{E}) \subset \operatorname{code}(G, \mathcal{E})$. Furthemore, if for any $\sigma \subset \mathcal{E}, t(\sigma) \subset \sigma$, then $t^m(\sigma) \subset \sigma$ where t^m is an *m*-fold application of the *t* operator to σ , for any $m \ge 1$. Thus, $t_c(\sigma) \subset \sigma$. Therefore $\operatorname{code}(G, \mathcal{E})$.

Remark 5.19. This whole construction is common and known in finitely generated topologies, or equivalently Alexandroff topological spaces [2]. Note that $\emptyset, \mathcal{E} \in C$ and C being a lattice, means that C is a collection of closed sets of a topology on \mathcal{E} . The digraph we obtained is also known as the specialization preorder associated to the topology of C. Technically, the digraph we would obtain would have all loops, but we can ignore them as there is a one-to-one functorial correspondence of graphs with no loops and graphs with all loops.

Finally, we can prove Theorem 3.10.

Proof of Theorem 3.10. By definition, $\operatorname{code}(G_c, \mathcal{E}) = \{\sigma \subset \mathcal{E} \mid t_c(\sigma) \cap \mathcal{E} = t_c(\sigma) \subset \sigma\}$. Thus, we need to show that $\{\sigma \subset \mathcal{E} \mid t_c(\sigma) \cap \mathcal{E} = t_c(\sigma) \subset \sigma\} = \{\sigma \subset \mathcal{E} \mid c(\sigma) = \sigma\}$. Let G be the digraph on \mathcal{E} be defined by $i \to j$ if and only if $j \in c(i)$ for all $i, j \in \mathcal{E}$. Let $t(\sigma)$ be the targets of $\sigma \subset \mathcal{E}$ in G. Note that by construction, $t(\sigma) = t_c(\sigma) \cup \sigma$. Therefore for all $\sigma \subset \mathcal{E}$, $t(\sigma) = \sigma$ if and only if $t_c(\sigma) \subset \sigma$. Furthermore, by construction t(i) = c(i). Hence by Lemma 5.16 for all $\sigma \subset \mathcal{E}$ we have

$$t(\sigma) = \bigcup_{i \in \sigma} t(i) = \bigcup_{i \in \sigma} c(i) = c\left(\bigcup_{i \in \sigma} i\right) = c(\sigma).$$

Therefore for all $\sigma \subset \mathcal{E}$, $c(\sigma) = \sigma$ if and only if $t_c(\sigma) \subset \sigma$.

6 Appendix: Stability of Linear-Threshold Rate Dynamics

Here we expose the necessary results on the stability of the linear threshold dynamics from [19, 20] that we used in the proofs of Section 5. For a matrix A let ||A|| be its 2-norm and let |A| denote the matrix $|A|_{ij} = |A_{ij}|$.

Definition 6.1. An $n \times n$ matrix A is totally- \mathcal{L} stable, written $A \in \mathcal{L}$, if there exists $P = P^T > 0$ such that $(-I + A^T \Sigma)P + P(-I + \Sigma A) < 0$ for all $\Sigma = \text{diag}(\sigma)$ and $\sigma \in \{0, 1\}^n$.

Lemma 6.2. [19, Lemma 2.3]

- 1. $\rho(|W|) < 1 \Longrightarrow W \in \mathcal{L}.$
- 2. $||W|| < 1 \longrightarrow W \in \mathcal{L}.$

Given a $\sigma \in \{0,1\}^n$ let $\Sigma = \text{diag}(\sigma)$, that is the diagonal matrix with the elements of σ on the diagonal. In the statements of the following results the authors in [19] had a hypothesis that W is non-singular and that for all $\sigma \in \{0,1\}^n$, $(I - \Sigma W)$ is non-singular as well. By the standard correspondence between finite binary sequences and finite subsets of $2^{[n]}$, it is not hard to observe that for a given $\sigma \in \{0,1\}^n$, $(I - \Sigma W)$ is non-singular if and only if $(I - W)_{\sigma}$ is non-singular. Thus, the second hypothesis is equivalent to the Ground Assumption. We thus continue not writing out the Ground Assumption in the statement of the following theorem as is the case in the rest of the paper, but the reader should note that it is indeed necessary for the statement to be true.

Proposition 6.3. [19, Proposition 4.9] Consider the network dynamics in (1). If $\rho(|W|) < 1$ or ||W|| < 1, then for all $b \in \mathbb{R}^n$, the network has a unique fixed point x^* and it is globally exponentially stable relative to x^* .

Note that it has been shown in [19], that a more general condition [19, Theorem 4.8] can also guarantee the above result. However, checking if the matrix W satisfies those conditions is significantly harder in our context, thus we used the statement above instead.

Acknowledgments

This work was supported by the NSF Next Generation Networks for Neuroscience Program (award 2014217).

References

- Yashar Ahmadian, Francesco Fumarola, and Kenneth D Miller. "Properties of networks with partially structured and partially random connectivity". *Physical Review E* 91.1 (2015), p. 012820.
- [2] P. Alexandroff. "Diskrete Räume". Mat. Sbornik N.S. 2.44 (1937), pp. 501–519.
- [3] Abraham Berman and Robert J Plemmons. Nonnegative matrices in the mathematical sciences. SIAM, 1994.
- [4] Joshua Cruz, Chad Giusti, Vladimir Itskov, and Bill Kronholm. "On open and closed convex codes". Discrete & computational geometry 61.2 (2019), pp. 247–270.
- [5] C. Curto, A. Degeratu, and V. Itskov. "Encoding binary neural codes in networks of thresholdlinear neurons". *Neural Comput.* 25 (2013), pp. 2858–2903.
- [6] C Curto, V Itskov, A Veliz-Cuba, and N Youngs. "The neural ring: an algebraic tool for analyzing the intrinsic structure of neural codes." *Bulletin of mathematical biology* 75.9 (2013), pp. 1571–1611.
- [7] Carina Curto, Anda Degeratu, and Vladimir Itskov. "Flexible memory networks". Bulletin of mathematical biology 74.3 (2012), pp. 590–614.
- [8] Carina Curto, Elizabeth Gross, Jack Jeffries, Katherine Morrison, Mohamed Omar, Zvi Rosen, Anne Shiu, and Nora Youngs. "What makes a neural code convex?" SIAM Journal on Applied Algebra and Geometry 1.1 (2017), pp. 222–238.
- [9] Carina Curto and Katherine Morrison. "Pattern Completion in Symmetric Threshold-Linear Networks". Neural Computation 28 (Dec. 2015), pp. 2825–2852.

- [10] John Carew Eccles. "From electrical to chemical transmission in the central nervous system: the closing address of the sir henry dale centennial symposium cambridge, 19 september 1975". Notes and records of the Royal Society of London 30.2 (1976), pp. 219–230.
- [11] Chad Giusti and Vladimir Itskov. "A no-go theorem for one-layer feedforward networks". *Neural computation* 26.11 (2014), pp. 2527–2540.
- [12] R. H. Hahnloser, R. Sarpeshkar, M.A. Mahowald, R.J. Douglas, and H.S. Seung. "Digital selection and analogue amplification coexist in a cortex-inspired silicon circuit". *Nature* 405 (2000), pp. 947–951.
- [13] R. H. Hahnloser, H.S. Seung, and J.J. Slotine. "Permitted and forbidden sets in symmetric threshold-linear networks". *Neural Comput.* 15.3 (2003), pp. 621–638.
- [14] Roger A. Horn and Charles R. Johnson. *Matrix Analysis*. Cambridge University Press, 1990.
- [15] Jesper R Ipsen and Andre DH Peterson. "Consequences of Dale's law on the stabilitycomplexity relationship of random neural networks". *Physical Review E* 101.5 (2020), p. 052412.
- [16] V Itskov, A Kunin, and Z Rozen. "Hyperplane neural codes and the polar complex". The Abel Symposia proceedings 15 (2020), pp. 343–369.
- [17] Caitlin Lienkaemper, Anne Shiu, and Zev Woodstock. "Obstructions to convexity in neural codes". Advances in Applied Mathematics 85 (2017), pp. 31–59.
- [18] Katherine Morrison, Anda Degeratu, Vladimir Itskov, and Carina Curto. Diversity of emergent dynamics in competitive threshold-linear networks. 2022. URL: https://arxiv.org/ abs/1605.04463.
- [19] Erfan Nozari and Jorge Cortés. "Hierarchical selective recruitment in linear-threshold brain networks—Part I: Single-layer dynamics and selective inhibition". *IEEE Transactions on Automatic Control* 66.3 (2020), pp. 949–964.
- [20] Alexey Pavlov, Nathan van de Wouw, and Henk Nijmeijer. "Convergent piecewise affine systems: analysis and design part i: continuous case". Proceedings of the 44th IEEE Conference on Decision and Control. IEEE. 2005, pp. 5391–5396.
- [21] Kanaka Rajan and Larry F Abbott. "Eigenvalue spectra of random matrices for neural networks". *Physical review letters* 97.18 (2006), p. 188104.
- [22] S. Roman. Lattices and Ordered Sets. Springer New York, 2010.
- [23] Magnus Tournoy and Brent Doiron. "A Step Towards Uncovering The Structure of Multistable Neural Networks". arXiv:2210.03241 (2022).
- [24] X. Xie, R. H. Hahnloser, and H.S. Seung. "Selectively grouping neurons in recurrent networks of lateral inhibition". *Neural Comput.* 14 (2002), pp. 2627–2646.