# THE MEAN NUMBER OF 2-TORSION ELEMENTS IN THE CLASS GROUPS OF CUBIC ORDERS

#### ASHVIN A. SWAMINATHAN

ABSTRACT. We determine the mean number of 2-torsion elements in class groups of cubic orders, when such orders are enumerated by discriminant. Specifically, we prove that when isomorphism classes of totally real (resp., complex) cubic orders are enumerated by discriminant, the average 2-torsion in the class group is  $1 + \frac{1}{4} \times \frac{\zeta(2)}{\zeta(4)}$  (resp.,  $1 + \frac{1}{2} \times \frac{\zeta(2)}{\zeta(4)}$ ). In particular, we find that the average 2-torsion in the class group increases when one ranges over all orders in cubic fields instead of restricting to the subfamily of rings of integers of cubic fields, where the average 2-torsion in the class group was first determined in work of Bhargava to be  $\frac{5}{4}$  (resp.,  $\frac{3}{2}$ ).

By work of Bhargava–Varma, proving this result amounts to obtaining an asymptotic count of the number of "reducible"  $SL_3(\mathbb{Z})$ -orbits on the space  $\mathbb{Z}^2 \otimes_{\mathbb{Z}} Sym^2 \mathbb{Z}^3$  of  $3 \times 3$  symmetric integer matrices having bounded invariants and satisfying local conditions. In this paper, we resolve the generalization of this orbit-counting problem where the dimension 3 is replaced by any fixed odd integer  $N \geq 3$ . More precisely, we determine asymptotic formulas for the number of reducible  $SL_N(\mathbb{Z})$ -orbits on  $\mathbb{Z}^2 \otimes_{\mathbb{Z}} Sym^2 \mathbb{Z}^N$  satisfying general infinite sets of congruence conditions.

#### Contents

1.	Introduction		]
	1.1.	Main results	1
	1.2.	Historical context and related work	F
2.	. Algebraic preliminaries		(
	2.1.	Action of $SL_N$ on $W_N$	(
	2.2.	Action of subgroups of $SL_N$ on hyperplanes in $W_N$	7
	2.3.	Proof of Theorem 3	13
3.	Asymptotic formulae for the count of reducible orbits		13
	3.1.	Proof of Theorem 5	13
	3.2.	Change-of-variables formulas, and proof of Theorem 6	16
4.	Evaluation of local volumes for applications		18
	4.1.	Proof of Theorem 7	18
	4.2.	Proofs of Theorems 8.A and 8.B	19
Acknowledgments			21
References			91

### 1. Introduction

1.1. Main results. A striking result of Bhargava's thesis was the determination of the average size of the 2-torsion subgroup in the class groups of cubic number fields, enumerated by discriminant (see [1, Theorem 5.4] and [4, Theorem 5]). Specifically, Bhargava proved that when totally real (resp., complex) cubic fields are enumerated by the absolute values of their discriminants, the average 2-torsion in the class group is equal to  $\frac{5}{4}$  (resp.,  $\frac{3}{2}$ ). This result remains one of just a handful of cases of the heuristics on class groups of number fields, formulated in the foundational works of Cohen–Lenstra [15], Cohen–Martinet [16], and Malle [22], that have ever been proven.

The main result of this paper constitutes a generalization of Bhargava's breakthrough to the full family of all *orders* in cubic fields. We prove:

Date: November 17, 2022.

<sup>2020</sup> Mathematics Subject Classification. 11R29, 11R45 (primary), 11H55, 11E76 (secondary).

**Theorem 1.A.** When irreducible cubic orders  $\mathcal{O}$  over  $\mathbb{Z}$  are enumerated by the absolute values of their discriminants, the average size of  $Cl(\mathcal{O})[2]$  is:

(a) 
$$1 + \frac{1}{4} \times \frac{\zeta(2)}{\zeta(4)}$$
 for the family of totally real cubic orders  $\mathcal{O}$ , and

(b) 
$$1 + \frac{1}{2} \times \frac{\zeta(2)}{\zeta(4)}$$
 for the family of complex cubic orders  $\mathcal{O}$ .

Remark. Note that  $\frac{\zeta(2)}{\zeta(4)} = \frac{15}{\pi^2} \approx 1.51982 > 1$ .

Let  $N=2n+1\geq 3$  be an odd integer, and let  $U_N$  be the affine space over  $\mathbb{Z}$  whose R-points are given by binary N-ic forms over R for any  $\mathbb{Z}$ -algebra R. When N=3, the Delone–Faddeev–Levi correspondence [20] states that the map sending an irreducible binary cubic form  $f\in U_3(\mathbb{Z})$  to the ring  $R_f$  of global sections of the subscheme of  $\mathbb{P}^1_{\mathbb{Z}}$  cut out by f defines a bijection between the irreducible orbits of  $\mathrm{GL}_2(\mathbb{Z})$  on  $U_3(\mathbb{Z})$  and the set of isomorphism classes of orders in cubic number fields. In light of this, studying cubic orders amounts to studying irreducible integral binary cubic forms that lie in a fundamental region  $\mathcal{F}$  for the action of  $\mathrm{GL}_2(\mathbb{Z})$  on  $U_3(\mathbb{R})$ .

The region  $\mathcal{F}$  may be chosen so that the set of  $f \in \mathcal{F}$  with discriminant  $\operatorname{disc}(f) \ll X$  is approximately the set of  $f \in \mathcal{F}$  whose coefficients are all  $\ll X$ . Thus, we expect that the averages in Theorem 1.A should remain the same if we replace the family of cubic orders, enumerated by discriminant, with the family of cubic orders of the form  $R_f$ , where f runs through irreducible integral binary cubic forms enumerated by height; here we define the height H(f) of a binary form to be the maximum of the absolute values of its coefficients. Indeed, by modifying the proof of Theorem 1.A, we obtain the following variant:

**Theorem 1.B.** When forms  $f \in U_3(\mathbb{Z})$  are enumerated by height, the average size of  $Cl(R_f)[2]$  is:

(a) 
$$1 + \frac{1}{4} \times \frac{\zeta(2)}{\zeta(4)}$$
 for the family of totally real binary cubic forms  $f$ , and

(b) 
$$1 + \frac{1}{2} \times \frac{\zeta(2)}{\zeta(4)}$$
 for the family of complex binary cubic forms  $f$ .

To prove Theorems 1.A and 1.B, we utilize an orbit parametrization, discovered by Bhargava when N=3 (see [2, Theorem 4]), and generalized to all odd N by Wood (see [30, Theorem 1.3]). Fix an irreducible binary N-ic form  $f \in U_N(\mathbb{Z})$ , and assume f is primitive if N>3. Let  $K_f:=\mathbb{Q}\otimes_{\mathbb{Z}}R_f$ , and let  $W_N$  denote the affine space over  $\mathbb{Z}$  whose R-points are given by pairs of symmetric  $N\times N$  matrices over R. Then the Bhargava–Wood parametrization takes as input a pair  $(I,\delta)$ , where I is a 2-torsion ideal class of the order  $R_f$  and  $\delta \in K_f^\times/K_f^{\times 2}$  is a generator of  $I^2$  having square norm, and it produces as output the  $\mathrm{SL}_N(\mathbb{Z})$ -orbit of a pair  $(A,B)\in W_N(R)$  such that  $\mathrm{inv}(A,B):=(-1)^n\det(xA-yB)=f(x,y)$ .

The set of pairs  $(I, \delta)$  corresponding to a form  $f \in U_N(\mathbb{Z})$  naturally partitions into two subsets depending on whether or not  $\delta \equiv 1 \in K_f^{\times}/K_f^{\times 2}$ . Via the parametrization, pairs  $(I, \delta)$  with  $\delta \not\equiv 1$  (resp.,  $\delta \equiv 1$ ) correspond to so-called *irreducible* (resp., *reducible*)  $\mathrm{SL}_N(\mathbb{Z})$ -orbits on  $W_N(\mathbb{Z})$ . Here, (the  $\mathrm{SL}_N(\mathbb{Z})$ -orbit of) a pair  $(A, B) \in W_N(\mathbb{Z})$  is said to be reducible if, when A and B are viewed as symmetric bilinear forms over  $\mathbb{Q}$ , they share an isotropic space over  $\mathbb{Q}$  of maximal dimension. Geometrically, A and B may be viewed as defining a pair of quadric hypersurfaces in  $\mathbb{P}^{N-1}$ , and the condition that (A, B) is reducible is equivalent to stipulating that the (finite) Fano scheme parametrizing maximal linear spaces contained in the intersection of these two quadrics has a  $\mathbb{Q}$ -rational point.

<sup>&</sup>lt;sup>1</sup>Note that each cubic order  $\mathcal{O}$  in this family occurs infinitely many times, once for every f such that  $\mathcal{O} \simeq R_f!$ 

Note that if  $(I, \delta)$  is a pair with  $\delta \equiv 1$ , then I is the class of a 2-torsion ideal of  $R_f$  (i.e., a fractional ideal of  $R_f$  that squares to the unit ideal). Thus, the set of reducible  $\mathrm{SL}_N(\mathbb{Z})$ -orbits of pairs  $(A, B) \in \mathrm{inv}^{-1}(f) \cap W_N(\mathbb{Z})$  is in bijection with the 2-torsion subgroup of the ideal group  $\mathcal{I}(R_f)$ . If  $R_f$  is the maximal order in  $K_f$  (i.e.,  $R_f$  is integrally closed in  $K_f$ ), then  $\mathcal{I}(R_f)$  is torsion-free, and the only 2-torsion ideal is the trivial one. Thus, there is exactly one reducible orbit corresponding to f via the parametrization, and the problem of determining the average size of the 2-torsion in the class groups of maximal orders  $R_f$  amounts to counting just the irreducible orbits. A systematic method for counting irreducible orbits of representations (where the definition of "irreducible" depends on the representation) was developed by Bhargava in his thesis (see, e.g., [6]) and was later vastly streamlined in the seminal work of Bhargava and Shankar on Selmer groups of elliptic curves (see, e.g., [10]). Using this method to count just the irreducible orbits, Bhargava and Varma proved the following precursor to Theorem 1.A:

**Theorem 2.A** ([13, Theorem 2]). When irreducible cubic orders  $\mathcal{O}$  are enumerated by the absolute values of their discriminants, the average size of  $Cl(\mathcal{O})[2]$  is:

(a) 
$$1 + \frac{1}{4} \times \underset{\operatorname{disc}(\mathcal{O})>0}{\operatorname{Avg}} \# \mathcal{I}(\mathcal{O})[2]$$
 for the family of totally real cubic orders  $\mathcal{O}$ , and

(b) 
$$1 + \frac{1}{2} \times \underset{\text{disc}(\mathcal{O}) < 0}{\text{Avg}} \# \mathcal{I}(\mathcal{O})[2]$$
 for the family of complex cubic orders  $\mathcal{O}$ .

Just as Theorem 2.A was a precursor to Theorem 1.A, we have the following result of Ho, Shankar, and Varma, which holds for any N and was a precursor to Theorem 1.B in the case N = 3:

**Theorem 2.B** ([21, Theorem 6]). Let  $U_N(\mathbb{Z})^{(r)}$  be the set of irreducible forms  $f \in U_N(\mathbb{Z})$  having r real roots and  $\frac{N-r}{2}$  pairs of complex roots. When forms  $f \in U_N(\mathbb{Z})^{(r)}$ , primitive if N > 3, are enumerated by height, the average size of  $Cl(R_f)[2]$  is

$$1 + 2^{1 - \frac{N+r}{2}} \times \operatorname{Avg}_{\substack{f \in U_N(\mathbb{Z})^{(r)} \\ f \text{ prim. if } N > 3}} \# \mathcal{I}(R_f)[2].$$

To determine precise numerical values of the average 2-torsion in the class group for the families of orders considered in Theorems 2.A and 2.B, one needs to determine the average 2-torsion in their ideal groups. Via the Bhargava–Wood parametrization, this amounts to counting reducible  $\mathrm{SL}_N(\mathbb{Z})$ -orbits on  $W_N(\mathbb{Z})$ . Systematic methods for counting reducible orbits were not available until recently, when the author, in joint work with Shankar, Siad, and Varma, developed a new technique that applies to reducible orbits of many representations of importance in arithmetic statistics [25].

In the context of the action of  $\operatorname{SL}_N$  on  $W_N$ , our new technique proceeds according to the following series of steps. First, we prove in §3 (see Proposition 23) that the count of the reducible  $\operatorname{SL}_N(\mathbb{Z})$ -orbits on  $W_N(\mathbb{Z})$  is asymptotically equivalent to the count of the  $G_N(\mathbb{Z})$ -orbits on  $W_N^0(\mathbb{Z})$ , where  $G_N \subset \operatorname{SL}_N$  is a certain closed subgroup and  $W_N^0 \subset W_N$  is a certain hyperplane all of whose  $\mathbb{Q}$ -points are reducible (see §§2.2.4–2.2.5 for the precise definitions of  $G_N$  and  $W_N^0$ ). Second, we prove that the action of  $G_N$  on  $W_N^0$  satisfies the following strong local-to-global principle:

**Theorem 3.** Let  $f \in U_N(\mathbb{Z})$  be a binary N-ic form with nonzero discriminant. For each prime p, choose  $(A_p, B_p) \in \operatorname{inv}^{-1}(f) \cap W_N^0(\mathbb{Z}_p)$ . Then there exists  $(A, B) \in W_N^0(\mathbb{Z})$ , unique up to the action of  $G_N(\mathbb{Z})$ , such that (A, B) is  $G_N(\mathbb{Z}_p)$ -equivalent to  $(A_p, B_p)$  for each prime p.

To prove Theorem 3, we rely on [25, Theorem 22], which is a general result that provides criteria under which the integral orbits of a representation satisfy a local-to-global principle. Theorem 3 follows by verifying that the representation of  $G_N$  on  $W_N^0$  satisfies these criteria.

As a consequence of Theorem 3, counting  $G_N(\mathbb{Z})$ -orbits on  $W_N^0(\mathbb{Z})$  amounts to counting  $G_N(\mathbb{Z}_p)$ -orbits on  $W_N^0(\mathbb{Z}_p)$  for every prime p. In §3.1 we combine this local-to-global principle with a squarefree sieve for binary forms developed in work of Bhargava–Shankar–Wang [12] to obtain an asymptotic for the count of reducible  $\mathrm{SL}_N(\mathbb{Z})$ -orbits on  $W_N(\mathbb{Z})$  satisfying certain infinite families of congruence conditions, which we define precisely as follows:

**Definition 4.** We call a subset  $\mathfrak{S} \subset W_N^0(\mathbb{Z})$  a big family in  $W_N^0(\mathbb{Z})$  if  $\mathfrak{S} = \operatorname{inv}^{-1}(U_N(\mathbb{Z})^{(r)}) \cap W_N^0(\mathbb{Z}) \cap \bigcap_p \mathfrak{S}_p$ , where the sets  $\mathfrak{S}_p \subset W_N^0(\mathbb{Z}_p)$  satisfy the following properties:

- (1)  $\mathfrak{S}_p$  is  $G_N(\mathbb{Z}_p)$ -invariant and is the preimage under reduction modulo  $p^j$  of a nonempty subset of  $W_N^0(\mathbb{Z}/p^j\mathbb{Z})$  for some j > 0 for each p; and
- (2)  $\mathfrak{S}_p$  contains all elements  $(A, B) \in W_N^0(\mathbb{Z}_p)$  such that, for all  $p \gg 1$ , we have that Q(A, B) is a p-adic unit, where Q is a certain relative invariant for the action of  $G_N$  on  $W_N^0$  over  $\mathbb{Z}$  (see Proposition 10 for the precise definition of Q).

We call a subset  $S \subset W_N(\mathbb{Z})$  a big family in  $W_N(\mathbb{Z})$  if  $S = \operatorname{inv}^{-1}(U_N(\mathbb{Z})^{(r)}) \cap \bigcap_p S_p$ , where the sets  $S_p \subset W_N(\mathbb{Z}_p)$  are  $\operatorname{SL}_n(\mathbb{Z}_p)$ -invariant, and the intersections  $\mathfrak{S}_p = S_p \cap W_N^0(\mathbb{Z}_p)$  define a big family  $\mathfrak{S} = \operatorname{inv}^{-1}(U_N(\mathbb{Z})^{(r)}) \cap W_N^0(\mathbb{Z}) \cap \bigcap_p \mathfrak{S}_p$  in  $W_N^0(\mathbb{Z})$ .

Define the *height* of (A, B) by H(A, B) := H(inv(A, B)). The following result gives an asymptotic for the count of reducible  $SL_N(\mathbb{Z})$ -orbits of bounded height in a big family in  $W_N(\mathbb{Z})$ :

**Theorem 5.** Let X > 0, and let  $N_N^{(r)}(X)$  be the number of binary forms in  $U_N(\mathbb{Z})^{(r)}$  whose coefficients are of size at most X. Let  $S \subset \operatorname{inv}^{-1}(U_N(\mathbb{Z})^{(r)})$  be a big family in  $W_N(\mathbb{Z})$ . Then the number of reducible  $\operatorname{SL}_N(\mathbb{Z})$ -orbits on S of height less than X is given by

$$N_N^{(r)}(X) \times \prod_p \int_{f \in U_N(\mathbb{Z}_p)} \#\left(\frac{\operatorname{inv}^{-1}(f) \cap S_p \cap W_N^0(\mathbb{Z}_p)}{G_N(\mathbb{Z}_p)}\right) df + o(X^{N+1}),$$

where df is the Euclidean measure on  $U_N(\mathbb{Z}_p)$ , normalized so that  $\operatorname{Vol}(U_N(\mathbb{Z}_p)) = 1$ .

Theorem 5 expresses the asymptotic count of reducible  $\mathrm{SL}_N(\mathbb{Z})$ -orbits on S of bounded height in terms of a product of local integrals. The integrand of the integral at p — namely, the number of  $G_N(\mathbb{Z}_p)$ -orbits of pairs  $(A,B) \in S_p \cap W_N^0(\mathbb{Z}_p)$  with  $\mathrm{inv}(A,B) = f$  — appears to be quite difficult to evaluate in general, even for small degrees N and simple choices of the sets  $S_p$ . On the other hand, the integral at p can be rendered more tractable by performing a suitable change-of-variables, where instead of integrating over  $f \in U_N(\mathbb{Z}_p)$ , one integrates over  $(A,B) \in W_N^0(\mathbb{Z}_p)$ . Upon performing this change-of-variables at each prime p, we obtain the following variant of Theorem 5:

**Theorem 6.** Let X > 0, and let  $S \subset \operatorname{inv}^{-1}(U_N(\mathbb{Z})^{(r)})$  be a big family in  $W_N(\mathbb{Z})$ . Then the number of reducible  $\operatorname{SL}_N(\mathbb{Z})$ -orbits on S of height less than X is given by

$$N_N^{(r)}(X) \times \prod_p \xi_{p,n} \int_{w \in S_p \cap W_N^0(\mathbb{Z}_p)} |Q(w)|_p dw + o(X^{N+1}),$$

where  $\xi_{p,n} := (1-p^{-1})^{-1}(1-p^{-n-1})^{-1} \times \prod_{i=2}^n (1-p^{-i})^{-1}$  is a convergence factor,  $|-|_p$  denotes the usual p-adic absolute value, and where dw denotes the Euclidean measure on  $W_N^0(\mathbb{Z}_p)$ , normalized so that  $\operatorname{Vol}(W_N^0(\mathbb{Z}_p)) = 1$ .

The formulation of the asymptotic given in Theorem 6 is far more conducive to evaluation in specific examples. For instance, taking  $S_p = W_N(\mathbb{Z}_p)$  for every prime p, we obtain the following asymptotic formula for the total count of reducible  $\mathrm{SL}_N(\mathbb{Z})$ -orbits on  $W_N(\mathbb{Z})$ :

**Theorem 7.** Let X > 0. Then the number of reducible  $SL_N(\mathbb{Z})$ -orbits on  $inv^{-1}(U_N(\mathbb{Z})^{(r)})$  of height less than X is given by

$$N_N^{(r)}(X) \times \prod_{i=2}^N \zeta(i) + o(X^{N+1}).$$

Strikingly, Theorem 7 implies that the average number of reducible orbits lying above an integral binary N-ic form is equal to  $\prod_{i=2}^N \zeta(i)$ , which is simply the fundamental volume of the group  $\mathrm{SL}_N$  (i.e., the volume of  $\mathrm{SL}_N(\mathbb{Z}) \backslash \mathrm{SL}_N(\mathbb{R})$  with respect to Haar measure on  $\mathrm{SL}_N(\mathbb{R})$ , suitably normalized). This phenomenon — that the average number of reducible orbits lying above a given set of invariants equals the fundamental volume of the group — holds for many other representations of interest in arithmetic statistics. Indeed, as proven in [25, Theorem 1], this holds for the representation of the split orthogonal group on the space of N-ary quadratic forms for every integer  $N \geq 3$ , odd or even; furthermore, as explained in [25, Question 2 and subsequent discussion], this also holds for the representations of  $\mathrm{GL}_2$  on binary cubic and quartic forms.

We now turn our attention to applying Theorem 6 to prove our main results on class group statistics, namely Theorems 1.A and 1.B. We say that a pair  $(A, B) \in W_N(\mathbb{Z}_p)$  is projective if it corresponds to a 2-torsion ideal class under the Bhargava–Wood parametrization. As explained in Definition 29 (see §4.2, to follow), the condition of projectivity is given by a congruence condition modulo p for each prime p.

It is possible to determine the proportion of projective elements of  $W_N(\mathbb{Z}_p)$  for each prime p; this was essentially achieved in [13, §6] for N=3 and [21, §6.1] for  $N\geq 3$ . On the other hand, to apply Theorem 6 it is necessary to determine the proportion of projective elements of  $W_N^0(\mathbb{Z}_p)$  having a specified Q-invariant, and this appears to be intractable in general. Nonetheless, when N=3, the condition of being projective is not too complicated, and the relevant p-adic integrals can be evaluated. We thus obtain the following result giving the average 2-torsion in the ideal groups of cubic orders enumerated by discriminant:

**Theorem 8.A.** When either totally real or complex irreducible cubic orders  $\mathcal{O}$  are enumerated by the absolute values of their discriminants, the average 2-torsion in the ideal group is equal to  $\frac{\zeta(2)}{\zeta(4)}$ .

We also simultaneously obtain the following variant of Theorem 8.A for the family of cubic orders defined by binary cubic forms enumerated by height:

**Theorem 8.B.** When forms  $f \in U_3(\mathbb{Z})^{(r)}$  are enumerated by height, the average 2-torsion in the ideal group is equal to  $\frac{\zeta(2)}{\zeta(4)}$ .

Theorem 1.A (resp., Theorem 1.B) now follows immediately upon combining Theorem 2.A (resp., Theorem 2.B in the case N=3) with Theorem 8.A (resp., Theorem 8.B). As for orders defined by binary forms of higher odd degree, we obtain the following weaker result by simply combining Theorem 7 with Theorem 2.B:

**Theorem 9.** Let N > 3. When primitive forms  $f \in U_N(\mathbb{Z})^{(r)}$  are enumerated by height, the average sizes of  $\mathcal{I}(R_f)[2]$  and  $\mathrm{Cl}(R_f)[2]$  are bounded.

1.2. Historical context and related work. The problem of studying the arithmetic statistics of non-maximal orders in number fields dates back to the seminal work of Davenport [17,18], who determined the density of discriminants of orders in cubic fields (i.e., he obtained an asymptotic formula for the number of cubic orders having bounded discriminant). This result was generalized in work of Bhargava, Shankar, and Tsimerman [11, Theorem 8], who determined the density of discriminants of cubic orders satisfying general infinite sets of local specifications. As for orders of higher degree, Bhargava used the parametrizations of quartic/quintic orders that he developed in his thesis [3,5] to determine asymptotic formulas for the number of orders having bounded discriminant in quartic/quintic fields with Galois group equal to the full symmetric group [4,6].

More recently, progress has been made toward understanding the distribution of class groups of orders in number fields. The first result in this direction is due to Bhargava and Varma [14], who determined the average 3-torsion in the class groups of quadratic orders. Specifically, they showed that when real (resp., complex) quadratic orders are enumerated by discriminant, the average 3-torsion in the class group is given by  $1 + \frac{1}{3} \times \frac{\zeta(2)}{\zeta(3)}$  (resp.,  $1 + \frac{\zeta(2)}{\zeta(3)}$ ). This extends an earlier result of Davenport and Heilbronn [19], who determined the corresponding average to be  $\frac{4}{3}$  (resp., 2) for the family of maximal quadratic orders (i.e., rings of integers of quadratic number fields).

As explained in §1.1, Bhargava and Varma proved Theorem 2.A, which describes the average 2-torsion in the class groups of cubic orders in terms of the average 2-torsion in their ideal groups [13], and their result was generalized to orders defined by binary forms of any odd degree  $N \geq 3$  by Ho, Shankar, and Varma, who proved Theorem 2.B. A similar result was proven for monogenic orders — i.e., orders defined by monic binary forms — of odd degree by Siad in [26, Theorem 9]. In the case of monogenic orders of odd degree N, the problem of counting 2-torsion ideals boils down to a problem of counting reducible orbits of the aforementioned representation of the split orthogonal group acting on the space of N-ary quadratic forms. As explained in §1.1, asymptotics akin to Theorems 5–7 were proven for reducible orbits of this representation in [25]; these asymptotics were then applied in [29, §5.6] with N=3 to determine the average 2-torsion in the ideal groups of monogenic cubic orders. Combining this with the aforementioned result of Siad, we deduced (see [29, Theorem 173]) that when monogenic cubic orders are enumerated by height, the average 2-torsion in the class group is given by  $\frac{5}{4} + \frac{1}{4} \times \frac{\zeta(2)}{\zeta(3)}$  (resp.,  $\frac{3}{2} + \frac{1}{2} \times \frac{\zeta(2)}{\zeta(3)}$ ). This extends an earlier result of Bhargava, Hanke, and Shankar [9], who determined the corresponding average to be  $\frac{3}{2}$  (resp., 2) for the family maximal monogenic cubic orders.

We note that the problem of counting reducible  $SL_N(\mathbb{Z})$ -orbits on  $W_N(\mathbb{Z})$  was first considered by Bhargava, Shankar, and Wang in [12], where they used geometry-of-numbers methods to determine upper bounds of roughly the correct order of magnitude on the number of reducible orbits having large Q-invariant. They applied these upper bounds to prove the squarefree sieve for binary forms alluded to in §1.1; this sieve serves as a key ingredient in the proofs of our main theorems. Thus, we indirectly use *upper bounds* for the count of reducible orbits to obtain *precise asymptotics* for this count!

It is natural to ask what can be said for binary forms of even degree. We expect that the methods of [27, 28] can be adapted to determine the average 2-torsion in the class group of orders defined by binary forms of even degree in terms of the average 2-torsion in the ideal group; however, in the even-degree setting, the parametrization of 2-torsion ideal classes is significantly more complicated (e.g., it depends on the leading coefficient of the binary forms under consideration, and one must count reducible orbits in multiple families of representations). It is also natural to ask whether analogues of Theorems 5–7 can be proven for the action of  $SL_N$  on pairs of  $N \times N$  symmetric matrices, where  $N = 2n \ge 4$  is even. This representation was studied by Bhargava in [7] (cf. the closely related work of Bhargava, Gross, and Wang [8]), where he shows that this representation does *not* possess an analogous notion of reducibility and is thus able to determine asymptotics for the count of its integral orbits.

## 2. Algebraic preliminaries

In this section, we introduce the representation of  $SL_N$  on  $W_N$  and prove several useful results about the action of various subgroups of  $SL_N$  on various hyperplanes in  $W_N$ . In particular, we define and study the subgroup  $G_N$  and the hyperplane  $W_N^0$  referenced in §1.1. We conclude by proving Theorem 3, which gives a local-to-global principle for the action of  $G_N$  on  $W_N^0$ .

2.1. Action of  $SL_N$  on  $W_N$ . For integers m, m' > 0, let  $Mat_{m \times m'}$  denote the affine scheme over  $\mathbb{Z}$  whose R-points are given by the set of  $m \times m'$  matrices with entries in R for any  $\mathbb{Z}$ -algebra R.

As in §1.1, let  $N=2n+1\geq 3$  be an odd integer, and let  $W_N$  be the space of pairs of  $N\times N$  symmetric matrices (i.e., we have  $W_N(R)=R^2\otimes_R\operatorname{Sym}^2R^N$  for any  $\mathbb{Z}$ -algebra R). The space  $W_N$  has a natural structure of  $\operatorname{SL}_N$ -representation given as follows: for any  $g\in\operatorname{SL}_N(R)$  and any  $(A,B)\in W_N(R)$ , let

$$g \cdot (A, B) = (gAg^T, gBg^T) \in W_N(R),$$

where for a matrix M, we denote by  $M^T$  its transpose.

Let  $U_N$  be the affine scheme over  $\mathbb{Z}$  whose R-points are binary N-ic forms over R for any  $\mathbb{Z}$ -algebra R. Define a map inv:  $W_N \to U_N$  as follows: given a pair  $(A, B) \in W_N(R)$ , we set

$$inv(A, B) := (-1)^n \det(xA - yB) \in U_N(R).$$

One readily checks that the coefficients of  $\operatorname{inv}(A,B)$  are  $\operatorname{SL}_N$ -invariant; in fact, it is known [24] that if  $f_i \colon W_N \to \mathbb{A}^1$  is the map that takes  $(A,B) \in W_N(R)$  and returns the  $x^{N-i}y^i$ -coefficient of  $\operatorname{inv}(A,B)$ , then the ring of polynomial invariants for the action of  $\operatorname{SL}_N$  on  $W_N$  is freely generated by the functions  $f_i$  for  $i \in \{0,\ldots,N\}$ . In [21, §4.1], an explicit algebraic section  $\sigma_0$  defined over  $\mathbb{Z}$  was constructed for the map inv:  $W_N \to U_N$  (by "algebraic" and "defined over  $\mathbb{Z}$ ," we mean that the matrix entries of  $\sigma_0$  are polynomials with integer coefficients).

Let K be a field. As mentioned in §1.1, a pair  $(A, B) \in W_N(K)$  is said to be reducible over K if the symmetric bilinear forms defined by A and B share a maximal (i.e., n-dimensional) isotropic space, and irreducible otherwise. The condition of being (ir)reducible in evidently  $SL_N(K)$ -invariant, so we may speak of the (ir)reducible  $SL_N(K)$ -orbits on  $W_N(K)$ . If R is an integral domain with field of fractions K, we say that an  $SL_N(R)$ -orbit on  $W_N(R)$  is (ir)reducible if the corresponding property holds for the  $SL_N(K)$ -orbit containing it.

In the case N=3, we will have occasion to combine the action of  $\operatorname{SL}_3$  on  $W_3$  with the action of  $\operatorname{GL}_2$  on  $U_3$ . Recall that  $\operatorname{GL}_2$  acts on  $U_3$  via linear change-of-variable; i.e., given  $\gamma \in \operatorname{GL}_2(R)$  and  $f \in U_3(R)$ , we have  $(\gamma \cdot f)(x,y) = f((x,y) \cdot \gamma)$ . The ring of polynomial invariants for the action of  $\operatorname{GL}_2$  on  $U_3$  is freely generated by a single element known as the discriminant. We have an action of  $\operatorname{GL}_2 \times \operatorname{SL}_3$  on  $W_3$  defined as follows: given  $\gamma = \left[ \begin{smallmatrix} a & b \\ c & d \end{smallmatrix} \right] \in \operatorname{GL}_2(R)$ ,  $g \in \operatorname{SL}_3(R)$ , and  $(A,B) \in W_N(R)$ , we set

$$(\gamma, g) \cdot (A, B) = (g(aA - bB)g^T, g(cA - dB)g^T).$$

The map inv:  $W_3 \to U_3$  is equivariant for the  $GL_2$ -action, and the ring of polynomial invariants for the action of  $GL_2 \times SL_3$  on  $W_3$  is generated by the function obtained by postcomposing the map inv with the discriminant map  $U_3 \to \mathbb{A}^1$ .

- 2.2. Action of subgroups of  $SL_N$  on hyperplanes in  $W_N$ . For a matrix M, denote its row-i, column-j entry by  $M_{ij}$ . We now introduce three hyperplanes in  $W_N$  and study the actions of certain subgroups of  $SL_N$  on them.
- 2.2.1. Action of  $\operatorname{SL}_n \times \operatorname{SL}_{n+1}$  on  $W_N^{\operatorname{top}}$ . Let  $W_N^{\operatorname{top}} \subset W_N$  denote the hyperplane whose R-points are given by

$$W_N^{\text{top}}(R) := \{ (A, B) \in W_N(R) : A_{ij} = B_{ij} = 0 \text{ if } i, j \le n \text{ or } i, j \ge n+1 \}$$

for any  $\mathbb{Z}$ -algebra R. When it is convenient, we think of elements of  $W_N^{\text{top}}$  as being pairs of  $n \times (n+1)$  matrices, by considering just the top-right  $n \times (n+1)$  blocks.

Let  $\operatorname{SL}_n \times \operatorname{SL}_{n+1} \subset \operatorname{SL}_N$  be the subgroup consisting of block-diagonal matrices, where the diagonal consists of one  $n \times n$  block followed by one  $(n+1) \times (n+1)$  block, both having determinant 1. The action of  $\operatorname{SL}_N$  on  $W_N$  restricts to an action of  $\operatorname{SL}_n \times \operatorname{SL}_{n+1}$  on  $W_N^{\text{top}}$ .

We now describe the action of  $\mathrm{SL}_n \times \mathrm{SL}_{n+1}$  on  $W_N^{\mathrm{top}}$  over subfields  $K \subset \mathbb{C}$ :

**Proposition 10.** The ring of polynomial invariants of the action of  $SL_n \times SL_{n+1}$  on  $W_N^{\text{top}}$  is generated by a single polynomial Q such that for any subfield  $K \subset \mathbb{C}$  and any  $Q_0 \in K^{\times}$ , the set

 $Q^{-1}(Q_0) \subset W_N^{\text{top}}(K)$  is nonempty and consists of a single  $(\operatorname{SL}_n \times \operatorname{SL}_{n+1})(K)$ -orbit. Moreover, the stabilizer of any element of this orbit is trivial.

Proof. We first consider the case  $K = \mathbb{C}$ . Here, the proposition amounts to stating that the representation of  $(\mathrm{SL}_n \times \mathrm{SL}_{n+1})(\mathbb{C})$  on  $W_N^{\mathrm{top}}(\mathbb{C})$  is "prehomogeneous," and the proof is identical to that of [12, Proposition 3.1]. The main observation is that, for any integer  $n_0 \geq 2$ , the representation of  $(\mathrm{SL}_n \times \mathrm{SL}_{n+1})(\mathbb{C})$  on  $W_N^{\mathrm{top}}(\mathbb{C})$  in the case  $n = n_0$  is related to the representation of  $(\mathrm{SL}_n \times \mathrm{SL}_{n+1})(\mathbb{C})$  on  $W_N^{\mathrm{top}}(\mathbb{C})$  in the case  $n = n_0 - 1$  by what Sato and Kimura call a "castling transform" (see [23, §2]). Since the property of being prehomogeneous is preserved under castling transforms, it suffices to prove the lemma in the case n = 1, where the action of  $(\mathrm{SL}_n \times \mathrm{SL}_{n+1})(\mathbb{C})$  on  $W_N^{\mathrm{top}}(\mathbb{C})$  may be identified with the action of  $\mathrm{SL}_2(\mathbb{C})$  on  $\mathrm{Mat}_{2\times 2}(\mathbb{C})$  by right-multiplication. The desired polynomial invariant function Q for this action is given simply by the determinant.

We now describe the invariant Q for n>1. Given a pair  $(A,B)\in W_N^{\mathrm{top}}(R)$ , we take Q(A,B) to be the "hyperdeterminant" of (A,B), which is defined explicitly as follows. For each  $i\in\{1,\ldots,n+1\}$ , let  $A^{(i)}$  and  $B^{(i)}$  be the  $n\times n$  matrices obtained from A and B, respectively, by considering the top-right  $n\times (n+1)$  block and deleting the  $i^{\mathrm{th}}$  column. Then Q(A,B) is the determinant of the  $(n+1)\times (n+1)$  matrix whose row-i, column-j entry is the  $x^{n-j+1}y^{j-1}$ -entry of the binary n-ic form  $(-1)^{i+1}\det(xA^{(i)}-yB^{(i)})$ . That  $Q^{-1}(Q_0)$  is nonempty for any  $Q_0\in K^\times$  follows by examining the following pair of matrices in  $W_N^{\mathrm{top}}(K)$ , where empty entries are used to denote zeros:

$$\begin{pmatrix}
\begin{bmatrix}
& & & 1 \\
& & 1 \\
& & \ddots & \\
& 1 & & 
\end{bmatrix}, \begin{bmatrix}
& & \pm Q_0 \\
& & 1 \\
& & \ddots & \\
& 1 & & 
\end{bmatrix}
\end{pmatrix}$$

The pair in (1) evidently has Q-invariant  $Q_0$  or  $-Q_0$ , depending on the choice of sign of the entry " $\pm Q_0$ ." In fact, this pair gives an explicit section of the map  $Q: W_N^{\text{top}} \to \mathbb{A}^1$ , defined over  $\mathbb{Z}$ .

As for the claim about stabilizers, the action of  $\mathrm{SL}_2(\mathbb{C})$  by right-multiplication on matrices of nonzero determinant in  $\mathrm{Mat}_{2\times 2}(\mathbb{C})$  obviously has trivial stabilizers, and stabilizers are preserved under castling transforms. Lastly, the case where  $K\subset\mathbb{C}$  is any subfield follows from the case where  $K=\mathbb{C}$  by [25, Lemma 23].

2.2.2. Action of  $L_N$  on  $W_N^{\text{top},0}$ . Let  $W_N^{\text{top},0} \subset W_N^{\text{top}}$  be the hyperplane whose R-points are given by  $W_N^{\text{top},0}(R) := \{(A,B) \in W_N^{\text{top}}(R) : A_{ij} = 0 \text{ if } i+j \leq 2n+1 \text{ and } B_{ij} = 0 \text{ if } i+j \leq 2n\}$ 

for any  $\mathbb{Z}$ -algebra R. Let  $L_N \subset \operatorname{SL}_n \times \operatorname{SL}_{n+1}$  be the lower-triangular subgroup. Then the action of  $\operatorname{SL}_n \times \operatorname{SL}_{n+1}$  on  $W_N^{\operatorname{top}}$  restrict to an action of  $L_N$  on  $W_N^{\operatorname{top},0}$ . We now describe this restricted action over subfields  $K \subset \mathbb{C}$ :

**Proposition 11.** Let  $K \subset \mathbb{C}$  be a subfield, and let  $Q_0 \in K^{\times}$ . Then the set  $Q^{-1}(Q_0) \cap W_N^{\text{top},0}(K)$  is nonempty and consists of a single  $L_N(K)$ -orbit. Moreover, the stabilizer of any element of this orbit is trivial.

*Proof.* The nonemptiness statement follows immediately from the existence of the explicit section (1), which has image contained in  $W_N^{\text{top,0}}$ . The statement about stabilizers is obvious.

As for the transitivity statement, take  $(A, B) \in Q^{-1}(Q_0) \cap W_N^{\text{top},0}(K)$ . It suffices to show that (A, B) is  $L_N(K)$ -equivalent to (1). First, by replacing (A, B) with a suitable translate under the action of a diagonal element in  $L_N(K)$ , we may assume that  $A_{ij} = 1$  for all i + j = N + 1 and that  $B_{ij} = 1$  for all i + j = N, except when (i, j) = (1, N - 1), in which case  $B_{1(N-1)} = \pm Q_0$ . Now, call a lower-triangular unipotent element  $g \in \text{SL}_N(K)$  elementary if  $g_{ij} = 0$  for all but one pair (i, j) with  $i \geq j$ . It is easy to verify by inspection that, by hitting (A, B) with a suitable sequence of

elementary unipotent elements of  $L_N(K)$ , we can successively clear out the values of the following matrix entries:

(2) 
$$B_{1(2n+1)}, A_{2(2n+1)}, B_{2(2n)}, B_{2(2n+1)}, \dots, A_{k(2n+3-k)}, \dots, A_{k(2n+1)}, B_{k(2n+2-k)}, \dots, B_{k(2n+1)}, \dots, A_{n(2n+1)}, B_{n(n+2)}, \dots, B_{n(2n+1)}.$$

We may thus assume that the matrix entries of (A, B) that are listed in (2) are all equal to (1). But then (A, B) is equal to (1), which is sufficient.

**Corollary 12.** If two elements of  $Q^{-1}(Q_0) \cap W_N^{\text{top},0}(K)$  are equivalent under the action of  $g \in (\operatorname{SL}_n \times \operatorname{SL}_{n+1})(K)$ , then we have  $g \in L_N(K)$ .

Proof. Suppose for some  $g_1 \in (\operatorname{SL}_n \times \operatorname{SL}_{n+1})(K)$  and elements  $(A_1, B_1), (A_2, B_2) \in Q^{-1}(Q_0) \cap W_N^{\operatorname{top},0}(K)$  we have  $g_1 \cdot (A_1, B_1) = (A_2, B_2)$ . By Proposition 11, there exists  $g_2 \in L_N(K)$  such that  $g_2 \cdot (A_1, B_1) = (A_2, B_2)$ , so  $g_2^{-1}g_1$  stabilizes  $(A_1, B_1)$ , but by Proposition 10, the stabilizer of  $(A_1, B_1)$  in  $(\operatorname{SL}_n \times \operatorname{SL}_{n+1})(K)$  is trivial, so  $g_1 = g_2$ , as necessary.

2.2.3. Fundamental domain for  $(\operatorname{SL}_n \times \operatorname{SL}_{n+1})(\mathbb{Z}_p) \curvearrowright W_N^{\operatorname{top}}(\mathbb{Z}_p)$ . Using the results of §§2.2.1–2.2.2, we now construct a fundamental domain for the action of  $(\operatorname{SL}_n \times \operatorname{SL}_{n+1})(\mathbb{Z}_p)$  on  $W_N^{\operatorname{top}}(\mathbb{Z}_p)$ ; this fundamental domain plays a crucial role in the proof of Theorem 7 (see §4.1).

We start by choosing a convenient partition of  $W_N^{\text{top}}(\mathbb{Z}_p)$  into subsets indexed by pairs of nonnegative-integer-vectors of length n. Given  $\vec{a} = (a_1, \ldots, a_n), \vec{b} = (b_1, \ldots, b_n) \in \mathbb{N}^n$ , define a subset  $\mathcal{L}_{\vec{a}\vec{b}}(p)$  as follows:

$$\mathcal{L}_{\vec{a},\vec{b}}(p) := (\mathrm{SL}_n \times \mathrm{SL}_{n+1})(\mathbb{Z}_p) \times \left\{ (A,B) \in W_N^{\mathrm{top},0}(\mathbb{Z}_p) : \begin{array}{l} \nu_p(A_{ij}) = a_{N+1-j} \text{ for } (i,j) \text{ with } i+j=N+1, \\ \nu_p(B_{ij}) = b_i \text{ for } (i,j) \text{ with } i+j=N \end{array} \right\}$$

A calculation shows that every  $(A, B) \in \mathcal{L}_{\vec{a}, \vec{b}}(p)$  has Q-invariant  $|Q(A, B)|_p^{-1} = \prod_{i=1}^n p^{(n+1-i)a_i+ib_i}$ . The following result gives the desired fundamental domain in terms of the sets  $\mathcal{L}_{\vec{a}, \vec{b}}(p)$ :

**Proposition 13.** A fundamental domain for the action of  $(\operatorname{SL}_n \times \operatorname{SL}_{n+1})(\mathbb{Z}_p)$  on the set  $\{(A, B) \in W_N^{\operatorname{top}}(\mathbb{Z}_p) : Q(A, B) \neq 0\}$  is given by

(3) 
$$\bigsqcup_{\vec{a}, \vec{b} \in \mathbb{N}^n} \left\{ (A, B) \in \mathcal{L}_{\vec{a}, \vec{b}}(p) \cap W_N^{\text{top}, 0}(\mathbb{Z}_p) : \begin{array}{c} \nu_p(A_{ij}) \leq a_{N+1-j} \text{ for } (i, j) \text{ with } i+j > N+1, \\ \nu_p(B_{ij}) \leq b_i \text{ for } (i, j) \text{ with } i+j > N \end{array} \right\}$$

*Proof.* The first step is to show that the action of  $(\mathrm{SL}_n \times \mathrm{SL}_{n+1})(\mathbb{Z}_p)$  can be used to move points in  $W_N^{\mathrm{top}}(\mathbb{Z}_p)$  into the hyperplane  $W_N^{\mathrm{top},0}(\mathbb{Z}_p)$ :

**Lemma 14.** Let  $Q_0 \in \mathbb{Z}_p \setminus \{0\}$ . Then every  $(\operatorname{SL}_n \times \operatorname{SL}_{n+1})(\mathbb{Z}_p)$ -orbit on  $Q^{-1}(Q_0) \subset W_N^{\operatorname{top}}(\mathbb{Z}_p)$  meets  $W_N^{\operatorname{top},0}(\mathbb{Z}_p)$ .

Proof of Lemma 14. The proof is similar to that of [29, Lemma 206]. Take  $(A, B) \in Q^{-1}(Q_0) \cap W_N^{\text{top}}(\mathbb{Z}_p)$ . Then there exists  $g \in (\mathrm{SL}_n \times \mathrm{SL}_{n+1})(\mathbb{Q}_p)$  such that  $g \cdot (A, B) \in W_N^{\text{top},0}(\mathbb{Z}_p)$ , because the section (1) of Q is defined over  $\mathbb{Z}$  with image contained in  $W_N^{\text{top},0}$ , and because the elements of the set  $Q^{-1}(Q_0) \cap W_N^{\text{top}}(\mathbb{Z}_p)$  belong to the same  $(\mathrm{SL}_n \times \mathrm{SL}_{n+1})(\mathbb{Q}_p)$ -orbit by Proposition 10.

By the p-adic Iwasawa decomposition, we have that

$$(\mathrm{SL}_n \times \mathrm{SL}_{n+1})(\mathbb{Q}_p) = L_N(\mathbb{Q}_p)(\mathrm{SL}_n \times \mathrm{SL}_{n+1})(\mathbb{Z}_p),$$

so there exists  $g_1 \in L_N(\mathbb{Q}_p)$  and  $g_2 \in (\mathrm{SL}_n \times \mathrm{SL}_{n+1})(\mathbb{Z}_p)$  such that  $g = g_1g_2$ . But  $L_N$  acts on  $W_N^{\mathrm{top},0}$  (see §2.2.2 for further details), so  $g_2 \cdot (A,B) \in W_N^{\mathrm{top},0}(\mathbb{Q}_p)$ . Since  $g_2 \in (\mathrm{SL}_n \times \mathrm{SL}_{n+1})(\mathbb{Z}_p)$  and  $(A,B) \in W_N^{\mathrm{top},0}(\mathbb{Z}_p)$ , we must in fact have that  $g_2 \cdot (A,B) \in W_N^{\mathrm{top},0}(\mathbb{Z}_p)$ .

Now, by Lemma 14 and Corollary 12, it suffices to show that (3) is a fundamental domain for the action of  $L_N(\mathbb{Z}_p)$  on  $W_N^{\text{top,0}}(\mathbb{Z}_p)$ , but this follows by adapting the proof of Proposition 11 to work over  $\mathbb{Z}_p$ . (I.e., instead of using the elementary unipotent transformations to make certain matrix entries zero, we use these transformations to reduce these matrix entries modulo  $p^{a_1}, \ldots, p^{a_n}, p^{b_1}, \ldots, p^{b_n}$ .)

2.2.4. The subgroup  $G_N \subset \operatorname{SL}_N$ . Let  $G_N \subset \operatorname{SL}_N$  be the subgroup whose R-points are given by

$$G_N(R) := \{g \in \operatorname{SL}_N(R) : g_{ij} = 0 \text{ for all } (i,j) \text{ such that } i \leq n \text{ and } j \geq n+1 \}$$

for any  $\mathbb{Z}$ -algebra R.

An algebraic group G is said to have class number 1 over  $\mathbb{Q}$  if the group  $G(\mathbb{A}_{\mathbb{Q}})$  of adelic rational points is the "Frobenius" product of the subgroup  $G(\mathbb{Q})$  of rational points with the subgroup  $G(\mathbb{A}_{\mathbb{Z}})$  of adelic integral points —i.e., we have  $G(\mathbb{A}_{\mathbb{Q}}) = G(\mathbb{Q})G(\mathbb{A}_{\mathbb{Z}})$ . The following result establishes that this property holds for  $G = G_N$ :

**Proposition 15.** The algebraic group  $G_N$  has class number 1 over  $\mathbb{Q}$ .

*Proof.* The idea of the proof is to realize  $G_N$  as the Frobenius product of two subgroups, each of which has class number 1 over  $\mathbb{Q}$ , and then to use this product structure to deduce that  $G_N$  itself has class number 1 over  $\mathbb{Q}$ .

Let  $H_1$  be the lower-triangular unipotent subgroup of  $G_N$  whose R-points are given by

$$H_1(R) := \left\{ h \in G_N(R) : \begin{array}{l} h_{ii} = 1 \text{ for all } i \text{ and} \\ h_{jk} = 0 \text{ if } j < k \text{ or } j > k \ge n+1 \end{array} \right\}$$

for any  $\mathbb{Z}$ -algebra R, and let  $H_2$  be the block-diagonal subgroup of  $G_N$  whose R-points are given by

$$H_2(R) := \{ g \in \operatorname{GL}_n(R) \times \operatorname{GL}_{n+1}(R) : \det g = 1 \} \subset G_N(R).$$

**Lemma 16.** The algebraic group  $G_N$  is the Frobenius product of its subgroups  $H_1$  and  $H_2$ ; i.e., for any  $\mathbb{Z}$ -algebra R we have  $G_N(R) = H_1(R)H_2(R)$ .

Proof of Lemma 16. Take  $g \in G_N(R)$ , and write it in box form as follows:

$$g = \left[ \begin{array}{c|c} g' & 0 \\ \hline g''' & g'' \end{array} \right],$$

where  $g' \in GL_n(R)$ ,  $g'' \in GL_{n+1}(R)$ , and  $g''' \in Mat_{(n+1)\times n}(R)$ . Then it is easy to see that

$$g = \begin{bmatrix} \operatorname{id} & 0 \\ g'''g'^{-1} & \operatorname{id} \end{bmatrix} \times \begin{bmatrix} g' & 0 \\ 0 & g'' \end{bmatrix},$$

where by "id" (resp., "0") we mean the identity (resp., zero) matrix of the relevant dimensions.

It is easy to verify that the group  $H_1$  has class number 1 over  $\mathbb{Q}$ ; that  $H_2$  has class number 1 over  $\mathbb{Q}$  follows immediately from the fact that the same holds for the groups  $GL_n$  and  $GL_{n+1}$ . The next lemma gives a criterion under which the Frobenius product of two groups of class number 1 over  $\mathbb{Q}$  is itself a group of class number 1 over  $\mathbb{Q}$ :

**Lemma 17.** Let  $\Gamma$  be an algebraic group over  $\mathbb{Z}$  that is the Frobenius product of two sub-algebraic-groups  $\Gamma_1$  and  $\Gamma_2$ , both of which have class number 1 over  $\mathbb{Q}$ , and suppose that  $\Gamma_1(\mathbb{A}_{\mathbb{Z}})\Gamma_2(\mathbb{Q}) \subset \Gamma_2(\mathbb{Q})\Gamma_1(\mathbb{A}_{\mathbb{Q}})$ . Then  $\Gamma$  has class number 1 over  $\mathbb{Q}$ .

Proof of Lemma 17. Clearly,  $\Gamma(\mathbb{A}_{\mathbb{Q}}) \supset \Gamma(\mathbb{Q})\Gamma(\mathbb{A}_{\mathbb{Z}})$ . As for the reverse inclusion, we have that

$$\begin{split} \Gamma(\mathbb{A}_{\mathbb{Q}}) &= \Gamma_{1}(\mathbb{A}_{\mathbb{Q}})\Gamma_{2}(\mathbb{A}_{\mathbb{Q}}) = \Gamma_{1}(\mathbb{Q})\Gamma_{1}(\mathbb{A}_{\mathbb{Z}})\Gamma_{2}(\mathbb{Q})\Gamma_{2}(\mathbb{A}_{\mathbb{Z}}) \\ &\subset \Gamma_{1}(\mathbb{Q})\Gamma_{2}(\mathbb{Q})\Gamma_{1}(\mathbb{A}_{\mathbb{Q}})\Gamma_{2}(\mathbb{A}_{\mathbb{Z}}) \\ &= \Gamma_{1}(\mathbb{Q})\Gamma_{2}(\mathbb{Q})\Gamma_{1}(\mathbb{Q})\Gamma_{1}(\mathbb{A}_{\mathbb{Z}})\Gamma_{2}(\mathbb{A}_{\mathbb{Z}}) = \Gamma(\mathbb{Q})\Gamma(\mathbb{A}_{\mathbb{Z}}). \end{split}$$

By Lemma 17, it now suffices to check that the criterion  $H_1(\mathbb{A}_{\mathbb{Z}})H_2(\mathbb{Q}) \subset H_2(\mathbb{Q})H_1(\mathbb{A}_{\mathbb{Q}})$  holds. This is an immediate consequence of the following matrix identity:

$$\left[\begin{array}{c|c} \operatorname{id} & 0 \\ \hline g''' & \operatorname{id} \end{array}\right] \times \left[\begin{array}{c|c} g' & 0 \\ \hline 0 & g''' \end{array}\right] = \left[\begin{array}{c|c} g' & 0 \\ \hline 0 & g''' \end{array}\right] \times \left[\begin{array}{c|c} \operatorname{id} & 0 \\ \hline g''^{-1}g'''g' & \operatorname{id} \end{array}\right].$$

2.2.5. Action of  $G_N$  on  $W_N^0$ . Let  $W_N^0 \subset W_N$  be the hyperplane whose R-points are defined by

$$W_N^0(R) := \{(A, B) \in W_N(R) : A_{ij} = B_{ij} = 0 \text{ if } i, j \le n\}$$

for any  $\mathbb{Z}$ -algebra R, and notice that, when R is an integral domain, every pair  $(A, B) \in W_N^0(R)$  is reducible. The action of  $SL_N$  on  $W_N$  restricts to an action of  $G_N$  on  $W_N^0$ .

We extend the definition of the Q-invariant to any pair  $(A, B) \in W_N^0(R)$  by defining Q(A, B) to be the Q-invariant of the projection of (A, B) onto the hyperplane  $W_N^{\text{top}}(R)$ . The function Q is notably not  $G_N$ -invariant, despite being invariant under the action of the subgroup  $\text{SL}_n \times \text{SL}_{n+1} \subset G_N$ ; see the proof of Proposition 24 (to follow). We remark that the explicit section  $\sigma_0$  referenced in §2.1 has image contained in the locus of points in  $W_N^0$  with Q-invariant equal to 1.

We now describe the action of  $G_N$  on  $W_N^0$  over subfields  $K \subset \mathbb{C}$ :

**Proposition 18.** Let  $K \subset \mathbb{C}$  be a subfield, and let  $f \in U_N(K)$  with nonzero discriminant. Then the set  $\operatorname{inv}^{-1}(f) \cap W_N^0(K)$  is nonempty and consists of a single  $G_N(K)$ -orbit. Moreover, the stabilizer of any element of this orbit is trivial.

*Proof.* That inv<sup>-1</sup> $(f) \cap W_N^0(R)$  is nonempty holds over any  $\mathbb{Z}$ -algebra R and without the nonzero-discriminant hypothesis, because of the existence of the aforementioned explicit section  $\sigma_0$ .

For the remaining claims, we first consider the case  $K = \mathbb{C}$ . To see that  $\operatorname{inv}^{-1}(f) \cap W_N^0(\mathbb{C})$  consists of a single  $G_N(\mathbb{C})$ -orbit, take  $(A_0, B_0) \in \operatorname{inv}^{-1}(f) \cap W_N^0(\mathbb{C})$ , and set  $Q_0 = Q(A_0, B_0) \in \mathbb{C}^{\times}$ . We claim that there exists  $g_1 \in (\operatorname{SL}_n \times \operatorname{SL}_{n+1})(\mathbb{C}) \subset G_N(\mathbb{C})$  such that  $(A, B) := g_1 \cdot (A_0, B_0)$  has the following shape, where empty entries are used to denote zeros, and star entries are used to denote complex numbers whose particular values are irrelevant:

In the above, we have inserted horizontal and vertical lines immediately after row and column n+1 in the matrix A, and immediately after row and column n in the matrix B. The claim follows from Proposition 10 upon observing that by choosing the sign of the entry " $\pm Q_0$ " appropriately, we can arrange for  $Q(A, B) = Q_0$ . In fact, by postcomposing  $g_1$  with a suitable diagonal element in  $G_N(\mathbb{C})$ , we may take (A, B) to be of the form (5), with the entries " $\pm Q_0$ " replaced by "1."

Now, just as in the proof of Proposition 11, by hitting (A, B) with a suitable sequence of elementary unipotent elements of  $G_N(K)$ , we can successively clear out the values of the following matrix entries:

$$A_{(n+1)(n+2)}, \ldots, A_{(n+1)(2n+1)}, B_{(n+1)(n+2)}, \ldots, B_{(n+1)(2n+1)}, A_{(n+2)(n+3)}, \ldots, A_{(n+2)(2n+1)}, \ldots, B_{k(k+1)}, \ldots, B_{k(2n+1)}, A_{(k+1)(k+2)}, \ldots, A_{(k+1)(2n+1)}, \ldots, B_{(2n-1)2n}, B_{(2n-1)(2n+1)}, A_{(2n)(2n+1)}, B_{(2n)(2n+1)}.$$

We may thus assume that the matrix entries of (A, B) that are listed in (6) are all equal to 0. But then (A, B) lies in the image of the section  $\sigma_0$  (see [21, §4.1] for the definition of this section), which is sufficient.

As for the claim about stabilizers, let  $(A_0, B_0) \in \operatorname{inv}^{-1}(f) \cap W_N^0(\mathbb{C})$  be as above, and suppose  $g \in G_N(\mathbb{C})$  stabilizes  $(A_0, B_0)$ . By Lemma 16, we may write  $g = g_1g_2$ , where  $g_i \in H_i(\mathbb{C})$  for each  $i \in \{1, 2\}$ . Since the projection map  $W_N^0 \to W_N^{\text{top}}$  is invariant under the action of  $H_1(\mathbb{C})$ , it follows that  $g_2$  must stabilize the projection of  $(A_0, B_0)$  onto  $W_N^{\text{top}}(\mathbb{C})$ . Thus, by Proposition 10, we have  $g_2 = \operatorname{id}$ . Now, it is clear by inspection that the action of  $H_1(\mathbb{C})$  on  $W_N^0(\mathbb{C})$  has trivial stabilizers, so  $g_1 = \operatorname{id}$ , and thus  $g = g_1g_2 = \operatorname{id}$  too.

Lastly, just as in the proof of Proposition 10, the case where  $K \subset \mathbb{C}$  is any subfield follows from the case where  $K = \mathbb{C}$  by [25, Lemma 23].

We now work over  $\mathbb{Z}_p$  and describe the action of  $G_N(\mathbb{Z}_p)$  on the locus of pairs in  $W_N^0(\mathbb{Z}_p)$  with p-adic unit Q-invariant:

**Proposition 19.** Let  $f \in U_N(\mathbb{Z}_p)$  with nonzero discriminant. Then the set  $\{(A, B) \in \operatorname{inv}^{-1}(f) \cap W_N^0(\mathbb{Z}_p) : |Q(A, B)|_p = 1\}$  is nonempty and consists of a single  $G_N(\mathbb{Z}_p)$ -orbit. Moreover, the stabilizer of any element of this orbit is trivial.

*Proof.* The nonemptiness statement follows immediately from the existence of the explicit section  $\sigma_0$ , which has image contained in the locus of points in  $W_N^0$  with Q-invariant 1.

Next, let  $W_N^{00} \subset W_N^0$  be the hyperplane whose R-points are given by

$$W_N^{00}(R) := \{(A, B) \in W_N^0(R) : A_{ij} = 0 \text{ if } i + j \le 2n + 1 \text{ and } B_{ij} = 0 \text{ if } i + j \le 2n\}.$$

We remark that the explicit section  $\sigma_0$  referenced in §2.1 has image contained in  $W_N^{00}$ . For the transitivity statement, by Lemma 14 it suffices to show that  $\{(A,B) \in \operatorname{inv}^{-1}(f) \cap W_N^{00}(\mathbb{Z}_p) : |Q(A,B)|_p = 1\}$  is contained in a single  $G_N(\mathbb{Z}_p)$ -orbit, and the proof of Proposition 18 can be easily adapted to verify this.

Finally, the statement about stabilizers follows immediately from Proposition 18, which implies that the stabilizer in  $G_N(\mathbb{Q}_p)$ , and hence in  $G_N(\mathbb{Z}_p)$ , of any  $(A, B) \in \operatorname{inv}^{-1}(f) \cap W_N^0(\mathbb{Z}_p)$  is trivial. This concludes the proof of Proposition 19.

We conclude by giving a rather crude bound on the number of  $G_N(\mathbb{Z}_p)$ -orbits on  $W_N^0(\mathbb{Z}_p)$  of given Q-invariant lying above a given binary N-ic form; this bound will be crucial in the proof of Theorem 5 (see §3.1).

**Lemma 20.** Let  $f \in U_N(\mathbb{Z}_p)$  with nonzero discriminant, and let  $e_p \geq 0$  be an integer. Then there exist integers  $\gamma_1, \gamma_2 \geq 1$  such that the number of  $G_N(\mathbb{Z}_p)$ -orbits on  $\operatorname{inv}^{-1}(f) \cap W_N^0(\mathbb{Z}_p)$  with Q-invariant having p-adic valuation  $e_p$  is at most  $(p^{e_p})^{\gamma_1 n^{\gamma_2}}$ .

*Proof.* Let  $\pi \colon W_N^0 \to W_N^{\text{top}}$  denote the projection map. By Proposition 13, every  $G_N(\mathbb{Z}_p)$ -orbit on  $\text{inv}^{-1}(f) \cap W_N^0(\mathbb{Z}_p)$  meets the set

(7) 
$$\left\{ (A,B) \in \pi^{-1}(\mathcal{L}_{\vec{a},\vec{b}}(p)) \cap W_N^{00}(\mathbb{Z}_p) : \begin{array}{l} \nu_p(A_{ij}) \leq a_{N+1-j} \text{ for } (i,j) \text{ with } i+j > N+1, \ i \leq n, \\ \nu_p(B_{ij}) \leq b_i \text{ for } (i,j) \text{ with } i+j > N, \ i \leq n \end{array} \right\}$$

for some  $\vec{a}, \vec{b} \in \mathbb{N}^n$ . Now, take an element (A, B) of the set (7). Just as in the proof of Proposition 13, we can use elementary unipotent transformations in  $G_N(\mathbb{Z}_p)$  to reduce the matrix entries in (6) modulo  $p^{a_1}, \ldots, p^{a_n}, p^{b_1}, \ldots, p^{b_n}$  (where the modulus depends on the matrix entry in question). All that remains to determine are the diagonal matrix entries  $A_{ii}$  and  $B_{ii}$  for  $i \geq n+1$ . But since all the other matrix entries are determined, these diagonal entries are uniquely determined by the condition that  $\operatorname{inv}(A, B) = f$ .

It follows that for every  $G_N(\mathbb{Z}_p)$ -orbit on  $\operatorname{inv}^{-1}(f) \cap W_N^0(\mathbb{Z}_p)$ , there exists a pair  $\vec{a}, \vec{b} \in \mathbb{N}^n$  and a set  $\Sigma$  of order  $p^{\max\{a_1,\dots,a_n,b_1,\dots,b_n\}}$  such that the orbit admits a representative (A,B) all of

whose matrix entries lie in  $\Sigma$ . To prove the lemma, it suffices to get some control on the pairs  $(\vec{a}, \vec{b})$  such that inv<sup>-1</sup> $(f) \cap \pi^{-1}(\mathcal{L}_{\vec{a}, \vec{b}}(p)) \neq \emptyset$ . As noted in §2.2.3, for any  $(A, B) \in \mathcal{L}_{\vec{a}, \vec{b}}(p)$  we have  $|Q(A, B)|_p^{-1} = \prod_{i=1}^n p^{(n+1-i)a_i+ib_i}$ , so each  $a_i, b_i$  is at most  $e_p + 1 \leq p^{e_p+1}$ , and this implies the claimed bound.

- 2.3. **Proof of Theorem 3.** By [25, Theorem 22], the integral orbits of  $G_N$  on  $W_N^0$  satisfy the desired local-to-global principle as long as the following four properties hold:
  - (1) The algebraic group  $G_N$  has class number 1 over  $\mathbb{Q}$ ;
  - (2) For every  $f \in U_N(\mathbb{C})$  with nonzero discriminant, the set  $\operatorname{inv}^{-1}(f) \cap W_N^0(\mathbb{C})$  is nonempty and consists of a single  $G_N(\mathbb{C})$ -orbit; and
  - (3) For every  $f \in U_N(\mathbb{C})$  with nonzero discriminant, each element of  $\operatorname{inv}^{-1}(f) \cap W_N^0(\mathbb{C})$  has trivial stabilizer in  $G_N(\mathbb{C})$ .
  - (4) For every  $f \in U_N(\mathbb{Z})$  with nonzero discriminant, the set  $\operatorname{inv}^{-1}(f) \cap W_N^0(\mathbb{Z})$  is nonempty.

The first three properties have already been verified in Propositions 15 and 18, and the fourth property follows immediately from the existence of the explicit section  $\sigma_0$ . This completes the proof of Theorem 3.

As an immediate consequence of Theorem 3 along with Proposition 19, we have the following result concerning global integral orbits having unit Q-invariant:

**Corollary 21.** Let  $f \in U_N(\mathbb{Z})$  with nonzero discriminant. Then the set  $\{(A, B) \in \operatorname{inv}^{-1}(f) \cap W_N^0(\mathbb{Z}) : |Q(A, B)| = 1\}$  is nonempty and consists of a single  $G_N(\mathbb{Z})$ -orbit.

#### 3. Asymptotic formulae for the count of reducible orbits

In this section, we use the local-to-global principle in Theorem 3 to deduce Theorem 5, which gives an asymptotic formula for the count of reducible  $SL_N(\mathbb{Z})$ -orbits on  $W_N(\mathbb{Z})$  in terms of a product of local integrals. We then perform a change-of-variables argument to rewrite each of these integrals in a more convenient form, thus proving Theorem 6.

3.1. **Proof of Theorem 5.** Let  $\mathfrak{S}$  be a big family in  $W_N^0(\mathbb{Z})$ . We start by proving the following asymptotic formula for the count of  $G_N(\mathbb{Z})$ -orbits on  $\mathfrak{S}$  of bounded height.

**Theorem 22.** The number of  $G_N(\mathbb{Z})$ -orbits on  $\mathfrak{S}$  of height up to X is given by

(8) 
$$N_N^{(r)}(X) \times \prod_p \int_{f \in U_N(\mathbb{Z}_p)} \#\left(\frac{\operatorname{inv}^{-1}(f) \cap \mathfrak{S}_p}{G_N(\mathbb{Z}_p)}\right) df + o(X^{N+1}).$$

Moreover, when N=3, the number of  $(GL_2 \times G_3)(\mathbb{Z})$ -orbits on  $\mathfrak{S}$  with discriminant having absolute value less than X is given by

(9) 
$$N_{\Delta}^{(r)}(X) \times \prod_{p} \int_{f \in U_3(\mathbb{Z}_p)} \#\left(\frac{\operatorname{inv}^{-1}(f) \cap \mathfrak{S}_p}{G_3(\mathbb{Z}_p)}\right) df + o(X),$$

where  $N_{\Delta}^{(r)}(X)$  is the number of  $GL_2(\mathbb{Z})$ -orbits of irreducible binary cubic forms in  $U_3(\mathbb{Z})^{(r)}$  of discriminant up to X in absolute value.

*Proof.* We first prove (8), and then we explain how the proof of (9) differs. The proof is analogous to that of [25, Theorem 24], but we include the details for the sake of completeness. Fix an integer  $\mathfrak{b} \geq 1$ , and factorize it into primes as  $\mathfrak{b} = \prod_p p^{e_p}$ . We first prove an analogue of Theorem 22 with  $\mathfrak{S}$  replaced by the subfamily  $\mathfrak{S}(\mathfrak{b}) := \{w \in \mathfrak{S} : |Q(w)| = \mathfrak{b}\};$  note that  $\mathfrak{S}(\mathfrak{b})$  is itself a big family in  $W_N^0(\mathbb{Z})$ , where  $\mathfrak{S}(\mathfrak{b})_p = \{w \in \mathfrak{S}_p : |Q(w)|_p = |\mathfrak{b}|_p\}.$ 

If  $\mathfrak{S}(\mathfrak{b}) = \emptyset$ , then there is nothing to prove, so assume that  $\mathfrak{S}(\mathfrak{b}) \neq \emptyset$ . For each prime  $p \mid \mathfrak{b}$ , we partition  $U_N(\mathbb{Z}_p)$  as  $U_N(\mathbb{Z}_p) = \bigsqcup_{j=1}^{m_p} U_{p,j}$ , where each  $U_{p,j}$  is a level set for the function that sends  $f \in U_N(\mathbb{Z}_p)$  to  $\#(G_N(\mathbb{Z}_p) \setminus (\operatorname{inv}^{-1}(f) \cap \mathfrak{S}(\mathfrak{b})_p))$ . The quantity  $\#(G_N(\mathbb{Z}_p) \setminus (\operatorname{inv}^{-1}(f) \cap \mathfrak{S}(\mathfrak{b})_p))$  is

independent of the choice of  $f \in U_{p,j}$  (by the definition of a level set) and it follows from Lemma 20 that there exist integers  $\gamma_1, \gamma_2 \geq 1$  such that this quantity is bounded by  $(p^{e_p})^{\gamma_1 n^{\gamma_2}}$ , independent of both f and  $\mathfrak{S}_p$ .

Now for each prime  $p \nmid \mathfrak{b}$ , Proposition 19 tells us that  $\#(G_N(\mathbb{Z}_p) \setminus (\operatorname{inv}^{-1}(f) \cap \mathfrak{S}(\mathfrak{b})_p)) = 1$  for each  $f \in \operatorname{inv}(\mathfrak{S}(\mathfrak{b})_p)$ . It then follows from Theorem 3 that the quantity

$$(10) \qquad \#\left(\frac{\operatorname{inv}^{-1}(f)\cap\mathfrak{S}(\mathfrak{b})}{G_N(\mathbb{Z})}\right) = \prod_p \#\left(\frac{\operatorname{inv}^{-1}(f)\cap\mathfrak{S}(\mathfrak{b})_p}{G_N(\mathbb{Z}_p)}\right) = \prod_{p\mid\mathfrak{b}} \#\left(\frac{\operatorname{inv}^{-1}(f)\cap\mathfrak{S}(\mathfrak{b})_p}{G_N(\mathbb{Z}_p)}\right)$$

is independent of the choice of  $f \in \operatorname{inv}(\mathfrak{S}(\mathfrak{b})) \cap \bigcap_p U_{p,j_p}$ . Therefore, for each tuple  $(j_p)_{p|\mathfrak{b}} \in \prod_{p|\mathfrak{b}} \{1,\ldots,m_p\}$ , we have

(11) 
$$\sum_{\substack{f \in U_N(\mathbb{Z}) \cap \bigcap_p U_{p,j_p} \\ \mathrm{H}(f) < X}} \# \left( \frac{\mathrm{inv}^{-1}(f) \cap \mathfrak{S}(\mathfrak{b})}{G_N(\mathbb{Z})} \right) = \# \left( \frac{\mathrm{inv}^{-1}(f^*) \cap \mathfrak{S}(\mathfrak{b})}{G_N(\mathbb{Z})} \right) \times \sum_{\substack{f \in \mathrm{inv}(\mathfrak{S}(\mathfrak{b})) \cap \bigcap_p U_{p,j_p} \\ \mathrm{H}(f) < X}} 1$$

where  $f^* \in \operatorname{inv}(\mathfrak{S}(\mathfrak{b})) \cap \bigcap_{p|\mathfrak{b}} U_{p,j_p}$  is any fixed element. Since  $\mathfrak{S}$  is a big family, and since the aforementioned explicit section  $\sigma_0$  has image contained in the locus of pairs (A,B) with Q(A,B) = 1, it follows that  $\operatorname{inv}(\mathfrak{S}(\mathfrak{b})_p) \supset U_N(\mathbb{Z}_p)$  for every  $p \gg 1$ . This allows us to apply the squarefree sieve on  $U_N(\mathbb{Z})$  given in [12, Theorem 5], which yields the following:

(12) 
$$\sum_{\substack{f \in \text{inv}(\mathfrak{S}(\mathfrak{b})) \cap \bigcap_{p} U_{p,j_{p}} \\ \text{H}(f) < X}} 1 = N_{N}^{(r)}(X) \times \prod_{p \mid \mathfrak{b}} \int_{f \in U_{p,j_{p}}} df \times \prod_{p \nmid \mathfrak{b}} \int_{f \in \text{inv}(\mathfrak{S}(\mathfrak{b})_{p})} df + O(X^{N+1-\delta})$$

for some  $\delta > 0$ . Substituting the asymptotic (12) into the right-hand side of (11), applying (10) to the resulting expression, and summing that over tuples  $(j_p)_{p|\mathfrak{b}} \in \prod_{p|\mathfrak{b}} \{1, \ldots, m_p\}$  gives the following:

(13) 
$$\sum_{\substack{f \in U_N(\mathbb{Z}) \\ \mathrm{H}(f) < X}} \# \left( \frac{\mathrm{inv}^{-1}(f) \cap \mathfrak{S}(\mathfrak{b})}{G_N(\mathbb{Z})} \right) = \mathrm{N}_N^{(r)}(X) \times \prod_p \int_{f \in U_N(\mathbb{Z}_p)} \# \left( \frac{\mathrm{inv}^{-1}(f) \cap \mathfrak{S}(\mathfrak{b})_p}{G_N(\mathbb{Z}_p)} \right) df + O(\mathfrak{b}^{\gamma_1 n^{\gamma_2}} X^{N+1-\delta}).$$

Next, we prove that the theorem holds with "=" replaced by " $\geq$ ." For any real number M > 1, let  $\mathfrak{S}[M] := \{ w \in \mathfrak{S} : |Q(w)| < M \}$ . Summing (13) over  $\mathfrak{b} < M$  gives:

(14) 
$$\sum_{\substack{f \in U(\mathbb{Z}) \\ \mathcal{H}(f) < X}} \# \left( \frac{\operatorname{inv}^{-1}(f) \cap \mathfrak{S}[M]}{G(\mathbb{Z})} \right) = \mathcal{N}_{N}^{(r)}(X) \times \sum_{\mathfrak{b} < M} \prod_{p} \int_{f \in U(\mathbb{Z}_{p})} \# \left( \frac{\operatorname{inv}^{-1}(f) \cap \mathfrak{S}(\mathfrak{b})_{p}}{G(\mathbb{Z}_{p})} \right) df + O(M^{2\gamma_{1}n^{\gamma_{2}}}X^{N+1-\delta}).$$

Dividing through by  $N_N^{(r)}(X)$ , letting  $X \to \infty$ , and replacing  $\mathfrak{S}$  with  $\mathfrak{S}[M]$ , it follows from (14) that

(15) 
$$\lim_{X \to \infty} \frac{\sum_{f \in U_N(\mathbb{Z})} \# \left( \frac{\operatorname{inv}^{-1}(f) \cap \mathfrak{S}}{G_N(\mathbb{Z})} \right)}{\operatorname{N}_N^{(r)}(X)} \ge \sum_{\mathfrak{b} < M} \prod_p \int_{f \in U_N(\mathbb{Z}_p)} \# \left( \frac{\operatorname{inv}^{-1}(f) \cap \mathfrak{S}(\mathfrak{b})_p}{G_N(\mathbb{Z}_p)} \right) df.$$

Now, letting  $M \to \infty$  on the right-hand side of (15) and factoring the sum into an Euler product, we obtain the following:

$$\sum_{\mathfrak{h}=1}^{\infty} \prod_{p} \int_{f \in U_{N}(\mathbb{Z}_{p})} \# \left( \frac{\mathrm{inv}^{-1}(f) \cap \mathfrak{S}(\mathfrak{b})_{p}}{G_{N}(\mathbb{Z}_{p})} \right) df = \prod_{p} \sum_{e=0}^{\infty} \int_{f \in U_{N}(\mathbb{Z}_{p})} \# \left( \frac{\mathrm{inv}^{-1}(f) \cap \mathfrak{S}(p^{e})_{p}}{G_{N}(\mathbb{Z}_{p})} \right) df$$

(16) 
$$= \prod_{p} \int_{f \in U_{N}(\mathbb{Z}_{p})} \# \left( \frac{\operatorname{inv}^{-1}(f) \cap \mathfrak{S}_{p}}{G_{N}(\mathbb{Z}_{p})} \right) df.$$

Combining (15) with (16), we find that Theorem 22 holds with "=" replaced by "≥."

It thus remains to prove the theorem with "=" replaced by " $\leq$ ." Let  $\mathfrak{S}[M]' := \mathfrak{S} \setminus \mathfrak{S}[M]$ . Then for each  $w \in \mathfrak{S}[M]'$ , we have that  $|Q(w)| \geq M$ . In [12], Bhargava, Shankar, and Wang determined bounds for the number of reducible  $\mathrm{SL}_N(\mathbb{Z})$ -orbits on  $W_N(\mathbb{Z})$  having large Q-invariant (for those orbits on which the notion of Q-invariant can be extended naturally). In particular, by [12, Theorem 5] along with Proposition 23 (to follow), we have that

(17) 
$$\sum_{\substack{f \in U_N(\mathbb{Z}) \\ \text{H}(f) < X}} \# \left( \frac{\text{inv}^{-1}(f) \cap \mathfrak{S}[M]'}{G_N(\mathbb{Z})} \right) = O\left(X^{N+1}/M\right) + O\left(X^{N+1-\delta}\right).$$

On the other hand, it follows from (14) that

(18) 
$$\sum_{\substack{f \in U_N(\mathbb{Z}) \\ \mathcal{H}(f) < X}} \# \left( \frac{\operatorname{inv}^{-1}(f) \cap \mathfrak{S}[M]}{G_N(\mathbb{Z})} \right) \le \mathcal{N}_N^{(r)}(X) \times \prod_p \int_{f \in U_N(\mathbb{Z}_p)} \# \left( \frac{\operatorname{inv}^{-1}(f) \cap \mathfrak{S}_p}{G_N(\mathbb{Z}_p)} \right) df + O(M^{2\gamma_1 n^{\gamma_2}} X^{N+1-\delta}).$$

Taking  $M = X^{\delta/(4\gamma_1 n^{\gamma_2})}$  and combining (17) with (18) yields (8).

As for the proof of (9), the only steps that differ are the deductions of (12) and (17). For (12), one simply applies the asymptotics for counting  $GL_2(\mathbb{Z})$ -orbits of binary cubic forms satisfying local conditions obtained by Bhargava, Shankar, and Tsimerman in [11, Theorem 26]. For (17), one simply applies the estimate proven by Bhargava in [4, Proposition 23].

To deduce Theorem 5 from Theorem 22, we require the following result relating the asymptotics for  $G_N(\mathbb{Z})$ -orbits on  $W_N^0(\mathbb{Z})$  with the asymptotics for reducible  $\mathrm{SL}_N(\mathbb{Z})$ -orbits on  $W_N(\mathbb{Z})$ :

**Proposition 23.** The number of  $G_N(\mathbb{Z})$ -orbits on  $W_N^0(\mathbb{Z})$  of height less than X differs from the number of reducible  $\mathrm{SL}_n(\mathbb{Z})$ -orbits on  $W_N(\mathbb{Z})$  of height less than X by an amount that is  $o(X^{N+1})$ .

Analgously, the number of  $(GL_2 \times G_3)(\mathbb{Z})$ -orbits on  $W_3^0(\mathbb{Z})$  with discriminant having absolute value less than X differs from the number of reducible  $(GL_2 \times SL_3)(\mathbb{Z})$ -orbits on  $W_3(\mathbb{Z})$  with discriminant having absolute value less than X by an amount that is o(X).

Proof. Let  $f \in U_N(\mathbb{Z})$  be irreducible with nonzero discriminant. We first claim that if we have  $(A_1, B_1), (A_2, B_2) \in \operatorname{inv}^{-1}(f) \cap W_N^0(\mathbb{Z})$  and  $g_1 \in \operatorname{SL}_N(\mathbb{Z})$  such that  $g_1 \cdot (A_1, B_1) = (A_2, B_2)$ , then  $g_1 \in G_N(\mathbb{Z})$ . Indeed, by Proposition 18, there exists  $g_2 \in G_N(\mathbb{Q})$  such that  $g_2 \cdot (A_1, B_1) = (A_2, B_2)$ , so  $g_1^{-1}g_2 \in \operatorname{SL}_N(\mathbb{Q})$  stabilizes  $(A_1, B_1)$ . But by [28, Proposition 14], the stabilizer in  $\operatorname{SL}_N(\mathbb{Z})$  of any element of  $\operatorname{inv}^{-1}(f)$  is trivial, so  $g_1 = g_2 \in \operatorname{SL}_N(\mathbb{Z}) \cap G_N(\mathbb{Q}) = G_N(\mathbb{Z})$ , as claimed.

It follows from the above claim that, up to an error of  $o(X^{n+1})$ , there are at least as many reducible  $\mathrm{SL}_N(\mathbb{Z})$ -orbits on  $W_N(\mathbb{Z})$  of height up to X as there are  $G_N(\mathbb{Z})$ -orbits  $W_N^0(\mathbb{Z})$  of height up to X. It therefore suffices to show that the number of reducible  $\mathrm{SL}_N(\mathbb{Z})$ -orbits on  $W_N(\mathbb{Z})$  that fail to meet the hyperplane  $W_N^0(\mathbb{Z})$  is  $o(X^{N+1})$ , but this follows immediately from [21, proof of Proposition 4.6].

The proof of the analogous claim about  $(GL_2 \times G_3)(\mathbb{Z})$ -orbits on  $W_3^0(\mathbb{Z})$  is essentially identical, with [28, Proposition 14] replaced by [4, §2.1, p. 1039] and [21, proof of Proposition 4.6] replaced by [4, Lemma 12].

Theorem 5 now follows from Theorem 22 and Proposition 23 by taking  $\mathfrak{S} = S \cap W_N^0(\mathbb{Z})$ , where S is a big family in  $W_N(\mathbb{Z})$ .

3.2. Change-of-variables formulas, and proof of Theorem 6. The purpose of this section is to deduce Theorem 6 from Theorem 5. The objective is, for each prime p, to reexpress the integral over  $U_N(\mathbb{Z}_p)$  that occurs in asymptotic given by Theorem 5 in terms of an integral over  $W_N^0(\mathbb{Z}_p)$ . To do this, we first write the integral over  $U_N(\mathbb{Z}_p)$  as an integral over  $G_N(\mathbb{Z}_p) \times U_N(\mathbb{Z}_p)$ ; we then perform a change-of-variables to relate the natural measure on  $G_N \times U_N$  with the natural measure on  $W_N^0$  (note that  $\dim G_N \times U_N = \dim G_N + \dim U_N = \dim W_N^0$ ).

We also prove a second change-of-variables formula relating the natural measure on  $W_N^{\text{top}}$  with the natural measure on  $(\operatorname{SL}_n \times \operatorname{SL}_{n+1}) \times \mathbb{A}^1$ , where  $\mathbb{A}^1$  parametrizes the Q-invariant (note that  $\dim ((\operatorname{SL}_n \times \operatorname{SL}_{n+1}) \times \mathbb{A}^1) = \dim \operatorname{SL}_n + \dim \operatorname{SL}_{n+1} + \dim \mathbb{A}^1 = \dim W_N^{\text{top}})$ . This second change-of-variables plays a crucial role in the proof of Theorem 7 (see §4.1, to follow).

3.2.1. Explicit choices of volume forms. We start by making explicit choices of volume forms on  $U_N$ ,  $W_N^0$ , and  $G_N$ . Take df, dw,  $dh_1$ , and  $dh_2$  to be generators of the 1-dimensional  $\mathbb{Z}$ -modules of top-degree differentials on  $U_N$ ,  $W_N^0$ ,  $H_1$ , and  $H_2$ , respectively. On  $G_N$  we take the measure  $dg = dh_1 dh_2$  (recall that  $G_N$  is the Frobenius product of  $H_1$  and  $H_2$  by Lemma 16).

We now give an explicit formula for dg on an open subscheme of  $G_N$ . For a square matrix M, we denote the minor obtained by deleting the first row and column by  $M_{(1,1)}$ ; if M is 1-dimensional, then we set  $M_{(1,1)} := 1$ . Let  $G_N^{\circ}$  be the open subscheme of  $G_N$  whose R-points are given by matrices

$$g = \left[ \begin{array}{c|c} g' & 0 \\ \hline g''' & g''' \end{array} \right] \in G_N(R)$$

such that  $g'_{(1,1)} \in R^{\times}$  for any  $\mathbb{Z}$ -algebra R. Then we may realize  $G_N^{\circ}$  as an open subscheme of the affine space

$$\mathcal{M} := \operatorname{Spec} \mathbb{Z} \left[ \left\{ M_{ij} : \begin{array}{c} 1 \leq i, j \leq n, \text{ where} \\ (i, j) \neq (1, 1) \text{ and } j \leq n \text{ if } i \leq n \end{array} \right\} \right]$$

via the map that sends a matrix  $g \in G_N(R)$  to its list of matrix entries  $g_{ij}$ , excluding the entries  $g_{ij}$  for any (i,j) such that  $i \leq n$  and  $j \geq n+1$  or (i,j)=(1,1).

Let  $\prod dM_{ij}$  be the Euclidean measure on  $\mathcal{M}$ , normalized so that  $\mathcal{M}(\mathbb{Z})$  has covolume 1 in  $\mathcal{M}(\mathbb{R})$ . We denote by  $\prod dg_{ij}$  the restriction of this measure to  $G_N^{\circ}$  via the embedding  $G_N^{\circ} \subset \mathcal{M}$  defined above, and by abuse of notation, we also denote by dg the restriction to  $G_N^{\circ}$  of the volume form on  $G_N$ . Then a calculation shows that the measure  $(g'_{(1,1)})^{-1}(\det g'')^{n-1} \times \prod dg_{ij}$  is invariant under the right-action of  $G_N$ , and so on  $G_N^{\circ}$  we have  $dg = (g'_{(1,1)})^{-1}(\det g'')^{n-1} \times \prod dg_{ij}$ , up to sign. As for the left-action of  $G_N$ , if we take

(19) 
$$h = \begin{bmatrix} h' & 0 \\ \overline{h'''} & \overline{h''} \end{bmatrix} \in G_N(R),$$

then one readily verifies that

(20) 
$$d(hg) = \rho(h)dg, \text{ where } \rho(h) := |\det h''|^{N}.$$

Note that the formula (20) holds on all of  $G_N$ , not just on the open subscheme  $G_N^{\circ}$ .

We finish by making explicit choices of the volume forms on  $\mathbb{A}^1$  and  $\mathrm{SL}_n \times \mathrm{SL}_{n+1}$ . Take dq to be a top form on  $\mathbb{A}^1$  defined over  $\mathbb{Z}$ , and as for the group  $\mathrm{SL}_n \times \mathrm{SL}_{n+1}$ , we take any generator of the 1-dimensional  $\mathbb{Z}$ -module of top-degree differentials defined over  $\mathbb{Z}$  (such a differential will necessarily be both left- and right-invariant).

3.2.2. Stating the change-of-variables formulas. We are now in position to state our change-of-variables formula relating the measure dw on  $W_N^0$  with the measure dgdf on  $G_N \times U_N$ .

**Proposition 24.** Let  $R = \mathbb{R}$  or  $\mathbb{Z}_p$  for a prime p. Let  $\phi \colon W_N^0(R) \to \mathbb{R}$  be a  $G_N(R)$ -invariant measurable function. Then there exists a nonzero rational number  $\mathcal{J} \in \mathbb{Q}^{\times}$ , possibly depending on N, such that

$$\frac{1}{|\mathcal{J}|} \times \frac{1}{\operatorname{Vol}(G_N(R))} \int_{w \in W_N^0(R)} \phi(w) |Q(w)| dw = \int_{\substack{f \in U_N(R) \\ \Delta(f) \neq 0}} \left( \sum_{w \in \frac{\operatorname{inv}^{-1}(f) \cap W_N^0(R)}{G_N(R)}} \phi(w) \right) df,$$

where |-| denotes the usual absolute value on R.

*Proof.* We follow the general strategy used in the proof of [25, Proposition 14]. Let  $\mathcal{U} \subset U_N(\mathbb{R})$  be an open set and let  $\sigma \colon \mathcal{U} \to W_N^0(\mathbb{R})$  be a continuous section of inv (note that such a section exists, namely the aforementioned explicit section  $\sigma_0$ ). We first claim that for some  $\mathcal{J} \in \mathbb{Q}^{\times}$  we have

(21) 
$$\int_{w \in G_N(\mathbb{R}) \cdot \sigma(\mathcal{U})} \phi(w) |Q(w)| dw = |\mathcal{J}| \int_{f \in \mathcal{U}} \int_{g \in G_N(\mathbb{R})} dg df.$$

By the Stone–Weierstrass theorem, it suffices to treat the case where  $\sigma$  is piecewise analytic. In that case, we have

$$\int_{w \in G_N(\mathbb{R}) \cdot \sigma(\mathcal{U})} \phi(w) |Q(w)| dw = \int_{f \in \mathcal{U}} \int_{g \in G_N(\mathbb{R})} |\mathcal{J}_{\sigma}(g, f)| \phi(g \cdot \sigma(f)) dg df,$$

where  $\mathcal{J}_{\sigma}(g, f)$  is the determinant of the Jacobian matrix coming from the change-of-variables taking the measure Q(w)dw on  $W_N^0$  to the product measure dgdf on  $G_N \times U_N$ .

We now show that  $\mathcal{J}_{\sigma}(g,f)$  is independent of g. Take  $h \in G_N(\mathbb{R})$ , and consider the transformation on  $W_N^0(\mathbb{R})$  that sends  $w \mapsto h \cdot w$ . Then there exists a function  $\chi_Q \colon G_N(\mathbb{R}) \to \mathbb{R}_{>0}$  such that  $Q(h \cdot w)d(h \cdot w) = \chi_Q(h)Q(w)dw$ ; indeed, one checks that if h is expressed as in (19), we have  $Q(h \cdot w) = |\det h''|^{-1}Q(w)$  and  $d(h \cdot w) = |\det h''|^{N+1}$ , so  $\chi_Q(h) = |\det h''|^N$ . On the other hand, the transformation  $w \mapsto h \cdot w$  acts on  $G_N(\mathbb{R}) \times \mathcal{U}$  by sending  $(g, f) \mapsto (hg, f)$ . Letting  $\rho \colon G_N(\mathbb{R}) \to \mathbb{R}_{>0}$  be as in (20), we have that

$$\mathcal{J}_{\sigma}(hg, f)d(hg)df = \rho(h)\mathcal{J}_{\sigma}(hg, f)dgdf.$$

But  $\rho(h) = |\det h''|^N = \chi_Q(h)$ , so we have that

$$\mathcal{J}_{\sigma}(hg, f)d(hg)df = Q(hw)dw = \chi_{Q}(h)Q(w)dw = \chi_{Q}(h)\mathcal{J}_{\sigma}(g, f)dgdf.$$

Upon comparing the above two displayed equations, we deduce that the function  $\mathcal{J}_{\sigma}(g, f) = \mathcal{J}(f)$  is independent of g.

That  $\mathcal{J}_{\sigma}(g, f)$  is independent of  $\sigma$  follows from an argument identical to Step 2 in the proof of [10, Proposition 3.10] (this step requires that the measure dg be right-invariant). Thus, we can take  $\sigma$  to be the polynomial section  $\sigma_0$  referred to in the proof of Proposition 18. With this choice of section, that  $\mathcal{J}_{\sigma_0}(g, f)$  is independent of f and equal to a nonzero rational constant follows from an argument identical to Steps 3 and 4 in the proof of [10, Proposition 3.10].

We have thus proven (21). Proposition 24 now follows from (21) and the principle of permanence of identities, just as [10, Proposition 3.7] is deduced from [10, Proposition 3.10].  $\Box$ 

We shall also require the following change-of-variables formula relating the pushforward of the measure dw from  $W_N^0$  to  $W_N^{\text{top}}$  with the product of the Haar measure on  $\text{SL}_n \times \text{SL}_{n+1}$  and the measure dq on  $\mathbb{A}^1$ . The proof is analogous to that of Proposition 24, so we omit it.

**Proposition 25.** Let  $R = \mathbb{R}$  or  $\mathbb{Z}_p$  for a prime p. Let  $\phi \colon W_N^{\mathrm{top}}(R) \to \mathbb{R}$  be an  $(\mathrm{SL}_n \times \mathrm{SL}_{n+1})(R)$ -invariant measurable function, and extend  $\phi$  to a function on  $W_N^0$  by precomposing with the natural

projection map  $W_N^0 \to W_N^{\text{top}}$ . Then there exists a nonzero rational number  $\mathcal{J}' \in \mathbb{Q}^{\times}$ , possibly depending on N, such that

$$\int_{w \in W_N^0(R)} \frac{\phi(w)}{|Q(w)|} dw = |\mathcal{J}'| \times \operatorname{Vol}\left(\left(\operatorname{SL}_n \times \operatorname{SL}_{n+1}\right)(R)\right) \int_{q \in R \setminus \{0\}} \left(\sum_{w \in \frac{Q^{-1}(q) \cap W_N^{\operatorname{top}}(R)}{\left(\operatorname{SL}_n \times \operatorname{SL}_{n+1}\right)(R)}} \phi(w)\right) dq,$$

where |-| denotes the usual absolute value on R.

We now turn our attention to the proof of Theorem 6. For this, let S be a big family in  $W_N(\mathbb{Z})$ . An application of Proposition 24 with  $R = \mathbb{Z}_p$  and  $\phi$  equal to the indicator function of  $S_p \cap W_N^0(\mathbb{Z}_p)$  yields that

(22) 
$$\int_{f \in U_N(\mathbb{Z}_p)} \# \left( \frac{\operatorname{inv}^{-1}(f) \cap S_p \cap W_N^0(\mathbb{Z}_p)}{G_N(\mathbb{Z}_p)} \right) = \frac{1}{\operatorname{Vol}(G_N(\mathbb{Z}_p))} \int_{w \in S_p \cap W_N^0(\mathbb{Z}_p)} |Q(w)|_p dw.$$

In the next lemma, we determine  $Vol(G_N(\mathbb{Z}_p))$ :

**Lemma 26.** We have that  $Vol(G_N(\mathbb{Z}_p)) = \xi_{p,n}^{-1} = (1-p^{-1})(1-p^{-n-1}) \times \prod_{i=2}^n (1-p^{-i})^2$ , and also that  $Vol\left((\operatorname{SL}_n \times \operatorname{SL}_{n+1})(\mathbb{Z}_p)\right) = (1-p^{-n-1}) \times \prod_{i=2}^n (1-p^{-i})^2$ .

*Proof.* We prove only the claimed formula for  $\operatorname{Vol}(G_N(\mathbb{Z}_p))$ , as the other formula can be proven similarly. Since  $G_N$  is smooth over  $\mathbb{Z}$ , we have that  $\operatorname{Vol}(G_N(\mathbb{Z}_p)) = \#G_N(\mathbb{F}_p)/p^{\dim G_N}$ . It is easy to see that  $\dim G_N = 3n^2 + 3n$ , and by Lemma 16, we have that

$$#G_N(\mathbb{F}_p) = #H_1(\mathbb{F}_p) \times #H_2(\mathbb{F}_p) = #\mathbb{A}^{n^2+n}(\mathbb{F}_p) \times (\#\operatorname{SL}_n(\mathbb{F}_p) \times \#\operatorname{GL}_{n+1}(\mathbb{F}_p))$$
$$= p^{n^2+n} \times (p-1)^{-1} \times \prod_{i=0}^{n-1} (p^n - p^i) \times \prod_{i=0}^{n} (p^{n+1} - p^i).$$

Finally, substituting the formula for  $Vol(G_N(\mathbb{Z}_p))$  given by Lemma 26 into (22) and combining the result with Theorem 5 yields Theorem 6.

#### 4. Evaluation of local volumes for applications

In this section, we evaluate the local integrals in Theorem 6 in two cases: (1) where  $S_p = W_N(\mathbb{Z}_p)$  for each prime p; and (2) where  $S_p$  is the set of projective elements in  $W_3(\mathbb{Z}_p)$  for each prime p. As a consequence of (1), we deduce Theorem 7, and as a consequence of (2), we deduce Theorems 8.A and 8.B.

4.1. **Proof of Theorem 7.** To deduce Theorem 7 from Theorem 6, we must take  $S = W_N(\mathbb{Z})$  and evaluate the integral over  $\mathbb{Z}_p$  for each prime p. We do this as follows:

**Proposition 27.** Fix a prime p. Then we have that

$$\frac{1}{\text{Vol}(G_N(\mathbb{Z}_p))} \int_{w \in W_N^0(\mathbb{Z}_p)} |Q(w)|_p dw = |\mathcal{J}|_p' \times \prod_{i=2}^N \frac{1}{1 - p^{-i}}.$$

Proof. Our strategy is to partition  $W_N^0(\mathbb{Z}_p)$  into the union over  $\vec{a}, \vec{b} \in \mathbb{N}^n$  of the preimage under projection  $\pi \colon W_N^0(\mathbb{Z}_p) \to W_N^{\text{top}}(\mathbb{Z}_p)$  of the set  $\mathcal{L}_{\vec{a},\vec{b}}(p)$  defined in §2.2.2. Taking  $\phi$  to be the indicator function of  $\pi^{-1}(\mathcal{L}_{\vec{a},\vec{b}}(p))$  and applying Proposition 25 yields that

$$(23) \int_{w \in \pi^{-1}(\mathcal{L}_{\vec{a},\vec{b}}(p))} |Q(w)|_{p} dw =$$

$$|\mathcal{J}'|_{p} \times \frac{\operatorname{Vol}\left((\operatorname{SL}_{n} \times \operatorname{SL}_{n+1})(\mathbb{Z}_{p})\right)}{\left(\prod_{i=1}^{n} p^{(n+1-i)a_{i}+ib_{i}}\right)^{2}} \int_{\nu_{p}(q) = \sum_{i=1}^{n} (n+1-i)a_{i}+ib_{i}} \#\left(\frac{Q^{-1}(q) \cap \mathcal{L}_{\vec{a},\vec{b}}(p)}{(\operatorname{SL}_{n} \times \operatorname{SL}_{n+1})(\mathbb{Z}_{p})}\right) dq.$$

Now, let  $\nu_p$  denote the usual *p*-adic valuation. Proposition 13 implies that we have for all *q* with  $\nu_p(q) = \sum_{i=1}^n (n+1-i)a_i + ib_i$  that

(24) 
$$\#\left(\frac{Q^{-1}(q)\cap\mathcal{L}_{\vec{a},\vec{b}}(p)}{(\mathrm{SL}_n\times\mathrm{SL}_{n+1})(\mathbb{Z}_p)}\right) = \prod_{i=1}^n p^{(n-i)a_i+ib_i},$$

so substituting (24) along with the calculation of Vol  $((SL_n \times SL_{n+1}(\mathbb{Z}_p)))$  from Lemma 26 into the right-hand side of (23) yields

(25) 
$$\int_{w \in \mathcal{L}_{\vec{a}, \vec{b}}(p)} |Q(w)|_p dw = |\mathcal{J}'|_p \times \frac{(1 - p^{-1})(1 - p^{-n-1}) \times \prod_{i=2}^n (1 - p^{-i})^2 \times \prod_{i=1}^n p^{(n-i)a_i + ib_i}}{\left(\prod_{i=1}^n p^{(n+1-i)a_i + ib_i}\right)^3}.$$

Summing (25) over all  $\vec{a}, \vec{b} \in \mathbb{N}^n$  and using the calculation of  $Vol(G_N(\mathbb{Z}_p))$  from Lemma 26, we have

$$\frac{1}{\text{Vol}(G_N(\mathbb{Z}_p))} \int_{w \in W_N^0(\mathbb{Z}_p)} |Q(w)|_p dw = \frac{1}{\text{Vol}(G_N(\mathbb{Z}_p))} \sum_{\vec{a}, \vec{b} \in \mathbb{N}^n} \int_{w \in \pi^{-1}(\mathcal{L}_{\vec{a}, \vec{b}}(p))} |Q(w)|_p dw$$

$$= |\mathcal{J}'|_p \times \prod_{i=1}^n \sum_{\vec{a} \in \mathbb{N}^n} \frac{1}{p^{(2n+3-2i)a_i}} \times \sum_{\vec{b} \in \mathbb{N}^n} \frac{1}{p^{2ib_i}}$$

$$= |\mathcal{J}'|_p \times \prod_{i=1}^n \frac{1}{1 - p^{-(N-(2i-2))}} \times \frac{1}{1 - p^{-2i}} = |\mathcal{J}'|_p \times \prod_{i=2}^N \frac{1}{1 - p^{-i}},$$

which is the desired result.

Up to a factor of  $|\mathcal{J}/\mathcal{J}'|$ , Theorem 7 now follows by combining Proposition 27 with Theorem 6 and evaluating the resulting Euler product. We now finish the proof of Theorem 7 by showing that the factor of  $|\mathcal{J}/\mathcal{J}'|$  does not in fact contribute to the asymptotic:

**Lemma 28.** We have that  $|\mathcal{J}/\mathcal{J}'| = 1$ .

Proof. By Corollary 21, there is exactly one  $G_N(\mathbb{Z})$ -orbit on  $W_N^0(\mathbb{Z})$  with unit Q-invariant lying above a binary form  $f \in U_N(\mathbb{Z})$  having nonzero discriminant. In particular, the average number of  $G_N(\mathbb{Z})$ -orbits on  $W_N^0(\mathbb{Z})$  with unit Q-invariant is equal to 1. On the other hand, combining Theorem 22 with Propositions 24 and 25 and Lemma 26 and using the formula (25), we find that the average number of  $G_N(\mathbb{Z})$ -orbits on  $W_N^0(\mathbb{Z})$  with unit Q-invariant is equal to

$$|\mathcal{J}| \times \prod_{p} \frac{1}{\operatorname{Vol}(G_N(\mathbb{Z}_p))} \int_{w \in \mathcal{L}_{\vec{0},\vec{0}}(p)} dw = \left| \frac{\mathcal{J}}{\mathcal{J}'} \right|. \qquad \Box$$

# 4.2. Proofs of Theorems 8.A and 8.B. Let R be a principal ideal domain.

**Definition 29.** We say that (the  $(GL_2 \times SL_3)(R)$ -orbit of) a pair  $(A, B) \in W_3(R)$  is projective if the following property is satisfied. For each  $k \in \{0, 1, 2\}$ , let  $C^{(k)}$  be the  $3 \times 3$  matrix over R defined as follows:

(26) 
$$C^{(0)} = f_0 B A^{-1} B + f_1 B + f_2 A, \quad C^{(1)} = B, \quad C^{(2)} = A.$$

Let  $M \in \operatorname{Mat}_{3 \times 6}(R)$  be the matrix whose  $k^{\operatorname{th}}$  row consists of the entries of  $C^{(k)}$  lying on or above the diagonal, written as a list in some order that is uniform over k. Then the pair (A, B) is projective if and only if the greatest common divisor of the  $3 \times 3$  minors of M is equal to 1.

One can check from the above definition that: (1) projectivity is a  $(GL_2 \times SL_3)(R)$ -invariant condition; (2) projectivity over  $\mathbb{Z}$  is equivalent to projectivity over  $\mathbb{Z}_p$  for every prime p; (3) projectivity over  $\mathbb{Z}_p$  is a mod-p condition (i.e., whether or not a pair (A, B) is projective is determined by the residue class of (A, B) modulo p); and (4) any pair  $(A, B) \in W_3^0(\mathbb{F}_p)$  with  $Q(A, B) \neq 0$  is projective. In particular, the set of projective elements of  $W_3(\mathbb{Z})$  is a big family in  $W_3(\mathbb{Z})$ .

The motivation to introduce the notion of projectivity is the following parametrization result, which relates 2-torsion ideals of rings defined by binary cubic forms to projective reducible  $SL_3(\mathbb{Z})$ -orbits on  $W_3(\mathbb{Z})$ :

**Theorem 30** ([13, Theorem 17] and [29, Proposition 204]). The elements of the group  $\mathcal{I}(R_f)[2]$  are in natural bijection with the projective reducible  $\mathrm{SL}_3(\mathbb{Z})$ -orbits of pairs  $(A, B) \in W_3(\mathbb{Z})$  with  $-\det(xA - yB) = f(x, y)$ .

By Theorem 30, proving Theorems 8.A and 8.B amounts to determining asymptotics for the number of projective reducible  $SL_3(\mathbb{Z})$ -orbits on  $W_3(\mathbb{Z})$ . An argument entirely analogous to the proof of Proposition 23 implies that these asymptotics are the same as the asymptotics for the number of projective  $G_3(\mathbb{Z})$ -orbits on  $W_3^0(\mathbb{Z})$ . By Theorem 22, Proposition 24, and Lemma 28, this amounts to evaluating a certain p-adic integral for each prime p, which we do as follows:

**Proposition 31.** Fix a prime p. Then we have that

$$\frac{1}{\operatorname{Vol}(G_3(\mathbb{Z}_p))} \int_{\substack{w \in W_3^0(\mathbb{Z}_p) \\ w \text{ is proj.}}} |Q(w)|_p dw = 1 + p^{-2}.$$

Proof. Our strategy is to slice up the set of projective elements of  $W_3^0(\mathbb{Z}_p)$  into level sets for the function  $w \mapsto \nu_p(Q(w))$ , to evaluate the integral on each level set, and to sum up the results. To this end, given an integer  $k \geq 1$ , let  $L_k := \{(A,B) \in W_3^0(\mathbb{Z}_p) : \nu_p(Q(A,B)) = k-1\}$ , given  $a,b,c,d \in \mathbb{F}_p$ , let

 $S_{a,b,c,d} := \{(A,B) \in W_3^0(\mathbb{F}_p) : A_{12} = a, A_{13} = b, B_{12} = c, B_{13} = d, \text{ and } (A,B) \text{ is proj.} \},$  and given  $m \in \mathbb{Z}/p^k\mathbb{Z}$ , denote by  $\overline{m}$  the mod-p reduction of m. Then we have that

(27) 
$$\int_{\substack{w \in L_k \\ w \text{ is proj.}}} |Q(w)|_p dw = p^{1-k} \times p^{-4k} \times \sum_{\substack{M \in \text{Mat}_{2 \times 2}(\mathbb{Z}/p^k\mathbb{Z}) \\ \nu_p(\det M) = k-1}} p^{-6} \times \#S_{\overline{M}_{11}, \overline{M}_{12}, \overline{M}_{21}, \overline{M}_{22}}.$$

We now determine the size of the set  $S_{a,b,c,d}$  for each choice of  $a,b,c,d \in \mathbb{F}_p$ :

- Let  $(A, B) \in W_3^0(\mathbb{F}_p)$ . As mentioned above, if  $Q(A, B) \neq 0$ , then (A, B) is projective. Thus, if  $ad bc \neq 0$ , then  $\#S_{a,b,c,d} = p^6$ .
- Now suppose that Q(A,B)=0. A calculation reveals that if p divides all of  $A_{12}$ ,  $A_{13}$ ,  $B_{12}$ , and  $B_{13}$ , then (A,B) is not projective. Thus, for (A,B) to be projective, at least one of these four matrix entries must be a unit. We now fiber over these four matrix entries and determine the number of possibilities for the pair (A,B) in each fiber. Fix four elements  $a,b,c,d \in \mathbb{F}_p$  with ad-bc=0 and  $\{0\} \neq \{a,b,c,d\}$ . We claim that  $\#S_{a,b,c,d}$  is independent of the choice of a,b,c,d. Indeed, if  $a',b',c',d' \in \mathbb{F}_p$  with a'd'-b'c'=0 and  $\{0\} \neq \{a',b',c',d'\}$ , then there exists  $\gamma \in H_2(\mathbb{F}_p)$  such that if we set  $(A',B') := \gamma \cdot (A,B)$ , then  $(A'_{12},A'_{13},B'_{12},B'_{13})=(a,b,c,d)$ . Thus,  $\gamma$  induces a bijection between  $S_{a,b,c,d}$  and  $S_{a',b',c',d'}$ , so it suffices to compute  $\#S_{0,1,0,0}$ . A calculation reveals that a pair (A,B) with  $(A_{12},A_{13},B_{12},B_{13})=(0,1,0,0)$  is projective if and only if  $B_{22}B_{33}-B_{23}^2\neq 0$ . Using this characterization, it is easy to check that  $\#S_{0,1,0,0}=p^5(p-1)$ .

Substituting the formulas for  $\#S_{a,b,c,d}$  obtained above into the right-hand side of (27), we find that

(28) 
$$\int_{\substack{w \in L_k \\ w \text{ is proj}}} |Q(w)|_p dw = p^{1-5k} \times \begin{cases} c(k), & \text{if } k = 1, \\ p^{-1}(p-1) \times c(k), & \text{if } k \ge 2. \end{cases}$$

where  $c(k) := \#\{M \in \operatorname{Mat}_{2\times 2}(\mathbb{Z}/p^k\mathbb{Z}) : \nu_p(\det M) = k-1 \text{ and } M \not\equiv 0 \pmod{p}\}$ . Let  $c'(k) := \#\{M \in \operatorname{Mat}_{2\times 2}(\mathbb{Z}/p^k\mathbb{Z}) : \nu_p(\det M) = k-1\}$ . Then we have

(29) 
$$c(k) = c'(k) - p^4 c'(k-2),$$

where for convenience we set c'(m) = 0 if  $m \le 0$ . The next lemma computes c'(k):

**Lemma 32.** Let  $k \ge 1$  be as above. Then  $c'(k) = p^{2k-1}(p^2 - 1)(p^k - 1)$ .

Proof of Lemma 32. For  $d \in \mathbb{Z}/p^k\mathbb{Z}$ , let  $M(d) := \{M \in \operatorname{Mat}_{2 \times 2}(\mathbb{Z}/p^k\mathbb{Z}) : M_{11}M_{22} = M_{12}M_{21} = d\}$ , and let  $N(d) := \{(a,b) \in (\mathbb{Z}/p^k\mathbb{Z})^2 : ab = d\}$ . Then  $M(d) = N(d)^2$ , and so if we set  $c''(k) := \#\{M \in \operatorname{Mat}_{2 \times 2}(\mathbb{Z}/p^k\mathbb{Z}) : \det M = 0\}$  of matrices over  $\mathbb{Z}/p^k\mathbb{Z}$  with determinant 0, then we have

(30) 
$$c''(k) = \sum_{d \in \mathbb{Z}/p^k \mathbb{Z}} M(d) = \sum_{d \in \mathbb{Z}/p^k \mathbb{Z}} N(d)^2.$$

Let  $\phi$  denote Euler's totient function. First suppose  $\nu_p(d) < k$ , and fix a number  $j \in \{0, \dots, \nu_p(d)\}$ . If d factors as d = ab where  $\nu_p(a) = j$ , then we have that there are  $\phi(p^{k-j})$  choices for a and  $\phi(p^{k-\nu_p(d)+j})\phi(p^{k-\nu_p(d)})^{-1}$  choices for b. Summing over all j, we find that

(31) 
$$N(d) = \sum_{j=0}^{\nu_p(d)} \phi(p^{k-\nu_p(d)})^{-1} \phi(p^{k-j}) \phi(p^{k-\nu_p(d)+j}) \quad \text{if } \nu_p(d) < k.$$

Now suppose  $d \equiv 0 \pmod{p^k}$ . Suppose d factors as d = ab, let  $i = \nu_p(a)$  if  $\nu_p(a) < k$  and i = k otherwise. Then there are  $\phi(p^{k-i})$  choices for a, and for each  $j \in \{k-i,\ldots,k\}$ , there are  $\phi(p^{k-j})$  choices for b. Summing over all i and j, we find that

(32) 
$$N(0) = \sum_{i=0}^{k} \sum_{j=k-i}^{k} \phi(p^{k-i})\phi(p^{k-j}).$$

Substituting the results of (31) and (32) into (30) and evaluating the sum, we deduce that

(33) 
$$c''(k) = p^{2k-1}(p^k(p+1) - 1).$$

Finally, note that we have  $c'(k) = p^4 c''(k-1) - c''(k)$ . Substituting in the formula (33) for c''(k) yields the lemma.

Substituting the formula for c'(k) given by Lemma 32 into (29) yields

$$c(k) = c'(k) - p^{4}c'(k-2) = \begin{cases} p(p-1)(p^{2}-1), & \text{if } k = 1, \\ p^{3k-3}(p^{2}-1)^{2}, & \text{if } k \ge 2. \end{cases}$$

Substituting this expression for c(k) into the right-hand side of (28), summing up over all positive integers k, and dividing by the volume  $Vol(G_3(\mathbb{Z}_p))$  as given in Lemma 26 yields the proposition.  $\square$ 

#### ACKNOWLEDGMENTS

It is a pleasure to thank Manjul Bhargava, Andrew Granville, Arul Shankar, Artane Siad, Ila Varma, and Melanie Matchett Wood for helpful discussions. This material is based in part upon work supported by the National Science Foundation, under the Graduate Research Fellowship as well as Award No. 2202839.

#### References

- [1] M. Bhargava. Higher composition laws. ProQuest LLC, Ann Arbor, MI, 2001. Thesis (Ph.D.)—Princeton University.
- [2] M. Bhargava. Higher composition laws. II. On cubic analogues of Gauss composition. Ann. of Math. (2), 159(2):865–886, 2004.
- [3] M. Bhargava. Higher composition laws. III. The parametrization of quartic rings. Ann. of Math. (2), 159(3):1329–1360, 2004.
- [4] M. Bhargava. The density of discriminants of quartic rings and fields. Ann. of Math. (2), 162(2):1031–1063, 2005.

- [5] M. Bhargava. Higher composition laws. IV. The parametrization of quintic rings. Ann. of Math. (2), 167(1):53–94, 2008
- [6] M. Bhargava. The density of discriminants of quintic rings and fields. Ann. of Math. (2), 172(3):1559–1591, 2010.
- [7] M. Bhargava. Most hyperelliptic curves over  $\mathbb Q$  have no rational points. arXiv preprint 1308.0395, 2013.
- [8] M. Bhargava, B. H. Gross, and X. Wang. A positive proportion of locally soluble hyperelliptic curves over  $\mathbb{Q}$  have no point over any odd degree extension. *J. Amer. Math. Soc.*, 30(2):451–493, 2017. With an appendix by Tim Dokchitser and Vladimir Dokchitser.
- [9] M. Bhargava, J. Hanke, and A. Shankar. The mean number of 2-torsion elements in class groups of n-monogenized cubic fields. arXiv preprint 2010.15744, 2020.
- [10] M. Bhargava and A. Shankar. Binary quartic forms having bounded invariants, and the boundedness of the average rank of elliptic curves. *Ann. of Math.* (2), 181(1):191–242, 2015.
- [11] M. Bhargava, A. Shankar, and J. Tsimerman. On the Davenport-Heilbronn theorems and second order terms. Invent. Math., 193(2):439–499, 2013.
- [12] M. Bhargava, A. Shankar, and X. Wang. Squarefree values of polynomial discriminants II. arXiv preprint 2207.05592, 2022.
- [13] M. Bhargava and I. Varma. On the mean number of 2-torsion elements in the class groups, narrow class groups, and ideal groups of cubic orders and fields. Duke Math. J., 164(10):1911–1933, 2015.
- [14] M. Bhargava and I. Varma. The mean number of 3-torsion elements in the class groups and ideal groups of quadratic orders. Proc. Lond. Math. Soc. (3), 112(2):235–266, 2016.
- [15] H. Cohen and H. W. Lenstra, Jr. Heuristics on class groups of number fields. In *Number theory, Noordwijkerhout* 1983 (Noordwijkerhout, 1983), volume 1068 of *Lecture Notes in Math.*, pages 33–62. Springer, Berlin, 1984.
- [16] H. Cohen and J. Martinet. Class groups of number fields: numerical heuristics. Math. Comp., 48(177):123–137, 1987.
- [17] H. Davenport. On the class-number of binary cubic forms. I. J. London Math. Soc., 26:183–192, 1951.
- [18] H. Davenport. Corrigendum: On the Class-Number of Binary Cubic Forms (I). J. London Math. Soc., 27(4):512, 1952.
- [19] H. Davenport and H. Heilbronn. On the density of discriminants of cubic fields. II. Proc. Roy. Soc. London Ser. A, 322(1551):405–420, 1971.
- [20] B. N. Delone and D. K. Faddeev. The theory of irrationalities of the third degree. Translations of Mathematical Monographs, Vol. 10. American Mathematical Society, Providence, R.I., 1964.
- [21] W. Ho, A. Shankar, and I. Varma. Odd degree number fields with odd class number. Duke Math. J., 167(5):995– 1047, 2018.
- [22] G. Malle. On the distribution of class groups of number fields. Experiment. Math., 19(4):465–474, 2010.
- [23] M. Sato and T. Kimura. A classification of irreducible prehomogeneous vector spaces and their relative invariants. Nagoya Math. J., 65:1–155, 1977.
- [24] G. W. Schwarz. Representations of simple Lie groups with a free module of covariants. *Invent. Math.*, 50(1):1–12, 1978/79.
- [25] A. Shankar, A. Siad, A. Swaminathan, and I. Varma. Geometry-of-numbers methods in the cusp. arXiv preprint arXiv:2110.09466, 2021.
- [26] A. Siad. Monogenic fields with odd class number part I: odd degree. arXiv preprint 2011.08834, 2020.
- [27] A. Siad. Monogenic fields with odd class number part II: even degree. arXiv preprint 2011.08842, 2020.
- [28] A. Swaminathan. Average 2-torsion in class groups of rings associated to binary n-ic forms. arXiv preprint 2011.13578, 2020.
- [29] A. Swaminathan. 2-Selmer Groups, 2-Class Groups, and the Arithmetic of Binary Forms. ProQuest LLC, Ann Arbor, MI, 2022. Thesis (Ph.D.)—Princeton University.
- [30] M. M. Wood. Parametrization of ideal classes in rings associated to binary forms. J. Reine Angew. Math., 689:169–199, 2014.

DEPARTMENT OF MATHEMATICS, HARVARD UNIVERSITY, CAMBRIDGE, MA 02138

E-mail address, Ashvin A. Swaminathan: ashvins@alumni.princeton.edu