# ALGEBRAIC RESULTS ON SP NUMBERS ALONG WITH A GENERALIZATION

RAGHAVENDRA N BHAT, SUNDARRAMAN MADHUSUDANAN

ABSTRACT. We defined numbers of the form  $p \cdot a^2$  as **SP numbers (Square-Prime numbers)**  $(a \neq 1, p \text{ prime})$  in [1] along with proofs on their distribution. Some examples of SP Numbers :  $75 = 3 \cdot 25$ ;  $108 = 3 \cdot 36$ ;  $45 = 5 \cdot 9$ . These numbers are listed in the OEIS as A228056 [2]. In this paper, we will prove a few algebraic theorems and generalize the definition of SP Numbers to allow factors of arbitrary natural number powers.

#### 1. INTRODUCTION AND SUMMARY OF PREVIOUS RESULTS

In our previous paper titled 'Distribution of square-prime numbers' [1], we introduced the notion of a product of a prime and a square, with the square not being 1. We call these SP numbers.

Here are the first 25 SP numbers: 8, 12, 18, 20, 27, 28, 32, 44, 45, 48, 50, 52, 63, 68, 72, 75, 76, 80, 92, 98, 99, 108, 112, 116, 117.

In the paper we mainly proved three results. Firstly, SP numbers have an asymptotic distribution similar to prime numbers, meaning, for a large natural number n, the number of SP numbers smaller than n is asymptotic to  $\frac{n}{\log n}$ , as with the case of primes. We denote the number of SP numbers smaller than n by SP(n). The result derived in the paper is  $SP(n) = (\zeta(2) - 1) \cdot \frac{n}{\log n} + O\left(\frac{n}{\log n^2}\right)$ . We see that although asymptotically similar to the distribution of primes, the actual value of SP(n) is lower than  $\pi(n)$  for large n, since  $\zeta(2)-1 < 1$ .

Secondly, there are infinitely many pairs of SP numbers that are consecutive natural numbers. We call these SP twins. An example is (27,28). More generally, we showed that if  $SP_1$  and  $SP_2$ are SP Numbers, the equation  $SP_1 - SP_2 = g$ , where  $g \in \mathbb{N}$  has infinite solutions if it has one solution. Here g is the gap between the two SP Numbers.

Thirdly, we analyzed techniques to give asymptotic estimates for number of SP numbers having a certain last digit. For example, the number of SP numbers ending in 1 approaches  $\frac{1}{400} \frac{n}{\log n} (\zeta(2, 1/10) + \zeta(2, 9/10) + \zeta(2, 3/10) + \zeta(2, 7/10) - 4)$ . In this paper, our goal is to study algebraic properties of SP numbers, their gaps and also their

In this paper, our goal is to study algebraic properties of SP numbers, their gaps and also their structure with regard to square-free numbers. Analogies to famous prime number conjectures

1

Date: September 2022.

like the  $x^2 + 1$  conjecture and primes between squares are studied. We then generalize the idea of a square multiplied by a prime to any power of a natural number multiplied by a prime.

#### 2. Algebraic SP Theorems

**Theorem 2.1:** For any positive integer x, there exist at least one pair of SP numbers (a, b) such that a - b = x.

*Proof.* We will divide all positive integers into five groups. We will then prove the existence of one pair of SP numbers with difference x in each group.

- x=1: 28 27 = 1. Thus, there exists an SP pair with difference 1.
- x is prime: Consider the following Pell's equation

$$m^2 - pxn^2 = 1 \tag{3}$$

where  $p \in \mathbf{P} \neq x$ . We know from Chakravala that (3) has at least 1 positive integer solution. Let that solution be (M, N). Thus,

$$M^2 - pxN^2 = 1 \tag{4}$$

$$\Rightarrow xM^2 - px^2N^2 = x \Rightarrow xM^2 - p(xN)^2 = x \tag{5}$$

Thus, we have generated two SP numbers,  $xM^2$  and  $p(xN)^2$  that differ by x. Since x was an arbitrary prime, this case holds.

• **x** is an odd composite square free number : Thus we have  $x = p_1 \cdot p_2 \cdot \ldots \cdot p_n$ , where  $p_1, p_2 \ldots, p_n$  are all odd prime numbers. Let  $p_1$  be the smallest prime factor of x. Thus  $p_1 \geq 3$  and  $p_2, p_3, \ldots, p_n > 3$ . Thus we can write

$$x = p_1 \cdot y$$

where  $y = p_2 \times p_3 \times ... \times p_n$ . And since  $p_2, p_3, ..., p_n > 3$ , y > 3. Furthermore, since y is a product of odd prime numbers, y has to be odd. Thus, we can also write the above equation as

$$x = p_1 \cdot (2k+1)$$

And again, since y > 3, we have k > 1.

$$x = p_1 \cdot [(k+1)^2 - k^2] \Rightarrow x = p_1 \cdot (k+1)^2 - p_1 \cdot k^2$$

And since k > 1 and  $p_1$  is prime, by definition  $p_1 \cdot (k+1)^2$  and  $p_1 \cdot k^2$  are both *SP* numbers with difference x. This case holds.

• **x** is an even composite square free number : This means  $x = p_1 \cdot p_2 \cdot ... \cdot p_n$ , where  $p_1 = 2$  and  $p_2, p_3...p_n$  are all odd prime numbers. Thus  $p_2, p_3, ...p_n \ge 3$ . Thus, as previously written,

$$x = 2 \times y$$

where  $y = p_2 \times p_3 \times ... \times p_n$ . And since  $p_2, p_3, ..., p_n \ge 3$ ,  $y \ge 3$ . Furthermore, since y is a product of odd prime numbers, y is odd. Thus, we can also write the above equation as

$$x = 2 \cdot (2k+1)$$

And again, since  $y \ge 3$ , we have  $k \ge 1$ . The above equation simplifies to

$$x = 2 \cdot [(k+1)^2 - k^2] \Rightarrow x = 2 \cdot (k+1)^2 - 2 \cdot k^2$$

Note that in the previous case we had k > 1 and here, we have  $k \ge 1$ . This is an issue because, when  $k = 1, 2 \cdot k^2$  will be 2 which is not an SP number. Note that

$$k = 1 \Rightarrow y = 3 \Rightarrow p_2 \cdot p_3 \cdot \ldots \cdot p_n = 3 \Rightarrow p_2 = 3$$

since 3 is prime and has only 1 factor. Thus, except in the case where  $p_1 = 2, p_2 = 3$ , this proof works exactly like the previous case.

Now, let us look at  $p_1 = 2, p_2 = 3$ , or in other words,  $x = p_1 \cdot p_2 = 6$ . 18 - 12 = 6 and  $18 = 2 \cdot 3^2, 12 = 3 \cdot 2^2$ .

Since 6 was the only integer in this group for which the general method did not work, we can combine the previous 2 paragraphs to jointly say that for every even composite square free integer x, there exists a pair of SP numbers (a, b) such that a - b = x.

• **x** is a non square free number : By the property of non square free numbers, we can write  $x = t^2 \cdot s$  where s is a square free number. Now, using the result from the four cases above, we can say that for all s, there exists a pair of SP numbers (j, k) such that

$$s = j - k \Rightarrow t^2 \cdot s = t^2 \cdot j - t^2 \cdot k \Rightarrow x = t^2 \cdot j - t^2 \cdot k$$

Since (j, k) are SP numbers,  $(t^2 \cdot j \text{ and } t^2 \cdot k)$  are both SP numbers having difference x. Thus, for every non square free integer x, there exists a pair of SP numbers (a, b) such that a - b = x.

Combining all the groups above, we can say that for any positive integer x, there exist at least one pair of SP numbers (a, b) such that a - b = x.

**Theorem 2.2:** There are infinitely many SP Numbers of the form  $x^2 + 1$ .

*Proof.* Our goal is to show that the equation  $x^2 + 1 = pa^2$  has infinitely many solutions where  $a, x \in \mathbb{N}$  and p is prime. We start the proof with the knowledge that at least one solution already exists (we have 50, 325 and 1025 as the first 3 SP Numbers of the form  $x^2 + 1$ ).

$$m^2 - 2n^2 = -1$$

has a solution, namely (7,5). Let (A,B) = (7,5). Now, consider the equation

$$m^2 - 2n^2 = 1$$

We know the above Pell's equation has infinitely many solutions. Let (X, Y) be an arbitrary solution to the above equation. Thus, we have the following 2 equations:

$$A^2 - 2B^2 = -1 \tag{1}$$

$$X^2 - 2Y^2 = 1 (2)$$

Multiplying (1) and (2), we get

$$(A^{2} - 2B^{2})(X^{2} - 2Y^{2}) = -1 \Rightarrow (AX + 2BY)^{2} - 2(AY + BX)^{2} = -1$$

Thus, for every one of the infinite solutions of the Pell's equation (X, Y), we have a new solution for the equation:

$$m^2 - 2n^2 = -1 \Rightarrow 2n^2 = m^2 + 1$$

Thus, there are infinitely many SP numbers of the form  $x^2 + 1$ .

**Theorem 2.3 :** For any positive integer x, there exists at least one SP number between  $x^2$  and  $(x+2)^2$ .

*Proof.* Consider the sub-sequence of SP numbers 8, 18, 32, 50, 72, 98, 128, 162, .... All of them are of the form  $2n^2$ . Notice that  $2n^2$  lies between  $x^2$  and  $(x+2)^2$  if and only if  $n\sqrt{2}$  lies between x and x+2.

Thus, we now need to prove that for any positive integer x, there exists at least one positive integer n such that  $n\sqrt{2}$  lies between x and (x + 2). We will prove this using contradiction.

Assume that there exists a positive integer x such that the smallest multiple of  $\sqrt{2}$  greater than x is greater than x + 2. Let this multiple be  $a\sqrt{2}$  This means there is no multiple of  $\sqrt{2}$  in [x, x + 2]. But then,

$$(a-1)\sqrt{2} < x < x+2 < a\sqrt{2} \Rightarrow a\sqrt{2} - (a-1)\sqrt{2} > x+2-x \Rightarrow \sqrt{2} > 2$$

which is a contradiction.

**Theorem 2.4 :** All SP numbers of the form  $pa^2$ , where p is a prime and a has at least one prime factor of the form 4k + 1 can be written as a sum of 2 SP numbers.

*Proof.* By the 2 squares theorem, if 4k + 1 is a prime,  $(4k + 1)^2$  can be written as  $x^2 + y^2$ , where x, y > 1. Now, since 4k + 1 is a factor of a, we can write

$$a = n \cdot (4k+1) \Rightarrow pa^2 = pn^2 \cdot (4k+1)^2$$
$$\Rightarrow pa^2 = pn^2 \cdot (x^2 + y^2) \Rightarrow pa^2 = p \cdot (nx)^2 + p \cdot (ny)^2$$

We now have our original SP number  $pa^2$  as a sum of two SP numbers  $p \cdot (nx)^2$  and  $p \cdot (ny)^2$ 

## 3. SP Numbers of the form $x^3 + 1$

**Conjecture 3.1**: There are infinitely many SP Numbers of the form  $x^3 + 1$ , if the Bunyakovsky [3] conjecture is true.

 $\square$ 

*Proof.* Let  $x = t^2 - 1$  for  $t \in \mathbb{N}$ . We have  $x^3 + 1 = (x + 1)(x^2 - x + 1) \Rightarrow x^3 + 1 = t^2 \cdot (t^4 - 3t^2 + 3)$ Consider  $f(t) = t^4 - 3t^2 + 1$ . It satisfies the following 3 properties of the Bunyakovsky conjecture: 1) The leading coefficient is positive.

2) The polynomial is irreducible over integers.

3) GCD(f(2), f(3)) is 1 and the coefficients of f(x) are relatively prime.

Using the above three conditions, it follows from the Bunyakovsky conjecture that f(t) is prime for infinitely many positive integers t. This implies  $t^2 \cdot (t^4 - 3t^2 + 3)$  is of the form  $prime \cdot t^2$  for infinitely many t which further implies there exists infinitely many  $x = t^2 - 1$  such that  $x^3 + 1$ is an SP number.

### 4. A GENERALIZATION OF SP NUMBERS

SP Numbers are those numbers that are of the form  $pa^2$ . We could extend the definition to numbers of the form  $pa^k$  and get asymptotic estimates on their distribution. Let us call these numbers KP<sub>k</sub> Numbers. SP Numbers are a special case of KP numbers and can be defined to be KP<sub>2</sub>. Trivially, the prime numbers and semi-prime numbers are KP<sub>0</sub> and KP<sub>1</sub> respectively. Let  $KP_k(n)$  denotes the number of KP<sub>k</sub> Numbers smaller than n. As proved in [1],

$$KP_2(n) = (\zeta(2) - 1)\frac{n}{\log n} + O\left(\frac{n}{\log^2 n}\right)$$

**Theorem 4.1 :** For any k greater than 1,

$$KP_k(n) = (\zeta(k) - 1)\frac{n}{\log n} + O\left(\frac{n}{\log^2 n}\right)$$

Proof.

$$KP_k(n) = \sum_{a=2}^{\sqrt{n/2}} \pi\left(\frac{n}{a^k}\right).$$

The error term remains to be  $O\left(\frac{n}{\log^2 n}\right)$ , as in [1]. The main term simplifies to

$$n\sum_{a=2}^{\log^2 n} \frac{1}{a^k (\log n - \log(a^k))} = \frac{n}{\log n} \sum_{a=2}^{\log^2 n} \frac{1}{a^k (1 - \frac{k \log a}{\log n})}$$
$$\Rightarrow \frac{n}{\log n} \sum_{a=2}^{\log^2 n} \frac{1}{a^k} \left( 1 + O\left(\frac{\log a}{\log n}\right) \right)$$
$$= \frac{n}{\log n} \sum_{a=2}^{\log^2 n} \frac{1}{a^k} + O\left(\frac{n}{\log^2 n}\right) \le \frac{n}{\log n} \sum_{a=2}^{\infty} \frac{1}{a^k} + O\left(\frac{n}{\log^2 n}\right)$$

$$= (\zeta(k) - 1)\frac{n}{\log n} + O\left(\frac{n}{\log^2 n}\right)$$

#### 5. PSP Numbers

A special type of SP Numbers can be defined to be the product of a prime and a square such that the square is a prime square. An approach similar to the proof of Theorem 4.1 can be employed to get the asymptotic estimate of the count of PSP Numbers.

**Theorem 5.1 :** The number of PSP numbers smaller than n is

$$P(2)\frac{n}{\log n} + O\left(\frac{n}{\log^2 n}\right),$$

where P(2) is the prime zeta function of 2.

Proof.

$$PSP(n) = \sum_{p=2}^{\sqrt{n/2}} \pi\left(\frac{n}{p^2}\right), \text{ where } p \text{ is prime.}$$
$$\leq \frac{n}{\log n} \sum_{p=2}^{\infty} \frac{1}{p^2} + O\left(\frac{n}{\log^2 n}\right) = P(2)\frac{n}{\log n} + O\left(\frac{n}{\log^2 n}\right).$$

As a corollary (owing to Theorem 4.1), we have

$$PKP_k(n) = P(k)\frac{n}{\log n} + O\left(\frac{n}{\log^2 n}\right)$$

where PKP Numbers are numbers of the form  $p_1 p_2^k$ , where  $p_1$  and  $p_2$  are arbitrary prime numbers and P(k) is the prime zeta function of k.

#### 6. FUTURE WORK AND CONCLUDING REMARKS

Conjecture 3.1 can be explored further via the use of elliptic curves. Trying to find an SP Number of the form  $x^3 + 1$  is analogous to finding integer solutions to

$$x^3 + 1 = pa^2 \Rightarrow px^3 + p = (pa)^2$$

Letting y = pa, we have the elliptic curve equation

$$y^2 = px^3 + p.$$

Computational experiments have given us over 25 integer solutions to this equation. Potential future work can involve proving the existence of infinite integer solutions to the same.

#### References

- Bhat, R. "Distribution of Square-Prime Numbers." Missouri J. Math. Sci. 34 (1) 121 126, May 2022. https://doi.org/10.35834/2022/3401121 https://arxiv.org/pdf/2109.10238.pdf
- [2] The On-Line Encyclopedia of Integer Sequences, published electronically at https://oeis.org. Sequence A228056. https://oeis.org/A228056
- [3] Pegg, Ed Jr. "Bouniakowsky Conjecture." From MathWorld-A Wolfram Web Resource, created by Eric W. Weisstein. https://mathworld.wolfram.com/BouniakowskyConjecture.html