ORDERING OF NUMBER FIELDS AND DISTRIBUTION OF CLASS GROUPS: ABELIAN EXTENSIONS

WEITONG WANG

ABSTRACT. When p divides the ordering of Galois group, the distribution of $\mathbb{Z}_p \otimes \operatorname{Cl}_K$ is closely related to the problem of counting fields with certain specifications. Moreover, different orderings of number fields affect the answers of such questions in a nontrivial way. So, in this paper, we set up an invariant of number fields with parameters, and consider field counting problems with specifications while the parameters change as a variable. The case of abelian extensions shows that the result of counting abelian fields has a main term with parameters. The estimate of counting cubic fields with a parameter shows that infinite moment is true for some ordering but not very likely for the others.

1. INTRODUCTION

In this paper, we show some results on the distribution of ideal class groups of number fields that could be compared with the *Cohen-Lenstra-Martinet Heuristics* (see Cohen and Martinet [6, Hypothèse 6.6]). We first try to give an overview of the heuristics in this section. Let Γ be a finite group, and let Γ_{∞} be a fixed subgroup of Γ . For each prime $p \nmid |\Gamma|$, we can construct a random $\mathbb{Z}_{(p)}[\Gamma]$ -module X such that for each finite $\mathbb{Z}_{(p)}[\Gamma]$ -module M, we have

$$\mathbb{P}(X \cong M) = \frac{c}{|\operatorname{Aut}_{\Gamma}(M)||M^{\Gamma_{\infty}}|},$$

where c > 0 is a constant (normalizing the measure so that it is a probability space). Let \mathcal{S} be the set of Γ -extensions K/\mathbb{Q} such that its decomposition group at infinity is Γ_{∞} . For any finite Γ -module N, let $N[p^{\infty}] := \mathbb{Z}_{(p)}[\Gamma] \otimes_{\Gamma} N$ be the Sylow p-subgroup of N with its Γ -module structure. For each finite $\mathbb{Z}_{(p)}[\Gamma]$ -module M, we define

$$\mathbb{P}(\operatorname{Cl}_K[p^{\infty}] \cong M) := \lim_{x \to \infty} \frac{|\{K \mid P(K) < x \text{ and } \operatorname{Cl}_K[p^{\infty}] \cong M\}|}{|\{K \mid P(K) < x|\}|}$$

where K runs over all fields in \mathcal{S} . When M runs over all finite $\mathbb{Z}_{(p)}[\Gamma]$ -module, we obtain the distribution of ideal class group $\operatorname{Cl}_K[p^{\infty}]$. The Cohen-Lenstra-Martinet Heuristics predict that for each finite $\mathbb{Z}_{(p)}[\Gamma]$ -module M, we have

(1.1)
$$\mathbb{P}(\operatorname{Cl}_K[p^{\infty}] \cong M) = \mathbb{P}(X \cong M)$$

See also [20, Section 3] for details. Since the random module X is specifically constructed, it is much better understood than ideal class groups. For example, for a finite $\mathbb{Z}_{(p)}[\Gamma]$ -module M, let's define the *M*-moment of ideal class groups as follows:

$$\mathbb{E}(|\operatorname{Hom}_{\Gamma}(\operatorname{Cl}_{K}, M)|) := \frac{\sum_{P(K) < x} |\operatorname{Hom}_{\Gamma}(\operatorname{Cl}_{K}, M)|}{|\{K \mid P(K) < x|\}}$$

Key words and phrases. Cohen-Lenstra heuristics, distribution of class groups, field counting.

where K runs over all fields in S. It is not easy to compute it in general, but according to [20, Theorem 1.2], we have the following

(1.2)
$$\mathbb{E}(|\operatorname{Sur}_{\Gamma}(X,M)|) = \frac{1}{|M^{\Gamma_{\infty}}|}$$

where Sur means surjective homomorphisms. Note that

$$|\operatorname{Hom}_{\Gamma}(\operatorname{Cl}_{K}, M)| = \sum_{N \subseteq M} |\operatorname{Sur}_{\Gamma}(X, N)|.$$

The question whether the heuristics is true or how to prove it could be described as widely open. See the work of Davenport and Heilbronn [8] and Datskovsky and Wright [7] on the $\mathbb{Z}/3\mathbb{Z}$ -moment of class groups of quadratic extensions, and the work of Bhargava [3] on $\mathbb{Z}/2\mathbb{Z}$ -moment of class groups of cubic fields.

The heuristics not only predict the distribution of ideal class groups in the case of Galois number fields, but also for non-Galois extensions. Let Γ' be a fixed subgroup of Γ , and let p be a prime not dividing $|\Gamma|$. Let S' be the set of fields K' such that its Galois closure K/\mathbb{Q} is a Γ -extension with decomposition group Γ_{∞} at infinity and that $K' = K^{\Gamma'}$. We can first construct a maximal $\mathbb{Z}_{(p)}$ -order \mathfrak{o} as a subset of $\mathbb{Z}_{(p)}[\Gamma]$ in the algebra $\mathbb{Q}[\Gamma' \setminus \Gamma/\Gamma']$. If the heuristics hold for $\operatorname{Cl}_K[p^{\infty}]$ when K runs over all fields in S, i.e., we can construct some random $\mathbb{Z}_{(p)}[\Gamma]$ -module X such that for each finite $\mathbb{Z}_{(p)}[\Gamma]$ -module M, we have

$$\mathbb{P}(\mathrm{Cl}_K[p^\infty] \cong M) = \mathbb{P}(X \cong M).$$

Then, we can construct a random \mathfrak{o} -module Y out of X, such that for each finite \mathfrak{o} -module N, we have

$$\mathbb{P}(\mathrm{Cl}_{K'}[p^{\infty}] \cong N) = \mathbb{P}(Y \cong N)$$

where K' runs over fields in S'. See [20, Section 7-8] for details. It is worth mentioning here that when Γ' is a normal subgroup, the random \mathfrak{o} -module Y will coincide with a particular random Γ/Γ' -module given by the heuristics (see [20, Section 9]). So, the heuristics work uniformly for number fields.

Since we will talk about primes $q \mid |\Gamma|$, let's here emphasize the following fact. If A is an abelian group, then let $\operatorname{rk}_p A$ denote the *p*-rank of A where p is a given prime number. Using this notation, the heuristics imply that for all $r = 0, 1, 2, \ldots$

(1.3)
$$\mathbb{P}(\operatorname{rk}_{p}\operatorname{Cl}_{K'} \leq r) := \lim_{x \to \infty} \frac{\sum_{P(K') < x} \mathbf{1}_{\operatorname{rk}_{p} \leq r}(\operatorname{Cl}_{K'}[p^{\infty}])}{\sum_{P(K') < x} 1} > 0,$$
and
$$\lim_{r \to \infty} \mathbb{P}(\operatorname{rk}_{p}\operatorname{Cl}_{K'} \leq r) = 1,$$

where K' runs fields in S', and $\mathbf{1}_{\mathrm{rk}_p \leq r}(A) = 1$ if $\mathrm{rk}_p A \leq r$ and $\mathbf{1}_{\mathrm{rk}_p \leq r}(A) = 0$ otherwise. This can be deduced from (1.1). In addition, the formula of moments (1.2) implies the following. The heuristics implies that for each finite abelian *p*-group *A*, we have

(1.4)
$$\mathbb{E}(|\mathrm{Hom}(\mathrm{Cl}_{K'}, A)|) < \infty.$$

Note that here we forget the Γ -module structure for the convenience of our discussion.

Remark. In the original Cohen-Lenstra-Martinet heuristics, fields are ordered by discriminant, which was an obvious ordering for number fields. Now we have the question of what kind of ordering one should put on the fields. In some cases, ordering fields by discriminant will contradict what is predicted by the heuristics. See Bartel [2] for example. The invariants of number fields that have been used for ordering all rely on the combination of ramified primes of different inertia, and we are mainly focused on the product of ramified primes, which is denoted by P(K). See Wood [21] for more discussion on different orderings.

In the above discussions, we restrict the choice of the prime p to the case when $p \nmid |\Gamma|$. Now let's give an example when this prime divides the order of the Galois group. Recall the genus theory for quadratic number fields, which says that

(1.5)
$$\omega(P(K)) - 1 \le \operatorname{rk}_2 \operatorname{Cl}_K \le \omega(P(K)),$$

where P(K) is the product of ramified primes of K/\mathbb{Q} and $\omega(n)$ is the number of distinct prime divisors for $n \in \mathbb{Z}$. The distribution of the group $\operatorname{Cl}_{K}[2^{\infty}]$ then cannot be described by the approach of the Cohen-Lenstra-Martinet Heuristics. Because, first, the inequality (1.5) basically means that the number of ramified primes determines $\operatorname{rk}_{2}\operatorname{Cl}_{K}$ (up to 1). This phenomenon could be thought of as "predictable", hence contradicting the spirit of the Cohen-Lenstra-Martinet Heuristics. Second, we can show that, for all $r \in \mathbb{N}$,

$$\mathbb{P}(\mathrm{rk}_2 \operatorname{Cl}_K \leq r) = 0$$
 and $\mathbb{E}(|\operatorname{Hom}(\operatorname{Cl}_K, C_2)|) = +\infty$

which is qualitatively different from what is predicted by the heuristics, i.e., (1.3) and (1.4). According to this example, the statistical behaviour of $\operatorname{Cl}_K[2^\infty]$ should be considered more carefully than other parts. See [11, 19] for example. On the other hand, we can also try to generalize the genus theory for quadratic number fields to general ones, whose details are given in Section 3 following the idea of Ishida [12, Chapter 4]. Here we present the main result in a brief way. Let K/\mathbb{Q} be a number field, and let q be a prime. If we have ideal factorization $p\mathcal{O}_K = \mathfrak{p}_1^{e_1} \cdots \mathfrak{p}_m^{e_m}$ such that $\operatorname{gcd}(e_1, \ldots, e_m) \equiv 0 \mod q$, then we call pa ramified prime of type q. For any number field K/\mathbb{Q} , define its genus group \mathcal{G} to be the Galois group of the maximal unramified extension Kk/K obtained by composing with an abelian extension k/\mathbb{Q} . Then, genus theory implies the following.

Theorem 1.1. Let K/\mathbb{Q} be a number field with maximal abelian subextension K_0/\mathbb{Q} . Fix a rational prime q dividing $n := [K : \mathbb{Q}]$. Then, the q-rank of the genus group \mathcal{G} admits the following inequality

 $\operatorname{rk}_q \mathcal{G} \geq |\{p \text{ is a ramified prime of type } q \text{ and } p \equiv 1 \mod q\}| - \operatorname{rk}_q \operatorname{Gal}(K_0/\mathbb{Q}).$

The genus group of K, by definition, is the quotient of the ideal class group. P.Roquette and H.Zassenhaus [18] construct a subgroup of the class group that is associated to ramified primes. The main theorem of their paper says the following.

Theorem 1.2. Let K be a number field of degree n over \mathbb{Q} and q be a given prime number, then

$$\operatorname{rk}_q \operatorname{Cl}_K \geq |\{p \text{ is a ramified prime of type } q\}| - 2(n-1)$$

The obvious difference compared to genus theory is that we do not require that p be 1 mod q in the condition. For more details and comparison between these two theories, see Section 3.

Since the splitting type of a prime p in a field extension K/\mathbb{Q} is determined by its decomposition group G_p and inertia subgroup I_p and so on, it is reduced to a problem of group theory to find out all primes q so that we can apply the above Theorems 1.1 and 1.2 and expect to get nontrivial estimate for $\mathrm{rk}_q \operatorname{Cl}_K$.

Definition 1.3. Let $1 \leq G \leq S_n$ be a finite transitive permutation group. Let $\sigma \in G$ be any permutation. Define $e(\sigma) := \gcd(|\langle \sigma \rangle \cdot 1|, \ldots, |\langle \sigma \rangle \cdot n|)$, i.e., the greatest common divisor of the size of orbits. We call q a non-random prime for G if $q \mid e(\sigma)$ for some $\sigma \in G$. On the other hand, for a permutation $\sigma \in G$, if $q^l \parallel e(\sigma)$, then we call σ an element inertia of type q^l . Define $\Omega(G, q^l)$ to be the subset of G consisting of all elements inertia of type q^l . We denote $\bigcup_{l=1}^{\infty} \Omega(G, q^l)$ by $\Omega(G, q^{\infty})$.

Example 1.4. First let $G = S_3$. If K is a non-Galois cubic number field, then the permutation action of G on $K \to \mathbb{C}$ is exactly the conventional action of S_3 on $\{1, 2, 3\}$, which induces the isomorphism $G \cong \text{Gal}(\hat{K}/\mathbb{Q})$. By checking the elements of S_3 , we see that 3 is a non-random prime for G, e.g., 3 divides the length of (123). Using the language in Theorem 1.2, totally ramified primes satisfy the condition that $e_K(p) \equiv 0 \mod 3$, in other words totally ramified primes are just ramified prime of type 3, hence

 $\operatorname{rk}_3 \operatorname{Cl}_K \ge |\{p \mid p \text{ is totally ramified}\}| - 4.$

Given a transitive permutation group $1 \leq G \leq S_n$, and a non-random prime q for G, we first make the following conjecture on counting fields based on the Malle-Bhargava Heuristics (see [14, 4], [10, p. 291-339] for example) for counting fields with fixed number of ramified primes. For an extension K/k of number fields, we denote its Galois closure by \hat{K} .

Definition 1.5. Let $1 \leq G \leq S_n$ be a transitive permutation group, and let k be a number field. Let S be the set of all number fields $(K/k, \psi)$ such that its Galois closure $(\hat{K}/k, \psi)$ is a G-extension (see Definition 2.1), and that $K = \hat{K}^{G_1}$ where G_1 is the stabilizer of 1. Suppose that Ω is a (nonempty) subset of G that is closed under invertible powering, i.e., if $g^a = h$, $h^b = g$, then $g \in \Omega$ if and only if $h \in \Omega$. Define for the set Ω , and for all $r = 0, 1, 2, \ldots$,

$$\mathbf{1}_{(\Omega,r)}(K) := \begin{cases} 1 & \text{if there are exactly } r \text{ primes } p \nmid |G| \\ & \text{s.t. } I(p) \cap \Omega \neq \emptyset; \\ 0 & \text{otherwise.} \end{cases}$$

where I(p) here means the inertia subgroup of p.

Conjecture 1.6. Keep G, k, S as above. Suppose that $id \notin \Omega$ is a (nonempty) subset of G that is closed under invertible powering.

(1) For all r = 0, 1, 2, ..., there exists some r', such that

$$\sum_{\substack{K \in \mathcal{S} \\ P(K) < x}} \mathbf{1}_{(\Omega, r)}(K) = o\left(\sum_{\substack{K \in \mathcal{S} \\ P(K) < x}} \mathbf{1}_{(\Omega, r')}(K)\right),$$

In this case we say that the conjecture 1 holds for the pair (S, Ω) . (2) For all r = 0, 1, 2, ...,

$$\sum_{\substack{K \in \mathcal{S} \\ P(K) < x}} \mathbf{1}_{(\Omega, r)}(K) = o\left(\sum_{\substack{K \in \mathcal{S} \\ P(K) < x}} 1\right)$$

In this case we say that the conjecture 2 holds for the pair (\mathcal{S}, Ω) .

Using the conjecture on counting fields, we can present our result on the distribution of class groups. Recall that for a number field K, if L/K is an algebraic extension, then let $\operatorname{Cl}(L/K) := \operatorname{ker}(\operatorname{Nm}_{L/K}(\operatorname{Cl}_L \to \operatorname{Cl}_K))$ be the *relative class group*. We see that $\operatorname{Cl}(K/\mathbb{Q}) = \operatorname{Cl}_K$ is just the usual class group.

Theorem 1.7. Let $1 \leq G \leq S_n$ be a transitive permutation group, and let k be a number field. Let S be the set of all number fields $(K/k, \psi)$ such that its Galois closure $(\hat{K}/k, \psi)$ is a G-extension, and that $K = \hat{K}^{G_1}$. Let $H \subseteq G$ be a subgroup such that $\hat{K}^H \subseteq K$ for $K \in S$. If q is a non-random prime for G such that $q \mid [K : \hat{K}^H]$, and Conjecture 1.6(2) holds for (S, Ω) , where $\Omega := \Omega(G, q^{\infty})$, then

 $\mathbb{P}(\mathrm{rk}_q \operatorname{Cl}(K/\hat{K}^H) \le r) = 0 \quad and \quad \mathbb{E}(|\mathrm{Hom}(\mathrm{Cl}(K/\hat{K}^H), C_q)|) = +\infty,$

where K runs over fields in S for the product of ramified primes in K/\mathbb{Q} , and $\operatorname{Cl}(K/K^H)$ denotes the relative class group.

These two phenomena described in the above theorem, zero-probability and infinite moment, justify the notion "non-random prime" from the aspect of statistics. With the help of Class Field Theory and Tauberian Theorems, we can prove the Conjecture 1.6(1) for abelian extensions. To be precise, we have the following.

Theorem 1.8. Let Γ be a finite abelian group with a subgroup Λ , and let S be the set of all abelian Γ -extensions K/\mathbb{Q} . If q is a prime number such that $q \mid |\Gamma/\Lambda|$, then the Conjecture 1 holds for (S, Ω) , where $\Omega := \Omega(\Gamma, q^{\infty})$. In addition, we have

$$\mathbb{P}(\mathrm{rk}_{q} \operatorname{Cl}(K/K^{\Lambda}) \leq r) = 0 \quad and \quad \mathbb{E}(|\mathrm{Hom}(\mathrm{Cl}(K/K^{\Lambda}), C_{q})|) = +\infty,$$

where K runs over fields in S for the product of ramified primes in K/\mathbb{Q} .

For non-abelian extensions, the first obstacle is counting fields. We present here an example, D_4 -fields. Let D_4 be the dihedral group of order 8, and we are going to consider quartic number fields L/\mathbb{Q} whose Galois closure M/\mathbb{Q} are D_4 -fields. According to the work of S.A.Altug, A.Shankar, I.Varma, K.H.Wilson [1], the result of counting such fields by the Artin conductor of 2-dimensional irreducible representation of D_4 is proven. The main result in this case can be summarized as follows.

Theorem 1.9. Let S be the set of quartic number fields L/\mathbb{Q} whose Galois closure are D_4 -extensions M/\mathbb{Q} . We have

$$\mathbb{E}_C(|\mathrm{Hom}(\mathrm{Cl}_L, C_2)|) = +\infty,$$

where the subscript C means that the fields $L \in S$ are ordered by the Artin conductor of 2-dimensional irreducible representation of D_4 .

To summarize this section, the notion "good prime" in [6, Définition 6.1] (see also [20, Section 7]) gives a criterion for us to apply the Cohen-Martinet-Lenstra Heuristics so that we can predict the statistical behaviour of the class groups. In this paper, the notion "non-random prime" predicts the cases when we expect nontrivial subgroup of Cl_K associated to ramified primes and qualitatively different statistical results from "good prime" cases. However, Example 4.2 in Section 4 will show that there are primes p that are neither good in the sense of the heuristics nor non-random in this paper. So, such situations may suggest more ideas on the study of class groups.

ACKNOWLEDGEMENT

I would like to thank Melanie Wood for many useful conversations related to this work. I would like to thank Yuan Liu for comments that helped improve the study such that the related results could be generalized to relative class groups. I would like to thank Alex Bartel, Aaron Landesman, Fabian Gundlach, and Michael Kural for many useful discussions related to this paper. This work was done with the support of National Science Foundation grants DMS-2052036 and DMS-2140043.

2. Basic notations

In this section we introduce some of the notations that will be used in the paper. We use some standard notations coming from analytic number theory. For example, write a complex number as $s = \sigma + it$. The notation $q^l || n$ means that $q^l || n$ but $q^{l+1} \nmid n$. Denote the Euler function by $\phi(n)$. Let $\omega(n)$ counts the number of distinct prime factors of n and so on.

We also follow the notations of inequalities with unspecified constants from Iwaniec and Kowalski [13, Introduction, p.7]. Let's just write down the ones that are important for us. Let X be some space, and let f, g be two functions. Then $f(x) \ll g(x)$ for $x \in X$ means that $|f(x)| \leq Cg(x)$ for some constant $C \geq 0$. Any value of C for which this holds is called an implied constant. Note that we use Vinogradov's $f \ll g$ and Landau's f = O(g) as synonyms. We use $f(x) \approx g(x)$ for $x \in X$ if $f(x) \ll g(x)$ and $g(x) \ll f(x)$ both hold with possibly different implied constants. We say that f = o(g) as $x \to x_0$ if for any $\epsilon > 0$ there exists some (unspecified) neighbourhood U_{ϵ} of x_0 such that $|f(x)| \leq \epsilon g(x)$ for $x \in U_{\epsilon}$.

Since there are multiple ways to describe field extensions, we give the following two definitions to make the term like "the set of all non-Galois cubic number fields" precise.

Definition 2.1. For a field k, by a Γ -extension of k, we mean an isomorphism class of pairs (K, ψ) , where K is a Galois extension of k, and $\psi : \operatorname{Gal}(K/k) \cong \Gamma$ is an isomorphism. An isomorphism of pairs $(\alpha, m_{\alpha}) : (K, \psi) \to (K', \psi')$ is an isomorphism $\alpha : K \to K'$ such that the map $m_{\alpha} : \operatorname{Gal}(K/k) \to \operatorname{Gal}(K'/k)$ sending σ to $\alpha \circ \sigma \circ \alpha^{-1}$ satisfies $\psi' \circ m_{\alpha} = \psi$. We sometimes leave the ψ implicit, but this is always what we mean by a Γ -extension. We also call Γ -extensions of \mathbb{Q} Γ -fields.

Definition 2.2. Let $G \subseteq S_n$ whose action on $\{1, 2, ..., n\}$ is transitive. Let k be a number field. Let $\mathcal{S}(G, k)$ be the set of pairs (K, ψ) such that the Galois closure (\hat{K}, ψ) is a Gextension of k and that $K = \hat{K}^{G_1}$, where G_1 is the stabilizer of 1. In other words ψ defines the Galois action of G on K over k and [K : k] = n. If the base field $k = \mathbb{Q}$, then we just omit it and write $\mathcal{S}(G) := \mathcal{S}(G, \mathbb{Q})$.

Then, we give the notation of counting number fields.

Definition 2.3. Let S = S(G, k) where G is a transitive permutation group, and let $d : S \to \mathbb{R}^+$ be an invariant of number fields (e.g. discriminant or product of ramified primes). Let S(d) be the set S of fields ordered by d. Define

$$N(\mathcal{S}(d); x) := \sum_{\substack{K \in \mathcal{S} \\ d(K) < x}} 1.$$

If f is a function defined over \mathcal{S} , then we define

$$N(\mathcal{S}(d); f; x) := \sum_{\substack{K \in \mathcal{S} \\ d(K) < x}} f(K).$$

In particular, if $f = \mathbf{1}_{(\Omega,r)}$ (see Definition 1.5), then just write

$$N(\mathcal{S}(d); (\Omega, r); x) := N(\mathcal{S}(d); \mathbf{1}_{(\Omega, r)}; x).$$

If there is no danger of confusion, we will use abbreviations:

$$N(x) := N(\mathcal{S}(d); x)$$
 and $N_r(x) := N(\mathcal{S}(d); (\Omega, r); x).$

For example, recall that if q is a rational prime, r is a natural number, then for any finite abelian group A we can define the following function:

$$\mathbf{1}_{\mathrm{rk}_q \leq r}(A) = \begin{cases} 1 & \text{if } \mathrm{rk}_q A \leq r \\ 0 & \text{otherwise.} \end{cases}$$

It could be viewed as a function defined on S, because every $K \in S$ admits an ideal class group Cl_K . Using the notations introduced in this section, if S = S(G) is a set of number fields with G a transitive permutation group, and P means the product of ramified primes, then we can define for a rational prime q and an integer $r \ge 0$ the following notation, which could be called the probability of the p-rank of Cl_K less than or equal to r where K runs over all fields in S for the product of ramified primes:

$$\mathbb{P}(\mathrm{rk}_q \operatorname{Cl}_K \le r) := \lim_{x \to \infty} \frac{N(\mathcal{S}(P); \mathbf{1}_{\mathrm{rk}_q \le r}; x)}{N(\mathcal{S}(P); x)}.$$

3. Non-randomness

The term non-randomness here refers to the case where we can obtain some nontrivial ideal classes in Cl_K associated to the ramified primes. There are two main reasons why we are interested in the relation between ideal classes and ramified primes. The first one is that we have the famous example of genus theory for quadratic number fields as in Section 1, which implies that the 2-rank is determined by the number of ramified primes (up to 1). We of course want to see if similar phenomena also happen when we look at higher degree fields. The second reason is that product of ramified primes is the main ordering we currently use for number fields. Even in the case people use some other orderings (discriminant, conductor etc.), they are still related to ramified primes. It then makes the results, providing information of the relation between nontrivial ideal classes and ramified primes, more valuable. First recall that we've defined the notation ramified prime of type q for a fixed prime q in Section 1. Here let's generalize it a little bit as follows.

Definition 3.1. Let K/\mathbb{Q} be a number field, and let p, q be two rational primes. If $p\mathcal{O}_K = \mathfrak{p}_1^{e_1} \cdots \mathfrak{p}_m^{e_m}$, then let

$$e_K(p) := \gcd(e_1, \ldots, e_m).$$

If $e_K(p) \equiv 0 \mod q^l$ for some $l \ge 1$, then we call p a ramified prime of type q^l .

Because of the fundamental identity

$$[K:\mathbb{Q}] = \sum_{i=1}^{m} e_i f_i$$

where e_i is the ramification index and f_i is the inertia degree for a fixed prime, we only need to discuss ramified primes of type q for $q \mid [K : \mathbb{Q}]$.


Example 3.2. If K/\mathbb{Q} is Galois itself, i.e., $\operatorname{Gal}(K/\mathbb{Q}) \cong \Gamma$, then for any rational prime p, one has $p\mathcal{O}_K = \prod_i^m \mathfrak{p}_i^e$. So it is just a question whether q divides the ramification index e or not. An example would be a quadratic extension K/\mathbb{Q} with q = 2. In this case, p is a ramified prime of type q if and only if p is ramified in K/\mathbb{Q} .

For non-Galois extensions, things become a little more complicated. Let K/\mathbb{Q} be a non-Galois cubic extension with q = 3. Then p is a ramified prime of type 3 if and only if p is totally ramified in K/\mathbb{Q} . Note that there are partially ramified primes for non-Galois cubic extensions, i.e., $p\mathcal{O}_K = \mathfrak{p}_1^2\mathfrak{p}_2$, which are not ramified primes of type 3.

3.1. Genus theory. The goal of this section is to prove Theorem 1.1, which requires a brief introduction on genus theory for number fields. The basic question of genus theory is to find out the maximal unramified abelian extension of a number field K obtained by composing with an absolute abelian number field k. To be precise, we have the following definition.

Definition 3.3. let K/\mathbb{Q} be a number field of degree n, and let k/\mathbb{Q} be the maximal abelian extension such that Kk/K is an unramified extension. We call such a field the *genus field* Kk over K, and call the Galois group Gal(Kk/K) the *genus group* \mathcal{G} .

Since Kk/K is an unramified abelian extension, it is a subextension of the Hilbert extension of K whose Galois group is isomorphic to the class group Cl_K of K, hence the genus group is a quotient group of the class group Cl_K . According to Ishida [12, p.33-39], we make a summary of the results on the genus group. Fix a rational prime q such that $q^t || n$ with $t \geq 1$. Let $A \cong \operatorname{Gal}(k/\mathbb{Q})$ be the Galois group of the abelian extension k/\mathbb{Q} , and let K_0 be the intersection of k and K which is also the maximal abelian subextension of K/\mathbb{Q} . See the diagram for summary below.



Definition 3.4. Let K/\mathbb{Q} be a number field, and let p be a prime number. Let k(p) be the unique subfield of $\mathbb{Q}(\zeta_p)$ with degree $gcd(p-1, e_K(p))$ where ζ_p is a primitive pth root of unity.

Note that k(p) is nontrivial if and only if $gcd(p-1, e_K(p))$ is nontrivial. Moreover the Galois group of $k(p)/\mathbb{Q}$ is cyclic of order $gcd(p-1, e_K(p))$. The first result is to describe the extension k/\mathbb{Q} given the ramified primes of K/\mathbb{Q} .

Theorem 3.5. [12, Chapter IV, Theorem 3] Let K be a number field, and let k be the abelian field such that Kk is the genus field. Let k_1/\mathbb{Q} be the composite of k(p) where p runs through all rational prime numbers such that $p \nmid e_K(p)$, and let k_2/\mathbb{Q} be the intersection of all inertia subfields of k at p where p runs through all rational prime numbers such that $p \nmid e_K(p)$. Then $k = k_1k_2$ and $k_1 \cap k_2 = \mathbb{Q}$. In particular, A admits a subgroup

$$\operatorname{Gal}(k/k_2) \cong \operatorname{Gal}(k_1/\mathbb{Q}) \cong \prod_{p \nmid e_K(p)} \mathbb{Z}/\operatorname{gcd}(p-1, e_K(p)).$$

Now that we have a result on the group A, the description of \mathcal{G} follows from $A/\operatorname{Gal}(K_0/\mathbb{Q}) \cong \mathcal{G}$. In the sense of estimation for $\operatorname{rk}_q q^r \mathcal{G}$, we give the following statement.

Theorem 3.6. Let K/\mathbb{Q} be a number field with maximal abelian subextension K_0/\mathbb{Q} . Fix a rational prime q dividing $n := [K : \mathbb{Q}]$ and some integer $l \ge 1$, the q-rank of the group $q^{l-1}\mathcal{G}$ admits the following inequality

 $\operatorname{rk}_{q} q^{l-1} \mathcal{G} \geq |\{p \text{ is a ramified prime of type } q^{l} \text{ and } p \equiv 1 \mod q\}| - \operatorname{rk}_{q} \operatorname{Gal}(K_{0}/\mathbb{Q}).$

Remark. Theorem 1.1 just follows from this result. Moreover we can see that if $\mathcal{G}[q^{\infty}]$ admits higher torsion part, then $\operatorname{Cl}_K[q^{\infty}]$ must also have higher torsion part. Also, if there is no prime $p \mid e_K(p)$, then $k_2 = \mathbb{Q}$ and $A \cong \prod_{p \nmid e_K(p)} \mathbb{Z}/\operatorname{gcd}(p-1, e_K(p))$, hence $\mathcal{G} \cong (\prod_{p \nmid e_K(p)} \mathbb{Z}/\operatorname{gcd}(p-1, e_K(p)))/\operatorname{Gal}(K_0/\mathbb{Q}).$

Example 3.7. Let K/\mathbb{Q} be a non-Galois cubic field, and q equal 3. Then the requirement $gcd(p-1, e_K(p)) \equiv 0 \mod 3$ is equivalent to p totally ramified in K/\mathbb{Q} and $p \equiv 1 \mod 3$. In other words, we have

 $\operatorname{rk}_3 \operatorname{Cl}_K \ge |\{p \text{ is a totally ramified prime and } p \equiv 1 \mod 3\}|.$

This clearly generalizes genus theory for the quadratic case. See also [12, Chapter 5] for more discussions on the case of odd prime degree.

3.2. The class rank estimate on the invariant part of the class group. In the paper of Roquette and Zassenhaus [18], there is another result on the estimate of the q-rank of the class group with respect to ramified primes whose idea is totally different from genus theory. As seen in previous part, what genus theory really cares about is the genus group, which is the quotient of the class group, while the invariant part of the class group we will talk about is a subgroup of the class group.

We will explain briefly the work of Roquette and Zassenhaus by showing the proof of Theorem 3.11. More importantly, we want to show the construction, which is more useful in this paper. Let's first introduce some notations to present their precise statement. If K is a number field, let rk_K be its rank of global units. Let $v_q(n)$ denote the normalized exponential valuation associated to the rational prime q, i.e., we compute the exponent of q showing up in the factorization of n.

Definition 3.8. Let K/\mathbb{Q} be a number field whose Galois closure is a Γ -extension L/\mathbb{Q} . By viewing the group of fractional ideals \mathscr{I}_K of K as a subgroup of \mathscr{I}_L , we define the *invariant* part of the class group, denoted by C_K^{Γ} , of K under the action of Γ as the image of $\mathscr{I}_L^{\Gamma} \cap \mathscr{I}_K$ in Cl_K , i.e.,

$$C_K^{\Gamma} := \operatorname{im}(\mathscr{I}_L^{\Gamma} \cap \mathscr{I}_K \to \operatorname{Cl}_K).$$

To give a precise statement on the estimation for the invariant part C_K^{Γ} of the class group, we in addition need the following definition of q-radical subfields of K.

Definition 3.9 (q-radical subfields). Let K_q be the subfield of K generated by all elements $\xi \in K$ which are q-radicals over \mathbb{Q} , i.e., $\xi^q \in \mathbb{Q}$. Let $n_q = [K_q : \mathbb{Q}]$ be its degree.

It is clear that $n_q \mid n$, hence one has

$$v_q(n_q) \le v_q(n).$$

But if one wants to ask when $v_q(n_q)$ is strictly less than $v_q(n)$, then the following notation actually gives a condition for it.

Definition 3.10. Define the notation $\delta_K^{(q)}$ to be 1 or 0 according to whether or not the following conditions are *simultaneously* satisfied:

- (1) q > 2;
- (2) K contains a primitive q-th root of unity ζ ;
- (3) There exists some η in K^* such that $\zeta^{-1}\eta^q \in \mathbb{Q}$.

Remark. If all of the above three conditions hold, then one can show that $[K_q(\eta) : K_q] = q$ so that $n_q \cdot q \mid n$ and hence $v_q(n_q) + 1 \leq v_q(n)$. See [18, Section 6].

Using the notations introduced above, we can first state an estimation for C_K due to Roquette and Zassenhaus. Though in the paper [18], the main goal is to prove the result for the *q*-rank of the class group. We here instead describe the structure of the subgroup $C_K^{\Gamma}[q^{\infty}] \subseteq \operatorname{Cl}_K[q^{\infty}]$, which is also due to Roquette and Zassenhaus.

Theorem 3.11. Let K/\mathbb{Q} be a (not necessarily Galois) extension of degree n whose Galois group is Γ , and let q be a given prime number, and let $l \ge 1$ be an integer, then

$$\begin{aligned} &|\{p \text{ is a ramified prime of type } q^l\}| - \left(\mathrm{rk}_K + v_q(n_q) + \delta_K^{(q)}\right) \\ &\leq \mathrm{rk}_q \, q^{l-1} C_K^{\Gamma} \leq \\ &|\{p \text{ is a ramified prime of type } q^l\}|. \end{aligned}$$

We prove the theorem by two lemmas. The first lemma gives a detailed description of the elements in C_K^{Γ} in the sense of fractional ideals.

Lemma 3.12. [18, Equation (8)] Let K/\mathbb{Q} be a number field whose Galois closure a Γ extension L/\mathbb{Q} . For each prime p, define $\mathfrak{a}(p)$ to be the ideal such that $p\mathcal{O}_K = \mathfrak{a}(p)^{e_K(p)}$. Then $\mathscr{I}_K \cap \mathscr{I}_L^{\Gamma}$ is a free abelian group generated by $\{\mathfrak{a}(p)\}$. Let \mathscr{P}_k be the group of principal ideals of a number field k. Then the group $\tilde{C}_K^{\Gamma} := \mathscr{I}_K \cap \mathscr{I}_L^{\Gamma}/\mathscr{P}_{\mathbb{Q}}$ is given by

$$\tilde{C}_K^{\Gamma} \cong \prod_p \mathbb{Z}/e_K(p)$$

The second lemma is to estimate the difference between principal ideals of K and \mathbb{Q} .

Lemma 3.13. [18, Equation(11)] Let K/\mathbb{Q} be a number field whose Galois closure a Γ -extension L/\mathbb{Q} , and let \mathscr{P}_k be the group of principal ideals of a number field k. Then

(3.1) $rk_q \mathscr{P}_K^{\Gamma} / \mathscr{P}_{\mathbb{Q}} \leq \mathrm{rk}_K + v_q(n_q) + \delta_K^{(q)},$

where $\mathscr{P}_{K}^{\Gamma} := \mathscr{P}_{K} \cap \mathscr{I}_{L}^{\Gamma}$.

Now let's prove the theorem.

Proof of Theorem 3.11. Lemma 3.12 gives the upper bound of $\operatorname{rk}_q q^{l-1} C_K^{\Gamma}$, because $\mathscr{P}_{\mathbb{Q}} \subseteq \mathscr{P}_K$. According to the short exact sequence

$$0 \to \mathscr{P}_{K}^{\Gamma}/\mathscr{P}_{\mathbb{Q}} \to \mathscr{I}_{K}^{\Gamma}/\mathscr{P}_{\mathbb{Q}} \to C_{K}^{\Gamma} \to 0,$$

the inequality (3.1) tells us the lower bound directly.

We can apply the theorem to the class group Cl_K , for $C_K^{\Gamma} \subseteq \operatorname{Cl}_K$.

Corollary 3.14. Let K be a number field of degree n over \mathbb{Q} , and let q be a prime number, and let $l \geq 1$ be an integer, then

$$\operatorname{rk}_{q} q^{l-1} \operatorname{Cl}_{K} \geq |\{p \text{ is a ramified prime of type } q^{l}\}| - \left(\operatorname{rk}_{K} + v_{q}(n_{q}) + \delta_{K}^{(q)}\right).$$

Remark. One can show that the number $\operatorname{rk}_{K} + v_q(n_q) + \delta_{K}^{(q)}$ is always smaller than 2(n-1), i.e., the above inequality has a weaker but shorter expression

 $\operatorname{rk}_{q} q^{l-1} \operatorname{Cl}_{K} \ge |\{ \text{ramified primes of type } q^{l} \}| - 2(n-1),$

which proves the statement of Theorem 1.2.

One of the advantages of this theory, as mentioned above, is that C_K^{Γ} is a subgroup of Cl_K . So, we can even try to discuss the relative class group here, i.e., for a subfield $K' \subseteq K$, we want to give a description for $C_K^{\Gamma} \cap \operatorname{Cl}(K/K')$.

Theorem 3.15. Let K/\mathbb{Q} be a number field of degree n whose Galois group is Γ . If $K' \subseteq K$ such that $q^l \mid [K : K']$ where q is a rational prime, and $l \geq 1$, then

 $\operatorname{rk}_{q} q^{l-1} \operatorname{Cl}(K/K') \geq |\{p \text{ is a ramified prime of type } q^{l}\}| - 2(n-1).$

Proof. For each prime p, recall that $\mathfrak{a}(p)$ is the ideal of K such that $p\mathscr{O}_K = \mathfrak{a}(p)^{e_K(p)}$. We've shown in Lemma 3.12 that $\mathfrak{a}(p)$ is fixed by the action of Γ , viewed as an element of \mathscr{I}_L . So, we have the following computation

$$\operatorname{Nm}_{K/K'}(\mathfrak{a}(p)) = \mathfrak{a}(p)^{[K:K']},$$

where $\mathfrak{a}(p)^{[K:K']}$ is treated as an ideal of K'. If $q^l | e_K(p)$, and $\mathfrak{b}(p) := (\mathfrak{a}(p))^{e_K(p)/q^l}$, then $\operatorname{Nm}_{K/K'}(\mathfrak{b}(p))$ becomes a power of $p\mathcal{O}_{K'}$, hence a principal ideal of K'. So, $\mathfrak{b}(p)$ represents an ideal class of $\operatorname{Cl}(K/K')$. By Lemma 3.12, the group C_K^{Γ} is generated by ideals of the form $\mathfrak{a}(p)$, we therefore know that the subgroup $B_K^{\Gamma} := \langle \mathfrak{b}(p) | p$ is a ramified prime of type $q^l \rangle / \mathscr{P}_K \subseteq C_K^{\Gamma}[q]$ is contained in $\operatorname{Cl}(K/K')$. In particular, $\langle \mathfrak{b}(p) \rangle / \mathscr{P}_{\mathbb{Q}} \cong \prod_{i=1}^s \mathbb{Z}/q^i$, where $s = |\{p \text{ is a ramified prime of type } q^l\}|$. Then by Lemma 3.13, we know that $\operatorname{rk}_q \mathscr{P}_K^{\Gamma} / \mathscr{P}_{\mathbb{Q}}$ is bounded above by a constant $\operatorname{rk}_K + v_q(n_q) + \delta_K^{(q)} \leq 2(n-1)$, hence the result.

4. Non-random primes

In Theorem 1.1 and Theorem 1.2, we see that the q-rank $\operatorname{rk}_q \operatorname{Cl}_K$ of class group is related to ramified primes of type q. If we fix the Galois group, a question we can ask is for which primes q we can get ramified primes of type q? In a G-extension (see Definition 2.1), let's first answer this question by the following lemma.

Lemma 4.1. Let G be a finite transitive permutation group, and let q be a fixed prime. If $(K/\mathbb{Q}, \psi)$ is an extension of number fields such that its Galois closure (\hat{K}, ψ) is a G-field and that $K = \hat{K}^{G_1}$, then a prime $p \nmid |G|$ is a ramified prime of type q if and only if $I(p) \cap \Omega(G, q^{\infty}) \neq \emptyset$ where I(p) is the inertia subgroup of p.

Proof. If \mathfrak{P} is a prime of \hat{K} lying above p, then $\mathfrak{P} \cap K$ is of course a prime of K above p. Since G acts on all primes of \hat{K} above p, we know that $\sigma \mathfrak{P} \cap K$ will also run over all primes of K above p when σ runs over all elements in G. But if $\sigma \in G_1$, then $\sigma \mathfrak{P} \cap K = \mathfrak{P} \cap K$. So, we obtain a G-set $G_1 \setminus G$, i.e., $\sigma_1 \sim \sigma_2$ if

$$\sigma_1 \mathfrak{P} \cap K = \sigma_2 \mathfrak{P} \cap K,$$

which means that $G_1\sigma_1 = G_1\sigma_2$. This can also be induced by the permutation action of G on the embeddings $K \to \mathbb{C}$.

One can check that if $G(\mathfrak{P})$ is the decomposition group of \mathfrak{P} , then the orbits of $G(\mathfrak{P})$ on G/G_1 correspond to the primes \mathfrak{p} of K above p, and the size of orbits of $I(\mathfrak{P})$ corresponds to the ramification index respectively.

If p is a tamely ramified prime, then $I(\mathfrak{P})$ is just cyclic. So, p is a ramified prime of type q if and only if the size of orbits of the inertia generator has common divisor q, i.e., $q \mid e(\sigma)$ using the notations in Definition 1.3, which means that $I(\mathfrak{P}) \cap \Omega(G, q^{\infty}) \neq \emptyset$. Note that I(p) is well-defined up to conjugation. And we are done.

Example 4.2 ("Counter-example"). We here show an example where the prime q divides the order of the extension K/\mathbb{Q} but it is *not* a non-random prime. Consider the permutation group G defined as the image of the following morphism $A_4 \to S_6$, $(12)(34) \mapsto (1)(2)(34)(56)$, and $(123) \mapsto (134)(265)$. In the sense of number theory, $\mathcal{S}(G)$ contains number fields K/\mathbb{Q} of degree 6, whose Galois closure \hat{K}/\mathbb{Q} are A_4 -extensions, so that the action of $\text{Gal}(\hat{K}/\mathbb{Q})$ on K/\mathbb{Q} induces a map $A_4 \to S_6$. The stabilizer of 1 is the image of $\{1, (12)(34)\}$ in this case. One can check that the group G has no element required in Definition 1.3 to turn the prime 2 into a non-random prime, despite the fact that 2 divides the order of $[K : \mathbb{Q}]$. Note that according to Cohen and Martinet [6], the prime 2 is not "good", i.e., 2 is not good but not non-random. See also [20, Section 7] for details on the concept "good primes".

Our most important goal in this section is to prove Theorem 1.7, which can justify the notion non-random prime from the view of statistics. Let $\mathcal{S} := \mathcal{S}(G)$ be the set of *G*-fields (see Definition 2.2). If $f : \mathcal{S} \to \mathbb{R}$ is a function, then

$$N_f(x) := N(\mathcal{S}(d); f; x) = \sum_{\substack{K \in \mathcal{S} \\ d(K) < x}} f(K)$$

is the notation for the sum of f over G-fields ordered by the invariant d (see Definition 2.3). In particular, let $N(x) := N(\mathcal{S}(d); x)$ and $N_r(x) := N(\mathcal{S}(d); (\Omega, r); x)$. We reach the theorem in several steps. Recall that we have the Conjectures 1.6 in Section 1. Step zero, as a note of the conjecture itself, is to prove the relation between these two conjectures, i.e., Conjecture 1.6(1) implies Conjecture 1.6(2).

Lemma 4.3. Let G be a transitive permutation group and k be a fixed number field, and let S := S(G, k). Let d be an invariant of number fields in S. Suppose that Ω is a (nonempty)

subset of G that is closed under invertible powering. If for all r = 0, 1, 2, ..., there exists some r' such that

$$N_r(x) = o\left(N_{r'}(x)\right)$$

then for all $r = 0, 1, 2, \ldots$, we have

$$N_r(x) = o\left(N(x)\right).$$

Proof. Since all fields counted by $N_r(x)$ and $N_{r'}(x)$ are also counted by N(x), we have $N_{r'}(x) \leq N(x)$ for all x > 0. The lemma then follows.

The first step is to show that Conjecture 1.6(2), the weaker one, implies zero-probability.

Theorem 4.4. Let $1 \leq G \leq S_n$ be a transitive permutation group, and let S := S(G). Let $G_1 \subseteq H \subseteq G$, where G_1 is the stabilizer of 1 that fixes $K \in S$. Let q be a non-random prime for G such that $q^l \mid [H : G_1]$, where $l \geq 1$, and let $\Omega := \bigcup_{j=l}^{\infty} \Omega(G, q^j)$. Let d be an invariant of the number fields in S. If for all $r = 0, 1, 2, \ldots$, we have

$$N_r(x) = o(N(x)),$$

then for all $r = 0, 1, 2, \ldots$, we have

$$\mathbb{P}(\operatorname{rk}_{q} q^{l-1}\operatorname{Cl}(K/\hat{K}^{H}) \leq r) := \lim_{x \to \infty} \frac{N(\mathcal{S}(d); \mathbf{1}_{\operatorname{rk}_{q} \leq r}(q^{l-1}\operatorname{Cl}(K/K^{H})); x)}{N(x)} = 0,$$

where K runs over fields in S ordered by d, and \hat{K} is the Galois closure of K.

Proof. First of all, recall that we have for all finite abelian group A and for all r = 0, 1, 2, ...,

$$\mathbf{1}_{\mathrm{rk}_q \le r}(A) = \begin{cases} 1 & \text{if } \mathrm{rk}_q A \le r \\ 0 & \text{otherwise.} \end{cases}$$

According to Theorem 3.15, if there are at least r + 2n ramified primes p for K/\mathbb{Q} contained in the set $\{p \nmid |G| : I(p) \cap \Omega \neq \emptyset\}$, then

$$\operatorname{rk}_{q} q^{l-1} \operatorname{Cl}(K/\hat{K}^{H}) > r.$$

If $N_r(x) = o(N(x))$ for all r, then this implies that

$$\mathbb{P}(\mathrm{rk}_{q} q^{l-1} \operatorname{Cl}(K/\hat{K}^{H}) \leq r) = \lim_{x \to \infty} \frac{N(\mathcal{S}(d); \mathbf{1}_{\mathrm{rk}_{q} \leq r}(q^{l-1} \operatorname{Cl}(K/\hat{K}^{H})); x)}{N(\mathcal{S}, d; x)}$$
$$\leq \lim_{x \to \infty} \frac{N(\mathcal{S}(d); \sum_{i=0}^{r+2n} \mathbf{1}_{(\Omega,i)}; x)}{N(x)} = 0.$$

Then let's prove that bounded probability, hence also zero-probability, implies infinite moment.

Theorem 4.5. Let $1 \leq G \leq S_n$ be a transitive permutation group, and let S := S(G). Let $G_1 \subseteq H \subseteq G$, where G_1 is the stabilizer of 1 that fixes $K \in S$. Let d be an invariant of the number fields in S. Let q be a rational prime and $l \geq 1$ be an integer. If there exists some constant $0 \leq c < 1$ such that for all r = 0, 1, 2, ..., we have

$$\mathbb{P}(\operatorname{rk}_{q} q^{l-1} \operatorname{Cl}(K/\hat{K}^{H}) \leq r) \leq c,$$
13

then,

$$\mathbb{E}(|\operatorname{Hom}(q^{l-1}\operatorname{Cl}(K/\hat{K}^{H}), C_{q})|) := \lim_{x \to \infty} \frac{N(\mathcal{S}(d); |\operatorname{Hom}(q^{l-1}\operatorname{Cl}(K/\hat{K}^{H}), C_{q})|; x)}{N(x)}$$
$$= +\infty,$$

where K runs over fields in S ordered by d, and \hat{K} is the Galois closure of K, and C_q is the cyclic group of order q.

Proof. According to similar idea, $\mathbf{1}_{\mathrm{rk}_q=r}(A)$ to be the indicator that tells us if $\mathrm{rk}_q A = r$. By definition of the C_q -moment and the probability of the q-rank, for all $r \geq 0$, we have

$$\mathbb{E}(|\operatorname{Hom}(q^{l-1}\operatorname{Cl}(K/\hat{K}^{H}), C_{q})|)$$

$$= \lim_{x \to \infty} \frac{N(\mathcal{S}(d); |\operatorname{Hom}(q^{l-1}\operatorname{Cl}(K/\hat{K}^{H}), C_{q})|; x)}{N(x)}$$

$$= \lim_{x \to \infty} \sum_{n=0}^{\infty} q^{n} \frac{N(\mathcal{S}(d); \mathbf{1}_{\operatorname{rk}_{q}=n}(q^{l-1}\operatorname{Cl}(K/\hat{K}^{H})); x)}{N(x)}$$

$$\geq q^{r} \lim_{x \to \infty} \left(1 - \frac{N\left(\mathcal{S}(d); \mathbf{1}_{\operatorname{rk}_{q} \leq r-1}(q^{l-1}\operatorname{Cl}(K/\hat{K}^{H}))\right); x)}{N(x)}\right)$$

$$= q^{r} \cdot (1 - \mathbb{P}(\operatorname{rk}_{q} q^{l-1}\operatorname{Cl}(K/\hat{K}^{H}) < r)) \geq q^{r}(1 - c).$$

For any number N > 0, by taking a large enough r > 0, we have

$$\mathbb{E}(|\mathrm{Hom}(q^{l-1}\operatorname{Cl}(K/\hat{K}^H), C_q)|) > N,$$

hence $\mathbb{E}(|\operatorname{Hom}(q^{l-1}\operatorname{Cl}(K/\hat{K}^H), C_q)|) = +\infty.$

In short, we can just say that whenever the Conjecture 1.6 holds, then we have zeroprobability and infinite moment, i.e., Theorem 1.7 is true. We will show in later sections that for abelian extensions, the Conjecture 1.6(1) holds (see Theorem 6.2). Here we discuss an example where the fields are not ordered by product of ramified primes.

Example 4.6 (Ordering by discriminant). Let's consider the set $S := S(S_3)$ of non-Galois cubic extensions. Let $\Omega := \Omega(S_3, 3) = \{(123), (132)\}$. In this case we want to show that the analogous statement of Conjecture 1.6(1) is *false* for (S, Ω) when the fields ordered by *discriminant*, despite the fact that 3 is non-random for S_3 .

According to the work of Bhargava, Shankar, and Tsimerman [5, Theorem 8], there exists some positive constants c_1, c_2 such that

$$N_0(x) \sim c_1 x$$
 and $N(x) \sim c_2 x$,

when the cubic fields are ordered by discriminant. So, counting nowhere totally ramified cubic fields ordered by discriminant admits a main term c_1x and counting all non-Galois cubic fields gives a main term c_2x . This already contradicts the analogous statement of Conjecture 1.6(2) when ordering fields by discriminant, the weaker one. On the other hand, Proposition 7.9 shows that, under some hypothesis, Conjecture 1.6(2) holds for (\mathcal{S}, Ω) . This shows that the conjecture, and possibly the statistical behaviors of non-random primes are dependent on the ordering of number fields.

5. DIRICHLET SERIES AND TAUBERIAN THEOREM

In this section, we discuss the analytic properties of some Dirichlet series, which could be treated as useful tools in later sections.

5.1. Tauberian Theorem. Let's first present a Tauberian Theorem that is used in the paper repeatedly. Recall that for the complex variable $s \in \mathbb{C}$, we write $s = \sigma + it$.

Theorem 5.1 (Delange-Ikehara). [16, Appendix II Theorem I] Assume that the coefficients of a Dirichlet series are real and non-negative, and that it converges in the half-plane $\sigma > 1$, defining a holomorphic function f(s). Assume, moreover, that in the same half-plane one can write

$$f(s) = \sum_{j=0}^{q} g_j(s) \log^{b_j} \left(\frac{1}{s-1}\right) (s-1)^{-\alpha_j} + g(s),$$

where functions g, g_0, \ldots, g_q are holomorphic in the closed half plane $\sigma \geq 1$, the b_j -s are non-negative rational integers, α_0 is a positive real number, $\alpha_1, \ldots, \alpha_q$ are complex numbers with $\Re \alpha_j < \alpha_0$, and $g_0(1) \neq 0$.

Then for the summatory function $S(x) = \sum_{n \le x} a_n$ we have, for x tending infinity,

$$S(x) = \left(\frac{g_0(1)}{\Gamma(\alpha_0)} + o(1)\right) x(\log x)^{\alpha_0 - 1} (\log \log x)^{b_0}.$$

If f satisfies the same assumptions, except that $\alpha_0 = 0$ and $b_0 \ge 1$, then

$$S(x) = (b_0 g_0(1) + o(1))x \frac{(\log \log x)^{b_0 - 1}}{\log x}.$$

See the paper of Delange [9] for more results on Tauberian Theorems.

5.2. Dirichlet L-series. For any Dirichlet character

$$\chi: (\mathbb{Z}/m\mathbb{Z})^* \to S^1 = \{ s \in \mathbb{C} \mid |s| = 1 \},\$$

we obtain a multiplicative function $\chi : \mathbb{Z} \to \mathbb{C}$ by

$$\chi(n) = \begin{cases} \chi(n \mod m) & \text{ for } \gcd(n,m) = 1, \\ 0 & \text{ for } \gcd(n,m) \neq 1. \end{cases}$$

Define the *Dirichlet L-series* to be the form

$$L(\chi, s) := \sum_{n=1}^{\infty} \frac{\chi(n)}{n^s}$$

It admits many analytic properties similar to those of Riemann zeta function $\zeta(s)$.

Proposition 5.2. [17, VII.2.1] The series $L(\chi, s)$ converges absolutely and uniformly in the domain $\Re(s) \ge 1 + \delta$, for any $\delta > 0$. It therefore represents a holomorphic function on the half-plane $\Re(s) > 1$. We have Euler's identity

$$L(\chi, s) = \prod_{\substack{p \\ 15}} \frac{1}{1 - \chi(p)p^{-s}}.$$

It also admits a holomorphic continuation to the complex plane (with a pole at s = 1 when χ is the trivial character modulo m). To present the result, we need some notations. Define the *exponent* $p \in \{0, 1\}$ of χ by

$$\chi(-1) := (-1)^p \chi(1).$$

Define

$$\Gamma(\chi, s) := \Gamma\left(\frac{s+p}{2}\right) = \int_0^\infty e^{-y} y^{(s+p)/2} \frac{\mathrm{d}\, y}{y}.$$

Define

$$L_{\infty}(\chi, s) := \left(\frac{m}{\pi}\right)^{s/2} \Gamma(\chi, s).$$

Define the *completed L-series* of the character χ to be

$$\Lambda(\chi, s) := L_{\infty}(\chi, s) L(\chi, s).$$

For $n \in \mathbb{Z}$, the Gauss sum $\tau(\chi, n)$ associated to the Dirichlet character χ is defined to be the complex number

$$\tau(\chi, n) := \sum_{\nu=0}^{m-1} \chi(\nu) e^{2\pi i \nu n/m}.$$

When n = 1, write $\tau(\chi) := \tau(\chi, 1)$. Finally, we can give the statement of the functional equation of the completed L-series $\Lambda(\chi, s)$.

Theorem 5.3. [17, VII.2.8] If χ is a nontrivial primitive Dirichlet character, then the completed L-series $\Lambda(\chi, s)$ admits a holomorphic continuation to the whole complex plane \mathbb{C} and satisfies the functional equation

$$\Lambda(\chi, s) = W(\chi)\Lambda(\bar{\chi}, 1-s)$$

with the factor $W(\chi) = \frac{\tau(\chi)}{i^p \sqrt{m}}$. This factor has absolute value 1.

Using the functional equation, we can obtain the following result on $L(\chi, s)$ itself.

Proposition 5.4. If χ is a nontrivial primitive Dirichlet character, then $L(\chi, s)$ admits a holomorphic continuation to the whole complex plane \mathbb{C} which is zero only at -2n - p where p is the exponent.

Proof. Let $A(x) := \sum_{n < x} \chi(n)$. When χ is nontrivial, we have

$$|A(x)| \le m,$$

for all x > 0. So when $\Re(s) > 0$, the integral form

$$L(\chi, s) = s \int_0^\infty x^{-s-1} A(x) \,\mathrm{d} x$$

shows that $L(\chi, s)$ is absolutely convergent. The Gamma function

$$\Gamma(s) = \int_0^\infty e^{-y} y^{-s} \frac{\mathrm{d}\, y}{y}$$

is holomorphic and nowhere zero when $\Re(s) > 0$. Its meromorphic continuation to \mathbb{C} admits simple poles at s = -n, $n = 0, 1, 2, \ldots$ So we see that $L_{\infty}(\chi, s)$ admits a meromorphic continuation to the whole complex plane, which is nowhere zero and admits simple poles at $(s+p)/2 = -n, n = 0, 1, 2, \dots$ By the functional equation, we see that

$$L(\chi, s)L_{\infty}(\chi, s) = W(\chi)L(\bar{\chi}, 1-s)L_{\infty}(\bar{\chi}, 1-s).$$

By comparing the poles, we have the following. If the exponent p = 0, then we see that $L(\chi, s)$ admits a simple zero at s = -2n, n = 0, 1, 2, ... If the exponent p = 1, then $L(\chi, s)$ admits a simple zero at s = -2n - 1, n = 0, 1, 2, ...

Remark. Actually we have

$$L(\chi, 1-k) = -\frac{B_{k,\chi}}{k}$$

where $B_{k,\chi}$ is the generalized Bernoulli number. See Neukirch [17, VII.2.9] for more details.

5.3. **Some applications.** Let's introduce a notation, which is inspired by problems of counting fields.

Definition 5.5. Let \mathcal{P} be the set of all rational primes. For a pair (m, n) of relatively prime numbers, let

$$\mathcal{P}(m,n) := \{ p \in \mathcal{P} \mid p \equiv n \bmod m \}.$$

Define

$$\zeta(m,n;s) = \prod_{p \in \mathcal{P}(m,n)} (1-p^{-s})^{-1}$$

when $\Re(s) > 1$.

Let's show that a suitable power of the function $\zeta(m, n; s)$ admits a meromorphic extension to the closed half-plane $\sigma \geq 1$. Recall that ϕ is the Euler function, i.e.,

$$\phi(n) = |(\mathbb{Z}/n)^*|.$$

Lemma 5.6. Let (m, n) be a pair of relatively prime numbers. The function $\zeta(m, n; s)^{\phi(m)}$ admits a meromorphic continuation to the half plane $\sigma > 1/2$ with a simple pole at s = 1. To be precise, $\zeta(m, n; s)$ defines a holomorphic function in the open half-plane $\sigma > 1$, and there exists some holomorphic function $f_0(s)$ in the closed half plane $\sigma \ge 1$ such that

$$\zeta(m,n;s)^{\phi(m)} = f_0(s) \frac{1}{s-1}.$$

Proof. Let $\langle \mathcal{P}(m,n) \rangle$ denote the semi-subgroup of $\langle \mathbb{Z}_{>0}, \cdot \rangle$ generated by $\mathcal{P}(m,n)$. Since

$$\sum_{\substack{n \le x \\ n \in \langle \mathcal{P}(m,n) \rangle}} \frac{1}{|n^s|} \le \sum_{n \le x} \frac{1}{|n^s|}$$

we know that $\zeta(m, n; s)$ converges absolutely and uniformly in $\sigma > 1 + \delta$ for any $\delta > 0$. Then for any $\sigma > 1$, we have

(5.1)
$$\log \zeta(m,n;s) = \sum_{p \in \mathcal{P}(m,n)} p^{-s} + \frac{1}{2} \sum_{p \in \mathcal{P}(m,n)} p^{-2s} + \cdots$$

Similarly for any Dirichlet character $\chi \mod m$, we have

$$\log L(s,\chi) = \sum_{p \in \mathcal{P}} \frac{\chi(p)}{p^s} + \frac{1}{2} \sum_{p \in \mathcal{P}} \frac{\chi(p^2)}{p^{2s}} + \cdots$$

Therefore

(5.2)
$$\log \zeta(m,n;s) = \frac{1}{\phi(m)} \sum_{\chi} \overline{\chi}(n) \log L(s,\chi) + g(s)$$

for all $\sigma > 1$, where χ runs over all Dirichlet characters modulo m, and g(s) is given by

$$g(s) = \sum_{n=2}^{\infty} \frac{1}{n} \sum_{p \in \mathcal{P}(m,n)} \left(p^{-ns} - \frac{1}{\phi(m)} \sum_{\chi} \overline{\chi}(n) \chi(p^n) p^{-ns} \right)$$
$$= \sum_{n=2}^{\infty} \frac{1}{n} \sum_{p^n \neq n \bmod m} p^{-ns}$$

Note that g(s) is absolutely convergent in $\sigma > 1/2$. By taking the exponent, we have

$$\zeta(m,n;s)^{\phi(m)} = \prod_{\chi} L(s,\chi)^{\overline{\chi}(n)} \cdot h(s)$$

where h(s) is a non-vanishing function in $\sigma > 1/2$. Therefore the pole behaviour of $\zeta(m, n; s)$ is the same as $\prod_{\chi} L(s, \chi)^{\overline{\chi}(n)}$, hence the same as $L(s, \chi_0)$ where χ_0 is trivial Dirichlet character modulo m (see for example [15, 4.8,4.9]), which concludes the proof for the formula. It is clear that $h(s) \neq 0$ in the half-plane $\sigma > 1/2$, hence also f(s).

It is a classical result saying that

$$\sum_{p \in \mathcal{P}(m,n)} p^{-s} = \frac{1}{\phi(m)} \log\left(\frac{1}{s-1}\right) + O(1),$$

where gcd(m, n) = 1 and $\sigma > 1$, see [13, Equation (2.25)] for example. We prove a statement of analytic continuation so that it satisfies the condition of the Tauberian Theorem 5.1.

Lemma 5.7. For a given pair of coprime numbers (m, n), the series $\sum_{p \in \mathcal{P}(m,n)} p^{-s}$ defines a holomorphic function in the open half-plane $\Re(s) > 1$, and there is some holomorphic function g(s) in the closed half plane $\sigma \geq 1$, such that

$$\sum_{p \in \mathcal{P}(m,n)} p^{-s} = \frac{1}{\phi(m)} \log\left(\frac{1}{s-1}\right) + g(s).$$

Proof. Compared to Riemann zeta function, we know that the series $\sum_{p \in \mathcal{P}(m,n)} p^{-s}$ converges in the open half-plane $\sigma > 1$, hence defines a holomorphic function. Let's consider the formula coming from (5.1):

(5.3)
$$\sum_{p \in \mathcal{P}(m,n)} p^{-s} = \log \zeta(m,n;s) - \frac{1}{2} \sum_{p \in \mathcal{P}(m,n)} p^{-2s} - \cdots$$

By Proposition 5.4, if χ is a nontrivial Dirichlet character, then the Dirichlet L-function $L(\chi, s)$ is non-zero in the half-plane $\Re(s) > 0$. So the formula (5.2) says that $\log \zeta(m, n; s)$ admits a holomorphic continuation to the closed half-plane $\Re(s) \ge 1$ (a neighbourhood of $\Re(s) > 1$) except on the interval $\sigma \le 1$ of the real line. The other terms on the right-hand

side of (5.3) defines a holomorphic function in the closed half-plane $\sigma \ge 1$. So, there exists some holomorphic function g(s) in the closed half-plane $\sigma \ge 1$ such that

$$\sum_{p \in \mathcal{P}(m,n)} p^{-s} = \frac{1}{\phi(m)} \log\left(\frac{1}{s-1}\right) + g(s).$$

Then let's show the main result of this section.

Definition 5.8. We call a function $b : \mathcal{P} \to \mathbb{Z}_{\geq 0}$ a function modulo m, if b(p) = b(q) for all $p \equiv q \mod m$. In this case b is also a function $(\mathbb{Z}/m\mathbb{Z})^* \to \mathbb{Z}_{\geq 0}$. Let $U := \{s \in \mathcal{C} \mid \Re(s) > 0\}$ be the open half-plane of \mathbb{C} . We call $a : \mathcal{P} \times U \to \mathbb{C}$ a holomorphic function if for each $p \in \mathcal{P}$, the function $a_p(s) := a(p, s) : U \to \mathbb{C}$ is holomorphic with $a_p(\sigma)$ being real for all $\sigma \in \mathbb{R}_+$. We say that a holomorphic function $a : \mathcal{P} \times U \to \mathbb{C}$ is a coefficient function of main term $b : \mathcal{P} \to \mathbb{R}_{>0}$ if there exists some real number $\delta > 0$ independent of p such that

$$a_p(s) - b(p) \ll p^{-\delta\sigma}$$

when $\Re(s) > 0$, where the implied constant is independent of p.

An example of such coefficient function a could be constructed as follows. Let $l(x) = ax \in \mathbb{C}[x]$ be a linear polynomial where a is a positive real number. Define $a(p,s) := l(p^{-\delta s}) + b(p) = ap^{-\delta s} + b(p)$, where $\delta > 0$ a real number, and $b : \mathcal{P} \to \mathbb{R}_{\geq 0}$ is a function modulo m. Then a(p,s) is a coefficient function of main term b_1 . Moreover, one can show that $a(p,s)^{-1}$ is also a coefficient function.

Theorem 5.9. If $b : \mathcal{P} \to \mathbb{Z}_{\geq 0}$ is a function modulo m, and $a : \mathcal{P} \times U \to \mathcal{C}$ is a coefficient function of main term b, then the infinite product

$$D(s) := \prod_{p \in \mathcal{P}} (1 + a(p, s)p^{-s})$$

defines a holomorphic function in the open half-plane $\Re(s) > 1$, and there exists some holomorphic function f in the closed half-plane $\Re(s) \ge 1$ such that $f(s) \ne 0$ for all $\Re(s) \ge 1$ and such that $D(s)^{\phi(m)} = f(s)(s-1)^{\beta}.$

$$\frac{1}{2} \sum_{i=1}^{n} \frac{1}{i} \left(\cdot \right)$$

$$\beta = \sum_{\alpha \in (\mathbb{Z}/m\mathbb{Z})^*} b(\alpha).$$

Proof. First of all, let $b : \mathcal{P} \to \mathbb{Z}_{\geq 0}$ be the function such that b(p) = 1 if $p \equiv \alpha \mod m$ and b(p) = 0 otherwise, where $\alpha \in (\mathbb{Z}/m\mathbb{Z})^*$ is a fixed element. Then by comparing with $\zeta(s)$, we see that D(s) is a holomorphic function in the open half-plane $\Re(s) > 1$. Moreover, when $\Re(s) > 1$, we have

$$\begin{aligned} \left|\log \zeta(\alpha, m; s)/D(s)\right| &= \left|\sum_{\substack{p \in \mathcal{P} \\ p \equiv \alpha \bmod m}} \sum_{n=1}^{\infty} \frac{1}{n} p^{-ns} (1 - (-1)^{n-1})\right| \\ &\leq \sum_{n=2}^{\infty} \frac{2}{n} p^{-n\sigma}. \end{aligned}$$

This shows that $\zeta(\alpha, m; s)/D(s)$ admits a holomorphic continuation to the closed half-plane $\Re(s)$, and we admit the desired meromorphic continuation for $D(s)^{\phi(m)}$ by Lemma 5.6.

Now for any $b: \mathcal{P} \to \mathbb{Z}_{>0}$, we can write it as

$$b = b_{\alpha_1} + \dots + b_{\alpha_l},$$

where b_{α} is the function such that $b_{\alpha}(p) = 1$ if $p \equiv \alpha \mod m$ and $b_{\alpha}(p) = 0$ otherwise. In other words, if $b(\alpha) = n$, the there are exactly n functions b_{α} showing up in the linear combination. So, we have

$$D(s) = \prod_{p \in \mathcal{P}} (1 + \sum_{i=1}^{l} b_{\alpha_i}(p) p^{-s}).$$

Define

$$\tilde{D}(s) := \prod_{p \in \mathcal{P}} \prod_{i=1}^{l} (1 + b_{\alpha_i}(p)p^{-s}).$$

It is clear that D(s) is a holomorphic function in the open half-plane $\Re(s) > 1$ and its $\phi(m)$ -power admits a meromorphic continuation to the closed half-plane $\Re(s) \geq 1$. When $\Re(s) > 1$, we have

$$\begin{aligned} |\log \tilde{D}(s)/D(s)| &= \Big| \sum_{p \in \mathcal{P}} \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} p^{-ns} \Big((\sum_{i=1}^{l} b_{\alpha_i}(p))^n - \sum_{i=1}^{l} b_{\alpha_i}^n(p) \Big) \Big| \\ &\leq \sum_{p \in \mathcal{P}} \sum_{n=2}^{\infty} \frac{2}{n} p^{-n\sigma} b(p)^n. \end{aligned}$$

Since b(p) is bounded, we see that $\tilde{D}(s)/D(s)$ admits a holomorphic continuation to the closed half-plane $\Re(s) > 1$. This shows the analytic properties, in particular the meromorphic continuation, of $D(s)^{\phi(m)}$.

Finally let a be a coefficient function of main term b. If $B = \max_{p} \{b(p)\}$, then by comparing with

$$\prod_{p \in \mathcal{P}} (1 + (B+1)p^{-s}),$$

we see that D(s) is absolutely and uniformly convergent in $\Re(s) > 1 + \delta$ for any $\delta > 0$. This shows that D(s) is a holomorphic function in the open half-plane $\Re(s) > 1$. Let

$$\tilde{D}(s) := \prod_{p \in \mathcal{P}} (1 + b(p)p^{-s}).$$

By similar idea, we see that $\tilde{D}(s)$ is also absolutely and uniformly convergent in $\Re(s) > 1 + \delta$ for any $\delta > 0$. When $\Re(s) > 1$, there exist some $\delta > 0$ such that

$$|\log D(s)/\tilde{D}(s)| = \left|\sum_{p\in\mathcal{P}}\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} p^{-ns} (a^n - b^n)\right| \\ \ll \sum_{p\in\mathcal{P}} \left(p^{-(1+\delta)\sigma} + \sum_{n=2}^{\infty} \frac{1}{n} p^{-n\sigma} ((b+1)^n + b^n)\right).$$

This shows that $D(s)/\tilde{D}(s)$ not only is convergent in $\Re(s) > 1$, but also admits a holomorphic continuation to $\Re(s) \ge 1$. So the meromorphic continuation of $D(s)^{\phi(m)}$ follows from the one of $\tilde{D}(s)^{\phi(m)}$.

Theorem 5.10. For a finite set of positive integers $\{n_1, \ldots, n_l\}$, viewed as a vector \underline{n} , let $c(\underline{n}) := \max\{n_1^{-1}, \ldots, n_l^{-1}\}$, and let $r(\underline{n}) := \#\{1 \le i \le l \mid n_i^{-1} = c(\underline{n})\}$. Let r be a positive integer. If b_1, \ldots, b_r are non-zero functions modulo m, and if a_i is a coefficient function of main term b_i , for all $i = 1, \ldots, r$, then the series

$$D_r(s) := \sum_{\substack{\underline{p} \in \mathcal{P}^r \\ p_1 < \dots < p_r}} \prod_{i=1}^r a_i(p_i, s) p_i^{-n_i s},$$

where $\underline{n} = (n_1, \ldots, n_r)$ are positive integers, defines a holomorphic function in the open halfplane $\Re(s) > c(\underline{n})$, and there exists some holomorphic functions $f_0, \ldots, f_{r(\underline{n})}$ in the closed half-plane $\Re(s) \ge c(\underline{n})$ such that f_0 could be taken to be a positive constant and such that

$$D_r(s) = \sum_{i=0}^{r(c)} f_i(s) \left(\log \frac{1}{s - c(\underline{n})}\right)^{r(\underline{n}) - i}.$$

Proof. We prove the statement by induction on r. If r = 1, then the series is just

$$D_1(s) = \sum_{p \in \mathcal{P}} a(p, s) p^{-ns}.$$

It suffices to prove the statement for n = 1. By comparing with Since a(p, s) is bounded for all $p \in \mathcal{P}$ and $s \in \Re(s) > 0$, we see that D(s) defines a holomorphic function in the open half-plane $\Re(s) > 1$. Let

$$\tilde{D}_1(s) := \sum_{p \in \mathcal{P}} b(p) p^{-s},$$

then $D_1(s)$ is also a holomorphic function in the open half-plane $\Re(s) > 1$ by similar argument. There exist some $\delta > 0$ such that

$$|D_1(s) - \tilde{D}_1(s)| = \left| \sum_{p \in \mathcal{P}} (a - b) p^{-s} \right|$$
$$\ll \sum_{p \in \mathcal{P}} p^{-(1+\delta)\sigma}.$$

This shows that the difference $D_1(s) - \tilde{D}_1(s)$ could be extended to $\Re(s) \ge 1$ analytically. So the continuation of $D_1(s)$ follows from the one of $\tilde{D}_1(s)$, which is a corollary of Lemma 5.7.

Now assume that the statement holds for the cases $1, 2, \ldots, r-1$ with $r \ge 2$. Let $\underline{n}^r = (n_1, \ldots, n_r)$ and $\underline{n}^{r-1} = (n_1, \ldots, n_{r-1})$. For any coefficient function a of main term b, write

$$D_a(s) := \sum_{p \in \mathcal{P}} a(p, s) p^{-s}.$$

Then

$$D_{r-1}(s) \cdot D_{a_r}(n_r s) = D_r(s) + \sum_{i=1}^{r-1} D_{r-1}^{(i)}(s),$$

where

$$D_{r-1}^{(i)}(s) = \sum_{\substack{\underline{p} \in \mathcal{P}^{r-1} \\ p_1 < \dots < p_{r-1}}} a_1(p_1, s) p_1^{-n_1 s} \cdots a_{i}(p_i, s) a_r(p_i, s) p_i^{-(n_i + n_r)s} \cdots a_{r-1}(p_{r-1}, s) p_{r-1}^{-n_{r-1}s}$$

Note that the function $\tilde{a}_i(p,s) := a_i(p,s)a_r(p,s)$ is a coefficient function of main term $b_i \cdot b_r$. So, $D_{r-1}^{(i)}(s)$ satisfies the induction assumption. Let $\underline{n}_i^{r-1} := (n_1, \ldots, n_i + n_r, \ldots, n_{r-1})$, then there exists holomorphic functions $f_{i0}, \ldots, f_{ir(\underline{n}_i)}$ in the closed half-plane $\Re(s) \ge 1$ such that

$$D_{r-1}^{(i)}(s) = \sum_{j=0}^{r(\underline{n}_i^{r-1})} f_{ij}(s) \left(\log \frac{1}{s - c(\underline{n}_i^{r-1})}\right)^{r(\underline{n}_i^{r-1}) - j}$$

By reordering the indices if necessary, we may assume without loss of generality that $n_r^{-1} = c(\underline{n}^r)$. Then we see that $c(\underline{n}_i^{r-1}) < c(\underline{n}^r)$, and that $r(\underline{n}_i^{r-1}) < r(\underline{n}^r)$. And the continuation of $D_{r-1}(s)D_{a_r}(n_rs)$ is given by the form

(5.4)
$$h(s)\log\frac{1}{s-n_r^{-1}}\cdot\sum_{i=0}^{r(\underline{n}^{r-1})}g_{r-i}(s)\left(\log\frac{1}{s-c(\underline{n}^{r-1})}\right)^i,$$

where the function g_i is holomorphic in the closed half-plane $\Re(s) \ge c(\underline{n}^{r-1})$, and h is a positive constant.

If $n_r^{-1} = c(\underline{n}^r) > c(\underline{n}^{r-1})$, then clearly we obtain the following continuation

$$D(s) = f_1(s) \log \frac{1}{s - c(\underline{n})} + f_0(s),$$

where f could be take to be the positive constant equal to

$$\lim_{s \to 1^+} \frac{D(s)}{-\log(s-1)} = D_{r-1}(1),$$

because all other factors and terms showing up are holomorphic in the closed half-plane $\Re(s) \ge 1$. Otherwise, we see that (5.4) becomes

$$\sum_{i=0}^{r(\underline{n}^r)} \tilde{f}_i(s) \left(\log \frac{1}{s - c(\underline{n}^r)}\right)^{r(\underline{n}^r) - i},$$

with \tilde{f}_0 a positive constant. Since $r(\underline{n}^r) > r(\underline{n}_i^{r-1})$ for all $i = 1, \ldots, r-1$, we finally obtain the desired continuation of $D_r(s)$.

6. Abelian extensions

In this section, let G be a finite abelian group. Let G act on itself, we see that G is also a transitive permutation group. So S := S(G) is just the set of abelian G-fields. In this section we will prove some results on counting fields with specifications related to Theorem 3.11 and then on the distribution of the q-part class groups with $q \mid |G|$.

6.1. Main results. We make the following notations. The first one comes from Wood [21]. Definition 6.1. For any $g \in G$, let |g| denote its order. For any subset S of G, let

$$\beta(S) := \sum_{\substack{g \in S \\ g \neq 1}} \phi(|g|),$$

where ϕ is the Euler's phi function.

Theorem 6.2. Let Ω be a non-empty subset of G that is closed under invertible powering and let $\Omega^c =: \{g \in G \mid \langle g \rangle \cap \Omega = \emptyset\}$. Then define $\beta := \beta(\Omega^c)$. For any integer $r \ge 0$, we have

$$N_r(x) := N(\mathcal{S}; (\Omega, r); x) \ll \begin{cases} x(\log x)^{\beta-1}(\log \log x)^r & \text{if } \beta > 0, \\ \frac{x}{\log x}(\log \log x)^{r-1} & \text{else if } \beta = 0. \end{cases}$$

If r > |G|, then

$$N_r(x) \gg \begin{cases} x(\log x)^{\beta-1}(\log \log x)^{r-|G|} & \text{if } \beta > 0\\ x(\log x)^{\beta-1}(\log \log x)^{r-|G|-1} & \text{else if } \beta = 0. \end{cases}$$

From the theorem, we immediately obtain the following.

Corollary 6.3. When $q \mid |G|$, Conjecture 1.6(1) is true for $(\mathcal{S}, \Omega(q^{\infty}))$.

Proof. First of all, $\Omega(q^{\infty})$ is non-trivial, for $q \mid |G|$. Once the subset $\Omega = \Omega(q^{\infty})$ is fixed, the number $\beta = \beta(\Omega^c)$ is fixed. So, for each integer $r \geq 0$, let r' := r + |G|, we have

$$N_r(x) = o(N_{r'}(x))$$

as $x \to \infty$, which concludes the proof.

6.2. Generating series. Using Class Field Theory (see Lemma A.2), we know that Artin reciprocity maps $\chi : \mathbb{C}_{\mathbb{Q}} \to G$ admits a one-to-one correspondence to surjective homomorphisms $\rho : \prod_{p \nmid \infty} \mathbb{Z}_p^* \to G$ given by $\rho = \prod_{p \nmid \infty} \chi|_p$. So let's define the invariant of morphisms.

Definition 6.4. (1) Let p be a finite rational prime. If $\rho : \mathbb{Z}_p^* \to G$ is a local homomorphism, then define

$$P(\rho) := \begin{cases} p & \text{if } \rho(\mathbb{Z}_p^*) \neq \{1\}\\ 1 & \text{otherwise.} \end{cases}$$

For any lift $\tilde{\rho} : \mathbb{Q}_p^* \to G$, we just define $P(\tilde{\rho}) = P(\rho)$. (2) For any homomorphism $\rho : \prod_{p \nmid \infty} \mathbb{Z}_p^* \to G$, let

$$P(\rho) := \prod_{p \nmid \infty} P(\rho|_p).$$

If $\tilde{\rho} : \mathbb{C}_{\mathbb{Q}} \to G$ is any lift of ρ , then just let $P(\tilde{\rho}) := P(\rho)$.

Similarly, we can compute $\mathbf{1}_{(\Omega,r)}(\rho)$ for $\rho \in \operatorname{Hom}(\prod_{p \nmid \infty} \mathbb{Z}_p^*, G)$.

Definition 6.5. (1) Let p be a finite rational prime, If $\rho : \mathbb{Z}_p^* \to G$ is a local homomorphism, then define

$$\mathbf{1}_{\Omega}(\rho) = \begin{cases} 1 & \text{if } p \nmid |G| \\ & \text{and } \rho(\mathbb{Z}_p^*) \cap \Omega \neq \emptyset, \\ 0 & \text{otherwise.} \end{cases}$$

(2) If $\rho: \prod_{p \nmid \infty} \mathbb{Z}_p^* \to G$ is a homomorphism, then let

$$\mathbf{1}_{(\Omega,r)}(\rho) := \delta(\sum_{p \nmid \infty} \mathbf{1}_{\Omega}(\rho|_p) - r)$$

where $\delta(0) = 1$ and $\delta(x) = 0$ for all $x \neq 0$. If $\tilde{\rho} : \mathbb{C}_{\mathbb{Q}} \to G$ is a lift of ρ , then define $\mathbf{1}_{(\Omega,r)}(\tilde{\rho}) := \mathbf{1}_{(\Omega,r)}(\rho)$.

Let $\chi: \prod_{p \nmid \infty} \mathbb{Z}_p^* \to G$ be a fixed homomorphism. Let S_{χ} be the set of primes p such that $\chi|_p$ is non-trivial, then define

$$D_{r,\chi}(s) := \sum_{\substack{\rho: \prod_{p \nmid \infty} \mathbb{Z}_p^* \to G \\ \rho|_p = \chi|_p \ \forall p \in S_{\chi}}} \mathbf{1}_{(\Omega,r)}(\rho) (P(\rho))^{-s}.$$

If χ is trivial, then just denote it by $D_r(s)$. We first show the arithmetic meaning of $D_{\chi,r}(s)$. Lemma 6.6. If we write

$$D_r(s) = \sum_{n=1}^{\infty} a_n n^{-s},$$

then

$$\tilde{N}_r(x) := \sum_{n < x} a_n = \sum_{\substack{\rho: \prod_{p \nmid \infty} \mathbb{Z}_p^* \to G \\ P(\rho) < x}} \mathbf{1}_{(\Omega, r)}(\rho).$$

Proof. The proof follows from the definition directly. Namely

$$a_n = \sum_{\substack{\rho: \prod_{p \nmid \infty} \mathbb{Z}_p^* \to G \\ P(\rho) < n}} \mathbf{1}_{(\Omega, r)}(\rho).$$

Take the sum over all n < x and we get the conclusion.

Lemma 6.7. There exists at least one surjective $\chi : \prod_{p \nmid \infty} \to G$ such that $\mathbf{1}_{(\Omega,r_0)}(\chi) = 1$ with $r_0 \leq |G|$. Let r > |G|. If we write

$$D_{r,\chi}(s) = \sum_{n=1}^{\infty} a_n n^{-s}$$

then for all x > 0, we have

$$N_r(x) \ge \sum_{\substack{n < x \\ 24}} a_n.$$

Proof. For each $g \in G \setminus \{1\}$, choose a rational prime p_g such that $p_g \equiv 1 \mod |G|$. Define $\rho : \mathbb{Z}_{p_g}^* \to G$ to be the homomorphism determined by $\rho(\xi_{p_g}) = g$, where ξ_{p_g} is a generator of the roots of unity of \mathbb{Q}_{p_g} . Let $\chi := \prod_{g \in G} \rho$. It is clear that χ is a surjective. Since $|\Omega| < |G|$, we see that $r_0 \leq |G|$.

The inequality

$$N_r(x) \ge \sum_{n < x} a_n$$

follows from the fact that if $\rho : \prod \mathbb{Z}_p^* \to G$ is a homomorphism such that $\rho|_{p_g} = \chi|_{p_g}$, then ρ must be surjective.

Lemma 6.8. Let $\chi : \prod_{p \nmid \infty} \mathbb{Z}_p^* \to G$ be a homomorphism such that $\mathbf{1}_{(\Omega,r_0)}(\chi) = 1$. If the set S_{χ} contains the primes $p \mid |G|$, then we have

$$\begin{split} D_{r_0+r,\chi}(s) = & P(\chi)^{-s} \sum_{\substack{\underline{p} \in (\mathcal{P} \setminus S_\chi)^r \\ p_1 < \dots < p_r}} \prod_{i=1}^r (\sum_{\substack{\rho : \mathbb{Z}_{p_i}^* \to G}} \mathbf{1}_\Omega(\rho) P(\rho)^{-s}) \\ & \cdot \prod_{\substack{p \in \mathcal{P} \setminus S_\chi \\ p \nmid p_1 \dots p_r}} (\sum_{\substack{\rho : \mathbb{Z}_p^* \to G}} (1 - \mathbf{1}_\Omega(\rho)) P(\rho)^{-s}). \end{split}$$

Proof. Write $D_{r_0+r,\chi}(s) = \sum_{n=1}^{\infty} a_n n^{-s}$. Then

$$a_{n} = \sum_{\substack{\rho:\prod_{p\nmid\infty}\mathbb{Z}_{p}^{*}\to G \\ P(\rho)=n}} \sum_{\substack{P(\rho)=n}} \mathbf{1}_{\{\Omega,r_{0}+r\}}(\rho)$$
$$= \sum_{n/P(\chi)=p_{1}\cdots p_{r}q_{1}\cdots q_{l}} \prod_{i=1}^{r} (\sum_{\rho:\mathbb{Z}_{p_{i}}^{*}\to G} \mathbf{1}_{\Omega}(\rho)) \prod_{j=1}^{l} (\sum_{\rho:\mathbb{Z}_{q_{j}}^{*}\to G} (1-\mathbf{1}_{\Omega}(\rho)))$$

where the decomposition $n/P(\chi) = p_1 \cdots p_r q_1 \cdots q_l$ runs over all primes $p_i, q_i \in \mathcal{P} \setminus S_{\chi}$. \Box

Proposition 6.9. For each r = 0, 1, 2, ... and for each homomorphism $\chi : \prod_{p \nmid \infty} \mathbb{Z}_p^* \to G$ with $\mathbf{1}_{(\Omega,r_0)}(\chi) = 1$, the series $D_{r_0+r,\chi}(s)$ defines a holomorphic function in the open halfplane $\Re(s) > 1$. And there exists holomorphic functions f_0, \ldots, f_r in the closed half-plane $\Re(s) \geq 1$ such that $f_0(1)$ is non-zero and such that

$$D_{r_0+r,\chi}(s) = (s-1)^{\beta} \sum_{i=0}^r f_{r-i}(s) \left(\log \frac{1}{s-1}\right)^i.$$

Proof. Write $\tilde{S}_{\chi} := S_{\chi} \cup \{p \in \mathcal{P} : p \mid |G|\}$. By the identity

$$D_{r,\chi}(s) = \sum_{\substack{\chi: \prod_{p \in \tilde{S}_{\chi}} \mathbb{Z}_{p}^{*} \to G \\ \tilde{\chi}|_{p} = \chi|_{p} \ p \in S_{\chi} \\ 25}} D_{r,\tilde{\chi}}$$

we see that it suffices to prove the case when S_{χ} contains the set of primes $p \mid |G|$. Since $S_{\chi} \supseteq \{p \in \mathcal{P} : p \mid |G|\}$, we can write

$$D_{r_0+r,\chi}(s) = P(\chi)^{-s} \sum_{\substack{\underline{p} \in (\mathcal{P} \setminus S_{\chi})^r \\ p_1 < \dots < p_r}} \prod_{i=1}^r (\sum_{\substack{\rho : \mathbb{Z}_{p_i}^* \to G \\ p_i < \dots < p_r}} \mathbf{1}_{\Omega}(\rho) P(\rho)^{-s})$$

$$\cdot \prod_{\substack{p \in \mathcal{P} \setminus S_{\chi} \\ p \nmid p_1 \dots p_r}} (\sum_{\substack{\rho : \mathbb{Z}_p^* \to G \\ p \in \mathcal{P} \setminus S_{\chi}}} (1 - \mathbf{1}_{\Omega}(\rho)) P(\rho)^{-s})$$

$$= P(\chi)^{-s} \prod_{\substack{p \in \mathcal{P} \setminus S_{\chi} \\ p \in \mathcal{P} \setminus S_{\chi}}} (1 + a(p)p^{-s})$$

$$\cdot \sum_{\substack{\underline{p} \in (\mathcal{P} \setminus S_{\chi})^r \\ p_1 < \dots < p_r}} \prod_{i=1}^r b(p, s)p^{-s}$$

where $a(p) = \#\{\rho : \mathbb{Z}_p^* \to G \mid \operatorname{im}(\rho) \cap \Omega = \emptyset\}$, and

$$b(p,s) = \frac{\sum_{\rho:\mathbb{Z}_p^* \to G} \mathbf{1}_{\Omega}(\rho)}{1 + a(p)p^{-s}}.$$

Then a(p) is a function modulo |G|, and b(p, s) is a coefficient function with main term

$$b(p) := \#\{\rho : \mathbb{Z}_p^* \to G \mid \operatorname{im}(\rho) \cap \Omega \neq \emptyset\},\$$

which is also a function modulo |G|. It is clear that b(p) is non-zero for primes $p \equiv 1 \mod |G|$. So by Theorem 5.10 and 5.9, we see that the following continuation exists:

$$\prod_{p \in \mathcal{P} \setminus S_{\chi}} (1 + a(p)p^{-s})^{\phi(|G|)} = g(s)(s-1)^{\beta'},$$

and
$$\sum_{\substack{\underline{p} \in (\mathcal{P} \setminus S_{\chi})^r \\ p_1 < \dots < p_r}} \prod_{i=1}^r b(p,s)p^{-s} = \sum_{i=0}^r g_{r-i}(s) \left(\log \frac{1}{s-1}\right)^i,$$

where

$$\begin{split} \beta' &= \sum_{\alpha \in (\mathbb{Z}/|G|)^*} a(\alpha) \\ &= \sum_{\substack{g \in G \\ \langle g \rangle \cap \Omega = \emptyset}} \phi(|g|) \frac{\phi(|G|)}{\phi(|g|)} = \phi(|G|)\beta. \end{split}$$

Since g(s) is nowhere zero in $\Re(s) \ge 1$, we obtain the desired continuation for $D_{r,\chi}(s)$. \Box

6.3. **Proof of Theorem 6.2.** Let's just prove the theorem in this section. First let χ : $\prod_{p \nmid \infty} \mathbb{Z}_p^* \to G$ be the trivial homomorphism. According to Proposition 6.9, we see that for each $r = 0, 1, 2, \ldots$, the series $D_r(s)$ defines a holomorphic function in the open half-plane $\Re(s) > 1$ and there exists some holomorphic functions f_0, \ldots, f_r in the closed half-plane

 $\Re(s) \ge 1$ such that

$$D_r(s) = (s-1)^{\beta} \sum_{i=0}^r f_{r-i}(s) \left(\log \frac{1}{s-1}\right)^i.$$

Write $D_r(s) = \sum_{n=1}^{\infty} a_n n^{-s}$, and let $\tilde{N}_r(x) := \sum_{n < x} a_n$. By Tauberian Theorem 5.1, we obtain the following result:

$$\tilde{N}_r(x) \sim \begin{cases} \frac{f_0(1)}{\Gamma(\beta)} x (\log x)^{\beta-1} (\log \log x)^r & \text{if } \beta > 0\\ rg_0(1) \frac{x}{\log x} (\log \log x)^{r-1} & \text{if } \beta = 0 \text{ and } r > 1. \end{cases}$$

Since $\beta = 0$ if $\Omega = G \setminus \{1\}$. We see that the case when r = 0 simply means that we are counting homomorphisms $\operatorname{Hom}(\prod_{p||G|} \mathbb{Z}_p^*, G)$, which is a finite number. Then by Lemma 6.6, we see that

$$N_r(x) \le \tilde{N}_r(x),$$

which proves the estimate on the upper bound of $N_r(x)$. Now let r > |G|, and let $\chi : \prod_{p \nmid \infty} \mathbb{Z}_p^* \to G$ be any surjective homomorphism such that $\mathbf{1}_{(\Omega,r)}(\chi) = r_0 \leq |G|$. Write $D_{r,\chi}(s) = \sum_{n=1}^{\infty} b_n n^{-s}$, and let $N_{r,\chi}(x) := \sum_{n < x} b_n$. Then by Proposition 6.9 again, we know that $D_{r,\chi}(s)$ defines a holomorphic function in the open half-plane $\Re(s) > 1$ and there exists some holomorphic functions g_0, \ldots, g_{r-r_0} such that

$$D_{r,\chi}(s) = (s-1)^{\beta} \sum_{i=0}^{r-r_0} g_{r-r_0-i} \left(\log \frac{1}{s-1}\right)^i.$$

By similar method, we can obtain the estimate of the lower bound.

6.4. **Proof of Theorem 1.8.** In this section, we give the proof of Theorem 1.8. It suffices to show that the Conjecture 1.6(2) holds for (\mathcal{S}, Ω) where $\Omega = \Omega(\Gamma, q^{\infty})$. According to the condition that $q \mid |\Gamma/\Lambda|$, we see that Ω is non-empty. By Theorem 6.2, for each $r = 0, 1, 2, \ldots$ we have

$$N_r(x) \ll x(\log x)^{\beta-1}(\log \log x)^r.$$

On the other hand, according to Melanie [21, Theorem 3.1], we see that

$$N(x) \sim Cx(\log x)^{\beta(G)-1},$$

for some constant C > 0. This shows that

$$N_r(x) = o(N(x))$$

for all r = 0, 1, 2, ..., i.e., the conjecture holds. So the results on zero-probability and infinite moments are true, by Theorem 4.4 and 4.5.

Of course, one can also show that the Conjecture 1.6(1) is true in this case using the estimate given by Theorem 6.2.

7. DIHEDRAL EXTENSIONS

In this section, we first try to estimate dihedral extensions with local specifications, and then use examples to illustrate their applications towards distribution of class groups. We use the following notation to explain what we mean by dihedral extensions in this section.

Definition 7.1. Let G be a finite group. We call G a dihedral group if there exists a splitting short exact sequence

$$1 \to C_m \to G \to C_2 \to 1,$$

of groups, where C_n is the cyclic group of order n, and $m \ge 3$ is an integer, such that for the generator $g_2 \in C_2$ and for a generator $g_m \in C_m$ we have $g_2 g_m g_2^{-1} = g_m^{-1}$. In this case we just write $G = D_m$

For example, $G = D_q$ where q is an odd prime. Let $S := S(D_m)$. Then S contains fields number fields K_m such that $[K_m : \mathbb{Q}] = m$ with Galois closure K_{2m} that is a Galois D_m -field. Let K_2 be the fixed field of C_m .

7.1. Algebraic theory. Let K be any number field, and let \mathscr{O} be its ring of integers. For each rational prime p, let $\mathscr{O}_p^* := \prod_{\mathfrak{p}|p} \mathscr{O}_{\mathfrak{p}}^*$ where \mathfrak{p} are primes of K. Let ζ_p be the generator of $\mu(K_{\mathfrak{p}}^*)$ for some $\mathfrak{p} \mid p$.

Proposition 7.2. Let $K_2 = \mathbb{Q}(\sqrt{d})$ be a quadratic number field. Let $p \nmid 2md\infty$ be a rational prime. We have the following.

(1) If $p\mathcal{O}_{K_2} = \mathfrak{p}_1\mathfrak{p}_2$, then we have a bijective map

$$\operatorname{Hom}_{C_2}(\mathscr{O}_p^*, C_m) \to \{g \in C_m \mid p \equiv 1 \bmod |g|\}$$

given by $\rho \mapsto \rho(\zeta_p)$.

(2) If $p\mathcal{O}_{K_2} = \mathfrak{p}$, then we have a bijective map

 $\operatorname{Hom}_{C_2}(\mathscr{O}_p^*, C_m) \to \{g \in C_m \mid p \equiv -1 \bmod |g|\},\$

given by $\rho \mapsto \rho(\zeta_p)$.

Proof. If p splits in K_2/\mathbb{Q} , then a local C_2 -morphism $\rho : \mathscr{O}_p^* \to C_m$ reduces to a homomorphism $\rho : \mathscr{O}_{\mathfrak{p}_1} \to C_m$, i.e.,

$$\operatorname{Hom}_{C_2}(\mathscr{O}_p^*, C_m) = \{\operatorname{Ind}_1^{C_2} \rho \mid \rho : \mathscr{O}_{\mathfrak{p}_1}^* \to C_m\}$$

Because under an isometry $\mathscr{O}_{\mathfrak{p}_1} \cong \mathscr{O}_{\mathfrak{p}_2}$, we have

$$g_2 \cdot (\alpha_1, \alpha_2) = (\alpha_2, \alpha_1),$$

where g_2 is the generator of C_2 . A homomorphism $\rho : \mathscr{O}_{\mathfrak{p}_1}^* \to C_m$ is totally determined by the image of $\zeta_{\mathfrak{p}_1}$ in this case, which gives the desired result.

Else if p is inert in K_2/\mathbb{Q} , then a local C_2 -morphism $\rho : \mathscr{O}_p^* \to C_m$ is equivalent to some abelian extension over K_p . The Frobenius acts on the inertia $\rho(\zeta_p)$ in a non-trivial way because of the Galois action from K_p/\mathbb{Q} . Therefore if we let x be the Frobenius element, then we can write $x = hg_2$ with $h \in C_m$. The action of x on $\rho(\zeta_p)$ says that

$$x\rho(\zeta_p)x^{-1} = \rho(\zeta_p)^{-1}.$$

Therefore $p \equiv -1 \mod |g|$. Conversely, if g is an element of C_m such that $p \equiv -1 \mod |g|$, then we define ρ by setting $\rho(\zeta_p) = g$. Note that g_2 acts on the residue field as the Frobenius automorphism. The action of g_2 says that

$$g^{-1} = g_2 g g_2^{-1}$$

= $g_2 \rho(\zeta_{\mathfrak{p}}) g_2^{-1}$
= $\rho(g_2 \cdot \zeta_{\mathfrak{p}}) = \rho(\zeta_{\mathfrak{p}}^p) = g^p.$

And we are done.

7.2. Main results. We present our results as follows.

1

Theorem 7.3. Assume that the Cohen-Lenstra Heuristics holds for quadratic number fields, or to be precise, assume that

$$\sum_{\substack{K_2 = \mathbb{Q}(\sqrt{d}) \\ P(K_2) < x}} |\operatorname{Hom}(\operatorname{Cl}_{K_2}, C_m)| \ll x.$$

Let $\Omega = C_m \setminus \{1\}$ be the non-empty subset of D_m . Then for each r = 1, 2, ...

$$N_r(x) := \sum_{\substack{K_m \in \mathcal{S} \\ P(K_m) < x}} \mathbf{1}_{(\Omega, r)}(K_m) \ll x (\log \log x)^r.$$

7.3. Generating series. Let Ω be a non-empty subset of C_m that is closed under invertible powering. It is also closed under conjugation in D_m . First, we need a notation.

Definition 7.4. Let K_2 be a quadratic number field. Let $p \nmid \infty$ be a rational prime, and let $\rho : \mathscr{O}_p^* \to C_m$ be a homomorphism.

(1) Define

$$P(\rho) := \begin{cases} p & \text{if } \operatorname{im}(\rho) \neq \{1\};\\ 1 & \text{otherwise.} \end{cases}$$

(2) Define

$$\mathbf{1}_{\Omega}(\rho) = \begin{cases} 1 & \text{if } p \nmid 2mP(K_2) \text{ and } \rho(\zeta_p) \cap \Omega \neq \emptyset \\ 0 & \text{otherwise.} \end{cases}$$

For each $\chi : \prod_{p \nmid \infty} \mathscr{O}_p^* \to C_m$, (or $\chi : \mathcal{C}_{K_2} \to C_m$), define

$$P(\chi) := \prod_{p \nmid \infty} P(\chi|_p).$$

And for each $r = 0, 1, 2, \ldots$, define

$$\mathbf{1}_{(\Omega,r)}(\rho) = \delta(r - \sum_{p} \mathbf{1}_{\Omega}(\rho)),$$

where $\delta(x) = 1$ when x = 0 and $\delta(x) = 0$ elsewhere.

Let's construct a generating series as follows. For each quadratic number field K_2 , and for each $r = 0, 1, 2, \ldots$, let

$$D_{K_2,r}(s) := |\operatorname{Hom}(\operatorname{Cl}_{K_2}, C_3)| P(K_2)^{-s} \sum_{\rho: \prod_{p \nmid \infty} \mathscr{O}_p^* \to C_m} \mathbf{1}_{(\Omega,r)}(\rho) (P(\rho))^{-s},$$

where ρ only runs over C_2 -morphisms.

Lemma 7.5. Let $K_2 = \mathbb{Q}(\sqrt{d})$ be a quadratic number field. If we write

$$D_{K_2,r}(s) = \sum_{n=1}^{\infty} a_n n^{-s},$$

and let $\tilde{N}_{d,r}(x) := \sum_{n < x} a_n$, then

$$\sum_{\substack{K_m \in \mathcal{S} \\ P(K_m) < x}} \sum_{K_2 \subseteq K_{2m}} \mathbf{1}_{(\Omega, r)}(K_m) \le \tilde{N}_{d, r}(x)$$

for all x > 0.

Proof. According to Proposition A.7, we know that a D_m -field K_{2m} over K_2 corresponds to a surjective C_2 -morphisms $C_{K_2} \to C_m$. Conversely, Lemma A.6 says that a surjective C_2 -morphism $C_{K_2} \to C_m$ corresponds to a D_m -field K_{2m} containing K_2 . Finally, Proposition A.5 says that for a fixed surjective C_2 -morphism $\rho : \prod_p \mathcal{O}_p^* \to C_m$, there are at most $|\text{Hom}(\text{Cl}_{K_2}, C_m)|$ many lift to $C_{K_2} \to C_m$. And by the computation:

$$a_n = |\operatorname{Hom}(\operatorname{Cl}_{K_2}, C_m)| \sum_{\substack{\rho: \prod_p \mathcal{O}_p^* \to C_m \\ P(\rho) = n}} \mathbf{1}_{(\Omega, r)}(\rho)$$

we see that the statement is true.

In addition to Definition 7.4, we need a condition to help us count dihedral extensions. Recall that for a rational number a that is coprime to an odd prime p, we have the Legendre symbol $\left(\frac{a}{p}\right)$: its value is 1 if $x^2 \equiv a \mod p$ admits a solution and -1 otherwise. In particular, we have the following:

Proposition 7.6. [17, I.8.5] For squarefree d and gcd(p, 2d) = 1, we have the equivalence

$$\left(\frac{d}{p}\right) = 1 \iff p \text{ is totally split in } \mathbb{Q}(\sqrt{d}).$$

By Quadratic Reciprocity Law, we know that the ramification of p in $\mathbb{Q}(\sqrt{d})$ is totally determined by p modulo 4d. Of course, this is not always the case. One can consider $\mathbb{Q}(\sqrt{-3})$ for example.

Definition 7.7. For each square-free number d, and for each $g \in C_m$, let

$$d(g) := \lim_{x \to \infty} \frac{\#\{p < x \mid (\frac{d}{p}) = i \text{ and } p \equiv i \mod |g|, i = \pm 1\}}{\#\{p < x\}}.$$

For each subset $S \subseteq C_m$, define

$$\beta_d(S) := \sum_{g \in S} d(g).$$

When gcd(m, 4d) = 1, by Chinese Remainder Theorem, we know that

$$\mathbb{Z}/(4dm) \cong \mathbb{Z}/4d \times \mathbb{Z}/m$$

so, $d(g) = \phi^{-1}(|g|)$ for all $g \in C_m$. However, this is not always the case. For example, consider m = 3, and the cubic fields K_3 associated to $K_2 = \mathbb{Q}(\sqrt{-3})$. One can show that

d(g) = 1 for $g \in C_3 \setminus \{1\}$. Loosely speaking, the number of S_3 -fields above $\mathbb{Q}(\sqrt{-3})$ is "much more" than other quadratic fields. So, in general, we exclude nontrivial roots of unity when counting fields. In this section, the issue caused by roots of unity does not affect the results because of the conditions we will set up, hence we do not emphasize this issue.

Proposition 7.8. The series $D_{K_2,r}(s)$ defines a holomorphic function in the open half-plane $\Re(s) > 1$. Moreover, there exists holomorphic functions f_0, \ldots, f_r in the closed half-plane such that $f_0(1) \neq 0$ and such that

$$D_{K_{2},r}(s) = (s-1)^{-\beta} \sum_{i=0}^{r} f_{r-i}(s) \left(\log \frac{1}{s-1}\right)^{i},$$

where

$$\beta = \sum_{\substack{g \in C_m \setminus \{1\} \\ \langle g \rangle \cap \Omega = \emptyset}} d(g).$$

Proof. Since there are only finitely many prime $p \mid 2mP(K_2)$, we see that if

$$\tilde{D}_{K_2,r}(s) := \sum_{\rho:\prod_{p \nmid 2mP(K_2)} \mathscr{O}_p^* \to C_m} \mathbf{1}_{(\Omega,r)}(\rho)(P(\rho))^{-s}$$

admits similar analytic properties, then so is $D_{K_2,r}(s)$. Because any homomorphism ρ : $\prod_{p \nmid \infty} \mathscr{O}_p^* \to C_m$ could be written as the product of $\rho|_{p|2mP(K_2)}$ and $\rho|_{p \nmid 2mP(K_2)}$. We can rewrite $\tilde{D}_{K_2,r}(s)$ as follows:

$$\tilde{D}_{K_2,r}(s) = \sum_{\substack{p \in \mathcal{P}^r \\ p_1 < \dots < p_r}} \prod_{i=1}^r \left(\sum_{\rho: \mathscr{O}_p^* \to C_m} \mathbf{1}_{\Omega}(\rho) P(\rho)^{-s} \right)$$
$$\cdot \prod_{p \nmid 2m P(K_2) p_1 \dots p_r} \left(\sum_{\rho: \mathscr{O}_p^* \to C_m} (1 - \mathbf{1}_{\Omega}(\rho)) P(\rho)^{-s} \right),$$

where all the homomorphisms are compatible with C_2 -action. This is just the decomposition of a C_2 -morphism $\rho : \prod_{p \nmid 2mP(K_2)} \mathscr{O}_p^* \to C_m$ with local factors. By comparing with a large enough power of Riemann zeta function, we know that it is a holomorphic function in the open half-plane $\Re(s) > 1$. Recall that for all $g \in C_m$, and for almost all K_2 , either |g| = 2or $\gcd(|g|, 4\operatorname{Disc}(K_2)) = 1$. Then write $K_2 = \mathbb{Q}(\sqrt{d})$ and we have

$$\operatorname{Hom}_{C_2}(\mathscr{O}_p^*, C_m) = \begin{cases} \{g \in C_m \mid p \equiv 1 \mod |g|\} & \text{if } \left(\frac{d}{p}\right) = 1; \\ \{g \in C_m \mid p \equiv -1 \mod |g|\} & \text{if } \left(\frac{d}{p}\right) = -1. \end{cases}$$

We see that $\operatorname{Hom}_{C_2}(\mathscr{O}_p^*, C_m)$ is dependent only on p modulo 4dm. Then we have

$$\tilde{D}_{K_{2,r}}(s) = \prod_{\substack{p \nmid 2mP(K_{2})}} (1 + a(p)p^{-s})$$
$$\cdot \sum_{\substack{\underline{p} \in \mathcal{P}^{r} \\ p_{1} < \dots < p_{r} \\ 31}} \prod_{i=1}^{r} b(p_{i}, s)p_{i}^{-s},$$

where a(p) is a function modulo 4md and b(p, s) is a coefficient function with main term b(p) which is also a function modulo 4md. The continuation of $\tilde{D}_{K_2,r}(s)$ is then given by Theorem 5.10 and Theorem 5.9. In other words, there exists holomorphic functions g(s) and h_0, \ldots, h_r in the closed half-plane $\Re(s) \ge 1$ such that g(s) nowhere zero in $\Re(s) \ge 1$ and $h_0(1) \ne 0$, and such that

$$\prod_{\substack{p \nmid 2mP(K_2)}} (1 + a(p)p^{-s})^{\phi(4dm)} = g(s)(s-1)^{\beta'}$$

and
$$\sum_{\substack{p \in \mathcal{P}^r \\ p_1 < \dots < p_r}} \prod_{i=1}^r b(p_i, s)p_i^{-s} = \sum_{i=0}^r h_{r-i}(s) \left(\log \frac{1}{s-1}\right)^i,$$

where β' is given by

$$\sum_{\alpha \in (\mathbb{Z}/4dm)^*} a(\alpha) = \sum_{\substack{\alpha \in (\mathbb{Z}/4dm)^* \\ p \equiv \alpha \mod 4dm}} \#\{1 \neq \rho \in \operatorname{Hom}_{C_2}(\mathscr{O}_p^*, C_m) \mid \operatorname{im}(\rho) \cap \Omega = \emptyset\}$$
$$= \sum_{\substack{g \in C_m \setminus \{1\} \\ \langle g \rangle \cap \Omega = \emptyset}} d(g)\phi(4dm) = \phi(4dm)\beta.$$

So we obtain the desired continuation for $D_{K_2,r}(s)$.

7.4. **Proof of Theorem.** For a fixed quadratic number field K_2 , since $\Omega = C_m \setminus \{1\}$, we can consider the following series

$$D_{K_2,r}(s) := |\operatorname{Hom}(\operatorname{Cl}_{K_2}, C_3)| P(K_2)^{-s} \Big(\sum_{\substack{\rho: \prod_{p \mid 2m} \mathcal{O}_p^* \to C_m \\ p_1 < \dots < p_r}} P(\rho)^{-s} \Big)$$
$$\cdot \sum_{\substack{p \in \mathcal{P}^r \\ p_1 < \dots < p_r}} \prod_{i=1}^r \Big(\sum_{\substack{\rho: \mathcal{O}_{p_i}^* \to C_m \\ \rho_i \to C_m}} P(\rho)^{-s} \Big),$$

where ρ is taken to be *any* homomorphisms. It is clear that if we write $D_r(s) = \sum_{n=1}^{\infty} a_n n^{-s}$, and $D_{K_2,r}(s) = \sum_{n=1}^{\infty} b_n n^{-s}$, then $b_n \leq a_n$ for all $n = 1, 2, \ldots$ Since $|\text{Hom}(\mathscr{O}_p^*, C_m)|$ is bounded above by a number c for any rational prime p and for any quadratic number field K_2 , we can even consider the series

$$\tilde{D}_r(s) = \sum_{\substack{\underline{p} \in \mathcal{P}^r \\ p_1 < \dots < p_r}} \prod_{i=1}^r cp_i^s$$

Then we see that it is a holomorphic function in the open half-plane $\Re(s) > 1$, and there exists some holomorphic functions g_0, \ldots, g_r in the closed half-plane $\Re(s) \ge 1$ such that $g_0(1) \ne 0$ and such that

$$\tilde{D}_r(s) = \sum_{i=0}^r g_{r-i}(s) \left(\log \frac{1}{s-1} \right)^i.$$
32

Write $\tilde{D}_r(s) = \sum_{n=1}^{\infty} \tilde{a}_n n^{-s}$ and let $\tilde{A}_r(x) := \sum_{n < x} \tilde{a}_n$. Then Tauberian Theorem 5.1 implies the following: when $r \ge 1$, we have

$$\sum_{n < x} a_n \le \tilde{A}_r(x) \sim rg(1) \frac{x}{\log x} (\log \log x)^{r-1}.$$

Now let

$$D_r(s) := \prod_{p|2m} (1 + cp^{-s}) \sum_{K_2 = \mathbb{Q}(\sqrt{d})} |\operatorname{Hom}(\operatorname{Cl}_{K_2}, C_m)| P(K_2)^{-s} \tilde{D}_r^{-s}$$

Then by Lemma 7.5, we see that if we write $D_r = \sum_{n=1}^{\infty} c_n n^{-s}$ and let $\tilde{N}_r(x) = \sum_{n < x} c_n$, then

$$N_r(x) \le N_r(x)$$

for all x > 0. Write $H(x) := \sum_{d < x} |\text{Hom}(\text{Cl}_{K_2}, C_m)|$. By Riemann-Stieltjes integral, we have the following.

$$N_{r}(x) \leq \int_{1^{-}}^{x} H(x/t) \,\mathrm{d}\,\tilde{A}_{r}(t)$$

$$\ll \int_{1^{-}}^{x} \frac{x}{t} \,\mathrm{d}\,\tilde{A}_{r}(t)$$

$$= \frac{x}{t} \tilde{A}_{r}(t)|_{1}^{x} + x \int_{1^{-}}^{x} \frac{\tilde{A}_{r}(t)}{t} \,\mathrm{d}\,t$$

$$\ll \frac{x}{\log x} (\log\log x)^{r-1} + \int_{e}^{x} \frac{(\log\log t)^{r-1}}{t\log t} \,\mathrm{d}\,t$$

$$= \frac{x}{\log x} (\log\log x)^{r-1} + \frac{1}{r} x (\log\log x)^{r}.$$

This shows that $N_r(x) \ll x(\log \log x)^r$ as required.

7.5. Applications.

Proposition 7.9. Let q be an odd prime, and let $\Omega := \Omega(D_q, q)$, which is exactly $(C_q \setminus \{1\}) \subseteq D_q$. If the Cohen-Lenstra Heuristics hold for quadratic number fields, or to be precise, for x > 1, we have

$$\sum_{\substack{K_2 = \mathbb{Q}(\sqrt{d}) \\ P(K_2) < x}} |\operatorname{Hom}(\operatorname{Cl}_{K_2}, C_q)| \ll x,$$

and the Malle-Bhargava Heuristics hold for D_q -fields, or to be precise, for x > 1, we have

$$N(x) = \sum_{\substack{K_q \in S \\ P(K_q) < x}} \gg x \log x,$$

then, we have

$$N_r(x) = o(N(x))$$

for all r = 0, 1, ..., Also, for all <math>r = 0, 1, 2, ..., we have

$$\mathbb{P}(\operatorname{rk}_q \operatorname{Cl}_K \leq r) = 0 \quad and \quad \mathbb{E}(|\operatorname{Hom}(\operatorname{Cl}_K, C_q)|) = +\infty,$$

where K runs over all fields in S for the product of ramified primes in K/\mathbb{Q} .

Proof. Theorem 7.3 shows that

$$N_r(x) = o(N(x))$$

for all $r = 1, 2, \ldots$ The case when r = 0 follows from Cohen-Lenstra Heuristics itself. To be precise, if K_{2q}/K_2 is an abelian C_q -extension that is unramified, then it is a subextension of the Hilbert class field of K_2 . So K_{2q}/K_2 corresponds to a homomorphism $\operatorname{Hom}(\operatorname{Cl}_{K_2}, C_q)$. Conversely, the set of such homomorphisms is embedded into the set of Artin reciprocity maps. In other words, any abelian C_q -extension that is a subextension of the Hilbert class field could be lifted to a homomorphism $\operatorname{Cl}_{K_2} \to C_q$. On the other hand, there are only finitely many primes that divides $|D_q|$, namely 2 and q, so there exists a constant C such that

$$\#\{\rho \in \operatorname{Hom}(J_{K_2}^{\infty}K_2^*/K_2^*, C_q) \mid \rho|_p = 1 \text{ for all } p \nmid 2q\} \le C$$

for all K_2 . By (A.1), we see that

$$N_0(x) \ll \sum_{\substack{K_2 = \mathbb{Q}(\sqrt{d}) \\ P(K_2) < x}} |\operatorname{Hom}(\operatorname{Cl}_{K_2}, C_q)|.$$

Then the two heuristics in the conditions show that $N_0(x) = o(N(x))$ as required. We see that actually the Conjecture 1.6(2) holds under the hypothesis of the statement. And the rest just follows from Theorem 4.4 and Theorem 4.5.

8. D_4 EXTENSIONS

In this section, let $D_4 = \langle \sigma, \tau \mid \sigma^4 = 1 = \tau^2, \tau^{-1}\sigma\tau = \sigma^3 \rangle$ be the dihedral group of order 8. Define $\mathcal{S} := \mathcal{S}(D_4)$ with D_4 being a subgroup of S_4 , i.e., we are focused on the quartic extensions whose Galois closure are D_4 -fields (see also Definition 2.1 and 2.2). According to the Definition 1.3 of non-random primes, we know by checking the cycles in $D_4 \to S_4$ that the prime 2 is the only non-random prime for the permutation group D_4 .

8.1. Main results. We first introduce the definition of the *conductor* for a quadratic extension of a quadratic number field, which will be used here as the invariant of the number fields.

Definition 8.1 (Conductor). If K is a quadratic number field and L is a quadratic extension of K, define the *conductor* of the pair (L, K) as

$$C(L,K) := \frac{\operatorname{Disc}(L)}{\operatorname{Disc}(K)}.$$

If L is a D_4 -field and K denotes its (unique) quadratic subfield, then C(L, K) = C(L) (the conductor of L).

Note that the notation given by the above definition agrees with the Artin conductor for the irreducible 2-dimensional representation of D_4 , if the quartic field has D_4 -Galois closure. See also [1, Section 2.3] for details. We here follow this definition for the convenience of both computation and generalization to other quartic fields. Note that L admits a unique quadratic subfield K. Let q = 2, which is the only non-random prime for D_4 . Define the following notations:

$$N(x) := N(\mathcal{S}(C); x) = \sum_{\substack{L \in \mathcal{S} \\ C(L) < x}} 1$$
$$H(x) := N(\mathcal{S}(C); |\text{Hom}(\text{Cl}_L, C_2)|; x) = \sum_{\substack{L \in \mathcal{S} \\ C(L) < x}} |\text{Hom}(\text{Cl}_L, C_2)|$$

In this section we are going to prove the following main result.

Theorem 8.2. For each $L \in S$, let K be its unique quadratic subfield. We have

$$\mathbb{E}_C(|\mathrm{Hom}(\mathrm{Cl}_L, C_2)|) = \lim_{x \to \infty} \frac{H(x)}{N(x)} = +\infty,$$

where L runs over all fields in S ordered by the conductor C.

8.2. Preliminary tools. We want to study the statistical behaviour of $\operatorname{Cl}_L[2^{\infty}]$ where $L \in \mathcal{S}$ for the conductor. Let's first prove a lemma whose statement is similar to [1, Lemma 5.1]. Recall that $\omega(n)$ means the number of distinct prime factors dividing n.

Lemma 8.3. For any $0 < \epsilon < \frac{1}{2}$, we have

$$\sum_{\substack{0 < D < x \\ D \ squarefree}} \frac{2^{\omega(D)}}{D} \cdot \left(\sum_{m=1}^{\infty} \sum_{\substack{n=1 \\ mn \neq \Box}}^{D^{\frac{1}{2}+\epsilon}} \frac{\mu(m)}{m^2 n} \left(\frac{D}{mn} \right) \right) = o(x),$$

where $(\frac{\cdot}{\cdot})$ means Legendre symbol here.

Proof. The proof itself is straightforward. First of all, the Riemann zeta function admits a meromorphic continuation to the complex plane with a pole of order 1 at s = 1. Similarly, the Dirichlet series

$$\prod_{p \in \mathcal{P}} (1 + 2p^{-s})$$

admits a meromorphic continuation to the closed half-plane $\Re(s) \ge 1$ with a pole of order 2 at s = 1. Then, according to [15, Theorem 5.11], we have

$$\sum_{n < x} \frac{1}{n} \sim c_1 \log x \quad \text{and} \quad \sum_{\substack{0 < D < x \\ D \text{ squarefree}}} \frac{2^{\omega(D)}}{D} \sim c_2 \log^2 x$$

for some constant $c_1, c_2 > 0$. So, we know that

$$\Big|\sum_{\substack{D \text{ squarefree}\\D \text{ squarefree}}} \frac{2^{\omega(D)}}{D} \cdot \Big(\sum_{m=1}^{\infty} \sum_{\substack{n=1\\mn\neq\square}}^{D^{\frac{1}{2}+\epsilon}} \frac{\mu(m)}{m^2 n} \Big(\frac{D}{mn}\Big)\Big)$$
$$\leq \sum_{\substack{0 < D < x\\D \text{ squarefree}}} \frac{2^{\omega(D)}}{D} \cdot \Big(\sum_{m=1}^{\infty} \sum_{n=1}^{x} \frac{1}{m^2 n}\Big)$$
$$\ll (\log x)^3 = o(x), \text{ as } x \to \infty.$$

Using the above lemma, we can prove the following proposition, which is similar to [1, Proposition 5.2].

Proposition 8.4. We have:

(8.1)

$$\sum_{\substack{[K:\mathbb{Q}]=2\\0<\operatorname{Disc}(K)< x}} \frac{L(1, K/\mathbb{Q})}{L(2, K/\mathbb{Q})} \cdot \frac{2^{\omega(\operatorname{Disc}(K))}}{\operatorname{Disc}(K)}}{\operatorname{Disc}(K)}$$

$$= \sum_{\substack{[K:\mathbb{Q}]=2\\0<\operatorname{Disc}(K)< x}} \frac{2^{\omega(\operatorname{Disc}(K))}}{\operatorname{Disc}(K)} \cdot \sum_{\substack{0

$$and \sum_{\substack{[K:\mathbb{Q}]=2\\-x<\operatorname{Disc}(K)<0}} \frac{L(1, K/\mathbb{Q})}{L(2, K/\mathbb{Q})} \cdot \frac{2^{\omega(\operatorname{Disc}(K))}}{\operatorname{Disc}(K)}}{\operatorname{Disc}(K)}$$

$$= \sum_{\substack{[K:\mathbb{Q}]=2\\-x<\operatorname{Disc}(K)<0}} \frac{2^{\omega(\operatorname{Disc}(K))}}{\operatorname{Disc}(K)} \cdot \sum_{\substack{0$$$$

where $L(s, K/\mathbb{Q}) = \sum_{n} \frac{\chi_{K}(n)}{n^{s}}$ and χ_{K} is the quadratic character associated to K. Proof. The proof is similar to that of [1, Proposition 5.2]. According to [1, (17)], we have

$$\frac{L(1, K/\mathbb{Q})}{L(2, K/\mathbb{Q})} = \frac{1}{L(2, K/\mathbb{Q})} \cdot \sum_{n=1}^{|\operatorname{Disc}(K)|^{\frac{1}{2}+\epsilon}} \frac{\chi_K(n)}{n} + O_\epsilon \left(\frac{\log(|\operatorname{Disc}(K)|)}{|\operatorname{Disc}(K)|^{\epsilon}}\right).$$

Let's rewrite the left-hand sides of (8.1) as

(8.2)
$$\frac{\sum_{\substack{[K:\mathbb{Q}]=2\\0<\operatorname{Disc}(K)$$

and similar expression for imaginary quadratic fields. Note that

$$\sum_{\substack{[K:\mathbb{Q}]=2\\0<|\operatorname{Disc}(K)|< x}} \frac{2^{\omega(\operatorname{Disc}(K))}}{\operatorname{Disc}(K)} \frac{\log(|\operatorname{Disc}(K)|)}{|\operatorname{Disc}(K)|^{\epsilon}} = O_{\epsilon}(1),$$

we focus on the remaining terms. Then by [1, (20)], we know that

(8.3)
$$\frac{1}{L(2, K/\mathbb{Q})} \sum_{n=1}^{|\operatorname{Disc}(K)|^{\frac{1}{2}+\epsilon}} \frac{\chi_K(n)}{n}$$
$$= \sum_{\substack{0 < a, b < \infty \\ (\operatorname{Disc}(K), ab) = 1}} \frac{\mu(a)}{a^3 b^2} + \sum_{n=1}^{|\operatorname{Disc}(K)|^{\frac{1}{2}+\epsilon}} \frac{\chi_K(n)}{n} \sum_{\substack{m=1 \\ mn \neq \square}} \frac{\mu(m)\chi_K(m)}{m^2}$$

By substituting (8.3) back to (8.2), we have

(8.4)
$$\sum_{\substack{[K:\mathbb{Q}]=2\\0<\operatorname{Disc}(K)< x}} \frac{2^{\omega(\operatorname{Disc}(K))}}{\operatorname{Disc}(K)} \left(\sum_{\substack{0< a, b<\infty\\(\operatorname{Disc}(K), ab)=1}} \frac{\mu(a)}{a^3b^2} + \sum_{n=1}^{|\operatorname{Disc}(K)|^{\frac{1}{2}+\epsilon}} \frac{\chi_K(n)}{n} \sum_{\substack{m=1\\mn\neq\square}} \frac{\mu(m)\chi_K(m)}{m^2}\right),$$

and similar expression for imaginary quadratic fields. By Lemma 8.3, we know that

$$\sum_{\substack{[K:\mathbb{Q}]=2\\0<\operatorname{Disc}(K)< x}} \frac{2^{\omega(\operatorname{Disc}(K))}}{\operatorname{Disc}(K)} \sum_{n=1}^{|\operatorname{Disc}(K)|^{\frac{1}{2}+\epsilon}} \frac{\chi_K(n)}{n} \sum_{\substack{m=1\\mn\neq\square}} \frac{\mu(m)\chi_K(m)}{m^2} = o(x).$$

Similarly for the case when Disc(K) < 0. And we are done for the proof.

8.3. **Proof of Theorem 8.2.** Recall that for any function $f : S \to \mathbb{R}$, we have defined the notation

$$N_f(x) := N(\mathcal{S}(C); f; x) = \sum_{L \in \mathcal{S}, C(L) < x} f(L),$$

where C means the conductor of $L \in S$ (see also Definition 2.3). In particular, let N(x) := N(S(C); x). In addition, here we can even generalize the usual counting fields notation as follows.

Definition 8.5. For any function $f : S \to \mathbb{R}$, define

$$N_f(x, y) := \sum_{n < x} \sum_{\substack{L \in \mathcal{S} \\ C(L) = n, \text{Disc}(K) < y}} f(L).$$

In other words we put some restrictions on the discriminant of K/\mathbb{Q} by this notation. See [1, Theorem 4.3] for its usage in counting D_4 -fields.

Proof. The basic idea of the proof is similar to that of [1, Theorem 2]. First of all, by Theorem 3.11, it suffices to prove that

$$\mathbb{E}_C(f) = \lim_{x \to \infty} \frac{N_f(x)}{N(x)} = +\infty$$

where $f(L) = 2^{\omega(\text{Disc}(K))}$ for each $L \in S$, because $|\text{Hom}(\text{Cl}_L, C_2)| \ge 2^{-6} \cdot f(L)$. Second, let $\chi_{[0,1]}$ is the characteristic function of [0, 1]. Then we can write

$$N_f(x, x^{\beta}) = \sum_{\substack{[K:\mathbb{Q}]=2\\|\operatorname{Disc}(K)| < x^{\beta}}} \sum_{\substack{[L:\mathbb{Q}]=2\\L \in \mathcal{S}}} f(L)\chi_{[0,1]}\left(\frac{|\operatorname{Disc}(K)\operatorname{Nm}_{K/\mathbb{Q}}(\operatorname{Disc}(L/K))|}{x}\right),$$

where $0 < \beta < 1$. For $\epsilon > 0$, there exists some compactly supported smooth functions $\varphi^{\pm} : \mathbb{R}_{\geq 0} \to \mathbb{R}_{\geq 0}$ such that $\varphi^{\pm} - \chi_{[0,1]}$ takes value in \mathbb{R}^{\pm} and that $\operatorname{Vol}(\varphi^{\pm}) = 1 \pm \epsilon$. Then [1, Lemma 4.5 and (13)] implies that

$$\sum_{\substack{[K:\mathbb{Q}]=2\\|\mathrm{Disc}(K)|< x^{\beta}}} \sum_{\substack{[L:\mathbb{Q}]=2\\L\in\mathcal{S}}} f(L)\varphi^{\pm} \Big(\frac{|\mathrm{Disc}(K)\operatorname{Nm}_{K/\mathbb{Q}}(\mathrm{Disc}(L/K))|}{x}\Big)$$
$$= \sum_{\substack{[K:\mathbb{Q}]=2\\|\mathrm{Disc}(K)|< x^{\beta}}} \frac{1\pm\epsilon}{\zeta(2)} \cdot \frac{L(1,K/\mathbb{Q})}{L(2,K/\mathbb{Q})} \cdot \frac{2^{-r_2(K)}}{|\mathrm{Disc}(K)|} f(L)x$$
$$+ O_{\epsilon} \Big(\sum_{\substack{[K:\mathbb{Q}]=2\\|\mathrm{Disc}(K)|< x^{\beta}}} |\mathrm{Disc}(K)|^{-\frac{1}{4}+\epsilon} x^{\frac{1}{2}+\epsilon}\Big),$$

where $r_2(K)$ means the number of complex embeddings of K. Here φ^{\pm} gives upper bound, resp. lower bound, of $N_f(x, x^{\beta})$. Note that the error term is bounded by $O_{\epsilon}(x^{1/2+3\beta/4+\epsilon})$. So, when $\beta < 2/3$, the error term is bounded by o(x). By letting ϵ goes to 0, we are reduced to proving that

$$\sum_{\substack{[K:\mathbb{Q}]=2\\|\operatorname{Disc}(K)|< y}} \frac{L(1,K/\mathbb{Q})}{L(2,K/\mathbb{Q})} \frac{f(L)}{|\operatorname{Disc}(K)|} \sim c(\log y)^2.$$

Now Proposition 8.4 says that it suffices to prove that

(8.5)

$$\sum_{\substack{[K:\mathbb{Q}]=2\\0<\operatorname{Disc}(K)< y}} \frac{L(1,K/\mathbb{Q})}{L(2,K/\mathbb{Q})} \cdot \frac{f(L)}{\operatorname{Disc}(K)}$$

$$= \sum_{\substack{[K:\mathbb{Q}]=2\\0<\operatorname{Disc}(K)< y}} \frac{f(L)}{\operatorname{Disc}(K)} \cdot \sum_{\substack{0

$$\operatorname{and} \sum_{\substack{[K:\mathbb{Q}]=2\\-y<\operatorname{Disc}(K)<0}} \frac{L(1,K/\mathbb{Q})}{L(2,K/\mathbb{Q})} \cdot \frac{f(L)}{\operatorname{Disc}(K)}$$

$$= \sum_{\substack{[K:\mathbb{Q}]=2\\0<\operatorname{Disc}(K)< y}} \frac{f(L)}{\operatorname{Disc}(K)} \cdot \sum_{\substack{0$$$$

Using [1, (21)],

$$\sum_{\substack{0 < a, b < \infty \\ (\text{Disc}(K), ab) = 1}} \frac{\mu(a)}{a^3 b^2} = \frac{\zeta(2)}{\zeta(3)} \prod_{p \mid \text{Disc}(K)} \frac{1 - \frac{1}{p^2}}{1 - \frac{1}{p^3}}$$

we get

$$\lim_{y \to \infty} \frac{\zeta(2)}{\zeta(3) \log(y)^2} \sum_{\substack{1 < D < y \\ D \text{ squarefree}}} \frac{2^{\omega(D)}}{D} \prod_{p \mid D} \frac{1 - \frac{1}{p^2}}{1 - \frac{1}{p^3}} = \zeta(2) \prod_{p \in \mathcal{P}} (1 - \frac{3}{p^2} - \frac{1}{p^3} + \frac{6}{p^4} - \frac{3}{p^5}).$$

We can do the similar computation when the squarefree D takes $D \equiv \pm 1 \mod 4$ and so on, i.e., discriminant of quadratic number fields belongs to different conjugacy classes (see [1, (23)] for example). Similar result holds when Disc(K) < 0. This proves (8.5), hence the theorem.

8.4. Further discussion with heuristics. In this section, let S := S(P) be the set of quartic fields ordered by product of ramified primes. The Malle-Bhargava Heuristics predicts that

(8.6)
$$N(x) \gg x(\log x)^3.$$

On the other hand, Alex Smith [19] shows that the Gerth's conjecture is true for quadratic number fields. So, in this case we have

(8.7)
$$\sum_{\substack{[K:\mathbb{Q}]=2\\|\operatorname{Disc}(K)|< x}} |\operatorname{Sur}(\operatorname{Cl}_K, C_4)| \sim cx,$$

for some constant c > 0. This implies that

$$\sum_{\substack{[K:\mathbb{Q}]=2\\\text{Disc}(K)|< x}} |\text{Hom}(\text{Cl}_K, C_4)| \ll x \log x,$$

i.e., it is determined by the 2-rank of Cl_K , which is given by the genus theory. According to Theorem 3.15, we know that $\operatorname{rk}_2 2 \operatorname{Cl}_L$ admits a lower bound given by the number of primes p such that $e_L(p) \equiv 0 \mod 4$. So let $\Omega := \{\sigma, \sigma^3\}$, and let's count L with r (tamely ramified) primes such that its inertia generator lies in Ω . Define

$$N_r(x) := N(\mathcal{S}(P); (\Omega, r); x) = \sum_{\substack{L \in \mathcal{S} \\ P(L) < x}} \mathbf{1}_{(\Omega, r)}(L).$$

Given a quadratic number field K, for each rational prime p, let $\mathscr{O}_p^* := \prod_{\mathfrak{p}|p} \mathscr{O}_{\mathfrak{p}}^*$ where \mathfrak{p} is a prime of K. Though according to the isomorphism $D_4 \cong C_4 \rtimes C_2$, we see that C_4 is a non-trivial C_2 -module, however the submodule $2C_4$ is trivial. So, we obtain the following.

Lemma 8.6. For each quadratic number field K, and for each odd prime p, we have

$$|\operatorname{Hom}_{C_2}(\mathscr{O}_p^*, 2C_4)| = 2.$$

In other words, it consists of a trivial morphism and a surjective one.

Proof. Let \mathfrak{p} be a prime of K lying above p. If $\rho_1, \rho_2 : \mathscr{O}_p^* \to 2C_4$ are the same when restricted to \mathfrak{p} , then the Galois action forces $\rho_1 = \rho_2$. This shows that $\rho : \mathscr{O}_p^* \to 2C_4$ is totally determined by the group homomorphism $\rho : \mathscr{O}_p^* \to 2C_4$. But p is an odd prime, and \mathscr{O}_p^* factors through $\mu(K_p)$, i.e., it is totally determined by ζ_p . So, there are only two different C_2 -morphisms.

Proposition 8.7. There exists a constant C > 0, for each quadratic number field K such that for each r = 0, 1, 2, ..., we have

$$N_{K,r}(x) := \sum_{\substack{L \in \mathcal{S} \\ K \subseteq L, P(L) < x}} \mathbf{1}_{(\Omega,r)}(L) \le C |\operatorname{Hom}(\operatorname{Cl}_K, C_4)| \frac{x}{P(K)} (\log \log x)^r.$$

Proof. Let

$$a_2 := \max_{[\Sigma_2:\mathbb{Q}_2]=2} |\operatorname{Hom}(\Sigma_2, C_4)|.$$

Define a generating series as follows:

$$D_r(s) := a_2 \prod_{p \in \mathcal{P}} (1 + p^{-s}) \sum_{\substack{\underline{p} \in \mathcal{P}^r \\ 2 < p_1 < \dots < p_r}} 4^r p_1^{-s} \cdots p_r^{-s}$$

Claim: for each quadratic number field, let $D_{K,r}(s) := |\text{Hom}(\text{Cl}_K, C_4)| P(K)^{-s} D_r(s)$ and write $D_{K,r}(s) = \sum_{n=1}^{\infty} a_n n^{-s}$, then we have

$$N_{K,r}(x) \le \sum_{n < x} a_n$$

for all x > 0. According to Proposition A.5, we know that

$$N_{K,r}(x) \le |\operatorname{Hom}(\operatorname{Cl}_K, C_4)| \sum_{\substack{\rho: \prod_{p \nmid \infty} \mathcal{O}_p^* \to C_4 \\ P(\rho) < x/P(K)}} \mathbf{1}_{(\Omega, r)}(\rho).$$

(See Definition 7.4 for $P(\rho)$, $\mathbf{1}_{(\Omega,r)}(\rho)$.) For each quadratic number field K, and for each odd prime p, we have

$$|\operatorname{Hom}_{C_2}(\mathscr{O}_p^*, 2C_4)| = 2$$
 and $|\operatorname{Sur}(\mathscr{O}_p^*, C_4)| \le 4.$

Then for each natural number n, we have

$$\sum_{\substack{L\in\mathcal{S}\\K\subseteq L,P(L)=n}} \mathbf{1}_{(\Omega,r)}(L) \leq \sum_{\substack{[K:\mathbb{Q}]=2\\n/P(K)=p_1\cdots p_rq_1\cdots q_l}} \prod_{i=1}^r \left(\sum_{\substack{\rho:\mathscr{O}_{p_i}^*\to C_4\\\mathrm{im}(\rho)=C_4}} 1\right) \prod_{j=1}^i \left(\sum_{\substack{\rho:\mathscr{O}_{q_j}^*\to C_4\\\mathrm{im}(\rho)=2C_4}} 1\right)$$
$$\leq \sum_{\substack{[K:\mathbb{Q}]=2\\n/P(K)=p_1\cdots p_rq_1\cdots q_l}} a_2 4^r.$$

So $D_r(s)$ gives an estimate of $N_{K,r}(x)$. The continuation of $D_r(s)$ is guaranteed by Theorem 5.10 and 5.9. To be precise, there exists some holomorphic functions f_0, \ldots, f_r in the closed half-plane $\Re(s) \geq 1$ such that $f_0(1) \neq 0$ and such that

$$D_r(s) = (s-1)^{-1} \sum_{\substack{i=0\\40}}^r f_{r-i}(s) \left(\log \frac{1}{s-1}\right)^i.$$

Therefore, by Tauberian Theorem 5.1, we have

$$N_{K,r}(s) \le C |\operatorname{Hom}(\operatorname{Cl}_K, C_4)| \frac{x}{P(K)} (\log \log x)^r.$$

Theorem 8.8. If (8.6) holds, then the Conjecture 1.6(2) is true for $(\mathcal{S}(P), \Omega)$.

Proof. First we give an estimate of $N_r(x)$. Write $D_r(x) = \sum_{n=1}^{\infty} a_n n^{-s}$ and let $A(x) := \sum_{n \le x} a_n$. According to Proposition 8.7, we have

$$N_r(x) = \sum_{[K:\mathbb{Q}]=2} N_{K,r}(x)$$

$$\leq \int_{1^-}^x H(x/t) \,\mathrm{d}\,A(t)$$

$$\ll \int_{1^-}^x \frac{x}{t} \log x \,\mathrm{d}\,A(t)$$

$$= \frac{x}{t} \log x A(t) \Big|_1^x + x \log x \int_{1^-}^x \frac{A(t)}{t^2} \,\mathrm{d}\,t$$

$$\ll x \log x (\log \log x)^r + x \log x \int_{e^+}^x \frac{(\log \log t)^r}{t} \,\mathrm{d}\,t$$

$$\ll x (\log x)^2 (\log \log x)^r.$$

This shows that

$$N_r(x) = o(N(x))$$

for all $r = 0, 1, 2, \ldots$ We therefore see that the Conjecture 1.6(2) holds in this case.

By Theorem 4.4 and 4.5, we get the following.

Corollary 8.9. If (8.6) holds, then we have

$$\mathbb{P}(\operatorname{rk}_2 2\operatorname{Cl}_L \leq r) = 0 \quad and \quad \mathbb{E}(|\operatorname{Hom}(2\operatorname{Cl}_L, C_2)|) = +\infty.$$

A. CLASS FIELD THEORY

For the convenience of the reader, we write down here the main results used from Class Field Theory. Let K be any number field. Denote by $\mathbb{A}_K = \prod'_{\mathfrak{p}} K_{\mathfrak{p}}$ the ring of adèles of K, and by $J_K := \mathbb{A}_K^*$ the *idèle group* of K. There exists a diagonal embedding $K^* \to J_K$, and we call elements of K^* the *principal idèles*. The quotient group $C_K = J_K/K^*$ is the *idèle class* group of K. There is a map $(\alpha_{\mathfrak{p}})_{\mathfrak{p}} \mapsto \prod_{\mathfrak{p} \neq \infty} \mathfrak{p}^{v_{\mathfrak{p}}(\alpha_{\mathfrak{p}})}$, whose kernel is $J_K^{S_{\infty}} := \prod_{v \mid \infty} K_v^* \times \prod_{v \nmid \infty} \mathscr{O}_v^*$. So, we get the following short exact sequence

(A.1)
$$1 \to J_K^{S_\infty} K^* / K^* \to \mathcal{C}_K \to \mathcal{C}_K \to 1$$

More generally, for any finite set S of primes containing infinity, we obtain

(A.2)
$$1 \to J_K^S K^* / K^* \to \mathcal{C}_K \to \mathcal{C}l_K^S \to 1.$$

by the similar idea. Using the above short exact sequence, we have the following result.

Lemma A.1. If S is a finite set of primes containing infinity such that $\operatorname{Cl}_K^S = 1$, then $J_K^S K^* / K^* \cong \operatorname{C}_K$.

Since J_K^S in general is easier to work with, let's consider the following short exact sequence.

(A.3)
$$1 \to K^S \to J_K^S \to J_K^S K^*/K^* \to 1,$$

where K^S is the S-units of K, which could be defined as $K^* \cap J_K^S$. Since $J_K^S K^*/K^* = J_K^S/J_K^S \cap K^*$, the notation itself justifies the sequence. When $K = \mathbb{Q}$, we take $S = S_{\infty}$, and the short exact sequence says that

$$1 \to \{\pm 1\} \to J_{\mathbb{Q}}^{S_{\infty}} \to J_{\mathbb{Q}}^{S_{\infty}} \mathbb{Q}^* / \mathbb{Q}^* \to 1.$$

If G is any finite abelian group, we have

 $1 \to \operatorname{Hom}(J^{S_{\infty}}_{\mathbb{Q}}\mathbb{Q}^*/\mathbb{Q}^*, G) \to \operatorname{Hom}(J^{S_{\infty}}_{\mathbb{Q}}, G) \to \operatorname{Hom}(\{\pm 1\}, G) \to \cdots$

Since for any map $\rho: J_{\mathbb{Q}}^{S_{\infty}} \to G$, either $\rho(-1) = 1$ or $\rho(-1) \neq 1$, we see that a homomorphism $\rho: \prod_{p} \mathbb{Z}_{p}^{*} \to G$ could always be extended to a homomorphism $\tilde{\rho}: J_{\mathbb{Q}}^{S_{\infty}} \to G$ such that it is an image of an element of $\operatorname{Hom}(J_{\mathbb{Q}}^{S_{\infty}}\mathbb{Q}^{*}/\mathbb{Q}^{*}, G)$, by setting up a suitable value of $\rho_{\infty}(-1)$. Clearly such a value is unique. In other words, there exists a unique map $\tilde{\rho}: J_{\mathbb{Q}}^{S_{\infty}}\mathbb{Q}^{*}/\mathbb{Q}^{*} \to G$ such that $\tilde{\rho}|_{p} = \rho|_{p}$ for all $p \nmid \infty$. Based on this correspondence, we have the following result.

Lemma A.2. Let G be an abelian group. There is a one-to-one correspondence between $\operatorname{Hom}(C_{\mathbb{Q}},G)$ and $\operatorname{Hom}(\prod_{p\nmid\infty}\mathbb{Z}_p^*,G)$, given by $\tilde{\rho}\mapsto\prod_{p\restriction\infty}\tilde{\rho}|_p$. In particular, $\operatorname{Hom}(\prod_p\mathbb{Z}_p^*,G)$ corresponds to the set of Artin reciprocity maps, i.e., abelian G-fields, under this map.

Proof. The only thing we need to prove here is that if $\tilde{\rho}$ is an Artin reciprocity map, then $\rho := \prod_{p \nmid \infty} \tilde{\rho}|_p$ is already surjective. Since $\tilde{\rho}$ corresponds to a class field L, we know that $G \cong \operatorname{Gal}(L/\mathbb{Q})$ must be generated by all the inertia subgroups I(p) where p runs over all rational primes. Using Global-to-Local principal, we know that I(p) is just given by $\rho(\mathbb{Z}_p^*)$. Therefore, $\operatorname{im}(\rho) = G$. So the image of the set of Artin reciprocity maps under the correspondence $\tilde{\rho} \mapsto \prod_{p \mid \infty} \tilde{\rho}|_p$ is just $\operatorname{Sur}(\prod_{p \nmid \infty} \mathbb{Z}_p^*, G)$.

When Class Field Theory is applied to general number fields, we need to consider the Galois action carefully. In general, we could prove the following.

Lemma A.3. Let K be a fixed Galois number field with an isomorphism $\operatorname{Gal}(K/\mathbb{Q}) \cong H_1$, and let H_2 be a finite abelian group. If S is a finite set of primes of K containing the ones at infinity such that \mathscr{O}_K^S has class number 1, then there is a one-to-one correspondence between $\operatorname{Hom}_{H_2}(J_K^S K^*/K^*, H_2)$ and $\operatorname{Hom}_{H_1}(C_K, H_2)$, where J_K is the group of idèles, and C_K is the idèles class group, and H_2 is equipped with a fixed H_1 -action.

Proof. The correspondence is induced by the isomorphism $J_K^S K^*/K^* \cong C_K$. Note that since C_K is an H_1 -module, we see that $J_K^S K^*/K^*$ is also a well-defined H_1 -module.

Lemma A.4. Let H_1, H_2 be two finite abelian groups. Let K be a fixed Galois number field with an isomorphism $\operatorname{Gal}(K/\mathbb{Q}) \cong H_1$, and let H_2 be a finite abelian group, and let L/K be an H_2 -extension. The field extension L/\mathbb{Q} is Galois with $G := \operatorname{Gal}(L/\mathbb{Q})$ if and only if the corresponding Artin reciprocity map $\rho : C_K \to H_2$ is an H_1 -morphism, in which case H_2 is equipped with the H_1 -action satisfying the short exact sequence

(A.4)
$$1 \to H_2 \to G \to H_1 \to 1.$$

Proof. Let L/K be an H_2 -extension, and let M be the Galois closure of L/\mathbb{Q} . We treat C_K, C_L as subgroups of C_M .

First assume that L/\mathbb{Q} is a Galois extension, which means that we already have the short exact sequence (A.4). And the action of H_1 on H_2 is exactly the action of conjugation. By Class Field Theory we know that $C_K/N C_L \cong \text{Gal}(L/K) \cong H_2$, where N means norm $\text{Nm}_{L/K}$. Since L is Galois, we see that $N C_L$ is a well-defined H_1 -module, which means that the Artin reciprocity map $\rho : C_K \to H_2$ is an H_1 -morphism. All the H_1 -actions showing up are consistent with the short exact sequence (A.4), which gives the Galois action.

Conversely, let $\rho : C_K \to H_2$ be the Artin reciprocity map corresponding to L/K. If $L' \neq L$ is a conjugate, then we see that $N C_L \neq N C_{L'}$, because they corresponds to different abelian extensions of K. In particular, if $g \in \text{Gal}(M/\mathbb{Q})$ is an element such that gL = L', then we see that $g|_K$ induces the map $N C_L \to N C_{L'}$. But this already shows that ρ cannot be an H_1 -morphism. And we are done.

In general, we obtain the following results.

Proposition A.5. Let G be a finite abelian group. If $\rho : J_K^{S_{\infty}} \to G$ be a surjective homomorphism, then there exists at most $|\text{Hom}(\text{Cl}_K, G)|$ many different Artin reciprocity maps $\chi : \text{C}_K \to G$ such that $\chi|_{\mathfrak{p}} = \rho|_{\mathfrak{p}}$ for all $\mathfrak{p} \nmid \infty$.

Proof. Using (A.1), we see that

$$1 \to \operatorname{Hom}(\operatorname{Cl}_K, G) \to \operatorname{Hom}(\operatorname{C}_K, G) \to \operatorname{Hom}(J_K^{S_\infty} K^*/K^*, G) \to \cdots$$

If a homomorphism $\psi : J_K^{S_{\infty}} K^*/K^*$ could be lifted to $\operatorname{Hom}(C_K, G)$, then size of the preimages is determined by $|\operatorname{Hom}(\operatorname{Cl}_K, G)|$ according to this long exact sequence. Similarly, by (A.3), we have

$$1 \to \operatorname{Hom}(J_K^{S_{\infty}} K^*/K^*, G) \to \operatorname{Hom}(J_K^{S_{\infty}}, G) \to \operatorname{Hom}(K^S, G) \to \cdots$$

The long exact sequence says that if $\rho : J_K^{S_{\infty}} \to G$ is a homomorphism with a lift in $\operatorname{Hom}(J_K^{S_{\infty}}K^*/K^*, G)$, then the preimage is unique. Combining the two parts, we obtain the result immediately.

In the case of dihedral extensions, we also have the following.

Lemma A.6. If $m \ge 3$, and C_m is a C_2 -module with the nontrivial action $g_2 \cdot g_m = g_m^{-1}$ where g_m , resp. g_2 , is a generator of C_m , resp. C_2 , then $H^2(C_2, C_m) = 0$. In other words, there is only one group $G = D_m$, the dihedral group, satisfying the short exact sequence

$$1 \to C_m \to D_m \to C_2 \to 1,$$

up to isomorphisms between sequences.

Proof. Since C_2 is a cyclic group, the computation of $H^2(C_2, C_m)$ reduces to the Tate cohomology $\hat{H}^0(C_2, C_m) = C_m^{C_2}/N_{C_2}C_m$. An element $g \in C_m$ is fixed by g_2 if and only if g = 1, so $C_m^{C_2} = 1$. And we get $H^2(C_2, C_m) = \hat{H}^0(C_2, C_m) = 0$. Since the zero element in $H^2(C_2, C_m) = \text{Ext}(C_2, C_m)$ is exactly $G = C_m \rtimes C_2$, we obtain the short exact sequence

$$1 \to C_m \to G \to C_2 \to 1.$$

Then we see that $G = \langle g_m, g_2 | g_m^m = g_2^2 = 1, g_2 g_m g_2^{-1} = g_m^{-1} \rangle$, which is exactly the dihedral group D_m .

Proposition A.7. If K_2 is a quadratic number field, C_m is the non-trivial C_2 -module, then a extension K_{2m}/K_2 of degree 2m gives rise to a Galois D_m -field extension K_{2m}/\mathbb{Q} if and only if the Artin reciprocity map $\rho : C_{K_2} \to C_m$ is a C_2 -morphism.

Proof. According to Lemma A.6, the field extension K_{2m}/\mathbb{Q} is Galois if and only if the Artin reciprocity map $\rho : \mathbb{C}_{K_2} \to \mathbb{C}_m$ is a \mathbb{C}_2 -morphism. The Galois group $G := \operatorname{Gal}(K_{2m}/\mathbb{Q})$ satisfies the short exact sequence

$$1 \to C_m \to G \to C_2 \to 1.$$

If C_m is a trivial C_2 -module, then clearly G is isomorphic to C_{2m} . Else if C_m is nontrivial, then according to Lemma, we know that $G \cong C_m \rtimes C_2$ with the action of C_2 on C_m the same as in the condition, which is just D_m .

References

- Salim Ali Altug, A. I. Vijaya Shankar, Ila Varma, and Kevin H. Wilson. The number of d₄-fields ordered by conductor. *Journal of the European Mathematical Society*, 23:2733–2785, 2021.
- [2] Alex Bartel and Hendrik W Lenstra Jr. On class groups of random number fields. Proceedings of the London Mathematical Society, 121(4):927–953, 2020.
- [3] Manjul Bhargava. The density of discriminants of quartic rings and fields. Annals of mathematics, ISSN 0003-486X, Vol. 162, № 2, 2005, pags. 1031-1062, 162:1031-1063, 09 2005.
- [4] Manjul Bhargava. Mass formulae for extensions of local fields, and conjectures on the density of number field discriminants. *International Mathematics Research Notices*, 2007, 2007.
- [5] Manjul Bhargava, Arul Shankar, and Jacob Tsimerman. On the davenport-heilbronn theorems and second order terms. *Inventiones mathematicae*, 193(2):439–499, 2013.
- [6] Henri Cohen and Jacques Martinet. Étude heuristique des groupes de classes des corps de nombres. Journal für die reine und angewandte Mathematik, 404:39–76, 1990.
- [7] Boris Datskovsky and David J Wright. Density of discriminants of cubic extensions. 1988.
- [8] H. Davenport and H. Heilbronn. On the Density of Discriminants of Cubic Fields. II. Proceedings of the Royal Society of London Series A, 322(1551):405–420, May 1971.
- [9] Hubert Delange. Généralisation du théorème de Ikehara. Annales scientifiques de l'École Normale Supérieure, 3e série, 71(3):213-242, 1954.
- [10] Ellen Eischen, Ling Long, Rachel Pries, and Katherine Stange. Directions in Number Theory: Proceedings of the 2014 WIN3 Workshop, volume 3. Springer, Cham, 01 2016.
- [11] III Frank Gerth. Extension of conjectures of cohen and lenstra. Expositiones Mathematicae, 5(2):181 184, 1987.
- [12] Makoto Ishida. The genus fields of algebraic number fields. Lecture notes in mathematics. Springer, Berlin, Heidelberg, 1 edition, 1976.
- [13] Henryk Iwaniec and Emmanuel Kowalski. Analytic Number Theory. American Mathematical Society Colloquium Publications. American Mathematical Society, Providence, Rhode Island, 2004.
- [14] Gunter Malle. On the distribution of galois groups, ii. Experimental Mathematics, 13(2):129–135, 2004.
- [15] H.L. Montgomery and R.C. Vaughan. Multiplicative Number Theory I: Classical Theory. Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 2006.
- [16] W. Narkiewicz. Elementary and Analytic Theory of Algebraic Numbers. Springer, Berlin Heidelberg, 2014.
- [17] J. Neukirch and N. Schappacher. Algebraic Number Theory. Grundlehren der mathematischen Wissenschaften. Springer, Berlin Heidelberg, 2013.
- [18] P. Roquette and H. Zassenhaus. A class rank estimate for algebraic number fields. Journal of the London Mathematical Society, s1-44(1):31–38, 1969.
- [19] Alexander Smith. The distribution of l[∞]-Selmer groups in degree l twist families. arXiv e-prints, page arXiv:2207.05674, July 2022.
- [20] Weitong Wang and Melanie Matchett Wood. Moments and interpretations of the cohen–lenstra–martinet heuristics. Commentarii Mathematici Helvetici, 96(2):339–387, 2021.

[21] Melanie Matchett Wood. On the probabilities of local behaviors in abelian field extensions. Compositio Mathematica, 146(1):102–128, 2010.

SHING-TUNG YAU CENTER OF SOUTHEAST UNIVERSITY, 15TH FLOOR YIFU ARCHITECTURE BUILDING, 2 SIPAILOU, NANJING, JIANGSU PROVINCE 210096 CHINA Email address: wangweitong@seu.edu