

Article

Entanglement dynamics and classical complexity

Jiaozi Wang^{1,*}, Barbara Dietz² , Dario Rosa^{2,3} , and Giuliano Benenti^{4,5,6} 

¹ Department of Physics, University of Osnabrück, D-49069 Osnabrück, Germany

² Center for Theoretical Physics of Complex Systems, Institute for Basic Science (IBS), Daejeon - 34126, Korea

³ Basic Science Program, Korea University of Science and Technology (UST), Daejeon - 34113, Korea

⁴ Center for Nonlinear and Complex Systems, Dipartimento di Scienza e Alta Tecnologia, Università degli Studi dell'Insubria, via Valleggio 11, 22100 Como, Italy

⁵ Istituto Nazionale di Fisica Nucleare, Sezione di Milano, via Celoria 16, 20133 Milano, Italy

⁶ NEST, Istituto Nanoscienze-CNR, I-56126 Pisa, Italy

* Correspondence: jiaozi.wang@uos.de

Version November 22, 2022 submitted to Journal Not Specified

Abstract: We study the dynamical generation of entanglement for a two-body interacting system, starting from a separable coherent state. We show analytically that in the quasiclassical regime the entanglement growth rate can be simply computed by means of the underlying classical dynamics. Furthermore, this rate is given by the Kolmogorov-Sinai entropy, which characterizes dynamical complexity of classical motion. Our results, illustrated by numerical simulations on a model of coupled rotators, establish in the quasiclassical regime a link between the generation of entanglement, a purely quantum phenomenon, and classical complexity.

Keywords: quantum complexity, quantum to classical transition

1. Introduction

The characterization of complexity in quantum systems is a key problem, not only for fundamental reasons but also for the development of quantum technologies [1–4]. While for classical dynamical systems a well-established notion of complexity exists, based on Kolmogorov-Sinai (KS) entropy [5], which in turn is related to the exponential instability of orbits, in the quantum realm the measure of complexity has proven to be an elusive problem.

First, we cannot *sic and simpliciter* use trajectories, due the Heisenberg uncertainty principle. To circumvent such problem, phase-space approaches have been proposed [6–21], based on the evolution of phase space distributions. Second, entanglement, the key resource in the quest for quantum advantage, is peculiar to quantum composite systems, and therefore is a source of quantum complexity without classical analogue. Since for pure bipartite systems the reduced von Neumann entropy, known as entanglement entropy, is the well-established measure of entanglement [22], it is interesting to investigate whether its growth in a dynamical system is related to the KS entropy of the underlying classical dynamics.

For bosonic systems with an unstable quadratic Hamiltonian, entanglement entropy grows linearly in time, with a rate upper bounded by the KS entropy, the bound being saturated under suitable conditions on the size of the bipartitions [23]. The question then arises, whether the entanglement growth of chaotic quantum systems in the quasiclassical regime is also determined by the KS entropy. This issue was investigated more than two decades ago, with numerical results suggesting that the entanglement generation rate is given by the KS entropy [24]. On the other hand, such results were obtained in the

28 weakly chaotic regime, with coexistence of chaotic seas and tori, while another study in the strongly chaotic
 29 region, where the effect of tori is negligible, showed instead no increase of the entanglement production
 30 rate upon increase of the maximum Lyapunov exponents [25]. This apparent contradiction was explained
 31 by a quasiclassical calculation for the linear entropy, approximating the entanglement entropy, under the
 32 condition of weak coupling between the subsystems in the underlying classical dynamics [26]. This work
 33 showed that the entanglement growth rate is determined by the minimal value of the three rates given
 34 by the standard one deduced from the interaction term and the largest Lyapunov exponents of the two
 35 subsystems, respectively.

36 In this paper, we remove the above restriction on the coupling strength and compare the quantum
 37 evolution starting from separable coherent states with the classical evolution of initially Gaussian
 38 distributions, of size determined by the effective Planck constant of the corresponding quantum dynamics.
 39 We show that in the quasiclassical regime quantum and classical linear entropy are in agreement and grow
 40 with rate given by the KS entropy of classical dynamics. Our analytical results are illustrated by numerical
 41 simulations for a model of kicked coupled rotators.

42 This work is dedicated to our friend and colleague Giulio Casati, who has always had a deep interest
 43 in understanding the complexity of quantum motion.

44 2. Analytical results

In this section, we connect, for an overall pure bipartite system, the growth rate of linear entropy to
 the KS entropy of the classical underlying dynamics. We consider a two body system, whose Hamiltonian
 reads

$$\hat{H} = \hat{H}_1(\hat{q}_1, \hat{p}_1) + \hat{H}_2(\hat{q}_2, \hat{p}_2) + \hat{H}_{12}(\hat{q}_1, \hat{p}_1, \hat{q}_2, \hat{p}_2). \quad (1)$$

The corresponding classical Hamiltonian is written as

$$H(q_1, p_1, q_2, p_2) = H_1(q_1, p_1) + H_2(q_2, p_2) + H_{12}(q_1, p_1, q_2, p_2). \quad (2)$$

We compute as entanglement measure the linear entropy (also known as *second Rényi entropy*) of a
 subsystem (for example, system 1), which is defined as

$$S(\hat{\rho}_1) = -\ln(\text{Tr}(\hat{\rho}_1^2)). \quad (3)$$

45 Here $\hat{\rho}_1$ is the reduced density matrix of the system 1, $\hat{\rho}_1 = \text{Tr}_2(\hat{\rho})$, where the partial trace is taken over
 46 system 2 and $\hat{\rho}$ is the density matrix of the composite system. Note that equivalently we could have
 47 considered system 2, since $S(\hat{\rho}_2) = S(\hat{\rho}_1)$, with $\hat{\rho}_2 = \text{Tr}_1(\hat{\rho})$.

In order to obtain the classical analog of the linear entropy, we make use of the Husimi function [27]
 of the density matrix $\hat{\rho}$, given by

$$W_H(\gamma) = \frac{1}{(2\pi\hbar)^2} \langle \gamma | \hat{\rho} | \gamma \rangle, \quad (4)$$

where $\gamma = (q_1, p_1, q_2, p_2)$, $|\gamma\rangle$ denotes the coherent state of the composite system centered at γ , and \hbar is
 the effective Planck constant. In the quasiclassical limit $\hbar \rightarrow 0$, the trace of $\hat{\rho}_1^2$ can be carried out by making
 use of the Husimi function W_H^1 of $\hat{\rho}_1$ as

$$\text{Tr}(\hat{\rho}_1^2) = \int d\gamma_1 [W_H^1(\gamma_1)]^2, \quad (5)$$

where $\gamma_1 = (q_1, p_1)$, $|\gamma_1\rangle$ denotes the coherent state of system 1 centered at γ_1 , and

$$W_H^1(\gamma_1) = \frac{1}{2\pi\hbar} \langle \gamma_1 | \hat{\rho}_1 | \gamma_1 \rangle. \quad (6)$$

Furthermore, the reduced density matrix $\hat{\rho}_1$ can also be obtained in terms of the coherent states of the system 2, denoted by $|\gamma_2\rangle$, as

$$\hat{\rho}_1 = \text{Tr}_2(\hat{\rho}) = \frac{1}{2\pi\hbar} \int d\gamma_2 \langle \gamma_2 | \hat{\rho} | \gamma_2 \rangle. \quad (7)$$

Substituting Eq.(7) into Eq.(6), we have

$$W_H^1(\gamma_1) = \int d\gamma_2 W_H(\gamma), \quad (8)$$

yielding with Eq. (5)

$$\text{Tr}(\hat{\rho}_1^2) = \int d\gamma_1 \left| \int W_H(\gamma) d\gamma_2 \right|^2. \quad (9)$$

Hence, we obtain

$$S(\hat{\rho}_1) = -\ln \left[\int d\gamma_1 \left(\int d\gamma_2 W_H(\gamma) \right)^2 \right]. \quad (10)$$

After replacing the Husimi function $W_H(\gamma)$ with the classical distribution function $\rho(\gamma)$, the classical analog of linear entropy can be written as

$$S_{cl}(\rho_1) = -\ln \left[\int d\gamma_1 (\rho_{re}^1(\gamma_1))^2 \right], \quad (11)$$

where $\rho_{re}^1(\gamma_1)$ indicates the marginal distribution function of γ_1 ,

$$\rho_{re}^1(\gamma_1) = \int d\gamma_2 \rho(\gamma). \quad (12)$$

It is expected that

$$S(\hat{\rho}_1) \approx S_{cl}(\rho_1) \quad (13)$$

48 holds in the quasiclassical limit in which the effective Planck constant $\hbar \rightarrow 0$.

An explicit expression can be derived for the classical entropy $S_{cl}(\rho_1)$ as follows. We consider the initial state as the “most classical” state, that is, a coherent state $|\gamma\rangle$, whose corresponding classical distribution function can be written as

$$\rho_0(\gamma) = \frac{1}{(\pi\hbar_c)^2} \exp\left(-\frac{1}{\hbar_c} |\gamma - \gamma^0|^2\right), \quad (14)$$

which has a Gaussian form whose center is denoted by γ^0 , $\hbar_c = \hbar$ is chosen to be the same as the effective Planck constant in the quantum case, and $|\gamma - \gamma^0|$ indicates the norm of the vector $\delta\gamma = \gamma - \gamma^0$. In the quasiclassical limit, one has $\hbar_c \rightarrow 0$, which means that, for times smaller than the Ehrenfest time scale t_E (with $t_E \rightarrow \infty$ as $\hbar_c \rightarrow 0$), almost all the states in the ensemble remain close to the center $\gamma^0(t)$. This

implies that the distribution of states at time t , $\rho_t(\gamma)$, is significantly different from zero only for small $|\delta\gamma|$. In this case, the time evolution of $\delta\gamma$ is determined by the so-called stability matrix

$$\mathbf{M}_t^{ij} = \left. \frac{\partial(\delta\gamma_i(t))}{\partial(\delta\gamma_j(0))} \right|_{\delta\gamma(0)=0}, \quad (15)$$

with

$$\delta\gamma(t) = \mathbf{M}_t \delta\gamma(0). \quad (16)$$

49 As the classical linear entropy is independent of the coordinates origin, for the convenience of the following
50 discussion, we choose the position of the center $\gamma^0(t)$ as the origin of coordinates. In this local coordinate
51 system along $\gamma^0(t)$, we can replace $\delta\gamma(t) = \gamma(t) - \gamma^0(t)$ by $\gamma(t)$.

Then making use of Liouville's theorem, the distribution at time t can be written as

$$\rho_t(\gamma) = \rho_0(\mathbf{M}_t^{-1}\gamma), \quad (17)$$

and therefore

$$\rho_t(\gamma) = \frac{1}{(\pi\hbar_c)^2} \exp\left(-\frac{1}{\hbar_c} |\mathbf{M}_t^{-1}\gamma|^2\right). \quad (18)$$

Using the positive definite symmetric matrix

$$\mathbf{A}_t \equiv (\mathbf{M}_t^{-1})^T \mathbf{M}_t^{-1}, \quad (19)$$

the density distribution at time t can be written as

$$\rho_t(\mathbf{x}) = \frac{1}{(\pi\hbar_c)^2} \exp\left(-\frac{1}{\hbar_c} \sum_{i,j=1}^4 x_i A_t^{ij}(t) x_j\right), \quad (20)$$

which is a Gaussian distribution, with \mathbf{x} corresponding to γ , that is, $(x_1, x_2, x_3, x_4) = (q_1, p_1, q_2, p_2)$. In order to calculate the classical linear entropy, we first calculate the marginal distribution function of $\rho_t(\mathbf{x})$ for system 1:

$$\rho_t^1(x_1, x_2) = \int \rho_t(x_1, x_2, x_3, x_4) dx_3 dx_4. \quad (21)$$

Then the classical linear entropy at time t can be written as

$$S_{cl}(\rho_t) = -\ln \left[\int dx_1 dx_2 \left(\rho_t^1(x_1, x_2) \right)^2 \right]. \quad (22)$$

As outlined in the Appendix, by writing \mathbf{A}_t in block form,

$$\mathbf{A}_t = \begin{pmatrix} \hat{\mathbf{a}} & \hat{\mathbf{b}} \\ \hat{\mathbf{b}}^T & \hat{\mathbf{d}} \end{pmatrix}, \quad (23)$$

we obtain our first main result

$$S_{cl}(\rho_t) = \ln(2\pi\hbar) + \frac{1}{2} \ln[\det(\hat{\mathbf{d}})]. \quad (24)$$

In order to compute $\det \hat{d}$, we sum the eigenvalues of the operator \hat{d} (denoted by d_k , in order of descending energy), which are in close relation to the eigenvalues of A_t . We diagonalize the symmetric matrix A_t as

$$A_t = V \text{diag}\{A_1, A_2, A_3, A_4\} V^T, \quad (25)$$

52 where diag indicates a diagonal matrix, A_k is the k -th eigenvalue of A_t , and V is an orthogonal matrix. If
 53 the system is chaotic, $A_k \propto e^{2\lambda_k t}$, where λ_k is the k -th Lyapunov exponent, with $\lambda_1 > \lambda_2 > 0 > \lambda_3 > \lambda_4$,
 54 and $\lambda_3 = -\lambda_2, \lambda_4 = -\lambda_1$.

Hence, in the typical case in which the eigenvectors (denoted by $|A_k\rangle, k = 1, 2$) of A_t corresponding to the eigenvalues A_1 and A_2 have non-zero components within the Hilbert space of system 2, we have

$$d_1 \propto e^{2\lambda_1 t}, \quad d_2 \propto e^{2\lambda_2 t}. \quad (26)$$

As a result,

$$\det \hat{d} \propto \exp 2(\lambda_1 + \lambda_2)t, \quad (27)$$

which directly leads to

$$S_{cl}(\rho_t) - S_{cl}(\rho_0) = (\lambda_1 + \lambda_2)t, \quad (28)$$

55 indicating that the growth rate of the linear entropy is given by the Kolmogorov-Sinai entropy of the
 56 overall system, which is the second main result of our work.

57 3. Numerical results

In this section, we numerically illustrate the prediction of equivalence between the classical and quantum growth of linear entropies, Eq. (13), as well as the growth as predicted in Eq. (28), by means of a two-body system which has a clearly defined classical counterpart. More specifically, we consider two coupled rotators (or coupled tops)[28,29], with respective angular momentum operators $\hat{S} = (\hat{S}_x, \hat{S}_y, \hat{S}_z)^T$ and $\hat{L} = (\hat{L}_x, \hat{L}_y, \hat{L}_z)^T$, and a time-dependent Hamiltonian with kicked interaction:

$$\hat{H} = \frac{a}{j}(\hat{S}_z + \hat{L}_z) + \frac{c}{j^2} \hat{S}_x \hat{L}_x \sum_{n=-\infty}^{\infty} \delta(t-n), \quad (29)$$

where j is the (half-integer or integer) total angular momentum quantum number of both tops. The Hamiltonian possesses constants of the motion, \hat{S}^2 and \hat{L}^2 . The Hilbert space is expanded by making use of $|s, m_s, l, m_l\rangle \equiv |s, m_s\rangle \otimes |l, m_l\rangle$, which are the joint eigenvectors of $\hat{S}^2, \hat{S}_z, \hat{L}^2, \hat{L}_z$,

$$\begin{aligned} \hat{S}^2 |s, m_s, l, m_l\rangle &= s(s+1) |s, m_s, l, m_l\rangle, \\ \hat{S}_z |s, m_s, l, m_l\rangle &= m_s |s, m_s, l, m_l\rangle, \\ \hat{L}^2 |s, m_s, l, m_l\rangle &= l(l+1) |s, m_s, l, m_l\rangle, \\ \hat{L}_z |s, m_s, l, m_l\rangle &= m_l |s, m_s, l, m_l\rangle. \end{aligned} \quad (30)$$

58 where $m_s \in \{-s, -s+1, \dots, s-1, s\}$, and $m_l \in \{-l, -l+1, \dots, l-1, l\}$. Here we choose $s = l = j$.

The Floquet operator, that is the unitary evolution operator between consecutive kicks, can be written as

$$\hat{F} = \exp[-ia(\hat{S}_z + \hat{L}_z)] \exp[-i\frac{c}{j} \hat{S}_x \hat{L}_x]. \quad (31)$$




Figure 1. Three-dimensional Poincaré surface of section for the chaotic case (a): $a = 5, c = 3$ and the near-integrable case (b): $a = 5, c = 0.5$, where we fix $\phi_2 = 0$. Here we only consider a single trajectory start from $(\theta_1, \phi_1, \theta_2, \phi_2) = (\frac{\pi}{4}, 0, \frac{3\pi}{4}, 0)$ (see text for the definition of the angles θ_k and $\phi_k, k = 1, 2$).

The classical counterpart can be obtained by taking the quasiclassical limit $\hbar = \frac{1}{j} \rightarrow 0$. Introducing the rescaled angular momenta $\hat{S}_k = \frac{\hat{S}_k}{j}$ and $\hat{L}_k = \frac{\hat{L}_k}{j}$, and considering the quasiclassical limit $j \rightarrow \infty$, yields the classical analog of the model,

$$\mathcal{H}_c = a(\mathcal{S}_z + \mathcal{L}_z) + c\mathcal{S}_x\mathcal{L}_x \sum_{n=-\infty}^{\infty} \delta(t-n), \quad (32)$$

59 where $\mathcal{S}_x^2 + \mathcal{S}_y^2 + \mathcal{S}_z^2 = \mathcal{L}_x^2 + \mathcal{L}_y^2 + \mathcal{L}_z^2 = 1$. Depending on the coupling strength the classical motion can
 60 be either chaotic or nearly-integrable, as shown by the three-dimensional Poincaré surfaces of sections of
 61 Fig. 1.

In the numerical simulations of both the quantum and classical cases, the linear entropy is averaged over N_p different initial states. In the quantum case, we consider the initial states $|\theta_1, \phi_1, \theta_2, \phi_2\rangle \equiv |\theta_1, \phi_1\rangle \otimes |\theta_2, \phi_2\rangle$, where $|\theta_1, \phi_1\rangle$ and $|\theta_2, \phi_2\rangle$ indicate the spin coherent state of the first rotor,

$$|\theta_1, \phi_1\rangle = e^{i\theta_1\hat{S}_z} e^{i\phi_1\hat{S}_y} |j, j\rangle, \quad (33)$$

and an analogous expression holds for the second rotor. Then, the quantum averaged linear entropy is calculated as follows,

$$\bar{S}_q(t) = \frac{1}{N_p} \sum_p \text{Tr}((\hat{\rho}_1^p(t))^2), \quad (34)$$

where

$$\hat{\rho}_1^p(t) = \text{Tr}_2(\hat{F}^t |\theta_1^p, \phi_1^p, \theta_2^p, \phi_2^p\rangle \langle \theta_1^p, \phi_1^p, \theta_2^p, \phi_2^p| (\hat{F}^\dagger)^t), \quad (35)$$

and $(\theta_1^p, \phi_1^p, \theta_2^p, \phi_2^p)$ are chosen randomly. In the classical case, we consider an initial ensemble of Gaussian states,

$$\begin{aligned} \rho_0(\theta'_1, \phi'_1, \theta'_2, \phi'_2) &= A \exp\left(-\frac{(\theta'_1 - \theta_1)^2}{\hbar_c} - \frac{\sin^2(\theta_1)(\phi'_1 - \phi_1)^2}{\hbar_c}\right) \\ &\times \exp\left(-\frac{(\theta'_2 - \theta_2)^2}{\hbar_c} - \frac{\sin^2(\theta_2)(\phi'_2 - \phi_2)^2}{\hbar_c}\right), \end{aligned} \quad (36)$$

which in case of $\hbar_c \rightarrow 0$ can be written in terms of canonical variables $(q_1, p_1, q_2, p_2) = (\phi_1, \cos \theta_1, \phi_2, \cos \theta_2)$ as

$$\begin{aligned} \rho_0(q'_1, p'_1, q'_2, p'_2) &= A' \exp\left(-\frac{(q'_1 - q_1)^2}{\hbar_c(1 - p_1^2)} - \frac{(1 - p_1^2)(p'_1 - p_1)^2}{\hbar_c}\right) \\ &\times \exp\left(-\frac{(q'_2 - q_2)^2}{\hbar_c(1 - p_2^2)} - \frac{(1 - p_2^2)(p'_2 - p_2)^2}{\hbar_c}\right). \end{aligned} \quad (37)$$

Here A and A' are normalization constants. Then the classical averaged linear entropy is calculated as

$$\bar{S}_{cl}(t) = \frac{1}{N_p} \sum_p S_{cl}(\rho_t^p), \quad (38)$$

62 where $S_{cl}(\rho_t^p)$ indicates the classical linear entropy (defined in Eq.(11)), starting from the initial ensemble,
 63 centered at $(q_1^k, p_1^k, q_2^k, p_2^k)$. In our numerical simulations, we considered 10^7 trajectories for each initial
 64 ensemble, and the integral in Eq. (11) is calculated by summing over the whole phase space with respect
 65 to system 1, which is divided into 4×10^6 phase cells.

66 Results for the chaotic regime are shown in Fig. 2. Note that λ_2 is comparable to λ_1 , and the behavior
 67 $S(t) - S(0) = (\lambda_1 + \lambda_2)t$ predicted in Eq.(28) on the basis of a purely classical calculation, can be clearly
 68 seen both for quantum and classical simulations. The growth rate, in very good agreement with the KS
 69 entropy $\lambda_1 + \lambda_2$, is clearly distinguished from the growth rate given by the largest Lyapunov exponent
 70 λ_1 alone. Note that by increasing the coupling strength c the entanglement growth rate increases, in
 71 accordance with the increase of the classical KS entropy. Moreover, it can be clearly seen that the agreement
 72 between the classical and quantum linear entropy extends to longer times as $\hbar = \hbar_c$ is reduced.

73 In Fig. 3, we show data in the regular regime with weaker coupling strength, for which invariant tori
 74 of the integrable model at $c = 0$ are deformed but survive. The volume occupied by tori is the largest
 75 portion of the phase space and this affects the growth of the linear entropy, which is logarithmic rather than
 76 linear. Our numerical results show that, for large enough \hbar , the entropy $\bar{S}(t) \propto \log t^\alpha$, with $\alpha \approx 1$, while α
 77 slowly increases with reducing \hbar . Note that the separation between nearby trajectories increases linearly
 78 in time for integrable dynamics. Therefore, the number of cells of area \hbar occupied in the two-dimensional
 79 phase-space for system 1 is proportional to t^2 , leading to the expected growth $\bar{S}(t) \propto \log t^2$. We therefore
 80 conjecture that such growth would be achieved in the limit $\hbar \rightarrow 0$.

81 4. Conclusions

82 We have shown that in the quasiclassical regime the entanglement growth rate is given by the
 83 Kolmogorov-Sinai entropy of the underlying classical dynamics. Note that we are considering initial
 84 separable coherent states, so that the quantum wave packet closely follows the underlying classical phase
 85 space distribution up to the Ehrenfest time, which diverges as the effective Planck constant $\hbar \rightarrow 0$. In




Figure 2. Quantum (circles with solid line) and classical (triangles with solid line) averaged linear entropy for different \hbar and \hbar_c in the kicked coupled tops model defined in Eq. (29), for (a): $a = 5, c = 3$ and (b): $a = 5, c = 5$. The dashed lines indicate the functions $\overline{S} = (\lambda_1 + \lambda_2)t$ (red) and $\overline{S} = \lambda_1 t$ (light blue). The initial values of $S(t = 0)$ are subtracted.




Figure 3. Same as in Fig. 2, but for weaker coupling strength $c = 0.5$, for which motion is quasi-integrable. The lines $\bar{S}(t) \propto \log t$ and $\bar{S}(t) \propto \log t^2$ are also drawn.

86 spite of the lack of entanglement in classical mechanics, our results prove, in the quasiclassical regime, the
 87 close connection between entanglement generation and complexity of classical motion. Moreover, our
 88 derivation based on purely classical grounds provides an intuitive picture that could hardly be obtained
 89 on the basis of purely quantum calculations. Finally, the entanglement growth is linear in the classically
 90 chaotic and logarithmic in the regular regime, thus showing the entangling power of chaos.

91 Appendix A Derivation of Eq. (24)

We write A_t in block form,

$$A_t = \begin{pmatrix} \hat{a} & \hat{b} \\ \hat{b}^T & \hat{d} \end{pmatrix}, \quad (\text{A1})$$

where $\hat{a} = \hat{a}^T$, $\hat{d} = \hat{d}^T$, and \hat{b} are 2×2 matrices. Furthermore, we introduce two-dimensional vectors $q_1^T = (x_1, x_2)$ and $q_2^T = (x_3, x_4)$. The matrix Eq. (A1) can be brought to the form

$$(q_1^T, q_2^T) \begin{pmatrix} \hat{a} & \hat{b} \\ \hat{b}^T & \hat{d} \end{pmatrix} \begin{pmatrix} q_1 \\ q_2 \end{pmatrix} = (q_1^T, \tilde{q}_2^T) \begin{pmatrix} \hat{a} & \hat{0} \\ \hat{0} & \hat{d} \end{pmatrix} \begin{pmatrix} q_1 \\ \tilde{q}_2 \end{pmatrix}, \quad (\text{A2})$$

with

$$\tilde{q}_2 = \hat{Y} q_1 + q_2, \quad \hat{Y} = \hat{d}^{-1} \hat{b}^T, \quad (\text{A3})$$

and

$$\hat{a} = \hat{a} - \hat{b} \hat{d}^{-1} \hat{b}^T. \quad (\text{A4})$$

Then, Eq. (20) becomes

$$\rho_i^1(q_1) = \frac{1}{(\pi\hbar_c)^2} \exp\left(-\frac{1}{\hbar_c} q_1^T \hat{a} q_1\right) \int d\tilde{q}_2 \exp\left(-\frac{1}{\hbar_c} \tilde{q}_2^T \hat{d} \tilde{q}_2\right). \quad (\text{A5})$$

The integrals can be performed after introducing the integration variable transformation $\tilde{q}_2 \rightarrow \hat{R}_2 \tilde{q}_2$, with

$$\hat{d} = \hat{R}_2^T \Lambda_2 \hat{R}_2, \quad \Lambda_2 = \begin{pmatrix} d_1 & 0 \\ 0 & d_2 \end{pmatrix}, \quad (\text{A6})$$

yielding

$$\rho_i^1(q_1) = \frac{1}{\pi\hbar_c} \frac{1}{\sqrt{\det(\hat{d})}} \exp\left(-\frac{1}{\hbar_c} q_1^T \hat{a} q_1\right). \quad (\text{A7})$$

Using that $\det(A_i) = 1 = \det(\hat{d}) \det(\hat{a})$, one has

$$\int dq_1 \left(\rho_i^1(q_1)\right)^2 = \frac{\det(\hat{a})}{(\pi\hbar_c)^2} \int dq_1 \exp\left(-\frac{2}{\hbar_c} q_1^T \hat{a} q_1\right) = \frac{1}{2\pi\hbar_c} \sqrt{\det(\hat{a})} = \frac{1}{2\pi\hbar_c} \frac{1}{\sqrt{\det(\hat{d})}}, \quad (\text{A8})$$

92 which leads to Eq.(24).

93 **Author Contributions:** J.W. developed analytical calculations and performed numerical simulations. The work was
94 supervised by G.B., with inputs from B.D and D.R.. All authors discussed the results and contributed to writing and
95 revising the manuscript. All authors have read and agreed to the published version of the manuscript

96 **Funding:** J.W. is supported by the Deutsche Forschungsgemeinschaft (DFG) within the Research Unit FOR 2692
97 under Grant No. 397107022 (GE 1657/3-2) and No. 397067869 (STE 2243/3-2), B.D. and D.R. acknowledge support
98 from the Institute for Basic Science in Korea (IBSR024-D1). G.B. acknowledges the financial support of the INFN
99 through the project QUANTUM.

100 **Conflicts of Interest:** The authors declare no conflict of interest. The funders had no role in the design of the study; in
101 the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the
102 results.

103

- 104 1. Dowling, J.P.; Milburn, G.J. Quantum technology: the second quantum revolution. *Phil. Trans. R. Soc. A.* **2003**,
105 361, 1655–1674.
- 106 2. Acín, A.; Bloch, I.; Buhrman, H.; Calarco, T.; Eichler, C.; Eisert, J.; Esteve, D.; Gisin, N.; Glaser, S.J.; Jelezko,
107 F.; Kuhr, S.; Lewenstein, M.; Riedel, M.F.; Schmidt, P.O.; Thew, R.; Wallraff, A.; Walmsley, I.; Wilhelm, F.K.
108 The quantum technologies roadmap: a European community view. *New Journal of Physics* **2018**, *20*, 080201.
109 doi:10.1088/1367-2630/aad1ea.
- 110 3. Wang, J.; Sciarrino, F.; Laing, A.; Thompson, M.G. Integrated photonic quantum technologies. *Nature Photonics*
111 **2020**, *14*, 273–284. doi:10.1038/s41566-019-0532-1.
- 112 4. Benenti, G.; Casati, G.; Rossini, D.; Strini, G. *Principles of Quantum Computation and Information (A Comprehensive*
113 *Textbook)*; World Scientific Singapore, 2019.
- 114 5. Cornfeld, I.P.; Fomin, S.V.; Sinai, Y.G. *Ergodic Theory*; Springer New York, NY, 1982.
115 doi:https://doi.org/10.1007/978-1-4615-6927-5.
- 116 6. Gu, Y. Evidences of classical and quantum chaos in the time evolution of nonequilibrium ensembles. *Physics*
117 *Letters A* **1990**, *149*, 95–100. doi:https://doi.org/10.1016/0375-9601(90)90532-S.
- 118 7. Ford, J.; Mantica, G.; Ristow, G.H. The Arnol'd cat: Failure of the correspondence principle. *Physica D: Nonlinear*
119 *Phenomena* **1991**, *50*, 493–520. doi:https://doi.org/10.1016/0167-2789(91)90012-X.

- 120 8. Gu, Y.; Wang, J. Time evolution of coarse-grained entropy in classical and quantum motions of strongly chaotic
121 systems. *Physics Letters A* **1997**, *229*, 208–216. doi:https://doi.org/10.1016/S0375-9601(97)00194-1.
- 122 9. Pattanayak, A.K.; Brumer, P. Chaos and Lyapunov exponents in classical and quantal distribution dynamics.
123 *Phys. Rev. E* **1997**, *56*, 5174–5177. doi:10.1103/PhysRevE.56.5174.
- 124 10. Sokolov, V.V.; Zhirnov, O.V.; Benenti, G.; Casati, G. Complexity of quantum states and reversibility of quantum
125 motion. *Phys. Rev. E* **2008**, *78*, 046212. doi:10.1103/PhysRevE.78.046212.
- 126 11. Benenti, G.; Casati, G. How complex is quantum motion? *Phys. Rev. E* **2009**, *79*, 025201.
127 doi:10.1103/PhysRevE.79.025201.
- 128 12. Balachandran, V.; Benenti, G.; Casati, G.; Gong, J. Phase-space characterization of complexity in quantum
129 many-body dynamics. *Phys. Rev. E* **2010**, *82*, 046216. doi:10.1103/PhysRevE.82.046216.
- 130 13. Prosen, T. Complexity and nonseparability of classical Liouvillian dynamics. *Phys. Rev. E* **2011**, *83*, 031124.
131 doi:10.1103/PhysRevE.83.031124.
- 132 14. Benenti, G.; Carlo, G.G.; Prosen, T. Wigner separability entropy and complexity of quantum dynamics. *Phys.*
133 *Rev. E* **2012**, *85*, 051129. doi:10.1103/PhysRevE.85.051129.
- 134 15. Qin, P.; Wang, W.g.; Benenti, G.; Casati, G. Complexity and instability of quantum motion near a quantum
135 phase transition. *Phys. Rev. E* **2014**, *89*, 032120. doi:10.1103/PhysRevE.89.032120.
- 136 16. Rozenbaum, E.B.; Ganeshan, S.; Galitski, V. Lyapunov Exponent and Out-of-Time-Ordered Correlator's Growth
137 Rate in a Chaotic System. *Phys. Rev. Lett.* **2017**, *118*, 086801. doi:10.1103/PhysRevLett.118.086801.
- 138 17. Rammensee, J.; Urbina, J.D.; Richter, K. Many-Body Quantum Interference and the Saturation of
139 Out-of-Time-Order Correlators. *Phys. Rev. Lett.* **2018**, *121*, 124101. doi:10.1103/PhysRevLett.121.124101.
- 140 18. García-Mata, I.; Saraceno, M.; Jalabert, R.A.; Roncaglia, A.J.; Wisniacki, D.A. Chaos Signatures in the
141 Short and Long Time Behavior of the Out-of-Time Ordered Correlator. *Phys. Rev. Lett.* **2018**, *121*, 210601.
142 doi:10.1103/PhysRevLett.121.210601.
- 143 19. Bergamasco, P.D.; Carlo, G.G.; Rivas, A.M.F. Out-of-time ordered correlators, complexity, and entropy in
144 bipartite systems. *Phys. Rev. Research* **2019**, *1*, 033044. doi:10.1103/PhysRevResearch.1.033044.
- 145 20. Prakash, R.; Lakshminarayan, A. Scrambling in strongly chaotic weakly coupled bipartite systems: Universality
146 beyond the Ehrenfest timescale. *Phys. Rev. B* **2020**, *101*, 121108. doi:10.1103/PhysRevB.101.121108.
- 147 21. Wang, J.; Benenti, G.; Casati, G.; Wang, W.g. Complexity of quantum motion and
148 quantum-classical correspondence: A phase-space approach. *Phys. Rev. Research* **2020**, *2*, 043178.
149 doi:10.1103/PhysRevResearch.2.043178.
- 150 22. Bennett, C.H.; Bernstein, H.J.; Popescu, S.; Schumacher, B. Concentrating partial entanglement by local
151 operations. *Phys. Rev. A* **1996**, *53*, 2046–2052. doi:10.1103/PhysRevA.53.2046.
- 152 23. Bianchi, E.; Hackl, L.; Yokomizo, N. Linear growth of the entanglement entropy and the Kolmogorov-Sinai
153 rate. *Journal of High Energy Physics* **2018**, *2018*, 25. doi:10.1007/JHEP03(2018)025.
- 154 24. Miller, P.A.; Sarkar, S. Signatures of chaos in the entanglement of two coupled quantum kicked tops. *Phys. Rev.*
155 *E* **1999**, *60*, 1542–1550. doi:10.1103/PhysRevE.60.1542.
- 156 25. Fujisaki, H.; Miyadera, T.; Tanaka, A. Dynamical aspects of quantum entanglement for weakly coupled kicked
157 tops. *Phys. Rev. E* **2003**, *67*, 066201. doi:10.1103/PhysRevE.67.066201.
- 158 26. Petitjean, C.; Jacquod, P. Lyapunov Generation of Entanglement and the Correspondence Principle. *Phys. Rev.*
159 *Lett.* **2006**, *97*, 194103. doi:10.1103/PhysRevLett.97.194103.
- 160 27. Husimi, K. Some formal properties of the density matrix. *Proc. Phys. Math. Soc. Jpn.* **1940**, *22*, 264.
- 161 28. Emerson, J.; Ballentine, L. Characteristics of quantum-classical correspondence for two interacting spins. *Phys.*
162 *Rev. A* **2001**, *63*, 052103. doi:10.1103/PhysRevA.63.052103.
- 163 29. Haake, F.; Gnutzmann, S.; Kuś, M. *Quantum Signatures of Chaos*; Springer-Verlag: Heidelberg, 2018.

166 © 2022 by the authors. Submitted to *Journal Not Specified* for possible open access publication under the terms and
167 conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).