

The hadronic running of the electroweak couplings from lattice QCD

Marco Cè,^{*a*,*} Antoine Gérardin,^{*b*} Georg von Hippel,^{*c*} Harvey B. Meyer,^{*c*,*d*} Kohtaroh Miura,^{*c*,*d*,*e*} Konstantin Ottnad,^{*c*} Andreas Risch,^{*f*} Teseo San José^{*g*} and Hartmut Wittig^{*c*,*d*}

- ^bAix-Marseille Université, Université de Toulon, CNRS, CPT, Marseille, France
- ^c PRISMA⁺ Cluster of Excellence and Institut für Kernphysik, Johannes Gutenberg-Universität Mainz, 55099 Mainz, Germany
- ^d Helmholtz-Institut Mainz, Johannes Gutenberg-Universität Mainz, 55099 Mainz, Germany and GSI Helmholtzzentrum für Schwerionenforschung, Planckstraße 1, 64291 Darmstadt, Germany
- ^eKEK Theory Center, High Energy Accelerator Research Organization, Tsukuba, Japan
- ^f John von Neumann-Institut für Computing NIC, Deutsches Elektronen-Synchrotron DESY, Platanenallee 6, 15738 Zeuthen, Germany
- ^gLaboratoire de Physique des 2 Infinis Irène Joliot-Curie, CNRS/IN2P3, Université Paris-Saclay, 91405 Orsay, France

E-mail: marcoce@itp.unibe.ch

The energy dependency (running) of the strength of electromagnetic interactions α plays an important role in precision tests of the Standard Model. The running of the former to the Z pole is an input quantity for global electroweak fits, while the running of the mixing angle is susceptible to the effects of Beyond Standard Model physics, particularly at low energies. We present a computation of the hadronic vacuum polarization (HVP) contribution to the running of these electroweak couplings at the non-perturbative level in lattice QCD, in the space-like regime up to Q^2 momentum transfers of 7 GeV². This quantity is also closely related to the HVP contribution to the muon g - 2. We observe a tension of up to 3.5 standard deviation between our lattice results for $\Delta \alpha_{had}^{(5)}(-Q^2)$ and estimates based on the *R*-ratio for Q^2 in the 3 to 7 GeV² range. The tension is, however, strongly diminished when translating our result to the Z pole, by employing the Euclidean split technique and perturbative QCD, which yields $\Delta \alpha_{had}^{(5)}(M_Z^2) = 0.02773(15)$. This value agrees with results based on the *R*-ratio within the quoted uncertainties, and can be used as an alternative to the latter in global electroweak fits.

41st International Conference on High Energy physics - ICHEP20226-13 July, 2022Bologna, Italy

arXiv:2211.11401v2 [hep-lat] 16 Dec 2022

^aAlbert Einstein Center for Fundamental Physics (AEC) and Institut für Theoretische Physik, Universität Bern, Sidlerstrasse 5, 3012 Bern, Switzerland

^{*}Speaker

[©] Copyright owned by the author(s) under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License (CC BY-NC-ND 4.0).

1. Introduction

In the Standard Model (SM) of particle physics, the strength of electromagnetic interactions is represented by the quantum electrodynamics (QED) coupling α that depends on the energy at which electromagnetism is probed. At low energies the value of $\alpha^{-1} = 137.035\,999\,139(31)$ is known experimentally to better than a part per billion. At higher energies, such as at the Z pole, the most precise determination is obtained by applying the renormalization group running. Choosing to work in the *on-shell* scheme, one has

$$\alpha(q^2) = \frac{\alpha}{1 - \Delta \alpha(q^2)},\tag{1}$$

where the major contribution to the error on the running $\Delta \alpha(q^2)$ comes from the low-energy hadronic component $\Delta \alpha_{had}(q^2)$, which is proportional to the subtracted hadronic vacuum polarization (HVP) function of an off-shell photon propagator

$$\Delta \alpha_{\rm had}(q^2) = 4\pi \alpha \,{\rm Re}\,\bar{\Pi}(q^2) \qquad \bar{\Pi}(q^2) = \Pi(q^2) - \Pi(0). \tag{2}$$

A precise determination of the HVP function and in turn of $\Delta \alpha_{had}(q^2)$ can be obtained from experimental hadronic cross-section data encoded in the well-known ratio R(s) [1–3]. Alternatively, the HVP function $\Pi(-Q^2)$ at space-like $Q^2 > 0$ momentum transfers can be computed on the lattice [4–7]. In combination with the Euclidean split technique [8], this can be used to provide a value of $\Delta \alpha_{had}(M_Z^2)$ independent of R(s) data, which is the main result presented here [7].

The interests in computing $\overline{\Pi}(-Q^2)$ on the lattice is further motivated by the connection with the anomalous magnetic moment (g - 2) of the muon, and the long-standing tension between the muon g - 2 theory prediction and experimental results [9]. In particular, the leading-order muon g - 2 HVP contribution can be expressed as an integral over $\overline{\Pi}(-Q^2)$ at space-like Q^2 with a positive-definite integration kernel

$$a_{\mu}^{\text{HVP,LO}} = 4\alpha^2 \int_0^1 dx \, (1-x)\bar{\Pi}(-Q^2), \qquad Q^2 = \frac{x^2 m_{\mu}^2}{1-x},$$
 (3)

a fact that is at the core of the MUonE experiment proposal for an independent measurement of $a_{\mu}^{\text{HVP,LO}}$ in *t*-channel scattering experiments [10]. The precise lattice determination of $a_{\mu}^{\text{HVP,LO}}$ by the Budapest-Marseille-Wuppertal collaboration (BMWc) [11] points to a larger value—in tension with the data-driven determination and closer to solving the g - 2 puzzle—that according to eq. (3) has to be correlated with an increase of $\overline{\Pi}(-Q^2)$ and $\Delta \alpha_{\text{had}}(-Q^2)$. The evidence is even stronger when lattice data for the so-called HVP window is considered [12, 13]. At the same time, this increase implies an increase of R(s) that puts these lattice results in tension with the experimental measurements of the hadronic cross-section.

Computing $\overline{\Pi}(-Q^2)$ from first principles on the lattice can help shed light on this tension, albeit with limited energy resolution on R(s) due to the lattice data being limited to space-like Q^2 . Moreover, the lattice determination of $\Delta \alpha_{had}(M_Z^2)$ can be used as an alternative input to the data-driven estimate in global EW fits.

2. Lattice calculation

The lattice computation is performed using the time-momentum representation (TMR) method [14]. For each selected value of Q^2 up to 7 GeV², we compute the integral

$$\bar{\Pi}(-Q^2) = \int_0^\infty dt \, G(t) \left[t^2 - \frac{4}{Q^2} \sin^2\left(\frac{Qt}{2}\right) \right],$$
(4)

where the continuum expression for the kernel is used and G(t) is the Euclidean-space correlator of two electromagnetic currents at imaginary time separation t computed on the lattice

$$G(t) = -\frac{1}{3} \int d^3x \sum_{k=1}^{3} \langle j_k(t, \vec{x}) j_k(0) \rangle, \qquad j_\mu = \frac{2}{3} \bar{u} \gamma_\mu u - \frac{1}{3} \bar{d} \gamma_\mu d - \frac{1}{3} \bar{s} \gamma_\mu s + \frac{2}{3} \bar{c} \gamma_\mu c + \dots$$
(5)

We use a set of $N_f = 2 + 1$ dynamical quark ensembles of gauge field configurations from the Coordinated Lattice Simulations (CLS) effort [15]. On these ensembles, the up and down quarks are mass degenerate and the charm quark contribution is present in the valence sector only. The lattice current is renormalized and both the action and the observables are non-perturbatively O(a)-improved to ensure discretization effects starting at $O(a^2)$, where *a* is the lattice spacing. We refer the reader to ref. [7] for more details.

Our set of ensembles covers four lattice spacings from $a \approx 0.086$ fm to ≈ 0.050 fm and a range of pion and kaon masses from the SU(3)-flavour-symmetric point at $m_{\pi} = m_K \approx 415$ MeV to physical ones along a trajectory that keeps $m_K^2 + m_{\pi}^2/2$ approximately constant. This allow us to reliably extrapolate $\Delta \alpha_{had}(-Q^2) = 4\pi \alpha \overline{\Pi}(-Q^2)$ for a set of values of Q^2 to the "physical" point corresponding to an isospin-symmetric world.

At $Q^2 = 5 \text{ GeV}^2$, the result at the physical point is

$$\Delta \alpha_{\rm had}(-5\,{\rm GeV}^2) = 0.007\,16(9). \tag{6}$$

On top of the statistical error on the lattice data, we include in the final error budget a systematic uncertainty. This is composed of the systematic error from varying the fit model, the scale-setting error, the systematics from missing charm sea-quark loops, and the one from missing isospin-breaking effects.

3. Comparison with phenomenology

In figure 1 our lattice results are compared to other lattice results from BMWc (upper part) [6, 11], as well as with three data-driven estimates (lower part) [1–3]. We observe a small tension with the BMWc results, especially in the I = 1 component at small value of Q^2 , while at larger Q^2 the results are mostly compatible. The tension is instead significant when our result is compared with the data-driven estimates, up to 3.5 standard deviations.

To estimate $\Delta \alpha_{had}^{(5)}(M_Z^2)$ we employ the so-called Euclidean split technique (or Adler function approach) [8], which consists in splitting the contribution to the running to M_Z as

$$\Delta \alpha_{\rm had}^{(5)}(M_Z^2) = \Delta \alpha_{\rm had}^{(5)}(-Q_0^2) + [\Delta \alpha_{\rm had}^{(5)}(-M_Z^2) - \Delta \alpha_{\rm had}^{(5)}(-Q_0^2)] + [\Delta \alpha_{\rm had}^{(5)}(M_Z^2) - \Delta \alpha_{\rm had}^{(5)}(-M_Z^2)].$$
(7)

In the r.h.s., the first term is evaluated on the lattice and it is the main result presented in the previous section. The result at $Q_0^2 = 5 \text{ GeV}^2$ is given in eq. (6). The second term involves the running at




Figure 1: Left, upper panel: ratio of the hadronic running $\Delta \alpha_{had}$ computed by BMWc [6] divided by our results, for five different momenta. Left, lower panel: the total hadronic running $\Delta \alpha_{had}^{(5)}$ from various phenomenological estimates [1–3] and the lattice result of ref. [6], normalized by the result of this work. Right: Compilation of results for the four-flavor $\Delta \alpha_{had}$ lattice computations [6, 11] (above) and the five-flavor $\Delta \alpha_{had}^{(5)}$ phenomenological estimates (below) at selected values of Q^2 .

space-like momenta up to the Z pole. This can either be estimated using the data-driven method from the hadronic cross section [1–3], or for large enough Q_0^2 it can be computed in perturbative QCD (pQCD). The result based on pQCD is our preferred choice since it allows us to be independent of experimental R(s) input. Choosing the threshold energy $Q_0^2 = 5 \text{ GeV}^2$, the estimate obtained using the pQCDAdler code¹ is $[\Delta \alpha_{had}^{(5)}(-M_Z^2) - \Delta \alpha_{had}^{(5)}(-Q_0^2)]_{pQCD} = 0.020528(107)$. Finally, the third term in the r.h.s. of eq. (7) has been estimated in pQCD in ref. [3] and amounts to a small and precise contribution, $[\Delta \alpha_{had}^{(5)}(M_Z^2) - \Delta \alpha_{had}^{(5)}(-M_Z^2)]_{pQCD} = 0.000045(2)$.

Combining these results, the hadronic contribution to the running of the electromagnetic coupling α at the Z pole is

$$\Delta \alpha_{\rm had}^{(5)}(M_Z^2)|_{\rm lat. + pQCD} = 0.027\,73(9)_{\rm lat}(2)_b(12)_{\rm pQCD} = 0.027\,73(15),\tag{8}$$

where the first error is the uncertainty of our lattice estimate, while the second error accounts for the neglected contribution form bottom quark effects, and the third error is the one on the second term in the r.h.s. of eq. (7). The total error is augmented by the maximum deviation obtained by varying the threshold energy Q_0^2 in the interval (5 ± 2) GeV², which has a negligible effect.

Our results are compared in figure 2 with other estimates using the standard dispersive approach [1–3] and global EW fits [17–21]. We observe that the tension with the data-driven results is strongly diminished, mostly due to the major contribution to the error of the running from Q_0^2 to the Z pole. Of course, this contribution is fully correlated between lattice and data-driven results, and the tension in the $Q^2 \leq 5 \text{ GeV}^2$ contribution is only hidden. However, the results from global EW fits show a pull of a bit more than one standard deviation between the indirect determination of $\Delta \alpha_{had}^{(5)}(M_Z^2)$ and our result in eq. (8), indicating no inconsistency between lattice results and the global EW fits, as also observed in ref. [21].

http://www-com.physik.hu-berlin.de/~fjeger/software.html




Figure 2: Compilation of results for $\Delta \alpha_{had}^{(5)}(M_Z^2)$. The first two data points (red symbols) are the lattice results of ref. [7]. Green circles denote results based the data-driven method in, from top to bottom, refs. [1, 16], [2], and [3]. The estimate based on the Adler function in ref. [3] is shown as a green diamond. Blue symbols represent the results from global EW fits, published in refs. [17–21]. The upper triangle point from ref. [20] does not use the Higgs mass. The gray band represents our final result quoted in eq. (8).

4. Conclusions

We presented a computation of the hadronic contribution to the running of the electromagnetic coupling α . Our result is obtained on the lattice for space-like Q^2 up to $\approx 7 \text{ GeV}^2$ and it is slightly larger but still compatible with an earlier calculation by BMWc. However, there is a significant tension with the predictions based on the data driven method.

Combining our result obtained in the $Q^2 = (5 \pm 2) \text{ GeV}^2$ range with pQCD, we obtain an estimate for $\Delta \alpha_{had}^{(5)}(M_Z^2)$ that does not rely on experimental hadronic cross section data as input. This result is consistent with and of similar precision as estimates employing the data-driven approach. Moreover, we observe no significant tensions between our lattice result and global EW fits.

References

- [1] A. Keshavarzi, D. Nomura and T. Teubner, Muon g 2 and $\alpha(M_Z^2)$: a new data-based analysis, Phys. Rev. D 97 (2018) 114025 [1802.02995].
- [2] M. Davier, A. Hoecker, B. Malaescu and Z. Zhang, A new evaluation of the hadronic vacuum polarisation contributions to the muon anomalous magnetic moment and to $\alpha(m_Z^2)$, Eur. Phys. J. C 80 (2020) 241 [1908.00921].
- [3] F. Jegerlehner, α_{QED,eff}(s) for precision physics at the FCC-ee/ILC, CERN Yellow Reports: Monographs 3 (2020) 9.
- [4] F. Burger, K. Jansen, M. Petschlies and G. Pientka, *Leading hadronic contributions to the running of the electroweak coupling constants from lattice QCD*, *JHEP* 1511 (2015) 215 [1505.03283].
- [5] A. Francis, V. Gülpers, G. Herdoíza, H. Horch, B. Jäger, H.B. Meyer et al., Study of the

hadronic contributions to the running of the QED coupling and the weak mixing angle, PoS LATTICE2015 (2015) 110 [1511.04751].

- [6] Sz. Borsányi, Z. Fodor, C. Hoelbling, T. Kawanai, S. Krieg, L. Lellouch et al., (BUDAPEST-MARSEILLE-WUPPERTAL COLLABORATION), *Hadronic vacuum polarization contribution to the anomalous magnetic moments of leptons from first principles, Phys. Rev. Lett.* **121** (2018) 022002 [1711.04980].
- [7] M. Cè, A. Gérardin, G. von Hippel, H.B. Meyer, K. Miura, K. Ottnad et al., *The hadronic running of the electromagnetic coupling and the electroweak mixing angle from lattice QCD*, *JHEP* 2208 (2022) 220 [2203.08676].
- [8] F. Jegerlehner, *The running fine structure constant* $\alpha(E)$ *via the Adler function, Nucl. Phys. B Proc. Suppl.* **181-182** (2008) 135 [0807.4206].
- [9] T. Aoyama, N. Asmussen, M. Benayoun, J. Bijnens, T. Blum, M. Bruno et al., *The anomalous magnetic moment of the muon in the Standard Model*, *Phys. Rep.* 887 (2020) 1 [2006.04822].
- [10] G. Abbiendi, C.M. Carloni Calame, U. Marconi, C. Matteuzzi, G. Montagna, O. Nicrosini et al., *Measuring the leading hadronic contribution to the muon g – 2 via μe scattering, Eur. Phys. J. C* 77 (2017) 139 [1609.08987].
- [11] Sz. Borsányi, Z. Fodor, J.N. Guenther, C. Hoelbling, S.D. Katz, L. Lellouch et al., (BUDAPEST-MARSEILLE-WUPPERTAL COLLABORATION), *Leading hadronic contribution to the muon magnetic moment from lattice QCD*, *Nature* **593** (2021) 51 [2002.12347].
- [12] M. Cè, A. Gérardin, G. von Hippel, R.J. Hudspith, S. Kuberski, H.B. Meyer et al., Window observable for the hadronic vacuum polarization contribution to the muon g – 2 from lattice QCD, Phys. Rev. D 106 (2022) 114502 [2206.06582].
- [13] C. Alexandrou, S. Bacchio, P. Dimopoulos, J. Finkenrath, R. Frezzotti, G. Gagliardi et al., Lattice calculation of the short and intermediate time-distance hadronic vacuum polarization contributions to the muon magnetic moment using twisted-mass fermions, 2206.15084.
- [14] D. Bernecker and H.B. Meyer, Vector correlators in lattice QCD: Methods and applications, Eur. Phys. J. A 47 (2011) 148 [1107.4388].
- [15] M. Bruno, D. Djukanovic, G.P. Engel, A. Francis, G. Herdoiza, H. Horch et al., *Simulation of QCD with* $N_{\rm f} = 2 + 1$ *flavors of non-perturbatively improved Wilson fermions*, *JHEP* **1502** (2015) 043 [1411.3982].
- [16] A. Keshavarzi, D. Nomura and T. Teubner, g 2 of charged leptons, $\alpha(M_Z^2)$, and the hyperfine splitting of muonium, Phys. Rev. D 101 (2020) 014029 [1911.00367].
- [17] J. Haller, A. Hoecker, R. Kogler, K. Mönig, T. Peiffer and J. Stelzer, (THE GFITTER GROUP), Update of the global electroweak fit and constraints on two-Higgs-doublet models, Eur. Phys. J. C 78 (2018) 675 [1803.01853].
- [18] A. Crivellin, M. Hoferichter, C.A. Manzari and M. Montull, *Hadronic vacuum polarization:* $(g 2)_{\mu}$ versus global electroweak fits, *Phys. Rev. Lett.* **125** (2020) 091801 [2003.04886].
- [19] A. Keshavarzi, W.J. Marciano, M. Passera and A. Sirlin, *Muon* g 2 and $\Delta \alpha$ connection, *Phys. Rev. D* **102** (2020) 033002 [2006.12666].
- [20] B. Malaescu and M. Schott, Impact of correlations between a_{μ} and α_{QED} on the EW fit, Eur. *Phys. J. C* **81** (2021) 46 [2008.08107].
- [21] J. de Blas, M. Ciuchini, E. Franco, A. Goncalves, S. Mishima, M. Pierini et al., *Global analysis of electroweak data in the standard model*, *Phys. Rev. D* **106** (2021) 033003 [2112.07274].