

# ON THE PRIME SELMER RANKS OF CYCLIC PRIME TWIST FAMILIES OF ELLIPTIC CURVES OVER GLOBAL FUNCTION FIELDS

SUN WOO PARK

**ABSTRACT.** Fix a prime number  $p$ . Let  $\mathbb{F}_q$  be a finite field of characteristic coprime to 2, 3, and  $p$ , which also contains the primitive  $p$ -th root of unity  $\mu_p$ . Based on the works by Swinnerton-Dyer [SD08] and Klagsbrun, Mazur, and Rubin [KMR14], we prove that the probability distribution of the sizes of prime Selmer groups over a family of cyclic prime twists of non-isotrivial elliptic curves over  $\mathbb{F}_q(t)$  satisfying a number of mild constraints conforms to the distribution conjectured by Bhargava, Kane, Lenstra, Poonen, and Rains [PR12, BKLJ<sup>+</sup>15] with explicit error bounds. The key tools used in proving these results are the Riemann hypothesis over global function fields, the Erdős-Kac theorem, and the geometric ergodicity of Markov chains.

## CONTENTS

1. Introduction	1
2. Remarks, outlines, and illustrations	3
3. Effective theorems from the Riemann hypothesis	9
4. Local cyclic prime twists of elliptic curves	14
5. Equidistribution of unramified characters and auxiliary primes	25
6. Proof of the main theorem	31
Acknowledgements	39
References	39

## 1. INTRODUCTION

The objective of this paper is to compute the probability distribution of the dimensions of prime Selmer groups over a family of cyclic prime twists of non-isotrivial elliptic curves over  $\mathbb{F}_q(t)$  satisfying a number of mild constraints.

**Theorem 1.1** (Main Theorem over  $\mathbb{F}_q(t)$ ). *Fix a prime number  $p$ . Let  $K = \mathbb{F}_q(t)$  be a global function field whose characteristic is coprime to 2, 3, and  $p$ . Let  $E : y^2 = F(x) = x^3 + Ax + B$  be an elliptic curve over  $K$  which satisfies the following conditions.*

- (1)  *$E$  is non-isotrivial.*
- (2)  *$E$  contains a place  $\infty$  of split multiplicative reduction.*
- (3) *The constant field  $\mathbb{F}_q$  contains the  $p$ -th root of unity  $\mu_p$ , i.e.  $q \equiv 1 \pmod{p}$ .*
- (4) *The Galois group  $\text{Gal}(K(E[p])/K)$  contains  $\text{SL}_2(\mathbb{F}_p)$ , i.e. it is isomorphic to  $\text{SL}_2(\mathbb{F}_p) \rtimes T$  for some finite cyclic subgroup  $T$ .*

Let  $F_n(\mathbb{F}_q)$  be the set of polynomials of degree  $n$  over  $\mathbb{F}_q$ . Given a polynomial  $f \in F_n(\mathbb{F}_q)$ , let  $\chi_f \in \text{Hom}(\text{Gal}(\overline{\mathbb{F}_q(t)}/\mathbb{F}_q(t)), \mu_p)$  be a cyclic order  $p$  character, with  $L^{\chi_f}/\mathbb{F}_q(t) := \mathbb{F}_q(\sqrt[p]{f}, t)/\mathbb{F}_q(t)$  the associated cyclic  $p$ -extension of  $\mathbb{F}_q(t)$ . Denote by  $E^{\chi}$ , a cyclic order  $p$  twist of  $E$ , the kernel of the norm map  $N : \text{Res}_{\mathbb{F}_q(t)}^{L^{\chi}} E \rightarrow E$ , where  $\text{Res}_K^L E$  denotes the Weil restriction of scalars of  $E/K$  with

respect to the field extension  $L/K$ . For each non-negative integer  $j$ , denote by

$$\mathbb{P} \left[ \dim_{\mathbb{F}_p} \text{Sel}_{(1-\zeta_p)}(E^{X_f}) = j \mid f \in F_n(\mathbb{F}_q) \right] := \frac{\#\{f \in F_n(\mathbb{F}_q) \mid \dim_{\mathbb{F}_p} \text{Sel}_{(1-\zeta_p)}(E^{X_f}) = j, L^{X_f} := \mathbb{F}_q(\sqrt[p]{f}, t)\}}{\#F_n(\mathbb{F}_q)} \quad (1)$$

the probability that the dimensions of  $(1 - \zeta_p)$ -Selmer groups of finitely many cyclic order- $p$  twists of  $E$  over  $f \in F_n(\mathbb{F}_q)$  are equal to  $j$ . Fix a positive constant  $0 < \rho < 1$ . Then for sufficiently large  $n$ , there exist fixed constants  $B > 0$  and  $0 < \alpha < \min\left(\rho \ln \rho + 1 - \rho, -\rho \ln\left(1 - \frac{1}{|T|} \frac{p}{p^2-1}\right)\right)$  such that

$$\left| \mathbb{P} \left[ \dim_{\mathbb{F}_p} \text{Sel}_{(1-\zeta_p)}(E^{X_f}) = j \mid f \in F_n(\mathbb{F}_q) \right] - \left( \prod_{m \geq 0} \frac{1}{1 + p^{-m}} \right) \left( \prod_{m=1}^j \frac{p}{p^m - 1} \right) \right| < Bn^{-\alpha}.$$

The main theorem, in particular, allows us to compute the distribution of 2-Selmer ranks of quadratic twist families of non-isotrivial elliptic curves  $E$  over any global function field  $K = \mathbb{F}_q(t)$  under certain mild conditions, with maximum attainable error bounds of order given by approximately  $n^{-0.3339976579}$ .

**1.1. Preliminary.** Given a universal family of elliptic curves over a global field  $K$ , Bhargava, Kane, Lenstra, Poonen, and Rains made a conjecture on the distribution of  $p$ -selmer groups of elliptic curves for some prime number  $p$ .

**Conjecture 1.2** (Conjecture 1.1, [PR12] [BKLJ<sup>+</sup>15]). *Let  $K$  be a fixed global field of characteristic coprime to 2 and 3. Let  $p$  be a prime number coprime to the characteristic of  $K$ . Then as  $E$  varies over all elliptic curves over  $K$ , ordered by height,*

$$\mathbb{P} \left[ \dim_{\mathbb{F}_p} \text{Sel}_p(E) = d \right] = \left( \prod_{j \geq 0} (1 + p^{-j})^{-1} \right) \left( \prod_{j=1}^d \frac{p}{p^j - 1} \right).$$

*In particular, the average size of  $\text{Sel}_p E$  over all elliptic curves  $E/K$  is  $p + 1$ . (Here, the probability is computed over finitely many elliptic curves  $y^2 = x^3 + Ax + B$ , where  $A, B \in K$  have bounded height  $B$ , and calculating the limit of the probability as  $B$  grows arbitrarily large.)*

Let  $K$  be a global field whose characteristic is coprime to 2 and 3. Fix a non-isotrivial abelian variety  $A/K$ . Let  $m \in \text{End}(A/K)$  be an isogeny of the abelian variety whose degree is coprime to the characteristic of  $K$ . The short exact sequence of group schemes

$$0 \rightarrow A[m] \rightarrow A \xrightarrow{m} A \rightarrow 0$$

induces the following commutative diagram,

$$\begin{array}{ccccccc} 0 & \longrightarrow & A(K)/mA(K) & \longrightarrow & H_{\text{ét}}^1(K, A[m]) & \longrightarrow & H_{\text{ét}}^1(K, A)[m] \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \prod_v A(K_v)/mA(K_v) & \longrightarrow & \prod_v H_{\text{ét}}^1(K_v, A[m]) & \longrightarrow & \prod_v H_{\text{ét}}^1(K_v, A)[m] \longrightarrow 0, \end{array}$$

where  $v$  varies over all places of  $K$ . The  $m$ -Selmer group of the abelian variety  $A$  is given by

$$\text{Sel}_m(A) := \text{Ker} \left( H_{\text{ét}}^1(K, A[m]) \rightarrow \prod_v H_{\text{ét}}^1(K_v, A)[m] \right). \quad (2)$$

The main theorem of this manuscript confirms the Bhargava-Kane-Lenstra-Poonen-Rains heuristics for certain cyclic order- $p$  twist families of  $p - 1$  dimensional abelian varieties over global function fields  $\mathbb{F}_q(t)$  such that  $q \equiv 1 \pmod{p}$ .

$$\mathbb{P}[\dim_{\mathbb{F}_2} \text{Sel}_2(E_f) = j \mid f \in \mathbb{F}_q[t], \deg f = n,] \approx \underbrace{\left[ \underbrace{\left( \frac{1}{3} + \frac{1}{2}M + \frac{1}{6}M^2 + \epsilon \right)}_{\text{Effective Chebotarev}} \right]}_{\substack{\text{Equidistribution} \\ \text{Auxiliary Place}}} \underbrace{(\delta)}_{\substack{\text{Effective Erdős-Kac} \\ \ln n}} \approx \underbrace{PR(j) + \epsilon'}_{\substack{\text{Geom. Ergodicity} \\ \text{Markov Chain}}}$$

FIGURE 1. A simplification of the ideas of the proof of Theorem 1.1 for the case where  $p = 2$ . The Markov chain  $M$  is constructed in Definition 4.8.

## 2. REMARKS, OUTLINES, AND ILLUSTRATIONS

**2.1. Some Notations.** We briefly introduce some (but not all) of the notations which will appear throughout the whole manuscript.

- We denote by  $F_n(\mathbb{F}_q)$  the set of polynomials of degree  $n$  over  $\mathbb{F}_q$ .
- Let  $E$  be an elliptic curve over the global function field  $\mathbb{F}_q(t)$  with minimal Weierstrass model given by  $E : y^2 = x^3 + Ax + B$ . Throughout this manuscript, we suppose that  $\text{Gal}(\mathbb{F}_q(t)(E[p])/\mathbb{F}_q(t)) \supset SL_2(\mathbb{F}_p)$ .
- Let  $\chi \in \text{Hom}(\text{Gal}(\overline{\mathbb{F}_q(t)}/\mathbb{F}_q(t)), \mu_p)$  be a cyclic order  $p$  character, with  $L^\chi/\mathbb{F}_q(t)$  the associated cyclic  $p$ -extension of  $\mathbb{F}_q(t)$ .
- Denote by  $E^\chi$  the kernel of the norm map  $N : \text{Res}_{\mathbb{F}_q(t)}^{L^\chi} E \rightarrow E$ , where  $\text{Res}_K^L E$  denotes the Weil restriction of scalars of  $E/K$  with respect to the field extension  $L/K$ . In particular,  $E^\chi$  is an abelian variety over  $K$  of dimension  $p - 1$ .
- Let  $\pi = (1 - \zeta_p)$  be the unique prime ideal of  $\mathbb{Z}[\text{Gal}(L^\chi/\mathbb{F}_q(t))]$  lying above the prime ideal  $(p) \subset \mathbb{Z}$ . The endomorphism ring  $\text{End}(E^\chi/K)$  contains the group ring  $\mathbb{Z}[\text{Gal}(L^\chi/\mathbb{F}_q(t))] \cong \mathbb{Z}[\zeta_p]$ . Therefore, the prime ideal  $\pi$  induces an endomorphism of the abelian variety  $E^\chi$ .
- If  $\chi$  is a quadratic character, then  $E^\chi$  corresponds to the quadratic twist of  $E$  twisted by  $\chi$ , and the ideal  $\pi$  defines the multiplication-by-2 isogeny over the quadratic twist of  $E$ .
- By using (2), we define the  $\pi$ -Selmer group  $\text{Sel}_\pi(E^\chi)$  of the cyclic order- $p$  twist of  $E$  (For  $p = 2$ , the Selmer group is the usual 2-Selmer group  $\text{Sel}_2(E^\chi)$  of the quadratic twist of  $E$ ).

**2.2. Key Ingredients.** The three key ingredients utilized in proving the main theorem are as follows, stated along with respective theorems in the manuscript.

- (1) **Effective Chebotarev Density Theorem** (Theorem 3.1, Proposition 4.5, Corollary 5.6): We use the effective Chebotarev density theorem to **(1)** construct a Markov chain  $M$  which governs the variation of  $\pi$ -Selmer groups with respect to cyclic prime twists (as shown in [KMR14]), and **(2)** establish a correspondence between variations of  $\pi$ -Selmer groups twisted by global cyclic characters and variations of those twisted by local cyclic characters by utilizing equidistribution of restriction of global characters to cartesian products of local characters.
- (2) **Effective Erdős-Kac Theorem** (Theorem 3.7, Proposition 5.4): We use the effective Erdős-Kac Theorem to determine the number of steps one needs to apply the constructed Markov chain to obtain the probability distribution of  $\pi$ -Selmer groups. In fact, the number of distinct irreducible factors of a polynomial  $f \in \mathbb{F}_q[t]$  of degree at least  $n^{\frac{1}{3}}$  is the number of steps we will apply the constructed Markov chain to compute the desired distribution.
- (3) **Geometric Convergence of Markov Chains** (Corollary 4.14): We explicitly compute the geometric rate of convergence of the constructed Markov chain to observe how fast the Markov chain converges to the desired Bhargava-Kane-Lenstra-Poonen-Rains distribution of  $\pi$ -Selmer groups.

Figure 1 illustrates a simplified strategy to prove the main theorem, along with respective locations on where the main ingredients required for proving the theorem will be used.

**2.3. Some Remarks.** Let us recall the statement of Theorem 1.1 and the constraints imposed over the choice of the elliptic curve  $E/\mathbb{F}_q(t)$ .

**[Conditions]**

- (1)  $E$  is non-isotrivial.
- (2)  $E$  contains a place  $\infty$  of split multiplicative reduction.
- (3) The constant field  $\mathbb{F}_q$  contains the  $p$ -th root of unity  $\mu_p$ , i.e.  $q \equiv 1 \pmod{p}$ .
- (4) The Galois group  $\text{Gal}(K(E[p])/K)$  contains  $\text{SL}_2(\mathbb{F}_p)$ , i.e. it is isomorphic to  $\text{SL}_2(\mathbb{F}_p) \rtimes T$  for some finite cyclic subgroup  $T$ .

**[Result]**

Let  $D$  be a constant

$$D := \min \left( \rho \ln \rho + 1 - \rho, -\rho \ln \left( 1 - \frac{1}{|T|} \frac{p}{p^2 - 1} \right) \right) \quad (3)$$

For sufficiently large  $n$ , there exist fixed constants  $B > 0$  and  $0 < \alpha < D$  such that

$$\left| \mathbb{P} \left[ \dim_{\mathbb{F}_p} \text{Sel}_{(1-\zeta_p)}(E^{\chi_f}) = j \mid f \in F_n(\mathbb{F}_q) \right] - \left( \prod_{m \geq 0} \frac{1}{1 + p^{-m}} \right) \left( \prod_{m=1}^j \frac{p}{p^m - 1} \right) \right| < Bn^{-\alpha}.$$

**Remark 2.1.** Note that for each prime  $p$ , the constant

$$D := \min \left( \rho \ln \rho + 1 - \rho, -\rho \ln \left( 1 - \frac{1}{|T|} \frac{p}{p^2 - 1} \right) \right) \quad (4)$$

achieves its maximum if one chooses a constant  $0 < \rho < 1$  such that

$$\rho \ln \rho + 1 - \rho = -\rho \ln \left( 1 - \frac{1}{|T|} \frac{p}{p^2 - 1} \right). \quad (5)$$

We will explore in this manuscript that the error bounds are determined from the application of the effective Chebotarev density theorem, the effective Erdős-Kac theorem (LHS), and the geometric convergence rate of Markov chains (RHS). In fact, the two error terms outlined in (5) originate from the effective Erdős-Kac theorem and the geometric convergence rate of Markov chains, whereas the error terms originating from the effective Chebotarev density theorem is negligible compared to the other two error terms.

For  $p = 2$ , the probability distribution of the size of 2-Selmer groups of quadratic twists of  $E$  over  $\mathbb{F}_q(t)$  converges to the Bhargava-Kane-Lenstra-Poonen-Rains distribution [PR12, BKLJ<sup>+</sup>15] with maximum attainable error bounds of order given by approximately  $n^{-0.3339976579}$ .

**Remark 2.2.** The condition that  $E$  is non-isotrivial further implies that conditions (2) and (3) are obtainable after taking finite separable extension of any global function field  $K = \mathbb{F}_q(t)$  [BLV09, Proposition 3.4]. We note that the dimension of  $\pi$ -Selmer groups of an elliptic curve twisted by a cyclic order- $p$  character associated to a polynomial  $f \in F_n(\mathbb{F}_q)$  provides an upper bound on the difference of the rank of  $K(\sqrt[p]{f})$ -rational points of  $E$  and the rank of  $K$ -rational points of  $E$ :

$$\text{rk}(E(K(\sqrt[p]{f}))) - \text{rk}(E(K)) \leq \dim_{\mathbb{F}_p} \text{Sel}_{\pi}(E^{\chi_f}). \quad (6)$$

Theorem 1.1 hence implies that for any cyclic order- $p$  abelian extension over  $K$  satisfying the given conditions,

$$\lim_{n \rightarrow \infty} \mathbb{P} \left[ \text{rk}(E(K(\sqrt[p]{f}))) - \text{rk}(E(K)) \leq j \mid f \in F_n(\mathbb{F}_q) \right] \leq \sum_{j'=0}^j \left( \prod_{m \geq 0} \frac{1}{1 + p^{-m}} \right) \left( \prod_{m=1}^j \frac{p}{p^m - 1} \right). \quad (7)$$

In particular, for large enough values of prime  $p$ , the upper bound on the probability distribution of the differences in ranks behaves in a similar manner to Goldfeld's conjecture (or the minimalist conjecture), which states that half of the quadratic twist families of elliptic curves achieves rank 0, the other half achieves rank 1, and it is very unlikely that the quadratic twist families of elliptic curves achieves rank at least 2.

We warn the readers, however, that the given upper bound is not binding for any values of  $p \geq 3$  unlike the case for quadratic twist families of elliptic curves, as the  $p$ -torsion subgroup of the Tate-Shafarevich group of the abelian variety  $E^{\chi_f}$  is not of an even dimensional  $\mathbb{F}_p$ -vector space, as explicitly constructed by William Stein [Ste02] and discussed in detail by Howe [How01]. Specific conditions which can guarantee the Tate-shafarevich groups to be of even dimension are provided in [MR07, Chapter 6]. Indeed, there are conjectural statements by David, Fearnley, and Kisilevsky [DFK07] and Mazur and Rubin [MR19] who suggested that it is very unlikely that the ranks of the elliptic curves will increase by at least 1 with respect to cyclic order- $p$  extensions over  $\mathbb{Q}$ . The function field analogue was carefully studied in a recent work by Comeau-Lapointe, David, Lalin, and Li [CLDLL22], where they show that the conjecture fails for isotrivial cyclic twist families of elliptic curves, whereas numerical data suggests that the conjecture may hold for non-isotrivial cyclic twist families of elliptic curves.

**Remark 2.3.** The statements of the conjecture are known for certain large families of elliptic curves, such as the universal family of elliptic curves ordered by height, or quadratic twist families of elliptic curves ordered by the norm of the twist.

Suppose  $K = \mathbb{Q}$ . We list some previous studies which focused on computing the probability distribution of Selmer groups over certain families of elliptic curves.

- Bhargava and Shankar compute the first moments of 2,3,4 and 5-Selmer groups over the universal family of elliptic curves, see for example [BS15].
- Heath-Brown, Swinnerton-Dyer, and Kane compute the probability distribution of 2-Selmer groups over the quadratic twist families of elliptic curves with full 2-torsions and no cyclic subgroup of order 4 over  $\mathbb{Q}$  [HB94, SD08, Kan13].
- Klagsbrun, Mazur, and Rubin generalized the construction of Markov chains suggested by Swinnerton-Dyer [SD08] to compute the probability distribution of 2-Selmer groups over the quadratic twist families of elliptic curves with  $\text{Gal}(K(E[2])/K) = S_3$ . Note that the elliptic curves are ordered in a non-canonical manner using Fan structures. They obtain the probability distribution of prime Selmer groups over non-canonically ordered cyclic order- $p$  twist families of elliptic curves with  $\text{Gal}(K(E[p])/K) = SL_2(\mathbb{F}_p)$  as well [KMR14].
- Smith successfully calculates the probability distribution of 2-Selmer groups over quadratic twist families of elliptic curves of bounded height  $H$  except for some cases where  $E[2](\mathbb{Q}) = \mathbb{Z}/2\mathbb{Z}$ . As the upper bound on the height  $H$  grows to infinity, the error bounds of the probability distribution is given by an order of  $O(e^{-c(\log \log \log H)^{\frac{1}{4}}})$  for some constant  $c > 0$ . Smith utilizes Markov chains which govern the variations of kernel ranks of alternating square matrices comprised induced from Cassels-Tate pairings. Note that the Markov chains Smith utilized are different from those constructed by Swinnerton-Dyer and Klagsbrun, Mazur, and Rubin [Smi17, Smi20, Smi22b, Smi22a].
- The Markov chains suggested by Smith can be utilized to prove the Cohen-Lenstra heuristics on  $l^\infty$ -torsion subgroups of class groups of cyclic  $l$ -extensions of  $\mathbb{Q}$  (assuming the generalized

Riemann hypothesis) [KP21], and Stevenhagen's conjecture on the asymptotic behavior of the solubility of negative Pell equations [KP22].

Consider the case where  $K = \mathbb{F}_q(t)$  of characteristic coprime to 2 and 3. Previous studies computed the probability distribution of  $p$ -Selmer groups of families of elliptic curves over global function fields  $\mathbb{F}_q(t)$  under different conditions. Denote by  $\mathcal{M}_n(\mathbb{F}_q)$  a finite subfamily of elliptic curves  $E$  over  $\mathbb{F}_q(t)$  of a fixed height  $n$ . The height of an elliptic curve is determined by the degrees of coefficient terms of  $E$ . (Of course, the choice of the height depends on over which families of elliptic curves the probability distribution of 2-Selmer groups is computed.)

Given a non-negative integer  $j$ , denote by  $\mathbb{P}[\dim_{\mathbb{F}_2} \text{Sel}_p(E) = j \mid E \in \mathcal{M}_n(\mathbb{F}_q)]$  the probability that the dimensions of 2-Selmer groups of finitely many elliptic curves of fixed height  $n$  are equal to  $j$ . Below we list three probability distributions of 2-Selmer groups of elliptic curves that can be computed over global function fields:

$$\lim_{n \rightarrow \infty} \mathbb{P}[\dim_{\mathbb{F}_2} \text{Sel}_p(E) = j \mid E \in \mathcal{M}_n(\mathbb{F}_q)] \quad (8)$$

$$\lim_{q \rightarrow \infty} \lim_{n \rightarrow \infty} \mathbb{P}[\dim_{\mathbb{F}_2} \text{Sel}_p(E) = j \mid E \in \mathcal{M}_n(\mathbb{F}_q)] \quad (9)$$

$$\lim_{n \rightarrow \infty} \lim_{q \rightarrow \infty} \mathbb{P}[\dim_{\mathbb{F}_2} \text{Sel}_p(E) = j \mid E \in \mathcal{M}_n(\mathbb{F}_q)] \quad (10)$$

As before, we list some previous studies which focused on computing the desired probability distribution over  $\mathbb{F}_q(t)$ .

- For the second limit (large-height, then large- $q$  limit), Ho, Le Hung, and Ngo [QH14] compute the average size of 2-Selmer groups over the universal family of elliptic curves, whereas de Jong [dJ02] computes that of 3-Selmer groups over the same family.
- For the third limit (large-1 limit, then large-height), Feng, Landesman, and Rains [FLR20] prove that the third limit (large- $q$ , then large-height limit) is equal to the Poonen-rains distribution for any  $m$ -Selmer groups over universal families of elliptic curves, under the condition that  $q$  is coprime to  $2m$ . They propose a Markov chain constructed from random kernel models, which governs the variation of  $n$ -Selmer groups over global function fields  $\mathbb{F}_q(t)$ . Using this Markov chain, they successfully prove the Poonen-Rains conjecture for  $n$ -Selmer groups of universal families of elliptic curves under the large  $q$ -limit.
- The average size of  $m$ -Selmer groups of universal families of elliptic curves under the third limit were computed by Landesman [Lan21] over universal families of elliptic curves.
- The average size of  $p$ -Selmer groups of quadratic twist families of non-isotrivial elliptic curves under the third limit were computed by the author of this paper and Wang [PW21].
- The key ingredient behind computing these distributions is a careful and rigorous determination of images of monodromy over algebraic spaces whose geometric fibers parametrize  $p$ -Selmer groups over a prescribed family of elliptic curves [EVW16, Hal06].

Theorem 1.1 proves that the first limit (large-height limit) is equal to the Poonen-rains distribution for  $p = 2$  over quadratic twist families of elliptic curves.

**Remark 2.4.** We finally note that it is not always the case that the probability distribution of 2-Selmer groups over quadratic twist families of elliptic curves over a global field  $K$  can be formulated. For example, Klagsbrun and Lemke Oliver showed that more than half the quadratic twists of elliptic curves over number fields  $K$  with partial  $K$ -rational 2-torsion points (i.e.  $E[2](K) = \mathbb{Z}/2\mathbb{Z}$ ) and without any cyclic 4-isogeny over  $K$  have arbitrarily large 2-Selmer ranks [KLO15]. Wang extends their results to global function fields  $K = \mathbb{F}_q(t)$  in his Ph.D. thesis for arbitrary number of elements of the constant field  $\mathbb{F}_q$  [Wan21].

**2.4. Outline.** We present the paper as provided in Figure 2.4. Section 3 states the effective Chebotarev density theorem and Erdős-Kac theorem over global function fields. Section 4 constructs




FIGURE 2. An outline of the organization of the paper.

the Markov chain from [SD08, KMR14], prove its geometric ergodicity, and show the parity of the distribution of  $\pi$ -Selmer groups over cyclic order-twist families of elliptic curves. Section 5 reviews the equidistribution results on restriction of global characters to products of unramified local characters. Along the way, we will introduce the notion of splitting partition of polynomials and auxiliary places, which simplifies the correspondence between global cyclic twists and local cyclic twists of elliptic curves. Section 6 concludes the manuscript by incorporating the aforementioned ingredients to compute the desired probability distribution and its error bounds.

We first utilize the effective Erdős-Kac theorem (Section 3.2) and splitting partition of polynomials (Section 5.1) to demonstrate that computing the distribution of  $\pi$ -Selmer groups of elliptic curves twisted by cyclic order- $p$  characters associated to  $p$ -th power free polynomials of fixed degree  $n$  can be reduced to computing the moments of those twisted by local order  $p$  characters satisfying the following two constraints:

- (1) The number of distinct irreducible factors of a polynomial  $f$  of degree  $n$  is approximately equal to  $\ln n$ .
- (2) All distinct irreducible factors of such  $f$  are of sufficiently large degree (e.g., at least  $n^{\frac{1}{3}}$ .)

The latter problem of computing the variations of  $\pi$ -Selmer groups of elliptic curves twisted by certain global characters can be reformulated as a problem of computing those twisted by products of local characters using the equidistribution results for quadratic characters (Section 5.2). The problem on local quadratic twists of elliptic curves can be solved by constructing a Markov model governing the variation of  $\pi$ -Selmer groups with respect to a twist by a local quadratic character, as carefully studied in previous studies by Klagsbrun, Mazur, and Rubin [KMR14], see also Section 4.1 of the manuscript. A rigorous calculation on the error bounds of Markov model is obtainable from the effective Chebotarev density theorem (Section 3.1). To compute the variation of  $\pi$ -Selmer groups over elliptic curves twisted by products of local characters, we carefully study the composition of the previously constructed Markov model. The associated propagation of error bounds can be demonstrated to behave well by the geometric ergodicity of the Markov model (Section 4.2).




FIGURE 3. An illustration of the key insights presented in [KMR14]. All the three examples above are computed in Magma [BCP97].

**2.5. Exemplary Illustrations.** The contributions of Theorem 1.1 can be summarized as follows. Under the assumptions from Theorem 1.1:

- (1) Given a fixed prime  $p$ , the probability distribution converges to the Poonen-Rains distribution for any value of  $q \equiv 1 \pmod{p}$ , as long as  $q$  is coprime to 2 and 3.
- (2) The explicit error bounds of the probability distributions are obtainable.

The crucial insight of using Markov chains to obtain the probability distribution originates from the previous work by Swinnerton-Dyer [SD08]. Klagsbrun, Mazur, and Rubin [KMR14] further proved that there exists a Markov chain governing the variation of  $\pi$ -Selmer ranks under local order- $p$  twists, whose stationary distribution is equal to the Poonen-Rains distribution. We illustrate with three examples from Figure 3 to give a sense of what Klagsbrun, Mazur, and Rubin demonstrated in [KMR14]. Figure 3 demonstrates the percentage of primes with a given number of prime factors over the cubic extension  $L/\mathbb{Q}$  and a given dimension of the 2-Selmer groups of the associated twist of the elliptic curve  $E$ . We note that all three example are computed in Magma [BCP97], and one can compute examples of a similar manner by using elliptic curves  $E : y^2 = f(x)$  given an irreducible cubic polynomial with negative square-free discriminant  $f \in \mathbb{Q}[x]$ , see for example [Li19] for further discussions on the relations between 2-Selmer groups and 2-torsion subgroup of the class group of the cubic field extension  $L = \mathbb{Q}[x]/(f(x))$ . A naïve observation shows that the percentage of primes with 2 or 3 prime factors over the cubic extension  $L/\mathbb{Q}$  whose dimension of the 2-Selmer groups of the associated twist of the elliptic curve  $E$  can be obtained from the respective percentage of primes with 1 or 2 prime factors over  $L$ . For example, for the elliptic curve  $E^{[1]} : y^2 = x^3 + x - 1$ , the percentage of inert primes  $p$  in  $L$  such that  $\dim_{\mathbb{F}_2} \text{Sel}_2(E_p) = 1$ , which is close to 75%, seems to split into two equal percentages,  $\sim 37.5\%$ , to the percentage of primes  $p$  with 2 prime factors in  $L$  such that  $\dim_{\mathbb{F}_2} \text{Sel}_2(E_p) = 0$  or 2. Similar phenomena can be observed for different elliptic curves as well, regardless of the dimension of 2-Selmer groups or the rank of the elliptic curve. Klagsbrun, Mazur, and Rubin modeled these phenomena using a Markov chain shown on the right hand side of Figure 3 (which will be described in full generality in Section 4.2), which corresponds to twisting an elliptic curve by a non-inert prime factor over  $L$ . The color scheme shown on the right hand



side of Figure 3 heuristically illustrates the diffusion of probability distribution after applying the Markov chain.

Using this observation, Klagsbrun, Mazur, and Rubin computed the distribution of 2-Selmer groups over quadratic twist families of elliptic curves over the “fan structure”, a non-canonically ordered set of global characters whose restriction to cartesian products of local characters are ramified over finite set of places whose norms, when rearranged in an increasing order, strictly increases at a certain rate. The condition that the rearranged finite set of places must have strictly increasing norm is a crucial supposition required to ensure that one can utilize Markov chains to analyze the distribution of 2-Selmer groups. Indeed, one can observe that the three examples would have produced results deviating from those predicted by the Markov chain  $M$  had not the upper bound for the prime  $p$  been large enough. Even with the condition that  $p < 500,000$ , one can still observe the error terms between all three examples from Figure 3 and the probability distribution provided by the chain  $M$ . The inability to effectively quantify and bound the numerical deviations and the growth of the norms of consecutive places over number fields prevents us from extending the result obtained over the “fan structure” to the canonical order defined over the twist families of elliptic curves, see [KMR14, Section 9-11] for further details.

Nevertheless, the effective Chebotarev density theorem and Erdős-Kac theorem allow the Markov chain to govern the variation of  $\pi$ -Selmer ranks twisted by local characters with norm of order at least  $n^{\frac{1}{3}}$  (Proposition 4.5). For sufficiently large  $n$ , such conditions are satisfied for most proportions (but not all) of irreducible factors of a randomly chosen  $p$ -th power-free polynomial  $f \in F_n(\mathbb{F}_q)$  (Proposition 5.4). Using these two observations, one can show how much the large- $n$  limit of the probability distribution of  $\pi$ -Selmer groups perturbs from the stationary distribution of the Markov chain, which is the Bhargava-Kane-Lenstra-Poonen-Rains distribution. The validity of the generalized Riemann hypothesis is a key arithmetic difference between global function fields  $K = \mathbb{F}_q(t)$  and rational numbers  $K = \mathbb{Q}$ . This determines the error bounds for both the Chebotarev density theorem and the Erdős-Kac theorem over global function fields.

### 3. EFFECTIVE THEOREMS FROM THE RIEMANN HYPOTHESIS

We review some of the preliminary results on global function fields  $K$  which will be utilized in computing the probability distribution of prime Selmer groups associated to cyclic prime twists of elliptic curves. Given a place  $v$  over  $K$ , we denote by  $\text{Frob}_v$  the Frobenius element at  $v$ . Denote by  $g_L$  the genus of a finite separable field extension  $L/K$ .

**3.1. Effective Chebotarev density theorem.** The effective version of Chebotarev density theorem over global function fields can be formulated as follows:

**Theorem 3.1** (Effective Chebotarev density theorem). [FJ08, Proposition 6.4.8]

*Let  $L/K$  be a Galois extension of global function fields over  $\mathbb{F}_q(t)$ . Pick a conjugacy class  $C \subset G = \text{Gal}(L/K)$ . If the constant fields of  $L$  and  $K$  are both equal to  $\mathbb{F}_q$ , then*

$$\left| \left\{ v \text{ a place over } K \mid \text{Frob}_v \in C, \dim_{\mathbb{F}_q}(\mathcal{O}_K/v) = n \right\} - \frac{|C| q^n}{|G| n} \right| < \frac{2|C|}{n|G|} \left[ (|G| + g_L) q^{\frac{n}{2}} + |G|(2g_K + 1) q^{\frac{n}{4}} + (|G| + g_L) \right]$$

The constraint that the constant fields of  $L$  and  $K$  are identical allows one to reconstruct the counterpart of the Chebotarev density theorem with explicit error bounds for function fields. Suppose the constant field of  $L$ , say  $\mathbb{F}_{q'}$ , is a non-trivial extension of the constant field  $\mathbb{F}_q$  of  $K$ . Then to compute the equation stated in Theorem 3.1, one is required to compare whether the restriction of the conjugacy class  $C$  to  $\text{Gal}(\mathbb{F}_{q'}/\mathbb{F}_q)$  agrees with the  $n$ -th power of the arithmetic Frobenius

$\tau : x \mapsto x^q$  as a cyclic generator of  $\text{Gal}(\mathbb{F}_{q^t}/\mathbb{F}_q)$ . If not, then there are no places of degree  $n$  whose Frobenius element lives inside the conjugacy class  $C$ . For a detailed description of the effective Chebotarev density theorem over function fields, see for example [BSGKS20]. We note that Galois extensions of global function fields with non-trivial constant field extensions also satisfy the following equation:

$$\lim_{s \rightarrow 1^+} \frac{\sum_{v \text{ a place over } K} |\{\mathcal{O}_K/v\}|^{-s}}{\sum_{v \text{ a place over } K} |\{\mathcal{O}_K/v\}|^{-s}} = \frac{|C|}{|G|} \quad (11)$$

where  $s \rightarrow 1^+$  implies that  $s$  approaches 1 from above over the real values.

Using the explicit bounds obtained above, the density theorem can be obtained for any two conjugacy classes of the Galois group of the extension  $L/K$  of function fields.

**Corollary 3.2.** *Let  $L/K$  be a Galois extension of global function fields over  $\mathbb{F}_q(t)$ . Pick two subsets  $S, S' \subset G = \text{Gal}(L/K)$  stable under conjugation. If the constant fields of  $L$  and  $K$  are both equal to  $\mathbb{F}_q$ , then*

$$\begin{aligned} & \left| \frac{|\{v, \text{ a place over } K \mid \text{Frob}_v \in S, \dim_{\mathbb{F}_q}(\mathcal{O}_K/v) = n\}|}{|\{v, \text{ a place over } K \mid \text{Frob}_v \in S', \dim_{\mathbb{F}_q}(\mathcal{O}_K/v) = n\}|} - \frac{|S|}{|S'|} \right| \\ & < 4 \frac{|S|}{|S'|} (|G| + g_L + 2g_K) \left[ \frac{1}{q^{\frac{n}{2}} - q^{\frac{n}{4}} - 2(|G| + g_L + 2g_K)} \right] \end{aligned}$$

In particular, for  $n \geq 2 \frac{\ln 8 + \ln(|G| + g_L + 2g_K)}{\ln q}$ , we have

$$\left| \frac{|\{v, \text{ a place over } K \mid \text{Frob}_v \in S, \dim_{\mathbb{F}_q}(K/v) = n\}|}{|\{v, \text{ a place over } K \mid \text{Frob}_v \in S', \dim_{\mathbb{F}_q}(K/v) = n\}|} - \frac{|S|}{|S'|} \right| < 16 \frac{|S|}{|S'|} (|G| + g_L + 2g_K) q^{-\frac{n}{2}}$$

**Remark 3.3.** We note that Deligne's proof of the Weil conjectures determine the error bounds of the effective Chebotarev density theorem. We refer to [Ros02, Theorem 9.13B] for further discussions.

**3.2. Erdős-Kac Theorem.** Let  $m$  be an integer. We denote by  $w(m)$  the number of distinct irreducible factors of  $m$ . The Erdős-Kac Theorem states that the normal order of  $w(m)$  is  $\ln \ln m$ . Its analogue over global function fields  $K$  can be formulated as follows.

**Theorem 3.4** (Erdős-Kac Theorem for Function Fields). [Liu04, Theorem 1]

Denote by  $w(f)$  the number of distinct irreducible factors dividing a polynomial  $f \in F_n(\mathbb{F}_q)$  of degree  $n$ . Then for any  $a \in \mathbb{R}$ ,

$$\lim_{n \rightarrow \infty} \frac{\#\left\{f \in F_n(\mathbb{F}_q) \mid \frac{w(f) - (\ln n + \ln \ln q)}{\sqrt{\ln n + \ln \ln q}} \leq a\right\}}{\#F_n(\mathbb{F}_q)} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^a e^{-\frac{t^2}{2}} dt$$

For sufficiently large  $n$  and a positive number  $0 < \rho < 1$ , the number of distinct prime divisors  $w(f)$  for almost every polynomial  $f \in F_n(\mathbb{F}_q)$  satisfies

$$\rho \ln n < w(f) < (2 - \rho) \ln n.$$

Fix positive integers  $\alpha, \beta$ . We denote by

$$\mathbb{P}[\alpha < w(f) < \beta \mid f \in F_n(\mathbb{F}_q)]$$

the probability that the number of irreducible factors of a square-free polynomial  $f$  of degree  $n$  over  $\mathbb{F}_q$  is greater than  $\alpha$  and less than  $\beta$ . In other words,

$$\mathbb{P}[\alpha < w(f) < \beta \mid f \in F_n(\mathbb{F}_q)] := \frac{\#\{f \in F_n(\mathbb{F}_q) \mid \alpha < w(f) < \beta\}}{\#\{f \in F_n(\mathbb{F}_q)\}} \quad (12)$$

A more rigorous formulation of the above observation is demonstrated in the theorem below.

**Theorem 3.5** (Large deviation principles for Erdős-Kac Theorem). *Denote by  $w(f)$  the number of distinct irreducible factors dividing  $f \in F_n(\mathbb{F}_q)$ . Denote by  $m_n$  the quantity*

$$m_n := \ln n + \ln \ln q.$$

*Then the following limit converges:*

$$\lim_{n \rightarrow \infty} \left( m_n^{-\frac{1}{3}} \ln \mathbb{P} \left[ w(f) > m_n + m_n^{\frac{2}{3}} \mid f \in F_n(\mathbb{F}_q) \right] + \frac{(m_n + m_n^{\frac{2}{3}})^2}{2} \right) = 0.$$

*Proof.* The theorem follows immediately from [FWY20, Theorem 2]. For any positive  $\epsilon > 0$ ,

$$\lim_{n \rightarrow \infty} \left( m_n^{-2\epsilon} \ln \mathbb{P} \left[ w(f) > m_n + m_n^{\frac{1}{2}+\epsilon} \mid f \in F_n(\mathbb{F}_q) \right] + \frac{(m_n + m_n^{\frac{1}{2}+\epsilon})^2}{2} \right) = 0$$

Choose  $\epsilon = \frac{1}{6}$  to prove the theorem.  $\square$

The above theorem shows that for sufficiently large  $n$  and a positive  $0 < \rho < 1$ , there exists a constant  $C > 0$  such that

$$\mathbb{P} [w(f) > (2 - \rho) \ln n \mid f \in F_n(\mathbb{F}_q)] < \mathbb{P} \left[ w(f) > m_n + m_n^{\frac{2}{3}} \mid f \in F_n(\mathbb{F}_q) \right] < C \cdot e^{-\frac{(\ln n)^2}{2}} = C \cdot n^{-\frac{1}{2} \ln n} \quad (13)$$

To verify that the probability  $\mathbb{P} [w(f) < \rho \ln n \mid f \in F_n(\mathbb{F}_q)]$  is arbitrarily small for sufficiently large  $n$ , we first prove the following technical lemma.

**Lemma 3.6.** *Fix positive integers  $n$  and  $1 \leq w \leq 2 \ln n$ . Let  $I$  be a function in variable  $k$  defined as*

$$I := \int_1^k \int_{x_1}^{\frac{n}{w} - \frac{x_1}{w}} \dots \int_{x_i}^{\frac{n}{w-i+1} - \sum_{j=1}^i \frac{x_j}{w-i+1}} \dots \int_{x_{w-1}}^{\frac{n}{2} - \sum_{j=1}^{w-1} \frac{x_j}{2}} \frac{1}{n - x_1 - \dots - x_w} \prod_{j=1}^w \frac{1}{x_j} dx_w dx_{w-1} \dots dx_1.$$

*Note that the bounds of the integration of  $I$  is a subset of  $\mathbb{R}_{\geq 0}^w$ . If  $1 \leq k \leq \frac{n}{w+1}$ , then*

$$I \leq \frac{1}{n} \frac{1}{w!} ((\ln(n-w))^w - (\ln|n-wk| - \ln|k|)^w)$$

*Proof.* Observe that

$$\begin{aligned} & \int_{x_{w-1}}^{\frac{n}{2} - \sum_{j=1}^{w-1} \frac{x_j}{2}} \frac{1}{n - x_1 - \dots - x_w} \prod_{j=1}^w \frac{1}{x_j} dx_w \\ &= \left( \prod_{j=1}^{w-1} \frac{1}{x_j} \right) \int_{x_{w-1}}^{\frac{n}{2} - \sum_{j=1}^{w-1} \frac{x_j}{2}} \frac{1}{n - x_1 - \dots - x_w} \frac{1}{x_w} dx_w \\ &= \left( \prod_{j=1}^{w-1} \frac{1}{x_j} \right) \int_{x_{w-1}}^{\frac{n}{2} - \sum_{j=1}^{w-1} \frac{x_j}{2}} \frac{1}{n - x_1 - \dots - x_{w-1}} \left( \frac{1}{x_w} + \frac{1}{n - x_1 - \dots - x_w} \right) dx_w \\ &= \frac{1}{n - x_1 - \dots - x_{w-1}} \left( \prod_{j=1}^{w-1} \frac{1}{x_j} \right) \int_{x_{w-1}}^{\frac{n}{2} - \sum_{j=1}^{w-1} \frac{x_j}{2}} \frac{1}{x_w} + \frac{1}{n - x_1 - \dots - x_w} dx_w \\ &= \frac{1}{n - x_1 - \dots - x_{w-1}} \left( \prod_{j=1}^{w-1} \frac{1}{x_j} \right) [\ln|x_w| - \ln|n - x_1 - \dots - x_w|]_{x_{w-1}}^{\frac{n}{2} - \sum_{j=1}^{w-1} \frac{x_j}{2}} \\ &= \frac{1}{n - x_1 - \dots - x_{w-1}} \left( \prod_{j=1}^{w-1} \frac{1}{x_j} \right) (\ln|n - x_1 - \dots - x_{w-2} - 2x_{w-1}| - \ln|x_{w-1}|). \end{aligned}$$

For any  $1 \leq i \leq w-2$ , we also deduce that

$$\begin{aligned}
& \int_{x_i}^{\frac{n}{w-i+1} - \sum_{j=1}^i \frac{x_j}{w-i+1}} \frac{1}{n - x_1 - \cdots - x_{i+1}} \left( \prod_{j=1}^{i+1} \frac{1}{x_j} \right) \frac{(\ln |n - x_1 - \cdots - x_i - (w-i)x_{i+1}| - \ln |x_{i+1}|)^{w-i-1}}{(w-i-1)!} dx_{i+1} \\
&= \frac{1}{(w-i-1)!} \left( \prod_{j=1}^i \frac{1}{x_j} \right) \int_{x_i}^{\frac{n}{w-i+1} - \sum_{j=1}^i \frac{x_j}{w-i+1}} \frac{1}{x_{i+1}} \frac{1}{n - x_1 - \cdots - x_{i+1}} \\
&\quad \times (\ln |n - x_1 - \cdots - x_i - (w-i)x_{i+1}| - \ln |x_{i+1}|)^{w-i-1} dx_{i+1} \\
&= \frac{1}{n - x_1 - \cdots - x_i} \frac{1}{(w-i-1)!} \left( \prod_{j=1}^i \frac{1}{x_j} \right) \int_{x_i}^{\frac{n}{w-i+1} - \sum_{j=1}^i \frac{x_j}{w-i+1}} \left( \frac{1}{x_{i+1}} + \frac{1}{n - x_1 - \cdots - x_{i+1}} \right) \\
&\quad \times (\ln |n - x_1 - \cdots - x_i - (w-i)x_{i+1}| - \ln |x_{i+1}|)^{w-i-1} dx_{i+1} \\
&\leq \frac{1}{n - x_1 - \cdots - x_i} \frac{1}{(w-i-1)!} \left( \prod_{j=1}^i \frac{1}{x_j} \right) \int_{x_i}^{\frac{n}{w-i+1} - \sum_{j=1}^i \frac{x_j}{w-i+1}} \left( \frac{1}{x_{i+1}} + \frac{1}{n - x_1 - \cdots - x_i - (w-i)x_{i+1}} \right) \\
&\quad \times (\ln |n - x_1 - \cdots - x_i - (w-i)x_{i+1}| - \ln |x_{i+1}|)^{w-i-1} dx_{i+1} \\
&= \frac{1}{n - x_1 - \cdots - x_i} \frac{1}{(w-i)!} \left[ -(\ln |n - x_1 - \cdots - x_i - (w-i)x_{i+1}| - \ln |x_{i+1}|)^{w-i} \right]_{x_i}^{\frac{n}{w-i+1} - \sum_{j=1}^i \frac{x_j}{w-i+1}} \\
&= \frac{1}{n - x_1 - \cdots - x_i} \left( \prod_{j=1}^i \frac{1}{x_j} \right) \frac{(\ln |n - x_1 - \cdots - x_{i-1} - (w-i+1)x_i| - \ln |x_i|)^{w-i}}{(w-i)!}
\end{aligned}$$

Therefore, the integral can be abbreviated as:

$$\begin{aligned}
I &\leq \int_1^k \frac{1}{n - x_1} \frac{1}{x_1} \frac{(\ln |n - wx_1| - \ln |x_1|)^{w-1}}{(w-1)!} dx_1 \\
&\leq \frac{1}{n} \left[ -\frac{(\ln |n - wx_1| - \ln |x_1|)^w}{w!} \right]_1^k \\
&= \frac{1}{n} \frac{1}{w!} ((\ln(n-w))^w - (\ln |n - wk| - \ln |k|)^w)
\end{aligned}$$

□

We can now prove the effective version of the Erdős-Kac theorem.

**Theorem 3.7** (Effective Erdős-Kac Theorem). *Denote by  $w(f)$  the number of distinct irreducible factors dividing  $f \in F_n(\mathbb{F}_q)$ . Then for sufficiently large  $n$  and a fixed positive number  $0 < \rho < 1$ , there exists a constant  $C > 0$  and a fixed number  $0 < \alpha_{ek} < \rho \ln \rho + 1 - \rho$  such that*

$$\mathbb{P}[\rho \ln n < w(f) < (2 - \rho) \ln n \mid f \in F_n(\mathbb{F}_q)] > \left(1 - C \frac{1}{n^{\alpha_{ek}}}\right)$$

*Proof.* It suffices to show that

$$\mathbb{P}[w(f) < \rho \ln n \mid f \in F_n(\mathbb{F}_q)] < C \frac{1}{n^{\alpha_{ek}}}.$$

Given such a polynomial  $f \in F_n(\mathbb{F}_q)$ , we can write the factorization of  $f$  as  $f = f_1 f_2 \cdots f_w$ , where  $\deg f_i \leq \deg f_{i+1}$  for all  $1 \leq i \leq w$ . The mobius inversion formula implies that the number of monic irreducible polynomials of degree  $i$  is at most  $\frac{q^i}{i}$ . We also recall that the number of monic squarefree polynomials of degree  $i$  is equal to  $q^i - q^{i-1}$ . This implies that the number of monic polynomials of

degree  $n$  with exactly  $w$  distinct irreducible factors is at most:

$$\begin{aligned}
 & \#\{f \in F_n(\mathbb{F}_q) \mid w(f) = w, f \text{ monic}\} \\
 & \leq \sum_{\substack{(m_1, m_2, \dots, m_{w-1}) \in \mathbb{N}^{w-1} \\ m_1 \leq m_2 \leq \dots \leq m_{w-1} \leq n - m_1 - \dots - m_{w-1}}} \left( \sum_{k=1}^{n-m_1-\dots-m_{w-1}} \frac{q^k}{k} \right) \prod_{i=1}^{w-1} \frac{q^{m_i}}{m_i} \\
 & \leq 2 \cdot \sum_{\substack{(m_1, m_2, \dots, m_{w-1}) \in \mathbb{N}^{w-1} \\ m_1 \leq m_2 \leq \dots \leq m_{w-1} \leq n - m_1 - \dots - m_{w-1}}} \frac{q^{n-m_1-\dots-m_{w-1}}}{n - m_1 - \dots - m_{w-1}} \prod_{i=1}^{w-1} \frac{q^{m_i}}{m_i} \\
 & = 2 \cdot \sum_{\substack{(m_1, m_2, \dots, m_{w-1}) \in \mathbb{N}^{w-1} \\ m_1 \leq m_2 \leq \dots \leq m_{w-1} \leq n - m_1 - \dots - m_{w-1}}} \frac{q^n}{m_1 m_2 \dots m_{w-1} (n - m_1 - \dots - m_{w-1})} \\
 & \leq 4q^n \int_1^{\frac{n}{w}} \int_{x_1}^{\frac{n}{w-1} - \frac{x_1}{w-1}} \dots \int_{x_i}^{\frac{n}{w-i} - \sum_{j=1}^i \frac{x_j}{w-i}} \dots \int_{x_{w-2}}^{\frac{n}{2} - \sum_{j=1}^{w-2} \frac{x_j}{2}} \frac{1}{n - x_1 - \dots - x_{w-1}} \prod_{j=1}^{w-1} \frac{1}{x_j} dx_{w-1} dx_{w-2} \dots dx_1 \\
 & \leq \frac{4q^n}{n} \frac{1}{(w-1)!} (\ln |n|)^{w-1}.
 \end{aligned}$$

We note that

$$\sum_{k=1}^{\infty} \frac{q^n}{n} \frac{1}{k!} (\ln |n|)^k = \frac{q^n}{n} e^{\ln |n|} = q^n.$$

We recall that for any  $1 \leq k \leq \ln n$ ,

$$\frac{(\ln n)^k}{k!} < \frac{(\ln n)^{k+1}}{(k+1)!}$$

Pick  $M_n$  to be an integer which satisfies

$$\rho \ln n \leq M_n \leq \rho \ln n + 1 \tag{14}$$

Then the number of monic polynomials of degree  $n$  with at most  $\rho \ln n$  irreducible factors is bounded above by:

$$\#\{f \in F_n(\mathbb{F}_q) \mid f \text{ monic}, w(f) \leq \rho \ln n\} \leq \sum_{k=1}^{M_n} \frac{4q^n}{n} \frac{1}{k!} (\ln |n|)^k$$

Note that

$$\begin{aligned}
 \mathbb{P}[w(f) \leq \rho \ln n \mid f \in F_n(\mathbb{F}_q)] & \leq \frac{1}{q^n - q^{n-1}} \sum_{k=1}^{M_n} \frac{4q^n}{n} \frac{(\ln |n|)^k}{k!} \\
 & = \frac{4q}{q-1} \frac{1}{n} \left( \sum_{k=1}^{M_n} \frac{(\ln |n|)^k}{k!} \right) \\
 & \leq \frac{4q}{q-1} \frac{(\ln |n|)^{M_n}}{n(M_n - 1)!} \\
 & = \frac{4q}{q-1} \frac{(\ln n)^{M_n}}{n \Gamma(M_n)} \\
 & \leq \frac{4q}{q-1} \frac{(\ln n)^{(\rho \ln n)}}{n \Gamma(\rho \ln n)}
 \end{aligned}$$

where  $\Gamma(x)$  is the Gamma function.

Consider the following function

$$H(x) := \frac{\left(\frac{1}{\rho}x\right)^x}{e^{\frac{1}{\rho}x}\Gamma(x)} \quad (15)$$

We recall that Stirling's approximation of Gamma functions is given by (see [Imp03] for a detailed treatment of the subject):

$$x \ln x - x + \frac{1}{2} \ln \frac{2\pi}{x} < \ln \Gamma(x) < x \ln x - x + \frac{1}{2} \ln \frac{2\pi}{x} + \frac{1}{12x} \quad (16)$$

Hence, we obtain

$$\begin{aligned} \ln H(x) &:= x \ln\left(\frac{1}{\rho}x\right) - \frac{1}{\rho}x - \ln \Gamma(x) \\ &< x \ln \frac{1}{\rho} + x \ln x - \frac{1}{\rho}x - x \ln x + x - \frac{1}{2} \ln \frac{2\pi}{x} \\ &< (1 - \ln \rho - \frac{1}{\rho})x + \frac{1}{2} \ln x. \end{aligned}$$

Because  $0 < \rho < 1$ , we have  $1 - \ln \rho - \frac{1}{\rho} < 0$ . This implies that there exists a fixed constant  $0 < \alpha_{ek} < \rho \ln \rho + 1 - \rho$  such that for sufficiently large  $n$ ,

$$\frac{(\ln n)^{\rho \ln n}}{n \Gamma(\rho \ln n)} \leq \frac{4q}{q-1} \frac{1}{n^{\alpha_{ek}}}. \quad (17)$$

Therefore, there exists a fixed constant  $4 < C < 5$  such that

$$\mathbb{P}[w(f) < \rho \ln n \mid f \in F_n(\mathbb{F}_q)] < C \frac{1}{n^{\alpha_{ek}}}.$$

□

**Remark 3.8.** We note that Theorem 3.7 can be obtained from using the results by Cohen, see for instance [Coh69, Theorem 6] and [CLNY22, Theorem 1.1]. Nevertheless, to make the manuscript self-contained, we included an elementary proof of the Theorem. We will use the ideas from Theorem 3.7 to obtain a result that for almost all polynomials  $f \in \mathbb{F}_q[t]$  of degree  $n$ , a non-trivial proportion of irreducible factors of  $f$  is of degree greater than  $n^{\frac{1}{3}}$ , see in particular Proposition 5.4.

#### 4. LOCAL CYCLIC PRIME TWISTS OF ELLIPTIC CURVES

**4.1. Local twists.** The objective of this section is to analyze the variations of 2-Selmer ranks under local quadratic twists of  $E$ . Klagsbrun, Mazur, and Rubin [KMR14] carefully analyzes these variations over number fields using Galois cohomology groups and Poitou-Tate duality theorem. Hence, the analogous results on the variation of 2-Selmer ranks under local quadratic twists of  $E$  over  $\mathbb{F}_q(t)$  can also be proven using the identical argument, see for instance Chapter 1 of [Mil06] for a rigorous treatment of Poitou-Tate duality theorems for global function fields. We further enrich these results by using the effective Chebotarev density theorem and Erdős-Kac theorem over  $\mathbb{F}_q(t)$ .

Let  $p$  be any prime number that is coprime to the characteristic of the global function field  $K = \mathbb{F}_q(t)$  of characteristic coprime to 2 and 3. Throughout this section, we assume that the following properties hold, where  $F(x) \in K[x]$  is a cubic polynomial.

- $E : y^2 = F(x)$  is a non-isotrivial elliptic curve over  $K$ .
  - $E$  contains a place  $\infty$  of split multiplicative reduction.
  - The constant field  $\mathbb{F}_q$ , of characteristic coprime to  $2, 3, p$ , and contains  $\mu_p$ .
  - The image of  $\text{Gal}(\overline{K}/K) \rightarrow \text{Aut}(E[p])$  contains  $\text{SL}_2(\mathbb{F}_p)$ .
- (18)

By Igusa's theorem, for any non-isotrivial elliptic curve  $E$ , there exists a prime  $p$  and a finite separable extension of  $K = \mathbb{F}_q(t)$  such that  $E$  satisfies all the four conditions [Igu59, BLV09].

**Definition 4.1.** We introduce the following notations, as stated in [KMR14, Sections 5, 7, and 9].

- $\Sigma$ : the set of places of  $K$  consisting of places of bad reduction of  $E$  and  $\infty$ .
- $\sigma$ : a square-free product of places  $v$  of  $K$  such that  $v \notin \Sigma$ .
- $\deg \sigma$ : the sum of degrees of places  $v \mid \sigma$ , i.e.  $\deg \sigma = \sum_{v \mid \sigma} \deg v$ .
- $\Omega_\sigma$ : the set of finite cartesian products of local characters

$$\chi := (\chi_v)_v \in \prod_{\substack{v \in \Sigma \text{ or} \\ v \mid \sigma}} \text{Hom}(\text{Gal}(\overline{K}_v/K_v), \mu_p)$$

such that the component  $\chi_v$  is ramified if  $v \mid \sigma$ . For the sake of convenience, we will denote by  $\text{Hom}_{\text{unr}}(\text{Gal}(\overline{K}_v/K_v), \mu_p)$  the set of unramified local characters at place  $v$ , and by  $\text{Hom}_{\text{ram}}(\text{Gal}(\overline{K}_v/K_v), \mu_p)$  the set of ramified local characters at place  $v$ . Assuming that  $\mu_p \subset K_v$ , there are  $p$  distinct unramified local characters at  $v$ , and  $p(p-1)$  distinct ramified local characters at  $v$ .

- Given a global character  $\chi \in \text{Hom}(\text{Gal}(\overline{K}/K), \mu_p)$ , the twist  $E^\chi$  is defined as follows (see [MRS07, Definition 5.1] for further details):

Let  $F/K$  be the cyclic Galois extension of degree  $p$  corresponding to the character  $\chi$ . Denote by  $L/K$  the Galois closure of  $F$  with  $G := \text{Gal}(L/K)$ . Denote by  $K[G]$  the group ring of  $G$  with coefficients in  $K$ . Then  $K[G]$  admits a decomposition into a direct sum of minimal two-sided ideals

$$K[G] = \bigoplus_\rho K[G]_\rho$$

where  $\rho$  spans the set of irreducible  $K$ -representations of  $G$ , and  $K[G]_\rho$  is the sum of all left ideals of  $K[G]$  isomorphic to  $\rho$ .

Let  $I_F$  be the sum of integral left ideals of  $K[G]$  isomorphic to  $\chi$ , i.e.

$$I_F := K[G]_\chi \cap \mathcal{O}_K[G]$$

Then the twist of  $E$  by  $\chi$  is given by

$$E^\chi := I_F \otimes E.$$

Note that if  $\chi$  is a quadratic character, then  $E^\chi$  is the quadratic twist of  $E$  over  $K$ . If  $\chi$  is a cyclic order- $p$  character, one can take  $E^\chi$  to be the kernel of the norm map

$$E^\chi := \text{Ker} \left( \text{Res}_K^{L^\chi} E \rightarrow E \right) \quad (19)$$

from the Weil restriction of scalars of  $E/K$  associated to the cyclic  $p$ -extension  $L^\chi/K$  associated to the character  $\chi$ . We note that the  $K$ -rational points of  $\text{Res}_K^{L^\chi} E$  are the  $L^\chi$ -rational points of  $E$ . The twist  $E^\chi/K$  is a  $p-1$  dimensional abelian variety over  $K$ .

- Given a cyclic order- $p$  character  $\chi$ , the endomorphism ring of  $E^\chi$ , denoted as  $\text{End}(E^\chi)$ , contains the group ring  $\mathbb{Z}[\text{Gal}(F^\chi/K)] \cong \mathbb{Z}[\zeta_p]$ . We denote by  $\pi$  the unique prime ideal of  $\mathbb{Z}[\zeta_p]$  lying above the ideal  $(p) \subset \mathbb{Z}$ . Note that  $\pi$  defines an isogeny over the twist  $E^\chi$ .
- Given a local character  $\chi \in \Omega_\sigma$ , the  $\pi$ -Selmer group of the twist  $E^\chi$  is given by

$$\text{Sel}_\pi(E^\chi) := \text{Ker} \left( H_{\text{ét}}^1(K, E[p]) \rightarrow \bigoplus_v H_{\text{ét}}^1(K_v, E[p]) / \text{im} \delta_v^\chi \right), \quad (20)$$

where  $\delta_v^\chi : E^{\chi_v}(K_v) / \pi E^{\chi_v}(K_v) \rightarrow H_{\text{ét}}^1(K_v, E[p])$  is the local Kummer map at  $v$ . Under all but the third assumption stated in (18), we use the isomorphism

$$\begin{aligned} H_{\text{ét}}^1(K, E[p]) &\cong H_{\text{ét}}^1(K, E^\chi[\pi]), \\ H_{\text{ét}}^1(K_v, E[p]) &\cong H_{\text{ét}}^1(K_v, E_v^\chi[\pi]) \end{aligned}$$

to define the Selmer group  $\text{Sel}_\pi(E^\chi)$ , see in particular [MR07, Proposition 4.1, Definition 4.3]. Even though the reference particularly states about elliptic curves over number fields, the ideas of the proofs of relevant results are extendable to global function fields. Note that if  $p = 2$ , the  $\pi$ -Selmer groups of  $E^\chi$  correspond to 2-Selmer groups of  $E$  twisted by a quadratic character  $\chi$ .

- For  $1 \leq i \leq 2$ , define the set

$$\mathcal{P}_i := \{v \mid v \notin \Sigma, \mu_p \subset K_v, \text{ and } \dim_{\mathbb{F}_p} E(K_v)[p] = i\}$$

We also define the set

$$\mathcal{P}_0 := \{v \mid v \notin \Sigma \cup \mathcal{P}_1 \cup \mathcal{P}_2\}.$$

The set  $\mathcal{P}$  is the set

$$\mathcal{P} := \{v \mid v \notin \Sigma\} = \mathcal{P}_0 \cup \mathcal{P}_1 \cup \mathcal{P}_2.$$

- Given a positive number  $d \in \mathbb{N}$ , the set  $\mathcal{P}_i(d)$  is defined as

$$\mathcal{P}_i(d) := \{v \in \mathcal{P}_i \mid \deg v = d\}.$$

Likewise, the set  $\mathcal{P}(d)$  is defined as

$$\mathcal{P}(d) := \{v \in \mathcal{P} \mid \deg v = d\}.$$

- Given a local character  $\chi \in \Omega_\sigma$ , we denote by  $\text{rk}(\chi)$  the dimension of  $\text{Sel}_\pi(E^\chi)$  as an  $\mathbb{F}_p$ -vector space.

Using the assumption (18), we recall the following statement from [KMR13, Lemma 4.3] that the Frobenius elements of certain primes lying above a place  $v$  over  $K$  determines which classes of  $\mathcal{P}_i$  the place  $v$  lives in. Again, the original statement of the lemma is shown for arbitrary number fields, which can be extended to the case for global function fields.

**Lemma 4.2.** [KMR13, Lemma 4.3] *Fix an elliptic curve  $E/K$  satisfying the conditions stated in (18). Let  $v$  be a place over  $K$  such that  $v \notin \Sigma$ . Denote by  $\text{Frob}_v \in \text{Gal}(K(E[p])/K)$  the Frobenius element associated to  $v$ . Then*

- (1)  $v \in \mathcal{P}_2 \iff \text{Frob}_v = 1$
- (2)  $v \in \mathcal{P}_1 \iff \text{Frob}_v \text{ has order exactly } p$
- (3)  $v \in \mathcal{P}_0 \iff \text{Frob}_v^p \neq 1$

**Remark 4.3.** Igusa's theorem implies that any non-isotrivial elliptic curve satisfying conditions (18) satisfies the condition that  $\text{Gal}(K(E[p])/K) \cong \text{SL}_2(\mathbb{F}_p) \rtimes T$ , where  $T$  is a cyclic subgroup of order prime to  $p$  corresponding to the Galois group of the constant field extension of  $K(E[p])/K$ . With the condition that  $\mu_p \subset K$ , one may assume without loss of generality that  $|T| = 1$ . Nevertheless, the proofs of the results outlined in the manuscript are shown for any such finite cyclic group  $T$ .

Computing the conjugacy classes of  $\text{SL}_2(\mathbb{F}_p)$  and Theorem 3.1 show that there exists a constant  $C > 0$  such that for sufficiently large  $d$ ,

$$\max \left\{ \left| \frac{\#\mathcal{P}_0(d)}{\#\mathcal{P}(d)} - \left(1 - \frac{p}{|T|(p^2 - 1)}\right) \right|, \left| \frac{\#\mathcal{P}_1(d)}{\#\mathcal{P}(d)} - \frac{1}{|T|p} \right|, \left| \frac{\#\mathcal{P}_2(d)}{\#\mathcal{P}(d)} - \frac{1}{|T|(p^3 - p)} \right| \right\} < Cq^{-\frac{d}{2}}. \quad (21)$$

Suppose in particular that  $p = 2$ . Given a Weierstrass equation of an elliptic curve  $E : y^2 = F(x)$  satisfying the conditions from Theorem 1.1, denote by  $L$  the cubic field extension  $L = K[x]/(F(x))$ . Note that the constant field of  $L$  is equal to  $\mathbb{F}_q$ . The sets  $\mathcal{P}_0, \mathcal{P}_1$ , and  $\mathcal{P}_2$  correspond to set of unramified places over  $K$  not in  $\Sigma$  which are inert, split into two places, or totally split in  $L$ . Theorem 3.1 shows that there exists a constant  $C > 0$  such that for sufficiently large  $d$ ,

$$\max \left\{ \left| \frac{\#\mathcal{P}_0(d)}{\#\mathcal{P}(d)} - \frac{1}{3} \right|, \left| \frac{\#\mathcal{P}_1(d)}{\#\mathcal{P}(d)} - \frac{1}{2} \right|, \left| \frac{\#\mathcal{P}_2(d)}{\#\mathcal{P}(d)} - \frac{1}{6} \right| \right\} < Cq^{-\frac{d}{2}} \quad (22)$$

Note that (22) immediately follows from (21) by setting  $p = 2$  and  $|T| = 1$ .



**Definition 4.4.** Fix a square-free product of places  $\sigma$  coprime to elements in  $\Sigma$ .

Fix a local character  $\chi \in \Omega_\sigma$ . Given a single place  $\mathfrak{v}$  over  $K$  such that  $\mathfrak{v} \nmid \sigma$  and  $\mathfrak{v} \notin \Sigma$ , let  $\chi' \in \Omega_{\sigma\mathfrak{v}}$  be a local character such that

- For any  $v \mid \sigma$  or  $v \in \Sigma$ ,  $\chi'_v = \chi_v$ .
- At  $\mathfrak{v}$ ,  $\chi'_\mathfrak{v}$  is ramified.

Denote by  $\Omega_{\chi,\mathfrak{v}}$  the set of local characters  $\chi'$  satisfying the two conditions above. Note that

$$\Omega_{\sigma\mathfrak{v}} = \bigsqcup_{\chi \in \Omega_\sigma} \Omega_{\chi,\mathfrak{v}}.$$

Denote by  $\Omega_{\chi,\mathfrak{v}}^\eta$  the set of local characters  $\chi'$  given a fixed ramified character  $\eta \in \text{Hom}(\text{Gal}(\overline{K}_v/K_v), \mu_p)$  which satisfy the following conditions:

- For any  $v \mid \sigma$  or  $v \in \Sigma$ ,  $\chi'_v = \chi_v$ .
- At  $\mathfrak{v}$ , the local character  $\frac{\chi'_\mathfrak{v}}{\eta}$  is an unramified order- $p$  local character of  $\text{Gal}(\overline{K}_v/K_v)$ .

**Proposition 4.5** (Local twists of  $\pi$ -Selmer groups). *Let  $E$  be a non-isotrivial elliptic curve over  $K$  satisfying the conditions from equation (18). Fix a square-free product of places  $\sigma$  coprime to elements in  $\Sigma$ . Fix a local character  $\chi \in \Omega_\sigma$ . Let  $\Omega_{\chi,\mathfrak{v}}$  be the set of local characters  $\chi'$  satisfying the first two enumerated conditions from 4.4.*

Let  $c_{i,j}$  be given by the following table:

$c_{i,j}$	$j = -2$	$j = -1$	$j = 0$	$j = 1$	$j = 2$
$i = 0$	$\times$	$\times$	1	$\times$	$\times$
$i = 1$	$\times$	$1 - p^{-rk(\chi)}$	$\times$	$p^{-rk(\chi)}$	$\times$
$i = 2$	$1 - (p+1)p^{-rk(\chi)} + p^{1-2rk(\chi)}$	$\times$	$(p+1)p^{-rk(\chi)} - (p + \frac{1}{p})p^{-2rk(\chi)}$	$\times$	$p^{-1-2rk(\chi)}$

Here, the term " $\times$ " denotes the case where such a difference of ranks cannot occur.

(1) Let  $d$  be an integer which satisfies the condition that

$$d > (8 \ln p + 8 \ln 2 + 2 \ln \Delta_E) \deg \sigma \quad (23)$$

Then there exists a constant  $C > 0$  such that

$$\left| \frac{\sum_{\substack{\mathfrak{v} \in \mathcal{P}_i(d) \\ \mathfrak{v} \nmid \sigma, \mathfrak{v} \notin \Sigma}} \#\{\chi' \in \Omega_{\chi,\mathfrak{v}} \mid rk(\chi') - rk(\chi) = j\}}{\sum_{\substack{\mathfrak{v} \in \mathcal{P}_i(d) \\ \mathfrak{v} \nmid \sigma, \mathfrak{v} \notin \Sigma}} \#\Omega_{\chi,\mathfrak{v}}} - c_{i,j} \right| < C \frac{\deg \sigma}{q^{\frac{\deg \sigma}{2}}} \quad (24)$$

(2) Suppose  $w(\sigma) \leq 2 \ln(\deg \sigma) - 2$ . Let  $d$  be an integer which satisfies the condition that

$$d > \frac{4(\ln \deg \sigma)^2}{\ln q}. \quad (25)$$

Then for  $\sigma$  of sufficiently large degree, there exists a constant  $C > 0$  such that

$$\left| \frac{\sum_{\substack{\mathfrak{v} \in \mathcal{P}_i(d) \\ \mathfrak{v} \nmid \sigma, \mathfrak{v} \notin \Sigma}} \#\{\chi' \in \Omega_{\chi,\mathfrak{v}} \mid rk(\chi') - rk(\chi) = j\}}{\sum_{\substack{\mathfrak{v} \in \mathcal{P}_i(d) \\ \mathfrak{v} \nmid \sigma, \mathfrak{v} \notin \Sigma}} \#\Omega_{\chi,\mathfrak{v}}} - c_{i,j} \right| < C(\deg \sigma)^{-\ln \deg \sigma}.$$

In particular, there exists a constant  $C > 0$  and a cyclic finite group  $T$  such that

$$\left| \frac{\sum_{\substack{\mathfrak{v} \in \mathcal{P}(d) \\ \mathfrak{v} \nmid \sigma, \mathfrak{v} \notin \Sigma}} \#\{\chi' \in \Omega_{\chi,\mathfrak{v}} \mid rk(\chi') - rk(\chi) = j\}}{\sum_{\substack{\mathfrak{v} \in \mathcal{P}(d) \\ \mathfrak{v} \nmid \sigma, \mathfrak{v} \notin \Sigma}} \#\Omega_{\chi,\mathfrak{v}}} - \left( \left(1 - \frac{p}{|T|(p^2-1)}\right) c_{0,j} + \frac{1}{|T|p} c_{1,j} + \frac{1}{|T|(p^3-p)} c_{2,j} \right) \right| < C(\deg \sigma)^{-\ln \deg \sigma}.$$

(3) Let  $\Omega_{\chi,\mathfrak{v}}^\eta$  be the set of local characters  $\chi'$  given a ramified local character  $\eta \in \text{Hom}(\text{Gal}(\overline{K}_v/K_v), \mu_p)$  satisfying the last two enumerated conditions from Definition 4.4. Then both statements (1) and (2) hold for the set of characters  $\Omega_{\chi,\mathfrak{v}}^\eta$  given any choice of a ramified local character  $\eta \in \text{Hom}(\text{Gal}(\overline{K}_v/K_v), \mu_p)$ .

*Proof.* The theorem can be proved in an analogous way to how [KMR14, Proposition 7.2] and [KMR14, Proposition 9.4] were proved over number fields. Nevertheless, it is necessary to apply the effective Chebotarev density theorem and Erdős-Kac theorem to calculate the explicit error bounds.

### A review on local twists of $p$ -Selmer groups

We recall some of the key ideas used in proving both propositions from [KMR14]. Consider the groups

$$\begin{aligned}\mathrm{Sel}_\pi(E^\chi)^{(\mathfrak{v})} &:= \mathrm{Ker} \left( H_{\mathrm{\acute{e}t}}^1(K, E[p]) \rightarrow \bigoplus_{v \neq \mathfrak{v}} H_{\mathrm{\acute{e}t}}^1(K_v, E[p]) / \mathrm{im} \delta_v^\chi \right) \\ \mathrm{Sel}_\pi(E^\chi)_{(\mathfrak{v})} &:= \mathrm{Ker} \left( \mathrm{Sel}_\pi(E^\chi)^{(\mathfrak{v})} \rightarrow H_{\mathrm{\acute{e}t}}^1(K_{\mathfrak{v}}, E[p]) \right)\end{aligned}$$

where the morphisms are induced from local restrictions of cohomology groups. The group  $\mathrm{Sel}_\pi(E^\chi)^{(\mathfrak{v})}$ , which contains  $\mathrm{Sel}_\pi(E^\chi)$ , is obtained by imposing no local conditions at  $\mathfrak{v}$ . On the other hand, the group  $\mathrm{Sel}_\pi(E^\chi)_{(\mathfrak{v})}$ , which is contained in  $\mathrm{Sel}_\pi(E^\chi)$ , is obtained by imposing the strongest local condition at  $\mathfrak{v}$ .

The conditions on  $\chi' \in \Omega_{\chi, \mathfrak{v}}$  implies that the pair of sequences

$$\begin{aligned}0 \rightarrow \mathrm{Sel}_\pi(E^\chi)_{(\mathfrak{v})} \rightarrow \mathrm{Sel}_\pi(E^\chi) \rightarrow \mathrm{im} \delta_{\mathfrak{v}}^\chi \rightarrow 0 \\ 0 \rightarrow \mathrm{Sel}_\pi(E^\chi)_{(\mathfrak{v})} \rightarrow \mathrm{Sel}_\pi(E^{\chi'}) \rightarrow \mathrm{im} \delta_{\mathfrak{v}}^{\chi'} \rightarrow 0\end{aligned}$$

are short exact. One can then use the relation

$$\mathrm{rk}(\chi') - \mathrm{rk}(\chi) = \dim_{\mathbb{F}_p} \mathrm{im} \delta_{\mathfrak{v}}^{\chi'} - \dim_{\mathbb{F}_p} \mathrm{im} \delta_{\mathfrak{v}}^\chi$$

to deduce that the cases where  $c_{i,j}$  are notated as  $\times$  in the table are unobtainable. As stated in [KMR14, Proposition 7.2],

$$\mathrm{rk}(\chi') - \mathrm{rk}(\chi) = \begin{cases} 0 & \text{if } \mathfrak{v} \in \mathcal{P}_0(d) \\ -1 & \text{if } \dim_{\mathbb{F}_p} \mathrm{im} \delta_{\mathfrak{v}}^\chi = 1 \text{ and } \mathfrak{v} \in \mathcal{P}_1(d) \\ 1 & \text{if } \dim_{\mathbb{F}_p} \mathrm{im} \delta_{\mathfrak{v}}^\chi = 0 \text{ and } \mathfrak{v} \in \mathcal{P}_1(d) \\ -2 & \text{if } \dim_{\mathbb{F}_p} \mathrm{im} \delta_{\mathfrak{v}}^\chi = 2 \text{ and } \mathfrak{v} \in \mathcal{P}_2(d) \\ 0 & \text{if } \dim_{\mathbb{F}_p} \mathrm{im} \delta_{\mathfrak{v}}^\chi = 1 \text{ and } \mathfrak{v} \in \mathcal{P}_2(d) \\ 0 & \text{if } \dim_{\mathbb{F}_p} \mathrm{im} \delta_{\mathfrak{v}}^\chi = 0 \text{ and } \mathfrak{v} \in \mathcal{P}_2(d) \text{ with conditional probability } 1 - \frac{1}{p} \\ 2 & \text{if } \dim_{\mathbb{F}_p} \mathrm{im} \delta_{\mathfrak{v}}^\chi = 0 \text{ and } \mathfrak{v} \in \mathcal{P}_2(d) \text{ with conditional probability } \frac{1}{p} \end{cases} \quad (26)$$

where  $\chi' \in \Omega_{\chi, \mathfrak{v}}$ . Note that the case where  $\mathrm{rk}(\chi') - \mathrm{rk}(\chi) = 2$  with  $v \in \mathcal{P}_2(d)$  bijectively corresponds to the elements of the orbit of a cyclic order- $p$  character  $\chi'_v \in \mathrm{Hom}(\mathrm{Gal}(\overline{K}/K), \mu_p)$  under the action of the automorphism group  $\mathrm{Aut}(\mu_p)$ , as shown in [KMR14, Definition 5.10, Proposition 7.2]. This observation hence implies that the desired probability distributions obtained over the set of characters  $\Omega_{\chi, \mathfrak{v}}$  and  $\Omega_{\chi, \mathfrak{v}}^\eta$  given any ramified order- $p$  local character  $\eta \in \mathrm{Hom}(\mathrm{Gal}(\overline{K}_v/K_v), \mu_p)$  are identical.

For the remainder of the proof, we focus on computing the desired probability distributions obtained over the set of characters  $\Omega_{\chi, \mathfrak{v}}$ . The case for  $\Omega_{\chi, \mathfrak{v}}^\eta$  follows analogously. Denote by  $\mathrm{Res}_p$  the restriction morphism of cohomology groups:

$$H_{\mathrm{\acute{e}t}}^1(K, E[p]) \rightarrow H_{\mathrm{\acute{e}t}}^1(K(E[p]), E[p])^{\mathrm{Gal}(K(E[p])/K)} = \mathrm{Hom}(\mathrm{Gal}(\overline{K(E[p])}/K(E[p])), E[p])^{\mathrm{Gal}(K(E[p])/K)}.$$

Let  $F_{\sigma, \chi}$  be the fixed field of the following subgroup of  $\mathrm{Gal}(\overline{K(E[p])}/K(E[p]))$ :

$$\bigcap_{c \in \mathrm{Sel}_\pi(E^\chi)} \mathrm{Ker} \left( \mathrm{Res}_p(c) : \mathrm{Gal}(\overline{K(E[p])}/K(E[p])) \rightarrow E[p] \right)$$

The field  $F_{\sigma, \chi}$  satisfies the following properties, as shown in [KMR14, Proposition 9.3]:

- (1)  $F_{\sigma, \chi}$  is Galois over  $K$ .

- (2) There is a  $\text{Gal}(K(E[p])/K)$ -module isomorphism  $\text{Gal}(F_{\sigma,\chi}/K(E[p])) \cong (E[p])^{\text{rk}(\chi)}$ .
- (3)  $F_{\sigma,\chi}/K$  is unramified outside of places dividing  $\sigma$  or in  $\Sigma$ .

Observe that if  $K(E[p])$  is a geometric Galois extension over  $K$ , then condition (2) implies that  $F_{\sigma,\chi}$  is a geometric Galois extension over  $K$ . The aforementioned condition holds for  $p = 2$  whenever  $E$  is a non-isotrivial elliptic curve such that  $\text{Gal}(K(E[2])/K) \cong S_3$ .

### Constant field of $F_{\sigma,\chi}$

Note that the constant field of  $F_{\sigma,\chi}$  is equal to  $\mathbb{F}_q$ , regardless of whether  $K(E[p])$  is a geometric Galois extension over  $K$  or not. It suffices to show that any basis element  $c \in \text{Sel}_\pi(E^\chi)$  maps the arithmetic Frobenius  $\tau \in \text{Gal}(\overline{\mathbb{F}_q}/\mathbb{F}_q)$  to the identity element of  $E[p]$ . Consider the local Kummer map  $\text{im}\delta_\infty^\chi$  at the place  $\infty$ . Because  $E$  has split multiplicative reduction at  $\infty$ ,  $E$  is a Tate curve at  $\infty$ . This implies that there exists an element  $q \in K_\infty^\times$  with positive valuation such that the  $\overline{K_\infty}$ -rational points of  $E$  is given by

$$E(\overline{K_\infty}) \cong \overline{K_\infty}^\times / \langle q \rangle,$$

which implies for any positive number  $n$ ,

$$E[n](\overline{K_\infty}) \cong \langle q^{\frac{1}{n}}, \mu_n \rangle / \langle q \rangle,$$

see for example [Section 3.3][BLV09] for a detailed discussion on these results. To analyze the condition that the basis element  $c \in \text{Sel}_\pi(E^\chi)$  maps the arithmetic Frobenius  $\tau \in \text{Gal}(\overline{\mathbb{F}_q}/\mathbb{F}_q)$  to the identity element of  $E[p]$ , it suffices to verify that  $Q^\tau - Q = O$  for  $Q \in E[p](\overline{K_\infty})$ . This follows from the assumption that the constant field of  $\overline{K_\infty}$  contains the primitive  $p$ th-root of unity.

### Frobenius conjugacy class

Let  $d_{i,j}$  be the constant given by the following:

$d_{i,j}$	$j = 0$	$j = 1$	$j = 2$
$i = 0$	1	$\times$	$\times$
$i = 1$	$p^{-\text{rk}(\chi)}$	$1 - p^{-\text{rk}(\chi)}$	$\times$
$i = 2$	$p^{-2\text{rk}(\chi)}$	$(p+1)(p^{-\text{rk}(\chi)} - p^{-2\text{rk}(\chi)})$	$1 - (p+1)p^{-\text{rk}(\chi)} + p^{1-2\text{rk}(\chi)}$

Using the techniques of the proof from [KMR14, Proposition 9.4], one can show that the non-zero values of  $d_{i,j}$  from the table above are in fact ratios between two subsets  $S_{i,j}, S'_i \subset \text{Gal}(F_{\sigma,\chi}/K)$  stable under conjugation, i.e.  $d_{i,j} = \frac{\#S_{i,j}}{\#S'_i}$ . These subsets satisfy the condition that

$$\begin{cases} \mathfrak{v} \in \mathcal{P}_i(d) & \iff \text{Frob}_{\mathfrak{v}} \in S'_i \\ \dim_{\mathbb{F}_p} \text{im}\delta_{\mathfrak{v}}^\chi = j \text{ and } \mathfrak{v} \in \mathcal{P}_i(d) & \iff \text{Frob}_{\mathfrak{v}} \in S_{i,j} \end{cases} \quad (27)$$

We refer to [KMR14, Proposition 9.4] for a detailed description of what these subsets are in  $\text{Gal}(F_{\sigma,\chi}/K)$ .

### Effective error bounds

Because the constant field of  $F_{\sigma,\chi}$  is  $\mathbb{F}_q$ , we can use Theorem 3.1 to bound the error terms of the following equation:

$$\left| \frac{\#\{\mathfrak{v} \in \mathcal{P}_i(d) \mid \mathfrak{v} \nmid \sigma, \mathfrak{v} \notin \Sigma, \dim_{\mathbb{F}_p} \text{im}\delta_{\mathfrak{v}}^\chi = j\}}{\#\{\mathfrak{v} \in \mathcal{P}_i(d) \mid \mathfrak{v} \nmid \sigma, \mathfrak{v} \notin \Sigma\}} - d_{i,j} \right|. \quad (28)$$

To prove the first statement of the proposition, choose large enough  $d$  which satisfies

$$d > (8 \ln p + 8 \ln 2 + 2 \ln \Delta_E) \deg \sigma$$

Applying Theorem 3.1 gives (24).

To prove the second statement of the proposition, suppose that the number of prime factors of  $\sigma$  satisfy the condition that

$$w(\sigma) \leq 2 \ln(\deg \sigma) - 2,$$

One caveat of naively applying Theorem 3.1 is that the degree of the field extension  $F_{\sigma, \chi}/K$  may grow arbitrarily large as  $\deg \sigma$  grows arbitrarily large. This implies that the error bounds from Corollary 3.2 may be unbounded.

Nevertheless, the condition on  $w(\sigma)$  implies that the error bounds from Corollary 3.2 are bounded given any choice of the local character  $\chi$  and a square-free product of places  $\sigma$ . Recall that  $F_{\sigma, \chi}/K$  is unramified away from  $v \in \Sigma$  or  $v \mid \sigma$ . Then

$$\text{rk}(\chi) \leq 4 \ln(\deg \sigma).$$

This implies that there exists a fixed constant  $C > 0$  independent of  $\deg \sigma$  such that

$$|\text{Gal}(F_{\sigma, \chi}/K)| = (p^2)^{\text{rk}(\chi)} |T|(p^3 - p) \leq Cp^{8 \ln(\deg \sigma)} = C(\deg \sigma)^{8 \ln p}.$$

We can also use Riemann-Hurwitz theorem to show that

$$g_{F_{\sigma, \chi}} \leq \deg \Delta_E \cdot (p^2)^{\text{rk}(\chi)} |T|(p^3 - p) \cdot \deg \sigma \leq C \deg \Delta_E \cdot (\deg \sigma)^{8 \ln p + 1}.$$

Then Corollary 3.2 implies that for any  $d$  satisfying

$$d > 2 \frac{\ln 8 + \ln(2C \deg \Delta_E \deg \sigma p^{2w(\sigma)})}{\ln q} = 2 \frac{4 \ln 2 + \ln C + \ln \deg \Delta_E + \ln \deg \sigma + 2w(\sigma) \ln p}{\ln q} \quad (29)$$

there exists a fixed constant  $\alpha > 0$  such that

$$\left| \frac{\#\{\mathfrak{v} \in \mathcal{P}_i(d) \mid \mathfrak{v} \nmid \sigma, \mathfrak{v} \notin \Sigma, \dim_{\mathbb{F}_p} \text{im} \delta_{\mathfrak{v}}^{\chi} = j\}}{\#\{\mathfrak{v} \in \mathcal{P}_i(d) \mid \mathfrak{v} \nmid \sigma, \mathfrak{v} \notin \Sigma\}} - d_{i,j} \right| < \alpha \frac{|S_{i,j}|}{|S'_{i,j}|} 2 \deg \Delta_E \bullet \deg \sigma \bullet p^{2 \text{rk}(\chi)} \bullet q^{-\frac{d}{2}}.$$

Pick  $i$  and  $j$  such that  $d_{i,j} \leq (p+1)p^{-\text{rk}(\chi)}$ . These cases occur except for  $(i, j) = (1, 1)$  and  $(2, 2)$ . Then for each fixed  $i$ , we obtain that

$$\begin{aligned} \left| \frac{\#\{\mathfrak{v} \in \mathcal{P}_i(d) \mid \mathfrak{v} \nmid \sigma, \mathfrak{v} \notin \Sigma, \dim_{\mathbb{F}_p} \text{im} \delta_{\mathfrak{v}}^{\chi} = j\}}{\#\{\mathfrak{v} \in \mathcal{P}_i(d) \mid \mathfrak{v} \nmid \sigma, \mathfrak{v} \notin \Sigma\}} - d_{i,j} \right| &< 2\alpha \deg \Delta_E (p+1) p^{\text{rk}(\chi)} \bullet \deg \sigma \bullet q^{-\frac{d}{2}} \\ &< 2\alpha \deg \Delta_E (p+1) p^{w(\sigma)} \bullet \deg \sigma \bullet q^{-\frac{d}{2}}. \end{aligned}$$

Choose  $d$  to be any number satisfying

$$d > \frac{4(\ln \deg \sigma)^2}{\ln q}.$$

We note that for any large enough  $\deg \sigma$ , any  $d$  satisfying the above inequality also satisfies (29) because  $w(\sigma) \leq 2 \ln(\deg \sigma) - 2$ . Therefore,

$$\begin{aligned} \left| \frac{\#\{\mathfrak{v} \in \mathcal{P}_i(d) \mid \mathfrak{v} \nmid \sigma, \mathfrak{v} \notin \Sigma, \dim_{\mathbb{F}_p} \text{im} \delta_{\mathfrak{v}}^{\chi} = j\}}{\#\{\mathfrak{v} \in \mathcal{P}_i(d) \mid \mathfrak{v} \nmid \sigma, \mathfrak{v} \notin \Sigma\}} - d_{i,j} \right| &< 2\alpha \deg \Delta_E (p+1) p^{2 \ln \deg \sigma} \bullet \deg \sigma \bullet q^{-\frac{d}{2}} \\ &< 2\alpha \deg \Delta_E (p+1) \frac{(\deg \sigma)^{2 \ln p + 1}}{q^{\frac{2(\ln \deg \sigma)^2}{\ln q}}} \\ &< 2\alpha \deg \Delta_E (p+1) (\deg \sigma)^{-2 \ln \deg \sigma + 2 \ln p + 1} \end{aligned}$$

We can then use (26) to prove the first equation of the second statement of the proposition.

### Local quadratic twists

Let  $d_i$  be the rational numbers given by

$$d_i = \begin{cases} 1 - \frac{p}{|T|(p^2-1)} & \text{if } i = 0 \\ \frac{1}{|T|p} & \text{if } i = 1 \\ \frac{1}{|T|(p^3-p)} & \text{if } i = 2 \end{cases}$$

for some cyclic finite group  $T$ . Note that these probabilities correspond to those from Remark 4.3.

For any fixed  $j$ , we can use Remark 4.3 to deduce the following equation:

$$\begin{aligned} & \left| \frac{\sum_{\substack{\mathbf{v} \in \mathcal{P}(d) \\ \mathbf{v} \nmid \sigma, \mathbf{v} \notin \Sigma}} \#\{\chi' \in \Omega_{\chi, \mathbf{v}} \mid \text{rk}(\chi') - \text{rk}(\chi) = j\}}{\sum_{\substack{\mathbf{v} \in \mathcal{P}(d) \\ \mathbf{v} \nmid \sigma, \mathbf{v} \notin \Sigma}} \#\Omega_{\chi, \mathbf{v}}} - (d_0 c_{0,j} + d_1 c_{1,j} + d_2 c_{2,j}) \right| \\ & \leq \sum_{i=0}^2 \left| \frac{\sum_{\substack{\mathbf{v} \in \mathcal{P}_i(d) \\ \mathbf{v} \nmid \sigma, \mathbf{v} \notin \Sigma}} \#\{\chi' \in \Omega_{\chi, \mathbf{v}} \mid \text{rk}(\chi') - \text{rk}(\chi) = j\}}{\sum_{\substack{\mathbf{v} \in \mathcal{P}_i(d) \\ \mathbf{v} \nmid \sigma, \mathbf{v} \notin \Sigma}} \#\Omega_{\chi, \mathbf{v}}} \frac{\sum_{\substack{\mathbf{v} \in \mathcal{P}_i(d) \\ \mathbf{v} \nmid \sigma, \mathbf{v} \notin \Sigma}} \#\Omega_{\chi, \mathbf{v}}}{\sum_{\substack{\mathbf{v} \in \mathcal{P}(d) \\ \mathbf{v} \nmid \sigma, \mathbf{v} \notin \Sigma}} \#\Omega_{\chi, \mathbf{v}}} - d_i c_{i,j} \right| \\ & = \sum_{i=0}^2 \left| \frac{\sum_{\substack{\mathbf{v} \in \mathcal{P}_i(d) \\ \mathbf{v} \nmid \sigma, \mathbf{v} \notin \Sigma}} \#\{\chi' \in \Omega_{\chi, \mathbf{v}} \mid \text{rk}(\chi') - \text{rk}(\chi) = j\}}{\sum_{\substack{\mathbf{v} \in \mathcal{P}_i(d) \\ \mathbf{v} \nmid \sigma, \mathbf{v} \notin \Sigma}} \#\Omega_{\chi, \mathbf{v}}} \frac{\#\{\mathbf{v} \in \mathcal{P}_i(d) \mid \mathbf{v} \nmid \sigma, \mathbf{v} \notin \Sigma\}}{\#\{\mathbf{v} \in \mathcal{P}(d) \mid \mathbf{v} \nmid \sigma, \mathbf{v} \notin \Sigma\}} - d_i c_{i,j} \right| \\ & \leq \sum_{i=0}^2 \left( \frac{\#\{\mathbf{v} \in \mathcal{P}_i(d) \mid \mathbf{v} \nmid \sigma, \mathbf{v} \notin \Sigma\}}{\#\{\mathbf{v} \in \mathcal{P}(d) \mid \mathbf{v} \nmid \sigma, \mathbf{v} \notin \Sigma\}} \cdot \left| \frac{\sum_{\substack{\mathbf{v} \in \mathcal{P}_i(d) \\ \mathbf{v} \nmid \sigma, \mathbf{v} \notin \Sigma}} \#\{\chi' \in \Omega_{\chi, \mathbf{v}} \mid \text{rk}(\chi') - \text{rk}(\chi) = j\}}{\sum_{\substack{\mathbf{v} \in \mathcal{P}_i(d) \\ \mathbf{v} \nmid \sigma, \mathbf{v} \notin \Sigma}} \#\Omega_{\chi, \mathbf{v}}} - c_{i,j} \right| + \left| \frac{\#\{\mathbf{v} \in \mathcal{P}_i(d) \mid \mathbf{v} \nmid \sigma, \mathbf{v} \notin \Sigma\}}{\#\{\mathbf{v} \in \mathcal{P}(d) \mid \mathbf{v} \nmid \sigma, \mathbf{v} \notin \Sigma\}} - d_i \right| \cdot c_{i,j} \right) \\ & < C(\deg \sigma)^{-\ln \deg \sigma} + Cq^{-\frac{2(\ln \deg \sigma)^2}{\ln q}} < 2C(\deg \sigma)^{-\ln \deg \sigma}. \end{aligned}$$

□

**Remark 4.6.** The technical condition on the degree of the place  $\mathbf{v}$  from (24) and (25) will be used in the upcoming sections when we compute the probability distribution of  $\pi$ -Selmer ranks of elliptic curves twisted by cyclic order- $p$  characters associated to  $p$ -th power free polynomials  $f$  of large enough degree  $n$ . For almost every  $f \in F_n(\mathbb{F}_q)$ , the degrees of distinct irreducible factors of  $f$  satisfy (25) by Theorem 3.4. This will allow us to compute the probability distribution of  $\pi$ -Selmer rank of the cyclic order- $p$  twists of  $E$  from  $\pi$ -Selmer ranks of cyclic twists  $E^{X_{g_i}}$  for every irreducible factor  $g_i$  of  $f$ .

**Remark 4.7.** Proposition 4.5 states that if  $\text{Gal}(K(E[p])/K) \supset \text{SL}_2(\mathbb{F}_p)$ , then the Chebotarev density theorem completely determines the variations of  $\pi$ -Selmer groups of elliptic curves twisted by local cyclic order- $p$  characters. This is not the case if the Galois group  $\text{Gal}(K(E[p])/K)$  does not contain  $\text{SL}_2(\mathbb{F}_p)$ , as carefully studied in [FIMR13] and [Smi22b]. For example, suppose that  $p = 2$  and  $\text{Gal}(K(E[p])/K) = \mathbb{Z}/3\mathbb{Z}$ . Friedlander, Iwaniec, Mazur, and Rubin showed that the variation of 2-Selmer groups of certain subfamilies of quadratic twists of elliptic curves are governed by the spin of odd principal prime ideals defined over totally real cyclic Galois extensions [FIMR13, Chapter 3, Chapter 10]. Smith uses a generalized notion of spin of prime ideals called “symbols of prime ideals” [Smi22b, Definition 3.11, Proposition 3.14] to classify which classes of prime ideals equivalently varies the Selmer groups of twistable modules, a generalized notion of quadratic twist families of abelian varieties [Smi22b, Chapter 4]. Thankfully, Proposition 4.5 demonstrates that one does not require to use the spin of prime ideals to determine the variations of  $\pi$ -Selmer groups of cyclic prime twist families of non-isotrivial elliptic curves.

**4.2. Governing Markov operator.** We will use the Markov operator constructed from [KMR14], known as the mod  $p$  Lagrangian operator, to analyze variations of  $\pi$ -Selmer ranks of a subfamily of global quadratic twists of elliptic curves over  $K$  satisfying the conditions from Theorem 1.1.

**Definition 4.8.** Let  $M_L = [p_{r,s}]$  be the operator over the state space of non-negative integers  $\mathbb{Z}_{\geq 0}$  given by

$$p_{r,s} = \begin{cases} 1 - p^{-r} & \text{if } s = r - 1 \geq 0 \\ p^{-r} & \text{if } s = r + 1 \\ 0 & \text{else} \end{cases}$$

**Remark 4.9.** The construction of the mod  $p$  Lagrangian Markov operator dates back to previous works by [SD08] and [KMR14]. Other references such as [Smi17], [Smi20], and [FLR20] also use Markov chains to obtain the probability distribution of  $l$ -Selmer groups of certain families of elliptic curves.

We list some crucial properties the operator  $M_L$  satisfies, the proof of which can be found in [KMR14, Section 2].

**Definition 4.10.** Let  $\mu : \mathbb{Z}_{\geq 0} \rightarrow [0, 1]$  be a probability distribution over the state space of non-negative integers  $\mathbb{Z}_{\geq 0}$ . The parity of  $\mu$  is the sum of probabilities at odd state spaces, i.e.

$$\rho(\mu) := \sum_{n \text{ odd}} \mu(n)$$

**Proposition 4.11.** [KMR14, Proposition 2.4]

Let  $E^+, E^- : \mathbb{Z}_{\geq 0} \rightarrow [0, 1]$  be probability distributions such that

$$E^+(n) = \begin{cases} \prod_{j=1}^{\infty} (1 + p^{-j})^{-1} \prod_{j=1}^n \frac{p}{p^j - 1} & \text{if } n \text{ even} \\ 0 & \text{if } n \text{ odd} \end{cases}$$

$$E^-(n) = \begin{cases} 0 & \text{if } n \text{ even} \\ \prod_{j=1}^{\infty} (1 + p^{-j})^{-1} \prod_{j=1}^n \frac{p}{p^j - 1} & \text{if } n \text{ odd} \end{cases}$$

Let  $\mu : \mathbb{Z}_{\geq 0} \rightarrow [0, 1]$  be a probability distribution. Then

$$\lim_{k \rightarrow \infty} M_L^{2k}(\mu) = (1 - \rho(\mu))E^+ + \rho(\mu)E^-$$

$$\lim_{k \rightarrow \infty} M_L^{2k+1}(\mu) = \rho(\mu)E^+ + (1 - \rho(\mu))E^-.$$

In particular, if  $\rho(\mu) = \frac{1}{2}$ , then

$$\lim_{k \rightarrow \infty} M_L^k(\mu)(n) = \prod_{j \geq 0} (1 + p^{-j})^{-1} \prod_{j=1}^n \frac{p}{p^j - 1} \quad (30)$$

**Remark 4.12.** Note that  $M_L^2$  is an aperiodic, irreducible, and positive-recurrent Markov chain over the state space of positive odd integers  $\mathbb{Z}_{\text{odd}, \geq 0}$  and non-negative even integers  $\mathbb{Z}_{\text{even}, \geq 0}$ . The unique stationary distributions of the Markov chain are  $E^-(n)$  and  $E^+(n)$ , respectively.

Given that  $M_L^2$  is aperiodic, irreducible, and positive-recurrent, it is natural to ask what the rate of convergence of  $M_L$  is. Assuming certain conditions on the initial probability distribution over the state space and the stationary distribution of  $M$ , the geometric rate of convergence of  $M$  can be verified using the following theorem.

**Theorem 4.13** (Geometric ergodic theorem for Markov chains). [MT93, Theorem 15.0.1]

Let  $M$  be an irreducible, aperiodic, and positive-recurrent Markov chain over a countable state space  $\mathcal{X} := (x_n)_{n \in \mathbb{Z}}$ . Let  $X_1, X_2, \dots, X_n, \dots : \mathcal{X} \rightarrow [0, 1]$  be a sequence of random variables which satisfies

$$X_{n+1} = M(X_n) \quad (31)$$

for all  $n$ . Let  $\pi$  be an invariant probability distribution of  $M$  (not necessarily unique). Suppose  $V : X \rightarrow [1, \infty)$  is a function such that  $\lim_{n \rightarrow \infty} V(x_n) = \infty$ . If there exists  $0 < \rho < 1$  and  $\kappa > 0$  such that for all but finitely many  $x_n \in \mathcal{X}$ ,

$$\mathbb{E}[V(M(X_n)) \mid X_n = x_k] - V(x_k) \leq -\rho V(x_k) + \kappa \quad (32)$$

then there exists a constant  $0 \leq \gamma < 1$  and a constant  $c > 0$  such that for any probability distribution  $\mu$  over  $X$  and every  $n \in \mathbb{N}$ ,

$$\sup_{z \in X} |M^n(\mu)(z) - \pi| < c\gamma^n(\mu V + 1)$$

where the term  $\mu V$  is the expected value of  $V$  under the probability distribution  $\mu$ , i.e.  $\mu V := \mathbb{E}[V(x) \mid x \in \mu]$ . In fact, one can choose  $\gamma = 1 - \rho$ .

Given a cyclic finite group  $T$ , Proposition 4.5 implies that the Markov chain

$$\left(1 - \frac{p}{|T|(p^2 - 1)}\right) + \frac{1}{|T|p}M_L + \frac{1}{|T|(p^3 - p)}M_L^2 \quad (33)$$

over the state space  $\mathbb{Z}_{\geq 0}$  governs the variation of  $\pi$ -Selmer ranks of families of elliptic curves twisted by local cyclic order- $p$  characters. To elaborate, the sequence of random variables  $X_n$  corresponds to the empirical probability distribution of dimensions of  $\pi$ -Selmer groups of a non-isotrivial elliptic curve  $E$  consecutively twisted by  $n$  local characters at places  $v$  satisfying the conditions from Proposition 4.5. The markov chain  $M_L$  provides a mechanism on how the empirical probability distribution of dimensions of  $\pi$ -Selmer groups of  $E$  consecutively twisted by  $n + 1$  local characters can be obtained from that computed over families of  $E$  consecutively twisted by  $n$  local characters.

Proposition 4.11 also shows that regardless of the parity of the initial probability distribution over the state space  $\mathbb{Z}_{\geq 0}$ , the stationary distribution of the Markov chain from (33) is given by the Poonen-Rains distribution as stated in (30). One can also show that given a fixed prime number  $p$ , the Markov chain of our interest is an irreducible aperiodic Markov chain over the countably infinite state space  $\mathbb{Z}_{\geq 0}$ . In fact, it is geometrically ergodic over  $\mathbb{Z}_{\geq 0}$  (without requiring the restriction that  $p = 2$ ).

**Corollary 4.14.** *Let  $\mu : \mathbb{Z}_{\geq 0} \rightarrow [0, 1]$  be a probability distribution over the state space  $\mathbb{Z}_{\geq 0}$ . Denote by  $\pi$  the stationary probability distribution of the Markov operator given by*

$$M := \left(1 - \frac{p}{|T|(p^2 - 1)}\right) + \frac{1}{|T|p}M_L + \frac{1}{|T|(p^3 - p)}M_L^2. \quad (34)$$

for some fixed prime number  $p$  and a finite cyclic group  $T$ . Fix any positive number  $\beta > 1$ . Then for every  $n \in \mathbb{N}$ , there exists a constant  $0 \leq \gamma < 1$  and a constant  $c > 0$  such that

$$\sup_{z \in X} \left| \left( \left(1 - \frac{p}{|T|(p^2 - 1)}\right) + \frac{1}{|T|p}M_L + \frac{1}{|T|(p^3 - p)}M_L^2 \right)^n (\mu) - \pi \right| < c\gamma^n(\beta^\mu + 1) \quad (35)$$

where the term  $\beta^\mu$  is the expected value of the function  $V(x) = \beta^x$  under the probability distribution  $\mu$ , i.e.  $\beta^\mu := \mathbb{E}[\beta^x \mid x \in \mu]$ . Explicitly, the rate of convergence  $\gamma$  satisfies

$$1 - \frac{1}{|T|} \frac{p}{p^2 - 1} < \gamma < 1 \quad (36)$$

*Proof.* Fix any positive number  $\beta > 1$ . Set  $V(x) = \beta^x$ . Computational results then show that there exists a fixed constant  $\kappa > 0$  such that for every  $x \in \mathbb{Z}_{\geq 0}$ ,

$$\mathbb{E}[\beta^{X_n} \mid X_{n-1} = x] - \beta^x = - \left( \frac{p}{|T|(p^2 - 1)} - \frac{1}{|T|p\beta} - \frac{1}{|T|(p^3 - p)\beta^2} \right) \cdot \beta^x + \kappa.$$

Setting  $\rho = 1 - \frac{p}{|T|(p^2 - 1)} + \frac{1}{|T|p\beta} + \frac{1}{|T|(p^3 - p)\beta^2}$ , the corollary follows from Theorem 4.13.  $\square$

In particular, the above corollary implies that the rate of convergence of the markov chain  $M$  from (33) is given by  $1 - \frac{p}{|T|(p^2-1)} + \epsilon$  for some positive number  $\epsilon > 0$ . A common choice of  $\beta$  is to take  $\beta = p$ , which yields the rate of convergence to be equal to  $\gamma = 1 - \frac{1}{|T|} \frac{p^2-p+1}{p^3}$ . For example, when  $p = 2$  and  $|T| = 1$ , the rate of convergence of  $M$  is greater than  $\frac{1}{3}$ , and is given by  $\frac{5}{8}$  when one chooses  $\beta = 2$ .

It now remains to show that the stationary distribution of the desired Markov chain (33) is the probability distribution conjectured by Bhargava-Kane-Lenstra-Poonen-Rains [BKLJ<sup>+</sup>15].

**Lemma 4.15.** *Let  $p$  be any fixed value of prime, and let  $T$  be a finite cyclic group. Then the probability distribution*

$$PR(j) := \prod_{j \geq 0}^{\infty} (1 + p^{-j})^{-1} \prod_{j=1}^n \frac{p}{p^j - 1} \quad (37)$$

*is the unique stationary distribution of the Markov chain*

$$M := \left(1 - \frac{p}{|T|(p^2-1)}\right) + \frac{1}{|T|p} M_L + \frac{1}{|T|(p^3-p)} M_L^2. \quad (38)$$

*Proof.* Note that the operators  $\text{Id}$  and  $M_L^2$  are parity preserving Markov operators, whereas  $M_L$  is a parity reversing Markov operator. Because the Markov chain of our interest is aperiodic and irreducible, it follows that the Markov chain has a unique stationary distribution  $\pi$ . The following relation holds for the parity of  $\pi$ , which is obtainable by comparing the parity between  $\pi$  and  $M(\pi)$ .

$$\rho(\pi) = \left(1 - \frac{1}{|T|p}\right) \rho(\pi) + \frac{1}{|T|p} (1 - \rho(\pi)) = \left(1 - \frac{2}{|T|p}\right) \rho(\pi) + \frac{1}{|T|p}. \quad (39)$$

Therefore, we obtain that  $\rho(\pi) = \frac{1}{2}$ . Using Proposition 4.11 and the fact that the Markov chain  $M$  is aperiodic and irreducible, we immediately obtain the statement of the lemma.  $\square$

**Remark 4.16.** One crucial result from using Corollary 4.14 and Lemma 4.15 is that the stationary distribution of applying the Markov chain from (33) is equal to the Bhargava-Kane-Lenstra-Poonen-Rains distribution regardless of the initial probability distribution. Furthermore, as long as the initial probability distribution is finitely supported, we can also ensure that the Markov chain converges to the stationary distribution at a geometric convergence rate. These two observations will be of crucial importance when we analyze the variations of  $\pi$ -Selmer groups with respect to global cyclic order- $p$  twists of elliptic curves.

**Remark 4.17.** We note that the Markov chain constructed from Smith's work is different from the Markov chain presented in this manuscript [Smi22b, Smi22a]. The sequence of random variables  $X_n$  Smith considers correspond to the empirical probability distribution of the subspace

$$\dim_{\mathbb{F}_p} \pi^{n-1} \text{Sel}_{\pi^n}(E^\chi) \subset \text{Sel}_\pi(E) \quad (40)$$

where  $\chi$  ranges over grids of twists [Smi22b, Chapter 6]. Here, the grids of twists are defined as a finite Cartesian product of collections of prime ideals, where each collection contains prime ideals whose symbols are equal to each other [Smi22b, Definition 4.13].

To elaborate, this manuscript regards the variable  $n$  from a sequence of random variables  $\{X_n\}_{n \in \mathbb{Z}}$  as the number of distinct irreducible places, whereas Smith's work regards the variable  $n$  from a sequence of random variables  $\{X_n\}_{n \in \mathbb{Z}}$  as a quantifier for detecting elements inside higher  $\pi^n$ -Selmer groups which also lie inside the  $\pi$ -Selmer group of  $E$ .



## 5. EQUIDISTRIBUTION OF UNRAMIFIED CHARACTERS AND AUXILIARY PRIMES

We now present how one can use variations of  $\pi$ -Selmer groups over local cyclic order- $p$  twist families of elliptic curves  $E$  to obtain those over global twist families of  $E$ . To rigorously model the effects of twisting  $E$  by global quadratic characters of bounded height on the dimension of  $\pi$ -Selmer groups, we define what is called the splitting partition of a  $p$ -th power-free polynomial of degree  $n$ .

## 5.1. Splitting partition of polynomials over finite fields.

**Definition 5.1** (Splitting partition given  $N$ ). Fix an integer  $N < n$  and an elliptic curve  $E$  over  $\mathbb{F}_q(t)$ . Let  $f \in F_n(\mathbb{F}_q)$  be a  $p$ -th power-free polynomial of degree  $n$  coprime to  $\Delta_E$ . Denote by  $\mathcal{F}_{q,i}$  the set of monic irreducible polynomials of degree  $i$  over  $\mathbb{F}_q$  coprime to  $\Delta_E$ .

A list of non-negative integers  $[\mu, \lambda_1, \lambda_2, \lambda_3]$  is called a splitting partition of  $f$  given  $N$  if it defines corresponding  $p$ -th power-free polynomials  $f_\mu, f_{\lambda_1}, f_{\lambda_2}, f_{\lambda_3}$  such that:

- (1)  $\mu + \lambda_1 + \lambda_2 + \lambda_3 = n$  and  $f = f_\mu f_{\lambda_1} f_{\lambda_2} f_{\lambda_3}$ .
- (2) Denote by  $f_\mu, f_{\lambda_1}, f_{\lambda_2}, f_{\lambda_3}$  the  $p$ -th power-free polynomials:

$$f_\mu = u \times \prod_{\substack{v|f \\ v \in \cup_{i=1}^N \mathcal{F}_{q,i}(\mathbb{F}_q)}} v^{l_v}, \quad f_{\lambda_1} = \prod_{\substack{v|f \\ v \in \cup_{i>N} \mathcal{F}_{q,i}(\mathbb{F}_q)}} v^{l_v}, \quad f_{\lambda_2} = \prod_{\substack{v|f \\ v \in \cup_{i>N} \mathcal{F}_{q,i}(\mathbb{F}_q)}} v^{l_v}, \quad f_{\lambda_3} = \prod_{\substack{v|f \\ v \in \cup_{i>N} \mathcal{F}_{q,i}(\mathbb{F}_q)}} v^{l_v},$$

where  $u \in \mathbb{F}_q^\times / (\mathbb{F}_q^\times)^p$  and  $1 \leq l_v \leq p-1$  for each corresponding place  $v$ . The polynomial  $f_\mu$  is the product of bounded powers of all monic irreducible factors of  $f$  of degree at most  $N$ , whereas the polynomial  $f_{\lambda_i}$  is the product of all bounded powers of monic irreducible factors of  $f$  of degree greater than  $N$  which lives inside  $\mathcal{P}_i$ 's.

- (3) Denote by  $\mu, \lambda_1, \lambda_2, \lambda_3$  to be the degrees of the polynomials  $f_\mu, f_{\lambda_1}, f_{\lambda_2}, f_{\lambda_3}$ , respectively.

**Definition 5.2.** Fix an integer  $N < n$ . Given a fixed square-free polynomial  $g \in \mathbb{F}_q[t]$ , denote by  $F_{n,g,N}$  the set of polynomials of degree  $n$  over  $\mathbb{F}_q$  defined as

$$F_{n,g,N} := \{f \in F_n(\mathbb{F}_q) \mid f_\mu \text{ divides } g, \deg h \leq N \text{ for any irreducible factor } h \text{ of } f_\mu\} \quad (41)$$

One can take  $g$  to be a constant in  $\mathbb{F}_q$ . The fixed elements of  $F_{n,g,N}$  with respect to the Galois action  $\text{Gal}(\overline{\mathbb{F}_q}/\mathbb{F}_q)$  are the respective square-free polynomials over  $\mathbb{F}_q$ . With abuse of notation, we denote by  $F_{n,g,N}(\mathbb{F}_q)$  the set of such square-free polynomials in  $F_{n,g,N}$ .

For example,  $F_{n,1,N}(\mathbb{F}_q)$  consists of monic square-free polynomials over  $\mathbb{F}_q$  all of whose irreducible factors are of degree greater than  $N$ .

**Remark 5.3.** Given a fixed integer  $n$ , suppose  $f \in F_{n,g,N}(\mathbb{F}_q)$ . Let  $N$  be the smallest integer  $N$  that satisfies

$$N \geq \frac{4(\ln n)^2}{\ln q}.$$

The splitting partition of  $f$  gives rise to the factorization  $f = f_\mu f_{\lambda_1} f_{\lambda_2} f_{\lambda_3}$ , where  $f_\mu$  is the product of all irreducible factors of  $f$  of degree at most  $N$ . Let  $g$  be any polynomial dividing  $f$  such that  $f_\mu$  divides  $g$ . Let  $h$  be any irreducible factor of  $f_{\lambda_i}$ 's. We first show that  $g$  cannot be the product of all irreducible factors of degree at most  $N$ . Given an integer  $k$ , the polynomial  $x^{q^k} - x$  is the product of all irreducible factors of degree dividing  $k$ . This implies that

$$\deg g \geq q^N.$$

Therefore, as long as  $n \geq 2$ ,

$$\deg g \geq q^{\frac{4(\ln n)^2}{\ln q}} = q^{\log_q n \times 4(\ln n)} = n^{4 \ln n}$$

which contradicts the condition that  $\deg g < \deg f$ . Thus, it cannot be the case that  $g$  is the product of all irreducible factors of degree at most  $N$ .

Since every irreducible factor of  $f_{\lambda_i}$ 's are of degree at least  $N + 1$ , the polynomials  $h$  satisfy

$$\deg h \geq N + 1 > \frac{4(\ln n)^2}{\ln q} > \frac{4(\ln(\deg g))^2}{\ln q}.$$

which is the condition for Proposition 4.5.

It is then natural to ask how the probability distribution of  $\pi$ -Selmer groups vary as we consecutively twist the elliptic curve  $E$  with global cyclic order- $p$  characters associated to an irreducible polynomial  $f$ . We note that Proposition 4.5 unfortunately only specifies how the irreducible factors of  $f$  of degree greater than  $N$  vary the probability distribution of  $\pi$ -Selmer groups. Nevertheless, given that  $w$ , the number of distinct irreducible factors of  $f$ , is between  $\rho \ln n$  and  $(2 - \rho) \ln n$  for some positive  $0 < \rho < 1$ , the probability that  $f_\mu$  is a product of more than any positive fraction  $0 < \epsilon < 1$  of the number of distinct irreducible factors of  $f$  converges to 0 if  $N = n^{\frac{1}{3}}$ . (We note that the proposition still holds for the case where  $N > \frac{4(\ln n)^2}{\ln q}$ , the bound we obtained from the aforementioned remark.)

**Proposition 5.4.** *Let  $N$  be the smallest integer which is at least  $n^{\frac{1}{3}}$ . Fix a positive number  $0 < \epsilon < 1$  and a positive number  $0 < \rho < 1$ . Then for sufficiently large  $n$ , there exists a fixed constant  $C > 0$  such that*

$$\mathbb{P} \left[ w(f_\mu) > \epsilon w(f) \mid \substack{f = f_\mu f_{\lambda_1} f_{\lambda_2} f_{\lambda_3} \in F_n(\mathbb{F}_q) \\ \rho \ln n \leq w(f) \leq (2 - \rho) \ln n} \right] < C n^{-\frac{\rho}{2} \epsilon \ln \ln n}, \quad (42)$$

where

$$\mathbb{P} \left[ w(f_\mu) > \epsilon w(f) \mid \substack{f = f_\mu f_{\lambda_1} f_{\lambda_2} f_{\lambda_3} \in F_n(\mathbb{F}_q) \\ \rho \ln n \leq w(f) \leq (2 - \rho) \ln n} \right] := \frac{\#\{f = f_\mu f_{\lambda_1} f_{\lambda_2} f_{\lambda_3} \in F_n(\mathbb{F}_q) \mid \substack{w(f_\mu) > \epsilon w(f) \\ \rho \ln n \leq w(f) \leq (2 - \rho) \ln n}\}}{\#\{f \in F_n(\mathbb{F}_q) \mid \rho \ln n \leq w(f) \leq (2 - \rho) \ln n\}} \quad (43)$$

*Proof.* Fix a positive integer  $w > 0$  such that  $\rho \ln n \leq w \leq (2 - \rho) \ln n$ . Fix a positive number  $0 < \epsilon < 1$ .

Denote by  $B_{low}$  the lower bound of the number of distinct factors of  $f_\mu$ . Recall that we used the notation  $\mu$  to denote the degree of  $f_\mu$ . Then the following relation holds.

$$\begin{aligned} & \#\{f \in F_n(\mathbb{F}_q) \mid w(f_\mu) \geq B_{low}, w(f) = w, f \text{ monic}\} \\ & \leq \sum_{i=B_{low}}^w \sum_{\mu=i}^{iN} \left( \sum_{1 \leq m_1 \leq \dots \leq m_{i-1} \leq \mu - m_1 - \dots - m_{i-1}} \left( \sum_{k=1}^{\mu - m_1 - \dots - m_{i-1}} \frac{q^k}{k} \right) \prod_{j=1}^{i-1} \frac{q^{m_j}}{m_j} \right) \\ & \quad \times \left( \sum_{N \leq m_{i+1} \leq \dots \leq m_{w-1} \leq n - \mu - m_{i+1} - \dots - m_{w-1}} \left( \sum_{k=1}^{n - \mu - m_{i+1} - \dots - m_{w-1}} \frac{q^k}{k} \right) \prod_{j=i+1}^{w-1} \frac{q^{m_j}}{m_j} \right) \\ & \leq \sum_{i=B_{low}}^w \sum_{\mu=i}^{iN} \left( 2 \cdot \sum_{1 \leq m_1 \leq \dots \leq m_{i-1} \leq \mu - m_1 - \dots - m_{i-1}} \frac{q^{\mu - m_1 - \dots - m_{i-1}}}{\mu - m_1 - \dots - m_{i-1}} \prod_{j=1}^{i-1} \frac{q^{m_j}}{m_j} \right) \\ & \quad \times \left( 2 \cdot \sum_{N \leq m_{i+1} \leq \dots \leq m_{w-1} \leq n - \mu - m_{i+1} - \dots - m_{w-1}} \frac{q^{n - \mu - m_{i+1} - \dots - m_{w-1}}}{n - \mu - m_{i+1} - \dots - m_{w-1}} \prod_{j=i+1}^{w-1} \frac{q^{m_j}}{m_j} \right) \end{aligned} \quad (44)$$

Using Lemma 3.6 and the proof of Proposition 3.7, we obtain

$$\begin{aligned} (44) & \leq \sum_{i=B_{low}}^w \sum_{\mu=i}^{iN} 16q^n \frac{1}{\mu(n - \mu)} \frac{1}{(i - 1)!(w - i - 1)!} (\ln(\mu - (i - 1)))^{i-1} \left( \ln \frac{n - \mu - N(w - i - 1)}{N} \right)^{w-i-1} \\ & \leq \sum_{i=B_{low}}^w \sum_{\mu=i}^{iN} 16q^n \frac{1}{\mu(n - \mu)} \frac{1}{\Gamma(i - 2)\Gamma(w - i - 2)} (\ln \mu)^{i-1} \left( \ln \frac{n}{N} \right)^{w-i-1} \end{aligned} \quad (45)$$

Using the Stirling's approximation formula for Gamma functions and the fact that the Gamma function is a strict log-convex function, we obtain:

$$\begin{aligned}
 (44) &\leq \sum_{i=B_{low}}^w \sum_{\mu=i}^{iN} 16q^n \frac{1}{\mu(n-\mu)} \frac{1}{\Gamma(\frac{w}{2}-2)^2} (\ln \mu)^{i-1} \left(\ln \frac{n}{N}\right)^{w-i-1} \\
 &\leq \sum_{i=B_{low}}^w \sum_{\mu=i}^{iN} \frac{8q^n}{\pi} \frac{1}{\mu(n-\mu)} \frac{e^{w-4}}{\left(\frac{w}{2}-2\right)^{w-\frac{9}{2}}} (\ln \mu)^{i-1} \left(\ln \frac{n}{N}\right)^{w-i-1} \\
 &\leq \sum_{i=B_{low}}^w \sum_{\mu=i}^{iN} \frac{8q^n}{\pi} \frac{1}{B_{low}(n-B_{low})} \frac{e^{w-4}}{\left(\frac{w}{2}-2\right)^{w-\frac{9}{2}}} \left(\frac{\ln wN}{\ln \frac{n}{N}}\right)^{i-1} \left(\ln \frac{n}{N}\right)^w \\
 &\leq \sum_{i=B_{low}}^w w(N-1) \frac{8q^n}{\pi} \frac{1}{B_{low}(n-B_{low})} \frac{e^{w-4}}{\left(\frac{w}{2}-2\right)^{w-\frac{9}{2}}} \left(\frac{\ln wN}{\ln \frac{n}{N}}\right)^{i-1} \left(\ln \frac{n}{N}\right)^w
 \end{aligned} \tag{46}$$

Note that the sum of the geometric sequence is bounded above by

$$\sum_{i=B_{low}}^w \left(\frac{\ln wN}{\ln \frac{n}{N}}\right)^{i-1} \leq 2 \left(\frac{\ln wN}{\ln \frac{n}{N}}\right)^{B_{low}-1} \tag{47}$$

This allows us to further reduce (44) as follows:

$$\begin{aligned}
 (44) &\leq \frac{16q^n}{\pi} \frac{w(N-1)}{B_{low}(n-B_{low})} \frac{e^{w-4}}{\left(\frac{w}{2}-2\right)^{w-\frac{9}{2}}} \left(\ln \frac{n}{N}\right)^w \left(\frac{\ln wN}{\ln \frac{n}{N}}\right)^{B_{low}-1} \\
 &\leq \frac{16q^n}{\pi} \frac{w(N-1)}{B_{low}(n-B_{low})} \frac{e^{w-4}}{\left(\frac{w}{2}-2\right)^{w-\frac{9}{2}}} \left(\ln \frac{n}{N}\right)^{w-B_{low}+1} \left(\frac{\ln wN}{\ln \frac{n}{N}}\right)^{B_{low}-1} \\
 &\leq \frac{16e^{\frac{1}{2}}q^n}{\pi} \frac{w(N-1)}{B_{low}(n-B_{low})} \left(\ln \frac{n}{N}\right)^{\frac{9}{2}} \left(\frac{2e \ln \frac{n}{N}}{w-4}\right)^{w-B_{low}-\frac{7}{2}} \left(\frac{2e \left(\frac{\ln wN}{\ln \frac{n}{N}}\right)}{w-4}\right)^{B_{low}-1}
 \end{aligned} \tag{48}$$

We now use the constraints that  $0 < \epsilon < 1$  and  $\frac{1}{2} \ln n < w < \frac{3}{2} \ln n$ . Fix  $N = n^{\frac{1}{3}}$ . Then for sufficiently large  $n$ ,

$$\frac{\ln \frac{n}{N}}{w-4} \leq \frac{2+1}{3\rho} = \frac{1}{\rho}, \quad \frac{2e \frac{\ln wN}{\ln \frac{n}{N}}}{w-4} \leq \frac{12}{w-4} \leq \frac{13}{\rho \ln n}.$$

Using the fact that  $B_{low} = \epsilon w$ , there exists a fixed constant  $\tilde{C} > 0$  such that for sufficiently large  $n$ ,

$$\begin{aligned}
 (44) &\leq \frac{16e^{\frac{1}{2}}q^n}{\pi} \frac{2n^{\frac{1}{3}} \ln n}{\epsilon w(n-\epsilon w)} \left(\frac{1}{\rho}\right)^{w-\epsilon w-\frac{7}{2}} \left(\frac{13}{\rho \ln n}\right)^{\epsilon w-1} \\
 &\leq \frac{16e^{\frac{1}{2}}q^n}{\pi} \left(\frac{1}{\rho}\right)^{(2-\rho) \ln n} \left(\frac{13}{\rho \ln n}\right)^{\rho \epsilon \ln n} \\
 &\leq \tilde{C} q^n n^{-\frac{\rho}{2} \epsilon \ln \ln n}.
 \end{aligned} \tag{49}$$

□

Proposition 5.4 shows that for sufficiently large  $n$ , the majority (but not almost every) of the distinct irreducible factors of  $f \in F_n(\mathbb{F}_q)$  have degrees at least  $N = n^{\frac{1}{3}}$ . As  $n$  grows sufficiently large, the number of such irreducible factors of degree at least  $N$  also grows sufficiently large. We

will observe that computing the variation of  $\pi$ -Selmer groups with respect to twisting the elliptic curve by global characters associated to these irreducible polynomials is enough to obtain the Bhargava-Kane-Lenstra-Poonen-Rains distribution.

**5.2. Equidistribution of local characters.** In Section 4.1, we made a crucial observation that the Markov chains constructed in Section 4.2 governs the variation of  $\pi$ -Selmer groups of families of an elliptic curve  $E$  twisted by local cyclic order- $p$  characters. To compute the probability distribution of  $\pi$ -Selmer groups of families of twists of elliptic curves, one needs to relate the variation of  $\pi$ -Selmer groups of global twist families of  $E$  with that of local twist families of  $E$ .

In this subsection, we prove that for sufficiently large  $n$ , the probability distribution that the set of global cyclic order- $p$  characters induced from the set of irreducible polynomials of degree  $n$  forms a uniform distribution when restricted to the set of finite cartesian products of local unramified cyclic order- $p$  characters at finitely many places of degree strictly less than  $n$ . This observation in particular will serve us to show that consecutively twisting the elliptic curve  $E$  by a randomly chosen polynomial  $f$  can be approximated by consecutively twisting  $E$  by a product of local cyclic characters ramified at places dividing  $f$  uniformly chosen at random over a certain subset of local ramified characters, with error bounds which converge to 0 as  $n$  grows sufficiently large. The probabilistic behavior of the restrictions of global characters over global function fields are well-studied, for instance as seen from the following Theorem from [Hsu98].

**Theorem 5.5** (Theorem 2.1, [Hsu98]). *Let  $h$  be any square-free polynomial over  $\mathbb{F}_q$ . Let  $\chi_h$  be a non-trivial character  $\chi : (\mathbb{F}_q[t]/h)^\times \rightarrow \mathbb{C}^\times$ . Recall that we denote by  $\mathcal{F}_{q,i}(\mathbb{F}_q)$  the set of monic irreducible polynomials of degree  $i$  over  $\mathbb{F}_q$  coprime to  $\Delta_E$ . Then*

$$\sum_{v \in \mathcal{F}_{q,i}(\mathbb{F}_q)} \chi(v) \leq (\deg h + 1) \frac{q^{\frac{i}{2}}}{i}. \quad (50)$$

An immediate corollary of the theorem above is that the effective error bounds of the density of whether the restriction of a global cyclic order- $p$  character associated to an irreducible polynomial forms a uniform distribution over the set of finite cartesian products of local unramified cyclic characters is given by the order of  $q^{-\frac{n}{2}}$ .

**Corollary 5.6.** *Fix a prime number  $p$ . Let  $K = \mathbb{F}_q(t)$  be a global function field such that  $\mu_p \subset \mathbb{F}_q$ . Fix  $w$  to be any integer such that  $w \leq 2 \ln n$ . Let  $h_1, h_2, \dots, h_w$  be irreducible polynomials over  $\mathbb{F}_q$ , all of degree at most  $n$ . Pick an integer  $i$  such that  $i > \max\{\frac{4(\ln n)^2}{\ln q}, \max_{i=1}^w \{\deg h_i\}\}$ . Given a place  $v$  of degree  $i$ , denote by  $(\frac{v}{h_i})_p \in \mu_p$  the  $p$ -th power residue symbol. Then for any element  $a \in \prod_{i=1}^w \mu_p$ ,*

$$\left| \frac{\#\{v \in \mathcal{F}_{q,i}(\mathbb{F}_q) \mid \prod_{i=1}^w (\frac{v}{h_i})_p = a \in \prod_{k=1}^w \mu_p\}}{\#\mathcal{F}_{q,i}(\mathbb{F}_q)} - \frac{1}{p^w} \right| < n^{-2 \ln n + 2}. \quad (51)$$

*Proof.* We outline an elementary proof using Theorem 5.5 for the case where  $p = 2$ . The cases for general prime numbers follow analogously. Let  $H := \{h_1, h_2, \dots, h_w\}$  be a given set of irreducible polynomials over  $\mathbb{F}_q$ , all of degree at most  $n$ . Then there are  $2^w$  square-free polynomials (including the trivial polynomial 1), all of whose irreducible factors lie in the given set. Denote by  $\tilde{H}$  the set of such square-free polynomials determined by irreducible polynomials in  $H$ . Fix an integer  $i \geq \frac{4(\ln n)^2}{\ln q}$ , a variable which denotes the degree of the irreducible polynomials  $v \in \mathcal{F}_{q,i}(\mathbb{F}_q)$ . To prove the statement of the theorem, it suffices to show that

$$\left| \frac{\#\{v \in \mathcal{F}_{q,i}(\mathbb{F}_q) \mid ((\frac{v}{h_k}))_{k=1}^w = a \in \prod_{k=1}^w \{\pm 1\}\}}{\#\mathcal{F}_{q,i}(\mathbb{F}_q)} - \frac{1}{2^w} \right| < n^{-2 \ln n + 2}. \quad (52)$$

where  $\left(\frac{v}{\cdot}\right)$  is the Legendre symbol. Denote by  $H_{a,i}$  the set

$$H_{a,i} := \{v \in \mathcal{F}_{q,i}(\mathbb{F}_q) \mid \left(\left(\frac{v}{h_k}\right)\right)_{k=1}^w = a \in \prod_{k=1}^w \{\pm 1\}\},$$

and by  $p_{a,i}$  the probability

$$p_{a,i} := \frac{\#\{v \in \mathcal{F}_{q,i}(\mathbb{F}_q) \mid \left(\left(\frac{v}{h_i}\right)\right)_{i=1}^w = a \in \prod_{i=1}^w \{\pm 1\}\}}{\#\mathcal{F}_{q,i}(\mathbb{F}_q)}.$$

We prove that

$$\left|p_{a,i} - \frac{1}{2^w}\right| < n^{-2 \ln n + 2}$$

for any  $a \in \prod_{k=1}^w \{\pm 1\}$ .

For each  $\tilde{h} \in \tilde{H}$ , one can associate a quadratic character  $\chi_{\tilde{h}} := \left(\frac{\cdot}{\tilde{h}}\right)$ . Theorem 5.5 states that for any  $\tilde{h} \in \tilde{H}$ , which is of degree at most  $n$ , there exists a fixed constant  $0 < C < 1$  such that

$$\frac{\sum_{v \in \mathcal{F}_{q,i}(\mathbb{F}_q)} \chi(v)}{\#\mathcal{F}_{q,i}(\mathbb{F}_q)} \leq C(\deg \tilde{h} + 1)q^{-\frac{i}{2}} \leq 2n^{-2 \ln n + 1}.$$

Meanwhile, given any nontrivial  $a \in \prod_{k=1}^w \{\pm 1\}$ , there exist exactly  $2^{w-1}$  elements  $\tilde{h} \in \tilde{H}$  such that  $\left(\frac{v}{\tilde{h}}\right) = 1$  for any  $v \in H_{a,i}$ .

For the trivial polynomial  $1 \in \tilde{H}$ , we have

$$\sum_{a \in \prod_{k=1}^w \{\pm 1\}} p_{a,i} = 1. \quad (53)$$

For any other non-trivial polynomial  $\tilde{h} \in \tilde{H}$ , we have

$$-2n^{-2 \ln n + 1} < \sum_{\substack{a \in \prod_{k=1}^w \{\pm 1\} \\ \left(\frac{v}{\tilde{h}}\right)=1 \text{ for any } v \in H_{a,i}}} p_{a,i} - \sum_{\substack{a \in \prod_{k=1}^w \{\pm 1\} \\ \left(\frac{v}{\tilde{h}}\right)=-1 \text{ for any } v \in H_{a,i}}} p_{a,i} < 2n^{-2 \ln n + 1}. \quad (54)$$

We note the fact that both summations from (53) have  $2^{w-1}$  terms. Taking the summation of (54) and (54) over all  $\tilde{h} \in \tilde{H}$  yields:

$$1 - (2^w - 1)n^{-2 \ln n + 2} < 2^w p_{0,i} < 1 + (2^w - 1)n^{-2 \ln n + 2}$$

Hence, we obtain that  $p_{0,i}$  satisfies

$$\left|p_{0,i} - \frac{1}{2^w}\right| < n^{-2 \ln n + 2}.$$

Likewise, we can compute other probabilities  $p_{a,i}$  given any non-trivial  $a \in \prod_{k=1}^w \{\pm 1\}$ . Suppose  $a, a'$  are two non-trivial distinct elements in  $\prod_{k=1}^w \{\pm 1\}$ . For each  $a \in \prod_{k=1}^w \{\pm 1\}$ , let  $\tilde{H}_a$  be the subset of  $\tilde{H}$  defined as

$$\tilde{H}_a := \{\tilde{h} \in \tilde{H} \mid \left(\frac{v}{\tilde{h}}\right) = 1 \text{ for any } v \in H_{a,i}\}.$$

Then  $|\tilde{H}_a \cap \tilde{H}_{a'}| = \frac{|\tilde{H}_a|}{2} = \frac{|\tilde{H}_{a'}|}{2}$ . (If  $a'$  is trivial, then the intersection is equal to  $\tilde{H}_a$ .) Using this fact, we can take the summation of equations as follows: We first take the summation of (53) and (54) over all  $\tilde{h} \in \tilde{H}_a$ , and take the subtraction of (54) over all  $\tilde{h} \in \tilde{H} - \tilde{H}_a$  afterwards. This yields:

$$1 - (2^w - 1)n^{-2 \ln n + 2} < 2^w p_{a,i} < 1 + (2^w - 1)n^{-2 \ln n + 2}$$

Hence, we obtain that  $p_{0,i}$  satisfies

$$\left|p_{a,i} - \frac{1}{2^w}\right| < n^{-2 \ln n + 2}.$$

□

**Remark 5.7.** We note that one can also prove the statement of the Corollary using the effective Chebotarev density theorem (Theorem 3.1), as suggested in [EH91, Chapter 5] or [Ros02, Chapter 3, Chapter 10]. Suppose that  $K = \mathbb{F}_q(t)$  is a global function field which contains the primitive  $p$ -th root of unity  $\mu_p$ . Consider the cyclic order- $p$  abelian extension  $K(\sqrt[p]{h})/K$  for some irreducible polynomial  $h$ . We note that if  $v$  is coprime to  $h$ , then the  $p$ -th power residue symbol  $\left(\frac{v}{h}\right)_p$  defines the action of the Frobenius element  $\text{Frob}_v$  on  $\sqrt[p]{h}$  via

$$\text{Frob}_v(\sqrt[p]{h}) = \left(\frac{v}{h}\right)_p \sqrt[p]{h}$$

which in fact originates from the definition of the Artin reciprocity map, see [Ros02, Chapter 3, Chapter 10] for a detailed description. The  $p$ -th power residue symbols therefore generates all possible unramified local cyclic order- $p$  characters inside  $\text{Hom}(\text{Gal}(\overline{K_h}/K), \mu_p)$ . Furthermore, the statement of the corollary implies that the restriction of global cyclic order- $p$  characters induces a uniform distribution over the set of finite cartesian products of unramified cyclic order- $p$  characters.

**5.3. Auxiliary places.** Now that we showed how global characters can be restricted to form a uniform distribution over the set of finite cartesian products of local characters, it remains to analyze how one can rearrange global characters ordered by degrees to align with the set of local characters  $\Omega_{\chi, \mathfrak{v}}$  or  $\Omega_{\chi, \mathfrak{v}}^\eta$  introduced from Section 4.1. To do so, we introduce what is called the auxiliary place  $\mathfrak{w}$ , which will play a key role in rearranging the global characters to a desired order which allows one to easily compute the variations of the ranks of  $\pi$ -Selmer groups with respect to consecutive local cyclic characters. Throughout this section we assume that the constant field of  $K$  contains the primitive  $p$ -th root of unity for a given prime  $p$ .

**Definition 5.8.** Fix integers  $n, N$ , an elliptic curve  $E/K$  satisfying the conditions from (18), and a prime  $p$ . Pick a polynomial  $g \in F_{n', 1, N}(\mathbb{F}_q)$  such that the degrees of irreducible factors of  $g$  lying in  $\mathcal{P}_0$  are at most  $n - \deg(g)$  such that  $n - \deg(g) \geq N$ . Denote by  $F_{n, 1, N}^{g, p, l}$  the set of polynomials

$$F_{n, 1, N}^{g, p, l}(\mathbb{F}_q) := \left\{ g \cdot f_*^l \in F_n(\mathbb{F}_q) \mid f_* \in \mathcal{P}_0 \left( \frac{n - \deg(g)}{l} \right) \text{ s.t. } \deg f_* \geq \deg v \text{ for all other irred. } v \mid g_{\lambda_1} \right\} \quad (55)$$

Note that we have the following decomposition for monic polynomials in  $F_{n, 1, N}(\mathbb{F}_q)$  with at least one irreducible factor of degree at least  $N$  in  $\mathcal{P}_0$ :

$$\#\{f \in F_{n, 1, N}(\mathbb{F}_q) \mid \deg f_{\lambda_1} \geq N\} = \bigsqcup_{n' \geq N} \bigsqcup_{\substack{g \in F_{n', 1, N}(\mathbb{F}_q) \\ \forall \text{ place } v \mid g \text{ and } v \in \mathcal{P}_0, \deg v < \frac{n-n'}{l}}} \bigsqcup_{l \geq 1} F_{n, 1, N}^{g, p, l}(\mathbb{F}_q) \quad (56)$$

The above composition can be obtained from picking the irreducible factor of a polynomial  $f \in F_{n, 1, N}(\mathbb{F}_q)$ , denoted as  $f_*$ , with  $\deg(f_{\lambda_1}) \geq N$  whose degree is maximal among all other irreducible factors of  $f$  which lives in  $\mathcal{P}_0$ . The polynomial  $g$  corresponds to the polynomial  $\frac{f}{f_*^{v_{f_*}(f)}}$ , where  $v_{f_*}$  is the valuation with respect to  $f_*$ .

Note the above decomposition is a disjoint union of sets  $F_{n, 1, N}^{g, p, l}(\mathbb{F}_q)$ . For the case where there are at least 2 irreducible polynomial factors of  $f$  in  $\mathcal{P}_0$  with maximal degree, one can endow a natural total order over the set of polynomials of degree  $n$  over  $\mathbb{F}_q$  and pick the irreducible polynomial factor  $v$  of  $f$  that is maximal among all other irreducible factors of  $f$ .

**Definition 5.9** (Auxiliary restriction). Fix integers  $n, n', N$  and a prime  $p$ . Fix an elliptic curve  $E/K$  satisfying the conditions from (18). Denote by  $\Sigma$  the set of bad places of reduction of  $E$ , including the place  $\infty$  at infinity. Pick a place  $v$  of  $K$  such that  $v \in \mathcal{P}_0$ ,  $\deg v > N$  and is coprime to  $\Delta_E$ . We will call such a place an auxiliary place of  $K$ .

Let  $g \in F_{n',1,N}(\mathbb{F}_q)$  be a polynomial satisfying the conditions as stated in Definition 5.8. Given a polynomial  $f \in F_{n,1,N}^{g,p,l}(\mathbb{F}_q)$ , let  $\chi^{[f]}$  be the global character associated to the cyclic order- $p$  field extension  $K(\sqrt[p]{f})/K$ . Consider the restriction of  $\chi^{[f]}$  to the set of cartesian products of local characters over the set  $\Sigma \cup \{w \text{ place} \mid w \text{ divides } f\}$ , which are ramified at places dividing  $f$ :

$$\left( \prod_{w' \in \Sigma} \chi^{[f]}|_{w'} \right) \times \left( \prod_{w|f} \chi^{[f]}|_w \right) \in \left( \prod_{w' \in \Sigma} \text{Hom}(\text{Gal}(\overline{K_{w'}}/K_{w'}), \mu_p) \right) \times \left( \prod_{w|f} \text{Hom}_{\text{ram}}(\text{Gal}(\overline{K_w}/K_w), \mu_p) \right)$$

Then by Proposition 4.5, there exists a cartesian product of local characters  $\chi^{\tilde{[f]}}$ , as an element of

$$\chi^{\tilde{[f]}} \in \left( \prod_{w' \in \Sigma} \text{Hom}(\text{Gal}(\overline{K_{w'}}/K_{w'}), \mu_p) \right) \times \left( \prod_{w|g} \text{Hom}_{\text{ram}}(\text{Gal}(\overline{K_w}/K_w), \mu_p) \right) \times \text{Hom}_{\text{ram}}(\text{Gal}(\overline{K_v}/K_v), \mu_p) \quad (57)$$

which is defined as

$$\begin{aligned} \chi^{\tilde{[f]}}|_w &= \chi^{[f]}|_w \text{ for any } w \in \Sigma \text{ or } w \mid g \\ \dim_{\mathbb{F}_p} \text{Sel}_{\pi}(E^{\chi^{\tilde{[f]}}}) &= \dim_{\mathbb{F}_p} \text{Sel}_{\pi}(E^{\chi^{[f]}}). \end{aligned} \quad (58)$$

for any auxiliary place  $v$ . One is allowed to do this because as suggested in Proposition 4.5, any twists by local characters over places  $v \in \mathcal{P}_0$  do not alter the dimension of  $\pi$ -Selmer groups. Pick an auxiliary place  $v \in \mathcal{P}_0$  which is coprime to  $g$ . We denote by  $\text{Res}_{\Sigma,g,v}$  the set-theoretic restriction map associated to auxiliary prime  $v$ , which represents elements in  $F_{n,1,N}^{g,p,l}$  as local characters supported over the finite set of primes  $\Sigma \cup \{w \text{ place} \mid w \text{ divides } g\} \cup \{v\}$ :

$$\begin{aligned} \text{Res}_{\Sigma,g,v} : F_{n,1,N}^{g,p,l}(\mathbb{F}_q) &\rightarrow \left( \prod_{w \in \Sigma} \text{Hom}_{\text{unr}}(\text{Gal}(\overline{K_w}/K_w), \mu_p) \right) \times \left( \prod_{w|g \text{ or } w=v} \text{Hom}_{\text{ram}}(\text{Gal}(\overline{K_w}/K_w), \mu_p) \right) \\ f &\mapsto \chi^{\tilde{[f]}}. \end{aligned} \quad (59)$$

## 6. PROOF OF THE MAIN THEOREM

Using all the key ingredients and constructions from previous sections, we can finally prove the statement of the main theorem. Before we proceed to the proof, we present a diagram which illustrates the key ideas of the proof of Propositions 6.1 and 6.3.

**6.1. Strategy of the proof.** Figure 4 demonstrates a summary of strategies to be implemented in computing the probability distribution of  $\pi$ -Selmer groups of cyclic order- $p$  Selmer groups of non-isotrivial elliptic curves over  $K$ . Proposition 6.3 allows one to show that it is suffice to compute the probability distribution of the dimensions of  $\pi$ -Selmer groups of families of Elliptic curves twisted by polynomials in  $F_{n,1,N}(\mathbb{F}_q)$ . Proposition 6.1 demonstrates that the set  $F_{n,1,N}(\mathbb{F}_q)$  can be decomposed into disjoint unions of sets of form  $F_{n,1,N}^{g,p,l}(\mathbb{F}_q)$  using auxiliary places, as defined from Section 5.3. These sets are of great importance to us, because the restriction of elements in  $F_{n,1,N}^{g,p,l}(\mathbb{F}_q)$  induces a distribution which can be regarded as a uniform distribution over a subspace of finite Cartesian products of sets of local cyclic order- $p$  characters. This allows us to apply Proposition 4.5 to compute the probability distribution of the sizes of  $\pi$ -Selmer groups of families of local twists of elliptic curves using Markov chains.

**6.2. Polynomials with irreducible factors of large enough degrees.** As before, fix  $N$  to be the smallest integer which is at least  $n^{\frac{1}{3}}$ . Consider the set  $F_{n,1,N}(\mathbb{F}_q) \cup F_{n,u,N}(\mathbb{F}_q)$  where  $u$  is a non-trivial element in  $\mathbb{F}_q^{\times}/(\mathbb{F}_q^{\times})^2$ . Assuming that  $n$  is sufficiently large, the Markov chain constructed from Proposition 4.5 and Section 4.2 governs the distribution of  $\pi$ -Selmer groups over the family of




FIGURE 4. An illustration of the key ideas and related concepts utilized in the proof of the main theorem. In this diagram, we assume that the polynomials of degree  $n$  over  $\mathbb{F}_q$  are coprime to the discriminant of  $E$ , and the numbers of their distinct irreducible factors are between  $\rho \ln n$  and  $(2 - \rho) \ln n$  given a fixed value  $0 < \rho < 1$ .

elliptic curves  $E$  twisted by  $f \in F_{n,1,N}(\mathbb{F}_q)$ . The same fact holds for  $\pi$ -Selmer groups over the family of elliptic curves  $E$  twisted by  $f \in F_{n,u,N}(\mathbb{F}_q)$  for any unit  $u \in \mathbb{F}_q^\times$ .

**Proposition 6.1.** *Fix a prime number  $p$ . Let  $K = \mathbb{F}_q(t)$  be a global function field whose characteristic is coprime to  $2, 3$ , and  $p$ . Let  $E : y^2 = F(x)$  be an elliptic curve over  $K$  such that*

- (1)  $E$  is non-isotrivial
- (2)  $E$  has a place  $\infty$  of split multiplicative reduction.
- (3) The constant field  $\mathbb{F}_q$  contains the primitive  $p$ -th root of unity  $\mu_p$ .
- (4) The Galois group  $\text{Gal}(K(E[p])/K)$  contains  $\text{SL}_2(\mathbb{F}_p)$ , i.e. is isomorphic to  $\text{SL}_2(\mathbb{F}_p) \rtimes T$  for some finite cyclic group  $T$ .

Fix  $N$  to be the smallest integer which is at least  $n^{\frac{1}{3}}$ . Denote by  $w(f)$  the number of irreducible factors of  $f \in F_n(\mathbb{F}_q)$ . Then for every non-negative integer  $j$ , a fixed integer  $w$  which satisfies  $\rho \ln n \leq w \leq (2 - \rho) \ln n$  for some positive  $0 < \rho < 1$ , a positive  $\epsilon' > 0$ , and sufficiently large  $n$ , there exists a fixed constant  $C > 0$  such that

$$\left| \mathbb{P} [\dim_{\mathbb{F}_p} \text{Sel}_{\pi}(E^{X_f}) = j \mid f \in F_{n,1,N}(\mathbb{F}_q), w(f) = w] - \prod_{m \geq 0} \frac{1}{1 + p^{-m}} \prod_{m=1}^j \frac{p}{p^m - 1} \right| < C n^{\rho \ln \left( 1 - \frac{1}{|T|} \frac{p}{p^2 - 1} + \epsilon' \right)}$$

where the probability of our interest is defined as

$$\mathbb{P} [\dim_{\mathbb{F}_p} \text{Sel}_{\pi}(E^{X_f}) = j \mid f \in F_{n,1,N}(\mathbb{F}_q), w(f) = w] := \frac{\#\{f \in F_{n,1,N}(\mathbb{F}_q) \mid w(f) = w, \dim_{\mathbb{F}_p} \text{Sel}_{\pi}(E^{X_f}) = j \text{ where } \chi_f \text{ assoc. to } K(\sqrt[p]{f})/K\}}{\#\{f \in F_{n,1,N}(\mathbb{F}_q) \mid w(f) = w\}}$$



**Proof. Decomposition of  $F_{n,1,N}(\mathbb{F}_q)$  using auxiliary places**

By Remark 4.3, we have for sufficiently large  $n$ ,

$$\frac{\#\{f \in F_{n,1,N}(\mathbb{F}_q) \mid \deg f_{\lambda_1} = 0, w(f) = w\}}{\#\{f \in F_{n,1,N}(\mathbb{F}_q) \mid w(f) = w\}} = \left( \frac{p}{|T|(p^2 - 1)} + \epsilon' \right)^w < n^{\rho \ln \left( \frac{p}{|T|(p^2 - 1)} + \epsilon' \right)} \quad (60)$$

for some error term  $\epsilon'$  such that  $|\epsilon'| < Cq^{-\frac{N}{2}}$  for some constant  $C > 0$ . We recall from Definition 5.8 and 5.9 that the set of elements in  $F_{n,1,N}(\mathbb{F}_q)$  with at least one irreducible factor in  $\mathcal{P}_0$  decomposes as follows:

$$\#\{f \in F_{n,1,N}(\mathbb{F}_q) \mid \deg f_{\lambda_1} \geq N\} = \bigsqcup_{n' \geq N} \bigsqcup_{g \in F_{n',1,N}(\mathbb{F}_q)} \bigsqcup_{l \geq 1} F_{n,1,N}^{g,p,l}(\mathbb{F}_q) \quad (61)$$

$\forall \text{ place } v|g \text{ and } v \in \mathcal{P}_0, \deg v < \frac{n-n'}{l}$

**Probability distribution over  $F_{n,1,N}^{g,p,l}(\mathbb{F}_q)$**

For each set  $F_{n,1,N}^{g,p,l}(\mathbb{F}_q)$ , one can consider the restriction map  $\text{Res}_{\Sigma,g,v}$  associated to an auxiliary place  $v$  from Definition 5.9. Let  $g = g_{\mu}g_{\lambda_1}g_{\lambda_2}g_{\lambda_3}$  be the splitting partition of  $g$ .

Let  $\chi^{[g]}$  be the restriction of the global character associated to the cyclic order- $p$  field extension  $K(\sqrt[p]{g})/K$  over the finite set of places  $w \mid g$  or  $w \in \Sigma$ . Denote by  $\text{Orbit}(\chi^{[g]}, \mu_p)$  the subset of the following cartesian product of local characters

$$\text{Orbit}(\chi^{[g]}, \mu_p) \subset \prod_{w|g} \text{Hom}_{ram}(\text{Gal}(\overline{K_w}/K_w), \mu_p) \quad (62)$$

comprised of finite cartesian products of local characters obtained by multiplying the local restriction of the character  $\chi^{[g]}|_w$  with any unramified character of  $\text{Hom}(\text{Gal}(\overline{K_w}/K_w), \mu_p)$  for each place  $w \mid g$ . Then by definition, the image of the restriction map  $\text{Res}_{\Sigma,g,v}$  is equal to:

$$\text{Res}_{\Sigma,g,v}(F_{n,1,N}^{g,p,l}(\mathbb{F}_q)) = \left( \prod_{w \in \Sigma} \text{Hom}_{unr}(\text{Gal}(\overline{K_w}/K_w), \mu_p) \right) \times \mathcal{S}_v \times \text{Orbit}(\chi^{[g]}, \mu_p). \quad (63)$$

for some subset  $\mathcal{S}_v \subset \text{Hom}_{ram}(\text{Gal}(\overline{K_v}/K_v), \mu_p)$ . Note that because  $v \in \mathcal{P}_0$  does not contribute to variations of the dimensions of  $\pi$ -Selmer groups, it suffices to fix  $\mathcal{S}_v = \{\chi_v\}$  to be a singleton set.

Let  $\phi_v$  be the projection map from the finite cartesian product of certain local cyclic characters to the local ramified character of order  $p$  at the auxiliary place  $v$  and places inside  $\Sigma$ :

$$\begin{aligned} \phi_v : & \left( \prod_{w \in \Sigma} \text{Hom}_{unr}(\text{Gal}(\overline{K_w}/K_w), \mu_p) \right) \times \text{Hom}_{ram}(\text{Gal}(\overline{K_v}/K_v), \mu_p) \times \text{Orbit}(\chi^{[g]}, \mu_p) \\ & \rightarrow \left( \prod_{w \in \Sigma} \text{Hom}_{unr}(\text{Gal}(\overline{K_w}/K_w), \mu_p) \right) \times \text{Hom}_{ram}(\text{Gal}(\overline{K_v}/K_v), \mu_p) \end{aligned} \quad (64)$$

Let us denote by  $\delta : \mathbb{Z}_{\geq 0} \rightarrow [0, 1]$  the following probability distribution obtained from local cyclic order- $p$  twists of  $\pi$ -Selmer groups:

$$\delta(j) := \frac{\#\{\chi \in \text{Res}_{\Sigma,g,v}(F_{n,1,N}^{g,p,l}(\mathbb{F}_q)) \mid \dim_{\mathbb{F}_p} \text{Sel}_{\pi}(E^{\phi_v(\chi)}) = j\}}{\#\text{Res}_{\Sigma,g,v}(F_{n,1,N}^{g,p,l}(\mathbb{F}_q))} \quad (65)$$

Recall that  $\pi$ -Selmer groups are determined by local cohomological conditions. Furthermore, Corollary 5.6 implies that the image of  $\text{Res}_{\Sigma,g,v}$  restricts to a uniform distribution over the set  $\text{Orbit}(\chi^{[g]}, \mu_p)$ . This shows that one can deduce the desired probability distribution

$$\frac{\#\{f \in F_{n,1,N}^{g,p,l}(\mathbb{F}_q) \mid \dim_{\mathbb{F}_p} \text{Sel}_{\pi}(E^{\chi_f}) = j, \chi_f \text{ assoc. to } K(\sqrt[p]{f})/K, w(f) = w\}}{\#F_{n,1,N}^{g,p,l}(\mathbb{F}_q)} \quad (66)$$

from the initial probability distribution  $\delta : \mathbb{Z}_{\geq 0} \rightarrow [0, 1]$  from (65) by iteratively applying Proposition 4.5 over the collections of sets of form  $\Omega_{\chi, \mathfrak{w}}^\eta$ , as defined from Definition 4.4. Indeed, every character of  $\text{Orbit}(\chi^{[g]}, \mu_p)$  corresponds to the set of finite cartesian products of local characters obtained from multiplying the ramified character  $\chi^{[g]}|_{\mathfrak{w}}$  with any unramified character of  $\text{Hom}(\text{Gal}(\overline{K_{\mathfrak{w}}}/K_{\mathfrak{w}}), \mu_p)$  for each place  $\mathfrak{w} \mid g$ . Thus, we achieve that

$$\begin{aligned}
(66) &= \frac{\#\{f \in F_{n,1,N}^{g,p,l}(\mathbb{F}_q) \mid \dim_{\mathbb{F}_p} \text{Sel}_{\pi}(E^{\chi_f}) = j, \chi_f \text{ assoc. to } K(\sqrt[p]{f})/K, w(f) = w\}}{\#\text{Res}_{\Sigma,g,v}(\{f \in F_{n,1,N}^{g,p,l}(\mathbb{F}_q) \mid w(f) = w\})} \times \frac{\#\text{Res}_{\Sigma,g,v}(\{f \in F_{n,1,N}^{g,p,l}(\mathbb{F}_q) \mid w(f) = w\})}{\#\{f \in F_{n,1,N}^{g,p,l}(\mathbb{F}_q) \mid w(f) = w\}} \\
&= \frac{\#\{f \in F_{n,1,N}^{g,p,l}(\mathbb{F}_q) \mid \dim_{\mathbb{F}_p} \text{Sel}_{\pi}(E^{\text{Res}_{\Sigma,g,v}(f)}) = j, w(f) = w\}}{\#\text{Res}_{\Sigma,g,v}(\{f \in F_{n,1,N}^{g,p,l}(\mathbb{F}_q) \mid w(f) = w\})} \times \frac{\#\text{Res}_{\Sigma,g,v}(\{f \in F_{n,1,N}^{g,p,l}(\mathbb{F}_q) \mid w(f) = w\})}{\#\{f \in F_{n,1,N}^{g,p,l}(\mathbb{F}_q) \mid w(f) = w\}} \\
&= \frac{\#\{f \in F_{n,1,N}^{g,p,l}(\mathbb{F}_q) \mid \dim_{\mathbb{F}_p} \text{Sel}_{\pi}(E^{\text{Res}_{\Sigma,g,v}(f)}) = j, w(f) = w\}}{\#\{\chi \in \text{Res}_{\Sigma,g,v}(\{f \in F_{n,1,N}^{g,p,l}(\mathbb{F}_q) \mid w(f) = w\}) \mid \dim_{\mathbb{F}_p} \text{Sel}_{\pi}(E^{\chi}) = j\}} \times \frac{\#\{f \in F_{n,1,N}^{g,p,l}(\mathbb{F}_q) \mid w(f) = w\}}{\#\text{Res}_{\Sigma,g,v}(\{f \in F_{n,1,N}^{g,p,l}(\mathbb{F}_q) \mid w(f) = w\})} \\
&\quad \times \frac{\#\{\chi \in \text{Res}_{\Sigma,g,v}(\{f \in F_{n,1,N}^{g,p,l}(\mathbb{F}_q) \mid w(f) = w\}) \mid \dim_{\mathbb{F}_p} \text{Sel}_{\pi}(E^{\chi}) = j\}}{\#\text{Res}_{\Sigma,g,v}(\{f \in F_{n,1,N}^{g,p,l}(\mathbb{F}_q) \mid w(f) = w\})}
\end{aligned} \tag{67}$$

One can apply the Chebotarev density theorem (Theorem 3.1) over the compositum of field extensions  $K(E[p])$  (Lemma 4.2) and  $K(\sqrt[p]{f})$  over all irreducible factors of the polynomial  $g$  (discussions following Corollary 5.6) to obtain that

$$\left| \frac{\#\{v \in \mathcal{P}_0(i) \mid \prod_{i=1}^w \left(\frac{v}{h_i}\right)_p = a \in \prod_{k=1}^w \mu_p\}}{\#\mathcal{P}_0(i)} - \frac{1}{p^w} \right| < C n^{-2 \ln n + 2 + 2 \ln p} \tag{68}$$

for some fixed constant  $C > 0$ . This implies that the image of  $\text{Res}_{\Sigma,g,v}$  restricts to a uniform distribution over the set  $\text{Orbit}(\chi^{[g]}, \mu_p) \times \left(\prod_{w \in \Sigma} \text{Hom}_{\text{unr}}(\text{Gal}(\overline{K_w}/K), \mu_p)\right)$ . We thus obtain that there exists an error term  $\epsilon_1$  satisfying

$$|\epsilon_1| < p^{w(f)} \cdot n^{-2 \ln n + 2 + 2 \ln p} < n^{-2 \ln n + 2 + 4 \ln p} \tag{69}$$

for sufficiently large  $n$  such that the first term of (67) can be expressed as

$$\frac{\#\{f \in F_{n,1,N}^{g,p,l}(\mathbb{F}_q) \mid \dim_{\mathbb{F}_p} \text{Sel}_{\pi}(E^{\text{Res}_{\Sigma,g,v}(f)}) = j, w(f) = w\}}{\#\{\chi \in \text{Res}_{\Sigma,g,v}(\{f \in F_{n,1,N}^{g,p,l}(\mathbb{F}_q) \mid w(f) = w\}) \mid \dim_{\mathbb{F}_p} \text{Sel}_{\pi}(E^{\chi}) = j\}} \times \frac{\#\{f \in F_{n,1,N}^{g,p,l}(\mathbb{F}_q) \mid w(f) = w\}}{\#\text{Res}_{\Sigma,g,v}(\{f \in F_{n,1,N}^{g,p,l}(\mathbb{F}_q) \mid w(f) = w\})} = 1 + \epsilon_1. \tag{70}$$

By Corollary 5.6, one can use Proposition 4.5 iteratively over the collections of sets of form  $\Omega_{\chi,v}^\eta$  to simplify the second term of (67) as

$$\frac{\#\{\chi \in \text{Res}_{\Sigma,g,v}(\{f \in F_{n,1,N}^{g,p,l}(\mathbb{F}_q) \mid w(f) = w\}) \mid \dim_{\mathbb{F}_p} \text{Sel}_{\pi}(E^{\chi}) = j\}}{\#\text{Res}_{\Sigma,g,v}(\{f \in F_{n,1,N}^{g,p,l}(\mathbb{F}_q) \mid w(f) = w\})} = M_L^{\deg(g_{\lambda_2}) + 2 \deg(g_{\lambda_3})}(\delta)(j) + \epsilon_{Ch} \tag{71}$$

where we recall that  $g = g_\mu g_{\lambda_1} g_{\lambda_2} g_{\lambda_3}$  is the splitting partition of  $g$ , and the error term  $\epsilon_{Ch}$  is obtained from the effective Chebotarev density theorem (i.e. Proposition 4.5 with  $\deg \sigma \geq N = n^{\frac{1}{3}}$ ) satisfying

$$|\epsilon_{Ch}| \leq C n^{-\frac{1}{9} \ln n}. \tag{72}$$

for some fixed constant  $C > 0$ . This implies that

$$\frac{\#\{f \in F_{n,1,N}^{g,p,l}(\mathbb{F}_q) \mid \dim_{\mathbb{F}_p} \text{Sel}_{\pi}(E^{\chi_f}) = j, \chi_f \text{ assoc. to } K(\sqrt[p]{f})/K, w(f) = w\}}{\#\{f \in F_{n,1,N}^{g,p,l}(\mathbb{F}_q) \mid w(f) = w\}} = M_L^{\deg(g_{\lambda_2}) + 2 \deg(g_{\lambda_3})}(\delta)(j) + C_g(\epsilon_1 + \epsilon_{Ch}) \tag{73}$$

for some fixed constant  $0 < C_g \leq 2$ .

### Probability Distribution over $F_{n,1,N}(\mathbb{F}_q)$

Given a polynomial  $g \in F_{n',1,N}(\mathbb{F}_q)$ , denote by  $\lambda_g := [\deg(g_\mu), \deg(g_{\lambda_1}), \deg(g_{\lambda_2}), \deg(g_{\lambda_3})]$  the list of degrees of the factors of the splitting partition of  $g = g_\mu g_{\lambda_1} g_{\lambda_2} g_{\lambda_3}$ .

Fix a list of 4 integers  $\lambda := [\mu, \lambda_1, \lambda_2, \lambda_3]$ . Denote by  $l(\lambda)$  the sum of 4 integers, i.e.  $l(\lambda) = \mu + \lambda_1 + \lambda_2 + \lambda_3$ . Consider the set of polynomials

$$F_{n,1,N}^{\lambda,l}(\mathbb{F}_q) := \left\{ g \cdot f_*^l \in F_n(\mathbb{F}_q) \mid \begin{array}{l} \lambda_g = \lambda, n' = l(\lambda) \\ f_* \in \mathcal{P}_0\left(\frac{n-n'}{l}\right) \text{ s.t. } \deg f_* \geq \deg v \text{ for all other } v|_{g_{\lambda_1}} \end{array} \right\}. \quad (74)$$

Then it holds that

$$F_{n,1,N}^{\lambda,l}(\mathbb{F}_q) = \bigsqcup_{\substack{g \in F_{n',1,N}(\mathbb{F}_q) \\ \lambda_g = \lambda, l(\lambda) = n'}} F_{n,1,N}^{g,p,l}(\mathbb{F}_q). \quad (75)$$

Without loss of generality, we may assume that the disjoint subsets  $F_{n,1,N}^{g,p,l}(\mathbb{F}_q)$  of  $F_{n,1,N}^{\lambda,l}(\mathbb{F}_q)$  are all of the same cardinality, which is the case as long as  $\frac{n-n'}{l} > \deg v$  for all  $v \mid g_{\lambda_1}$ . For polynomials  $g$  such that there exists a prime factor  $v \mid g_{\lambda_1}$  such that  $\frac{n-n'}{l} = \deg v$ , the difference in cardinality between such  $F_{n,1,N}^{g,p,l}(\mathbb{F}_q)$ 's are negligible from other disjoint subsets  $F_{n,1,N}^{g',p,l}(\mathbb{F}_q)$  for polynomials  $g'$  with  $\frac{n-n'}{l} > \deg v$  for all  $v \mid g_{\lambda_1}$  in that

$$\left| 1 - \frac{\#F_{n,1,N}^{g,p,l}(\mathbb{F}_q)}{\#F_{n,1,N}^{g',p,l}(\mathbb{F}_q)} \right| < \frac{2 \ln n}{q^{n^{\frac{1}{3}}}} \quad (76)$$

because we assumed that all irreducible factors of  $f \in F_{n,1,N}^{\lambda,l}(\mathbb{F}_q)$  are at least  $N = n^{\frac{1}{3}}$ . For any polynomial  $g \in F_{n',1,N}$  with  $\lambda_g = \lambda$  and  $n' = l(\lambda)$ , we will use the constant  $G_{n,1,N}^{\lambda,l}$  to denote

$$G_{n,1,N}^{\lambda,l} := \frac{\#F_{n,1,N}^{g,p,l}(\mathbb{F}_q)}{\#F_{n,1,N}^{\lambda,l}(\mathbb{F}_q)}. \quad (77)$$

Using the effective Chebotarev density theorem to Lemma 4.2, we obtain that

$$\begin{aligned} & \left| \frac{\#\{f \in F_{n,1,N}^{\lambda,l}(\mathbb{F}_q) \mid \dim_{\mathbb{F}_p} \text{Sel}_{\pi}(E_f^X) = j, w(f) = w\}}{\#\{f \in F_{n,1,N}^{\lambda,l}(\mathbb{F}_q) \mid w(f) = w\}} - \left( \left(1 - \frac{p}{|T|(p^2-1)}\right) + \frac{1}{|T|p} M_L + \frac{1}{|T|(p^3-p)} M_L^2 \right)^{w-1} (\delta)(j) \right| \\ &= \left| \frac{\sum_{\substack{g \in F_{n',1,N}(\mathbb{F}_q) \\ \lambda_g = \lambda, l(\lambda) = n' \\ w(g) = w-1}} \#\{f \in F_{n,1,N}^{g,p,l}(\mathbb{F}_q) \mid \dim_{\mathbb{F}_p} \text{Sel}_{\pi}(E^{Xf}) = j, w(f) = w\}}{\#\{f \in F_{n,1,N}^{\lambda,l}(\mathbb{F}_q) \mid w(f) = w\}} - \left( \left(1 - \frac{p}{|T|(p^2-1)}\right) + \frac{1}{|T|p} M_L + \frac{1}{|T|(p^3-p)} M_L^2 \right)^{w-1} (\delta)(j) \right| \\ &= \left| \sum_{\substack{g \in F_{n',1,N}(\mathbb{F}_q) \\ \lambda_g = \lambda, l(\lambda) = n' \\ w(g) = w-1}} \left( \frac{\#\{f \in F_{n,1,N}^{g,p,l}(\mathbb{F}_q) \mid \dim_{\mathbb{F}_p} \text{Sel}_{\pi}(E^{Xf}) = j, w(f) = w\}}{\#\{f \in F_{n,1,N}^{g,p,l}(\mathbb{F}_q) \mid w(f) = w\}} \times \frac{\#\{f \in F_{n,1,N}^{g,p,l}(\mathbb{F}_q) \mid w(f) = w\}}{\#\{f \in F_{n,1,N}^{\lambda,l}(\mathbb{F}_q) \mid w(f) = w\}} \right) \right. \\ & \quad \left. - \left( \left(1 - \frac{p}{|T|(p^2-1)}\right) + \frac{1}{|T|p} M_L + \frac{1}{|T|(p^3-p)} M_L^2 \right)^{w-1} (\delta)(j) \right|. \end{aligned} \quad (78)$$

for some finite cyclic subgroup  $T$  and an arbitrary initial finitely supported probability distribution  $\delta : \mathbb{Z}_{\geq 0} \rightarrow [0, 1]$ . Using the substitution from (77), we obtain:

$$\begin{aligned}
&= \left| \sum_{\substack{g \in F_{n',1,N}(\mathbb{F}_q) \\ \lambda_g = \lambda, l(\lambda) = n' \\ w(g) = w-1}} G_{n,1,N}^{\lambda,l} \frac{\#\{f \in F_{n,1,N}^{g,p,l}(\mathbb{F}_q) \mid \dim_{\mathbb{F}_p} \text{Sel}_{\pi}(E^{X_f}) = j, w(f) = w\}}{\#\{f \in F_{n,1,N}^{g,p,l}(\mathbb{F}_q) \mid w(f) = w\}} \right. \\
&\quad \left. - \left( \left(1 - \frac{p}{|T|(p^2-1)}\right) + \frac{1}{|T|p} M_L + \frac{1}{|T|(p^3-p)} M_L^2 \right)^{w-1} (\delta)(j) \right| \\
&\leq \left| G_{n,1,N}^{\lambda,l} \sum_{\substack{g \in F_{n',1,N}(\mathbb{F}_q) \\ \lambda_g = \lambda, l(\lambda) = n' \\ w(g) = w-1}} \left( M_L^{\deg(g\lambda_2)+2\deg(g\lambda_3)} (\delta)(j) + 2(\epsilon_1 + \epsilon_{Ch}) \right) \right. \\
&\quad \left. - \left( \left(1 - \frac{p}{|T|(p^2-1)}\right) + \frac{1}{|T|p} M_L + \frac{1}{|T|(p^3-p)} M_L^2 \right)^{w-1} (\delta)(j) \right| \\
&\leq C'(\epsilon_1 + \epsilon_{Ch} + \epsilon'_{Ch}).
\end{aligned} \tag{79}$$

for some fixed constant  $0 < C' < 2$ , and the error terms  $\epsilon_1, \epsilon_{Ch}, \epsilon'_{Ch}$  independent from the choice of  $g \in F_{n',1,N}(\mathbb{F}_q)$ . All the error terms are obtained from the effective Chebotarev density theorem (i.e. Corollary 5.6, Proposition 4.5 with  $\deg \sigma \geq N$  and Remark 4.3) satisfying

$$|\epsilon_1| \leq Cn^{-2\ln n + 2 + 4\ln p}, \tag{80}$$

$$|\epsilon_{Ch}| \leq Cn^{-\frac{1}{9}\ln n}, \tag{81}$$

$$|\epsilon'_{Ch}| \leq Cq^{-\frac{1}{2}n^{\frac{1}{3}}}. \tag{82}$$

for some fixed constant  $C > 0$ . Using the geometric ergodicity of Markov chains from Corollary 4.14, we obtain that there exists an error term  $\epsilon_2$  satisfying

$$|\epsilon_2| < C\gamma^{w-1} < Cn^{\rho \ln \gamma} \tag{83}$$

for some fixed constant  $C > 0$  and  $0 < \gamma < 1$  such that

$$\left| \left( \left(1 - \frac{p}{|T|(p^2-1)}\right) + \frac{1}{|T|p} M_L + \frac{1}{|T|(p^3-p)} M_L^2 \right)^{w-1} (\delta)(j) - PR(j) \right| \leq \epsilon_2 + C'(\epsilon_1 + \epsilon_{Ch} + \epsilon_{Ch'}) \tag{84}$$

where  $PR : \mathbb{Z}_{\geq 0} \rightarrow [0, 1]$  is the Bhargava-Kane-Lenstra-Poonen-Rains distribution. Note that  $\gamma$  is the geometric rate of convergence of the Markov chain  $M$  which satisfies

$$1 - \frac{1}{|T|} \frac{p}{p^2-1} < \gamma < 1 \tag{85}$$

as verified from Corollary 4.14. Therefore, we obtain that there exists an error term  $\epsilon$  with

$$|\epsilon| < |\epsilon_2| + 2|\epsilon_1| + 2|\epsilon_{Ch}| + 2|\epsilon'_{Ch}| < 5Cn^{\rho \ln \gamma}. \tag{86}$$

such that

$$\frac{\#\{f \in F_{n,1,N}^{\lambda,l}(\mathbb{F}_q) \mid \dim_{\mathbb{F}_p} \text{Sel}_{\pi}(E_f^X) = j, w(f) = w\}}{\#\{f \in F_{n,1,N}^{\lambda,l}(\mathbb{F}_q) \mid w(f) = w\}} = PR(j) + \epsilon. \tag{87}$$

Using the decomposition from (61), we immediately obtain that

$$\frac{\#\{f \in F_{n,1,N}(\mathbb{F}_q) \mid \dim_{\mathbb{F}_p} \text{Sel}_{\pi}(E_f^X) = j, \deg f_{\lambda_1} \geq N, w(f) = w\}}{\#\{f \in F_{n,1,N}(\mathbb{F}_q) \mid \deg f_{\lambda_1} \geq N, w(f) = w\}} = PR(j) + \epsilon. \tag{88}$$

By Proposition 4.5, the maximal dimension of  $\pi$ -Selmer group twisted by characters associated to  $f \in F_{n,1,N}(\mathbb{F}_q)$  with  $\deg f_{\lambda_1} = 0$  is at most  $4\ln n + C_E$  for some fixed constant  $C_E > 0$  which

depends solely on the choice of the elliptic curve  $E$ . Therefore, for some fixed constant  $C > 0$  and a positive  $\epsilon' > 0$ ,

$$\begin{aligned}
 & \left| \mathbb{P} \left[ \dim_{\mathbb{F}_p} \text{Sel}_{\pi}(E^{\chi_f}) = j \mid f \in F_{n,1,N}(\mathbb{F}_q), w(f) = w \right] - PR(j) \right| \\
 &= \left| \frac{\#\{f \in F_{n,1,N}(\mathbb{F}_q) \mid \dim_{\mathbb{F}_p} \text{Sel}_{\pi}(E^{\chi_f}) = j, \deg f_{\lambda_1} = 0, w(f) = w\}}{\#\{f \in F_{n,1,N}(\mathbb{F}_q) \mid \deg f_{\lambda_1} = 0, w(f) = w\}} \cdot \frac{\#\{f \in F_{n,1,N}(\mathbb{F}_q) \mid \deg f_{\lambda_1} = 0, w(f) = w\}}{\#\{f \in F_{n,1,N}(\mathbb{F}_q) \mid w(f) = w\}} \right. \\
 &+ \left. \frac{\#\{f \in F_{n,1,N}(\mathbb{F}_q) \mid \dim_{\mathbb{F}_p} \text{Sel}_{\pi}(E^{\chi_f}) = j, \deg f_{\lambda_1} \geq N, w(f) = w\}}{\#\{f \in F_{n,1,N}(\mathbb{F}_q) \mid \deg f_{\lambda_1} \geq N, w(f) = w\}} \cdot \frac{\#\{f \in F_{n,1,N}(\mathbb{F}_q) \mid \deg f_{\lambda_1} \geq N, w(f) = w\}}{\#\{f \in F_{n,1,N}(\mathbb{F}_q) \mid w(f) = w\}} - PR(j) \right| \quad (89) \\
 &< n^{\rho \ln \left( \frac{p}{|T|(p^2-1)} + \epsilon' \right)} + |\epsilon| < C n^{\rho \ln \left( \frac{p}{|T|(p^2-1)} + \epsilon' \right)}
 \end{aligned}$$

□

**Remark 6.2.** Proposition 6.1 implies that the Markov chain constructed from Corollary 4.14 governs the probability distribution of  $\pi$ -Selmer ranks of cyclic order- $p$  twists of  $E$  by elements in  $F_{n,1,N}(\mathbb{F}_q)$  with  $w$  irreducible factors. The same argument can be used for the the probability distribution of  $\pi$ -Selmer ranks of quadratic twists of  $E$  by elements in  $F_{n,f_{\Delta_E},N}(\mathbb{F}_q)$  with  $w$  irreducible factors, where  $f_{\Delta_E}$  is either an element of  $\mathbb{F}_q^{\times}$  or a polynomial whose set of irreducible factors are contained in the set of bad places of reduction of  $E$ . For the latter case, the local cyclic characters over the places  $w \in \Sigma$  are ramified, but the overall argument of the proof remains the same. We hence omit the proof for such classes of polynomials. Note that the error bounds from Proposition 6.1 are independent from the number of irreducible factors  $w(f)$  of the polynomial  $f$ .

**6.3. Other polynomials.** Using Proposition 6.1, we now show that the probaibility distribution of  $\pi$ -Selmer groups of cyclic order- $p$  twist families of elliptic curves over  $F_n(\mathbb{F}_q)$  also converges to the Bhargava-Kane-Lenstra-Poonen-Rains distribution for sufficiently large  $n$ . For the statements written in the next proposition, we omit the fact that the order- $p$  characters  $\chi_f$  are associated to the field extension  $K(\sqrt[p]{f})/K$ .

**Proposition 6.3.** *Let  $E$  be a non-isotrivial elliptic curve over  $K$  satisfying the aforementioned conditions from Proposition 6.1. Let  $N = n^{\frac{1}{3}}$ . Then for sufficiently large  $n$  and a fixed constant  $0 < \rho < 1$ , there exists a fixed constant  $C > 0$  and a positive number  $0 < \alpha_{mc} < -\rho \ln \left( 1 - \frac{1}{|T|} \frac{p}{p^2-1} \right)$  such that*

$$\left| \mathbb{P} \left[ \dim_{\mathbb{F}_p} \text{Sel}_{\pi}(E^{\chi_f}) = j \mid \substack{f \in F_n(\mathbb{F}_q) \\ \rho \ln n \leq w(f) \leq (2-\rho) \ln n} \right] - \prod_{m \geq 0} \frac{1}{1 + p^{-m}} \prod_{m=1}^j \frac{p}{p^m - 1} \right| < C n^{-\alpha_{mc}},$$

where

$$\mathbb{P} \left[ \dim_{\mathbb{F}_p} \text{Sel}_{\pi}(E^{\chi_f}) = j \mid \substack{f \in F_n(\mathbb{F}_q) \\ \rho \ln n \leq w(f) \leq (2-\rho) \ln n} \right] := \frac{\#\{f \in F_n(\mathbb{F}_q) \mid \substack{\rho \ln n \leq w(f) \leq (2-\rho) \ln n \\ \dim_{\mathbb{F}_p} \text{Sel}_{\pi}(E^{\chi_f}) = j\}}{\#\{f \in F_n(\mathbb{F}_q) \mid \rho \ln n \leq w(f) \leq (2-\rho) \ln n\}}$$

*Proof.* Fix a small enough positive number  $\epsilon > 0$  such that

$$0 < \epsilon < -\frac{\ln \left( 1 - \frac{1}{|T|} \frac{p}{p^2-1} \right)}{\ln p} \quad (90)$$

The numerator term originates from the tight lower bound of the convergence rate of the Markov chain from (33) and from applying Corollary 4.14. Given a number  $w$ , denote by  $\mathcal{F}_{N,w}(\mathbb{F}_q)$  the set of  $p$ -th power -free polynomials of at most  $\epsilon w$  irreducible factors whose degrees are at most  $N$ .

$$\mathcal{F}_{N,w}(\mathbb{F}_q) := \{f \in \cup_{i=1}^{(2-\rho) \cdot (p-1) \cdot n^{\frac{1}{3}}} F_i(\mathbb{F}_q) \mid f = \prod_{i=1}^{\tilde{w}} v_i^{l_{v_i}} \text{ such that } v_i \in \cup_{i=1}^N \mathcal{F}_{q,i}(\mathbb{F}_q), \tilde{w} < \epsilon w, 1 \leq l_{v_i} \leq p-1\}. \quad (91)$$

For any  $f \in F_n(\mathbb{F}_q)$ , by Definition 5.1, the polynomial admits a factorization  $f = f_{\mu} f_{\lambda_1} f_{\lambda_2} f_{\lambda_3}$ . By construction and Proposition 5.4, there exist fixed constants  $C > 0$  and  $C' > 0$  such that for

sufficiently large  $n$ ,

$$\frac{\#\{f \in F_n(\mathbb{F}_q) \mid f_\mu \in \mathcal{F}_{N,w}(\mathbb{F}_q)\}}{\#F_n(\mathbb{F}_q)} \geq \left(1 - Cn^{-\frac{1}{3}\epsilon \ln \ln n}\right). \quad (92)$$

Note that the very choice of the constant  $\epsilon$  implies that the dimension  $\dim_{\mathbb{F}_p} \text{Sel}_\pi(E^{\chi_{f_\mu}})$  is bounded, and not as large as to prevent consecutive twists  $\chi_{f_{\lambda_1}} \chi_{f_{\lambda_2}} \chi_{f_{\lambda_3}}$  from preventing the dimension  $\dim_{\mathbb{F}_p} \text{Sel}_\pi(E^{\chi_f})$  to be equal to 0. Furthermore, the choice of  $\epsilon$  which satisfies (90) allows one to utilize Corollary 4.14 to still guarantee the Markov chain of our interest to converge to the desired probability distribution at a geometric, though significantly weaker, rate. Proposition 6.1 proves that the following equation holds as well for any such  $f_\mu \in \mathcal{F}_N(\mathbb{F}_q)$  for some fixed constant  $C, C_1, C_2 > 0$ :

$$\begin{aligned} \left| \frac{\#\{f \in F_{n,f_\mu,N}(\mathbb{F}_q) \mid \dim_{\mathbb{F}_p} \text{Sel}_\pi(E^{\chi_f}) = j, w(f) = w\}}{\#\{f \in F_{n,f_\mu,N}(\mathbb{F}_q) \mid w(f) = w\}} - PR(j) \right| &< C_1 \left(1 - \frac{1}{|T|} \frac{p}{p^2-1} + \epsilon\right)^w \cdot p^{2\epsilon w} + C_2 n^{\frac{1}{2} \ln \left(\frac{p}{|T|(p^2-1)} + \epsilon\right)} \\ &< C \left(1 - \frac{1}{|T|} \frac{p}{p^2-1} + \epsilon\right)^{w(1-2\epsilon)} + C_2 n^{\frac{1}{2} \ln \left(\frac{p}{|T|(p^2-1)} + \epsilon\right)} \\ &< C n^{\rho(1-2\epsilon) \ln \left(1 - \frac{1}{|T|} \frac{p}{p^2-1} + \epsilon\right)}. \end{aligned} \quad (93)$$

Denote by  $\mathcal{F}_{\Delta_E}$  the set of  $p$ -th power free polynomials whose irreducible factors are contained in the set of bad places of reduction of the elliptic curve  $E$ . Then by Remark 6.2, we obtain that for sufficiently large  $n$  there exists a fixed constant  $C, C_1, C_2 > 0$  and a positive number  $0 < \epsilon_{mc} < -\rho \ln \left(1 - \frac{1}{|T|} \frac{p}{p^2-1}\right)$  such that

$$\begin{aligned} &\left| \frac{\#\{f \in F_n(\mathbb{F}_q) \mid \dim_{\mathbb{F}_p} \text{Sel}_\pi(E^{\chi_f}) = j, \rho \ln n \leq w(f) \leq (2-\rho) \ln n\}}{\#\{f \in F_n(\mathbb{F}_q) \mid \rho \ln n \leq w(f) \leq (2-\rho) \ln n\}} - PR(j) \right| \\ &= \left| \frac{\sum_{w \geq \rho \ln n}^{(2-\rho) \ln n} \#\{f \in F_n(\mathbb{F}_q) \mid \dim_{\mathbb{F}_p} \text{Sel}_\pi(E^{\chi_f}) = j, w(f) = w\}}{\#\{f \in F_n(\mathbb{F}_q) \mid \rho \ln n \leq w(f) \leq (2-\rho) \ln n\}} - PR(j) \right| \\ &< \frac{\#\{f \in F_n(\mathbb{F}_q) \mid \rho \ln n \leq w(f) \leq (2-\rho) \ln n, f_\mu \notin \mathcal{F}_{\Delta_E}\}}{\#\{f \in F_n(\mathbb{F}_q) \mid \rho \ln n \leq w(f) \leq (2-\rho) \ln n\}} + \left| \sum_{w \geq \rho \ln n}^{(2-\rho) \ln n} \sum_{f_{\mu,1} \in \mathcal{F}_N(\mathbb{F}_q) \setminus \mathcal{F}_{\Delta_E}} \sum_{f_{\mu,2} \in \mathcal{F}_{\Delta_E}} \frac{\#\{f \in F_{n,f_{\mu,1}f_{\mu,2},N}(\mathbb{F}_q) \mid w(f) = w\}}{\#\{f \in F_n(\mathbb{F}_q) \mid \rho \ln n \leq w(f) \leq (2-\rho) \ln n\}} \right. \\ &\quad \times \left. \frac{\#\{f \in F_{n,f_{\mu,1}f_{\mu,2},N}(\mathbb{F}_q) \mid \dim_{\mathbb{F}_p} \text{Sel}_\pi(E^{\chi_f}) = j, w(f) = w\}}{\#\{f \in F_{n,g,N}(\mathbb{F}_q) \mid w(f) = w\}} - PR(j) \right| \\ &< C_1 n^{-\frac{1}{3}\epsilon \ln \ln n} + \sum_{w \geq \rho \ln n}^{(2-\rho) \ln n} C_2 n^{\rho(1-2\epsilon) \ln \left(1 - \frac{1}{|T|} \frac{p}{p^2-1} + \epsilon\right)} < C n^{\rho(1-3\epsilon) \ln \left(1 - \frac{1}{|T|} \frac{p}{p^2-1} + \epsilon\right)} < C n^{-\alpha_{mc}} \end{aligned}$$

The last equation follows from choosing arbitrarily small  $\epsilon > 0$ . □

The main theorem now follows immediately from Theorem 3.7 and Proposition 6.3.

*Theorem 1.1.* Recall that

$$\mathbb{P} \left[ \dim_{\mathbb{F}_p} \text{Sel}_\pi(E^{\chi_f}) = j \mid f \in F_n(\mathbb{F}_q) \right] := \frac{\#\{f \in F_n(\mathbb{F}_q) \mid \dim_{\mathbb{F}_p} \text{Sel}_\pi(E^{\chi_f}) = j\}}{\#F_n(\mathbb{F}_q)}. \quad (94)$$

By Theorem 3.7, there exists a fixed constant  $C_1 > 0$  and a constant  $0 < \alpha_{ek} < \rho \ln \rho + 1 - \rho$  such that

$$\left| (94) - \frac{\#\{f \in F_n(\mathbb{F}_q) \mid \dim_{\mathbb{F}_p} \text{Sel}_\pi(E^{\chi_f}) = j, \rho \ln n \leq w(f) \leq (2-\rho) \ln n\}}{\#\{f \in F_n(\mathbb{F}_q) \mid \rho \ln n \leq w(f) \leq (2-\rho) \ln n\}} \right| < C_1 n^{-\alpha_{ek}}.$$

By Proposition 6.3, there exists a fixed constant  $C_2 > 0$  and  $0 < \alpha_{mc} < -\rho \ln \left(1 - \frac{1}{|T|} \frac{p}{p^2-1}\right)$  such that

$$\left| (94) - \left( \prod_{m \geq 0} \frac{1}{1 + p^{-m}} \right) \left( \prod_{m=1}^j \frac{p}{p^m - 1} \right) \right| < C_1 n^{-\alpha_{ek}} + C_2 n^{-\alpha_{mc}}.$$

□

## ACKNOWLEDGEMENTS

My apologies in advance that the acknowledgement section may be longer than what one may read from other manuscripts.

This research was conducted during my compulsory military service as a research personnel in National Institute for Mathematical Sciences (NIMS) and during my graduate studies at the University of Wisconsin-Madison after returning from leave of absence. The progress I made on this project during my military service (which comprises most of the ideas covered in this manuscript) was primarily done at the end of the day after my usual working hours as a research personnel.

I would like to express my sincerest and most immense gratefulness to my PhD advisor Jordan Ellenberg for suggesting various helpful references, giving crucial and enlightening insights, and investing valuable time and effort for having regular Skype meetings during my leave of absence. I would also like to thank him for suggesting to understand the role of generalized Riemann hypothesis in computing the desired probability distributions. Without his continuous and immensely enthusiastic and patient support, this work would not have led to its fruition.

I would like to thank Zev Klagsbrun, Aaron Landesman, Jungin Lee, Wanlin Li, Mark Shusterman, Alex Smith, and Niudun Wang for carefully explaining their works on probability distributions of families of elliptic curves, suggesting valuable references which are relevant to this paper [KMR14, FLR20, Wan21, CLDLL22, Smi22b], and for engaging in very helpful discussions. I would like to thank Jungin Lee and Zev Klagsbrun for comments on improving the exposition of this paper. I would like to thank U Jin Choi and Yongsul Won for helpful discussions on Markov chains, and many other researchers at NIMS for shaping enjoyable experiences during my service.

Lastly, I would like to sincerely thank my parents for their wholehearted continuous support throughout my military service and in the midst of the COVID-19 global pandemic. Without their unconditional love and continuous support, the experiences I had with preparing this manuscript would not have been as enjoyable.

## REFERENCES

- [AP58] A. Rényi and P. Turán. On a theorem of erdos-kac. *Acta Arithmetica*, 4(1):71–84, 1958.
- [BCP97] Wieb Bosma, John Cannon, and Catherine Playoust. The Magma algebra system. I. The user language. *J. Symbolic Comput.*, 24(3-4):235–265, 1997. Computational algebra and number theory (London, 1993).
- [BK77] Armand Brumer and Kenneth Kramer. The rank of elliptic curves. *Duke Mathematical Journal*, 44(4):715–743, 1977.
- [BKLJ<sup>+</sup>15] Manjul Bhargava, Daniel M. Kane, Hendrik W. Lenstra Jr., Bjorn Poonen, and Eric Rains. Modelign the distribution of ranks, selmer groups, and shafarevich-tate groups of elliptic curves. *Cambridge Journal of Mathematics*, 3(3):275–321, 2015.
- [BLV09] Andrea Bandini, Ignazio Longhi, and Stefano Vigni. Torsion points on elliptic curves over function fields and a theorem of igusa. *Expositiones Mathematicae*, 27(3):175–209, 2009.
- [BS15] Manjul Bhargava and Arul Shankar. Binary quadratic forms having bounded invariants, and the boundedness of the average rank of elliptic curves. *Annals of Mathematics*, 181:1–52, 2015.
- [BSGKS20] Lior Bary-Soroker, Ofir Gorodetsky, Taelin Karidi, and Will Sawin. Chebotarev density theorem in short intervals for extensions of  $\mathbb{F}_q(t)$ . *Transactions of the American Mathematical Society*, 373:597–628, 2020.
- [CCH05] Brian Conrad, Keith Conrad, and Harald Helfgott. Root numbers and ranks in positive characteristic. *Advances in Mathematics*, 198(2):684–731, 2005.
- [CLDLL22] Antoine Comeau-Lapointe, Chantal David, Matilde Lalin, and Wanlin Li. On the vanishing of twisted l-functions of elliptic curves over rational function fields. *Preprint*, 2022.
- [CLNY22] Gilyoung Cheong, Jungin Lee, Hayan Nam, and Myungjun Yu. Jordan–landau theorem for matrices over finite fields. *Linear Algebra and its Applications*, 655:100–128, 2022.
- [Coh69] Stephen D. Cohen. Further arithmetical functions in finite fields. *Proceedings of the Edinburgh Mathematical Society*, 16:349–363, 1969.
- [Del80] Pierre Deligne. La conjecture de weil, ii. *Publications Mathématiques de l’IHES*, 52:137–252, 1980.

- [DFK07] Chantal David, Jack Fearnley, and Hershy Kisilevsky. Vanishing of  $L$ -functions of elliptic curves over number fields, ranks of elliptic curves and random matrix theory. *London Math. Soc. Lecture Note Series*, 341:247–259, 2007.
- [DH85] G. Dueck and H.C. Williams. Computation of the class number and class group of a complex cubic field. *Mathematics of Computation*, 45(171):223–231, 1985.
- [dJ02] A.J. de Jong. Counting elliptic surfaces over finite fields. *Moscow Mathematical Journal*, 2:281–311, 2002.
- [EH91] Gove Effinger and David Hayes. Additive number theory of polynomials over a finite fields. *Oxford Mathematical Monographs*, pages 75–88, 1991.
- [EK40] Paul Erdos and Mark Kac. The gaussian laws of errors in the theory of additive number theoretic functions. *American journal of mathematics*, 62(1/4):738–742, 1940.
- [EVW16] Jordan Ellenberg, Akshay Venkatesh, and Craig Westerland. Homological stability for hurwitz spaces and the cohen-lenstra conjecture over function fields. *Annals of Mathematics*, 183:729–786, 2016.
- [FIMR13] J.B. Friedlander, H. Iwaniec, B. Mazur, and K. Rubin. The spin of prime ideals. *Inventiones Mathematicae*, pages 697–749, 2013.
- [FJ08] Michael D. Fried and Moshe Jarden. *Field Arithmetic*. Springer, Berlin, Germany, 2008.
- [FJ20] Daniel Fiorilli and Florent Jouve. Distribution of frobenius elements in families of galois extensions. *Preprint available at <https://arxiv.org/abs/2001.05428>*, 2020.
- [FLR20] Tony Feng, Aaron Landesman, and Eric Rains. The geometric distribution of selmer groups of elliptic curves over function fields. *Preprint available at <https://arxiv.org/abs/2003.07517>*, 2020.
- [FWY20] Tingting Feng, Shaochen Wang, and Guangyu Yang. Large and moderate deviation principles for the erdos-kac theorem in function fields. *Statistics and probability letters*, 163, 2020.
- [Hal06] Chris Hall. Big symplectic or orthogonal monodromy modulo  $l$ . *Duke mathematics journal*, 141(1):179–203, 2006.
- [HB94] D.R. Heath-Brown. The size of selmer groups for the congruent number problem, ii. *Inventiones mathematicae*, 118:331–370, 1994.
- [How01] Everett W. Howe. Isogeny classes of abelian varieties with no principal polarizations. *Moduli of Abelian Varieties (Texel Island, 1999)*, *Birkhauser Progress in Mathematics*, pages 203–216, 2001.
- [Hsu98] Chih-Nung Hsu. On certain character sums over  $\mathbb{F}_q[t]$ . *Proceedings of the American Mathematical Society*, 126(3):647–652, 1998.
- [Igu59] J.-I. Igusa. Fibre systems of jacobian varieties (iii. fibre systems of elliptic curves). *American Journal of Mathematics*, 81:453–476, 1959.
- [Imp03] Chris Impens. Stirling’s series made easy. *The American Mathematical Monthly*, 110(8):730–735, 2003.
- [Kan13] Daniel Kane. On the ranks of the 2-selmer groups of twists of a given elliptic curve. *Algebra and Number Theory*, 7(5):1253–1279, 2013.
- [KLO15] Zev Klagsbrun and Robert J. Lemke Oliver. The distribution of 2-selmer ranks of quadratic twists of elliptic curves with partial two-torsion. *Mathematika*, 62(1):67–78, 2015.
- [KMR13] Zev Klagsbrun, Barry Mazur, and Karl. Rubin. Disparity in selmer ranks of quadratic twists of elliptic curves. *Annals of Mathematics*, 178(1):287–320, 2013.
- [KMR14] Zev Klagsbrun, Barry Mazur, and Karl Rubin. A markov model for selmer ranks in families of twists. *Compositio Mathematica*, 150:1077–1106, 2014.
- [KP21] Peter Koymans and Carlo Pagano. On the distribution of  $cl(k)[l^\infty]$  for degree  $l$  cyclic fields. *Journal of the European Mathematical Society*, 24(4):1189–1283, 2021.
- [KP22] Peter Koymans and Carlo Pagano. On stevenhagen’s conjecture. *Preprint available at <https://arxiv.org/pdf/2201.13424.pdf>*, 2022.
- [Lan21] Aaron Landesman. The geometric average size of selmer groups over function fields. *Algebra and Number Theory*, 15(3):673–709, 2021.
- [Li19] Chao Li. 2-selmer groups, 2-class groups and rational points on elliptic curves. *Transactions of the American Mathematical Society*, 371:4631–4653, 2019.
- [Liu04] Yu-Ru Liu. A generalization of the erdos-kac theorem and its applications. *Canadian Mathematical Bulletin*, 48(4):589–606, 2004.
- [LO75] J. Lagarias and A. Odlyzko. Effective versions of the chebotarev density theorem. *Proceedings of Symposia in Pure Mathematics*, pages 409–464, 1975.
- [Mil06] J.S. Milne. *Arithmetic Duality Theorems*. BookSurge, LLC, second edition, 2006.
- [Mon96] P. Monsky. Generalizing the birch-stephens theorem. i. modular curves. *Mathematische Zeitschrift*, 221(3):415–420, 1996.
- [MP08] Adam Morgan and Ross Paterson. On 2-selmer groups of twists after quadratic extension. *Preprint available at <https://arxiv.org/abs/2011.04374>*, 145(3):513–526, 2008.



- [MR07] Barry Mazur and Karl Rubin. Finding large selmer rank via an arithmetic theory of local constants. *Annals of Mathematics*, 166:579–612, 2007.
- [MR19] Barry Mazur and Karl Rubin. Arithmetic conjectures suggested by the statistical behavior of modular symbols. *To appear in Experimental Mathematics*, 2019.
- [MRS07] Barry Mazur, Karl Rubin, and Alice Silverberg. Twisting commutative algebraic groups. *Journal of Algebra*, 314(1):419–438, 2007.
- [MT93] Sean Meyn and Richard Tweedie. *Markov chains and stochastic stability*. Springer-Verlag, Berlin, Germany, 1993.
- [MZ16] Behzad Mehrdad and Lingjiao Zhu. Moderate and large deviations for erdos-kac theorem. *Quarterly Journal of Mathematics*, 67(1):147–160, 2016.
- [Pit74] J.W. Pitman. Uniform rates of convergence for markov chain transition probabilities. *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete*, 29:193–227, 1974.
- [Poo15] Bjorn Poonen. Average rank of elliptic curves, after manjul bhargava and arul shankar. *Seminaire Bourbaki*, 1049:1–17, 2015.
- [PR12] Bjorn Poonen and Eric Rains. Random maximal isotropic subspaces and selmer groups. *Journal of the AMS*, 25(1):245–269, 2012.
- [PTBW20] Lilian B. Pierce, Caroline L. Turnage-Butterbaugh, and Melanie Matchett Wood. An effective chebotarev density theorem for families of number fields, with an application to  $l$ -torsion in class groups. *Inventiones mathematicae*, 219:701–778, 2020.
- [PW21] Sun Woo Park and Niudun Wang. On the average of  $p$ -selmer rank in quadratic twist families of elliptic curves over function field. *Preprint available at <https://arxiv.org/abs/2102.00549?context=math>*, 2021.
- [QH14] B.C. Ngo Q.P. Ho, V.B. Le Hung. Average size of 2-selmer groups of elliptic curves over function fields. *Mathematical research letters*, 21:1305–1339, 2014.
- [Riz03] Ottavio G. Rizzo. Average root numbers for a nonconstant family of elliptic curves. *Compositio Mathematica*, 136:1–23, 2003.
- [Ros02] Michael Rosen. *Number theory in function fields*. Springer, Berlin, Germany, 2002.
- [SD08] Peter Swinnerton-Dyer. The effect of twisting on the 2-selmer group. *Mathematical Proceedings of the Cambridge Philosophical Society*, 145(3):513–526, 2008.
- [Ser82] J.-P. Serre. Quelques applications du théorème de densité de chebotarev. *Publications MATHématiques de l’IHES*, 54:123–201, 1982.
- [Smi17] Alexander Smith.  $2^\infty$ -selmer groups,  $2^\infty$ -class groups, and goldfeld’s conjecture. *Preprint available at <https://arxiv.org/abs/1702.02325>*, 2017.
- [Smi20] Alexander Smith.  $l^\infty$ -selmer groups in degree  $l$  twist families. *Doctoral dissertation, Harvard University*, 2020.
- [Smi22a] Alexander Smith. The distribution of fixed point selmer groups in twist families. *Preprint available at <https://arxiv.org/pdf/2207.05143.pdf>*, 2022.
- [Smi22b] Alexander Smith. The distribution of  $l^\infty$  selmer groups in degree  $l$  twist families. *Preprint available at <https://arxiv.org/abs/2207.05674>*, 2022.
- [SPT08] Daniel Barrera Salazar, Ariel Pacetti, and Gonzalo Tornaria. On 2-selmer groups and quadratic twists of elliptic curves. *To appear in Mathematical Research Letters*. *Preprint available at <https://arxiv.org/abs/2001.02263>*, 145(3):513–526, 2008.
- [Ste02] William Stein. Shafarevich-tate groups of nonsquare order. *Modular Curves and Abelian Varieties 2002 Barcelona Conference Proceedings, Birkhauser Progress in Mathematics*, 224:277–289, 2002.
- [TY14] Fabien Trihan and Seidai Yasuda. The  $l$ -parity conjecture for abelian varieties over function fields of characteristic  $p > 0$ . *Compositio Mathematica*, 150(4):507–522, 2014.
- [Wan21] Niudun Wang. 2-selmer groups of quadratic twists of elliptic curves. *Ph.D. Thesis available at <https://depot.library.wisc.edu/repository/fedora/1711.dl:ITQ5IQCZ72U6I8H/datastreams/REF/content>*, 2021.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN – MADISON, 480 LINCOLN DR., MADISON, WI 53706, USA

NATIONAL INSTITUTE FOR MATHEMATICAL SCIENCES, 463-1 JEONMIN-DONG, YUSEONG-GU, DAEJEON, 34047, REPUBLIC OF KOREA.

Email address: [spark483@math.wisc.edu](mailto:spark483@math.wisc.edu)