arXiv:2211.11648v3 [math.NT] 5 Dec 2022

Sums of powers of integers and generalized Stirling numbers of the second kind

José Luis Cereceda
Collado Villalba, 28400 (Madrid), Spain
jl.cereceda@movistar.es

Abstract

By applying the Newton-Gregory expansion to the polynomial associated with the sum of powers of integers $S_k(n) = 1^k + 2^k + \cdots + n^k$, we derive a couple of infinite families of explicit formulas for $S_k(n)$. One of the families involves the r-Stirling numbers of the second kind $\left\{ {k \atop j} \right\}_r$, $j = 0, 1, \ldots, k$, while the other involves their duals $\left\{ {k \atop j} \right\}_{-r}$, with both families of formulas being indexed by the non-negative integer r. As a by-product, we obtain three additional formulas for $S_k(n)$ involving the numbers $\left\{ {k \atop j} \right\}_{n+m}$, $\left\{ {k \atop j} \right\}_{n-m}$ (where m is any given non-negative integer), and $\left\{ {k \atop j} \right\}_{k-j}$, respectively. Moreover, we provide a formula for the Bernoulli polynomials $B_k(x-1)$ in terms of $\left\{ {k \atop j} \right\}_x$ and the harmonic numbers.

1 Introduction

Following Broder [5, Equation 57] (see also Carlitz [6, Equation (3.2)]) we define the generalized (or weighted) Stirling numbers of the second kind by

$$R_{k,j}(x) = \sum_{i=0}^{k-j} \binom{k}{i} \left\{ {k-i \atop j} \right\} x^i, \quad \text{integers } 0 \le j \le k,$$

where x stands for any arbitrary real or complex value, and where the $\left\{ {k \atop j} \right\}$'s are the ordinary Stirling numbers of the second kind [3]. Note that $R_{k,j}(x)$ is a polynomial in x of degree $k-j$ with leading coefficient $\binom{k}{j}$ and constant term $R_{k,j}(0) = \left\{ {k \atop j} \right\}$. Furthermore, we have that $R_{k,j}(1) = \left\{ {k+1 \atop j+1} \right\}$. In general, when x is the non-negative integer r, $R_{k,j}(r)$ becomes the r-Stirling number of the second kind $\left\{ {k+r \atop j+r} \right\}_r$ [5]. A combinatorial interpretation of the polynomial $R_{k,j}(x)$ is given in [5, Theorem 27] (see also the definition provided by Bényi and Matsusaka in [1, Definition 2.13]).

For convenience and notational simplicity, in this paper we employ the notation $\left\{ {k \atop j} \right\}_r$ to refer to Broder's r-Stirling numbers of the second kind $\left\{ {k+r \atop j+r} \right\}_r$. The former notation has been used recently by Ma and Wang in [14] (see also [1] and [15]). The numbers $\left\{ {k \atop j} \right\}_r$ are then given by

$$\left\{ {k \atop j} \right\}_r = \sum_{i=0}^{k-j} \binom{k}{i} \left\{ {k-i \atop j} \right\} r^i, \quad \text{integer } r \ge 0.$$

Likewise, adopting the notation in [14], we define the counterpart or dual of $\left\{ {k \atop j} \right\}_r$ for negative integer r as

$$\left\{ {k \atop j} \right\}_{-r} = \sum_{i=0}^{k-j} (-1)^i \binom{k}{i} \left\{ {k-i \atop j} \right\} r^i, \quad \text{integer } r \ge 0.$$

Alternatively, $\left\{ {k \atop j} \right\}_r$ and $\left\{ {k \atop j} \right\}_{-r}$ can be equivalently expressed in the form

$$\left\{ {k \atop j} \right\}_r = \frac{1}{j!} \sum_{i=0}^{j} (-1)^{j-i} \binom{j}{i} (i+r)^k, \quad \text{integer } r \geq 0, \tag{1}$$

and

$$\left\{ {k \atop j} \right\}_{-r} = \frac{1}{j!} \sum_{i=0}^{j} (-1)^{j-i} \binom{j}{i} (i-r)^k, \quad \text{integer } r \geq 0, \tag{2}$$

respectively. Clearly, both $\left\{ {k \atop j} \right\}_r$ and $\left\{ {k \atop j} \right\}_{-r}$ reduce to $\left\{ {k \atop j} \right\}$ when $r = 0$. It is to be noted that the numbers $\left\{ {k \atop j} \right\}_{-r}$ were introduced and studied by Koutras under the name of non-central Stirling numbers of the second kind and denoted by $S_r(k, j)$ (see [13, Equations (2.5) and (2.6)]).

On the other hand, for non-negative integer k, let $S_k(n)$ denote the sum of k-th powers of the first n positive integers

$$S_k(n) = 1^k + 2^k + \cdots + n^k,$$

with $S_k(0) = 0$ for all k. As is well known, $S_k(n)$ can be expressed in terms of the Stirling numbers of the second kind as (see, e.g., [18])

$$S_k(n) = -\delta_{k,0} + \sum_{j=0}^{k} j! \binom{n+1}{j+1} \left\{ {k \atop j} \right\}, \tag{3}$$

where $\delta_{k,0}$ is the Kronecker delta, which ensures that $S_0(n) = n$. Additionally, $S_k(n)$ admits the following variant of (3):

$$S_k(n) = \sum_{j=1}^{k+1} (j-1)! \binom{n}{j} \left\{ {k+1 \atop j} \right\} = \sum_{j=0}^{k} j! \binom{n}{j+1} \left\{ {k+1 \atop j+1} \right\}, \tag{4}$$

(see, e.g., [20, Equation (9)], [7], [11, Corollary 2], and [8, Theorem 5]). The first expression in (4) can be readily obtained from the exponential generating function [4, Equation (11)]

$$\sum_{n=1}^{\infty} (1^k + 2^k + \cdots + n^k) \frac{x^n}{n!} = e^x \sum_{j=1}^{k+1} \frac{1}{j} \left\{ {k+1 \atop j} \right\} x^j.$$

Of course, (3) and (4) are equivalent formulas. Indeed, it is a simple exercise to convert (3) into (4), and vice versa, by means of the recursion $\left\{ {k \atop j} \right\} = j \left\{ {k-1 \atop j} \right\} + \left\{ {k-1 \atop j-1} \right\}$ and the well-known combinatorial identity $\binom{n}{j+1} + \binom{n}{j} = \binom{n+1}{j+1}$.

Incidentally, it is worthwhile to mention that, in his 1928 Monthly article [10], Ginsburg wrote down explicitly the first few instances of (4) for $k = 2, 3, 4, 5$ in terms of the binomial coefficients

$\binom{n}{j+1}$, where $j = 0, 1, \ldots, k$, namely

$$\begin{aligned}
S_2(n) &= \binom{n}{1} + 3\binom{n}{2} + 2\binom{n}{3},\\
S_3(n) &= \binom{n}{1} + 7\binom{n}{2} + 12\binom{n}{3} + 6\binom{n}{4},\\
S_4(n) &= \binom{n}{1} + 15\binom{n}{2} + 50\binom{n}{3} + 60\binom{n}{4} + 24\binom{n}{5},\\
S_5(n) &= \binom{n}{1} + 31\binom{n}{2} + 180\binom{n}{3} + 390\binom{n}{4} + 360\binom{n}{5} + 120\binom{n}{6}.
\end{aligned}$$

As noted by Ginsburg, the above formulas appeared on page 88 of the book by Schwatt, *Introduction to Operations with Series* (Philadelphia, The Press of the University of Pennsylvania, 1924).

In this paper, we obtain a unifying formula for $S_k(n)$ giving (3) and (4) as particular cases. Indeed, we derive a couple of infinite families of explicit formulas for $S_k(n)$, one of them involving the numbers $\left\{ {k \atop j} \right\}_r$ and the other the numbers $\left\{ {k \atop j} \right\}_{-r}$, with $j = 0, 1, \ldots, k$. Specifically, we establish the following theorem which constitutes the main result of this paper.

Theorem 1. *Let k and n be any non-negative integers and let $\left\{ {k \atop j} \right\}_r$ and $\left\{ {k \atop j} \right\}_{-r}$ be the numbers defined in (1) and (2), respectively, where r stands for any arbitrary but fixed non-negative integer. Then*

$$S_k(n) = \sum_{j=0}^{k} j! \left[\binom{n+1-r}{j+1} + (-1)^j \binom{r+j-1}{j+1} \right] \left\{ {k \atop j} \right\}_r, \tag{5}$$

and

$$S_k(n) = \sum_{j=0}^{k} j! \left[\binom{n+1+r}{j+1} - \binom{r+1}{j+1} \right] \left\{ {k \atop j} \right\}_{-r}. \tag{6}$$

Before we prove Theorem 1 in the next section, a few observations are in order.

Remark 1. *It is easily seen that both (5) and (6) reduce to (3) when $r = 0$. Furthermore, (5) reduces to (4) when $r = 1$. Moreover, setting $r = n$ in (5) leads to*

$$S_k(n) = n^{k+1} + \sum_{j=1}^{k} (-1)^j j! \binom{n+j-1}{j+1} \left\{ {k \atop j} \right\}_n. \tag{7}$$

Similarly, setting $r = n+1$ in (5) yields

$$S_k(n) = \sum_{j=0}^{k} (-1)^j j! \binom{n+j}{j+1} \left\{ {k \atop j} \right\}_{n+1}, \tag{8}$$

retrieving the result obtained in [12, Equation (4.8)].

Remark 2. *It should be stressed that both (5) and (6) hold irrespective of the value taken by the non-negative integer parameter r. This means that, actually, the right-hand side of (5) and (6) provides us with an infinite supply of explicit formulas for $S_k(n)$, one for each value of r. For example, for $r = 2$, and noting that $S_k(1) = 1$ for all k, we have from (5)*

$$S_k(n) = 1 + \sum_{j=0}^{k} j! \binom{n-1}{j+1} \left\{ {k \atop j} \right\}_2,$$

where

$$\left\{ {k \atop j} \right\}_2 = \frac{1}{j!} \sum_{i=0}^{j} (-1)^{j-i} \binom{j}{i} (i+2)^k.$$

Analogously, for $r = 2$, we have from (6)

$$S_k(n) = -\delta_{k,0} + (-1)^{k+1}(1 + 2^k) + \sum_{j=0}^{k} j! \binom{n+3}{j+1} \left\{ {k \atop j} \right\}_{-2},$$

where

$$\left\{ {k \atop j} \right\}_{-2} = \frac{1}{j!} \sum_{i=0}^{j} (-1)^{j-i} \binom{j}{i} (i-2)^k.$$

Remark 3. *In the last section, we obtain a more general formula for $S_k(n)$ involving the numbers $\left\{ {k \atop j} \right\}_{n+m}$ and $\left\{ {k \atop j} \right\}_{n-m}$, where m is any given non-negative integer (see equations (19) and (20)). Furthermore, we provide another formula for $S_k(n)$ involving the numbers $\left\{ {k \atop j} \right\}_{k-j}$ (see equation (21)), as well as a formula for the Bernoulli polynomials $B_k(x-1)$ in terms of $\left\{ {k \atop j} \right\}_x$ and the harmonic numbers (see equation (22)).*

2 Proof of Theorem 1

The proof of Theorem 1 is based on the following lemma.

Lemma 1. *For x a real or complex variable, let $S_k(x)$ denote the unique interpolating polynomial in x of degree $k+1$ such that $S_k(x) = 1^k + 2^k + \cdots + x^k$ whenever x is a positive integer (with $S_k(0) = 0$). Then*

$$S_k(x) = S_k(a-1) + \sum_{j=0}^{k} j! \binom{x+1-a}{j+1} R_{k,j}(a), \tag{9}$$

where a is a parameter taking any arbitrary but fixed real or complex value.

Proof. As is well known (see, e.g., [9, Equation (15]), $S_k(x)$ can be expressed in terms of the Bernoulli polynomials $B_k(x)$ as follows

$$S_k(x) = \frac{1}{k+1}[B_{k+1}(x+1) - B_{k+1}(1)]. \tag{10}$$

Recall further that the forward difference operator Δ acting on the function $f(x)$ is defined by $\Delta f(x) = f(x+1) - f(x)$. Then, the following elementary result

$$\Delta S_k(x) = (x+1)^k \tag{11}$$

follows immediately from (10) and the difference equation $\Delta B_{k+1}(x) = (k+1)x^k$ [9, Equation (12)].

On the other hand, the Newton-Gregory expansion of the function $f(x)$ is given by (see, e.g., [17, Equation (A.9), p. 230])

$$f(x) = \sum_{j=0}^{\infty} \binom{x-a}{j} \Delta^j f(a),$$

where, for any integer $j \geq 1$, the j-th order difference operator Δ^j is defined by $\Delta^j f(x) = \Delta(\Delta^{j-1} f(x)) = \Delta^{j-1}(\Delta f(x))$ and $\Delta^0 f(x) = f(x)$, and where $\Delta^j f(a) = \Delta^j f(x)|_{x=a}$. Hence, applying the Newton-Gregory expansion to the power sum polynomial $S_k(x)$ and using (11) yields

$$S_k(x) = S_k(a) + \sum_{j=0}^{k} \binom{x-a}{j+1} \Delta^j (a+1)^k, \tag{12}$$

where the terms in the summation with index j greater than k have been omitted because $\Delta^j (x+1)^k = 0$ for all $j \geq k+1$ [17, Equation (6.16), p. 68].

The connection between (12) and the generalized Stirling numbers $R_{k,j}(x)$ stems from the fact that (see, e.g., [5, Theorem 29] and [6, Equation (3.8)])

$$R_{k,j}(x) = \frac{1}{j!} \Delta^j x^k. \tag{13}$$

Therefore, combining (12) and (13), and making $a \to a-1$, we get (9). □

When x and a are the non-negative integers n and r, respectively, (9) becomes

$$S_k(n) = S_k(r-1) + \sum_{j=0}^{k} j! \binom{n+1-r}{j+1} \left\{ {k \atop j} \right\}_r, \tag{14}$$

with $S_k(-1) = 0$ for all $k \geq 1$, and $S_0(-1) = -1$. Now, letting $n = 0$ in (14) and using the relation

$$\binom{-x}{k} = (-1)^k \binom{x+k-1}{k} \tag{15}$$

gives

$$S_k(r-1) = \sum_{j=0}^{k} (-1)^j j! \binom{r+j-1}{j+1} \left\{ {k \atop j} \right\}_r. \tag{16}$$

Then, substituting (16) in (14), we get (5).

Moreover, making the transformation $r \to -r$ in (14) and invoking the symmetry property of the power sum polynomial (see, e.g., [16, Theorem 10])

$$S_k(-r-1) = -\delta_{k,0} + (-1)^{k+1} S_k(r),$$

it follows that

$$S_k(n) = -\delta_{k,0} - (-1)^k S_k(r) + \sum_{j=0}^{k} j! \binom{n+1+r}{j+1} \left\{ {k \atop j} \right\}_{-r}, \tag{17}$$

from which, upon setting $n = 0$, we further obtain

$$(-1)^k S_k(r) = -\delta_{k,0} + \sum_{j=0}^{k} j! \binom{r+1}{j+1} \left\{ {k \atop j} \right\}_{-r}. \tag{18}$$

Finally, substituting (18) in (17), we get (6).

We conclude this section with the following two remarks.

Remark 4. *By renaming r as n in (18) we find that*

$$S_k(n) = (-1)^k \left(-\delta_{k,0} + \sum_{j=0}^{k} j! \binom{n+1}{j+1} \left\{ {k \atop j} \right\}_{-n} \right),$$

which may be compared with (3).

Remark 5. *It is to be noted that the equation (14) above is equivalent to the equation appearing in [2, Corollary 2.2] in which $d = 1$ and $a = r$.*

3 Concluding remarks

Let us observe that, by letting $r = n + m$ in (5), where m is any given non-negative integer, and using (15), we obtain

$$S_k(n) = \sum_{j=0}^{k} (-1)^j j! \left[\binom{n+m+j-1}{j+1} - \binom{m+j-1}{j+1} \right] \left\{ {k \atop j} \right\}_{n+m}. \tag{19}$$

Of course, (19) reduces to (7) and (8) when $m = 0$ and $m = 1$, respectively. Similarly, putting $r = n - m$ in (5), where m is any given non-negative integer, we obtain

$$S_k(n) = \sum_{j=0}^{k} j! \left[\binom{m+1}{j+1} + (-1)^j \binom{n+j-m-1}{j+1} \right] \left\{ {k \atop j} \right\}_{n-m}. \tag{20}$$

Note that, when $m = n$, (20) reduces to (3).

On the other hand, applying the formula $\left\{ {k \atop j} \right\}_{-r} = (-1)^{k-j} \left\{ {k \atop j} \right\}_{r-j}$ (see [14, Equation (2.4)]) and taking $r = k$ in (6) yields

$$S_k(n) = \sum_{j=0}^{k} (-1)^{k-j} j! \left[\binom{n+k+1}{j+1} - \binom{k+1}{j+1} \right] \left\{ {k \atop j} \right\}_{k-j}. \tag{21}$$

Incidentally, for $n = 1$, (21) gives us the identity

$$\sum_{j=0}^{k} (-1)^{k-j} j! \binom{k+1}{j} \left\{ {k \atop j} \right\}_{k-j} = 1.$$

Moreover, from (10) and (16), we obtain the following formula for the Bernoulli polynomials evaluated at the non-negative integer r

$$B_{k+1}(r) = B_{k+1}(1) + (k+1)\sum_{j=0}^{k}(-1)^j j!\binom{r+j-1}{j+1}\left\{ {k \atop j} \right\}_r, \quad \text{integer } r \geq 0.$$

Likewise, making $r \to -r$ in the preceding equation and using (15) gives the following formula for the Bernoulli polynomials evaluated at the negative integer $-r$

$$B_{k+1}(-r) = B_{k+1}(1) - (k+1)\sum_{j=0}^{k} j!\binom{r+1}{j+1}\left\{ {k \atop j} \right\}_{-r}, \quad \text{integer } r \geq 0.$$

One can naturally extend the above formula for $B_{k+1}(r)$ to apply to any real or complex variable x as follows

$$B_{k+1}(x) = B_{k+1}(1) + (k+1)\sum_{j=0}^{k}(-1)^j j!\binom{x+j-1}{j+1}\left\{ {k \atop j} \right\}_x,$$

where, using the notation in [1], $\left\{ {k \atop j} \right\}_x$ refers to the Stirling polynomial of the second kind $R_{k,j}(x)$, namely

$$\left\{ {k \atop j} \right\}_x = \frac{1}{j!}\sum_{i=0}^{j}(-1)^{j-i}\binom{j}{i}(i+x)^k.$$

Finally, we point out that $B_k(x-1)$ can be expressed in the form

$$B_k(x-1) = \sum_{j=0}^{k}(-1)^j j! H_{j+1}\left\{ {k \atop j} \right\}_x, \tag{22}$$

where $H_j = 1 + \frac{1}{2} + \cdots + \frac{1}{j}$ is the j-th harmonic number. In particular, setting $x = 1$ in (22) yields the following known formula for the Bernoulli numbers (see, e.g., [19, Equation (5.9)])

$$B_k = \sum_{j=0}^{k}(-1)^j j! H_{j+1}\left\{ {k+1 \atop j+1} \right\}.$$

Furthermore, from (10) and (22), we arrive at the following formula for the sum of powers of integers

$$S_{k-1}(n) = \frac{1}{k}\sum_{j=0}^{k}(-1)^j j! H_{j+1}\left(\left\{ {k \atop j} \right\}_{n+2} - \left\{ {k \atop j} \right\}_2\right),$$

which holds for any integers $n \geq 0$ and $k \geq 1$, with $S_{k-1}(0) = 0$ for all $k \geq 1$.

References

[1] Bényi, B., Matsusaka, T. (2021). Combinatorial aspects of poly-Bernoulli polynomials and poly-Euler numbers, preprint. Available at: arXiv:2106.05585v2 [math.CO].

[2] Bounebirat, F., Laissaoui, D., Rahmani, M. (2018). Several explicit formulae of sums and hyper-sums of powers of integers. *Online J. Anal. Comb.* 13, Article #4, 9 pp.

[3] Boyadzhiev, K. N. (2012). Close encounters with the Stirling numbers of the second kind. *Math. Mag.* 85(4):252–266.

[4] Boyadzhiev, K. N. (2020). Sums of powers and special polynomials. *Discuss. Math. Gen. Algebra Appl.* 40(2):275–283.

[5] Broder, A. Z. (1984). The r-Stirling numbers. *Discrete Math.* 49(3):241–259.

[6] Carlitz, L. (1980). Weighted Stirling numbers of the first and second kind—I. *Fibonacci Quart.* 18(2):147–162.

[7] Cereceda, J. L. (2015). Newton's interpolation polynomial for the sums of powers of integers. *Amer. Math. Monthly.* 122(10):1007.

[8] Chrysafi, L., Marques, C. A. (2021). Sums of powers via matrices. *Math. Mag.* 94(1):43–52.

[9] El-Mikkawy, M., Atlan, F. (2013). Derivation of identities involving some special polynomials and numbers via generating functions with applications. *Appl. Math. Comput.* 220:518–535.

[10] Ginsburg, J. (1928). Note on Stirling's numbers. *Amer. Math. Monthly.* 35(2):77–80.

[11] Kargın, L., Dil, A., Can, M. (2020). Formulas for sums of powers of integers and their reciprocals, preprint. Available at: arXiv:2006.01132v1 [math.CO].

[12] Kargın, L., Cenkci, M., Dil, A., Can, M. (2022). Generalized harmonic numbers via poly-Bernoulli polynomials. *Publ. Math. Debrecen.* 100(3-4):365–386.

[13] Koutras, M. (1982). Non-central Stirling numbers and some applications. *Discrete Math.* 42(1):73–89.

[14] Ma, Q., Wang, W. (2023). Riordan arrays and r-Stirling number identities. *Discrete Math.* 346(1):113211.

[15] Merris, R. (2000). The p-Stirling numbers. *Turk. J. Math.* 24(4):379–399.

[16] Newsome, N. J., Nogin, M. S., Sabuwala, A. H. (2017). A proof of symmetry of the power sum polynomials using a novel Bernoulli number identity. *J. Integer Seq.* 20, Article 17.6.6, 10 pp.

[17] Quaintance, J., Gould, H. W. (2016). *Combinatorial Identities for Stirling Numbers. The Unpublished Notes of H W Gould*. Singapore: World Scientific Publishing.

[18] Shirali, S. (2018). Stirling set numbers & powers of integers. *At Right Angles.* 7(1):26–32.

[19] Wang, W. (2010). Riordan arrays and harmonic number identities. *Comput. Math. Appl.* 60(5):1494–1509.

[20] Wituła, R., Kaczmarek, K., Lorenc, P., Hetmaniok, E., Pleszczyński. (2014). Jordan numbers, Stirling numbers and sums of powers. *Discuss. Math. Gen. Algebra Appl.* 34(2):155–166.