

Counting atypical black hole microstates from entanglement wedges

Zixia Wei^{a,b} and Yasushi Yoneta^c

^a*Center for Gravitational Physics and Quantum Information,
Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto 606-8502, Japan*

^b*Kavli Institute for Theoretical Physics,
University of California, Santa Barbara, CA 93106, USA*

^c*Department of Basic Science,
The University of Tokyo, 3-8-1 Komaba, Meguro, Tokyo 153-8902, Japan*

Abstract

Disentangled black hole microstates are atypical states in holographic CFTs whose gravity duals do not have smooth horizons. If there exist sufficiently many disentangled microstates to account for the entire Bekenstein-Hawking entropy, then any black hole microstate can be written as a superposition of states without smooth horizons. We show that there exist sufficiently many disentangled microstates to account for almost the entire Bekenstein-Hawking entropy of a large AdS black hole at the semiclassical limit $G_N \rightarrow 0$. In addition, we also argue that in generic quantum many-body systems with short-ranged interactions, there exist sufficiently many area law states in the microcanonical subspace to account for almost the entire thermodynamic entropy in the standard thermodynamic limit. Area law states are atypical since a typical state should contain volume law entanglement. Furthermore, we also present an explicit way to construct such a set of area law states, and argue that the same construction may also be used to construct disentangled states.

Contents

| | | |
|----------|--|-----------|
| 1 | Introduction | 2 |
| 2 | Preliminaries | 6 |
| 2.1 | Entanglement of formation and other correlation measures | 6 |
| 2.2 | Entanglement of purification in AdS/CFT | 9 |
| 2.3 | Holographic dual of the microcanonical ensemble | 11 |
| 3 | Disentangled black hole microstates in AdS/CFT | 13 |
| 3.1 | Upper bounding E_F from holography | 13 |
| 3.2 | General results for microcanonical ensembles | 15 |
| 3.3 | Existence of $e^{S_{BH}}$ disentangled microstates | 19 |
| 4 | Area law states in quantum many-body systems | 21 |
| 4.1 | Conjecture for E_{RE} | 21 |
| 4.2 | Existence of e^S area law microstates | 22 |
| 5 | Constructing area law states | 23 |
| 5.1 | Constructing an ensemble of states with low entanglement | 23 |
| 5.2 | Constructing e^S area law states | 26 |
| 5.3 | Constructing $e^{S_{BH}}$ disentangled states in holographic CFTs? | 27 |
| 6 | Summary and Discussions | 27 |
| 6.1 | Connections to additivity conjectures | 28 |
| 6.2 | Atypical microstate counting for black holes | 28 |
| 6.3 | Why atypicality is important in studying quantum gravity? | 29 |
| A | Axioms for entanglement measures | 30 |
| B | Alternative proofs of corollary 4 and 6 | 31 |

1 Introduction

The notion of typicality [1–4] lies in the most fundamental part of statistical mechanics, which provides an explanation for the basic principles of equilibrium statistical mechanics [5]. On the other hand, generic or collective features of atypical states are much less investigated. One reason for the lack of interest is because generic atypical states in generic quantum many-body systems will eventually thermalize and become a typical state under time evolution, and hence are difficult to be related to experimental observations.

However, studying atypical states appear to be useful in a seemingly totally different context: black hole physics. General relativity predicts the existence of a smooth horizon as well as a black hole interior, which leads to mysteries such as the information paradox [6,7] and the AMPS firewall paradox [8,9] when combined with fundamental principles in quantum theory. A long-standing yet crucial question in the study of quantum gravity is whether microstates of black holes also have smooth horizons, or the existence of smooth horizons and black hole interior is merely an emergent illusion of (semi-)classical gravity after coarse-graining from a UV complete quantum gravitational theory [8–12].

In this context, it is enlightening to consider a class of atypical states called disentangled states [13] in holographic conformal field theories. These are states which have a large entanglement deficit compared to typical states. According to the AdS/CFT correspondence [14] and the holographic entanglement entropy formulae [15–20], a disentangled state on the CFT side cannot be dual to AdS geometry which has a smooth horizon [13]. In other words, a disentangled state either corresponds to a geometry that has nontrivial structures at or outside the horizon, or does not have a geometrical description at all [13].

More precisely, for a typical AdS black hole state $|\text{typ}\rangle$, when dividing the corresponding CFT into two regions A and B where B is smaller than A , the von Neumann entropy of the reduced density matrix on B is the same as that of a canonical density matrix $\rho^{(\text{can})}$ with the corresponding temperature, i.e.

$$S(\text{Tr}_A |\text{typ}\rangle\langle\text{typ}|) \approx S(\text{Tr}_A \rho^{(\text{can})}) \quad (1.1)$$

at the leading order of $1/G_N$. Here, G_N is the Newton constant in the AdS gravity, and the $G_N \rightarrow 0$ limit corresponds to the large degrees of freedom limit (large gauge rank number N limit, large central charge c limit, etc.) on the CFT side. Therefore, the von Neumann entropy of $\rho_B^{(\text{typ})} \equiv \text{Tr}_A |\text{typ}\rangle\langle\text{typ}|$, i.e. the entanglement entropy between A and B , can be

evaluated by a Ryu-Takayanagi surface [15,16] or quantum extremal surface [19,20] wrapping the black hole horizon like the one shown in the left of figure 1.




Figure 1: The geometry of an AdS black hole. The asymptotic boundary where the dual CFT lives is shown in gray. The spatial region of the CFT is divided into A and B where B is slightly smaller than A . (Left) The orange surface $\Gamma_B^{(\min)}$ is the Ryu-Takayanagi surface that gives the entanglement entropy for a typical microstate and it wraps the black hole horizon. (Right) The blue surface $\Sigma_{AB}^{(\min)}$ gives $\frac{1}{2}S(\text{Tr}_{AA^*}|\text{TFD}\rangle\langle\text{TFD}|)$ and coincides the entanglement wedge cross section for a mixed thermal state.

On the other hand, if a pure state has way much less entanglement between A and B , then it will be atypical. Such an atypical state is called a disentangled state. To be more concrete for later discussions, let us define disentangled states in a rigorous way, following the one given in [13].

Definition (Disentangled States). *A disentangled state $|\text{disent}\rangle$ is a state which satisfies*

$$S(\text{Tr}_A|\text{disent}\rangle\langle\text{disent}|) \lesssim \frac{1}{2}S(\text{Tr}_{AA^*}|\text{TFD}\rangle\langle\text{TFD}|), \quad (1.2)$$

where “ \lesssim ” means “less than or equal to at the leading order of $1/G_N$ ”.

Here, $|\text{TFD}\rangle$ is the thermofield double state

$$|\text{TFD}\rangle \propto \sum_i e^{-\beta E_i/2} |E_i\rangle |E_i^*\rangle, \quad (1.3)$$

where $|E_i\rangle$ is the energy eigenstate associated to the eigenvalue E_i , and $|E_i^*\rangle$ is its CPT conjugate living in a copy of the original Hilbert space. $A^*(B^*)$ is the copy of A (B). Noting

that $|\text{TFD}\rangle$ is a purification of the canonical density matrix $\rho^{(\text{can})}$,

$$\rho^{(\text{can})} = \text{Tr}_{A^*B^*} |\text{TFD}\rangle\langle\text{TFD}|, \quad (1.4)$$

it directly follows from the positivity of the mutual information that

$$S(\text{Tr}_A \rho^{(\text{can})}) - \frac{1}{2} S(\text{Tr}_{AA^*} |\text{TFD}\rangle\langle\text{TFD}|) \geq 0. \quad (1.5)$$

In a holographic CFT, the gap is $O(1/G_N)$ when the corresponding AdS black hole is sufficiently large. In this case, the entanglement held by a disentangled state is significantly smaller than a typical state, which is not sufficient to support an existence of a smooth horizon in its gravity dual [13]. In a black hole geometry, the surface which gives $\frac{1}{2} S(\text{Tr}_{AA^*} |\text{TFD}\rangle\langle\text{TFD}|)$ ends on the black hole horizon as shown in the right of figure 1.

Based on these, it is natural to ask how many disentangled states one can find in the sub-Hilbert space of the black hole microstates, which is spanned by the energy eigenstates in the energy window¹ $[E - \Delta E, E]$. Let us denote this subspace as $\mathcal{H}_{[E-\Delta E, E]}$ and call it the microcanonical subspace. In [13], Hayden and Penington showed that if extensive violations of additivity conjectures do not exist, then one can find $\exp[S_{BH} - O(1)]$ orthogonal disentangled states in $\mathcal{H}_{[E-\Delta E, E]}$, which is sufficient to account for the Bekenstein-Hawking entropy S_{BH} at $G_N \rightarrow 0$. Here, additivity conjectures are a series of equivalent conjectures in quantum information and communication theory [21]. However, small violations of additivity conjectures have already been found [22], and there is so far no upper bound for violations of the additivity conjectures. In other words, their extensive violation may be even expectable. Possibilities of extensive violations are further discussed in, for example, [23, 24]. On the other hand, the existence of $\exp[S_{BH} - O(1)]$ orthogonal disentangled states in $\mathcal{H}_{[E-\Delta E, E]}$ is surprising since it means that any black hole microstate can be written as a superposition of states without smooth horizons at $G_N \rightarrow 0$. Therefore, one may argue that, since this is hard to believe, one may be able to construct an explicit example of extensive violations of additivity conjectures from black holes. From this point of view, it is important to know how far one can go without any assumptions on additivity conjectures.

In this paper, we show that one can find at least $\exp[S_{BH}(1 - o(1))]$ orthogonal disentangled states in $\mathcal{H}_{[E-\Delta E, E]}$, which can account for almost the entire Bekenstein-Hawking entropy S_{BH} at $G_N \rightarrow 0$. Our proof has no assumptions on additivity conjectures. We start

¹Our arguments in this paper hold for any choice of the energy window as long as one can get a valid microcanonical ensemble from it. We will take $[E - \Delta E, E]$ just for convenience.

from the observation that the surface which gives $\frac{1}{2}S(\text{Tr}_{AA^*}|\text{TFD}\rangle\langle\text{TFD}|)$ (sketched in the right of figure 1) is exactly the same as a geometrical quantity called entanglement wedge cross section [25, 26] in a black hole background, which can be related to a quantum informational quantity called entanglement of purification [27] in the corresponding holographic CFT. This allows us to give an upper bound to the entanglement of formation [28] of the microcanonical ensemble $\rho^{(\text{micro})}$ defined on $\mathcal{H}_{[E-\Delta E, E]}$. This upper bound guarantees at least one disentangled state exists in $\mathcal{H}_{[E-\Delta E, E]}$. Then we can exclude this disentangled state, consider its orthogonal complement and try to find another disentangled state in it. We will show that such a procedure can be repeated for at least $\exp[S_{BH}(1 - o(1))]$ times, which leads to $\exp[S_{BH}(1 - o(1))]$ orthogonal disentangled states. The preliminaries for entanglement measures and holography used in the proof will be reviewed in section 2, and the main statements will be presented as propositions and proven in section 3.

In section 4, we apply the techniques developed in section 3 to an alternative situation. We consider an alternative type of atypical states, the area law states, at the standard thermodynamic limit² $V \rightarrow \infty$, where V is the system volume. Let us denote the system length as L so that we have $V = O(L^d)$ where d is the spatial dimension of the system. We will see that for any quantum many-body system, if the reflected entropy [33] of its microcanonical ensemble follows an area law, i.e. scales as $O(L^{d-1})$, then one can find at least $\exp[S(1 - o(1))]$ orthogonal area law states in $\mathcal{H}_{[E-\Delta E, E]}$. They can account for the thermodynamic entropy S at the thermodynamic limit $L \rightarrow \infty$. Holographic CFTs³ are such examples whose reflected entropy satisfies the area law [12, 33].

²It is known that the $G_N \rightarrow 0$ limit in AdS/CFT shares many similar features with the standard thermodynamic limit (the large volume V limit) in generic quantum many-body systems. For example, it is known that an $O(V^{1/2})$ correction [29] appears at the phase transition point of the entanglement entropy found in generic quantum many-body systems satisfying the eigenstate thermalization hypothesis [30], and one can find a similar $O(G_N^{-1/2})$ correction in AdS/CFT [31, 32]. However, while being similar, they are two different limits and should be distinguished from each other. In a lattice regularization, one can intuitively regard $G_N \rightarrow 0$ as the limit where the degrees of freedom on each site go to infinite, while the standard thermodynamic limit corresponds to the limit where the number of lattice itself goes to infinity.

³In holographic CFTs, the thermodynamic entropy S coincides the Bekenstein-Hawking entropy S_{BH} . However, we would like to note that, when we talk about $\exp[S_{BH}(1 - o(1))]$ disentangled states, we are considering the $G_N \rightarrow 0$ limit with the volume V fixed. On the other hand, when we talk about $\exp[S(1 - o(1))]$ area law states, we are considering the $V \rightarrow \infty$ limit (with the Newton constant G_N fixed if for holographic CFTs). While sharing similar features, these two limits, as well as the Landau symbols associated with them, should not be confused with each other.

In section 5, we present an explicit construction of $\exp[S(1 - o(1))]$ area law states. This is one of the crucial points in our paper: not only do we prove the existence, we even present an explicit construction! Moreover, we also discuss the possibility to apply the same method for constructing disentangled states.

In section 6, we summarize our results, discuss their possible relations with additivity conjectures and atypical microstate counting, and show some future directions.

2 Preliminaries

As we have introduced in the previous section, upper bounding the entanglement of formation of microcanonical ensembles in holographic CFTs will be an important piece of our jigsaw puzzle. In this section, we review several preliminaries for this purpose. We will firstly review entanglement of formation and other relevant correlation measures in general quantum systems in section 2.1. Then we will review entanglement of purification in holography and gravity duals of microcanonical ensembles in section 2.2 and 2.3 respectively. Readers who are familiar with these topics can skip this section.

2.1 Entanglement of formation and other correlation measures

Consider a bipartite system composed by A and B , and denote the Hilbert space associated with it as $\mathcal{H}_A \otimes \mathcal{H}_B$. For a (possibly mixed) state $\rho \in \mathcal{D}(\mathcal{H}_A \otimes \mathcal{H}_B)$, there are many quantities to measure the entanglement and correlation between A and B .

Since entanglement is an important resource which classical systems does not have, one may want to construct a function which counts only the amount of entanglement but not any classical correlations between A and B . Such a function is called an entanglement measure. More precisely, an entanglement measure must satisfy some axioms [34–37], which guarantees that it can quantitatively measure the amount of entanglement and does not contain any classical correlations, which can be generated via LOCC operations. See appendix A for a set of axioms required for an entanglement measure. On the other hand, functions which do not fully satisfy all the axioms are often called correlation measures. In the following, let us list some entanglement measures and correlation measures which will be used in this paper.

Entanglement entropy

When ρ is a pure state, the entanglement entropy is defined as

$$E_{vN}(\rho) = S(\rho_A) = S(\rho_B), \quad (2.1)$$

where

$$\rho_{A(B)} \equiv \text{Tr}_{B(A)}(\rho), \quad (2.2)$$

is the reduced density matrix on $\mathcal{H}_{A(B)}$ and

$$S(\sigma) = -\text{Tr}(\sigma \log \sigma) \quad (2.3)$$

is the von Neumann entropy. The reduced von Neumann entropy on pure states is the only function which satisfy all the axioms for entanglement measures [34]. Therefore, the entanglement entropy is an entanglement measure when restricted to pure states.

On the other hand, constructing a computable entanglement measure for a mixed state which has a good operational meaning is much more challenging.

Entanglement of formation

One way to define a correlation measure for a mixed state ρ is to regard ρ as an ensemble of pure states

$$\rho = \sum_i p_i |\psi_i\rangle\langle\psi_i|, \quad (2.4)$$

and then consider the average entanglement entropy over this ensemble:

$$\sum_i p_i E_{vN}(|\psi_i\rangle\langle\psi_i|). \quad (2.5)$$

The entanglement of formation [28] of ρ is defined as the minimal one of such average entanglement entropies over all variations of possible ensemble decomposition:

$$E_F(\rho) \equiv \inf \left\{ \sum_i p_i E_{vN}(|\psi_i\rangle\langle\psi_i|) : \rho = \sum_i p_i |\psi_i\rangle\langle\psi_i| \right\}. \quad (2.6)$$

The entanglement of formation is not an entanglement measure, and it is usually not computable since the definition includes a minimization. However, it has a good operational meaning related to quantum measurements [28] which motivates people to consider it.

In this paper, the scaling behavior of E_F in holographic CFTs and quantum many-body systems will play a central role. Therefore, we will review two correlation measures which gives rigorous upper bounds to $E_F(\rho)$.

Entanglement of purification

Another idea to measure the correlation between A and B is to first uplift it into a pure state and then consider entanglement entropy on that pure state. Such an uplifting procedure is called purification. More precisely, for an arbitrary ancilla C and a pure state $|\psi_{ABC}\rangle$ in $\mathcal{H}_A \otimes \mathcal{H}_B \otimes \mathcal{H}_C$, if it reduces to $\rho \in \mathcal{D}(\mathcal{H}_A \otimes \mathcal{H}_B)$ after tracing out C , i.e.

$$\rho = \text{Tr}_C |\psi_{ABC}\rangle\langle\psi_{ABC}|, \quad (2.7)$$

then $|\psi_{ABC}\rangle$ is called a purification of ρ . By artificially dividing \mathcal{H}_C into two parts

$$\mathcal{H}_C = \mathcal{H}_{A'} \otimes \mathcal{H}_{B'}, \quad (2.8)$$

and assigning $A'(B')$ to $A(B)$, we can get $A \cup A'$ ($B \cup B'$) as an extension of $A(B)$. It is natural to expect that the entanglement entropy between $A \cup A'$ and $B \cup B'$, i.e. $S(\text{Tr}_{BB'} |\psi_{ABC}\rangle\langle\psi_{ABC}|)$ can measure the correlation between A and B .

Obviously, different choices of $|\psi_{ABC}\rangle$ and different ways of factorizing \mathcal{H}_C give different results. Among all these choices, the minimal entanglement entropy evaluated in this way is called the entanglement of purification [27],

$$E_P(\rho) \equiv \inf \left\{ S(\text{Tr}_{BB'} |\psi_{ABC}\rangle\langle\psi_{ABC}|) : \rho = \text{Tr}_C |\psi_{ABC}\rangle\langle\psi_{ABC}| \text{ and } C = A' \cup B' \right\}. \quad (2.9)$$

$E_P(\rho)$ gives a rigorous upper bound of entanglement of formation $E_F(\rho)$ as [27]

$$E_F(\rho) \leq E_P(\rho). \quad (2.10)$$

On the other hand, while $E_P(\rho)$ is not computable in general due to the minimization, it can be evaluated in a class of states in holographic CFTs [25, 26]. We will apply these two properties when discussing black hole microstates later.

Reflected entropy

Since the entanglement of purification is not computable in general, one may want to consider a specific way of purification with which one can get a computable quantity. For example, let us consider a mixed state $\rho \in \mathcal{D}(\mathcal{H}_A \otimes \mathcal{H}_B)$ and diagonalize it as

$$\rho = \sum_{\nu} p_{\nu} |\varphi_{\nu}\rangle\langle\varphi_{\nu}|. \quad (2.11)$$

For each $|\varphi_\nu\rangle \in \mathcal{H}_A \otimes \mathcal{H}_B$, we can consider its CPT conjugate and denote it as $|\varphi_\nu^*\rangle \in \mathcal{H}_{A^*} \otimes \mathcal{H}_{B^*}$. Then, obviously, the state

$$|\sqrt{\rho}\rangle = \sum_\nu \sqrt{p_\nu} |\varphi_\nu\rangle |\varphi_\nu^*\rangle \quad (2.12)$$

defined on $\mathcal{H}_A \otimes \mathcal{H}_B \otimes \mathcal{H}_{A^*} \otimes \mathcal{H}_{B^*}$ serves as a purification of ρ . Such a purification is called the canonical purification [33].

Associated with the canonical purification, the reflected entropy [33] is defined as the entanglement entropy between $A \cup A^*$ and $B \cup B^*$:

$$E_{RE}(\rho) = S(\text{Tr}_{BB^*} |\sqrt{\rho}\rangle \langle \sqrt{\rho}|). \quad (2.13)$$

By definition, it satisfies

$$E_P(\rho) \leq E_{RE}(\rho). \quad (2.14)$$

Therefore, combining with (2.10), we have

$$E_F(\rho) \leq E_P(\rho) \leq E_{RE}(\rho). \quad (2.15)$$

The best advantage of $E_{RE}(\rho)$ is that it is usually easy to compute. We will apply these features to discuss generic quantum many-body systems in the thermodynamic limit.

2.2 Entanglement of purification in AdS/CFT

While the entanglement of purification is in general very hard to compute, it can be easily evaluated for geometric states in holographic CFTs. Here, a holographic CFT is a CFT which admits a semiclassical gravity dual via the AdS/CFT correspondence [14]. However, not all the states in \mathcal{H}_{CFT} necessarily admit a geometrical dual⁴. A state which admits a geometrical dual is called a geometric state.

For geometric states, the entanglement entropy and the entanglement of purification correspond to geometric objects in the AdS gravity. Consider a static geometric state ρ in the AdS _{$d+2$} /CFT _{$d+1$} correspondence and divide the whole d -dimensional spatial region of the

⁴For example, it is believed a superposition of a geometric state corresponding to the vacuum AdS and another state corresponding to an AdS black hole does not admit a geometrical dual.

CFT into A and B . The von Neumann entropy of the reduced entropy on A is given by the Ryu-Takayanagi formula [15, 16]

$$S(\text{Tr}_B(\rho)) = \min_{\Gamma_A} \frac{\text{Area}(\Gamma_A)}{4G_N}, \quad (2.16)$$

where Γ_A is a d -dimensional surface which is homologous to A . The Γ_A with the minimal area is called the RT surface. See the left of figure 2 for a sketch.




Figure 2: A time slice of a global AdS_3 . The asymptotic boundary where the dual CFT lives is shown in gray. The spatial region of the CFT is divided into A and B where A is the two segments on the left and right, and B is the two segments on the top and the bottom. (Left) The orange surface $\Gamma_A^{(\min)}$ shows the Ryu-Takayanagi surface. (Right) The entanglement wedge of $A = A_1 \cup A_2$ is shaded in yellow. The surface $\Sigma_{A_1 A_2}^{(\min)}$ which gives the entanglement wedge cross section $E_W(\rho_A, A_1 : A_2)$ is shown in blue.

The region surrounded by A and $\Gamma_A^{(\min)}$ is called the entanglement wedge⁵ [18, 38, 39] of A and it is considered to be the gravity dual of the reduced density matrix

$$\rho_A \equiv \text{Tr}_B(\rho). \quad (2.17)$$

If we divide A into $A = A_1 \cup A_2$, we can consider a quantity called the entanglement wedge cross section⁶

$$E_W(\rho_A, A_1 : A_2) \equiv \min_{\Sigma_{A_1 A_2}} \frac{\text{Area}(\Sigma_{A_1 A_2})}{4G_N} \quad (2.18)$$

⁵More precisely, what we are considering here is the time-reflection symmetric time slice of the entanglement wedge.

⁶In the following, we will sometimes explicitly show the bipartition if there might be confusion.

where $\Sigma_{A_1 A_2}$ is a d -dimensional surface which divide the entanglement wedge into two parts with A_1 and A_2 living in each part.

It is conjectured in [25, 26] that the entanglement purification between A_1 and A_2 of state ρ_A equals the entanglement wedge cross section $E_W(\rho_A, A_1 : A_2)$:

$$E_P(\rho_A, A_1 : A_2) = E_W(\rho_A, A_1 : A_2). \quad (2.19)$$

This conjecture can be justified by a two-step argument. According to the surface/state correspondence [40] or the tensor network description of holography [41–43], the entanglement wedge with some ancilla degrees of freedom put on the RT surface can be regarded as an “optimal geometric purification”, and hence finding the minimal $\Gamma_{A_1 A_2}$ dividing the RT surface Γ_A into two parts corresponds to finding the optimal way to divide the ancilla in this purification. This first step suggests that E_W at least gives an upper bound of E_P , i.e. $E_P \leq E_W$. As the second step, one can geometrically prove E_W satisfy all the known inequalities satisfied by E_P [25, 26]. Therefore, it is plausible to believe that $E_P = E_W$. However, in fact, only the first step $E_P \leq E_W$ is sufficient for the arguments in this paper.

2.3 Holographic dual of the microcanonical ensemble

As we have explained above, not all the states in a holographic CFT are geometric states. Since the standard AdS/CFT dictionary [44, 45] is an equivalence between the partition functions of the gravity path integral and the CFT path integral, when a state can be easily prepared via a path integral, then one can use the semiclassical approximation of the gravity path integral, by picking up the configuration which minimizes the gravitational action I_{grav} .

Geometric states constructed in this way include the ground state, canonical ensembles

$$\rho^{(\text{can})}(\beta) \propto \sum_i e^{-\beta E_i} |E_i\rangle \langle E_i|, \quad (2.20)$$

thermofield double states (i.e. canonical purifications of canonical ensembles)

$$|\text{TFD}(\beta)\rangle \propto \sum_i e^{-\beta E_i/2} |E_i\rangle |E_i^*\rangle, \quad (2.21)$$

and so on. Here, E_i is the i -th eigenvalue of the Hamiltonian H and $|E_i\rangle$ is the corresponding eigenstate. These three examples correspond to global AdS, one-sided AdS black holes, and two sided AdS black holes [46] respectively.

In this paper, we would like to focus on microcanonical ensembles

$$\rho^{(\text{micro})}(E) \propto \sum_{i \text{ s.t. } E_i \in [E-\Delta E, E]} |E_i\rangle\langle E_i|, \quad (2.22)$$

which is more complicated to prepare with a standard path integral. Fortunately, the gravity dual of the canonical purifications of (2.22), named microcanonical thermofield double states,

$$|\text{MCTFD}(E)\rangle \propto \sum_{i \text{ s.t. } E_i \in [E-\Delta E, E]} |E_i\rangle|E_i^*\rangle \quad (2.23)$$

has been already discussed in [47]⁷. This is again given by a two-sided black hole in the semiclassical limit $G_N \rightarrow 0$. Accordingly, the gravity dual of a microcanonical ensemble is given by a one-sided black hole.

The difference between the gravity duals of a canonical ensemble and a microcanonical ensemble is that, while we need to find the classical geometry which minimize the gravitational action I_{grav} on the AdS side when using a standard path integral to prepare a canonical ensemble or a thermofield double state, we need to find the one which maximize another functional, which coincides the black hole entropy for black hole geometries, when considering the microcanonical counterparts [47].

The results explained above can be naturally understood from the view point of the relation between statistical mechanics and thermodynamics. In statistical mechanics, different ensembles including the canonical ensemble and the microcanonical ensemble are equivalent to each other in reproducing the macroscopic quantities at the limit of large degrees of freedom (i.e. the thermodynamic limit). In the current case, the semiclassical limit $1/G_N \rightarrow \infty$, where G_N is the Newton constant, plays a similar role of the thermodynamic limit. At the thermodynamic limit, the natural thermodynamic potential associated with the canonical ensemble is the Helmholtz free energy, which holds temperature $1/\beta$ as a natural variable⁸.

⁷In fact, the exact setup analyzed in [47] is $|\psi\rangle \propto \sum_{E'} e^{-\beta E'} f(E' - E) |E'\rangle |E'^*\rangle$ where $f(E' - E) = \exp(-(E' - E)^2/4\sigma^2)$. However, as commented in [47], the results hold for any $f(E' - E)$ concentrated around $E' = E$.

⁸Here, “a thermodynamic potential A holds x, y, \dots as natural variables” means that all the thermodynamic quantities can be determined from A as a function of x, y, \dots . For example, when expressing the Helmholtz free energy F as a function of the temperature $1/\beta$ and the volume V , all the thermodynamic quantities can be determined from partial derivatives of $F[1/\beta, V]$. However, when expressing F as a function of the energy E and the volume V , some information is lost in $F(E, V)$. A thermodynamic function expressed by its natural variables is called a complete thermodynamic function in some textbooks. Complete thermodynamic functions are equivalent to each other under Legendre transformations.

On the other hand, the natural thermodynamic potential associated with the microcanonical ensemble is the thermodynamic entropy, which holds energy E as the natural variable. In the current case, the gravitational action I_{grav} parameterized by β corresponds to the Helmholtz free energy, and the functional which coincides the black hole entropy in black hole geometries [47] parameterized by E corresponds to the thermodynamic entropy. The thermodynamic equilibrium is realized when the Helmholtz free energy is minimized (or equivalently, when the thermodynamic entropy is maximized).

Moreover, it is also shown in [47] that the Ryu-Takayanagi formula [15,16] and its covariant extension, the Hubeny-Rangamani-Takayanagi formula [17] still hold in the microcanonical counterparts. Therefore, it is plausible that we can also apply the $E_P = E_W$ conjecture (or at least $E_P \leq E_W$) for microcanonical ensembles.

3 Disentangled black hole microstates in AdS/CFT

In this section, we show that, for a holographic CFT, there exists an orthogonal basis of the microcanonical subspace $\mathcal{H}_{[E-\Delta E, E]}$ such that at least $\exp[S_{BH}(1 - o(1))]$ elements are disentangled states, where S_{BH} is the Bekenstein-Hawking entropy of the corresponding black hole on the AdS side.

We will firstly give an upper bound to the entanglement of formation of the microcanonical ensemble $\rho^{(\text{micro})}$ using holography in section 3.1. This bound indicates that there is at least one disentangled state in $\mathcal{H}_{[E-\Delta E, E]}$. Then we can exclude this state and consider its orthogonal complement (and a new microcanonical density matrix associated to it). It turns out the entanglement of formation of the new microcanonical ensemble can also be upper bounded. We will show some results which are crucial in this argument in section 3.2 merely from linear algebra. This means results in section 3.2 are universal for any quantum systems. According as the new upper bound behaves, we may repeat the above procedure again and again. We will finally show this procedure can be repeated $\exp[S_{BH}(1 - o(1))]$ times in section 3.3.

3.1 Upper bounding E_F from holography

As the first step, we will use the preliminaries reviewed in section 2 to give an upper bound to E_F of $\rho^{(\text{micro})}$. We consider dividing the spatial region of the CFT into two parts A and




Figure 3: AdS black hole as the gravity dual of the microcanonical density matrix $\rho^{(\text{micro})}$. (Left) The orange surface, which coincides the black hole horizon shows the Ryu-Takayanagi surface of the whole system. (Right) The entanglement wedge of the whole system is the outside region of the black hole shown in yellow. Dividing the spatial region of the CFT into A and B , the blue surface $\Sigma_{AB}^{(\text{min})}$ gives the entanglement wedge cross section $E_W(\rho^{(\text{micro})}, A : B)$.

B , where B is slightly smaller than A . See figure 1 for a sketch.

As we have reviewed in section 2.3, the gravity dual of $\rho^{(\text{micro})}$ is a black hole geometry. The entanglement wedge of the whole system $A \cup B$ is the exterior of the black hole, as shown in figure 3. Accordingly, the entanglement wedge cross section E_W is given by the area of the minimal surface connecting the interface $\partial A = \partial B$ and the black hole horizon. Let us denote this minimal surface as $\Sigma_{AB}^{(\text{min})}$. Combining the holographic formula for entanglement of purification (2.19) and the general relation between entanglement of formation and entanglement of purification (2.10), we have

$$E_F \leq E_P = E_W = \frac{\text{Area}(\Sigma_{AB}^{(\text{min})})}{4G_N} + O(1). \quad (3.1)$$

Therefore, by definition of E_F , it is straightforward to say that there exists an ensemble decomposition of $\rho^{(\text{micro})}$ such that the averaged entanglement entropy is less than or equal to $\text{Area}(\Sigma_{AB}^{(\text{min})})/4G_N$ at leading order. Let us summarize this as a proposition for later reference.

Proposition 1. *In a holographic CFT, let $\rho^{(\text{micro})}$ be a microcanonical ensemble and divide the spatial region into A and B where B is slightly smaller than A . Then there exists a classical ensemble of states $\{(p_i, |\psi_i\rangle)\}_{i=1,2,\dots}$ such that*

$$\rho^{(\text{micro})} = \sum_i p_i |\psi_i\rangle\langle\psi_i|, \quad (3.2)$$

and the averaged entanglement entropy

$$\overline{E_{vN}(\{(p_i, |\psi_i\rangle)\}_{i=1,2,\dots})} = \sum_i p_i E_{vN}(|\psi_i\rangle\langle\psi_i|) \lesssim \frac{\text{Area}(\Sigma_{AB}^{(\text{min})})}{4G_N}, \quad (3.3)$$

where “ \lesssim ” means “less than or equal to at leading order”.

Noting that $\text{Area}(\Sigma_{AB}^{(\text{min})})/4G_N$ and $\frac{1}{2}S(\text{Tr}_{AA^*}|\text{TFD}\rangle\langle\text{TFD}|)$ matches at the leading order of $1/G_N$, according to the definition of disentangled states (1.2), it straightforwardly follows from the positivity of probability and von Neumann entropy that at least one element $|\psi_j\rangle$ with $p_j \neq 0$ in $\{|\psi_i\rangle\}_{i=1,2,\dots}$ is a disentangled state, i.e. it satisfies

$$E_{vN}(|\psi_j\rangle\langle\psi_j|) \lesssim \frac{\text{Area}(\Sigma_{AB}^{(\text{min})})}{4G_N} \approx \frac{1}{2}S(\text{Tr}_{AA^*}|\text{TFD}\rangle\langle\text{TFD}|). \quad (3.4)$$

However, to this point, it is nontrivial whether this $|\psi_j\rangle$ is in the microcanonical subspace or not.

3.2 General results for microcanonical ensembles

In this subsection, we show some general properties of microcanonical density matrices from linear algebra, and comment on their applications on the system we are considering. Note that the results presented in this subsection are general, and no properties of specific physical systems will be used to show them.

Proposition 2. *Let ρ be a density matrix which can be written as $\rho = \Pi_{\mathcal{E}}/W$, where $\Pi_{\mathcal{E}}$ is the projection operator onto a W -dimensional subspace $\mathcal{E} \cong \mathbb{C}^W$. If an ensemble of states $\{(p_i, |\psi_i\rangle)\}_{i=1,2,\dots}$ generates ρ , i.e.,*

$$\rho = \sum_i p_i |\psi_i\rangle\langle\psi_i|, \quad (3.5)$$

then for every $|\psi_j\rangle \in \{|\psi_i\rangle\}_{i=1,2,\dots}$, $|\psi_j\rangle \in \mathcal{E}$ if $p_j \neq 0$.

Proof. Note that

$$p_j \|(1 - \Pi_{\mathcal{E}})|\psi_j\rangle\|^2 = p_j |\langle \psi_j | (1 - \Pi_{\mathcal{E}}) |\psi_j\rangle|^2 \leq \sum_i p_i |\langle \psi_i | (1 - \Pi_{\mathcal{E}}) |\psi_j\rangle|^2. \quad (3.6)$$

Therefore, using (3.5), we have

$$p_j \|(1 - \Pi_{\mathcal{E}})|\psi_j\rangle\|^2 \leq \langle \psi_j | (1 - \Pi_{\mathcal{E}}) \rho (1 - \Pi_{\mathcal{E}}) |\psi_j\rangle = \langle \psi_j | (1 - \Pi_{\mathcal{E}}) \frac{\Pi_{\mathcal{E}}}{W} (1 - \Pi_{\mathcal{E}}) |\psi_j\rangle = 0. \quad (3.7)$$

This implies that $|\psi_j\rangle = \Pi_{\mathcal{E}}|\psi_j\rangle \in \mathcal{E}$. \square

Based on proposition 2, we can say at least one $|\psi_j\rangle$ appearing in proposition 1 is a disentangled state in $\mathcal{H}_{[E-\Delta E, E]}$. Therefore, we can consider excluding $|\psi_j\rangle$ from $\mathcal{H}_{[E-\Delta E, E]}$. The next step is to consider if we can pick up another disentangled state in the orthogonal complement of $|\psi_j\rangle$ with respect to $\mathcal{H}_{[E-\Delta E, E]}$. To answer this question, we will need to prove the following theorem.

Theorem 3. *Let ρ be a density matrix which can be written as $\rho = \Pi_{\mathcal{E}}/W$, where $\Pi_{\mathcal{E}}$ is the projection operator onto a W -dimensional subspace $\mathcal{E} \cong \mathbb{C}^W$. Let $\{(p_i, |\psi_i\rangle)\}_{i=1,2,\dots}$ be an ensemble of states which generates ρ with the average entanglement entropy*

$$\overline{E_{vN}} = \sum_i p_i E_{vN}(|\psi_i\rangle\langle\psi_i|). \quad (3.8)$$

Pick up an arbitrary set of k orthogonal states $\{|\varphi_1\rangle, |\varphi_2\rangle, \dots, |\varphi_k\rangle\} \subset \mathcal{E}$ and consider its orthogonal complement

$$\mathcal{E}' = \mathcal{E} \cap (\text{span}\{|\varphi_1\rangle, |\varphi_2\rangle, \dots, |\varphi_k\rangle\})^\perp. \quad (3.9)$$

Let $\Pi'_{\mathcal{E}}$ denote the projection operator onto \mathcal{E}' . Then there exists an ensemble of states $\{(p'_i, |\psi'_i\rangle)\}_{i=1,2,\dots}$ generating the density matrix $\rho' \propto \Pi'_{\mathcal{E}}$ whose average entanglement entropy

$$\overline{E'_{vN}} = \sum_i p'_i E_{vN}(|\psi'_i\rangle\langle\psi'_i|). \quad (3.10)$$

can be bounded from two sides as

$$\overline{E'_{vN}} < \frac{1}{1 - k/W} \left[\overline{E_{vN}} + (k/W)^{1/3} \left\{ \frac{1}{3} \log W/k + 5 \log \dim \mathcal{H}_B \right\} + k/W \log \dim \mathcal{H}_B \right], \quad (3.11)$$

$$\overline{E'_{vN}} > \frac{1}{1 - k/W} \left[(1 - 2k/W) \overline{E_{vN}} - (k/W)^{1/3} \left\{ \frac{1}{3} \log W/k + 5 \log \dim \mathcal{H}_B \right\} - k/W \log \dim \mathcal{H}_B \right]. \quad (3.12)$$

Proof. Let us show the theorem in a constructive way. It is crucial to note that, since $|\varphi_i\rangle \in \mathcal{E}$,

$$\sum_i p_i \langle \psi_i | P | \psi_i \rangle = \text{Tr} \rho P = \text{Tr} \left[\frac{\Pi_{\mathcal{E}}}{W} \sum_{i=1}^k |\varphi_i\rangle \langle \varphi_i| \right] = k/W. \quad (3.13)$$

Here P is the projection operator onto the subspace spanned by the excluded states in (3.9). This implies $\langle \psi_i | P | \psi_i \rangle$ is small with a large probability when W is large, which motivates us to take

$$p'_i = \frac{1 - \langle \psi_i | P | \psi_i \rangle}{1 - k/W} p_i, \quad (3.14)$$

$$|\psi'_i\rangle = \frac{(1 - P)|\psi_i\rangle}{\sqrt{1 - \langle \psi_i | P | \psi_i \rangle}}. \quad (3.15)$$

It can be easily checked that this $\{(p'_i, |\psi'_i\rangle)\}$ generates ρ' as follows:

$$\begin{aligned} \Pi_{\mathcal{E}'} &= \Pi_{\mathcal{E}} - P = (\Pi_{\mathcal{E}} - P)\Pi_{\mathcal{E}}(\Pi_{\mathcal{E}} - P) \\ &= (\Pi_{\mathcal{E}} - P) \left[W \sum_i p_i |\psi_i\rangle \langle \psi_i| \right] (\Pi_{\mathcal{E}} - P) = (W - k) \sum_i p'_i |\psi'_i\rangle \langle \psi'_i|. \end{aligned} \quad (3.16)$$

Then we proceed to consider how the average entanglement entropy changes when moving from $\{(p_i, |\psi_i\rangle)\}$ to $\{(p'_i, |\psi'_i\rangle)\}$. Using the triangle inequality, we have

$$\begin{aligned} |\overline{E_{vN}'} - \overline{E_{vN}}| &= \left| \sum_i p'_i E_{vN}(|\psi'_i\rangle \langle \psi'_i|) - \sum_i p_i E_{vN}(|\psi_i\rangle \langle \psi_i|) \right| \\ &= \left| \sum_i (p'_i - p_i) E_{vN}(|\psi_i\rangle \langle \psi_i|) + \sum_i p'_i (E_{vN}(|\psi'_i\rangle \langle \psi'_i|) - E_{vN}(|\psi_i\rangle \langle \psi_i|)) \right| \\ &\leq \underbrace{\sum_i |p'_i - p_i| E_{vN}(|\psi_i\rangle \langle \psi_i|)}_{=(*1)} + \underbrace{\sum_i p'_i |E_{vN}(|\psi'_i\rangle \langle \psi'_i|) - E_{vN}(|\psi_i\rangle \langle \psi_i|)|}_{=(*2)}. \end{aligned} \quad (3.17)$$

First we give an upper bound for (*1). By definition, we have

$$p'_i - p_i = \frac{k/W}{1 - k/W} p_i - \frac{1}{1 - k/W} p_i \langle \psi_i | P | \psi_i \rangle. \quad (3.18)$$

Substituting this into (*1), we obtain

$$\begin{aligned} (*1) &\leq \frac{k/W}{1 - k/W} \sum_i p_i E_{vN}(|\psi_i\rangle \langle \psi_i|) + \frac{1}{1 - k/W} \sum_i p_i \langle \psi_i | P | \psi_i \rangle E_{vN}(|\psi_i\rangle \langle \psi_i|) \\ &\leq \frac{k/W}{1 - k/W} \sum_i p_i E_{vN}(|\psi_i\rangle \langle \psi_i|) + \frac{1}{1 - k/W} \log \dim \mathcal{H}_B \sum_i p_i \langle \psi_i | P | \psi_i \rangle \\ &= \frac{k/W}{1 - k/W} \{ \overline{E_{vN}} + \log \dim \mathcal{H}_B \}. \end{aligned} \quad (3.19)$$

Here we have used Eqs. (3.8) and (3.13).

As the next step, we consider bounding (*2) from above. It would be useful to note the following facts:

- Since for any state $|\phi\rangle \in \mathcal{H}$, $0 \leq E_{v_N}(|\phi\rangle\langle\phi|) \leq \log \dim \mathcal{H}_B$, we have

$$|E_{v_N}(|\psi'_i\rangle\langle\psi'_i|) - E_{v_N}(|\psi_i\rangle\langle\psi_i|)| \leq \log \dim \mathcal{H}_B. \quad (3.20)$$

- Using Eq. (3.15), we have

$$\|\text{Tr}_A |\psi'_i\rangle\langle\psi'_i| - \text{Tr}_A |\psi_i\rangle\langle\psi_i|\|_1 \leq \| |\psi'_i\rangle\langle\psi'_i| - |\psi_i\rangle\langle\psi_i| \|_1 = 2\sqrt{1 - |\langle\psi'_i|\psi_i\rangle|^2} = \sqrt{4\langle\psi_i|P|\psi_i\rangle}. \quad (3.21)$$

Therefore, by applying Fannes' inequality, we find

$$|E_{v_N}(|\psi'_i\rangle\langle\psi'_i|) - E_{v_N}(|\psi_i\rangle\langle\psi_i|)| \leq \sqrt{4\langle\psi_i|P|\psi_i\rangle} \log \dim \mathcal{H}_B - \sqrt{4\langle\psi_i|P|\psi_i\rangle} \log \sqrt{4\langle\psi_i|P|\psi_i\rangle}. \quad (3.22)$$

This gives a better bound compared to (3.20) when $\langle\psi_i|P|\psi_i\rangle$ is sufficiently small.

- For any $\epsilon > 0$, one finds

$$\sum_{\substack{i \text{ s.t.} \\ 4\langle\psi_i|P|\psi_i\rangle > W^{-2\epsilon}}} p_i < 4kW^{-1+2\epsilon}. \quad (3.23)$$

This can be seen as follows.

$$4^{-1}W^{-2\epsilon} \sum_{\substack{i \text{ s.t.} \\ 4\langle\psi_i|P|\psi_i\rangle > W^{-2\epsilon}}} p_i < \sum_{\substack{i \text{ s.t.} \\ 4\langle\psi_i|P|\psi_i\rangle > W^{-2\epsilon}}} p_i \langle\psi_i|P|\psi_i\rangle \leq \sum_i p_i \langle\psi_i|P|\psi_i\rangle = k/W, \quad (3.24)$$

where we used (3.13).

Based on these, it is straightforward to see

$$(*2) \leq \frac{1}{1 - k/W} \sum_i p_i |E_{vN}(|\psi'_i\rangle\langle\psi'_i|) - E_{vN}(|\psi_i\rangle\langle\psi_i|)| \quad (3.25)$$

$$\begin{aligned} &= \frac{1}{1 - k/W} \sum_{\substack{i \text{ s.t.} \\ 4\langle\psi_i|P|\psi_i\rangle > W^{-2\epsilon}}} p_i |E_{vN}(|\psi'_i\rangle\langle\psi'_i|) - E_{vN}(|\psi_i\rangle\langle\psi_i|)| \\ &+ \frac{1}{1 - k/W} \sum_{\substack{i \text{ s.t.} \\ 4\langle\psi_i|P|\psi_i\rangle \leq W^{-2\epsilon}}} p_i |E_{vN}(|\psi'_i\rangle\langle\psi'_i|) - E_{vN}(|\psi_i\rangle\langle\psi_i|)| \end{aligned} \quad (3.26)$$

$$\begin{aligned} &\leq \frac{1}{1 - k/W} \log \dim \mathcal{H}_B \sum_{\substack{i \text{ s.t.} \\ 4\langle\psi_i|P|\psi_i\rangle > W^{-2\epsilon}}} p_i \\ &+ \frac{1}{1 - k/W} \{W^{-\epsilon} \log \dim \mathcal{H}_B + \epsilon W^{-\epsilon} \log W\} \sum_{\substack{i \text{ s.t.} \\ 4\langle\psi_i|P|\psi_i\rangle \leq W^{-2\epsilon}}} p_i \end{aligned} \quad (3.27)$$

$$< \frac{1}{1 - k/W} \{4kW^{-1+2\epsilon} \log \dim \mathcal{H}_B + W^{-\epsilon} \log \dim \mathcal{H}_B + \epsilon W^{-\epsilon} \log W\}, \quad (3.28)$$

where we used (3.14) in the first line and (3.23) in the last line. By taking

$$\epsilon = \frac{1}{3} \left(1 - \frac{\log k}{\log W}\right), \quad (3.29)$$

we get

$$(*2) < \frac{(k/W)^{1/3}}{1 - k/W} \left\{ \frac{1}{3} \log W/k + 5 \log \dim \mathcal{H}_B \right\}. \quad (3.30)$$

Combining (*1) and (*2), we obtain

$$|\overline{E_{vN}'} - \overline{E_{vN}}| < \frac{(k/W)^{1/3}}{1 - k/W} \left\{ \frac{1}{3} \log W/k + 5 \log \dim \mathcal{H}_B \right\} + \frac{k/W}{1 - k/W} \{\overline{E_{vN}} + \log \dim \mathcal{H}_B\}. \quad (3.31)$$

This gives both the upper bound and the lower bound shown in the theorem. \square

3.3 Existence of $e^{S_{BH}}$ disentangled microstates

In this subsection, we combine the results presented in the previous subsections to get our final claim. As the first step, let us consider the microcanonical ensemble $\rho^{(\text{micro})}$ as ρ , the associated microcanonical subspace

$$\mathcal{H}_{[E-\Delta E, E]} = \text{span} \{ |E_i\rangle \in \mathcal{H} | E_i \in [E - \Delta E, E] \}, \quad (3.32)$$

as \mathcal{E} , and a disentangled state $|\psi_j\rangle$ appearing in an ensemble of states in proposition 1 as $|\varphi_1\rangle$ (which is guaranteed by proposition 2) in theorem 3. Let us consider the $k = 1$ case of theorem 3. Then $W \sim e^{S_{BH}} = e^{A_{\text{horizon}}/4G_N}$, where A_{horizon} is the horizon area of the AdS black hole corresponding to $\rho^{(\text{micro})}$. In this case, since $\overline{E_{vN}} = O(1/G_N) = O(S_{BH})$ and $\log \dim \mathcal{H}_B = O(1/G_N) = O(S_{BH})$, according to the upper bound (3.11) given in theorem 3, $\rho' \propto \Pi_{\mathcal{E}'}$ can be generated by an ensemble of states whose average entanglement entropy is changed by at most $O(e^{-S_{BH}/3} S_{BH})$ from $\overline{E_{vN}}$.

Then we can take another disentangled state $|\psi'_{j'}\rangle$ from the ensemble that generates ρ' obtained in the first step, and regard it as $|\varphi_2\rangle$. Applying the $k = 2$ case of theorem 3, we can find an ensemble of states generating the microcanonical ensemble associated with

$$\mathcal{H}_{[E-\Delta E, E]} \cap (\text{span}\{|\varphi_1\rangle, |\varphi_2\rangle\})^\perp, \quad (3.33)$$

with average entanglement entropy changed by at most $O(e^{-S_{BH}/3} S_{BH})$ from $\overline{E_{vN}}$.

According to theorem 3, one can repeat this procedure for at least $\exp[S_{BH}(1 - o(1))]$ times without changing the average entanglement entropy in the leading order, i.e. $O(1/G_N)$. Therefore one can obtain $\exp[S_{BH}(1 - o(1))]$ disentangled states as

$$|\varphi_1\rangle, |\varphi_2\rangle, |\varphi_3\rangle, \dots \quad (3.34)$$

By construction, these states are orthogonal. To summarize, we obtain the following statement.

Corollary 4. *In a holographic CFT, let $\mathcal{H}_{[E-\Delta E, E]}$ be the microcanonical subspace whose dimension is given by $e^{S_{BH}}$. There exists an orthogonal basis of $\mathcal{H}_{[E-\Delta E, E]}$ such that at least $\exp[S_{BH}(1 - o(1))]$ elements are disentangled states.*

Note that we have not only proven the existence of a basis satisfying requirements mentioned in corollary 4, but also given an algorithm to construct such a basis from an ensemble of states satisfying requirements mentioned in proposition 1 in the proof of theorem 3. We will come back to discuss if one can give an ensemble of states satisfying requirements mentioned in proposition 1 in section 5.3. Besides, we also note that if one only wants to prove the existence of $\exp[S_{BH}(1 - o(1))]$ disentangled states but does not care about how to construct them, there actually exists a more simple proof. We will leave it in appendix B.

4 Area law states in quantum many-body systems

In section 3, we discussed disentangled states in holographic CFT at the semiclassical limit $G_N \rightarrow 0$. In this section, we see that we can get a similar result for generic short-ranged Hamiltonian with $L \gg 1$. Here, L is the linear scale of the system. Accordingly, the volume of the system is $O(L^d)$, where d is the system dimension. Again, let us divide the system into two parts A and B , where B is slightly smaller than A . A typical state in the microcanonical subspace $\mathcal{H}_{[E-\Delta E, E]}$ would be a volume law state, i.e. its entanglement entropy would be $O(L^d)$. Below in this section we show that there exists an orthogonal basis of the microcanonical subspace $\mathcal{H}_{[E-\Delta E, E]}$ such that at least $\exp[S(1 - o(1))]$ elements are area law states, where S is the thermodynamic entropy.

We take a similar (but slightly different) strategy as in section 3. We will firstly make a conjecture⁹ about the entanglement entropy in microcanonical TFD states and give an upper bound to the entanglement of formation of the microcanonical ensemble $\rho^{(\text{micro})}$ under the conjecture. Then we can apply proposition 2 and theorem 3 in a similar way as in section 3.3 to show there exist at least $\exp[S(1 - o(1))]$ orthogonal area law states in $\mathcal{H}_{[E-\Delta E, E]}$.

4.1 Conjecture for E_{RE}

We would like to first give an upper bound to the entanglement of formation of the microcanonical ensemble $\rho^{(\text{micro})}$. Different from the holographic case, now the entanglement of purification is not accessible. However, since we are now looking at the scaling behavior with respect to L , we can give it a looser upper bound using the reflected entropy. The canonical purification of the microcanonical ensemble is simply given by the microcanonical TFD state defined in (2.23). Both the canonical ensemble and microcanonical ensemble are statistical mixtures of energy eigenstates, whose probability distribution has a sharp peak around the energy expectation value in the thermodynamic limit ($L \rightarrow \infty$). Therefore, for sufficiently large L , their canonical purifications are expected to have similar entanglement properties. In the canonical TFD states (2.21), the entanglement entropy is extensively studied for various systems [12, 48]. In particular for $d = 1$, it has been shown that the canonical TFD state obeys an area law by the field-theoretical analysis [48] and rigorously proven for sufficiently high temperature [49]. Besides, this has also been shown in holographic CFTs from gravita-

⁹It is worth noting that holographic CFTs are explicit examples where this conjecture can be shown as we will see later.

tional computations [12]. From our knowledge about TFD states, it is plausible to make the following conjecture.

Conjecture 5. *Consider a d -dimensional quantum many-body system with local interactions, and divide the spatial region into A and B where B is slightly smaller than A . Consider a microcanonical TFD state defined on $\mathcal{H}_A \otimes \mathcal{H}_B \otimes \mathcal{H}_{A^*} \otimes \mathcal{H}_{B^*}$. Then the entanglement entropy between AA^* and BB^* is $O(L^{d-1})$, i.e. it obeys the area law.*

It is worth noting one can show conjecture 5 is true in holographic CFTs. First, according to [47], microcanonical TFD states are also geometric states corresponding to two-sided eternal black holes and the RT formula hold in them. Then one can perform the same computation in [12] to find that the entanglement entropy indeed obeys an area law.

4.2 Existence of e^S area law microstates

Combining this conjecture and the general relation between entanglement of formation and reflected entropy (2.15), we have

$$E_F \leq E_{RE} = O(L^{d-1}). \quad (4.1)$$

Therefore, by definition of E_F , it is straightforward to say that there exists an ensemble of states generating $\rho^{(\text{micro})}$ such that the averaged entanglement entropy is $O(L^{d-1})$. Thus, there must exist at least one area law state $|\varphi_1\rangle$ in the microcanonical subspace according to proposition 2. Then we proceed to consider if we can pick up another disentangled state in the orthogonal complement of $|\varphi_1\rangle$. The discussion here is parallel with that in section 5.3.

As the first step, consider the microcanonical ensemble $\rho^{(\text{micro})}$ as ρ , the associated microcanonical subspace $\mathcal{H}_{[E-\Delta E, E]}$ as \mathcal{E} , and an area law state appearing in an ensemble of states, whose average entanglement entropy is $O(L^{d-1})$, as $|\varphi_1\rangle$ in theorem 3. Again, let us first consider the $k = 1$ case of theorem 3. Then $W \sim e^S = e^{O(L^d)}$. In this case, since $\overline{E_{vN}} = O(L^{d-1}) = O(S^{\frac{d-1}{d}})$ and $\log \dim \mathcal{H}_B = O(L^d) = O(S)$, according to the upper bound (3.11) given in theorem 3, $\rho' \propto \Pi_{\mathcal{E}'}$ can be generated by an ensemble of states whose average entanglement entropy is changed by at most $O(e^{-S/3}S)$ from $\overline{E_{vN}}$.

Then we can take another area law state from the ensemble that generates ρ' obtained in the first step as $|\varphi_2\rangle$. Again, applying the $k = 2$ case of theorem 3, we can find an ensemble of states generating the microcanonical ensemble associated with

$$\mathcal{H}_{[E-\Delta E, E]} \cap (\text{span}\{|\varphi_1\rangle, |\varphi_2\rangle\})^\perp, \quad (4.2)$$

with average entanglement entropy changed by at most $O(e^{-S/3}S)$ from the original $\overline{E_{vN}}$.

Then one can repeat this procedure for at least $\exp[S(1 - o(1))]$ times to get orthogonal area law states in $\mathcal{H}_{[E-\Delta E, E]}$. As a result, we have the following corollary.

Corollary 6. *In a d -dimensional quantum many-body system with local interactions, let $\mathcal{H}_{[E-\Delta E, E]}$ be the microcanonical subspace whose dimension is given by e^S , and divide the spatial region into A and B where B is slightly smaller than A . Assuming conjecture 5, then there exists an orthogonal basis of $\mathcal{H}_{[E-\Delta E, E]}$ such that at least $\exp[S(1 - o(1))]$ elements are area law states.*

Here, we have applied the cumbersome theorem 3 to prove corollary 6. In fact, if combined with another theorem, this proof can even let us construct an explicit example of such $\exp[S(1 - o(1))]$ orthogonal area law states, as we will see in section 5. However, if one merely wants to prove corollary 6 without giving any explicit construction, there actually exists a more straightforward way. We will present such an alternative proof in appendix B.

5 Constructing area law states

In section 4, we have shown that there exists an orthogonal basis of the microcanonical subspace $\mathcal{H}_{[E-\Delta E, E]}$ such that at least $\exp[S(1 - o(1))]$ elements are area law states under a plausible conjecture. In this section, we provide a concrete method for constructing $\exp[S(1 - o(1))]$ area law states.

In section 5.1, we present an explicit example of an ensemble of states generating the microcanonical ensemble whose average entanglement entropy is bounded by the reflected entropy. Therefore, according to conjecture 5, the average entanglement entropy is at most $O(L^{d-1})$. General results got in this section will be summarized as a theorem. Then in section 5.2, we will show that we can construct at least $\exp[S(1 - o(1))]$ orthogonal area law states from the ensemble of states constructed in section 5.1 using the algorithm presented in the proof of theorem 3.

5.1 Constructing an ensemble of states with low entanglement

In this subsection, we construct an ensemble of states $\{(p_i, |\psi_i\rangle)\}_{i=1,2,\dots}$ that generates $\rho^{(\text{micro})}$ and show that the average entanglement entropy associated with this ensemble is bounded by the reflected entropy and hence follows an area law under conjecture 5.

To be concrete, let us consider a discrete system defined on a lattice with L^d sites. For a continuous quantum field theory, we can consider its lattice regularization. Then we have $O(e^{L^d})$ orthogonal product states which span the Hilbert space of the whole system. Let us denote the set of these product states as $\{|P_i\rangle\}_{i=1,2,\dots}$. Since the microcanonical ensemble $\rho^{(\text{micro})}$ can be decomposed as

$$\begin{aligned}
\rho^{(\text{micro})} &= e^{-S} \Pi_{[E-\Delta E, E]} \\
&= e^{-S} \Pi_{[E-\Delta E, E]} \left(\sum_i |P_i\rangle\langle P_i| \right) \Pi_{[E-\Delta E, E]} \\
&= \sum_i (e^{-S/2} \Pi_{[E-\Delta E, E]} |P_i\rangle) (\langle P_i| \Pi_{[E-\Delta E, E]} e^{-S/2}) \\
&= \sum_i e^{-S} \langle P_i| \Pi_{[E-\Delta E, E]} |P_i\rangle \left(\frac{\Pi_{[E-\Delta E, E]} |P_i\rangle}{\sqrt{\langle P_i| \Pi_{[E-\Delta E, E]} |P_i\rangle}} \right) \left(\frac{\langle P_i| \Pi_{[E-\Delta E, E]}}{\sqrt{\langle P_i| \Pi_{[E-\Delta E, E]} |P_i\rangle}} \right), \quad (5.1)
\end{aligned}$$

it seems plausible to expect that

$$p_i = e^{-S} \langle P_i| \Pi_{[E-\Delta E, E]} |P_i\rangle, \quad (5.2)$$

$$|\psi_i\rangle = \frac{\Pi_{[E-\Delta E, E]} |P_i\rangle}{\sqrt{\langle P_i| \Pi_{[E-\Delta E, E]} |P_i\rangle}}, \quad (5.3)$$

gives a small average entanglement. This is indeed the case that the average entanglement entropy in this case can be bounded by the reflected entropy of $\rho^{(\text{micro})}$. In fact, we can show a more general theorem presented below.

Theorem 7. *Let ρ be a density matrix defined on a bipartite system $\mathcal{H} = \mathcal{H}_A \otimes \mathcal{H}_B$. Take any set of orthogonal states $\{|P_i\rangle_{A(B)}\}$ for which $\sum_i |P_i\rangle\langle P_i|_{A(B)}$ gives the unity matrix $I_{A(B)}$ on $\mathcal{H}_{A(B)}$. Then, using product states $|P_{ij}\rangle = |P_i\rangle_A \otimes |P_j\rangle_B$, one can decompose ρ as*

$$\rho = \sum_{i,j} \sqrt{\rho} |P_{ij}\rangle\langle P_{ij}| \sqrt{\rho} = \sum_{i,j} p_{ij} |\psi_{ij}\rangle\langle\psi_{ij}|, \quad (5.4)$$

where

$$p_{ij} = \langle P_{ij}| \rho |P_{ij}\rangle, \quad (5.5)$$

$$|\psi_{ij}\rangle = \frac{\sqrt{\rho} |P_{ij}\rangle}{\sqrt{\langle P_{ij}| \rho |P_{ij}\rangle}}. \quad (5.6)$$

For this ensemble of states, the average entanglement entropy is upper bounded by the reflected entropy of ρ :

$$\overline{E_{vN}} = \sum_{i,j} p_{ij} E_{vN} (|\psi_{ij}\rangle\langle\psi_{ij}|) \leq E_{RE}(\rho). \quad (5.7)$$

Proof. A key observation in our proof is that

$$I_A \otimes I_B \otimes |P_i^*\rangle\langle P_i^*|_{A^*} \otimes |P_j^*\rangle\langle P_j^*|_{B^*} |\sqrt{\rho}\rangle = \sqrt{p_{ij}} |\psi_{ij}\rangle |P_{ij}^*\rangle, \quad (5.8)$$

where $|\sqrt{\rho}\rangle$ is the canonical purification of ρ . Indeed, by definition of the canonical purification (2.12), we have

$$\begin{aligned} |\sqrt{\rho}\rangle &= \sqrt{\rho} \otimes I \sum_{\nu} |\varphi_{\nu}\rangle |\varphi_{\nu}^*\rangle = \sqrt{\rho} \otimes I \sum_{\nu} \left(\sum_{i,j} |P_{ij}\rangle \langle P_{ij}| \right) |\varphi_{\nu}\rangle |\varphi_{\nu}^*\rangle \\ &= \sqrt{\rho} \otimes I \sum_{\nu} \sum_{i,j} \langle P_{ij} | \varphi_{\nu} \rangle |P_{ij}\rangle |\varphi_{\nu}^*\rangle = \sqrt{\rho} \otimes I \sum_{\nu} \sum_{i,j} \langle \varphi_{\nu}^* | P_{ij}^* \rangle |P_{ij}\rangle |\varphi_{\nu}^*\rangle \\ &= \sqrt{\rho} \otimes I \sum_{i,j} |P_{ij}\rangle \left(\sum_{\nu} |\varphi_{\nu}^*\rangle \langle \varphi_{\nu}^*| \right) |P_{ij}^*\rangle = \sqrt{\rho} \otimes I \sum_{i,j} |P_{ij}\rangle |P_{ij}^*\rangle = \sum_{i,j} \sqrt{p_{ij}} |\psi_{ij}\rangle |P_{ij}^*\rangle, \end{aligned} \quad (5.9)$$

where in the second line, we have used the antiunitarity of the CPT operator. Eq. (5.8) motivates us to evaluate the changes in the entanglement entropy due to the local projective measurements on A^* and B^* .

Consider the projection

$$K_j = I_A \otimes I_B \otimes I_{A^*} \otimes |P_j^*\rangle\langle P_j^*|_{B^*}. \quad (5.10)$$

From the completeness of K_j , we get

$$\text{Tr}_{BB^*} |\sqrt{\rho}\rangle\langle\sqrt{\rho}| = \sum_j \text{Tr}_{BB^*} [K_j |\sqrt{\rho}\rangle\langle\sqrt{\rho}| K_j^\dagger]. \quad (5.11)$$

Therefore, we have

$$E_{RE}(\rho) = S(\text{Tr}_{BB^*} |\sqrt{\rho}\rangle\langle\sqrt{\rho}|) = S\left(\sum_j s_j \text{Tr}_{BB^*} |K_j \sqrt{\rho}\rangle\langle K_j \sqrt{\rho}|\right), \quad (5.12)$$

where

$$|K_j \sqrt{\rho}\rangle = \frac{K_j |\sqrt{\rho}\rangle}{\sqrt{\langle \sqrt{\rho} | K_j | \sqrt{\rho} \rangle}}, \quad (5.13)$$

$$s_j = \langle \sqrt{\rho} | K_j | \sqrt{\rho} \rangle. \quad (5.14)$$

Further, using the matrix concavity of the von Neumann entropy, we obtain

$$E_{RE}(\rho) \geq \sum_i s_i S(\text{Tr}_{BB^*} |K_i \sqrt{\rho}\rangle\langle K_i \sqrt{\rho}|). \quad (5.15)$$

In the same way, we also obtain

$$\begin{aligned}
& S(\text{Tr}_{BB^*} |K_j \sqrt{\rho}\rangle \langle K_j \sqrt{\rho}|) \\
&= S(\text{Tr}_{AA^*} |K_j \sqrt{\rho}\rangle \langle K_j \sqrt{\rho}|) \\
&\geq \sum_j r_{ij} S(\text{Tr}_{AA^*} |J_i K_j \sqrt{\rho}\rangle \langle J_i K_j \sqrt{\rho}|) \\
&= \sum_j r_{ij} S(\text{Tr}_{BB^*} |J_i K_j \sqrt{\rho}\rangle \langle J_i K_j \sqrt{\rho}|),
\end{aligned} \tag{5.16}$$

where

$$J_i = I_A \otimes I_B \otimes |P_i^*\rangle \langle P_i^*|_{A^*} \otimes I_{B^*}, \tag{5.17}$$

$$|J_i K_j \sqrt{\rho}\rangle = \frac{J_i K_j |\sqrt{\rho}\rangle}{\sqrt{\langle \sqrt{\rho} | J_i K_j | \sqrt{\rho} \rangle}}, \tag{5.18}$$

$$r_{ij} = \frac{\langle \sqrt{\rho} | J_i K_j | \sqrt{\rho} \rangle}{\langle \sqrt{\rho} | K_j | \sqrt{\rho} \rangle}. \tag{5.19}$$

Thus, we have

$$E_{RE}(\rho) \geq \sum_{i,j} r_{ij} s_j S(\text{Tr}_{BB^*} |J_i K_j \sqrt{\rho}\rangle \langle J_i K_j \sqrt{\rho}|). \tag{5.20}$$

Using Eq. (5.8), we immediately see that

$$r_{ij} s_j = p_{ij}, \tag{5.21}$$

$$|J_i K_j \sqrt{\rho}\rangle = |\psi_{ij}\rangle |P_{ij}^*\rangle. \tag{5.22}$$

Then, from additivity and non-negativity of the von Neumann entropy, we finally obtain

$$E_{RE}(\rho) \geq \sum_{i,j} p_{ij} \{S(\text{Tr}_B |\psi_{ij}\rangle \langle \psi_{ij}|) + S(\text{Tr}_{B^*} |P_{ij}^*\rangle \langle P_{ij}^*|)\} \geq \sum_{i,j} p_{ij} E_{vN}(|\psi_{ij}\rangle \langle \psi_{ij}|) = \overline{E_{vN}}. \tag{5.23}$$

Hence Eq. (5.7) is shown. \square

5.2 Constructing e^S area law states

Proposition 2 and theorem 3 are universal for any quantum system since no properties of specific physical systems are used in their proofs. Therefore, we can follow the algorithm presented in the proof of theorem 3 to construct orthogonal area law states. Starting from

the ensemble of states (5.3), we can pick up one area law state from them, according to theorem 7 and conjecture 5. Then we can construct another orthogonal area law state with the procedure presented in the proof of theorem 3. Again, we can repeat this procedure for at least $\exp[S(1 - o(1))]$ times without changing area law scaling of the average entanglement entropy.

Following this procedure, one can explicitly construct $\exp[S(1 - o(1))]$ orthogonal area law states from the states given in (5.3).

5.3 Constructing $e^{S_{BH}}$ disentangled states in holographic CFTs?

One may wonder if the state ensemble in (5.3) provides an example satisfying requirements in proposition 1, from which we can construct $\exp[S_{BH}(1 - o(1))]$ orthogonal disentangled states in holographic CFTs.

For this purpose, theorem 7 turns out to be useless, since in a holographic theory, $E_{RE} = 2E_P$ [33] and the difference between them $E_{RE} - E_P$ is $O(1/G_N) = O(S_{BH})$, i.e. at the leading order of $1/G_N$. On the other hand, numerical results in various one-dimensional critical quantum spin chains (including both integrable ones and chaotic ones) [50] show that $\overline{E_{vN}}$ over the ensemble (5.3) is given by $E_{RE}/2$. This indicates that we may expect (5.3) satisfies requirements in proposition 1. We leave confirming $\overline{E_{vN}} \approx E_{RE}/2$ in more general dimensions and systems (hopefully even in holographic CFTs) as a future problem.

6 Summary and Discussions

In this paper, we have studied two types of atypical states in holographic CFTs and generic quantum many-body systems. Our results can be summarized as follows:

- There exist at least $\exp[S_{BH}(1 - o(1))]$ orthogonal disentangled black hole microstates in the microcanonical subspace $\mathcal{H}_{[E-\Delta E, E]}$ in holographic CFTs at the semiclassical limit $G_N \rightarrow 0$.
- There exist at least $\exp[S(1 - o(1))]$ orthogonal area law states in the microcanonical subspace in generic quantum many-body systems, under the assumption that the microcanonical TFD obeys an area law of entanglement. Holographic CFTs and generic one-dimensional quantum many-body systems with short-ranged interactions satisfy this assumption.

- We have presented an explicit algorithm to construct $\exp[S(1 - o(1))]$ orthogonal area law states in generic quantum many-body systems and discussed its potential to be used to construct disentangled states in holographic CFTs.

Let us discuss possible improvements of our arguments, relations with other topics including additivity conjectures and black hole microstate counting, and future directions below.

6.1 Connections to additivity conjectures

In [13], the authors argue that the nonexistence of extensive violations of additivity conjectures leads to the existence $\exp[S_{BH} - O(1)]$ orthogonal disentangled microstates. Our results, the existence $\exp[S_{BH}(1 - o(1))]$ orthogonal disentangled microstates, is slightly weaker than this, but the physics associated with them seems to be essentially the same. Therefore, it seems that trying to find extensive violations of additivity conjectures from the black hole states might be a less fruitful direction than people expected [13, 24].

However, our arguments have the potential to give crucial insights into additivity conjectures from an alternative point of view. The reason is that bounding the entanglement of formation E_F plays a significant role in our argument, and there exists a version of additivity conjecture which is profoundly related to E_F [21]. The entanglement of formation E_F is lower bounded by the entanglement cost E_C [28], i.e. $E_F \geq E_C$, in any bipartite systems. Here, the entanglement cost of a bipartite state is defined as the minimal number of Bell pairs one needs to prepare the state with LOCC. It is known that additivity conjectures lead to $E_F = E_C$. Also in this case, while small violations are known [22, 51], extensive violations have not been found yet [52]. Therefore, it would be interesting to find possible relations between our arguments and this version of additivity conjectures.

6.2 Atypical microstate counting for black holes

Although it has long been known that one can find $e^{S_{BH}}$ black hole microstates from string theory since [53], less knowledge is known about the atypical state counting. One of the most successful atypical microstate counting from string theory is that of superstrata. Superstrata [54, 55] are horizonless geometries which are solutions of supergravity and describe microstates of the D1-D5-P black hole. These geometries deviate from the black hole and cap off smoothly. In this sense, the Ryu-Takayanagi surface in it would look very similar to the blue surface shown in the right of figure 1. This suggests superstrata are likely to be

explicit examples of disentangled states. However, so far, only about $O\left(\exp\left[S_{BH}^{3/4}\right]\right)$ superstrata have been found. This is parametrically smaller than our lower bound of disentangled states $\exp[S_{BH}(1 - o(1))]$. Although not every disentangled state is guaranteed to have a geometrical dual, this comparison is expected to give hints to atypical microstate counting in string theory.

6.3 Why atypicality is important in studying quantum gravity?

Let us get some hints from statistical mechanics and argue why studying atypicality is important in understanding quantum gravity. In statistical mechanics, although one knows almost every state in a microscopic theory is typical, it is usually hard to explicitly construct a typical state by experience [56], otherwise, we would just pick up such a typical state to get macroscopic information rather than use canonical ensembles or microcanonical ensembles.

An intuitive reason is as follows. In order to pick up a microstate from the energy shell in such a complicated many-body system, one usually needs to design a sophisticated algorithm. As a result, the state will be special in the sense it is picked up by a sophisticated algorithm, and hence tend to be atypical. From this point of view, it is natural why solving supergravity induced from string theory can give so many atypical geometries. This might also be able to help us understand why the chaotic nature of gravity seems to disappear when supersymmetry is involved in AdS₂ holography [57, 58].

On the other hand, this also suggests that studying generic and collective features of atypical states should be important in understanding the microscopic structure of quantum gravity since the microstates which are easier to get access to are expected to be atypical.

Acknowledgements

We are grateful to Yuya Kusuki, Yasunori Nomura, Pratik Rath, Masaki Shigemori, Hiroyasu Tajima, Tadashi Takayanagi and Kotaro Tamaoka for useful discussions and comments. ZW is supported by Grant-in-Aid for JSPS Fellows No. 20J23116. ZW was also supported in part by the National Science Foundation under Grant No. NSF PHY-1748958 and by the Heising-Simons Foundation. YY is supported by Grant-in-Aid for JSPS Fellows No. 20J22982.

A Axioms for entanglement measures

In this appendix, we summarize the axioms (or postulates) [34–37] that are expected for an entanglement measure $E(\rho)$. We consider a bipartite system $\mathcal{H} = \mathcal{H}_A \otimes \mathcal{H}_B$, and $\rho \in \mathcal{D}(\mathcal{H})$. We assume $\dim \mathcal{H}_B \leq \dim \mathcal{H}_A$ without loss of generality. The axioms are as follows.

- (i) If ρ is a separable state, then $E(\rho) = 0$.
- (ii) *Normalization* - If ρ is a maximally entangled state, then $E(\rho) = \log \dim \mathcal{H}_B$.
- (iii) *Monotonicity under LOCC* - $E(\rho)$ is monotonically decreasing under LOCC, i.e., for any LOCC operation $\Gamma_{\text{LOCC}}(\cdot)$,

$$E(\Gamma_{\text{LOCC}}(\rho)) \leq E(\rho). \quad (\text{A.1})$$

- (iv) *Asymptotic Continuity* - For any sequence of Hilbert space $\{\mathcal{H}_n\}$ and any sequence of states $\{\rho_n\}$ and $\{\sigma_n\}$ with $\rho_n, \sigma_n \in \mathcal{D}(\mathcal{H}_n)$,

$$\|\rho_n - \sigma_n\|_1 \rightarrow 0 \quad \Rightarrow \quad \frac{E(\rho_n) - E(\sigma_n)}{1 + \log \dim \mathcal{H}_n} \rightarrow 0. \quad (\text{A.2})$$

- (v) *Additivity* - For any state $\rho \in \mathcal{D}(\mathcal{H}) = \mathcal{D}(\mathcal{H}_A \otimes \mathcal{H}_B)$, consider its n -copy $\rho^{\otimes n} \in \mathcal{D}(\mathcal{H}^{\otimes n})$. Then for the bipartition $(\mathcal{H}_{A_1} \otimes \dots \otimes \mathcal{H}_{A_n}) \otimes (\mathcal{H}_{B_1} \otimes \dots \otimes \mathcal{H}_{B_n})$,

$$E(\rho^{\otimes n}) = nE(\rho). \quad (\text{A.3})$$

- (vi) *Convexity* - For any $\rho, \sigma \in \mathcal{D}(\mathcal{H})$ and any $\lambda \in [0, 1]$,

$$E(\lambda\rho + (1 - \lambda)\sigma) \leq \lambda E(\rho) + (1 - \lambda)E(\sigma). \quad (\text{A.4})$$

- (vii) *Subadditivity* - For any $\rho \in \mathcal{D}(\mathcal{H}) = \mathcal{D}(\mathcal{H}_A \otimes \mathcal{H}_B)$ and $\sigma \in \mathcal{D}(\mathcal{H}') = \mathcal{D}(\mathcal{H}_{A'} \otimes \mathcal{H}_{B'})$, consider divide $\mathcal{H} \otimes \mathcal{H}'$ into $(\mathcal{H}_A \otimes \mathcal{H}_{A'}) \otimes (\mathcal{H}_B \otimes \mathcal{H}_{B'})$, then

$$E(\rho \otimes \sigma) \leq E(\rho) + E(\sigma). \quad (\text{A.5})$$

B Alternative proofs of corollary 4 and 6

In this appendix, we present alternative proofs of corollary 4 and 6. These proofs avoid using the cumbersome theorem 3, and therefore turn out to be more simple if one merely wants to show the existence of $\exp[S_{BH}(1 - o(1))]$ disentangled states or $\exp[S(1 - o(1))]$ area law states and does not care about how to construct them. Since the proofs for both cases are similar, we will only present the proof for corollary 6. We will explain how corollary 4 can be proved in a similar way at the end of this appendix.

We firstly show the following theorem which bounds the change of the reflected entropy of microcanonical ensembles under projections. Note that a more general version of this theorem has already been shown in [59].

Theorem 8. *Consider an arbitrary W -dimensional subspace $\mathcal{E} \cong \mathbb{C}^W$ and a density matrix ρ which can be written as $\rho \propto \Pi_{\mathcal{E}}$, where $\Pi_{\mathcal{E}}$ is the projection operator onto \mathcal{E} . Then, for any $(W - k)$ -dimensional subspace $\mathcal{E}' \cong \mathbb{C}^{W-k}$ of \mathcal{E} , the reflected entropy of $\rho' \propto \Pi_{\mathcal{E}'}$ between A and B can be bounded from two sides as*

$$E_{RE}(\rho') \leq E_{RE}(\rho) + \sqrt{\frac{k}{W}} \left[\log \frac{W}{k} + 4 \log \dim \mathcal{H}_B - 2 \log 2 \right], \quad (\text{B.1})$$

$$E_{RE}(\rho') \geq E_{RE}(\rho) - \sqrt{\frac{k}{W}} \left[\log \frac{W}{k} + 4 \log \dim \mathcal{H}_B - 2 \log 2 \right], \quad (\text{B.2})$$

where $\Pi_{\mathcal{E}'}$ are the projection operators onto \mathcal{E}' .

Proof. Take an orthogonal basis $\{|\varphi_{\nu}\rangle\}_{\nu=1, \dots, W}$ of \mathcal{E} such that

$$\mathcal{E}' = \text{span}\{|\varphi_{\nu}\rangle\}_{\nu=k+1, \dots, W}. \quad (\text{B.3})$$

Then we have

$$\rho = \frac{1}{W} \sum_{\nu=1}^W |\varphi_{\nu}\rangle \langle \varphi_{\nu}|, \quad (\text{B.4})$$

$$\rho' = \frac{1}{W - k} \sum_{\nu=k+1}^W |\varphi_{\nu}\rangle \langle \varphi_{\nu}|. \quad (\text{B.5})$$

Therefore, according to the definition of the canonical purification in Eq. (2.12), we have

$$|\sqrt{\rho}\rangle = \frac{1}{\sqrt{W}} \sum_{\nu=1}^W |\varphi_{\nu}\rangle |\varphi_{\nu}^*\rangle, \quad (\text{B.6})$$

$$|\sqrt{\rho'}\rangle = \frac{1}{\sqrt{W - k}} \sum_{\nu=k+1}^W |\varphi_{\nu}\rangle |\varphi_{\nu}^*\rangle. \quad (\text{B.7})$$

Hence,

$$\langle \sqrt{\rho'} | \sqrt{\rho} \rangle = \sqrt{1 - \frac{k}{W}}. \quad (\text{B.8})$$

Since

$$\begin{aligned} & \| \text{Tr}_{AA^*} | \sqrt{\rho'} \rangle \langle \sqrt{\rho'} | - \text{Tr}_{AA^*} | \sqrt{\rho} \rangle \langle \sqrt{\rho} | \|_1 \\ & \leq \| | \sqrt{\rho'} \rangle \langle \sqrt{\rho'} | - | \sqrt{\rho} \rangle \langle \sqrt{\rho} | \|_1 \\ & = 2\sqrt{1 - |\langle \sqrt{\rho'} | \sqrt{\rho} \rangle|^2} \\ & = \sqrt{\frac{4k}{W}}, \end{aligned} \quad (\text{B.9})$$

applying Fannes' inequality, we obtain

$$\begin{aligned} |E_{RE}(\rho') - E_{RE}(\rho)| & = |S(\text{Tr}_{AA^*} | \sqrt{\rho'} \rangle \langle \sqrt{\rho'} |) - S(\text{Tr}_{AA^*} | \sqrt{\rho} \rangle \langle \sqrt{\rho} |)| \\ & \leq \sqrt{\frac{4k}{W}} \log \dim \mathcal{H}_{BB^*} - \sqrt{\frac{4k}{W}} \log \sqrt{\frac{4k}{W}} \\ & = \sqrt{\frac{k}{W}} \left[\log \frac{W}{k} + 4 \log \dim \mathcal{H}_B - 2 \log 2 \right]. \end{aligned} \quad (\text{B.10})$$

□

Let us then combine conjecture 5 and theorem 8. As \mathcal{E} , let us consider the microcanonical subspace

$$\mathcal{H}_{[E-\Delta E, E]} = \text{span} \{ |E_i\rangle \in \mathcal{H} | E_i \in [E - \Delta E, E] \}. \quad (\text{B.11})$$

Then ρ is the associated microcanonical ensemble $\rho^{(\text{micro})}$. According to the Boltzmann formula, $W \sim e^S = O(L^d)$, where S is the thermodynamic entropy. By using conjecture 5 and $E_F \leq E_{RE}$, we can say that there exists at least one area law state $|\varphi_1\rangle$ in $\mathcal{H}_{[E-\Delta E, E]}$.

Then we exclude $|\varphi_1\rangle$ from $\mathcal{H}_{[E-\Delta E, E]}$ and consider $\mathcal{H}_{[E-\Delta E, E]} \cap (\text{span}\{|\varphi_1\rangle\})^\perp$ as a new subspace \mathcal{E}' . In this case, since $\log \dim \mathcal{H}_B = O(L^d) = O(S)$, according to the upper bound (B.1) given in theorem 8, the reflected entropy of $\rho' \propto \Pi_{\mathcal{E}'}$ is changed by at most $O(e^{-S/2}S)$ from that of ρ . Again, using $E_F \leq E_{RE}$, we can say that there exists at least one area law state $|\varphi_2\rangle$ in $\mathcal{H}_{[E-\Delta E, E]} \cap (\text{span}\{|\varphi_1\rangle\})^\perp$.

According to the bound given in theorem 8, one can at least repeat this procedure for $\exp[S(1 - o(1))]$ times. As a result, one can obtain $\exp[S(1 - o(1))]$ area law states. By construction, these states are orthogonal. As a result, one can get corollary 6.

The key point of theorem 8 is that the change of the reflected entropy when excluding a state from a microcanonical ensemble is sufficiently small. Therefore, if one can show that the change of the entanglement of purification when excluding a state from a microcanonical ensemble is also sufficiently small, the corollary 4 can be shown in the same manner. In fact, such a statement follows directly from the continuity of the entanglement of purification shown in [27, 59]. Alternatively, one can also use the continuity of the entanglement of formation shown in [60] to show corollary 4 and 6.

References

- [1] S. Goldstein, J. L. Lebowitz, R. Tumulka and N. Zanghì, *Canonical typicality*, *Phys. Rev. Lett.* **96** (2006) 050403 [[0511091](#)].
- [2] S. Popescu, A. J. Short and A. Winter, *Entanglement and the foundations of statistical mechanics*, *Nat. Phys.* **2** (2006) 754 [[0511225](#)].
- [3] P. Reimann, *Typicality for generalized microcanonical ensemble*, *Phys. Rev. Lett.* **99** (2007) 160404 [[0710.4214](#)].
- [4] M. P. Müller, E. Adlam, L. Masanes and N. Wiebe, *Thermalization and Canonical Typicality in Translation-Invariant Quantum Lattice Systems*, *Commun. Math. Phys.* **340** (2015) 499 [[1312.7420](#)].
- [5] T. Mori, T. Ikeda, E. Kaminishi and M. Ueda, *Thermalization and prethermalization in isolated quantum systems: a theoretical overview*, *J. Phys. B* **51** (2018) 112001 [[1712.08790](#)].
- [6] S. W. Hawking, *Particle Creation by Black Holes*, *Commun. Math. Phys.* **43** (1975) 199.
- [7] D. N. Page, *Information in black hole radiation*, *Phys. Rev. Lett.* **71** (1993) 3743 [[hep-th/9306083](#)].
- [8] A. Almheiri, D. Marolf, J. Polchinski and J. Sully, *Black Holes: Complementarity or Firewalls?*, *JHEP* **02** (2013) 062 [[1207.3123](#)].

- [9] A. Almheiri, D. Marolf, J. Polchinski, D. Stanford and J. Sully, *An Apologia for Firewalls*, *JHEP* **09** (2013) 018 [[1304.6483](#)].
- [10] S. D. Mathur, *The Fuzzball proposal for black holes: An Elementary review*, *Fortsch. Phys.* **53** (2005) 793 [[hep-th/0502050](#)].
- [11] Y. Nomura, J. Varela and S. J. Weinberg, *Black Holes, Information, and Hilbert Space for Quantum Gravity*, *Phys. Rev. D* **87** (2013) 084050 [[1210.6348](#)].
- [12] T. Hartman and J. Maldacena, *Time Evolution of Entanglement Entropy from Black Hole Interiors*, *JHEP* **05** (2013) 014 [[1303.1080](#)].
- [13] P. Hayden and G. Penington, *Black hole microstates vs. the additivity conjectures*, [2012.07861](#).
- [14] J. M. Maldacena, *The Large N limit of superconformal field theories and supergravity*, *Int. J. Theor. Phys.* **38** (1999) 1113 [[hep-th/9711200](#)].
- [15] S. Ryu and T. Takayanagi, *Holographic derivation of entanglement entropy from AdS/CFT*, *Phys. Rev. Lett.* **96** (2006) 181602 [[hep-th/0603001](#)].
- [16] S. Ryu and T. Takayanagi, *Aspects of Holographic Entanglement Entropy*, *JHEP* **08** (2006) 045 [[hep-th/0605073](#)].
- [17] V. E. Hubeny, M. Rangamani and T. Takayanagi, *A Covariant holographic entanglement entropy proposal*, *JHEP* **07** (2007) 062 [[0705.0016](#)].
- [18] A. C. Wall, *Maximin Surfaces, and the Strong Subadditivity of the Covariant Holographic Entanglement Entropy*, *Class. Quant. Grav.* **31** (2014) 225007 [[1211.3494](#)].
- [19] T. Faulkner, A. Lewkowycz and J. Maldacena, *Quantum corrections to holographic entanglement entropy*, *JHEP* **11** (2013) 074 [[1307.2892](#)].
- [20] N. Engelhardt and A. C. Wall, *Quantum Extremal Surfaces: Holographic Entanglement Entropy beyond the Classical Regime*, *JHEP* **01** (2015) 073 [[1408.3203](#)].
- [21] P. W. Shor, *Equivalence of additivity questions in quantum information theory*, *Commun. Math. Phys.* **246** (2004) 453 [[0305035](#)].

- [22] M. B. Hastings, *Superadditivity of communication capacity using entangled inputs*, *Nature Physics* **5** (2009) 255–257.
- [23] Z. Ji, A. Natarajan, T. Vidick, J. Wright and H. Yuen, *MIP*=RE*, [2001.04383](#).
- [24] J. Wang, *Beyond islands: A free probabilistic approach*, [2209.10546](#).
- [25] T. Takayanagi and K. Umemoto, *Entanglement of purification through holographic duality*, *Nature Phys.* **14** (2018) 573 [[1708.09393](#)].
- [26] P. Nguyen, T. Devakul, M. G. Halbasch, M. P. Zaletel and B. Swingle, *Entanglement of purification: from spin chains to holography*, *JHEP* **01** (2018) 098 [[1709.07424](#)].
- [27] B. M. Terhal, M. Horodecki, D. W. Leung and D. P. DiVincenzo, *The entanglement of purification*, *Journal of Mathematical Physics* **43** (2002) 4286–4298 [[quant-ph/0202044](#)].
- [28] C. H. Bennett, D. P. DiVincenzo, J. A. Smolin and W. K. Wootters, *Mixed-state entanglement and quantum error correction*, *Physical Review A* **54** (1996) 3824–3851 [[quant-ph/9604024](#)].
- [29] C. Murthy and M. Srednicki, *Structure of chaotic eigenstates and their entanglement entropy*, *Phys. Rev. E* **100** (2019) 022131 [[1906.04295](#)].
- [30] M. Srednicki, *Chaos and quantum thermalization*, *Phys. Rev. E* **50** (1994) 888 [[cond-mat/9403051](#)].
- [31] D. Marolf, S. Wang and Z. Wang, *Probing phase transitions of holographic entanglement entropy with fixed area states*, *JHEP* **12** (2020) 084 [[2006.10089](#)].
- [32] X. Dong and H. Wang, *Enhanced corrections near holographic entanglement transitions: a chaotic case study*, *JHEP* **11** (2020) 007 [[2006.10051](#)].
- [33] S. Dutta and T. Faulkner, *A canonical purification for the entanglement wedge cross-section*, [1905.00577](#).
- [34] M. J. Donald, M. Horodecki and O. Rudolph, *The uniqueness theorem for entanglement measures*, *Journal of Mathematical Physics* **43** (2002) 4252–4272 [[quant-ph/0105017](#)].

- [35] M. B. Plenio and S. Virmani, *An Introduction to entanglement measures*, *Quant. Inf. Comput.* **7** (2007) 1 [[quant-ph/0504163](#)].
- [36] R. Horodecki, P. Horodecki, M. Horodecki and K. Horodecki, *Quantum entanglement*, *Rev. Mod. Phys.* **81** (2009) 865 [[quant-ph/0702225](#)].
- [37] G. Benenti, G. Casati, D. Rossini and G. Strini, *Principles of Quantum Computation and Information*. WORLD SCIENTIFIC, 2018, [10.1142/10909](#).
- [38] B. Czech, J. L. Karczmarek, F. Nogueira and M. Van Raamsdonk, *The Gravity Dual of a Density Matrix*, *Class. Quant. Grav.* **29** (2012) 155009 [[1204.1330](#)].
- [39] M. Headrick, V. E. Hubeny, A. Lawrence and M. Rangamani, *Causality & holographic entanglement entropy*, *JHEP* **12** (2014) 162 [[1408.6300](#)].
- [40] M. Miyaji and T. Takayanagi, *Surface/State Correspondence as a Generalized Holography*, *PTEP* **2015** (2015) 073B03 [[1503.03542](#)].
- [41] B. Swingle, *Entanglement Renormalization and Holography*, *Phys. Rev. D* **86** (2012) 065007 [[0905.1317](#)].
- [42] F. Pastawski, B. Yoshida, D. Harlow and J. Preskill, *Holographic quantum error-correcting codes: Toy models for the bulk/boundary correspondence*, *JHEP* **06** (2015) 149 [[1503.06237](#)].
- [43] P. Hayden, S. Nezami, X.-L. Qi, N. Thomas, M. Walter and Z. Yang, *Holographic duality from random tensor networks*, *JHEP* **11** (2016) 009 [[1601.01694](#)].
- [44] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, *Gauge theory correlators from noncritical string theory*, *Phys. Lett.* **B428** (1998) 105 [[hep-th/9802109](#)].
- [45] E. Witten, *Anti-de Sitter space and holography*, *Adv. Theor. Math. Phys.* **2** (1998) 253 [[hep-th/9802150](#)].
- [46] J. M. Maldacena, *Eternal black holes in anti-de Sitter*, *JHEP* **04** (2003) 021 [[hep-th/0106112](#)].
- [47] D. Marolf, *Microcanonical Path Integrals and the Holography of small Black Hole Interiors*, *JHEP* **09** (2018) 114 [[1808.00394](#)].

- [48] T. Barthel, *One-dimensional quantum systems at finite temperatures can be simulated efficiently on classical computers*, [1708.09349](#).
- [49] T. Kuwahara, Á. M. Alhambra and A. Anshu, *Improved Thermal Area Law and Quasilinear Time Algorithm for Quantum Gibbs States*, *Phys. Rev. X* **11** (2021) 011047 [[2007.11174](#)].
- [50] Y. Kusuki, K. Tamaoka, Z. Wei and Y. Yoneta, *to appear*, .
- [51] M. Fukuda and M. M. Wolf, *Simplifying additivity problems using direct sum constructions*, *J. Math. Phys.* **48** (2007) 072101.
- [52] S. T. Belinschi, B. Collins and I. Nechita, *Almost one bit violation for the additivity of the minimum output entropy*, *Commun. Math. Phys.* **341** (2016) 885 [[1305.1567](#)].
- [53] A. Strominger and C. Vafa, *Microscopic origin of the Bekenstein-Hawking entropy*, *Phys. Lett. B* **379** (1996) 99 [[hep-th/9601029](#)].
- [54] I. Bena, S. Giusto, R. Russo, M. Shigemori and N. P. Warner, *Habemus Superstratum! A constructive proof of the existence of superstrata*, *JHEP* **05** (2015) 110 [[1503.01463](#)].
- [55] M. Shigemori, *Counting Superstrata*, *JHEP* **10** (2019) 017 [[1907.03878](#)].
- [56] S. Sugiura and A. Shimizu, *Thermal Pure Quantum States at Finite Temperature*, *Phys. Rev. Lett.* **108** (2012) 240401 [[1112.0740](#)].
- [57] H. W. Lin, J. Maldacena, L. Rozenberg and J. Shan, *Holography for people with no time*, [2207.00407](#).
- [58] H. W. Lin, J. Maldacena, L. Rozenberg and J. Shan, *Looking at supersymmetric black holes for a very long time*, [2207.00408](#).
- [59] C. Akers and P. Rath, *Entanglement Wedge Cross Sections Require Tripartite Entanglement*, *JHEP* **04** (2020) 208 [[1911.07852](#)].
- [60] M. A. Nielsen, *Continuity bounds for entanglement*, *Physical Review A* **61** (2000) [[quant-ph/9908086](#)].