Embedding of the Georgi-Glashow SU(5) model in the superconformal algebra

P. D. Alvarez^{*1}, R. A. Chavez^{†1}, and J. Zanelli^{‡2}

¹Departamento de Física, Universidad de Antofagasta, Aptdo. 02800, Chile

²Centro de Estudios Científicos (CECs), Arturo Prat 514, Valdivia, Chile, Universidad San Sebastián, General Lagos 1163, Valdivia,

Chile

December 6, 2022

Abstract

We present a scheme to construct grand unified models based on the superconformal algebra and the inclusion of matter fields in the adjoint representation of supersymmetry. As an illustration, we implemented the Georgi-Glashow SU(5) model. The model predicts the existence of a hidden $(\mathbf{1}, \mathbf{24}, 0) + (\mathbf{5}, \mathbf{5}^*, -y') + (\mathbf{5}^*, \mathbf{5}, y')$ sector and an anomalous $U(1)_Z$.

Contents

Inti	roduction	2	
The	e model	3	
$3.1 \\ 3.2$	Fermionic sector	10	
4 Conclusions			
ppen	dices	14	
	The Gau 3.1 3.2 3.3 Con	Introduction The model Gauge couplings 3.1 Fermionic sector 3.2 Gauge potentials 3.3 Comments on anomalies Conclusions ppendices	

^{*}E-mail: pedro.alvarez@uantof.cl

[†]E-mail: rafael.chavez.linares@uantof.cl

[‡]E-mail: z@cecs.cl

A Fundamental representation of $SU(2,2 n)$	14
B Bi-fundamental representation of $SU(2,2 n)$	17

 $\mathbf{18}$

C SU(2,2|n) curvatures

1 Introduction

The simplest grand unified theory (GUT) is the Georgy-Glashow (GG) SU(5) model, where the fifteen leptons and quarks of the Standard Model are included in the **5**^{*} and a **10** representation of SU(5) [1, 2] (in the one-family approximation). Among the welcomed features we have unification of coupling constants, quark charge quantization and a rationale for anomaly cancellation. The price to pay for such beauty is the presence of proton decay channels and the doublettriplet splitting problems on top of the usual quadratic mass renormalization of the scalars.

There are improved models that address some of the issues of the GG model. To protect the Higgs doublets from quadratic mass renormalization, an SU(5) model with softly broken supersymmetry was proposed [3]. To avoid proton decay and the doublet-triplet splitting problems of the GG SU(5) model, the flipped SU(5) model was proposed [4]. The supersymmetric flipped SU(5) model, that produces hierarchical neutrino masses, was proposed [5, 6, 7, 8]. Flipped SU(5) can also be embedded in string theory [9, 10, 11].

Minimal flipped SU(5) GUT was shown to survive experimental electroweak limits [12]. The flipped SU(5) GUT has also been embedded in no-scale supergravity models [13]. Such model can generate no-scale inflation, similar to Starobinsky's inflation, [14], and makes predictions for cosmic microwave background observables that can be used to constrain significantly the parameters of the model.

One shortcoming of the flipped SU(5) model is that the hypercharge is in the factor of the gauge group proposed for unification $(SU(5) \times U(1)_{\chi})/\mathbb{Z}_5$. The fact that the structure $SU(n) \times U(1)$ can be found in the superconformal algebra is suggestive. However, the embedding of supersymmetric SU(n) GUT models in the superconformal algebra faces technical difficulties [15, 16].

In this paper we present a model that circumvents those difficulties by using a novel implementation of supersymmetry with matter in the adjoint representation of the superconformal algebra. We use two basic assumptions, firstly we use a diagonal symmetry group

$$SU(2,2|n)_{\text{diag}} = [SU(2,2|n) \times SU(2,2|d_n)]_{\text{diag}}$$
(1)

where $d_n = n(n-1)/2$. Such a value comes from the embedding of supercharges in the superconformal algebra carrying a bi-fundamental index [17]. Secondly, we define a dual operator which is naturally embedded in SU(2, 2|n) [18]. This dual operator induces a grading in the conformal algebra. This grading allows defining actions with gauge symmetry $G^+ \subset SU(2, 2) \times SU(n) \times U(1)$, where G^+ is the group generated by the grading-even generators [18]. The symmetries generated by the grading-odd generators, and the supersymmetries become on-shell symmetries [19]. For the sake of simplicity we focus on the implementation of the GG model.

2 The model

Let us define a model for gauge potential for the group (1). The gauge connection is

$$\mathbb{A} = \Omega + \overline{\mathbb{Q}}^{i} \not e \psi_{i} + \overline{\psi}^{i} \not e \mathbb{Q}_{i} , \qquad (2)$$

$$\mathbb{A}' = \Omega' + \frac{1}{2} \overline{\mathbb{Q}}^{ij} \not\in \chi_{ij} + \frac{1}{2} \overline{\chi}^{ij} \not\in \mathbb{Q}_{ij} , \qquad (3)$$

where

$$\Omega = \frac{1}{2}\omega^{ab}\mathbb{J}_{ab} + f^a\mathbb{J}_a + g^a\mathbb{K}_a + h\mathbb{D} + A^I\mathbb{T}_I + A\mathbb{Z}, \qquad (4)$$

$$\Omega' = \frac{1}{2} \omega'^{ab} \mathbb{J}_{ab} + f'^a \mathbb{J}_a + g'^a \mathbb{K}_a + h' \mathbb{D} + A'^X \mathbb{T}_X + A' \mathbb{Z} , \qquad (5)$$

and $a = 0, \dots, 3$, $I = 1, \dots, n^2 - 1$, $X = 1, \dots, d_n^2 - 1$. We are interested in the embedding of the Georgi-Glashow model into the superconformal algebra. For this purpose we will consider (1) with n = 5. Note that, in the present model, the gauge bosons A, A' and $A'^{\widetilde{X}}$ with $\widetilde{X} = n^2, n^2 + 1, \dots, d_n^2 - 1$, are new with respect to the Georgi-Glashow model. The spinors transform in the appropriate representations of SU(5),

$$(\psi^i)_L = \mathbf{5}^*, \quad (\chi_{ij})_L = \mathbf{10}.$$
 (6)

with $\chi_{ij} = -\chi_{ji}$, and the fifteen left-handed quarks and lepton of the standard model are placed in a reducible $5^* + 10$ representation,

$$\mathbf{5}^* = (\mathbf{3}^*, \mathbf{1}, \frac{1}{3}) + (\mathbf{1}, \mathbf{2}^*, -\frac{1}{2}), \tag{7}$$

$$\mathbf{10} = (\mathbf{3}^*, \mathbf{1}, -\frac{2}{3}) + (\mathbf{3}, \mathbf{2}, \frac{1}{6}) + (\mathbf{1}, \mathbf{1}, 1), \qquad (8)$$

where $(\mathbf{n}_3, \mathbf{n}_2, y)$ give the representations under SU(3), SU(2) and the weak hypercharge [2].

The field assignment is as follows

$$(\psi^{i})_{L} = \begin{pmatrix} d^{c1} \\ d^{c2} \\ d^{c3} \\ e^{-} \\ -\nu_{e} \end{pmatrix}_{L}, \qquad (9)$$

and

$$(\chi_{ij})_L = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & u^{c3} & -u^{c2} & -u_1 & -d_1 \\ -u^{c3} & 0 & u^{c1} & -u_2 & -d_2 \\ u^{c2} & -u^{c1} & 0 & -u_3 & -d_3 \\ u^1 & u^2 & u^3 & 0 & -e^+ \\ d^1 & d^2 & d^3 & e^+ & 0 \end{pmatrix}_L$$
(10)

The action principle is based on a relative of the MacDowell-Mansouri action [20],

$$\mathcal{S} = -\int \left(\langle \xi \mathbb{F} \circledast \mathbb{F} \rangle + \langle \xi' \mathbb{F}' \circledast \mathbb{F}' \rangle \right) \,. \tag{11}$$

The field strength is defined as usual by $\mathbb{F} = d\mathbb{A} + \mathbb{A} \wedge \mathbb{A}$, where wedge products are assumed. The explicit expressions for the curvature components are in appendix C. In the action (11) we use a generalized dual operator defined by¹

and similarly for \mathbb{F}' . *S* is a symmetry breaking operator that allows to obtain general relativity as the gravity action, see appendix A. The parameters ε_s , ε_1 , ε_2 , ε_3 and ε_{ψ} can take values +1 or -1 only, so that $\circledast^2 = -1$ and therefore the correct sign of the kinetic terms of bosons is ensured. For the present paper we will fix $\varepsilon_s = +1 = \varepsilon_1 = \varepsilon_2$ and $\varepsilon_3 = -1$.

Demanding the diagonal symmetry group to be su(2,2|5) means that not all the fields in A and A' are independent,

$$\omega^{\prime ab} = \omega^{ab} \,, \tag{13}$$

$$g^{\prime a} = g^{a} , \tag{15}$$

$$h' = h, \tag{16}$$

$$A' = A, \tag{17}$$

and

$$A^{\prime X}\big|_{X=I} = A^I \,. \tag{18}$$

Therefore (11) has only two adjustable parameters in front of the unprimed and primed contributions, ξ and ξ' respectively. The present model is a unified model in the sense that the ratios of *all* the couplings are determined from the superconformal algebra and the way in which the embedding of quarks and leptons is implemented. The weights of the unprimed and primed sector are not

 $^{^1\}mathrm{It}$ would be interesting to see how this Hodge operator relates to the one defined on a supermanifold [21, 22].

determined a priori though, and they can be treated as phenomenological parameters. These parameters define two dimensional parameter space of allowed relative strengths for the gauge couplings $g^{(SU(5))}$ and $g^{(U(1))}$. The relations between gauge couplings will not be arbitrary though as they will be consequence of the superconformal algebra and the particular embedding of bosons and fermions. We discuss this in details in section 3

3 Gauge couplings

In order to obtain an explicit expansion of (11), we use the invariant traces defined in the superalgebra, given in appendix A, see eqs. (117) - (120) and (121) - (124). After imposing the diagonal symmetry (13)-(17),

$$\mathcal{L} = \frac{1}{4} \varepsilon_s (\xi + \xi') \epsilon_{abcd} \mathcal{F}^{ab} \mathcal{F}^{cd} - \varepsilon_1 (\xi + \xi') \mathcal{H} * \mathcal{H} - \frac{1}{2} \varepsilon_2 \left(\xi \mathcal{F}^I * \mathcal{F}^I + \xi' (n-2) \mathcal{F}'^X * \mathcal{F}'^X \right) - 4 \varepsilon_3 \left[\xi (4/n-1) + \xi' (4/d_n-1) \right] \mathcal{F} * \mathcal{F} , - 2i \varepsilon_{\psi} \overline{\mathcal{X}} \gamma_5 \mathcal{X} - \frac{i}{2} \varepsilon_{\chi} \overline{\mathcal{Y}} \gamma_5 \mathcal{Y} .$$
(19)

Here we have restored the label n of the superalgebra su(2, 2|n) in order to draw general conclusions about the validity of the model. In the bosonic sector, $\psi_i = 0 = \chi_{ij}$, and imposing the diagonal symmetry along the su(n) generators as well, yields

$$\mathcal{L}_{\rm b} = \frac{1}{4} \varepsilon_s(\xi + \xi') \epsilon_{abcd} \mathcal{R}^{ab} \mathcal{R}^{cd} - \varepsilon_1(\xi + \xi') H * H - \frac{1}{2} \varepsilon_2 \left(\xi + \xi'(n-2)\right) F^I * F^I - 4 \varepsilon_3 \left[\xi(4/n-1) + \xi'(4/d_n-1)\right] F * F - \frac{(n-2)}{2} \varepsilon_2 \xi' \left[2F^I * F_1^I + F_1^I * F_1^I + F^{\tilde{X}} * F^{\tilde{X}}\right],$$
(20)

where

$$F^{I} = dA^{I} + \frac{1}{2} f_{JK}{}^{I} A^{J} A^{K} , \qquad (21)$$

$$F_1^I = f_{J\widetilde{Y}}{}^I A^J A^{\widetilde{Y}} + \frac{1}{2} f_{\widetilde{Y}\widetilde{Z}}{}^I A^{\widetilde{Y}} A^{\widetilde{Z}} , \qquad (22)$$

$$F^{\widetilde{X}} = dA^{\widetilde{X}} + \frac{1}{2} \left(f_{JK}{}^{\widetilde{X}}A^{J}A^{K} + 2f_{J\widetilde{Y}}{}^{\widetilde{X}}A^{J}A^{\widetilde{Y}} + f_{\widetilde{Y}\widetilde{Z}}{}^{\widetilde{X}}A^{\widetilde{Y}}A^{\widetilde{Z}} \right) .$$
(23)

In (22) and (23) we removed the primed to the $A'^{\widetilde{X}}$ fields to simplify the notation.

Let us now discuss the values of the coupling constants implied by the model. The gravity theory, in the sector defined by

$$f^a = \rho e^a , \qquad g^a = \sigma e^a , \quad |\rho| > |\sigma|$$
(24)

corresponds to general relativity with cosmological constant

$$\frac{1}{16\pi G_N} \int \epsilon_{abcd} \left(\frac{1}{2} R^{ab} e^c e^d - \frac{\Lambda}{12} e^a e^b e^c e^d \right) \subset \mathcal{S} \,, \tag{25}$$

under the following identification of constants

$$\frac{1}{16\pi G_N} = (\rho^2 - \sigma^2)(\xi + \xi'), \qquad \Lambda = -3(\rho^2 - \sigma^2) = -\frac{3}{2(\xi + \xi')}M_P^2.$$
(26)

The positivity of the kinetic terms for the gauge fields requires

$$\xi + \xi' > 0, \qquad (27)$$

$$\xi(4/n-1) + \xi'(4/d_n - 1) < 0, \qquad (28)$$

$$\xi + \xi'(n-2) > 0.$$
⁽²⁹⁾

Such conditions are satisfied in a two dimensional region of the parameter space for any $n \ge 5$, see fig. 1




Figure 1: Allowed regions of the (ξ, ξ') plane, The left panel corresponds to the case n = 5. The right panel is the plot for n = 6, which illustrates the qualitative behavior for n > 5. In green, blue and red are the null conditions for eqs. (27), (28) and (29) respectively. Conditions (28) and (29) are degenerate for n = 5.

The values of the coupling constants are defined with respect to canonically normalized gauge fields,

$$-aF * F = -\frac{1}{2}F^{\text{phys}} * F^{\text{phys}}, \qquad (30)$$

where the physical gauge potential is $A^{\text{phys}} = \sqrt{2a}A$. Assuming the covariant derivative on the spinor to be of the form

$$D = d - ig_0 \ \rho(T_r)A^r \,, \tag{31}$$

where $\rho(T_r)$ stands for the corresponding representation of the generators T_r . Then, the physical values of the coupling constants, $g = g_0/\sqrt{2a}$ can be inferred to be

$$g_{(SU(n))} = g_{(SU(d_n))} = \frac{1}{\sqrt{\xi + \xi'(n-2)}},$$
(32)

$$g_{(U(1))}^{(\text{rank 1})} = \frac{4/n - 1}{\sqrt{-8(\xi(4/n - 1) + \xi'(4/d_n - 1))}},$$
(33)

$$g_{(U(1))}^{(\text{rank }2)} = \frac{4/d_n - 1}{\sqrt{-8(\xi(4/n - 1) + \xi'(4/d_n - 1))}} \,.$$
(34)

The fact that $g_{(SU(n))} = g_{(SU(d_n))}$ allows for a natural embedding of the representations in the GG model (6) and it will also ensure gauge coupling unification. To simplify the notation we will denote such coupling constants by g. The normalization of the gauge field h gives $g^{(U_D(1))} = 1/\sqrt{2(\xi + \xi')}$, however, careful evaluation of the fermion kinetic terms in the last two terms of (19), shows that such coupling cancels out and therefore the field h becomes a hidden photon [23]. This is in fact necessary for physical consistency of the model since the transformations generated by \mathbb{D} are non compact.

3.1 Fermionic sector

The fermion kinetic terms are contained in the last two terms of (19),

$$-2i\overline{\mathcal{X}}\gamma_5\mathcal{X} - \frac{i}{2}\overline{\mathcal{Y}}\gamma_5\mathcal{Y}.$$
(35)

The fact that such a terms contain the right expressions for the kinetic terms of ψ and χ can be seen as a consequence of a few assumptions:

- (a) The fundamental fermionic fields are in the spin 1/2 representation of the fermionic field in the gauge potential, $\Psi = \not e \psi$ and $X = \not e \chi$, that are in the kernel of the the spin 3/2 projector, $P_{(3/2)}\Psi = 0$ and $P_{(3/2)}X = 0$.
- (b) The fermionic components of the curvature are the exterior covariant derivatives of Ψ and X with the connection for the corresponding bosonic gauge groups $\mathcal{X} = D_{G_1}\psi$ and $\mathcal{Y} = D_{G_2}\psi$, where $G_1 = SU(2,2) \times SU(n) \times U(1)$ and $G_2 = SU(2,2) \times SU(d_n) \times U(1)$ respectively.
- (c) The dual operator acting on the fermionic components of the curvature is a grading operator of the bosonic gauge symmetries that splits the symmetry between broken an unbroken symmetries. Therefore the covariant derivatives in $\overline{\mathcal{X}} D_{\Omega} \otimes D_{\Omega} \mathcal{X}$ split naturally into $D_{\Omega} = D_{+} + \Omega_{-}$, where $\Omega = \Omega_{+} + \Omega_{-}$ (for more details see [18]).
- (d) The fields in Ω_{-} are dual to generators that anticommute with \circledast , in the present case

$$\Omega_{-} = \frac{1}{2} f^a \gamma_a + \frac{1}{2} g^a \tilde{\gamma}_a \,. \tag{36}$$

Furthermore, these fields are related to the orthonormal frames by (24).

With these assumptions, it can be seen that the terms containing second order derivatives of the fermionic fields are boundary terms, while the first order terms give rise to the gauge covariant kinetic terms. The resulting Lagrangian for the spinor ψ with a connection $\mathcal{W} \in so(1,3) \times su(n) \times u(1)$ is

$$\frac{\mathcal{L}(\psi)}{\xi} = -2i\overline{\psi}(\overleftarrow{D}_{\mathcal{W}}\notin\gamma_5\Omega_-\notin-\notin\Omega_-\gamma_5\notin D_{\mathcal{W}})\psi
-2i\overline{\psi}(\overrightarrow{T}\gamma_5\Omega_-\notin+\notin\Omega_-\gamma_5\overrightarrow{T})\psi
+2i\overline{\psi}\gamma_5\notin \left[R^{ab}\Sigma_{ab}-2i(4/n-1)F\right]\notin\psi
+2id\left[\overline{\psi}\gamma_5\notin D_+(\notin\psi)\right],$$
(37)

where

and P_R (P_L) is the chiral right-handed projector (left-handed projector). Note here that the term $(1/2)h\mathbb{D}$ in (37) cancels out from \mathcal{W} thanks to the fact that \mathbb{J}_a and \mathbb{K}_a anticommute with \mathbb{D} . The constants α and β are linear combinations of ρ and σ

$$\alpha = \frac{\rho + \sigma}{2}, \quad \beta = \frac{\rho - \sigma}{2}. \tag{39}$$

Similarly for the spinor χ with a connection $\mathcal{W}' \in so(1,3) \times su(d_n) \times u(1)$,

$$\frac{\mathcal{L}(\chi)}{\xi'} = -\frac{i}{2} \overline{\chi} (\overleftarrow{D}_{\mathcal{W}'} \notin \gamma_5 \Omega_- \not\!\!\!/ e - \not\!\!/ \Theta_- \gamma_5 \not\!\!/ D_{\mathcal{W}'}) \chi
- \frac{i}{2} \overline{\chi} (\overrightarrow{T} \gamma_5 \Omega_- \not\!\!/ e + \not\!\!/ \Theta_- \gamma_5 \not\!\!/ T) \chi
+ \frac{i}{2} \overline{\chi} \gamma_5 \not\!\!/ e \left[R^{ab} \Sigma_{ab} - 2i(4/d_n - 1) F \right] \not\!\!/ e \chi ,
+ \frac{i}{2} d [\overline{\chi} \gamma_5 \not\!\!/ e D_+ (\not\!\!/ \chi)] ,$$
(40)

The first lines in (37) and (40) give the gauge covariant kinetic terms

$$\frac{\mathcal{L}_{K}(\psi) + \mathcal{L}_{K}(\chi)}{|e|d^{4}x} = 12\xi\overline{\psi}\left[\alpha P_{R}(\overleftarrow{\nabla} - \nabla)P_{L} + \beta P_{L}(\overleftarrow{\nabla} - \nabla)P_{R}\right]\psi + 3\xi'\overline{\chi}\left[\alpha P_{R}(\overleftarrow{\nabla}' - \nabla')P_{L} + \beta P_{L}(\overleftarrow{\nabla}' - \nabla')P_{R}\right]\chi.$$
(41)

Hence, the physical Weyl spinors are defined by

$$\psi_L^{\text{phys}} = \sqrt{24\alpha\xi}\psi_L \,, \quad \psi_R^{\text{phys}} = \sqrt{24\beta\xi}\psi_R \,, \tag{42}$$

$$\overline{\psi}_L^{\text{phys}} = \sqrt{24\alpha\xi}\overline{\psi}_L \,, \quad \overline{\psi}_R^{\text{phys}} = \sqrt{24\beta\xi}\overline{\psi}_R \,, \tag{43}$$

and

$$\chi_L^{\rm phys} = \sqrt{6\alpha\xi'}\chi_L \,, \quad \chi_R^{\rm phys} = \sqrt{6\beta\xi'}\chi_R \,, \tag{44}$$

$$\overline{\chi}_L^{\text{phys}} = \sqrt{6\alpha\xi'} \overline{\chi}_L \,, \quad \overline{\chi}_R^{\text{phys}} = \sqrt{6\beta\xi'} \overline{\chi}_R \,. \tag{45}$$

In order to reproduce a chiral Lagrangian from (41) there are two possible scenarios. Firstly, one could assume the spinors in the gauge connection to be chiral,

$$P_R \psi = 0, \quad P_R \chi = 0, \tag{46}$$

then

$$\frac{\mathcal{L}_{K}(\psi) + \mathcal{L}_{K}(\chi)}{|e|d^{4}x} = \frac{1}{2}\overline{\psi}_{L}^{\text{phys}}(\overleftarrow{\nabla} - \nabla)\psi_{L}^{\text{phys}} + \frac{1}{2}\overline{\chi}_{L}^{\text{phys}}(\overleftarrow{\nabla}' - \nabla')\chi_{L}^{\text{phys}}, \quad (47)$$

where

$$\psi_L^{\text{phys}} = \sqrt{24\alpha\xi}\psi_L \,, \tag{48}$$

$$\overline{\psi}_L^{\text{phys}} = \sqrt{24\alpha\xi}\overline{\psi}_L\,,\tag{49}$$

and

$$\chi_L^{\rm phys} = \sqrt{6\alpha\xi'}\chi_L\,,\tag{50}$$

$$\overline{\chi}_L^{\rm phys} = \sqrt{6\alpha\xi'}\overline{\chi}_L \,. \tag{51}$$

Another possibility is to assume that $\beta \to 0$, then the gauge covariant kinetic term takes the same form, (47) and the physical Weyl spinors are

$$\psi_L^{\text{phys}} = \sqrt{24\alpha\xi}\psi_L \,, \quad \psi_R^{\text{phys}} = 0 \,, \tag{52}$$

$$\overline{\psi}_L^{\text{phys}} = \sqrt{24\alpha\xi}\overline{\psi}_L \,, \quad \overline{\psi}_R^{\text{phys}} = 0 \,, \tag{53}$$

and

$$\chi_L^{\text{phys}} = \sqrt{6\alpha\xi'}\chi_L \,, \quad \chi_R^{\text{phys}} = 0 \,, \tag{54}$$

$$\overline{\chi}_L^{\text{phys}} = \sqrt{6\alpha\xi'}\overline{\chi}_L \,, \quad \overline{\chi}_R^{\text{phys}} = 0 \,, \tag{55}$$

It can be seen that $\psi_R^{\rm phys} = 0$ and $\chi_R^{\rm phys} = 0$ is a consistent choice such that the bilinears that mix chiralities are well defined

$$\phi = \frac{1}{24\sqrt{\alpha\beta}} \phi^{\text{phys}} \,, \tag{56}$$

$$\phi_5 = \frac{1}{24\sqrt{\alpha\beta}}\phi_5^{\text{phys}} \tag{57}$$

$$\phi_{ab} = \frac{1}{24\sqrt{\alpha\beta}} \phi_{ab}^{\text{phys}}, \qquad (58)$$

where $\phi_X = \overline{\psi} \gamma_X \psi$. It remains an interesting question to address in the future to determine whether or not there are consistent alternatives to $\psi_R^{\text{phys}} = 0$ and $\chi_R^{\rm phys}=0\ {\rm when}\ \beta\to 0.$ Let us now show the connection to the SU(5) GG model. The covariant

derivatives in (47) expand as

$$\nabla \psi_L^{\text{phys}} = \nabla_{su(5)} \psi_L^{\text{phys}} - i g_{(U(1))}^{(\text{rank 1})} \mathcal{A} \psi_L^{\text{phys}}, \qquad (59)$$

$$\nabla' \chi_L^{\text{phys}} = \nabla_{su(5)} \chi_L^{\text{phys}} - ig A^{\tilde{X}} t_{\tilde{X}} \chi_L^{\text{phys}} - ig_{(U(1))}^{(\text{rank }2)} A \chi_L^{\text{phys}}, \qquad (60)$$

where the coupling constants are given in (32)-(34). The generators t_X in (60) carry a rank 2 antisymmetric representation of su(10), where t_X with X = I = $1, \dots, 24$ form a su(5) algebra. The index X runs for the range $= 25, \dots, 99$, see reference [17] or Appendix B for more details.

From the first terms in (59) and (60), the couplings between fermions and the gauge sector of the GG model can be recovered. We also get new gauge couplings not present in the GG model coming from the last terms in (59) and (60).

3.2Gauge potentials

In the GG model the gauge bosons of SU(5) transform in the adjoint representation that decomposes as

$$\mathbf{24} = (\mathbf{8}, \mathbf{1}, 0) + (\mathbf{1}, \mathbf{3}, 0) + (\mathbf{1}, \mathbf{1}, 0) + (\mathbf{3}, \mathbf{2}^*, -\frac{5}{6}) + (\mathbf{3}^*, \mathbf{2}, \frac{5}{6}).$$
(61)

The first three representations are the gluons, the W-bosons and the B-boson. The last two representations carry both color and flavor and are associated to colored and electrically charged gauge bosons X_i and Y_i (here $i \in SU(3)$) that form an SU(2) doublet. The hypercharge and electric charge generators are

$$Y = \text{diag}(-1/3, -1/3, -1/3, 1/2, 1/2), \qquad (62)$$

$$Q = \text{diag}(-1/3, -1/3, -1/3, 1, 0), \qquad (63)$$

when embedded in the SU(5) block of the superalgebra. When Q is embedded in the superalgebra, the charge assignments of the fermions can be verified using

$$[Q, \upsilon] = q(Q)\upsilon, \qquad (64)$$

where v is

$$\Psi = \mathbb{Q}_i \psi^i \quad \text{or} \quad \Phi = \overline{\mathbb{Q}}^{ij} \chi_{ij} \,. \tag{65}$$

It is convenient to visualize the full su(5) matrix in the following form

$$A^{I}(-i(t_{I})_{i}{}^{j}) = \frac{1}{\sqrt{2}}A_{i}{}^{j}, \qquad (66)$$

where

$$A_{i}{}^{j} = \begin{pmatrix} G_{1}{}^{1} - \frac{2B}{\sqrt{30}} & G_{1}{}^{2} & G_{1}{}^{3} & \overline{X}_{1} & \overline{Y}_{1} \\ G_{2}{}^{1} & G_{2}{}^{2} - \frac{2B}{\sqrt{30}} & G_{2}{}^{3} & \overline{X}_{2} & \overline{Y}_{2} \\ G_{3}{}^{1} & G_{3}{}^{2} & G_{3}{}^{3} - \frac{2B}{\sqrt{30}} & \overline{X}_{3} & \overline{Y}_{3} \\ \hline X^{1} & X^{2} & X^{3} & \frac{W^{3}}{\sqrt{2}} + \frac{3B}{\sqrt{30}} & W^{+} \\ Y^{1} & Y^{2} & Y^{3} & W^{-} & -\frac{W^{3}}{\sqrt{2}} + \frac{3B}{\sqrt{30}} \end{pmatrix}.$$
(67)

The electric charges of particles X and Y can also be verified using their embedding in the superconformal algebra, obtaining q(X) = 4/3 and q(Y) = 1/3.

The gauge bosons $A^{\tilde{X}}$ generate gauge kinetic couplings that are not present in the GG model (see (76)). Before carrying out the actual computation it is convenient to split the SU(10) block into the SU(5) block of the GG model and to also split the U(1) blocks of the superconformal algebra,

$$\overline{\psi}_{L} \nabla \psi_{L} + \overline{\chi}_{L} \nabla \chi_{L} = \overline{\psi}_{L} \nabla g_{su(5)} \psi_{L} + \overline{\chi}_{L} \nabla g_{su(5)} \chi_{L} - ig\overline{\chi}_{L} \mathcal{A}^{X} t_{\widetilde{X}} \chi_{L} - i\overline{\psi}_{L} g_{(U(1))}^{(\text{rank }1)} \mathcal{A} \psi_{L} - i\overline{\chi}_{L} g_{(U(1))}^{(\text{rank }2)} \mathcal{A} \chi_{L} ,$$
(68)

where we dropped the label of the physical spinor to simplify the notation. The gauge coupling in the first two terms of (68) has the standard form of the GG model,

$$\mathcal{L}_{G}^{GG} = g \left[\overline{u} \mathcal{G} u + \overline{d} \mathcal{G} d \right] + g \left[\left(\overline{u} \, \overline{d} \right)_{L} \mathcal{W} \left(\frac{\overline{u}}{d} \right)_{L} + \left(\overline{e}^{-} \, \overline{\nu}_{e} \right)_{L} \mathcal{W} \left(\frac{\overline{e}^{-}}{\overline{\nu}_{e}} \right)_{L} \right] + g \sqrt{\frac{3}{5}} \left[-\frac{1}{2} \left(\overline{e}_{L}^{-} \mathcal{B} e_{L}^{-} + \overline{\nu}_{eL} \mathcal{B} \nu_{eL} \right) + \frac{1}{6} \left(\overline{u}_{L} \mathcal{B} u_{L} + \overline{d}_{L} \mathcal{B} d_{L} \right) + \frac{2}{3} \overline{u}_{R} \mathcal{B} u_{R} - \frac{1}{3} \overline{d}_{R} \mathcal{B} d_{R} - \overline{e}_{R}^{+} \mathcal{B} e_{R}^{+} \right] + \frac{g}{\sqrt{2}} \left[\overline{d}_{R}^{i} \overline{X}_{i} e_{R}^{+} + \overline{d}_{L}^{i} \overline{X}_{i} e_{L}^{+} + \epsilon^{ijk} \overline{u}_{iL}^{c} \overline{X}_{j} u_{kL} \right] + \text{H.C.} + \frac{g}{\sqrt{2}} \left[-\overline{d}_{R}^{i} \overline{Y}_{i} \nu_{R}^{c} - \overline{u}_{L}^{i} \overline{Y}_{i} e_{L}^{+} + \epsilon^{ijk} \overline{u}_{iL}^{c} \overline{Y}_{j} d_{kL} \right] + \text{H.C.}$$
(69)

where

$$\mathcal{G} = \gamma^{\mu} G^{I}_{\mu} \left(-i\frac{\lambda_{I}}{2} \right) , \quad I \in SU(3)$$

$$\tag{70}$$

$$\Psi = \gamma^{\mu} W^{I}_{\mu} \left(-i\frac{\tau_{I}}{2} \right) , \quad I \in SU(2)$$
(71)

$$\not B = \gamma^{\mu} B_{\mu} \,, \tag{72}$$

$$\overline{X}_{i} = \gamma^{\mu} \overline{X}_{\mu i}, \quad \overline{Y}_{i} = \gamma^{\mu} \overline{Y}_{\mu i}, \qquad (73)$$

and λ^{I} and τ^{I} are Gell-Mann and Pauli matrices respectively.² One of the

²The extra minus sign in the definitions (70) and (71) w.r.t. reference [2] comes from the fact that we are using antihermitian generators $\lambda_I = i \lambda_I^{\text{(Gell-Mann)}}$, etc., which is motivated by the conventions in supergravity.

beautiful aspects of the GG model is the correct assignment of fractional quark charges, and that is also valid in the present model.

From (68) we also get new couplings that are not present in the GG model.

$$\mathcal{L}_{G}^{\text{new}} = -ig_{(U(1))}^{(\text{rank 1})} \left[\overline{d}_{L}^{c} \mathcal{A} d_{L}^{c} + \overline{e}_{L}^{-} \mathcal{A} \overline{e}_{L}^{-} + \overline{\nu}_{eL} \mathcal{A} \nu_{eL} \right]$$
(74)

$$-ig_{(U(1))}^{(\operatorname{rank} 2)} \left[\overline{u}_{L}^{c} \mathcal{A} u_{L}^{c} + \overline{u}_{L} \mathcal{A} u_{L} + \overline{d}_{L} \mathcal{A} d_{L} + \overline{e}_{L}^{+} \mathcal{A} e_{L}^{+} \right]$$
(75)

$$+ ig\overline{\chi}_L A^X t_{\widetilde{X}} \chi_L , \qquad (76)$$

The last line in (76) has the interactions mediated by the SU(10) gauge bosons that are not in the SU(5) block (67). The matrix elements of $t_{\tilde{X}}$ have been defined in [17] in no particular basis, as obtained by a Gramsh-Schmidt procedure. From the point of view of phenomenological applications it is useful to consider the following decomposition of the adjoint representation of SU(10)

$$99 = (24, 1, 0) + (1, 24, 0) + (1, 1, 0) + (5, 5^*, -y') + (5^*, 5, y').$$
(77)

The fields in (67) are uncharged with respect to the new (1, 24, 0) representation. There is also a quintet-quintet set of bosons charged respect to the (24, 1, 0) and the (1, 24, 0) representation. In addition, there is a new hypercharge-like operator associated to a new *B*-boson.

3.3 Comments on anomalies

As it happens in the GG model, anomaly cancellation of the cubic SU(3), SU(2)and hypercharge are preserved and received a rationale within the SU(5) block [2]. What seems to be problematic now are the extra $U(1)_Z$ groups produced by the generators along the diagonal in the superconformal algebra.

The $U(1)_Z$ charges are computed according to

$$[Z^{(\text{rank 1})}, \Psi] = x_1 \Psi, \quad [Z^{(\text{rank 2})}, \mathbf{X}] = x_2 \mathbf{X},$$
 (78)

where

$$x_1 = \left(\frac{-iz_1}{5}\right), \quad x_2 = \left(\frac{3iz_2}{5}\right), \tag{79}$$

and z_1 and z_2 are the overall constants of the $U(1)_Z$ generator in the fundamental and bi-fundamental representation, (91) and (129), respectively. In the following table we have a list of the quarks and leptons, hypercharges and Zcharges. Besides the cancellation of the mixed $(Y SU(3)^2)$, $(Y SU(2)^2)$ anomalies, and of the cancellation of the cubic anomaly Y^3 , we observe that the cubic Z^3 and the mixed anomalies $(Z SU(3)^2)$, $(Z SU(3)^2)$, cannot possibly all cancel simultaneously in the present model:

$$A_Z = 6x_2^3 + 2x_1^3 - \left(3x_2^3 + 3x_1^3 + x_2^3\right), \qquad (80)$$

$$A_{SU(2)} = 3x_2 + x_1 \,, \tag{81}$$

$$A_{SU(3)} = 2x_2 - (x_2 + x_1) . (82)$$

	SU(3)	SU(2)	Y	Z
Q_L	3	2	+1/6	x_2
L_L	1	2	-1/2	x_1
u_R	3	1	+2/3	x_2
d_R	3	1	-1/3	x_1
e_R	1	1	-1	x_2

Therefore, in order to avoid the anomaly induced by the Z symmetry we would have to invoke anomaly cancellation either between families, or by considering of left-right symmetric models such as the Pati-Salam, or by including new matter content [24].

4 Conclusions

In the present paper we have constructed a grand unified model using the superconformal algebra and matter in the adjoint representation of supersymmetry. The model reproduces the main features, charges and families of the Georgi-Glashow model, starting from an unconventional form of supersymmetry that includes gravity from the start [25, 18]. In this construction, local SUSY is not an invariance of the action but could be feature the ground state [19]. The model predicts a hidden sector $(1, 24, 0) + (5, 5^*, -y') + (5^*, 5, y')$ sector and an anomalous $U(1)_Z$.

A very interesting issue to address as a future work is the computation of the anomaly from the point of view of the embedding of the superconformal algebra, in terms of the symmetric tensor

$$d^{MNL} = \frac{1}{2} \langle T^M \{ T^N, T^L \} \rangle , \qquad (83)$$

where $\langle X \rangle$ stands for the supertrace of X, and also including the gradding associated to the operator \circledast [26, 27]. Such study could shed light into the question of existence of representations of the superconformal algebra that, perhaps adding new matter fields, could be anomaly free.

Another aspect of the model that we leave for a future study is the introduction of spontaneous symmetry breaking with a Higgs potential.

Acknowlegements

We thank Biancha Cerchiai, Laura Andrianopoli, Mario Trigiante and Lucrezia Ravera for useful discussions. P. A. acknowledges MINEDUC-UA project ANT 1755 and Semillero de Investigación project SEM18-02 from Universidad de Antofagasta, Chile. This research has been partially funded by Fondecyt Grant 1220862.

Appendices

A Fundamental representation of SU(2,2|n)

Let us consider the following representation of SU(2,2|n)

$$\mathbb{J}_{a} = \begin{bmatrix} \frac{\underline{s}}{2} \gamma_{a} & 0_{4 \times n} \\ \hline 0_{n \times 4} & 0_{n \times n} \end{bmatrix}, \quad \text{or} \quad (\mathbb{J}_{a})^{A}{}_{B} = \frac{\underline{s}}{2} (\gamma_{a})^{\alpha}{}_{\beta} \delta^{A}{}_{\alpha} \delta^{\beta}{}_{B} = \frac{\underline{s}}{2} (\gamma_{a})^{A}{}_{B}, \quad (84)$$

$$\mathbb{J}_{ab} = \begin{bmatrix} \frac{1}{4} [\gamma_a, \gamma_b] & 0_{4 \times n} \\ \hline & 0_{n \times 4} & 0_{n \times n} \end{bmatrix}, \quad \text{or} \quad (\mathbb{J}_{ab})^A_{\ B} = \frac{1}{4} [\gamma_a, \gamma_b]^A_{\ B} = (\Sigma_{ab})^A_{\ B}, \quad (85)$$

$$\mathbb{K}_{a} = \begin{bmatrix} \frac{1}{2}\tilde{\gamma}_{a} & 0_{4\times n} \\ \hline 0_{n\times 4} & 0_{n\times n} \end{bmatrix}, \quad \text{or} \quad (\mathbb{K}_{a})^{A}_{\ B} = \frac{1}{2}(\tilde{\gamma}_{a})^{A}_{\ B}, \tag{86}$$

$$\mathbb{D} = \begin{bmatrix} \frac{1}{2}\gamma_5 & 0_{4\times n} \\ \hline 0_{n\times 4} & 0_{n\times n} \end{bmatrix}, \quad \text{or} \quad (\mathbb{D})^A_{\ B} = \frac{1}{2}(\gamma_5)^A_{\ B}, \tag{87}$$

$$\mathbb{T}_{I} = \begin{bmatrix} 0_{4 \times 4} & 0_{4 \times n} \\ \hline 0_{n \times 4} & \frac{i}{2} \lambda_{I}^{t} \end{bmatrix}, \quad \text{or} \quad (\mathbb{T}_{I})_{B}^{A} = \frac{i}{2} (\lambda_{I}^{t})_{B}^{A},$$
(88)

$$\left(\mathbb{Q}_{i}^{\alpha}\right)_{B}^{A} = \left[\begin{array}{c|c} 0_{4\times4} & 0_{4\times n} \\ \hline \delta_{i}^{A}\delta_{B}^{\alpha} & 0_{n\times n} \end{array}\right] = \delta_{i}^{A}\delta_{B}^{\alpha} , \qquad (89)$$

$$(\overline{\mathbb{Q}}^{i}_{\alpha})^{A}_{\ B} = \begin{bmatrix} 0_{4\times4} & \delta^{A}_{\alpha}\delta^{i}_{B} \\ \hline & 0_{n\times4} & 0_{n\times n} \end{bmatrix} = \delta^{A}_{\alpha}\delta^{i}_{B},$$
(90)

$$\mathbb{Z}^{A}_{\ B} = z_1 \left[\frac{i\delta^{\alpha}_{\beta} \quad 0_{4 \times n}}{0_{n \times 4} \quad \frac{4}{n} i\delta^{i}_{j}} \right] = z_1 \left(i\delta^{A}_{\alpha}\delta^{\alpha}_{B} + \frac{4i}{n}\delta^{A}_{i}\delta^{i}_{B} \right) , \qquad (91)$$

where $\gamma_5 = i\gamma^0\gamma^1\gamma^2\gamma^3$, $(\gamma_5)^2 = \mathbb{1}$,

$$\gamma_{abc} = \gamma_{[a} \gamma_b \gamma_{c]} = \frac{1}{3!} \sum_{\Pi(a,b,c)} \operatorname{sign}(\Pi(a,b,c)) \gamma_a \gamma_b \gamma_c = i \epsilon_{abcd} \gamma_5 \gamma^d \,.$$

and

$$\tilde{\gamma}_a = rac{i}{3!} \epsilon_{abcd} \gamma^{bcd} = -\gamma_5 \gamma_a \,,$$

The γ -matrices are in a 4×4 spinor-representation (α, β, \cdots) run from 1 to 4). The indices of the tangent space a, b = 0, 1, 2, 3. Indices in the adjoint representation of su(n) take values $I, J = 1, 2, \ldots, n^2 - 1$, and in the fundamental take the values $i, j = 1, 2, \ldots, n$. The γ -matrices are endomorphisms and they act on spinors

$$\psi^{\alpha} \xrightarrow{\gamma_{a}} (\gamma_{a})^{\alpha}{}_{\beta}\psi^{\beta} .$$
(92)

These γ -matrices satisfy $\{\gamma^a, \gamma^b\} = 2\eta^{ab}$, where the metric η is given by $\eta = \text{diag}(-, +, +, +)$. The spinor indices will be often omitted.

In a similar way the $\lambda\text{-matrices}$ are also endomorphisms and they act on spinors as

$$\psi_i^{\alpha} \xrightarrow{\lambda_I} (\lambda_I)_i^{\ j} \psi_j^{\alpha} \,. \tag{93}$$

The λ -matrices satisfy $[\lambda_I, \lambda_J] = f^{IJK} \lambda_K$.

Indices of the representation are $A, B = 1, \dots, n+4$, so we have a $(n+4) \times (n+4)$ representation. We find it convenient to use the splitting $A = (\alpha, i)$. All the possible products that mix tensors from different spaces, like $p^i_A q^A_{\alpha}$, are trivial. The following relations are understood

$$(\gamma_a)^A_{\ B} = \delta^A_\alpha(\gamma_a)^\alpha_{\ \beta} \delta^\beta_B \,, \tag{94}$$

$$C_{\alpha A} = C_{\alpha \beta} \delta_A^\beta \,. \tag{95}$$

The generators J_a and J_{ab} form an adS_4 algebra,

$$[\mathbf{J}_a, \mathbf{J}_b] = s^2 \mathbf{J}_{ab} \,, \tag{96}$$

$$[\mathbf{J}_a, \mathbf{J}_{bc}] = \eta_{ab} \mathbf{J}_c - \eta_{ac} \mathbf{J}_b \,, \tag{97}$$

$$[\mathbf{J}_{ab}, \mathbf{J}_{cd}] = -(\eta_{ac} \mathbf{J}_{bd} - \eta_{ad} \mathbf{J}_{bc} - \eta_{bc} \mathbf{J}_{ad} + \eta_{bd} \mathbf{J}_{ac}).$$
(98)

The parameter s^2 can take values $s^2 = +1, -1$ for anti-de Sitter or de Sitter algebras, respectively.

Among \mathbb{D} and \mathbb{K}_a they form the conformal algebra,

$$[\mathbb{K}_a, \mathbb{K}_b] = -\mathbb{J}_{ab} \,, \tag{99}$$

$$[\mathbf{J}_a, \mathbf{K}_b] = s\eta_{ab} \mathbb{D} \,, \tag{100}$$

$$[\mathbb{K}_a, \mathbb{J}_{bc}] = \eta_{ab} \mathbb{K}_c - \eta_{ac} \mathbb{K}_b \,, \tag{101}$$

$$[\mathbb{D}, \mathbb{K}_a] = -s^{-1} \mathbb{J}_a \,, \tag{102}$$

$$[\mathbb{D}, \mathbb{J}_a] = -s\mathbb{K}_a \,. \tag{103}$$

For the internal generators we have the su(n) algebra

$$[\mathbb{T}_I, \mathbb{T}_J] = f_{IJ}{}^K \mathbb{T}_K \,, \tag{104}$$

and they are anti-hermitian $\mathbb{T}_{I}^{\dagger} = -\mathbb{T}_{I}$ (also $\mathbb{Z}^{\dagger} = -\mathbb{Z}$).

Including \mathbb{Q}_i^{α} and $\overline{\mathbb{Q}}_{\alpha}^i$ the commutators close in a su(2,2|n) superalgebra

$$[\mathbf{J}_{a}, \overline{\mathbf{Q}}_{\alpha}^{i}] = \frac{s}{2} \overline{\mathbf{Q}}_{\beta}^{i} (\gamma_{a})^{\beta}{}_{\alpha}, \quad [\mathbf{J}_{a}, \mathbf{Q}_{i}^{\alpha}] = -\frac{s}{2} (\gamma_{a})^{\alpha}{}_{\beta} \mathbf{Q}_{i}^{\beta}, \tag{105}$$

$$[\mathbf{J}_{ab}, \overline{\mathbb{Q}}_{\alpha}^{i}] = \overline{\mathbb{Q}}_{\beta}^{i} (\Sigma_{ab})_{\alpha}^{\beta}, \quad [\mathbf{J}_{ab}, \mathbb{Q}_{i}^{\alpha}] = -(\Sigma_{ab})_{\beta}^{\alpha} \mathbb{Q}_{i}^{\beta}, \tag{106}$$

$$[\mathbb{K}_{a}, \overline{\mathbb{Q}}_{\alpha}^{i}] = \frac{1}{2} \overline{\mathbb{Q}}_{\beta}^{i} (\tilde{\gamma}_{a})_{\alpha}^{\beta}, \quad [\mathbb{K}_{a}, \mathbb{Q}_{i}^{\alpha}] = -\frac{1}{2} (\tilde{\gamma}_{a})_{\beta}^{\alpha} \mathbb{Q}_{i}^{\beta}, \quad (107)$$

$$\left[\mathbb{D}, \overline{\mathbb{Q}}_{\alpha}^{i}\right] = \frac{1}{2} \overline{\mathbb{Q}}_{\beta}^{i} (\gamma_{5})^{\beta}{}_{\alpha}, \quad \left[\mathbb{D}, \mathbb{Q}_{i}^{\alpha}\right] = -\frac{1}{2} (\gamma_{5})^{\alpha}{}_{\beta} \mathbb{Q}_{i}^{\beta}, \quad (108)$$

$$[\mathbb{T}_I, \overline{\mathbb{Q}}^i_{\alpha}] = -\frac{i}{2} \overline{\mathbb{Q}}^j_{\alpha} (\lambda_I)_j^{\ i}, \quad [\mathbb{T}_I, \mathbb{Q}^{\alpha}_i] = \frac{i}{2} (\lambda_I)_i^{\ j} \mathbb{Q}^{\alpha}_j, \tag{109}$$

$$[\mathbb{Z}, \overline{\mathbb{Q}}_{\alpha}^{i}] = -iz_{1}(4/n-1)\overline{\mathbb{Q}}_{\alpha}^{i}, \quad [\mathbb{Z}, \mathbb{Q}_{i}^{\alpha}] = iz_{1}(4/n-1)\mathbb{Q}_{i}^{\alpha}, \qquad (110)$$

$$\{\mathbb{Q}_{i}^{\alpha}, \overline{\mathbb{Q}}_{\beta}^{j}\} = (\gamma^{C})^{\alpha}{}_{\beta}\delta_{i}^{j}\mathbb{J}_{C} + \delta_{\beta}^{\alpha}\left(-i(\lambda_{I})_{i}{}^{j}\mathbb{T}_{I} - \frac{i}{4z_{1}}\delta_{i}^{j}\mathbb{Z}\right), \qquad (111)$$

where we have used the short-hand

$$(\gamma^C)^{\alpha}{}_{\beta}\mathbb{J}_C = \left(\frac{1}{2s}(\gamma^a)^{\alpha}{}_{\beta}\mathbb{J}_a - \frac{1}{2}(\Sigma^{ab})^{\alpha}{}_{\beta}\mathbb{J}_{ab} - \frac{1}{2}(\tilde{\gamma}^a)^{\alpha}{}_{\beta}\mathbb{K}_a + \frac{1}{2}(\gamma_5)^{\alpha}{}_{\beta}\mathbb{D}\right).$$
(112)

Traces:

The gradding operator is given by

$$\mathcal{G}^{A}_{\ B} = \delta^{A}_{\alpha} \delta^{\alpha}_{\ B} - \delta^{A}_{i} \delta^{i}_{\ B} \,, \tag{113}$$

which classifies generators in bosonic $B = \{ \mathbb{J}_a, \mathbb{J}_{ab}, \mathbb{K}_a, \mathbb{D}, \mathbb{T}_I, \mathbb{Z} \}$ or fermionic $F = \{ Q_i^{\alpha}, \overline{\mathbb{Q}}_{\alpha}^i \}$, by $[B, \mathcal{G}] = 0 = \{F, \mathcal{G}\}$, and squares to one, $\mathcal{G}^2 = 1$. The gradding operator defines an invariant supertrace

$$\langle G \rangle \equiv Tr(\mathcal{G}G) = 0. \tag{114}$$

The supertrace has the following properties

$$\langle B_1 B_2 \rangle = \langle B_2 B_1 \rangle, \quad \langle BF \rangle = \langle FB \rangle, \quad \langle F_1 F_2 \rangle = -\langle F_2 F_1 \rangle.$$
 (115)

All generators ${\cal G}$ in the representation are super-traceless

$$\langle G \rangle = 0, \quad G = \{ \mathbf{J}_a, \mathbf{J}_{ab}, \mathbf{K}_a, \mathbb{D}, \mathbb{T}_I, \mathbb{Z}, Q_i^{\alpha}, \overline{\mathbb{Q}}_{\alpha}^{\imath} \}.$$
 (116)

The quadratic combinations that give nontrivial traces are

$$\langle \mathbf{J}_a \mathbf{J}_b \rangle = s^2 \eta_{ab} , \qquad \langle \mathbf{J}_{ab} \mathbf{J}_{cd} \rangle = -(\eta_{ac} \eta_{bd} - \eta_{bc} \eta_{ad}) , \qquad (117)$$

$$\langle \mathbb{K}_a \mathbb{K}_b \rangle = -\eta_{ab} , \qquad \langle \mathbb{D}^2 \rangle = +1 , \qquad (118)$$

$$\langle \mathbb{T}_I \mathbb{T}_J \rangle = \frac{1}{2} \delta_{IJ} , \qquad \langle \mathbb{Z}^2 \rangle = 4z_1^2 (4/n - 1) , \qquad (119)$$

$$\langle \mathbb{Q}_i^{\alpha} \overline{\mathbb{Q}}_{\beta}^{j} \rangle = -\delta_{\beta}^{\alpha} \delta_i^{j} = -\langle \overline{\mathbb{Q}}_{\beta}^{j} \mathbb{Q}_i^{\alpha} \rangle \,. \tag{120}$$

S-grading operator

The S operator is essential for the generalized \circledast dual opperator, as in (12). The nontrivial traces involving S are

$$\langle S\mathbb{D}\rangle = 2i\varepsilon_s\,,\tag{121}$$

$$\langle S \mathbb{J}_{ab} \mathbb{J}_{cd} \rangle = -\varepsilon_s \epsilon_{abcd} = \langle \mathbb{J}_{ab} S \mathbb{J}_{cd} \rangle, \qquad (122)$$

$$\langle \mathbf{J}_a S \mathbb{K}_b \rangle = -i\varepsilon_s s \eta_{ab} = -\langle \mathbb{K}_a S \mathbf{J}_b \rangle, \qquad (123)$$

$$\langle \mathbb{Z}S\mathbb{D} \rangle = -2z_1 \varepsilon_s = \langle \mathbb{D}S\mathbb{Z} \rangle.$$
 (124)

These traces are instrumental in producing the usual expressions for the standard kinetic terms in the action.

B Bi-fundamental representation of SU(2,2|n)

The generators of the conformal group SU(2,2), $\{\mathbb{J}_C\} = \{\mathbb{J}_a, \mathbb{J}_{ab}, \mathbb{K}_a, \mathbb{D}\}$, embedded in the superalgebra are:

$$\mathbf{J}_{a} = \begin{bmatrix} \frac{s}{2}\gamma_{a} & 0_{4\times d_{n}} \\ \hline 0_{d_{n}\times 4} & 0_{d_{n}\times d_{n}} \end{bmatrix}, \quad \text{or} \quad (\mathbf{J}_{a})^{A}_{\ B} = \frac{s}{2}(\gamma_{a})^{\alpha}_{\ \beta}\delta^{A}_{\ \alpha}\delta^{\beta}_{\ B} = \frac{s}{2}(\gamma_{a})^{A}_{\ B},$$
(125)

$$\mathbf{J}_{ab} = \begin{bmatrix} \frac{1}{4} [\gamma_a, \gamma_b] & 0_{4 \times d_n} \\ \hline 0_{d_n \times 4} & 0_{d_n \times d_n} \end{bmatrix}, \quad \text{or} \quad (\mathbf{J}_{ab})^A_{\ B} = \frac{1}{4} [\gamma_a, \gamma_b]^A_{\ B} = (\Sigma_{ab})^A_{\ B},$$
(126)

$$\mathbb{K}_{a} = \begin{bmatrix} \frac{1}{2}\tilde{\gamma}_{a} & 0_{4\times d_{n}} \\ \hline 0_{d_{n}\times 4} & 0_{d_{n}\times d_{n}} \end{bmatrix}, \quad \text{or} \quad (\mathbb{K}_{a})_{\ B}^{A} = \frac{1}{2}(\tilde{\gamma}_{a})_{\ B}^{A}, \tag{127}$$

$$\mathbb{D} = \begin{bmatrix} \frac{1}{2}\gamma_5 & 0_{4\times d_n} \\ \hline 0_{d_n\times 4} & 0_{d_n\times d_n} \end{bmatrix}, \quad \text{or} \quad (\mathbb{D})^A_{\ B} = \frac{1}{2}(\gamma_5)^A_{\ B}, \qquad (128)$$

$$\mathbb{Z}^{A}_{\ B} = iz_2 \left(\delta^A_\alpha \delta^\alpha_B + \frac{4}{d_n} \Delta^A_{ij} \Delta^{ij}_B \right) \,. \tag{129}$$

The following commutators get modified with respect to the fundamental representation, and they close in $su(2, 2|d_n)$:

$$[\mathbb{G}_{X'},\mathbb{G}_{Y'}] = f_{X'Y'}{}^{Z'}\mathbb{G}_{Z'}, \qquad (130)$$

and

$$[\mathbb{G}_{X'}, \mathbb{Q}_{ij}^{\alpha}] = i(g_{X'})_{ij}{}^{kl}\mathbb{Q}_{kl}^{\alpha}, \quad [\mathbb{G}_{X'}, \overline{\mathbb{Q}}_{\alpha}^{ij}] = -i\overline{\mathbb{Q}}_{\alpha}^{kl}(g_{X'})_{kl}{}^{ij}, \qquad (131)$$

$$\left[\mathbb{Z}, \mathbb{Q}_{ij}^{\alpha}\right] = i z_2 \left(4/d_n - 1\right) \mathbb{Q}_{ij}^{\alpha}, \quad \left[\mathbb{Z}, \overline{\mathbb{Q}}_{\alpha}^{ij}\right] = -i z_2 \left(4/d_n - 1\right) \overline{\mathbb{Q}}_{\alpha}^{ij}, \quad (132)$$

$$\{\mathbb{Q}_{ij}^{\alpha}, \overline{\mathbb{Q}}_{\beta}^{kl}\} = (\gamma^C)^{\alpha}{}_{\beta}\Delta_{ij}^{kl}\mathbb{J}_C - \left(\frac{4i}{n-2}(g_{X'})_{ij}{}^{kl}\mathbb{G}_{X'} + \frac{i}{2z_2}\Delta_{ij}^{kl}\mathbb{Z}\right)\delta_{\beta}^{\alpha}.$$
 (133)

(134)

The generators $g_{X'}$ and v_X , that belong to the bifundamental rep. of SU(n)and the fundamental rep. of $SU(d_n)$, are related to each other by means of the change of basis matrix defined by

$$g_{X'} = C_{X'}{}^X v_X \,. \tag{135}$$

We will define the inverse matrix by

$$v_X = C_X{}^{X'}g_{X'}, (136)$$

which satisfies

$$C_X{}^{X'}C_X{}^{Y'} = \frac{1}{n-2}\delta^{X'Y'}.$$
(137)

Let us define the structure constants of $\mathfrak{su}(d_n)$ in the fundamental representation by

$$[v_X, v_Y] = iF_{XY}{}^Z v_Z \,. \tag{138}$$

Using the matrix $C_X{}^{X'}$ and its inverse we can determine the structure constants $f_{X'Y'}{}^{Z'}$, that are defined in the $g_{X'}$ basis

$$[g_{X'}, g_{Y'}] = i f_{X'Y'} {}^{Z'} g_{Z'}, \qquad (139)$$

where

$$f_{X'Y'}{}^{Z'} = C_{X'}{}^{X}C_{Y'}{}^{Y}F_{XY}{}^{Z}C_{Z}{}^{Z'}.$$
(140)

The quadratic supertraces are given by (117)-(118) and

$$\langle \mathbb{G}_{X'} \mathbb{G}_{Y'} \rangle = \frac{n-2}{2} \delta_{X'Y'}, \qquad \langle \mathbb{Z}^2 \rangle = 4z_2^2 (4/d_n - 1),$$
(141)

$$\langle \mathbb{Q}_{ij}^{\alpha} \overline{\mathbb{Q}}_{\beta}^{kl} \rangle = -\delta_{\beta}^{\alpha} \Delta_{ij}^{kl} = -\langle \overline{\mathbb{Q}}_{\beta}^{j} \mathbb{Q}_{i}^{\alpha} \rangle.$$
(142)

C SU(2,2|n) curvatures

Here we give the explicity expressions of the curvatures.

The field strength is defined by

$$\mathbb{F} = \frac{1}{2} \mathcal{F}^{ab} \mathbb{J}_{ab} + \mathcal{F}^a \mathbb{J}_a + \mathcal{G}^a \mathbb{K}_a + \mathcal{H} \mathbb{D} + \mathcal{F}^I \mathbb{T}_I + \mathcal{F} \mathbb{Z} + \overline{\mathbb{Q}}^i_\alpha \mathcal{X}^\alpha_i + \overline{\mathcal{X}}^i_\alpha \mathbb{Q}^\alpha_i, \quad (143)$$

$$\mathcal{F}^{a} = Df^{a} + \frac{1}{s}g^{a}h + \frac{1}{2s}\overline{\psi}^{i}\phi\gamma^{a}\phi\psi_{i}, \qquad (145)$$

$$\mathcal{G}^{a} = Dg^{a} + sf^{a}h - \frac{1}{2}\overline{\psi}^{i} \not\in \tilde{\gamma}^{a} \not\in \psi_{i} , \qquad (146)$$

$$\mathcal{H} = H + sf^a g_a + \frac{1}{2} \overline{\psi}^i \not e \gamma_5 \not e \psi_i , \qquad (147)$$

$$\mathcal{F} = F - \frac{i}{4z} \overline{\psi}^i \phi \phi_i \,, \tag{149}$$

$$\mathcal{X}_{i}^{\alpha} = D(\not\!\!/\psi_{i})^{\alpha} + \frac{s}{2} f^{a} (\gamma_{a} \not\!/\psi_{i})^{\alpha} + \frac{1}{2} g^{a} (\tilde{\gamma}_{a} \not\!/\psi_{i})^{\alpha} + \frac{1}{2} h (\gamma_{5} \not\!/\psi_{i})^{\alpha}, \qquad (150)$$

where

$$H = dh, \qquad (152)$$

$$\mathcal{R}^{ab} = R^{ab} + s^2 f^a f^b - g^a g^b \,, \tag{153}$$

$$R^{ab} = d\omega^{ab} + \omega^a_{\ c} \omega^{cb} \,, \tag{154}$$

$$F^{I} = dA^{I} + \frac{1}{2} f^{I}_{JK} A^{J} A^{K} , \qquad (155)$$

$$F = dA. (156)$$

The covariant derivative on a Lorentz vector V^a is

$$DV^a = dV^a + \omega^a_{\ b} V^b \,, \tag{157}$$

and on a spinor

$$D\psi^{\alpha} = d\psi^{\alpha} + \frac{1}{2}\omega^{ab}(\Sigma_{ab}\psi)^{\alpha} - \frac{i}{2}A^{I}(\lambda_{I}\psi)^{\alpha}_{i} - iz(4/N-1)A\psi^{\alpha}_{i}.$$
 (158)

The covariant derivative D is defined for the $SO(1,3) \times SU(n) \times U(1)$ connection. The left-acting exterior derivative satisfies $\Omega^m \overleftarrow{d} = (-1)^m d\Omega^m$ for an m-form, in the spinor representation yields the relation (158).

References

- H. Georgi and S. L. Glashow, "Unity of All Elementary Particle Forces," *Phys. Rev. Lett.*, vol. 32, pp. 438–441, 1974.
- [2] P. Langacker, "Grand Unified Theories and Proton Decay," *Phys. Rept.*, vol. 72, p. 185, 1981.
- [3] S. Dimopoulos and H. Georgi, "Softly Broken Supersymmetry and SU(5)," Nucl. Phys. B, vol. 193, pp. 150–162, 1981.
- [4] S. M. Barr, "A New Symmetry Breaking Pattern for SO(10) and Proton Decay," *Phys. Lett. B*, vol. 112, pp. 219–222, 1982.
- [5] I. Antoniadis, J. R. Ellis, J. S. Hagelin, and D. V. Nanopoulos, "Supersymmetric Flipped SU(5) Revitalized," *Phys. Lett. B*, vol. 194, pp. 231–235, 1987.
- [6] J. R. Ellis, J. L. Lopez, and D. V. Nanopoulos, "The Prospects for CHO-RUS and NOMAD in the light of COBE and GALLEX," *Phys. Lett. B*, vol. 292, pp. 189–194, 1992.
- [7] J. R. Ellis, J. L. Lopez, D. V. Nanopoulos, and K. A. Olive, "Flipped angles and phases: A Systematic study," *Phys. Lett. B*, vol. 308, pp. 70–78, 1993.

- [8] J. R. Ellis, D. V. Nanopoulos, and K. A. Olive, "Flipped heavy neutrinos: From the solar neutrino problem to baryogenesis," *Phys. Lett. B*, vol. 300, pp. 121–127, 1993.
- [9] I. Antoniadis, J. R. Ellis, J. S. Hagelin, and D. V. Nanopoulos, "An Improved SU(5) x U(1) Model from Four-Dimensional String," *Phys. Lett. B*, vol. 208, pp. 209–215, 1988. [Addendum: Phys.Lett.B 213, 562 (1988)].
- [10] B. A. Campbell, J. R. Ellis, J. S. Hagelin, D. V. Nanopoulos, and R. Ticciati, "Flipped SU(5) From Manifold Compactification of the Tendimensional Heterotic String," *Phys. Lett. B*, vol. 198, pp. 200–204, 1987.
- [11] J. L. Lopez, D. V. Nanopoulos, and K.-j. Yuan, "The Search for a realistic flipped SU(5) string model," Nucl. Phys. B, vol. 399, pp. 654–690, 1993.
- [12] J. R. Ellis, D. V. Nanopoulos, and J. Walker, "Flipping SU(5) out of trouble," *Phys. Lett. B*, vol. 550, pp. 99–107, 2002.
- [13] J. Ellis, M. A. G. Garcia, N. Nagata, D. V. Nanopoulos, and K. A. Olive, "Starobinsky-like Inflation, Supercosmology and Neutrino Masses in No-Scale Flipped SU(5)," JCAP, vol. 07, p. 006, 2017.
- [14] S. Ferrara, R. Kallosh, A. Linde, and M. Porrati, "Minimal Supergravity Models of Inflation," *Phys. Rev. D*, vol. 88, no. 8, p. 085038, 2013.
- [15] S. Ferrara, M. Kaku, P. K. Townsend, and P. van Nieuwenhuizen, "Gauging the Graded Conformal Group with Unitary Internal Symmetries," *Nucl. Phys. B*, vol. 129, pp. 125–134, 1977.
- [16] M. Kaku, P. K. Townsend, and P. van Nieuwenhuizen, "Superconformal Unified Field Theory," *Phys. Rev. Lett.*, vol. 39, p. 1109, 1977.
- [17] P. D. Alvarez, R. A. Chavez, and J. Zanelli, "Embedding of rank two supercharges in the superconformal algebra," *J. Math. Phys.*, vol. 63, no. 4, p. 042304, 2022.
- [18] P. D. Alvarez, R. A. Chavez, and J. Zanelli, "Gauging the superconformal group with a graded dual operator," *JHEP*, vol. 02, p. 111, 2022.
- [19] P. D. Alvarez, L. Delage, M. Valenzuela, and J. Zanelli, " $\mathcal{N} = 2$ extended MacDowell-Mansouri supergravity," *JHEP*, vol. 07, p. 176, 2021.
- [20] S. W. MacDowell and F. Mansouri, "Unified Geometric Theory of Gravity and Supergravity," *Phys. Rev. Lett.*, vol. 38, p. 739, 1977. [Erratum: Phys.Rev.Lett. 38, 1376 (1977)].
- [21] L. Castellani, R. Catenacci, and P. A. Grassi, "The Geometry of Supermanifolds and New Supersymmetric Actions," *Nucl. Phys. B*, vol. 899, pp. 112– 148, 2015.

- [22] L. Castellani, R. Catenacci, and P. A. Grassi, "Hodge Dualities on Supermanifolds," Nucl. Phys. B, vol. 899, pp. 570–593, 2015.
- [23] P. Arias, D. Cadamuro, M. Goodsell, J. Jaeckel, J. Redondo, and A. Ringwald, "WISPy Cold Dark Matter," *JCAP*, vol. 06, p. 013, 2012.
- [24] A. Bilal, "Lectures on Anomalies," 2 2008.
- [25] P. D. Alvarez, P. Pais, and J. Zanelli, "Unconventional supersymmetry and its breaking," *Phys. Lett. B*, vol. 735, pp. 314–321, 2014.
- [26] Z. Wan and J. Wang, "Beyond Standard Models and Grand Unifications: Anomalies, Topological Terms, and Dynamical Constraints via Cobordisms," JHEP, vol. 07, p. 062, 2020.
- [27] J. Davighi, B. Gripaios, and N. Lohitsiri, "Global anomalies in the Standard Model(s) and Beyond," JHEP, vol. 07, p. 232, 2020.