

On Dynamical Parameter Space of Cubic Polynomials with a Parabolic Fixed Point

Runze Zhang

Abstract

This article focus on the connected locus of the cubic polynomial slice $Per_1(\lambda)$ with a parabolic fixed point of multiplier $\lambda = e^{2\pi i \frac{p}{q}}$. We first show that any parabolic component, which is a parallel notion of hyperbolic component, is a Jordan domain. Moreover, a continuum \mathcal{K}_λ called the central part in the connected locus is defined. This is the natural analogue to the closure of the main hyperbolic component of $Per_1(0)$. We prove that \mathcal{K}_λ is almost a double covering of the filled-in Julia set of the quadratic polynomial $P_\lambda(z) = \lambda z + z^2$.

1 Introduction

The dynamics of iteration of degree $d \geq 2$ complex polynomials acting on the complex plane is very rich. As a consequence, the connected locus $\mathcal{C}^d := \{f \in \mathcal{P}^d; \text{ the Julia set of } f \text{ is connected.}\}$ presents complicate fractal structures, where $\mathcal{P}^d \cong \mathbb{C}^{d-1}$ is the parameter space of degree d polynomials. When $d = 2$, $\mathcal{C}^2 \subset \mathbb{C}$ is known as the Mandelbrot set \mathbf{M} . The famous MLC conjecture asserting that \mathbf{M} is locally connected is a central problem in the study of dynamics of quadratic polynomials, since it is equivalent to the combinatorial rigidity conjecture and will imply the density of hyperbolicity conjecture. It also implies that \mathbf{M} has a topological model of a certain quotient of the closed unit disk. However when d augments, \mathcal{C}^d exhibits much more sophisticated structures and the analogue of MLC conjecture for \mathcal{C}^3 turns out to be false [11]. For this reason, Milnor suggested to investigate \mathcal{C}^3 by restricting the parameter space to complex 1-dimensional algebraic curves by adding dynamical conditions. In fact, when restricting to 1-dimensional slices, the local connectivity usually holds for non-renormalisable or finitely renormalisable parameters ([16], [20], [2]). This paper will focus on $Per_1(\lambda)$, the 1-dimensional slice of the cubic polynomials which have a parabolic fixed point with multiplier $\lambda = e^{2\pi i \frac{p}{q}}$ (p, q coprime). We aim at investigating local connectivity at certain parameters in $\mathcal{C}_\lambda := \mathcal{C}^3 \cap Per_1(\lambda)$ and giving a global description of \mathcal{C}_λ . As a byproduct, we prove that \mathcal{C}_λ presents a big patch called "the central part" which is almost a double covering of the filled-in Julia set of $P_\lambda(z) = \lambda z + z^2$.

More precisely, consider the space of unitary cubic polynomials fixing the origin 0:

$$f_{\lambda,a}(z) = \lambda z + az^2 + z^3, \quad (\lambda, a) \in \mathbb{C}^2. \quad (1)$$

Notice that every cubic polynomial is affinely conjugated to one of the polynomials of form (1). By fixing λ , one gets the slice

$$Per_1(\lambda) := \{f_{\lambda,a}; a \in \mathbb{C}\} \cong \mathbb{C}$$

and its corresponding connected locus $\mathcal{C}_\lambda := \{a \in \mathbb{C}; J_{\lambda,a} \text{ is connected}\}$, where $J_{\lambda,a}$ is the Julia set of $f_{\lambda,a}$. It is a classical result that the Julia set of a polynomial is connected if and only if none of its critical points escapes to infinity. Thus one also has

$$\mathcal{C}_\lambda = \{a \in \mathbb{C}; \text{both critical points of } f_{\lambda,a} \text{ do not escape to } \infty\}.$$

When $|\lambda| < 1$. (resp. $\lambda = e^{\frac{2\pi ip}{q}}$), 0 is an attracting (resp. parabolic) fixed point. By classical results in holomorphic dynamics, 0 attracts at least one of the two critical points of $f_{\lambda,a}$. For each λ fixed, it is natural to consider the following attracting locus:

$$\mathcal{H}^\lambda := \{a \in \mathbb{C}; \text{ both critical points of } f_{\lambda,a} \text{ are attracted by } 0\}. \quad (2)$$

Definition. Let $|\lambda| < 1$. resp. $\lambda = e^{\frac{2\pi ip}{q}}$. The **central part** \mathcal{K}_λ of \mathcal{C}_λ is defined to be the connected component of $\overline{\mathcal{H}_\lambda}$ containing $a = 0$.

Our main result is the following:

Main Theorem. Let $\lambda = e^{2\pi i \frac{p}{q}}$. Every connected component of $\mathring{\mathcal{H}}^\lambda$ is a Jordan domain. The central part \mathcal{K}_λ is a full continuum and is locally connected. Moreover, there exists a dynamically defined double covering:

$$\mathfrak{G} : \mathcal{K}_\lambda \setminus \mathcal{I} \longrightarrow K_\lambda \setminus \bigcup_{i=0}^{q-1} \overline{P_i}$$

where $\mathcal{I} \subset \mathcal{K}_\lambda$ is a curve passing $a = 0$; K_λ the filled-in Julia set of $P_\lambda(z) = \lambda z + z^2$ and $(P_i)_i$ are q petals contained respectively in the q immediate basins of $P_\lambda(z)$.

The Main Theorem can also be stated for $|\lambda| < 1$ and is proved in [16] ($\lambda = 0$) and [19] ($0 < |\lambda| < 1$). Our result generalizes theirs to all parabolic slices. In a recent paper by A. Blokh, L. Oversteegen and V. Timorin [1], a global description of \mathcal{C}_λ for $|\lambda| \leq 1$ is given by decomposing it into a full continuum \mathcal{CU}_λ called "main cuboid" and the limbs attached to it, where they define \mathcal{CU}_λ to be the collection of all $f_{\lambda,a}$ verifying the following property:

$f_{\lambda,a}$ has a non-repelling fixed point, $f_{\lambda,a}$ has no repelling periodic cutpoints in $J_{\lambda,a}$, and all non-repelling periodic points of $f_{\lambda,a}$, except at most one fixed point, have multiplier 1.

As a byproduct of the proof of our Main Theorem, we are able to obtain a more visualisable description of \mathcal{CU}_λ :

Corollary I. For $\lambda = e^{2\pi ip/q}$ we have $\mathcal{K}_\lambda = \mathcal{CU}_\lambda$.

Before illustrating the strategy of the proof, let us say a few more words about \mathcal{H}^λ . Let c_1, c_2 be the two critical points of $f_{\lambda,a}$. Following Milnor [15], parameters in \mathcal{H}^λ are divided into four different types:

- (A) Adjacent: c_1, c_2 are contained in the same Fatou component.
- (B) Bitransitive: c_1, c_2 belong to two different periodic Fatou components in the same cycle.
- (C) Capture: one of c_1, c_2 eventually hits a periodic Fatou component.
- (D) Disjoint: c_1, c_2 belong to different cycles of periodic Fatou components.
- (MP) Misiurewicz parabolic: one of c_1, c_2 eventually hits the parabolic fixed point $z = 0$.

Definition. Let $\lambda = e^{2\pi i \frac{p}{q}}$. Type (D) parameters are exactly those $a \in \mathcal{C}_\lambda$ such that $z = 0$ becomes parabolic degenerate. We also call them **double parabolic** parameters, and denote by $\mathcal{A}_{p/q}$ the collection of them. A connected component of Type (A) (B) or (C) is usually called a **parabolic component**.

Strategy of the proof. Let $\lambda = e^{2\pi i \frac{p}{q}}$. To prove the Main Theorem, it is natural to begin with building an identification between $\mathring{\mathcal{K}}_\lambda$ and \mathring{K}_λ : intuitively, Type (A) and (B) corresponds to the immediate basins of P_λ ; Type (C) in $\mathring{\mathcal{K}}_\lambda$ corresponds to infinite towers of preimages of the immediate basins attached on their boundaries (see Figure 1 to have a global picture of \mathcal{K}_λ in mind). This can be done by giving dynamical parametrisations to $\mathring{\mathcal{H}}_\lambda$. The hard part of the Main Theorem is to extend this identification continuously to

$\partial\mathcal{K}_\lambda$, where the need of local connectivity naturally arises. The strategy for the proof of local connectivity of $\partial\mathcal{K}_\lambda$ is to use the "dynamical-parameter puzzle" technique: transfer the shrink property of dynamical puzzles to that of parameter puzzles. However, comparing to the super-attracting case ([7], [16]) where this technique is applied, there are several difficulties that we need to overcome:

- A priori, the existence of Type (A) (B) (D) parameters with given combinatorics (namely, the dynamical gap between two critical points) is not obvious. Based on Cui-Tan's pinching deformation theory [5] (Appendix B), we are able to fully solve this question.
- The parametrisation of Type (A) (B) components is more complicate. We need to deal with $f_{\lambda,a}$ such that the boundary of its maximal Fatou petal contains two critical points of the return map $f_{\lambda,a}^q$, since at such a , the parametrisation by locating the free critical value fails. We will show that the collection of such parameters in each Type (A) (B) components is a curve with end points at Type (D) parameters. Then the parametrisation will work in the complement of this curve.
- \mathcal{C}_λ can be viewed as a "pinched model" of \mathcal{C}_0 , where the pinched points are exactly Type (D) (MP) parameters. At such parameters there are no longer nested puzzles surrounding it and thus Grötzsch inequality can not be applied. Instead, we will use an argument based on holomorphic motion to conclude local connectivity. See 6.1. Moreover, in order to construct puzzle pieces adapted to Type (D) parameters, we need to investigate the landing external rays at them, which is a rather subtle subject, see 3.6.




Figure 1: \mathcal{C}_λ and the Julia set of $\lambda z + z^2$, with $\lambda = 1$ on the left and $e^{\frac{2\pi i}{3}}$ on the right. Different types of parameters in \mathcal{H}^λ are marked out. Type (A) (B) and (C) are in green. $0, a_0, a_1, a_2$ are of Type (D). Black parts are copies of the Mandelbrot set or the parabolic Mandelbrot set, correspond respectively to polynomial-like or parabolic-like maps [13].

Outline of the paper. In Section 2 we recall some known results of parameter external rays, and the description of \mathcal{C}_0 given by [16]. In Section 3, we first prove the existence of Type (A) (B) components in 3.1 and prove that there are exactly q Type (D) parameters in 3.2. Then we parametrize Type (A) (B) components in 3.3 and prove their uniqueness (Corollary 3.4.7). We describe relative positions among Type (A) (B) (D) parameters and investigate landing parameter external rays at Type (D) parameters. (Proposition I and II). These results have their own interests and will simplify the construction of parameter puzzles later. Section 4 focus on the construction of admissible dynamical graphs, so that one can apply

Yoccoz's Theorem to obtain shrinking puzzle pieces. In Section 5, we construct parameter puzzles and show that dynamical graphs moves holomorphically when the parameter varies in parameter puzzles. Finally Section 6 is devoted to the proof of the Main Theorem and Corollary I.

Acknowledgements. I would like to thank Pascale Roesch for comments on the the manuscript. The pictures are generated by It of Mannes Technology and a personnel computer program of Arnaud Chéritat.

2 Preparations

2.1 Families with marked out critical points

Consider the following two families

$$g_{\lambda,c}(z) = \lambda z \left(1 - \frac{1+1/c}{2}z + \frac{1/c}{3}z^2 \right), \quad (\lambda, c) \in (\mathbb{C}^*)^2 \quad (3)$$

$$\hat{g}_{\lambda,s}(z) = \lambda z \left(1 - \frac{s+1/s}{2}z + \frac{1}{3}z^2 \right), \quad (\lambda, s) \in (\mathbb{C}^*)^2 \quad (4)$$

The advantage of $g_{\lambda,c}$ (or $\hat{g}_{\lambda,s}$) is that the two critical points $1, c$ (resp. $s, 1/s$) are marked out. Fix λ -slice, denote by $\tilde{\mathcal{C}}_\lambda$ $\hat{\mathcal{C}}_\lambda$ respectively the connected locus for these two families. The attracting locus $\tilde{\mathcal{H}}^\lambda, \hat{\mathcal{H}}^\lambda$ are defined likewise as in (2). For $\lambda \neq 0$, the relation between $f_{\lambda,a}$, $g_{\lambda,c}$ and $\hat{g}_{\lambda,s}$ is given by

Lemma 2.1.1. *$\hat{g}_{\lambda,s}$ is conjugate to $f_{\lambda,a}$ by $z \mapsto \sqrt{3/\lambda} \cdot z$ with $a = \sigma(s) = -\sqrt{3\lambda} \cdot \frac{s+1/s}{2}$ and is conjugate to $g_{\lambda,c}$ by $z \mapsto \frac{1}{s} \cdot z$ with $c = \iota(s) = s^2$.*

2.2 Parameter external rays in $Per_1(\lambda)$, $|\lambda| \leq 1$

Let f be a polynomial with connected Julia set and z a (pre-)periodic point. It is a classical result by Yoccoz that z admits an external ray landing (cf. [10]). The collection of all the angles of external rays landing at z is called the **portrait** at z . For family (1), the Böttcher coordinate $\phi_{\lambda,a}^\infty$ at ∞ depends analytically on (λ, a) by taking the normalization $\phi_{\lambda,a}^\infty(z) = z + o(1)$. So there is no ambiguity of angles when (λ, a) varies.

Now suppose $|\lambda| \leq 1$. It is a classical result (cf. [22]) that \mathcal{C}_λ is a full continuum containing $\pm\sqrt{3\lambda}$. Thus one can define analytically the two critical points of $f_{\lambda,a}$ for $a \in \mathbb{C} \setminus \mathcal{C}_\lambda$: $c_{\lambda,a}^\pm = \frac{-a \pm \sqrt{a^2 - 3\lambda}}{3}$ such that $c_{\lambda,a}^+ \in K_{\lambda,a}$ and $c_{\lambda,a}^- \in \mathbb{C} \setminus K_{\lambda,a}$. Set $v_{\lambda,a}^\pm = f_a(c_\pm(a))$. One can parametrize $\mathbb{C} \setminus \mathcal{C}_\lambda$ by looking at the position of $v_{\lambda,a}^-$ in the Böttcher coordinate:

Proposition 2.2.1 ([22]). *Let $|\lambda| \leq 1$. The mapping $\Phi_\infty^\lambda : \mathbb{C} \setminus \mathcal{C}_\lambda \longrightarrow \mathbb{C} \setminus \overline{\mathbb{D}}$ defined by $a \mapsto \phi_a^\infty(v_{\lambda,a}^-)$ is a degree 3 covering.*

A parameter external ray with angle t is defined to be a connected component of $(\Phi_\infty^\lambda)^{-1}(\{re^{2\pi it}; r > 1\})$. In most cases, we denote a parameter external ray by $\mathcal{R}_\infty^\lambda$ without precising the component. When we say " $\mathcal{R}_\infty^\lambda$ lands at $a_0 \in \mathcal{C}_\lambda$ ", it means that one of the three components of $(\Phi_\infty^\lambda)^{-1}(\{re^{2\pi it}; r > 1\})$ accumulates at a_0 . We use the notation $\mathcal{R}_\infty^\lambda(t) * (r_1, r_2)$ to represent the set $(\Phi_\infty^\lambda)^{-1}(\{re^{2\pi it}; r \in (r_1, r_2)\})$.

Remark 2.2.2. *Let us just mention that in [22] the proposition above is stated only for $\lambda = e^{2\pi i\theta}$ with θ of Brjuno type. However the proof there works without any change for all $\lambda \in \overline{\mathbb{D}} \setminus \{0\}$.*

Proposition 2.2.3 can be passed to family (4) by Lemma 2.1.1:

Proposition 2.2.3. For $\lambda \in \overline{\mathbb{D}} \setminus \{0\}$, $\mathbb{C}^* \setminus \hat{\mathcal{C}}_\lambda$ has exactly two connected components ($\tau : s \mapsto 1/s$)

$$\hat{\mathcal{H}}_\infty = \{s \in \mathbb{C}^*; \hat{g}_{\lambda,s}^n(s) \rightarrow \infty \text{ as } n \rightarrow \infty\}$$

$$\tau\hat{\mathcal{H}}_\infty = \{s \in \mathbb{C}^*; \hat{g}_{\lambda,s}^n(1/s) \rightarrow \infty \text{ as } n \rightarrow \infty\}$$

which are punctured neighborhoods of $\infty, 0$ respectively and are homeomorphic to punctured disk. Moreover the mapping $\hat{\Phi}_\infty : \hat{\mathcal{H}}_{\infty,0} \rightarrow \mathbb{C} \setminus \overline{D}$ given by $\hat{\Phi}_\infty(s) = \hat{\phi}_s^\infty(\hat{g}_{\lambda,s}(s))$ is a degree 3 covering, where $\hat{\phi}_s^\infty(z) = \phi_{\sigma(s)}^\infty(\frac{\lambda}{3} \cdot z)$.

By Lemma 2.1.1 we can write $\mathbb{C}^* \setminus \check{\mathcal{C}}_\lambda = \check{\mathcal{H}}_\infty^\lambda \cup \tau\check{\mathcal{H}}_\infty^\lambda$, where $\tau : c \mapsto \frac{1}{c}$ and

$$\check{\mathcal{H}}_\infty^\lambda = \{c \in \mathbb{C}^*; g_{\lambda,c}^n(c) \rightarrow \infty \text{ as } n \rightarrow \infty\}$$

$$\tau\check{\mathcal{H}}_\infty^\lambda = \{c \in \mathbb{C}^*; g_{\lambda,c}^n(1) \rightarrow \infty \text{ as } n \rightarrow \infty\}$$

by noticing that $cg_{\lambda,c}(\frac{z}{c}) = g_{\lambda,1/c}(z)$. These two escaping regions are simply connected.

Lemma 2.2.4. Let $\check{\mathcal{U}} \subset \check{\mathcal{H}}^\lambda$ be an open component. Then $\check{\mathcal{U}}$ is simply connected and $\partial\check{\mathcal{U}} \subset \partial\check{\mathcal{C}}_\lambda$. In particular, if $\partial\check{\mathcal{U}}$ is locally connected, then it is a Jordan curve.

Proof. Simply connectivity can be shown easily by applying maximum principle to $g_{\lambda,c}^n(c)$ or $g_{\lambda,c}^n(1)$; for $\partial\check{\mathcal{U}} \subset \partial\check{\mathcal{C}}_\lambda$, see [23, Lem. 2.1.5]. We prove the last statement. Suppose $\partial\check{\mathcal{U}}$ is locally connected, then any conformal representation $\Psi : \mathbb{D} \rightarrow \check{\mathcal{U}}$ can be extended continuously and surjectively to the boundary. Moreover $\Psi : \partial\mathbb{D} \rightarrow \partial\check{\mathcal{U}}$ is injective: if not, then there exists $a \in \partial\check{\mathcal{U}}$ accessible by two rays $\Psi(re^{it_1}), \Psi(re^{it_2})$ which bound a simply connected region containing part of $\partial\check{\mathcal{U}}$. This contradicts $\partial\check{\mathcal{U}} \subset \check{\mathcal{C}}_\lambda$. \square

Definition 2.2.5. Let $|\lambda| \leq 1$. $a_0 \in \mathcal{C}_\lambda$ is a **Misiurewicz parameter** if one of the critical points is repelling pre-periodic. For $\lambda = e^{\frac{2\pi ip}{q}}$, $a_0 \in \mathcal{C}_\lambda$ is a **Misiurewicz parabolic parameter** if one of the critical points is pre-periodic to $z = 0$. The same notions are defined for $g_{\lambda,c}$ and $\hat{g}_{\lambda,s}$.

Lemma 2.2.6. Let $|\lambda| \leq 1$, $t \in \mathbb{Q}/\mathbb{Z}$. Then $\mathcal{R}_\infty^\lambda(t)$ land at some $a_0 \in \mathcal{C}_\lambda$ which is geometrically finite.

Proof. It suffices to prove the accumulation set of $\mathcal{R}_\infty^\lambda(t)$ is finite, since the accumulation set is connected. It will be convenient to work in family (4) since the critical points are marked out and by Proposition 2.2.3, external rays with angle t , denoted by $\hat{\mathcal{R}}_\infty^\lambda(t)$, in $\hat{\mathcal{H}}_\infty$ are well-defined. Suppose s_0 is accumulated by $\hat{\mathcal{R}}_\infty^\lambda(t)$. Moreover if $\lambda = e^{\frac{2\pi i p}{q}}$ suppose s_0 is not double parabolic (it is not hard to see that $\#\mathcal{A}_{p/q} < \infty$). Suppose $\hat{R}_{\lambda,s_0}^\infty(t)$ (the dynamical external ray of \hat{g}_{λ,s_0}) lands at x_{λ,s_0} .

- If x_{λ,s_0} is repelling pre-periodic or if $\lambda = e^{\frac{2\pi ip}{q}}$ and x_{λ,s_0} is pre-periodic to $z = 0$, then by stability of repelling Koenigs coordinate or repelling Fatou coordinate one concludes that $x_{\lambda,s_0} = \hat{v}(s_0) := \hat{g}_{\lambda,s_0}(s_0)$. Therefore s_0 satisfies a non-trivial algebraic equation: $\hat{g}^{lk}(\hat{v}(s_0)) = \hat{g}^l(\hat{v}(s_0))$ which has only finitely many solutions.
- If x_{λ,s_0} is parabolic pre-periodic and when $\lambda = e^{\frac{2\pi ip}{q}}$ it is not pre-periodic to 0. Then there exists $l \geq 0, k \geq 1$ (only depend on t) such that $\hat{g}_{\lambda,s_0}^l(\hat{R}_{\lambda,s_0}^\infty(t))$ is fixed by \hat{g}_{λ,s_0}^k . By the snail lemma, (s_0, x_{λ,s_0}) verifies

$$\hat{g}_{\lambda,s}^{k+l}(z) = \hat{g}_{\lambda,s}^l(z), (\hat{g}_{\lambda,s}^k)'(\hat{g}_{\lambda,s}^l(z)) = 1. \quad (5)$$

(5) defines a non-trivial algebraic variety (not equal to $\mathbb{C}^* \times \mathbb{C}$) and hence consists of only finitely many irreducible components of dimension 1 and 0 (i.e. points). We claim that when $\lambda \neq e^{\frac{2\pi ip}{q}}$, there are no dimension 1 component; when $\lambda = e^{\frac{2\pi ip}{q}}$, the only dimension 1 component is $z = 0$. Indeed, suppose there is another such component X . Then X is unbounded (goes to the boundary of $\mathbb{C}^* \times \mathbb{C}$).

Consider the projections $\pi_1 : X \rightarrow \mathbb{C}$, $\pi(s, z) = s$ and $\pi_2 : X \rightarrow \mathbb{C}$, $\pi(s, z) = z$. Then at least one of $\pi_i(X)$, $i = 1, 2$ is unbounded. If $\pi_1(X)$ is unbounded, let $(s_n, z_n) \subset X$ with $s_n \rightarrow \infty$ or 0 , then λ has to be $e^{\frac{2\pi i p}{q}}$ and for n large enough, $z_n = 0$, so X has to be $z = 0$. If $\pi_2(X)$ is unbounded, let $(s_n, z_n) \subset X$ with $z_n \rightarrow \infty$. If (s_n) also tends to ∞ or 0 , then we are in the precedent case; if not, then for all n , the basin at infinity of \hat{g}_{λ, s_0} contains a common neighborhood of ∞ . Hence for n large enough, z_n escapes to ∞ , contradicting the assumption that (s_n, z_n) verifies (5).

To conclude, the above analysis shows that the accumulation set of $\mathcal{R}_0(t)$ is finite and the possible accumulations are geometrically finite maps \square

For our purpose, we extract from the lemma above the case of $\lambda = e^{2\pi i \frac{p}{q}}$:

Lemma 2.2.7. *Let $\lambda = e^{2\pi i \frac{p}{q}}$, $t \in \mathbb{Q}/\mathbb{Z}$. Then $\mathcal{R}_\infty^\lambda(t)$ lands at some a_0 . In the dynamical plan of f_{a_0} , $R_{\lambda, a_0}^\infty(t)$ lands at a (pre-)periodic point $x(a_0)$. Moreover*

- *If $x(a_0)$ is repelling, then $x(a_0)$ is the free critical value.*
- *If $a_0 \notin \mathcal{A}_{p/q}$, and $x(a_0)$ is in the inverse orbit of 0 , then $x(a_0)$ is the free critical value.*
- *If $a_0 = \mathbf{a}_m \in \mathcal{A}_{p/q}$ and t is not pre-periodic to the portrait at $z = 0$ of f_{λ, a_0} , then $\mathcal{R}_\infty^\lambda(t)$ does not land at a_0 .*

Lemma 2.2.8. *Suppose $|\lambda| \leq 1$. Let a_0 be a Misiurewicz parameter or parabolic Misiurewicz parameter, $t \in \mathbb{Q}/\mathbb{Z}$. Then $R_{\lambda, a_0}^\infty(t)$ lands at one of the critical values of $f_{\lambda, a}$ if and only if $\mathcal{R}_\infty^\lambda(t)$ lands at a_0 .*

Proof. First of all $a_0 \neq \pm\sqrt{3\lambda}$, i.e. the two critical points are not the same. This is clear when $|\lambda| < 1$ or $\lambda = e^{\frac{2\pi i p}{q}}$ since 0 attracts at least one critical point. If λ is Siegel, then the boundary of the Siegel disk is contained in the accumulation set of critical orbits; if λ is Cremer, then the Julia set is not locally connected. Therefore $c_{\lambda, a}^\pm$ can be analytically defined near a_0 . Then one uses stability of repelling Koenigs coordinate or repelling Fatou coordinate to conclude the proof. \square

Lemma 2.2.9. *Let $t \in \mathbb{Q}/\mathbb{Z}$.*

- *Suppose $|\lambda| \leq 1$ but $\lambda \neq e^{2\pi i \frac{p}{q}}$. Let $\mathcal{O} \subset \mathbb{C}$ be the set of parameters a such that $R_{\lambda, a}^\infty(t)$ lands at a repelling (pre-)periodic point, Then $\mathbb{C} \setminus \bigcup_{k \geq 1} \overline{\mathcal{R}_\infty^\lambda(3^k t)} \subset \mathcal{O}$. Moreover $\bigcup_{k \geq 0} \overline{R_{\lambda, a}^\infty(3^k t)}$ admits a dynamical holomorphic motion for a in any component of \mathcal{O} .*
- *Suppose $\lambda = e^{2\pi i \frac{p}{q}}$. Let $\mathcal{O} \subset \mathbb{C}$ be the set of parameters a such that $R_{\lambda, a}^\infty(t)$ lands at a repelling (pre-)periodic point or the inverse orbit of 0 . Then $(\mathbb{C} \setminus \bigcup_{k \geq 1} \overline{\mathcal{R}_\infty^\lambda(3^k t)}) \setminus \mathcal{A}_{p/q} \subset \mathcal{O}$. Moreover $\bigcup_{k \geq 0} \overline{R_{\lambda, a}^\infty(3^k t)}$ admits a dynamical holomorphic motion for a in any component of $\mathcal{O} \setminus \mathcal{A}_{p/q}$ ($\mathcal{A}_{p/q}$ is the set of double parabolic parameters).*

Proof. For the first point, see [16, Lem. 3.8]. For the second point, for $a \in \mathbb{C} \setminus \bigcup_{k \geq 1} \overline{\mathcal{R}_\infty^\lambda(3^k t)}$, $R_{\lambda, a}^\infty(3^k t)$ land for $k \geq 0$ since they do not crash on the free critical point. It suffices to prove that the landing point $x(a)$ of $R_{\lambda, a}^\infty(t)$ is either repelling pre-periodic or in the inverse orbit of 0 . Suppose the contrary that it is parabolic (pre-)periodic beyond $z = 0$, then the portrait at $z = 0$ of $f_{\lambda, a}$ does not intersect $\{3^k t\}_{k \geq 0}$. Since the portrait at $z = 0$ is stable by the stability of repelling Fatou coordinate at $z = 0$ (notice that $a \notin \mathcal{A}_{p/q}$), so for a' in a neighborhood of a , the portrait at $z = 0$ of $f_{\lambda, a'}$ does not intersect $\{3^k t\}_{k \geq 0}$. However on the other hand, from the second point in the proof of Lemma 2.2.6, the set of parabolic parameters with a given period is discrete. So for a' in a punctured neighborhood of a , $x(a')$ is either repelling or sent to $z = 0$ with portrait containing the cycle in $\{3^k t\}_{k \geq 0}$. So the second case is impossible. The first case contradicts the maximum principle for $a \mapsto 1/(f^n)'(x(a))$ for n large enough. \square

2.3 Results for \mathcal{C}_0

When $\lambda = 0$, $z = 0$ is a super-attracting fixed point of $f_{0,a}$, $-\frac{2a}{3}$ is the free critical point. \mathcal{C}_0 has a unique hyperbolic component \mathcal{H}_0^0 containing $a = 0$. (The upper index means $\lambda = 0$, the lower means it is of depth 0, i.e. $-\frac{2a}{3}$ is in the immediate basin of 0). Similarly as $\mathbb{C} \setminus \mathcal{C}_0$, one can parametrize \mathcal{H}_0^0 :

Proposition 2.3.1 ([16]). *The mapping $\Phi_0^0 : \mathcal{H}_0^0 \rightarrow \mathbb{D}$ given by $\Phi_0^0(a) = \phi_{0,a}^0(f_{0,a}(-\frac{2a}{3}))$ is a degree 4 holomorphic covering ramified at 0.*

The internal ray of angle t in \mathcal{H}_0^0 is defined to be a connected component of

$$(\Phi_0^0)^{-1}(\{re^{2\pi it}; r \in (0, 1)\}).$$

Remark 2.3.2. *It is easy to verify by symmetric that*

$$\Phi_0^0(-a) = \Phi_0^0(a), \Phi_0^0(\bar{a}) = \overline{\Phi_0^0(a)}; \Phi_\infty^0(-a) = -\Phi_\infty^0(a), \Phi_\infty^0(\bar{a}) = \overline{\Phi_\infty^0(a)}$$

Remark 2.3.3. *By Proposition 2.3.1 and 2.2.1, there are 4 (resp. 3) internal (resp. external) rays associated to a given angle. Let $\mathcal{S} = \{x + iy, x \geq 0, y > 0\}$. These four rays are contained in four sectors $\mathcal{S}, i\mathcal{S}, -\mathcal{S}, -i\mathcal{S}$ respectively. Hence we precise a ray contained in a sector by adding this sector in the above index, for example $\mathcal{R}_0^{0,\mathcal{S}}(t)$ (resp. $\mathcal{R}_\infty^{0,\mathcal{S}}(t)$) denotes the internal (resp. external) ray with angle t contained in \mathcal{S} . Nevertheless in most time we omit for simplicity this index if the sector to which this ray belong is clear or if there is no need to precise, according to the context.*

The main results in [16] can be summarized as the following proposition and theorem:

Proposition 2.3.4. [16, Lem 2.29]. *For all $t \in \mathbb{R}/\mathbb{Z}$, $\mathcal{R}_0^{0,\mathcal{S}}(t)$ lands at some $a(t) \in \partial\mathcal{H}_0^0$. In the dynamical plan, $\mathcal{R}_{a(t)}^0(t)$ lands at a parabolic k -periodic point if and only if $\frac{t+1}{2}$ is k -periodic under multiplication by 2.*

Theorem 2.3.5. *For all $n \geq 0$, $\partial\mathcal{H}_n^0$ is a union of Jordan curve. If $a_0 \in \partial\mathcal{H}_n^0$ is renormalizable, then there are exactly two external rays landing at it; otherwise there is only one.*

If $a_0 \in \partial\mathcal{H}_0^0$ is renormalizable, then the two corresponding rays landing at it separate \mathbb{C} into two connected components U_{a_0} and W_{a_0} with $\mathcal{H}_0^0 \subset U_{a_0}$ and $\overline{W_{a_0}}$ containing a copy of Mandelbrot set rooted at a_0 . For $a \in \partial\mathcal{H}_0^0$, define limbs by $\mathcal{L}_a := W_a \cap \mathcal{C}_0$ if a is renormalisable; otherwise $\mathcal{L}_a := \{a\}$. Then $\mathcal{C}_0 = \mathcal{H}_0^0 \cup (\bigcup_{a \in \partial\mathcal{H}_0^0} \mathcal{L}_a)$.

Let us give three lemmas which will be used later in the proof of Proposition 3.6.1:

Lemma 2.3.6. *Let $a_1, a_2 \in \partial\mathcal{H}_0^0$ be two different renormalizable parameters. By Theorem 2.3.5, suppose that $\mathcal{R}_{0,\infty}^0(t_i), \mathcal{R}_{0,\infty}^0(t'_i)$ land at a_i , $i = 1, 2$. Then $\{t_1, t'_1\} \neq \{t_2, t'_2\}$.*

Proof. Suppose the contrary that $\{t_1, t'_1\} = \{t_2, t'_2\}$. Then the portrait of the co-critical point \tilde{c}_{a_i} will be the same, i.e. the angles of the two external rays bounding \tilde{c}_{a_i} equal those for \tilde{c}_{a_1} . On the other hand, since the parametrisation of \mathcal{H}_∞^0 defined by $a \mapsto \phi_{0,a}^\infty(\tilde{c}_a)$ is an isomorphism, we conclude that the angles of parameter rays landing at a_i are different. Thus in the dynamical plan, the portrait at \tilde{c}_{a_i} should also be different, a contradiction. \square

Lemma 2.3.7. *There are exactly $4q$ external rays whose angles are of rotation number p/q (under multiplication by 3) landing at $\partial\mathcal{H}_0^0$.*

Proof. By Theorem A.2, the periodic cycle of rotation number p/q under multiplication by 2 is unique. Denote this cycle by $T_{\frac{p}{q}}$.

Claim. *Let $t \in T_{\frac{p}{q}}$ such that $\frac{t+1}{2}$ is **not** q -periodic. Then $1-t \in T_{1-\frac{p}{q}}$ and $\frac{(1-t)+1}{2}$ is q -periodic.*

proof of the claim. Let $t = \frac{m}{2^q-1}$. By hypothesis $2^q \cdot \frac{t+1}{2} \neq \frac{t+1}{2}$, which by an elementary computation is equivalent to $\frac{m}{2} \neq \frac{1}{2}$, i.e. m is even. Hence $1-t = \frac{\tilde{m}}{2^q-1}$ where $\tilde{m} = 2^q - 1 - m$ is odd. Hence $\frac{(1-t)+1}{2}$ is q -periodic. \square

Let $a(t)$ be the landing point of $\mathcal{R}_0^{0,S}(t)$. Suppose there are k element in $T_{\frac{p}{q}}$ such that $R_{a(t)}^0(t)$ lands at a parabolic periodic point, then from Proposition 2.3.4 and the claim we see that there are $q-k$ elements in $T_{\frac{p}{q}}$ such that $R_{a(1-t)}^0(1-t)$ lands at a parabolic periodic point. Moreover, if we name these k elements and $q-k$ elements by $t_1, \dots, t_k, \tilde{t}_1, \dots, \tilde{t}_{q-k}$, the claim tells us that $T_{p/q} = \{t_1, \dots, t_k, 1-\tilde{t}_1, \dots, 1-\tilde{t}_{q-k}\}$. Denote by $a(t_i)$ the landing point of $\mathcal{R}_0^{0,S}(t_i)$ for $1 \leq i \leq k$ and $a(\tilde{t}_i)$ the landing point of $\mathcal{R}_0^{0,-iS}(\tilde{t}_i)$ for $1 \leq i \leq q-k$. In the dynamical plan, let $(x_i)_{1 \leq i \leq k}$ and $(\tilde{x}_i)_{1 \leq i \leq q-k}$ be the landing point of $R_{a(t_i)}^0(t_i)$ and $R_{a(\tilde{t}_i)}^0(\tilde{t}_i)$ respectively. Then $(x_i), (\tilde{x}_i)$ are all parabolic periodic points with rotation number p/q . By Theorem 2.3.5, for any $t \in T_{p/q}$, there are exactly two parameter external rays $\mathcal{R}_\infty(\eta), \mathcal{R}_\infty(\eta')$ landing at $a(t)$ with $\eta, \eta' \in \bigcup_k \Theta_k$. Therefore we have found $2q$ external rays landing at parameters on $\partial\mathcal{H}_0^0$ in the right-half plane. By symmetric, there are in total $4q$ rays. \square

Lemma 2.3.8. *Suppose $t \in \mathbb{Q}/\mathbb{Z}$ has rotation number p/q under multiplication by 3. Suppose that a_0 the landing point of $\mathcal{R}_\infty^0(t)$ is a parabolic parameter, i.e. $R_{0,a_0}^\infty(t)$ lands at a parabolic periodic point. Then there are exactly two external rays $\mathcal{R}_\infty^0(\theta^+), \mathcal{R}_\infty^0(\theta^-)$ landing at a_0 (t is one of θ^+, θ^-). Moreover, $R_{0,a_0}^\infty(\theta^+), R_{0,a_0}^\infty(\theta^-)$ bound the critical value v_{0,a_0} , separating it from the immediate basin of 0.*

Proof. Let $\mathcal{P}_n^0(a_0)$ be the puzzle piece ($a_0 \in \mathcal{P}_n^0(a_0)$) constructed in [16]. Since a_0 is renormalisable, by [16, Prop. 3.26], $\mathbf{M}_{a_0} := \bigcap_n \overline{\mathcal{P}_n^0(a_0)}$ is a copy of Mandelbrot set and let $\chi : \mathbf{M}_{a_0} \rightarrow \mathbf{M}$ be the homeomorphism. Let θ_n^- (resp. θ_n^+) be the smallest/largest angle of the external rays involved in $\partial\mathcal{P}_n^0(a_0)$.

Step 1. For n large enough, $3^k \theta_n^\pm = \theta_{n-k}^\pm$.

By [16, Lem. 3.17, Prop. 3.22], for n large enough, there is a natural homeomorphism preserving equipotentials and rays between $\partial\mathcal{P}_n^0(a_0)$ and $\partial P_{0,n}^{a_0}$. Thus θ_n^\pm are also the largest/smallest angle involved in $\partial P_{0,n}^{a_0}$. Notice that $\mathcal{R}_\infty^0(\theta_n^\pm)$ land at the same hyperbolic component since adjacent and capture components have disjoint boundaries (cf. [16, Lem. 1]). Thus the two corresponding dynamical rays $R_{0,a_0}^\infty(\theta_n^\pm)$ land at the boundary of some Fatou component U . Let x^\pm be their landing points and y^\pm their intersection with the equipotential in $\partial P_{0,n}^{a_0}$. Then y^+, y^- are linked by a curve L consisting of two segments of $R_{0,a_0}^\infty(\theta_n^\pm)$ linking y^\pm to x^\pm and a segment contained in $\overline{U} \cap \partial P_{0,n}^{a_0}$ linking x^+, x^- . Suppose the contrary that $3^k \theta_n^\pm \neq \theta_{n-k}^\pm$. Then there is a segment of equipotential γ with angles between (θ_n^-, θ_n^+) linking $f_{0,a_0}^k(y^+), f_{0,a_0}^k(y^-)$. Thus $\gamma \cup f_{0,a_0}^k(L)$ bounds a simply connected domain P which do not contain the free critical value of f_{0,a_0} . Let Q be the component of $f_{0,a_0}^{-k}(P)$ whose boundary contains L . Then $f_{0,a_0}^k : \overline{Q} \rightarrow \overline{P}$ is injective. Hence ∂Q contains the segment of equipotential consisting of angles in $\mathbb{S}^1 \setminus (\theta_n^-, \theta_n^+)$, which in particular contains the angle 0. While $R_{0,a_0}^\infty(0)$ is fixed by f_{0,a_0} , this leads to a contradiction.

Step 2. Two external rays land at $\chi^{-1}(\frac{1}{4})$.

Since $\overline{P_{0,n}^{a_0}} \subset P_{0,n-k}^{a_0}$, we obtain a sequence of decreasing/increasing angles $\{\theta_{n_0+lk}^\pm\}_{l \geq 0}$ for some n_0 large enough. Let l tend to infinity we get two limits θ^\pm which satisfy $3^k \theta^\pm = \theta^\pm$ by Step 1. Let a^\pm be the landing point of $\mathcal{R}_{0,\infty}(\theta^\pm)$. Clearly $a^\pm \in \mathbf{M}_{a_0}$. We claim that $a^+ = a^- = \chi^{-1}(\frac{1}{4})$. We prove for a^+ and the left one will be the same. If not, then $R_{0,a^+}^\infty(\theta^+)$ land at a repelling fixed point for f_{0,a^+}^k . By [16, Lem. 2.24], this point is the free critical value of f_{0,a^+} . Let $P_{c^+}(z) = z^2 + c^+$ be the corresponding quadratic polynomial to which $f_{0,a^+}^k|_{P_{c^+}^{a^+}}$ is conjugate. Then we have $P_{c^+}(c^+) = c^+$, i.e. $c^+ = 0$. While $c^+ \in \partial\mathbf{M}$, a contradiction. Thus we have proved that $\mathcal{R}_\infty^0(\theta^\pm)$ land at $\chi^{-1}(\frac{1}{4})$.

Step 3. Two external rays land at $\chi^{-1}(c)$ for $c \neq \frac{1}{4}$ a parabolic parameter on the cardioid.

Let $a = \chi^{-1}(c)$ and $R_{0,a}^\infty(t), R_{0,a}^\infty(3^k t), \dots, R_{0,a}^\infty(3^{(s-1)k} t)$ be the cycle of external rays landing at the parabolic fixed point of $f_{0,a}^k$. Notice that c is satellite, hence it is the intersection of the closure of the main

hyperbolic component \mathbf{H}_0 and the closure of the periodic hyperbolic component \mathbf{H}_1 attached at c . Thus there are two pinching paths $\gamma_0 \subset \chi^{-1}(\mathbf{H}_0)$ and $\gamma_1 \subset \chi^{-1}(\mathbf{H}_1)$ converging to $f_{0,a}$ such that

1. if $a' \in \gamma_0$, then $\{R_{0,a'}^\infty(3^{kl}t)\}_{l \geq 0}$ land at a repelling s -periodic cycle of $f_{0,a'}^k$.
2. if $a' \in \gamma_1$, then $\{R_{0,a'}^\infty(3^{kl}t)\}_{l \geq 0}$ land at a common repelling fixed point of $f_{0,a'}^k$.

Since landing at a repelling periodic cycle is an open property, there must exist two external rays among $\{\mathcal{R}_\infty^0(3^l t)\}_{l \geq 0}$ landing at a .

Step 4. The two rays landing are unique. The proof of [16, Thm. 3] can be adapted.

Step 5. a_0 is on the cardioid of \mathbf{M}_{a_0} .

Let k be the period of renormalisation of a_0 . Since the cycle $t, 3t, \dots, 3^{q-1}t$ has rotation number p/q under multiplication by 3, then so does the cycle $t, 3^k t, \dots, 3^{sk} t$ under multiplication by 3^k . By [16, Prop. 3.22], $f_{0,a_0}^k : P_{0,n}^{a_0} \rightarrow P_{0,n-k}^{a_0}$ is hybrid conjugate to a quadratic polynomial $P_c(z) = z^2 + c$ for n large enough, where $P_{0,n}^{a_0}$ is the dynamical puzzle piece containing the free critical value of f_{0,a_0} . Thus the cycle of the parabolic periodic point of P_c has a cycle of access, which also admits a rotation number. This cycle of access is homotopic to a cycle of external rays with the same rotation number. Since by Theorem A.2, there is only one cycle of angles for a given rotation number under multiplication by 2, this implies that the parabolic periodic point of P_c is actually fixed, hence c is on the cardioid, so is a_0 .

From Step 1 and 3 we see that $R_{0,a_0}^\infty(\theta^+), R_{0,a_0}^\infty(\theta^-)$ bound a sector containing v_{0,a_0} , separating it from $z = 0$. \square

Corollary 2.3.9. *Under the same hypothesis and notations of Lemma 2.3.8, the interval (θ^-, θ^+) does not contain any angle in $\{3^k \theta^+; k \geq 0\} \cup \{3^k \theta^-; k \geq 0\}$.*

Proof. It suffices to prove for the case when a_0 is the root of \mathbf{M}_{a_0} .

First we prove that in the dynamical plan of f_{0,a_0} , the wake W bounded by $R_{0,a_0}^\infty(\theta^\pm)$ (that is also the wake containing the critical value $v_{0,a}$) does not contain the critical point $-\frac{2a}{3}$. Let \tilde{W} be the wake bounding $-\frac{2a}{3}$ and bounded by $R_{0,a_0}^\infty(\tilde{\theta}^\pm)$. Suppose the contrary, then we have $\tilde{W} \subset W$. Hence $f_{0,a}^k|_{W \setminus \tilde{W}}$ is injective and proper, where k is the period of the parabolic cycle. Since $3^{k-1}\theta^\pm = \tilde{\theta}^\pm$, we conclude that $f_{0,a}^k(W \setminus \tilde{W}) \subset \tilde{W}$. However on the other hand since $\tilde{W} \subset W$, the wake bounded by $R_{0,a_0}^\infty(3^{k-1}\theta^\pm)$ contains the wake bounded by $R_{0,a_0}^\infty(3^{k-1}\tilde{\theta}^\pm)$. Thus $f_{0,a}(f_{0,a}^{k-1}(W \setminus \tilde{W}))$ will not intersect W , a contradiction.

Next we prove that W contains no other point in the parabolic periodic cycle. Suppose the contrary, then W contains another wake W' bounded by some $R_{0,a_0}^\infty(3^l \theta^\pm)$ with $1 \leq l \leq k-2$. Thus $f_{0,a_0}^l(W) = W'$. Apply Denjoy-Wolff theorem we obtain $f_{0,a_0}^{nl}(W)$ converges to ∞ , a contradiction. \square

3 Description of \mathcal{H}^λ , $\lambda = e^{\frac{2\pi i p}{q}}$

The main goal of this section is to prove the following:

Proposition I. *There are exactly q Type (D) parameters, i.e. $\#\mathcal{A}_{p/q} = q$. The different $\mathbf{a}_m \in \mathcal{A}_{p/q}$ ($0 \leq m \leq q-1$) are characterized by the portrait at $z = 0$. Moreover each \mathbf{a}_m admits 4 parameter external rays landing, cutting $\text{Per}_1(\lambda)$ with respect to the portrait at $z = 0$. Moreover, these four rays are unique among all rays with rational angle.*

Proposition II. *\mathcal{H}^λ has exactly 2 adjacent components $\mathcal{B}_0, \mathcal{B}_q$ symmetric with respect to $a \mapsto -a$ and $q-1$ bitransitive components $\mathcal{B}_1, \dots, \mathcal{B}_{q-1}$. These components are characterized by the portrait at $z = 0$. Moreover,*

$$\overline{\mathcal{B}_m} \cap \mathcal{A}_{p/q} = \mathbf{a}_m \cup \mathbf{a}_{m-1}, \quad 1 \leq m \leq q-1; \quad \overline{\mathcal{B}_0} \cap \mathcal{A}_{p/q} = \mathbf{a}_0, \quad \overline{\mathcal{B}_q} \cap \mathcal{A}_{p/q} = \mathbf{a}_{q-1}.$$

Our construction of \mathbf{a}_m and \mathcal{B}_m is based on the technique of pinching deformation developed by Cui-Tan [5]. Its advantage is that the combinatorial information is explicitly prescribed. See a different approach using transversality method applied to the same problem for quadratic rational maps [3]. Notice that the uniqueness of the 4 external rays in Proposition I is not obvious: this is the cubic analogue to the ray landing problem at parabolic parameters in the Mandelbrot set, which has been solved by Douady-Hubbard [10]. See also [12] for an easier presentation. Essentially, the "Tour de valse" argument should work, but should also be more delicate since the fixed point is parabolic degenerate. Instead, here we use a "ray counting" argument based on pinching deformation to avoid the complicate analysis in [10]. Let us mention that the landing problem at double parabolic parameters has been solved recently by [1, Thm 11.7]. However our method are different from theirs.

3.1 Existence of adjacent and bitransitive components

In this subsection we prove existence of adjacent and bitransitive components satisfying any combinatorics given. The main tool is pinching deformation.

For $(\lambda, a) \in (0, 1) \times (-\sqrt{3\lambda}, \sqrt{3\lambda})$, $z = 0$ is an attracting fixed point for $f_{\lambda, a}$. Denote by $\Omega_{\lambda, a}$ the maximal linearization domain at $z = 0$. Since $\overline{f_{\lambda, a}} = f_{\lambda, a}$, $\Omega_{\lambda, a}$ is symmetric with respect to x -axis and therefore $\partial\Omega_{\lambda, a}$ contains both critical points of $f_{\lambda, a}$. Denote by $c_{\lambda, a}^+$ (resp. $c_{\lambda, a}^-$) the one contained in the upper-half (resp. lower-half) plane. Let $\phi_{\lambda, a} : \overline{\Omega_{\lambda, a}} \rightarrow \mathbb{D}$ be the Koenigs coordinate normalised by $\phi_{\lambda, a}(c_{\lambda, a}^+) = 1$. Take a branch of $\log(\cdot)$ such that $\log(1) = 0$. Set $\varphi_{\lambda, a} = -\log(\phi_{\lambda, a})$. Then clearly $\varphi_{\lambda, a}(\partial\Omega_{\lambda, a}) = i(-2\pi, 0]$. Define $I_\lambda(a) = \Im(-\varphi_{\lambda, a}(c_{\lambda, a}^-))$. Then by definition I is strictly positive. It is not hard to prove the following lemma by quasiconformal deformation:

Lemma 3.1.1. *For any $x \in (0, 2\pi)$ there exists $(\lambda, a) \in (0, 1) \times (-\sqrt{3\lambda}, \sqrt{3\lambda})$ such that $x = I_\lambda(a)$.*




Figure 2: An illustration for $\frac{p}{q} = \frac{2}{3}$. The bands in orange are \tilde{S}_k .

Let $L_{p/q}^0$ be the line passing 0 and $q \log \lambda + 2\pi(q-p)i$. For $k = 0, 1, \dots, q-1$, let $L_{p/q}^k = L_{p/q}^0 - k \log \lambda \pmod{2\pi i}$ be the line intersecting $i(-2\pi, 0]$. The open Strip S bounded by $L_{p/q}^0, L_{p/q}^0 - 2\pi i$ is divided into q sub-strips by $L_{p/q}^k$. Let S_k be the open sub-strip whose upper boundary is $L_{p/q}^k$. Let $\tilde{L}_{p/q}^k$ be the central line of S_k and suppose that it intersects $i\mathbb{R}$ at $i\tilde{y}_k$. Notice that \tilde{y}_k does not depend on (λ, a) . For any $0 \leq m \leq \lfloor \frac{q}{2} \rfloor$, choose λ and $a_m \in (-\sqrt{3\lambda}, \sqrt{3\lambda})$ such that $-I_\lambda(a) \notin \{\tilde{y}_0, \dots, \tilde{y}_{q-1}\}$ and the interval $(-I_\lambda(a), 0)$ contains exactly m elements in $\{\tilde{y}_0, \dots, \tilde{y}_{q-1}\}$. Such λ and a_m necessarily exist by Lemma 3.1.1. For each k pick an open strips \tilde{S}_k centered at $\tilde{L}_{p/q}^k$ such that $\forall k, \varphi_{\lambda, a_m}(c_{\lambda, a_m}^\pm) \notin \tilde{S}_k$. Then $\{\tilde{S}_k\}_{0 \leq k \leq q-1}$ defines a non-separating multi-annulus \mathcal{A} in the quotient space of f_{λ, a_m} by setting

$$\mathcal{A} = \pi\phi_{\lambda,a_m}^{-1} \left(\exp \left(- \bigcup_{k=0}^{q-1} \tilde{S}_k \right) \right)$$

which by Theorem B.8 gives a converging pinching path $\psi_t \circ f_{\lambda,a_m} \circ \psi_t^{-1} \in \text{Per}_1(\lambda_t)$. Here we choose a different normalisation for ψ_t by setting $\psi_t(0) = 0$ and $\psi_t(z) = z + o(1)$ at ∞ . Therefore the resulting limit when $t \rightarrow \infty$

$$f_{\bar{a}_m} := \psi_\infty \circ f_{\lambda,a} \circ \psi_\infty^{-1} \in \text{Per}_1(e^{2\pi i p/q})$$

is of adjacent type if $m = 0$ and bitransitive type if $0 < m \leq \lfloor \frac{q}{2} \rfloor$.

Now we give a more detailed description for the pinching limit in terms of the portrait at $z = 0$. Notice that the 0-level skeleton for \mathcal{A} is a q -cycle of rays with rotation number p/q starting from 0 and landing at $\partial B_{\lambda,a}^*(0)$. Denote by $\{x_i(a_m)\}$ this corresponding landing cycle of points. Then the period of $\{x_i(a_m)\}$ is q since the cycle of external rays landing at $\{x_i(a_m)\}$ has rotation number p/q . Denote by $\{\theta_i(a_m)\}$ the angles for these external rays, then these angles form a cycle with rotation number p/q under multiplication by 3. The external rays with the same angles, denoted by $R_{\bar{a}_m}^\infty(\theta_i(a_m))$, for the pinching limit $f_{\bar{a}_m}$ land at $z = 0$.

Lemma 3.1.2. *The mapping $m \mapsto \{\theta_i(a_m)\}$ is injective ($0 \leq m \leq \lfloor \frac{q}{2} \rfloor$).*

Proof. Rearranging the index we may assume that $f_{\bar{a}_m}^m(c_m^+)$ belongs to the immediate basin containing c_m^- , where c_m^\pm are the critical points of $f_{\bar{a}_m}^m$. Let $\{\theta_0(a_m), \dots, \theta_{q-1}(a_m)\}$ be such that $3 \cdot \theta_i(a_m) = \theta_{i+1}(a_m)$ and c_m^+ is in the sector bounded by $R_{\bar{a}_m}^\infty(\theta_0(a_m)), R_{\bar{a}_m}^\infty(\theta_1(a_m))$. Let d_i be the length of $[\theta_i(a_m), \theta_{i+1}(a_m)]$. Then $(d_i)_i$ satisfy

$$\begin{cases} 3d_0 = \sum_{k=0}^q d_k + d_1, & 3d_m = \sum_{k=0}^q d_k + d_{m+1} \\ 3d_i = d_{i+1}, & i \neq 0, m \end{cases} \quad \text{if } m \neq 0, \quad (6)$$

$$3d_0 = 2 \sum_{k=0}^q d_k + d_1, \quad 3d_i = d_{i+1}, \quad i \neq 0 \quad \text{if } m = 0. \quad (7)$$

Notice that $\sum d_i = 1$, one can solve the equations above to get $d_0 = \frac{3^q}{3^q-1} \cdot (\frac{1}{3^{m+1}} + \frac{1}{3})$ and $d_m = \frac{3^q}{3^q-1} \cdot (\frac{1}{3^{q+1-m}} + \frac{1}{3})$ if $m \neq 0$; $d_0 = \frac{3^q}{3^q-1} \cdot \frac{2}{3}$ if $m = 0$. Thus the $\{\theta_i(a_m)\}$ are different for $0 \leq m \leq \lfloor \frac{q}{2} \rfloor$. \square




Figure 3: An illustration for pinching bands of depth 0 in $B_{\lambda,a}^*$ for $\frac{p}{q} = \frac{2}{3}$. The resulting pinching limit is in an 1-component.

Definition 3.1.3. By Lemma 3.1.2, we get $\lfloor \frac{q}{2} \rfloor + 1$ different cycles of angles $\{\theta_i(a_m)\}$ ($0 \leq m \leq \lfloor \frac{q}{2} \rfloor$) with rotation number p/q under multiplication by 3. Notice that $\{\theta_i(a_m) + \frac{1}{2}\}$ is also a cycle with rotation number p/q and is different from $\{\theta_i(a_m)\}$ if $m \neq \frac{q}{2}$. Therefore we get $q+1$ different cycles of rotation number p/q . By Theorem A.2, there are exactly $q+1$ cycles of rotation number p/q under multiplication by 3. Hence $\{\theta_i(a_m)\}, \{\theta_i(a_m) + \frac{1}{2}\}$ only depend on m . Denote by Θ_m the cycle $\{\theta_i(a_m)\}$ if $0 \leq m \leq \lfloor \frac{q}{2} \rfloor$; the cycle $\{\theta_i(a_{q-m}) + \frac{1}{2}\}$ if $\lfloor \frac{q}{2} \rfloor < m \leq q$.

Remark 3.1.4. The condition that $\tilde{L}_{p/q}^k$ is the central line of S_k is not essential: suppose $\tilde{L}_{p/q}^0 \subset S_0$ is any line parallel to $L_{p/q}^0$ and $\tilde{L}_{p/q}^k = \tilde{L}_{p/q}^0 - k \log \lambda \pmod{2\pi i}$ be the line intersecting $i(-2\pi, 0]$ at \tilde{y}_k . For each $0 \leq m \leq \lfloor \frac{q}{2} \rfloor$, similarly choose (λ, a_m) so that $(-I_\lambda(a_m), 0)$ contains m elements in $\{\tilde{y}_0, \dots, \tilde{y}_{q-1}\}$. Then the portrait at $z=0$ for the resulting pinching limit is still Θ_m .

Definition 3.1.5. Let $0 \leq m \leq q$. We call an adjacent or bitransitive component of the family f_a a **m -component** if in which all the parameters have portrait Θ_m at $z=0$. In particular, an adjacent component is either a 0-component or a q -component.

Definition 3.1.6. Let $0 \leq m \leq q-1$. For the family $g_{\lambda,c}$, an adjacent or bitransitive component \tilde{B} is called a **m -component** if for all $g_{\lambda,c} \in \tilde{B}$, there are m repelling axis between 1 and c in the counterclockwise direction. In particular, an adjacent component is a 0-component.

From the discussion above, we see that m -component exists with every $0 \leq m \leq q$ for the family $f_{\lambda,a}$ and with every $0 \leq m \leq q-1$ for the family $g_{\lambda,c}$.

3.2 Double parabolic parameters

Let $\lambda = e^{2\pi i \frac{p}{q}}$. By Fatou-Leau Theorem we have Taylor expansions near 0:

$$f_{\lambda,a}^q(z) = z + A_{p/q}(a)z^{q+1} + \mathcal{O}(z^{q+2}). \quad (8)$$

$$g_{\lambda,c}^q(z) = z + C_{p/q}(c)z^{q+1} + \mathcal{O}(z^{q+2}). \quad (9)$$

Definition 3.2.1. We say that $f_{\lambda,a}$ (resp. $g_{\lambda,c}$) is **double parabolic** if $A_{p/q}(a) = 0$ (resp. $C_{p/q}(c) = 0$). For the family $f_{\lambda,a}$, denote by $\mathcal{A}_{p/q}$ the collection of these parameters.

In this subsection we will show that there are exactly q double parabolic parameters for families $f_{\lambda,a}$ and $g_{\lambda,c}$.

Lemma 3.2.2 ([4]). $C_{p/q}(\frac{1}{c})$ is a polynomial in c of degree q .

Proof. Since $C_{p/q}(c)$ is a polynomial in $\frac{1}{c}$, $C_{p/q}(\frac{1}{c})$ is a polynomial in c . When c tends to 0, the map $g_c(z) = c^{-1}g_{\lambda,c}(cz)$ converges uniformly on compact subsets of \mathbb{C} to $P_\lambda(z) = \lambda z(1 - \frac{z}{2})$. Let $P_\lambda^q(z) = z + C_0 z^q + 1 + \mathcal{O}(z^{q+2})$. Then $C_0 \neq 0$ because P_λ has only one parabolic basin. While from (9) we see that $g_c^q(z) = z + c^q C_{p/q}(c) z^{q+1} + \mathcal{O}(z^{q+2})$. Therefore $c^q C_{p/q}(c)$ converges to C_0 when $c \rightarrow 0$. Hence $C_{p/q}(\frac{1}{c})$ has degree q . \square

By the above lemma and the relation between $f_{\lambda,a}, g_{\lambda,c}$ given by Lemma 2.1.1, it suffices to find q different double parabolic parameters for the family $f_{\lambda,a}$. Similarly we construct such parameters by pinching deformation.

Consider the family $f_{\lambda,a}$ for $0 < \lambda < 1$. Recall in Subsection 3.1 the construction of lines $L_{p/q}^k$, the corresponding strips S_k and the corresponding central lines $\tilde{L}_{p/q}^k$, $0 \leq k \leq q-1$. Recall that $\tilde{L}_{p/q}^k$ intersect $i\mathbb{R}$ at \tilde{y}_k . For $0 \leq m \leq \lfloor \frac{q}{2} \rfloor$, choose $(\lambda, a'_m) \in (0, 1) \times (-\sqrt{3\lambda}, \sqrt{3\lambda})$ such that $-I_\lambda(a'_m) = \tilde{y}_m$ (Lemma 3.1.2). Let

S_k^+, S_k^- be the two components of $S_k \setminus \tilde{L}_{p/q}^k$. Let $(\tilde{L}_{p/q}^k)^+, (\tilde{L}_{p/q}^k)^-$ be the central line respectively. For each k pick two narrow strips $\tilde{S}_k^+, \tilde{S}_k^-$ centred at $(\tilde{L}_{p/q}^k)^+, (\tilde{L}_{p/q}^k)^-$. Define a non-seperating annulus for f_{λ, \tilde{a}_m} :

$$\tilde{\mathcal{A}} = \pi \phi_{\lambda, \tilde{a}_m}^{-1} \left(\exp \left(- \bigcup_{k=0}^{q-1} \tilde{S}_k^+ \cup \tilde{S}_k^- \right) \right)$$

For $0 \leq m \leq \lfloor \frac{q}{2} \rfloor$, the corresponding pinching limit yields a double parabolic parameter $\mathbf{a}_m \in \text{Per}_1(e^{2\pi i \frac{p}{q}})$. Its portrait at $z = 0$ is $\Theta_m \cup \Theta_{m+1}$ by Remark 3.1.4. Therefore the portrait for $-\mathbf{a}_m$ at $z = 0$ is $\Theta_{q-m} \cup \Theta_{q-m-1}$. Thus for every $0 \leq m \leq q-1$, we obtain a double parabolic parameter \mathbf{a}_m with portrait Θ_m, Θ_{m+1} at $z = 0$.

Definition 3.2.3. Let $0 \leq m \leq q-1$. \mathbf{a}_m is called the double parabolic parameter of ***m*-type**.

Lemma 3.2.4. A m -component for f_a can only have double parabolic parameters of type $m, m-1$ on its boundary.

Proof. Suppose the contrary that on the boundary there is a double parabolic parameter a_0 of type n , $n \neq m, m-1$. Then the cycle with angle Θ_m will land a repelling q -cycle of f_{a_0} . Since the landing property is stable (cf.[10]), we conclude that for all a in this m -component, $R_{\lambda, a}^\infty(t)$ with $t \in \Theta_m$ lands at a repelling periodic point, a contradiction since these rays should land at $z = 0$. \square

3.3 Parametrisation of parabolic components

In this subsection always fix $\lambda = e^{2\pi i \frac{p}{q}}$. We parametrize m -components $\check{\mathcal{B}}_m$ of family (3) by locating the free critical value in the immediate basins of the quadratic model $P_\lambda(z) = \lambda z + z^2$.

The critical point of P_λ is $-\frac{\lambda}{2}$. Denote by $B_0^*(0)$ the immediate basin of P_λ containing $-\frac{\lambda}{2}$. There are exacty q immediate basins attached at 0 in the cyclic order $B_0^*(0), \dots, B_{q-1}^*(0)$. Let Ω_0^0 be an maximal admissible petal of $P_\lambda^q|_{B_0^*(0)}$. Let $\phi : \overline{\Omega_0^0} \rightarrow \overline{\mathbb{H}}$ (\mathbb{H} is the right half plan) be the Fatou coordiante normalised by $\phi(-\frac{\lambda}{2}) = 0$. Let $1 \leq m \leq q$. Define $\tilde{\Omega}_{c,m}^0 := g_{\lambda, c}^s(\Omega_{c,0}^0)$. For any $0 \leq k \leq q-1$, $n \in \mathbb{Z}$ such that $np+k=0 \pmod{q}$, define Ω_k^n and $\tilde{\Omega}_k^n$ as in (34). For simplicity we will omit index λ for all terms related to family $g_c := g_{\lambda, c}$.

Now consider family g_c . Let $\Omega_{c,0}^0 \subset B_{c,0}^*(0)$ be the standard maximal petal of $g_c^q|_{B_{c,0}^*(0)}$. For $1 \leq m \leq q-1$, let $\tilde{\Omega}_{c,m}^0 := g_c^s(\Omega_{c,0}^0)$ where $1 \leq s \leq q-1$ is the smallest integer such that $sp = m \pmod{q}$. For any $0 \leq k \leq q-1$, $n \in \mathbb{Z}$ such that $np+k=0 \pmod{q}$, $\tilde{\Omega}_{c,k}^n$ is defined as in (34). The degree of $g_c^q|_{B_{c,m}^*(0)}$ is 3 if $m=0$ and is 4 if $1 \leq m \leq q-1$. In the latter case, denote by $1, cr_1, cr_2$ respectively the three critical points of $g_c^q|_{B_{c,m}^*(0)}$; denote by $c, \tilde{cr}_1, \tilde{cr}_2$ respectively the three critical points of $g_c^q|_{B_{c,m}^*(0)}$. Notice that $g_c^r(cr_{1,2}) = c$, $g_c^{q-r}(\tilde{cr}_{1,2}) = 1$.

We introduce the following loci in order to distinguish which critical point is on the maximal petal:

$$\begin{aligned} \check{\mathcal{D}}_0 &= \{c \in \check{\mathcal{B}}_0; 1 \in \partial\Omega_{c,0}^0 \text{ but } c \notin \partial\Omega_{c,0}^0\} \\ \check{\mathcal{I}}_0 &= \{c \in \check{\mathcal{B}}_0; 1, c \in \partial\Omega_{c,0}^0\} \\ \check{\mathcal{D}}_m &= \{c \in \check{\mathcal{B}}_m; 1 \in \partial\Omega_{c,0}^0 \text{ but } cr_1, cr_2 \notin \partial\Omega_{c,0}^0\}, \text{ for } 1 \leq m \leq q-1 \\ \check{\mathcal{I}}_m &= \{c \in \check{\mathcal{B}}_m; 1 \text{ and one of } cr_1, cr_2 \in \partial\Omega_{c,0}^0\}, \text{ for } 1 \leq m \leq q-1 \end{aligned} \tag{10}$$

Let $\tilde{\mathcal{D}}_m = \check{\mathcal{B}}_m \setminus (\check{\mathcal{I}}_m \cup \check{\mathcal{D}}_m)$. Clearly for $c \in \tilde{\mathcal{D}}_m$, $1 \notin \partial\Omega_{c,0}^0$. In particular, if $1 \leq m \leq q$, then $1 \notin \partial\Omega_{c,0}^0$, hence one of $cr_1, cr_2 \in \partial\Omega_{c,0}^0$, and $\tilde{\Omega}_{c,m}^0 \subset B_{c,m}^*(0)$ is a maximal petal for $g_c^q|_{B_{c,m}^*(0)}$, having c on its boundary

but not containing $\tilde{c}r_{1,2}$. Thus we have

$$\begin{aligned}\tilde{\mathcal{D}}_0 &= \{c \in \tilde{\mathcal{B}}_0; c \in \partial\Omega_{c,0}^0 \text{ but } 1 \notin \partial\Omega_{c,0}^0\} \\ \tilde{\mathcal{D}}_m &\subset \{c \in \tilde{\mathcal{B}}_m; c \in \partial\tilde{\Omega}_{c,m} \text{ but } \tilde{c}r_1, \tilde{c}r_2 \notin \partial\tilde{\Omega}_{c,m}\}, \text{ for } 1 \leq m \leq q-1\end{aligned}\tag{11}$$

Remark 3.3.1. By Proposition D.3, $\tilde{\mathcal{D}}_m, \tilde{\mathcal{D}}_m$ are open for $0 \leq m \leq q-1$.

Lemma 3.3.2. Let $c \in \tilde{\mathcal{B}}_m$ and $\tilde{\phi}_c$ any Fatou coordiante for $g_c^q|_{B_{c,0}^*(0)}$. For $1 \leq m \leq q-1$, if $g_c^q(cr_1, 2), g_c^q(1)$ are contained $\Omega_{c,0}^0$ and $\tilde{\phi}_c(cr_{1,2}) = \tilde{\phi}_c(1)$, then $1 = cr_1 = cr_2$; for $m = 0$, $g_c^q(c), g_c^q(1)$ are contained $\Omega_{c,0}^0$ and $\tilde{\phi}_c(c) = \tilde{\phi}_c(1)$, then $1 = c$.

Proof. We only do the proof for $1 \leq m \leq q-1$, the case $m = 0$ is similar. By hypothesis, since $\tilde{\phi}_c$ is injective on $\Omega_{c,0}^0$, we have $g_c^q(cr_1, 2) = g_c^q(1)$. If $cr_1, cr_2, 1$ are not distinct, then $g_c^q(1)$ has at least 5 preimages counting multiplicity, contradicting $\deg(g_c^q|_{B_{c,0}^*(0)}) = 4$. \square

For $0 \leq m \leq q-1$ and $c \in \tilde{\mathcal{D}}_m$ (resp. $c \in \tilde{\mathcal{D}}_0$), let $\phi_c : \overline{\Omega_{c,0}^0} \rightarrow \overline{\mathbb{H}}$ be the Fatou coordiante of $g_c^q|_{B_{c,0}^*(0)}$ normalised by $\phi_c(1) = 0$ (resp. $\phi_c(c) = 0$). Define $h_c : \Omega_{c,0}^0 \rightarrow \Omega_0^0$ by $h_c = \phi^{-1} \circ \phi_c$. Pull back h_c by g_c and P_λ until we reach the critical value $g_c(c)$ (resp. $g_c(1)$):

$$\begin{array}{ccccccc} \Omega_{c,m+p}^{n_c} & \xrightarrow{g_c} & \Omega_{c,m+2p}^{n_c-1} & \xrightarrow{g_c} & \dots & \xrightarrow{g_c} & \Omega_{c,q-p}^1 \xrightarrow{g_c} \Omega_{c,0}^0 \\ \downarrow h_c & & \downarrow h_c & & \downarrow h_c & & \downarrow h_c \\ \Omega_{m+p}^{n_c} & \xrightarrow{P_\lambda} & \Omega_{m+2p}^{n_c-1} & \xrightarrow{P_\lambda} & \dots & \xrightarrow{P_\lambda} & \Omega_{q-p}^1 \xrightarrow{P_\lambda} \Omega_0^0 \end{array}\tag{12}$$

where n_c is the smallest integer such that $\Omega_{c,m+p}^{n_c}$ contains $g_c(c)$ (resp. $g_c(1)$). Moreover at each step h_c is conformal.

Similarly, for $1 \leq m \leq q-1$ and $c \in \tilde{\mathcal{D}}_m$, let $\tilde{\phi}_c : \overline{\tilde{\Omega}_{c,m}} \rightarrow \overline{\tilde{\mathbb{H}}}$ be the Fatou coordinate of $g_c^q|_{B_{c,m}^*(0)}$ normalised by $\tilde{\phi}_c(c) = 0$, where $\tilde{\mathbb{H}} = \tilde{\phi}_c(\tilde{\Omega}_{c,m})$. Define $\tilde{h}_c : \tilde{\Omega}_{c,m} \rightarrow \Omega_0^0$ by $\tilde{h}_c = \phi^{-1} \circ \tilde{\phi}_c$. Pull back \tilde{h}_c by g_c and P_λ until we reach the critical value $g_c(1)$.

$$\begin{array}{ccccccc} \tilde{\Omega}_{c,p}^{n_c} & \xrightarrow{g_c} & \tilde{\Omega}_{c,2p}^{n_c-1} & \xrightarrow{g_c} & \dots & \xrightarrow{g_c} & \tilde{\Omega}_{c,m-p}^1 \xrightarrow{g_c} \tilde{\Omega}_{c,m}^0 \\ \downarrow \tilde{h}_c & & \downarrow \tilde{h}_c & & \downarrow \tilde{h}_c & & \downarrow \tilde{h}_c \\ \Omega_{p-m}^{n_c} & \xrightarrow{P_\lambda} & \Omega_{2p-m}^{n_c-1} & \xrightarrow{P_\lambda} & \dots & \xrightarrow{P_\lambda} & \Omega_{q-p}^1 \xrightarrow{P_\lambda} \Omega_0^0 \end{array}\tag{13}$$

Define four holomorphic mappings

$$\begin{aligned}\Phi_0^{adj} : \tilde{\mathcal{D}}_0 &\longrightarrow B_p^*(0), \quad c \mapsto h_c(g_c(c)) \\ \tilde{\Phi}_0^{adj} : \tilde{\mathcal{D}}_0 &\longrightarrow B_p^*(0), \quad c \mapsto h_c(g_c(1)) \\ \Phi_m^{bit} : \tilde{\mathcal{D}}_m &\longrightarrow B_{m+p}^*(0) \setminus \overline{\Omega_{m+p}^{-s}}, \quad c \mapsto h_c(g_c(c)) \\ \tilde{\Phi}_m^{bit} : \tilde{\mathcal{D}}_m &\longrightarrow B_{p-m}^*(0) \setminus \overline{\Omega_{p-m}^l}, \quad c \mapsto \tilde{h}_c(g_c(1))\end{aligned}\tag{14}$$

where $2 \leq s \leq q$, $0 \leq l \leq q-1$ are the unique integers such that $-sp + m + p = 0 \pmod{q}$ and $lp + p - m = 0 \pmod{q}$.

Lemma 3.3.3. *Let Φ be one of the four holomorphic maps in (14). Then Φ satisfies the following property: let $(c_n) \subset \check{\mathcal{B}}_m$ be a sequence contained in the domain of definition of Φ which converges to $\partial\check{\mathcal{B}}_m$ (resp. $\check{\mathcal{I}}_m$), then $\Phi(c_n)$ converges to the boundary of the corresponding immediate basin (resp. $\partial\Omega_p^{-1} \setminus \{0\}, \partial\Omega_{m+p}^{-s} \setminus \{0\}$ or $\partial\Omega_{p-m}^l \setminus \{0\}$).*

Proof. We only do the proof for Φ_m^{bit} , the others are similar. First we justify that $\Phi_m^{bit}(\check{\mathcal{D}}_m) \subset B_{m+p}^*(0) \setminus \overline{\Omega_{m+p}^{-s}}$. Since Φ_m^{bit} is open, it suffices to prove that its image does not intersect $\overline{\Omega_{m+p}^{-s}}$. If not, then there exists $c \in \check{\mathcal{D}}_m$ such that $g_c(c) \in \Omega_{c,m+p}^{-s}$, hence $g_c^q(cr_1) = g_c^q(cr_2) \in \Omega_{c,0}^{-q}$. Let $\Lambda = g_c^{-q}(\Omega_{c,0}^{-q}) \cap B_{c,0}^*(0)$. Then $g^q : \Lambda \rightarrow \Omega_{c,0}^{-q}$ is of degree 4. Since g_c is a polynomial, each component of Λ is simply connected. Applying Riemann-Hurwitz formula one easily sees that Λ should contain 3 critical points of g_c^q . However $1 \notin \Lambda$ since $1 \in \partial\Omega_{c,0}^0$, so $g_c^q(1) \in \partial\Omega_{c,0}^{-q}$.

Next we verify properness of Φ_m^{bit} . Clearly $\partial\check{\mathcal{D}}_m \subset \check{\mathcal{I}}_m \cup \partial\check{\mathcal{B}}_m$. Let $(c_n) \subset \check{\mathcal{D}}_m$ be a sequence converging to some $c_0 \in \partial\check{\mathcal{D}}_m$. If $c_0 \in \check{\mathcal{I}}_m$, then by Corollary D.2, $g_{c_n}^k(c_n)$ is compactly contained in $\Omega_{c_0,0}^0$ for n large enough, where k is the smallest integer such that $g_{c_n}^k(c_n) \in B_{c_n,0}^*(0)$. Moreover $h_{c_n}(g_{c_n}^k(c_n))$ converges to $\partial\Omega_0^{-q} \setminus \{0\}$ since $\Re\{\phi_{c_n}(g_{c_n}^k(c_n)) - \phi_{c_n}(g_{c_n}^q(1))\}$ converges to 0 and $\Im\{\phi_{c_n}(g_{c_n}^k(c_n)) - \phi_{c_n}(g_{c_n}^q(1))\}$ remains bounded. (In fact we have $h_{c_n}(g_{c_n}^k(c_n)) \rightarrow z_0 \in \partial\Omega_0^{-q} \setminus \{0\}$ with $\phi(z_0) = I_m(c_0)$). This implies that $\Phi_m^{bit}(c_n) = h_{c_n}(c_n) = P_\lambda^{-k} h_{c_n}(g_{c_n}^k(c_n))$ converges to $\partial\Omega_{m+p}^{-s} \setminus \{0\}$.

So let $c_0 \in \partial\check{\mathcal{B}}_m$. We want to prove that $\Phi_m^{bit}(c_n)$ converges to $\partial B_{m+p}^*(0)$. Suppose the contrary that, up to taking a subsequence, $\Phi_m^{bit}(c_n)$ converges to $z_0 \in B_{m+p}^*(0)$. Clearly $\{(h_{c_n}|_{\Omega_{c_n,0}^0})^{-1}\}$ is a normal family on Ω_0^0 . Up to taking a subsequence, suppose $h_{c_n}^{-1}$ converges uniformly to ψ . We claim that $\psi(\Omega_0^0)$ does not intersect J_{c_0} , the Julia set of g_{c_0} . Indeed, if not, then there exists a repelling periodic point $x_{c_0} \in J_{c_0} \cap \psi(\Omega_0^0)$ (since the repelling cycles are dense in J_{c_0}). Notice that x_{c_0} moves holomorphically for c in a small neighborhood of c_0 , which gives a repelling periodic point x_c of g_c . However for n large enough, $x_c \in h_{c_n}^{-1}(\Omega_0^0) \subset B_{c_n,0}^*(0)$, a contradiction. Therefore $\psi(\Omega_0^0)$ and $g_{c_0}^q(1)$ are contained in the same Fatou component of g_{c_0} . On the other hand, take $s \geq 0$ such that $P_\lambda^s(z_0) \in \Omega_0^0$, then by diagram (12), $g_{c_n}^s(c_n) \in \Omega_{c_n,0}^0$ and $h_{c_n}(g_{c_n}^s(c_n))$ converges to $P_\lambda^s(z_0)$. Therefore

$$\psi(P_\lambda^s(z_0)) = \lim_{n \rightarrow \infty} h_{c_n}^{-1}(h_{c_n}(g_{c_n}^s(c_n))) = g_{c_0}^s(c_0) \in \psi(\Omega_0^0)$$

which means that $1, c_0$ are eventually attracted by the same Fatou component, contradicts $c_0 \in \partial\check{\mathcal{B}}_m$. \square

Proposition 3.3.4. *The mappings $\Phi_m^{bit}, \tilde{\Phi}_m^{bit}$ in (14) are bijective.*

Proof. We only do for Φ_m^{bit} , the other is similar. By Lemma 3.3.3, it suffices to verify injectivity. Suppose $\Phi_m^{bit}(c_1) = \Phi_m^{bit}(c_2)$. Starting from the conjugacy $g_{c_2} \circ \phi_{c_2}^{-1} \circ \phi_{c_1} = \phi_{c_2}^{-1} \circ \phi_{c_1} \circ g_{c_1}$ on $\Omega_{c_1,0}^0$, lift $\phi_{c_2}^{-1} \circ \phi_{c_1}$ by g_{c_1}, g_{c_2} to $\Omega_{c_1,k}^n \subset B_{c_1,k}^*$ (recall definition of $\Omega_{c,k}^n$ at the beginning of 3.3). Denote by φ the lifting of $\phi_{c_2}^{-1} \circ \phi_{c_1} \circ g_{c_1}$. The only possible issue that might stop the lifting is when $k = m+p$ and $g_{c_1}(c_1) \in \Omega_{c_1,k}^n$. However $\Phi_m^{bit}(c_1) = \Phi_m^{bit}(c_2)$ implies $\varphi(g_{c_1}(c_1)) = g_{c_2}(c_2)$, which ensures that the lifting is still valid (also notice that $g_{c_1}|_{B_{c_1,m}^*}, g_{c_1}|_{B_{c_1,m}^*}$ are of degree 2). So φ is extended to $\bigcup_i B_{c_1,i}^*$ and hence to all filled-in Julia set K_{c_1} .

On the other hand, take a connected open set $\check{\mathcal{V}} \subset \check{\mathcal{B}}_m$ linking c_1, c_2 , on which the Böttcher coordinate at ∞ depends analytically on c . Therefore for $c \in \check{\mathcal{V}}$, $\psi_c := (\phi_c^\infty)^{-1} \circ \phi_{c_1}^\infty$ is a dynamical holomorphic motion on $C \setminus \check{K}_{c_1}$, and can be quasiconformally extend to \mathbb{C} (Słodkowski's theorem). In particular, ψ_{c_2} conjugates g_{c_1} to g_{c_2} on $\mathbb{C} \setminus \check{K}_{c_1}$, coincides with φ on ∂K_{c_1} . Applying Rickman's lemma, the global conjugacy defined by sewing φ and ψ_{c_2} is conformal, i.e. identity, so $c_1 = c_2$. \square

The injectivity for $\Phi_0^{adj}, \tilde{\Phi}_0^{adj}$ is more subtle:

Proposition 3.3.5. *Let $c_1, c_2 \in \check{\mathcal{D}}_0$ be such that $\Phi_0^{adj}(c_1) = \Phi_0^{adj}(c_2)$.*

1. If $\Phi_0^{\text{adj}}(c_1) \in \overline{\Omega_p^{-1}}$, then $c_1 = c_2$;
2. if not, then for $i = 1, 2$, $(g_{c_i}^{-1})^{-1}(\overline{\Omega_{c_i,p}^{-1}})$ has two connected components, one of which has the figure of a filled eight, denoted by H_{c_i} . H_{c_i} cuts $B_{c_i,0}^*(0)$ into two connected components $\Delta_{c_i}^+, \Delta_{c_i}^-$ (+ is on the right-hand side of -). If $c_i \in \Delta_{c_i}^+$ for $i = 1, 2$ or $c_i \in \Delta_{c_i}^-$ for $i = 1, 2$, then $c_1 = c_2$.

The same results hold for $\tilde{\Phi}_0^{\text{adj}}$.

Proof. We do the proof for the second point. The first is similar. Without loss of generality suppose $c_i \in \Delta_{c_i}^-$ for $i = 1, 2$. Like what we did at the beginning of the proof of Proposition 3.3.4, lift $\varphi = \phi_{c_2}^{-1} \circ \phi_{c_1}$ by g_{c_1}, g_{c_2} . Similarly, when we lift φ to $\Omega_{c_1,0}^n$ with n_0 the smallest integer such that $g_{c_1}^q(c_1) \in \Omega_{c_1,0}^{n_0}$, it is not clear whether φ can be lifted once more to $\Omega_{c_1,0}^{n_0+q}$, where $n_0 = k_0 q$ for some $k_0 \geq 1$. Notice that for $i = 1, 2$, $1 \leq k \leq k_0 + 1$, $\Omega_{c_i,0}^{kq}$ is simply connected with piecewise smooth boundary intersecting $\partial B_{c_i,0}^*(0)$ at 2^k points, and $\Omega_{c_i,0}^{kq} \setminus \Omega_{c_i,0}^{(k-1)q}$ has 2^{k-1} connected components $D_{\epsilon_0 \dots \epsilon_{k-2}}^i$ with

$$\epsilon_j = \begin{cases} 0, & \text{if } g_{c_i}^{jq}(D_{\epsilon_0 \dots \epsilon_{k-2}}^i \cap \overline{\Omega_{c_i,0}^{kq}}) \subset \overline{\Delta_{c_i}^-} \\ 1, & \text{if } g_{c_i}^{jq}(D_{\epsilon_0 \dots \epsilon_{k-2}}^i \cap \overline{\Omega_{c_i,0}^{kq}}) \subset \overline{\Delta_{c_i}^+} \end{cases}$$

Since $\Phi_0^{\text{adj}}(c_1) = \Phi_0^{\text{adj}}(c_2)$, there exists $\tilde{\epsilon}_0 \dots \tilde{\epsilon}_{k-2}$ such that $g_{c_i}(c_i) \in D_{\tilde{\epsilon}_0 \dots \tilde{\epsilon}_{k-2}}^i$. So $c_i \in D_{\delta_i \tilde{\epsilon}_0 \dots \tilde{\epsilon}_{k-2}}^i$. But by hypothesis $c_i \in \Delta_{c_i}^-$, so $\delta_i = 0$ for $i = 1, 2$. Therefore we can extend φ to $\Omega_{c_1,0}^{(k_0+1)q}$ by assigning injectively $D_{\epsilon_0 \dots \epsilon_{k_0-1}}^1$ onto $D_{\tilde{\epsilon}_0 \dots \tilde{\epsilon}_{k_0-1}}^2$. Notice that $\Omega_{c_1,0}^{(k_0+1)q}$ contain the two critical points and the two cocritical points of $g_{c_1}^q|_{B_{c_1,0}^*(0)}$, thus the lifting process is valid for all $k \geq k_0 + 1$, so φ can be extended to $B_{c_1,0}^*(0)$, and hence to the filled-in Julia set K_{c_1} . Now apply the same strategy as in the second paragraph of the proof of Proposition 3.3.4, we conclude that $c_1 = c_2$. \square

3.4 Describing the special locus $\tilde{\mathcal{I}}_m$

Recall the definition of $\tilde{\mathcal{I}}_m$ in (10). For $m = 0$, $c \in \tilde{\mathcal{I}}_0$, define $I_0(c) = \Im(\phi_c(c) - \phi_c(1))$; for $1 \leq m \leq q-1$, $c \in \tilde{\mathcal{I}}_m$, define $I_m(c) = \Im(\phi_c(cr) - \phi_c(1))$ where $cr = cr_1$ or cr_2 , since $\phi_c(cr_1) = \phi_c(cr_2)$.

Lemma 3.4.1. *For every m -component $\check{\mathcal{B}}_m$, $I_m^{-1}(0) \neq \emptyset$.*

Proof. Let Φ be the corresponding mapping in (14) defined on $\tilde{\mathcal{D}}_m$. Clearly $P_\lambda^s(-\frac{\lambda}{2}) \in \partial\Phi(\mathcal{N}) \cap B_{\frac{m+p}{m+p}}^*(0)$. Thus by Lemma 3.3.3, there exists a sequence of parameters $(c_k) \subset \mathcal{N}$ converging to $c_0 \in \tilde{\mathcal{I}}_m$ such that $\Phi(c_k)$ converges to $P_\lambda^s(-\frac{\lambda}{2})$. Recall that $\Phi(c) = h_c(g_c(c))$, thus $P_\lambda^{q-s} \circ h_{c_k}(g_{c_k}(c_k)) = h_c(g_{c_k}^{q-s+1}(c_k))$ converges to $P_\lambda^q(-\frac{\lambda}{2})$, which implies that $\phi_{c_k}(g_{c_k}^q(c)) - \phi_{c_k}(g_{c_k}^q(1)) \rightarrow 0$ if $m = 0$ and $\phi_{c_k}(g_{c_k}^q(cr_{1,2})) - \phi_{c_k}(g_{c_k}^q(1)) \rightarrow 0$ if $1 \leq m \leq q-1$. Thus at c_0 we have respectively $\phi_{c_0}(c) = \phi_{c_0}(1)$ and $\phi_{c_0}(cr_{1,2}) = \phi_{c_0}(1)$. By Lemma 3.3.2, $c_0 = 1$ resp. $cr_{1,2} = 1$, i.e. $I_m(c_0) = 0$. \square

Corollary 3.4.2. *For family g_c the 0-component is unique. It contains $c = 1$ and is symmetric with respect to $\tau : c \mapsto 1/c$.*

Proof. By Lemma 3.4.1, $I_0^{-1}(0) \subset \check{\mathcal{B}}_0$, while by Lemma 3.3.2, $I_0^{-1}(0) = 1$. Thus $\check{\mathcal{B}}_0$ is unique. Since $\tau(\check{\mathcal{B}}_0)$ is also a 0-component, hence $\tau(\check{\mathcal{B}}_0) = \check{\mathcal{B}}_0$. \square

Lemma 3.4.3. *Let $0 \leq m \leq q-1$, $\check{\mathcal{B}}_m$ be a m -component, $\tilde{\mathcal{I}}_m$ be as in (10). Then $I_m : \tilde{\mathcal{I}}_m \rightarrow (-\infty, +\infty)$ is injective.*

Proof. For $1 \leq m \leq q-1$, adapt the proof of Proposition 3.3.4; for $m = 0$ adapt that of Proposition 3.3.5. \square

Lemma 3.4.4. *If there exists $c_{t_0} \in \check{\mathcal{I}}_m$ such that $t_0 := I_m(c_0) > 0$ (resp. < 0), then for any $t > 0$ (resp. < 0) there is $c_t \in \check{\mathcal{I}}_m$ with $I_m(c_t) = t$. Moreover $I_m^{-1}((-\infty, 0))$, $I_m^{-1}((0, +\infty))$ are curves parametrized by I_m^{-1} .*

Proof. This can be shown by quasiconformal deformation. \square

Lemma 3.4.5. *Let $(c_k) \subset \check{\mathcal{I}}_m$ be a sequence of parameters. If*

1. $I_m(c_k) \rightarrow \pm\infty$, then c_k converges to a double parabolic parameter;
2. $I_m(c_k) \rightarrow 0$, then c_k converges to $I_m^{-1}(0)$.

Proof. Let c_0 be any accumulation point of c_k .

1. If $I_m(c_k) \rightarrow \pm\infty$, then clearly $c_0 \in \partial\check{\mathcal{B}}_m$. If c_0 is not double parabolic, then only one critical point is in the immediate basins, say 1. One can then apply Proposition D.3, which implies that for c near c_0 , 1 is always on the boundary of the maximal petal of g_c^q while the other critical points of g_c^q are not, contradicting the definition of $\check{\mathcal{I}}_m$.

2. If $I_m(c_k) \rightarrow 0$. It suffices to prove that $c_0 \in \check{\mathcal{B}}_m$, since then by taking limit in k , we get $I_m(c_0) = 0$. For the same reason as above, if c_0 is not double parabolic, then $c_0 \notin \partial\check{\mathcal{B}}_m$ and $c_0 \in \check{\mathcal{B}}_m$. So it remains to show that c_0 is not double parabolic. Suppose the contrary. Without loss of generality we assume that $I_m(c_k) > 0$. Then by the first point and Lemma 3.4.4, $I_m^{-1}((0, +\infty))$ is a curve separating $\check{\mathcal{B}}_m$ into two simply connected components ($\check{\mathcal{B}}_m$ is simply connected, Lemma 2.2.4). Let \mathcal{N} be the one not containing $I_m^{-1}(0)$. This \mathcal{N} exists since $I_m^{-1}(0)$ is a single point by Lemma 3.4.3. Then $\mathcal{N} \setminus \check{\mathcal{I}}_m$ is $\check{\mathcal{D}}_m$ or $\check{\check{\mathcal{D}}}_m$. But recall that $\partial\check{\mathcal{D}}_m \cap \check{\mathcal{B}}_m = \partial\check{\check{\mathcal{D}}}_m \cap \check{\mathcal{B}}_m = \check{\mathcal{I}}_m$, so $I_m^{-1}(0) \in \partial(\mathcal{N} \setminus \check{\mathcal{I}}_m)$, a contradiction. \square

Proposition 3.4.6. *For $0 \leq m \leq q-1$, $\check{\mathcal{I}}_m$ is a curve parametrized by $I_m^{-1} : (-\infty, +\infty) \rightarrow \check{\mathcal{I}}_m$. Moreover, when $q > 1$, $\check{\mathcal{I}}_m$ has two different end points; when $q = 1$, $\check{\mathcal{I}}_0$ has only one end point -1 .*

Proof. First we prove that I_m^{-1} is a parametrisation. By Lemma 3.4.4 and 3.4.5, it suffices to prove that $I_m^{-1}((-\infty, 0))$ and $I_m^{-1}((0, +\infty))$ are not empty. Consider the case $m = 0$: by Corollary 3.4.2, $\check{\mathcal{B}}_0$ is unique and symmetric with respect to $\tau : c \mapsto 1/c$. Moreover it is easy to see that $\tau(\check{\mathcal{D}}_0) = \check{\check{\mathcal{D}}}_0$, which are non empty. If both $I_0^{-1}((-\infty, 0))$, $I_0^{-1}((0, +\infty))$ are empty, then one of $\check{\mathcal{D}}_0$, $\check{\check{\mathcal{D}}}_0$ is empty, a contradiction. So suppose $I_0^{-1}((-\infty, 0)) \neq \emptyset$, hence $\tau(I_0^{-1}((-\infty, 0))) = I_0^{-1}((0, +\infty)) \neq \emptyset$.

Next we prove for $1 \leq m \leq q-1$. By Proposition 3.3.4, $\Phi_m^{bit} : \check{\mathcal{D}}_m \rightarrow B_{m+p}^*(0) \setminus \overline{\Omega_{m+p}^{-s}}$ is conformal, so for any $z \in \partial\Omega_{m+p}^{-s} \setminus \{0\}$, there exists $c_n \rightarrow c_0 \in \check{\mathcal{I}}_m$ such that $\Phi_m^{bit}(c_n) \rightarrow z$. Thus $I_m(c_0)$ can take arbitrary value in \mathbb{R} by choosing properly z .

Now we investigate the end points of $\check{\mathcal{I}}_m$. First consider the case $m = 0$. Notice that $\check{\mathcal{I}}_0$ is symmetric with respect to $\tau : c \mapsto \frac{1}{c}$. If $q = 0$, it is easy to see that there is only one double parabolic parameter -1 , hence by Lemma 3.4.5 the end point of $\check{\mathcal{I}}_0$ is -1 . If $q > 1$, we prove that $\check{\mathcal{I}}_0$ ends at two different points. Suppose the contrary, then $\check{\mathcal{I}}_0$ must land at -1 since $\check{\mathcal{I}}_0 = \tau(\check{\mathcal{I}}_0)$. By Lemma 2.1.1, $\sigma\iota^{-1}(\check{\mathcal{I}}_0)$ is a curve symmetric with respect to $z \mapsto -z$, starting from $\sqrt{3}e^{2\pi i \frac{p}{q}}$ and ending at $-\sqrt{3}e^{2\pi i \frac{p}{q}}$. Moreover $0 = \sigma\iota^{-1}(-1)$ is a double parabolic parameter for the family f_a . By Lemma 3.2.4, 0 is of type 0 and $q-1$, i.e. $q = 1$, a contradiction.

Now we consider the case $1 \leq m \leq q-1$. If not, then $\check{\mathcal{I}}_m$ is a simple closed curve. We claim that $\check{\mathcal{I}}_m$ must separate $0, \infty$. Suppose not, then $\check{\mathcal{I}}_m$ will bound a simply connected region $\mathcal{O} \subset \check{\mathcal{B}}_m$, since by Proposition 2.2.3 and Lemma 2.1.1 there are only two connected components $\tau\check{\mathcal{H}}_\infty, \check{\mathcal{H}}_\infty$ of $\mathbb{C}^* \setminus \check{\mathcal{C}}_\lambda$, which are punctured neighborhoods of 0 and ∞ respectively. Thus \mathcal{O} is either $\check{\mathcal{D}}_m$ or $\check{\check{\mathcal{D}}}_m$ and Φ_m^{bit} or $\check{\Phi}_m^{bit}$ is well-defined on \mathcal{O} . But this contradicts Lemma 3.3.3 since $\partial\mathcal{O} = \check{\mathcal{I}}_m$. So $\check{\mathcal{I}}_m$ is a closed curve separating $0, \infty$. Now we claim that $\check{\mathcal{B}}_m$ is invariant under $\tau : c \mapsto 1/c$. Suppose not, then $\tau(\check{\mathcal{B}}_m) \cap \check{\mathcal{B}}_m = \emptyset$. In particular, $\tau(\check{\mathcal{I}}_m) \cap \check{\mathcal{I}}_m = \emptyset$

and both their closures separate $0, \infty$. Therefore $\mathbb{C}^* \setminus (\overline{\check{\mathcal{I}}_m \cup \tau(\check{\mathcal{I}}_m)})$ has a connected component \mathcal{V} which do not intersect $\tau\check{\mathcal{H}}_\infty, \check{\mathcal{H}}_\infty$. By MSS J -stability theorem, g_c is stable on \mathcal{V} , hence $\check{\mathcal{I}}_m, \tau(\check{\mathcal{I}}_m)$ are in fact in the same parabolic component $\check{\mathcal{B}}_m$, a contradiction. Since $\check{\mathcal{B}}_m$ is invariant under τ , so is $\overline{\check{\mathcal{B}}_m}$. Write $\partial\check{\mathcal{B}}_m = c_0 \cup \mathcal{K}_0 \cup \mathcal{K}_\infty$, where c_0 is the end point of $\check{\mathcal{I}}_m$, \mathcal{K}_0 is the connected component of $\partial\check{\mathcal{B}}_m \setminus \{c_0\}$ contained in the bounded component of $\mathbb{C} \setminus \overline{\check{\mathcal{I}}_m}$ and \mathcal{K}_∞ the one contained in the unbounded component of $\mathbb{C} \setminus \overline{\check{\mathcal{I}}_m}$. Clearly $\mathcal{K}_0 \subset \partial\tau\check{\mathcal{H}}_\infty \setminus \partial\check{\mathcal{H}}_\infty$ and $\mathcal{K}_\infty \subset \partial\check{\mathcal{H}}_\infty \setminus \partial\tau\check{\mathcal{H}}_\infty$. By the relation $\frac{1}{c}g_c(cz) = g_{1/c}(z)$ we conclude that $\tau(\mathcal{K}_0) = \mathcal{K}_\infty$ and $\tau(\mathcal{K}_\infty) = \mathcal{K}_0$, hence $\tau(c_0) = c_0$, $c_0 = \pm 1$, while $1 \in \check{\mathcal{B}}_0$, so $c_0 = -1$. Thus by Lemma 2.1.1, $\sigma\iota^{-1}(-1) = 0$ and $\sigma\iota^{-1}(\check{\mathcal{I}}_m)$ is a closed curve separating \mathbb{C} into two connected components, each of which intersects $\sigma(\hat{\mathcal{H}}_\infty \cup \tau\hat{\mathcal{H}}_\infty)$, the complementary of the connected locus for the family f_a . But this contradicts Proposition 2.2.3, which says that $\sigma(\hat{\mathcal{H}}_\infty \cup \tau\hat{\mathcal{H}}_0)$ has only one connected component. \square

Corollary 3.4.7. *For $0 \leq m \leq q-1$, the m -component for the family g_c is unique.*

Proof. The case $m = 0$ has already been justified in Corollary 3.4.2. We prove for $m \geq 1$. For every $0 \leq k \leq q-1$ pick a k -component. By the above proposition and Lemma 3.2.4, $\mathcal{Z} = \bigcup_{k=0}^{q-1} \check{\mathcal{I}}_k$ is a simple closed curve surrounding 0. If for some $m \geq 1$ there exists another m -component $\check{\mathcal{B}}'_m$ with corresponding $\check{\mathcal{I}}'_m$, then by the above proposition, $\mathbb{C}^* \setminus (\mathcal{Z} \cup \overline{\check{\mathcal{I}}'_m})$ has a component which do not intersect $\check{\mathcal{H}}_\infty, \tau\check{\mathcal{H}}_\infty$ while it intersects $\partial\check{\mathcal{C}}_\lambda$, a contradiction. \square

3.5 Parametrizations transferred for family f_a

The first thing to do here is to find a dynamically defined curve $\mathcal{I} \subset \text{Per}_1(e^{2\pi i \frac{p}{q}})$ symmetric with respect to $a \mapsto -a$ linking $-\sqrt{3}\lambda, \sqrt{3}\lambda$, so that on $\mathbb{C} \setminus \mathcal{I}$ we can define the two critical points of f_a such that they vary analytically for $a \in \mathbb{C} \setminus \mathcal{I}$.

$\sigma\iota^{-1}(\bigcup_{m=0}^{q-1} \check{\mathcal{I}}_m)$ is a "good" candidate, but it is not necessarily symmetric with respect to $a \mapsto -a$. In order to solve this problem, let us first notice that if q is odd, then $c = -1$ is double parabolic; if q is even, then $c = -1 \in \check{\mathcal{D}}_{\frac{q}{2}}$. Let $\rho = \Re\{\phi_{-1}(g_{-1}^{\frac{q}{2}}(-1))\}$, then $0 < \rho < 1$. Let $\Lambda \subsetneq \Omega_0^0 \subset B_0^*(0)$ be the petal of $P = P_\lambda$ of level ρ . Since

$$\Phi_{\frac{q}{2}}^{bit} : \check{\mathcal{D}}_{\frac{q}{2}} \longrightarrow B_{\frac{q}{2}+p}^*(0) \setminus \overline{\Omega_{\frac{q}{2}+p}^{-\frac{q}{2}-1}}, \quad c \mapsto h_c(g_c(c))$$

is an isomorphism (Lemma 3.3.3), $(\Phi_{\frac{q}{2}}^{bit})^{-1}(P^{-\frac{q}{2}+1}(\partial\Lambda))$ is a curve linking the two double parabolic parameters on $\partial\check{\mathcal{B}}_{\frac{q}{2}}$. Let $\check{\gamma}$ be the subcurve of it linking the double parabolic parameter on $\partial\check{\mathcal{B}}_{\frac{q}{2}} \cap \partial\check{\mathcal{B}}_{\frac{q}{2}-1}$ and $c = -1$.

Let $\mathcal{Z} = \bigcup_{m=0}^{\lfloor \frac{q}{2} \rfloor} \check{\mathcal{I}}_m$ if q is odd or $\mathcal{Z} = \bigcup_{m=0}^{\lfloor \frac{q-1}{2} \rfloor} \check{\mathcal{I}}_m \cup \check{\gamma}$ if q even. Then \mathcal{Z} is a curve linking $c = 1$ and $c = -1$. Now $\iota^{-1}(\mathcal{Z})$ (in the s -plane, recall in (4)) has two connected component. Take the one containing $s = -1$ and denote its image under σ by \mathcal{G} . Set $\mathcal{I} = \mathcal{G} \cup -\mathcal{G}$. Thus \mathcal{I} is a curve passing $a = 0$, symmetric under $a \mapsto -a$. Set $\mathcal{I}_m := \mathcal{I} \cap \mathcal{B}_m$.

Remark 3.5.1. *When q is even and $m = \frac{q}{2}$, we have $\mathcal{I}_m \cap \sigma\iota^{-1}(\check{\mathcal{I}}_m) = \emptyset$. To see this, it suffices to prove that $\iota\sigma^{-1}(\mathcal{I}_m) \cap \check{\mathcal{B}}_m \cap \check{\mathcal{I}}_m = \emptyset$. Indeed, by construction $\iota\sigma^{-1}(\mathcal{I}_m) \cap \check{\mathcal{B}}_m = \check{\gamma} \cup \tau\check{\gamma}$ ($\tau : c \mapsto 1/c$) and $\check{\gamma} \cap \check{\mathcal{I}}_m = \emptyset$. To see that $\tau\check{\gamma} \cap \check{\mathcal{I}}_m = \emptyset$, it suffices to justify that for $c \in \check{\gamma}$, $\Re\{\phi_c(g_c^{\frac{q}{2}}(1)) - \phi_c(g_c^q(c))\} < 0$. This is clear since $g_c^{\frac{q}{2}}(c) \in \Omega_{c,0}^0$, while $1 \in \partial\Omega_{c,0}^0$.*

Let $a \mapsto \sqrt{a^2 - 3\lambda}$ be the inverse branch defined on $\mathbb{C} \setminus \mathcal{I}$ such that $(a - \sqrt{a^2 - 3\lambda}) \rightarrow 0$ as $|a| \rightarrow +\infty$. Define $c_\pm(a) = \frac{-a \pm \sqrt{a^2 - 3\lambda}}{3}$ and let $v_\pm(a) = f_a(c_\pm(a))$.

Proof of Proposition II. It is just a summary of what we have obtained for the family f_a :

- For every $0 \leq m \leq q$, there is a unique m -component \mathcal{B}_m (Definition 3.1.5). This is direct from Corollary 3.4.7 and the relation between families g_c and f_a (Lemma 2.1.1). Moreover, since \mathcal{C}_λ is symmetric under $a \mapsto -a$, we have $\mathcal{B}_m = -\mathcal{B}_{q-m}$.
- For every $0 \leq m \leq q-1$, there is a unique double parabolic parameter \mathbf{a}_m of m -type (Definition 3.2.3). Moreover $\mathbf{a}_m \in \partial\mathcal{B}_m \cap \partial\mathcal{B}_{m+1}$ (Lemma 3.2.4).
- From Lemma 3.2.4, 3.4.5 and Proposition 3.4.6, the special curve \mathcal{I} defined above passes $\bigcup_{m=0}^q \mathcal{B}_m$ in the following order:

$$\mathcal{B}_0, \mathbf{a}_0, \mathcal{B}_1, \mathbf{a}_1, \mathcal{B}_2, \dots, \mathcal{B}_{q-2}, \mathbf{a}_{q-2}, \mathcal{B}_{q-1}, \mathbf{a}_{q-1}, \mathcal{B}_q.$$

Moreover for $a \in \mathbb{C} \setminus \mathcal{I}$, $c_+(a)$ is always on the boundary of the maximal petal for $f_a^q|_{B_{a,0}^*(0)}$.

□




Figure 4: The curves \mathcal{I} for \mathcal{C}_λ with $\lambda = e^{\frac{2\pi i}{3}}$ and $e^{\frac{2\pi i}{4}}$.

The parametrization of m -components for the family g_c (recall (14)) can be transferred by $\sigma\iota^{-1}$ to the family f_a . More precisely:

- For $m = 0$, define $\Psi_0^{adj} : \mathcal{B}_m \setminus \mathcal{I}_0 \rightarrow B_p^*(0)$ by $\Psi_0^{adj} = \Phi_0^{adj} \circ \iota \circ \sigma^{-1}$, where σ^{-1} the inverse branch such that $\iota \circ \sigma^{-1}(\mathcal{B}_m \setminus \mathcal{I}_0) = \tilde{\mathcal{D}}_0$.
- For $1 \leq m < \lfloor \frac{q}{2} \rfloor$, define $\mathcal{D}_m = \sigma\iota^{-1}(\tilde{\mathcal{D}}_m)$ and $\tilde{\mathcal{D}}_m = \sigma\iota^{-1}(\tilde{\tilde{\mathcal{D}}}_m)$, where ι^{-1} is the inverse branch such that $\sigma\iota^{-1}(\tilde{\mathcal{B}}_m) = \mathcal{B}_m$. Define $\Psi_m^{bit} : \mathcal{D}_m \rightarrow B_{\frac{m+p}{2}}^*(0) \setminus \overline{\Omega_{m+p}^{-s}}$ by $\Psi_m^{bit} = \Phi_m^{bit} \circ \iota \circ \sigma^{-1}$ and $\tilde{\Psi}_m^{bit} : \tilde{\mathcal{D}}_m \rightarrow B_{\frac{p-m}{2}}^*(0) \setminus \overline{\Omega_{p-m}^l}$ by $\tilde{\Psi}_m^{bit} = \tilde{\Phi}_m^{bit} \circ \iota \circ \sigma^{-1}$, where σ^{-1} is the inverse branch such that $\iota\sigma^{-1}(\mathcal{B}_m) = \tilde{\mathcal{B}}_m$.
- For $m = \frac{q}{2}$, \mathcal{B}_m is divided by $\sigma\iota^{-1}(\tilde{\mathcal{I}}_m)$ into two components. Let \mathcal{D}_m be the one that is contained in a connected component of $\mathcal{B}_m \setminus \mathcal{I}_m$ (Remark 3.5.1). Define similarly Ψ_m^{bit} on \mathcal{D}_m .

From Proposition 3.3.4, we have

Proposition 3.5.2. For $1 \leq m \leq \lfloor \frac{q}{2} \rfloor$, $\Psi_m^{bit}, \tilde{\Psi}_m^{bit}$ are isomorphisms. For $m = \frac{q+1}{2}$, Ψ_m^{bit} is an isomorphism.

Capture components can also be parametrized by locating $v_-(a)$:

Proposition 3.5.3. *Let \mathcal{U}_k be a capture component of depth $k \geq 1$. For $a \in \mathcal{U}_k$, suppose $f_a^k(U_a) = B_{a,l}^*(0)$, where U_a is the Fatou component containing $c_-(a)$. Then $\Psi_{\mathcal{U}_k} : \mathcal{U}_k \rightarrow B_l^*(0)$ defined by $a \mapsto h_a(f^k(c_-(a)))$ is an isomorphism, where h_a is the conjugating map between $f_a^q|_{B_{a,l}^*(0)}$ et $P^q|_{B_l^*(0)}$.*

Proof. The proof goes exactly the same as Proposition 3.3.4. \square

Proposition 3.5.4. *For $m = 0$, $(\Psi_0^{adj})^{-1}(\Omega_p^{-1})$ is a topological disk whose boundary is a piecewise smooth closed curve passing $\sqrt{3\lambda}, 2\sqrt{\lambda}$. Let $\mathcal{D}_0, \tilde{\mathcal{D}}_0$ be the two connected component of $(\mathcal{B}_0 \setminus \mathcal{I}_0) \setminus \Psi_0^{adj}(\overline{\Omega_p^{-1}})$. Then $\Psi_0^{adj}|_{\mathcal{D}_0}, \Psi_0^{adj}|_{\tilde{\mathcal{D}}_0}$ are isomorphisms with image $B_p^*(0) \setminus \overline{\Omega_p^{-1}}$. See Figure 5.*

Proof. First of all $\Psi_0^{adj}(\Omega_p^{-1})$ is not empty. Indeed, there is a holomorphic motion induced by $\phi_a^{-1} \circ \phi_{a_0}$ of $\overline{\Omega_{a,p}^{-1}}$ in a small neighborhood of $a_0 = 2\sqrt{\lambda}$. Notice that if $a_0 = 2\sqrt{\lambda}$, then $v_-(a_0) = 0$. Let $z_h \in \partial\Omega_{a,0}^{-q}$ be such that $\phi_{a_0}(z_h) = 1/h \in \mathbb{R}$. Apply Rouché's Theorem to $F(a, h) = v_-(a) - f_a^{1-q}\phi_a^{-1}\phi_{a_0}(z_h)$ for h near 0 (take f_a^{1-q} to be the inverse branch of $f_a^{q-1}|_{\overline{\partial\Omega_{a,p}^{-1}}}$), there is a sequence of $a_n \in \mathcal{B}_m \setminus \mathcal{I}_0$ converging to $2\sqrt{\lambda}$ such that $v_-(a_n) \in \overline{\Omega_{a,p}^{-1}}$, i.e. $a_n \in (\Psi_0^{adj})^{-1}(\partial\Omega_p^{-1})$ and $a_n \rightarrow 2\sqrt{\lambda}$. So $\Psi_0^{adj}(\Omega_p^{-1}) \neq \emptyset$.

Next we investigate the end points of $\partial[(\Psi_0^{adj})^{-1}(\Omega_p^{-1})]$. Notice that by properness of Ψ_0^{adj} (Lemma 3.3.3) and stability of Fatou coordinate,

$$\partial[(\Psi_0^{adj})^{-1}(\Omega_p^{-1})] \subset \{\mathbf{a}_0, 2\sqrt{\lambda}\} \cup (\Psi_0^{adj})^{-1}(\partial\Omega_p^{-1}) \cup \mathcal{I}_0.$$

By Proposition 3.3.5 and a quasiconformal deformation argument, $\partial[(\Psi_0^{adj})^{-1}(\partial\Omega_p^{-1})]$ is the closure of the union of two curves γ_1, γ_2 having $2\sqrt{\lambda}$ as a common end point. We parametrize γ_1, γ_2 by $I(a) = \Im\phi[c_-(a)]$. By the above analysis, we see that as $|I_a| \rightarrow \infty$, $a \rightarrow 2\sqrt{\lambda}$. We want to prove that as $|I_a| \rightarrow 0$, $a \rightarrow \sqrt{3\lambda}$. By Lemma 3.3.2, it suffices to show that a do not accumulate at $\partial\mathcal{B}_0$. Suppose the contrary. If $a \in \gamma_1$ accumualtes at $a_0 \in \partial\mathcal{B}_0 \setminus \overline{\mathcal{A}_{p/q}}$, then by stability of Fatou coordinate, $a_0 = 2\sqrt{\lambda}$, then γ_1 surround a topological disk U such that $\overline{\Psi_0^{adj}(U)} \cap \partial\mathcal{B}_0 = \{2\sqrt{\lambda}\}$. But $\Psi_0^{adj}(\partial U \setminus \{2\sqrt{\lambda}\}) = \Psi_0^{adj}(\gamma_1)$ is a semi-arc of $\partial\Omega_p^{-1}$, which does not separate $B_p^*(0)$, a contradiction. So it remains to exclude the case where both γ_1, γ_2 land at \mathbf{a}_0 as $|I(a)| \rightarrow \infty$. Suppose we have this, Then $\gamma_1 \cup \gamma_2$ bounds a topological disk V such that $V \setminus \mathcal{I}_0$ is sent conformally onto Ω_p^{-1} by Ψ_0^{adj} (Proposition 3.3.5). Since $\partial V \cup \mathcal{I}_0$ is locally connected, $(\Psi_0^{adj})|_{\Omega_p^{-1}}^{-1}$ can be continuously extended to $\partial\Omega_p^{-1}$ and in particular $(\Psi_0^{adj})^{-1}(0) = 2\sqrt{\lambda}$. On the other hand, by Lemma 3.3.3, when $a \in \mathcal{B}_0 \setminus \mathcal{I}_0$ tends to \mathbf{a}_0 , $\Psi_0^{adj}(a) \rightarrow \partial B_p^*(0)$. Hence $(\Psi_0^{adj})^{-1}(0) = \mathbf{a}_0$, a contradiction.

The rest of the proposition is immediate by Lemma 3.3.3 and Proposition 3.3.5. \square




Figure 5: A zoom of \mathcal{B}_0 in \mathcal{C}_λ with $\lambda = e^{2\pi i \frac{1}{4}}$. The union of curves in red, blue and orange is equipotentials and external rays of level 0,1,2 in \mathcal{D}_0 . $W = (\Psi_0^{adj})^{-1}(\Omega_p^{-1})$. $a = \sqrt{3\lambda}$ is the parameter such that $c_+(a) = c_-(a)$; $a = \sqrt{3\lambda}$ is the parameter such that $v_-(a) = 0$.




Figure 6: Equipotentials in $\mathcal{D}_1 \subset \mathcal{C}_\lambda$ and $\mathcal{D}_2 \subset \mathcal{C}_\lambda$, $\lambda = e^{2\pi i \frac{1}{4}}$

3.6 Landing properties at double parabolic parameters

Proposition 3.6.1. *Let $\mathbf{a}_m \in \mathcal{A}_{p/q}$ be the double parabolic parameter of m -type. Then among all the external rays with angles in $\bigcup_k \Theta_k$, there are exactly four landing at \mathbf{a}_m .*

Proof. First we prove the existence of four such rays. For $0 \leq m \leq q$, let $\mathcal{O}_m \subset (\mathbb{C} \setminus \mathcal{A}_{p/q})$ be the set of parameters such that the dynamical external rays with angles Θ_m (recall in Definition 3.1.3) land at 0. Then \mathcal{O}_m is open. Its boundary is contained in $\bigcup_{\theta_i \in \Theta_m} \overline{\mathcal{R}_\infty(\theta_i)}$. From the discussion in 3.1, we have $\mathcal{B}_m \subset \mathcal{O}_m$ and $\mathcal{B}_m \cap \mathcal{O}_k = \emptyset$ for $k \neq m$. Therefore for $0 \leq m \leq q-1$, there are at least four parameter external rays $\mathcal{R}_\infty(t_{1,2})$ with $t_{1,2} \in \Theta_m$ and two rays $\mathcal{R}_\infty(t'_{1,2})$ with $t'_{1,2} \in \Theta_{m+1}$ landing at the double parabolic parameter of m -type. Moreover both $\mathcal{R}_\infty(t_{1,2})$ and $\mathcal{R}_\infty(t'_{1,2})$ separate $\mathcal{B}_m, \mathcal{B}_{m+1}$.

The harder part is the uniqueness. Since the parametrisation Φ_∞ is of degree 3, there are in total $Q := 3q(q+1)$ parameter external rays whose angles belong to $\bigcup_k \Theta_k$. So in order to prove uniqueness, it suffices to find $Q - 4q$ rays not landing at $\mathcal{A}_{p/q}$ among these Q rays. By Lemma 2.3.7, in $Per_1(0)$ there are

$Q - 4q$ rays landing at $\mathcal{C}_0 \setminus \overline{\mathcal{H}_0^0}$. Let A be the set of landing points of these $Q - 4q$ rays. Write $A = \text{Mis} \cup \text{Par}$ where "Mis" (resp. "Par") means that a_0 is Misiurewicz (resp. parabolic).

Claim. *Let $a_1, a_2 \notin \mathcal{H}$ be two different geometrically finite parameters. let $a'_1, a'_2 \in \text{Per}_1(e^{2\pi i \frac{p}{q}})$ be their pinching limit. If $a'_1, a'_2 \notin \mathcal{A}_{p/q}$, then $a'_1 \neq a'_2$.*

Admitting the claim, we finish the proof of the proposition:

- for $a_0 \in \text{Mis}$, its pinching limit $a'_0 \in \text{Per}_1(e^{2\pi i \frac{p}{q}})$ is also Misiurewicz (since $a_0 \notin \partial \mathcal{H}_0^0$ and $v_{0,a}$ is periodic, v_{0,a_0} does not belong to any skeleton). Recall that the pinching deformation preserves external rays, thus the portrait at v_{0,a_0} is the same as that of $v_{a'_0}$. Suppose there are r external rays among the $Q - 4q$ rays landing at a_0 . Then by Lemma 2.2.8, there are also r external rays with angles in $\bigcup_k \Theta_k$ landing at a'_0 .
- If $a_0 \in \text{Par}$, then for the same reason its pinching limit $a'_0 \in \text{Per}_1(e^{2\pi i \frac{p}{q}})$ is also parabolic. We want to prove that there exist two external rays with angles in $\bigcup_k \Theta_k$ landing at a'_0 . Let θ_+, θ_- be the two angles given in Lemma 2.3.8. Since pinching preserves external rays, $R_{a'_0}^\infty(\theta_+), R_{a'_0}^\infty(\theta_-)$ land at the parabolic periodic point, bounding the critical value $v_{a'_0}$. By a simple plumbing surgery (see [5]), for any neighborhood U of a_0 , there exists $a'_1 \in U$ such that $R_{a'_1}^\infty(t_+), R_{a'_1}^\infty(t_-)$ land at the same repelling periodic point and bound $v_-(a'_1)$. By Lemma 2.2.9, there is at least one ray $\mathcal{R}_\infty(\theta)$ with $\theta \in \bigcup_k \Theta_k$ landing at a'_0 . Suppose the contrary that this is the only ray landing at a'_0 . Then the holomorphic motion given by Lemma 2.2.9 implies that $R_a^\infty(\theta_+), R_a^\infty(\theta_-)$ land at the same repelling periodic point and bound $v_-(a)$ when a is close to $\mathcal{R}_\infty(\theta)$. Since $a \in \mathcal{R}_\infty(t)$ is equivalent to $v_-(a) \in R_a^\infty(t)$ (definition of parameter external rays), therefore $\theta \in (t_-, t_+)$. This contradicts Corollary 2.3.9.

To conclude, notice that by Lemma 2.3.8, the $Q - 4q$ rays in $\text{Per}_1(0)$ are decomposed into

$$\left(\bigcup_{a_0 \in \text{Mis}} \bigcup_{i=0}^r \mathcal{R}_\infty^0(s_i) \right) \cup \left(\bigcup_{a_0 \in \text{Par}} (\mathcal{R}_\infty^0(t) \cup \mathcal{R}_\infty^0(t')) \right).$$

While by the above discussion and the claim, we obtain $Q - 4q$ external rays with angles in $\bigcup_m \Theta_m$. This finishes the proof. \square

Proof of the claim. Suppose the contrary that we have $a'_1 = a'_2$. Notice that in the dynamical plans of a_1, a_2 , their critical values v_{0,a_1}, v_{0,a_2} are bounded respectively by wakes W_1, W_2 attached at the boundary of the immediate basin of 0. Since the pinching deformation preserves external rays, we conclude that the angles of the two rays defining W_1 are the same to those defining W_2 . By Lemma 2.3.6, this implies that a_1, a_2 belongs to the same wake (in the parameter plan) attached at $\partial \mathcal{H}_0^0$, which in particular is contained in a quadrant. Thus the angles of external rays landing at them are distinct, since the parametrisation $\Phi_\infty^0 : \mathbb{C} \setminus \mathcal{C}_0 \rightarrow \mathbb{C} \setminus \overline{\mathbb{D}}$ is injective on each quadrant. Thus the pinching limits a'_1, a'_2 are distinct, a contradiction. \square

Definition 3.6.2. *Let $\mathbf{a}_m \in \mathcal{A}_{p/q}$ be of type m . The four rays landing at \mathbf{a}_m is separated by \mathcal{I} into two groups α_m^+, β_m^+ and α_m^-, β_m^- with $\alpha_m^\pm \in \Theta_m, \beta_m^\pm \in \Theta_{m+1}$. They bound 2 open regions separated by \mathcal{I} . These two regions are called **double parabolic wakes** attached to \mathbf{a}_m , denoted by $\mathcal{W}^\pm(\mathbf{a}_m)$ respectively.*




Figure 7: The external rays landing at double parabolic parameters for \mathcal{C}_λ , $\lambda = e^{2\pi i \frac{1}{3}}$.

Denote by \mathcal{S}_m the connected component of \mathcal{V} containing \mathcal{B}_m , where

$$\mathcal{V} := \mathbb{C} \setminus \bigcup_{m=0}^{q-1} \overline{(\mathcal{R}_\infty(\alpha_m^+) \cup \mathcal{R}_\infty(\alpha_m^-) \cup \mathcal{R}_\infty(\beta_m^+) \cup \mathcal{R}_\infty(\beta_m^-))} \quad (15)$$

Definition 3.6.3. For $1 \leq m \leq \lfloor \frac{q}{2} \rfloor$, denote by \mathcal{S}_m^+ the connected component of $\mathcal{S}_m \setminus \mathcal{I}_m$ intersecting \mathcal{D}_m , \mathcal{S}_m^- the other component. For $m = 0$, set $\mathcal{S}_0^\pm = \mathcal{S}_0 \setminus \mathcal{I}_0$.

Now as a corollary of Proposition 3.6.1, we can give a description of the portrait at the parabolic fixed point 0 for the family f_a :

Corollary 3.6.4. Let $a \in \mathbb{C} \setminus \mathcal{A}_{p/q}$. If $a \in \mathcal{W}^\pm(\mathbf{a}_m)$, the portrait of f_a at the parabolic fixed point 0 is Θ_m, Θ_{m+1} ; if $a \in \mathcal{S}_m$, the portrait is Θ_m .

Proof. First we prove that when we go through \mathcal{I} in the direction $\mathcal{B}_0, \dots, \mathcal{B}_q$, α_m^\pm is on the left-hand side of β_m^\pm . Suppose the contrary. Then By Proposition 3.6.1, there exists $a \in \mathcal{W}^\pm(\mathbf{a}_m) \cap \mathcal{C}_\lambda$ such that a and \mathcal{B}_m (resp. \mathcal{B}_{m+1}) are contained in the same connected component of

$$\mathbb{C} \setminus \bigcup_{\theta \in \hat{\Theta}_{m+1}} \overline{\mathcal{R}_\infty(\theta)}, \text{ resp. } \mathbb{C} \setminus \bigcup_{\theta \in \hat{\Theta}_m} \overline{\mathcal{R}_\infty(\theta)},$$

where $\hat{\Theta}_k = (\bigcup_i \Theta_i) \setminus \Theta_k$. Since the portrait at $z = 0$ for a in \mathcal{B}_k is Θ_k , Lemma 2.2.9 implies that the portrait at $z = 0$ of f_a can not contain Θ_m, Θ_{m+1} . But it can neither contain other Θ_k with rotation number p/q for the same reason. But $z = 0$ should at least admits a cycle of landing rays with rotation number p/q , a contradiction.

Let $\mathcal{O}_m^\pm \subset \mathcal{W}^\pm(\mathbf{a}_m)$ be the collection of parameters whose portrait at $z = 0$ is **exactly** Θ_m, Θ_{m+1} . Then clearly \mathcal{O}_m^\pm is open. By the above analysis, $\mathcal{O}_m^\pm \neq \emptyset$ and

$$\overline{\mathcal{R}_\infty(\alpha_m^\pm) \cup \mathcal{R}_\infty(\beta_m^\pm)} \subset \partial \mathcal{O}_m^\pm.$$

Suppose the inclusion above is strict. Then for any $a_0 \in \partial\mathcal{O}_m^\pm \setminus \overline{\mathcal{R}_\infty(\alpha_m^\pm) \cup \mathcal{R}_\infty(\beta_m^\pm)}$, there exists $t_{a_0} \in \Theta_m \cup \Theta_{m+1}$ such that $R_{a_0}^\infty(t_{a_0})$ crashes on $c_-(a_0)$. So $a_0 \in \mathcal{R}_\infty(3t_{a_0})$. By stretching external rays we can show that for all $a \in \mathcal{R}_\infty(3t_{a_0})$, $R_a^\infty(t_{a_0})$ crashes on $c_-(a)$ and also $a \in \partial\mathcal{O}_m^\pm$. Therefore $\mathcal{R}_\infty(3t_{a_0})$ must land at \mathbf{a}_m . By Proposition 3.6.1, $3t_{a_0}$ is one of $\alpha_m^\pm, \beta_m^\pm$, a contradiction. Hence $\mathcal{O}_m^\pm = \mathcal{W}^\pm(\mathbf{a}_m)$.

Let $\tilde{\mathcal{O}}_m \subset \mathbb{C}$ be the collection of parameters whose portrait at $z = 0$ is **exactly** Θ_m . By a similar argument as above we can show that $\tilde{\mathcal{O}}_m = \mathcal{S}_m$. □

The following finishes the proof of Proposition I.

Corollary 3.6.5. *For any m , there are no other external rays with rational angles landing at $\mathbf{a}_m \in \mathcal{A}_{p/q}$.*

Proof. By Lemma 2.2.7, it suffices to treat the case $3^n t = \alpha_m^+$ for some $n \geq 1$. Since t is rational, the three external rays with angle t in $Per_1(0)$ land at 3 different parameters a_1, a_2, a_3 . Now we prove that a_1, a_2, a_3 are all Misiurewicz parameters. Suppose the contrary that a_1 is a parabolic parameter, then by hypothesis on t , $R_\infty^0(\alpha_m^+)$ lands at the parabolic point of f_{0,a_1} . Then by stability of rays landing at repelling point, there exists $\theta \in \Theta_m$ such that $\mathcal{R}_\infty^0(\theta)$ lands at a_1 . This contradicts Lemma 2.3.8. Notice that by the claim in Proposition 3.6.1, the corresponding pinching limits (clearly not double parabolic) $a'_1, a'_2, a'_3 \in Per_1(e^{2\pi i \frac{p}{q}})$ a'_1, a'_2, a'_3 are distinct. By Lemma 6.4.1, they admit in total three external rays with angle t landing. □

4 Dynamical graphes and puzzles

This section mainly aims to construct admissible graph for family (1) that infinitely rings the free critical point $c_-(a)$ in order to apply Yoccoz' Theorem (Theorem 4.3.1) (For basic knowledge of admissible graphs, we refer the readers to [16, Appendix]). More precisely, under the following assumption on f_a

Assumption (\diamond). $a \in \mathcal{C}_\lambda \setminus \mathcal{H}^\lambda$ is not Misiurewicz parabolic and is not contained in any double parabolic wake.

we prove the following technical lemma (Proposition 4.3.8):

Key Lemma. *If a satisfies Assumption (\diamond), then there is an admissible graph infinitely ringing $c_-(a)$.*

Let us fix some notations. Let $f_a \in \mathcal{C}_\lambda \setminus \mathcal{H}^\lambda$ and f_a not Misiurewicz parabolic. In the sequel always fix such an f_a and omit the index a . The critical point in the parabolic basin of 0 will be denoted by c_+ , the free critical point by c_- . Let $B_0^*(0)$ be the immediate basin containing c_+ and $B_1^*(0), \dots, B_{q-1}^*(0)$ the other immediate basins in cycle order. Let $\{R^\infty(\theta_i); \theta_i \in \Theta\}$ be a cycle of external rays landing at 0 and $R^\infty(\theta_0^\pm)$ be the two external rays landing at 0 and bounding $B_0^*(0)$ with $\theta_0^\pm \in \Theta$. These q rays together with $\{0\}$ divide \mathbb{C} into q sectors and let S_i be the sector containing $B_i^*(0)$. Let $\Omega \subset B_0^*(0)$ be the maximal petal, $\tilde{\Omega} = f^q(\Omega)$. For any $r > 1$, denote by $E^\infty(r)$ the external equipotential of potential r .

4.1 Local connectivity at $f^{-n}(0)$

Take some $r_0 > 1$. Define the graph of depth n by

$$Y_0 = \left(\bigcup_{i=1}^q (f^i(\partial\tilde{\Omega}) \cup R^\infty(\theta_i)) \right) \cup E^\infty(r_0), \quad Y_n = f^{-n}(Y_0). \quad (16)$$

A puzzle piece Q_n of depth n is a connected component of $\mathbb{C} \setminus Y_n$ intersecting the Julia set. Let Q_n^\pm be the puzzle piece whose boundary intersects $R^\infty(\theta_\pm)$ respectively.

Proposition 4.1.1. *Let $X_{\pm} = \bigcap_{k \geq 0} \overline{Q_{kq}^{\pm}}$. Then either $X_{\pm} = \{0\}$, or it is a continuum and $f^q : X_{\pm} \rightarrow X_{\pm}$ is a degree two ramified covering. Moreover in the second case, there are two cycles of external rays landing at 0 with angle cycles Θ, Θ' and there exists $\zeta^{\pm} \in \Theta'$ such that $\gamma_{\pm} = R^{\infty}(\theta_0^{\pm}) \cup R^{\infty}(\zeta^{\pm}) \cup \{0\}$ separates $X_{\pm} \setminus \{0\}$ from $\overline{B_0^*(0)} \setminus \{0\}$.*

Proof. We only prove for $X = X_+$ and the other is similar. Set $\theta = \theta_0^+$. First notice that since $f^q(\overline{Q_{(k+1)q}}) = \overline{Q_{kq}}$, hence $f^q(X) = X$. Suppose that $X \neq \{0\}$, then there exists an isomorphism $\phi : \mathbb{C} \setminus X \rightarrow \mathbb{C} \setminus \overline{\mathbb{D}}$. Take a small enlargement U of X with U simply connected and open, such that $f^q : U' \rightarrow U$ has degree $d \geq 2$ and $U \setminus X$ does not contain $f^q(c_-)$, where U' is the connected component of $f^{-q}(U)$ containing X . Hence $f^{-q}(X) \cap U' = X$. Let $W = \phi(U)$ and $W' = \phi(U')$: Then the map $F = \phi \circ f \circ \phi^{-1}$ is a ramified covering of degree d . By Schwarz reflection principle, F extends holomorphically to a neighborhood of \mathbb{S}^1 . Let $g = F|_{\mathbb{S}^1}$. Adapting the same strategy in the proof of [17, Prop. 2.4], we can show that $g : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ is a degree d covering and has $d - 1$ fixed points. Since in the dynamical plane of f , there is an access $\delta \in \mathbb{C} \setminus X$ to 0 fixed by f^q (for example one may take δ to be one of the external rays landing at 0), then $\phi(\delta)$ lands at \mathbb{S}^1 and gives a fixed point of g (see [9]). Thus $d = 2$.

Now we prove that if $X \neq \{0\}$, there is another cycle of external rays landing at 0 and satisfying the conclusion stated in the proposition. Let $R^{\infty}(\theta_k)$ be the external ray landing at $\partial B_0^*(0) \setminus \{0\}$ involved in ∂Q_{kq} . Then $3^q \theta'_{k+1} = \theta'_k$ since $f^q(R^{\infty}(\theta_{k+1})) = R^{\infty}(\theta_k)$. Clearly θ_k has a limit θ' when $k \rightarrow \infty$ since θ_k is monotone. Thus $3^q \theta = \theta$ and $R^{\infty}(\theta')$ enters every puzzle piece Q_{kq} . This implies that $R^{\infty}(\theta')$ lands at a fixed point $x \in X$ of f^q . Notice that $R^{\infty}(\theta')$ is also fixed by f^q , thus x corresponds to the unique fixed point of g , i.e. $x = 0$. Since $X \neq \{0\}$ and is contained in every Q_{kq} , there exists η between θ, θ_k for all k such that $R^{\infty}(\eta)$ landing at $x' \in X$ with $x' \neq 0$. Hence $\theta' \neq \theta$. Clearly γ separates $X_{\pm} \setminus \{0\}$ from $\overline{B_0^*(0)} \setminus \{0\}$. \square

We get immediately

Corollary 4.1.2. *$\partial B_0^*(0)$ is locally connected at 0 and at its inverse orbit.*

4.2 Wakes attached to $\partial B_i^*(0)$

Let $T_0 = \left(\bigcup_{i=1}^q (f^i(\tilde{\Omega}) \cup R^{\infty}(\theta_i)) \right) \cup \{0\}$ and T_n the connected component of $f^{-n}(T_0)$ containing 0. Let Δ_{\pm} be the two unbounded connected components of $S_0 \setminus T_q$, such that $R^{\infty}(\theta_{\pm}) \subset \partial \Delta_{\pm}$. Hence any $z \in \partial B_0^*(0)$ which is not in the inverse orbit of 0 has a unique dyadic representation $(\epsilon_n)_{n \geq 0}$ encoding its orbit position under f^q . More precisely, $\epsilon_n = 1$ if $f^{nq}(z) \in \Delta_+$, $\epsilon_n = 0$ if $f^{nq}(z) \in \Delta_-$. For z in the inverse orbit of 0, we take the convention that ϵ_n is zero for all but finitely many n . Then the following mapping is well defined:

$$\kappa : \partial B_0^*(0) \rightarrow \mathbb{S}^1, z \mapsto (\epsilon_n)_{n \geq 0}$$

For $i \geq 1$, κ is naturally extended to $\partial B_i^*(0)$ by $\kappa(z) = \kappa(w)$ where $w \in \partial B_0^*(0)$ is the first iterated image of z . Notice that there are exactly q external rays involved in T_n at any $u \in \bigcup_i \partial B_i^*(0)$ which is in the inverse orbit of 0. Suppose $u \neq 0$, then among these q rays there are two of them bounding a open region separating all the other $q - 2$ rays with $\bigcup_i \partial B_i^*(0)$. Let $U(u)$ be the closure of this region. Now κ is extended to $\mathbb{C} \setminus \bigcup_i B_0^*(0)$ in the following way: if z belongs to some $U(u)$, then $\kappa(z) := \kappa(u)$ (which is dyadic); if $z \in T_0$, then $\kappa(z) := 0$; else, for every $n \geq 0$, let U_n be the component of $\mathbb{C} \setminus T_n$ containing z , then set $z \in U(z) := \bigcap_n \overline{U_n}$ and define $\kappa(z) = \kappa(u)$ for any $u \in (\bigcup_i \partial B_i^*(0)) \cap U(z)$. Notice that in the second case $\kappa(z)$ does not depend on the choice of u . $U(z)$ is called a **wake** attached to $\partial B_i^*(0)$. Define the corresponding **limb** by $L(z) = U(z) \cap K_f$. Denote by $U(t)$ resp. $L(t)$ the union of $U(z)$ resp. $L(z)$ with $\kappa(z) = t$.

Remark 4.2.1. *By construction, $\mathbb{C} = \bigcup_i B_i^*(0) \cup \bigcup_t U(t)$, $K_f = \bigcup_i B_i^*(0) \cup \bigcup_t L(t)$.*

Lemma 4.2.2. *$L(0) = 0$ if and only if f does not belong to wakes attached at double parabolic parameters.*

Proof. If $L(0) = 0$ but f belongs to some wake attached at a double parabolic parameter. Then by Corollary 3.6.4, there are two cycles of external rays landing at 0. Suppose the corresponding cycle of angles are Θ, Θ' respectively, then there exists $\theta \in \Theta, \theta' \in \Theta'$ such that $R^\infty(\theta), R^\infty(\theta')$ bound an open sector S not intersecting any immediate basin at 0, hence $S \cap K_f \subset L(0)$, contradicting $L(0) = 0$.

If f does not belong to wakes at double parabolic parameters, then one may deduce by the same argument in the second part of the proof of Proposition 4.1.1 that $L(0) = 0$. \square

4.3 Finding infinitely ringed puzzle pieces around c_-

Theorem 4.3.1 (Yoccoz). *Let $f : U' \rightarrow U$ be a quadratic rational-like map and x_0 be its unique critical point. Let $x \in K_f$. For any admissible graph Γ that rings x_0 and rings infinitely x , we have the following alternative:*

- if the tableau of x_0 is k -periodic, then $f^k : P_{l+k}(x_0) \rightarrow P_l(x_0)$ is quadratic-like for l large enough. $\text{Imp}(x)$ is either x or a conformal copy of $\text{Imp}(x_0)$, depending on whether the forward orbit of x intersects $\text{Imp}(x_0)$.
- if the tableau of x_0 is not periodic, then $\text{Imp}(x) = x$.

Observation. If Γ rings infinitely x_0 , then $\Gamma \cap K_f$ contains no parabolic cycle. Indeed, since a parabolic cycle (say period k) must attract a critical point, we may suppose that $f^{kn}(x_0)$ are close to each other when n large enough. Fix such n , pick n_i large enough such that $f^{n_i k}(x_0)$ is ringed at depth n_i , then $P_{n_i}(f^{n_i k}(x_0))$ is compactly contained in $f^{kN}(P_{n_i}(f^{n_i k}(x_0))) = P_{n_i - kN}(f^{(n_i + N)k}(x_0))$ for N large enough. In particular $\Gamma \cap K_f$ does not contain the parabolic cycle.

Now we start to construct admissible graphs for $f = f_a$ satisfying Assumption (\diamond) until the end of this subsection. Firstly we work in the quadratic model $P_\lambda(z) = e^{2\pi i \frac{p}{q}} z + z^2$ with $\lambda = e^{2\pi i \frac{p}{q}}$. Recall the notations at the beginning of Subsection 3.3.

Fix $0 \leq m \leq q - 1$. Suppose $m = lp \pmod{q}$ with $1 \leq l \leq q$. Let H be the connected component of $P_\lambda^{-2q+l}(\Omega_0^{-q})$ contained in $\overline{B_m^*(0)}$. Notice that \mathbb{C} is separated by the q external rays landing at 0 into q sectors S_0, \dots, S_{q-1} (written in cyclic order) with $B_i^*(0) \subset S_i$. Since $P^q : B_m^*(0) \rightarrow B_m^*(0)$ is of degree 2, the dynamics of $P^q|_{B_m^*(0)}$ can be identified with $P_1|_{B(0)}$ by their Fatou coordinates. Thus the construction of "jigsawed" internal rays for P_1 ([17], [23]) can be transferred to P_λ . For $k \geq 1$, let $R(\theta_\pm) \subset B_m^*(0)$ be the internal ray with angle $\theta_\pm = \frac{\pm 1}{2^{k-1}}$. For $0 \leq i \leq k - 1$, define $R(2^i \theta_\pm) = f^{qi}(R(\theta_\pm)) \setminus P^q(H)$. The rays $R(\theta)$ with θ strictly pre-periodic to some $2^i \theta_\pm$ are then defined to be the iterated preimages of $R(2^i \theta_\pm)$ in $B_m^*(0)$. Set $\Theta_\pm = \{\theta; \exists i \text{ s.t. } 2\theta = 2^i \theta_\pm\}$. Define the equipotential of depth $n \geq 0$ by

$$E_0^m = \bigcup_{i=1}^q P_\lambda^i(\partial H), \quad E_n^m = P_\lambda^{-n}(E_0^m). \quad (17)$$

Define the union of internal rays of depth $n \geq 0$ by

$$R_0^m = \bigcup_{i=1}^q \left(\bigcup_{\theta \in \Theta_\pm} P_\lambda^i(R(\theta) \setminus H) \right), \quad R_n^m = P_\lambda^{-n}(R_0^m). \quad (18)$$

Now return to f . We abuse the notations of $S_i, B_i^*(0)$, etc.. Suppose $c_- \in S_m$. Since the dynamics on the immediate basins of f is equivalent to that of P_λ , E_0 in (17) and R_0 in (18) can be transferred to the immediate basins of f . Define similarly E_n, R_n for f to be the n -th preimage. Let $\theta \in \Theta_\pm$, then the landing point x_θ of $R(\theta)$ is pre-periodic, hence there is at least an external ray $R^\infty(t)$ landing at it. For each x_θ we choose a $R^\infty(t)$ and denote by T the collection of these t such that $3^q T \subset T$.

Pick $r \leq 1$. Define the graph of depth $n \geq 0$ by

$$X_n^m = E_n^m \cup R_n^m \cup \left(\bigcup_{t \in T} R^\infty(3^i t) \right) \cup E^\infty(r). \quad (19)$$

Notice that if $m = 0$, the graph above can also be written as

$$X_0^0 = \left(\bigcup_{i=1}^q f^i(\partial\Omega) \right) \cup \bigcup_{i=1}^q \left(\bigcup_{\theta \in \Theta_\pm} f^i(R(\theta) \setminus \Omega) \cup \bigcup_{t \in T} R^\infty(3^i t) \right) \cup E^\infty(r) \quad (20)$$

$$X_n^0 = f^{-n}(X_0^\pm).$$

Recall that $\Omega \subset B_0^*(0)$ is the maximal petal for the return map $f^q|_{B_0^*(0)}$.




Figure 8: An admissible graph for $f \in \text{Per}_1(e^{\frac{2\pi i}{3}})$ with $m = 1$ and θ . The graph in red is X_0^1 , the graph in blue is the first pre-image of X_0^1 that is not contained in X_0^1 . For this graph, c_- is ringed at depth 0.

Lemma 4.3.2. *If $z \in (K_f \setminus \bigcup_i B_i^*(0)) \cap S_m$ satisfies $\kappa(z) \in [\frac{1}{4}, \frac{1}{2}] \cup (\frac{1}{2}, \frac{3}{4}]$, then z is ringed at depth 0 by one of the graphs (19) for all k large enough.*

Proof. By construction, the annulus $P_0 \setminus \overline{P_1}$ of depth 0 is non-degenerated if P_1 is a piece whose boundary contains two of the internal rays $R^0(2^j \theta_\pm + \frac{1}{2})$, $j = 0, \dots, k-2$. The lemma follows by noticing that

$$\theta_+ + \frac{1}{2} < 2\theta_+ + \frac{1}{2} < \dots < 2^{k-2}\theta_+ + \frac{1}{2}$$

$$\theta_- + \frac{1}{2} > 2\theta_- + \frac{1}{2} > \dots > 2^{k-2}\theta_- + \frac{1}{2}$$

and $2^{k-2}\theta_+ + \frac{1}{2} > \frac{3}{4}$ tends to $\frac{3}{4}$ as $k \rightarrow \infty$; $2^{k-2}\theta_- + \frac{1}{2} < \frac{1}{4}$ tends to $\frac{1}{4}$ as $k \rightarrow \infty$. \square

Lemma 4.3.3. *Suppose that $t_0 \neq \frac{1}{2}$. Then the critical value $v_- = f(c_-) \in S_{m+p}$ and $f^q(c) \in L(2t_0) \cap S_m$. Moreover if $z \in L(t_0) \cap S_m$ such that $f(z)$ is not in the puzzle piece (defined by (19)) of depth 0 containing v_- , then z is ringed at depth 0.*

Proof. Suppose $v_- \in S_{m'}$ and let $L'(t_0) \cap S_{m'}$ be the limb containing v_- . Notice that $f^{-1}(L'(t_0))$ has only two connected components, one containing c_- . If $m = 0$, then $m' = p$, otherwise $f^{-1}(L'(t_0))$ will have three components: two in S_0 and one in $S_{m'-p}$. If $m \neq 0$, then $m' = m + p \pmod{q}$, otherwise $f^{-1}(L'(t_0))$ will have three components: one in S_m , one in S_0 and one in $S_{m'-p}$. It is then clear that $f^q(c) \in L(2t_0) \cap S_m$.

Now suppose $z \in L(t_0) \cap S_m$ such that $f(z)$ is not in the puzzle piece of depth 0 containing v_- . Notice that in $L(t_0) \cap S_m$ there is a preimage of $\bigcup_i B_i^*(0)$. Hence z is bounded by a puzzle piece P_1 of depth 1 whose boundary does not intersect puzzle pieces of depth 0, since the internal/external rays in ∂P_1 are all stemming from $f^{-1}(\bigcup_i B_i^*(0)) \cap L(t_0)$, which are different from the rays involved in depth 0. \square

In the next lemma we do **not** suppose $L(0) = 0$.

Lemma 4.3.4. *Suppose $z \in K_f$ is not in the inverse orbit of 0. If $f^n(z)$ eventually hits $\partial B_0^*(0)$. Then z is infinitely ringed by one of the graphs (20) for all k large enough.*

Proof. The proof goes the same as in [17, Lem. 6.1]. The orbit of z must satisfy one of the following conditions, and the result follows by Lemma 4.3.2:

- If there exists a subsequence n_j such that $2^{n_j} \kappa(z) \rightarrow \frac{1}{4}$ resp. $\frac{3}{4}$, then z is infinitely ringed by (19) defined by θ_- resp. θ_+ .
- Otherwise z is ringed by (19) defined by θ_+ and θ_- .

\square

Lemma 4.3.5. *Suppose $z \in K_f$ is not in the inverse orbit of 0 and not in any basin of 0. If z satisfies $f^n(z) \notin L(t_0)$ for n large enough, then z is infinitely ringed by one of the graphs (20) for all k large enough.*

Proof. The proof goes the same as in [17, Lem. 6.1] if $f^n(z)$ avoid $L(\frac{1}{2})$ for n large enough. If $f^n(z)$ meets infinitely many times $L(\frac{1}{2})$, then we treat by two subcases:

- If $f^n(z) \in L(\frac{1}{2})$ for n large enough. Let U be a connected component of $f^{-j}(L(\frac{1}{2}))$ ($1 \leq j \leq q$) contained in $L(\frac{1}{2}) \cap S_0$ that intersects infinitely the orbit of z . Notice that $L(\frac{1}{2}) \cap S_0$ is divided into $q-1$ connected components by $q-2$ external rays landing at x_0 , the first preimage of 0 (other than 0) on $\partial B_0^*(0)$. Suppose U belongs to one of these $q-1$ components V , which is bounded by $R^\infty(\eta_1), R^\infty(\eta_2)$. Then $U \subset\subset V$ and the angles of external rays involved in ∂U are between η_1, η_2 . Recall graph (16) and its corresponding puzzle pieces Q_n . Let the Q_n^\pm be the two puzzle pieces of depth n at x_0 . Since $L(0) = 0$, all the angles of external rays involved in Q_n^\pm converge to η_1 or η_2 . Thus if we take k large enough in $\theta_\pm = \frac{\pm 1}{2^k - 1}$, U will be included by a puzzle piece P_1^\pm of depth 1 defined by graph (20) such that P_1^\pm is compactly contained in some puzzle piece of depth 0.
- There is a subsequence such that $f^{n_j}(z) \notin L(\frac{1}{2})$. By hypothesis, z returns infinitely many times to $L(\frac{1}{2}) \cap S_0$. Thus $\{f^n(z); n \geq 0\} \cap (L(\frac{1}{2}) \cap S_0)^c$ will meet infinitely $(L(\frac{1}{4}) \cup L(\frac{3}{4})) \cap S_0$. By Lemma 4.3.2, if $f^n(z)$ is contained in $(L(\frac{1}{4}) \cup L(\frac{3}{4})) \cap S_0$, then it is ringed at depth 0.

Thus in both cases, we have found a graph that rings infinitely z . \square

Proposition 4.3.6. *Suppose $t_0 \neq \frac{1}{2}$ and that $z \in K_f$ returns infinitely many times to $L(c_-) = L(t_0) \cap S_m$ but never hits $\partial B_m^*(0)$. Then for all k large enough, there exists one of the two graphs (19) that rings infinitely z .*

Proof. For simplicity we only do the proof for the case $m = 0$. The case $m \neq 0$ will be similar. We adapt a similar strategy in the proof of [17, Lem. 8.2]. By hypothesis there is a subsequence $\{f^{n_i}(z)\}_i \subset L(t_0)$. Take any $w \in \{f^{n_i}(z)\}_i$.

- $t_0 \in [\frac{1}{4}, \frac{1}{2})$. Then by Lemma 4.3.2, w is ringed at depth 0 by graph (20) defined by θ_- for k large enough.
- $t_0 \in (\frac{1}{8}, \frac{1}{4})$. By Lemma 4.3.3, the limb containing the critical value v_- is $L(2t_0) \cap S_p$ with $2t_0 \in (\frac{1}{4}, \frac{1}{2})$. Take k large enough and the corresponding graph X_0^+ such that $L(2t_0) \cap S_p$ is bounded by the puzzle piece of depth 0 whose boundary intersects $f(R^0(2^{k-2}\theta_+)), f(R^0(2^{k-1}\theta_+))$. Thus by Lemma 4.3.3, any w satisfying

$$\kappa(f(w)) \in (0, 2^{k-2}\theta_+) \cup (2^{k-1}\theta_+, 1)$$

is ringed by graph (20) defined by θ_+ at depth 0. Hold the same k and take $\theta_- = \frac{-1}{2^{k-1}}$. If w satisfies

$$\kappa(f(w)) \in [2^{k-2}\theta_+, \frac{1}{2} + \theta_-),$$

then by Lemma 4.3.2, $f^j(f(w))$ is ringed at depth 0 by graph (20) defined by θ_- , where $j \leq q-1$ is the smallest integer such that $f^j(f(w)) \in S_0$. Now suppose that w satisfies

$$\kappa(f(w)) \in [\frac{1}{2} + \theta_-, 2^{k-1}\theta_+].$$

Take k' slightly smaller than k such that $\theta'_- = \frac{-1}{2^{k'-1}}$ satisfies $2t_0 < 1 + 2^{k'-1}\theta'_- < \frac{1}{2} + \theta_-$. (adjust k, k' to be larger if necessary). Thus we can apply Lemma 4.3.3 to graph (20) defined by θ'_- and hence w is ringed at depth 0. To conclude, we have proved that for any w , there exists a graph (20) defined by θ_+, θ_- or θ'_- (not depending on w) that rings w at some depth less than q .

- $t_0 \in (\frac{1}{2^{n+1}}, \frac{1}{2^n}]$, $n \geq 3$. The strategy is quite similar as above. In this case we have $2t_0 \in (\frac{1}{2^{n-2}}, \frac{1}{2^{n-1}}]$. For the same reason, w is ringed by graph defined by θ_+ for k large if $\kappa(f(w)) \in (0, 2^{k-n}\theta_+) \cup (2^{k-n+1}\theta_+, 1)$. Take $\epsilon > 0$ small enough such that $2^{n-2}(2^{k-n+1}\theta_+ - \epsilon) > \frac{1}{2} + \theta_-$. Thus for w satisfying

$$\kappa(f(w)) \in [2^{k-n}\theta_+, 2^{k-n+1}\theta_+ - \epsilon)$$

and j the smallest integer such that $f^j(f(w)) \in S_0$, we have $\kappa(f^N(w)) \in [\frac{1}{4}, \frac{1}{2} + \theta_-)$, where $N = (n-2)q + j + 1$. Thus $f^N(w)$ is ringed at depth 0 by graph defined by θ_- . If w satisfies

$$\kappa(f(w)) \in [2^{k-n+1}\theta_+ - \epsilon, 2^{k-n+1}\theta_+),$$

and $j \leq q-1$ is the smallest integer such that $f^j(f(w)) \in S_0$, then

$$\kappa(f^{N'}(w)) \in [2^{k-2}\theta_+ - 2^{n-3}\epsilon, 2^{k-2}\theta_+) \subset [\frac{1}{4}, \frac{1}{2} + \theta_-)$$

(take ϵ smaller if necessary), where $N' = (n-3)q + j + 1$. Hence $f^{N'}(w)$ is ringed at depth 0 by graph defined by θ_- .

- $t_0 \in (\frac{1}{2}, 1)$. The argument is symmetric to the case $t_0 \in (0, \frac{1}{2})$, which is already handled above.

□

Lemma 4.3.7. *Let $t_0 = \frac{1}{2}$ and z satisfy the hypothesis in Proposition 4.3.6. Then z is infinitely ringed by some graph (20).*

Proof. The prove goes exactly the same as Lemma 4.3.5. □

To summarize all the cases above, Lemma 4.3.4, 4.3.5, 4.3.7 and Proposition 4.3.6 gives

Proposition 4.3.8. *Suppose $L(0) = 0$. Let $z \in K_f$ not in the inverse orbit of 0 and not in any basin of 0, then for all k large enough, z is infinitely ringed by one of the graphs (19).*

Applying Yoccoz's Theorem to $v_- = f(c_-)$, $f = f_a$, we get

Theorem 4.3.9. *Let $\mathcal{U} \subset \mathcal{C}_\lambda$ be a parabolic component of adjacent, bitransitive or capture type that is not in any $\mathcal{W}^\pm(\mathbf{a}_m)$. Suppose $a \in \partial\mathcal{U}$, $f_a^n(c_-(a)) \neq 0, \forall n \geq 1$ and $a \notin \mathcal{A}_{p/q}$. Then there exists a graph (19) and a sequence of non-degenerated annuli $A_{n_i}^a, i \geq 0$, such that*

1. $A_{n_i}^a = P_{n_i}^{a,v} \setminus \overline{P_{n_i+1}^{a,v}}$, $i \geq 1$, where $P_n^{a,v}$ is the puzzle piece of depth n containing $v_-(a)$.
2. $f_a^{n_i-n_0} : A_{n_i}^a \rightarrow A_{n_0}^a$ is a non-ramified covering.
3. either $\sum_i \text{mod}(A_{n_i}^a) = \infty$ or there exists $k \geq 1$ such that $f_a^k : P_{m+k}^{a,v} \rightarrow P_m^{a,v}$ is quadratic-like for all m large enough and $\bigcap P_m^{a,v}$ is the filled Julia set of the renormalized map f_a^k .

The following lemma allows us to apply Yoccoz's Theorem to other points of the Julia set.

Lemma 4.3.10. *Let $L(t_n)$ be the limb containing $f^n(c_-)$, $n \geq 0$. If*

- t_n is not dyadic for n large enough,
- or there exists n_1, n_2 such that $t_{n_1} \in [\frac{1}{4}, \frac{1}{2})$, $t_{n_2} \in (\frac{1}{2}, \frac{3}{4}]$,

*then there exists N such that c_- is ringed at depth N for **both** graphs (19) for all k large enough.*

Proof. The proof for the first case is just a repeat of [17, Lem. 6.2]. The second case is deduced directly from Lemma 4.3.2 \square

Theorem 4.3.11. *Let f satisfy Assumption (\diamond) . For all $n \geq 0$, $f^{-n}(B_m^*(0))$ is locally connected.*

Proof. Let $z \in B_m^*(0)$. Corollary 4.1.2 treat the case when z is in the inverse orbit of 0. For other z , suppose first that $c_- \in L(t_n)$ satisfies one of the hypothesis in Lemma 4.3.2, then Lemma 4.3.4 allows us to use Yoccoz's Theorem to get the dichotomy: if the intersection of the puzzle pieces containing z shrink to z , then we are done; if the intersection is a quadratic copy, then this copy is separated from $B_m^*(0)$ by two external rays landing at z . Let W be the open region bounded by the two rays containing the quadratic copy, then $(P_n(z) \setminus W)_n$ form a connected basis of z . For details of this part, see the proof of [17, Thm. 1].

The only case left not covered by Lemma 4.3.10 is that t_n is dyadic for all n , but either $t_n \notin [\frac{1}{4}, \frac{1}{2})$ for all n or $t_n \notin (\frac{1}{2}, \frac{3}{4}]$ for all n . For example if it is the first alternative, we treat it furthermore by two subcases:

- $t_n = \frac{1}{2}$ for all n . By Lemma 4.3.7, we may suppose (19) defined by θ_+ rings c_- for all k large enough. By the proof of Lemma 4.3.4, z might not be infinitely ringed by the graph of θ_+ only when $\kappa(z)$ is recurrent to $\frac{1}{4}$. However, in this case there exists k large enough such that the graph with θ_+ does not contain z for all n . There is N such that the puzzle piece P_0 containing $w \in \partial B_m^*(0)$ with $\kappa(w) = \frac{1}{4}$ contains $f^N(z)$. Then $\overline{P_0}$ does not contain forward orbit of c_- . Hence by the shrinking lemma, the pieces around $f^N(z)$ shrink to $f^N(z)$.
- t_n is not always $\frac{1}{2}$. Without loss of generality, suppose $t_{n_0} = \frac{3}{4}$. Then by Lemma 4.3.2, c_- is ringed at depth 0 by graph of θ_+ . Similarly as the above case we can apply the shrinking lemma for $\kappa(z)$ recurrent to $\frac{1}{4}$.

\square

4.4 Wakes attached to end points

Suppose f satisfy Assumption (\diamond) . Set $B_0 := \bigcup_m B_m^*(0)$. Define B_n to be the union of Fatou components not in B_{n-1} but attached to ∂B_{n-1} at a preimage of 0. Set $B = \overline{\bigcup_n B_n}$. For each component $U \subset B_n$, \overline{U} is eventually, firstly and bijectively sent to some $\overline{B_m^*(0)}$ by a certain iteration of f . So any $z \in \partial U$ is associated to an angle ω_n . Recall that $B_m^*(0) \subset S_m$, so we associate U with $\epsilon_n = m$. Therefore we can using the following address to encode the position of z :

$$[(\epsilon_0, \omega_0); (\epsilon_1, \omega_1); \dots; (\epsilon_n, \omega_n)] \quad (21)$$

where $\omega_0, \dots, \omega_{n-1}$ are dyadic, $\omega_n \in \mathbb{R}/\mathbb{Z}$; $0 \leq \epsilon_k \leq q-1$ is chosen so that $f^k(z) \in S_{\epsilon_k}$. In the same way we can encode the position of components of B_n .

Define wakes $W_U(\omega)$ and limbs $L_U(\omega)$ attached to ∂U similarly as 4.2 Let W be the open region bounded by the two external rays separating U with B_{n-1} . Likewise we have $W = U \cup \bigcup_{\omega \neq 0} W_U(\omega)$, $K_f \cap W = U \cup \bigcup_{\omega \neq 0} L_U(\omega)$. Notice that the wakes and limbs defined here include those in 4.2: $L_{B_m^*(0)}(\omega) = L(\omega) \cap S_m$.

Proposition 4.4.1. *Suppose ω is not dyadic. Let $\{z\} = L_U(\omega) \cap \partial U$. If $L_U(\omega)$ is non trivial, then either z is in the inverse orbit of c_- or z is (pre)-periodic. Moreover, if $L_U(\omega)$ is non trivial, then there are exactly two external rays landing at z separating $L_U(\omega)$ with U , otherwise there is only one.*

Proof. Suppose $L_U(\omega)$ is non trivial and z not in the inverse orbit of c_- . Then $L_U(\omega)$ is sent injectively to the critical limb by some f^N (since the width of $W_U(\omega) > 0$ and it is multiple by 3 by every iteration of f). So we may suppose $L_U(\omega)$ is the critical limb. Then $U = B_m^*(0)$, since for otherwise $f^n(L_{f(U)}(f(z)))$ will never contain $L_U(\omega)$. Hence z is periodic.

Since in the proof of Theorem 4.3.11, we see that the pieces shrink to z if the limb is trivial, so it is direct that there is only one external ray landing at z if furthermore z does not hit c_- . Existence of two external rays landing at z if the tableau of z is (pre)-periodic comes from the shrinking property of $P_n \setminus W$, where W is as in the begining of the proof of Theorem 4.3.11. For the proof of the uniqueness of these two rays, see [17, Lem. 7.2]. \square

More generally, consider $z \in B \setminus \bigcup_n \overline{B_n}$, and suppose $z_k \in \bigcup_n \overline{B_n}$ converging to z . Then for all n , there exists z_k such that $z \in \overline{B_n}$. Otherwise, there exists N such that for $k \geq N$, z_k belongs to the closure of some Fatou component in B_{N+1} attached at some $\partial U \subset B_N$. Clearly z_k converges to some $L_U(t)$ with $z_k \notin L_U(t)$ since $z \in B \setminus \bigcup_n \overline{B_n}$. Then $L_U(t)$ must be non trivial since z is not the root of $L_U(t)$. By Proposition 4.4.1, $L_U(t)$ is separated from all other limbs by two external rays, so z_k can not converge to z . Therefore there exists a unique sequence of Fatou components $U_n \subset B_n$ such that $z \in \bigcap_n L_{U_n}(\omega_n)$ with ω_n dyadic. Thus any $z \in \partial B \setminus \{0\}$ can be represented by a unique infinite sequence (if $z \in \partial \overline{B_n}$, add zeros after (ϵ_n, ω_n)):

$$[z] := [(\epsilon_0, \omega_0); (\epsilon_0, \omega_1); \dots; (\epsilon_n, \omega_n)); \dots].$$

where $\omega_k \in (0, 1)$, $0 \leq \epsilon_k \leq q-1$. The coordinate for $f(z)$ is

$$[f(z)] = \begin{cases} [(\epsilon_0 + p, 2\omega_0); (\epsilon_1, \omega_1); \dots; (\epsilon_n, \omega_n)) \dots], & \text{if } \epsilon_0 = 0, \omega_0 \neq \frac{1}{2} \\ [(\epsilon_1, \omega_1); (\epsilon_2, \omega_2); \dots; (\epsilon_n, \omega_n)) \dots], & \text{if } \epsilon_0 = 0, \omega_0 = \frac{1}{2} \\ [(\epsilon_0 + p, \omega_0); (\epsilon_1, \omega_1); \dots; (\epsilon_n, \omega_n)) \dots], & \text{if } \epsilon_0 \neq 0 \end{cases} \quad (22)$$

Lemma 4.4.2. *$z \in B$ has at least 2 preimages in B .*

Proof. This is clear for $z \in \overline{B_n}$. Take $z \in B \setminus \bigcup_n \overline{B_n}$, then z is the limit of some (z_n) with $z_n \in \overline{B_{N_n}}$. Each z_n has two preimages x_n, y_n , which, up to taking a subsequence, have different limits x, y . Clearly $f(x) = f(y) = z$. \square

Remark 4.4.3. The construction of the sequence $[(\epsilon_0, \omega_0); (\epsilon_0, \omega_1); \dots; (\epsilon_n, \omega_n); \dots]$ is also valid for the quadratic model $P_\lambda(z) = e^{2\pi i \frac{p}{q}} z + z^2$. Moreover the following map is bijective:

$$\Xi : J_\lambda \setminus \{0\} \longrightarrow [\mathbb{Z}/q\mathbb{Z} \times (0, 1)]^\mathbb{N}, \quad z \mapsto [z]. \quad (23)$$

Indeed, the surjectivity is clear; the injectivity is also clear for $[z]$ finite. Notice that J_λ can be regarded as a pinching limit of the Julia set of $\lambda z + z^2$ with $|\lambda| < 1$ and the pinching map ϕ is injective beyond the skeletons, while the skeletons are $\phi^{-1}(y)$ with y preimage of 0 (cf. Theorem B.8). Thus if $[z]$ is infinite, then clearly $\phi^{-1}(z)$ does not intersect any skeleton, hence $\#\phi^{-1}(z) = 1$.

Lemma 4.4.4. Suppose f is non renormalisable. If $v_- = f(c_-) \in B \setminus \bigcup_n \overline{B_n}$, then there is only one external ray landing at v_- .

Proof. The existence of external ray landing at v_- comes from the shrinking property of puzzles pieces around it given by Theorem 4.3.9. Suppose the contrary that there are two rays landing. Let W be the open region separated from B by these two rays. Then there exists N such that W is sent injectively onto $f^N(W)$ with $c_- \in f^N(W)$. But this is impossible, since by Lemma 4.4.2 $f^{-1}(v_-)$ has two preimages in B , so c_- must belong to B because f has degree 3. \square

5 Passing to parameter plan

In this section we pass the dynamical combinatorics to parameter ones. To simplify the redaction, from now on we will mainly work in the "fundamental region" \mathcal{S}_m^+ for $0 \leq m \leq \lfloor \frac{q}{2} \rfloor$ (recall Definition 3.6.3).

5.1 Parameter equipotentials and rays

For $m = 0$, define parameter equipotentials and union of internal rays in \mathcal{D}_0 and $\tilde{\mathcal{D}}_0$ by

$$\mathcal{E}_n^0 = (\Psi_0^{adj})^{-1}(E_n^0 \cap B_p^*(0)), \quad \mathcal{R}_n^0 = (\Psi_0^{adj})^{-1}(R_n^0 \cap B_p^*(0)), \quad n \geq 0. \quad (24)$$

For $1 \leq m \leq \lfloor \frac{q}{2} \rfloor$, define parameter equipotentials and union of internal rays in \mathcal{D}_m resp. $\tilde{\mathcal{D}}_m$ by

$$\mathcal{E}_n^m = (\Psi_m^{bit})^{-1}(E_n^m \cap B_{m+p}^*(0)), \quad \mathcal{R}_n^m = (\Psi_m^{bit})^{-1}(R_n^m \cap B_{m+p}^*(0)), \quad n \geq 0. \quad (25)$$

Let $k \geq 1$, $\mathcal{U}_k \subset \tilde{\mathcal{S}}_m$ be a capture component. Suppose for $a \in \mathcal{U}_k$, $c_-(a)$ firstly hit $B_{a,l}^*(0)$ after k iterations. Define the parameter equipotentials and internal rays by

$$\mathcal{E}_n^{\mathcal{U}_k} = (\Psi_{\mathcal{U}_k})^{-1}(E_n^m \cap B_l^*(0)), \quad \mathcal{R}_n^{\mathcal{U}_k} = (\Psi_{\mathcal{U}_k})^{-1}(R_n^m \cap B_l^*(0)), \quad n \geq 0. \quad (26)$$

Next we investigate landing properties for equipotentials in \mathcal{B}_m . We only state the result for \mathcal{E}_n^m in \mathcal{D}_m since it will be the same for $\tilde{\mathcal{E}}_n^m$.

Proposition 5.1.1. Let $n, k \geq 0$. Let $1 \leq l \leq q$ be such that $m = lp \pmod{q}$. Then $\mathcal{E}_n^m \cap \partial \mathcal{D}_m$ consists of finitely many points. Denote by \mathcal{E}_n^m / \sim the quotient space of $\mathcal{E}_n^m \cap \partial \mathcal{D}_m$ by

- gluing the double parabolic parameter on $\partial \mathcal{B}_0$ and $2\sqrt{\lambda}$ if $m = 0$;
- gluing the two parabolic parameters on $\partial \mathcal{B}_m$ if $1 \leq m \leq \frac{q}{2}$.

Then \mathcal{E}_n^m / \sim is homeomorphic to $E_n^m \cap B_{m+p}^*$. In particular $\mathcal{E}_{kq}^m = \mathcal{E}_{kq+1}^m = \dots = \mathcal{E}_{(k+1)q-1}^m$ and $(\mathcal{E}_{kq}^m \cap \partial \mathcal{D}_m) \subsetneq (\mathcal{E}_{(k+1)q}^m \cap \partial \mathcal{D}_m)$. Moreover for $a \in \mathcal{E}_{kq}^m \cap \partial \mathcal{D}_m$ not double parabolic, $f_a^{(k+1)q-l}(v_-(a)) = 0$.

Proof. The proof has no difference to the case $p/q = 1$. See [23, Prop. 3.2.10]. \square

By the above proposition, we see that \mathcal{D}_m is cut "binarily" by \mathcal{E}_{kq}^m into 2^k pieces if $m = 0$ and 2^{k+1} pieces otherwise. For a double parabolic parameter $b \in \partial\mathcal{B}_m$, let $\{a_k\}_{k \geq 0}$ be the sequence such that $a_k \in \partial\mathcal{B}_m$ be the closest summit to b among $\mathcal{E}_{kq}^m \cap \partial\mathcal{B}_m$. By Lemma 2.2.8, there are q parameter external rays landing at a_k and q dynamical rays with the same angles landing at $v_-(a_k)$. The angles are preimages of 0 under multiplication by 3. Denote by T_k the set of these angles.

Lemma 5.1.2. $3^q T_{k+1} = T_k$. Moreover all angles in T_k converges to α , the angle of the external ray $\mathcal{R}_\infty(\alpha)$ landing at b with $\mathcal{R}_\infty(\alpha) \subset \partial\mathcal{S}_m$.

Proof. We only do the proof for $m = 0$ to illustrate the idea. In this case, $b = \mathbf{a}_0$ and hence $\theta = \beta_0^+$ (Definition 3.6.2). For $n \geq 0$, let \mathcal{U}_k be the component bounded by $\mathcal{R}_\infty(\alpha_0^+) \cup \mathcal{E}_{kq}^0 \cup \mathcal{R}_\infty(t_k)$, where $\mathcal{R}_\infty(t_k)$ is the external ray closest to $\mathcal{R}_\infty(\beta_0^+)$ landing at $a_k \in \mathcal{E}_{kq}^0 \cap \partial\mathcal{B}_0$, the point closest to \mathbf{a}_0 (in the binary sense). By Corollary 3.6.5, external rays $\mathcal{R}_\infty(t)$ with $3^{kq}t \in \Theta_0$ do not land at \mathbf{a}_0 . Hence

$$\mathcal{U}_k \setminus \bigcup_{\{t; 3^{kq}t \in \Theta_0\}} \mathcal{R}_\infty(t)$$

contains $\mathcal{W}^+(\mathbf{a}_0)$, and denote by $\tilde{\mathcal{U}}_k$ the component containing it. Notice that for $a \in \tilde{\mathcal{U}}_{k-1}$, there is a dynamical holomorphic motion of $\bigcup_{t \in T_k} R_a^\infty(t) \cup \overline{\Omega_{a,p}^{kq-1}}$. Therefore T_k is exactly the set of angle of external rays landing at $f_a^{-kq}(0) \cap \partial B_{a,p}^*(0)$ which is closest to β_0^+ . By the proof of Proposition 4.1.1 we see that T_k satisfies $3^q T_{k+1} = T_k$ and tends to β_0^+ as $k \rightarrow +\infty$. □

Proposition 5.1.3. Let \mathcal{U}_k be a capture component. Then $\partial\mathcal{U}_k$ contains no double parabolic parameter.

Proof. Suppose the contrary that there exists $b \in \mathcal{A}_{p/q} \cap \partial\mathcal{U}_k$. Consider respectively two cases:

- \mathcal{U}_k is contained in some \mathcal{S}_m . Since $\partial\mathcal{U}_k \subset \partial\mathcal{C}_\lambda$ (Lemma 2.2.4), there is an external ray contained in \mathcal{S}_m accumulating to $\partial\mathcal{U}_k$. This contradicts Lemma 5.1.2.
- \mathcal{U}_k is contained in some double parabolic wake. Consider equation of z , $f_a^q(z) = z$. When $a = b$, 0 is a multiple solution of order $2q + 1$, while for $a \neq b$ near b , 0 is has order $q + 1$. All the other solution besides 0 of $f_b^q(z) = z$ gives a repelling periodic cycle for b , which admits a holomorphic motion for a near b . Thus there are only q non-zero solutions left of $f_a^q(z) = z$ for $a \neq b$ (a near b) which do not come from the holomorphic motion. Among these q solutions, there is at least a repelling cycle C of period dividing q for f_a . Now suppose furthermore $a \in \mathcal{U}_k$. Let Θ be the set of angle of external rays for f_a landing at C . Notice that Θ does not depend on a since in \mathcal{U}_k there is a holomorphic motion of $B_a(\infty)$ induced by Böttcher coordinate. Θ can not be Θ_m, Θ_{m+1} since C is not 0. Hence for f_b , the external rays with angles in Θ land at a repelling cycle, which in turn gives a repelling cycle for f_a by holomorphic motion. This contradicts how we choose C . □

We get immediately

Corollary 5.1.4. $\overline{\mathcal{E}_n^{\mathcal{U}_k}}$ is homeomorphic to $\overline{\Psi_{\mathcal{U}_k}(\mathcal{E}_n^{\mathcal{U}_k})}$. Moreover $a \in \overline{\mathcal{E}_n^{\mathcal{U}_k}} \cap \partial\mathcal{U}_k$ is Misiurewicz parabolic.

Corollary 5.1.5. Let \mathcal{U}_k be a capture component contained in some double parabolic wake $\mathcal{W}^\pm(\mathbf{a}_m)$, then for $a_0 \in \partial\mathcal{U}_k$, $f_{a_0}^k(c_-(a_0))$ is on the boundary of some $B_{a_0,m}^*$. Moreover $\partial\mathcal{U}_k$ is a Jordan curve.

Proof. By Proposition 5.1.3, there is an open neighborhood of \mathcal{U}_k , on which the union of immediate basins $\bigcup_i B_{a,i}^*(0)$ admits an holomorphic motion h_a induced by Fatou coordinate. One can choose the base point a'_0 of h_a in \mathcal{U}_k . By λ -lemma, the motion is extended to $\overline{\bigcup_i B_{a,i}^*(0)}$. For $a \in \mathcal{U}_k$, by definition of capture component there exists m such that $f_a^k(c_-(a)) \in B_{a,m}^*$. Let a tend to a_0 , we get $f_{a_0}^k(c_-(a_0)) \in \partial B_{a,m}^*$ since $\overline{B_{a,m}^*}$ moves holomorphically.

Define $H : \partial\mathcal{U}_k \rightarrow \partial B_{a'_0,m}^*$ by $a \mapsto h_a^{-1}(f_a^k(c_-(a)))$. H is locally regular, hence $\partial\mathcal{U}_k$ is locally connected since $\partial B_{a'_0,m}^*$ is (Julia set of $f_{a'_0}$ is locally connected since it is geometrically finite [21]). $\partial\mathcal{U}_k$ is then a Jordan curve by Lemma 2.2.4. \square

From Proposition 5.1.1 and 5.1.3, we see that $\mathcal{R}_n^m, \mathcal{R}_n^{\mathcal{U}_k}$ can not land at double parabolic parameters. We have the following landing property of internal rays (the proof is similar to that for external rays) :

Lemma 5.1.6. *Let $0 \leq m \leq \lfloor \frac{q}{2} \rfloor$ and \mathcal{U}_k be a capture component not in double parabolic wake. Then any component of \mathcal{R}_n^m resp. $\mathcal{R}_n^{\mathcal{U}_k}$ land at some $a_0 \in \partial\mathcal{B}_m$ resp. $\partial\mathcal{U}_k$ which is neither Misiurewicz parabolic nor double parabolic. The landing points of $f_{a_0}^{-k}(R_{a_0}^m)$ are (pre-)periodic to the same cycle, where $R_{a_0}^m$ is just R_0^m in (18). Moreover if this cycle is repelling, then some component of $f_{a_0}^{-k}(R_{a_0}^m)$ will land at $v_-(a_0)$.*

We also have the analogue to Lemma 6.4.1:

Lemma 5.1.7. *Suppose $a_0 \in \mathcal{C}_\lambda \setminus \bigcup_m \mathcal{W}^\pm(\mathbf{a}_m)$ is a Misiurewicz parameter. Then a_0 is the landing point of some \mathcal{R}_n^m or $\mathcal{R}_n^{\mathcal{U}_k}$, if and only if $v_-(a_0)$ is the landing point of some $f_{a_0}^{-k}(R_{a_0}^m)$.*

5.2 Dynamical objects move holomorphically

Definition 5.2.1. *Let $A_0 \subset \mathbb{S}^1$ be finite and satisfy $3A_0 = A_0$, A_n be the n -th preimage of A_0 under multiplication by 3. A parameter object of depth $n \geq 0$ is the intersection of one of the following four sets and $\tilde{\mathcal{S}}_m$.*

$$\bigcup_{\eta \in A_n} \overline{\mathcal{R}_\infty(\eta)}, \bigcup_{k=1}^n \left(\bigcup_{\mathcal{U}_k \subset \mathcal{H}_k} \overline{\mathcal{E}_{n-k}^{\mathcal{U}_k}} \right) \cup \overline{\mathcal{E}_n^m}, \bigcup_{k=1}^n \left(\bigcup_{\mathcal{U}_k \subset \mathcal{H}_k} \overline{\mathcal{R}_{n-k}^{\mathcal{U}_k}} \right) \cup \overline{\mathcal{R}_n^m}, \mathcal{E}_\infty(r^{1/3^n}). \quad (27)$$

Let \mathcal{O}_n be a parameter object of depth n . Let $\tilde{\mathcal{O}}_n$ be a connected component of $\mathcal{S}_m^+ \setminus \mathcal{O}_n$ intersecting $\partial\mathcal{C}_\lambda$. Let $a \in \tilde{\mathcal{O}}_n$.

When \mathcal{O}_n is $\bigcup_{\eta \in A_n} \overline{\mathcal{R}_\infty(\eta)}$ or $\mathcal{E}_\infty(r^{1/3^n})$, $\bigcup_{\eta \in A_{n+1}} \overline{R_a^\infty(\eta)}$ resp. $E_a^\infty(r^{1/3^{n+1}})$ does not contain $c_-(a)$. Hence each external ray $R_a^\infty(\eta)$ is well defined and lands at J_a , resp. $E_a^\infty(r^{1/3^{n+1}})$ is homeomorphic to \mathbb{S}^1 . Moreover there is a dynamical holomorphic motion of $\bigcup_{\eta \in A_{n+1}} \overline{R_a^\infty(\eta)}$ resp. $E_a^\infty(r^{1/3^{n+1}})$ induced by Böttcher coordinate ϕ_a^∞ .

Now let $\mathcal{O}_n = \bigcup_{k=1}^n \left(\bigcup_{\mathcal{U}_k \subset \mathcal{H}_k} \overline{\mathcal{E}_{n-k}^{\mathcal{U}_k}} \right) \cup \overline{\mathcal{E}_n^m}$. We want to find a dynamical holomorphic motion for equipotentials in $B_a(0)$. Recall that Ω_a is the maximal petal contained in $B_{a,0}^*(0)$ and $\tilde{\Omega}_a$. Let H_a be the connected component of $P^{-2q+l}(\tilde{\Omega}_a)$ contained in $\overline{B_{a,m}^*(0)}$. Then by definition of $\mathcal{E}_n^m, \mathcal{E}_{n-k}^{\mathcal{U}_k}$, $f_a^{-n-1}(E_0^a)$ does not contain $c_-(a)$, where $E_0^a := \bigcup_{i=1}^q f_a^i(\partial H_a)$. Hence there is a dynamical holomorphic motion of $E_{n+1}^a := f_a^{-n-1}(E_0^a)$ induced by Fatou coordinate and the pull back of f_a .

Now let $\mathcal{O}_n = \bigcup_{k=1}^n \left(\bigcup_{\mathcal{U}_k \subset \mathcal{H}_k} \overline{\mathcal{R}_{n-k}^{\mathcal{U}_k}} \right) \cup \overline{\mathcal{R}_n^m}$. Notice that here the construction of internal rays is more subtle, since $c_-(a)$ might be in $B_{a,m}^*(0)$, as a consequence we can not identify completely the dynamics in $B_{a,m}^*(0)$ with the quadratic model (but still partially, up to certain depth, see below the rewritten diagram of (12)). The idea is to pull back the beginning of the internal rays so long as they do not contain $v_-(a)$. Similar to the quadratic model, one can define internal rays $R_a(\theta) \subset B_{a,m}^*(0)$ with $\theta \in \Theta_\pm = \{\theta; \exists i \text{ s.t. } 2\theta = 2^i\theta_\pm\}$ where $\theta_\pm = \frac{\pm 1}{2^{k-1}}$. Define similarly R_0^a to R_0^m in (18). There is also a dynamical holomorphic motion of $R_{n+1}^a := f_a^{-n-1}(R_0^a)$. See [23, 3.1] for more details.

$$\begin{array}{ccccccccc}
\Omega_{a,m+p}^{n_a} & \xrightarrow{f_a} & \Omega_{a,m+2p}^{n_a-1} & \xrightarrow{f_a} & \dots & \xrightarrow{f_a} & \Omega_{a,q-p}^1 & \xrightarrow{f_a} & \Omega_{a,0}^0 \\
\downarrow h_a & & \downarrow h_a & & \downarrow h_a & & \downarrow h_a & & \downarrow h_a \\
\Omega_{m+p}^{n_a} & \xrightarrow{P_\lambda} & \Omega_{m+2p}^{n_a-1} & \xrightarrow{P_\lambda} & \dots & \xrightarrow{P_\lambda} & \Omega_{q-p}^1 & \xrightarrow{P_\lambda} & \Omega_0^0
\end{array} \tag{28}$$

In the above diagram, $\phi_a : \overline{\Omega_{a,0}^0} \rightarrow \overline{\mathbb{H}}$ is the Fatou coordiante of $f_a^q|_{B_{a,0}^*(0)}$ normalised by $\phi_a(c_+(a)) = 0$, n_a is the smallest integer such that $\Omega_{a,m+p}^{n_a}$ contains $v_-(a)$.

Definition 5.2.2. Let $a \in \tilde{\mathcal{O}}_n$. The dynamical objects O_n^a of depth n corresponding to the four parameter ones are

$$\bigcup_{\eta \in A_{n+1}} \overline{R_a^\infty(\eta)}, f_a^{-n-1}(E_0^a), f_a^{-n-1}(\overline{R_0^a}), E_a^\infty(r^{1/3^{n+1}}) \tag{29}$$

To conclude from the discussion in this subsection:

Proposition 5.2.3. Let \mathcal{O}_n be a parameter object of depth n , $a, a_0 \in \tilde{\mathcal{O}}_n$. Let $O_{n+1}^a, O_{n+1}^{a_0}$ be the corresponding dynamical object of depth $n+1$. There exists a dynamical holomorphic motion $L_n : \mathcal{O}_n \times O_{n+1}^{a_0} \rightarrow \mathbb{C}$ with $L_n(a, O_{n+1}^{a_0}) = O_{n+1}^a$. Moreover, if O_{n+1}^a is the first one or the third one in (29), $O_{n+1}^a \cap J_a$ is either repelling or pre-periodic to $z = 0$.

5.3 Parameter graphs and puzzles

Parameter graphs \mathcal{Y}_n and puzzles \mathcal{Q}_n in $\overline{\mathcal{S}_m^+}$

Fix some $r > 1$, for each $n \geq 0$ define the graph adapted for parameters of Misiurewicz parabolic type.

$$\mathcal{Y}_n = \bigcup_{k=1}^n \left(\bigcup_{\mathcal{U}_k \subset \mathcal{H}_k \cap \mathcal{S}_m^+} \overline{\mathcal{E}_{n-k}^{\mathcal{U}_k}} \right) \cup \left(\bigcup_{t \in T_n} \overline{\mathcal{R}_\infty(t)} \right) \cup \mathcal{E}_n^m \cup (\mathcal{E}_\infty(r^{1/3^n}) \cap \mathcal{S}_m^+)$$

where

$$T_n = \{t; 3^{n+l}t \in \Theta_m\} \cap [\alpha_{m-1}^+, \beta_m^+], \quad 0 \leq l \leq q-1 \text{ s.t. } lp+m=0 \pmod{q}$$

By Proposition 5.1.1, Corollary 5.1.4, the landing points of equipotentials in parabolic components are Misiurewicz parabolic. By Lemma 6.4.1 these points are also landing points of external rays with angles in T_n . Hence \mathcal{Y}_n is connected, and every connected component of $\mathcal{S}_m^+ \setminus \mathcal{Y}_n$ is simply connected. We call such a connected component \mathcal{Q} a puzzle piece associated to \mathcal{Y}_n if it is bounded and $\partial \mathcal{Q} \cap \mathcal{E}_\infty(r^{1/3^n}) \neq \emptyset$. We denote it by \mathcal{Q}_n in the sequel. By construction, every \mathcal{Q}_{n+1} is contained in a unique puzzle piece \mathcal{Q}_n .

Parameter graphs \mathcal{X}_n and puzzles $\mathcal{P}_n(a_0)$ in $\overline{\mathcal{S}_m^+}$

Next we define the graph adapted for parameters which are **not** of Misiurewicz parabolic type. Let $a_0 \in \mathcal{S}_m^+$ such that f_{a_0} satisfies Assumption (\diamond) . By Theorem 4.3.9, there is a graph (19) infinitely ringing $v_-(a_0)$. Recall that this graph is associated to the angle $\theta_l = \frac{\pm 1}{2^l - 1}$ of an internal ray. Let H be the collection of angles of all external rays in (19) of depth 0. For each $n \geq 0$ consider

$$\mathcal{X}_n = \bigcup_{k=1}^n \left(\bigcup_{\mathcal{U}_k \subset \mathcal{H}_k \cap \mathcal{S}_m^+} \overline{\mathcal{E}_{n-k}^{\mathcal{U}_k} \cup \mathcal{R}_{n-k}^{\mathcal{U}_k}} \right) \cup \left(\bigcup_{\eta \in A_n} \overline{\mathcal{R}_\infty(\eta)} \right) \cup \mathcal{E}_n^m \cup \mathcal{R}_n^m \cup (\mathcal{E}_\infty(r^{1/3^n}) \cap \mathcal{S}_m^+)$$

where

$$A_n = \{\eta; 3^n \eta \in H\} \cap [\alpha_{m-1}^+, \beta_m^+].$$

We call a connected component \mathcal{P} of $\mathcal{S}_m^+ \setminus \mathcal{X}_n$ a puzzle piece associated to \mathcal{X}_n if it is bounded and $\partial\mathcal{P} \cap \mathcal{E}_\infty(r^{1/3^n}) \neq \emptyset$. We denote it by \mathcal{P}_n in the sequel. Clearly every \mathcal{P}_{n+1} is contained in a unique \mathcal{P}_n . Define $\mathcal{P}_n(a_0)$ to be the puzzle piece containing a_0 . This is well-defined since $a_0 \notin \mathcal{X}_n$, for otherwise some external ray $R_{a_0}^\infty(\eta)$ or internal ray will land at a parabolic pre-periodic point or $v_-(a_0)$, contradicting with the construction of the dynamical graph for f_{a_0} (see the observation at the beginning of 4.3).

Lemma 5.3.1. *Let $n \geq 1$. Any $a \in \mathcal{X}_n \cap \partial\mathcal{H} \cap \mathcal{P}_0(a_0)$ is a Misiurewicz parameter. It is the landing point of an internal ray $\mathcal{R} \subset \mathcal{X}_n$ and an external ray $\mathcal{R}_\infty(t) \subset \mathcal{X}_n$. In particular \mathcal{X}_n is connected.*

Proof. By Proposition 5.2.3, landing points of R_a^0 are repelling. Therefore by Lemma 5.1.6, f_a is Misiurewicz. Also each landing point is landed by some external ray $R_{a_0}^\infty(\eta)$ with $R_{a_0}^\infty(\eta)$ involved in the admissible graph of f_{a_0} . By Lemma 6.4.1, a is the landing point of $\mathcal{R}_\infty(\eta)$. Hence \mathcal{X}_n is connected. \square

Dynamics graphs defined up to certain depth

From the discussion in Subsection 5.2, we see that for a satisfying the condition in Proposition 5.2.3, E_0^a, R_0^a are well defined and homeomorphic to the corresponding objects in the quadratic model (17), (18). Set $E_n^a = f_a^{-n}(E_0^a), R_n^a = f_a^{-n}(R_0^a)$. Define

$$Y_n^a = E_n^a \cup \left(\bigcup_{t \in T_n} \overline{R_a^\infty(3^i t)} \right) \cup E_a^\infty(r).$$

$$X_n^a = E_n^a \cup R_n^a \cup \left(\bigcup_{\eta \in A_n} \overline{R_a^\infty(3^i \eta)} \right) \cup E_a^\infty(r).$$

Proposition 5.2.3 gives immediately

Lemma 5.3.2. *Let $a_0 \in \mathcal{Q}_n$. Then for $0 \leq k \leq n+1$ there exists a dynamical holomorphic motion $L_k : \mathcal{Q}_n \times Y_k^{a_0} \rightarrow \mathbb{C}$ with base point a_0 such that $L_k(a, Y_k^{a_0}) = Y_k^a$.*

Lemma 5.3.3. *Let $a_0 \in \mathcal{S}_m^+$ such that f_{a_0} satisfies the Assumption (\diamond) . For $0 \leq k \leq n+1$, there is a dynamical holomorphic motion $L_k : \mathcal{P}_n(a_0) \times X_k^{a_0} \rightarrow \mathbb{C}$ with base point a_0 such that $L_k(a, X_k^{a_0}) = X_k^a$.*

6 Local connectivity of the central part $\partial\mathcal{K}_\lambda$, $\lambda = e^{2\pi i \frac{p}{q}}$

By Yoccoz's Theorem, parameters in

$$\mathcal{C}_\lambda \setminus \left[\mathcal{H}^\lambda \cup \left(\bigcup_{m=0}^{q-1} \mathcal{W}^+(\mathbf{a}_m) \cup \mathcal{W}^-(\mathbf{a}_m) \right) \right]$$

are divided into four cases:

1. Double parabolic: $a \in \mathcal{A}_{p/q}$.
2. Misiurewicz parabolic: $\exists n \geq 1$ s.t. $f_a^n(c_-(a)) = 0$.
3. Renormalisable (in the sense of Douady-Hubbard [6]): there exists $U' \subset \subset U$ surrounding $c_-(a)$ and $k \geq 1$ such that $f^k|_{U'} : U' \rightarrow U$ is quadratic like, and the copy of the quadratic Julia set is connected.
4. Non renormalisable.

In Section 4, we have constructed dynamical admissible graph infinitely ring the free critical point (critical value) for f_a and obtain the dichotomy given by Yoccoz's Theorem. Now we pass this dichotomy to the parameter plane:

Theorem I. *If a satisfies Assumption (\diamond) , i.e. is of case 3 and 4, then there is a dynamically defined nest of puzzles $\{\mathcal{P}_n(a)\}_n$ surrounding a , such that $\bigcap_n \mathcal{P}_n(a) = \{a\}$ if a is non renormalisable; $\bigcap_n \mathcal{P}_n(a) = \mathbf{M}_a$ is a copy of the Mandelbrot set containing a if a is renormalisable. In particular, \mathcal{C}_λ is locally connected at a if a is non renormalisable.*

In general, we do not get local connectivity at all renormalisable parameters, since essentially this needs the MLC conjecture to be true. However for certain a renormalisable (for instance the cusp of the main cardioid or the tips of \mathbf{M}_a), we can separate \mathbf{M}_a from \mathcal{C}_λ by two external rays landing at a . Hence \mathcal{C}_λ is also locally connected at such a . If we are in case 1 or 2, then the parameter puzzles we construct no longer surround a , but have a on their boundaries. But still there is a decreasing sequence of parameter puzzles shrinking to a , since the corresponding decreasing dynamical puzzles shrink to a certain preimage of $z = 0$, and this shrinking property can be passed to parameter puzzles by holomorphic motion.

To summarize, we obtain

Theorem II. *The boundary of every component of \mathcal{H}^λ is a Jordan curve.*

Theorem I actually gives more than local connectivity: the shrinking property of decreasing parameter puzzle pieces gives landing parameter external rays, which provide clean cuts of \mathcal{C}_λ . This enables us to give a global description of \mathcal{C}_λ (see Theorem 6.4.8 for a precise statement):

Theorem III. *\mathcal{C}_λ is decomposed into the central part \mathcal{K}_λ and the limbs attached to them at end points.*

Section 6 is organised as follows: Theorem I together with Theorem II is a synthesis of 6.1, 6.2 and 6.3; 6.4 is devoted to the proof of Theorem III, the Main Theorem and Corollary I.

6.1 Misiurewicz parabolic and double parabolic case

Proposition 6.1.1. *Let $\mathcal{U} \subset \mathcal{S}_m^+$ be a parabolic component. Then $\partial\mathcal{U}$ is locally connected at all Misiurewicz parabolic parameters.*

Proof. Let $a_0 \in \partial\mathcal{U}_k$ be Misiurewicz parabolic. By Lemma 6.4.1, let $t_0^1, t_0^2, \dots, t_0^{q-1}$ be the q angles such that $\mathcal{R}_\infty(t_0^i)$ lands at a_0 . Then $\mathcal{R}_\infty(t_0)$ is part of \mathcal{Y}_n for all $n \geq N$, where N is some fixed integer. Therefore there are exactly $q+1$ adjacent puzzle pieces of depth n $\mathcal{Q}_n^0, \mathcal{Q}_n^1, \dots, \mathcal{Q}_n^q$ attached at a_0 . Clearly $\mathcal{Q}_{n+1}^i \subset \mathcal{Q}_n^i$. Now we claim that for $a \in \mathcal{Q}_n^i$, $v_-(a) \in (Q_n^a)^i$. Indeed, for any $\epsilon > 0$, there exists $n' > n$ large enough and t'_0 such that $3^{n'} t'_0 = 1$, $\mathcal{R}_\infty(t'_0) \cap \mathcal{Q}_n^i \neq \emptyset$ and $|t'_0 - t_0| < \epsilon$. Hence for $a' \in \mathcal{R}_\infty(t'_0) \cap \mathcal{Q}_n^i$, $v_-(a') \in (Q_n^{a'})^\pm$. By Lemma 5.3.2, Y_n^a moves holomorphically for $a \in \mathcal{Q}_n^\pm$. Notice that $v_-(a)$ also moves holomorphically and it does not belong to Y_n^a when $a \in \mathcal{Q}_n^i$, we deduce that $v_-(a) \in (Q_n^a)^i$ since $v_-(a') \in (Q_n^{a'})^i$. Now if $a \in \bigcap_n \mathcal{Q}_n^i$, then a is in \mathcal{C}_λ but does not belong to any parabolic component since a is excluded by every equipotentials in \mathcal{H}_∞ and \mathcal{H} . Moreover $f_a^n(c_-(a)) \neq 0, \forall n \geq 1$ by definition of \mathcal{Q}_n . Therefore by Proposition 4.1.1, $\bigcap_{m \geq 1} (Q_m^a)^i = \emptyset$. But on the other hand $v_-(a) \in \bigcap_n (Q_n^a)^\pm$, a contradiction. Hence $\bigcap_n \mathcal{Q}_n^\pm = \emptyset$. This means that if we set

$$\mathcal{O}_n := \bigcup_{i=0}^q \mathcal{Q}_n^i \cup \bigcup_{i=0}^{q-1} \mathcal{R}_\infty(t_0^i) * (0, r^{1/3^n}) \cup \{a_0\}.$$

then $(\mathcal{O}_n \cap \partial\mathcal{U})_n$ form a basis of connected neighborhood of a_0 . □

Notice that $a_0 = \mathbf{a}_m \in \mathcal{A}_{p/q}$ is not covered by Proposition 6.1.1. But still we can verify the local connectivity with a similar argument.

Proposition 6.1.2. *Let $\mathcal{Q}_n^\pm \subset \mathcal{S}_m$ be the unique two puzzle pieces of depth n containing \mathbf{a}_m on their boundaries. Without loss of generality suppose that their boundaries all contain $\mathcal{R}_\infty(\alpha_m^\pm) * (0, r^{1/3^n})$. Then $\bigcap_n \mathcal{Q}_n^\pm = \emptyset$.*

Proof. We only prove for the sequence $\{\mathcal{Q}_n^+\}_n$, the case left is similar. Suppose the contrary $\bigcap \mathcal{Q}_n^+ \neq \emptyset$. Recall that in Lemma 5.1.2 there is a sequence of external rays $\mathcal{R}_\infty(t_n)$ landing at $\partial\mathcal{B}_m$ with t_k in the preimage of Θ_m and t_n converging monotonously to β_m^+ as $n \rightarrow \infty$. In particular $\mathcal{R}_\infty(t_n) * (0, r^{1/3^n}) \subset \partial\mathcal{Q}_n^+$. Therefore for $a \in \mathcal{Q}_n^+$, the critical value $v_-(a)$ is contained in \mathcal{Q}_n^+ , the puzzle piece whose boundary contains $R_a^\infty(\alpha_m^+) * (0, r^{1/3^n}), R_a^\infty(t_n) * (0, r^{1/3^n})$. Now if $a \in \bigcap_n \mathcal{Q}_n^+$, then $a \in \mathcal{C}_\lambda$ is not Misiurewicz parabolic and not in any parabolic component. This contradicts Proposition 4.1.1 which tells us that $\bigcap_n \mathcal{Q}_n^+ = \emptyset$. \square

6.2 Non renormalizable case

In this subsection, we always fix some $a_0 \in \mathcal{S}_m^+$ such that f_{a_0} satisfies the Assumption (\diamond) . Consider the corresponding graphs \mathcal{X}_n and puzzle pieces $\mathcal{P}_n(a_0)$.

We give the relation between para-puzzles and dynamical puzzles:

Lemma 6.2.1. *Let $n \geq 0$. The mapping $H_n : \mathcal{P}_n(a_0) \cap \mathcal{X}_{n+1} \rightarrow P_n^{a_0, v} \cap X_{n+1}^{a_0}$ defined by $H_n(a) = (L_{n+1}^a)^{-1}(v_-(a))$ is injective. Moreover there exists $N \geq 0$ such that $\forall n \geq N$, H_n is surjective.*

Proof. First we prove that H_n is well-defined. By definition of \mathcal{X}_n , we have $v_-(a) \in P_n^{a, v} \cap X_{n+1}^a$. Consider the holomorphic motion starting at a : $\tilde{L}_{n+1}(a', z) := L_{n+1}(a', (L_{n+1}^a)^{-1}(z))$. By continuity there exists a disk $B(a, r)$ on which $v_-(a')$ is surrounded by $\tilde{L}_{n+1}(a', \partial P_n^{a, v})$. Pick any a' on $\partial B(a, r)$, if $a' \notin \partial\mathcal{P}_n(a_0)$, then for any a'' in a small disk $B(a', r')$, $v_-(a'')$ is surrounded by $\tilde{L}_{n+1}(a'', \partial P_n^{a, v})$. Hence we can extend this property step by step until we reach $\partial\mathcal{P}_n(a_0)$. In particular $H_n(a) \in P_n^{a_0, v}$. By Lemma 5.3.3, $\partial P_{n+1}^{a_0, v}$ moves holomorphically when $a \in \mathcal{P}_n(a_0)$, hence $H_n(a) \in X_{n+1}^{a_0}$.

Next we verify injectivity. The injectivity is clear when a belongs to parameter external rays and equipotentials since in the dynamical plan, angles of external rays and equipotentials are preserved by L_{n+1}^a . We will only prove injectivity for a belongs to parameter internal rays. The proof for internal equipotentials is similar. Suppose there are two distinct parameters a, a' such that $H_n(a) = H_n(a')$. Then clearly a, a' belong to different components $\mathcal{U}, \mathcal{U}'$. Set $a \in \overline{\mathcal{R}_\mathcal{U}}, a' \in \overline{\mathcal{R}_{\mathcal{U}'}}$.

- First suppose that the landing point b, b' of $\mathcal{R}_\mathcal{U}, \mathcal{R}_{\mathcal{U}'}$ do not coincide. Consider the external rays $\mathcal{R}_\infty(s), \mathcal{R}_\infty(s')$ involved in \mathcal{X}_{n+1} landing at b, b' respectively. Then the two rays $H_n(\mathcal{R}_\infty(s)), H_n(\mathcal{R}_\infty(s'))$ land at a common point $x(a_0)$ since $H_n(a) = H_n(a')$. But this is impossible since f_{a_0} is injective near the forward orbit of $x(a_0)$.
- Next if $b = b'$. There are two possibilities: $H_n(\mathcal{R}_\infty(s)) = H_n(\mathcal{R}_\infty(s'))$ or they are two different rays landing at a common point $x(a_0)$. While in the graph $X_{n+1}^{a_0}$, only one internal ray lands at $x(a_0)$, so the second is impossible. Then there exists two other internal rays $\tilde{\mathcal{R}}_\mathcal{U} \subset \mathcal{U}, \tilde{\mathcal{R}}_{\mathcal{U}'} \subset \mathcal{U}'$ landing at \tilde{b}, \tilde{b}' respectively which have the same image under H_n . Then $\tilde{b} \neq \tilde{b}'$, for otherwise one finds a loop in $\overline{\mathcal{U} \cup \mathcal{U}'}$ surrounding points in \mathcal{H}_∞ . We repeat the argument above to \tilde{b}, \tilde{b}' and get a contradiction.

Finally we verify surjectivity. First we prove

Claim. $\partial\mathcal{P}_n(a_0) \cap (\mathcal{R}_\infty(\alpha_{m-1}^+) \cup \mathcal{R}_\infty(\beta_m^+)) = \emptyset$ for n large enough.

Proof of the claim. Suppose for example $\partial\mathcal{P}_0(a_0)$ contains parts of β_m^+ (the proof proceeds the same way if it is α_{m-1}^+). Recall in Lemma 5.1.2, there is a sequences of external rays $\mathcal{R}_\infty(s_n)$ landing at $a_n \in \partial\mathcal{B}_m$ with $s_n \rightarrow \beta_m^+$ and a_n Misiurewicz parabolic. For n large enough, let $\mathcal{R}_n \subset \mathcal{B}_m \cap \mathcal{X}_n \cap \mathcal{P}_0(a_0)$ be a sequence of internal rays landing at $b_n \in \partial\mathcal{B}_m$. By Lemma 5.3.1, there exists $\mathcal{R}_\infty(t_n) \subset \mathcal{X}_n$ landing at b_n with $|t_n - \beta_m^+| < |s_n - \beta_m^+|$. Thus if $\partial\mathcal{P}_n(a_0) \cap \mathcal{R}_\infty(\beta_m^+) \neq \emptyset$, then $a_0 \in \mathcal{Q}_n^+$, where \mathcal{Q}_n^+ is as defined in Proposition 6.1.2. Recall there we have shown that $\bigcap_n \mathcal{Q}_n^+ = \emptyset$. Hence $\partial\mathcal{P}_n(a_0) \cap \mathcal{R}_\infty(\beta_m^+) = \emptyset$ for n large enough. \square

H_n is therefore surjective on the part of external equipotential and external rays by its injectivity and the claim. So we verify surjectivity for internal rays and for internal equipotentials will be similar. Let

$z_0 \in X_{n+1}^{a_0} \cap B_{a_0}(0) \cap P_n^{a_0, v}$. Suppose that z_0 is on some connected component ζ of $f_{a_0}^{-k}(R_0^{a_0})$. Let x_0 be the landing point of ζ , then there is an external ray $R_{a_0}^\infty(t) \subset X_{n+1}^{a_0}$ landing at x_0 . By Lemma 5.3.1, $\mathcal{R}_\infty(t)$ lands at a Misiurewicz parameter a' which is accessible by some internal ray \mathcal{R}_U (Lemma 5.1.7). Clearly $H_n(\mathcal{R}_U) = \zeta$. \square

Corollary 6.2.2. *Let $a \in \mathcal{P}_{n-1}(a_0)$. Let C_n^a be the puzzle piece bounded by $L_n(\partial P_n^{a_0, v})$. Then $a \notin \overline{\mathcal{P}_n(a_0)}$ if and only if $v_-(a) \notin \overline{C_n^a}$.*

Proof. If $a \notin \overline{\mathcal{P}_n(a_0)}$, then take a simple path $a_t \subset \mathcal{P}_{n-1}(a_0)$ connecting a_0, a such that $a_t \cap \mathcal{P}_n(a_0)$ only contains one point. Thus Lemma 6.2.1 ensures that once a_t goes out of $\mathcal{P}_n(a_0)$, $f_{a_t}(c_-(a_t))$ will never enter again $\overline{C_n^a}$. Conversely if $a \in \mathcal{P}_n(a_0)$, then clearly $v_-(a) \in C_n^a$ since $\partial P_n^{a_0, v}$ moves holomorphically and $v_-(a)$ does not intersect X_n^a . \square

Corollary 6.2.3. *For n large enough, if $\overline{P_{n+1}^{a_0, v}} \subset P_n^{a_0, v}$, then $\overline{\mathcal{P}_{n+1}(a_0)} \subset \mathcal{P}_n(a_0)$. Moreover*

$$H_n : \partial \mathcal{P}_{n+1}(a_0) \longrightarrow \partial P_{n+1}^{a_0, v}$$

is a homeomorphism.

Proof. By Lemma 6.2.1 $H_n^{-1}(\overline{\partial P_{n+1}^{a_0, v}}) \subset \mathcal{P}_n(a_0)$ bounds a puzzle piece \mathcal{P} . So it suffices to prove that $a_0 \in \mathcal{P}$. Suppose not, then there is $a' \in \partial \mathcal{P}$ but $a' \notin \overline{\mathcal{P}_{n+1}(a_0)}$. By Corollary 6.2.2, $v_-(a') \notin \overline{C_{n+1}^{a'}}$, a contradiction. \square

Corollary 6.2.4. *Let $\mathcal{U} \subset \mathcal{S}_m^+$ be a capture component. If $a_0 \in \partial \mathcal{U}$ or $\partial \mathcal{B}_m$ is non renormalisable, then $v_-(a_0)$ eventually hit the boundary of immediate basins.*

Proof. By Corollary 6.2.3, $\partial P_n^{a_0, v}$ intersects a Fatou component U (preimage to some immediate basin) for all n . Hence by Theorem 4.3.9 $\bigcap_n \overline{P_n^{a_0, v}}$ intersects ∂U at $v_-(a_0)$. \square

By Theorem 4.3.9, there exists graph (19), such that in the dynamical plane of f_{a_0} one has a sequence of non-degenerated annuli $A_{n_i}^{a_0} := P_{n_i}^{a_0, v} \setminus \overline{P_{n_i+1}^{a_0, v}}$. The above corollary implies that in the parameter plane, $\mathcal{A}_{n_i}^{a_0} := \mathcal{P}_{n_i}(a_0) \setminus \overline{\mathcal{P}_{n_i+1}(a_0)}$ is also non-degenerated for i large enough. Applying Shishikura's trick [18], we can get the distortion control between the moduli of para-annuli and dynamical annuli:

Lemma 6.2.5. *There exists $K > 1$ such that for i large enough*

$$\frac{1}{K} \text{mod}(A_{n_i}^{a_0}) \leq \text{mod}(\mathcal{A}_{n_i}) \leq K \text{mod}(A_{n_i}^{a_0}).$$

Corollary 6.2.6. \mathcal{C}_λ *is locally connected at non renormalisable parameters.*

Proof. This follows immediately from the above lemma and Grötzsch's inequality. \square

6.3 Renormalizable case

In this subsection, we always fix some $a_0 \in \mathcal{S}_m^+$ such that f_{a_0} satisfies the Assumption (\diamond) and is renormalisable. Recall that a set $\mathbf{M}' \subset \mathbb{C}$ is called a copy of the Mandelbrot set (cf. [6]) \mathbf{M} if there exists $k \geq 1$ and a homeomorphism $\chi : \mathbf{M}' \longrightarrow \mathbf{M}$ such that f_a is k -renormalisable and f_a^k is quasiconformally conjugated to $z^2 + \chi(a)$.

Proposition 6.3.1. $\mathbf{M}_{a_0} := \bigcap \mathcal{P}_n(a_0)$ *is a copy of the Mandelbrot set. Moreover, there are exactly 2 external rays landing at the cusp $\chi^{-1}(\frac{1}{4})$, separating \mathbf{M}_{a_0} with \mathcal{B}_m . In particular, if $a_0 \in \partial \mathcal{B}_m$, then $a_0 = \chi^{-1}(\frac{1}{4})$, $\partial \mathcal{B}_m$ is locally connected at a_0 .*

Proof. For the first statement, see [16, Prop. 3.26]. As for the second statement, the proof for the existence is similar to Step 1, Step 2 in Lemma 2.3.8; for the uniqueness, see the proof of [16, Thm. 3] \square

Remark 6.3.2. By the above proposition, the wake $\mathcal{W}(a)$ and limb $\mathcal{L}(a)$ can be defined in the classical sense.

Proposition 6.3.3. Let $\mathcal{U} \subset \mathcal{S}_m^+$ be a capture component. Suppose $a_0 \in \partial\mathcal{U}$ is renormalisable. Then a_0 is a tip of \mathcal{M}_{a_0} , with $\mathcal{M}_{a_0} \cap \mathcal{B}_m = a_1$. Moreover there are exactly two external rays landing at a_0 , separating \mathcal{U} with \mathcal{M}_{a_0} . In particular, $\partial\mathcal{U}$ is locally connected at a_0 .

Proof. See [16, Prop. 4.15, Thm. 4]. \square

Proof of Theorem II. Let \mathcal{U} be an open connected component of \mathcal{H}^λ . Corollary 5.1.5 solves already the case where \mathcal{U} is contained in a double parabolic wake. So suppose \mathcal{U} is in the complement. From the discussion in 6.1, 6.2 and 6.3 we see that $\partial\mathcal{U}$ is locally connected. It is then a Jordan curve by Lemma 2.2.4. \square

6.4 Global descriptions

Lemma 6.4.1. Let $a_0 \in \mathcal{C}_\lambda$ be Misiurewicz parabolic. Suppose moreover that a_0 is not in any double parabolic wake. Then there are q external rays landing at it, each two adjacent rays bound a parabolic component.

Proof. One can prove the existence of parabolic component attached to a_0 by showing that there are parameter equipotential rays landing at a_0 . To prove the existence of landing rays, one uses holomorphic motion and Rouché's Theorem, similar to the proof of Proposition 3.5.4. \square

Let $a_0 \in \partial\mathcal{B}_m$ be Misiurewicz parabolic. By Lemma 6.4.1, there are q external rays (whose angles are preimages of angles in Θ_m) landing at a_0 each two adjacent rays separate a capture component attached at a_0 with \mathcal{B}_m . Applying a similar argument as in the proof of Lemma 5.1.2, one can show that when a sequence of Misiurewicz parabolic parameters $a_n \in \partial\mathcal{B}_m$ converges to a_0 , the angles of external rays landing at a_n converges to the biggest or smallest angle among the angles of the q external rays landing at a_0 . Therefore there are **exactly** q external rays landing at a_0 and **exactly** $q - 1$ capture components attached at a_0 . A similar result holds for the capture components attached at $\partial\mathcal{B}_m$, also for the next "generation" of capture components attached, and so on.

Definition 6.4.2. Let $0 \leq m \leq \lfloor \frac{q}{2} \rfloor$. A capture component $\mathcal{U}_1 \subset \mathcal{S}_m^+$ is of level 1, if it is attached to $\partial\mathcal{B}_m$ at a Misiurewicz parabolic parameter. A capture component \mathcal{U}_n is of level $n \geq 2$ if it is not of level $n - 1$, and is attached to some $\partial\mathcal{U}_{n-1}$ at a Misiurewicz parabolic parameter with \mathcal{U}_{n-1} of level $n - 1$. The point $\mathbf{r}_n = \partial\mathcal{U}_n \cap \partial\mathcal{U}_{n-1}$ is called the root of \mathcal{U}_n . The open region bounded by the two external rays containing \mathcal{U}_n is called the wake of \mathcal{U}_n , denoted by $\mathcal{W}_{\mathcal{U}_n}$. The corresponding limb is defined to be $\mathcal{L}_{\mathcal{U}_n} = \overline{\mathcal{W}_{\mathcal{U}_n}} \cap \mathcal{C}_\lambda$.

Definition 6.4.3. Set $\mathcal{K}_0^m = \mathcal{B}_m \cap \mathcal{S}_m^+$, \mathcal{K}_n^m to be the union of all level $n \geq 1$ capture components in \mathcal{S}_m^+ , $\mathcal{K}^m := \bigcup_{n \geq 0} \mathcal{K}_n^m$ is called a **central component**. Let $\mathcal{K} := \bigcup_{m=0}^{\lfloor q/2 \rfloor} \mathcal{K}^m$. Similarly, if we work in \mathcal{S}_m^- , we can construct $\tilde{\mathcal{K}}_n^m, \tilde{\mathcal{K}}^m, \tilde{\mathcal{K}}$.

Construction of \mathfrak{G} in the Main Theorem. Recall $P_\lambda(z) = \lambda z + z^2$, K_λ its filled-in Julia set, Ω_0^0 its maximal petal contained in $B_0^*(0)$, Ω_l^k defined as in (34). Let D_0, \tilde{D}_0 be the two connected components of $B_0^*(0) \setminus \overline{P_\lambda^{-1}(\Omega_p^{-1})}$, such that D_0 is on the left-hand side of \tilde{D}_0 . Let $\tilde{\Omega} = P_\lambda^{-1}(\Omega_p^{-1}) \setminus \Omega_0^0$. Recall also the parametrisations in 3.5: $\Psi_0^{adj} : \mathcal{B}_m \setminus \mathcal{I}_0 \rightarrow B_p^*(0)$, $\Psi_m^{bit} : \mathcal{D}_m \setminus \mathcal{I}_m \rightarrow B_{m+p}^*(0) \setminus \overline{\Omega_{m+p}^{-s}}$ and $\Psi_{\mathcal{U}_k} : \mathcal{U}_k \rightarrow B_l^*(0)$. By Proposition 3.3.5 and 3.5.4, the following mapping is an isomorphism from $\mathcal{B}_0 \setminus \mathcal{I}_0$ to $B_0^*(0) \setminus \overline{\Omega_0^0}$:

$$a \mapsto \begin{cases} (P_\lambda|_{D_0})^{-1} \circ \Psi_0^{adj}(a), & \text{if } a \in \mathcal{D}_0, \\ (P_\lambda|_{\tilde{D}_0})^{-1} \circ \Psi_0^{adj}(a), & \text{if } a \in \tilde{\mathcal{D}}_0, \\ [(P_\lambda|_{\overline{\Omega} \setminus \{-\frac{\lambda}{2}\}})^{-1} \circ \Psi_0^{adj}(a), & \text{if } a \in \overline{W} \setminus \{\sqrt{3\lambda}\} \end{cases} \quad (30)$$

Next, by Proposition 3.5.2, the following mapping is an isomorphism from $\mathcal{D}_m \setminus \mathcal{I}_m$ to $B_m^*(0) \setminus \overline{\Omega_m^{-s+1}}$, where $2 \leq s \leq q$ is the unique integer such that $-sp + m + p = 0 \pmod{q}$:

$$a \mapsto (P_\lambda|_{B_m^*(0)})^{-1} \circ \Psi_m^{bit}(a) \quad (31)$$

Finally, for a capture component \mathcal{U}_n of \mathcal{K}_n^m , we define its address $[\mathcal{U}_n]$ as the address of the Fatou component containing $c_-(a)$ for $a \in \mathcal{U}_n$ (recall 21). By Proposition 3.5.3 the mapping defined below

$$a \mapsto (P_\lambda|_{U_n})^{-1} \circ \Psi_{\mathcal{U}_n}(a) \quad (32)$$

is an isomorphism from \mathcal{U}_n to U_n , where U_n is the Fatou component of P_λ with address $[\mathcal{U}_n]$. Combining (30) (31) (32), we have constructed an bijection ($s = 1$ if $m = 0$)

$$\mathfrak{G}^m : \bigcup_{n \geq 0} \mathcal{K}_n^m \longrightarrow (K_\lambda^\circ \cap S_m) \setminus \overline{\Omega_m^{-s+1}}.$$

For $1 \leq m \leq \lfloor \frac{q}{2} \rfloor$, if we work in \mathcal{S}_m^- , then similarly we can define $\tilde{\mathcal{K}}_n^m$, $\tilde{\mathcal{K}}^m$ and construct an bijection

$$\tilde{\mathfrak{G}}^m : \bigcup_{n \geq 0} \tilde{\mathcal{K}}_n^m \longrightarrow (K_\lambda^\circ \cap S_{q-m}) \setminus \overline{\Omega_{q-m}^{l+1}}.$$

where $0 \leq l \leq q-1$ is the unique integer such that $lp + p - m = 0 \pmod{q}$. So to prove the Main Theorem, it remains to extend \mathfrak{G}^m (resp. $\tilde{\mathfrak{G}}^m$) to $\mathcal{K}^m \cap \mathcal{S}_m^+$ (resp. $\tilde{\mathcal{K}}^m \cap \mathcal{S}_m^-$). We will only do the extension for \mathfrak{G}^m , since it is the same for $\tilde{\mathfrak{G}}^m$.

The set $\mathcal{K}^m \setminus \bigcup_{n \geq 0} \mathcal{K}_n^m$. By Proposition 6.3.3, there are no renormalisable parameter on $\partial\mathcal{U}_n$ for $n \geq 1$. By Corollary 6.2.4, if $a \in \partial\mathcal{U}_n$, $c_-(a)$ eventually hits the boundary of immediate basins for $a \in \partial\mathcal{U}_n$. For $a \in \partial\mathcal{U}_n$, define its address by

$$[[a]] := [c_-(a_n)] = [(m, \omega_0); (\epsilon_1, \omega_1); \dots; (\epsilon_n, \omega_n)]. \quad (33)$$

where $\omega_0, \dots, \omega_{n-1}$ are dyadic, $\omega_n \in (0, 1)$; $0 \leq \epsilon_k \leq q-1$ is chosen so that $f_{a_n}^k(c_-(a_n)) \in S_{a, \epsilon_k}$ (S_{a, ϵ_k} is the sector S_{ϵ_k} in 4.4). If $a \in \partial\mathcal{B}_m \setminus \mathcal{A}_{p/q}$, define $[[a]]$ to be $[r_a]$ if a is renormalisable, where $r_a \in \partial B_{a,m}^*(0) \setminus \{0\}$ is the parabolic periodic point (Proposition 6.3.1); otherwise $[[a]] := [c_-(a_n)]$.

Lemma 6.4.4. *For $a_0 \in \partial\mathcal{U}_n$ or $\partial\mathcal{B}_m$ non renormalisable, there is a unique external ray landing at it.*

Proof. This comes from Proposition 4.4.1 and the homeomorphism between para-puzzles and dynamical puzzles, cf. Corollary 6.2.3. \square

Lemma 6.4.5. *We have decompositions*

$$\begin{aligned} \mathcal{S}_m^+ \cap \mathcal{C}_\lambda &= (\overline{\mathcal{B}_m} \cap \mathcal{S}_m^+) \cup \left(\bigcup_{\mathcal{U}_1 \subset \mathcal{K}_1^m} \mathcal{L}_{\mathcal{U}_1} \cup \bigcup_{\substack{a \in \partial\mathcal{B}_m \cap \mathcal{S}_m^+ \\ \text{parabolic}}} \mathcal{L}(a) \right) \\ \mathcal{L}_{\mathcal{U}_n} &= \overline{\mathcal{U}_n} \cup \bigcup_{\substack{\mathcal{U}_{n+1} \subset \\ \mathcal{K}_{n+1}^m \cap \mathcal{W}_{\mathcal{U}_n}}} \mathcal{L}_{\mathcal{U}_{n+1}}. \end{aligned}$$

As a direct consequence, $\overline{\mathcal{K}_n^m} \setminus \mathcal{A}_{p/q} = \bigcup_{\mathcal{U}_n \subset \mathcal{K}_n^m} \overline{\mathcal{U}_n}$.

Proof. This is a direct consequence of the above lemma. Let us do the proof for $\mathcal{S}_m^+ \cap \mathcal{C}_\lambda$. For any $n \geq 0$, consider all the Misiurewicz parabolic parameters of depth n (that is, $f_a^n(v_-(a)) = 0$) on $\partial\mathcal{B}_m \cap \mathcal{S}_m^+$ and the q unique landing external rays. These rays together with $\partial\mathcal{B}_m$ separate $\mathcal{S}_m^+ \setminus \overline{\mathcal{B}_m}$ into several open sectors of depth n . Take $b \in \mathcal{C}_{p/q} \cap (\overline{\mathcal{U}_0})^c \cap \mathcal{S}_m^+$ not in any $\mathcal{L}_{\mathcal{U}_1}$. Let $\mathcal{S}_n(b)$ be the sector of depth n containing b . Let $\mathcal{R}_\infty(t_n), \mathcal{R}_\infty(t'_n)$ be the two external rays bounding $\mathcal{S}_n(b)$ and a_n, a'_n their landing point respectively. Clearly a_n, a'_n converges to some $a \in \partial\mathcal{B}_m$ since $\partial\mathcal{B}_m$ is a Jordan curve. Then a must be renormalisable. If not, then by the previous lemma, t_n, t'_n converge to the same angle, which implies that $b = a$, a contradiction since we take $b \notin \overline{\mathcal{B}_m}$. \square

Proposition 6.4.6. *Let $a_0 \in \mathcal{K}^m \setminus \bigcup_{n \geq 0} \overline{\mathcal{K}_n^m}$ not be double parabolic. Then a_0 is contained in a infinite sequence of limbs, i.e. there exist $\mathcal{U}_n \subset \mathcal{K}_n$ such that $a_0 \in \bigcap_n \mathcal{L}(\mathcal{U}_n)$. Moreover a_0 can not be Misiurewicz parabolic. If a_0 is non renormalisable, there is only one external ray landing at it; if it is renormalisable, then it is the cusp of \mathbf{M}_{a_0} and exactly two external rays land at it.*

Proof. The existence of \mathcal{U}_n is just a direct consequence of Lemma 6.4.5. Next we prove that a_0 is not Misiurewicz parabolic. Suppose the contrary that $f_{a_0}^N(v_-(a_0)) = 0$. Let $a_n \in \mathcal{U}_{N_n} \subset \mathcal{K}_{N_n}^m$ be a sequence converging to a_0 and suppose $f_{a_n}^{M_n}(v_-(a_n)) = 0$. Clearly $M_n \geq n$. Fix $N' \gg N$, set $N'' := M_{N'} - N$. Let $\tilde{B}_{N''}$ be the connected component containing 0 of $f_{a_0}^{-N''}(\bigcup_i B_{a_0, i}^*(0))$. Notice that there exists a neighborhood \mathcal{V} of a_0 on which there is a dynamical holomorphic motion h_a of $\tilde{B}_{N''}$. By construction, $f_{a_n}^N(v_-(a_n))$ is bounded by a wake attached at a preimage of 0 (not equal to 0) on $\partial(h_a(\tilde{B}_{N''}))$. Moreover the angles of the two external rays $R_{a_n}^\infty(t_1), R_{a_n}^\infty(t_2)$ determining this wake do not depend on n . Shrink \mathcal{V} if necessary so that there is a holomorphic motion of $R_{a_0}^\infty(t_1) \cup R_{a_0}^\infty(t_2)$. This implies that $f_{a_n}^N(v_-(a_n))$ does not converge to 0 as $n \rightarrow \infty$, contradicting $a_n \rightarrow a_0$.

So a_0 is either non renormalisable or renormalisable. Suppose the first case. The existence of external ray landing comes from the shrinking property of parameter puzzles around a_0 (Theorem 4.3.9 and Lemma 6.2.5). The uniqueness comes from Lemma 6.2.1 and Lemma 4.4.4. Next suppose a_0 is renormalisable. By Proposition 6.3.1, \mathbf{M}_{a_0} is separated from \mathcal{K}^m by two external rays landing at the cusp. While $a_0 \in \mathcal{K}^m$, it has to be the cusp. \square

We conclude that $\partial\mathcal{K}^m$ is combinatorically rigid:

Corollary 6.4.7. *Suppose $a_0 \in \mathcal{K}^m \setminus \bigcup_{n \geq 0} \overline{\mathcal{K}_n^m}$. Let $\{\mathcal{U}_n\}_n$ be as in Proposition 6.4.6. Then $\bigcap_n \mathcal{L}(\mathcal{U}_n) \cap \mathcal{K}^m = \{a_0\}$.*

Proof. This comes from the shrinking property of puzzle pieces $\mathcal{P}_n(a)$ when a is non renormalisable; for a renormalisable, it comes from the shrinking property of $\mathcal{P}_n(a) \setminus \overline{\mathcal{W}(a)}$. \square

Theorem 6.4.8. *We have the decomposition*

$$\mathcal{S}_m^+ \cap \mathcal{C}_\lambda = \mathcal{K}^m \cap \mathcal{S}_m^+ \cup \bigcup_{\substack{a \in \partial\mathcal{K}^m \\ \text{renormalisable}}} \mathcal{L}(a).$$

In particular, $\mathcal{K}_\lambda = \mathcal{K} \cup -\mathcal{K} \cup \tilde{\mathcal{K}} \cup -\tilde{\mathcal{K}}$.

Proof. Here \mathcal{K} is parallel to $\overline{\mathcal{B}_m}$ in Lemma 6.4.5, and instead of considering sectors defined by rays landing at $\partial\mathcal{B}_m$, one should consider sectors defined by rays landing at $\partial\mathcal{K}^m$. One then concludes by using the landing property given in Proposition 6.4.6. \square

Extension of \mathfrak{G}^m . By Proposition 6.4.6, every $a \in \partial\mathcal{K}^m \setminus \bigcup_n \partial\mathcal{K}_n^m$ can be represented by a infinite sequence

$$[[a]] := [(m, \omega_0); (\epsilon_1, \omega_1); (\epsilon_2, \omega_2); \dots; (\epsilon_n, \omega_n), \dots]$$

with ω_k dyadic, $0 \leq \epsilon_k \leq q-1$ are chosen so that $f_{a_n}^k(c_-(a_n)) \in S_{a, \epsilon_k}$. Compare with Remark 4.4.3 the way we encode $J_\lambda \setminus \{0\}$, where J_λ is the Julia set of the quadratic model $P_\lambda = \lambda z + z^2$.

End of the proof of the Main Theorem. First of all, \mathfrak{G}^m extends continuously to $\bigcup_n \overline{\mathcal{K}_n^m}$, since by Lemma 6.4.5, $\overline{\mathcal{K}_n^m} \cap S_m^+ = \bigcup_{\mathcal{U}_n \subset \mathcal{K}_n^m} \overline{\mathcal{U}_n}$, and by Theorem II, every component of \mathcal{K}_n^m is a Jordan curve. Let $\{a_k\}_k \subset \bigcup_n \overline{\mathcal{K}_n^m}$ be a sequence converging to some $b \in (\mathcal{K}^m \cap S_m^+) \setminus \bigcup_n \overline{\mathcal{K}_n^m}$. Let \mathcal{U}_k be the parabolic component whose closure containing a_k , and $\{\mathcal{U}_n(b)\}_n$ the sequence of parabolic components associated to b given by Proposition 6.4.6. Then by Corollary 6.4.7, for any n , the first n coordinates for $[[a_k]]$ will be the same for all k large enough, i.e. equal to $[[\mathcal{U}_n(b)]]$. Define $\mathfrak{G}^m(b) := \lim_{n \rightarrow \infty} \Xi^{-1}([[\mathcal{U}_n(b)]]]$. Clearly $\mathfrak{G}^m|_{\partial\mathcal{K}^m \cap S_m^+}$ coincides with the following mapping

$$\mathfrak{H} : \partial\mathcal{K}^m \cap S_m^+ \longrightarrow J_\lambda \cap S_m, \quad a \mapsto \Xi^{-1}([a]),$$

where Ξ^{-1} is defined in (23). Now we prove that \mathfrak{H} is a homeomorphism. Clearly \mathfrak{H} is continuous and surjective on $\partial\mathfrak{H}^m \cap S_m^+$. Moreover it is injective on $\partial\mathcal{K}^m \cap \mathcal{K}_n^m$ for all n . The injectivity on $\partial\mathcal{K}^m \setminus \bigcup_n \partial\mathcal{K}_n^m$ comes from the combinatorial rigidity (Corollary 6.4.7) i.e. the mapping $a \mapsto [[a]]$ is injective. It remains to verify continuity of \cdot . Take any sequence of $\{z\}_k \subset J_\lambda$ converging to some $w \in J_\lambda$. If w is on the boundary of some Fatou Component, then the continuity of \mathfrak{H}^{-1} at w is guaranteed by Lemma 6.4.5; if w an end point, then the continuity is given by Corollary 6.4.7. \square

Proof of Corollary I. Clearly $\mathring{\mathcal{K}}_\lambda \subset \mathcal{CU}_\lambda$. Without loss of generality we may suppose $a \in \partial\mathcal{K}^m$. Then a is either Misiurewicz parabolic, either non renormalisable, either double parabolic or renormalisable with a parabolic periodic point of multiplier 1 (Proposition 6.3.1 and 6.4.6). Moreover there is only one external ray landing at every repelling periodic point of f_a (Proposition 4.4.1 Lemma 4.4.4). Thus there are no repelling cut point in J_a , $a \in \mathcal{CU}_\lambda$.

Conversely, take $a \in \mathcal{CU}_\lambda$. Without loss of generality let $a \in S_m^+$. By Theorem 6.4.8, $a \in \mathcal{K}^m \subset \mathcal{K}_\lambda$. \square




Figure 9: $\mathcal{K}^0 \cap S_0^+$ and $K_\lambda \cap S_0$ with $\lambda = e^{2\pi i \frac{1}{3}}$.

Appendices

A Rotation number

Definition A.1. Let $f : \mathbb{R}/\mathbb{Z} \rightarrow \mathbb{R}/\mathbb{Z}$ be the d -fold covering map, i.e. $f(\theta) = d\theta$. A periodic cycle $\Theta = \{\theta_0, \dots, \theta_{q-1}\}$ has rotation number $\frac{p}{q}$ if $\forall i, f(\theta_i) = \theta_{i+p \bmod q}$.

Theorem A.2 ([8]). The d -fold mapping $\theta \mapsto d\theta$ has $\binom{d+q-2}{q}$ periodic cycles of rotation number p/q .

Lemma A.3. If Θ has rotation number p/q , then $\tilde{\Theta} = \{\tilde{\theta}_0, \dots, \tilde{\theta}_{q-1}\}$ has rotation number $1 - \frac{p}{q}$, where $\tilde{\theta}_i = 1 - \theta_{q-i-1}$.

Proof. By definition, it suffices to prove that $f(\tilde{\theta}_i) = \tilde{\theta}_{i+q-p \bmod q}$:

$$f(\tilde{\theta}_i) = d(1 - \theta_{q-i-1}) = -d\theta_{q-i-1} = -\theta_{q-i-1+p \bmod q} = -\theta_{-i-1+p \bmod q} = \tilde{\theta}_{i+q-p \bmod q}.$$

□

B Pinching deformation

This part is a resume of the theory of pinching deformation developed by Cui-Tan in [5].

Let $f : \hat{\mathbb{C}} \rightarrow \hat{\mathbb{C}}$ be a rational map. Denote by $\mathcal{F}_f, \mathcal{J}_f$ the Fatou, Julia set respectively. Denote by Crit_f the set of critical points of f . Define the post-critical set of f by

$$\text{Post}_f := \overline{\bigcup_{n>0} f^n(\text{Crit}_f)}.$$

f is called **geometrically finite** if the accumulation points of Post_f is finite. Define $\tilde{\mathcal{R}}_f \subset \mathcal{F}_f$ as follows: $z \in \tilde{\mathcal{R}}_f$ if and only if

1. z does not belong to super-attracting basins,
2. $\#(\{f^n(z); n \geq 0\}) = +\infty$ and $\{f^n(z); n \geq 0\} \cap \text{Post}_f = \emptyset$.

Definition B.1. The **quotient space** \mathcal{R}_f of f is defined as the set $\tilde{\mathcal{R}}_f$ quotient the grand orbit equivalence relation \sim , i.e. $z_1 \sim z_2$ if and only if $\exists m, n \geq 0, f^m(z_1) = f^n(z_2)$. Denote by $\pi_f : \tilde{\mathcal{R}}_f \rightarrow \mathcal{R}_f$ the natural projection.

Remark B.2. \mathcal{R}_f has only finitely many components, each of them is either an at least 1-punctured torus, corresponding to an attracting basin, of an at least 1-punctured infinite cylinder, corresponding to a parabolic basin. Refer to [14] for details.

In the sequel we always assume that $\tilde{\mathcal{R}}_f \neq \emptyset$.

Lemma B.3. Let γ be a Jordan curve. Then either each component of $\pi_f^{-1}(\gamma)$ is a Jordan curve or each component of $\pi_f^{-1}(\gamma)$ is an eventually periodic arc. Moreover, every eventually periodic arc β lands at both ends.

Definition B.4. Let β be a k -periodic arc and $b \in \beta$. The limit point of the forward (resp. backward) orbit of b under f^k is called the **attracting end** (resp. **repelling end**). Notice that the two ends might coincide if both of them are parabolic.

Definition B.5. A **multi-annulus** $\mathcal{A} \subset \mathcal{R}_f$ is a finite disjoint union of annuli whose boundaries are pairwise disjoint simple closed curves such that each component of $\pi_f^{-1}(e(\mathcal{A}))$ is an eventually periodic arc, where $e(\mathcal{A})$ is the union of the equators of the annuli in \mathcal{A} . A multi-annulus \mathcal{A} is called **non-separating** if for any choice of finitely many components of $\pi_f^{-1}(e(\mathcal{A}))$, the closure of their union (denote by K) does not separate \mathcal{J}_f , i.e. there is only one component of $\hat{\mathbb{C}} \setminus K$ intersecting \mathcal{J}_f .

Definition B.6. A component B of $\pi_f^{-1}(\mathcal{A})$ is called a **band**. Notice that B is a topological open disk bounded by two eventually periodic arcs, and hence itself is also eventually periodic and lands at both ends. The **core arc** β of B is the lift of the equator of $\pi_f(B)$ to B , whose two (or maybe one) end points are exactly those of B . A band B (resp. its core arc β) is of **level** n ($n \geq 0$) if n is the smallest integer such that $f^n(B)$ (resp. $f^n(\beta)$) is periodic. A **skeleton of level** n is a connected component of $\bigcup_{\text{level } n} \beta$. The **fill-in** of a skeleton S , denote by \hat{S} , is the union of S with all its complement components disjoint from \mathcal{J}_f . \hat{S} is called a **fill-in skeleton**.

For $r > 1$, denote by $\mathbb{A}(r) := \{z; \frac{1}{r} < |z| < r\}$, for $r > 1$. For $t \geq 0$, let $w_{t,r} : \mathbb{A}(r) \rightarrow \mathbb{A}(r^{1+t})$ be the pinching model defined as in [5], 5.1. Let $\mathcal{A} = \cup A_i$ be a non-separating multi-annulus. Let $\chi_i : A_i \rightarrow \mathbb{A}(r_i)$ be a conformal representation and $\mu_{i,t}$ be the Beltrami differential of $w_{t,r_i} \circ \chi_i$. Set $\mu_t = \mu_{i,t}$ on A_i and $\mu_t = 0$ elsewhere. Define $\tilde{\mu}_t$ to be the pullback of μ_t under π_f , i.e. $\tilde{\mu}_t = \pi_f^* \mu_t$ on $\tilde{\mathcal{A}}_f$ and $\tilde{\mu}_t = 0$ elsewhere. Integrate μ_t with a choice of normalization (not depending on t) by fixing 3 distinct points in Post_f , we get a quasiconformal mapping ϕ_t .

Definition B.7. The path $f_t = \phi_t \circ f \circ \phi_t^{-1}$ ($t \geq 0$) is called a **pinching path** starting from f supported on \mathcal{A} .

The following result in [5] affirms that the pinching path is converging:

Theorem B.8. Let f be a geometrically finite rational map. Let $f_t = \phi_t \circ f \circ \phi_t^{-1}$ be a pinching path starting from f supported on a non-separating multi-annulus. Then the following properties hold:

1. f_t converges uniformly to a geometrically finite rational map g as $t \rightarrow \infty$.
2. ϕ_t converges uniformly to a continuous surjective map φ with $\varphi(\mathcal{J}_f) = \mathcal{J}_g$.
3. For each fill-in skeleton \hat{S} , $\varphi(\hat{S})$ is a parabolic (pre-)periodic point. Moreover φ is injective in the complement of the union of all fill-in skeletons.

C Admissible petals

Let $R : \mathbb{C} \rightarrow \mathbb{C}$ be a rational map with $R_a(0) = 0$, $R'_a(0) = e^{2\pi i \frac{p}{q}}$. Write the Taylor expansion near 0: $R^q(z) = z + \omega(a)z^{vq+1} + o(z^{vq+1})$. There exactly v cycles of immediate basins with rotation number p/q around 0. Let $B_0^*(0), \dots, B_{q-1}^*(0)$ be one of such cycle, written in cycle order. On every $B_m^*(0)$ there is a unique Fatou coordinate up to translation $\phi_k : B_m^*(0) \rightarrow \mathbb{C}$ semi-conjugating R to $z \mapsto z + 1$.

Definition C.1. An **admissible petal** $P^\gamma \subset B_m^*(0)$ is a petal such that the boundary of $\phi_k(P)$ is a smooth curve γ (well-defined up to translation) intersecting every horizontal line $y = b$ at exactly one point. We say that P^γ is **standard** if γ is a vertical line. P^γ is called **maximal** if ∂P^γ contains at least one critical point of $R^q|_{B_m^*(0)}$. Clearly once γ is chosen, the corresponding maximal petal P^γ is unique.

Lemma C.2. If $P^\gamma \subset B_m^*(0)$ is an admissible petal, then so is $R^n(P^\gamma)$ for any $n \geq 0$. Moreover $R^n(P^\gamma)$ are also associated to γ .

Let $P_m^0 := P \subset B_m^*(0)$ be a maximal admissible petal (we omit the index of associated curve γ). For any $0 \leq k \leq q-1$, $n \in \mathbb{Z}$ such that $np + k = m \pmod{q}$, P_k^n is defined to be the (unique) connected component of $R^{-n}(P_m^0)$ satisfying $P_k^n \subset B_k^*(0)$ and $0 \in \partial P_k^n$. See the sequence of mappings below, where the bar means \pmod{q} :

$$P_{m-np}^n \xrightarrow{R} \dots \xrightarrow{R} P_{m-p}^1 \xrightarrow{R} P_m^0 \xrightarrow{R} P_{m+p}^{-1} \xrightarrow{R} \dots \xrightarrow{R} P_{m+np}^{-n} \quad (34)$$

D Holomorphic dependence of Fatou coordinate

Let $R_a : \mathbb{C} \rightarrow \mathbb{C}$ be an analytic family (parametrized by $a \in \Lambda \subset \mathbb{C}$, Λ open) with $R_a(0) = 0$, $R'_a(0) = e^{2\pi i \frac{p}{q}}$. Write the Taylor expansion near 0: $R_a^q(z) = z + \omega(a)z^{vq+1} + o(z^{vq+1})$. In this section we always suppose that at some a_0 , $\omega(a_0) \neq 0$. Then for every attracting (resp. repelling) axis of R_{a_0} , there exists

- a small neighborhood D_{a_0} of a_0 ; a topological disk $V_{att} = \{x + iy; x > c - b|y|\}$ (resp. $V_{rep} = \{x + iy; x < c - b|y|\}$), where the constants b, c are positive and do not depend on a ; a family of attracting (resp. repelling) petals $(P_a)_{a \in B(a_0)}$
- a family of Fatou coordinates $(\phi_a)_{a \in D_{a_0}}$ of R_a such that $\phi_a : P_a \rightarrow V_{att}$ (resp. V_{rep}) is conformal and ϕ_a^{-1} is analytic in a .

An immediate consequence from this is that $h_a = \phi_a^{-1} \circ \phi_{a_0}$ (parametrized by $a \in U_{a_0}$) defines a dynamical holomorphic motion of P_{a_0} such that $h_a(P_{a_0}) = P_a$.

Next we prove that h_a can be extended to "sub-petal" arbitrarily close to the maximal petal Ω_{a_0} of R_{a_0} . Suppose that $\phi_{a_0} : \Omega_{a_0} \rightarrow \mathbb{H}_\rho$ is conformal, where $\mathbb{H}_\rho = \{z; \Re(z) > \rho\}$. For any $\delta > 0$, define $\Omega_{a_0}^\delta = \phi_{a_0}^{-1}(\mathbb{H}_{\rho+\delta})$. Notice that there exists $k \geq 1$ such that h_a is well-defined on $\tilde{P}_{a_0} := R_{a_0}^{kq}(\Omega_{a_0}^\delta)$. Set $\tilde{P}_a := h_a(\tilde{P}_{a_0})$.

Lemma D.1. *Let W_a be the connected component of $R_a^{-kq}(\tilde{P}_a)$ containing \tilde{P}_a . Then for a close enough to a_0 , $R_a^{kq} : W_a \rightarrow \tilde{P}_a$ is conformal (injective, surjective).*

Proof. Let $a_n \rightarrow a_0$. First we prove that

$$\forall \epsilon > 0, \exists N \text{ such that } \forall n \geq N, \overline{W_{a_n}} \subset_\epsilon \overline{\Omega_{a_0}^\delta}$$

where for two sets $A, B \subset \mathbb{C}$, $A \subset_\epsilon B$ means that A is contained in the ϵ -neighborhood of B . Suppose the contrary, then up to taking a subsequence we may suppose that

$$\exists \epsilon_0 > 0, \text{ a sequence } x_n \in \overline{W_{a_n}} \text{ with } x_n \rightarrow x_0, \text{ such that } \text{dist}(x_n, \overline{\Omega_{a_0}^\delta}) > \epsilon_0. \quad (35)$$

Clearly $R_{a_0}^{kq}(x_0) \in \overline{\tilde{P}_{a_0}}$ since $R_{a_n}^{kq}(x_n) \in \overline{\tilde{P}_{a_n}}$. For every n take a path $\gamma_n \subset \overline{W_{a_n}}$ connecting $0, x_n$ such that $\gamma \cap \tilde{P}_{a_0}$ does not depend on n . This is possible since $\overline{\tilde{P}_{a_0}}$ moves holomorphically. Moreover since the Hausdorff topology on the space of compact sets is sequentially compact, we may suppose that γ_n converges to a connected compact set γ_0 containing 0 and x_0 . This implies that $\text{dist}(x_0, \overline{\Omega_{a_0}^\delta}) = 0$, contradicting (35). Hence for a close enough to a_0 , W_a contains no critical points of R_a^{kq} and the lemma is proven. \square

Pull back h_a by $R_{a_0}^{kq}, R_a^{kq}$ and apply λ -Lemma we get immediately:

Corollary D.2. *For any $\delta > 0$, the holomorphic motion $h_a : \tilde{P}_{a_0} \rightarrow \tilde{P}_a$ can be dynamically extended to $\overline{\Omega_{a_0}^\delta}$ when a is close enough to a_0 .*

Proposition D.3. *Suppose for a in a neighborhood of a_0 , R_a^q has l (does not depend on a) distinct critical points $c_1(a), \dots, c_l(a)$ which vary analytically on a . Suppose moreover that the immediate basin of R_a is always simply connected. If $c_1(a_0)$ is the unique critical point on $\partial\Omega_{a_0}$, then for a close enough to a_0 , $c_1(a)$ is also the unique critical point on $\partial\Omega_{a_0}$.*

Proof. Clearly for a close enough to a_0 , $c_1(a)$ is in the immediate basin. Indeed, first take a path γ_{a_0} in the immediate basin, linking $c_1(a_0)$ and some z_{a_0} in a small petal P_{a_0} . One can connect it with $h_a(z_{a_0}) \in P_a = h_a(P_{a_0})$ by a compact path γ_a which is very close to γ_{a_0} . Hence for a very close to a_0 , γ_a is attracted by $z = 0$.

Hence we can normalise ϕ_a by taking $\phi_a(c_1(a)) = 0$. By Corollary D.2, $\phi_a^{-1} : \mathbb{H}_1 \rightarrow \tilde{\Omega}_a$ is conformal such that $R_a^q(c_1(a)) \in \partial\tilde{\Omega}_a$. Now we prove that for a close enough to a_0 , the component of $(R^q)_a^{-1}(\tilde{\Omega}_a)$ containing $\tilde{\Omega}_a$, denoted by Ω_a , contains no critical point of $R_a^q(\tilde{\Omega}_a)$. Suppose the contrary, then there is a sequence $a_n \rightarrow a_0$ such that, without loss of generality, $c_2(a_n) \in \Omega_{a_n}$. Up to taking a subsequence, $\overline{\Omega_{a_n}}$ has a limit Ω^* in the Hausdorff topology. Moreover Ω^* is compact, connected and Ω^* is the closure of the union of several components of $(R^q)_{a_0}^{-1}(\tilde{\Omega}_{a_0})$. Clearly $\overline{\Omega_{a_0}} \subset \Omega^*$. Suppose $c_1(a)$ has multiplicity m , then besides Ω_a , there are m distinct components of $(R_a^q)^{-1}(\tilde{\Omega}_a)$ attached at $c_1(a)$ since the basin is simply connected. Denote by $U_{a,1}, \dots, U_{a,m}$ these components. Then up to taking subsequences, $\overline{U_{a_n,i}}$ admits a limit U_i^* containing $\overline{U_{a_0,i}}$. Since $\Omega_{a_n} \cap U_{a_n,i} = \emptyset$, so $\Omega^* \cap U_i^* \subset \partial\Omega^*$, $\Omega^* = \Omega_{a_0}$. Hence $c_2(a_0) \in \partial\Omega_{a_0}$, a contradiction. \square

References

- [1] V. Timorin A. Blokh, L. Oversteegen. Slices of the parameter space of cubic polynomials. *Trans. Amer. Math. Soc.*, Number 8:5313–5359, 2022.
- [2] Artur Avila, Jeremy Kahn, Mikhail Lyubich, and Weixiao Shen. Combinatorial rigidity for unicritical polynomials. *Annals of Mathematics*, 170, 08 2005.
- [3] Xavier Buff, Jean Ecalte, and Adam Epstein. Limits of degenerate parabolic quadratic rational maps. *Geometric and Functional Analysis*, 23, 02 2013.
- [4] Arnaud Chéritat. On the size of siegel disks with fixed multiplier for cubic polynomials. <http://arxiv.org/abs/2003.13337>, 2020.
- [5] G. Cui and L. Tan. Hyperbolic-parabolic deformations of rational maps[j]. *SCIENCE CHINA-MATHEMATICS*, pages 61(12):2157–2220, 2018.
- [6] Adrien Douady and John Hamal Hubbard. On the dynamics of polynomial-like mappings. *Annales scientifiques de l'École Normale Supérieure*, Ser. 4, 18(2):287–343, 1985.
- [7] D. Faught. *Local connectivity in a family of cubic polynomials*. Ph. D. Thesis of Cornell University, 1992.
- [8] Lisa Goldberg. Fixed points of polynomial maps. I. Rotation subsets of the circles. *Annales scientifiques de l'École Normale Supérieure*, Ser. 4, 25(6):679–685, 1992.
- [9] Lisa Goldberg and John Willard Milnor. Fixed points of polynomial maps. Part II. Fixed point portraits. *Annales scientifiques de l'École Normale Supérieure*, Ser. 4, 26(1):51–98, 1993.
- [10] A. Douady, J. Hubbard. *Étude dynamique des polynômes complexes*. Publ. Math. d'Orsay, 1984.
- [11] Pierre Lavaurs. Systemes dynamiques holomorphes: explosion de points periodiques paraboliques. *PhD thesis, These de doctrat de l'Universite de Paris-Sud, Orsay, France*, 1989.
- [12] Tan Lei. *Local properties of the Mandelbrot set at parabolic points*, page 133–160. London Mathematical Society Lecture Note Series. Cambridge University Press, 2000.
- [13] L. Lomonaco. Parabolic-like mappings. *Ergodic Theory and Dynamical Systems*, 35(7):2171–2197, 2015.
- [14] Curtis T. McMullen and Dennis Sullivan. Quasiconformal homeomorphisms and dynamics iii. the teichmüller space of a holomorphic dynamical system. *Advances in Mathematics*, 135:351–395, 1998.
- [15] J. Milnor. Cubic polynomial maps with periodic critical orbit, part i. *Complex Dynamics: Families and Friends*, pages 333–412, 01 2009.

- [16] P. Roesch. Hyperbolic components of polynomials with a fixed critical point of maximal order. *Annales scientifiques de l'École Normale Supérieure*, Ser. 4, 40(6):901–949, 2007.
- [17] P. Roesch. *Cubic polynomials with a parabolic point*. Ergodic Theory and Dynamical Systems, 30(6), 1843–1867. doi:10.1017/S0143385709000820, 2010.
- [18] Pascale Roesch. *Holomorphic motions and puzzles (following M. Shishikura)*, page 117–132. London Mathematical Society Lecture Note Series. Cambridge University Press, 2000.
- [19] L. Tan and C. L. Petersen. *Branner-Hubbard Motions and attracting dynamics*. Dynamics on the Riemann sphere, 45–70, Eur. Math. Soc., Zurich, 2006.
- [20] Xiaoguang Wang. Hyperbolic components and cubic polynomials. *Advances in Mathematics*, 379:107554, 2021.
- [21] Tan Lei, Yin Yongcheng. *Local connectivity of the Julia set for geometrically finite rational maps*. Science in China (Serie A) 39, 39–47, 1996.
- [22] Saeed Zakeri. Dynamics of cubic siegel polynomials. *Communications in Mathematical Physics*, 206:185–233, 1999.
- [23] R. Zhang. Parabolic components in cubic polynomial slice $\text{Per}_1(1)$. <http://arxiv.org/abs/2210.14305>, 2022.

runze.zhang@math.univ-toulouse.fr