

# Frequency space derivation of linear and non-linear memory gravitational wave signals from eccentric binary orbits

Arpan Hait,<sup>1,\*</sup> Subhendra Mohanty,<sup>2,†</sup> and Suraj Prakash<sup>1,‡</sup>

<sup>1</sup>*Indian Institute of Technology Kanpur, Kalyanpur, Kanpur 208016, India*

<sup>2</sup>*Theoretical Physics Division, Physical Research Laboratory, Ahmedabad - 380009, India*

The memory effect in gravitational wave (GW) signals is the phenomenon, wherein the relative position of two inertial GW detectors undergoes a permanent displacement owing to the passage of GWs through them. Measurement of the memory signal is an important target for future observations as it establishes a connection between observations with field-theoretic results like the soft-graviton theorems. Theoretically, the memory signal is predicted at the leading order quadrupole formula for sources like binaries in hyperbolic orbits. This can be in the realm of observations by Advanced-LIGO, Einstein-Telescope, or LISA for stellar mass black-holes scattered by the super-massive black-hole at the galactic center. Apart from the direct memory component there is a non-linear memory signal in the secondary GW emitted from the primary GW chirp-signals emitted by coalescing binaries. In this paper, we compute the gravitational wave signals and their energy spectrum using the field-theoretic method by computing the scattering amplitudes for eccentric elliptical and hyperbolic binary orbits. The field theoretic calculation gives us the gravitational waveforms of linear and non-linear memory signals directly in the frequency space. The frequency domain templates are useful for extracting signals from the data. We compare our results with other calculations of linear and non-linear memory signals in literature and point out novel features we find in our calculations like the presence of  $\log(\omega)$  terms in the linear memory from hyperbolic orbits.

## I. INTRODUCTION

The memory effect in gravitational wave (GW) signals is the phenomenon, wherein the relative position of two inertial GW detectors undergoes a permanent displacement owing to the passage of GWs through them [1–3]. The memory effect is categorised broadly according to the nature of the source of GW. Linear memory arises from sources which have unbound components, like hyperbolic binary orbits [4, 5], neutrinos from supernova [6–12], gamma-ray bursts [13–15] and exotic objects like cosmic strings [16].

Another type of memory signal is the non-linear memory [17–20], where secondary gravitational waves are produced by the primary gravitational waves from sources like coalescing binaries. The importance of non-linear memory is that it will be an experimental proof of graviton-graviton coupling or the non-linear nature of the gravitational field equations in Einsteinian gravity.

The memory signal has a significance in field theory as it follows from the soft-graviton theorems [21–23] where the amplitude of a low energy graviton emission from a scattering process can be related, by a multiplicative kinematic factor, to the hard scattering amplitude without the graviton emission. The zero-graviton frequency amplitude of the soft-amplitude has a pole in the frequency space which in Fourier space is a step function in time which is the characteristic of the memory signal. Weinberg's soft theorem amplitude has been generalised to include higher order terms in graviton momenta using the gravitational gauge invariance and angular momentum conservation [24]. Calculations of graviton emission amplitudes show that there are non-analytic logarithmic terms in graviton frequencies even in tree level scattering [25–27].

---

\* arpan20@iitk.ac.in

† mohanty@prl.res.in

‡ surajprk@iitk.ac.in

The non-linear memory signal, for binaries in a quasi-circular orbit, already occurs at the 0-PN order [18, 28]. In [29, 30] the non-linear memory for the quasi-circular orbit was computed at the 3-PN (Post-Newtonian) order. The non-linear memory signal from eccentric binary orbits was calculated at 3-PN order in [31] and the 3-PN calculation for eccentric orbits including the tail contributions is done in [32].

Binaries (like the Hulse-Taylor) can have large initial eccentricities but by the time their frequencies enter the threshold of detectors like Advanced LIGO with a threshold of  $\sim 10$  Hz, they lose their eccentricities due to gravitational radiation reaction. However, there can be other initial configurations of binary star orbits which could give rise to large eccentricity orbits by the time the frequency enters the Advanced LIGO threshold [33, 34]. An eccentric binary can be formed by capture from an unbound orbit [35, 36]. An initial three body system may eject one of the bodies and result in an eccentric binary at coalescence [37]. A three body system in which the orbit of one body is at a larger radius, may cause the distant body to perturb the orbit of the tight binary system. This perturbation may give rise to Kozai-Lidov oscillations [38, 39] in the binary orbit which can drive the eccentricity to large values [40, 41]. Constructing gravitational wave templates for coalescent eccentric binaries is therefore of importance for observations [34] at future detectors like Advanced-LIGO. A determination of eccentricity of the orbits of the coalescent binary will be possible at Advanced-LIGO by improving the low frequency sensitivity [42]. Experimental determination of both the memory signal and the eccentricity of the orbit affect the signal near the low-frequency threshold of the detector, therefore construction of memory signal templates must accurately take into account the eccentricity of the orbit.

The non-linear gravitational wave memory from binary mergers may be discernible through next generation ground based detectors like Advanced LIGO [43] and Advanced Virgo [44] by improving the sensitivity in the 5-10 Hz band [45–48]. The memory effect for individual binaries may be resolvable by forthcoming interferometers such as LISA [49], Cosmic Explorer [50] and Einstein Telescope [51]. Gravitational-wave memory from binary mergers may also be seen in Advanced LIGO by combining the signals from multiple events [52]. The memory signal from several unresolved binary event mergers may be observed as a cumulative change over time in the pulsar timing residuals and may be observed in pulsar timing arrays (PTAs) [53–57]. Ground-based detectors are sensitive to the frequency of the GW in the  $5 - 10^3$  Hz band. Space-based detector LISA will probe frequencies as low as  $10^{-3}$  Hz, and pulsar timing arrays may measure frequency as low as  $10^{-9}$  Hz.

Black-holes in hyperbolic orbits in the gravitational field of super-massive black-holes at the galactic center radiate bursts of gravitational waves [58–64] which may be detectable at Advanced LIGO or Einstein Telescope [65, 66].

For hyperbolic orbits the nonlinear memory appears at the 2.5-PN level in the waveform. This is contrary to the case of elliptical and quasicircular orbits, where the nonlinear memory appears at the Newtonian (0-PN) order [18, 31]. This is because the radiation reaction effects accumulate over time in the closed orbits while for open orbits the radiation reaction is maximum at the closest approach and is zero at asymptotic past and future times.

The Bayesian inference of specific signals from the data is most efficiently achieved by match filtering of signals in the frequency domain with the data [69, 70]. With this aim, we compute the wave-forms for the memory signals for eccentric elliptical and hyperbolic orbits in the frequency domain. We compute the waveforms in using the tree level graviton emission amplitude with the stress tensor of the binary orbits as sources. Using this graviton emission amplitude we compute the frequency spectrum of the energy radiated following [71–74]. For elliptical orbits, the energy radiation spectrum is the source of the non-linear gravitational memory which we thus obtain directly in the frequency space.

The anatomy of this article can be described as follows: In section II, we outline how the gravitational waveform can be constructed in a field-theoretic approach and we also highlight the computation of the rate of energy loss associated with gravitational wave radiation. In section III, we compute components of the stress tensor, in frequency domain, for binaries in hyperbolic orbits. This is followed by the evaluation of the same quantities in the limit of vanishing frequency and the associated linear memory in section IV. Next, we elucidate the general formalism for the non-linear memory associated with the radiation from binaries in closed orbits in section V. We delve into the details for the special case of elliptical orbits in section VI, where we compute the transverse-traceless memory waveform. We summarize our conclusions in section VII.

## II. GRAVITATIONAL WAVEFORM AND ENERGY RADIATED FROM SCATTERING AMPLITUDES

The probability amplitude of emitting a graviton of polarisation  $\epsilon_{\mu\nu}^\lambda(\vec{n})$  from a source with stress tensor (in the momentum space)  $\tilde{T}^{\mu\nu}(k)$  is given by

$$\mathcal{A}_\lambda(k_0, \vec{n}; k_0) = -i \frac{\kappa}{2} \epsilon_{\mu\nu}^{*\lambda}(\vec{n}) \tilde{T}^{\mu\nu}(k_0, \vec{n}; k_0). \quad (1)$$

We can express the gravitational wave metric observed at the detector in terms of the probability amplitude of a graviton emission by a source at a distance  $r$  as,

$$h_{\alpha\beta}(\vec{x}, t) = \frac{1}{4\pi r} \int \frac{dk_0}{(2\pi)} \sum_{\lambda=1}^2 \epsilon_{\alpha\beta}^\lambda(\vec{n}) \mathcal{A}_\lambda(k_0, \vec{n}; k_0) e^{-ik_0(t-r)}. \quad (2)$$

The graviton field in Eq. (2) is a canonical spin-2 field with mass dimension 1 as it is defined as an expansion of the metric  $g_{\mu\nu} = \eta_{\mu\nu} + \kappa h_{\mu\nu}$  where  $\kappa = \sqrt{32\pi G}$ . The metric perturbation identified as gravitational wave is the dimensionless quantity  $\tilde{h}_{\mu\nu} \equiv g_{\mu\nu} - \eta_{\mu\nu} = \kappa h_{\mu\nu}$ . The expression for the dimensionless gravitational wave in terms of the amplitude is therefore from Eq. (2) given by

$$\tilde{h}_{\alpha\beta}(\vec{x}, t) = \frac{\kappa}{4\pi r} \int \frac{dk_0}{(2\pi)} \sum_{\lambda=1}^2 \epsilon_{\alpha\beta}^\lambda(\vec{n}) \mathcal{A}_\lambda(k_0, \vec{n}; k_0) e^{-ik_0(t-r)}. \quad (3)$$

This relates the waveform at the detector to the probability amplitude of graviton emission by the source. To relate the waveform at the detector to the source stress tensor we substitute in Eq. (3), the expression for the amplitude given in Eq. (1), to obtain

$$\begin{aligned} \tilde{h}_{\alpha\beta}(\vec{x}, t) &= -\frac{\kappa^2}{8\pi r} \int \frac{dk_0}{2\pi} \sum_{\lambda=1}^2 \epsilon_{\alpha\beta}^\lambda(\vec{n}) \epsilon_{\mu\nu}^{*\lambda}(\vec{n}) \tilde{T}^{\mu\nu}(k_0, \vec{n}; k_0) e^{-ik_0(t-r)} \\ &= -\frac{4G}{r} \int \frac{dk_0}{2\pi} \left( \tilde{T}_{\alpha\beta}(k_0, \vec{n}; k_0) - \frac{1}{2} \eta_{\alpha\beta} \tilde{T}^\mu{}_\mu(k_0, \vec{n}; k_0) \right) e^{-ik_0(t-r)}. \end{aligned} \quad (4)$$

where we made use of the completeness relation

$$\sum_{\lambda=1}^2 \epsilon_{\mu\nu}^\lambda(k) \epsilon_{\alpha\beta}^{*\lambda}(k) = \frac{1}{2} (\eta_{\mu\alpha} \eta_{\nu\beta} + \eta_{\mu\beta} \eta_{\nu\alpha}) - \frac{1}{2} \eta_{\mu\nu} \eta_{\alpha\beta}. \quad (5)$$

To obtain the propagating degrees of freedom, we need to project the transverse traceless (TT) components of the wavefunction constructed in Eq. (4).

$$\left[ \tilde{h}_{ij} \right]^{\text{TT}}(\vec{x}, t) = -\frac{4G}{r} \Lambda_{ij,kl}(\vec{n}) \int \frac{dk_0}{2\pi} \left( \tilde{T}_{kl}(k_0, \vec{n}; k_0) - \frac{1}{2} \eta_{kl} \tilde{T}^\mu{}_\mu(k_0, \vec{n}; k_0) \right) e^{-ik_0(t-r)}. \quad (6)$$

where,  $\Lambda_{ij,kl}(\vec{n})$  is the transverse-traceless projection operator defined with respect to the direction,  $\hat{n}$ , of the emitted gravitational wave. The explicit form of the TT projection operator is

$$\Lambda_{ij,kl}(\hat{n}) = P_{ik}(\hat{n}) P_{jl}(\hat{n}) - \frac{1}{2} P_{ij}(\hat{n}) P_{kl}(\hat{n}) = (\delta_{ik} - n_i n_k) (\delta_{jl} - n_j n_l) - \frac{1}{2} (\delta_{ij} - n_i n_j) (\delta_{kl} - n_k n_l). \quad (7)$$

Since  $\Lambda_{ij,kl} \eta_{kl} = 0$ , the  $\tilde{T}^\mu{}_\mu$  term vanishes and we obtain the simpler result,

$$\left[ \tilde{h}_{ij} \right]^{\text{TT}}(\vec{x}, t) = -\frac{4G}{r} \Lambda_{ij,kl}(\vec{n}) \int \frac{dk_0}{2\pi} T_{kl}(k_0, \vec{n}; k_0) e^{-ik_0(t-r)}. \quad (8)$$

In frequency space, the observed gravitational wave and the stress-tensor of the source can be related as,

$$\left[\tilde{h}_{ij}\right]^{\text{TT}}(\vec{x}, k_0) = -\frac{4G}{r} \Lambda_{ij,kl}(\vec{n}) T_{kl}(k_0, \vec{n} k_0). \quad (9)$$

We shall use Eq. (9) to compute the gravitational waveform from various sources, like compact binaries in bound and unbound orbits, by computing the stress-tensor of the source in frequency space.

### A. Power spectrum of gravitational wave in field theoretic approach

The rate of graviton emission is given by the Fermi Golden Rule, and is the amplitude squared summed over the final state graviton polarisation and integrated over the phase space volume,

$$\begin{aligned} \Gamma &= \sum_{\lambda} \int \frac{|S_{fi}|^2}{T} \frac{d^3\vec{k}}{(2\pi)^3 2\omega} = \sum_n \sum_{\lambda} \int |\mathcal{A}_{\lambda}(\omega, \omega'_n)|^2 (2\pi) \delta(\omega - \omega'_n) \frac{d^3\vec{k}}{(2\pi)^3 2\omega} \\ &= \frac{\kappa^2}{4} \sum_n \sum_{\lambda} \int |T_{\mu\nu}(\vec{k}, \omega'_n) \epsilon_{\lambda}^{*\mu\nu}(\vec{k})|^2 (2\pi) \delta(\omega - \omega'_n) \frac{d^3\vec{k}}{(2\pi)^3 2\omega_k}. \end{aligned} \quad (10)$$

Here,  $\lambda$  accounts for the possible states of polarization and  $n$  denotes the harmonics corresponding to the emission. The rate of energy radiated is obtained from the probability of radiation given above by including an extra factor of  $\omega_k = |\vec{k}|$  in the integral,

$$\frac{dE_{\text{gw}}}{dt} = \frac{\kappa^2}{4} \sum_n \sum_{\lambda} \int |T_{\mu\nu}(\vec{k}, \omega'_n) \epsilon_{\lambda}^{*\mu\nu}(\vec{k})|^2 (2\pi) \delta(\omega - \omega'_n) \omega_k \frac{d^3\vec{k}}{(2\pi)^3 2\omega}. \quad (11)$$

We can open the modulus squared piece of the integrand in Eq. (11) and simplify using the polarisation sum relation, see Eq. (5), as follows:

$$\begin{aligned} \sum_{\lambda} |T_{\mu\nu}(\vec{k}, \omega'_n) \epsilon_{\lambda}^{*\mu\nu}(\vec{k})|^2 &= \sum_{\lambda} \left( T_{\mu\nu}(\vec{k}, \omega'_n) T_{\alpha\beta}^*(\vec{k}, \omega'_n) \right) \left( \epsilon_{\lambda}^{*\mu\nu}(\vec{k}) \epsilon_{\lambda}^{\alpha\beta}(\vec{k}) \right) \\ &= T_{\mu\nu}(\vec{k}, \omega'_n) T^{*\nu\mu}(\vec{k}, \omega'_n) - \frac{1}{2} |T^{\mu}_{\mu}(\vec{k}, \omega'_n)|^2. \end{aligned} \quad (12)$$

The  $T^{00}$  and  $T^{i0}$  components can be expressed in terms of the  $T^{ij}$  ones by utilizing the conserved current relation,  $k_{\mu} T^{\mu\nu} = 0$ . This allows us to write

$$T_{0j} = -\hat{k}^i T_{ij}, \quad T_{00} = \hat{k}^i \hat{k}^j T_{ij}. \quad (13)$$

Using these relations, we can rewrite Eq. (12) as

$$\begin{aligned} |T_{\mu\nu}(\vec{k}, \omega'_n)|^2 - \frac{1}{2} |T^{\mu}_{\mu}(\vec{k}, \omega'_n)|^2 &= T_{ij} T^{*ji} + T_{00} T^{*00} + T_{0i} T^{*i0} + T_{i0} T^{*0i} - \frac{1}{2} (T^0_0 + T^i_i) (T^{*0}_0 + T^{*j}_j) \\ &= \left( T_{ij} T^{*ji} - \frac{1}{2} T^i_i T^{*j}_j \right) + \frac{1}{2} \hat{k}^i \hat{k}^j \hat{k}^l \hat{k}^m T_{ij} T_{lm}^* - \left( \hat{k}^l \hat{k}^m T_{il} T_{mi}^* + \hat{k}^l \hat{k}^m T_{il} T_{mi}^* \right) \\ &\quad + \frac{1}{2} \left( \hat{k}^l \hat{k}^m T_{lm}^* T^i_i + \hat{k}^l \hat{k}^m T_{lm} T^{*j}_j \right). \end{aligned} \quad (14)$$

In the quadrupole approximation of the source, for sources smaller in size than the wavelength of the GWs,  $\vec{k} \cdot \vec{x} \ll 1$ , the stress tensor in momentum space  $T_{\mu\nu}(\vec{k}, \omega'_n)$  has no explicit  $\vec{k}$  dependence. Therefore, after substituting the contents of Eq. (14) in Eq. (11), one can perform the angular integrations using the following relations:

$$\int d\Omega_k = 4\pi, \quad \int d\Omega_k \hat{k}^i \hat{k}^j = \frac{4\pi}{3} \delta_{ij}, \quad \int d\Omega_k \hat{k}^i \hat{k}^j \hat{k}^l \hat{k}^m = \frac{4\pi}{15} (\delta_{ij} \delta_{lm} + \delta_{il} \delta_{jm} + \delta_{im} \delta_{jl}). \quad (15)$$

to obtain

$$\int d\Omega_k \left[ |T_{\mu\nu}(\vec{k}, \omega'_n)|^2 - \frac{1}{2} |T^\mu{}_\mu(\vec{k}, \omega'_n)|^2 \right] = \frac{8\pi}{5} \left( T_{ij}(\omega'_n) T_{ji}^*(\omega'_n) - \frac{1}{3} |T^i{}_i(\omega'_n)|^2 \right), \quad (16)$$

Finally, using the result of Eq. (16), the expression for the energy radiated by a source in terms of the source stress tensor can be obtained as a modification of Eq. (11) as

$$\frac{dE_{\text{gw}}}{dt} = \frac{\kappa^2}{4} \sum_n \int \frac{8\pi}{5} \left( T_{ij}(\omega'_n) T_{ji}^*(\omega'_n) - \frac{1}{3} |T^i{}_i(\omega'_n)|^2 \right) \omega^2 2\pi \delta(\omega'_n - \omega) \frac{d\omega}{(2\pi)^3 2\omega}. \quad (17)$$

We make use of this expression for computing the energy radiated by binaries in elliptical orbits which is the source of the non-linear memory signal. We will also show using this expression that in the memory signal from hyperbolic orbits, there is non-zero energy radiated around the zero-frequency band.

### III. HYPERBOLIC ORBITS

For the case of unbounded orbits, we consider a black-hole of mass  $m_2$  in a hyperbolic orbit around a larger black-hole of mass  $m_1$ . Similar to the case of the elliptical orbits, in the centre of mass frame, the motion is equivalently described by a single body having the reduced mass  $\mu = \frac{m_1 m_2}{m_1 + m_2}$ . The following quantities describe the system:

- The coordinates parametrized as,

$$x(\xi) = a(e - \cosh \xi), \quad y(\xi) = b \sinh \xi, \quad z(\xi) = 0, \quad \frac{\omega'}{\nu} t = \Omega t = (e \sinh \xi - \xi), \quad (18)$$

where,  $a$  and  $b$  ( $= a\sqrt{e^2 - 1}$ ,  $e > 1$ ) denote the semi-major and semi-minor axes and the variables  $e$  and  $\xi \in (-\infty, \infty)$  refer to the eccentricity and the hyperbolic anomaly of the orbit respectively.

- The angular frequency  $\omega'$  is proportional to the fundamental frequency  $\Omega$ , as  $\omega' = \nu \Omega$ , where  $\nu \in (0, \infty)$  is a non negative real number and  $\Omega = \left( \frac{G(m_1 + m_2)}{a^3} \right)^{1/2}$ .

We start with the computation of the stress-tensor components in frequency space. Focussing first on the  $xx$  component, the calculation proceeds as follows:

$$T_{xx}(\omega') = \int_{-\infty}^{\infty} dt e^{i\omega' t} \mu \dot{x}^2 = -\iota \mu \omega' \int_{-\infty}^{\infty} dt e^{i\omega' t} x \dot{x} = -\iota \mu \omega' \int_{-\infty}^{\infty} d\xi e^{i\nu(e \sinh \xi - \xi)} x \frac{dx}{d\xi}, \quad (19)$$

where, the second equality is obtained after implementing integration by parts and neglecting a term proportional to  $\dot{x}(t)$ . The final expression is obtained after incorporating a change of variables  $t \rightarrow \xi$ . Since, the integration is over  $\xi$ , the result will simply be a function of  $\nu$ . Also, since  $\omega' = \nu \Omega$ , it makes more sense to express  $T_{ij}$  as a function of  $\nu$

instead of  $\omega'$ . Thus, we obtain

$$\begin{aligned} T_{xx}(\nu) &= \iota \mu \nu \Omega a^2 \int_{-\infty}^{\infty} d\xi e^{\iota\nu(e \sinh \xi - \xi)} \sinh \xi (e - \cosh \xi) \\ &= \mu \nu \Omega a^2 \pi \left[ \frac{\iota}{\nu e^2} H_{\nu}^{(1)}(\iota e \nu) - \left( e - \frac{1}{e} \right) H_{\nu}^{(1)'}(\iota e \nu) \right]. \end{aligned} \quad (20)$$

We replaced  $\sinh \xi$ ,  $\cosh \xi$  by their exponential counterparts. Subsequently, we identified and replaced the integrals with Hankel functions, i.e.,

$$H_p^{(1)}(q) = \frac{1}{\iota \pi} \int_{-\infty}^{\infty} d\xi e^{q \sinh \xi - p \xi}. \quad (21)$$

Then, we utilized the following recurrence relations for Hankel functions to simplify the expression:

$$H_{p-1}^{(1)}(q) + H_{p+1}^{(1)}(q) = \frac{2p}{q} H_p^{(1)}(q) \quad \text{and} \quad H_{p-1}^{(1)}(q) - H_{p+1}^{(1)}(q) = 2 H_p^{(1)'}(q). \quad (22)$$

The remaining non-zero components can similarly be evaluated as:

$$\begin{aligned} T_{yy}(\omega') &= \int_{-\infty}^{\infty} dt e^{\iota\omega' t} \mu \dot{y}^2 = -\iota \mu \omega' \int_{-\infty}^{\infty} dt e^{\iota\omega' t} y \dot{y} = -\iota \mu \omega' \int_{-\infty}^{\infty} d\xi e^{\iota\nu(e \sinh \xi - \xi)} y \frac{dy}{d\xi} \\ T_{yy}(\nu) &= \mu \nu \Omega a^2 (e^2 - 1) \pi \left[ \frac{\iota}{\nu e^2} H_{\nu}^{(1)}(\iota e \nu) + \frac{1}{e} H_{\nu}^{(1)'}(\iota e \nu) \right], \\ T_{xy}(\omega') &= \int_{-\infty}^{\infty} dt e^{\iota\omega' t} \mu \dot{x} \dot{y} = -\iota \mu \omega' \int_{-\infty}^{\infty} dt e^{\iota\omega' t} y \dot{x} = -\iota \mu \omega' \int_{-\infty}^{\infty} d\xi e^{\iota\nu(e \sinh \xi - \xi)} x \frac{dx}{d\xi} \\ T_{xy}(\nu) &= -\mu \nu \Omega a^2 \sqrt{e^2 - 1} \pi \left[ \left( \frac{1}{e^2} - 1 \right) H_{\nu}^{(1)}(\iota e \nu) + \frac{\iota}{\nu e} H_{\nu}^{(1)'}(\iota e \nu) \right]. \end{aligned} \quad (23)$$

### A. Rate of emission of gravitational wave energy

Using the expressions for  $T_{xx}$ ,  $T_{yy}$  and  $T_{xy}$  derived in Eqs. (20) and (23) we obtain,

$$T_{ij}(\nu) T_{ji}^*(\nu) - \frac{1}{3} |T^i_i(\nu)|^2 = \mu^2 \nu^2 \Omega^2 a^4 \pi^2 \left[ f_1(\nu, e) \left( H_{\nu}^{(1)}(\iota e \nu) \right)^2 + f_2(\nu, e) \left( H_{\nu}^{(1)'}(\iota e \nu) \right)^2 \right]. \quad (24)$$

Here,

$$f_1(\nu, e) = \frac{2}{e^4} (e^2 - 1)^3 + \frac{6 - 6e^2 + 2e^4}{3\nu^2 e^4} \quad \text{and} \quad f_2(\nu, e) = 2 \left( \frac{e^2 - 1}{e} \right) \left( \frac{1}{\nu^2 e} + \frac{e^2 - 1}{e} \right). \quad (25)$$

After evaluating the integral in Eq. (17), we get:

$$P(\nu) = \frac{dE_{\text{gw}}}{dt} = \frac{\kappa^2}{20} \mu^2 \Omega^4 \nu^4 a^4 \pi \underbrace{\left[ f_1(\nu, e) \left( H_{\nu}^{(1)}(\iota e \nu) \right)^2 + f_2(\nu, e) \left( H_{\nu}^{(1)'}(\iota e \nu) \right)^2 \right]}_{f(\nu, e)}, \quad (26)$$

where  $f_{1,2}(\nu, e)$  are given in Eq. (25). A plot depicting the features of  $P(\nu)$  for different values of eccentricity has

been shown in Fig. 1. The total energy loss can be obtained from the power spectrum and is given by:

$$\Delta E = \int_{-\infty}^{\infty} dt P(t) = \frac{1}{\pi} \int_0^{\infty} d\nu P(\nu). \quad (27)$$

Explicit integration over products of Hankel functions is difficult, especially when the order  $p$  of the function  $H_p^{(1)}(q)$  also depends on the integration variable. Therefore, we have presented an estimate by doing a numerical integration with various choices of  $\nu_{\max}$  (the upper limit of the integral). The results for different eccentricities have been plotted in Fig. 2. We notice that, in each case, the curves become parallel to the  $x$ -axis and approach a constant, for fairly small values of  $\nu_{\max}$ .




FIG. 1. The power spectrum, in frequency space, of gravitational wave radiation from binaries in hyperbolic orbits with varying eccentricities.




FIG. 2. Numerical estimate of the total energy loss due to gravitational wave radiation from binaries in hyperbolic orbits. Individual curves correspond to varying eccentricities. The horizontal axis corresponds to the upper limit in the integral  $\Delta E = \frac{1}{\pi} \int_0^{\nu_{\max}} d\nu P(\nu)$ . It is evident that even for small enough values of  $\nu_{\max}$ , the energy loss attains a constant value. This suggests that with a large  $\nu_{\max}$  the numerical estimate is a good approximation.

#### IV. LINEAR MEMORY FROM HYPERBOLIC ORBITS

##### A. Power emitted at zero frequency

Computing the zero frequency limits of the stress tensors is equivalent to obtaining approximations for them in the  $\nu \rightarrow 0$  limit. As  $\nu e \rightarrow 0$ , the Hankel function and its first derivative assume the following form:

$$H_{\nu\nu'}^{(1)}(\nu e\nu) \simeq \frac{2\nu}{\pi} \ln(\nu e), \quad H_{\nu\nu'}^{(1)'}(\nu e\nu) \simeq \frac{2}{\pi\nu e}. \quad (28)$$

Substituting the above in the expressions for  $T_{xx}$ ,  $T_{yy}$  and  $T_{xy}$ , in Eqs. (20) and (23) yields,

$$\begin{aligned} T_{xx} &= -\frac{2\mu a^2 \Omega}{e^2} \left[ \ln(\nu e) + (e^2 - 1) \right], & T_{yy} &= -\frac{2\mu a^2 \Omega}{e^2} (e^2 - 1) \left[ \ln(\nu e) - 1 \right], \\ T_{xy} &= 2\mu\nu\Omega a^2 \sqrt{e^2 - 1} \left[ \frac{(e^2 - 1)}{e^2} \ln(\nu e) - \frac{1}{\nu^2 e^2} \right] \end{aligned} \quad (29)$$

Using these, we can rewrite,

$$T_{ij}(\nu) T_{ji}^*(\nu) - \frac{1}{3} |T^i{}_i(\nu)|^2 = 4\mu^2 \nu^2 a^4 \left[ \tilde{f}_1(\nu, e) (\ln(\nu e))^2 + \tilde{f}_2(\nu, e) \ln(\nu e) + \tilde{f}_3(\nu, e) \right]. \quad (30)$$

with

$$\begin{aligned} \tilde{f}_1(\nu, e) &= \frac{2}{e^4} (e^2 - 1)^3 + \frac{6 - 6e^2 + 2e^4}{3\nu^2 e^4}, & \tilde{f}_2(\nu, e) &= \frac{2(e^2 - 1)}{\nu^2 e^4} - \frac{6(e^2 - 1)^2}{\nu^2 e^4}, \\ \tilde{f}_3(\nu, e) &= \frac{2(e^2 - 1)}{\nu^4 e^4} + \frac{2(e^2 - 1)^2}{\nu^2 e^4}. \end{aligned} \quad (31)$$

The expression for the power spectrum of the gravitational wave, see Eq. (26), now becomes,

$$P(\nu) = \frac{dE}{dt} = \frac{\kappa^2}{5} \pi \mu^2 a^4 \Omega^4 \nu^4 f(\nu, e). \quad (32)$$

In the limit  $\nu \rightarrow 0$ ,

$$\lim_{\nu \rightarrow 0} \nu^4 f(\nu, e) = \frac{2(e^2 - 1)}{\pi^2 e^4}, \quad (33)$$

which is finite and different from zero, except for  $e = 1$  and  $e \rightarrow \infty$ . Then the power radiated by the GW of zero frequency is given by,

$$P(\nu = 0) = \frac{64G}{5} \mu^2 a^4 \Omega^4 \frac{(e^2 - 1)}{e^4}. \quad (34)$$

This result agrees with the expression derived in [61].

##### B. The memory waveform

The gravitational wave amplitude corresponding to polarisation  $\lambda$ , measured by a detector located at a distance  $r$ , is given (in the frequency domain) by

$$h_\lambda(\omega', r) = \frac{4G}{r} \epsilon_\lambda^{ij}(\vec{n}) T_{ij}(n, \omega') \quad (35)$$



The waveforms for the + and  $\times$  polarisations are

$$\begin{aligned}
h_+(\omega', r) &= \iota \frac{4G}{r} \epsilon_+^{ij}(\vec{n}) T_{ij}(\vec{n}, \omega') = \iota \frac{4G}{r} (\vec{e}_{\theta i} \vec{e}_{\theta j} - \vec{e}_{\phi i} \vec{e}_{\phi j}) T_{ij}(\vec{n}, \omega') \\
&= \iota \frac{4G}{r} \left( T_{xx} (\cos^2 \phi - \sin^2 \phi \cos^2 \theta) + T_{yy} (\sin^2 \phi - \cos^2 \phi \cos^2 \theta) - T_{zz} \sin^2 \theta \right. \\
&\quad \left. - T_{xy} \sin 2\phi (1 + \cos^2 \theta) + T_{xz} \sin \phi \sin 2\theta + T_{yz} \cos \phi \sin 2\theta \right), \\
h_\times(\omega', r) &= \iota \frac{4G}{r} \epsilon_\times^{ij}(\vec{n}) T_{ij}(\vec{n}, \omega') = \iota \frac{4G}{r} (\vec{e}_{\theta i} (\vec{e}_\phi)_j + (\vec{e}_\phi)_i (\vec{e}_\theta)_j) T_{ij}(\vec{n}, \omega') \\
&= \iota \frac{4G}{r} \left( (T_{xx} - T_{yy}) \sin 2\phi \cos \theta + 2T_{xy} \cos 2\phi \cos \theta \right) - 2T_{xz} \cos \phi \sin \theta + 2T_{yz} \sin \phi \sin \theta. \tag{36}
\end{aligned}$$

We have used the spherical coordinates to describe the polarisation of a GW traversing in the radial  $\vec{n}$  direction. The time domain waveforms can be obtained after evaluating the Fourier transforms of  $h_+(\omega', r)$  and  $h_\times(\omega', r)$ . This can be accomplished numerically and the results have been presented in Fig. 3




FIG. 3. Plots displaying the characteristics of  $h_+(t)$  and  $h_\times(t)$  for different eccentricities. Here,  $\phi = \theta = 0$ .

### C. Comparison with soft-theorems

According to the soft-graviton theorems, the amplitude of a graviton emission from the scattering of  $n$ -particles of momenta  $p_a$  can be written as the product of a kinematical factor and the amplitude  $\mathcal{A}_n(p_a)$  of the  $n$ -particle scattering as [24, 26],

$$\mathcal{A}_{n+1}(p_a, q) = \frac{\kappa}{2} \epsilon_{\lambda\mu\nu}^* \sum_{a=1}^n \left( \frac{p_{a\mu} p_{a\nu}}{p_a \cdot q} - i \frac{p_a^\mu q_\beta J_a^{\beta\nu}}{p_a \cdot q} - \frac{q_\alpha q_\beta J_a^{\alpha\mu} J_a^{\beta\nu}}{p_a \cdot q} \right) \mathcal{A}_n(p_a). \tag{37}$$

Here,  $J_a^{\alpha\beta} = x_a^\alpha p_a^\beta - x_a^\beta p_a^\alpha + S_a^{\alpha\beta}$  described the total angular momentum of particle "a". The series of the soft factors are at the same order in the gravitational coupling but in increasing powers of the graviton frequency  $q_0 = \omega$ . The leading term goes as  $\omega^{-1}$  while the sub-leading terms go as  $\sim \omega^0$  and  $\sim \omega^1$  respectively. The gravitational waveform at distance  $r$  of the radiated soft graviton of momentum  $q$  is given by

$$h_{\mu\nu}(r, q) = \frac{4G}{r} \left( \frac{p_{a\mu} p_{a\nu}}{p_a \cdot q} - i \frac{p_a^\mu q_\beta J_a^{\beta\nu}}{p_a \cdot q} - \frac{q_\alpha q_\beta J_a^{\alpha\mu} J_a^{\beta\nu}}{p_a \cdot q} \right). \tag{38}$$

There are logarithmic corrections to the leading order terms suppressed by  $G$  which give a tail contribution to the linear memory signal (which as a function of time goes as  $1/t$ ), [25, 26]. The low frequency graviton signal from a generic hard scattering can be written as [25],

$$h_{ij}(\omega) = i\omega^{-1}A_{ij} + B_{ij} \log \omega^{-1} + \dots \quad (39)$$

where  $A_{ij} \sim \mathcal{O}(G/r)$  and  $B_{ij} \sim \mathcal{O}(G^2/r)$ .

In this section we computed the linear memory for hyperbolic eccentric orbits and it is evident from (29) and (36) that there are logarithmic contributions at  $\mathcal{O}(G/r)$  also.

## V. NON-LINEAR MEMORY

The non-linear memory is due to the secondary gravitational waves which are emitted by the primary gravitational waves from an oscillating source, such as, a coalescing binary [3, 17, 18, 29, 30, 78].

The stress tensor of gravitational waves is related to the energy radiated as

$$\tau_{ij}^{\text{gw}} = \frac{dE_{\text{gw}}}{dt d\Omega} n_i n_j, \quad (40)$$

and the non-linear memory waveform is given by [3, 17, 18],

$$h_{ij}(t, r \hat{n}) = \frac{4G}{r} \int_{-\infty}^{t-r} dt' d\Omega' \frac{dE_{\text{gw}}}{dt' d\Omega'} \left[ \frac{n'_i n'_j}{1 - \vec{n}' \cdot \vec{n}} \right]^{\text{TT}}. \quad (41)$$

Here  $\vec{n}'$  is the unit vector from the source to the solid angle denoted by  $d\Omega'$  and  $\vec{n}$  is the unit vector along the line of sight from the source to the detector.

We outline the derivation of these results in this section. The source term of the non-linear gravitational waves is the Landau-Lifshitz stress tensor of the primary gravitational waves given by

$$\tau_{ij}^{\text{gw}} = \frac{1}{32\pi G} \langle \partial_i h^{ab} \partial_j h_{ab} \rangle, \quad (42)$$

where the angular brackets encode averaging over time longer than the time period and volumes larger than the wavelengths of the source gravitational waves  $h_{ab}$ . The source gravitational waves travel outward radially with speed of light and are functions of  $(t - r)$ , i.e  $h_{ab}(t, \vec{x}) = h_{ab}(t - r, \Omega)$ . This implies we can relate their spatial and time derivatives as,  $\partial_i h_{ab}(t - r) = -n_i \partial_0 h_{ab}(t - r)$ , where  $n_i = x_i/r$ . Therefore we can write

$$\tau_{ij}^{\text{gw}} = \frac{1}{32\pi G} \langle \partial_i h^{ab} \partial_j h_{ab} \rangle = n_i n_j \tau_{00}^{\text{gw}}. \quad (43)$$

We can model the energy density  $\tau_{00}^{\text{gw}}$  produced from a source and propagating radially outwards on null-rays as

$$\tau_{00}^{\text{gw}}(t, \vec{x}) = \frac{1}{r^2} \frac{dE_{\text{gw}}(t - r, \Omega)}{dt d\Omega}. \quad (44)$$

where  $dE_{\text{gw}}/dt$  is the luminosity of the source in gravitational waves and  $dE_{\text{gw}}/d\Omega$  denotes the angular distribution of the source luminosity. Therefore we can write  $\tau_{ij}^{\text{gw}}$  in terms of the energy flux as

$$\tau_{ij}^{\text{gw}} = n_i n_j \tau_{00}^{\text{gw}} = n_i n_j \frac{1}{r^2} \frac{dE_{\text{gw}}(t - r, \Omega)}{dt d\Omega}. \quad (45)$$

The secondary gravitational waves sourced by the gravitational wave stress tensor will obey the in-homogenous wave equation:

$$\square h_{ij} = -16\pi G \tau_{ij}^{\text{gw}}. \quad (46)$$

whose solution assumes the following form,

$$h_{ij}^{\text{mem}}(t, \vec{x}) = 4G \int dt' d^3x' \tau_{ij}^{\text{gw}}(t', \vec{x}') \frac{\delta(t' - (t - |\vec{x} - \vec{x}'|))}{|\vec{x} - \vec{x}'|}. \quad (47)$$

We can express the source term, Eq. (43), in terms of the null coordinate  $u = t' - r'$  as follows

$$\tau_{ij}^{\text{gw}}(t', \vec{x}') = \frac{n'_i n'_j}{r'^2} \frac{dE_{\text{gw}}(t' - r', \Omega')}{dt' d\Omega'} = \int du \frac{n'_i n'_j}{r'^2} \delta(u - (t' - r')) \frac{dE_{\text{gw}}(u, \Omega)}{dt' d\Omega'}. \quad (48)$$

Substituting the above in Eq. (47) we obtain

$$h_{ij}^{\text{mem}}(t, \vec{x}) = 4G \int du dt' dr' r'^2 d\Omega' \frac{n'_i n'_j}{r'^2} \frac{\delta(t' - (t - |\vec{x} - \vec{x}'|))}{|\vec{x} - \vec{x}'|} \delta(u - (t' - r')) \frac{dE_{\text{gw}}(u, \Omega)}{dt' d\Omega'}. \quad (49)$$

Since the distance to the observer is much larger than the source size,  $r \gg r'$ , we take the approximations

$$\frac{1}{|\vec{x} - \vec{x}'|} \simeq \frac{1}{r(1 - \vec{n}' \cdot \vec{n})}, \quad \delta(t' - (t - |\vec{x} - \vec{x}'|)) \simeq \delta(t' - (t - r)). \quad (50)$$

Now we can perform the integral over  $r'$  using the second delta function in Eq. (49) and then do the  $t'$  integration using the remaining delta function to obtain

$$h_{ij}^{\text{mem}}(t, \vec{x}) = \frac{4G}{r} \int_{-\infty}^{t-r} du \int_{4\pi} d\Omega' \frac{dE_{\text{gw}}(u, \Omega)}{du d\Omega'} \frac{n'_i n'_j}{(1 - \vec{n}' \cdot \vec{n})}. \quad (51)$$

Expression for the corresponding transverse-traceless wavefunction is obtained after multiplication by the projection operator  $\Lambda_{ij,kl}(\vec{n})$ , defined in Eq. (7),

$$[h_{ij}^{\text{mem}}]^{\text{TT}}(t, \vec{x}) = \frac{4G}{r} \int_{-\infty}^{t-r} du \int_{4\pi} d\Omega' \frac{dE_{\text{gw}}(u, \Omega)}{du d\Omega'} \frac{\Lambda_{ij,kl}(\vec{n}) n'_k n'_l}{(1 - \vec{n}' \cdot \vec{n})}, \quad (52)$$

### A. Non-linear memory from circular orbits

We will now compute the non-linear memory from binary orbits. The case of elliptical orbits is important from the point of observations as the non-linear memory signal arises at the same Newtonian order as the oscillatory signals. For the case of hyperbolic orbits the non-linear term is suppressed by a factor  $(v/c)^5$  compared to the quadrupole signal [18].

We will start by computing the non-linear memory from binaries in a circular orbit. The expression for rate of energy radiated in the direction  $d\Omega'$  is given by

$$\frac{dE_{\text{gw}}}{dt' d\Omega'} = \frac{\kappa^2}{4} \int \left( T_{ij}(\omega') T_{kl}^*(\omega') \Lambda_{ij,kl}(\vec{n}') \right) \omega'^3 2\pi \delta(\omega' - 2\omega_0) \frac{d\omega'}{(2\pi)^3 2\omega'}. \quad (53)$$

where  $\omega_0 = \sqrt{G(m_1 + m_2)}/a^3$  is the angular frequency of the Kepler orbit.




FIG. 4. The axis of rotation of the binary designated by  $\vec{L}$  makes an angle  $i$  with the  $z$ -axis and therefore lies in the  $y$ - $z$  plane. The primary graviton emits a secondary graviton at  $\vec{r}' = r' \hat{n}' = r'(\sin \theta' \cos \phi', \sin \theta' \sin \phi', \cos \theta')$  and the secondary graviton travels to the earth located at  $\vec{r} = r \hat{n} = (0, 0, 1)$ .

The stress-energy tensor  $T_{ij}$ , where  $i, j = x, y, z$ , can be written in matrix form as:

$$\mathbb{T}(\omega_0) = \frac{\mu a^2 \omega_0^2}{2} \begin{pmatrix} 1 & \iota & 0 \\ \iota & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}. \quad (54)$$

We consider the orientation defined in Fig. 4, where the axis of rotation of the binary system lies in the  $y$ - $z$  plane and is counter-clockwise rotated, by an angle  $i$ , with respect to the  $z$  axis, i.e.  $\vec{L} = (0, -\sin i, \cos i)$ . Rotation of the axis impacts the form of the stress-tensor matrix  $\mathbb{T}$  as follows:

$$\mathbb{T} \rightarrow \mathbb{T}' = \mathcal{R} \mathbb{T} \mathcal{R}^T, \quad \text{with } \mathcal{R} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos i & -\sin i \\ 0 & \sin i & \cos i \end{pmatrix} \Rightarrow \mathbb{T}' = \frac{\mu a^2 \omega_0^2}{2} \begin{pmatrix} 1 & \iota \cos i & \iota \sin i \\ \iota \cos i & -\cos^2 i & -\cos i \sin i \\ \iota \sin i & -\cos i \sin i & -\sin^2 i \end{pmatrix} \quad (55)$$

The product  $T_{ij} T_{kl}^* \Lambda_{ij,kl}(\hat{n}')$  which can be rewritten entirely as a product of  $T_{ij}$ 's and  $\hat{n}'$ , using Eq. (7), as follows:

$$T_{ij} T_{kl}^* \Lambda_{ij,kl}(\hat{n}') = T_{ij} T_{ji}^* - 2 T_{ij} T_{jl}^* n'_i n'_l + \frac{1}{2} T_{ij} T_{kl}^* n'_i n'_j n'_k n'_l + \frac{1}{2} (T_{ii} T_{kl}^* n'_k n'_l + T_{ij} T_{kk}^* n'_i n'_j - T_{ii} T_{kk}^*). \quad (56)$$

For the circular case both the stress-tensor matrices  $\mathbb{T}$  and  $\mathbb{T}'$  are already traceless. Therefore, the last three terms, within the parentheses, on the right hand side of Eq. (56) vanish. The remaining terms can be evaluated as products of matrices as shown below:

$$T_{ij} T_{kl}^* \Lambda_{ij,kl}(\hat{n}') \equiv \text{Tr}[\mathbb{T}' \mathbb{T}'^\dagger] - 2 (\hat{n}'^T \mathbb{T}') \cdot (\mathbb{T}'^\dagger \hat{n}') + \frac{1}{2} (\hat{n}'^T \mathbb{T}' \hat{n}') (\hat{n}'^T \mathbb{T}'^\dagger \hat{n}') \quad (57)$$

Substituting for  $\mathbb{T}'$ ,  $\hat{n}'$  in Eq. (57) and evaluating the matrix products and traces yields:

$$T_{ij} T_{kl}^* \Lambda_{ij,kl}(\hat{n}') \equiv \frac{\mu^2 a^4 \omega_0^4}{4} \frac{1}{2} (1 + 6 \cos^2 \theta + \cos^4 \theta). \quad (58)$$

Here,  $\theta$  is the angle between the axis of rotation of the binary  $\vec{L} = (0, -\sin i, \cos i)$  and the direction where the primary

graviton emits the second graviton:  $\hat{n}' = (\sin \theta' \cos \phi', \sin \theta' \sin \phi', \cos \theta')$ , i.e.,

$$\cos \theta = -\sin i \sin \theta' \sin \phi' + \cos i \cos \theta'. \quad (59)$$

Eq. (53) can now be expressed, using the result of Eq. (58) as

$$\frac{dE_{\text{gw}}}{dt' d\Omega'} = \frac{G}{2\pi} \omega_0^6 \mu^2 a^4 \left(1 + 6 \cos^2 \theta + \cos^4 \theta\right). \quad (60)$$

The Kepler orbital frequency can be written in terms of the semi-major axis  $a$ , i.e.  $\omega_0 = (GM/a^3)^{1/2}$  ( $M = m_1 + m_2$ ). This allows us to rewrite the expression for the rate of energy radiated as,

$$\frac{dE_{\text{gw}}}{dt' d\Omega'} = \frac{G^4 \mu^2 M^3}{2\pi a^5} \left(1 + 6 \cos^2 \theta + \cos^4 \theta\right). \quad (61)$$

Substituting the above in Eq. (51), we obtain the following expression for the memory signal:

$$\begin{aligned} h_{ij}^{\text{mem}}(t, \vec{x}) &= \frac{4G}{r} \int_{-\infty}^{t-r} dt' \int_{4\pi} d\Omega' \frac{dE_{\text{gw}}(t', \Omega)}{dt' d\Omega'} \frac{n'_i n'_j}{(1 - \hat{n}' \cdot \hat{n})} \\ &= \frac{4G}{r} \int_{-\infty}^t dt' \frac{G^4 \mu^2 M^3}{2\pi a^5} \underbrace{\int_{4\pi} d\Omega' \left(1 + 6 \cos^2 \theta + \cos^4 \theta\right)}_{A_{ij}} \frac{n'_i n'_j}{(1 - \hat{n}' \cdot \hat{n})}. \end{aligned} \quad (62)$$

where we have replaced  $t - r$  by  $t$  in the limit of the integrand as the waves are detected at a fixed  $r$  and the dependence in  $u$  then becomes a dependence on time of observation  $t$ . To obtain the transverse-traceless part of  $h_{ij}^{\text{mem}}$  we apply the TT projection operator, through multiplication with  $\Lambda_{ij,kl}(\vec{n})$ , see Eq. (52),

$$\left[h_{ij}^{\text{mem}}\right]^{\text{TT}}(t, \vec{x}) = \frac{4G^5}{2\pi r} \mu^2 M^3 \int_{-\infty}^t dt' \frac{1}{a^5} \underbrace{\int_{4\pi} d\Omega' \left(1 + 6 \cos^2 \theta + \cos^4 \theta\right) \frac{\Lambda_{ij,kl}(\hat{n}) n'_k n'_l}{(1 - \hat{n}' \cdot \hat{n})}}_{[A_{ij}]^{\text{TT}}}. \quad (63)$$

The tensor product within the angular integral can be further expressed as:

$$\Lambda_{ij,kl}(\hat{n}) n'_k n'_l = \left[ \left( n'_i n'_j - \frac{1}{2} \delta_{ij} + \frac{1}{2} n_i n_j \right) - (n_i n'_j + n'_i n_j) (\hat{n}' \cdot \hat{n}) + \frac{1}{2} (\delta_{ij} + n_i n_j) (\hat{n}' \cdot \hat{n})^2 \right]. \quad (64)$$

Since,  $\hat{n} = (0, 0, 1)$ ,  $\hat{n}' \cdot \hat{n} = \cos \theta'$  and substituting for the components of  $\hat{n}$ ,  $\hat{n}'$  and  $\delta_{ij}$ , the angular integrals can be evaluated, for  $i, j = 1, 2$  as follows:

$$\begin{aligned} [A_{11}]^{\text{TT}} &= \int_0^\pi \sin \theta' d\theta' \int_0^{2\pi} d\phi' (1 + 6 \cos^2 \theta + \cos^4 \theta) \times \frac{1}{2} (1 + \cos \theta') \cos 2\phi' = -\frac{2\pi}{15} (17 + \cos^2 i) \sin^2 i, \\ [A_{12}]^{\text{TT}} &= \int_0^\pi \sin \theta' d\theta' \int_0^{2\pi} d\phi' (1 + 6 \cos^2 \theta + \cos^4 \theta) \times \frac{1}{2} (1 + \cos \theta') \sin 2\phi' = 0. \end{aligned} \quad (65)$$

The other components are:  $[A_{22}]^{\text{TT}} = -[A_{11}]^{\text{TT}}$  and  $[A_{12}]^{\text{TT}} = [A_{21}]^{\text{TT}} = 0$ . The transverse-traceless gravitational wavefunction can be decomposed into + and  $\times$  modes:

$$\left[h_{ij}^{\text{mem}}\right]^{\text{TT}} = h_I \epsilon_{ij}^I = h_+^{\text{mem}} \epsilon_{ij}^+ + h_\times^{\text{mem}} \epsilon_{ij}^\times. \quad (66)$$

Based on this we can make the identification:

$$h_+^{\text{mem}}(t, \vec{x}) = \frac{4G^5}{2\pi r} \mu^2 M^3 \int_{-\infty}^{t-r} dt' \frac{1}{a^5} \times \left[ \frac{2\pi}{15} (17 + \cos^2 i) \sin^2 i \right], \quad h_\times^{\text{mem}}(t, \vec{x}) = 0. \quad (67)$$

The time dependence of the integrand in Eq. (62) is due to the change in the radius of the orbit which occurs due to the energy loss of the orbit by the primary gravitational waves. This change in the radius is given as  $da/dt = (dE/dt)(da/dE)$ , which using  $E = -(1/2)G\mu M/a$  is

$$\frac{da}{dt} = -\frac{64 G^3 \mu M^2}{5 a^3}. \quad (68)$$

Solving this equation gives us the time dependence of the separation distance  $a(t)$

$$a(t) = a_c \left[ 1 + \frac{256 G^3 \mu M^2}{5 a_c^4} (t_c - t) \right]^{1/4}. \quad (69)$$

where  $t_c$  is the time of coalescence of the black-holes. The growth of memory signal stops at the time of the coalescence  $t_c$  when the black-hole separation becomes smaller than the innermost stable circular orbit radius  $a_c = 6 GM$ . The frequency  $f = \omega_0/\pi = (1/\pi)\sqrt{(GM)/a^3}$  increases till the time of coalescence as

$$f(t') \simeq \left( \frac{5}{256} \right)^{3/8} \frac{1}{\pi} (G\mathcal{M}_c)^{-5/8} \left[ \frac{5}{256} \frac{a_c^4}{G^3 \mu M^2} + (t_c - t') \right]^{-3/8} \quad (70)$$

where  $\mathcal{M}_c = \mu^{3/5} M^{2/5}$  is the chirp mass of the binary pair.

Using Eq. (69) to substitute for  $a(t)$  in Eq. (67), we obtain the the non-linear memory wavefunction of binaries, whose rotation axis makes an angle  $i$  with respect to the earth-source direction, as

$$\begin{aligned} h_+^{\text{mem}}(t) &= \frac{1}{192 r} G (5G\mu^3 M^2)^{1/4} \frac{1}{\left[ 5 \left( \frac{3}{2} \right)^4 \frac{GM^2}{\mu} + (t_c - t) \right]^{1/4}} \sin^2 i (17 + \cos^2 i), \\ &= h_{\text{circ}} \frac{1}{[1 + (t'_c - t')]^{1/4}} \sin^2 i (17 + \cos^2 i), \end{aligned} \quad (71)$$

where  $h_{\text{circ}} = \frac{2}{3} \frac{1}{192} \frac{G\mu}{r}$  and  $(t'_c - t') = (t_c - t)/T'$  with  $T'$  being  $\left( \frac{3}{2} \right)^4 \frac{5GM^2}{\mu}$ . The memory effect will be strongest in edge on binaries  $i = \pi/2$  and will be zero in the face on binaries  $i = 0$ . This is demonstrated in Fig. 5.




FIG. 5. The variation in  $h_+^{\text{mem}}$  against time plotted for different choices of the angle  $i$  between the line of sight of observation and the binary axis.

## VI. ELLIPTICAL ORBITS

For eccentric elliptical orbits, Peters and Mathews [81, 82] calculated the average energy and angular momentum emission rates at Newtonian order. Their calculation has been improved to the 3-PN level [83, 84], including nonlinear tail-effect (which arises from the scattering of gravitational waves by the near-field potential) at 3-PN [85].

In this section we compute the frequency spectrum of the energy radiated following [71–74]. The radiated energy acts as the source term for secondary gravitational waves which carry the non-linear memory signal.

To describe a compact binary system, comprised of stars having masses  $m_1$  and  $m_2$ , in an elliptical orbit, the following quantities are of relevance:

- Since the orbital motion is around the common centre of mass, it is useful to describe the motion in terms of the reduced mass  $\mu = \frac{m_1 m_2}{m_1 + m_2}$  and the total mass  $M = (m_1 + m_2)$ .
- The elliptical Keplerian orbit can be parametrized as:

$$x(\xi) = a(\cos \xi - e), \quad y(\xi) = b \sin \xi, \quad z(\xi) = 0, \quad \frac{\omega'_n}{n} t = \Omega t = (\xi - e \sin \xi). \quad (72)$$

Here,  $a$  and  $b$  ( $= a\sqrt{1-e^2}$ ,  $e < 1$ ) denote the semi-major and semi-minor axes respectively. The variables  $e$  and  $\xi \in (0, 2\pi)$  refer to the eccentricity of the orbit and the eccentric anomaly respectively.

- The angular frequency corresponding to the  $n^{\text{th}}$  harmonic is denoted as  $\omega'_n = n\Omega$ , with  $n = \{0, 1, 2, \dots\}$  being a non-negative integer and the fundamental frequency  $\Omega$  can be related to the semi-major axis and the total mass of the system as:  $\Omega = \left(\frac{G(m_1 + m_2)}{a^3}\right)^{1/2}$ .

The first step in the computation of the rate of energy radiated involves the evaluation of the stress-tensor components in frequency space. Once again starting with the  $xx$  component, the calculation proceeds as follows:

$$T_{xx}(\omega'_n) = \frac{\mu}{T} \int_0^T dt \dot{x}^2(t) e^{i\omega'_n t} = \frac{-i\mu\omega'_n}{T} \int_0^T dt \dot{x}(t) x(t) e^{i\omega'_n t}. \quad (73)$$

In the above, we have used integration by parts. Next, the parametric form of the orbit coordinates, shown in Eq. (72), allows us to write,

$$\dot{x} dt = \frac{dx}{d\xi} d\xi = -a \sin \xi d\xi. \quad (74)$$

Using the above transformation and also substituting for  $x$  and  $\omega'_n t$  in terms of functions of  $\xi$ , we get

$$T_{xx}(\omega'_n) = \frac{i\mu a^2 \omega_n'^2}{2\pi n} \int_0^{2\pi} d\xi \sin \xi (\cos \xi - e) e^{in(\xi - e \sin \xi)} = -\frac{\mu a^2 \omega_n'^2}{n} \left[ \left(\frac{1-e^2}{e}\right) J'_n(ne) - \frac{1}{ne^2} J_n(ne) \right]. \quad (75)$$

To arrive at the second line of Eq. (75), we replaced the trigonometric functions by the corresponding exponential functions and identified the integral form of Bessel functions of first kind,

$$J_n(z) = \frac{1}{2\pi} \int_0^{2\pi} e^{in(\xi - e \sin \xi)} d\xi. \quad (76)$$

The final expression in terms of  $J_n(ne)$  and  $J'_n(ne)$  is obtained by utilizing the recurrence relations given below:

$$J_{n-1}(z) + J_{n+1}(z) = \frac{2n}{z} J_n(z), \quad J_{n-1}(z) - J_{n+1}(z) = 2 J'_n(z). \quad (77)$$

The other non-zero components, i.e  $T_{yy}$  and  $T_{xy}$  can similarly be obtained as:

$$\begin{aligned} T_{yy}(\omega'_n) &= \frac{\mu}{T} \int_0^T dt \dot{y}^2(t) e^{i\omega'_n t} = \frac{\mu \omega_n'^2 a^2 (1-e^2)}{n} \left[ \frac{1}{e} J'_n(ne) - \frac{1}{ne^2} J_n(ne) \right], \\ T_{xy}(\omega'_n) &= \frac{\mu}{T} \int_0^T dt \dot{x}(t) \dot{y}(t) e^{i\omega'_n t} = \frac{i\mu \omega_n'^2 a^2 \sqrt{(1-e^2)}}{n} \left[ - \left( \frac{1-e^2}{e^2} \right) J_n(ne) + \frac{1}{ne} J'_n(ne) \right]. \end{aligned} \quad (78)$$

### A. Rate of gravitational radiation from binaries in elliptical orbits

Substituting for  $T_{xx}$ ,  $T_{yy}$  and  $T_{xy}$  using Eqs. (75) and (78) gives,

$$T_{ij}(\omega') T_{ji}^*(\omega') - \frac{1}{3} |T^i_i(\omega')|^2 = g(n, e). \quad (79)$$

where we have defined  $g(n, e)$  as,

$$\begin{aligned} g(n, e) &= J_n(ne)^2 \left[ \frac{2n^2}{e^4} (1-e^2)^3 + \frac{6-6e^2+2e^4}{3e^4} \right] + J'_n(ne)^2 \left[ \frac{2n^2}{e^2} (1-e^2)^2 + \frac{2(1-e^2)}{e^2} \right] \\ &+ J_n(ne) J'_n(ne) \left[ \frac{(-8+14e^2-6e^4)n}{e^3} \right]. \end{aligned} \quad (80)$$

In this case, the energy loss due to gravitational radiation as given in Eq. (17) becomes,

$$\frac{dE_{\text{gw}}}{dt}(e) = \frac{\kappa^2}{8(2\pi)^2} \sum_{n=0}^{\infty} \frac{8\pi}{5} \left[ T_{ij}(\omega'_n) T_{ji}^*(\omega'_n) - \frac{1}{3} |T^i_i(\omega'_n)|^2 \right] \omega_n'^2 = \frac{32G}{20} \Omega^6 \mu^2 a^4 \sum_{n=0}^{\infty} n^2 g(n, e). \quad (81)$$

The series sum over products of Bessel functions and their derivative, weighted by powers of  $n$ , can be expressed completely as functions of  $e$ , using the identities derived in the appendix of [81] and reproduced in Eq. (112). This enables the identification of the eccentricity dependent part of the energy loss as

$$\tilde{g}(e) = \sum_{n=0}^{\infty} n^2 g(n, e) = \frac{4}{(1-e^2)^{7/2}} \left( 1 + \frac{73}{24} e^2 + \frac{37}{96} e^4 \right). \quad (82)$$

Therefore, the energy radiated as gravitational waves from the binary can be written as:

$$\frac{dE_{\text{gw}}}{dt} = \frac{32G}{5} \Omega^6 \left( \frac{m_1 m_2}{m_1 + m_2} \right)^2 a^4 \frac{1}{(1-e^2)^{7/2}} \left( 1 + \frac{73}{24} e^2 + \frac{37}{96} e^4 \right). \quad (83)$$

### B. Non-linear memory from elliptical orbits

Rate of energy radiated with respect to time and solid angle is described using the following formula:

$$\frac{dE_{\text{gw}}}{dt' d\Omega'} = \frac{\kappa^2}{8(2\pi)^2} \sum_{n=0}^{\infty} \left[ T_{ij}(\omega'_n) T_{kl}^*(\omega'_n) \Lambda_{ij,kl}(\hat{n}') \right] \omega_n'^2 \quad (84)$$



The stress-energy tensor  $T_{ij}$ , where  $i, j = x, y, z$ , can be written in matrix form as:

$$\mathbb{T}(n, e) = \mu a^2 \Omega^2 \begin{pmatrix} q_1(n, e) & \iota q_2(n, e) & 0 \\ \iota q_2(n, e) & q_3(n, e) & 0 \\ 0 & 0 & 0 \end{pmatrix}. \quad (85)$$

The non-zero elements of the matrix are functions of  $n$ , corresponding to the  $n$ th harmonic, and  $e$ , the eccentricity of the orbit and these are given as

$$\begin{aligned} q_1(n, e) &= -\frac{n(1-e^2)}{e} J'_n(ne) + \frac{1}{e^2} J_n(ne), \\ q_2(n, e) &= \frac{(1-e^2)^{1/2}}{e} J'_n(ne) - \frac{n(1-e^2)^{3/2}}{e^2} J_n(ne), \\ q_3(n, e) &= \frac{n(1-e^2)}{e} J'_n(ne) - \frac{(1-e^2)}{e^2} J_n(ne). \end{aligned} \quad (86)$$

We follow the orientation for the system as shown in Fig. 4. The stress-energy matrix for the rotated system is given as  $\mathbb{T}' = \mathcal{R} \mathbb{T} \mathcal{R}^T$ , with  $\mathcal{R}$  being the rotation matrix defined in Eq. (55). This allows us to write,

$$\mathbb{T}' = \mu a^2 \Omega^2 \begin{pmatrix} q_1(n, e) & \iota q_2(n, e) \cos i & \iota q_2(n, e) \sin i \\ \iota q_2(n, e) \cos i & q_3(n, e) \cos^2 i & q_3(n, e) \cos i \sin i \\ \iota q_2(n, e) \sin i & q_3(n, e) \cos i \sin i & q_3(n, e) \sin^2 i \end{pmatrix}. \quad (87)$$

Expanding the matrix product  $T_{ij} T_{kl}^* \Lambda_{ij,kl}(\hat{n}')$  in terms of products of  $T_{ij}$  and  $n'_i$ , yields the same result as Eq. (56),

$$T_{ij} T_{kl}^* \Lambda_{ij,kl}(\hat{n}') = T_{ij} T_{ji}^* - 2 T_{ij} T_{jl}^* n'_i n'_l + \frac{1}{2} T_{ij} T_{kl}^* n'_i n'_j n'_k n'_l + \frac{1}{2} (T_{ii} T_{kl}^* n'_k n'_l + T_{ij} T_{kk}^* n'_i n'_j - T_{ii} T_{kk}^*). \quad (88)$$

Unlike the case of circular orbits, none of the terms on the right hand side of Eq. (88) vanish. Therefore, the expression when rewritten as a sum over matrix products appears as,

$$\begin{aligned} T_{ij} T_{kl}^* \Lambda_{ij,kl}(\hat{n}') &\equiv \text{Tr}[\mathbb{T}' \mathbb{T}'^\dagger] - 2 (\hat{n}'^T \mathbb{T}') \cdot (\mathbb{T}'^\dagger \hat{n}') + \frac{1}{2} (\hat{n}'^T \mathbb{T}' \hat{n}') (\hat{n}'^T \mathbb{T}'^\dagger \hat{n}') + \\ &\frac{1}{2} (\hat{n}'^T \mathbb{T}' \hat{n}') \text{Tr}[\mathbb{T}'^\dagger] + \frac{1}{2} \text{Tr}[\mathbb{T}'] (\hat{n}'^T \mathbb{T}'^\dagger \hat{n}') - \frac{1}{2} \text{Tr}[\mathbb{T}'] \text{Tr}[\mathbb{T}'^\dagger]. \end{aligned} \quad (89)$$

Since,  $T_{ij}$ 's are functions of  $n$  and  $e$ , while  $\hat{n}'$  is parametrized in terms of the angles  $\theta'$ ,  $\phi'$ , the matrix product can be expressed as an overall function of  $n$ ,  $e$ ,  $\theta'$ , and  $\phi'$

$$T_{ij}(n, e) T_{kl}^*(n, e) \Lambda_{ij,kl}(\theta', \phi') = \mu^2 a^4 \Omega^4 \mathcal{I}(n, e, \theta', \phi') \quad (90)$$

with

$$\begin{aligned} \mathcal{I}(n, e, \theta', \phi') &= \frac{1}{2} q_1^2(n, e) [1 - p_1^2(\theta', \phi')]^2 + \frac{1}{2} q_3^2(n, e) [1 - p_2^2(\theta', \phi')]^2 - 2 q_1(n, e) q_3(n, e) \\ &+ 2 q_2^2(n, e) [1 - p_1^2(\theta', \phi')] [1 - p_2^2(\theta', \phi')] + q_1(n, e) q_3(n, e) [1 + p_1^2(\theta', \phi')] [1 + p_2^2(\theta', \phi')]. \end{aligned} \quad (91)$$

Here,

$$p_1(\theta', \phi') = \sin \theta' \cos \phi' \quad \text{and} \quad p_2(\theta', \phi') = \cos i \sin \theta' \sin \phi' + \sin i \cos \theta' \quad (92)$$

Computation of the transverse-traceless wavefunction involves (i) an angular integral over  $\theta'$ ,  $\phi'$  and (ii) a sum over

the orders ( $n$ ) of the Bessel functions. This will ultimately yield an eccentricity dependent function:

$$[\mathbf{A}_{ij}]^{\text{TT}}(e) = \sum_{n=0}^{\infty} \int_{4\pi} d\Omega' \mathcal{I}(n, e, \theta', \phi') \frac{\Lambda_{ij,kl}(\hat{n}) n'_k n'_l}{(1 - \hat{n}' \cdot \hat{n})}. \quad (93)$$

The tensor product within the angular integral can be further expanded as:

$$\Lambda_{ij,kl}(\hat{n}) n'_k n'_l = \left[ \left( n'_i n'_j - \frac{1}{2} \delta_{ij} + \frac{1}{2} n_i n_j \right) - (n_i n'_j + n'_i n_j) (\hat{n}' \cdot \hat{n}) + \frac{1}{2} (\delta_{ij} + n_i n_j) (\hat{n}' \cdot \hat{n})^2 \right]. \quad (94)$$

Since,  $\hat{n} = (0, 0, 1)$ ,  $\hat{n}' \cdot \hat{n} = \cos \theta'$  and substituting for the components of  $\hat{n}$ ,  $\hat{n}'$  and  $\delta_{ij}$ , the angular integrals can be evaluated, for  $i, j = 1, 2$  as follows:

$$\begin{aligned} [\mathbf{A}_{11}]^{\text{TT}}(e) &= \sum_{n=0}^{\infty} \int_0^\pi \sin \theta' d\theta' \int_0^{2\pi} d\phi' \mathcal{I}(n, e, \theta', \phi') \times \frac{1}{2} (1 + \cos \theta') \cos 2\phi' \\ &= \frac{2\pi}{15} \sum_{n=0}^{\infty} \left[ \left( 6 q_1(n, e) q_3(n, e) - 3 q_1^2(n, e) - 8 q_2^2(n, e) \right) \right. \\ &\quad \left. + \left( 8 q_2^2(n, e) + 2 q_3^2(n, e) - 6 q_1(n, e) q_3(n, e) \right) \cos^2 i + q_3^2(n, e) \cos^4 i \right] \\ &= \frac{2\pi}{15} (C_0(e) + C_2(e) \cos^2 i + C_4(e) \cos^4 i) \end{aligned} \quad (95)$$

The sum over  $n$  can be evaluated, using the identities given in Eq. (112), to obtain the coefficients  $C_i(e)$ ,  $i = 0, 1, 2$ :

$$\begin{aligned} C_0(e) &= \sum_{n=0}^{\infty} \left[ \frac{(6e^2 - 9)}{e^4} n^2 [J_n(ne)]^2 - \frac{8(1 - e^2)^3}{e^4} n^4 [J_n(ne)]^2 - \frac{8(1 - e^2)}{e^2} n^2 [J'_n(ne)]^2 - \frac{9(1 - e^2)^2}{e^2} n^4 [J'_n(ne)]^2 \right. \\ &\quad \left. + \frac{(1 - e^2)}{e^3} (34 - 22e^2) n^3 [J_n(ne) J'_n(ne)] \right] \\ &= -\frac{1}{4e^2(1 - e^2)^{7/2}} \left[ 68e^2 + \frac{1565}{8}e^4 + \frac{1071}{64}e^6 \right], \\ C_2(e) &= \sum_{n=0}^{\infty} \left[ \frac{2(4 - 5e^2 + e^4)}{e^4} n^2 [J_n(ne)]^2 + \frac{8(1 - e^2)^3}{e^4} n^4 [J_n(ne)]^2 + \frac{8(1 - e^2)}{e^2} n^2 [J'_n(ne)]^2 + \frac{8(1 - e^2)^2}{e^2} n^4 [J'_n(ne)]^2 \right. \\ &\quad \left. + \frac{2(1 - e^2)}{e^3} (-16 + 13e^2) n^3 [J_n(ne) J'_n(ne)] \right] \\ &= \frac{1}{4e^2(1 - e^2)^{7/2}} \left[ 64e^2 + \frac{399}{2}e^4 + \frac{101}{4}e^6 \right], \\ C_4(e) &= \sum_{n=0}^{\infty} \frac{(1 - e^2)^2}{e^4} \left[ n^2 [J_n(ne)]^2 + e^2 n^4 [J'_n(ne)]^2 - 2e n^3 [J_n(ne) J'_n(ne)] \right] \\ &= \frac{1}{4e^2(1 - e^2)^{9/2}} \left[ -1 - \frac{15}{4}e^2 + \frac{83}{4}e^4 + \frac{21}{8}e^6 - \frac{37}{64}e^8 \right]. \end{aligned} \quad (96)$$

The cross-component  $[\mathbf{A}_{12}]^{\text{TT}}(e)$  can similarly be evaluated as:

$$[\mathbf{A}_{12}]^{\text{TT}}(e) = \sum_{n=0}^{\infty} \int_0^\pi \sin \theta' d\theta' \int_0^{2\pi} d\phi' \mathcal{I}(n, e, \theta', \phi') \times \frac{1}{2} (1 + \cos \theta') \sin 2\phi' = 0. \quad (97)$$

The other components can be obtained using the above as:  $[A_{22}]^{\text{TT}} = -[A_{11}]^{\text{TT}}$  and  $[A_{12}]^{\text{TT}} = [A_{21}]^{\text{TT}} = 0$ . The decomposition of the transverse-traceless wavefunction into + and  $\times$  polarization modes, see Eq. (66), allows us to identify:

$$h_+^{\text{mem}}(t, \vec{x}) = \frac{4G}{r} \int_{-\infty}^{t-r} dt' \frac{1}{\pi} \frac{G^4 \mu^2 M^3}{a(t')^5} \times \frac{2\pi}{15} \left( C_0(e) + C_2(e) \cos^2 i + C_4(e) \cos^4 i \right), \quad h_\times^{\text{mem}}(t, \vec{x}) = 0. \quad (98)$$

Within the integrand, the time-dependence is encoded in both the semi-major axis  $a$  and the eccentricity  $e$  of the orbit. This is on account of the fact that energy loss due to primary gravitational waves alters the features of the orbit. The explicit time-dependence for  $a(t)$  and  $e(t)$  can be obtained by solving the following system of non-linear differential equations [82]:

$$\frac{da}{dt} = -\frac{64}{5} \frac{G^3 \mu M^2}{a^3 (1-e^2)^{7/2}} \left( 1 + \frac{73}{24} e^2 + \frac{37}{96} e^4 \right), \quad (99)$$

$$\frac{de}{dt} = -\frac{304}{15} \frac{G^3 \mu M^2}{a^4 (1-e^2)^{5/2}} \left( 1 + \frac{121}{304} e^2 \right). \quad (100)$$

The above can be used to eliminate  $t$  and obtain a differential equation involving  $a$  and  $e$ , i.e.,

$$\frac{da}{de} = \frac{12}{19} \frac{a}{e(1-e^2)} \frac{1 + (73/24)e^2 + (37/96)e^4}{1 + (121/304)e^2}, \quad (101)$$

whose solution expresses  $a$  in terms of  $e$ ,

$$a(e) = c_0 \frac{e^{12/19}}{(1-e^2)} \left( 1 + \frac{121}{304} e^2 \right)^{870/2299}, \quad (102)$$

where  $c_0 = a_0 e_0^{-12/19} (1-e_0^2) (1 + (121/304)e_0^2)^{-870/2299}$  and  $a_0 = a(e_0)$  defines the initial condition necessary for solving the differential equation. The time evolution can be given as a function of eccentricity as

$$t - t_c = -\frac{15}{304} \frac{c_0^4}{G^3 \mu M^2} \frac{19}{48} e^{48/19} F_1 \left( \frac{24}{19}, -\frac{1181}{2299}, \frac{3}{2}, \frac{43}{19}; -\frac{121}{304} e^2, e^2 \right), \quad (103)$$

where  $t_c$  corresponds to the instant of coalescence of the black-holes, i.e., when  $e$  becomes close to zero.  $F_1$  is the hypergeometric Appell function with the following integral representation:

$$F_1(a, b_1, b_2, c; x, y) = \frac{\Gamma(c)}{\Gamma(a)\Gamma(c-a)} \int_0^1 t^{a-1} (1-t)^{c-a-1} (1-xt)^{-b_1} (1-yt)^{-b_2} dt, \quad \Re c > \Re a > 0. \quad (104)$$

Eq. (103) can be expressed entirely in terms of dimensionless quantities by noting that  $\bar{T} = c_0^4 / (G^3 \mu M^2)$  has the dimensions of time. Therefore,

$$\frac{t - t_c}{\bar{T}} = \bar{t} - \bar{t}_c = -\frac{15}{304} \frac{19}{48} e^{48/19} F_1 \left( \frac{24}{19}, -\frac{1181}{2299}, \frac{3}{2}, \frac{43}{19}; -\frac{121}{304} e^2, e^2 \right), \quad (105)$$

The wavefunction in Eq. (98) can be rewritten, after a change of integration variable, from  $t'$  to  $e'$  as,

$$\begin{aligned} h_+^{\text{mem}}(e, \vec{x}) &= \frac{4G}{r} \int_e^0 \frac{de'}{\dot{e}(e') \pi} \frac{1}{\pi} \frac{G^4 \mu^2 M^3}{a(e')^5} \frac{2\pi}{15} [C_0(e') + C_2(e') \cos^2 i + C_4(e') \cos^4 i], \\ &= \frac{4G}{r} \frac{15}{304} \frac{G\mu M}{c_0} \frac{2}{15} [C_h^{(0)}(e) + C_h^{(2)}(e) \cos^2 i + C_h^{(4)}(e) \cos^4 i], \end{aligned} \quad (106)$$

or in terms of dimensionless quantities, we can write

$$\frac{h_+^{\text{mem}}(e, \vec{x})}{h_0} = \bar{h}_+^{\text{mem}}(e, \vec{x}) = \frac{1}{152} \left[ C_h^{(0)}(e) + C_h^{(2)}(e) \cos^2 i + C_h^{(4)}(e) \cos^4 i \right]. \quad (107)$$

Here,  $h_0 = (4G^2\mu M)/(rc_0)$ . The eccentricity dependent part is constituted of

$$\begin{aligned} C_h^{(0)}(e) &= 17 \frac{19 e^{-12/19}}{12} F_1 \left( -\frac{6}{19}, \frac{3169}{2299}, 0, \frac{13}{19}; -\frac{121}{304} e^2, 0 \right) - \frac{1565}{32} \frac{19 e^{26/19}}{26} F_1 \left( \frac{13}{19}, \frac{3169}{2299}, 0, \frac{32}{19}; -\frac{121}{304} e^2, 0 \right) \\ &\quad - \frac{1071}{256} \frac{19 e^{64/19}}{64} F_1 \left( \frac{32}{19}, \frac{3169}{2299}, 0, \frac{51}{19}; -\frac{121}{304} e^2, 0 \right), \\ C_h^{(2)}(e) &= -16 \frac{19 e^{-12/19}}{12} F_1 \left( -\frac{6}{19}, \frac{3169}{2299}, 0, \frac{13}{19}; -\frac{121}{304} e^2, 0 \right) + \frac{399}{8} \frac{19 e^{26/19}}{26} F_1 \left( \frac{13}{19}, \frac{3169}{2299}, 0, \frac{32}{19}; -\frac{121}{304} e^2, 0 \right) \\ &\quad + \frac{101}{16} \frac{19 e^{64/19}}{64} F_1 \left( \frac{32}{19}, \frac{3169}{2299}, 0, \frac{51}{19}; -\frac{121}{304} e^2, 0 \right), \\ C_h^{(4)}(e) &= \frac{1}{4} \frac{19 e^{-50/19}}{50} F_1 \left( -\frac{25}{19}, \frac{3169}{2299}, 1, -\frac{6}{19}; -\frac{121}{304} e^2, e^2 \right) + \frac{15}{16} \frac{19 e^{-12/19}}{12} F_1 \left( -\frac{6}{19}, \frac{3169}{2299}, 1, \frac{13}{19}; -\frac{121}{304} e^2, e^2 \right) \\ &\quad + \frac{83}{16} \frac{19 e^{26/19}}{26} F_1 \left( \frac{13}{19}, \frac{3169}{2299}, 1, \frac{32}{19}; -\frac{121}{304} e^2, e^2 \right) + \frac{21}{32} \frac{19 e^{64/19}}{64} F_1 \left( \frac{32}{19}, \frac{3169}{2299}, 1, \frac{51}{19}; -\frac{121}{304} e^2, e^2 \right) \\ &\quad - \frac{37}{256} \frac{19 e^{102/19}}{102} F_1 \left( \frac{51}{19}, \frac{3169}{2299}, 1, \frac{70}{19}; -\frac{121}{304} e^2, e^2 \right), \end{aligned} \quad (108)$$

The variation in the memory waveform with time can be obtained by exploiting the relation between the waveform and eccentricity of the orbit, as described in Eq. (107) and the implicit relation between time and eccentricity highlighted in Eq. (105). A plot depicting such variation is shown in Fig. 6.




FIG. 6. The variation in  $\bar{h}_+^{\text{mem}}$  with respect to the time interval  $\bar{t} - \bar{t}_c$  plotted for different choices of the angle between between the line of sight of observation and the binary axis. The first figure displays the  $\bar{h}_+^{\text{mem}}$  values on a logarithmic scale, whereas the second figure presents the same information but with the regular scaling of the axes.

An alternative description can be obtained in terms of the frequency  $\nu = (2\pi)^{-1} m^{1/2} a^{-3/2}$ , by noting that the

relation between  $\nu$  and  $e$  can be obtained by solving the following differential equation:

$$\frac{d\nu}{de} = -\frac{18}{19} \frac{\nu}{e(1-e^2)} \frac{1 + (73/24)e^2 + (37/96)e^4}{1 + (121/304)e^2}. \quad (109)$$

The solution can be evaluated as,

$$\nu(e) = \tilde{c}_0 e^{-18/19} (1-e^2)^{3/2} \left(1 + \frac{121}{304}e^2\right)^{-1305/2299}, \quad (110)$$

where  $\tilde{c}_0 = \nu_0 e_0^{-18/19} (1-e_0^2)^{-3/2} (1 + (121/304)e_0^2)^{1305/2299}$  with  $\nu_0 = \nu(e_0)$  defining the initial condition. In terms of dimensionless quantities, the same equation can be rewritten as,

$$\frac{\nu(e)}{\tilde{c}_0} = \bar{\nu}(e) = e^{-18/19} (1-e^2)^{3/2} \left(1 + \frac{121}{304}e^2\right)^{-1305/2299}, \quad (111)$$

The variation in the memory signal with respect to the frequency can be obtained based on their mutual dependence on eccentricity, as highlighted in Eqs. (107) and (111). A plot depicting such a relation is shown in Fig. 7.




FIG. 7. The variation in  $\bar{h}_+^{\text{mem}}$  with respect to the dimensionless frequency parameter  $\bar{\nu}$ , defined in Eq. (111), plotted for different choices of the angle between between the line of sight of observation and the binary axis.

## VII. CONCLUSIONS

In this paper we have computed the linear memory signal for eccentric hyperbolic encounters in both frequency and time domain. We performed a field theoretic calculation of the amplitude of graviton emission from a classical stress tensor and related this amplitude to the gravitational wave signal as a function of frequency. We took the zero-frequency limit of the waveform to identify the memory signals which we computed in the frequency space and taking the Fourier transform also in the time domain. We performed these calculations for hyperbolic orbits with large eccentricity while retaining terms of all orders in eccentricity. We find that in the eccentric hyperbolic orbits

the low-frequency memory component has terms which vary with the frequency  $\omega = \nu\Omega$  of the gravitational waves as  $\log(\omega e/\Omega)$ . This log dependence is due to the eccentric hyperbolic orbit and is different from the tail terms which occur for any unbound scattering at  $\mathcal{O}(G^2)$  [25–27]. Hyperbolic encounters of supermassive black-holes of mass  $10^7 M_\odot$  with closest approach of 1pc will produce gravitational waves with peak frequency  $\sim 0.2$  mHz and encounters between blackholes of mass  $10^3 M_\odot$  will produce gravitational waves at peak frequency 0.04 mHz . The time window for the observations is of the order of 1 day which makes such observations possible for third generation gravitational detectors.

The non-linear memory which occurs from GW radiated by GW is equally important as it would prove the non-linear nature of graviton-graviton interactions. We have computed the non-linear memory for eccentric elliptical orbits. Highly eccentric orbits are possible if the initial system has three bodies of which one is ejected [37] or by capture from an unbound orbit [35, 36]. We compute the frequency spectrum energy radiated by the binaries using the field theory technique which gives the energy spectrum directly in the frequency space. This is used as the source term for the secondary gravitational waves, which results in the memory waveform in the frequency space. We use the instantaneous eccentricity as the parameter for tracking the change of frequency and semi-major axis in time due to radiation reaction. We thus obtain the non-linear memory as a function of the instantaneous eccentricity. We then exploited the change in eccentricity with respect to both time and frequency to obtain the non-linear waveform as a function of both time and frequency. The calculation of the non-linear memory has been done up to all orders in eccentricity. These signal templates in frequency and time domain may be useful for extracting the memory signal from data by the future detectors like Advanced-LIGO, Einstein Telescope and LISA.

## VIII. APPENDIX

For evaluating the sum over  $n$  for elliptical binaries, we have used following formulae [81]:

$$\begin{aligned}
\sum_{n=0}^{\infty} n^2 J_n^2(ne) &= \frac{e^2}{4(1-e^2)^{7/2}} \left(1 + \frac{e^2}{4}\right), \\
\sum_{n=0}^{\infty} n^2 [J'_n(ne)]^2 &= \frac{e^2}{4(1-e^2)^{5/2}} \left(1 + \frac{3e^2}{4}\right), \\
\sum_{n=0}^{\infty} n^3 J_n(ne) J'_n(ne) &= \frac{e}{4(1-e^2)^{9/2}} \left(1 + 3e^2 + \frac{3}{8}e^4\right), \\
\sum_{n=0}^{\infty} n^4 J_n^2(ne) &= \frac{e^2}{4(1-e^2)^{13/2}} \left(1 + \frac{37}{4}e^2 + \frac{59}{8}e^4 + \frac{27}{64}e^6\right), \\
\sum_{n=0}^{\infty} n^4 [J'_n(ne)]^2 &= \frac{1}{4(1-e^2)^{11/2}} \left(1 + \frac{39}{4}e^2 + \frac{79}{8}e^4 + \frac{45}{64}e^6\right).
\end{aligned} \tag{112}$$

## ACKNOWLEDGMENTS

S.P. is supported by the MHRD, Government of India, under the Prime Minister's Research Fellows (PMRF) Scheme, 2020. A.H. would like to thank MHRD, Government of India for research fellowship.

---

[1] Y. B. Zel'dovich and A. G. Polnarev, "Radiation of gravitational waves by a cluster of superdense stars", *Soviet Astr.* **18**, 17 (1974); <https://adsabs.harvard.edu/full/1974SvA....18...17Z>

- [2] L. P. Grishchuk and A. G. Polnarev, “Gravitational wave pulses with ‘velocity coded memory.’,” *Sov. Phys. JETP* **69**, 653-657 (1989)
- [3] V. B. Braginskii and K. S. Thorne, “Gravitational-wave bursts with memory and experimental prospects,” *Nature* **327** 123–125 (1987).
- [4] M. Turner, “Gravitational radiation from point-masses in unbound orbits - Newtonian results” *Ap J.*, **216**, 610-619 (1977).
- [5] M. Turner and C.M. Will, “Post-Newtonian gravitational bremsstrahlung” *Ap J.*, **220**, 1107-1124 (1978).
- [6] R. Epstein, “The Generation of Gravitational Radiation by Escaping Supernova Neutrinos,” *Astrophys. J.* **223**, 1037-1045 (1978) doi:10.1086/156337
- [7] M.S. Turner, “Gravitational radiation from supernova neutrino bursts”, *Nature* **274** 565 ( 1978).
- [8] K. Kotake, K. Sato and K. Takahashi, “Explosion mechanism, neutrino burst, and gravitational wave in core-collapse supernovae,” *Rept. Prog. Phys.* **69**, 971-1144 (2006) doi:10.1088/0034-4885/69/4/R03 [arXiv:astro-ph/0509456 [astro-ph]].
- [9] H. Andresen, B. Müller, E. Müller and H. T. Janka, “Gravitational Wave Signals from 3D Neutrino Hydrodynamics Simulations of Core-Collapse Supernovae,” *Mon. Not. Roy. Astron. Soc.* **468**, no.2, 2032-2051 (2017) doi:10.1093/mnras/stx618 [arXiv:1607.05199 [astro-ph.HE]].
- [10] V. Morozova, D. Radice, A. Burrows and D. Vartanyan, “The gravitational wave signal from core-collapse supernovae,” *Astrophys. J.* **861**, no.1, 10 (2018) doi:10.3847/1538-4357/aac5f1 [arXiv:1801.01914 [astro-ph.HE]].
- [11] D. Vartanyan and A. Burrows, “Gravitational Waves from Neutrino Emission Asymmetries in Core-collapse Supernovae,” *Astrophys. J.* **901**, no.2, 108 (2020) doi:10.3847/1538-4357/abafac [arXiv:2007.07261 [astro-ph.HE]].
- [12] M. Mukhopadhyay, C. Cardona and C. Lunardini, “The neutrino gravitational memory from a core collapse supernova: phenomenology and physics potential,” *JCAP* **07**, 055 (2021) doi:10.1088/1475-7516/2021/07/055 [arXiv:2105.05862 [astro-ph.HE]].
- [13] N. Sago, K. Ioka, T. Nakamura and R. Yamazaki, “Gravitational wave memory of gamma-ray burst jets,” *Phys. Rev. D* **70**, 104012 (2004) doi:10.1103/PhysRevD.70.104012 [arXiv:gr-qc/0405067 [gr-qc]].
- [14] G. Urrutia, F. De Colle, C. Moreno and M. Zanolin, “Gravitational Waves from Long Gamma-Ray Bursts and Supernovae,” [arXiv:2208.00129 [astro-ph.HE]].
- [15] T. Piran, “Jet Gravitational Waves,” [arXiv:2210.02740 [astro-ph.HE]].
- [16] A. C. Jenkins and M. Sakellariadou, “Nonlinear gravitational-wave memory from cusps and kinks on cosmic strings,” *Class. Quant. Grav.* **38**, no.16, 165004 (2021) [arXiv:2102.12487 [gr-qc]].
- [17] D. Christodoulou, “Nonlinear nature of gravitation and gravitational wave experiments,” *Phys. Rev. Lett.* **67** , 1486–1489 (1991).
- [18] A. G. Wiseman and C. M. Will, “Christodoulou’s nonlinear gravitational wave memory: Evaluation in the quadrupole approximation,” *Phys. Rev.* **D44** no. 10, R2945–R2949 (1991) .
- [19] L. Blanchet and T. Damour, “Hereditary effects in gravitational radiation,” *Phys. Rev. D* **46**, 4304-4319 (1992) doi:10.1103/PhysRevD.46.4304
- [20] K. S. Thorne, “Gravitational-wave bursts with memory: The Christodoulou effect,” *Phys. Rev. D* **45**, no.2, 520-524 (1992) doi:10.1103/PhysRevD.45.520
- [21] S. Weinberg, “Photons and Gravitons in Perturbation Theory: Derivation of Maxwell’s and Einstein’s Equations”, *Phys. Rev.* **138**, B988 (1965).
- [22] S. Weinberg, “Infrared photons and gravitons,” *Phys. Rev.* **140**, B516 (1965)
- [23] S. Weinberg, “The Quantum theory of fields. Vol. 1: Foundations,” Cambridge, UK: Univ. Pr. (1995).
- [24] F. Cachazo and A. Strominger, “Evidence for a New Soft Graviton Theorem,” [arXiv:1404.4091 [hep-th]].
- [25] A. Laddha and A. Sen, “Logarithmic Terms in the Soft Expansion in Four Dimensions,” *JHEP* **10**, 056 (2018) doi:10.1007/JHEP10(2018)056 [arXiv:1804.09193 [hep-th]].
- [26] B. Sahoo and A. Sen, “Classical and Quantum Results on Logarithmic Terms in the Soft Theorem in Four Dimensions,” *JHEP* **02**, 086 (2019) [arXiv:1808.03288 [hep-th]].
- [27] A. Laddha and A. Sen, “Observational Signature of the Logarithmic Terms in the Soft Graviton Theorem,” *Phys. Rev. D* **100**, no.2, 024009 (2019) doi:10.1103/PhysRevD.100.024009 [arXiv:1806.01872 [hep-th]].
- [28] D. Kennefick, “Prospects for detecting the Christodoulou memory of gravitational waves from a coalescing compact binary and using it to measure neutron-star radii”, *Phys. Rev. D* **50**, 3587 (1994)
- [29] M. Favata, “Post-Newtonian corrections to the gravitational-wave memory for quasi-circular, inspiralling compact binaries,” *Phys. Rev. D* **80**, 024002 (2009) [arXiv:0812.0069 [gr-qc]].

- [30] M. Favata, *Class. Quant. Grav.* **27** (2010), 084036 doi:10.1088/0264-9381/27/8/084036 [[arXiv:1003.3486](#) [gr-qc]].
- [31] M. Favata, “The Gravitational-wave memory from eccentric binaries,” *Phys. Rev. D* **84**, 124013 (2011) [[arXiv:1108.3121](#) [gr-qc]].
- [32] M. Ebersold, Y. Boetzel, G. Faye, C. K. Mishra, B. R. Iyer and P. Jetzer, “Gravitational-wave amplitudes for compact binaries in eccentric orbits at the third post-Newtonian order: Memory contributions,” *Phys. Rev. D* **100**, no.8, 084043 (2019) doi:10.1103/PhysRevD.100.084043 [[arXiv:1906.06263](#) [gr-qc]].
- [33] J. Samsing, “Eccentric Black Hole Mergers Forming in Globular Clusters,” *Phys. Rev. D* **97**, no.10, 103014 (2018) doi:10.1103/PhysRevD.97.103014 [[arXiv:1711.07452](#) [astro-ph.HE]].
- [34] A. Tucker and C. M. Will, “Residual eccentricity of inspiralling orbits at the gravitational-wave detection threshold: Accurate estimates using post-Newtonian theory,” *Phys. Rev. D* **104**, no.10, 104023 (2021) doi:10.1103/PhysRevD.104.104023 [[arXiv:2108.12210](#) [gr-qc]].
- [35] G. D. Quinlan and S. L. Shapiro, “The collapse of dense star clusters to supermassive black holes - Binaries and gravitational radiation”, *Astrophys. J.* **321**, 199 (1987).
- [36] R. M. O’Leary, B. Kocsis and A. Loeb, “Gravitational waves from scattering of stellar-mass black holes in galactic nuclei,” *Mon. Not. Roy. Astron. Soc.* **395**, no.4, 2127-2146 (2009) doi:10.1111/j.1365-2966.2009.14653.x [[arXiv:0807.2638](#) [astro-ph]].
- [37] D. C. Heggie, “Binary evolution in stellar dynamics”, *Mon. Not. R. Astron. Soc.* **173**, 729 (1975).
- [38] Y. Kozai, “Secular perturbations of asteroids with high inclination and eccentricity”, *Astron. J.* **67**, 591 (1962).
- [39] M. L. Lidov, “The evolution of orbits of artificial satellites of planets under the action of gravitational perturbations of external bodies”, *Planetary Sp. Sci.* **9**, 719 (1962)
- [40] L. Wen, “On the eccentricity distribution of coalescing black hole binaries driven by the Kozai mechanism in globular clusters,” *Astrophys. J.* **598**, 419-430 (2003) doi:10.1086/378794 [[arXiv:astro-ph/0211492](#) [astro-ph]].
- [41] S. Naoz, B. Kocsis, A. Loeb and N. Yunes, “Resonant Post-Newtonian Eccentricity Excitation in Hierarchical Three-body Systems,” *Astrophys. J.* **773**, 187 (2013) doi:10.1088/0004-637X/773/2/187 [[arXiv:1206.4316](#) [astro-ph.SR]].
- [42] M. Favata, C. Kim, K. G. Arun, J. Kim and H. W. Lee, “Constraining the orbital eccentricity of inspiralling compact binary systems with Advanced LIGO,” *Phys. Rev. D* **105**, no.2, 023003 (2022) doi:10.1103/PhysRevD.105.023003 [[arXiv:2108.05861](#) [gr-qc]].
- [43] J. Aasi *et al.* [LIGO Scientific], “Advanced LIGO,” *Class. Quant. Grav.* **32**, 074001 (2015) doi:10.1088/0264-9381/32/7/074001 [[arXiv:1411.4547](#) [gr-qc]].
- [44] F. Acernese *et al.* [VIRGO], “Advanced Virgo: a second-generation interferometric gravitational wave detector,” *Class. Quant. Grav.* **32**, no.2, 024001 (2015) doi:10.1088/0264-9381/32/2/024001 [[arXiv:1408.3978](#) [gr-qc]].
- [45] H. Yu, D. Martynov, S. Vitale, M. Evans, D. Shoemaker, B. Barr, G. Hammond, S. Hild, J. Hough and S. Huttner, *et al.* “Prospects for detecting gravitational waves at 5 Hz with ground-based detectors,” *Phys. Rev. Lett.* **120**, no.14, 141102 (2018) doi:10.1103/PhysRevLett.120.141102 [[arXiv:1712.05417](#) [astro-ph.IM]].
- [46] H. Yang and D. Martynov, “Testing Gravitational Memory Generation with Compact Binary Mergers,” *Phys. Rev. Lett.* **121**, no.7, 071102 (2018) doi:10.1103/PhysRevLett.121.071102 [[arXiv:1803.02429](#) [gr-qc]].
- [47] C. Talbot, E. Thrane, P. D. Lasky and F. Lin, “Gravitational-wave memory: waveforms and phenomenology,” *Phys. Rev. D* **98**, no.6, 064031 (2018) [[arXiv:1807.00990](#) [astro-ph.HE]].
- [48] A. D. Johnson, S. J. Kapadia, A. Osborne, A. Hixon and D. Kennefick, “Prospects of detecting the nonlinear gravitational wave memory,” *Phys. Rev. D* **99**, no.4, 044045 (2019) doi:10.1103/PhysRevD.99.044045 [[arXiv:1810.09563](#) [gr-qc]].
- [49] K. G. Arun *et al.* [LISA], “New horizons for fundamental physics with LISA,” *Living Rev. Rel.* **25**, no.1, 4 (2022) doi:10.1007/s41114-022-00036-9 [[arXiv:2205.01597](#) [gr-qc]].
- [50] B. P. Abbott *et al.* [LIGO Scientific], “Exploring the Sensitivity of Next Generation Gravitational Wave Detectors,” *Class. Quant. Grav.* **34**, no.4, 044001 (2017) doi:10.1088/1361-6382/aa51f4 [[arXiv:1607.08697](#) [astro-ph.IM]].
- [51] S. Hild *et al.*, “Sensitivity studies for third-generation gravitational wave observatories”, *Class. Quant. Grav.* **28**, 094013 (2011).
- [52] P. D. Lasky, E. Thrane, Y. Levin, J. Blackman and Y. Chen, “Detecting gravitational-wave memory with LIGO: implications of GW150914,” *Phys. Rev. Lett.* **117**, no.6, 061102 (2016) doi:10.1103/PhysRevLett.117.061102 [[arXiv:1605.01415](#) [astro-ph.HE]].
- [53] M. Maiorano, F. De Paolis and A. A. Nucita, “Principles of Gravitational-Wave Detection with Pulsar Timing Arrays,” *Symmetry* **13**, 2418 (2021) doi:10.3390/sym13122418 [[arXiv:2112.08064](#) [astro-ph.GA]].



- [54] K. Aggarwal *et al.* [NANOGrav], “The NANOGrav 11 yr Data Set: Limits on Gravitational Wave Memory,” *Astrophys. J.* **889**, 38 (2020) doi:10.3847/1538-4357/ab6083 [arXiv:1911.08488 [astro-ph.HE]].
- [55] S. Burke-Spolaor, S. R. Taylor, M. Charisi, T. Dolch, J. S. Hazboun, A. M. Holgado, L. Z. Kelley, T. J. W. Lazio, D. R. Madison and N. McManis, *et al.* “The Astrophysics of Nanohertz Gravitational Waves,” *Astron. Astrophys. Rev.* **27**, no.1, 5 (2019) doi:10.1007/s00159-019-0115-7 [arXiv:1811.08826 [astro-ph.HE]].
- [56] P. A. Rosado, A. Sesana and J. Gair, “Expected properties of the first gravitational wave signal detected with pulsar timing arrays,” *Mon. Not. Roy. Astron. Soc.* **451**, no.3, 2417-2433 (2015) doi:10.1093/mnras/stv1098 [arXiv:1503.04803 [astro-ph.HE]].
- [57] M. S. Pshirkov, D. Baskaran and K. A. Postnov, “Observing gravitational wave bursts in pulsar timing measurements,” *Mon. Not. Roy. Astron. Soc.* **402**, 417 (2010) doi:10.1111/j.1365-2966.2009.15887.x [arXiv:0909.0742 [astro-ph.CO]].
- [58] S. Capozziello, M. De Laurentis, F. De Paolis, G. Ingrosso and A. Nucita, “Gravitational waves from hyperbolic encounters,” *Mod. Phys. Lett. A* **23**, 99-107 (2008) doi:10.1142/S0217732308026236 [arXiv:0801.0122 [gr-qc]].
- [59] L. De Vittori, P. Jetzer and A. Klein, “Gravitational wave energy spectrum of hyperbolic encounters,” *Phys. Rev. D* **86**, 044017 (2012) [arXiv:1207.5359 [gr-qc]].
- [60] L. De Vittori, A. Gopakumar, A. Gupta and P. Jetzer, “Gravitational waves from spinning compact binaries in hyperbolic orbits,” *Phys. Rev. D* **90**, no.12, 124066 (2014) doi:10.1103/PhysRevD.90.124066 [arXiv:1410.6311 [gr-qc]].
- [61] J. García-Bellido and S. Nesseris, “Gravitational wave energy emission and detection rates of Primordial Black Hole hyperbolic encounters,” *Phys. Dark Univ.* **21**, 61-69 (2018) doi:10.1016/j.dark.2018.06.001 [arXiv:1711.09702 [astro-ph.HE]].
- [62] J. Garcia-Bellido and S. Nesseris, “Gravitational wave bursts from Primordial Black Hole hyperbolic encounters,” *Phys. Dark Univ.* **18**, 123-126 (2017) doi:10.1016/j.dark.2017.10.002 [arXiv:1706.02111 [astro-ph.CO]].
- [63] M. Gröbner, P. Jetzer, M. Haney, S. Tiwari and W. Ishibashi, “A note on the gravitational wave energy spectrum of parabolic and hyperbolic encounters,” *Class. Quant. Grav.* **37**, no.6, 067002 (2020) doi:10.1088/1361-6382/ab6be2 [arXiv:2001.05187 [gr-qc]].
- [64] G. Cho, A. Gopakumar, M. Haney and H. M. Lee, “Gravitational waves from compact binaries in post-Newtonian accurate hyperbolic orbits,” *Phys. Rev. D* **98**, no.2, 024039 (2018) doi:10.1103/PhysRevD.98.024039 [arXiv:1807.02380 [gr-qc]].
- [65] S. Mukherjee, S. Mitra and S. Chatterjee, “Gravitational wave observatories may be able to detect hyperbolic encounters of black holes,” *Mon. Not. Roy. Astron. Soc.* **508**, no.4, 5064-5073 (2021) doi:10.1093/mnras/stab2721 [arXiv:2010.00916 [gr-qc]].
- [66] E. Codazzo, M. Di Giovanni, J. Harms, M. Dall’Amico and M. Mapelli, “Study on the detectability of gravitational radiation from single-binary encounters between black holes in nuclear star cluster: the case of hyperbolic flybys,” [arXiv:2207.01326 [astro-ph.HE]].
- [67] J. García-Bellido, S. Jaraba and S. Kuroyanagi, “The stochastic gravitational wave background from close hyperbolic encounters of primordial black holes in dense clusters,” *Phys. Dark Univ.* **36**, 101009 (2022) doi:10.1016/j.dark.2022.101009 [arXiv:2109.11376 [gr-qc]].
- [68] D. Bini and A. Geralico, “Frequency domain analysis of the gravitational wave energy loss in hyperbolic encounters,” *Phys. Rev. D* **104**, no.10, 104019 (2021) doi:10.1103/PhysRevD.104.104019 [arXiv:2108.02472 [gr-qc]].
- [69] S. Khan, S. Husa, M. Hannam, F. Ohme, M. Pürrer, X. Jiménez Forteza and A. Bohé, “Frequency-domain gravitational waves from nonprecessing black-hole binaries. II. A phenomenological model for the advanced detector era,” *Phys. Rev. D* **93**, no.4, 044007 (2016) doi:10.1103/PhysRevD.93.044007 [arXiv:1508.07253 [gr-qc]].
- [70] C. García-Quirós, M. Colleoni, S. Husa, H. Estellés, G. Pratten, A. Ramos-Buades, M. Mateu-Lucena and R. Jaume, “Multimode frequency-domain model for the gravitational wave signal from nonprecessing black-hole binaries,” *Phys. Rev. D* **102**, no.6, 064002 (2020) [arXiv:2001.10914 [gr-qc]].
- [71] S. Mohanty and P. Kumar Panda, “Particle physics bounds from the Hulse-Taylor binary,” *Phys. Rev. D* **53**, 5723-5726 (1996) doi:10.1103/PhysRevD.53.5723 [arXiv:hep-ph/9403205 [hep-ph]].
- [72] T. Kumar Poddar, S. Mohanty and S. Jana, “Constraints on ultralight axions from compact binary systems,” *Phys. Rev. D* **101**, no.8, 083007 (2020) [arXiv:1906.00666 [hep-ph]].
- [73] T. Kumar Poddar, S. Mohanty and S. Jana, “Vector gauge boson radiation from compact binary systems in a gauged  $L_\mu - L_\tau$  scenario,” *Phys. Rev. D* **100**, no.12, 123023 (2019) doi:10.1103/PhysRevD.100.123023 [arXiv:1908.09732 [hep-ph]].
- [74] T. K. Poddar, S. Mohanty and S. Jana, “Gravitational radiation from binary systems in massive graviton theories,” *JCAP* **03**, 019 (2022) doi:10.1088/1475-7516/2022/03/019 [arXiv:2105.13335 [gr-qc]].

- [75] A. Addazi, M. Bianchi and G. Veneziano, “Glimpses of black hole formation/evaporation in highly inelastic, ultra-planckian string collisions,” JHEP **02**, 111 (2017) [[arXiv:1611.03643](#) [hep-th]].
- [76] A. Addazi, M. Bianchi and G. Veneziano, “Soft gravitational radiation from ultra-relativistic collisions at sub- and sub-sub-leading order,” JHEP **05**, 050 (2019) [[arXiv:1901.10986](#) [hep-th]].
- [77] A. Addazi and K. A. Zeng, “Soft gravitational radiation from multi-body collisions,” JHEP **11**, 193 (2021) [[arXiv:2110.01194](#) [hep-th]].
- [78] A. Tolish and R. M. Wald, Phys. Rev. D **89** (2014) no.6, 064008 doi:10.1103/PhysRevD.89.064008 [[arXiv:1401.5831](#) [gr-qc]].
- [79] A. Tolish and R. M. Wald, “Cosmological memory effect,” Phys. Rev. D **94**, no.4, 044009 (2016) [[arXiv:1606.04894](#) [gr-qc]].
- [80] A. Kehagias and A. Riotto, “BMS in Cosmology,” JCAP **05**, 059 (2016) [[arXiv:1602.02653](#) [hep-th]].
- [81] P. C. Peters and J. Mathews, “Gravitational radiation from point masses in a Keplerian orbit,” Phys. Rev. **131**, 435-439 (1963)
- [82] P. C. Peters, “Gravitational Radiation and the Motion of Two Point Masses,” Phys. Rev. **136**, B1224-B1232 (1964)
- [83] L. Blanchet, T. Damour, G. Esposito-Farese and B. R. Iyer, “Gravitational radiation from inspiralling compact binaries completed at the third post-Newtonian order,” Phys. Rev. Lett. **93**, 091101 (2004) doi:10.1103/PhysRevLett.93.091101 [[arXiv:gr-qc/0406012](#) [gr-qc]].
- [84] K. G. Arun, L. Blanchet, B. R. Iyer and M. S. S. Qusailah, “Inspiralling compact binaries in quasi-elliptical orbits: The Complete 3PN energy flux,” Phys. Rev. D **77**, 064035 (2008) doi:10.1103/PhysRevD.77.064035 [[arXiv:0711.0302](#) [gr-qc]].
- [85] K. G. Arun, L. Blanchet, B. R. Iyer and M. S. S. Qusailah, “Tail effects in the 3PN gravitational wave energy flux of compact binaries in quasi-elliptical orbits,” Phys. Rev. D **77**, 064034 (2008) doi:10.1103/PhysRevD.77.064034 [[arXiv:0711.0250](#) [gr-qc]].