

ON THE ELLIPTICAL RANGE THEOREMS FOR THE DAVIS–WIELANDT SHELL, THE NUMERICAL RANGE, AND THE CONFORMAL RANGE

GYULA LAKOS

ABSTRACT. The conformal range, which is a horizontal projection of the Davis–Wielandt shell, can be considered as the hyperbolic version of the numerical range. Here we explain (the analogue of) the elliptical range theorem of 2×2 complex matrices for the conformal range. In that course, comparison to the Davis–Wielandt shell and the numerical range is made.

INTRODUCTION

The conformal range, which is a horizontal projection of the Davis–Wielandt shell, can be considered as the hyperbolic version of the numerical range. Here we explain (the analogue of) the elliptical range theorem of 2×2 complex matrices for the conformal range. In that course, comparison to the Davis–Wielandt shell and the numerical range is also made. We give a (very elementary) presentation of how the conformal range acts as an analogue of the numerical range in the simple case of 2×2 complex matrices.

The Davis–Wielandt shell, the numerical range, and the conformal range. Assume that \mathfrak{H} is a Hilbert space, with interior product $\langle \cdot, \cdot \rangle$ (linear in the first variable, skew-linear in the second variable); $\|\mathbf{x}\|_2 = \sqrt{\langle \mathbf{x}, \mathbf{x} \rangle}$. If A is a linear operator on \mathfrak{H} , then its Davis–Wielandt shell is defined as

$$DW_{\text{CKB(P)}}(A) = \left\{ \left(\frac{\text{Re}\langle \mathbf{y}, \mathbf{x} \rangle}{\|\mathbf{x}\|_2^2}, \frac{\text{Im}\langle \mathbf{y}, \mathbf{x} \rangle}{\|\mathbf{x}\|_2^2}, \frac{\|\mathbf{y}\|_2^2}{\|\mathbf{x}\|_2^2} \right) : A\mathbf{x} = \mathbf{y}, \mathbf{x} \neq 0 \right\} \subset \overline{H}_{\text{CKB(P)}}^3,$$

or

$$DW_{\text{CKB}}(A) = \left\{ \left(\frac{2\text{Re}\langle \mathbf{y}, \mathbf{x} \rangle}{\|\mathbf{y}\|_2^2 + \|\mathbf{x}\|_2^2}, \frac{2\text{Im}\langle \mathbf{y}, \mathbf{x} \rangle}{\|\mathbf{y}\|_2^2 + \|\mathbf{x}\|_2^2}, \frac{\|\mathbf{y}\|_2^2 - \|\mathbf{x}\|_2^2}{\|\mathbf{y}\|_2^2 + \|\mathbf{x}\|_2^2} \right) : A\mathbf{x} = \mathbf{y}, \mathbf{x} \neq 0 \right\} \subset \overline{H}_{\text{CKB}}^3.$$

These sets are interpreted as subsets of the asymptotically closed Cayley–Klein–Beltrami models (ordinary or parabolic) of the hyperbolic space. Between the CKB and CKB(P) models, there are (for us) canonical correspondences given by

$$(1) \quad \begin{bmatrix} x_{\text{CKB}} \\ y_{\text{CKB}} \\ z_{\text{CKB}} \\ 1 \end{bmatrix} = \frac{1}{z_{\text{CKB(P)}} + 1} \begin{bmatrix} x_{\text{CKB(P)}} \\ y_{\text{CKB(P)}} \\ z_{\text{CKB(P)}} - 1 \\ z_{\text{CKB(P)}} + 1 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} x_{\text{CKB(P)}} \\ y_{\text{CKB(P)}} \\ z_{\text{CKB(P)}} \\ 1 \end{bmatrix} = \frac{1}{1 - z_{\text{CKB}}} \begin{bmatrix} x_{\text{CKB}} \\ y_{\text{CKB}} \\ 1 + z_{\text{CKB}} \\ 1 - z_{\text{CKB}} \end{bmatrix};$$

with $(0, 0, 1)_{\text{CKB}}$ corresponding to $\infty_{\text{CKB(P)}}$ (although these latter points do not appear in the shell for linear operators). Other models of the (asymptotically closed) hyperbolic space can also be used for the shell, however, for the purposes of linear algebra, the projective models CKB and CKB(P) are particularly suitable.

2020 *Mathematics Subject Classification*. Primary: 15A60, Secondary: 51M10.

Key words and phrases. Davis–Wielandt shell, numerical range, conformal range of operators.

The real Davis–Wielandt shell $DW_{\text{CKB(P)}}^{\mathbb{R}}(A)$ or $DW_{\text{CKB}}^{\mathbb{R}}(A)$ can be defined in the same manner but with the second coordinate omitted. This latter process can be interpreted as an orthogonal projection in hyperbolic geometry. Thus, the real Davis–Wielandt shell yields a subset of the asymptotically closed hyperbolic plane. In turn, the real Davis–Wielandt shell can also be defined for linear operators acting on real Hilbert spaces. In what follows, we use the term ‘conformal range’ for the real Davis–Wielandt shell. (In [24], the conformal range was defined originally as $\text{CR}(A) \equiv DW_{\text{PH}}^{\mathbb{R}}(A)$, the real Davis–Wielandt shell in the Poincaré half-plane model.) On the other hand, the vertical projection of $DW_{\text{CKB(P)}}^{\mathbb{R}}(A)$ is the well-known numerical range $W(A)$. (It is obtained from $DW_{\text{CKB}}^{\mathbb{R}}(A)$ via a central projection from $(0, 0, 1)_{\text{CKB}}$.) Thus, the conformal range can be considered as a hyperbolic analogue of the numerical range, which is indeed confirmed by its generally quite similar behaviour, cf. [24].

The setting of the elliptical range theorems. The simplest yet not entirely trivial case for the numerical and conformal ranges is when A is a complex 2×2 matrix. In that case, the numerical range is described geometrically by the well-known elliptical range theorem. (See its discussion later.) One may wonder whether a similar description is valid with respect to the conformal range. Indeed, the answer is affirmative. The only slightly unusual feature is that we have to consider conics on the hyperbolic plane. Nevertheless, their study predates the one of numerical range:

The systematic study of hyperbolic conics was apparently started by Story [35] (1882). By 1909, Coolidge [3] already contains a very nice compendium on hyperbolic conics based on expositions of Killing [21], Liebmann [28], and, especially, D’Ovidio [9], [10], [11] (1891). A bit more refined classification is considered in Barbarin [1] (1901). Massau [30] (1905) also writes on the subject. A very detailed study of the conics is conducted in Vörös [38] / [39] (1909/10), [40]. Thus, it can be said that by 1910 the hyperbolic conics were rather well-explored; although not all of the sources cited above can be considered equally accessible. Numerical aspects of the classification are discussed further in Fladt [12], [13], [14]. For the sake of completeness, we also mention Molnár [31], where the conics are studied from an other viewpoint. For those less familiar with hyperbolic geometry, an introduction like of Izmistiev [19] might be a good starting point. Ultimately, the point is that h -conics are simplest to be defined as conics of the projective models (i. e. of CKB(P) or CKB). Then, they enjoy similar synthetic presentations and properties as in the Euclidean case. However, the general information above is primarily just to straighten the record: The reason is that beyond lineal objects and cycles we will consider only ordinary h -ellipses and h -elliptic parabolas; and the behaviour of those will be clear from the examples.

It is easy to prove that for a complex 2×2 matrix A , the Davis–Wielandt shell $DW_{\text{CKB(P)}}^{\mathbb{R}}(A)$ or $DW_{\text{CKB}}^{\mathbb{R}}(A)$ is a possibly degenerate ellipsoid. This is a kind of elliptical range theorem itself. This immediately implies that the numerical and conformal ranges yield possibly degenerate elliptical disks, hence disks of possibly degenerate h -ellipses. Thus, in this most qualitative form, the elliptical range theorems are quite trivial. Therefore, any interesting statement should involve more precise (in our case: metric and focal) information. The elliptical range theorem for the conformal range is slightly more complicated than for the numerical range. This can be attributed primarily to the fact that asymptotic points can act as focal points. Otherwise, the analogy is close.

Due to the elementary nature of the topic, I have considered writing a much shorter exposition for the conformal range, which is possible (cf. Remark 5.44). However, I

came to the opinion that proper justice is made only by a more comprehensive view. The opportunity was also taken up to illustrate certain computational methods which may not be the simplest here but prominent in the higher dimensional case. Hence, there are certain redundancies resulted from alternative approaches. Although our paper is not self-contained (we refer to the relevant parts of [24] several times), we also include certain elementary facts (cf. Appendix A) for greater accessibility. Nevertheless, most of our statements are formulas, which are easy to check individually.

On the layout of this paper. After some reviews and preparations we will consider the quantitative properties of the Davis–Wielandt shell, the numerical range, and the conformal range for complex 2×2 matrices. This will be done in Sections 2, 3, and 5, respectively. Although the development allows various selections, the paper itself is intended to be read linearly. The prerequisites to this paper are contained in Sections 2, 5, 7 and Appendices A, B of [24]. The reader will see that any essential references to [24] can be eliminated; that, however, might not leave the simplest or most insightful arguments. Thus, at least a passing familiarity with the relevant parts of [24] is needed.

On terminology and notation. In general, we will adopt the terminology from [24]. We will often consider natural transformations of the hyperbolic plane and space. Due to their nature, depending on their appearance, the terms ‘ h -collineations’, ‘ h -isometries’, ‘ h -congruences’, ‘conformal transformations’, and, ‘Möbius transformations’, ‘fractional linear transformations’ (rather synonymous in the orientation-preserving case) are used quite ecclectically. If we do not specify that in what model we consider the Davis–Wielandt shell of A , then we write $DW_*(A)$ where $*$ might mean CKB, CKB(P), or PH. Similar comment applies to the conformal range $DW_*^{\mathbb{R}}(A)$. A possibly asymptotic hyperbolic point \mathbf{x} is understood as a compatible collection of \mathbf{x}_* ’s; $\mathbf{x}_{\text{CKB(P)}}$ is the tuple of its coordinates in the CKB(P) model, etc. If a statement is meant in synthetic hyperbolic sense, then the $*$ ’s can be omitted altogether. The the euclidean (or affine) segment between a and b is denoted by $[a, b]_e$.

The ‘classical adjoint’ or adjugate matrix of \mathbf{M} will be denoted $\text{adj } \mathbf{M}$. The connection to the inverse will be used several times. The contravariant multiplicative property

$$(2) \quad \text{adj}(\mathbf{M}_1\mathbf{M}_2) = (\text{adj } \mathbf{M}_2)(\text{adj } \mathbf{M}_1)$$

is taken granted without special reference. The involutivity property

$$(3) \quad \text{adj}(\text{adj } \mathbf{M}) = \mathbf{M} \quad \text{for } 2 \times 2 \text{ matrices}$$

and the property

$$(4) \quad \text{adj}(\text{adj } \mathbf{M}) = (\det \mathbf{M}) \cdot \mathbf{M} \quad \text{for } 3 \times 3 \text{ matrices}$$

will also be understood. The similarity (i. e. conjugacy) of matrices \mathbf{M}_1 and \mathbf{M}_2 will be denoted by $\mathbf{M}_1 \sim \mathbf{M}_2$. If \mathbf{M}_1 and \mathbf{M}_2 are proportional to each other by a nonzero scalar, then it is denoted by $\mathbf{M}_1 \cong \mathbf{M}_2$.

If $a > 0$, then $\frac{a}{0} = +\infty$. However, $\frac{0}{0} = \pm\infty$, for which, in formulas, special evaluation rules apply, which will be indicated; typically, the values 0, 1, or ∞ may apply.

Line y of formula (X) will be referred as (X/y) .

Acknowledgements. The author thanks Ákos G. Horváth for help related to hyperbolic conics.

1. NORMALITY OF 2×2 MATRICES AND RELATED CONCEPTS

Here all matrices will be 2×2 complex matrices.

1.A. Spectral type (review).

As in [24], we use the notation

$$(5) \quad D_A = \det \left(A - \frac{\operatorname{tr} A}{2} \operatorname{Id}_2 \right) = (\det A) - \frac{(\operatorname{tr} A)^2}{4}.$$

It is essentially the discriminant of A , as the eigenvalues of A are $\frac{1}{2} \operatorname{tr} A \pm \sqrt{-D_A}$.

In the general complex case, there are two main categories for the spectral type: parabolic ($D_A = 0$) and non-parabolic ($D_A \neq 0$).

1.B. The metric discriminant.

In terms of trace,

$$-D_A = \frac{1}{2} \operatorname{tr} \left(\left(A - \frac{\operatorname{tr} A}{2} \operatorname{Id} \right)^2 \right) = \frac{\operatorname{tr} A^2}{2} - \frac{(\operatorname{tr} A)^2}{4}.$$

Thus, a similar notion is given by the metric discriminant

$$(6) \quad U_A = \frac{1}{2} \operatorname{tr} \left(\left(A - \frac{\operatorname{tr} A}{2} \operatorname{Id} \right)^* \left(A - \frac{\operatorname{tr} A}{2} \operatorname{Id} \right) \right) = \frac{\operatorname{tr} A^* A}{2} - \frac{|\operatorname{tr} A|^2}{4}.$$

(We will see, however, that U_A is not truly a counterpart of $-D_A$, but of $|D_A|$.)

1.C. Canonical triangular form.

It is elementary, but useful to keep in mind that any 2×2 complex matrix A can be brought by unitary conjugation into form

$$(7) \quad \begin{bmatrix} \lambda_1 & t \\ 0 & \lambda_2 \end{bmatrix}$$

where $t \geq 0$. This form is unique, up to the order of the eigenvalues λ_1 and λ_2 .

In the situation above,

$$(8) \quad U_A = \frac{t^2}{2} + \left| \frac{\lambda_1 - \lambda_2}{2} \right|^2, \quad \text{and} \quad D_A = - \left(\frac{\lambda_1 - \lambda_2}{2} \right)^2.$$

Conversely,

$$(9) \quad \begin{aligned} &\lambda_1, \lambda_2 \text{ are the two roots of} \\ &\lambda^2 - (\operatorname{tr} A)\lambda + \det A = 0; \end{aligned}$$

and the ‘‘canonical off-diagonal quantity’’, as (8) shows, is

$$(10) \quad t = \sqrt{2(U_A - |D_A|)}.$$

(One familiar with the elliptical range theorem can recognize that U_A and $|D_A|$ are quantities to play role there. In particular, (10) is the length of the minor axis.)

1.D. The five data.

For a 2×2 complex matrix A , the quantities

$$(11) \quad \operatorname{Re} \operatorname{tr} A, \operatorname{Im} \operatorname{tr} A, \operatorname{Re} \det A, \operatorname{Im} \det A, \operatorname{tr}(A^* A)$$

will be called as the ‘five data’. Note that D_A and U_A can be expressed by the five data.

Lemma 1.1. *The five data (11) characterizes the complex 2×2 matrix A up to conjugation by unitary matrices.*

Proof. It is trivial that conjugation by unitary matrices does not change the five data. Conversely, consider the possible canonical triangular form for A as in (7). Then (9) and (10) imply that the five data determines the canonical form. \square

1.E. Normality.

Lemma 1.2. *Suppose that A is a 2×2 complex matrix. Then*

$$U_A \geq |D_A|.$$

If A is normal, then $U_A = |D_A|$. If A is not normal, then $U_A > |D_A|$.

Proof. Using a conjugation by a unitary matrix (which leaves $U_A, |D_A|$ invariant), we can assume that A is of triangular form (7). Then (8) shows the statement. \square

Alternative proof. From (6) we already see that $U_A \geq 0$. By direct computation (as a consequence of $\operatorname{tr}[A, \mathbf{v}]^2 = 8 \left(\left(\frac{1}{2} \operatorname{tr} \left(\left(A - \frac{\operatorname{tr} A}{2} \operatorname{Id}_2 \right) \left(\mathbf{v} - \frac{\operatorname{tr} \mathbf{v}}{2} \operatorname{Id}_2 \right) \right) \right)^2 - D_A D_{\mathbf{v}} \right)$), one finds the identity

$$\operatorname{tr}(A^* A - AA^*)^*(A^* A - AA^*) = 8 \left((U_A)^2 - |D_A|^2 \right).$$

The LHS is the squared Frobenius norm of the commutator $[A^*, A]$. This is nonnegative, and vanishes if and only if A is normal. \square

Remark 1.3. $U_A = -D_A$ if A is self-adjoint, and $U_A = D_A$ if A is skew-adjoint. \triangle

1.F. Invariance for complex Möbius transformations.

Lemma 1.4. *For complex 2×2 matrices A , the ratio*

$$U_A : |D_A|$$

is invariant for complex Möbius transformations (whenever they are applicable to A).

Proof. Indeed, adding a scalar matrix to A does not alter either U_A or $|D_A|$. Regarding the inverse (if applicable),

$$U_{A^{-1}} : |D_{A^{-1}}| \equiv \frac{U_A}{|\det A|^2} : \left| \frac{D_A}{(\det A)^2} \right| = U_A : |D_A|$$

holds. This is already sufficient to establish the complex Möbius invariance property. \square

One can also think that this quantity, or any of its variants, $\frac{U_A}{|D_A|}$, $\frac{|D_A|}{U_A}$, or $\frac{U_A - |D_A|}{U_A + |D_A|}$, measure non-normality in a complex Möbius invariant way. (This foreshadows the hyperbolic geometric interpretation.)

1.G. The canonical representatives (complex Möbius).

Lemma 1.5. *For complex 2×2 matrices up to complex Möbius transformations and unitary conjugations (which, of course, commute with each other) it is sufficient to consider the following representatives:*

$$(12) \quad \mathbf{0}_2 = \begin{bmatrix} 0 & \\ & 0 \end{bmatrix}$$

(parabolic, normal) ;

$$(13) \quad S_0 = \begin{bmatrix} 0 & 1 \\ & 0 \end{bmatrix}$$

(parabolic non-normal);

$$(14) \quad L_t = \begin{bmatrix} 1 & 2t \\ & -1 \end{bmatrix} \quad t \geq 0$$

(non-parabolic case; $t = 0$: normal, $t > 0$: non-normal).

The representatives above are inequivalent.

Proof. It is sufficient to normalize the eigenvalues by applying Möbius transformations, and then achieve normal form by unitary conjugation. By this, we arrive to the parabolic cases

$$(15) \quad \widehat{\mathbf{0}}_t = \begin{bmatrix} 0 & 2t \\ & 0 \end{bmatrix} \quad t \geq 0,$$

and the non-parabolic cases (14). The cases $\widehat{\mathbf{0}}_t$ with $t > 0$ scale into each other linearly (hence, by Möbius transformations). Thus (15) reduces only to (12) and (13). The ratio $U_A : |D_A|$ distinguishes the various representatives, cf. the following example. \square

Example 1.6. (a) Regarding L_t :

$$U_{L_t} = 1 + 2t^2, \quad \text{and} \quad |D_{L_t}| = 1.$$

(b) Regarding S_0 :

$$U_{S_0} = \frac{1}{2}, \quad \text{and} \quad |D_{S_0}| = 0.$$

(c) Regarding $\mathbf{0}_2$:

$$U_{\mathbf{0}_2} = 0, \quad \text{and} \quad |D_{\mathbf{0}_2}| = 0. \quad \diamond$$

Corollary 1.7. *The ratio*

$$U_A : |D_A|$$

is a full invariant of 2×2 complex matrices with respect to equivalence by complex Möbius transformations and unitary conjugation.

Proof. It distinguishes the canonical representatives of Lemma 1.5. \square

1.H. Norms (review).

We also recall, using the terminology of [24],

Lemma 1.8. *Suppose that A is a real or complex 2×2 matrix. Then, for the norm,*

$$\begin{aligned} \|A\|_2 &= \sqrt{\frac{\operatorname{tr}(A^*A)}{2} + \sqrt{\frac{(\operatorname{tr}(A^*A))^2}{4} - |\det A|^2}} \\ &= \frac{\sqrt{\operatorname{tr}(A^*A) + 2|\det A|} + \sqrt{\operatorname{tr}(A^*A) - 2|\det A|}}{2}; \end{aligned}$$

and, for the co-norm,

$$\begin{aligned} \|A\|_2^- &= \|A^{-1}\|_2^{-1} = \sqrt{\frac{\operatorname{tr}(A^*A)}{2} - \sqrt{\frac{(\operatorname{tr}(A^*A))^2}{4} - |\det A|^2}} \\ &= \frac{\sqrt{\operatorname{tr}(A^*A) + 2|\det A|} - \sqrt{\operatorname{tr}(A^*A) - 2|\det A|}}{2}. \end{aligned}$$

In particular,

$$(16) \quad \|A\|_2 \cdot \|A\|_2^- = |\det A|.$$

In case of real matrices, the values are the same for the Hilbert spaces \mathbb{R}^2 and \mathbb{C}^2 . \square

2. THE DAVIS–WIELANDT SHELL OF 2×2 MATRICES

2.A. The qualitative elliptical range theorems for the shell (review).

For an introduction to the Davis–Wielandt shell, see Wielandt [42]; Davis [4], [5], [7]; Li, Poon, Sze [27]; Lin, Spitkovsky, Zhong [29]. (But especially [4] and [5].) A general review of hyperbolic geometry is contained in, for example, Berger [2]. Nevertheless, all the basic facts we need are already mentioned in parts of [24], thus, in practice, we use it as the principal reference. We quote only the following two theorems; for everything else, including the description of the effects of Möbius transformations, we refer to [24].

Theorem 2.1 (Davis). *Suppose that A is a 2-dimensional linear operator (or just linear relation) on a complex Hilbert space. Then $\operatorname{DW}_{\operatorname{CKB}}(A)$ is a possibly degenerate ellipsoid (in Euclidean view).*

Recall that the complex numbers are embedded to the $\operatorname{CKB}(\mathbb{P})$ and CKB models by

$$\iota_{\operatorname{CKB}(\mathbb{P})}(\lambda) = (\operatorname{Re} \lambda, \operatorname{Im} \lambda, |\lambda|^2) \quad \text{and} \quad \iota_{\operatorname{CKB}}(\lambda) = \left(\frac{2 \operatorname{Re} \lambda}{1 + |\lambda|^2}, \frac{2 \operatorname{Im} \lambda}{1 + |\lambda|^2}, \frac{1 - |\lambda|^2}{1 + |\lambda|^2} \right).$$

Theorem 2.2 (Wielandt, Davis). *Suppose that A is a linear operator on a 2-dimensional complex Hilbert space. We have the following possibilities:*

- (i) A has a double eigenvalue λ , and A is normal (thus $A = \lambda \operatorname{Id}$).
Then $\operatorname{DW}_*(A)$ contains only the point $\iota_*(\lambda)$.
- (ii) A has two different eigenvalues $\lambda_1 \neq \lambda_2$, and A is normal.
Then $\operatorname{DW}_*(A)$ is the asymptotically closed h -line connecting $\iota_*(\lambda_1)$ and $\iota_*(\lambda_2)$.
- (iii) A has a double eigenvalue λ , and A is not normal.

Then $DW_*(A)$ is an asymptotically closed h -horosphere with asymptotical point $\iota_*(\lambda)$. In the CKB model this is an ellipsoid, whose equation is linearly generated by the quadratic equation of the unit sphere and the equation of the double plane tangent to unit sphere at $\iota_*(\lambda)$.

(iv) A has two different eigenvalues $\lambda_1 \neq \lambda_2$, and A is not normal.

Then $DW_*(A)$ is the an asymptotically closed h -tube around the h -line connecting $\iota_*(\lambda_1)$ and $\iota_*(\lambda_2)$. In the CKB model this is an ellipsoid, whose equation is linearly generated by the quadratic equation of the unit sphere and the quadratic equation of the union of planes tangent to unit sphere at $\iota_*(\lambda_1)$ and $\iota_*(\lambda_2)$.

Proofs. See Wielandt [41], [42] (mainly the normal case), and Davis [4], [5] (generally); but also see the discussions in [24]. \square

The theorems above are the more or less detailed qualitative descriptions of the Davis–Wielandt shells of 2×2 matrices. The main point is Theorem 2.1 (which is rather simple); then Theorem 2.2 follows by applying geometric symmetry principles (which requires some basic knowledge concerning the congruences of the hyperbolic space).

Let us make two observations regarding the shell.

The first observation is that the shell as a closed compact set (in the CKB(P) or CKB models) is continuous with respect to the Hausdorff topology. This is a simple consequence of its construction as a continuous image of the a sphere. Consequently, limit arguments in the shell apply. (Remark: This observation naturally extends to the numerical and conformal ranges.)

The second observation is that if we obtain an equation for the shell from the pencil construction, say, in the case of a normal non-parabolic matrix, then the resulted equation is not just any line ellipse for the shell but the “infinitesimal tube”, i. e. degenerate elliptic *circular* cylinder. This also applies if we obtain the quadratic form of the shell as a nonzero limit of quadratic forms from non-normal matrices. In the case of normal parabolic matrices similar comment applies with respect to “double tangent planes”. (Remark: This observation does not extend to the numerical and conformal ranges. Although there are natural point circles, they are not necessarily the non-zero limits.)

2.B. The matrix of the shell.

Our general objective is to consider more quantitative descriptions of the Davis–Wielandt shell of 2×2 matrices. We start with the explicit equation. This was found by Lins, Spitkovsky, Zhong [29]. The equation has sort of the nicest shape in the CKB(P) model. We give a several proofs.

Theorem 2.3 (Lins, Spitkovsky, Zhong [29]). *Suppose that A is linear operator on a 2-dimensional Hilbert space. Then $DW_{\text{CKB(P)}}(A)$ is given by the quadratic equation*

$$(17) \quad \begin{bmatrix} x_{\text{CKB(P)}} \\ y_{\text{CKB(P)}} \\ z_{\text{CKB(P)}} \\ 1 \end{bmatrix}^{\top} \mathbf{Q}_{\text{CKB(P)}}(A) \begin{bmatrix} x_{\text{CKB(P)}} \\ y_{\text{CKB(P)}} \\ z_{\text{CKB(P)}} \\ 1 \end{bmatrix} = 0$$

where

$$\begin{aligned}
(18) \quad \mathbf{Q}_{\text{CKB}(\text{P})}(A) &= U_A 2 \underbrace{\begin{bmatrix} 1 & & & \\ & 1 & & \\ & & -\frac{1}{2} & \\ & & & -\frac{1}{2} \end{bmatrix}}_{\mathbf{Q}_{\text{CKB}(\text{P})}^{\text{bas}}(A):=} + \\
&+ \underbrace{\begin{bmatrix} \frac{|\text{tr } A|^2}{2} + 2 \text{Re det } A & 2 \text{Im det } A & -\text{Re tr } A & -\text{Re}((\text{det } A)\overline{(\text{tr } A)}) \\ 2 \text{Im det } A & \frac{|\text{tr } A|^2}{2} - 2 \text{Re det } A & -\text{Im tr } A & -\text{Im}((\text{det } A)\overline{(\text{tr } A)}) \\ -\text{Re tr } A & -\text{Im tr } A & 1 & \frac{|\text{tr } A|^2}{4} \\ -\text{Re}((\text{det } A)\overline{(\text{tr } A)}) & -\text{Im}((\text{det } A)\overline{(\text{tr } A)}) & \frac{|\text{tr } A|^2}{4} & |\text{det } A|^2 \end{bmatrix}}_{\mathbf{Q}_{\text{CKB}(\text{P})}^{\text{spec}}(A):=} = \\
&= \begin{bmatrix} (\text{tr } A^* A) + 2 \text{Re det } A & 2 \text{Im det } A & -\text{Re tr } A & -\text{Re}((\text{det } A)\overline{(\text{tr } A)}) \\ 2 \text{Im det } A & (\text{tr } A^* A) - 2 \text{Re det } A & -\text{Im tr } A & -\text{Im}((\text{det } A)\overline{(\text{tr } A)}) \\ -\text{Re tr } A & -\text{Im tr } A & 1 & \frac{|\text{tr } A|^2 - \text{tr}(A^* A)}{2} \\ -\text{Re}((\text{det } A)\overline{(\text{tr } A)}) & -\text{Im}((\text{det } A)\overline{(\text{tr } A)}) & \frac{|\text{tr } A|^2 - \text{tr}(A^* A)}{2} & |\text{det } A|^2 \end{bmatrix}.
\end{aligned}$$

Here

$$\det \mathbf{Q}_{\text{CKB}(\text{P})}(A) = -4 \left((U_A)^2 - |D_A|^2 \right)^2.$$

In the non-normal cases, the equation yields ellipsoids in the model space. In the normal non-parabolic case the equation yields a line ellipsoid (degenerate elliptic cylinder). In the normal parabolic case, the equation yields a double plane tangent at the corresponding asymptotic point. (Thus in the normal case one should restrict the solutions to the model space $\text{CKB}(\text{P})$.)

In the non-normal cases, the interior points $(x_{\text{CKB}(\text{P})}, y_{\text{CKB}(\text{P})}, z_{\text{CKB}(\text{P})})$ of the ellipses are characterized by replacing ' $= 0$ ' with ' < 0 ' in (17).

Remark. Considering the canonical correspondences between the various models (cf. [24]), we find that in (17), replacing

$$\begin{bmatrix} x_{\text{CKB}(\text{P})} \\ y_{\text{CKB}(\text{P})} \\ z_{\text{CKB}(\text{P})} \\ 1 \end{bmatrix} \text{ by } \begin{bmatrix} x_{\text{CKB}} \\ y_{\text{CKB}} \\ 1 + z_{\text{CKB}} \\ 1 - z_{\text{CKB}} \end{bmatrix}, \begin{bmatrix} x_{\text{P}} \\ y_{\text{P}} \\ \frac{(x_{\text{P}})^2 + (y_{\text{P}})^2 + (z_{\text{P}} + 1)^2}{2} \\ \frac{(x_{\text{P}})^2 + (y_{\text{P}})^2 + (z_{\text{P}} - 1)^2}{2} \end{bmatrix}, \begin{bmatrix} x_{\text{PH}} \\ y_{\text{PH}} \\ (x_{\text{PH}})^2 + (y_{\text{PH}})^2 + (z_{\text{PH}})^2 \\ 1 \end{bmatrix}$$

one obtains the corresponding equations in CKB , P , PH , respectively.

In the case of the CKB model, the transition from $\text{CKB}(\text{P})$ is projective / linear (cf. (1)). Thus another possible viewpoint is that

$$(19) \quad \begin{bmatrix} x_{\text{CKB}} \\ y_{\text{CKB}} \\ 1 + z_{\text{CKB}} \\ 1 - z_{\text{CKB}} \end{bmatrix}^{\top} \mathbf{Q}_{\text{CKB}(\text{P})}(A) \begin{bmatrix} x_{\text{CKB}} \\ y_{\text{CKB}} \\ 1 + z_{\text{CKB}} \\ 1 - z_{\text{CKB}} \end{bmatrix} \equiv \begin{bmatrix} x_{\text{CKB}} \\ y_{\text{CKB}} \\ z_{\text{CKB}} \\ 1 \end{bmatrix}^{\top} \mathbf{Q}_{\text{CKB}}(A) \begin{bmatrix} x_{\text{CKB}} \\ y_{\text{CKB}} \\ z_{\text{CKB}} \\ 1 \end{bmatrix};$$

where

$$(20) \quad \mathbf{Q}_{\text{CKB}}(A) = \begin{bmatrix} 1 & & & \\ & 1 & & \\ & & 1 & 1 \\ & & -1 & 1 \end{bmatrix}^\top \mathbf{Q}_{\text{CKB(P)}}(A) \begin{bmatrix} 1 & & & \\ & 1 & & \\ & & 1 & 1 \\ & & -1 & 1 \end{bmatrix}$$

is obtained canonically. Similarly, we adopt the scheme of (19)/(20) in order to transcribe $\mathbf{Q}_{\text{CKB(P)}}^{\text{spec}}(A)$ to $\mathbf{Q}_{\text{CKB}}^{\text{spec}}(A)$. Due to projectivity / linearity, similar comments apply concerning the shape of solution sets in the CKB model as in the CKB(P) model.

There are some satellite statements which are worth mentioning in connection to Theorem 2.3. For this reason, we set

$$(21) \quad \mathbf{Q}_{\text{CKB(P)}}^0 = \begin{bmatrix} 1 & & & \\ & 1 & & \\ & & -\frac{1}{2} & \\ & -\frac{1}{2} & & \end{bmatrix}, \quad \text{and} \quad \mathbf{Q}_{\text{CKB}}^0 = \begin{bmatrix} 1 & & & \\ & 1 & & \\ & & 1 & \\ & & & -1 \end{bmatrix}.$$

These are also connected as in (19)/(20). Consequently, $(\mathbf{Q}_{\text{CKB(P)}}^0)^{-1} \mathbf{Q}_{\text{CKB(P)}}(A)$, and $(\mathbf{Q}_{\text{CKB}}^0)^{-1} \mathbf{Q}_{\text{CKB}}(A)$ are conjugates of each other.

Theorem 2.4. *The eigenvalues of $(\mathbf{Q}_{\text{CKB(P)}}^0)^{-1} \mathbf{Q}_{\text{CKB(P)}}(A) \sim (\mathbf{Q}_{\text{CKB}}^0)^{-1} \mathbf{Q}_{\text{CKB}}(A)$ are*

$$2(U_A - |D_A|) \quad \times 2, \quad 2(U_A + |D_A|) \quad \times 2.$$

In particular, we can recover $U_A : |D_A|$ from the ratio of the eigenvalues.

Theorem 2.5. *In (18), replacing “ U_A ” by “ $|D_A|$ ” one obtains the “axis” of the h-tube in the complex non-parabolic case; replacing “ U_A ” by “ $-|D_A|$ ” one obtains the equation for the pair of planes tangent at the asymptotic points corresponding to the eigenvalues of A . In the parabolic case $D_A = 0$, one obtains the double plane tangent at the asymptotic point corresponding to the eigenvalue. After these replacements the associated quadratic function will be nonnegative on the points of the (asymptotically closed) CKB(P) model.*

The first approach we consider is the method of “brute force”.

First Proof of Theorem 2.3. Assume that A is not normal. Recall, in projective coordinates $\text{DW}_{\text{CKB(P)}}(\mathbf{x}, A\mathbf{x}) = (\text{Re}\langle A\mathbf{x}, \mathbf{x} \rangle : \text{Im}\langle A\mathbf{x}, \mathbf{x} \rangle : \langle A\mathbf{x}, A\mathbf{x} \rangle : \langle \mathbf{x}, \mathbf{x} \rangle)$. Assume that

$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ and $\mathbf{x} = \begin{bmatrix} z_1 \\ z_2 \end{bmatrix}$ with $|z_1|^2 + |z_2|^2 = 1$. By direct computation, we find

$$\begin{bmatrix} \text{Re}\langle A\mathbf{x}, \mathbf{x} \rangle \\ \text{Im}\langle A\mathbf{x}, \mathbf{x} \rangle \\ \langle A\mathbf{x}, A\mathbf{x} \rangle \\ \langle \mathbf{x}, \mathbf{x} \rangle \end{bmatrix} = \underbrace{\begin{bmatrix} \text{Re} \frac{b+c}{2} & \text{Im} \frac{b-c}{2} & \text{Re} \frac{a-d}{2} & \text{Re} \frac{a+d}{2} \\ \text{Im} \frac{b+c}{2} & \text{Re} \frac{c-b}{2} & \text{Im} \frac{a-d}{2} & \text{Im} \frac{a+d}{2} \\ \text{Re}(\bar{a}b + \bar{c}d) & \text{Im}(\bar{a}b + \bar{c}d) & \frac{|a|^2 - |b|^2 + |c|^2 - |d|^2}{2} & \frac{|a|^2 + |b|^2 + |c|^2 + |d|^2}{2} \\ 0 & 0 & 0 & 1 \end{bmatrix}}_{\mathbf{S}_{\text{CKB(P)}}(A) :=} \begin{bmatrix} 2 \text{Re}(z_1 \bar{z}_2) \\ 2 \text{Im}(z_1 \bar{z}_2) \\ |z_1|^2 - |z_2|^2 \\ |z_1|^2 + |z_2|^2 \end{bmatrix}.$$

As the last column parameterizes the unit sphere in affine extended coordinates (as $|z_1|^2 + |z_2|^2 \equiv 1$, it yields the affine extension of $\iota_{\text{CKB}}(z_1/z_2)$), and $\mathbf{S}_{\text{CKB(P)}}(A)$ is itself

an affine extended matrix, we see that we deal with an affine linear image of the unit sphere. One can check directly that

$$(22) \quad \det \mathbf{S}_{\text{CKB(P)}}(A) = \frac{1}{2} ((U_A)^2 - |D_A|^2),$$

which is non-zero in the non-normal case. In that case, the image is a proper ellipsoid with matrix

$$(23) \quad \mathbf{S}_{\text{CKB(P)}}(A)^{-1, \top} \begin{bmatrix} 1 & & & \\ & 1 & & \\ & & 1 & \\ & & & -1 \end{bmatrix} \mathbf{S}_{\text{CKB(P)}}(A)^{-1} = \frac{1}{(U_A)^2 - |D_A|^2} \mathbf{Q}_{\text{CKB(P)}}(A).$$

The nonzero scalar multiplier may be omitted, so we obtain the matrix indicated. Due to nature of the object obtained, both the determinant and the interior property is straightforward. The case when A is normal follows from limiting arguments. (The limit matrix is nonzero as the entry in the (3, 3) position is 1.)

(Remark: This proof is very computational as the simplifications in (22) and (23) are cumbersome even in the case of the canonical triangular form. Note, however, that this method does not require any knowledge about the Davis–Wielandt shell beyond its definition; and it can also be used effectively for any concrete example.) \square

First Proof of Theorem 2.4. It follows from computing the characteristic polynomial of $(\mathbf{Q}_{\text{CKB(P)}}^0)^{-1} \mathbf{Q}_{\text{CKB(P)}}(A)$. (Remark: This is long even in terms of the five data.) \square

First Proof of Theorem 2.5. From the Theorem 2.4, we see that the singular quadratic forms of the pencil generated by $\mathbf{Q}_{\text{CKB(P)}}^0$ and $\mathbf{Q}_{\text{CKB(P)}}(A)$ are given, up to nonzero scalar multiples, by $\pm |D_A| 2\mathbf{Q}_{\text{CKB(P)}}^0 + \mathbf{Q}_{\text{CKB(P)}}^{\text{spec}}(A)$. In the non-parabolic case, bringing A to canonical form by unitary conjugation and taking off the off-diagonal term turns U_A into $|D_A|$, and this results in $|D_A| 2\mathbf{Q}_{\text{CKB(P)}}^0 + \mathbf{Q}_{\text{CKB(P)}}^{\text{spec}}(A)$ as the matrix of the axial tube. This leaves $-|D_A| 2\mathbf{Q}_{\text{CKB(P)}}^0 + \mathbf{Q}_{\text{CKB(P)}}^{\text{spec}}(A)$ as the matrix of the biplanar quadratic form. \square

The second approach we consider uses the conformal invariance of the shell.

Second Proof of Theorem 2.3. We can consider the matrices

$$(24) \quad L_t = \begin{bmatrix} 1 & 2t \\ & -1 \end{bmatrix}$$

($t \geq 0$). In this case, ± 1 are eigenvalues, thus $\iota_{\text{CKB}}(\pm 1) = (\pm 1, 0, 0)$ will be the asymptotic points of the h -tube $\text{DW}_{\text{CKB}}(L_t)$. In Euclidean view, the shell will be an ellipsoid with rotation axis $[(-1, 0, 0), (1, 0, 0)]_e$. As

$$\text{DW}_{\text{CKB}} \left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}, L_t \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right) = \text{DW}_{\text{CKB}} \left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 2t \\ -1 \end{bmatrix} \right) = \left(\frac{-1}{1+t^2}, 0, \frac{2t^2}{1+t^2} \right),$$

it can be checked that the fitting ellipsoid, for $t > 0$, is

$$(x_{\text{CKB}})^2 + \frac{(y_{\text{CKB}})^2}{t^2} + \frac{(z_{\text{CKB}})^2}{t^2} - 1 = 0;$$

$$\frac{\quad}{1+t^2} \quad \frac{\quad}{1+t^2}$$

or, in order to accommodate the limiting normal case $t = 0$,

$$(25) \quad 4t^2(x_{\text{CKB}})^2 + 4(1+t^2)(y_{\text{CKB}})^2 + 4(1+t^2)(z_{\text{CKB}})^2 - 4t^2 = 0.$$

(This was the initial argument of the proof.)

Abstractly, up to conjugation by unitary matrices, this have dealt with the case of

$$\operatorname{tr} L = 0 \quad \text{and} \quad \det L = -1,$$

being

$$t^2 = \frac{1}{2} \left(\frac{\operatorname{tr}(L^*L)}{2} - 1 \right).$$

If A is non-parabolic, such L can be obtained by taking the conformal transform

$$L = \frac{A - \frac{\operatorname{tr} A}{2} \operatorname{Id}}{\sqrt{-\det \left(A - \frac{\operatorname{tr} A}{2} \operatorname{Id} \right)}}.$$

Then

$$(26) \quad t^2 = \frac{1}{2} \left(\frac{\frac{1}{2} \operatorname{tr} \left(A - \frac{\operatorname{tr} A}{2} \operatorname{Id} \right)^* \left(A - \frac{\operatorname{tr} A}{2} \operatorname{Id} \right)}{\left| \det \left(A - \frac{\operatorname{tr} A}{2} \operatorname{Id} \right) \right|} - 1 \right) = \frac{1}{2} \left(\frac{U_A}{|D_A|} - 1 \right).$$

Let us use the temporary notation

$$(27) \quad T = -\frac{\operatorname{tr} A}{2}, \quad \Delta = \sqrt{-D_A}$$

(one complex value chosen). Then the equation (25) for $DW_{\text{CKB}}(L)$ can be rewritten as

$$(28) \quad 2(U_A + |\Delta|^2)(x_{\text{CKB}})^2 + 2(U_A - |\Delta|^2)(y_{\text{CKB}})^2 \\ + 2(U_A + |\Delta|^2)(z_{\text{CKB}})^2 + 2(-U_A + |\Delta|^2) = 0.$$

Thus the corresponding quadratic matrix, is

$$(29) \quad \mathbf{Q}_{\text{CKB}}^{[A]}(L) = \underbrace{2U_A \begin{bmatrix} 1 & & & \\ & 1 & & \\ & & 1 & \\ & & & -1 \end{bmatrix}}_{\mathbf{Q}_{\text{CKB}}^{\text{bas}}(L)} + \underbrace{2|\Delta|^2 \begin{bmatrix} -1 & & & \\ & 1 & & \\ & & 1 & \\ & & & 1 \end{bmatrix}}_{\mathbf{Q}_{\text{CKB}}^{\text{spec}}(L)}.$$

(The notation indicates that the scaling was chosen with respect to A .)

The L was, however, obtained after a displacement by the conformal transformation f such that

$$f(\lambda) = \frac{\lambda - \frac{\operatorname{tr} A}{2}}{\sqrt{-\det \left(A - \frac{\operatorname{tr} A}{2} \operatorname{Id} \right)}} \equiv \frac{\lambda + T}{\Delta},$$

whose projective representation matrix with respect to the CKB model is

$$R_{\text{CKB}}(f) = \frac{1}{|\Delta|} \cdot \frac{1}{2} \begin{bmatrix} 1 & 1 & & \\ & -i & i & \\ 1 & & & -1 \\ & & & 1 \end{bmatrix} \begin{bmatrix} 1 & \frac{\bar{T}}{\Delta} & T & \frac{|T|^2}{\Delta T} \\ & \Delta & \bar{T}\Delta & |\Delta|^2 \end{bmatrix} \begin{bmatrix} & & 1 & 1 \\ 1 & i & & \\ 1 & -i & & \\ & & -1 & 1 \end{bmatrix}.$$

Thus a matrix for the quadratic equation for $DW_{\text{CKB(P)}}(A)$ should be
(30)

$$\mathbf{Q}_{\text{CKB(P)}}^{\text{candidate}}(A) = \begin{bmatrix} 1 & & & \\ & 1 & & \\ & & \frac{1}{2} & -\frac{1}{2} \\ & & \frac{1}{2} & \frac{1}{2} \end{bmatrix}^{\top} \cdot R_{\text{CKB}}(f)^{\top} \cdot \mathbf{Q}_{\text{CKB}}^{[A]}(L) \cdot R_{\text{CKB}}(f) \cdot \begin{bmatrix} 1 & & & \\ & 1 & & \\ & & \frac{1}{2} & -\frac{1}{2} \\ & & \frac{1}{2} & \frac{1}{2} \end{bmatrix}.$$

Here the middle term is quadratic matrix for $DW_{\text{CKB}}(L)$, then $R_{\text{CKB}}(f)$ takes care to the displacement in CKB, and the outer term translates between CKB and CKB(P). Expanded, it

$$= \begin{bmatrix} 2T\bar{T} - \Delta^2 + \bar{T}^2 - \bar{\Delta}^2 + T^2 + 2U_A & -i\bar{\Delta}^2 + i\Delta^2 - iT^2 + i\bar{T}^2 & T + \bar{T} & \bar{T}^2 T + T^2 \bar{T} - \Delta^2 \bar{T} - \bar{\Delta}^2 T \\ -i\bar{\Delta}^2 + i\Delta^2 - iT^2 + i\bar{T}^2 & 2T\bar{T} + \Delta^2 - \bar{T}^2 + \bar{\Delta}^2 - T^2 + 2U_A & i\bar{T} - iT & i\Delta^2 \bar{T} + i\bar{T}^2 T - iT^2 \bar{T} - i\bar{\Delta}^2 T \\ T + \bar{T} & i\bar{T} - iT & 1 & T\bar{T} - U_A \\ \bar{T}^2 T + T^2 \bar{T} - \Delta^2 \bar{T} - \bar{\Delta}^2 T & i\Delta^2 \bar{T} + i\bar{T}^2 T - iT^2 \bar{T} - i\bar{\Delta}^2 T & T\bar{T} - U_A & -\Delta^2 \bar{T}^2 + T^2 \bar{T}^2 + \Delta^2 \bar{\Delta}^2 - \bar{\Delta}^2 T^2 \end{bmatrix}.$$

Then, taking (27) into consideration, it yields $\mathbf{Q}_{\text{CKB(P)}}^{\text{candidate}}(A) \equiv \mathbf{Q}_{\text{CKB(P)}}(A)$ indeed. From (29) (where $|\Delta|^2 = |D_A|$) and (30) (where $\det R_{\text{CKB}}(f) = 1$) the determinant follows immediately. The characterization of the interior carries from the case of L_t again.

Now, the argument above establishes the non-parabolic case. Then the parabolic case follows by continuity. (The limits are nontrivial, as the entries at (3, 3) are 1.) \square

Second Proof for 2.4. Continuing the previous Second Proof of Theorem 2.3,

$$\begin{aligned} (\mathbf{Q}_{\text{CKB}}^0)^{-1} \mathbf{Q}_{\text{CKB}}(A) &= \left(R_{\text{CKB}}(f)^{\top} \cdot \mathbf{Q}_{\text{CKB}}^0 \cdot R_{\text{CKB}}(f) \right)^{-1} R_{\text{CKB}}(f)^{\top} \cdot \mathbf{Q}_{\text{CKB}}^{[A]}(L) \cdot R_{\text{CKB}}(f) \\ &= R_{\text{CKB}}(f)^{-1} \cdot (\mathbf{Q}_{\text{CKB}}^0)^{-1} \mathbf{Q}_{\text{CKB}}^{[A]}(L) \cdot R_{\text{CKB}}(f) \sim (\mathbf{Q}_{\text{CKB}}^0)^{-1} \mathbf{Q}_{\text{CKB}}^{[A]}(L). \end{aligned}$$

From this, the conjugacy type of $(\mathbf{Q}_{\text{CKB}}^0)^{-1} \mathbf{Q}_{\text{CKB}}(A)$ can be read off immediately in the non-parabolic case. Due to the continuity of the characteristic polynomial the eigenvalues extend (but the conjugacy type does not). \square

Second Proof of Theorem 2.5. In the previous Second Proof of Theorem 2.3, we obtain the corresponding matrices by making the appropriate replacements $\pm|D_A|$ to U_A in (28)/(29) in the non-parabolic case $|D_A| \neq 0$. The parabolic case $|D_A| = 0$ follows from limiting arguments. \square

Remark 2.6. (a) In the initial argument of the Second Proof of Theorem 2.3,

$$\begin{aligned} DW_{\text{CKB}} \left(\begin{bmatrix} \sqrt{1+t^2} - t \\ 1 \end{bmatrix}, L_t \begin{bmatrix} \sqrt{1+t^2} - t \\ 1 \end{bmatrix} \right) &= \\ &= DW_{\text{CKB}} \left(\begin{bmatrix} \sqrt{1+t^2} - t \\ 1 \end{bmatrix}, \begin{bmatrix} \sqrt{1+t^2} + t \\ -1 \end{bmatrix} \right) = \left(0, 0, \sqrt{\frac{t^2}{1+t^2}} \right) \end{aligned}$$

gives immediately the top point of the shell (although this choice for the testing point looks unmotivated). From this, the ellipsoid of the tube can be recovered.

(b) A more geometric alternative in order to find out the top point is as follows. Using Lemma 1.8, one can compute $\|L_t\|_2 = t + \sqrt{1+t^2}$. Then, by symmetry, the top point of $DW_{\text{CKB(P)}}(L_t)$ of $(0, 0, (t + \sqrt{1+t^2})^2)$. Transcribed to the CKB model, the top point of $DW_{\text{CKB}}(L_t)$ is $(0, 0, \sqrt{\frac{t^2}{1+t^2}})$.

(c) If one finds the earlier initial arguments of the Second Proof of Theorem 2.3 too geometric, then it can be replaced by the following computational argument, which is a special case of the brute force argument (in modified form):

Using projective coordinates, for $\mathbf{x} = \begin{bmatrix} z_1 \\ z_2 \end{bmatrix}$, $|z_1|^2 + |z_2|^2 = 1$, it yields

$$(31) \quad DW_{\text{CKB}}(\mathbf{x}, L_t \mathbf{x}) \sim \begin{bmatrix} \text{Re}\langle L_t \mathbf{x}, \mathbf{x} \rangle \\ \text{Im}\langle L_t \mathbf{x}, \mathbf{x} \rangle \\ \frac{\langle L_t \mathbf{x}, L_t \mathbf{x} \rangle - \langle \mathbf{x}, \mathbf{x} \rangle}{2} \\ \frac{\langle L_t \mathbf{x}, L_t \mathbf{x} \rangle + \langle \mathbf{x}, \mathbf{x} \rangle}{2} \end{bmatrix} = \underbrace{\begin{bmatrix} t & 0 & 1 & 0 \\ 0 & -t & 0 & 0 \\ t & 0 & -t^2 & t^2 \\ t & 0 & -t^2 & t^2 + 1 \end{bmatrix}}_{\mathbf{S}_{\text{CKB}}(L_t) :=} \begin{bmatrix} 2 \text{Re}(z_1 \bar{z}_2) \\ 2 \text{Im}(z_1 \bar{z}_2) \\ |z_1|^2 - |z_2|^2 \\ |z_1|^2 + |z_2|^2 \end{bmatrix} =$$

$$= \underbrace{\sqrt{1+t^2}}_{S_0} \underbrace{\begin{bmatrix} 1 & & & \\ & \frac{t}{\sqrt{1+t^2}} & & \\ & & \frac{t}{\sqrt{1+t^2}} & \\ & & & 1 \end{bmatrix}}_{S_1} \underbrace{\left(\lambda \mapsto \frac{\lambda + t + \sqrt{1+t^2}}{\lambda + t - \sqrt{1+t^2}} \right)}_{S_2} \underbrace{\begin{bmatrix} 2 \text{Re}(z_1 \bar{z}_2) \\ 2 \text{Im}(z_1 \bar{z}_2) \\ |z_1|^2 - |z_2|^2 \\ |z_1|^2 + |z_2|^2 \end{bmatrix}}_{S_3}.$$

Then, S_3 ranges over the unit sphere (in projective coordinates), S_2 leaves it invariant, S_1 shrinks it, and the scaling factor S_0 is irrelevant; from this one obtain $DW_{\text{CKB}}(L_t)$ immediately. In fact, we do not even have to follow through this computation completely. It is sufficient to show that the rows of $\mathbf{S}_{\text{CKB}}(L_t)$ are $(1, 1, 1, -1)$ -pseudoorthogonal, however not $(1, 1, 1, -1)$ -pseudoorthonormal but of square length $1+t^2, t^2, t^2, -(1+t^2)$ respectively. \triangle

Ultimately, the second (“conformal”) approach is quite similar to the first (“brute force”) approach but the computation is via familiar geometric elements, making the computations of the determinant and eigenvalues particularly transparent.

Example 2.7. (a) For L_t as in (24),

$$\mathbf{Q}_{\text{CKB(P)}}(L_t) = \begin{bmatrix} 4t^2 & & & \\ & 4+4t^2 & & \\ & & 1 & -1-2t^2 \\ & & -1-2t^2 & 1 \end{bmatrix},$$

and

$$\mathbf{Q}_{\text{CKB}}(L_t) = \begin{bmatrix} 4t^2 & & & \\ & 4+4t^2 & & \\ & & 4+4t^2 & \\ & & & -4t^2 \end{bmatrix}.$$

(b) The non-normal parabolic representative S_0 yields

$$\mathbf{Q}_{\text{CKB(P)}}(S_0) = \begin{bmatrix} 1 & & & \\ & 1 & & \\ & & 1 & -\frac{1}{2} \\ & & -\frac{1}{2} & \end{bmatrix}, \quad \text{and} \quad \mathbf{Q}_{\text{CKB}}(S_0) = \begin{bmatrix} 1 & & & \\ & 1 & & \\ & & 2 & 1 \\ & & 1 & \end{bmatrix}.$$

(c) The (normal, parabolic) zero matrix $\mathbf{0}_2$ yields

$$\mathbf{Q}_{\text{CKB}(\text{P})}(\mathbf{0}_2) = \begin{bmatrix} 0 & & & \\ & 0 & & \\ & & 1 & \\ & & & 0 \end{bmatrix}, \quad \text{and} \quad \mathbf{Q}_{\text{CKB}(\mathbf{0}_2)} = \begin{bmatrix} 0 & & & \\ & 0 & & \\ & & 1 & 1 \\ & & 1 & 1 \end{bmatrix}. \quad \diamond$$

The third approach we consider is the ‘‘pencil argument’’. This, of course, bases heavily on the geometric interpretation of the qualitative elliptical range theorem.

Lemma 2.8. *Assume that $\lambda_1, \lambda_2 \in \mathbb{C}$.*

(a) *The biplanar quadratic form tangent to the asymptotic boundary of the $\text{CKB}(\text{P})$ model at λ_1 and λ_2 is given, up to nonzero scalar multiples, by*

$$(z - 2x(\text{Re } \lambda_1) - 2y(\text{Im } \lambda_1) + |\lambda_1|^2)(z - 2x(\text{Re } \lambda_2) - 2y(\text{Im } \lambda_2) + |\lambda_2|^2).$$

(b) *If $\lambda_1 \neq \lambda_2$, then the quadratic form of the infinitesimal h -tube connecting $\iota_{\text{CKB}(\text{P})}(\lambda_1)$ and $\iota_{\text{CKB}(\text{P})}(\lambda_2)$ is given, up to nonzero scalar multiples, by*

$$(z - 2x(\text{Re } \lambda_1) - 2y(\text{Im } \lambda_1) + |\lambda_1|^2)(z - 2x(\text{Re } \lambda_2) - 2y(\text{Im } \lambda_2) + |\lambda_2|^2) \\ - (((\text{Re } \lambda_1) - (\text{Re } \lambda_2))^2 + ((\text{Im } \lambda_1) - (\text{Im } \lambda_2))^2)(z - x^2 - y^2).$$

Here x, y, z were understood as $x_{\text{CKB}(\text{P})}, y_{\text{CKB}(\text{P})}, z_{\text{CKB}(\text{P})}$, respectively, throughout.

Proof. (a) The equations of the tangent planes are easy to recover, then one should multiply them. (b) The previous quadratic form should be corrected by a multiple of $z - x^2 - y^2$ so that it should fit to $\frac{\iota_{\text{CKB}(\text{P})}(\lambda_1) + \iota_{\text{CKB}(\text{P})}(\lambda_2)}{2}$. \square

Third Proof of Theorem 2.5. Assume that A have eigenvalues λ_1 and λ_2 . Then, using the identities $|D_A| = \frac{1}{4}|\lambda_1 - \lambda_2|^2 = \frac{1}{4}(((\text{Re } \lambda_1) - (\text{Re } \lambda_2))^2 + ((\text{Im } \lambda_1) - (\text{Im } \lambda_2))^2)$, $\text{Re tr } A = (\text{Re } \lambda_1) + (\text{Re } \lambda_2)$, etc., one can check that the matrices of the quadratic forms in Lemma 2.8 (a) and (b) are the matrices $-|D_A|2\mathbf{Q}_{\text{CKB}(\text{P})}^0 + \mathbf{Q}_{\text{CKB}(\text{P})}^{\text{spec}}(A)$ and $|D_A|2\mathbf{Q}_{\text{CKB}(\text{P})}^0 + \mathbf{Q}_{\text{CKB}(\text{P})}^{\text{spec}}(A)$, respectively. \square

Now, we could prove Theorem 2.3 by fitting an additional point, in the manner of Lemma 2.8(b). Instead, we proceed using a different argument:

Lemma 2.9. *The center of $\text{DW}_{\text{CKB}(\text{P})}(A)$ is $\left(\frac{\text{Re tr } A}{2}, \frac{\text{Im tr } A}{2}, \frac{\text{tr}(A^*A)}{2}\right)$.*

Proof. We have a possibly degenerate ellipsoid which is centrally symmetric. In each coordinate the extremal values are given by the two-two eigenvalues of the self-adjoint operators $\frac{A+A^*}{2}$, $\frac{A-A^*}{2i}$, A^*A , respectively. The averages of those eigenvalues are given by $\frac{1}{2}$ times the traces. \square

Third Proof of Theorem 2.3. Let us assume that A is not normal, i. e. the ellipsoid is proper. We have to modify $-|D_A|2\mathbf{Q}_{\text{CKB}(\text{P})}^0 + \mathbf{Q}_{\text{CKB}(\text{P})}^{\text{spec}}(A)$ by a scalar multiple of $\mathbf{Q}_{\text{CKB}(\text{P})}^0$ in order to get the equation for the shell. Thus the candidate is of shape $\xi 2\mathbf{Q}_{\text{CKB}(\text{P})}^0 + \mathbf{Q}_{\text{CKB}(\text{P})}^{\text{spec}}(A)$. According to Lemma 2.9, this has the central symmetry

$$(32) \quad \xi 2\mathbf{Q}_{\text{CKB}(\text{P})}^0 + \mathbf{Q}_{\text{CKB}(\text{P})}^{\text{spec}}(A) =$$

$$= \begin{bmatrix} -1 & & \operatorname{Re} \operatorname{tr} A \\ & -1 & \operatorname{Im} \operatorname{tr} A \\ & & -1 & \operatorname{tr}(A^*A) \\ & & & 1 \end{bmatrix}^\top \left(\xi 2\mathbf{Q}_{\text{CKB(P)}}^0 + \mathbf{Q}_{\text{CKB(P)}}^{\text{spec}}(A) \right) \begin{bmatrix} -1 & & \operatorname{Re} \operatorname{tr} A \\ & -1 & \operatorname{Im} \operatorname{tr} A \\ & & -1 & \operatorname{tr}(A^*A) \\ & & & 1 \end{bmatrix}.$$

In fact, geometrically, we could only claim equality up to a nonzero scalar multiple, but the coefficients 1 in positions (3, 3) imply equality. Now, however, comparing positions (3, 4), we obtain $\xi = U_A$ immediately. (Actually, writing (32) into transpose-commutator form, it simplifies further.)

If A is normal, then we already know that $|D_A|2\mathbf{Q}_{\text{CKB(P)}}^0 + \mathbf{Q}_{\text{CKB(P)}}^{\text{spec}}(A)$ functions properly for the equation.

Having $\mathbf{Q}_{\text{CKB(P)}}(A)$, its determinant can be computed (but it is easier to recover from the following Third Proof of Theorem 2.4). The coefficient 1 in (3, 3) position of $\mathbf{Q}_{\text{CKB(P)}}(A)$ informs us that how the sign relation for the interior should hold. \square

Third Proof of Theorem 2.4. In the non-parabolic case, on geometrical grounds, we know from the pencil that the eigenvalues come in pairs. From the parameter of the singular members of the pencil ($U_A \rightsquigarrow \pm|D_A|$), we see that the eigenvalues are as indicated. \square

Remark 2.10. In their characterization of the shell of 2×2 complex matrices, Li, Poon, Sze [27] indicates the center $\left(\frac{\operatorname{Re} \operatorname{tr} A}{2}, \frac{\operatorname{Im} \operatorname{tr} A}{2}, \frac{\operatorname{tr}(A^*A)}{2} \right)$ of $\text{DW}_{\text{CKB(P)}}(A)$ correctly; however, the vertical diameter (but not principal axis) through the center is with endpoints $\left(\frac{\operatorname{Re} \operatorname{tr} A}{2}, \frac{\operatorname{Im} \operatorname{tr} A}{2}, \frac{\operatorname{tr}(A^*A)}{2} \pm \sqrt{(U_A)^2 - |D_A|^2} \right)$. \triangle

However, the cleanest approach to get (18) is via the dual approach. As we deal with ellipsoids, some knowledge of projective geometry is sufficient. There is no need for advanced algebra in order to pass between the ordinary and the dual picture but a simple matrix inversion:

Fourth Proof of Theorem 2.3. Assume that A is non-normal. We consider the projective homogeneous polynomial for the dual (i. e. tangent) surface:

$$\begin{aligned} K_A^{\text{DW}}(u, v, s, w) &\equiv \\ &\equiv \det \left(u \frac{A + A^*}{2} + v \frac{A - A^*}{2i} + sA^*A + w \operatorname{Id}_2 \right) \\ &= \frac{1}{2} \left(\operatorname{tr} \left(u \frac{A + A^*}{2} + v \frac{A - A^*}{2i} + sA^*A + w \operatorname{Id}_2 \right) \right)^2 \\ &\quad - \frac{1}{2} \operatorname{tr} \left(\left(u \frac{A + A^*}{2} + v \frac{A - A^*}{2i} + sA^*A + w \operatorname{Id}_2 \right)^2 \right) \\ &= \frac{|\operatorname{tr} A|^2 - \operatorname{tr}(A^*A)}{4} (u^2 + v^2) + \frac{\operatorname{Re} \det A}{2} (u^2 - v^2) + uv \operatorname{Im} \det A \\ &\quad + us \operatorname{Re}((\det A)(\overline{\operatorname{tr} A})) + vs \operatorname{Im}((\det A)(\overline{\operatorname{tr} A})) + s^2 |\det A|^2 \\ &\quad + uw \operatorname{Re} \operatorname{tr} A + vw \operatorname{Im} \operatorname{tr} A + sw \operatorname{tr}(A^*A) + w^2 \\ &= \begin{bmatrix} u \\ v \\ s \\ w \end{bmatrix}^\top \mathbf{G}_{\text{CKB(P)}}(A) \begin{bmatrix} u \\ v \\ s \\ w \end{bmatrix}, \end{aligned}$$

where

$$(33) \quad \mathbf{G}_{\text{CKB(P)}}(A) := \begin{bmatrix} \frac{|\text{tr } A|^2 - \text{tr}(A^*A) + 2 \text{Re det } A}{4} & \frac{\text{Im det } A}{2} & \frac{\text{Re}((\text{det } A)(\overline{\text{tr } A}))}{2} & \frac{\text{Re tr } A}{2} \\ \frac{\text{Im det } A}{2} & \frac{|\text{tr } A|^2 - \text{tr}(A^*A) - 2 \text{Re det } A}{4} & \frac{\text{Im}((\text{det } A)(\overline{\text{tr } A}))}{2} & \frac{\text{Im tr } A}{2} \\ \frac{\text{Re}((\text{det } A)(\overline{\text{tr } A}))}{2} & \frac{\text{Im}((\text{det } A)(\overline{\text{tr } A}))}{2} & |\text{det } A|^2 & \frac{\text{tr}(A^*A)}{2} \\ \frac{\text{Re tr } A}{2} & \frac{\text{Im tr } A}{2} & \frac{\text{tr}(A^*A)}{2} & 1 \end{bmatrix}.$$

By that, we have written down the matrix of the dual of the shell. One finds that

$$(34) \quad \det \mathbf{G}_{\text{CKB(P)}}(A) = -\frac{1}{4} ((U_A)^2 - |D_A|^2)^2,$$

and

$$(35) \quad \mathbf{Q}_{\text{CKB(P)}}(A) = -((U_A)^2 - |D_A|^2) (\mathbf{G}_{\text{CKB(P)}}(A))^{-1}.$$

According to elementary projective geometry this implies that a matrix of the dual of the dual conic is $\mathbf{Q}_{\text{CKB(P)}}(A)$.

The normal case follows from limiting arguments.

(Remark: An advantage of the proof is that it is entirely in terms of the five data. The intermediate quantity $\mathbf{G}_{\text{CKB(P)}}(A)$ is also of geometric significance. We have

$$(36) \quad \mathbf{S}_{\text{CKB(P)}}(A) \begin{bmatrix} 1 & & & \\ & 1 & & \\ & & 1 & \\ & & & -1 \end{bmatrix} \mathbf{S}_{\text{CKB(P)}}(A)^\top = \mathbf{G}_{\text{CKB(P)}}(A)$$

in terms of the first proof.) □

Then one can proceed with Theorem 2.4 and Theorem 2.5 as in the first proofs.

Alternative proof to Lemma 2.9. In the non-normal case, the projective coordinates of the center of the shell can read off from the last column of $\mathbf{G}_{\text{CKB(P)}}(A)$ (or any non-zero multiple of the inverse of $\mathbf{Q}_{\text{CKB(P)}}(A)$). From this, the center is immediate. The normal case follows by continuity. □

$\mathbf{G}_{\text{CKB(P)}}(A)$ as defined by (33), is a natural dual quantity to $\mathbf{Q}_{\text{CKB(P)}}(A)$. As a shorthand notation, we will use

$$(37) \quad \mathbf{G}_{\text{CKB(P)}}^0 = (\mathbf{Q}_{\text{CKB(P)}}^0)^{-1} \quad \text{and} \quad \mathbf{G}_{\text{CKB}}^0 = (\mathbf{Q}_{\text{CKB}}^0)^{-1}.$$

Remark 2.11. $\mathbf{G}_{\text{CKB(P)}}(A)$ still allows a decomposition to basic and spectral parts:

$$\mathbf{G}_{\text{CKB(P)}}(A) = U_A \underbrace{\begin{bmatrix} -\frac{1}{2} & & & \\ & -\frac{1}{2} & & \\ & & 1 & \\ & & & 1 \end{bmatrix}}_{\mathbf{G}_{\text{CKB(P)}}^{\text{bas}}(A)} +$$

$$+ \underbrace{\begin{bmatrix} \frac{|\operatorname{tr} A|^2 + 4 \operatorname{Re} \det A}{8} & \frac{\operatorname{Im} \det A}{2} & \frac{\operatorname{Re}((\det A)(\overline{\operatorname{tr} A}))}{2} & \frac{\operatorname{Re} \operatorname{tr} A}{2} \\ \frac{\operatorname{Im} \det A}{2} & \frac{|\operatorname{tr} A|^2 - 4 \operatorname{Re} \det A}{8} & \frac{\operatorname{Im}((\det A)(\overline{\operatorname{tr} A}))}{2} & \frac{\operatorname{Im} \operatorname{tr} A}{2} \\ \frac{\operatorname{Re}((\det A)(\overline{\operatorname{tr} A}))}{2} & \frac{\operatorname{Im}((\det A)(\overline{\operatorname{tr} A}))}{2} & |\det A|^2 & \frac{|\operatorname{tr} A|^2}{4} \\ \frac{\operatorname{Re} \operatorname{tr} A}{2} & \frac{\operatorname{Im} \operatorname{tr} A}{2} & \frac{|\operatorname{tr} A|^2}{4} & 1 \end{bmatrix}}{\mathbf{G}_{\operatorname{CKB}(P)}^{\operatorname{spec}}(A)} :=$$

It fits into a corresponding dual pencil, cf.

$$\left(\lambda 2\mathbf{Q}_{\operatorname{CKB}(P)}^0 + \mathbf{Q}_{\operatorname{CKB}(P)}^{\operatorname{spec}}(A) \right) \left(\lambda \left(-\frac{1}{2}\mathbf{G}_{\operatorname{CKB}(P)}^0 \right) + \mathbf{G}_{\operatorname{CKB}(P)}^{\operatorname{spec}}(A) \right) = -(\lambda^2 - |D_A|^2) \operatorname{Id}_4.$$

Setting $\lambda = \pm|D_A|$ we obtain the singular points of the pencils; cf.

$$\det \left(\lambda 2\mathbf{Q}_{\operatorname{CKB}(P)}^0 + \mathbf{Q}_{\operatorname{CKB}(P)}^{\operatorname{spec}}(A) \right) = -4(\lambda^2 - |D_A|^2)^2,$$

and

$$\det \left(\lambda \left(-\frac{1}{2}\mathbf{G}_{\operatorname{CKB}(P)}^0 \right) + \mathbf{G}_{\operatorname{CKB}(P)}^{\operatorname{spec}}(A) \right) = -\frac{1}{4}(\lambda^2 - |D_A|^2)^2.$$

We also remark that

$$\mathbf{Q}_{\operatorname{CKB}(P)}^{\operatorname{spec}}(A) = 4\mathbf{Q}_{\operatorname{CKB}(P)}^0 \mathbf{G}_{\operatorname{CKB}(P)}^{\operatorname{spec}}(A) \mathbf{Q}_{\operatorname{CKB}(P)}^0,$$

and

$$\mathbf{Q}_{\operatorname{CKB}(P)}^{\operatorname{bas}}(A) = -4\mathbf{Q}_{\operatorname{CKB}(P)}^0 \mathbf{G}_{\operatorname{CKB}(P)}^{\operatorname{bas}}(A) \mathbf{Q}_{\operatorname{CKB}(P)}^0.$$

One can see that not only $\mathbf{Q}_{\operatorname{CKB}(P)}(A)$ and $\mathbf{G}_{\operatorname{CKB}(P)}(A)$ are linear rearrangements of each other, but also the corresponding pencils. \triangle

We can define, compatibly to (37),

$$(38) \quad \mathbf{G}_{\operatorname{CKB}}(A) = \begin{bmatrix} 1 & & & \\ & 1 & & \\ & & 1 & 1 \\ & & -1 & 1 \end{bmatrix}^{-1} \mathbf{G}_{\operatorname{CKB}(P)}(A) \begin{bmatrix} 1 & & & \\ & 1 & & \\ & & 1 & 1 \\ & & -1 & 1 \end{bmatrix}^{-1, \top}.$$

With this convention, $\mathbf{G}_{\operatorname{CKB}}(A) \mathbf{Q}_{\operatorname{CKB}}^0$ and $\mathbf{G}_{\operatorname{CKB}(P)}(A) \mathbf{Q}_{\operatorname{CKB}(P)}^0$ will be similar matrices.

Theorem 2.12. *The eigenvalues of $\mathbf{G}_{\operatorname{CKB}(P)}(A) \mathbf{Q}_{\operatorname{CKB}(P)}^0 \sim \mathbf{G}_{\operatorname{CKB}}(A) \mathbf{Q}_{\operatorname{CKB}}^0$ are*

$$-\frac{U_A - |D_A|}{2} \times 2 \quad \text{and} \quad -\frac{U_A + |D_A|}{2} \times 2.$$

In particular, we can recover $U_A : |D_A|$ from the ratio of the eigenvalues.

Proof. One can compare the characteristic polynomials of $\mathbf{G}_{\operatorname{CKB}(P)}(A) \mathbf{Q}_{\operatorname{CKB}(P)}^0$ and $\mathbf{G}_{\operatorname{CKB}(P)}^0 \mathbf{Q}_{\operatorname{CKB}(P)}(A)$ (cf. Theorem 2.4) in order find that they are appropriately scaled versions of each other. **Alternatively**, the statement follows from (34) and (35) generically (in the non-normal case), then we can use the continuity of the characteristic polynomials. \square

Lemma 2.13. *There is a similarity of matrices*

$$(\mathbf{Q}_{\text{CKB}(\mathbb{P})}^0)^{-1} \mathbf{Q}_{\text{CKB}(\mathbb{P})}(A) \sim -4 \mathbf{G}_{\text{CKB}(\mathbb{P})}(A) \mathbf{Q}_{\text{CKB}(\mathbb{P})}^0.$$

Proof. Consider the invertible affine matrix

$$S = \begin{bmatrix} & & -1 & & \frac{1}{2} \operatorname{Re} \operatorname{tr} A + \frac{1}{2} \operatorname{Im} \operatorname{tr} A \\ & 1 & & & -\frac{1}{2} \operatorname{Re} \operatorname{tr} A + \frac{1}{2} \operatorname{Im} \operatorname{tr} A \\ \operatorname{Re} \operatorname{tr} A + \operatorname{Im} \operatorname{tr} A & -\operatorname{Re} \operatorname{tr} A + \operatorname{Im} \operatorname{tr} A & -1 & & \\ & & & & 1 \end{bmatrix}.$$

Then,

$$(\mathbf{Q}_{\text{CKB}(\mathbb{P})}^0)^{-1} \mathbf{Q}_{\text{CKB}(\mathbb{P})}(A) = S \left(-4 \mathbf{G}_{\text{CKB}(\mathbb{P})}(A) \mathbf{Q}_{\text{CKB}(\mathbb{P})}^0 \right) S^{-1}.$$

(Remark: This also shows the equivalence of Theorem 2.12 and Theorem 2.4.) \square

2.C. Decompositions of the quadratic forms.

Corollary 2.14. *For 2×2 complex matrices, the five data determines the Davis–Wielandt shell, and vice versa.*

Proof. The five data determines the matrix up to unitary conjugation, which, of course, determines shell, as unitary conjugation leaves it invariant.

Conversely, if $\text{DW}_{\text{CKB}(\mathbb{P})}(A)$ is given, then from the asymptotic points, we can recover the eigenvalues, and from the central vertical diameter $2\sqrt{(U_A)^2 - |D_A|^2}$, the value $\operatorname{tr}(A^*A)$ can also be recovered. **Alternatively**, if the matrix A is normal, then the endpoints determine the eigenvalues, which is sufficient to recover the diagonal matrix up to unitary conjugation. If the matrix A is non-normal and \mathbf{M} is a matrix of the quadric in the $\text{CKB}(\mathbb{P})$ model, then $\mathbf{Q}_{\text{CKB}(\mathbb{P})}(A) = \mathbf{M}/\mathbf{M}_{33}$, and from this matrix the five data can immediately be read off. In fact, the preceding argument also works if A is normal but \mathbf{M} is required to be an infinitesimal tube or a double tangent plane.

(Remark: Cf. Theorem 3.7.) \square

We define the ‘core matrices’

$$\mathbf{Q}^{\text{C}}(A) = \begin{bmatrix} \operatorname{tr}(A^*A) - (\operatorname{Re} \operatorname{tr} A)^2 + 2 \operatorname{Re} \det A & -(\operatorname{Re} \operatorname{tr} A)(\operatorname{Im} \operatorname{tr} A) + 2 \operatorname{Im} \det A \\ -(\operatorname{Re} \operatorname{tr} A)(\operatorname{Im} \operatorname{tr} A) + 2 \operatorname{Im} \det A & \operatorname{tr}(A^*A) - (\operatorname{Im} \operatorname{tr} A)^2 - 2 \operatorname{Re} \det A \end{bmatrix},$$

and

$$\mathbf{G}^{\text{C}}(A) = -\frac{1}{4} \operatorname{adj} \mathbf{Q}^{\text{C}}(A).$$

As adj is an involutive operation on 2×2 matrices, one also has

$$\mathbf{Q}^{\text{C}}(A) = -4 \operatorname{adj} \mathbf{G}^{\text{C}}(A).$$

We set

$$\mathbf{B}_{\text{CKB}(\mathbb{P})}(A) = \begin{bmatrix} 1 & & & \frac{1}{2} \operatorname{Re} \operatorname{tr} A \\ & 1 & & \frac{1}{2} \operatorname{Im} \operatorname{tr} A \\ \operatorname{Re} \operatorname{tr} A & \operatorname{Im} \operatorname{tr} A & 1 & \frac{1}{2} \operatorname{tr}(A^*A) \\ & & & 1 \end{bmatrix},$$

which is an invertible affine matrix.

Theorem 2.15. (a)

$$\mathbf{Q}_{\text{CKB}}(A) = \mathbf{B}_{\text{CKB}(\text{P})}(A)^{-1, \top} \underbrace{\begin{bmatrix} \mathbf{Q}^{\text{C}}(A) & & \\ & 1 & \\ & & -((U_A)^2 - |D_A|^2) \end{bmatrix}}_{Q_1} \mathbf{B}_{\text{CKB}(\text{P})}(A)^{-1}.$$

(b) The eigenvalues of the $(2|1|1)$ block matrix Q_1 are

$$2(U_D - |D_A|), \quad 2(U_D + |D_A|), \quad | \quad 1, \quad | \quad -((U_A)^2 - |D_A|^2);$$

the matrix is diagonalizable.

Theorem 2.16. (a)

$$\mathbf{G}_{\text{CKB}}(A) = \mathbf{B}_{\text{CKB}(\text{P})}(A) \underbrace{\begin{bmatrix} \mathbf{G}^{\text{C}}(A) & & \\ & -((U_A)^2 - |D_A|^2) & \\ & & 1 \end{bmatrix}}_{G_1} \mathbf{B}_{\text{CKB}(\text{P})}(A)^{\top}.$$

(b) The eigenvalues of the $(2|1|1)$ block matrix G_1 are

$$-\frac{1}{2}(U_D - |D_A|), \quad -\frac{1}{2}(U_D + |D_A|), \quad | \quad -((U_A)^2 - |D_A|^2), \quad | \quad 1;$$

the matrix is diagonalizable.

Proofs. Straightforward computations. □

Then it easy to see that

$$\text{rk } \mathbf{Q}_{\text{CKB}(\text{P})}(A) = \text{rk } \mathbf{G}_{\text{CKB}(\text{P})}(A) = \begin{cases} 4 & \text{if } A \text{ is non-normal,} \\ 2 & \text{if } A \text{ is normal non-parabolic,} \\ 1 & \text{if } A \text{ is normal parabolic.} \end{cases}$$

However, we may also know this from the pencil picture. Another possibility is to look up the canonical representatives from Example 2.7 for $\text{rk } \mathbf{Q}_{\text{CKB}(\text{P})}(A)$. (Möbius transformations may introduce only a scaling by nonzero scalar multiplier; this is also valid in the degenerate cases. The $\text{CKB}(\text{P})$ and CKB representatives can be used to determine the rank equally.) Then Lemma 2.13 shows $\text{rk } \mathbf{Q}_{\text{CKB}(\text{P})}(A) = \text{rk } \mathbf{G}_{\text{CKB}(\text{P})}(A)$.

(Taking the quotient of symmetric matrix by an indefinite symmetric matrix may lead to non-trivial Jordan blocks. Consequently, Theorems 2.4 and 2.12 cannot be used to find the ranks. Indeed, for scalar matrices the eigenvalues there are all zero, while $\mathbf{Q}_{\text{CKB}}(A)$ and $\mathbf{G}_{\text{CKB}}(A)$ have ranks ≥ 1 in any case, due to the entry ‘1’.)

2.D. The transformation properties of the matrix.

Definition 2.17. Suppose that f is a Möbius transformation $f : \lambda \mapsto \frac{a\lambda+b}{c\lambda+d}$, $a, b, c, d \in \mathbb{C}$, $ad - bc \neq 0$, and A is 2×2 complex matrix. Then we set

$$(39) \quad \mathcal{C}(f, A) = \frac{1}{|ad - bc|^2} \begin{bmatrix} -\text{Re}(\bar{c}d) \\ -\text{Im}(\bar{c}d) \\ |d|^2 \\ |c|^2 \end{bmatrix}^{\top} \mathbf{Q}_{\text{CKB}(\text{P})}(A) \begin{bmatrix} -\text{Re}(\bar{c}d) \\ -\text{Im}(\bar{c}d) \\ |d|^2 \\ |c|^2 \end{bmatrix}$$

(it is well-defined). Actually, this can be rewritten as

$$(40) \quad \mathcal{C}(f, A) = \frac{1}{|ad - bc|^2} \begin{bmatrix} 2 \operatorname{Re}(\bar{c}d) \\ 2 \operatorname{Im}(\bar{c}d) \\ |c|^2 \\ |d|^2 \end{bmatrix}^\top \mathbf{G}_{\text{CKB}(\mathbb{P})}(A) \begin{bmatrix} 2 \operatorname{Re}(\bar{c}d) \\ 2 \operatorname{Im}(\bar{c}d) \\ |c|^2 \\ |d|^2 \end{bmatrix}.$$

Lemma 2.18. *Suppose that f is a Möbius transformation $f : \lambda \mapsto \frac{a\lambda+b}{c\lambda+d}$, $a, b, c, d \in \mathbb{C}$, $ad - bc \neq 0$, and A is 2×2 complex matrix such that $-\frac{d}{c}$ is not an eigenvalue of A .*

Let $R_{\text{CKB}(\mathbb{P})}(f)$ be a matrix of determinant 1 representing the projective action of f in $\text{CKB}(\mathbb{P})$. Then, the following transformation rules hold:

$$(41) \quad \mathbf{Q}_{\text{CKB}(\mathbb{P})}(f(A)) \cdot \mathcal{C}(f, A) = (R_{\text{CKB}(\mathbb{P})}(f))^{-1, \top} \mathbf{Q}_{\text{CKB}(\mathbb{P})}(A) (R_{\text{CKB}(\mathbb{P})}(f))^{-1};$$

$$(42) \quad \mathbf{G}_{\text{CKB}(\mathbb{P})}^0 \mathbf{Q}_{\text{CKB}(\mathbb{P})}(f(A)) \cdot \mathcal{C}(f, A) = \\ = (R_{\text{CKB}(\mathbb{P})}(f)) \mathbf{G}_{\text{CKB}(\mathbb{P})}^0 \mathbf{Q}_{\text{CKB}(\mathbb{P})}(A) (R_{\text{CKB}(\mathbb{P})}(f))^{-1};$$

$$(43) \quad \mathbf{G}_{\text{CKB}(\mathbb{P})}(f(A)) \cdot \mathcal{C}(f, A) = (R_{\text{CKB}(\mathbb{P})}(f)) \mathbf{G}_{\text{CKB}(\mathbb{P})}(A) (R_{\text{CKB}(\mathbb{P})}(f))^\top;$$

$$(44) \quad \mathbf{G}_{\text{CKB}(\mathbb{P})}(f(A)) \mathbf{Q}_{\text{CKB}(\mathbb{P})}^0 \cdot \mathcal{C}(f, A) = \\ = (R_{\text{CKB}(\mathbb{P})}(f)) \mathbf{G}_{\text{CKB}(\mathbb{P})}(A) \mathbf{Q}_{\text{CKB}(\mathbb{P})}^0 (R_{\text{CKB}(\mathbb{P})}(f))^{-1}.$$

Proof. By Theorem 2.3 and Möbius invariance, we already know that $\mathbf{Q}_{\text{CKB}(\mathbb{P})}(A)$ transforms naturally. The scaling factor in (41) is easy to recover using the general observation $\mathbf{Q}_{\text{CKB}(\mathbb{P})}(B)_{33} = 1$. Indeed, (39) is set up according to this. Equivalence to (42) follows from $\mathbf{Q}_{\text{CKB}(\mathbb{P})}^0 = (R_{\text{CKB}(\mathbb{P})}(f))^{-1, \top} \mathbf{Q}_{\text{CKB}(\mathbb{P})}^0 (R_{\text{CKB}(\mathbb{P})}(f))^{-1}$ (inverted and multiplying with). Similar argument applies regarding $\mathbf{G}_{\text{CKB}(\mathbb{P})}(A)$. The scaling factor in (43) is easy to recover using the general observation $\mathbf{G}_{\text{CKB}(\mathbb{P})}(B)_{44} = 1$. This is in correspondence to (40). Equivalence to (44) is as previously. \square

Alternative proof for Lemma 1.4. We see that axial degeneration (making $U_A \rightsquigarrow |D_A|$ while keeping the spectral part fixed) commutes with conformal transformations. \square

Lemma 2.19. *Suppose that f is a Möbius transformation $f : \lambda \mapsto \frac{a\lambda+b}{c\lambda+d}$, $a, b, c, d \in \mathbb{C}$, $ad - bc \neq 0$, and A is 2×2 complex matrix.*

(a) *If $-\frac{d}{c}$ is not an eigenvalue of A , then*

$$\mathcal{C}(f, A) > 0.$$

(b) *If $-\frac{d}{c}$ is an eigenvalue of A , then*

$$\mathcal{C}(f, A) = 0.$$

Proof. (a) From the transformation formula (41) of Lemma 2.18, $\mathcal{C}(f, A) \in \mathbb{R} \setminus \{0\}$ follows (cf. $\mathbf{Q}_{\text{CKB}(\mathbb{P})}(A)_{33} = 1$). Due to the connectedness of complex Möbius transformations, the result is > 0 . (Another argument is that the near most asymptotic points the associated quadratic form is positive.)

(b) is sufficient to check for matrices of shape $\begin{bmatrix} * & * \\ -\frac{d}{c} & * \end{bmatrix}$, which is straightforward. \square

Lemma 2.20. *It is true that*

$$U_A = \frac{1}{8} \operatorname{tr} \left(\left(\mathbf{Q}_{\text{CKB(P)}}^0 \right)^{-1} \mathbf{Q}_{\text{CKB(P)}}(A) \right).$$

Proof. This follows from Theorem 2.4, or by direct computation. \square

Lemma 2.21. *Suppose that f is a Möbius transformation $f : \lambda \mapsto \frac{a\lambda+b}{c\lambda+d}$, $a, b, c, d \in \mathbb{C}$, $ad - bc \neq 0$, and A is 2×2 complex matrix such that $-\frac{d}{c}$ is not an eigenvalue of A .*

(a) *Then,*

$$U_{f(A)} \cdot \mathcal{C}(f, A) = U_A$$

holds.

(b) *Furthermore, the transformation formula*

$$|D_{f(A)}| \cdot \mathcal{C}(f, A) = |D_A|$$

also holds.

Proof. Lemma 2.18 and Lemma 2.20 imply (a). Lemma 1.4 and (a) imply (b) generically (when $U_A > 0$). This extends by continuity. \square

Corollary 2.22. (a) *In the $U_A \neq 0$ case, the matrix is given by*

$$(45) \quad \tilde{\mathbf{Q}}_{\text{CKB(P)}}(A) = \frac{1}{U_A} \mathbf{Q}_{\text{CKB(P)}}(A)$$

has the transformation property

$$\tilde{\mathbf{Q}}_{\text{CKB(P)}}(f(A)) = \left(R_{\text{CKB(P)}}(f) \right)^{-1, \top} \tilde{\mathbf{Q}}_{\text{CKB(P)}}(A) \left(R_{\text{CKB(P)}}(f) \right)^{-1}.$$

whenever $f(A)$ makes sense.

(b) *In the $|D_A| \neq 0$ case, similar statement holds with ' U_A ' replaced by ' $|D_A|$ '.*

Proof. This is an immediate consequence of Lemma 2.18 and Lemma 2.21. \square

Remark 2.23. In the argument above we do not need to identify the value of $\mathcal{C}(f, A)$ exactly but we can start with considering it as the scaling factor of U_A . \triangle

By this corollary, we have quantities perfectly well-transforming under complex conformal transformation. However, in (a) we lose the normal parabolic case (that is scalar matrices); and in (b) we lose the whole parabolic case (two equal eigenvalues).

2.E. The geometry of the shell.

Returning to the situation of Theorem 2.2, we can describe the picture in even more geometric terms. The natural hyperbolic metric is characterized by collineation-invariance and the property that the constant ratio between the area and the angular defect is exactly 1. In the CKB model it is given by

$$(46) \quad d^{\text{CKB}}((x_{\text{CKB}}, y_{\text{CKB}}, z_{\text{CKB}}), (\tilde{x}_{\text{CKB}}, \tilde{y}_{\text{CKB}}, \tilde{z}_{\text{CKB}})) = \\ = \operatorname{arcosh} \left(\frac{1 - x_{\text{CKB}}\tilde{x}_{\text{CKB}} - y_{\text{CKB}}\tilde{y}_{\text{CKB}} - z_{\text{CKB}}\tilde{z}_{\text{CKB}}}{\sqrt{1 - (x_{\text{CKB}})^2 - (y_{\text{CKB}})^2 - (z_{\text{CKB}})^2} \sqrt{1 - (\tilde{x}_{\text{CKB}})^2 - (\tilde{y}_{\text{CKB}})^2 - (\tilde{z}_{\text{CKB}})^2}} \right).$$

In the CKB(P) model it is given by

$$(47) \quad d^{\text{CKB(P)}}((x_{\text{CKB(P)}}, y_{\text{CKB(P)}}, z_{\text{CKB(P)}}), (\tilde{x}_{\text{CKB(P)}}, \tilde{y}_{\text{CKB(P)}}, \tilde{z}_{\text{CKB(P)}})) =$$

$$= \operatorname{arcosh} \left(\frac{\frac{1}{2}z_{\text{CKB}(P)} + \frac{1}{2}\tilde{z}_{\text{CKB}(P)} - x_{\text{CKB}(P)}\tilde{x}_{\text{CKB}(P)} - y_{\text{CKB}(P)}\tilde{y}_{\text{CKB}(P)}}{\sqrt{z_{\text{CKB}(P)} - (x_{\text{CKB}(P)})^2 - (y_{\text{CKB}(P)})^2} \sqrt{\tilde{z}_{\text{CKB}(P)} - (\tilde{x}_{\text{CKB}(P)})^2 - (\tilde{y}_{\text{CKB}(P)})^2}} \right).$$

We remark that, in particular,

$$(48) \quad d^{\text{CKB}}((0, 0, 0), (s, 0, 0)) = \operatorname{arcosh} \frac{1}{\sqrt{1-s^2}} = \operatorname{arsinh} \sqrt{\frac{s^2}{1-s^2}} = \operatorname{artanh} |s| \\ = \frac{1}{2} \operatorname{arcosh} \frac{1+s^2}{1-s^2} = \frac{1}{2} \operatorname{arsinh} \frac{2|s|}{1-s^2} = \frac{1}{2} \operatorname{artanh} \frac{2|s|}{1+s^2}.$$

Theorem 2.24. *Suppose that A is a linear operator on a 2-dimensional complex Hilbert space with two distinct eigenvalues (i. e. it is non-parabolic). Then the hyperbolic radius of the (possibly degenerate) h -tube $DW_*(A)$ is*

$$\operatorname{radius}_* DW_*(A) = \frac{1}{2} \operatorname{arcosh} \frac{U_A}{|D_A|}.$$

Remark. In the parabolic non-normal case the corresponding value can be considered to be $+\infty$. In the parabolic normal (i. e. scalar matrix) case it can be taken as 0.

Proof. Let us continue the argument of the Second Proof of Theorem 2.3. After some displacement we obtained L . Then,

$$\operatorname{radius}_* DW_*(A) = \operatorname{radius}_* DW_*(L) = d^{\text{CKB}} \left((0, 0, 0), \left(0, 0, \sqrt{\frac{t^2}{1+t^2}} \right) \right) \\ = \frac{1}{2} \operatorname{arcosh} (1 + 2t^2)$$

(cf. (48/2)). Substituting (26), we obtain the result. \square

Together with the description of asymptotic points, the result above provides a characterization of the Davis–Wielandt shell in the complex non-parabolic (i. e. possibly degenerate h -tube) case. Also note that in obtaining Theorem 2.24 we have used very little information about the shell; merely conformal invariance and the description of $DW^{\mathbb{R}}(L_t)$ (the initial argument of the second proof of Theorem 2.3) was used.

As a consequence, we see (again) that $U_A/|D_A|$ is an invariant for complex Möbius transformations (as long as the spectrum allows them). Moreover, we cannot really expect to have other (unitarily invariant) ones (except its functions in the non-parabolic case and the normal / non-normal distinction in the parabolic case). Indeed, what makes tubes apart isometrically in hyperbolic geometry is just their radius, and horospheres all are isometric to each other.

Theorem 2.24 can be considered as the elliptical range theorem for the shell (with two distinct asymptotical foci). However, the case of horospheres cannot be characterized by strictly focal and metric data. The horospheric case is very special, but it requires a characterization based on other principles.

Let

$$(49) \quad Q_{\text{CKB}(P)}^{\text{spec}, A}(x_{\text{CKB}(P)}, y_{\text{CKB}(P)}, z_{\text{CKB}(P)}) = \begin{bmatrix} x_{\text{CKB}(P)} \\ y_{\text{CKB}(P)} \\ z_{\text{CKB}(P)} \\ 1 \end{bmatrix}^{\top} \mathbf{Q}_{\text{CKB}(P)}^{\text{spec}}(A) \begin{bmatrix} x_{\text{CKB}(P)} \\ y_{\text{CKB}(P)} \\ z_{\text{CKB}(P)} \\ 1 \end{bmatrix}$$

denote the quadratic expression associated to the matrix $\mathbf{Q}_{\text{CKB}(\mathcal{P})}^{\text{spec}}(A)$. We also let

$$(50) \quad \mathbf{Q}_{\text{CKB}(\mathcal{P})}^*(x_{\text{CKB}(\mathcal{P})}, y_{\text{CKB}(\mathcal{P})}, z_{\text{CKB}(\mathcal{P})}) = \begin{bmatrix} x_{\text{CKB}(\mathcal{P})} \\ y_{\text{CKB}(\mathcal{P})} \\ z_{\text{CKB}(\mathcal{P})} \\ 1 \end{bmatrix}^\top \mathbf{Q}_{\text{CKB}(\mathcal{P})}^0 \begin{bmatrix} x_{\text{CKB}(\mathcal{P})} \\ y_{\text{CKB}(\mathcal{P})} \\ z_{\text{CKB}(\mathcal{P})} \\ 1 \end{bmatrix}.$$

The same thing can be played down, with respect to $\mathbf{Q}_{\text{CKB}}^{\text{spec}}(A)$ and $\mathbf{Q}_{\text{CKB}}^0$ and the coordinates $x_{\text{CKB}}, y_{\text{CKB}}, z_{\text{CKB}}$. Now, taking (19)/(20)/(21) and (1) into account, we find that $Q_{\text{CKB}(\mathcal{P})}^{\text{spec},A}(x_{\text{CKB}(\mathcal{P})}, y_{\text{CKB}(\mathcal{P})}, z_{\text{CKB}(\mathcal{P})}) = \frac{1}{(1 - z_{\text{CKB}})^2} Q_{\text{CKB}}^{\text{spec},A}(x_{\text{CKB}}, y_{\text{CKB}}, z_{\text{CKB}})$; and similar formula holds for Q^0 . Thus, the ‘‘quadratic forms’’ are not equal, but for the quotients,

$$\frac{Q_{\text{CKB}(\mathcal{P})}^{\text{spec},A}(x_{\text{CKB}(\mathcal{P})}, y_{\text{CKB}(\mathcal{P})}, z_{\text{CKB}(\mathcal{P})})}{-2Q_{\text{CKB}(\mathcal{P})}^*(x_{\text{CKB}(\mathcal{P})}, y_{\text{CKB}(\mathcal{P})}, z_{\text{CKB}(\mathcal{P})})} = \frac{Q_{\text{CKB}}^{\text{spec},A}(x_{\text{CKB}}, y_{\text{CKB}}, z_{\text{CKB}})}{-2Q_{\text{CKB}}^*(x_{\text{CKB}}, y_{\text{CKB}}, z_{\text{CKB}})}$$

holds. (The division by -2 is merely a convenient normalization factor). These quotients are undefined for asymptotic points, but they are well-defined for interior points.

Lemma 2.25. For $(x_{\text{CKB}(\mathcal{P})}, y_{\text{CKB}(\mathcal{P})}, z_{\text{CKB}(\mathcal{P})}) \in H_{\text{CKB}(\mathcal{P})}^3$,

$$(51) \quad \frac{Q_{\text{CKB}(\mathcal{P})}^{\text{spec},A}(x_{\text{CKB}(\mathcal{P})}, y_{\text{CKB}(\mathcal{P})}, z_{\text{CKB}(\mathcal{P})})}{-2Q_{\text{CKB}(\mathcal{P})}^*(x_{\text{CKB}(\mathcal{P})}, y_{\text{CKB}(\mathcal{P})}, z_{\text{CKB}(\mathcal{P})})} \geq |D_A|;$$

and

$$(52) \quad \frac{Q_{\text{CKB}(\mathcal{P})}^{\text{spec},A}(x_{\text{CKB}(\mathcal{P})}, y_{\text{CKB}(\mathcal{P})}, z_{\text{CKB}(\mathcal{P})})}{-2Q_{\text{CKB}(\mathcal{P})}^*(x_{\text{CKB}(\mathcal{P})}, y_{\text{CKB}(\mathcal{P})}, z_{\text{CKB}(\mathcal{P})})} > 0.$$

Proof. According Lemma 2.5, (51) holds. (52) is trivial in the complex non-parabolic case ($|D_A| > 0$), and follows from the situation regarding the tangent double plane in the parabolic case. \square

Lemma 2.26. Let

$$(53) \quad \mathcal{R}_{\text{CKB}(\mathcal{P})}^A(x_{\text{CKB}(\mathcal{P})}, y_{\text{CKB}(\mathcal{P})}, z_{\text{CKB}(\mathcal{P})}) = \frac{1}{U_A} \cdot \frac{Q_{\text{CKB}(\mathcal{P})}^{\text{spec},A}(x_{\text{CKB}(\mathcal{P})}, y_{\text{CKB}(\mathcal{P})}, z_{\text{CKB}(\mathcal{P})})}{-2Q_{\text{CKB}(\mathcal{P})}^*(x_{\text{CKB}(\mathcal{P})}, y_{\text{CKB}(\mathcal{P})}, z_{\text{CKB}(\mathcal{P})})}.$$

This quantity has the transformation property

$$\mathcal{R}_{\text{CKB}(\mathcal{P})}^{f(A)}(f^{\text{CKB}(\mathcal{P})}(\mathbf{x}_{\text{CKB}(\mathcal{P})})) = \mathcal{R}_{\text{CKB}(\mathcal{P})}^A(\mathbf{x}_{\text{CKB}(\mathcal{P})}),$$

whenever $f(A)$ is well-defined. I. e., it is a truly geometrical quantity.

Proof. This is an immediate consequence of Corollary 2.22. \square

Let $Q_{\text{CKB}(\mathcal{P})}(x_{\text{CKB}(\mathcal{P})}, y_{\text{CKB}(\mathcal{P})}, z_{\text{CKB}(\mathcal{P})})$ be the form associated to $\mathbf{Q}_{\text{CKB}(\mathcal{P})}(A)$ in the scheme of (49). We can note that the quantity in the Lemma 2.26 can be replaced by

$$\frac{1}{U_A} \cdot \frac{Q_{\text{CKB}(\mathcal{P})}^A(x_{\text{CKB}(\mathcal{P})}, y_{\text{CKB}(\mathcal{P})}, z_{\text{CKB}(\mathcal{P})})}{-2Q_{\text{CKB}(\mathcal{P})}^*(x_{\text{CKB}(\mathcal{P})}, y_{\text{CKB}(\mathcal{P})}, z_{\text{CKB}(\mathcal{P})})}$$

(which is (53) and -1 added), or by

$$(54) \quad \frac{1}{|D_A|} \cdot \frac{Q_{\text{CKB}(\text{P})}^{\text{spec}, A}(x_{\text{CKB}(\text{P})}, y_{\text{CKB}(\text{P})}, z_{\text{CKB}(\text{P})})}{-2Q_{\text{CKB}(\text{P})}^*(x_{\text{CKB}(\text{P})}, y_{\text{CKB}(\text{P})}, z_{\text{CKB}(\text{P})})}$$

(which is (53) multiplied by $U_A/|D_A|$). These are equally well-behaved under complex Möbius transformations; moreover, $\text{CKB}(\text{P})$ can be replaced by CKB .

Theorem 2.27. *Let A be a complex 2×2 matrix. Then the h -distance between the point $(x_{\text{CKB}(\text{P})}, y_{\text{CKB}(\text{P})}, z_{\text{CKB}(\text{P})}) \in H_{\text{CKB}(\text{P})}^3$ and the axis of $\text{DW}_{\text{CKB}(\text{P})}(A)$ is*

$$(55) \quad d^{\text{CKB}(\text{P})}((x_{\text{CKB}(\text{P})}, y_{\text{CKB}(\text{P})}, z_{\text{CKB}(\text{P})}), \text{axis}_{\text{CKB}(\text{P})} \text{DW}_{\text{CKB}(\text{P})}(A)) = \\ = \frac{1}{2} \operatorname{arcosh} \left(\frac{1}{|D_A|} \cdot \frac{Q_{\text{CKB}(\text{P})}^{\text{spec}, A}(x_{\text{CKB}(\text{P})}, y_{\text{CKB}(\text{P})}, z_{\text{CKB}(\text{P})})}{-2Q_{\text{CKB}(\text{P})}^*(x_{\text{CKB}(\text{P})}, y_{\text{CKB}(\text{P})}, z_{\text{CKB}(\text{P})})} \right).$$

(In the parabolic case the axis is just the asymptotic point, and the value above is $+\infty$.)

Proof. Consider (54). Due to geometricity, it is sufficient to consider $A = L_t$ as in (24). Even so, we will use the CKB model. Then we find

$$\frac{1}{|D_{L_t}|} \cdot \frac{Q_{\text{CKB}}^{\text{spec}, L_t}(x_{\text{CKB}}, y_{\text{CKB}}, z_{\text{CKB}})}{-2Q_{\text{CKB}}^*(x_{\text{CKB}}, y_{\text{CKB}}, z_{\text{CKB}})} \equiv \frac{1}{1} \cdot \frac{2}{2} \cdot \frac{-(x_{\text{CKB}})^2 + (y_{\text{CKB}})^2 + (z_{\text{CKB}})^2 + 1}{-(x_{\text{CKB}})^2 - (y_{\text{CKB}})^2 - (z_{\text{CKB}})^2 + 1}.$$

On the other hand, the distance of $(x_{\text{CKB}}, y_{\text{CKB}}, z_{\text{CKB}})$ of from the axis of the tube is

$$d^{\text{CKB}}((\tilde{x}_{\text{CKB}}, \tilde{y}_{\text{CKB}}, \tilde{z}_{\text{CKB}}), (\tilde{x}_{\text{CKB}}, 0, 0)) = \frac{1}{2} \operatorname{arcosh} \left(\frac{1 - (\tilde{x}_{\text{CKB}})^2 + (\tilde{y}_{\text{CKB}})^2 + (\tilde{z}_{\text{CKB}})^2}{1 - (\tilde{x}_{\text{CKB}})^2 - (\tilde{y}_{\text{CKB}})^2 - (\tilde{z}_{\text{CKB}})^2} \right).$$

Comparing this to the RHS of the first expression, this yields the statement in the non-parabolic case. The remark concerning the parabolic case follows from limiting arguments. \square

As a corollary, we obtain

Theorem 2.28. *Consider*

$$(56) \quad \frac{1}{U_A} \cdot \frac{Q_{\text{CKB}(\text{P})}^A(x_{\text{CKB}(\text{P})}, y_{\text{CKB}(\text{P})}, z_{\text{CKB}(\text{P})})}{-2Q_{\text{CKB}(\text{P})}^*(x_{\text{CKB}(\text{P})}, y_{\text{CKB}(\text{P})}, z_{\text{CKB}(\text{P})})}.$$

If A is non-parabolic, then (56)

$$= \frac{\cosh(2 \cdot d^{\text{CKB}(\text{P})}((x_{\text{CKB}(\text{P})}, y_{\text{CKB}(\text{P})}, z_{\text{CKB}(\text{P})}), \text{axis}_{\text{CKB}(\text{P})} \text{DW}_{\text{CKB}(\text{P})}(A)))}{\cosh(2 \cdot \text{radius}_{\text{CKB}(\text{P})} \text{DW}_{\text{CKB}(\text{P})}(A))} - 1.$$

If A is parabolic, then (56)

$$= \exp\left(2 \cdot \overleftarrow{\text{dis}}^{\text{CKB}(\text{P})}((x_{\text{CKB}(\text{P})}, y_{\text{CKB}(\text{P})}, z_{\text{CKB}(\text{P})}), \text{DW}_{\text{CKB}(\text{P})}(A))\right) - 1,$$

where ' $\overleftarrow{\text{dis}}$ ' means oriented distance, negative inside the horosphere $\text{DW}_{\text{CKB}(\text{P})}(A)$, positive outside the horosphere $\text{DW}_{\text{CKB}(\text{P})}(A)$.

Proof. In the non-parabolic case, this is just Theorem 2.24 and Theorem 2.27 combined. In the parabolic case, the statement follows from limiting arguments via the formula

$$\lim_{u \rightarrow +\infty} \frac{\cosh 2(d+u)}{\cosh 2u} = \exp 2d. \quad \square$$

Theorem 2.29. *Let A be a complex 2×2 matrix. Then the signed h -distance between $(x_{\text{CKB}(\text{P})}, y_{\text{CKB}(\text{P})}, z_{\text{CKB}(\text{P})}) \in H_{\text{CKB}(\text{P})}^3$ and $\text{DW}_{\text{CKB}(\text{P})}(A)$ is*

$$\begin{aligned} & \overleftarrow{\text{dis}}_{\text{CKB}(\text{P})}((x_{\text{CKB}(\text{P})}, y_{\text{CKB}(\text{P})}, z_{\text{CKB}(\text{P})}), \text{DW}_{\text{CKB}(\text{P})}(A)) = \\ & = \text{artanh} \frac{\sqrt{\left(\frac{Q_{\text{CKB}(\text{P})}^{\text{spec}, A}(x_{\text{CKB}(\text{P})}, y_{\text{CKB}(\text{P})}, z_{\text{CKB}(\text{P})})}{-2Q_{\text{CKB}(\text{P})}^*(x_{\text{CKB}(\text{P})}, y_{\text{CKB}(\text{P})}, z_{\text{CKB}(\text{P})})} \right)^2 - |D_A|^2 - \sqrt{(U_A)^2 - |D_A|^2}}}{\frac{Q_{\text{CKB}(\text{P})}^{\text{spec}, A}(x_{\text{CKB}(\text{P})}, y_{\text{CKB}(\text{P})}, z_{\text{CKB}(\text{P})})}{-2Q_{\text{CKB}(\text{P})}^*(x_{\text{CKB}(\text{P})}, y_{\text{CKB}(\text{P})}, z_{\text{CKB}(\text{P})})} + U_A}}, \end{aligned}$$

understood so that the value is negative if the point is inside the ellipsoid, and the value is positive if it is outside.

Proof. First we assume that A is non-parabolic. Then the signed distance in question is

$$\text{d}^{\text{CKB}(\text{P})}((x_{\text{CKB}(\text{P})}, y_{\text{CKB}(\text{P})}, z_{\text{CKB}(\text{P})}), \text{axis}_{\text{CKB}(\text{P})} \text{DW}_{\text{CKB}(\text{P})}(A)) - \text{radius}_* \text{DW}_*(A).$$

Using standard addition theorems the result follows. Then the expression obtained extends to parabolic case continuously. \square

Let $O_{\text{CKB}(\text{P})} = (0, 0, 1) \in H_{\text{CKB}(\text{P})}^3$, $O_{\text{CKB}} = (0, 0, 0) \in H_{\text{CKB}}^3$, $O_{\text{PH}} = (0, 0, 1) \in H_{\text{PH}}^3$, $O_{\text{P}} = (0, 0, 0) \in H_{\text{P}}^3$. This ‘‘central point’’ O_* is characterized by being the point in $\text{DW}_* \left(\begin{bmatrix} 1 & \\ & -1 \end{bmatrix} \right) \cap \text{DW}_* \left(\begin{bmatrix} & -1 \\ 1 & \end{bmatrix} \right)$. Also, let

$$O_A = \frac{|\det A|^2}{2} + \frac{|\text{tr } A|^2}{4} + \frac{1}{2}.$$

Theorem 2.30. *Suppose that A is a 2×2 complex matrix with eigenvalues λ_1, λ_2 . Then $\text{DW}_*(A)$ is the possibly degenerate h -ellipsoid, which is tangent to the asymptotic sphere at $\iota_*(\lambda_1), \iota_*(\lambda_2)$ (double tangency counts in the parabolic case) and such that the signed distance O_* from it is*

$$(57) \quad \overleftarrow{\text{dis}}_*(O_*, \text{DW}_*(A)) = \text{artanh} \frac{\sqrt{(O_A)^2 - |D_A|^2} - \sqrt{(U_A)^2 - |D_A|^2}}{O_A + U_A}.$$

Proof. This is an immediate corollary of the previous statement. \square

This provides a characterization of $\text{DW}_*(A)$ for 2×2 matrices in fairly general terms. A similar but simpler characterization can be given in terms of the distance from ∞_* .

Distance from ∞_* is, of course, not taken as such, but it is replaced by a choice of a distance function from ∞_* . In our case, this will be the oriented distance from the horosphere \mathcal{P}^∞ given by $z_{\text{PH}} = 1$ (positive outside the horosphere, negative inside the horosphere). It is easy to see that this distance is given by

$$\begin{aligned} (58) \quad & \overleftarrow{\text{dis}}_*((x_*, y_*, z_*), \mathcal{P}_*^\infty) = -\log z_{\text{PH}} \\ & = -\log \sqrt{z_{\text{CKB}(\text{P})} - (x_{\text{CKB}(\text{P})})^2 - (y_{\text{CKB}(\text{P})})^2} \\ & = \log \frac{1 - z_{\text{CKB}}}{\sqrt{1 - (x_{\text{CKB}})^2 - (y_{\text{CKB}})^2 - (z_{\text{CKB}})^2}}. \end{aligned}$$

When we consider the oriented distance of a set M from the horosphere, then this distance is computed for any point of M , then infimum is taken.

Theorem 2.31. *Suppose that A is a 2×2 complex matrix with eigenvalues λ_1, λ_2 . Then $DW_*(A)$ is the possibly degenerate h -ellipsoid, which is tangent to the asymptotic sphere at $\iota_*(\lambda_1), \iota_*(\lambda_2)$ (double tangency counts in the parabolic case) and such that its signed distance from \mathcal{P}_*^∞ is*

$$(59) \quad \begin{aligned} \overleftarrow{\text{dis}}_*(DW_*(A), \mathcal{P}_*^\infty) &= -\log \left\| A - \frac{\text{tr } A}{2} \text{Id}_2 \right\|_2 \\ &= -\log \left(\sqrt{\frac{U_A + D_A}{2}} + \sqrt{\frac{U_A - D_A}{2}} \right) \\ &= -\frac{1}{2} \log \left(U_A + \sqrt{(U_A)^2 - (D_A)^2} \right). \end{aligned}$$

Proof. The RHS's of (59) are easy to see to be equivalent, thus it is sufficient to prove equality of the LHS to any of them.

If one can visualize an h -tube or h -horosphere in the PH model, then (59/1) is quite obvious from (58/1). Alternatively, we can continue the Second Proof of Theorem 2.3. Then the oriented distance is $-\frac{1}{2} \text{arcosh} \frac{U_A}{D_A}$ for L . As this was obtained by a displacement appearing as a linear transform with linear coefficient $\frac{1}{\sqrt{-D_A}}$, the oriented distance for A is corrected to $\overleftarrow{\text{dis}}_*(DW_*(A), \mathcal{P}_*^\infty) = -\frac{1}{2} \left(\text{arcosh} \frac{U_A}{D_A} \right) + \log \frac{1}{\sqrt{|D_A|}}$. By simple arithmetics, this can be brought to (59/3). By continuity, this extends to the parabolic case. A statement is that this data, beyond the spectral data, is sufficient to characterize the shell. This can be seen easily in either of the projective models. (Algebraically, this is equivalent to the recovery of U_A beyond the spectral data.) \square

Remark 2.32. If \mathcal{P}_1 and \mathcal{P}_2 are both horospheres, then $\overleftarrow{\text{dis}}(\mathcal{P}_1, \mathcal{P}_2) = \overleftarrow{\text{dis}}(\mathcal{P}_2, \mathcal{P}_1)$. \triangle

An even simpler characterization is as follows. Let us consider the function

$$\begin{aligned} \overleftarrow{\text{dis}}^{0\infty}_*(x_*, y_*, z_*) &= -\log \sqrt{(x_{\text{PH}})^2 + (y_{\text{PH}})^2 + (z_{\text{PH}})^2} \\ &= -\frac{1}{2} \log z_{\text{CKB}(\text{P})} \equiv -\log \sqrt{z_{\text{CKB}(\text{P})}} \\ &= -\frac{1}{2} \log \frac{1 + z_{\text{CKB}}}{1 - z_{\text{CKB}}} \equiv -\text{artanh } z_{\text{CKB}}. \end{aligned}$$

It has several interpretations; let us call it 'norm distance' now.

Theorem 2.33. *Suppose that A is a 2×2 complex matrix with eigenvalues λ_1, λ_2 . Then $DW_*(A)$ is the possibly degenerate h -ellipsoid, which is tangent to the asymptotic sphere at $\iota_*(\lambda_1), \iota_*(\lambda_2)$ (double tangency counts in the parabolic case) and such that (the infimum of) its norm distance is*

$$(60) \quad \overleftarrow{\text{dis}}^{0\infty}_*(DW_*(A)) = -\log \|A\|_2.$$

Proof. From the definition of $DW_{\text{CKB}(\text{P})}$, it follows that (60) is valid for any linear operator A (irrespective to the dimension of A). Thus the only nontrivial statement here is that this data, beyond the spectral data, is sufficient to characterize the shell. This can be seen easily in either of the projective models. \square

3. NUMERICAL RANGE

In this section, we will be interested in the numerical range primarily through its relationship to the Davis–Wielandt shell. This relationship itself is rather straightforward, as the numerical range is the vertical projection of the Davis–Wielandt shell in the CKB(P) (or in the PH) model.

If A is a normal 2×2 matrix, then it is easy to see that its numerical range is the segment connecting its eigenvalues. That leaves mainly the non-normal case for study.

Theorem 3.1. *For a non-normal complex 2×2 matrix A , the numerical range $W(A)$ is given by*

$$(61) \quad \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}^\top \mathbf{Q}^W(A) \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} \leq 0,$$

where

$$\begin{aligned} \mathbf{Q}^W(A)_{11} &= \operatorname{tr}(A^*A) - (\operatorname{Re} \operatorname{tr} A)^2 + 2 \operatorname{Re} \det A, \\ \mathbf{Q}^W(A)_{12} = \mathbf{Q}^W(A)_{21} &= -(\operatorname{Re} \operatorname{tr} A)(\operatorname{Im} \operatorname{tr} A) + 2 \operatorname{Im} \det A, \\ \mathbf{Q}^W(A)_{22} &= \operatorname{tr}(A^*A) - (\operatorname{Im} \operatorname{tr} A)^2 - 2 \operatorname{Re} \det A, \\ \mathbf{Q}^W(A)_{13} = \mathbf{Q}^W(A)_{31} &= -(\operatorname{Re} \operatorname{tr} A) \frac{\operatorname{tr}(A^*A) - |\operatorname{tr} A|^2}{2} - \operatorname{Re}((\det A) \overline{(\operatorname{tr} A)}), \\ \mathbf{Q}^W(A)_{23} = \mathbf{Q}^W(A)_{32} &= -(\operatorname{Im} \operatorname{tr} A) \frac{\operatorname{tr}(A^*A) - |\operatorname{tr} A|^2}{2} - \operatorname{Im}((\det A) \overline{(\operatorname{tr} A)}), \\ \mathbf{Q}^W(A)_{33} &= |\det A|^2 - \left(\frac{\operatorname{tr}(A^*A) - |\operatorname{tr} A|^2}{2} \right)^2. \end{aligned}$$

The matrix $\mathbf{Q}^W(A)|_{\{1,2\}}$ is positive definite.

Proof. Here we will use the fact that the numerical range is the vertical projection of the Davis–Wielandt in the CKB(P) model. Yet, even this simple computation can be presented in several ways:

(i) We can compute the discriminant of $Q_{\text{CKB(P)}}^A(x_{\text{CKB(P)}}, y_{\text{CKB(P)}}, z_{\text{CKB(P)}})$ in $z_{\text{CKB(P)}}$. This yields a quadric with matrix $-4\mathbf{Q}^W(A)$. The scalar multiple -4 , of course, can be omitted.

(ii) Consider the equation $Q_{\text{CKB(P)}}^A(x_{\text{CKB(P)}}, y_{\text{CKB(P)}}, z_{\text{CKB(P)}}) = 0$ and the projection of the solution set by the vertical projection. The preimage of the boundary is identified by the condition

$$\frac{\partial Q_{\text{CKB(P)}}^A(x_{\text{CKB(P)}}, y_{\text{CKB(P)}}, z_{\text{CKB(P)}})}{\partial z_{\text{CKB(P)}}} = 0.$$

This allows to eliminate $z_{\text{CKB(P)}}$ from the equation in order to obtain the boundary. Indeed, elimination will manifest in the linear map corresponding to the matrix

$$\mathbf{L}^W(A) = \begin{bmatrix} 1 & & & \\ & 1 & & \\ \operatorname{Re} \operatorname{tr} A & \operatorname{Im} \operatorname{tr} A & \frac{\operatorname{tr}(A^*A) - |\operatorname{tr} A|^2}{2} & \\ & & & 1 \end{bmatrix}.$$

Then,

$$\mathbf{L}^W(A)^\top \mathbf{Q}_{\text{CKB(P)}}(A) \mathbf{L}^W(A) = \mathbf{Q}^W(A).$$

(iii) We can use the fact from projective geometry that the projection is the dual of the restriction. Thus we can get the matrix of the numerical range as the inverse of the restriction of the inverse of the shell. Then, indeed,

$$(62) \quad \left(\left(\mathbf{Q}_{\text{CKB(P)}}^A(A) \right)^{-1} |_{\{1,2,4\}} \right)^{-1} \equiv \left(\frac{-1}{(U_A)^2 - |D_A|^2} \mathbf{G}_{\text{CKB(P)}}(A) |_{\{1,2,4\}} \right)^{-1} = \mathbf{Q}^W(A).$$

(The latter equality is a bit computational but not more than (63) and (64) together.)

By this we have identified the equation for the boundary of the numerical range. As the numerical range (in this non-normal case) will be a proper elliptical disk, we know that $\mathbf{Q}^W(A) |_{\{1,2\}}$ will be positive or negative definite. Checking the trace,

$$\text{tr } \mathbf{Q}^W(A) |_{\{1,2\}} = 4U_A$$

shows that it is positive definite. (If approach (i) or (ii) was taken, then the positive definiteness also follows by direct means.) This positive definiteness implies that (61) is correct with respect to the interior of the numerical range. \square

Theorem 3.2. *Suppose that A is linear operator on a 2-dimensional Hilbert space. Let us take $\mathbf{G}^W(A) := \mathbf{G}_{\text{CKB(P)}}^A(A) |_{\{1,2,4\}}$. I. e.,*

$$\mathbf{G}^W(A) = \begin{bmatrix} \frac{\text{Re det } A}{2} - \frac{\text{tr}(A^*A) - |\text{tr } A|^2}{4} & \frac{\text{Im det } A}{2} & \frac{\text{Re tr } A}{2} \\ \frac{\text{Im det } A}{2} & -\frac{\text{Re det } A}{2} - \frac{\text{tr}(A^*A) - |\text{tr } A|^2}{4} & \frac{\text{Im tr } A}{2} \\ \frac{\text{Re tr } A}{2} & \frac{\text{Im tr } A}{2} & 1 \end{bmatrix}.$$

Then

$$(63) \quad \mathbf{Q}^W(A) = -4 \text{adj } \mathbf{G}^W(A),$$

and

$$(64) \quad \det \mathbf{G}^W(A) = \frac{1}{4}((U_A)^2 - |D_A|^2).$$

Furthermore,

$$(65) \quad \det \mathbf{Q}^W(A) = -4((U_A)^2 - |D_A|^2)^2,$$

and

$$(66) \quad \text{adj } \mathbf{Q}^W(A) = 4((U_A)^2 - |D_A|^2) \mathbf{G}^W(A).$$

Proof. Equations (63) and (64) can be checked arithmetically. Then (65) and (66) follow from of properties of adj. \square

Alternative proof. In conjunction to the proof of Theorem 3.1 we can proceed as follows. We can argue that after having method (ii) carried out, method (iii) must give *exactly* the same matrix, thus equality in (62) must hold. Now, as $\mathbf{Q}^W(A) \simeq \mathbf{G}^W(A)^{-1} \simeq \text{adj } \mathbf{G}^W(A)$, equation (63) is sufficient to check out for position (3, 3) only, which is simple. Taking the validity of (62) into consideration again, this proves (64). Then (65) and (66) are also implied. Altogether, this proves the statement in the non-normal case. It extends to the normal case by continuity. \square

Remark 3.3. (a) The simplest way to present the matrix $\mathbf{Q}^W(A)$ itself is (63).

(b) We can note that matrix of the Kippenhahnian quadric

$$K_A^W(u, v, w) \equiv \det \left(u \frac{A + A^*}{2} + v \frac{A - A^*}{2i} + w \text{Id}_2 \right)$$

is $\mathbf{G}_{\text{CKB}(P)}^A(A)|_{\{1,2,4\}} \equiv \mathbf{G}^W(A)$. Thus, the direct Kippenhahnian proof of Theorem 3.1 is represented by (63) again.

(c) The center of $W(A)$ can be read off from the last column of $\mathbf{G}^W(A)$. By the continuity of the numerical range as a set (or, by direct arguments), this is also valid in the normal case. \triangle

Theorem 3.4 (Uhlig, [37]). (a)

$$\mathbf{Q}^W(A) = \begin{bmatrix} 1 & \frac{\text{Re tr } A}{2} \\ & 1 \\ & \frac{\text{Im tr } A}{2} \\ & & 1 \end{bmatrix}^{-1, \top} \underbrace{\begin{bmatrix} \mathbf{Q}^W(A)|_{\{1,2\}} & \\ & -((U_A)^2 - |D_A|^2) \end{bmatrix}}_{\mathbf{Q}^{W_0}(A):=} \begin{bmatrix} 1 & \frac{\text{Re tr } A}{2} \\ & 1 \\ & \frac{\text{Im tr } A}{2} \\ & & 1 \end{bmatrix}^{-1}.$$

(b) The eigenvalues of the (2|1)-blockdiagonal matrix $\mathbf{Q}^{W_0}(A)$ are

$$2(U_A - |D_A|), \quad 2(U_A + |D_A|), \quad | -((U_A)^2 - |D_A|^2).$$

Theorem 3.5. (a)

$$\mathbf{G}^W(A) = \begin{bmatrix} 1 & \frac{\text{Re tr } A}{2} \\ & 1 \\ & \frac{\text{Im tr } A}{2} \\ & & 1 \end{bmatrix} \underbrace{\begin{bmatrix} -\frac{1}{4} \text{adj } \mathbf{Q}^W(A)|_{\{1,2\}} & \\ & 1 \end{bmatrix}}_{\mathbf{G}^{W_0}(A):=} \begin{bmatrix} 1 & \frac{\text{Re tr } A}{2} \\ & 1 \\ & \frac{\text{Im tr } A}{2} \\ & & 1 \end{bmatrix}^{\top}.$$

(b) The eigenvalues of the (2|1)-blockdiagonal matrix $\mathbf{G}^{W_0}(A)$ are

$$-\frac{1}{2}(U_A - |D_A|), \quad -\frac{1}{2}(U_A + |D_A|), \quad | 1.$$

Proofs. (a) can easily be checked arithmetically. (b) follows easily, as the fact that the determinant and the trace matches $\mathbf{Q}^W(A)|_{\{1,2\}}$ can be checked easily. \square

(Note that the matrices $\mathbf{Q}^W(A)|_{\{1,2\}}$ and $-\frac{1}{4} \text{adj } \mathbf{Q}^W(A)|_{\{1,2\}}$ are the same as the matrices $\mathbf{Q}^C(A)$ and $\mathbf{G}^C(A)$ before.)

Theorem 3.6 (Toeplitz [36]; Johnson [20], Uhlig [37]). *If A is a 2×2 complex matrix, then $W(A)$ is the disk of the possibly degenerate ellipse, whose foci are the eigenvalues of A ($\mathbb{R}^2 \simeq \mathbb{C}$), the focal distance is $2\sqrt{|D_A|}$; and the major and minor semi-axes are*

$$\sqrt{\frac{U_A + |D_A|}{2}} \quad \text{and} \quad \sqrt{\frac{U_A - |D_A|}{2}},$$

respectively.

Proof. Firstly, we examine the non-normal case. Then the semi-axes come as the square root of (-1) times the third of eigenvalue of $\mathbf{Q}^{W_0}(A)$ divided by an other one). The

foci are trickier: It is known that if a non-degenerate quadric has matrix \mathbf{M} , then its foci are given as $(\operatorname{Re} f_i, \operatorname{Im} f_i)$ where the f_i are the roots of the quadratic equation

$$\begin{bmatrix} 1 \\ i \\ -f \end{bmatrix}^\top \mathbf{M}^{-1} \begin{bmatrix} 1 \\ i \\ -f \end{bmatrix} = 0.$$

Cf. Sommerville [34] for details; but it was already known to Plücker [33], inspiring the curve-theoretic foci. In the present case, \mathbf{M}^{-1} can be replaced by $\mathbf{G}^{\mathbf{W}}(A)$, and then the focal equation reduces to

$$f^2 - f \operatorname{tr} A + \det A = 0,$$

which is the characteristic equation. (This focal computation may look trivial in the Kippenhahnian view, but it involves the fact that the curve-theoretic foci are indeed generalizations of the ordinary foci.)

The normal case extends by continuity, or can easily be checked directly. \square

Curiously, the matrix $\mathbf{Q}^{\mathbf{W}}(A)$ of Theorem 3.1 as such does not much appear in the literature, but $\mathbf{Q}^{\mathbf{W}}(A)$ of Theorem 3.4 appears in Uhlig [37]. Theorem 3.6 is the well-known elliptical range theorem. The harder part regarding foci is due to Toeplitz [36], but he is not very explicit with the details. The semi-axes are computed explicitly in Johnson [20] (real case) and Uhlig [37]; since then they are often incorporated to the statement of the theorem, cf. Horn, Johnson [18]. One, of course, does not have to go through all these steps in order to get Theorem 3.6; see, e. g., [25] for a reduced proof. It is instructive to compare various proofs due to Toeplitz [36], Murnaghan [32], Donoghue [8], Gustafson, Rao [15], Li [26], and related ideas from Hausdorff [17], Kippenhahn [22] / [23], Dekker, cf. Halmos [16], Davis [6].

Theorem 3.7. *For 2×2 complex matrices, the five data determines the numerical range, and vice versa. (In particular, then, there is no loss of information in passing from the Davis–Wielandt shell to the numerical range.)*

Proof. The five data determines the matrix up to unitary conjugation, thus the numerical range. Conversely, if $\mathbf{W}(A)$ is given, then one recover the spectral data from the foci, and using either the major or minor semi-axes, one can get U_A , and hence $\operatorname{tr} A^*A$. \square

From Theorem 3.4 and Theorem 3.5, one can conclude immediately that

$$\operatorname{rk} \mathbf{G}^{\mathbf{W}}(A) = \begin{cases} 3 & \text{if } A \text{ is non-normal,} \\ 2 & \text{if } A \text{ is normal non-parabolic,} \\ 1 & \text{if } A \text{ is normal parabolic;} \end{cases}$$

and

$$\operatorname{rk} \mathbf{Q}^{\mathbf{W}}(A) = \begin{cases} 3 & \text{if } A \text{ is non-normal,} \\ 1 & \text{if } A \text{ is normal non-parabolic,} \\ 0 & \text{if } A \text{ is normal parabolic.} \end{cases}$$

Lemma 3.8. *Assume that the 2×2 complex matrix A is non-normal, and \mathbf{M} is the matrix of the quadric of the boundary of $\mathbf{W}(A)$. Then*

(a)

$$(67) \quad \mathbf{G}^{\mathbf{W}}(A) = \frac{\mathbf{M}^{-1}}{(\mathbf{M}^{-1})_{33}} = \frac{\operatorname{adj} \mathbf{M}}{\det \mathbf{M}_{\{1,2\}}};$$

(b)

$$\mathbf{Q}^{\mathbf{W}}(A) = \frac{-4}{(\det \mathbf{M})((\mathbf{M}^{-1})_{33})^2} \cdot \mathbf{M} = \frac{-4 \det \mathbf{M}}{(\det \mathbf{M}|_{\{1,2\}})^2} \cdot \mathbf{M} = \frac{-4 \operatorname{adj} \operatorname{adj} \mathbf{M}}{\det \mathbf{M}|_{\{1,2\}}^2}.$$

Proof. (a) follows from $\mathbf{G}^{\mathbf{W}}(A) \cong \mathbf{M}^{-1}$ and $\mathbf{G}^{\mathbf{W}}(A)_{33} = 1$. Then (63) implies (b). \square

Lemma 3.9. *Assume that A is a normal complex 2×2 matrix with eigenvalues λ_1, λ_2 .*

(a)

$$(68) \quad \mathbf{G}^{\mathbf{W}}(A) = \frac{1}{2} \left(\begin{bmatrix} \operatorname{Re} \lambda_1 \\ \operatorname{Im} \lambda_1 \\ 1 \end{bmatrix} \begin{bmatrix} \operatorname{Re} \lambda_2 \\ \operatorname{Im} \lambda_2 \\ 1 \end{bmatrix}^{\top} + \begin{bmatrix} \operatorname{Re} \lambda_2 \\ \operatorname{Im} \lambda_2 \\ 1 \end{bmatrix} \begin{bmatrix} \operatorname{Re} \lambda_1 \\ \operatorname{Im} \lambda_1 \\ 1 \end{bmatrix}^{\top} \right).$$

(b)

$$\mathbf{Q}^{\mathbf{W}}(A) = \mathbf{v}\mathbf{v}^{\top}$$

where

$$\mathbf{v} = \begin{bmatrix} \operatorname{Im} \lambda_1 - \operatorname{Im} \lambda_2 \\ \operatorname{Re} \lambda_2 - \operatorname{Re} \lambda_1 \\ (\operatorname{Re} \lambda_1)(\operatorname{Im} \lambda_2) - (\operatorname{Re} \lambda_2)(\operatorname{Im} \lambda_1) \end{bmatrix} = \begin{bmatrix} \operatorname{Re} \lambda_1 \\ \operatorname{Im} \lambda_1 \\ 1 \end{bmatrix} \times \begin{bmatrix} \operatorname{Re} \lambda_2 \\ \operatorname{Im} \lambda_2 \\ 1 \end{bmatrix}$$

(with \times meaning the ordinary vector product of Gibbs).

Proof. We can compute the data from $A = \begin{bmatrix} \lambda_1 & \\ & \lambda_2 \end{bmatrix}$. \square

Alternative proof for Theorem 3.7. In the \Leftarrow direction: $\mathbf{G}^{\mathbf{W}}(A)$ can be recovered from (67) / (68), and from this matrix the five data can be read off immediately. \square

As we have seen, $\mathbf{G}^{\mathbf{W}}(A)$ involves no loss of data relative to the five data, not even in the normal case. Regarding $\mathbf{Q}^{\mathbf{W}}(A)$:

Lemma 3.10. *Assume that A is complex 2×2 matrix.*

(a) *If A is normal with two distinct eigenvalues, then $\mathbf{Q}^{\mathbf{W}}(A)$ is a matrix of rank strictly 1. Geometrically, it corresponds to the (double) line containing the segment which is the numerical range. Beyond that, $\mathbf{Q}^{\mathbf{W}}(A)$, as it is, also contains the information of the distance of the eigenvalues $2\sqrt{|D_A|}$, but essentially not more.*

(b) *If A is normal with two equal eigenvalues, then $\mathbf{Q}^{\mathbf{W}}(A) = 0$.*

Proof. Let us consider the description in Lemma 3.9(b) with $\mathbf{v} \equiv \mathbf{v}_{\mathbf{W}}(\lambda_1, \lambda_2)$.

(b) If λ_1 and λ_2 are equal, then $(\operatorname{Re} \lambda_1, \operatorname{Im} \lambda_1) = (\operatorname{Re} \lambda_2, \operatorname{Im} \lambda_2)$, and $\mathbf{v}_{\mathbf{W}}(\lambda_1, \lambda_2)$ vanishes. This leads to the statement.

(a) If λ_1 and λ_2 are not equal, then $(\operatorname{Re} \lambda_1, \operatorname{Im} \lambda_1) \neq (\operatorname{Re} \lambda_2, \operatorname{Im} \lambda_2)$, and $\mathbf{v}_{\mathbf{W}}(\lambda_1, \lambda_2) \neq 0$. This leads to a double line for $\mathbf{Q}^{\mathbf{W}}(A)$. Then, by the construction of $\mathbf{v}_{\mathbf{W}}(\lambda_1, \lambda_2)$, it is easy to see that the line fits to the distinct points $(\operatorname{Re} \lambda_1, \operatorname{Im} \lambda_1)$ and $(\operatorname{Re} \lambda_2, \operatorname{Im} \lambda_2)$. From $\mathbf{Q}^{\mathbf{W}}(A)$ one can reconstruct $\pm \mathbf{v}_{\mathbf{W}}(\lambda_1, \lambda_2)$. If $\mathbf{v}_{\mathbf{W}}(\lambda_1, \lambda_2) = \pm(u_1, u_2, u_3)$, then $2\sqrt{|D_A|} = |\lambda_1 - \lambda_2| = \sqrt{(u_1)^2 + (u_2)^2}$, and this information is specific with respect to the scaling of $\pm \mathbf{v}_{\mathbf{W}}(\lambda_1, \lambda_2)$. (In particular, in the normal setting, we have the formula

$$2\sqrt{|D_A|} = \sqrt{\operatorname{tr} \left(\begin{bmatrix} 1 & \\ & 1 \\ & & 0 \end{bmatrix} \mathbf{Q}^{\mathbf{W}}(A) \right)}. \quad \square$$

Remark 3.11. We can consider the assignment

$$\widehat{\mathbf{Q}}^W(A) = \frac{1}{2U_A} \mathbf{Q}^W(A)$$

defined (here originally) for non-scalar matrices A .

(a) If A is normal and non-parabolic, then $\widehat{\mathbf{Q}}^W(A)$ contains no more information than the line of the eigenvalues.

(b) If $A = \lambda \text{Id}_2$ is a scalar matrix, then it is reasonable to define

$$\widehat{\mathbf{Q}}^W(\lambda \text{Id}_2) = \begin{bmatrix} 1 & 0 & -\text{Re } \lambda \\ 0 & 1 & -\text{Im } \lambda \\ -\text{Re } \lambda & -\text{Im } \lambda & |\lambda|^2 \end{bmatrix}.$$

This yields a natural but non-continuous extension of the original assignment. Then, in general,

$$\text{adj } \widehat{\mathbf{Q}}^W(A) = \left(1 - \left(\frac{|D_A|}{U_A} \right)^2 \right) \mathbf{G}^W(A)$$

holds with $\frac{0}{0} = 0$.

△

4. SPECTRAL INVARIANTS OF 2×2 COMPLEX MATRICES

4.A. Spectral type.

We extend the elliptic/parabolic/hyperbolic classification of real 2×2 matrices to complex 2×2 matrices as follows. We distinguish the following classes:

- real-elliptic case: two conjugate, strictly complex eigenvalues,
- real-parabolic case: two equal real eigenvalues,
- real-hyperbolic case: two distinct real eigenvalues,
- non-real parabolic case: two equal strictly complex eigenvalues,
- semi-real case: a real and a strictly complex eigenvalue,
- quasielliptic case: two non-conjugate eigenvalues λ_1, λ_2 with $(\text{Im } \lambda_1)(\text{Im } \lambda_2) < 0$,
- quasihyperbolic case: two distinct eigenvalues λ_1, λ_2 with $(\text{Im } \lambda_1)(\text{Im } \lambda_2) > 0$.

One can see that the classes above are closed for conjugation by unitary matrices and for real fractional linear (i. e. Möbius) transformations, whenever they are applicable.

4.B. Spectral quantities.

A quantity corresponding to $|D_A|$ is

$$E_A = \left(\frac{(\text{Im } \text{tr } A)}{2} \right)^2 + \frac{|D_A| - \text{Re } D_A}{2}.$$

Indeed, if λ_1 and λ_2 are the eigenvalues of A , then it is easy to see that

$$|D_A| = \left| \frac{\lambda_1 - \lambda_2}{2} \right|^2$$

and

$$E_A = \left| \frac{\lambda_1 - \bar{\lambda}_2}{2} \right|^2;$$

i. e. one corresponds to the other after conjugating an eigenvalue.

A related quantity is

$$H_A = \left(\frac{(\operatorname{Im} \operatorname{tr} A)}{2} \right)^2 - \frac{|D_A| + \operatorname{Re} D_A}{2} = E_A - |D_A|.$$

It is easy to see that

$$H_A = (\operatorname{Im} \lambda_1)(\operatorname{Im} \lambda_2).$$

Another related quantity is

$$K_A = \left(\frac{(\operatorname{Im} \operatorname{tr} A)}{2} \right)^2 + \frac{|D_A| + \operatorname{Re} D_A}{2}.$$

It is easy to see that

$$K_A = \frac{(\operatorname{Im} \lambda_1)^2 + (\operatorname{Im} \lambda_2)^2}{2}.$$

Then one can characterize the 2×2 complex spectral classes in terms of the signs of the quantities $|D_A|$, E_A , K_A , and H_A :

type	$ D_A $	E_A	K_A	H_A
real-elliptic	+	0	+	-
real-parabolic	0	0	0	0
real-hyperbolic	+	+	0	0
non-real parabolic	0	+	+	+
semi-real	+	+	+	0
quasielliptic	+	+	+	-
quasihyperbolic	+	+	+	+

As the spectral classes are closed to real fractional transformations (whenever they are applicable), the signs of $|D_A|$, E_A , K_A , and H_A are also invariant. However, a finer invariant is shown by

Lemma 4.1. *The ratio*

$$E_A : |D_A| = |\lambda_1 - \bar{\lambda}_2|^2 : |\lambda_1 - \lambda_2|^2$$

is invariant for real fractional transformations (whenever they are applicable to A).

Proof. It is sufficient to prove that they are invariant to the transformations $f : \lambda \mapsto \lambda + b$ ($b \in \mathbb{R}$) and (whenever applicable) $f : \lambda \mapsto 1/\lambda$; and these are easy to see. \square

In fact, we also see that the quantity

$$\frac{E_A - |D_A|}{E_A + |D_A|}$$

will act as an improved, real Möbius invariant version of the classifier $(\operatorname{sgn})H_A$.

Remark 4.2. (a) If A is in canonical form (7), then the five data and U_A , $|D_A|$, E_A , K_A , H_A are all polynomial in $\operatorname{Re} \lambda_1$, $\operatorname{Im} \lambda_1$, $\operatorname{Re} \lambda_2$, $\operatorname{Im} \lambda_2$, t ; thus computation with all this data is very simple in that way.

(b) U_A , $|D_A|^2$, $2E_A - |D_A|$, $2H_A + |D_A|$, $2K_A - |D_A|$ are polynomial in the five data. Thus, in terms of the more invariant five data, computation is a bit less simple. \triangle

4.C. The canonical representatives (real Möbius).

Lemma 4.3. *For complex 2×2 matrices up to real Möbius transformations and unitary conjugation it is sufficient to consider the following cases:*

$$(69) \quad \mathbf{0}_2 = \begin{bmatrix} 0 & \\ & 0 \end{bmatrix}$$

(real-parabolic, normal) ;

$$(70) \quad S_\beta = \begin{bmatrix} 0 & \cos \beta \\ i \sin \beta & \end{bmatrix} \quad \beta \in \left[0, \frac{\pi}{2}\right]$$

($\beta = 0$: real-parabolic non-normal, $0 < \beta < \pi/2$: semi-real non-normal, $\beta = \pi/2$: semi-real normal);

$$(71) \quad L_{\alpha,t}^\pm = \begin{bmatrix} \cos \alpha + i \sin \alpha & 2t \\ -\cos \alpha \pm i \sin \alpha & \end{bmatrix} \quad \alpha \in \left[0, \frac{\pi}{2}\right], t \geq 0$$

but the cases $L_{0,t}^+$ and $L_{0,t}^-$ are identical

($\alpha = 0$: real-hyperbolic case, $0 < \alpha < \pi/2$: quasielliptic $[-]$ / quasihyperbolic case $[+]$, $\alpha = \pi/2$: real-elliptic $[-]$ / non-real parabolic case $[+]$; $t = 0$: normal, $t > 0$ non-normal).

Apart from the degeneracy for $\alpha = 0$, the representatives above are inequivalent.

Proof. It is sufficient to follow the image of the eigenvalues in the Poincaré model, and apply the normal form up to unitary conjugation. By this, we arrive to the cases

$$(72) \quad \widehat{0}_t = \begin{bmatrix} 0 & 2t \\ & 0 \end{bmatrix} \quad t \geq 0,$$

$$(73) \quad \widehat{S}_t^\pm = \begin{bmatrix} 0 & 2t \\ & \pm i \end{bmatrix} \quad t \geq 0,$$

$$(74) \quad \widehat{L}_{\alpha,t}^{\pm 1, \pm 2} = \begin{bmatrix} \cos \alpha \pm_1 i \sin \alpha & 2t \\ -\cos \alpha \pm_2 i \sin \alpha & \end{bmatrix} \quad \alpha \in \left[0, \frac{\pi}{2}\right], t \geq 0.$$

Some cases of signs can be eliminated: Multiplying by -1 and applying unitary conjugation, we see that \widehat{S}_t^- and \widehat{S}_t^+ are equivalent; we eliminate \widehat{S}_t^- . Similarly, taking inverse and applying unitary conjugation, we see that $\widehat{L}_{\alpha,t}^{\pm 1, \pm 2}$ and $\widehat{L}_{\alpha,t}^{\mp 1, \mp 2}$ are equivalent; we eliminate $\widehat{L}_{\alpha,t}^{-, \pm 2}$. Now, $\widehat{0}_0$ is just $\mathbf{0}_2$. $\widehat{0}_t$ with $t > 0$ scales into S_0 . \widehat{S}_t^+ scales into S_β with $\beta \in (0, \pi/2]$. $\widehat{L}_{\alpha,t}^{+, \pm 2}$ serves $L_{\alpha,t}^{\pm 2}$.

From the location of the eigenvalues, it is not hard to see that representatives from different families are inequivalent. Inside the families, the ratio $U_A : |D_A| : E_A$ distinguishes the various representatives as the following example shows. \square

Example 4.4. (a) Regarding $L_{\alpha,t}^\pm$:

$$U_{L_{\alpha,t}^+} = 2t^2 + (\cos \alpha)^2, \quad |D_{L_{\alpha,t}^+}| = (\cos \alpha)^2, \quad E_{L_{\alpha,t}^+} = 1, \quad E_{L_{\alpha,t}^+} \geq |D_{L_{\alpha,t}^+}|;$$

and

$$U_{L_{\alpha,t}^-} = 2t^2 + 1, \quad |D_{L_{\alpha,t}^-}| = 1, \quad E_{L_{\alpha,t}^-} = (\cos \alpha)^2, \quad E_{L_{\alpha,t}^-} \leq |D_{L_{\alpha,t}^-}|.$$

(b) Regarding S_β :

$$U_{S_\beta} = \frac{1}{4} (1 + (\cos \beta)^2), \quad |D_{S_\beta}| = E_{S_\beta} = \frac{1}{4} (1 - (\cos \beta)^2).$$

(c) In case of $\mathbf{0}_2$:

$$U_{\mathbf{0}_2} = |D_{\mathbf{0}_2}| = E_{\mathbf{0}_2} = 0.$$

Corollary 4.5. *The triple ratio*

$$(75) \quad U_A : |D_A| : E_A$$

and (when $|D_A| = E_A > 0$) the possible choice of

real-hyperbolic / semi-real type

together form a full invariant of 2×2 complex matrices with respect to equivalence by real Möbius transformations and unitary conjugation.

Proof. (75) almost distinguishes the canonical representatives of Lemma 4.3. The only ambiguity is between $L_t \equiv L_{0,t}$ and S_β when $\cos \beta = \frac{t}{\sqrt{1+t^2}}$. \square

The content of the following three subsections will be considered only in Remark 5.46.

4.D. The square root.

Let $\widetilde{\text{sgn}}$ denote the half-sided sign function, for which

$$\widetilde{\text{sgn}}(x) = \begin{cases} 1 & \text{if } x \geq 0, \\ -1 & \text{if } x < 0. \end{cases}$$

Then, according to our conventions, the canonical square root we use for complex numbers, i. e. the square root cut along $(-\infty, 0) - i\epsilon$, is given as

$$(76) \quad \sqrt{z} = \sqrt{\frac{|z| + \text{Re } z}{2}} + i \cdot \widetilde{\text{sgn}}(\text{Im } z) \sqrt{\frac{|z| - \text{Re } z}{2}}.$$

We remark that two trivial but sometimes useful identities are

$$\text{Re}(a\sqrt{z}) \text{Re}(b\sqrt{z}) = \text{Im}(a\sqrt{-z}) \text{Im}(b\sqrt{-z}),$$

and

$$\text{Re}(a\sqrt{z}) \text{Im}(b\sqrt{z}) = -\text{Im}(a\sqrt{-z}) \text{Re}(b\sqrt{-z}).$$

4.E. The spectrum.

Using (76), the points of the spectrum of the 2×2 matrix A can be described as

$$\frac{1}{2} \text{tr } A \pm \sqrt{-D_A} = \frac{1}{2} \text{tr } A \pm \left(\sqrt{\frac{|D_A| - \text{Re } D_A}{2}} + \widetilde{\text{sgn}}(-\text{Im } D_A) i \sqrt{\frac{|D_A| + \text{Re } D_A}{2}} \right).$$

4.F. Spectral quantities (one more).

Let us set

$$B_A = \frac{\text{Im tr } A}{2} + i \text{Im } \sqrt{D_A}.$$

It is easy to see that $|B_A|^2 = E_A$, but B_A is less naturally defined than E_A . (B_A is an analogue of $\sqrt{D_A}$.)

Lemma 4.6. *Assume that A has eigenvalues λ_1 and λ_2 such that $\text{Im } \lambda_1 > 0 > \text{Im } \lambda_2$. Then $D_A \not\leq 0$ and*

$$B_A = \frac{\lambda_1 - \bar{\lambda}_2}{2i}.$$

In this case $B_{A^} = -B_A$ also holds.*

Proof. This follows from the formulas

$$\begin{aligned} \sqrt{D_A} &= \sqrt{\frac{|D_A| + \text{Re } D_A}{2}} + \widetilde{\text{sgn}}(\text{Im } D_A) i \sqrt{\frac{|D_A| - \text{Re } D_A}{2}}, \\ \lambda_1 &= \frac{1}{2} \text{tr } A + \left(\widetilde{\text{sgn}}(-\text{Im } D_A) \sqrt{\frac{|D_A| - \text{Re } D_A}{2}} + i \sqrt{\frac{|D_A| + \text{Re } D_A}{2}} \right), \\ \lambda_2 &= \frac{1}{2} \text{tr } A - \left(\widetilde{\text{sgn}}(-\text{Im } D_A) \sqrt{\frac{|D_A| - \text{Re } D_A}{2}} + i \sqrt{\frac{|D_A| + \text{Re } D_A}{2}} \right); \end{aligned}$$

and the fact that here $\text{Im } D_A = 0$ implies $D_A > 0$. \square

5. THE CONFORMAL RANGE OF 2×2 COMPLEX MATRICES

5.A. The qualitative elliptical range theorem (review).

For a complex linear operator A , the conformal range / real Davis-Wielandt shell is related to the Davis–Wielandt shell by

$$\text{DW}_*^{\mathbb{R}}(A) = \pi_*^{[2]}(\text{DW}_*(A)),$$

where $*$ may mean any of the models, CKB, CKB(P), PH, P, and $\pi_*^{[2]} : \overline{H_*^3} \rightarrow \overline{H_*^2}$ is a canonical projection from the hyperbolic 3-space to the hyperbolic 2-space. In the CKB and CKB(P) models, $\pi_*^{[2]}$ means the elimination of the second coordinate, or, in other terms, setting it to 0. (It is slightly more complicated in the conformal models.) Geometrically, this is an h -orthogonal projection from the h -space $\overline{H_*^3}$ to the canonically embedded h -plane $i^{[2]}(\overline{H_*^2})$ (asymptotically closed). Let $\iota_*^{[2]} = \pi_*^{[2]} \circ \iota_*$, the mapping of the Riemann-sphere to the asymptotically closed hyperbolic plane. Then

$$\iota_{\text{CKB(P)}}^{[2]}(\lambda) = (\text{Re } \lambda, |\lambda|^2) \quad \text{and} \quad \iota_{\text{CKB}}^{[2]}(\lambda) = \left(\frac{2 \text{Re } \lambda}{1 + |\lambda|^2}, \frac{1 - |\lambda|^2}{1 + |\lambda|^2} \right).$$

(In terms of PH this is just possible conjugation into the upper half-plane.)

According to Theorem 2.2, the conformal range is just a projection of (a possible degenerate) h -tube or h -horosphere to the h -plane $i^{[2]}(\overline{H_*^2})$. The situation is easy to visualize in the CKB and CKB(P) models. In [24], we have seen that for a real 2×2 matrix A acting on \mathbb{R}^2 the conformal range $\text{DW}_*^{\mathbb{R}}(A^{\mathbb{R}})$ either a h -circle, an h -horosphere, or a pair of h -hypercycles (asymptotically closed, and possibly degenerate); corresponding to the elliptic, parabolic, and hyperbolic case, respectively. In the complexified setting, we obtain an h -disk, an h -horodisk, or a distance band, respectively. It is easy to see that in terms of the shell those correspond either to the case of an h -tube with axis perpendicular to $i^{[2]}(\overline{H_*^2})$, or an h -horosphere perpendicular to $i^{[2]}(\overline{H_*^2})$, or an h -tube with axis lying on $i^{[2]}(\overline{H_*^2})$ respectively.

5.B. General observations.

Complex 2×2 matrixes are determined, up to unitary conjugation, by the ‘five data’ (11). Thus, these determine the Davis-Wielandt shell, and, in particular, the conformal range. Now, in the case of the conformal range, slightly less is sufficient.

Lemma 5.1. *Suppose that A_1 and A_2 are two complex 2×2 matrices. We claim that*

$$\text{CR}(A_1) = \text{CR}(A_2)$$

holds if and only if for all $\lambda \in \mathbb{R}$,

$$(77) \quad |\det(A_1 - \lambda \text{Id}_2)|^2 = |\det(A_2 - \lambda \text{Id}_2)|^2,$$

and

$$(78) \quad \text{tr}(A_1 - \lambda \text{Id}_2)^*(A_1 - \lambda \text{Id}_2) = \text{tr}(A_2 - \lambda \text{Id}_2)^*(A_2 - \lambda \text{Id}_2).$$

hold.

Proof. According to the dual viewpoint explained in [24], the information $\text{CR}(A)$ is equivalent to the information $\lambda \in \mathbb{R} \mapsto (\|A - \lambda \text{Id}_2\|_2, \|A - \lambda \text{Id}_2\|_2^-)$. By Lemma 1.8, this is equivalent to the information $\lambda \in \mathbb{R} \mapsto (\text{tr}(A - \lambda \text{Id}_2)^*(A - \lambda \text{Id}_2), |\det(A - \lambda \text{Id}_2)|^2)$. \square

It is easy to see, that if A is a complex 2×2 matrix, $\lambda \in \mathbb{R}$, then

$$\begin{aligned} |\det(A - \lambda \text{Id}_2)|^2 &= \\ &= \lambda^4 - (2 \text{Re tr } A)\lambda^3 + (|\text{tr } A|^2 + 2 \text{Re det } A)\lambda^2 - 2(\text{Re}((\det A)\overline{\text{tr } A}))\lambda + |\det A|^2, \end{aligned}$$

and

$$\text{tr}(A - \lambda \text{Id}_2)^*(A - \lambda \text{Id}_2) = 2\lambda^2 - (2 \text{Re tr } A)\lambda + \text{tr}(A^*A).$$

Theorem 5.2. *For complex 2×2 matrices A , the information contained conformal range is in bijective correspondence to the ‘reduced five data’*

$$\text{Re tr } A, \quad |\text{tr } A|^2 + 2 \text{Re det } A, \quad \text{Re}((\det A)\overline{\text{tr } A}), \quad |\det A|^2, \quad \text{tr}(A^*A).$$

Proof. Considering the polynomials (77) and (78) in λ , the information contained in them is the same as in their list of coefficients. \square

5.C. The matrix of the conformal range.

If A is normal, then \mathbb{C}^2 is the orthogonal direct sum of its complex eigenspaces. Thus $\text{DW}_*^{\mathbb{R}}(A)$ is the (possibly degenerate) h -segment connecting the points corresponding to the eigenvalues of A . In the non-normal case:

Theorem 5.3. *Suppose that A is linear operator on a 2-dimensional Hilbert space. Assume that A is not normal. Then $\text{DW}_{\text{CKB(P)}}^{\mathbb{R}}(A)$ is given by the quadratic inequality*

$$(79) \quad \begin{bmatrix} x_{\text{CKB(P)}} \\ z_{\text{CKB(P)}} \\ 1 \end{bmatrix}^{\top} \mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A) \begin{bmatrix} x_{\text{CKB(P)}} \\ z_{\text{CKB(P)}} \\ 1 \end{bmatrix} \leq 0$$

such that

$$\mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A) = \begin{bmatrix} Z^2 - 4Y & 2X - VZ & 2VY - XZ \\ 2X - VZ & Z - W + V^2 & \frac{WZ}{2} - VX - \frac{Z^2}{2} \\ 2VY - XZ & \frac{WZ}{2} - VX - \frac{Z^2}{2} & X^2 - YW + YZ \end{bmatrix},$$

where

$$(80) \quad \begin{aligned} V &\equiv \operatorname{Re} \operatorname{tr} A, \\ W &\equiv |\operatorname{tr} A|^2 + 2(\operatorname{Re} \det A), \\ X &\equiv \operatorname{Re}((\det A)(\overline{\operatorname{tr} A})), \\ Y &\equiv |\det A|^2, \\ Z &\equiv \operatorname{tr} A^* A \end{aligned}$$

is the ‘reduced’ five data.

In this non-normal case, $\mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A)|_{\{1,2\}}$ is positive definite.

Remark. In (79), replacing

$$\begin{bmatrix} x_{\text{CKB(P)}} \\ z_{\text{CKB(P)}} \\ 1 \end{bmatrix} \text{ by } \begin{bmatrix} x_{\text{CKB}} \\ 1 + z_{\text{CKB}} \\ 1 - z_{\text{CKB}} \end{bmatrix}, \begin{bmatrix} x_{\text{P}} \\ \frac{(x_{\text{P}})^2 + (z_{\text{P}+1})^2}{2} \\ \frac{(x_{\text{P}})^2 + (z_{\text{P}-1})^2}{2} \end{bmatrix} \begin{bmatrix} x_{\text{PH}} \\ (x_{\text{PH}})^2 + (z_{\text{PH}})^2 \\ 1 \end{bmatrix},$$

one obtains the corresponding equations in CKB, P, PH, respectively.

In the case of the CKB model, this allows to take the viewpoint that

$$(81) \quad \begin{bmatrix} x_{\text{CKB}} \\ 1 + z_{\text{CKB}} \\ 1 - z_{\text{CKB}} \end{bmatrix}^{\top} \mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A) \begin{bmatrix} x_{\text{CKB}} \\ 1 + z_{\text{CKB}} \\ 1 - z_{\text{CKB}} \end{bmatrix} \equiv \begin{bmatrix} x_{\text{CKB}} \\ z_{\text{CKB}} \\ 1 \end{bmatrix}^{\top} \mathbf{Q}_{\text{CKB}}^{\mathbb{R}}(A) \begin{bmatrix} x_{\text{CKB}} \\ z_{\text{CKB}} \\ 1 \end{bmatrix}$$

where

$$(82) \quad \mathbf{Q}_{\text{CKB}}^{\mathbb{R}}(A) = \begin{bmatrix} 1 & & \\ & 1 & 1 \\ & -1 & 1 \end{bmatrix}^{\top} \mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A) \begin{bmatrix} 1 & & \\ & 1 & 1 \\ & -1 & 1 \end{bmatrix}$$

is obtained canonically. This is similar to the case of the Davis–Wielandt shell.

A proof of Theorem 5.3 via the projection approach. Here we will use the fact the conormal range is a projection of the Davis–Wielandt shell. Again, this projection can be obtained in various ways:

(i) We can take the discriminant of $Q_{\text{CKB(P)}}^A(x_{\text{CKB(P)}}, y_{\text{CKB(P)}}, z_{\text{CKB(P)}})$ in variable $y_{\text{CKB(P)}}$. Then obtain the quadric with matrix $-4\mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A)$.

(ii) Consider the equation $Q_{\text{CKB(P)}}^A(x_{\text{CKB(P)}}, y_{\text{CKB(P)}}, z_{\text{CKB(P)}}) = 0$ and the projection of the solution set by $\pi_{\text{CKB(P)}}^{[2]}$. The preimage of the boundary is identified by the condition

$$\frac{\partial Q_{\text{CKB(P)}}^A(x_{\text{CKB(P)}}, y_{\text{CKB(P)}}, z_{\text{CKB(P)}})}{\partial y_{\text{CKB(P)}}} = 0.$$

This allows to eliminate $y_{\text{CKB(P)}}$ from the equation in order to obtain the boundary. Indeed, elimination will manifest in the linear map corresponding the matrix

$$\mathbf{L}_{\text{CKB(P)}}^{\mathbb{R}}(A) = \begin{bmatrix} 1 & & & \\ -2 \operatorname{Im} \det A & \operatorname{Im} \operatorname{tr} A & \operatorname{Re}((\det A) \overline{(\operatorname{tr} A)}) & \\ \operatorname{tr}(A^* A) - 2 \operatorname{Re} \det A & \operatorname{tr}(A^* A) - 2 \operatorname{Re} \det A & \operatorname{tr}(A^* A) - 2 \operatorname{Re} \det A & \\ & 1 & & 1 \end{bmatrix}.$$

(Note that $\operatorname{tr}(A^* A) - 2 \operatorname{Re} \det A > 0$ if A is non-normal.) Then, we obtain

$$(83) \quad (\mathbf{L}_{\text{CKB(P)}}^{\mathbb{R}}(A))^{\top} \mathbf{Q}_{\text{CKB(P)}}(A) \mathbf{L}_{\text{CKB(P)}}^{\mathbb{R}}(A) = \frac{1}{\operatorname{tr}(A^* A) - 2 \operatorname{Re} \det A} \cdot \mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A).$$

After that, the scalar factor is omitted.

(iii) Again, we can use the fact from projective geometry that the projection is the dual of the restriction. Thus we can get the matrix of the conformal range as the inverse of the restriction of the inverse of the matrix of the shell. Then, indeed,

$$(84) \quad \left(\left(\mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A) \right)^{-1} |_{\{1,3,4\}} \right)^{-1} \equiv \left(\frac{-1}{(U_A)^2 - |D_A|^2} \mathbf{G}_{\text{CKB(P)}}(A) |_{\{1,3,4\}} \right)^{-1} \\ = \frac{1}{\operatorname{tr}(A^* A) - 2 \operatorname{Re} \det A} \mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A).$$

(This is somewhat computational but not more than (85) and (86) together.)

In this non-normal case we know that the conformal range is a non-degenerate ellipse, thus $\mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A) |_{\{1,2\} \times \{1,2\}}$ is either positive definite or negative definite. However, it is easy to see that $\mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A)_{11} = Z - 4Y = \operatorname{tr}(A^* A)^2 - 4|\det A|^2 \geq 0$, thus it will be positive definite. In particular, the specification of the interior in (79) is also correct. (In cases (i) and (ii) this latter fact is quite transparent anyway.) \square

Theorem 5.4. *Suppose that A is linear operator on a 2-dimensional Hilbert space. Let use take $\mathbf{G}_{\text{CKB(P)}}^{\mathbb{R}}(A) := \mathbf{G}_{\text{CKB(P)}}(A) |_{\{1,3,4\}}$. I. e., in terms of (80), we define*

$$\mathbf{G}_{\text{CKB(P)}}^{\mathbb{R}}(A) = \begin{bmatrix} \frac{W-Z}{4} & \frac{X}{2} & \frac{V}{2} \\ \frac{X}{2} & Y & \frac{Z}{2} \\ \frac{V}{2} & \frac{Z}{2} & 1 \end{bmatrix}.$$

Then

$$(85) \quad \mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A) = -4 \operatorname{adj} \mathbf{G}_{\text{CKB(P)}}^{\mathbb{R}}(A).$$

Here

$$(86) \quad \det \mathbf{G}_{\text{CKB(P)}}^{\mathbb{R}}(A) = \frac{1}{16} (Z^3 - WZ^2 + 4VXZ - 4YZ - 4V^2Y + 4WY - 4X^2) \\ = \frac{1}{4} ((\operatorname{tr} A^* A) - (\operatorname{Re} \det A)) ((U_A)^2 - |D_A|^2) \\ = \frac{1}{2} (U_A - |D_A| + 2E_A) ((U_A)^2 - |D_A|^2).$$

This is strictly positive if A is non-normal, and it is zero if A is normal.

Furthermore,

$$\begin{aligned}
(87) \quad \det \mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A) &= -64 \left(\det \mathbf{G}_{\text{CKB(P)}}^{\mathbb{R}}(A) \right)^2 \\
&= -\frac{1}{4} (Z^3 - WZ^2 + 4VXZ - 4YZ - 4V^2Y + 4WY - 4X^2)^2 \\
&= -4 \left((\text{tr } A^*A) - (\text{Re } \det A) \right)^2 \left((U_A)^2 - |D_A|^2 \right)^2 \\
&= -16 (U_A - |D_A| + 2E_A)^2 \left((U_A)^2 - |D_A|^2 \right)^2.
\end{aligned}$$

This is strictly negative if A is non-normal, and it is zero if A is normal.

Moreover,

$$(88) \quad \text{adj } \mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A) = \left(16 \det \mathbf{G}_{\text{CKB(P)}}^{\mathbb{R}}(A) \right) \cdot \mathbf{G}_{\text{CKB(P)}}^{\mathbb{R}}(A).$$

Proof for Theorem 5.4. Equation (85) is simple to compute, and so is the first line of (86). The rest of (86) is somewhat computational but straightforward: The point is that both sides can be transcribed to polynomials in the five data. Indeed,

$$(89) \quad U_A - |D_A| + 2E_A, (U_A + |D_A|) + (U_A - |D_A|), \text{ and } (U_A + |D_A|)(U_A - |D_A|),$$

are all polynomials in the five data.

Taking the determinant for (85) implies (87/1), which leads to the rest of (87). In the determinants the critical factors are $U_A - |D_A| + 2E_A, U_A + |D_A|, U_A - |D_A|$. These are all greater or equal to $U_A - |D_A| \geq 0$, which vanishes exactly in the normal case. Applying adj to (85), we obtain (88). \square

Alternative proof for Theorem 5.4. For a relatively computation-free argument, let us continue the previous proof of Theorem 5.3 by projections (for the non-normal case). After having method (ii) carried out, method (iii) must give *exactly* the same matrix (cf. Appendix A), thus equality in (84) must hold. By the relation to the inverse, $\mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A)$ and $\text{adj } \mathbf{G}_{\text{CKB(P)}}^{\mathbb{R}}(A)$ are proportional. After checking for position (3, 3), equality (85) follows. Taking determinant, this implies (87/1). Now, (87/1) and the equality in (84) imply (86/2). Then (86/3) is just a minor variant, while (86/1) follows from a simple direct computation. Equations (86) and (87/1) imply the rest of (87). Applying adj to (85), we obtain (88). Non-degeneracy and (87/1) imply $\det \mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A) < 0$. This and equality in (84) imply $\det \mathbf{G}_{\text{CKB(P)}}^{\mathbb{R}}(A) > 0$. In the normal case the equalities extend by continuity; the quadratic forms must be degenerate with determinants 0. \square

Next we will consider other arguments for the proof of Theorem 5.3. As there is a general argument for the location the interior, we will restrict only to arguments identifying the boundary of conformal range. The following arguments will use convexity, but which, in these circumstances, is not more than the qualitative Theorem 2.1.

A proof of Theorem 5.3 via the standard enveloping construction. One can use the enveloping construction as described in [24] in order to show that

$$(90) \quad \begin{bmatrix} x_{\text{CKB(P)}} \\ z_{\text{CKB(P)}} \\ 1 \end{bmatrix}^{\top} \mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A) \begin{bmatrix} x_{\text{CKB(P)}} \\ z_{\text{CKB(P)}} \\ 1 \end{bmatrix} = 0$$

$$= \frac{1}{1+Y+Z} \left(\begin{bmatrix} V+X \\ Y-1 \end{bmatrix} + \frac{\begin{bmatrix} -1 \\ 1 \end{bmatrix} \mathbf{S} \begin{bmatrix} \cos \omega \\ \sin \omega \end{bmatrix}}{\sqrt{\begin{bmatrix} \cos \omega \\ \sin \omega \end{bmatrix} \mathbf{S} \begin{bmatrix} \cos \omega \\ \sin \omega \end{bmatrix}}} \right)$$

(written in column vector form). This traces the quadric with matrix

$$\begin{bmatrix} 1 & \frac{V+X}{1+Y+Z} \\ & 1 \\ & \frac{Y-1}{1+Y+Z} \\ & & 1 \end{bmatrix}^{-1, \top} \begin{bmatrix} \mathbf{S} & \\ & -\frac{1}{(1+Y+Z)^2} \det \mathbf{S} \end{bmatrix} \begin{bmatrix} 1 & \frac{V+X}{1+Y+Z} \\ & 1 \\ & \frac{Y-1}{1+Y+Z} \\ & & 1 \end{bmatrix}^{-1}.$$

Expanded, this yields the matrix $\mathbf{Q}_{\text{CKB}}^{\mathbb{R}}(A)$. \square

Note that the previous two arguments are close variants of each other. Yet, the second one is much more computational than the first one.

A proof of Theorem 5.3 via the algebraic enveloping construction. Consider

$$\begin{aligned} F_A^{\text{CR}}(\lambda, \nu) &\equiv \det(\nu \text{Id}_2 - (A - \lambda \text{Id}_2)^*(A - \lambda \text{Id}_2)) = \\ &= \nu^2 - (2\lambda^2 - 2V\lambda + Z)\nu + (\lambda^4 - 2V\lambda^3 + W\lambda^2 - 2X\lambda + Y). \end{aligned}$$

This is irreducible, as otherwise it would decompose to two linear factors in ν contradicting to non-normality. Then

$$\begin{aligned} F_A^{\text{CR}}(\lambda, \lambda^2 - 2x\lambda + z) &= \\ &= (4x^2 - 4Vx + W - Z)\lambda^2 + (4xz + 2Zx + 2Vz - 2X)\lambda + (z^2 - Zz + Y). \end{aligned}$$

(Here x and z should be understood as $x_{\text{CKB}(P)}$ and $z_{\text{CKB}(P)}$.) Then one directly finds that the discriminant of this polynomial is

$$4 \cdot \begin{bmatrix} x_{\text{CKB}(P)} \\ z_{\text{CKB}(P)} \\ 1 \end{bmatrix}^{\top} \mathbf{Q}_{\text{CKB}(P)}^{\mathbb{R}}(A) \begin{bmatrix} x_{\text{CKB}(P)} \\ z_{\text{CKB}(P)} \\ 1 \end{bmatrix}. \quad \square$$

A proof of Theorem 5.3 in a more Kippenhahnian formalism. We consider the projective homogeneous polynomial for the dual curve:

$$\begin{aligned} K_A^{\text{CR}}(u, s, w) &\equiv \\ &\equiv \det \left(u \frac{A + A^*}{2} + sA^*A + w \text{Id} \right) = \\ &= \frac{1}{2} \left(\text{tr} \left(u \frac{A + A^*}{2} + sA^*A + w \text{Id} \right) \right)^2 - \frac{1}{2} \text{tr} \left(\left(u \frac{A + A^*}{2} + sA^*A + w \text{Id} \right)^2 \right) \\ &= \frac{W - Z}{4} u^2 + Xus + Vuw + Ys^2 + Zsw + w^2 \\ &\equiv \begin{bmatrix} u \\ s \\ w \end{bmatrix}^{\top} \mathbf{G}_{\text{CKB}(P)}^{\mathbb{R}}(A) \begin{bmatrix} u \\ s \\ w \end{bmatrix}. \end{aligned}$$

(Cf. $\mathbf{G}_{\text{CKB}(P)}^{\mathbb{R}}(A) = \mathbf{G}_{\text{CKB}(P)}(A)|_{\{1,3,4\}}$.) As it is known from projective geometry, the matrix of the dual of the conic is given by the adjugate matrix. In this case this is $\text{adj } \mathbf{G}_{\text{CKB}(P)}^{\mathbb{R}}(A) = -\frac{1}{4} \mathbf{Q}_{\text{CKB}(P)}^{\mathbb{R}}(A)$. The scalar factor, of course, can be modified. \square

From [24], we know that the previous two arguments are essentially synonymous. Yet, in practice, they look quite different. The first one looks like the algebraic variant of the differential geometric argument using discriminant instead of differentiation. The second one is the restriction of the Kippenhahnian argument from the case of the shell, or version (iii) of the projection argument; invoking only a moderate amount of projective geometry. (And, by this, we have constructed a connected graph of essential equivalences between the various proofs above.)

Further observations in the line of Theorem 5.4 are as follows. Compatible to the scheme of (82), we set

$$\mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R},0} = \begin{bmatrix} 1 & & \\ & -\frac{1}{2} & \\ & -\frac{1}{2} & \end{bmatrix}, \quad \text{and} \quad \mathbf{Q}_{\text{CKB}}^{\mathbb{R},0} = \begin{bmatrix} 1 & & \\ & 1 & \\ & & -1 \end{bmatrix}.$$

On the other hand, we define

$$(93) \quad \mathbf{G}_{\text{CKB}}^{\mathbb{R}}(A) = \begin{bmatrix} 1 & & \\ & 1 & 1 \\ & -1 & 1 \end{bmatrix}^{-1} \mathbf{G}_{\text{CKB(P)}}^{\mathbb{R}}(A) \begin{bmatrix} 1 & & \\ & 1 & 1 \\ & -1 & 1 \end{bmatrix}^{-1, \top}.$$

We also use the shorthand notation

$$\mathbf{G}_{\text{CKB(P)}}^{\mathbb{R},0} = (\mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R},0})^{-1}, \quad \text{and} \quad \mathbf{G}_{\text{CKB}}^{\mathbb{R},0} = (\mathbf{Q}_{\text{CKB}}^{\mathbb{R},0})^{-1}.$$

Then, it is easy to see that $\mathbf{G}_{\text{CKB(P)}}^{\mathbb{R},0} \mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A) \sim \mathbf{G}_{\text{CKB}}^{\mathbb{R},0} \mathbf{Q}_{\text{CKB}}^{\mathbb{R}}(A)$ and $\mathbf{G}_{\text{CKB}}^{\mathbb{R}}(A) \mathbf{Q}_{\text{CKB}}^{\mathbb{R},0} \sim \mathbf{G}_{\text{CKB(P)}}^{\mathbb{R}}(A) \mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R},0}$. Applying (2) to (93), we obtain as a variant of (85),

$$\mathbf{Q}_{\text{CKB}}^{\mathbb{R}}(A) = -16 \operatorname{adj} \mathbf{G}_{\text{CKB}}^{\mathbb{R}}(A);$$

moreover,

$$(94) \quad \mathbf{G}_{\text{CKB(P)}}^{\mathbb{R},0} \mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A) = 16 \operatorname{adj} \left(\mathbf{G}_{\text{CKB(P)}}^{\mathbb{R}}(A) \mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R},0} \right),$$

and

$$\mathbf{G}_{\text{CKB}}^{\mathbb{R},0} \mathbf{Q}_{\text{CKB}}^{\mathbb{R}}(A) = 16 \operatorname{adj} \left(\mathbf{G}_{\text{CKB}}^{\mathbb{R}}(A) \mathbf{Q}_{\text{CKB}}^{\mathbb{R},0} \right).$$

Theorem 5.5. (a) The eigenvalues of $\mathbf{G}_{\text{CKB(P)}}^{\mathbb{R}}(A) \mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R},0}$ or $\mathbf{G}_{\text{CKB}}^{\mathbb{R}}(A) \mathbf{Q}_{\text{CKB}}^{\mathbb{R},0}$ are

$$-\frac{1}{2}(U_A - |D_A|), \quad -\frac{1}{2}(U_A + |D_A|), \quad -\frac{1}{2}(U_A - |D_A| + 2E_A).$$

(b) The eigenvalues of $\mathbf{G}_{\text{CKB(P)}}^{\mathbb{R},0} \mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A)$ or $\mathbf{G}_{\text{CKB}}^{\mathbb{R},0} \mathbf{Q}_{\text{CKB}}^{\mathbb{R}}(A)$, are

$$\begin{aligned} 4(U_A - |D_A|)(U_A + |D_A|), \\ 4(U_A - |D_A|)(U_A - |D_A| + 2E_A), \\ 4(U_A + |D_A|)(U_A - |D_A| + 2E_A). \end{aligned}$$

Proof. (a) Direct computation, cf. (89). (b) follows from (94). \square

Example 5.6. (a) Regarding $L_{\alpha,t}^{\pm}$:

$$\mathbf{G}_{\text{CKB}}^{\mathbb{R}}(L_{\alpha,t}^{\pm}) \mathbf{Q}_{\text{CKB}}^{\mathbb{R},0} = \begin{bmatrix} -((\cos \alpha)^2 + t^2) & & \\ & -t^2 & \\ & & -(1 + t^2) \end{bmatrix},$$

and

$$\mathbf{G}_{\text{CKB}}^{\mathbb{R},0} \mathbf{Q}_{\text{CKB}}^{\mathbb{R}}(L_{\alpha,t}^{\pm}) = \begin{bmatrix} 16t^2(1+t^2) & & \\ & 16((\cos \alpha)^2 + t^2)(1+t^2) & \\ & & 16t^2((\cos \alpha)^2 + t^2) \end{bmatrix}.$$

(b) Regarding S_{β} :

$$\mathbf{G}_{\text{CKB(P)}}^{\mathbb{R}}(S_{\beta}) \mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R},0} = \begin{bmatrix} -\frac{1}{4}(\cos \beta)^2 & & \\ & -\frac{1}{4} & \\ & -\frac{1}{2} & -\frac{1}{4} \end{bmatrix},$$

and

$$\mathbf{G}_{\text{CKB(P)}}^{\mathbb{R},0} \mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(S_{\beta}) = \begin{bmatrix} 1 & & \\ & (\cos \beta)^2 & \\ & -2(\cos \beta)^2 & (\cos \beta)^2 \end{bmatrix}.$$

(c) In case of $\mathbf{0}_2$:

$$\mathbf{G}_{\text{CKB(P)}}^{\mathbb{R}}(\mathbf{0}_2) \mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R},0} = \begin{bmatrix} 0 & & \\ & 0 & \\ & -\frac{1}{2} & 0 \end{bmatrix},$$

and

$$\mathbf{G}_{\text{CKB(P)}}^{\mathbb{R},0} \mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(\mathbf{0}_2) = \mathbf{0}_3.$$

For all the matrices above, the eigenvalues can be read off from the diagonals. \diamond

5.7. Using the notation (80), we can define

$$U_1(A) = 3Z - W,$$

$$U_2(A) = 3Z^2 - 2WZ + 4VX - 4Y,$$

$$U_3(A) = Z^3 - WZ^2 + 4VXZ - 4YZ - 4V^2Y + 4WY - 4X^2.$$

Here, the expressions are related to

$$\det \left(\lambda \text{Id}_3 - \mathbf{G}_{\text{CKB(P)}}^{\mathbb{R}}(A) \mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R},0} \right) = \lambda^3 + \frac{U_1(A)}{4} \lambda^2 + \frac{U_2(A)}{16} \lambda + \frac{U_3(A)}{64},$$

and

$$\det \left(\lambda \text{Id}_3 - \mathbf{G}_{\text{CKB(P)}}^{\mathbb{R},0} \mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A) \right) = \lambda^3 - U_2(A) \lambda^2 + U_1(A) U_3(A) \lambda - (U_3(A))^2.$$

Then $U_1(A)$, $U_2(A)$, $U_3(A)$ are elementary symmetric polynomials of $2(U_A - |D_A|)$, $2(U_A + |D_A|)$, $2(U_A - |D_A| + 2E_A)$. It is easy to see that

$$(95) \quad U_1(A) = 6(U_A - |D_A|) + 4|D_A| + 4E_A \geq 0,$$

with equality if and only if A is normal and real-parabolic, i. e. it is a real scalar matrix. Similarly,

$$(96) \quad U_2(A) = 12(U_A - |D_A|)^2 + 16(U_A - |D_A|)(|D_A| + E_A) + 16|D_A|E_A \geq 0,$$

with equality if and only if A is normal with two equal or conjugate eigenvalues. Moreover,

$$(97) \quad U_3(A) = 8(U_A - |D_A|)(U_A + |D_A|)(U_A - |D_A| + 2E_A) \geq 0,$$

with equality if and only if A is normal.

5.D. Decompositions of the quadratic forms.

Lemma 5.8. *Consider*

$$\mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A)|_{\{1,2\}} = \begin{bmatrix} Z^2 - 4Y & 2X - VZ \\ 2X - VZ & V^2 - W + Z \end{bmatrix}.$$

Then

$$(98) \quad \det \mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A)|_{\{1,2\}} = 16 \det \mathbf{G}_{\text{CKB(P)}}^{\mathbb{R}}(A) \\ = Z^3 - WZ^2 + 4VXZ - 4YZ - 4V^2Y + 4WY - 4X^2 \\ = 8(U_A - |D_A| + 2E_A)((U_A)^2 - |D_A|^2).$$

Regarding its rank,

$$\text{rk } \mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A)|_{\{1,2\}} = \begin{cases} 2 & \text{if } A \text{ is non-normal,} \\ 1 & \text{if } A \text{ is normal without equal or conjugate eigenvalues,} \\ 0 & \text{if } A \text{ is normal with equal or conjugate eigenvalues.} \end{cases}$$

Proof. Specifying (88) to entry (3, 3), we obtain (98/1), or by direct computation (98/2). The rest of (98) follows from (86). This determinant is non-vanishing if and only if A is non-normal. In the normal case the rank can be 0 or 1. If the canonical form (7) is considered, then

$$Z^2 - 4Y = t^4 + 2t^2(|\lambda_1|^2 + |\lambda_2|^2) + (|\lambda_1|^2 - |\lambda_2|^2)^2, \\ 2X - VZ = -t^2(\text{Re } \lambda_1 + \text{Re } \lambda_2) - (\text{Re } \lambda_1 - \text{Re } \lambda_2)(|\lambda_1|^2 - |\lambda_2|^2), \\ V^2 - W + Z = t^2 + (\text{Re } \lambda_1 - \text{Re } \lambda_2)^2.$$

This shows that rank 0 is achieved if and only if $t = 0$ (normality) and $\text{Re } \lambda_1 - \text{Re } \lambda_2 = 0$ and $|\lambda_1|^2 - |\lambda_2|^2 = 0$ holds. The latter two conditions together mean equal or conjugate eigenvalues. \square

Lemma 5.9. (a)

$$\mathbf{G}_{\text{CKB(P)}}^{\mathbb{R}}(A) = \begin{bmatrix} 1 & & \frac{1}{2}V \\ & 1 & \frac{1}{2}Z \\ & & 1 \end{bmatrix} \begin{bmatrix} -\frac{1}{4} \text{adj} \left(\mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A)|_{\{1,2\}} \right) & & \\ & & 1 \end{bmatrix} \begin{bmatrix} 1 & & \frac{1}{2}V \\ & 1 & \frac{1}{2}Z \\ & & 1 \end{bmatrix}^{\top}.$$

(b)

$$\text{rk } \mathbf{G}_{\text{CKB(P)}}^{\mathbb{R}}(A) = \begin{cases} 3 & \text{if } A \text{ is non-normal,} \\ 2 & \text{if } A \text{ is normal without equal or conjugate eigenvalues,} \\ 1 & \text{if } A \text{ is normal with equal or conjugate eigenvalues.} \end{cases}$$

Lemma 5.10. (a)

$$\mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A) = \\ = \begin{bmatrix} 1 & & \frac{1}{2}V \\ & 1 & \frac{1}{2}Z \\ & & 1 \end{bmatrix}^{-1, \top} \begin{bmatrix} \mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A)|_{\{1,2\}} & & \\ & & -\frac{1}{4} \det \mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A)|_{\{1,2\}} \end{bmatrix} \begin{bmatrix} 1 & & \frac{1}{2}V \\ & 1 & \frac{1}{2}Z \\ & & 1 \end{bmatrix}^{-1}.$$

(b)

$$\operatorname{rk} \mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A) = \begin{cases} 3 & \text{if } A \text{ is non-normal,} \\ 1 & \text{if } A \text{ is normal without equal or conjugate eigenvalues,} \\ 0 & \text{if } A \text{ is normal with equal or conjugate eigenvalues.} \end{cases}$$

Proofs. (a) These are simple computations. (b) These follow from Lemma 5.8. \square

Lemma 5.11.

$$\begin{bmatrix} 2\lambda \\ -1 \\ -\lambda^2 \end{bmatrix}^{\top} \mathbf{G}_{\text{CKB(P)}}^{\mathbb{R}}(A) \begin{bmatrix} 2\lambda \\ -1 \\ -\lambda^2 \end{bmatrix} = \det(\lambda \operatorname{Id}_2 - A) \cdot \det(\lambda \operatorname{Id}_2 - A^*).$$

Proof. Direct computation. (See [24] for greater generality.) \square

Lemma 5.12. *Assume that the 2×2 complex matrix A is non-normal, and \mathbf{M} is the matrix of the quadric of the boundary of $\operatorname{DW}_{\text{CKB(P)}}^{\mathbb{R}}(A)$. Then*

$$(a) \quad \mathbf{G}_{\text{CKB(P)}}^{\mathbb{R}}(A) = \frac{\mathbf{M}^{-1}}{(\mathbf{M}^{-1})_{33}} = \frac{\operatorname{adj} \mathbf{M}}{\det \mathbf{M}_{\{1,2\}}};$$

(b)

$$\mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A) = \frac{-4}{(\det \mathbf{M})(\mathbf{M}^{-1})_{33}^2} \cdot \mathbf{M} = \frac{-4 \det \mathbf{M}}{(\det \mathbf{M}_{\{1,2\}})^2} \cdot \mathbf{M} = \frac{-4 \operatorname{adj} \operatorname{adj} \mathbf{M}}{\det \mathbf{M}_{\{1,2\}}^2}.$$

Proof. (a) follows from $\mathbf{G}_{\text{CKB(P)}}^{\mathbb{R}}(A) \simeq \mathbf{M}^{-1}$ and $\mathbf{G}_{\text{CKB(P)}}^{\mathbb{R}}(A)_{33} = 1$. Then (85) implies (b). \square

Lemma 5.13. *Assume that A is a normal complex 2×2 matrix with eigenvalues λ_1, λ_2 . Thus the endpoints of $\operatorname{DW}_{\text{CKB(P)}}^{\mathbb{R}}(A)$ are $\iota_{\text{CKB(P)}}^{[2]}(\lambda_1) = (\operatorname{Re} \lambda_1, |\lambda_1|^2)$ and $\iota_{\text{CKB(P)}}^{[2]}(\lambda_2) = (\operatorname{Re} \lambda_2, |\lambda_2|^2)$. Then, $\mathbf{G}_{\text{CKB(P)}}^{\mathbb{R}}(A)$ and $\mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A)$ can be expressed in terms of the endpoints as follows:*

$$(a) \quad \mathbf{G}_{\text{CKB(P)}}^{\mathbb{R}}(A) = \frac{1}{2} \left(\begin{bmatrix} \operatorname{Re} \lambda_1 \\ |\lambda_1|^2 \\ 1 \end{bmatrix} \begin{bmatrix} \operatorname{Re} \lambda_2 \\ |\lambda_2|^2 \\ 1 \end{bmatrix}^{\top} + \begin{bmatrix} \operatorname{Re} \lambda_2 \\ |\lambda_2|^2 \\ 1 \end{bmatrix} \begin{bmatrix} \operatorname{Re} \lambda_1 \\ |\lambda_1|^2 \\ 1 \end{bmatrix}^{\top} \right).$$

(b)

$$\mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A) = \mathbf{v} \mathbf{v}^{\top}$$

where

$$\mathbf{v} = \begin{bmatrix} |\lambda_1|^2 - |\lambda_2|^2 \\ \operatorname{Re} \lambda_2 - \operatorname{Re} \lambda_1 \\ (\operatorname{Re} \lambda_1)|\lambda_2|^2 - (\operatorname{Re} \lambda_2)|\lambda_1|^2 \end{bmatrix} = \begin{bmatrix} \operatorname{Re} \lambda_1 \\ |\lambda_1|^2 \\ 1 \end{bmatrix} \times \begin{bmatrix} \operatorname{Re} \lambda_2 \\ |\lambda_2|^2 \\ 1 \end{bmatrix}$$

(with \times meaning the ordinary vector product of Gibbs).

Proof. We can compute the data from $A = \begin{bmatrix} \lambda_1 & \\ & \lambda_2 \end{bmatrix}$. \square

Alternative proof for Theorem 5.2. $\mathbf{G}_{\text{CKB(P)}}^{\mathbb{R}}(A)$ can be obtained by (99) / (100); from which V, W, X, Y, Z can be recovered. \square

As we know, $\mathbf{G}_{\text{CKB(P)}}^{\mathbb{R}}(A)$ has no data loss relative to the reduced five data of A , not even in the normal case. Regarding $\mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A)$:

Lemma 5.14. *Assume that A is complex 2×2 matrix.*

(a) *If A is normal with two distinct, non-conjugate eigenvalues, then $\mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A)$ is a matrix of rank strictly 1. Geometrically, it corresponds to the (double) line containing the segment which is the conformal range. Beyond that, $\mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A)$, as it is, also contains the data ‘squared reduced eigendistance’ $4\sqrt{|D_A|}\sqrt{E_A}$ but essentially not more.*

(b) *If A is normal with two equal or conjugate eigenvalues, then $\mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A) = 0$.*

Proof. Let us consider the description in Lemma 5.13(b) with $\mathbf{v} \equiv \mathbf{v}_{\text{CKB(P)}}(\lambda_1, \lambda_2)$.

(b) If λ_1 and λ_2 are equal or conjugates to other, then $(\text{Re } \lambda_1, |\lambda_1|^2) = (\text{Re } \lambda_2, |\lambda_2|^2)$, and $\mathbf{v}_{\text{CKB(P)}}(\lambda_1, \lambda_2)$ vanishes. This leads to the statement.

(a) If λ_1 and λ_2 are neither equal nor conjugates to other, then $(\text{Re } \lambda_1, |\lambda_1|^2) \neq (\text{Re } \lambda_2, |\lambda_2|^2)$, and $\mathbf{v}_{\text{CKB(P)}}(\lambda_1, \lambda_2) \neq 0$. This leads to a double line for $\mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A)$. Then, by the construction of $\mathbf{v}_{\text{CKB(P)}}(\lambda_1, \lambda_2)$, it is easy to see that the line fits to the distinct points $\iota_{\text{CKB(P)}}^{[2]}(\lambda_1) = (\text{Re } \lambda_1, |\lambda_1|^2)$ and $\iota_{\text{CKB(P)}}^{[2]}(\lambda_2) = (\text{Re } \lambda_2, |\lambda_2|^2)$. From $\mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A)$ one can reconstruct $\pm \mathbf{v}_{\text{CKB(P)}}(\lambda_1, \lambda_2)$. If $\mathbf{v}_{\text{CKB(P)}}(\lambda_1, \lambda_2) = \pm(u_1, u_2, u_3)$, then $4\sqrt{|D_A|}\sqrt{E_A} = |\lambda_1 - \lambda_2| \cdot |\lambda_1 - \bar{\lambda}_2| = \sqrt{(u_1)^2 - 4u_2u_3}$, and this information is specific with respect to the scaling of $\pm \mathbf{v}_{\text{CKB(P)}}(\lambda_1, \lambda_2)$. (In particular, in the normal setting, we have the formula $4\sqrt{|D_A|}\sqrt{E_A} = \sqrt{\text{tr}((\mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R},0})^{-1}\mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A))}$.) \square

Remark 5.15. Assume that A is normal with two equal or conjugate eigenvalues; say, the eigenvalues are λ or $\bar{\lambda}$. Then it is reasonable to define

$$\check{\mathbf{Q}}_{\text{CKB(P)}}^{\mathbb{R}}(A) = \begin{bmatrix} 4|\lambda|^2 & -2\text{Re } \lambda & -2|\lambda|^2 \text{Re } \lambda \\ -2\text{Re } \lambda & 1 & 2(\text{Re } \lambda)^2 - |\lambda|^2 \\ -2|\lambda|^2 \text{Re } \lambda & 2(\text{Re } \lambda)^2 - |\lambda|^2 & |\lambda|^4 \end{bmatrix}.$$

This is natural in the sense that in this case,

$$\text{adj } \check{\mathbf{Q}}_{\text{CKB(P)}}^{\mathbb{R}}(A) = 4|\text{Im } \lambda|^2 \cdot \mathbf{G}_{\text{CKB(P)}}^{\mathbb{R}}(A). \quad \triangle$$

5.E. The transformation properties of the matrix.

Lemma 5.16. *Suppose that f is a Möbius transformation $f : \lambda \mapsto \frac{a\lambda+b}{c\lambda+d}$, $a, b, c, d \in \mathbb{R}$, $ad - bc \neq 0$, and A is 2×2 complex matrix such that $-\frac{d}{c}$ is not an eigenvalue of A .*

(a) *Then, the transformation rule*

$$E_{f(A)} \cdot \mathcal{C}(f, A) = E_A$$

holds.

(b) *For $j \in \{1, 2, 3\}$, the transformation rules*

$$U_j(f(A)) \cdot \mathcal{C}(f, A)^j = U_j(A)$$

hold.

Proof. (a) By Lemma 2.21 we have the same transformation rule for U_A and $|D_A|$ (even more generally). As $E_A/|D_A|$ is generically an invariant for real Möbius transformations, E_A must have the same rule for real Möbius transformations.

(b) We deal with j -homogeneous polynomials of U_A , $|D_A|$, E_A ; the transformation rules are corresponding to this fact. \square

Lemma 5.17. *Suppose that f is a Möbius transformation $f : \lambda \mapsto \frac{a\lambda+b}{c\lambda+d}$, $a, b, c, d \in \mathbb{R}$, $ad - bc \neq 0$, and A is 2×2 complex matrix. Then*

$$\mathcal{C}(f, A) = \frac{1}{(ad - bc)^2} \begin{bmatrix} 2cd \\ c^2 \\ d^2 \end{bmatrix}^\top \mathbf{G}_{\text{CKB}(P)}^{\mathbb{R}}(A) \begin{bmatrix} 2cd \\ c^2 \\ d^2 \end{bmatrix}.$$

In particular, for a real f , $\mathcal{C}(f, A)$ depends only on the reduced five data of A .

Proof. This is just the transcription of (40). Note that $\mathbf{G}_{\text{CKB}(P)}^{\mathbb{R}}(A)$ depends only on the reduced five data. \square

Lemma 5.18. *Suppose that f is a Möbius transformation $f : \lambda \mapsto \frac{a\lambda+b}{c\lambda+d}$, $a, b, c, d \in \mathbb{R}$, $ad - bc \neq 0$, and A is 2×2 complex matrix such that $-\frac{d}{c}$ is not an eigenvalue of A .*

Let $R_{\text{CKB}(P)}^{[2]}(f)$ be a matrix of determinant -1 or 1 representing the projective action of f in $\text{CKB}(P)$. Then the following transformation rules holds:

$$(101) \quad \mathbf{Q}_{\text{CKB}(P)}^{\mathbb{R}}(f(A)) \cdot \mathcal{C}(f, A)^2 = \left(R_{\text{CKB}(P)}^{[2]}(f)\right)^{-1, \top} \mathbf{Q}_{\text{CKB}(P)}^{\mathbb{R}}(A) \left(R_{\text{CKB}(P)}^{[2]}(f)\right)^{-1},$$

$$(102) \quad \mathbf{G}_{\text{CKB}(P)}^{\mathbb{R}, 0} \mathbf{Q}_{\text{CKB}(P)}^{\mathbb{R}}(f(A)) \cdot \mathcal{C}(f, A)^2 = \\ = \left(R_{\text{CKB}(P)}^{[2]}(f)\right) \mathbf{G}_{\text{CKB}(P)}^{\mathbb{R}, 0} \mathbf{Q}_{\text{CKB}(P)}^{\mathbb{R}}(A) \left(R_{\text{CKB}(P)}^{[2]}(f)\right)^{-1};$$

$$(103) \quad \mathbf{G}_{\text{CKB}(P)}^{\mathbb{R}}(f(A)) \cdot \mathcal{C}(f, A) = \left(R_{\text{CKB}(P)}^{[2]}(f)\right) \mathbf{G}_{\text{CKB}(P)}^{\mathbb{R}}(A) \left(R_{\text{CKB}(P)}^{[2]}(f)\right)^\top;$$

$$(104) \quad \mathbf{G}_{\text{CKB}(P)}^{\mathbb{R}}(f(A)) \mathbf{Q}_{\text{CKB}(P)}^{\mathbb{R}, 0} \cdot \mathcal{C}(f, A) = \\ = \left(R_{\text{CKB}(P)}^{[2]}(f)\right) \mathbf{G}_{\text{CKB}(P)}^{\mathbb{R}}(A) \mathbf{Q}_{\text{CKB}(P)}^{\mathbb{R}, 0} \left(R_{\text{CKB}(P)}^{[2]}(f)\right)^{-1}.$$

Proof. By Theorem 5.3, we already know that $\mathbf{Q}_{\text{CKB}(P)}^{\mathbb{R}}(A)$ transforms naturally. The

scaling factor in (101) can generically be written as $\sqrt[3]{\frac{\det \mathbf{Q}_{\text{CKB}(P)}^{\mathbb{R}}(A)}{\det \mathbf{Q}_{\text{CKB}(P)}^{\mathbb{R}}(f(A))}}$, but, due to

(87) and the scaling properties of U_A , $|D_A|$, E_A , this reduces generically to $\mathcal{C}(f, A)^2$. Then, by continuity, the equality (101) extends. Equivalence to (102) follows from the invariance property $\mathbf{Q}_{\text{CKB}(P)}^{\mathbb{R}, 0} = \left(R_{\text{CKB}(P)}^{[2]}(f)\right)^{-1, \top} \mathbf{Q}_{\text{CKB}(P)}^{\mathbb{R}, 0} \left(R_{\text{CKB}(P)}^{[2]}(f)\right)^{-1}$. Similar argument applies to the $\mathbf{G}_{\text{CKB}(P)}^{\mathbb{R}}(A)$. \square

Alternative proof. (103) and (104) follow from (43) and (44) by restriction. Then the transformation properties (101) and (102) follow from (85). \square

Alternative proof to Theorem 5.5. Comparing Example 5.6 and Example 4.4 (Example 5.26), the statement checks out for the canonical representatives of Lemma 4.3. By Lemma 5.18, real Möbius transformations, do not change the ratio of the eigenvalues; moreover the overall scaling factors are also correct. \square

In the manner of Corollary 2.22, we can prepare various naturally well-transforming versions of $\mathbf{G}_{\text{CKB}(P)}^{\mathbb{R}}(A)$ and $\mathbf{Q}_{\text{CKB}(P)}^{\mathbb{R}}(A)$. As we will see, it is useful to be opportunistic about the scaling factor. We only have to be careful that it should be (-1) -homogeneous

in $U_A, |D_A|, E_A$ for $\mathbf{G}_{\text{CKB}(P)}^{\mathbb{R}}(A)$, and it should be (-2) -homogeneous in $U_A, |D_A|, E_A$ for $\mathbf{Q}_{\text{CKB}(P)}^{\mathbb{R}}(A)$. In particular,

Corollary 5.19. (a) In the $U_1(A) \neq 0$ case, the matrix is given by

$$(105) \quad \tilde{\mathbf{G}}_{\text{CKB}(P)}^{\mathbb{R}}(A) = \frac{1}{U_1(A)} \mathbf{G}_{\text{CKB}(P)}^{\mathbb{R}}(A)$$

has the transformation property

$$\tilde{\mathbf{G}}_{\text{CKB}(P)}^{\mathbb{R}}(f(A)) = \left(R_{\text{CKB}(P)}^{[2]}(f) \right) \tilde{\mathbf{G}}_{\text{CKB}(P)}^{\mathbb{R}}(A) \left(R_{\text{CKB}(P)}^{[2]}(f) \right)^{\top},$$

whenever $f(A)$ makes sense.

(b) In the $U_2(A) \neq 0$ case, the matrix is given by

$$(106) \quad \hat{\mathbf{Q}}_{\text{CKB}(P)}^{\mathbb{R}}(A) = \frac{1}{U_2(A)} \mathbf{Q}_{\text{CKB}(P)}^{\mathbb{R}}(A)$$

has the transformation property

$$\hat{\mathbf{Q}}_{\text{CKB}(P)}^{\mathbb{R}}(f(A)) = \left(R_{\text{CKB}(P)}^{[2]}(f) \right)^{-1, \top} \hat{\mathbf{Q}}_{\text{CKB}(P)}^{\mathbb{R}}(A) \left(R_{\text{CKB}(P)}^{[2]}(f) \right)^{-1},$$

whenever $f(A)$ makes sense. □

Remark 5.20. (a) If A is normal with two distinct, non-conjugate eigenvalues, then $\hat{\mathbf{Q}}_{\text{CKB}(P)}^{\mathbb{R}}(A)$ contains no more information than the line of the conformal range.

(b) Assume that A is a normal matrix with eigenvalues λ or $\bar{\lambda}$. Then (106) allows a natural but not continuous extension by

$$\hat{\mathbf{Q}}_{\text{CKB}(P)}^{\mathbb{R}}(A) = \frac{1}{8(\text{Im } \lambda)^2} \check{\mathbf{Q}}_{\text{CKB}(P)}^{\mathbb{R}}(A).$$

This leaves only real scalar matrices without a particular choice for $\hat{\mathbf{Q}}_{\text{CKB}(P)}^{\mathbb{R}}(A)$. In that case, only $\mathbb{R}^+ \hat{\mathbf{Q}}_{\text{CKB}(P)}^{\mathbb{R}}(A)$ is a sufficiently invariant object. △

5.F. Canonical representatives and the geometry of the conformal range.

Although Theorem 5.3 is sufficiently explicit, it is useful to visualize certain particular cases. We use the canonical representatives from Lemma 4.3.

Example 5.21. (a)

$$\mathbf{Q}_{\text{CKB}}^{\mathbb{R}}(L_{\alpha, t}^{\pm}) = \begin{bmatrix} 16t^2(1+t^2) & & \\ & 16((\cos \alpha)^2 + t^2)(1+t^2) & \\ & & -16t^2((\cos \alpha)^2 + t^2) \end{bmatrix}.$$

(b)

$$\mathbf{Q}_{\text{CKB}}^{\mathbb{R}}(S_{\beta}) = \begin{bmatrix} 1 & & \\ & 2(\cos \beta)^2 & (\cos \beta)^2 \\ & (\cos \beta)^2 & \end{bmatrix}.$$

(c)

$$\mathbf{Q}_{\text{CKB}}^{\mathbb{R}}(\mathbf{0}_2) = \mathbf{0}_3. \quad \diamond$$

Theorem 5.22. *The conformal ranges of the canonical representatives in the CKB model are as follows:*

The zero matrix $\mathbf{0}_2$ yields the point ellipse

$$\{(0, -1)\};$$

S_β yields the ellipse with axes

$$\left[-\frac{\sqrt{2}}{2} \cos \beta, \frac{\sqrt{2}}{2} \cos \beta \right] \times \left\{ -\frac{1}{2} \right\} \quad \text{and} \quad \{0\} \times [-1, 0];$$

and $L_{\alpha,t}^\pm$ yields the ellipse with axes

$$\left[-\frac{\sqrt{(\cos \alpha)^2 + t^2}}{\sqrt{1+t^2}}, \frac{\sqrt{(\cos \alpha)^2 + t^2}}{\sqrt{1+t^2}} \right] \times \{0\} \quad \text{and} \quad \{0\} \times \left[-\frac{t}{\sqrt{1+t^2}}, \frac{t}{\sqrt{1+t^2}} \right];$$

(the ellipses may be degenerate) .

Proof. In the non-normal cases ($t > 0, \beta \in [0, \pi/2)$) the result follows from Theorem 5.3 via Example 5.21. The normal cases follow from the continuity of the conformal range as a set (but not from the continuity of $\mathbf{Q}_{\text{CKB}}^\mathbb{R}(A)$). \square

Remark 5.23. One can see that the conformal range in the CKB model yields all possible degenerate ellipses in the unit disk which do not contain the “infinity” point $(0, 1)$. Hence, they can be identified as possibly degenerate h -ellipses (in the asymptotically closed plane) but which avoid the distinguished asymptotical point $(0, 1)_{\text{CKB}}$.

Although there are recipes for reconstruction (see Discussion 5.45), the simplest way to see this, in the non-normal case, is to consider the pencil generated the quadric of the candidate conformal range and the quadric of the asymptotic points; then apply the canonical form of the pencil. (Cf. Berger [2] for an introduction to the pencils, and most of the literature on h -conics for their application.) \triangle

Proof of Theorem 5.22 via symmetry principles. One can compute the conformal range in the CKB model for $L_{\alpha,t}^\pm$ of Lemma 4.3 with $t \geq 0, \alpha \in (0, \pi/2]$ as follows: As we deal with projections of non-degenerate h -tubes, we already know that the results will be (h -)ellipses in the CKB model. As $A = L_{\alpha,t}^\pm$ is unitarily conjugate not only to $(A^{-1})^*$ but to $-A$ or $-A^*$, we see that the ellipses will be symmetric to the x_{CKB} and z_{CKB} axes, thus they will be aligned along those axes. Now, it is easy to obtain the norms

$$\|L_{\alpha,t}^\pm\|_2 = \left\| \begin{bmatrix} \cos \alpha + i \sin \alpha & 2t \\ -\cos \alpha \pm i \sin \alpha & \end{bmatrix} \right\|_2 = t + \sqrt{1+t^2},$$

and the norms of the Cayley transforms,

$$\left\| \frac{\text{Id}_2 - L_{\alpha,t}^\pm}{\text{Id}_2 + L_{\alpha,t}^\pm} \right\|_2 = \left\| \begin{bmatrix} -i \tan \frac{\alpha}{2} & \pm 2it e^{\pm \frac{1-i}{2} i \alpha} \csc \alpha \\ \mp i \cot \frac{\alpha}{2} & \end{bmatrix} \right\|_2 = \frac{\sqrt{(\cos \alpha)^2 + t^2} + \sqrt{1+t^2}}{\sin \alpha}.$$

(Computing only case $+$ is sufficient.) Then, applying the Lemma of “Extremal values in ranges” from [24] (there ‘ y_{CKB} ’ was used instead of ‘ z_{CKB} ’), we find that

$$\sup z_{\text{CKB}}(\text{DW}_{\text{CKB}}^\mathbb{R}(L_{\alpha,t}^\pm)) = \frac{t}{\sqrt{1+t^2}},$$

and

$$\inf x_{\text{CKB}}(\text{DW}_{\text{CKB}}^\mathbb{R}(L_{\alpha,t}^\pm)) = -\frac{\sqrt{(\cos \alpha)^2 + t^2}}{\sqrt{1+t^2}}.$$

Taking the axial symmetries in account, this proves that $DW_{\text{CKB}}^{\mathbb{R}}(L_{\alpha,t}^{\pm})$ is as indicated. The case $\alpha = 0$ follows by continuity.

$DW_{\text{CKB}}^{\mathbb{R}}(S_{\beta})$ can be computed as similarly. We have symmetry for the z_{CKB} axis. Furthermore,

$$\|S_{\beta}\|_2 = 1, \quad \|S_{\beta}\|_2^{-} = 0,$$

and

$$\left\| \frac{\text{Id}_2 - S_{\beta}}{\text{Id}_2 + S_{\beta}} \right\|_2 = \left\| \begin{bmatrix} 1 & \frac{-2 \cos \beta}{1+i \sin \beta} \\ \frac{1-i \sin \beta}{1+i \sin \beta} & 1 \end{bmatrix} \right\|_2 = \sqrt{\frac{\sqrt{2} + \cos \beta}{\sqrt{2} - \cos \beta}}.$$

Applying the Lemma of ‘‘Extremal values in ranges’’ from [24], we find that

$$\sup z_{\text{CKB}}(DW_{\text{CKB}}^{\mathbb{R}}(S_{\beta})) = 0,$$

$$\inf z_{\text{CKB}}(DW_{\text{CKB}}^{\mathbb{R}}(S_{\beta})) = -1,$$

and

$$\inf x_{\text{CKB}}(DW_{\text{CKB}}^{\mathbb{R}}(S_{\beta})) = -\frac{\sqrt{2}}{2} \cos \beta.$$

Taking the symmetry into account, this proves that $DW_{\text{CKB}}^{\mathbb{R}}(S_{\beta})$ is as indicated.

The case of $\mathbf{0}_2$ is trivial. \square

It might also be interesting to see how a raw, ‘‘unconceptualized’’ computation works using only the definition of the conformal range:

Proof of Theorem 5.22 via raw computation. Assume that $\mathbf{x} = z_1 \mathbf{e}_1 + z_2 \mathbf{e}_2$ with $|z_1|^2 + |z_2|^2 = 1$. By explicit computation, one find that

$$\begin{bmatrix} \text{Re} \langle L_{\alpha,t}^{\pm} \mathbf{x}, \mathbf{x} \rangle \\ 0 \\ \langle L_{\alpha,t}^{\pm} \mathbf{x}, L_{\alpha,t}^{\pm} \mathbf{x} \rangle \\ \langle \mathbf{x}, \mathbf{x} \rangle \end{bmatrix} = \underbrace{\begin{bmatrix} t & 0 & \cos \alpha & 0 \\ 0 & 0 & 0 & 0 \\ 2t \cos \alpha & -2t \sin \alpha & -2t^2 & 2t^2 + 1 \\ 0 & 0 & 0 & 1 \end{bmatrix}}_{S_T :=} \begin{bmatrix} 2 \text{Re}(z_1 \bar{z}_2) \\ 2 \text{Im}(z_1 \bar{z}_2) \\ |z_1|^2 - |z_2|^2 \\ |z_1|^2 + |z_2|^2 \end{bmatrix}.$$

Now S_T itself can be written as the product

$$S_{T1} \cdot S_{T2} \cdot S_{T3} \equiv \begin{bmatrix} \sqrt{(\cos \alpha)^2 + t^2} & & & \\ & 0 & & \\ & & 2t\sqrt{1+t^2} & 2t^2+1 \\ & & & 1 \end{bmatrix} \cdot \begin{bmatrix} 1 & & & \\ -\frac{\sqrt{(\cos \alpha)^2 + t^2}}{\sqrt{1+t^2}} & \frac{\sin \alpha}{\sqrt{1+t^2}} & & \\ -\frac{\sin \alpha}{\sqrt{1+t^2}} & -\frac{\sqrt{(\cos \alpha)^2 + t^2}}{\sqrt{1+t^2}} & & \\ & & & 1 \end{bmatrix} \cdot \begin{bmatrix} & & & \\ \frac{t}{\sqrt{(\cos \alpha)^2 + t^2}} & \frac{\cos \alpha}{\sqrt{(\cos \alpha)^2 + t^2}} & & \\ & 1 & & \\ -\frac{\cos \alpha}{\sqrt{(\cos \alpha)^2 + t^2}} & \frac{t}{\sqrt{(\cos \alpha)^2 + t^2}} & & \\ & & & 1 \end{bmatrix}.$$

(if $\cos \alpha = t = 0$, then the identity matrix can be taken as S_{T_3}). Now it is easy to see how S_T acts on the affinized unit sphere: S_{T_3} and S_{T_2} are just orthogonal transformations leaving it invariant, while S_{T_1} collapses it to a (possibly degenerate) elliptical disk (and translates it). If $t > 0$, then its equation is

$$\frac{(x_{\text{CKB(P)}})^2}{(\sqrt{(\cos \alpha)^2 + t^2})^2} + \frac{(z_{\text{CKB(P)}} - (2t^2 + 1))^2}{(2t\sqrt{1 + t^2})^2} \leq 1,$$

but the situation is also transparent in the $t = 0$ case. Transcription to the CKB model is straightforward.

Similarly, one finds

$$\begin{bmatrix} \text{Re}\langle S_\beta \mathbf{x}, \mathbf{x} \rangle \\ 0 \\ \langle S_\beta \mathbf{x}, S_\beta \mathbf{x} \rangle \\ \langle \mathbf{x}, \mathbf{x} \rangle \end{bmatrix} = \underbrace{\begin{bmatrix} \frac{1}{2} \cos \beta & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -\frac{1}{2} & \frac{1}{2} \\ 0 & 0 & 0 & 1 \end{bmatrix}}_{S_B :=} \begin{bmatrix} 2 \text{Re}(z_1 \bar{z}_2) \\ 2 \text{Im}(z_1 \bar{z}_2) \\ |z_1|^2 - |z_2|^2 \\ |z_1|^2 + |z_2|^2 \end{bmatrix}.$$

Here S_B can be written as

$$S_{B1} \cdot S_{B2} \equiv \begin{bmatrix} \frac{1}{2} \cos \beta & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{1}{2} & \frac{1}{2} \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & & & \\ & -1 & & \\ & & -1 & \\ & & & 1 \end{bmatrix}.$$

If $\cos \beta > 0$, then this yields the equation

$$\frac{(x_{\text{CKB(P)}})^2}{(\frac{1}{2} \cos \beta)^2} + \frac{(z_{\text{CKB(P)}} - \frac{1}{2})^2}{(\frac{1}{2})^2} \leq 1,$$

but the $\cos \beta = 0$ case is also transparent. Transcription to the CKB model is also straightforward.

The case of \mathbf{O}_2 is trivial.

(Remark: Finding $S_{T_1}, S_{T_2}, S_{T_3}$ may seem complicated, but it is quite simple due to the symmetries of the conformal range: Firstly, from the last column of S_T and the row lengths of 3×3 block S_T , one can reconstruct S_{T_1} . This also gives us $S_{T_2}S_{T_3}$ but its second row. However, a second row can be reconstructed from the first and third rows by taking ordinary vector product. $S_{T_2}S_{T_3}$ could have been left as it is, but an obvious decomposition was taken in order to make orthogonality even more transparent.) \square

5.24. Previously, ‘‘possibly degenerate h -ellipses’’ were defined / considered in terms of their data in the projective models. If the reader is sufficiently familiar with hyperbolic conics, then there is no problem in interpreting them in terms of synthetic hyperbolic constructions. It is legitimate to assume this knowledge on a certain level, as several sources are available. Still, as our stance is somewhat expository, we will prove the relevant basic facts ourselves. However, the knowledge of the h -cycles is assumed.

The conformal ranges of the canonical representatives are particularly suitable to clarify geometric descriptions. Firstly, we establish some terminology regarding major and minor axes. Unfortunately these terms, might refer to possibly infinite segments (primarily), but also to lines (of these segments), or to lengths (of these segments). Hopefully, this will not cause too much confusion.

Consider $DW_*(L_{\alpha,t}^\pm)$:

- If $\alpha = 0$, then the range is a distance band. The major axis [segment] can be defined as the single h -line contained in $DW_*(L_{\alpha,t}^\pm)$. It allows several minor axes [segments] which are intersections of the range with any line perpendicular to the major axis. Note, however, that minor axes [lines] are defined naturally, even if the minor axes [segments] degenerate to points.
- If $0 < \alpha < \pi/2$, then the range is bounded but not circular. It is easy to see that it can be included into a h -disk of minimal radius. Then one of the diameters of the disk is contained in the range. This diameter is the major axis [segment]. The intersection of the range and the perpendicular bisector of the major axis [segment] is the minor axis [segment]. Note that the line of minor axis well-defined, even if the minor axis [segment] reduces to a point.
- If $\alpha = \pi/2$, then the range is bounded and circular. As this is a familiar case, we simply note there are several possible major and minor axes [lines] perpendicular to each other.

Consider $DW_*(S_\beta)$:

- If $\beta = 0$, then we have a horodisk. Major axes [segments] are intersections of the horodisk with its axes. Minor axes [segments or lines] are not defined at all, but the “length of the minor axis” is set to be $+\infty$.
- If $0 < \beta \leq \pi/2$, then the range is unbounded, part of a distance band, but only with one asymptotical point. The major axis [segment] is its intersection with its single symmetry axis. Minor axes [segments or lines] are not defined at all, but the “length of the minor axis” is set as follows. We consider the intersections of the range with the lines perpendicular to the major axis. Intersections will yield segments whose lengths, in the present case, limit as

$$\begin{aligned} \lim_{z \searrow -1} d_{\text{CKB}} \left(\left(\sqrt{-2z(1+z)} \cos \beta, z \right), \left(-\sqrt{-2z(1+z)} \cos \beta, z \right) \right) &= \\ &= \lim_{z \searrow -1} \operatorname{arcosh} \frac{1-z-2z(\cos \beta)^2}{1-z+2z(\cos \beta)^2} = \operatorname{arcosh} \frac{1+(\cos \beta)^2}{1-(\cos \beta)^2} = 2 \operatorname{arcosh} \frac{1}{\sin \beta}. \end{aligned}$$

This limit, or even better, supremum, can be defined as the length of the minor axis. (If $\beta = \pi/2$, then the range is a half-line, and the “length of the minor axis” is 0.)

In this case one can define the vertex of the range as the non-asymptotical endpoint of the major axis [segment].

Consider $DW_*(\mathbf{0}_2)$:

- The major axis [segment] is the range itself. The major axis [line] can be considered as any line asymptotical to the range. Minor axes [segments or lines] are not defined at all. The lengths of the major and minor axes can be considered as 0.

Although we considered the canonical examples, the definitions above were set up in terms of hyperbolic geometry. Consequently, they make sense generally (i. e. up to h -congruences), without any reference to any theory of h -conics. In particular, in all cases we have defined the lengths of the major and minor axes.

Example 5.25. In the case of $L_{\alpha,t}^\pm$, we may say that its characteristic CKB values are

$$\frac{\sqrt{(\cos \alpha)^2 + t^2}}{\sqrt{1+t^2}} \quad \text{and} \quad \frac{t}{\sqrt{1+t^2}}.$$

They correspond to the major and minor semi-axes of the (possibly degenerate) h -ellipse. In terms of hyperbolic geometry, these values are

$$\operatorname{arcosh} \frac{\sqrt{1+t^2}}{\sin \alpha} \quad \text{and} \quad \operatorname{arcosh} \sqrt{1+t^2},$$

respectively. (The major semi-axis is $+\infty$ for $\alpha = \pi/2$.) If the minor semi-axis is 0, then it may yield an h -point, an h -segment, or an h -line. If the minor axis is non-zero, then it yields a h -distance band or an ordinary h -ellipse.

In the case of S_β , we may say that its characteristic CKB values are

$$1 \quad \text{and} \quad \cos \beta.$$

They correspond to the major and minor semi-axes

$$+\infty \quad \text{and} \quad \operatorname{arcosh} \frac{1}{\sin \beta}.$$

(The minor semi-axis is $+\infty$ for $\beta = \pi/2$.) The minor semi-axis can be 0, non-zero finite or $+\infty$. Then it yields an h -half-line, an h -elliptic parabola, or an h -horosphere, respectively.

The case $\mathbf{0}_2$ corresponds to an asymptotical point. The wisest choice is to assign the characteristic values and semi-axes

$$0 \quad \text{and} \quad 0. \quad \diamond$$

As the representatives in Theorem 5.22 cover all cases up to real Möbius transformations, i. e. h -isometries in the conformal range, the terminology ‘characteristic CKB values’ can also be applied to arbitrary 2×2 complex matrices.

Example 5.26. Checking the canonical representatives of Lemma 4.3, we find:

(a) For $A = L_{\alpha,t}^\pm$,

$$U_A - |D_A| = 2t^2, \quad \{U_A + |D_A|, U_A - |D_A| + 2E_A\} = \{2((\cos \alpha)^2 + t^2), 2(1 + t^2)\}.$$

(b) For $A = S_\beta$,

$$U_A - |D_A| = \frac{1}{2}(\cos \beta)^2, \quad \{U_A + |D_A|, U_A - |D_A| + 2E_A\} = \left\{\frac{1}{2}, \frac{1}{2}\right\}.$$

(c) For $A = \mathbf{0}_2$,

$$U_A - |D_A| = 0, \quad \{U_A + |D_A|, U_A - |D_A| + 2E_A\} = \{0, 0\}. \quad \diamond$$

Discussion 5.27. The triple ratio

$$(107) \quad U_A - |D_A| : U_A + |D_A| : U_A - |D_A| + 2E_A$$

is just as appropriate in Corollary 4.5 as (75).

As the previous example shows, making indistinction in the order of the last two entries of (107), with some abuse of notation,

$$U_A - |D_A| : \{U_A + |D_A| : U_A - |D_A| + 2E_A\}$$

captures the commonality of the classes of $L_{\alpha,t}^+$ and $L_{\alpha,t}^-$. Actually, as $U_A - |D_A|$ is distinguished as the entry of the smallest absolute value, the same can be said about the unordered ratio

$$\{U_A - |D_A| : U_A + |D_A| : U_A - |D_A| + 2E_A\}. \quad \square$$

Theorem 5.28. *Assume that A is a 2×2 complex matrix.*

(a) *If $E_A \geq |D_A|$ (e. g. quasihyperbolic case), then the characteristic CKB values are*

$$\sqrt{\frac{U_A + |D_A|}{U_A - |D_A| + 2E_A}} \geq \sqrt{\frac{U_A - |D_A|}{U_A - |D_A| + 2E_A}}.$$

The corresponding h -distances (the semi-axes) are

$$\frac{1}{2} \operatorname{arcosh} \frac{U_A + E_A}{E_A - |D_A|} \geq \frac{1}{2} \operatorname{arcosh} \frac{U_A - |D_A| + E_A}{E_A}.$$

(b) *If $|D_A| \geq E_A$ (e. g. quasielliptic case), then the characteristic CKB values are*

$$\sqrt{\frac{U_A - |D_A| + 2E_A}{U_A + |D_A|}} \geq \sqrt{\frac{U_A - |D_A|}{U_A + |D_A|}}.$$

The corresponding h -distances (the semi-axes) are

$$\frac{1}{2} \operatorname{arcosh} \frac{U_A + E_A}{|D_A| - E_A} \geq \frac{1}{2} \operatorname{arcosh} \frac{U_A}{|D_A|}.$$

(c) *In the case $E_A = |D_A|$, the characteristic CKB values are*

$$1 \geq \sqrt{\frac{U_A - |D_A|}{U_A + |D_A|}} \equiv \sqrt{\frac{U_A - E_A}{U_A + E_A}}.$$

The corresponding h -distances (the semi-axes) are

$$+\infty \geq \frac{1}{2} \operatorname{arcosh} \frac{U_A}{|D_A|} \equiv \frac{1}{2} \operatorname{arcosh} \frac{U_A}{E_A}.$$

(At the characteristic CKB values the convention $\frac{0}{0} = 0$ was used; at the h -distances, the convention $\frac{0}{0} = 1$ was used.)

Proof. Cf. Example 5.25 and Example 4.4, and the invariance of $U_A : |D_A| : E_A$ for the characteristic CKB values. From a characteristic CKB value u , the corresponding semi-axis is $\operatorname{artanh} u = \frac{1}{2} \operatorname{arcosh} \frac{1+u^2}{1-u^2}$, cf. (48). \square

Theorem 5.29. *The unordered ratio*

$$(108) \quad \{U_A - |D_A| : U_A + |D_A| : U_A - |D_A| + 2E_A\}$$

and (when $|D_A| = E_A > 0$) the possible choice of type of possibly degenerate

h -distance band / h -elliptic parabolic disk

together form a full invariant of the conformal range up to h -isometries.

The characteristic CKB values can be obtained as follows: We divide by the term of highest absolute value any other term, and take square root.

Proof. This is just the previous theorem rephrased in a more compact form. \square

Theorem 5.30. *The inhomogeneous ratio*

$$(109) \quad U_1(A)^{[1]} : U_2(A)^{[2]} : U_3(A)^{[3]}$$

and (when $|D_A| = E_A > 0$) the possible choice of type of possibly degenerate

h -distance band / h -elliptic parabolic disk

together form a full invariant of the conformal range up to h -isometries.

(The inhomogeneous ratio is understood such that it scales with $\lambda, \lambda^2, \lambda^3$ in the corresponding terms for $\lambda \neq 0$.)

Proof. (108) and (109) are equivalent by the corresponding cubic polynomials. \square

The previous statements can be formulated in relation to the quadratic forms of the range.

Theorem 5.31. (a) From the eigenvalues of $\mathbf{G}_{\text{CKB(P)}}^{\mathbb{R}}(A)\mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R},0}$ or $\mathbf{G}_{\text{CKB}}^{\mathbb{R}}(A)\mathbf{Q}_{\text{CKB}}^{\mathbb{R},0}$, the characteristic CKB values can be obtained as follows: We divide by the eigenvalue of highest absolute value any other eigenvalue, and take square root.

More generally, the Jordan form of the matrices above, up to non-zero scalar multiples, is a complete invariant of the conformal range up to h -isometries.

(b) In the case when A is non-normal, from the eigenvalues of $\mathbf{G}_{\text{CKB(P)}}^{\mathbb{R},0}\mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A)$ or $\mathbf{G}_{\text{CKB}}^{\mathbb{R},0}\mathbf{Q}_{\text{CKB}}^{\mathbb{R}}(A)$, the characteristic CKB values can be obtained as follows: We divide the lowest eigenvalue by any other eigenvalue, and take square root.

More generally, the Jordan form of the matrices above, up to non-zero scalar multiples, is a complete invariant of the conformal range up to h -isometries.

If A is normal, then only the rank of the matrix can be derived from the ratio of the eigenvalues or from Jordan form of the matrices, up to non-zero scalar multiples.

In the cases above the convention $\frac{0}{0} = 0$ is used.

Proof. This is a consequence of Theorem 5.28 and Theorem 5.5, cf. Example 5.6. \square

Remark 5.32. Thus the nature of the Jordan forms translates to the nature of the pencils generated by the conformal ranges and the base forms. \triangle

5.G. The synthetic geometry of the conformal range.

Consider, $\iota_*^{[2]} = \pi_*^{[2]} \circ \iota_*$, the embedding of the Riemann-sphere projected to the asymptotically closed hyperbolic plane. In terms of hyperbolic geometry, if the eigenvalues of A are λ_1, λ_2 , then $\iota_*^{[2]}(\lambda_1)$ and $\iota_*^{[2]}(\lambda_2)$ will be called as the h -eigenpoints of A .

Lemma 5.33. If f is a real Möbius transformation applicable to the complex 2×2 matrix A , then the h -eigenpoints of A are related to the h -eigenpoints of $f(A)$ by the h -isometries associated to f .

Proof. This is obvious in the Poincaré half-plane model, and this naturally transcribes to the other models. \square

Lemma 5.34. In terms of hyperbolic geometry, if the eigenvalues of A are λ_1, λ_2 , then

$$\begin{aligned}
 (110) \quad d^* \left(\iota_*^{[2]}(\lambda_1), \iota_*^{[2]}(\lambda_2) \right) &= \\
 &= d^{\text{PH}} \left((\text{Re } \lambda_1, |\text{Im } \lambda_1|), (\text{Re } \lambda_2, |\text{Im } \lambda_2|) \right) = d^{\text{CKB(P)}} \left((\text{Re } \lambda_1, |\lambda_1|^2), (\text{Re } \lambda_2, |\lambda_2|^2) \right) \\
 &= \log \left| \frac{\sqrt{\frac{E_A}{|D_A|}} + 1}{\sqrt{\frac{E_A}{|D_A|}} - 1} \right| = \begin{cases} 2 \operatorname{artanh} \sqrt{\frac{E_A}{|D_A|}} & \text{if } \frac{E_A}{|D_A|} \leq 1, \\ 2 \operatorname{arcoth} \sqrt{\frac{E_A}{|D_A|}} & \text{if } \frac{E_A}{|D_A|} \geq 1 \end{cases} = \arccos \left| \frac{E_A + |D_A|}{E_A - |D_A|} \right|.
 \end{aligned}$$

Proof. It can be computed by (47) but with the second coordinate omitted. \square

It is reasonable to call (110) as the h -eigendistance of A . We set

$$Q_{\text{CKB(P)}}^A(x_{\text{CKB(P)}}, z_{\text{CKB(P)}}) = \begin{bmatrix} x_{\text{CKB(P)}} \\ z_{\text{CKB(P)}} \\ 1 \end{bmatrix}^\top \mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R}}(A) \begin{bmatrix} x_{\text{CKB(P)}} \\ z_{\text{CKB(P)}} \\ 1 \end{bmatrix};$$

$$Q_{\text{CKB(P)}}^*(x_{\text{CKB(P)}}, z_{\text{CKB(P)}}) = \begin{bmatrix} x_{\text{CKB(P)}} \\ z_{\text{CKB(P)}} \\ 1 \end{bmatrix}^\top \mathbf{Q}_{\text{CKB(P)}}^{\mathbb{R},0} \begin{bmatrix} x_{\text{CKB(P)}} \\ z_{\text{CKB(P)}} \\ 1 \end{bmatrix}.$$

Theorem 5.35. *Assume that A is a complex 2×2 matrix with no real eigenvalues. Let Λ_1 and Λ_2 be its h -eigenpoints. Let \mathbf{x} be a hyperbolic point (with coordinates $\mathbf{x}_{\text{CKB(P)}} = (x_{\text{CKB(P)}}, z_{\text{CKB(P)}})$ in the CKB(P) model). We will use the abbreviations*

$$f_1 = d(\mathbf{x}, \Lambda_1), \quad f_2 = d(\mathbf{x}, \Lambda_2), \quad m^+ = \operatorname{arcosh} \frac{U_A + E_A}{||D_A| - E_A|}.$$

Then

$$(111) \quad (\cosh(f_1 + f_2) - \cosh m^+) (\cosh m^+ - \cosh(f_1 - f_2)) =$$

$$= \frac{1}{(|D_A| - E_A)^2} \cdot \frac{Q_{\text{CKB(P)}}^A(x_{\text{CKB(P)}}, z_{\text{CKB(P)}})}{-4Q_{\text{CKB(P)}}(x_{\text{CKB(P)}}, z_{\text{CKB(P}})}.$$

Proof. As the RHS of (111) is one of the naturally transforming quantities, it is sufficient to check to the canonical representatives. This means $L_{\alpha,t}^\pm$ with $0 < \alpha \leq \pi/2$. Also, ‘CKB(P)’ can be changed to ‘CKB’ throughout. Then

$$(\Lambda_1)_{\text{CKB}} = (\cos \alpha, 0) \quad \text{and} \quad (\Lambda_2)_{\text{CKB}} = (-\cos \alpha, 0)$$

can be assumed. Furthermore,

$$\frac{U_A + E_A}{||D_A| - E_A|} = \frac{1 + 2t^2 + (\cos \alpha)^2}{(\sin \alpha)^2}, \quad \text{and} \quad (|D_A| - E_A)^2 = (\sin \alpha)^4.$$

The LHS of (111) expands to

$$1 - (\cosh f_1)^2 - (\cosh f_2)^2 - (\cosh m^+)^2 + 2(\cosh f_1)(\cosh f_2)(\cosh m^+)$$

(which is a symmetric expression in f_1, f_2, m^+). This form is particularly suitable to substitute the concrete expressions of f_1, f_2, m^+ in the CKB model, which are all ‘‘arcosh’’es, cf. (46). Then the equality checks out. \square

Corollary 5.36. *Consider the setup of Theorem 5.35. Then the points \mathbf{x} of $\text{DW}^{\mathbb{R}}(A)$ are described as the solutions of the inequality*

$$f_1 + f_2 \leq m^+,$$

with equality for the boundary. Thus, the boundary is the synthetic h -ellipse with the h -eigenpoints as the foci and the length of the major axis as the distance sum.

Proof. This is trivial if A is normal, and the conformal range is a segment. If A is not normal, then $Q_{\text{CKB(P)}}^A(x_{\text{CKB(P)}}, z_{\text{CKB(P)}}) \leq 0$ describes the conformal range, with equality on the boundary. The equation is (111) but with the RHS replaced by ≤ 0 . Then $|f_1 - f_2|$ is less or equal than the h -eigendistance. The h -eigendistance is strictly less than the length of the major axis. (This is immediate for the canonical representatives in the CKB model). Consequently $\cosh m^+ - \cosh(f_1 - f_2) > 0$. That makes the inequality $\cosh(f_1 + f_2) - \cosh m^+ \leq 0$, which can be rewritten as in the statement. \square

The statement above can be considered as the closest analogue of the elliptical range theorem regarding the numerical range. If we are a bit more familiar with h -conics, then the statement above can be obtained in another way:

Lemma 5.37. *Assume that A is a complex 2×2 matrix. Consider the possibly degenerate h -ellipse $DW^{\mathbb{R}}(A)$. Let s^+ be the h -length of its major semi-axis, s^- be the h -length of its minor semi-axis, and let c be half of the h -eigendistance of A . Then*

$$(112) \quad \cosh s^+ = (\cosh s^-)(\cosh c)$$

(the hyperbolic Pythagorean law) holds.

Proof. It is sufficient to check the identity squared. That is

$$\frac{1 + \cosh 2s^+}{2} = \frac{1 + \cosh 2s^-}{2} \frac{1 + \cosh 2c}{2}.$$

This, however, is already transparent from Theorem 5.28 and Lemma 5.34. \square

Alternative proof for Corollary 5.36. We already know that the conformal range must be given by a synthetic h -ellipse. The h -eigenpoints are in symmetric positions and, by (112), they fit to role of the foci. So, they are the foci, etc. \square

Assume that C is a h -horocycle with asymptotic point P . Let d_C be the oriented distance from C (negative inside the horodisk, positive outside). Then d_C will be called a distance function from P . Distance functions from P differ from each other in an additive constant (they belong to parallel horocycles).

Theorem 5.38. *Suppose that A is a complex 2×2 matrix with a strictly complex eigenvalue λ and a real eigenvalue λ_0 . Let the corresponding h -eigenpoints be Λ and Λ_0 , respectively. Let \mathbf{x} be the hyperbolic point represented by coordinates $(x_{\text{CKB}(\text{P})}, z_{\text{CKB}(\text{P})})$ in the $\text{CKB}(\text{P})$ model. Let d_0 be a distance function from Λ_0 . Furthermore, let V be the vertex of the conformal range. Then*

$$(113) \quad \left(\cosh(d(\mathbf{x}, \Lambda)) - \cosh(d_0(\mathbf{x}) + d_0(\Lambda) - 2d_0(V)) \right) \cdot \exp(d_0(\mathbf{x}) - d_0(\Lambda)) = \\ = \frac{1}{4|D_A|(U_A + |D_A|)} \cdot \frac{Q_{\text{CKB}(\text{P})}^A(x_{\text{CKB}(\text{P})}, z_{\text{CKB}(\text{P})})}{-4Q_{\text{CKB}(\text{P})}^*(x_{\text{CKB}(\text{P})}, z_{\text{CKB}(\text{P})})}.$$

Proof. As the RHS of (113) is one of the naturally transforming quantities, it is sufficient to check for the canonical representatives S_β . Also, ‘ $\text{CKB}(\text{P})$ ’ can be changed to ‘ CKB ’ throughout. Then

$$V_{\text{CKB}} = (0, 0), \quad \Lambda_{\text{CKB}} = \left(0, \frac{(\cos \beta)^2}{(\cos \beta)^2 - 2} \right), \quad \text{and} \quad (\Lambda_0)_{\text{CKB}} = (0, -1).$$

Moreover,

$$d_0^{\text{CKB}}(x_{\text{CKB}}, z_{\text{CKB}}) = \log \left(\frac{1 + z_{\text{CKB}}}{\sqrt{1 - (x_{\text{CKB}})^2 - (z_{\text{CKB}})^2}} \right)$$

can be assumed. (Then $d_0^{\text{CKB}}(V_{\text{CKB}}) = 0$.) Furthermore,

$$U_A + D_A = \frac{1}{2} \quad \text{and} \quad |D_A| = \frac{1}{4}(\sin \beta)^2.$$

With this data, equality (113) checks out. \square

Corollary 5.39. *Consider the setup of Theorem 5.38. Then the points \mathbf{x} of the conformal range are given by the inequality*

$$d(\mathbf{x}, \Lambda) + d_0(\mathbf{x}) + d_0(\Lambda) - 2d_0(V) \leq 0,$$

with equality on the boundary. The boundary is the h -elliptic parabola, whose points are of equal distance from Λ and the horocycle with equation $d_0(\tilde{\mathbf{x}}) + d_0(\Lambda) - 2d_0(V) = 0$. In other terms, this is the synthetic h -elliptic parabola with vertex V and focus Λ .

Proof. Consider the horodisk with equation $d_0(\tilde{\mathbf{x}}) + d_0(\Lambda) - 2d_0(V) \leq 0$. Then it is easy to see that $\tilde{\mathbf{x}} = \Lambda$ is in the interior. Thus $d(\mathbf{x}, \Lambda) > |d_0(\mathbf{x}) + d_0(\Lambda) - 2d_0(V)|$ holds for any \mathbf{x} in the exterior of horocycle. Consequently, $d(\mathbf{x}, \Lambda) = |d_0(\mathbf{x}) + d_0(\Lambda) - 2d_0(V)|$ may hold only with $d(\mathbf{x}, \Lambda) = -(d_0(\mathbf{x}) + d_0(\Lambda) - 2d_0(V))$. Even $d(\mathbf{x}, \Lambda) \leq |d_0(\mathbf{x}) + d_0(\Lambda) - 2d_0(V)|$ may hold only with $d(\mathbf{x}, \Lambda) \leq -(d_0(\mathbf{x}) + d_0(\Lambda) - 2d_0(V))$. \square

Assuming more familiarity with h -conics, the statement above can be obtained in another way:

Lemma 5.40. *Consider the setup of Theorem 5.38. Suppose that s^- is the length of the minor semi-axis (or rather half of the length of the minor axis).*

$$(114) \quad \exp d(V, \Lambda) = \exp(d_0(V) - d_0(\Lambda)) = \cosh s^-.$$

Proof. It is sufficient to check it for S_β with $\beta \in (0, \pi/2]$. Then, indeed, the equality checks out. \square

Alternative proof for Corollary 5.39. We already know that the conformal range must be a synthetic h -elliptic parabola. We may also know that h -elliptic parabolas have the property (114) with the focus in the place of Λ . This makes Λ the focus, etc. \square

The remaining cases are similar to the ones of the Davis–Wielandt shell:

Theorem 5.41. *Assume that A has two distinct real eigenvalues λ_1, λ_2 . (Thus $0 < |D_A| = E_A$.)*

(a) *Then the conformal range is a distance band with asymptotic points $\iota^{[2]}(\lambda_1)$ and $\iota^{[2]}(\lambda_2)$; and whose radius is half of the minor axis,*

$$s^- = \frac{1}{2} \operatorname{arcosh} \frac{U_A}{|D_A|}.$$

(b) *Let \mathbf{x} be the hyperbolic point represented by coordinates $(x_{\text{CKB}(P)}, z_{\text{CKB}(P)})$ in the CKB(P) model. Let*

$$d = d\left(\mathbf{x}, \operatorname{axis} \operatorname{DW}^{\mathbb{R}}(A)\right).$$

Then

$$(\cosh 2d) - (\cosh 2s^-) = \frac{1}{|D_A|(U_A + |D_A|)} \cdot \frac{Q_{\text{CKB}(P)}^A(x_{\text{CKB}(P)}, z_{\text{CKB}(P)})}{-4Q_{\text{CKB}(P)}^*(x_{\text{CKB}(P)}, z_{\text{CKB}(P)})}.$$

Proof. Again, it is sufficient to check the conformal representatives $L_{0,t}^\pm = L_t$ in the CKB model, where it is a routine calculation. \square

Theorem 5.42. *Assume that A has two equal real eigenvalues, λ_0 . (Thus $0 = |D_A| = E_A$.)*

(a) *Then the conformal range is a possibly degenerate horodisk with asymptotic point $\iota^{[2]}(\lambda_0)$.*

(b) Let d_0 be the oriented distance from its boundary (positive outside the horodisk, negative inside, identically $+\infty$ in the normal case). Let \mathbf{x} be the hyperbolic point represented by coordinates $(x_{\text{CKB}(\mathbb{P})}, z_{\text{CKB}(\mathbb{P})})$ in the $\text{CKB}(\mathbb{P})$ model. Then

$$\exp(2d_0(\mathbf{x})) - 1 = \frac{1}{(U_A)^2} \cdot \frac{Q_{\text{CKB}(\mathbb{P})}^A(x_{\text{CKB}(\mathbb{P})}, z_{\text{CKB}(\mathbb{P})})}{-4Q_{\text{CKB}(\mathbb{P})}^*(x_{\text{CKB}(\mathbb{P})}, z_{\text{CKB}(\mathbb{P})})}.$$

Proof. Again, it is sufficient to check the conformal representatives S_0 and $\mathbf{0}_2$ in the CKB model, where it is a routine calculation. \square

For the sake of the next statement, see Theorem 2.30 and Theorem 2.31, and the preceding discussions. The following observation applies to all real types:

Theorem 5.43. *Suppose that A is a 2×2 complex matrix with*

- (i) *real eigenvalues λ_1, λ_2 ; or*
- (ii) *conjugate strictly complex eigenvalues $\lambda, \bar{\lambda}$.*

Then $\text{DW}_^{\mathbb{R}}(A)$ is the*

- (i) *possibly degenerate h -distance band or h -horodisk, which is tangent to the asymptotic circle at $\iota_*^{[2]}(\lambda_1), \iota_*^{[2]}(\lambda_2)$ (double tangency counts in the parabolic case); or*
- (ii) *h -disk with center $\iota_*^{[2]}(\lambda) = \iota_*^{[2]}(\bar{\lambda})$;*

such that the signed distance of $O_^{\mathbb{R}} = \iota_*^{[2]}(O_*)$ from $\text{DW}_*^{\mathbb{R}}(A)$ is given by formula*

$$(115) \quad \overleftarrow{\text{dis}}_*(O_*^{\mathbb{R}}, \text{DW}_*^{\mathbb{R}}(A)) = \text{RHS of (57)},$$

and the signed distance of $\text{DW}_^{\mathbb{R}}(A)$ from $\mathcal{P}_*^{\infty, \mathbb{R}} = \mathcal{P}_*^{\infty} \cap i^{[2]}(\overline{H_*^2})$ is given by formula*

$$(116) \quad \overleftarrow{\text{dis}}_*(\text{DW}_*^{\mathbb{R}}(A), \mathcal{P}_*^{\infty, \mathbb{R}}) = \text{RHS of (59)};$$

and the norm distance of $\text{DW}_^{\mathbb{R}}(A)$ is given by formula*

$$(117) \quad \overleftarrow{\text{dis}}_{*\infty}^{\mathbb{R}}(\text{DW}_*^{\mathbb{R}}(A)) = -\log \|A\|_2.$$

In conjunction to the focal properties (i)/(ii) either equality characterizes $\text{DW}_^{\mathbb{R}}(A)$.*

Proof. Here, the configuration is symmetric to the embedded h -plane $i^{[2]}(\overline{H_*^2})$, thus $\overleftarrow{\text{dis}}_*(O_*^{\mathbb{R}}, \text{DW}_*^{\mathbb{R}}(A)) = \overleftarrow{\text{dis}}_*(O_*, \text{DW}_*(A))$ and $\overleftarrow{\text{dis}}_*(\text{DW}_*^{\mathbb{R}}(A), \mathcal{P}_*^{\infty, \mathbb{R}}) = \overleftarrow{\text{dis}}_*(\text{DW}_*(A), \mathcal{P}_*^{\infty})$ hold. Then Theorem 2.30 and Theorem 2.31 can be invoked. (117) can be seen similarly or directly. Sufficiency to characterize the conformal range can be seen easily. \square

Corollary 5.36, Corollary 5.39, Theorem 5.41(a) and Theorem 5.42(a) provide the analogues of the elliptical range theorem for the conformal range. The only imperfection is that Theorem 5.42(a) in itself does not provide a complete description of the conformal range; then something like Theorem 5.43 is needed.

Remark 5.44. At this point we can review what would constitute a minimalistic presentation of the (analogue of) the elliptical range theorem for the conformal range:

In terms of linear algebra, we need subsections 1.A–1.F, and 4.A–4.C. An appropriate knowledge of hyperbolic geometry is required. This must include a good knowledge of h -cycles but may include basics of h -conics in order to avoid elementary discussions. Regarding the conformal range, we need its definition and conformal invariance, but not more. From subsection 5.F we need Theorem 5.22 with the raw computation as proof, the subsequent geometric discussion of the objects, and Theorem 5.28. From subsection 5.G, we need Lemma 5.33, Lemma 5.34; Corollary 5.36 with its alternative proof, Corollary 5.39 with its alternative proof. The remaining cases are of real type,

in particular h -cycles, whose overview is simple. In that way, there is no mention of the Davis–Wielandt shell, qualitative elliptical range theorems, or $\mathbf{Q}_*^{\mathbb{R}}$. This approach is analogous to that of [25]. \triangle

In general, we see that the numerical range and the conformal range derive from the Davis–Wielandt shell similarly, and they have parallel properties, but the conformal range is a bit more complicated. We also see that the synthetic geometric properties of the ranges (including the shell) can be derived from a few canonical representatives.

5.H. The inverse problem.

Discussion 5.45. Let us make some comments on the inverse problem of recovering the five data from the conformal range. From $DW_{\text{CKB(P)}}^{\mathbb{R}}(A)$, the matrix $\mathbf{G}_{\text{CKB(P)}}^{\mathbb{R}}(A)$ can be recovered by (99) / (100). Using Lemma 5.11, if λ_1, λ_2 are the eigenvalues of A , then $\lambda_1, \bar{\lambda}_1, \lambda_2, \bar{\lambda}_2$ are the solutions of the equation

$$\begin{bmatrix} 2\lambda \\ -1 \\ -\lambda^2 \end{bmatrix}^{\top} \mathbf{G}_{\text{CKB(P)}}^{\mathbb{R}}(A) \begin{bmatrix} 2\lambda \\ -1 \\ -\lambda^2 \end{bmatrix} = 0.$$

Having obtained the possible eigenvalue pairs, the h -eigenpoints are obvious. In order to recover A , we have to choose concrete complex numbers for the eigenpoints. Having them chosen, the ordering relation $|D_A| \stackrel{\leq}{\geq} E_A$ is clear. Now, from Theorem 5.5(a), the set $\{U_A - |D_A|, U_A + |D_A|, U_A - |D_A| + 2E_A\}$ can be recovered. Using the ordering relation, $U_A, |D_A|, E_A$ can be recovered individually. In particular, by (10), the canonical triangular form of A is also recovered. Consequently, the five data of A is also recovered. As $\mathbf{G}_{\text{CKB(P)}}^{\mathbb{R}}(A)$ is in direct relationship to the reduced five data; this process also solves the problem of the reconstruction of the five data from the reduced five data. (The main point is using Theorem 5.5(a); deciding $|D_A| \stackrel{\leq}{\geq} E_A$ may, in fact, come after that.) \square

Remark 5.46. One can recover the ‘five data’ $\text{Re tr } A, \text{Im tr } A, \text{Re det } A, \text{Im det } A, \text{tr } A^*A$ from the ‘reduced five data’ $\text{Re tr } A, |\text{tr } A|^2 + 2 \text{Re det } A, \text{Re}((\text{det } A)\overline{\text{tr } A}), |\text{det } A|^2, \text{tr } A^*A$ but up to finitely many choices. Geometrically this is equivalent to recovering a h -tube from its projection. This is not entirely straightforward as, in general, one has to solve a cubic equation to do this. (Cf. also Discussion 5.48.) However, if $\text{Re tr } A, \text{Im tr } A, \text{Re det } A, \text{Im det } A, \text{tr } A^*A$ are already given, then it is relatively easy to find all other possibilities to the same corresponding reduced data: The possibilities are given by the scheme

$$\left. \begin{array}{l} \text{Re tr } A \\ \pm_1 \text{Im tr } A \\ \text{Re det } A \\ \pm_1 \text{Im det } A \\ \text{tr } A^*A \end{array} \right\} \leftrightarrow \left\{ \begin{array}{l} \text{Re tr } A \\ \pm_2 \cdot 2 \text{Re } \sqrt{D_A} \quad \equiv \quad \pm_3 \cdot 2 \text{Im } \sqrt{-D_A} \\ \frac{|\text{tr } A|^2}{4} - |D_A| \\ \pm_2 \cdot \text{Re}((\sqrt{D_A})(\overline{\text{tr } A})) \quad \equiv \quad \pm_3 \cdot \text{Im}((\sqrt{-D_A})(\overline{\text{tr } A})) \\ \text{tr } A^*A \end{array} \right.$$

where \pm_1 and \pm_2 can be chosen arbitrarily.

Note that

$$\begin{aligned} |D_A| &\leftrightarrow E_A, \\ U_A - |D_A| &\leftrightarrow U_A - |D_A|. \end{aligned}$$

The ambiguity is ‘in the spectral data’, due to possible complex conjugation of the eigenvalues. The degeneracy is

- 1-fold in the real hyperbolic and real parabolic cases,

1 real hyperbolic case \circlearrowleft

1 real parabolic case \circlearrowleft

- 2-fold in the semi-real case,

2 semi-real cases ($\text{Im tr } A \geq 0$) \circlearrowleft

- 3-fold in the real elliptic / non-real parabolic cases,

2 non-real parabolic cases ($\text{Im tr } A \geq 0$) \leftrightarrow 1 real-elliptic case

- 4-fold in the quasielliptic/quasihyperbolic cases

2 quasihyperbolic cases ($\text{Im tr } A \geq 0$) \leftrightarrow 2 quasielliptic cases ($(B_A)^0$ distinguishes).

Although the various ambiguities are essentially of spectral nature; conjugation of both eigenvalues is realized by the operation $A \rightsquigarrow A^*$. \triangle

5.I. Invariants in terms of the reduced five data.

Lemma 5.47. *The quantities*

$$C_1(A) = \frac{U_2(A)}{(U_1(A))^2}$$

and

$$C_2(A) = \frac{U_3(A)}{(U_1(A))^3}$$

are invariants of complex 2×2 matrices for real Möbius transformations and unitary conjugation. (They are “chaotic” only for real scalar matrices A .) Moreover, they are rational in the reduced five data.

Taken together with the real-hyperbolic / semi-real distinction they are full invariants of conformal ranges up to h -isometries.

Proof. This is a consequence of Discussion 5.7 and Theorem 5.30. \square

Discussion 5.48. $C_1(A)$ and $C_2(A)$ are related to (105) in terms of characteristic polynomials by

$$(118) \quad \det \left(\lambda \text{Id}_3 - \tilde{\mathbf{G}}_{\text{CKB}(P)}^{\mathbb{R}}(A) \mathbf{Q}_{\text{CKB}(P)}^{\mathbb{R},0} \right) = \lambda^3 + \frac{1}{4} \lambda^2 + \frac{1}{16} C_1(A) \lambda + \frac{1}{64} C_2(A).$$

Now, from an open subset, $C_1(A)$ and $C_2(A)$ can be rational numbers, such that the roots of (118) require cubic roots. As $\frac{U_A - |D_A|}{-4(3U_A - |D_A| + 2E_A)}$, etc., requires a cubic root, the reconstruction of the usual conformal invariants requires a cubic root. In particular, the reconstruction of the five data from the reduced five data typically requires solving a cubic equation. \square

APPENDIX A. THE ALGEBRAIC PROJECTION OF QUADRATIC FORMS

The following are very elementary but useful to keep in mind:

Assume that $\mathbf{M} = \begin{bmatrix} \mathbf{A} & \mathbf{b} \\ \mathbf{b}^\top & c \end{bmatrix}$ is a symmetric block matrix of type $(n-1|1)$. We can

think that it represents the quadratic form $\begin{bmatrix} \mathbf{x} \\ y \end{bmatrix} \mapsto \begin{bmatrix} \mathbf{x} \\ y \end{bmatrix}^\top \begin{bmatrix} \mathbf{A} & \mathbf{b} \\ \mathbf{b}^\top & c \end{bmatrix} \begin{bmatrix} \mathbf{x} \\ y \end{bmatrix}$. Our objective is to project (not restrict!) it to its first $(n-1) \times (n-1)$ block / first $n-1$ variables. We may consider various methods. At this point we compare them only algebraically (although we may have strong reasons for their equivalence on geometrical grounds).

(iii) We can invert the matrix \mathbf{M} , take restriction to the $(n-1) \times (n-1)$ block, and invert back. (This follows the ideology that projections and restrictions are dual.) The feasibility of this process assumes not only that \mathbf{M} is invertible but that $c \neq 0$. (Indeed, if $\begin{bmatrix} \mathbf{U}^{-1} & \mathbf{v} \\ \mathbf{v}^\top & d \end{bmatrix}$ is the inverse of \mathbf{M} , then multiplication yields $\mathbf{U}^{-1}\mathbf{b} + \mathbf{v}^\top c = 0$; thus $c = 0$ would imply $\mathbf{b} = 0$ in contradiction to the invertibility of \mathbf{M} .) In that case,

$$\begin{aligned} \begin{bmatrix} \mathbf{U}^{-1} & \mathbf{v} \\ \mathbf{v}^\top & d \end{bmatrix} &\equiv \begin{bmatrix} \mathbf{A} & \mathbf{b} \\ \mathbf{b}^\top & c \end{bmatrix}^{-1} = \left(\begin{bmatrix} \text{Id} & \frac{1}{c}\mathbf{b} \\ & 1 \end{bmatrix} \begin{bmatrix} \mathbf{A} - \frac{\mathbf{b}\mathbf{b}^\top}{c} & \\ & c \end{bmatrix} \begin{bmatrix} \text{Id} & \\ \frac{1}{c}\mathbf{b}^\top & 1 \end{bmatrix} \right)^{-1} \\ &= \begin{bmatrix} \text{Id} & \\ -\frac{1}{c}\mathbf{b}^\top & 1 \end{bmatrix} \begin{bmatrix} (\mathbf{A} - \frac{\mathbf{b}\mathbf{b}^\top}{c})^{-1} & \\ & c^{-1} \end{bmatrix} \begin{bmatrix} \text{Id} & -\frac{1}{c}\mathbf{b} \\ & 1 \end{bmatrix} \\ &= \begin{bmatrix} (\mathbf{A} - \frac{\mathbf{b}\mathbf{b}^\top}{c})^{-1} & -\frac{1}{c}(\mathbf{A} - \frac{\mathbf{b}\mathbf{b}^\top}{c})^{-1}\mathbf{b} \\ -\frac{1}{c}\mathbf{b}^\top(\mathbf{A} - \frac{\mathbf{b}\mathbf{b}^\top}{c})^{-1} & \frac{1}{c} + \frac{1}{c^2}\mathbf{b}^\top(\mathbf{A} - \frac{\mathbf{b}\mathbf{b}^\top}{c})^{-1}\mathbf{b} \end{bmatrix}. \end{aligned}$$

Thus, inversion, restriction and inversion back results $\mathbf{U} =$

$$\mathbf{A} - \frac{\mathbf{b}\mathbf{b}^\top}{c}.$$

(ii) We may pass to the quadratic form and then reduce out y according to

$$\frac{\partial}{\partial y} \left(\begin{bmatrix} \mathbf{x} \\ y \end{bmatrix}^\top \begin{bmatrix} \mathbf{A} & \mathbf{b} \\ \mathbf{b}^\top & c \end{bmatrix} \begin{bmatrix} \mathbf{x} \\ y \end{bmatrix} \right) = 0.$$

Then the resulted quadratic form is may be converted into a matrix.

Here the quadratic form

$$\mathbf{x}^\top \mathbf{A} \mathbf{x} + 2\mathbf{x}^\top \mathbf{b} y + cy^2$$

yields the reducing equation

$$2\mathbf{x}^\top \mathbf{b} + 2cy = 0,$$

which, in turn, leads to the quadratic form

$$\mathbf{x}^\top \mathbf{A} \mathbf{x} + 2\mathbf{x}^\top \mathbf{b} \left(-\frac{\mathbf{x}^\top \mathbf{b}}{c} \right) + c \left(-\frac{\mathbf{x}^\top \mathbf{b}}{c} \right)^2,$$

whose matrix is

$$\mathbf{A} - \frac{\mathbf{b}\mathbf{b}^\top}{c} = \begin{bmatrix} \text{Id} \\ -\frac{1}{c}\mathbf{b}^\top \end{bmatrix}^\top \begin{bmatrix} \mathbf{A} & \mathbf{b} \\ \mathbf{b}^\top & c \end{bmatrix} \begin{bmatrix} \text{Id} \\ -\frac{1}{c}\mathbf{b}^\top \end{bmatrix}.$$

The result is the same as before, but this method assumes only that $c \neq 0$.

(i) We may simply take the discriminant of the relevant quadratic form in variable y , and use this in order to obtain the matrix. Then the discriminant is

$$(2\mathbf{x}^\top \mathbf{b})^2 - 4\mathbf{x}^\top \mathbf{A} \mathbf{x} \cdot c,$$

whose matrix is

$$-4c\mathbf{A} + 4\mathbf{b}\mathbf{b}^\top.$$

Thus this method gives $-4c$ times the result of the previous one but no assumption on the matrix is taken. Ultimately, in the case $c \neq 0$, only a nonzero scalar factor enters. In that case, through dividing the discriminant by $-4c$ we can recover the previous results.

Additionally, we obtain

$$\det\left(\mathbf{A} - \frac{\mathbf{b}\mathbf{b}^\top}{c}\right) \cdot c = \det\begin{bmatrix} \mathbf{A} & \mathbf{b} \\ \mathbf{b}^\top & c \end{bmatrix}.$$

This is obvious from the discussion in case (iii), but, by arithmeticity, it must also valid in case (ii).

Remark A.1. For higher codimensions, method (iii) is more natural than generalizing discriminants from method (i). However, in our case, for codimension 1, method (i) apparently works out better than method (iii).

REFERENCES

- [1] Barbarin, P.: *Études de la géométrie analytique non euclidienne*. Mémoires Couronnés et Autres Mémoires, L'Académie Royale de Belgique, Coll. in-8°, **60/1.**, Bruxelles, 1901. 168 pages.
- [2] Berger, Marcel: *Geometry I, II*. Universitext. Springer-Verlag, Berlin, 1987.
- [3] Coolidge, Julian Lowell: *The Elements of Non-Euclidean Geometry*. Clarendon Press, Oxford, 1909.
- [4] Davis, Chandler: The shell of a Hilbert-space operator. *Acta Sci. Math. (Szeged)* **29** (1968), 69–86.
- [5] Davis, Chandler: The shell of a Hilbert-space operator. II. *Acta Sci. Math. (Szeged)* **31** (1970) 301–318.
- [6] Davis, Chandler: The Toeplitz-Hausdorff theorem explained. *Canad. Math. Bull.* **14** (1971), 245–246.
- [7] Davis, Chandler: Matrix-valued shell of an operator or relation. *Integral Equations Operator Theory* **1** (1978), 334–363.
- [8] Donoghue, William F., Jr.: On the numerical range of a bounded operator. *Michigan Math. J.* **4** (1957), 261–263.
- [9] D'Ovidio, Enrico: Le proprietà focali delle coniche nella metrica proiettiva. *Atti della R. Accademia delle Scienze di Torino* **26** (1891), 237–264.
- [10] D'Ovidio, Enrico: Sulle coniche confocali nella metrica proiettiva. *Atti della R. Accademia delle Scienze di Torino* **26** (1891), 314–325.
- [11] D'Ovidio, Enrico: Teoremi sulle coniche nella metrica proiettiva. *Atti della R. Accademia delle Scienze di Torino* **26** (1891), 369–379.
- [12] Fladt, Kuno: Die allgemeine Kegelschnittgleichung in der ebenen hyperbolischen Geometrie. *J. Reine Angew. Math.* **197** (1957), 121–139.
- [13] Fladt, Kuno: Die allgemeine Kegelschnittgleichung in der ebenen hyperbolischen Geometrie. II. *J. Reine Angew. Math.* **199** (1958), 203–207.
- [14] Fladt, K.: Elementare Bestimmung der Kegelschnitte in der hyperbolischen Geometrie. *Acta Math. Acad. Sci. Hung.* **15** (1964), 247–257.
- [15] Gustafson, Karl E.; Rao, Duggirala K. M.: *Numerical range. The field of values of linear operators and matrices*. Universitext. Springer-Verlag, New York, 1997.
- [16] Halmos, Paul R.: *A Hilbert space problem book*. 2nd, rev. enl. ed. (Orig. 1967.) Graduate Texts in Mathematics 19. Springer Verlag, New York, 1982.
- [17] Hausdorff, Felix: Der Wertvorrat einer Bilinearform. *Math. Z.* **3** (1919), 314–316.
- [18] Horn, Roger A.; Johnson, Charles R.: *Topics in matrix analysis*. Corr. repr. of the 1991 orig. Cambridge University Press, Cambridge, 1994.
- [19] Izmestiev, Ivan: Spherical and hyperbolic conics. In: *Eighteen essays in non-Euclidean geometry*. IRMA Lect. Math. Theor. Phys., 29. Eur. Math. Soc., Zürich, 2019. Pages 263–320.
- [20] Johnson, Charles R.: Computation of the field of values of a 2×2 matrix. *J. Res. Nat. Bur. Standards Sect. B* **78B** (1974), 105–107.

- [21] Killing, Wilhelm: *Die Nicht-Euklidischen Raumformen in Analytischer Behandlung*. B. G. Teubner, Leipzig, 1885.
- [22] Kippenhahn, R.: Über den Wertevorrat einer Matrix. *Math. Nachr.* **6** (1951), 193–228.
- [23] Kippenhahn, R.: On the numerical range of a matrix. *Linear Multilinear Algebra* **56** (2008), 185–225; translated from the German by Paul F. Zachlin and Michiel E. Hochstenbach.
- [24] Lakos, Gyula: *Convergence estimates for the Magnus expansion II. C^* -algebras*. arXiv:1910.03328
- [25] Lakos, Gyula: *A short proof of the elliptical range theorem*. arXiv:2208.06248
- [26] Li, Chi-Kwong: A simple proof of the elliptical range theorem. *Proc. Amer. Math. Soc.* **124** (1996), 1985–1986.
- [27] Li, Chi-Kwong; Poon, Yiu-Tung; Sze, Nung-Sing: Davis-Wielandt shells of operators. *Oper. Matrices* **2** (2008), 341–355.
- [28] Liebmann, Heinrich: *Nichteuklidische Geometrie*. G. J. Göschen, Leipzig, 1905.
- [29] Lins, Brian; Spitkovsky, Ilya M.; Zhong, Siyu: The normalized numerical range and the Davis–Wielandt shell. *Linear Algebra Appl.* **546** (2018), 187–209.
- [30] Massau, J.: *Note sur les géométries non euclidiennes*. (Extrait des annales de l’Association des Ingénieurs sortis des Écoles spéciales de Gand, 1904, pp. 419–456, 1905, pp. 379–513.) Dequesne-Masquillier & Fils, Mons, 1906.
- [31] Molnár, E.: Kegelschnitte auf der metrischen Ebene. *Acta Math. Acad. Sci. Hung.* **31** (1978), 317–343.
- [32] Murnaghan, Francis, D.: On the field of values of a square matrix. *Proc. Natl. Acad. Sci. USA*, **18** (1932), 246–248.
- [33] Plücker [, Julius]: Über solche Punkte, die bei Curven einer höhern Ordnung als der zweiten den Brennpuncten der Kegelschnitte entsprechen. *J. Reine Angew. Math.* **10** (1833), 84–91.
- [34] Sommerville, D. M. Y.: *Analytical conics*. 3rd rev. ed. G. Bell and Sons, London, 1933.
- [35] Story, William E.: On Non-Euclidean Properties of Conics. *Amer. J. Math.* **5** (1882), 358–381.
- [36] Toeplitz, Otto: Das algebraische Analogon zu einem Satze von Fejér. *Math. Z.* **2** (1918) 187–197.
- [37] Uhlig, Frank: The field of values of a complex matrix, an explicit description in the 2×2 case. *SIAM J. Algebraic Discrete Methods* **6** (1985), 541–545.
- [38] Vörös, Cyrill: *Analitikus Bolyai-féle geometria. 1. köt.: Síkgeometria*. [Hungarian.] Kókai, Budapest, 1909.
- [39] Vörös, Cyrill: *Analitika geometrio absoluta. 1. vol.: La ebena Bolyaia*. [Esperanto.] Kókai, Budapest, 1910.
- [40] Vörös, Cyrill: *Analitika geometrio absoluta. 2. vol.: La spaco Bolyaia*. [Esperanto.] Kókai, Budapest, 1912.
- [41] Wielandt, Helmut: Die Einschließung von Eigenwerten normaler Matrizen. *Math. Ann.* **121** (1949), 234–241.
- [42] Wielandt, H.: Inclusion theorems for eigenvalues. In: *Simultaneous linear equations and the determination of eigenvalues*. National Bureau of Standards Applied Mathematics Series, No. 29. U. S. Government Printing Office, Washington, D. C., 1953. Pages 75–78.

DEPARTMENT OF GEOMETRY, INSTITUTE OF MATHEMATICS, EÖTVÖS UNIVERSITY, PÁZMÁNY PÉTER
 S. 1/C, BUDAPEST, H-1117, HUNGARY
 Email address: lakos@cs.elte.hu