A Chiral W-Algebra Extension of $\mathfrak{so}(2,3)$

Nishant Gupta and Nemani V. Suryanarayana

Institute of Mathematical Sciences, Taramani, Chennai 600113, India &

Homi Bhabha National Institute, Anushakti Nagar, Mumbai 400094, India

E-mail: nishantg, nemani@imsc.res.in

ABSTRACT: In this note we show by implementing the associativity conditions of OPEs in terms of modes that there exist exactly two chiral algebras consisting of one Virasoro operator T(z), one $\mathfrak{sl}(2,\mathbb{R})$ current $J_a(z)$ (for $a=0,\pm 1$), and a doublet $G_s(z)$ (for $s=\pm 1/2$) of current algebra primaries of dimension 3/2. The simplest of these is the recently uncovered chiral extension of the Poincare algebra $\mathfrak{iso}(1,3)$. The second is a chiral extension of the $\mathfrak{so}(2,3)$. The latter is a novel \mathcal{W} -algebra that is analogous to the famous Bershadsky-Polyakov algebra $\mathcal{W}_3^{(2)}$, and is expected to play a role in the AdS/CFT correspondence.

Contents		
1	Introduction	1
2	OPE associativity in terms of modes	3
3	The OPE Ansatz	5
4	Imposing Jacobi identities	7
5	Conclusion	12
\mathbf{A}	$\mathcal{W}_3^{(2)}$ algebra - a rederivation	13

1 Introduction

The role of symmetries in physics cannot be over emphasised. Particularly in the context of two dimensional conformal field theories the symmetry algebras being typically infinite dimensional lead to complete determination of correlation functions of primary operators. However, it is not always easy to find out the exact symmetry algebra of a given theory. On the other hand it is equally important to know what the possible symmetry algebras of a CFT are extending beyond the usual Virasoro algebra. Exploration of this question was initiated by Zamolodchikov long ago [1] in the context of chiral conformal field theories which led to the discovery of new (typically higher spin) extensions of conformal symmetries, called the W-algebras (see [2] for a comprehensive review of such algebras). They arise in various contexts in physics including the constrained WZW models. Some of these (at least their classical limits) have been re-discovered using AdS_3/CFT_2 in more recent times (see for instance the initial works [3], [4]).

In a completely different context a novel chiral algebra extension of the Poincare algebra $\mathfrak{iso}(1,3)$ has been uncovered recently from the studies of soft-theorems of MHV graviton amplitudes in 4d [5, 6] as well as from a study of the asymptotic symmetries of $\mathbb{R}^{1,3}$ gravity [7]. This algebra consisted of six chiral operators

$$\{T(z), J_a(z), G_s(z)\}\$$
for $a \in \{0, \pm 1\}\$ and $s \in \{\pm 1/2\}.$ (1.1)

The operator T(z) is the usual chiral stress tensor with conformal dimension h = 2 and central charge c. The $J_a(z)$ are level- κ $\mathfrak{sl}(2,\mathbb{R})$ currents as well conformal primaries with h = 1. Finally the $G_s(z)$ are conformal primaries with h = 3/2 as

well as a doublet of $\mathfrak{sl}(2,\mathbb{R})$ current algebra primaries, and the OPE of two $G_s(z)$'s is non-singular. This spectrum of operators is very similar to the one in the famous $\mathcal{W}_3^{(2)}$ -algebra of Polyakov [8] and Bershadsky [9] – with the only difference being that the current algebra is an $\mathfrak{so}(1,1)$ one. However, the $\mathcal{W}_3^{(2)}$ -algebra is a \mathcal{W} -algebra with a non-trivial OPE between two $G_s(z)$. Therefore we ask: what is the most general admissible algebra of the six chiral operators in (1.1)?

The essential tool in establishing the existence of a symmetry algebra of any 2d chiral CFT is the associativity of OPEs of the chiral operators [10]. An equivalent implementation of the constraints from OPE associativity has been developed by Nahm et al. ([14] and references therein) which imposes Jacobi identities on the commutators of the modes of the chiral (quasi-) primaries. We make use of this method to answer our question.

Our result is the following. There are exactly two chiral algebras with the currents in (1.1). One of them is the chiral extension of $\mathfrak{iso}(1,3)$ of [5-7], and the other is the following chiral W-algebra:

$$[L_{m}, L_{n}] = (m - n) L_{m+n} + \frac{c}{12} m(m^{2} - 1) \delta_{m+n,0},$$

$$[L_{m}, J_{a,n}] = -n J_{a,m+n}, \quad [L_{m}, G_{s,r}] = \frac{1}{2} (n - 2r) G_{s,n+r},$$

$$[J_{a,m}, J_{b,n}] = -\frac{1}{2} \kappa m \eta_{ab} \delta_{m+n,0} + f_{ab}{}^{c} J_{c,m+n}, \quad [J_{a,n}, G_{s,r}] = G_{s',n+r} (\lambda_{a})^{s'}{}_{s},$$

$$[G_{s,r}, G_{s',r'}] = \epsilon_{ss'} \left[\alpha \left(r^{2} - \frac{1}{4} \right) \delta_{r+r',0} + \beta L_{r+r'} + \gamma (J^{2})_{r+r'} \right] + \delta(r - r') J_{a,r+r'} (\lambda^{a})_{ss'}$$
with $c = -\frac{6\kappa (1 + 2\kappa)}{5 + 2\kappa}, \quad \alpha = -\frac{1}{4} \gamma \kappa (3 + 2\kappa), \quad \beta = \frac{1}{4} \gamma (5 + 2\kappa), \quad \delta = -\frac{1}{2} \gamma (3 + 2\kappa)$

$$(1.2)$$

where $\eta_{ab} = (1 - 3a^2)\delta_{a+b,0}$, $f_{ab}{}^c = (a - b)\delta_{a+b}^c$, $(\lambda^a)_{ss'} = \frac{1}{2}\delta_{s+s'}^a$, $(\lambda_a)_{ss'} = \eta_{ab}(\lambda^b)_{ss'}$, $\kappa \neq -5/2$, and γ can be fixed to be any non-zero function of κ by rescaling the $G_{s,r}$ appropriately. Finally $(J^2)_n$ are the modes of the normal ordered quasi-primary $\eta^{ab}(J_aJ_b)(z)$.

The rest of this note is organised as follows: In section 2 we summarise the method of Nahm et al. in implementing the OPE associativity constraints in terms of modes of the currents. In section 3 we arrive at the most general ansatz for the algebra we seek using the global conformal symmetry as well as the global part of the $\mathfrak{sl}(2,\mathbb{R})$ current algebra symmetry. In there our conventions are also laid out with a list of identities that will be used in the main computation of Jacobi identities and their solutions which are presented in section 4. We conclude in section 5.

¹See [11], [12] and also the comprehensive review [13] of everything to do with OPEs.

2 OPE associativity in terms of modes

The problem of finding extensions of conformal symmetry of a 2d chiral CFT to those with extended set of symmetry currents has a long history. Such algebras can be described by the operator product expansions of the set of all the conformal primary chiral fields. The singular parts of the OPE result in Lie brackets of the modes with the regular parts forming the normal ordered products. These OPEs are demanded to be consistent with conformal symmetry as well as to satisfy the OPE associativity constraints [10]. One can implement the associativity constraints in terms of modes ϕ_n in which each chiral quasi-primary $\phi(z)$ with dimension h_{ϕ} is written as

$$\phi(z) = \sum_{n \in \mathbb{Z}} z^{-n - h_{\phi}} \phi_n \text{ for } h_{\phi} \in \mathbb{Z}_{>0}$$

or

$$\phi(z) = \sum_{r \in \mathbb{Z} + \frac{1}{2}} z^{-r - h_{\phi}} \phi_r \text{ for } h_{\phi} \in \mathbb{Z}_{\geq 0} + \frac{1}{2}.$$

This procedure was laid out fully by Nahm et al. [14] (see also [12]) developing further on earlier works such as [15]. As we follow this method we briefly list the relevant steps involved.

- 1. One starts by listing out all simple Virasoro primary fields.² In our case these are, the identity operator \mathbb{I} (h = 0), the chiral stress tensor T(z) (h = 2), the $\mathfrak{sl}(2,\mathbb{R})$ currents $J_a(z)$ (h = 1) and the current algebra primary doublet $G_s(z)$ (h = 3/2).
- 2. Then one constructs all the quasi-primaries that are formed by considering normal ordered products of pairs of simple primaries, that can appear on the right hand side of the OPE of simple primaries with dimensions up to 2h-2 where h is the largest conformal dimension in the set of simple primaries. In our case the highest dimension is that of T(z) which has h=2. Therefore we have to consider quadratic normal ordered quasi-primaries of dimension up to 2.
- 3. Then one uses global conformal symmetry (and other symmetries postulated) to solve for the structure constants that appear between simple primaries and global descendants of quasi-primaries completely. However, those between three simple primaries remain free parameters at this stage.
- 4. Finally one needs to impose the Jacobi identities on the corresponding commutators of modes of the simple primaries. This, in principle, determines the remaining structure constants.

²A primary is said to be simple if the corresponding state is orthogonal (with overlap defined using the BPZ duals) to those of all the normal ordered quasi-primaries [14].

The theorems of Nahm et al. [14] then guarantee that the resultant chiral algebra satisfies all the necessary constraints of 2d chiral CFTs.

Normal ordered quasi-primaries of interest

We have already declared our set of simple primaries. Now we need the list of relevant quasi-primaries that are normal ordered product between any two of these primaries. In terms of the modes of a primary

$$\phi_{i,n} = \oint \frac{dz}{2\pi i} z^{n+h_i-1} \phi_i(z) \tag{2.1}$$

the normal ordered product is defined by

$$(\phi_i \phi_j)_n = \sum_{k > -h_i} \phi_{j,n-k} \phi_{i,k} + \sum_{k \le -h_i} \phi_{i,k} \phi_{j,n-k} . \tag{2.2}$$

We need to consider normal ordered products of six types

$$(LL)(w), (LJ_a)(w), (LG_s)(w), (J_aJ_b)(w), (J_aG_s)(w), (G_sG_{s'})(w)$$

and their derivatives. Among these there is exactly one chiral operator of dimension $h \leq 2$ and it is $(J_a J_b)(w)$ which has h = 2. It is easy to see that the symmetric tensor $(J_{(a}J_{b)})(w)$ is a quasi-primary (whereas the anti-symmetric combination is not).³ This quasi-primary belongs to a reducible representation of the $\mathfrak{sl}(2,\mathbb{R})$ algebra generated by the zero modes of $J_a(z)$; which can be decomposed into irreps by separating the trace (constructed using the Killing form η_{ab} of $\mathfrak{sl}(2,\mathbb{R})$) and the traceless parts. So the only normal ordered quasi-primaries we need to consider are:

$$(J^2)(w) := \eta^{ab}(J_aJ_b)(w)$$
 and $(J_{(a}J_{b)})(w) - \frac{1}{3}\eta_{ab}(J^2)(w)$.

Note that these are a singlet (j = 0) and a quintuplet (j = 2) respectively of the $\mathfrak{sl}(2,\mathbb{R})$ algebra.

$$A_{[ab]}(w) := (J_a J_b)(w) - (J_b J_a)(w) - f_{ab}{}^c J_c'(w)$$

which is actually Virasoro primary as well. Using the $\mathfrak{sl}(2,\mathbb{R})$ current algebra the corresponding state, assuming $J_{a,k}|0\rangle = 0$ for $k \geq 0$ and $a = 0, \pm 1$, is

$$\left[J_{a,-1}J_{b,-1} - J_{b,-1}J_{a,-1} - f_{ab}{}^{c}J_{c,-2} \right] |0\rangle$$

which vanishes identically, making $A_{[ab]}(w)$ a null operator and so we ignore it.

³There is another quasi-primary (and a triplet of the $\mathfrak{sl}(2,\mathbb{R})$)

3 The OPE Ansatz

To construct the algebra that we seek we can start with all the OPEs that are unambiguously fixed already by the nature of the chiral operators we have postulated, namely, one T(z) (h = 2), one $J_a(z)$ (h = 1) and one $G_s(z)$ (h = 3/2). These are:

$$T(z)T(w) \sim \frac{c}{2}(z-w)^{-4} + 2(z-w)^{-2}T(w) + (z-w)^{-1}\partial T(w),$$

$$T(z)J_a(w) \sim (z-w)^{-2}J_a(w) + (z-w)^{-1}\partial J_a(w),$$

$$T(z)G_s(w) \sim (3/2)(z-w)^{-2}G_s(w) + (z-w)^{-1}\partial G_s(w)$$
(3.1)

and, following the conventions of Polyakov in [16] for the $\mathfrak{sl}(2,\mathbb{R})$ current algebra

$$J_a(z)J_b(w) \sim -\frac{\kappa}{2}\eta_{ab}(z-w)^{-2} + (z-w)^{-1}f_{ab}{}^c J_c(w),$$

$$J_a(z)G_s(w) \sim (z-w)^{-1}G_{s'}(w)(\lambda_a)^{s'}{}_s$$
(3.2)

where $a, b, \dots = 0, \pm 1$ and $s, s', \dots = \pm 1/2$. The matrices λ_a form a 2×2 matrix representation of $\mathfrak{sl}(2, \mathbb{R})$ algebra which we take to be:

$$[\lambda_a, \lambda_b] = (a - b) \lambda_{a+b}. \tag{3.3}$$

We use the following explicit form of these matrices

$$(\lambda_a)_{s'}^s = \frac{1}{2}(a - 2s')\delta_{a+s'}^s . {(3.4)}$$

We define the $\mathfrak{sl}(2,\mathbb{R})$ Killing form η_{ab} via $\operatorname{Tr}(\lambda_a\lambda_b)=\frac{1}{2}\eta_{ab}$. Using the λ_a in (3.4) we have

$$\operatorname{Tr}(\lambda_a \lambda_b) = (\lambda_a)_{s'}^{s'} (\lambda_b)_{s'}^{s} = \frac{1}{2} (3a^2 - 1)\delta_{a+b,0}$$
 (3.5)

which means $\eta_{ab} = (3a^2 - 1)\delta_{a+b,0}$ along with its inverse $\eta^{ab} = (3a^2 - 1)^{-1}\delta_{a+b,0}$.

There is a unique invariant tensor in the tensor product of two spinor (j = 1/2) representations of $\mathfrak{sl}(2,\mathbb{R})$, namely the antisymmetric $\epsilon_{ss'}$ and we define it such that $\epsilon_{-\frac{1}{2},+\frac{1}{2}} = 1$. We can lower and raise the fundamental/spinor indices on $(\lambda_a)^s_{\tilde{s}}$ using $\epsilon_{s\tilde{s}}$ and its inverse $\epsilon^{s\tilde{s}}$, and the adjoint indices using η_{ab} and its inverse η^{ab} . We will also need the matrices $(\lambda^a)_{s\tilde{s}} = \eta^{ab}\epsilon_{ss'}(\lambda_b)^{s'}_{\tilde{s}}$ which have

$$(\lambda^a)_{ss'} = \eta^{ab}(\lambda_b)_{ss'} = \frac{1}{2}\delta^a_{s+s'} \tag{3.6}$$

and so are symmetric. Some more useful identities satisfied by the matrices λ_a are

$$(\lambda_a)^r{}_s(\lambda_b)^{r'}{}_{s'}\eta^{ab} = -\frac{1}{2}\epsilon^{rr'}\epsilon_{ss'} - \frac{1}{4}\delta^r_s\delta^{r'}_{s'},$$

$$(\lambda_a)^s_{\tilde{s}}(\lambda_b)^{\tilde{s}}_{s'} = -\frac{1}{4}\eta_{ab}\,\delta^s_{s'} + \frac{1}{2}f_{ab}{}^c(\lambda_c)^s_{s'}. \tag{3.7}$$

From these one also has $\eta^{ab}(\lambda_a)^s_{\tilde{s}}(\lambda_b)^{\tilde{s}}_{s'} = -(3/4)\,\delta^s_{s'}$. One more identity that λ_a matrices satisfy is:

$$(\lambda_a)^s_{\hat{s}} \epsilon_{s'\tilde{s}} + (\lambda_a)^s_{\tilde{s}} \epsilon_{\hat{s}s'} + (\lambda_a)^s_{s'} \epsilon_{\tilde{s}\hat{s}} = 0 \tag{3.8}$$

Ansatz for $G_s(z)G_{s'}(w)$ OPE:

The only remaining OPE is that of $G_s(z)G_{s'}(w)$ and using global conformal invariance we can write the right hand side in terms of the quasi-primaries of dimension up to (and including) h = 2 and their global descendants. This list of such quasi-primaries is

$$\left\{ \mathbb{I}, \ J_a(w), \ G_s(w), \ T(w), \ (J^2)(w), \ (J_{(a}J_{b)})(w) - (1/3)\eta_{ab}(J^2)(w) \right\}$$
 (3.9)

Apart from the (global) conformal symmetry this OPE has to be consistent with the $\mathfrak{sl}(2,\mathbb{R})$ symmetry (the global part of the $\mathfrak{sl}(2,\mathbb{R})$ current algebra) as well. From the OPE of $J_a(z)$ with $G_s(w)$ we know that the index s is a doublet (j=1/2) index of this $\mathfrak{sl}(2,\mathbb{R})$. Therefore the quasi-primaries that can appear in the OPE $G_s(z)G_{s'}(w)$ can only carry $\mathfrak{sl}(2,\mathbb{R})$ indices of irreps in the tensor product of two doublets, namely the singlet (j=0) and the triplet (j=1). Therefore, in our list (3.9) of quasi-primaries neither $G_s(w)$ nor $(J_{(a}J_{b)})(w) - (1/3)\eta_{ab}(J^2)(w)$ can appear as they correspond to j=1/2 and j=2 irreps of $\mathfrak{sl}(2,\mathbb{R})$ respectively.

- The coefficients in front of the j=0 quasi-primary operators $\{\mathbb{I}, T(w), (J^2)(w)\}$ have to be proportional to the Clebsch–Gordan (CG) coefficients of $\frac{1}{2} \otimes \frac{1}{2} \to \mathbf{0}$, namely $\epsilon_{ss'}$.
- The coefficient in front of the j=1 quasi-primary operator $J_a(w)$ has to be proportional to the CG coefficients of $\frac{1}{2} \otimes \frac{1}{2} \to 1$, namely $(\lambda^a)_{ss'}$.

Thus after using all the global symmetries we arrive at the ansatz:

$$G_s(z)G_{s'}(w) = 2\alpha \,\epsilon_{ss'}(z-w)^{-3} + \delta \,(z-w)^{-2} \,(\lambda^a)_{ss'} \left[2 \,J_a(w) + (z-w) \,J_a'(w)\right] + \beta \,\epsilon_{ss'}(z-w)^{-1} \,T(w) + \gamma \,(z-w)^{-1} \,\epsilon_{ss'}(J^2)(w) . \tag{3.10}$$

The parameters $(\alpha, \beta, \gamma, \delta)$ in (3.10) as well as the central charges (c, κ) in (3.1, 3.2) are to be constrained using OPE associativity, which we implement as Jacobi identities of commutators of the modes of our simple primaries. Writing the operators T(z), $J_a(z)$ and $G_r(z)$ in terms of their modes as

$$T_n = \oint \frac{dz}{2\pi i} z^{n+1} T(z), \quad J_{a,n} = \oint \frac{dz}{2\pi i} z^n J_a(z), \quad G_{s,r} = \oint \frac{dz}{2\pi i} z^{s+\frac{1}{2}} G_r(z) \quad (3.11)$$

with $n \in \mathbb{Z}$ and $r \in \mathbb{Z} + \frac{1}{2}$, the OPEs (3.1, 3.2, 3.10) translate to:

1. Virasoro algebra:

$$[L_m, L_n] = (m-n) L_{m+n} + \frac{c}{12} m(m^2 - 1) \delta_{m+n,0}$$
(3.12)

2. Virasoro primaries:

$$[L_m, J_{a,n}] = -n J_{a,m+n}, \quad [L_m, G_{s,r}] = \frac{1}{2}(n-2r) G_{s,n+r}$$
 (3.13)

3. $\mathfrak{sl}(2,\mathbb{R})$ current algebra:

$$[J_{a,m}, J_{b,n}] = -\frac{1}{2} \kappa \, m \, \eta_{ab} \, \delta_{m+n,0} + f_{ab}{}^{c} \, J_{c,m+n}$$
 (3.14)

4. Current algebra primary:

$$[J_{a,n}, G_{s,r}] = G_{s',n+r}(\lambda_a)^{s'}, \tag{3.15}$$

5. Commutator $[G_{s,r}, G_{s',r'}]$:

$$[G_{s,r},G_{s',r'}] = \epsilon_{ss'} \left[\alpha \left(r^2 - \frac{1}{4} \right) \delta_{r+r',0} + \beta L_{r+r'} + \gamma (J^2)_{r+r'} \right] + \delta(r-r') J_{a,r+r'}(\lambda^a)_{ss'}$$
(3.16)

Finally, as summarised in section 2, following [14] we have to impose the Jacobi identities on these commutation relations. We turn to this next.

4 Imposing Jacobi identities

There are three species of modes $\{L_m, J_{a,n}, G_{s,r}\}$ and so a total of ten classes of Jacobi identities to impose. We choose to denote them schematically by (LLL), (LLJ), (LJJ), (JJJ), etc. in a self-evident notation. Out of these there are seven that involve at most one G and they can all be checked easily to be satisfied without providing any constraints.

Next there are two Jacobi identities that have two G's each: (LGG) and (JGG). To impose these we need the commutators $[L_m, (J^2)_n]$ and $[(J^2)_m, J_{a,n}]$. These can be computed easily using the definition

$$\eta^{ab}(J_a J_b)_n = \eta^{ab} \left[\sum_{k>-1} J_{b,n-k} J_{a,k} + \sum_{k<-1} J_{a,k} J_{b,n-k} \right]$$
(4.1)

of $(J^2)_n$ and repeatedly using (3.12 - 3.16). We find

$$[L_m, (J^2)_n] = (m-n)(J^2)_{m+n} - \frac{\kappa}{4}m(m^2-1)\delta_{m+n,0}, \qquad (4.2)$$

$$[(J^2)_m, J_{a,n}] = n(\kappa + 2) J_{a,m+n}.$$
(4.3)

We first impose the (LGG) Jacobi identity:

$$[L_m, [G_{s,r}, G_{s',r'}]] + [G_{s',r'}, [L_m, G_{s,r}]] - [G_{s,r}, [L_m, G_{s',r'}]] = 0.$$

$$(4.4)$$

For this we find:

•
$$[L_m, [G_{s,r}, G_{s',r'}]] =$$

$$\beta \epsilon_{ss'} \left[(m - r - r') L_{m+r+r'} + \frac{c}{12} m(m^2 - 1) \delta_{m+r+r'} \right] + \gamma \epsilon_{ss'} \left[(m - r - r') (J^2)_{m+r+r'} - \frac{\kappa}{4} m(m^2 - 1) \delta_{m+r+r'} \right] - \delta (r - r') (r + r') J_{a,m+r+r'}(\lambda^a)_{ss'}$$

•
$$[G_{s',r'}, [L_m, G_{s,r}]] =$$

$$\frac{1}{2}(m-2r) \Big[-\epsilon_{ss'} \left(\alpha \left((m+r)^2 - \frac{1}{4} \right) \delta_{m+r+r'} + \beta L_{m+r+r'} + \gamma (J^2)_{m+r+r'} \right) + \delta (r'-r-m) J_{a,m+r+r'} (\lambda^a)_{s's} \Big]$$

•
$$-[G_{s,r}, [L_m, G_{s',r'}]] =$$

$$-\frac{1}{2}(m-2r')\Big[\epsilon_{ss'}\left(\alpha\left((m+r')^2 - \frac{1}{4}\right)\delta_{m+r+r'} + \beta L_{m+r+r'} + \gamma (J^2)_{m+r+r'}\right) + \delta (r-r'-m) J_{a,m+r+r'}(\lambda^a)_{ss'}\Big]$$

substituting these into (4.4) we find that the coefficient of $\epsilon_{ss'} \delta_{m+r+r'}$ is

$$\frac{1}{2}m(m^2 - 1)\left[\beta \frac{c}{6} - \gamma \frac{\kappa}{2} - \alpha\right]$$

which vanishes (for generic values of m) only if

$$\alpha = \frac{c}{6}\beta - \gamma \frac{\kappa}{2} \ . \tag{4.5}$$

The coefficients of $\beta L_{m+r+r'}$, $\delta J_{a,m+r+r'}(\lambda^a)_{ss'}$, and $\gamma \epsilon_{ss'}(J^2)_{m+r+r'}$ vanish without further conditions. Next we consider the (JGG) Jacobi identity:

$$[J_{a,n}, [G_{s,r}, G_{s',r'}]] + [G_{s',r'}, [J_{a,n}, G_{s,r}]] - [G_{s,r}, [J_{a,n}, G_{s',r'}]] = 0.$$
(4.6)

For this we find

•
$$[J_{a,n}, [G_{s,r}, G_{s',r'}]]$$
 evaluates to
$$\epsilon_{ss'} [\beta - \gamma(\kappa + 2)] n J_{a,n+r+r'} + (r - r') \delta \left[f_{ab}{}^c J_{c,n+r+r'} (\lambda^b)_{ss'} - \frac{\kappa}{2} n (\lambda_a)_{ss'} \delta_{n+r+r',0} \right]$$

• $[G_{s',r'},[J_{a,n},G_{s,r}]]$ evaluates to (using $(\lambda^b)_{s'\tilde{s}}(\lambda_a)^{\tilde{s}}_s = -\frac{1}{4}\delta^b_a\epsilon_{s's} + \frac{1}{2}f_{ad}{}^b(\lambda^d)_{s's}$):

$$\left[\alpha \left(r'^{2} - \frac{1}{4}\right) \delta_{n+r+r',0} + \beta L_{n+r+r'} + \gamma (J^{2})_{n+r+r'}\right] (\lambda_{a})_{s's} + \delta(r' - n - r) \left[\frac{1}{4} \epsilon_{ss'} J_{a,n+r+r'} + \frac{1}{2} f_{ad}{}^{b} (\lambda^{d})_{ss'} J_{b,n+r+r'}\right]$$

• $-[G_{s,r}, [J_{a,n}, G_{s',r'}]]$ evaluates to

$$-\left[\alpha \left(r^{2} - \frac{1}{4}\right) \delta_{n+r+r',0} + \beta L_{n+r+r'} + \gamma (J^{2})_{n+r+r'}\right] (\lambda_{a})_{ss'} - \delta(r - n - r') \left[-\frac{1}{4} J_{b,n+r+r'} \epsilon_{ss'} + \frac{1}{2} f_{ad}{}^{b} (\lambda^{d})_{ss'} J_{b,n+r+r'} \right]$$

Substituting these back into (4.6) and collecting terms, the coefficients of $L_{n+r+r'}$, $(J^2)_{n+r+r'}$ and $f_{ab}{}^c J_{c,n+r+r'}(\lambda^b)_{ss'}$ vanish.

• Coefficient of $\delta_{n+r+r',0}(\lambda_a)_{ss'}$

$$-\delta \frac{\kappa}{2} n(r-r') + \alpha \left[\left(r'^2 - \frac{1}{4} \right) - \left(r^2 - \frac{1}{4} \right) \right] = -n(r'-r) \left[\alpha - \delta \frac{\kappa}{2} \right]$$

which imposes

$$\alpha - \delta \frac{\kappa}{2} = 0 \ . \tag{4.7}$$

• Coefficient of $\epsilon_{ss'} J_{a,n+r+r'}$ is

$$n \left[\beta - \gamma(\kappa + 2) \right] + \frac{1}{4}\delta(r' - n - r) + \frac{1}{4}\delta(r - n - r') = n \left[\beta - \gamma(\kappa + 2) - \frac{1}{2}\delta \right]$$

which imposes

$$\beta - \gamma \left(\kappa + 2\right) - \frac{1}{2}\delta = 0. \tag{4.8}$$

The last Jacobi identity left is of (GGG) type:

$$[G_{s,r}, [G_{s',r'}, G_{\tilde{s},\tilde{r}}]] + [G_{\tilde{s},\tilde{r}}, [G_{s,r}, G_{s',r'}]] - [G_{s',r'}, [G_{s,r}, G_{\tilde{s},\tilde{r}}]] = 0.$$

$$(4.9)$$

To impose this we further need the commutator $[(J^2)_n, G_{s,r}]$. Using the standard methods we find two different expressions for this commutator, namely

$$[(J^{2})_{n}, G_{s,r}] = \eta^{ab} \left[(J_{a}G_{s'})_{n+r} (\lambda_{b})^{s'}_{s} + (G_{s'}J_{b})_{n+r} (\lambda_{a})^{s'}_{s} \right] + \frac{3}{4} \left(r + \frac{1}{2} \right) G_{s,n+r}$$
(4.10)

and

$$[(J^2)_n, G_{s,r}] = 2\eta^{ab} (J_a G_{s'})_{n+r} (\lambda_b)^{s'}_{s} - \frac{3}{4} (n+1) G_{s,n+r} . \tag{4.11}$$

Taking the linear combination $(4.11) + 2 \times (4.10)$ we arrive at:

$$[(J^{2})_{n}, G_{s,r}] = \eta^{ab} \left[\frac{4}{3} (J_{a}G_{s'})_{n+r} (\lambda_{b})^{s'}_{s} + \frac{2}{3} (G_{s'}J_{a})_{n+r} (\lambda_{b})^{s'}_{s} \right] - \frac{1}{4} (n-2r)G_{s,n+r}$$
(4.12)

Using (4.12) along with the other commutators we find

$$\begin{split} [G_{s,r},[G_{s',r'},G_{\tilde{s},\tilde{r}}]] &= \epsilon_{s'\tilde{s}} \, G_{s,R} \left[-\left(\frac{\beta}{2} + \frac{\gamma}{4}\right) (R - 3r) + \frac{\delta}{4} (r' - \tilde{r}) \right] \\ &- \frac{\delta}{2} (r' - \tilde{r}) \, G_{s',R} \epsilon_{s\tilde{s}} - \gamma \left[\frac{4}{3} \left(J_a G_{\hat{s}}\right)_R + \frac{2}{3} \left(G_{\hat{s}} J_a\right)_R \right] (\lambda^a)^{\hat{s}}{}_s \epsilon_{s'\tilde{s}} \end{split}$$

where $R = r + r' + \tilde{r}$. The Jacobi identity (4.9) is obtained by adding to this two more terms that are obtained by cyclic permutations of pairs of indices $(s,r) \to (s',r') \to (\tilde{s},\tilde{r}) \to (s,r)$. Luckily the terms containing $(J_aG_{\hat{s}})_R$, and $(G_{\hat{s}}J_a)_R$ do not survive after this sum because of the identity (3.8). The rest are proportional to $G_{s,R}$ and we find

$$\epsilon_{s'\tilde{s}} G_{s,R} \left[\left(\frac{\beta}{2} - \frac{\gamma - \delta}{4} \right) (R - 3r) \right]$$

$$+ \epsilon_{\tilde{s}s} G_{s',R} \left[\left(\frac{\beta}{2} - \frac{\gamma - \delta}{4} \right) (R - 3r') \right]$$

$$+ \epsilon_{ss'} G_{\tilde{s},R} \left[\left(\frac{\beta}{2} - \frac{\gamma - \delta}{4} \right) (R - 3\tilde{r}) \right] .$$

This can vanish, for generic values of the floating indices, only if

$$\beta = \frac{\gamma - \delta}{2}.\tag{4.13}$$

This completes our analysis of the Jacobi identities with the result that the parameters $(\alpha, \beta, \gamma, \delta, c, \kappa)$ must satisfy

$$\alpha - \frac{c}{6}\beta + \gamma \frac{\kappa}{2} = 0, \quad \alpha - \delta \frac{\kappa}{2} = 0, \quad \beta - \gamma (\kappa + 2) - \frac{1}{2}\delta = 0, \quad \beta - \frac{\gamma - \delta}{2} = 0.$$

$$(4.14)$$

These are linear and homogeneous equations in the variables $(\alpha, \beta, \gamma, \delta)$. So for some non-trivial solutions for $(\alpha, \beta, \gamma, \delta)$ to exist the determinant of the coefficient matrix

$$\begin{pmatrix}
1 - \frac{c}{6} & \frac{\kappa}{2} & 0 \\
1 & 0 & 0 & -\frac{\kappa}{2} \\
0 & 1 & -(\kappa + 2) - \frac{1}{2} \\
0 & 1 & -\frac{1}{2} & \frac{1}{2}
\end{pmatrix}$$

has to vanish. This gives $c(2\kappa + 5) + 6\kappa (2\kappa + 1) = 0$ which can be solved for c (for $\kappa \neq -5/2$) as

$$c = -\frac{6\kappa \left(1 + 2\kappa\right)}{5 + 2\kappa} \ . \tag{4.15}$$

When this holds, for generic values of κ the coefficient matrix is of rank 3 and so there exists one non-trivial solution, which can be written (taking γ to be the independent variable) as

$$\alpha = -\frac{1}{4}\gamma\kappa(3+2\kappa), \quad \beta = \frac{1}{4}\gamma(5+2\kappa), \quad \delta = -\frac{1}{2}\gamma(3+2\kappa).$$
 (4.16)

This is our final result valid for generic $(\neq -5/2)$ values of κ . Some observations are in order:

• When $c \neq -\frac{6\kappa (1+2\kappa)}{5+2\kappa}$ we have to set $\alpha = \beta = \gamma = \delta = 0$, and the corresponding algebra is a chiral extension of $\mathfrak{iso}(1,3)$ with arbitrary c and κ .

This case corresponds to the algebra found in [5, 6] as the central charges there remained undetermined.

• We may also consider the limit $\gamma \to 0$ in the non-trivial solution (4.16). This corresponds to a contraction of the algebra (3.16) to the chiral extension of $\mathfrak{iso}(1,3)$, but with the relation (4.15) still in place.

The one found in [7] from an analysis of chiral boundary conditions on $\mathbb{R}^{1,3}$ or $\mathbb{R}^{2,2}$ gravity appears to belong to this class.

• When $\gamma \neq 0$ its value can be fixed to be any non-zero function of κ by an appropriate normalisation of the generators $G_{s,r}$. Then the algebra is determined completely in terms of one parameter (say) κ .

Relation to $\mathfrak{so}(2,3)$

The non-trivial solution (4.15, 4.16) makes the algebra in (3.12 - 3.16) like the Walgebras of yesteryears, particularly close to the Bershadsky-Polyakov algebra $W_3^{(2)}$. Now we will argue that, just as the $W_3^{(2)}$ algebra is based on $\mathfrak{sl}(3,\mathbb{R})$ algebra, this
new algebra is based on (and is an extension of) $\mathfrak{so}(2,3)$. For this, first note that $\mathfrak{so}(2,3)$ algebra can be written in a more suggestive manner as follows:

$$[L_{m}, L_{n}] = (m-n)L_{m+n}, \quad [\bar{L}_{m}, \bar{L}_{n}] = (m-n)\bar{L}_{m+n}, \quad [L_{m}, \bar{L}_{n}] = 0$$

$$[L_{n}, G_{s,r}] = \frac{1}{2}(n-2r)G_{s,n+r}, \quad [\bar{L}_{n}, G_{s,r}] = \frac{1}{2}(n-2s)G_{n+s,r},$$

$$[G_{s,r}, G_{s',r'}] = 2\epsilon_{rr'}\bar{L}_{s+s'} + 2\epsilon_{ss'}L_{r+r'}$$
(4.17)

⁴As a check on our computational method we provide a re-derivation of $\mathcal{W}_3^{(2)}$ in Appendix A.

for $m, n \cdots = 0, \pm 1, r, s, \cdots = \pm 1/2$. Here one may think of L_n and \bar{L}_n to generate either the $\mathfrak{so}(2,2) = \mathfrak{sl}(2,\mathbb{R}) \oplus \mathfrak{sl}(2,\mathbb{R})$ subalgebra or $\mathfrak{so}(1,3) = \mathfrak{sl}(2,\mathbb{C}) \oplus \mathfrak{sl}(2,\mathbb{C})^c$ subalgebra of $\mathfrak{so}(2,3)$. With the identification: $\bar{L}_a = J_{a,0}$ and noticing that $f_{ab}{}^c = (a-b)\delta_{a+b}^c$, the first two lines have the same form as those of our algebra (now restricted to $m, n \cdots = 0, \pm 1, r, s, \cdots = \pm 1/2$). To compare the $[G_{s,r}, G_{s',r'}]$ let us first take the large- κ (the "classical") limit in (3.16). In this limit, from (4.15, 4.16) we have⁵

$$\alpha \to -\frac{1}{2}\gamma\kappa^2, \quad \beta \to \frac{1}{2}\gamma\kappa, \quad \delta \to -\gamma\kappa, \quad c \to -6\kappa.$$
 (4.18)

Further in this limit by choosing $\gamma \to -\frac{2}{\kappa}$ we arrive at

$$\alpha \to \kappa, \quad \beta \to -1, \quad \gamma \to -\frac{2}{\kappa}, \quad \delta \to 2, \quad c \to -6 \,\kappa.$$
 (4.19)

Finally noticing that in (3.16) the first term proportional to α drops out when restricted to $r, r' = \pm 1/2$. The term proportional to δ , for $r, r' = \pm 1/2$ can be rewritten, using $(r - r') = -\epsilon_{rr'}$ and $(\lambda^a)_{ss'} = \frac{1}{2}\delta^a_{s+s'}$, as

$$\delta(r - r') J_{a,r+r'}(\lambda^a)_{ss'} \to -\epsilon_{rr'} \bar{L}_{s+s'} . \tag{4.20}$$

Therefore we conclude that the commutator (3.16) in the large- κ limit and restricted to the global modes $(r, r', s, s' \cdots = \pm 1/2)$ reads

$$[G_{s,r}, G_{s',r'}] = 2 \epsilon_{rr'} \bar{L}_{s+s'} + 2 \epsilon_{ss'} L_{r+r'} + \mathcal{O}(1/\kappa).$$

Thus the full algebra (1.2) is therefore an extension (and a deformation around large- κ) of $\mathfrak{so}(2,3)$ algebra. This completes our analysis.

5 Conclusion

We have shown that there exists a one-parameter family of chiral W-algebras generated by six chiral operators $(T(z), J_a(z), G_s(z))$ for $a = 0, \pm 1$ and $s = \pm 1/2$ with dimensions (2, 1, 3/2). In the $\kappa \to \infty$ limit it admits the $\mathfrak{so}(2, 3)$ as a subalgebra (just as the $W_3^{(2)}$ does of $\mathfrak{sl}(3, \mathbb{R})$). Just as $\mathfrak{so}(2, 3)$ admits a contraction to $\mathfrak{iso}(1, 3)$ our W-algebra (1.2) also admits a contraction $(\gamma \to 0)$ for finite κ) down to the chiral extension of $\mathfrak{iso}(1, 3)$ that appeared in the studies of graviton soft-theorems [5, 6] and the asymptotic symmetries of $\mathbb{R}^{1,3}$ gravity [7].

Using the nomenclature of [2] this new algebra (1.2) may be referred to as $W(\mathbf{1}^3, (\mathbf{3}/\mathbf{2})^2, \mathbf{2})$. Even though this algebra has a chiral primary $G_s(z)$ of half-integer dimension, is not a superconformal algebra such as the one of Knizhnik [20]. The

⁵It is curious that in this classical limit the Virasoro central charge approaches the same value as in [17–19], namely $c = -6\kappa$.

reason for this is that even though the current G_s has h = 3/2 it still has an integer spin defined by $L_0 - J_0$. In fact our algebra (1.2) is more akin to the Bershadsky-Polyakov algebra $\mathcal{W}_3^{(2)}$ – where by considering a twisted stress tensor one can make $G_s(z)$ to appear bosonic (integral dimension).

It will be interesting and important to find various contexts that realise our W-algebra. Some possibilities are the following:

- The classical (large-k limit) of the Bershadsky-Polyakov algebra $\mathcal{W}_3^{(2)}$ was shown to arise [21] in a chiral sector of the 3d higher spin gravity, described by a Chern-Simons gauge theory based on gauge algebra $\mathfrak{sl}(3,\mathbb{R}) \oplus \mathfrak{sl}(3,\mathbb{R})$. One expects that (the classical, i.e., large- κ limit of) our chiral \mathcal{W} -algebra should similarly arise with appropriate boundary conditions in the 3d conformal gravity described as a 3d Chern-Simons theory with $\mathfrak{so}(2,3)$ gauge algebra.
- Originally the (finite-k) $\mathcal{W}_3^{(2)}$ algebra was realised in the constrained WZW model with $\mathfrak{sl}(3,\mathbb{R})$ algebra [9]. One expects that a similar realisation should exist for the algebra presented here as well. It will be interesting to exhibit this.
- In a previous paper [7] we showed how the chiral extension of $\mathfrak{iso}(1,3)$ gets realised as the asymptotic symmetry algebra of 4d gravity with appropriately defined asymptotically flat boundary conditions. In an upcoming article [22] we will show how (the classical limit of) our \mathcal{W} -algebra (1.2) can be realised as asymptotic symmetry algebra of AdS_4 gravity with a well defined chiral non-Dirichlet asymptotically locally AdS_4 boundary conditions.

Also it will be important to identify other instances of chiral CFTs that admit this W-algebra as their symmetry algebra. They should play some role in both the AdS_4/CFT_3 correspondence and the symmetries of 3d conformal gravity.

${f A} \quad {\cal W}_3^{(2)} \; {f algebra} \; ext{-} \; {f a} \; {f rederivation}$

Here we recover the $W_3^{(2)}$ algebra using the methods followed in the text for our algebra (1.2). Even though the computation is straightforward we include it here as a validation of the method we followed in our main computation in the text.

From the Drinfeld-Sokolov (or the Hamiltonian) reduction of $\mathfrak{sl}(3,\mathbb{R})$ WZW model one gets a non-linear W-algebra called the Bershadsky-Polyakov algebra $W_3^{(2)}$ [8, 9]. In this problem one considers the stress tensor T(z) (h = 2), an $\mathfrak{so}(1,1)$ current H(z) (h = 1) and a couple $G_s(z)$ ($s = \pm 1/2$) of current algebra primaries with $\mathfrak{so}(1,1)$ eigenvalues ± 1 , and with h = 3/2.

Just as we did in the text for the $\mathfrak{so}(2,3)$ case we arrive at the following ansatz for the corresponding chiral algebra:

$$[L_m, L_n] = (m-n)L_{m+n} + \frac{c}{12}m(m^2-1)\delta_{m+n,0},$$

$$[L_m, H_n] = -nH_{m+n}, \quad [L_m, G_{s,r}] = \frac{1}{2}(m-2r)G_{s,m+r},$$

$$[H_n, G_{s,r}] = (2s) G_{s,n+r}, \quad [H_m, H_n] = \kappa m \delta_{m+n,0},$$

$$[G_{s,r}, G_{s',r'}] = \epsilon_{ss'} \left[\alpha \left(r^2 - \frac{1}{4} \right) \delta_{r+r'} + \beta L_{r+r'} + \gamma (H^2)_{r+r'} \right] + \delta \eta_{ss'}(r - r') H_{r+r'}$$
(A.1)

where $\epsilon_{ss'} = s' - s$, $\eta_{ss'} = \delta_{s+s',0}$ for $s, s' = \pm 1/2$. For imposing the Jocobi identities we need $[L_m, (H^2)_n]$, $[(H^2)_m, H_n]$:

$$[L_m, (H^2)_n] = (m-n)(H^2)_{m+n} + \frac{\kappa}{6}m(m^2-1)\delta_{m+n,0}, \quad [H_m, (H^2)_n] = 2\kappa \, mH_{m+n},$$

and $[(H^2)_m, G_{s,r}]$. For this we find two equivalent expressions:

$$[(H^{2})_{m}, G_{s,r}] = (4s) (HG)_{n+r} + 4s^{2} (n+1) G_{s,n+r},$$

$$[(H^{2})_{m}, G_{s,r}] = (2s) [(HG)_{n+r} + (GH)_{n+r}] - 4s^{2} (r + \frac{1}{2}) G_{s,n+r}$$

which can be combined to write:

$$[(H^2)_m, G_{s,r}] = \frac{8s}{3}(HG)_{n+r} + \frac{4s}{3}(GH)_{n+r} + \frac{4s^2}{3}(n-2r)G_{s,n+r}.$$
 (A.2)

Apart from the (LGG), (HGG) and (GGG) Jacobi identities the rest can be seen to be satisfied. Imposing these three Jacobi identities impose the following conditions:

• The (LGG) Jacobi identity imposes

$$\alpha = \frac{c}{6}\beta + \frac{\kappa}{3}\gamma \ . \tag{A.3}$$

• The (HGG) Jacobi identity requires

$$\alpha + \delta \kappa = 0, \quad \beta + 2(\delta + \kappa \gamma) = 0.$$
 (A.4)

• Finally the (GGG) identity demands

$$\beta = \frac{2}{3}(\delta - \gamma) \ . \tag{A.5}$$

Solving these conditions (A.3, A.4, A.5) leads to the following two-parameter (chosen to be α and κ) result (for $\kappa \neq -1$):

$$\beta = -\frac{2\alpha(1+\kappa)}{\kappa(1-3\kappa)}, \quad \gamma = \frac{4\alpha}{\kappa(1-3\kappa)}, \quad \delta = -\frac{\alpha}{\kappa}, \quad c = \frac{\kappa(7-9\kappa)}{1+\kappa}$$
 (A.6)

To compare it with [9] we redefine $\kappa = 1 + \frac{2k}{3}$ and choose $\alpha = \frac{1}{2}(1+k)(3+2k)$, which give (for $k \neq -3$)

$$c = 25 - \frac{24}{k+3} - 6(k+3), \quad \beta = -(k+3), \quad \gamma = 3, \quad \delta = -\frac{3}{2}(1+k)$$
 (A.7)

which match precisely with those of [9]. One could have set $\alpha = 0$ (or taken the $\alpha \to 0$ limit) which results in commuting $G_{s,r}$ which is a valid contraction of the Bershadsky-Polyakov algebra in which $G_{s,r}$ form an Abelian subalgebra.

References

- [1] A. B. Zamolodchikov, "Infinite Additional Symmetries in Two-Dimensional Conformal Quantum Field Theory," Theor. Math. Phys. **65** (1985), 1205-1213 doi:10.1007/BF01036128
- P. Bouwknegt and K. Schoutens, "W symmetry in conformal field theory," Phys. Rept. 223 (1993), 183-276 doi:10.1016/0370-1573(93)90111-P [arXiv:hep-th/9210010 [hep-th]].
- [3] M. Henneaux and S. J. Rey, "Nonlinear $W_{infinity}$ as Asymptotic Symmetry of Three-Dimensional Higher Spin Anti-de Sitter Gravity," JHEP **12** (2010), 007 doi:10.1007/JHEP12(2010)007 [arXiv:1008.4579 [hep-th]].
- [4] A. Campoleoni, S. Fredenhagen, S. Pfenninger and S. Theisen, JHEP 11 (2010), 007 doi:10.1007/JHEP11(2010)007 [arXiv:1008.4744 [hep-th]].
- [5] S. Banerjee, S. Ghosh and P. Paul, "MHV graviton scattering amplitudes and current algebra on the celestial sphere," JHEP **02** (2021), 176 doi:10.1007/JHEP02(2021)176 [arXiv:2008.04330 [hep-th]].
- [6] S. Banerjee, S. Ghosh and P. Paul, "(Chiral) Virasoro invariance of the tree-level MHV graviton scattering amplitudes," JHEP 09 (2022), 236 doi:10.1007/JHEP09(2022)236 [arXiv:2108.04262 [hep-th]].
- [7] N. Gupta, P. Paul and N. V. Suryanarayana, "An \widehat{sl}_2 Symmetry of $\mathbb{R}^{1,3}$ Gravity," [arXiv:2109.06857 [hep-th]].
- [8] A. M. Polyakov, "Gauge Transformations and Diffeomorphisms," Int. J. Mod. Phys. A 5 (1990), 833 doi:10.1142/S0217751X90000386
- [9] M. Bershadsky, "Conformal field theories via Hamiltonian reduction," Commun. Math. Phys. 139 (1991), 71-82 doi:10.1007/BF02102729
- [10] A. A. Belavin, A. M. Polyakov and A. B. Zamolodchikov, "Infinite Conformal Symmetry in Two-Dimensional Quantum Field Theory," Nucl. Phys. B 241 (1984), 333-380 doi:10.1016/0550-3213(84)90052-X
- [11] P. Di Francesco, P. Mathieu and D. Senechal, "Conformal Field Theory," Springer-Verlag, 1997, ISBN 978-0-387-94785-3, 978-1-4612-7475-9 doi:10.1007/978-1-4612-2256-9
- [12] R. Blumenhagen and E. Plauschinn, "Introduction to conformal field theory: with applications to String theory," Lect. Notes Phys. 779 (2009), 1-256 doi:10.1007/978-3-642-00450-6
- [13] K. Thielemans, "An Algorithmic approach to operator product expansions, W algebras and W strings," [arXiv:hep-th/9506159 [hep-th]].
- [14] R. Blumenhagen, M. Flohr, A. Kliem, W. Nahm, A. Recknagel and R. Varnhagen, "W algebras with two and three generators," Nucl. Phys. B 361 (1991), 255-289 doi:10.1016/0550-3213(91)90624-7

- [15] P. Bowcock, "Quasi-primary Fields and Associativity of Chiral Algebras," Nucl. Phys. B 356 (1991), 367-386 doi:10.1016/0550-3213(91)90314-N
- [16] A. M. Polyakov, "Quantum Gravity in Two-Dimensions," Mod. Phys. Lett. A 2 (1987), 893 doi:10.1142/S0217732387001130
- [17] S. G. Avery, R. R. Poojary and N. V. Suryanarayana, "An sl $(2,\mathbb{R})$ current algebra from AdS_3 gravity," JHEP **01** (2014), 144 doi:10.1007/JHEP01(2014)144 [arXiv:1304.4252 [hep-th]].
- [18] L. Apolo and M. Porrati, "Free boundary conditions and the AdS_3/CFT_2 correspondence," JHEP **03** (2014), 116 doi:10.1007/JHEP03(2014)116 [arXiv:1401.1197 [hep-th]].
- [19] R. R. Poojary and N. V. Suryanarayana, "Holographic chiral induced W-gravities," JHEP **10** (2015), 168 doi:10.1007/JHEP10(2015)168 [arXiv:1412.2510 [hep-th]].
- [20] V. G. Knizhnik, "Superconformal Algebras in Two-dimensions," Theor. Math. Phys. 66 (1986), 68-72 doi:10.1007/BF01028940
- [21] A. Campoleoni, S. Fredenhagen and S. Pfenninger, "Asymptotic W-symmetries in three-dimensional higher-spin gauge theories," JHEP 09 (2011), 113 doi:10.1007/JHEP09(2011)113 [arXiv:1107.0290 [hep-th]].
- [22] N. Gupta and N. V. Suryanarayana, "A chiral W-algebra from AdS_4 gravity," To appear.